REPORT UAG-32

WORLD DATA CENTER A
for
Solar-Terrestrial Physics

SYNOPTIC RADIO MAPS OF THE SUN AT 3.3 mm FOR THE YEARS 1967-1969



APRIL 1974

WORLD DATA CENTER A

National Academy of Sciences

2101 Constitution Avenue, N. W. Washington, D. C., U.S.A., 20418

World Data Center A consists of the Coordination Office and eight subcenters:

World Data Center A
Coordination Office
National Academy of Sciences
2101 Constitution Avenue, N.W.
Washington, D. C., U.S.A. 20418
Telephone (202) 389-6478

Solar and Interplanetary Phenomena, Ionospheric Phenomena, Flare-Associated Events, Geomagnetic Variations, Magnetospheric and Interplanetary Magnetic Phenomena, Aurora, Cosmic Rays, Airglow:
World Data Center A
for Solar-Terrestrial Physics
National Oceanic and Atmospheric Administration
Boulder, Colorado, U.S.A. 80302
Telephone (303) 499-1000 Ext. 6467

Geomagnetism, Seismology, Gravity (and Upper Mantle Project Archives):
World Data Center A:
Geomagnetism, Seismology and Gravity
Environmental Data Service, NOAA
Boulder, Colorado, U.S.A. 80302
Telephone (303) 499-1000 Ext. 6311

Glaciology:
World Data Center A:
Glaciology
U.S. Geological Survey
1305 Tacoma Avenue South
Tacoma, Washington, U.S.A. 98402
Telephone (206) 593-6502

Longitude and Latitude:
World Data Center A:
Longitude and Latitude
U. S. Naval Observatory
Washington, D. C., U.S.A. 20390
Telephone (202) 254-4547

Meteorology (and Nuclear Radiation):
World Data Center A:
Meteorology
National Climatic Center
Federal Building
Asheville, North Carolina, U.S.A. 28801
Telephone (704) 254-0683

Oceanography:
World Data Center A:
Oceanography
National Oceanic and Atmospheric Administration
Rockville, Maryland, U.S.A. 20852
Telephone (202) 426-9052

Rockets and Satellites:
World Data Center A:
Rockets and Satellites
Goddard Space Flight Center
Code 601
Greenbelt, Maryland, U.S.A. 20771
Telephone (301) 982-6695

Tsunami:
World Data Center A:
Tsunami
National Oceanic and Atmospheric Administration
2525 Korrea Road
Honolulu, Hawaii, U.S.A. 96822
Telephone (808) 948-8083

Notes:

(1) World Data Centers conduct international exchange of geophysical observations in accordance with the principles set forth by the International Council of Scientific Unions. WDC-A is established in the United States under the auspices of the National Academy of Sciences.
(2) Communications regarding data interchange matters in general and World Data Center A as a whole should be addressed to: World Data Center A, Coordination Office (see address above).
(3) Inquiries and communications concerning data in specific disciplines should be addressed to the appropriate subcenter listed above.

WORLD DATA CENTER A
for
Solar-Terrestrial Physics

REPORT UAG-32

SYNOPTIC RADIO MAPS OF THE SUN AT 3.3mm FOR THE YEARS 1967-1969

by

Earle B. Mayfield and Kennon P. White III

San Fernando Observatory

Space Physics Laboratory

and

Fred I. Shimabukuro

Electronics Research Laboratory

Laboratory Operations

The Aerospace Corporation

El Segundo, California 90245



Prepared by World Data Center A for
Solar-Terrestrial Physics, NOAA, Boulder, Colorado
and published by

U.S. DEPARTMENT OF COMMERCE
NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION

ENVIRONMENTAL DATA SERVICE
Asheville, North Carolina, USA 28801

APRIL 1974

SUBSCRIPTION PRICE: $9.00 a year; $2.50 additional for foreign mailing; single copy price varies.* Checks and money orders should be made payable to the Department of Commerce, NOAA. Remittance and correspondence regarding subscriptions should be sent to the National Climatic Center, Federal Building, Asheville, NC 28801, Attn: Publications.

*Price of this issue 35 cents

SYNOPTIC RADIO MAPS OF THE SUN AT 3.3 mm FOR THE YEARS 1967-1969

by

Earle B. Mayfield and Kennon P. White III
San Fernando Observatory
Space Physics Laboratory

and

Fred I. Shimabukuro
Electronics Research Laboratory

Laboratory Operations
The Aerospace Corporation
El Segundo, California 90245

The Aerospace Corporation millimeter radio telescope was completed in 1963 and has been used since then for various observations in planetary, solar and stellar research. It is located in El Segundo, California with coordinates 33° 54' 53" N and +7h 53m 30s (118° 22' 30" W) at a height of 38 m. Since 1966 radio maps of the whole solar disk have been obtained routinely at 90 GHz (3.3 mm wavelength). An effort has been made to make a map at least once a day, except when conditions of very dense clouds or rain prevail. Consequently, an extensive library of solar data exists which may be of considerable interest to others.

A difficulty in making these observations available is in developing an appropriate format for publication. The method adopted is based on our investigations of variations of the 3.3 mm emission and correlations with other solar phenomena which have established relationships with photospheric magnetic fields, faculae and centers of activity. Further, the regions of enhanced millimeter emission are well-defined and stable features which can be readily identified and followed during disk passage. Transient phenomena associated with flares which have been discussed by *Shimabukuro* [1968] are usually at the resolution limit of the antenna and are relatively infrequent. As a result, we consider synoptic maps as the most informative presentation and the most concise. For those who may require more detailed data or individual maps for specific dates, separate maps can be provided.

Although the instrument has been previously described by *Jacobs and King* [1965] and by *King et al.* [1966], a brief description is appropriate. The antenna is a Cassegrain design of 15-foot diameter (4.57 meters) in a polar mount. Although the surface has been finished for operation at 400 GHz (1 mm), it is usually operated at 90 GHz (3.3 mm). At this wavelength the beamwidth at the half-power points is about 2.8 arc-min. Pointing accuracy and stability for the instrument is about 20 arc-sec. The antenna is controlled by an on-line digital computer through a closed loop servo which permits a variety of tracking modes including a mapping mode which is used for solar studies. The radiometer is of the Dicke type in which switching is done at 465 Hz between the on-axis beam and a wide beam sky temperature reference horn. The first stage of the radiometer is a single-ended gallium arsenide crystal mixer which provides for frequency conversion to a 3 GHz intermediate amplifier. The system sensitivity for a signal-to-noise ratio of one at the output is about 0.6°K for 1 second integration time.

For the results reported here, the antenna was programmed to obtain a matrix of emission temperatures centered on the solar disk. A 19x19 matrix was obtained by rastering the antenna between adjacent points. Four seconds settle time was allowed between readings and one second integration for temperature measurement. These individual values were normalized to an undisturbed region near the center of the disk to eliminate the requirement for absolute calibration of the data. This reference temperature used for normalization was selected by analysis of observations previously obtained of the mm radio emission together with the Fraunhofer Institut maps to select a quiet region free of plage or other disturbed emission. One limitation of the maps is that beyond 0.7 $R_\odot$ (normalized solar radius) the interpretation becomes difficult because of the finite beam width of the antenna at the solar limb. As a result, the isotherms are restricted to regions less than 0.7 $R_\odot$. Occasional pointing errors also occur in centering the temperature matrix on the disk. These errors are typically greater than 20 arc-sec but less than 1 arc-min. Occasional variations in cloud cover can also account for slight differences in the temperature matrix.

A typical map is shown in Figure 1 for 30 June 1971. This has been overlaid on an Hα picture and isotherms have been constructed on the temperature matrix at levels of 2% enhancement. The isotherms in Figure 1 indicate the presence of two regions of significant enhancement which can be identified with the Hα plage in centers of activity associated with magnetic fields in sunspot groups. The maximum enhancements of these mm regions usually occur near the neutral lines of primary bipolar magnetic regions and show daily variations associated with the changes in the magnetic fields. Previous analysis by *Mayfield et al.* [1970] has discussed these relationships of mm enhancement and magnetic fields in detail.



Fig. 1. The 3.3 mm solar radio temperature contour map made on 30 June 1971 superimposed on a concurrent Hα photograph. The contours are labeled in percent enhancement relative to the temperature of the undisturbed region denoted by the hatched contours.

To summarize the results of three years' observation from 1967 through 1969, a synoptic presentation was used based on the Zürich heliographic maps of the sun by *Waldmeier* [1968] and the magnetic field presentation of Mt. Wilson by Howard. These maps are arranged to present the 360° of the solar disk showing the mm isotherms on a day near central meridian passage. A typical synoptic map is shown in Figure 2 for Carrington rotation 1521. The Carrington longitude is given at the top of the figure and the date of central meridian passage is located by the filled circle and day number at the bottom of the figure. The vertical scale is the solar longitude between ±50°. Maps used in the synoptic presentation have been normalized to the quiet regions indicated by the crosses in the figure; times at which the maps were obtained are indicated with tic marks at the bottom of the figure. For presentation in this synoptic format, only isotherms from 4% enhancement up have been used. This is done to identify enhanced regions and to show the close correlation between the mm enhancement and white light faculae identified in the Zürich maps.



Fig. 2. Synoptic 3.3 mm radio temperature contour map for Carrington rotation 1521 superimposed on the corresponding heliographic map of the photosphere by Waldmeier. Carrington longitude appears at the top of the Figure and date of CMP is given at the bottom. Vertical scale is solar longitude. The contours are at levels of 4, 6, 8% etc., enhancement above the quiet regions indicated by crosses. Peak enhancements are shown by the filled circles. Note the close correspondence between the 4% contour levels and facular areas and between peak enhancements and sunspots.

Plotted in Figure 2 also is an overlay of sunspots and faculae taken from the Zürich map for rotation 1521. This shows a close correlation between the 4% isotherm contour and areas of faculae in sunspot regions. An interpretation of mm emission and the relation with white light faculae has been given by *Mayfield et al.* [1973] and will not be discussed here. Synoptic maps for the years 1967 through 1969 are given in the appendix. These have been plotted so that they can be readily compared with the Zürich photoheliographic maps and with the synoptic magnetic field maps published in the *Quarterly Bulletin of Solar Activity*.

The question of the longevity of active regions can be investigated by inspection of successive synoptic maps to note the recurrence of active regions. In this respect it is important to point out the sensitivity and precision of the data contained in the synoptic maps, because these parameters influence the visibility of the enhancements. The lowest contour level plotted corresponds to a 4% enhancement, so chosen because at this level individual active regions are usually discernable, whereas a 3% contour usually will run nearly the whole length of the map providing very little information. Additionally, the repeatability of measurements made on different daily maps either on the same day or different days can be as large as ±0.5%, though the average is probably closer to ±0.3%, with some outstanding examples showing variations of no more than ±0.2% over a five- or six-day period. To maintain a precision of about 10% (e.g. 4 ±0.4%), we felt it was safest to display the 4% contour as the lowest level.

Therefore, the longevity of an active region is defined as the time from when it first exceeded 4% enhancement to when it first disappeared below the 4% level. Based on this criterion, some regions are found to last less than a full solar rotation, while others can persist and be followed over five or six rotations. As will be shown below, this result is entirely consistent with the result that the 4% contour level coincides in shape and location with the facular areas, which have lifetimes that can be as short as a few days or as long as a few solar rotations.

Another feature of the synoptic maps whose longevity can be gauged is the quiet regions, as indicated by the clustering of crosses. The nature of these regions is still in question, but certain examples of them can be followed over three or four rotations.

Daily observations of sunspots and photospheric faculae have been presented in synoptic form by *Waldmeier* [1968] in *Heliographic Maps of the Photosphere*. An example of these maps in Figure 2 shows each sunspot group at the time of its maximum evolution. Umbrae are given as black dots, penumbrae are shown by their outlines, and facular regions are shown by dashed lines on the maps.

Comparison of the radio synoptic map with the heliographic map in Figure 2 shows how well the 4% enhancement contours correspond to the extensions of the facular regions. Differences in shape and extent can be attributed to a number of causes, which include: (1) slow, evolutionary changes in the active regions, since the photospheric maps pertain to the maximum development of the regions, while the millimeter maps reflect the appearance closest to central meridian passage; (2) rapid changes in the active regions, as caused by flares, whose residual effects can influence the radio contours (although, such has been allowed for and corrected when possible); (3) selection of an invalid normalization point for a particular daily radio map (e.g., a dark absorption filament), which would cause an apparent "growth" or "contraction" of the enhancement levels; and (4) the lack of a physical connection between the two phenomena.

We hope these results will be of benefit to other investigations of solar phenomena. Subsequent data will be published as they are reduced and analyzed. Individual maps of particular times, if available, may be obtained for the years 1970 to the present by communicating with the authors. Comments or recommendations for improving the format of the presentation are welcomed.

Acknowledgements

This research was supported in part by NASA Ames Research Center under Contract NAS2-7868 and in part by company funds of The Aerospace Corporation.

REFERENCES

JACOBS, E. and H. E. KING 1965 *IEEE International Convention Record*, Part 5.

KING, H. E., E. JACOBS and J. M. STACEY 1966 *IEEE Trans. on Antennas and Propagation, AP-14*, 82.

MAYFIELD, E. B., J. HIGMAN and C. SAMSON 1970 *Solar Phys., 13*, 372.

MAYFIELD, E. B., K. P. WHITE III and F. I. SHIMABUKURO 1973 IAU Symposium 57 on Coronal Disturbances, Surfers Paradise, Australia.

SHIMABUKURO, F. I. 1968 *Solar Phys.*, *5*, 498.

WALDMEIER, M. 1968 *Publikationen der Eidgenossischen Sternwarte, Zürich 13*, Heft 2.

APPENDIX

Synoptic Radio Maps of the Sun at 3.3 mm Wavelength for the Years 1967-1969






1967 - ROTATION 1518
FEB
MAR
E 0°
0°W


1967 - ROTATION 1519
NO DATA OBTAINED
MAR
APR
E 0°
0°W


1967 - ROTATION 1520
NO DATA OBTAINED
NO DATA OBTAINED
MAY
MAY
APR


1967 - ROTATION 1521
JUN
JUN
MAY

1967 – ROTATION 1522



1967 – ROTATION 1523





No Data Available for Rotation 1525


1967 - ROTATION 1526
NO DATA OBTAINED
OCT
SEP


1967 - ROTATION 1527
NOV
NOV
OCT


1967 - ROTATION 1528
NOV
DEC
DEC
E 0°
0°W


1968 - ROTATION 1529
DEC
JAN
1968
JAN
E 0°
0°W


1968 - ROTATION 1530
0° 20 40 60 80 100 120 140 160 180 200 220 240 260 280 300 320 340 360°
40 30 20 10 E 0° −10 −20 −30 −40
0°W
12 11 10 9 8 7 6 5 4 3 2 1 31 30 29 28 27 26 25 24 23 22 21 20 19 18 17 16
FEB
FEB
JAN


1968 - ROTATION 1531
0° 20 40 60 80 100 120 140 160 180 200 220 240 260 280 300 320 340 360°
40 30 20 10 E 0° −10 −20 −30 −40
0°W
10 9 8 7 6 5 4 3 2 1 29 28 27 26 25 24 23 22 21 20 19 18 17 16 15 14 13 12
MAR
MAR
FEB


1968 - ROTATION 1532
0° 20 40 60 80 100 120 140 160 180 200 220 240 260 280 300 320 340 360°
40 30 20 10 E 0° −10 −20 −30 −40
0°W
7 6 5 4 3 2 1 31 30 29 28 27 26 25 24 23 22 21 20 19 18 17 16 15 14 13 12 11
APR APR MAR
6.1 4 4.3 4 5.2 4 6.3 4 6 6.4 4 6.4 4.6 4 6.5 6 4 5.9 4.9 4 4
4 6 6.9 5.4 4.8 4.5 4 4.3 4 8.1 6 4.0 4 5.4


1968 - ROTATION 1533
0° 20 40 60 80 100 120 140 160 180 200 220 240 260 280 300 320 340 360°
40 30 20 10 E 0° −10 −20 −30 −40
0°W
4 3 2 1 30 29 28 27 26 25 24 23 22 21 20 19 18 17 16 15 14 13 12 11 10 9 8 7
MAY MAY APR
4.7 4 4 4.2 6.0 4 4.8 4.4 4 4.8 4 5.0 4 5.3 4 5.5 4
4 4.9 4.1 4 4.6 4.4 4 4 4.7


1968 - ROTATION 1534
0° 20 40 60 80 100 120 140 160 180 200 220 240 260 280 300 320 340 360°
40 30 20 10 E 0° −10 −20 −30 −40
0°W
31 30 29 28 27 26 25 24 23 22 21 20 19 18 17 16 15 14 13 12 11 10 9 8 7 6 5 4
MAY
MAY


1968 - ROTATION 1535
0° 20 40 60 80 100 120 140 160 180 200 220 240 260 280 300 320 340 360°
40 30 20 10 E 0° −10 −20 −30 −40
0°W
27 26 25 24 23 22 21 20 19 18 17 16 15 14 13 12 11 10 9 8 7 6 5 4 3 2 1
JUN
JUN

1968 ROTATION 1536



1968 ROTATION 1537




1968 ROTATION 1538
0° 20 40 60 80 100 120 140 160 180 200 220 240 260 280 300 320 340 360°
40 30 20 10 E 0° -10 -20 -30 -40
0°W -10 -20 -30 -40
4 7.1 6
4 7.5 6
4 4.6 4
5.9 4
4 6 7.2
5.4 4
4.1 4
4 4.8 4.8
4 4.4
5.8 4
4 4.6
4 4.3
17 16 15 14 13 12 11 10 9 8 7 6 5 4 3 2 1 31 30 29 28 27 26 25 24 23 22 21
SEP
SEP
AUG


1968 ROTATION 1539
0° 20 40 60 80 100 120 140 160 180 200 220 240 260 280 300 320 340 360°
40 30 20 10 E 0° -10 -20 -30 -40
0°W -10 -20 -30 -40
4.3 4
5.9
4 5.2
5.5 4
4 4.2
4.2
4.2
4.9 4
4.7 4 5.1
5.8 4
4 4.6
4 5.3
14 13 12 11 10 9 8 7 6 5 4 3 2 1 30 29 28 27 26 25 24 23 22 21 20 19 18 17
OCT
OCT
SEP


1968 ROTATION 1540
OCT
NOV
NOV
E 0°
0°W


1968 ROTATION 1541
NOV
DEC
DEC
E 0°
0°W


1968 ROTATION 1542
0°
360°
0°W
E 0°
JAN 1969 JAN
DEC


1969 ROTATION 1543
0°
360°
0°W
E 0°
JAN


1969 ROTATION 1544
0° 20 40 60 80 100 120 140 160 180 200 220 240 260 280 300 320 340 360°
40 30 20 10 E 0° −10 −20 −30 −40
0°W
28 27 26 25 24 23 22 21 20 19 18 17 16 15 14 13 12 11 10 9 8 7 6 5 4 3 2 1
FEB
FEB
10.3
6.4
5.5
4.9
10.5
8.2
6.2
5.8
4.5
4.6
6.1


1969 ROTATION 1545
0° 20 40 60 80 100 120 140 160 180 200 220 240 260 280 300 320 340 360°
40 30 20 10 E 0° −10 −20 −30 −40
0°W
27 26 25 24 23 22 21 20 19 18 17 16 15 14 13 12 11 10 9 8 7 6 5 4 3 2 1
MAR
MAR
10.1
9.7
4.0
9.1
8.2
6.2
6.1
5.7
4.3
8.3
10.3


1969 ROTATION 1546
0° 20 40 60 80 100 120 140 160 180 200 220 240 260 280 300 320 340 360°
40 30 20 10 E 0° −10 −20 −30 −40
0°W
9.3 9.7 7.7 6.2 6.2 6.7 8.6 8.6 4.6 6.4 7.1 5.8 7.7 4.5 4.5 4.7 5.7 5.8
23 22 21 20 19 18 17 16 15 14 13 12 11 10 9 8 7 6 5 4 3 2 1 31 30 29 28
APR APR MAR


1969 ROTATION 1547
0° 20 40 60 80 100 120 140 160 180 200 220 240 260 280 300 320 340 360°
40 30 20 10 E 0° −10 −20 −30 −40
0°W
10.8 7.3 10 8.7 9.9 5.5 5.9 5.2 5.2 5.3 4.7
20 19 18 17 16 15 14 13 12 11 10 9 8 7 6 5 4 3 2 1 30 29 28 27 26 25 24
MAY MAY APR


1969 ROTATION 1548
0° 20 40 60 80 100 120 140 160 180 200 220 240 260 280 300 320 340 360°
40 30 20 10 E 0° −10 −20 −30 −40
0°W
11.1
10
5.8
5.8
5.8
8.4
5.6
6.7
6.6
17 16 15 14 13 12 11 10 9 8 7 6 5 4 3 2 1 31 30 29 28 27 26 25 24 23 22 21
JUN
JUN
MAY


1969 ROTATION 1549
0° 20 40 60 80 100 120 140 160 180 200 220 240 260 280 300 320 340 360°
40 30 20 10 E 0° −10 −20 −30 −40
0°W
5.1
5.0
8.8
10.2
8.3
6.3
5.1
6.5
6.6
8.0
7.3
7.3
14 13 12 11 10 9 8 7 6 5 4 3 2 1 30 29 28 27 26 25 24 23 22 21 20 19 18
JUL
JUL
JUN


1969 ROTATION 1550
0° 20 40 60 80 100 120 140 160 180 200 220 240 260 280 300 320 340 360°
40 30 20 10 E 0° −10 −20 −30 −40
0°W
10 9 8 7 6 5 4 3 2 1 31 30 29 28 27 26 25 24 23 22 21 20 19 18 17 16 15
AUG
AUG
JUL


1969 ROTATION 1551
0° 20 40 60 80 100 120 140 160 180 200 220 240 260 280 300 320 340 360°
40 30 20 10 E 0° −10 −20 −30 −40
0°W
6 5 4 3 2 1 31 30 29 28 27 26 25 24 23 22 21 20 19 18 17 16 15 14 13 12 11
SEP
SEP
AUG


1969 ROTATION 1552
SEP
OCT
E 0°
0°W


1969 ROTATION 1553
OCT
E 0°
0°W

1969 ROTATION 1554



1969 ROTATION 1555




1969 ROTATION 1556
JAN
DEC
1970
E
0°W

UAG Series of Reports

Prepared by World Data Center A for Solar-Terrestrial Physics, NOAA, Boulder, Colorado, U.S.A.

These reports are for sale through the National Climatic Center, Federal Building, Asheville, NC 28801, Attn: Publications. Subscription price: $9.00 a year; $2.50 additional for foreign mailing; single copy price varies. These reports are issued on an irregular basis with 6 to 12 reports being issued each year. Therefore, in some years the single copy rate will be less than the subscription price, and in some years the single copy rate will be more than the subscription price. Make check or money order payable to: Department of Commerce, NOAA.

Some issues are now out of print and are available only on microfiche as indicated. Requests for microfiche should be sent to World Data Center A for Solar-Terrestrial Physics, NOAA, Boulder, Co 80302, with check or money order made payable to Department of Commerce, NOAA.

UAG-1 "IQSY Night Airglow Data", by L. L. Smith, F. E. Roach and J. M. McKennan of Aeronomy Laboratory, ESSA Research Laboratories, July 1968, 305 pages, price $1.75.

UAG-2 "A Reevaluation of Solar Flares, 1964-1966", by Helen W. Dodson and E. Ruth Hedeman of McMath-Hulbert Observatory, The University of Michigan, August 1968, 28 pages, microfiche only, price 45 cents.

UAG-3 "Observations of Jupiter's Sporadic Radio Emission in the Range 7.6-41 MHz, 6 July 1966 through 8 September 1968", by James W. Warwick and George A. Dulk, Department of Astro-Geophysics, University of Colorado, October 1968, 35 pages, microfiche only, price 45 cents.

UAG-4 "Abbreviated Calendar Record 1966-1967", by J. Virginia Lincoln, Hope I. Leighton and Dorothy K. Kropp of Aeronomy and Space Data Center, Space Disturbances Laboratory, ESSA Research Laboratories, January 1969, 170 pages, price $1.25.

UAG-5 "Data on Solar Event of May 23, 1967 and its Geophysical Effects", compiled by J. Virginia Lincoln, World Data Center A, Upper Atmosphere Geophysics, ESSA, February 1969, 120 pages, microfiche only, price 90 cents.

UAG-6 "International Geophysical Calendars 1957-1969", by A. H. Shapley and J. Virginia Lincoln, ESSA Research Laboratories, March 1969, 25 pages, microfiche only, price 45 cents.

UAG-7 "Observations of the Solar Electron Corona: February 1964-January 1968", by Richard T. Hansen, High Altitude Observatory, Boulder, Colorado and Kamuela, Hawaii, October 1969, 12 pages, price 15 cents.

UAG-8 "Data on Solar-Geophysical Activity October 24-November 6, 1968", Parts 1 and 2, compiled by J. Virginia Lincoln, World Data Center A, Upper Atmosphere Geophysics, ESSA, March 1970, 312 pages, price (includes Parts 1 and 2) $1.75.

UAG-9 "Data on Cosmic Ray Event of November 18, 1968 and Associated Phenomena", compiled by J. Virginia Lincoln, World Data Center A, Upper Atmosphere Geophysics, ESSA, April 1970, 109 pages, price 55 cents.

UAG-10 "Atlas of Ionograms", edited by A. H. Shapley, ESSA Research Laboratories, May 1970, 243 pages, price $1.50.

UAG-11 "Catalogue of Data on Solar-Terrestrial Physics", June 1970. (now obsolete).

UAG-12 "Solar-Geophysical Activity Associated with the Major Geomagnetic Storm of March 8, 1970", Parts 1, 2 and 3, compiled by J. Virginia Lincoln and Dale B. Bucknam, World Data Center A, Upper Atmosphere Geophysics, NOAA, April 1971, 466 pages, price (includes Parts 1-3) $3.00.

UAG-13 "Data on the Solar Proton Event of November 2, 1969 through the Geomagnetic Storm of November 8-10, 1969", compiled by Dale B. Bucknam and J. Virginia Lincoln, World Data Center A, Upper Atmosphere Geophysics, NOAA, May 1971, 76 pages, microfiche only, price 90 cents.

UAG-14 "An Experimental, Comprehensive Flare Index and Its Derivation for 'Major' Flares, 1955-1969", compiled by Helen W. Dodson and E. Ruth Hedeman, McMath-Hulbert Observatory, The University of Michigan, July 1971, 25 pages, price 30 cents.

UAG-15 "Catalogue of Data on Solar-Terrestrial Physics", July 1971. (now obsolete).

UAG-16 "Temporal Development of the Geographical Distribution of Auroral Absorption for 30 Substorm Events in each of IQSY (1964-65) and IASY (1969)", by F. T. Berkey, V. M. Driatskiy, K. Henriksen, D. H. Jelly, T. I. Shchuka, A. Theander and J. Yliniemi, September 1971, 131 pages, price 70 cents.

UAG-17 "Ionospheric Drift Velocity Measurements at Jicamarca, Peru (July 1967-March 1970)", by Ben B. Balsley, Aeronomy Laboratory, National Oceanic and Atmospheric Administration, Boulder, Colorado, and Ronald F. Woodman, Jicamarca Radar Observatory, Instituto Geofisico del Perú, Lima, Peru, October 1971, 45 pages, microfiche only, price 45 cents.

UAG-18 "A Study of Polar Cap and Auroral Zone Magnetic Variations", by K. Kawasaki and S. -I. Akasofu, Geophysical Institute, University of Alaska, June 1972, 21 pages, price 20 cents.

UAG-19 "Reevaluation of Solar Flares 1967", by Helen W. Dodson and E. Ruth Hedeman, McMath-Hulbert Observatory, The University of Michigan, and Marta Rovira de Miceli, San Miguel Observatory, Argentina, June 1972, 15 pages, price 15 cents.

UAG-20 "Catalogue of Data on Solar-Terrestrial Physics", October 1972. (now obsolete).

UAG-21 "Preliminary Compilation of Data for Retrospective World Interval July 26 - August 14, 1972", compiled by J. Virginia Lincoln and Hope I. Leighton, World Data Center A for Solar-Terrestrial Physics, November 1972, 128 pages, price 70 cents.

UAG-22 "Auroral Electrojet Magnetic Activity indices (AE) for 1970", by Joe Haskell Allen, National Geophysical and Solar-Terrestrial Data Center, Environmental Data Service, November 1972, 146 pages, price 75 cents.

UAG-23 "U.R.S.I. Handbook of Ionogram Interpretation and Reduction", Second Edition, November 1972, edited by W. R. Piggott, Radio and Space Research Station, Slough, U.K., and K. Rawer, Arbeitsgruppe für Physikalische Weltraumforschung, Freiburg, G.F.R., November 1972, 324 pages, price $1.75.

UAG-24 "Data on Solar-Geophysical Activity Associated with the Major Ground Level Cosmic Ray Events of 24 January and 1 September 1971", Parts 1 and 2, compiled by Helen E. Coffey and J. Virginia Lincoln, World Data Center A for Solar-Terrestrial Physics, December 1972, 462 pages, price (includes Parts 1 and 2) $2.00.

UAG-25 "Observations of Jupiter's Sporadic Radio Emission in the Range 7.6-41 MHz, 9 September 1968 through 9 December 1971", by James W. Warwick, George A. Dulk and David G. Swann, Department of Astro-Geophysics, University of Colorado, February 1973, 35 pages, price 35 cents.

UAG-26 "Data Compilation for the Magnetospherically quiet Periods February 19-23 and November 29 - December 3, 1970", compiled by Helen E. Coffey and J. Virginia Lincoln, World Data Center A for Solar-Terrestrial Physics, May 1973, 129 pages, price 70 cents.

UAG-27 "High Speed Streams in the Solar Wind", by D. S. Intriligator, University of Southern California, Department of Physics, Los Angeles, California, 90007, June 1973, 16 pages, price 15 cents.

UAG-28 "Collected Data Reports on August 1972 Solar-Terrestrial Events", Parts 1, 2 and 3, edited by Helen E. Coffey, World Data Center A for Solar-Terrestrial Physics, July 1973, 932 pages, price (includes Parts 1-3) $4.50.

UAG-29 "Auroral Electrojet Magnetic Activity Indices AE (11) for 1968", by Joe Haskell Allen, Carl C. Abston and Leslie D. Morris, National Geophysical and Solar-Terrestrial Data Center, Environmental Data Service, October 1973, 148 pages, price 75 cents.

UAG-30 "Catalog of Data on Solar-Terrestrial Physics", prepared by Environmental Data Service, NOAA, Boulder, Colorado, October 1973, 317 pages, price $1.75.

UAG-31 "Auroral Electrojet Magnetic Activity Indices AE (11) for 1969", by Joe Haskell Allen, Carl C. Abston and Leslie D. Morris, National Geophysical and Solar-Terrestrial Data Center, Environmental Data Service, February 1974, 142 pages.

USCOMM-NOAA-ASHEVILLE,NC--4-74--1325