U.S. DEPARTMENT OF COMMERCE
NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION
NATIONAL WEATHER SERVICE
NATIONAL METEOROLOGICAL CENTER

OFFICE NOTE 6

A TWO-LEVEL MODEL WITH EFFECTS OF VERTICAL VARIETY ADUCTION, SINGLES TO RAIN, AND VARIABLE STATIC STABILITY

Phillip Duncan Thompson
Joint Numerical Weather Prediction Unit

10 January 1957

This is an unreviewed manuscript, primarily intended for informal exchange of information among NMC staff members

p=400 or 4 somb

I $\omega_0 = \frac{10^3 g m^2}{4d^2}\left[\frac{l}{f}J(\hat{\psi}, P_g) - \frac{1}{f}\frac{P_g - 600}{p}J(h, P_g)\right]$ $\quad 2^{24}\omega_0 = 2^{20}\hat{\omega}_0 \quad \hat{\omega}_0 = 2^{-4}\omega_0$

II $\nabla^2\hat{\chi} = -\frac{\hat{\omega}_0}{10^3 p}\frac{d^2}{m^2}2^3 \cdot 2^{-30} \quad \chi = 2^{30}\hat{\chi}$

III $V_x = \frac{g l m}{2fd}\frac{\partial \hat{\psi}}{\partial y} + \frac{2^{30} m}{2d}\frac{\partial \hat{\chi}}{\partial x} \quad V_y = \frac{g l m^2}{2fd}\frac{\partial \psi}{\partial x} + \frac{2^{30} m}{2d}\frac{\partial \hat{\chi}}{\partial y} \quad \psi = \frac{g l}{f}\hat{\psi}$

IV $\frac{\partial \sigma^2}{\partial t} = -C\left[\frac{g l m^2}{4fd^2}J(\hat{\psi}, \sigma^2) + \frac{2^{30} m^2}{d^2}A(\hat{\chi}, \sigma^2)\right] - C_2\frac{6\hat{\omega}_{00}^2 \cdot 2^4}{10^3 \cdot 2p} \quad \chi = 2^{30}\hat{\chi}$

$\omega_0 = 2^4\hat{\omega}_0$

$P_{mb} = 10^3 P_{cgs}$

V $\eta = \frac{g l m^2}{4fd^2}\nabla^2\hat{\psi} + f$

VI $\frac{m^2}{d^2}\nabla^2\left(\frac{\partial h}{\partial t}\right) - \frac{2f\eta}{\sigma^2}\frac{\partial h}{\partial t} = \frac{2f\eta}{\sigma^2}\left[\frac{g l m^2}{4fd^2}J(\hat{\psi}, h) + \frac{2^{30} m^2}{4d^2}A(\hat{\chi}, h)\right.$

$\left. - \left[\frac{g l m^2}{4fd^2}J(\hat{\psi}, \frac{m^2}{d^2}\nabla^2 h) + \frac{2^{30} m^2}{4d^2}A(\hat{\chi}, \frac{m^2}{d^2}\nabla^2 h)\right] - \frac{m^2}{4d^2}J(h, \eta) - \frac{2^4 f \eta \hat{\omega}_0}{10^3 g p}\right.$

VII $\omega = \frac{10^3 g p}{\sigma^2}\left[\frac{\partial h}{\partial t} + \frac{g l m^2}{4^3 d}J(\hat{\psi}, h) + \frac{2^{30} m^2}{4d^2}A(\hat{\chi}, h)\right]$

VIII $\frac{g l m^2}{fd^2}\nabla^2\frac{\partial \psi}{\partial t} = -B\left[\frac{g l m^2}{4fd^2}J(\hat{\psi}, \eta) + \frac{2^{30} m^2}{4d^2}A(\hat{\chi}, \eta)\right] + \frac{g w m^2}{f p d^2 10^3}\nabla^2 h + \frac{2^4 \eta \hat{\omega}_0}{10^3 2p}$

1. Derive Z_{600} from Z_{400} and Z_{850} by means of

expression $Z_{600} = 0.53792\, Z_{850} + 0.46208\, Z_{400} + C$

Provide for C having a field. When Z is in feet C = 270 ft.
[80% of standard atmosphere static stability has been used]
Print out for inspection

2. Compute the two components U_o and V_o of

$$V_o = 1K \times \nabla\psi - \frac{P_g - 600}{P} \frac{g}{f} 1K \times \nabla h$$

V_o ?

where ψ is the 600 mb stream function, P_g the pressure at the ground,
$h = Z_{400} - Z_{850}$ and $P = 450$ mb
Print out for inspection the two components U_o & V_o
in appropriate units

3. $P = 450$ in eqn I
4. $P = 400$ mb in the remaining eqn
5. $U_o = -g\rho W_o = 1.3 W_o$ where W_o is in cm/sec
   1 mb = 10^3 c g s

6. In eqn 6 replace $V \cdot \nabla(\nabla^2 h)$ by $\frac{2f}{g} V \cdot \nabla\left(\frac{g}{2f}\nabla^2 h\right)$

7. In eqn 4 $C\frac{\omega_0 \sigma^2}{2P}$ may be replaced by $C\frac{\omega\sigma^2}{2P}$

8. The contribution of the various terms (fluxes) in the prognostic equations will for each time step be summed on tape for later use (separate prognosis)

9. Make provisions for running the prognostic equations without 1) the ω_0 terms 2) the $\nabla^2 h$ terms

10 program eqn IV in the form

$$\frac{\partial}{\partial t}(\sigma^2) + C_1 V \cdot \nabla(\sigma^2) + C_2 \frac{\omega_0 \sigma^2}{2P} = 0$$

where C_1 & C_2 are two constants, yet not determined

STD 850 = 4,780 ft

STD 400 = 23,564 ft

Use Note ...

... Time - Level Main ... Tide ... Datum

Vitality Curves, ... Tau, and

Various State Stability ...

... Ralph P. Thompson

... Cdr, USN

... Post Graduate Work ... Unit

10 June 1957

In this model, the ω-profile is assumed to have the simple pattern shown below. We suppose that ω remains



at 200 mb (see T(19, #11), and that ω vanishes at 1000 mb and at 400 mb. However, although ω_1 is not zero, as in the ω-profile is not symmetrical around 600 mb. On statistical grounds like those given above, we assume that symmetrical profiles are more likely. That is,

$$\frac{\omega_2 - \omega_1}{p_2 - p_1} = \frac{\omega_0}{2p}$$

Applying the vorticity equation at the 400 and 800 mb surfaces, and replacing $\partial\eta/\partial p$ and $\partial\omega/\partial p$ by finite differences,

$$\frac{\partial \eta_1}{\partial t} + V_1 \cdot \nabla \eta_1 + \omega_1 \frac{(\eta_2 - \eta_1)}{p} - \eta_1 \frac{\omega_0}{p} = 0$$

$$\frac{\partial \eta_2}{\partial t} + V_2 \cdot \nabla \eta_2 + \omega_2 \frac{(\eta_2 - \eta_1)}{p} - \eta_2 \frac{(\omega_0 - \omega_1)}{p} = 0$$

where the subscripts 1 and 2 refer to conditions at the "outer" and "inner" surfaces, respectively. Subtracting the second of these equations from the first, and rearranging,

$$-\frac{\partial}{\partial t}\left(\frac{\eta_1-\eta_2}{2}\right) + w^*\nabla\eta' + w'\nabla\eta^* + \left(\frac{\eta_2-\eta_1}{2}\right)\frac{\omega_1-\omega_2}{\rho}$$

$$-\frac{\omega}{2\rho}(\eta_1-\eta_2) + \frac{\eta_2\omega_0}{2\rho} = 0$$

in which

$$\eta' = \frac{\eta_1-\eta_2}{2} \qquad \eta^* = \frac{\eta_1+\eta_2}{2}$$

Also, since $\omega_2-\omega_1 = \omega_0/2$,

$$\frac{\partial\eta'}{\partial t} + w^*\cdot\nabla\eta' + w'\cdot\nabla\eta^* - \frac{\eta^*\omega}{\rho} + \frac{\eta'\omega_2}{2\rho} = 0 \qquad (1)$$

We must introduce the quasi-static approximation in evaluating η' and w' in the equation above. By definition

$$\eta' = \frac{\eta_1-\eta_2}{2} = \frac{1}{2g}\left(\nabla\chi_1^2 - \nabla\chi_2^2\right) = \frac{1}{2g}\nabla^2\psi$$

$$w' = \frac{w_1-w_2}{2} = \frac{1}{2g}\,ikR\left(\nabla\chi_1 - \nabla\chi_2\right) = \frac{1}{2g}\,ik\times\nabla\psi$$

where ψ is the stream function for the 400 and 600 mb surfaces. Thus, substituting in Eq. (1),

$$\frac{\partial}{\partial t}\nabla^2\psi + \vec{V}\cdot\nabla(\nabla^2\psi) + J(\psi,\eta^*) = \frac{2f\eta^*\omega}{g\delta p} + \frac{f\eta^*\omega_0}{g\delta p} = 0 \quad (2)$$

Now, the parcel equation may be written as

$$\frac{\partial}{\partial t}\left(\frac{1}{\rho}\right) + \vec{V}\cdot\nabla\left(\frac{1}{\rho}\right) + \left(\frac{1}{\rho\theta}\frac{\partial\theta}{\partial p}\right)\omega = 0$$

where the quantities apply at 500 mb. But, from the hydrostatic equation

$$\frac{1}{\rho} = -\frac{g(\beta_1 - \beta_2)}{\delta p} = \frac{g\psi}{\delta p}$$

Thus, substituting in the equation above,

$$\frac{\partial\psi}{\partial t} + \vec{V}\cdot\nabla\psi + \frac{\delta p}{\rho\theta}\frac{\partial\theta}{\partial p}\cdot\frac{\omega}{g} = 0$$

$$\omega = \frac{\omega}{g\delta p} = \frac{1}{\sigma g}\left[\frac{\partial\psi}{\partial t} + \vec{V}\cdot\nabla\psi\right] \quad (3)$$

where $\sigma = -\frac{\rho^2}{\delta p}\frac{\partial\theta}{\partial p} = \frac{p^2}{R^2}\frac{1}{\theta^2}\frac{\partial\theta}{\partial p}$

Finally, introducing the result into Eq. (2), and dividing by W^* with W (the 600 mb wind), and η^* with η,

$$\frac{\partial}{\partial t}\nabla^2\psi - \frac{2f_0}{\sigma_2^*}\frac{\partial h_*}{\partial t} = \frac{2f_0}{\sigma_2^*}\left[-V\cdot\nabla h_* - W\cdot\nabla(\sigma_1^* h_*)\right.$$

$$\left. - J(h,\eta) - \frac{f\eta\,\omega_0}{gp}\right] \qquad (4)$$

This is one of the two basic prognostic equations.

Applying the vorticity equation at 600 mb,

$$\frac{\partial \eta}{\partial t} + V\cdot\nabla\eta + \omega\left(\frac{\eta_2-\eta_1}{p}\right) - \frac{\eta\omega_0}{2p} = 0$$

Thus, computing $\eta_2-\eta_1$ from the geostrophic wind,

$$\frac{\partial}{\partial t}\nabla^2\psi + V\cdot\nabla(\nabla^2\psi+f) - \frac{2\omega}{fp}\nabla^2 h_* - \frac{\eta\omega_0}{2p} = 0 \qquad (5)$$

This is the other basic prognostic equation.

$$\omega_0 = J(\psi, p_0) - \left(\frac{p_0 - 500}{\rho}\right)\frac{f}{\rho} J(\psi, p_0) \qquad (6)$$

In this first step we neglect effect due to the variation of only the stretching effect of convergence in boundary flow, we put

$$V = k \times \nabla\psi + \nabla\chi \qquad (7)$$

$$\nabla \cdot V = \nabla^2 \chi = -\frac{\omega_0}{2\rho}$$

The last quantity required to complete the system is τ^2, which is proportional to $\frac{\partial \theta}{\partial p}$. We have, in θ coordinates:

$$\frac{\partial}{\partial t}\left(\frac{\partial \psi}{\partial \theta}\right) = \frac{\partial}{\partial \theta}\left(\frac{\partial \psi}{\partial t}\right) + V \cdot \nabla_\theta \left(\frac{\partial \psi}{\partial \theta}\right)$$

$$= \frac{\partial}{\partial \theta}\left(\frac{\partial \psi}{\partial t}\right) + \frac{\partial}{\partial \theta}(V \cdot \nabla_\theta \psi) - \frac{\partial V}{\partial \theta} \cdot \nabla_\theta \psi$$

But in quasi-geostrophic flow $\frac{\partial V}{\partial \theta}$ is very nearly perpendicular to $\nabla_\theta \psi$, so that

$$\frac{\partial}{\partial t}\left(\frac{\partial \psi}{\partial \theta}\right) \simeq \frac{\partial}{\partial \theta}\left(\frac{\partial \psi}{\partial t}\right)$$

$$\frac{\partial \theta}{\partial p}\,\frac{d}{dt}\left(\frac{\partial p}{\partial \theta}\right) = \frac{\partial \omega}{\partial p}$$

$$\frac{d}{dt}\,\omega\,\frac{\partial p}{\partial \theta} = \frac{\partial \omega}{\partial p}$$

$$-\frac{d}{dt}\,\omega\,\frac{\partial \theta}{\partial p} = \frac{\partial \omega}{\partial p}$$

$$\frac{d}{dt}\left(\frac{\partial \theta}{\partial p}\right) + \frac{\partial \theta}{\partial p}\,\frac{\partial \omega}{\partial p} = 0$$

Then, in p - coordinates,

$$\frac{\partial}{\partial t}\left(\frac{\partial \theta}{\partial p}\right) + V\cdot\nabla\frac{\partial \theta}{\partial p} + \frac{\partial}{\partial p}\left(\omega\,\frac{\partial \theta}{\partial p}\right) = 0$$

We now average each term of the equation with respect to p, integrating from 200 to 1000 mb. Differentiating term

$$\frac{\partial s}{\partial t} + V\cdot\nabla s + \frac{\omega}{\partial p}\,s_0 = 0$$

where s is a value of $\partial\theta/\partial p$, appropriate to the entire troposphere, and s_0 is its value at 1000 mb. Then, if we assume that $s_0 = .6\,s$,

$$\frac{\partial s}{\partial t} + V\cdot\nabla s + \frac{.6\,\omega\,s}{2p} = 0$$

Finally, since the remaining factors of σ^2 vary much more slowly (percentagewise) than $\frac{\partial \theta}{\partial p}$,

$$\frac{\partial}{\partial t}(\sigma^2) + \bar{V} \cdot \nabla \sigma^2 + \frac{g \omega_0 \sigma^2}{2p} = 0 \qquad (8)$$

Once σ^2 and $\frac{\partial h}{\partial t}$ have been graphed, it is a straightforward matter to graph ω.

Summary of equations in order of solution

$$\frac{\partial \zeta}{\partial t} = J(\psi, \beta_2) - \left(\frac{\beta_2 - 600}{P}\right)\frac{\partial}{\partial x} J(\psi, \beta_3) \qquad \text{I}$$

$$\nabla^2 \chi = -\frac{\omega_0}{2P} \qquad \text{II}$$

$$W = k \times \nabla\psi + \nabla\chi \qquad \text{III}$$

$$\frac{\partial}{\partial t}(\sigma^2) + W \cdot \nabla(\sigma^2) + \frac{4\omega_0 \sigma^2}{2P} = 0 \qquad \text{IV}$$

$$\eta = \nabla^2\psi + f \qquad \text{V}$$

$$\nabla^2\left(\frac{\partial \psi}{\partial t}\right) - \frac{2f\eta}{\sigma^2}\frac{\partial \psi}{\partial x} = \frac{2f\eta}{\sigma^2} W \cdot \nabla\psi - W \cdot \nabla(\zeta^2\psi)$$

$$= J(\psi, \eta) - \frac{f\eta\omega_0}{gP} \qquad \text{VI}$$

$$\omega = \frac{gP}{\sigma^2}\left(\frac{\partial \psi}{\partial x} + W \cdot \nabla\psi\right) \qquad \text{VII}$$

$$\nabla^2\left(\frac{\partial \psi}{\partial t}\right) + W \cdot \nabla\eta - \frac{g\omega}{fP}\nabla^2\psi - \frac{\eta\omega_0}{2P} = 0 \qquad \text{VIII}$$

ψ = stream function, at 600 mb

$\psi = \beta_{400} - \beta_{800}$

Solution

For ψ, h, σ^2

(1) Compute w_0 from I
(2) Compute X by inverting II, using output of (1) and (2)
(3) Compute W from III, using output of (2)
(5) Solve IV for $\frac{\partial h}{\partial t}$, using output of (1) and (3)
(6) Compute w from VII, using output of (3) and (5)
(7) Solve VIII for $\frac{\partial \psi}{\partial t}$, using output of (1), (3), and (6)
(8) Compute σ^2 from IV, using output of (1) and (3)