

# NOAA Technical Memorandum NMFS-PIFSC-60

February 2017

doi:10.7289/V5/TM-PIFSC-60

---

## Stock Assessment of the Coral Reef Fishes of Hawaii, 2016



Marc O. Nadon

Pacific Islands Fisheries Science Center  
National Marine Fisheries Service  
National Oceanic and Atmospheric Administration  
U.S. Department of Commerce

## About this document

The mission of the National Oceanic and Atmospheric Administration (NOAA) is to understand and predict changes in the Earth's environment and to conserve and manage coastal and oceanic marine resources and habitats to help meet our Nation's economic, social, and environmental needs. As a branch of NOAA, the National Marine Fisheries Service (NMFS) conducts or sponsors research and monitoring programs to improve the scientific basis for conservation and management decisions. NMFS strives to make information about the purpose, methods, and results of its scientific studies widely available.

NMFS' Pacific Islands Fisheries Science Center (PIFSC) uses the **NOAA Technical Memorandum NMFS** series to achieve timely dissemination of scientific and technical information that is of high quality but inappropriate for publication in the formal peer-reviewed literature. The contents are of broad scope, including technical workshop proceedings, large data compilations, status reports and reviews, lengthy scientific or statistical monographs, and more. NOAA Technical Memoranda published by the PIFSC, although informal, are subjected to extensive review and editing and reflect sound professional work. Accordingly, they may be referenced in the formal scientific and technical literature.

A **NOAA Technical Memorandum NMFS** issued by the PIFSC may be cited using the following format:

Nadon, M. O.

2017. Stock assessment of the coral reef fishes of Hawaii, 2016. U.S.

Dep. Commer., NOAA Tech. Memo., NOAA-TM-NMFS-PIFSC-60, 212

p. doi:10.7289/V5/TM-PIFSC-60.

Species photos for *C. argus*, *P. insularis*, *M. vanicolensis*, and *C. melampygu*s were taken by the author for NOAA-PIFSC. All other species photos are from Randall, J.E. 1997. Randall's underwater photos. Collection of almost 2,000 underwater photos. Unpublished. Accessed through fishbase.org.

---

### For further information direct inquiries to

Chief, Science Operations Division  
Pacific Islands Fisheries Science Center  
National Marine Fisheries Service  
National Oceanic and Atmospheric Administration  
U.S. Department of Commerce  
1845 Wasp Boulevard  
Honolulu, Hawaii 96818

Phone: 808-725-5331

Fax: 808-725-5532

---

Cover: Photograph taken by author for the Pacific Islands Fisheries Science Center.  
National Marine Fisheries Service.



## Pacific Islands Fisheries Science Center

National Marine Fisheries Service  
National Oceanic and Atmospheric Administration  
U.S. Department of Commerce

---

### Stock Assessment of the Coral Reef Fishes of Hawaii, 2016

<sup>1,2</sup>Marc O. Nadon

<sup>1</sup>Joint Institute for Marine and Atmospheric Research  
University of Hawaii  
1000 Pope Road, Honolulu, Hawaii 96822

<sup>2</sup>Pacific Islands Fisheries Science Center  
National Marine Fisheries Service  
1845 Wasp Boulevard  
Honolulu, Hawaii 96818

---

NOAA Technical Memorandum NMFS-PIFSC-60

February 2017

doi:10.7289/V5/TM-PIFSC-60

## ABSTRACT

This report contains single-species assessments of 27 reef-associated fish stocks around the main Hawaiian Islands using data from various sources collected during the 2003–2016 period. Previous management actions have set acceptable biological catches (ABCs) at the family level using either a percentile of historical catches or a catch-MSY approach. Here, we used fishery-independent size composition and abundance data from diver surveys combined with fishery-dependent catch estimates to calculate current fishing mortality rates ( $F$ ), spawning potential ratios ( $SPR$ ),  $SPR$ -based sustainable fishing rates ( $F_{30}$ :  $F$  resulting in  $SPR = 30\%$ ), and catch levels corresponding to these sustainable rates ( $C_{30}$ ). We used a length-based model to obtain mortality rates and a relatively simple age-structured population model to obtain the various stock status metrics.  $C_{30}$  were obtained by combining  $F_{30}$  estimates with current population biomass estimates derived directly from diver surveys or indirectly from the total catch. The overfishing limits ( $OFL$ ) corresponding to a 50% risk of overfishing was defined as the median of the  $C_{30}$  distribution. A novel data-poor approach was used to estimate life history parameters for 11 species with either no or inadequate published growth and maturity studies. We used Monte Carlo simulations to incorporate all sources of uncertainty (i.e., life history parameters, average length, abundance, and catch). Of the 27 assessed species, 11 had median  $F/F_{30}$  ratios greater than 1 and therefore median  $SPR$  values below the minimum overfished limit of 30%. Another two were close to this limit ( $30\% \leq SPR < 35\%$ ). This suggests some stocks may be experiencing overfishing and, if at equilibrium, may also be overfished. Surgeonfishes and parrotfishes were the families with the most species with low  $SPR$  values, while goatfishes generally had higher  $SPR$  values. Typically, species with low  $SPR$  were the ones with long lifespan (i.e., surgeonfishes) or highly targeted (i.e., jacks, snappers). Species with shorter lifespans (i.e., goatfishes) fared generally better. As a final step, overfishing probability distributions for a range of catch limits were generated for all 27 species.





## CONTENTS

ABSTRACT .....	i
INTRODUCTION .....	1
Description of the Fisheries .....	2
METHODS .....	3
Stock Area .....	3
Size Selectivity in the Fishery .....	4
Data Sources .....	5
Total and Natural Mortality Models .....	8
Population Simulation Model .....	10
Overfishing Limit Calculation .....	11
Decision Process for Multiple Data Sources .....	12
Analyses Work Flow .....	12
RESULTS SUMMARY .....	14
DISCUSSION.....	15
ACKNOWLEDGEMENTS.....	18
REFERENCES .....	19
TABLES AND FIGURES.....	25
SPECIES REPORTS .....	35
<i>Acanthurus blochii</i> .....	36
<i>Acanthurus dussumieri</i> .....	42
<i>Naso brevirostris</i> .....	49
<i>Naso hexacanthus</i> .....	55
<i>Naso lituratus</i> .....	62
<i>Naso unicornis</i> .....	69
<i>Carangoides orthogrammus</i> .....	75
<i>Caranx ignobilis</i> .....	81
<i>Caranx melampygus</i> .....	88
<i>Aprion virescens</i> .....	95
<i>Lutjanus fulvus</i> .....	102
<i>Lutjanus kasmira</i> .....	109
<i>Mulloidichthys flavolineatus</i> .....	116
<i>Mulloidichthys pfluegeri</i> .....	123
<i>Mulloidichthys vanicolensis</i> .....	130

<i>Parupeneus cyclostomus</i> .....	136
<i>Parupeneus insularis</i> .....	143
<i>Parupeneus porphyreus</i> .....	150
<i>Calotomus carolinus</i> .....	157
<i>Chlorurus perspicillatus</i> .....	163
<i>Chlorurus spilurus</i> .....	169
<i>Scarus dubius</i> .....	175
<i>Scarus psittacus</i> .....	181
<i>Scarus rubroviolaceus</i> .....	187
<i>Cephalopholis argus</i> .....	193
<i>Monotaxis grandoculis</i> .....	199
<i>Myripristis berndti</i> .....	206

## INTRODUCTION

The 2006 re-authorization of the Magnuson-Stevens Fishery Conservation and Management Act calls for annual catch limits (ACLs) to be set for all exploited stocks in the United States and its territories in order to, among other goals, insure sustainable harvesting practices. In the U.S. Pacific, exploited stocks include a multitude of coral reef-associated finfish species inhabiting shallow-water areas around a large number of islands and atolls. The high species diversity, the mixture of commercial and recreational fishing effort, and the spatially diffused nature of the fisheries result in a comparatively data-poor situation for these stocks. This has led the Western Pacific Regional Fishery Management Council (WPRFMC) to set ACLs using basic analytical methods such as using the 75<sup>th</sup> percentile of historical catches or using catch-based methods applied at the family level (Sabater & Kleiber, 2013). However, recent efforts in fisheries-independent surveys and life history research by the Pacific Islands Fisheries Science Center (PIFSC) have improved this situation to the point where simple age-based assessment approaches can now be applied to individual coral-reef fish stocks (Nadon et al., 2015).

In this report, the status of 27 of the most commonly exploited coral-reef fish species of Hawaii was assessed using a length-based mortality model and a relatively simple numerical population model. Using this approach, we obtained estimates of fishing mortality ( $F$ ) and spawning potential ratio ( $SPR$ ) over the recent time period,  $F$  at  $SPR = 30\%$  ( $F_{30}$ ), catch associated with  $F_{30}$  ( $C_{30}$ ), and overfishing risk tables for a range of catch limits (including a proxy overfishing limit,  $OFL$ , defined as the  $C_{30}$  level that results in a 50% chance of overfishing; see Table 1 for the definition of all parameters).

For the purposes of our  $SPR$ -based approach in this paper, we used a default definition of overfished and overfishing as recommended by Restrepo et al. (1998):

- Overfished limit:  $SPR = 30\%$ , with overfished defined as  $SPR < 30\%$
- Overfishing limit:  $F$  at  $SPR = 30\%$  ( $F_{30}$ ), with overfishing defined as  $F > F_{30}$  or  $F/F_{30} > 1$ .

It is important to note that since our analyses assumed populations were at equilibrium a stock that is experiencing overfishing was also considered overfished (and vice-versa).

For this approach, we used fishery-independent size composition and abundance data provided by NOAA diver surveys as well as fishery-dependent data from the State of Hawaii commercial fishing database. These 27 species, from 8 different families, were selected based on the availability of length data from either data sources and the availability of reliable life history parameters related to growth, maturity, and longevity. In the situation where no published life history parameters were available, a new data-poor estimation approach developed at PIFSC and the University of Miami was implemented (Nadon & Ault, 2016). Additional species could not be assessed due to data limitations, but this situation may change as new data sources become available. Although most of these species' depth range occurs within the 3 nautical mile

limit defining State waters, species inhabiting depths beyond 50 m have significant portions of their range (18–26%) in federal waters, mainly on Penguin Bank and in the channel waters between Maui, Molokai, and Lanai (see Table 3, Figure 2, and Figure 3).

### **Description of the Fisheries**

The Hawaiian Archipelago is a large island chain extending 2600 km along a SE-NW axis from 19°N, 155°W to 28°N, 178°W (Figure 1). The archipelago is composed of 18 islands and atolls which are typically divided into two broad regions: the inhabited main Hawaiian Islands (MHI; 1.4 million individuals; [dbedt.hawaii.gov/census](http://dbedt.hawaii.gov/census)) and the mostly un-inhabited Northwestern Hawaiian Islands (NWHI). The MHI consists of 8 geologically young, high (4,205 m maximum elevation) volcanic islands while the NWHI have low elevation (275 m max elevation). The MHI were settled by people around AD 1250, and reef fish communities in the MHI have been exploited since that time (Kittinger et al., 2011). The NWHI were never permanently inhabited. However, they were the focus of commercial fishing, especially in the 19th century (Kittinger et al., 2011). Presently, the coral reef ecosystem around the MHI are critical to both fishing and tourism activities (Cesar & Beukering, 2004).

The coral reef fishery around the MHI involves near-shore recreational/subsistence fishing combined with a small commercial fishing sector. Almost a third of Hawaii households are involved in recreational-subsistence fishing (Hamnett et al., 2006), and their catches are estimated to exceed those of the commercial sector significantly (Friedlander & Parrish, 1997; Zeller et al., 2005; Williams & Ma, 2013). The recreational-subsistence sector is composed of (mostly) shore-based fishers using a variety of gears such as spears, hook-and-line, traps, and small gill and cast nets. The primary targeted families include larger jacks and snappers, and, to a lesser extent, smaller reef-associated families such as surgeonfish, goatfish, soldierfish, and parrotfish (Friedlander & Parrish, 1997). Commercial marine landings in Hawaii are mostly composed of coastal-pelagic species (> 80% of catches; DeMello, 2004), but also include the reef fishes targeted by the recreational-subsistence sector. The direct monetary value of the near-shore fishery is only 10–20% of the pelagic fishery (Gulko et al., 2002), but it is culturally and socially important to the local population.

## METHODS

The general assessment approach used in the present document recognizes the relatively data-poor status of most of the coral-reef fish stocks in Hawaii. We specifically focused on the best available data sources which are the ongoing NOAA-PIFSC diver surveys (2005–2016) and the recent commercial data from the State of Hawaii (2003–2015). Since these two data sources contain relatively short time-series, our assessment approach was limited to equilibrium models that assume relatively constant fishing mortality and recruitment. The validity of this assumption was investigated for each species by looking at temporal trends in average length in the catch and abundance index from diver surveys as illustrated in the Species Report section at the end of this document. Furthermore, the lack of stock-recruitment relationships meant that we had to rely on a biological reference point (BRP) based on per-recruit spawning stock biomass (SSBR) and the ratio of current SSBR to un-exploited SSBR (spawning potential ratio, *SPR*, a measure of the spawning potential of a stock). The general assessment approach consisted of three main steps: 1) calculate current fishing mortality rates (*F*) using diver and commercial data derived average length estimates in a length-mortality model, 2) calculate *SPR* and  $F_{30}$  (the fishing mortality rate resulting in  $SPR = 30\%$ ) using published life history parameters in a population simulation model, and 3) calculate the catch limit associated with a given  $F_{30}$  ( $C_{30}$ ) by combining the  $F_{30}$  estimate with an estimate of current population biomass (derived directly from diver surveys or indirectly by relating the current total catch with the current *F*). A Monte Carlo procedure was used to integrate the uncertainty in each individual parameters related to length, population size, and life history. The median of the  $C_{30}$  distribution is the overfishing limit (*OFL*) value, by definition, since it corresponds to a 50% chance of overfishing. A schematic of these steps and decisions are presented in Figure 4. The following sections explain this approach in greater details and describe the various data sources.

### Stock Area

The first step of any stock assessment is typically to define the geographical extent of the stocks being analyzed. It is still not entirely clear to what level the reef fish populations around the MHI are connected and if significant larval exchanges or adult movements exist between the different Hawaiian Islands, although it is generally accepted that fish populations between the MHI and NWHI are disconnected (see Discussion section for details). In this report, all 27 stocks were analyzed at the MHI scale (Figure 1) mainly due to data limitations, as well as to follow current management stock definitions. Further stock connectivity studies may suggest that future stock assessments be conducted at different spatial scales for certain species.

Another consideration for these assessments was the extent of each species' geographical range that fell in federal vs. state waters. For each species, we obtained depth range estimates from Baited Remote Underwater Video (BRUV) and bottom-fish camera (BotCam) exploratory surveys conducted by PIFSC (J. Asher, pers. comm.) and the University of Hawaii (J. Drazen, pers. comm.). We also used the mesophotic deep diving exploratory work conducted by Pyle et al. (2016). We did not attempt to quantify abundance-at-depth given the limited coverage of

these surveys. We simply reported the maximum depth that individual species can inhabit, which may be fairly marginal in certain cases, and calculated the sea floor area to this depth. In the Species Report section, we provide both the maximum depth and the percentage of sea floor area in federal vs. state waters for all species (see Table 3, Figure 2, and Figure 3 for a summary).

### Size Selectivity in the Fishery

The State of Hawaii has minimum size regulations on some of the species included in this report. However, it is uncertain to what degree these are respected by fishermen. To infer on the selectivity pattern in the reef fish fishery, we used the indirect abundance-at-length data from the State commercial data set (see Data Source section below for details) and, to a lesser extent, the Hawaii Marine Recreational Fishing Survey (HMRFS). Fishing records were utilized from the 3 main fishing gear types used in inshore areas: hook-and-line (44% of fishing reports for the species targeted in this study, DAR gear code: 1–6, 8–10, 61–63, 70, and 91–93), spearfishing (40%, gear code: 13 and 14), and various nets (15%, gear code: 20–23, 25–27, 30, 32, 33, 40, 41, and 45). Other gears, such as traps, represented less than 1% of reports for the species selected in the current study and data from these reports were not used. From what is known about the reef fish fishery and the size composition in the catch, it was clear that some form of logistic selectivity curve was most appropriate (i.e., there was no indication of reduced selectivity at larger sizes). We therefore used the following formulation of logistic selectivity:

$$S = \frac{1}{1 + \exp\left(-\ln(19) \frac{L - L_{S50}}{L_{S95} - L_{S50}}\right)} \quad (1)$$

where  $L$  is the length at which selectivity is estimated,  $L_{S50}$  is the length at 50% selectivity, and  $L_{S95}$  is the length at 95% selectivity. To estimate these two parameters, we first searched for a discontinuous break in the size composition histogram obtained from the commercial data set (and HMRFS data set, if it had sufficient observations) to obtain an initial estimate of  $L_{S95}$ . We then looked at the smallest size bin in the size composition graph and split the difference between this size and  $L_{S95}$  to obtain an initial estimate of  $L_{S50}$ . For example, the commercial catch size structure for *Parupeneus porphyreus* did not have individuals below 20 cm, had a few individuals in the 20–22 cm and 22–24 cm ranges (~ 50), and a significantly higher number of individuals in the 24–26 cm range (close to 200 fish). The number of fish in size bins above 24–26 cm increased in a continuous manner. This discontinuous jump between 22–24 cm and 24–26 cm appeared to be related to selectivity; consequently a first estimate of  $L_{S95}$  was set at 26 cm. The smallest size bin was 20–22 cm and we therefore set  $L_{S50}$  at 23 cm (i.e. the midpoint between 20 and 26 cm). To test the validity of the  $L_{S95}$  and  $L_{S50}$  parameters, we simulated a size structure using our population simulation model with these parameters (see Population Simulation section) and compared it to the actual size structure data. If the size structures did not agree, we adjusted the  $L_{S50}$  and  $L_{S95}$  parameters and ran further simulations. Typically, one or two runs were necessary to obtain an appropriate estimate of the selectivity parameters. For most species, sensitivity analyses were also used with different selectivity parameter pairs.

Finally, the length-mortality model used to obtain an estimate of total mortality (see Total Mortality section below) requires an estimate of length at full selectivity (“ $L_{S100}$ ”). Since it is impossible to differentiate between  $L_{S95}$  and  $L_{S100}$ , given their similarity and the limited resolution of our length data, we simply used  $L_{S95}$  as our estimate of length at full selectivity for the length-mortality model. This made our total mortality estimate slightly lower than if  $L_{S100}$  was known.

## Data Sources

### *Size composition, density, and total biomass from diver surveys*

The main source of fisheries-independent data came from the diver surveys conducted by the Pacific Islands Fisheries Science Center’s Coral Reef Ecosystem Program (CREP). The diver surveys were used to obtain size structure data (from which average length in the exploited phase could be calculated -  $\bar{L}$ ) and abundance data (from which population biomass estimates could be derived). Below is a brief description of the survey protocol. An in-depth description is available in Ayotte et al. (2015).

Starting in 2005, trained divers from the NOAA Pacific Islands Fisheries Science Center (PIFSC) have been conducting visual surveys around the MHI. Survey sites were randomly selected within strata defined by depth-bins (shallow, 0–6 m; mid, 6–18 m; and deep, 18–30 m). All coastlines from all islands in the MHI were surveyed, except for the small, restricted island of Kaho’olawe. For practical and safety reasons, surveys were limited to depths above 30 m. During a typical CREP survey day, a NOAA ship deployed 3 to 4 small dive boats that sampled pre-assigned random sites along 10 to 12 miles of coastline. The daily starting location of the ship along different coastlines of the MHI was set in a systematic manner, with the goal of covering as much of the shoreline as possible. At each site, stationary point counts were implemented by two paired divers inside contiguous 15-m diameter cylinders that extended from the bottom to the surface (Brandt et al., 2009; Smith et al., 2011; Williams et al., 2011). Divers first listed all observed fish species during an initial 5-minute period. The divers then went through this list, one species at a time, recording the number of individuals and estimating sizes of all fish seen within the cylinder. Fish sizes were recorded as total lengths to the nearest cm. Fishes from species not listed during the initial 5-minute period, but observed later in the survey, were also recorded but classified in a different data category (i.e., non-instantaneous count). Divers were continuously trained between cruises in size estimation using fish cut-outs of various sizes. Diver performance during research cruises was evaluated by comparing size and count estimates between paired divers.

Average length in the exploited phase ( $\bar{L}$ ) was obtained from the abundance-at-size data by averaging all length observations (weighted by count) from 2005 to 2016 for a species inside each of the 4 subregions of the MHI (see Figure 1, Table 2 for subregion descriptions). Only lengths above size at full selectivity ( $L_{S95}$ ) were kept. The overall  $\bar{L}$  was obtained by averaging all 4 sub-regional  $\bar{L}$  weighted by the respective size of each subregion’s shallow reef area (Table 2). This was done to account for differences in size composition due to uneven fishing pressure



(inferred from humans per reef area values—Table 2) and uneven sampling effort between regions. The standard deviations of  $\bar{L}$  estimates were obtained by bootstrapping the diver survey data set by re-sampling survey sites within subregion (Figure 1) and applying the weighted mean procedure described above to generate a distribution of  $\bar{L}$ .

Numerical density estimates (fish per 100 m<sup>2</sup>) were obtained by dividing the fish counts in each survey by the area per survey (353 m<sup>2</sup> from two 15-m diameter survey cylinders) and multiplying by 100 (an individual survey consisted of the combined fish counts from the two divers deployed at a random site). We calculated selectivity for each observation using the observed length and Eq. (1), and multiplied each fish count by this value. The overall average numerical density was obtained by (1) averaging site-level density estimates within a coastline sector (see Figure 5 for sector map) and by (2) averaging all sector-level density estimates together, weighted by the amount of reef area in each sector. It is important to note here that we did not use the higher spatial resolution sectors for the  $\bar{L}$  bootstrapping procedure due to the relatively low length observation sample sizes (we use the subregions instead). The standard deviations of overall mean density estimates were obtained by bootstrapping the diver survey data set by re-sampling survey sites within sector (Figure 5) and applying the weighted mean procedure described above to generate a distribution of mean numerical density. These density estimates are presented as time series graphs in the Species Report section. Finally, it is important to note that only instantaneous fish counts were kept for this calculation in order to get an abundance estimate close to “true” density. The implied assumption here is that the “catchability” ( $q$ ) of an individual underwater survey is the fraction of the total hard-bottom population area (96,208 ha) covered by a single survey (353 m<sup>2</sup>;  $q = 3.67e-7$ ). We used hard-bottom area since all species in this report are heavily associated with this habitat type. Some species, like goatfishes, feed over soft-bottom areas but they are usually within range of hard-bottom areas which they rely on for refuge.

Biomass density estimates (kg per 100 m<sup>2</sup>) were obtained by using the same approach as for numerical density, but by first converting each individual fish length into weight using published length-weight conversion parameters as provided in the Species Report section. Fish biomass density per sector was used to estimate fish biomass per sector by multiplying biomass density by the amount of hard-bottom area in each sector (obtained from CREP and bathymetric data compiled by the Hawaii Mapping Research Group). Total stock biomass was obtained by summing all sector biomass together. The standard deviations of biomass density and total fish biomass were obtained through bootstrapping in a similar fashion as for numerical density.

One limitation of this data set was the potential impact of fish behavior on the assumed catchability coefficient ( $3.66e-7$ ) for population biomass calculations. Cryptic behavior and diver avoidance (or attraction) will have an effect on this assumed value and this will differ between species. Although the biomass calculations for all species were done assuming this value, we discuss potential biomass estimate biases for certain species in the Species Report section.

Another limitation of this data set is the potential mismatch between the survey domain (limited to 30-m depth) and the greater depth range of certain species. For species occurring at depths greater than 30 m, we did not attempt to assign a population abundance to the un-sampled

sea floor area, given our limited knowledge of the amount of suitable habitat at these depths. We do however discuss this potential bias and implications for the relevant species in the Species Report section.

#### *Size composition and total catch from fishing report data*

We used the commercial fishing report data from the State of Hawaii's Division of Aquatic Resources (DAR) to obtain an estimate of total commercial catch. Catch records for certain taxa that were not identified to the species level were not included in analyses that depended on this data set (all parrotfishes, the "kala" group of surgeonfishes composed of *Naso unicornis*, *N. annulatus*, and *N. brevirostris*, and the *Acanthurus blochii*/*A. xanthopterus* reporting group). We also used the Hawaii Marine Recreational Fishing Survey program (HMRFS) catch estimates provided by Williams and Ma (2013) to obtain an estimate of total recreational catch from 2004 to 2015. For the commercial data, we simply summed the reported catch by weight by year from 2003 to 2015 by species (for species reported at the species level). Given the uncertainty associated with both catch data sets, we assumed that annual total catch has been mostly stable from 2003 to 2015 and that the year to year variability is representative of the uncertainty around the total catch estimates. Thus it was not necessary that the time periods of commercial and recreational data sets matched perfectly. HMRFS data were not always available for every species in every year. Past reports (Williams & Ma, 2013) have shown that the nearshore fishery is dominated by recreational fishing, and there is no evidence that recreational fishing effort has been going dramatically up in recent years given that Hawaii's human population is increasing at a fairly low rate (~ 1% per year). Plotting the catch data by year for these species revealed mostly year-to-year variability, although these estimates were generally fairly variable (see Species Report section). In order to come up with a total catch estimate, we fitted a lognormal distribution to both the commercial and recreational yearly catch estimates and summed these together using a Monte Carlo procedure by randomly sampling both distributions, adding those values together, and fitting a new lognormal distribution to the resulting data set. Of note, when summing lognormal distributions, the median value of the final distribution will be different than the sum of the two median values of the original distribution (see Species Report section).

We also used the commercial fishing report data to obtain a second source of size composition data for species that have individual species codes (note: the HMRFS data set did not have sufficient length observation to be used consistently). As discussed in the Selectivity section, all 3 main fishing gears (hook-and-line, spearfishing, and nets) had a similar selectivity pattern, as inferred from the similar size composition of their catch (i.e. similar logistic shape with no indication of lower selectivity at larger sizes). We therefore combined catch records from all 3 fishing gears, providing close to 100,000 individual reports. Unfortunately, direct-length observations are not recorded in the commercial data set. However, each record report the total weight and the number of fish caught by species. It was possible to obtain an indirect measure of length by dividing the catch in weight by the number of fish caught to obtain average weight per individual report. Average weights per report were then converted to total lengths using published standard allometric weight ( $W$ )–length ( $L$ ) relationships (see individual species tables in the Species Report section),

$$W = \alpha \cdot L^\beta \quad (2)$$

which can be inverted to obtain

$$L = \left( \frac{W}{\alpha} \right)^{\left( \frac{1}{\beta} \right)} \quad (3)$$

where  $\alpha$  is a scaling and  $\beta$  is a volumetric model parameter. For each species, average length per report were combined across all years using numbers caught per report as a weighting variable, in a similar fashion as for diver-survey  $\bar{L}$ . Converting average weight per report to average length per report can theoretically lead to a biased estimate of average length (Jensen's inequality caused by the non-linear length-weight relationship; Ruel & Ayres, 1999). To test the degree to which this occurred, for each species, we compared the average length from catch records with only one recorded fish caught vs. the average calculated for all catch records regardless of how many fish per record were reported. We found minimal differences in average length between these two calculation methods. The resulting length observations for each report were checked and lengths that were greater than the maximum reported for each species were discarded. The standard deviations of the average length estimates derived from commercial report data were estimated by bootstrapping individual report data by subregion and running the above analyses.

#### *Life history parameter sources*

We reviewed the scientific literature for published life history parameters related to growth, longevity, and maturity. We did not restrict our search to local studies given the paucity of peer-reviewed literature on coral reef fish biology. If multiple growth or maturity studies were available for a species, we prioritized local studies, followed by the most recent, in-depth studies (even if from a different geographical area). If no life history study was available, for certain species, we used the data-poor life history estimation approach described in Nadon and Ault (2016). In short, this approach uses a local estimate of maximum length to provide family-specific probability distribution for all main life history parameters. The standard deviations of life history parameters were obtained by one of the following methods, presented in order of preference based on reliability: 1) bootstrapping the raw length, age, or maturity data, when available, 2) using the coefficient of variations at different sample sizes from Kritzer et al. (2001) for growth and Nadon (unpublished) for maturity (Table 4), or 3) from the stepwise approach itself, if it was used to generate life history parameters (Nadon and Ault, 2016). See Figure 4 for a summary of life history steps within this assessment framework.

### **Total and Natural Mortality Models**

As mortality rates increase, the probability of a fish reaching larger sizes decreases, and thus the mean of the size frequency distribution ( $\bar{L}$ ) decreases accordingly. Theoretically, the average length  $\bar{L}$  in the catch can be expressed as

$$\bar{L} = \frac{\int_{a_c}^{a_\lambda} N_a L_a da}{\int_{a_c}^{a_\lambda} N_a da} \quad (4)$$

where the exploitable phase is integrated from  $a_c$  (age at first capture) to  $a_\lambda$  (oldest age),  $N_a$  is the abundance at age class  $a$ , and  $L_a$  is the expected length at age  $a$ .

A formula for estimating mortality rates using estimates of  $\bar{L}$  was derived from Eq. (4) by Ehrhardt and Ault (1992). The first step in this derivation was to substitute  $L_a$  in Eq. (1) with the von Bertalanffy growth function and  $N_a$  with the exponential mortality model

$$\bar{N}_{a+\Delta a} = \bar{N}_a e^{-Z\Delta a} \quad (5)$$

where  $Z$  is the total instantaneous mortality rate and  $\Delta a$  is the age interval, normally one year. Step two was to integrate and algebraically solve for  $Z$ ,

$$\left( \frac{L_\infty - L_\lambda}{L_\infty - L_{S95}} \right)^{Z/K} = \frac{Z(L_{S95} - \bar{L}) + K(L_\infty - \bar{L})}{Z(L_\lambda - \bar{L}) + K(L_\infty - \bar{L})} \quad (6)$$

where  $K$  and  $L_\infty$  are parameters of the von Bertalanffy growth equation (assumed to be constant over time),  $L_{S95}$  is the size at full selectivity (see Selectivity section), and  $L_\lambda$  is the expected size at oldest known age, respectively. We selected Eq. (6) instead of the Beverton-Holt model (Beverton & Holt 1956) due to a reported bias in this model associated with the assumption of infinite longevity (Ehrhardt & Ault, 1992). It is important to note that Eq. (4) and (6) are valid in equilibrium conditions where fishing mortality has been relatively constant for a sufficient amount of time for the population to be in a stable state. For each individual species, we looked at temporal trends in average lengths and estimated population abundance to verify that this was the case (see the Species Report section for further details). Fishing mortality was obtained from  $F = Z - M$ , where  $M$  is the instantaneous natural mortality rate which was derived from longevity in our study (see paragraph below for more details). Estimates of total instantaneous mortality rates  $Z$  were computed from Eq. (6) using a numerical procedure. Values of  $L_\lambda$  (expected length at maximum age  $a_\lambda$ ) were estimated from the von Bertalanffy growth function using an observed maximum age.

To derive  $F$  from  $Z$ , as well as to parameterize our population simulation model, it is necessary to obtain an estimate of natural mortality ( $M$ ). To do so, we used the procedure of Alagaraja (1984), similar to Hoenig (1983) and Hewitt and Hoenig (2005), which assumes that 4% of a cohort survives to the observed maximum age ( $a_\lambda$ ):

$$M = \frac{-\ln(0.04)}{a_\lambda} \quad (7)$$

We used the 4% cohort survivorship value based on the analyses of Nadon et al. (2015) which showed that this is an appropriate survivorship value for coral reef fishes. We did not have independent estimates of *M per se* and had to rely on this longevity-based approach. Although there are other data-poor methods for estimating natural mortality, involving other parameters (e.g.,  $K$ ,  $L_{inf}$ ,  $L_{mat}$ , water temperature), two recent scientific papers on the subject clearly suggest that longevity-only methods are better performing (Kenchington, 2014; Then et al., 2015). It is important to consider the potential difficulty in obtaining a representative longevity value in heavily exploited stocks. To reduce this concern as much as possible, we always selected the oldest recorded age, regardless of geographical location, as our measure of longevity. It was, unfortunately, impossible to only select longevity estimates from un-exploited stocks given that there are few life history studies on such stocks. The Species Report section provides details of parameters selected and their sources, and output parameter values and associated uncertainty.

### Population Simulation Model

We built a population model in order to calculate various stock metrics ( $SPR$ ,  $F_{30}$ ,  $LC_{30}$ ). For exploited fish populations, a biological reference point for sustainability risks (spawning potential ratio,  $SPR$ ) was computed using a numerical model to simulate exploited fish populations. The computations were based on the mortality rates derived from  $\bar{L}$  estimates and life history parameters synthesized from the literature. Numerical abundance at age  $a$  was estimated through use of an exponential mortality function (Eq. 9). Length-at-age was estimated from the von Bertalanffy growth equation, and converted to weight-at-age using the allometric weight-length relationship (Eq. 1).

The numerical model was used to obtain spawning stock biomass per recruit ( $SSBR$ ), at given levels of fishing mortality by summing over individuals in the population between the age of sexual maturity ( $a_m$ ; age where 50% of individuals are mature, with knife-edge assumption) and 1.5 times the oldest recorded age ( $a_l$ ),

$$SSBR = \sum_{a_m}^{1.5a_l} \bar{N}_a \bar{W}_a \quad (8)$$

Where  $\bar{N}_a$  is the mean abundance at age  $a$  and  $\bar{W}_a$  is the mean weight of individuals at age  $a$  (derived from the von Bertalanffy equation combined with length-weight equation). The model was run using weekly time steps. Average abundance at age was modeled using the following equation:

$$\bar{N}_{a+\Delta a} = \bar{N}_a e^{-(S \cdot F + M) \Delta a} \quad (9)$$

where  $F$  and  $M$  are fishing and natural mortality, respectively, and  $S$  is selectivity which was defined by a logistic curve with parameters  $L_{S50}$  and  $L_{S95}$  (see Selectivity section).  $SPR$  was computed as the ratio of the current  $SSB$  relative to that of an unexploited stock:

$$SPR = \frac{SSB_{exploited}}{SSB_{unexploited}} \quad (10)$$

Estimated *SPR*s were compared to the recommended 30% *SPR* threshold below which a stock is likely no longer sustainable (i.e., is experiencing recruitment overfishing), a standard recommended for less well-known stocks (Gabriel et al., 1989; Restrepo et al., 1998; Clark, 2002).  $L_{C30}$ , the size at first capture required to obtain an  $SPR = 30\%$  was also estimated using this model. To do so, we used an iterative procedure which calculated *SPR* at incrementing  $L_c$  values (keeping all other parameters fixed) until the  $SPR = 30\%$  level was reached. An identical procedure was used to obtain  $F_{30}$ .

Calculated *SPR*, proportion of *SPR* iterations that resulted in  $SPR < 0.30$ ,  $L_{C30}$ ,  $F_{30}$ , and  $F/F_{30}$  values are provided in the Species Report section.

### Overfishing Limit Calculation

The sections above presented the data sources and models used to obtain various population parameters (mortality rates, *SPR*,  $F_{30}$ ,  $L_{C30}$ ). To calculate an overfishing limit (*OFL*) estimate (i.e. the catch that results in a 50% chance of overfishing), we first needed to obtain an estimate of standing population biomass ( $B$ ). This could be obtained in up to two ways depending on data reliability and availability for each species: 1) extrapolating total biomass from the diver-survey biomass density estimates, as explained earlier, and 2) by using the estimates of total catch ( $C$ ), natural mortality, and length-derived fishing mortality in the Baranov catch equation:

$$B = C \div \frac{F}{F + M} (1 - e^{-(F+M)}) \quad (11)$$

From one or both of these estimates of current population biomass, we derived the catch level corresponding to  $F_{30}$  ( $C_{30}$ ) by using the Baranov equation and our estimates of sustainable fishing mortality rate ( $F_{30}$ ):

$$C_{30} = B \cdot \frac{F_{30}}{F_{30} + M} (1 - e^{-(F_{30}+M)}) \quad (12)$$

The final distribution of  $C_{30}$  estimates and other derived values (e.g., *SPR*) were obtained by incorporating all sources of uncertainty (data and parameters) using a Monte Carlo approach. In short, we drew a random value from the probability distributions of each data source ( $\bar{L}$ , life history parameters, diver-derived population biomass, and total catch) and ran all the steps to calculate  $C_{30}$  using these random values (Figure 4). The Monte Carlo draws for parameters that could not be negative (e.g. catch,  $\bar{L}$ ) were bounded at zero (if they were drawn from probability distributions that allowed negative values). The Monte Carlo procedure was repeated 6,000 times to generate distributions of  $C_{30}$  and other derived values. The median of the  $C_{30}$  distribution represented the catch level with a 50% chance of overfishing (*OFL*).

It is important to note that randomly drawn combinations of life history parameters could lead to a biologically impossible scenario where an  $\bar{L}$  was larger than the pristine average length predicted by these parameters. For these random draws,  $F$  will be negative and  $SPR$  will go above 1. This can be a fairly common situation for lightly fished stocks with  $\bar{L}$  close to its pristine value and is not necessarily an indication of incorrect life history parameter distributions. Instead, it is the result of a lack of a proper *a priori* covariance structure between these parameters that should have limited certain parameter combinations. For example, a very high  $M$  value, combined with a low  $K$  and low  $L_{inf}$  values can lead to a pristine average length that is unrealistically low and below a randomly drawn  $\bar{L}$ . To correct this issue, for a randomly drawn  $\bar{L}$ , we rejected life history parameter combinations that led to this situation and re-drew life history parameters until a realistic combination was sampled.

### Decision Process for Multiple Data Sources

Throughout the process used to generate *OFL* estimates (Figure 4), there were several steps where decisions had to be made regarding data sources. To reduce the subjectivity of these decisions, we created a decision table presented in Figure 4. In short, there were 4 main decision steps: 1) whether to combine the  $\bar{L}$  estimates from diver and report data or keep only one source, 2) whether to use a local study, external study, or the Nadon and Ault (2016) approach as a source of life history parameters, 3) whether to use a bootstrap procedure on raw data or the meta-analysis of Kritzer et al. (2001) and Nadon (unpubl.) to generate uncertainty of life history parameters, and 4) whether to use  $C_{30}$  distribution generated from diver-survey biomass or from catch-based biomass to calculate an *OFL* (this final decision point is discussed for individual species in the Species Reports section but is ultimately left to managers).

### Analyses Work Flow

The raw diver survey data were provided by CREP (file named “all\_diver\_rea.rdata”) and the raw commercial data were extracted directly from the PIFSC Oracle database where they are stored. Two R scripts were used to process these raw data sets (“process\_uvc\_data.r” and “process\_dar\_data.r”). Other R scripts were used to obtain various metrics and their associated distributions: average length (“get\_lbar.r”), population biomass (“get\_diver\_biomass.r”), commercial and recreational catch (“get\_catch.r”), and time series graphs (“get\_abund\_timeseries.r” and “get\_lbar\_timeseries.r”).

The overall approach to generate population status metrics ( $F$ ,  $F_{30}$ , and  $SPR$ ) and the  $C_{30}$  distributions (Figure 4) was implemented in a Java-language computer program developed specifically for this purpose. This tool requires inputs in the form of probability distribution parameters (e.g., mean, standard deviation) for 1) the life history parameters, 2) average length, 3) total catch (if available), and 4) population biomass from surveys (if available). In the case of the stepwise approach (Nadon and Ault, 2016), this tool also requires parameters for an  $L_{max}$  distribution and a species’ family-level taxonomic group. Other required parameters are entered as fixed values: selectivity ( $L_{S50}$  and  $L_{S95}$ ),  $a_0$ , length-weight parameters (alpha and beta), number

of Monte Carlo iterations, the assumed survivorship at maximum age value ( $S$ ), and the spawning schedule (when spawning occurs throughout the year, set to monthly by default).

Once launched, the program will draw random samples from the input distributions and run the calculations showed in Figure 4: 1) generate an estimate of  $Z$  from the length-mortality model, 2) calculate  $M$  from longevity (if necessary) and  $F$ , 3) calculate  $SPR$  and  $F_{30}$  using the population simulation model, and 4) calculate  $C_{30}$  from the diver surveys (if available) and from the catch data (if available). The program outputs a comma-separated data file (.CSV) containing parameter values for all Monte Carlo iterations. This CSV file is processed with an R script to generate the standard suite of figures and tables displayed in each species report (“process\_mast.r”).



## RESULTS SUMMARY

It is beyond the scope of this summary section to discuss individual assessments of the 27 reef fish species in this report. In-depth results, comments, and specific concerns can be found in the Species Reports section at the end of this manuscript. Here, we provide a brief overview of the state of reef fishes in Hawaii, as can be inferred from the species analyzed in this report. Table 5 presents a summary of selected stock status metrics for each species.

Out of the 27 species in this report, 25 had depth ranges extending significantly (21 to 26%) into federal waters (i.e., beyond the 3 nautical mile limit of state waters). Two parrotfish species appeared to have limited presence in federal waters (*Chlorurus spilurus* and *Scarus psittacus*). A significant portion (~ 40%) of federal reef fish seafloor habitat above 250 m occurred at depths between 40 and 70 m, mainly on Penguin Bank, off Molokai (Figure 2).

We found local life history parameters for only 11 species and had to use parameters from studies conducted elsewhere in the Indo-Pacific region for 5 species. The remaining 11 species had either no (9) or inadequate (2) published life history studies and we used the data-poor approach presented in Nadon & Ault (2016) to obtain estimates (Table 5). As expected, the assessments conducted with these estimates were more variable than those conducted with life history parameters from actual studies.

Of the 27 assessed species, 11 had median *SPR* values below the minimum 0.30, which is the recommended limit we used as the default metric for overfishing in the current report (Restrepo et al., 1998). By this metric, *SPR* values lower than 0.30 indicate a stock may be experiencing overfishing (and due to the equilibrium assumption, may also be overfished). Two species had a median *SPR* values close to this limit (< 0.35; Table 5). Surgeonfishes had the most species with low *SPR* values, while goatfishes generally had higher *SPR* values. Typically, species with low *SPR*s were the ones with long lifespan (i.e., surgeonfishes, large parrotfishes, *A. virescens*) or highly targeted (i.e., jacks). Species with shorter lifespans (i.e., goatfishes) fared generally better.

To generate  $C_{30}$  from estimates of  $F_{30}$  (fishing mortality at  $SPR = 0.3$ ), we had to obtain estimates of current stock biomass, either directly from diver surveys or indirectly from dividing total catch by an estimate of current  $F$ . Biomass estimates derived from diver-survey biomass were usually much more precise than those obtained through the catch. Consequently,  $C_{30}$  distributions derived from diver-biomass were generally more precise as well. Table 5 presents both estimates of population biomass by species. For species where both biomass estimates were available, 6 out of 12 species had biomass estimates within an order of magnitude of each other (Table 5). Almost all biomass estimates derived from the catch were lower than those derived from diver surveys, suggesting some measure of under-reporting in the commercial catch and/or some bias from the HMRFS recreational fishing survey. The only 2 species with greater catch-derived biomass than diver-survey biomass were goatfish species, with a potential bias associated with an important fishery for juveniles (see Species Reports section).

## DISCUSSION

The assessment approach used in this report focused on fisheries-independent diver-survey data and recent estimates of both commercial and recreational catch. It used mortality and population models that are relatively simple, but well-tested and appropriate for the data-poor situation that characterize coral-reef fisheries (Ehrhardt & Ault, 1992; Ault et al., 2005; Hordyk et al., 2015; Nadon et al., 2015). Several assumptions and caveats apply to these models.

First, we assumed the stocks analyzed in the current study were at equilibrium in terms of both mortality rates and recruitment (i.e., relatively constant over the last decade or so). Ault et al. (2005) showed that mortality rates derived from average length are fairly robust to even moderate levels of recruitment variation. In the case of an extreme recruitment event (e.g., an annual ten-fold increase in the background recruitment level), we would have expected average lengths to decrease dramatically for a few years followed by a quick upward rebound before a return to the long-term equilibrium. In the case of a long-term increasing trend in fishing mortality, we would have expected a slow, constant decline in average length. We did not observe such patterns in average length over time in our study and this suggests that potential fluctuations in recruitment levels over time were not significant enough to affect our average length estimates and that fishing mortality was more or less constant. Furthermore, the fish abundance time series from diver surveys did not reveal major trends, although these observations were fairly variable and only started in 2005. Despite these issues, these time series observations also support the general equilibrium assumption. It is worth noting again that because we assumed the populations were at equilibrium, a finding that a stock may be experiencing overfishing also meant that the stock may be overfished (and vice versa).

A second key assumption was that size composition, abundance, and catch data were representative of the true population around the MHI. Both the underwater visual survey and commercial report data sets had strengths and weaknesses. The underwater surveys by scuba divers did not reach depths beyond 30 m due to safety and time constraints, thus underestimating total population size for species with depth range extending beyond this depth. However, diver surveys were able to sample remote and exposed areas of the MHI that are likely visited less frequently by fishers. The size composition and abundance data for the visual survey data set was thus more representative of nearshore (< 30-m deep) communities but encompassed the entire nearshore waters in the MHI, including remote, lightly-populated, and relatively inaccessible sections of coastlines. On the other hand, size composition data from commercial reports included information on deeper fish communities, but were less likely to be representative of inaccessible coastlines. Despite these potential biases, the size composition information from these two disparate data sets have been shown to be similar suggesting that the average lengths used in the current report were likely representative of the real values (Nadon et al., 2015). It is also important to note that the population abundance estimate from the diver surveys assumed a catchability coefficient equal to the area of a single survey divided by the total hard-bottom habitat area above 30 m. In other words, we did not assume any detectability bias which could have an impact on population biomass estimates for certain, more mobile species (jacks, snappers, larger parrotfishes). However, for more mobile species we did not use diver survey

abundance estimates for  $C_{30}$  and *OFL* estimation because of this potential bias, and instead used catch based abundance estimates.

The total catch estimates used in this report came from the commercial reports and the HMRFS program for recreational catches (which are the vast majority of the total catch), both of which have issues. The commercial reports may under-estimate the catch given that it is self-reported. The HMRFS catches are based on an interpolation of total fishing effort from telephone surveys, combined with creel surveys, and suffer from low sampling effort. These issues reduced our confidence in catch-derived  $C_{30}$ . However, for most species, diver-survey based  $C_{30}$  were also available, which helped verify catch-derived metrics.

Third, it is highly likely that segments of fish stocks located around more heavily populated islands (i.e., Oahu, Maui) face considerably higher fishing pressure than more isolated parts of the MHI. However, it is not entirely clear to what level reef fish populations are connected between islands, in terms of larval exchange and/or adult movement. For example, a tagging study failed to detect inter-island movements for tagged *Caranx ignobilis*, a large and highly mobile predator (Meyer et al., 2007). Conversely, a State of Hawaii tagging program did record a kahala jack (*Seriola dumerili*, amberjack) swimming hundreds of kilometers from the NWHI to the MHI in a 3-year timespan (Tagawa & Clayward, 2006). Genetic connectivity studies indicate that most reef fish species have no genetic structuring across the Hawaiian archipelago (Rivera et al., 2004; Craig et al., 2007; Gaither et al., 2010). However, the absence of genetic structure does not necessarily imply that stocks are well-connected at time scales relevant to population dynamic processes. More informatively, recent genetic parentage analyses of two coral reef fish species in Australia have found parent-offspring pairs at distances up to 250 km, with a median dispersal distance of 110 km and 190 km (Williamson et al., 2016). As shown in Table 2, the longest distance between islands in the MHI is 116 km (Oahu – Kauai), with most islands separate by much shorter distances. Another parentage study conducted on Hawaii Island found yellow tang surgeonfish parent-offspring pairs separated by up to 184 km, although they did not attempt to find cross-channel pairs (Christie et al., 2010). Furthermore, a recent study of passive pelagic particle connectivity in the MHI, based on a pelagic larval duration of 45 days found a median distance for successful settlements around 100 km (Wren et al., 2016) and that cross-channel dispersal can be common. Population connectivity within the MHI is still an open question and will require further research attention, however, based on current research, it appears that our MHI-scale analyses are appropriate. As a side note, it is generally well-accepted that the MHI and NWHI reef fishes form different stocks, and that little larval or adult exchange exists between these two regions given the dominant current direction and the large distances involved, with the exception of the kahala example mentioned above (Toonen et al., 2011; Wren et al., 2016).

Fourth, for many species, we had to use life history parameters from other Pacific areas. It is possible that these values change geographically and with environmental conditions (Choat & Robertson, 2002; Gust et al., 2002, although see Donovan et al., 2013). The availability of an extensive underwater visual survey data set for the relatively pristine NWHI allowed Nadon et al. (2015) to evaluate the validity of the length-based mortality model used in the current report, as well as the validity of our life history information. Nadon et al. (2015) used independent

estimates of  $M$  from the NWHI (where  $Z$  derived from average length is assumed to be equal to  $M$ ) to derive an estimate of average cohort survivorship ( $S$ ) to maximum age ( $a_\lambda$ ). They obtained a value close to 0.04. The exact survivorship value is linked to the sampling effort in the data set from which  $a_\lambda$  is obtained (i.e., the larger the number of aged individuals, the greater the chance of finding extremely old individuals that are not representative of a 5% or even 1% cohort survivorship value; Kenchington, 2014). Since the  $a_\lambda$  value for reef fishes generally comes from life history studies with less than 100 aged individuals, it is possible that these values represent cohort survivorship higher than 1.5%, which is what our analysis suggests. For species with no published life history parameters, we used the approach presented in Nadon & Ault (2016) to provide first-step estimates. This approach uses a local estimate of  $L_{\max}$  which may be biased downward in heavily fished stocks (and thus result in biased life history parameters). This is less of an issue in the current report given that length data in the pristine NWHI were available to generate  $L_{\max}$  estimates.

### *Future directions*

The Pacific Islands Fisheries Science Center is conducting research on a wide variety of subjects that will help address some of the concerns mentioned above. The Coral Reef Ecosystem Program, with assistance from the Stock Assessment Program, will continue collecting fisheries-independent diver data. This data set, in conjunction with longer HMRFS and DAR catch data sets, will eventually have a time-series of sufficient length to run more advanced, non-equilibrium models. The continuing efforts in deep-water surveys using underwater cameras will also provide abundance and size composition data for a section of the reef fish populations that is not accessible by diver survey and may be significantly different. These camera system deployments are also generating deep-water habitat information. The Life History Program at PIFSC is continuing their work on growth, maturity, and longevity of local stocks in the U.S. Pacific which will lead to more appropriate life history parameters and will allow further assessment for data-less species. Finally, new population genetic work, mainly at the University of Hawaii, can provide further information regarding the scale of reef fish population connectivity across the MHI and depending on the results may lead to future assessments being conducted at a different scale (e.g., island or smaller island group).

## ACKNOWLEDGEMENTS

This technical memo was made possible by the work of a large number of people across multiple NOAA offices in Hawaii. The Coral Reef Ecosystem Program at PIFSC collected much of the diver survey data under the supervision of R. Brainard and I. Williams, with the assistance of the crew and officers of the NOAA Ship *Hi'ialakai* and *Oscar Elton Sette*. The Life History Program at PIFSC generated several of the maturity, growth, and longevity information used in this document (E. DeMartini, B. Taylor, A. Andrews, and J. O'Malley). The work contained in this document was also greatly improved by a number of reviewers (K. Stokes, G. Pilling, C. Dichmont, H. Choat, and A. Yau).

## REFERENCES

- Alagaraja, K.  
1984. Simple methods for estimation of parameters for assessing exploited fish stocks. *Indian Journal of Fisheries* **31**:177–208.
- Allen, G. R.  
1985. Snappers of the world: an annotated and illustrated catalogue of Lutjanid species known to date. Food and Agriculture Organization of the United Nations.
- Andrews, A. H., E. E. DeMartini, J. A. Eble, B. M. Taylor, D. C. Lou, and R. L. Humphreys.  
2016. Age and growth of bluespine unicornfish (*Naso unicornis*): a half-century life-span for a keystone browser, with a novel approach to bomb radiocarbon dating in the Hawaiian Islands. *Canadian Journal of Fisheries and Aquatic Sciences* **73**:1575–1586.
- Ault, J. S., S. G. Smith, and J. A. Bohnsack.  
2005. Evaluation of average length as an estimator of exploitation status for the Florida coral-reef fish community. *ICES Journal of Marine Science* **62**:417–423.
- Ayotte, P., K. McCoy, A. Heenan, I. Williams, and J. Zamzow.  
2015. Coral Reef Ecosystem Program Standard Operating Procedures: Data Collection for Rapid Ecological Assessment Fish Surveys. Available from [http://www.pifsc.noaa.gov/library/pubs/admin/PIFSC\\_Admin\\_Rep\\_15-07.pdf](http://www.pifsc.noaa.gov/library/pubs/admin/PIFSC_Admin_Rep_15-07.pdf).
- Beverton, R. J. H., and S. J. Holt.  
1956. A review of methods for estimating mortality rates in exploited fish populations, with special reference to sources of bias in catch sampling. *Rapports et proces-verbaux des reunions du Conseil International pour l'Exploration de la Mer* **140**:67–83.
- Brandt, M. E. et al.  
2009. A cooperative multi-agency reef fish monitoring protocol for the Florida Keys coral reef ecosystem. Natural Resource Report NPS/SFCN/NRR - 2009/150, National Park Service, Fort Collins.
- Cesar, H. S. J., and P. van Beukering.  
2004. Economic valuation of the coral reefs of Hawai'i. *Pacific Science* **58**:231–242.
- Choat, J. H., and L. M. Axe.  
1996. Growth and longevity in acanthurid fishes; an analysis of otolith increments. *Marine Ecology-Progress Series* **134**:15–26.
- Choat, J. H., and D. R. Robertson.  
2002. Age-based studies on coral reef fishes. Pages 57–80 *Coral Reef Fishes: Dynamics and Diversity in a Complex Ecosystem*. Academic Press, San Diego.

Christie, M. R., B. N. Tissot, M. A. Albins, J. P. Beets, Y. Jia, D. M. Ortiz, S. E. Thompson, and M. A. Hixon.

2010. Larval Connectivity in an Effective Network of Marine Protected Areas. PLoS ONE **5**:e15715.

Clark, W. G.

2002. F35% Revisited Ten Years Later. North American Journal of Fisheries Management **22**:251–257.

Cole, K. S.

2009. Size-dependent and age-based female fecundity and reproductive output for three Hawaiian goatfish (Family Mullidae) species, *Mulloidichthys flavolineatus* (yellowstripe goatfish), *M. vanicolensis* (yellowfin goatfish), and *Parupeneus porphyreus* (whitesaddle goatfish). Report to the Division of Aquatic Resources Dingell-Johnson Sport Fish Restoration.

Craig, M. T., J. A. Eble, B. W. Bowen, and D. R. Robertson.

2007. High genetic connectivity across the Indian and Pacific Oceans in the reef fish *Myripristis berndti* (Holocentridae). Marine Ecology-Progress Series **334**:245–254.

Craig, M. T., and E. C. Franklin.

2008. Life history of Hawaiian “redfish”: a survey of age and growth in “aweoweo (*Priacanthus meeki*) and u”u (*Myripristis berndti*). Hawaii Institute of Marine Biology, Kaneohe, Hawaii.

DeMello, J. D.

2004. Commercial marine landings from fisheries on the coral reef ecosystem of the Hawaiian Archipelago. Proceedings of the 2001 Fisheries Symposium of Hawai’i. American Fisheries Society - Hawaii Chapter, Honolulu.

Donovan, M. K., A. M. Friedlander, E. E. DeMartini, M. J. Donahue, and I. D. Williams.

2013. Demographic patterns in the peacock grouper (*Cephalopholis argus*), an introduced Hawaiian reef fish. Environmental Biology of Fishes **96**:981–994.

Eble, J. A., R. Langston, and B. W. Bowen.

2009. Growth and reproduction of Hawaiian Kala, *Naso unicornis*. Division of Aquatic Resources report, Honolulu.

Ehrhardt, N. M., and J. S. Ault.

1992. Analysis of two length-based mortality models applied to bounded catch length frequencies. Transactions of the American Fisheries Society **121**:115–122.

- Everson, A. R., and A. Williams.  
1989. Maturation and Reproduction in Two Hawaiian Eteline Snappers, Uku, *Aprion virescens*, and Onaga, *Etelis cormicans*. Fishery Bulletin **87**: 877–888.
- Friedlander, A. M., and J. D. Parrish.  
1997. Fisheries harvest and standing stock in a Hawaiian Bay. Fisheries Research **32**:33–50.
- Gabriel, W. L., M. P. Sissenwine, and W. J. Overholtz.  
1989. Analysis of spawning stock biomass per recruit: an example for Georges Bank haddock. North American Journal of Fisheries Management **9**:383–391.
- Gaither, M. R., R. J. Toonen, D. R. Robertson, S. Planes, and B. W. Bowen.  
2010. Genetic evaluation of marine biogeographical barriers: perspectives from two widespread Indo-Pacific snappers (*Lutjanus kasmira* and *Lutjanus fulvus*). Journal of Biogeography **37**:133–147.
- Gulko, D., J. E. Maragos, A. M. Friedlander, and R. E. Brainard.  
2002. Status of coral reefs in the Hawaiian Archipelago. Pages 155–182 The state of coral reef ecosystems of the United States and Pacific Freely Associated States: 2002. National Oceanic and Atmospheric Administration report, Silver Spring.
- Gust, G., J. H. Choat, and A. Ackerman.  
2002. Demographic plasticity in tropical reef fishes. Marine Biology **140**:1039–1051.
- Hamnett, M., M. Lui, and D. Johnson.  
2006. Fishing, ocean recreation, and threats to Hawaii’s coral reefs. Hawaii Coral Reef Initiative, Honolulu.
- Hewitt, D. A., and J. M. Hoenig.  
2005. Comparison of two approaches for estimating natural mortality based on longevity. Fishery Bulletin **103**:433–437.
- Hoenig, J. M.  
1983. Empirical use of longevity data to estimate mortality rates. Fisheries Bulletin **82**:898–903.
- Holland, K. N., J. D. Peterson, C. G. Lowe, and B. M. Wetherbee.  
1993. Movements, distribution and growth rates of the white goatfish *Mulloidichthys flavolineatus* in a fisheries conservation zone. Bulletin of Marine Science **52**:982–992.
- Hordyk, A., K. Ono, S. Valencia, N. Loneragan, and J. Prince.  
2015. A novel length-based empirical estimation method of spawning potential ratio (SPR), and tests of its performance, for small-scale, data-poor fisheries. ICES Journal of Marine Science **72**:217–231.



- Howard, K. G.  
2008. Community structure, life history, and movement patterns of parrotfishes: large protogynous fishery species. PhD. University of Hawaii at Manoa, Honolulu.
- Jehangeer, M. I.  
2003. Some population parameters of the goatfish, *Mulloidichthys vanicolensis* from the lagoon of Mauritius. ACP-EU Fisheries Research Report (14).
- Kenchington, T. J.  
2014. Natural mortality estimators for information-limited fisheries. *Fish and Fisheries* **15**:533–562.
- Kittinger, J. N., J. M. Pandolfi, J. H. Blodgett, T. L. Hunt, H. Jiang, K. Maly, L. E. McClenachan, J. K. Schultz, and B. A. Wilcox.  
2011. Historical reconstruction reveals recovery in Hawaiian coral reefs. *PLoS ONE* **6**:e25460.
- Kritzer, J. P., C. R. Davies, and B. D. Mapstone.  
2001. Characterizing fish populations: effects of sample size and population structure on the precision of demographic parameter estimates. *Canadian Journal of Fisheries and Aquatic Sciences* **58**:1557–1568.
- Kulbicki, M., N. Guillemot, and M. Amand.  
2005. A general approach to length-weight relationships for New Caledonian lagoon fishes. *Cybiurn* **29**:235–252.
- Meyer, C. G., K. N. Holland, and Y. P. Papastamatiou.  
2007. Seasonal and diel movements of giant trevally *Caranx ignobilis* at remote Hawaiian atolls: implications for the design of marine protected areas. *Marine Ecology Progress Series* **333**:13–25.
- Moffitt, R. B.  
1979. Age, growth, and reproduction of the kumu, *Parupeneus porphyreus*. M.Sc. thesis. University of Hawaii, Honolulu.
- Morales-Nin, B., and S. Ralston.  
1990. Age and growth of *Lutjanus kasmira* (Forskaal) in Hawaiian waters. *Journal of Fish Biology* **36**:191–203.
- Murty, V. S.  
2002. Marine Ornamental Fish Resources of Lakshadweep. CMFRI, Spl. Pub. 72: 134 pp.

- Nadon, M. O., and J. S. Ault.  
2016. A stepwise stochastic simulation approach to estimate life history parameters for data-poor fisheries. *Canadian Journal of Fisheries and Aquatic Sciences* **73**:1874–1884.
- Nadon, M. O., J. S. Ault, I. D. Williams, S. G. Smith, and G. T. DiNardo.  
2015. Length-based assessment of coral reef fish populations in the Main and Northwestern Hawaiian Islands. *PLoS ONE* **10**:e0133960.
- Pyle, R. L. et al.  
2016. A comprehensive investigation of mesophotic coral ecosystems in the Hawaiian Archipelago. *PeerJ* **4**:e2475.
- Restrepo, V. R. et al.  
1998. Technical guidance on the use of precautionary approaches in implementing national standard 1 of the Magnuson-Stevens Fishery Conservation and Management Act. NOAA Technical Memorandum NMFSF/ SPO-031, NOAA, USA: 54 pp.
- Rivera, M. A. J., C. D. Kelley, and G. K. Roderick.  
2004. Subtle population genetic structure in the Hawaiian grouper, *Epinephelus quernus* (Serranidae) as revealed by mitochondrial DNA analyses. *Biological Journal of the Linnean Society* **81**:449–468.
- Ruel, J. J., and M. P. Ayres.  
1999. Jensen's inequality predicts effects of environmental variation. *Trends in Ecology & Evolution* **14**:361–366.
- Sabater, M. G., and P. Kleiber.  
2013. Improving specification of acceptable biological catches of data-poor reef fish stocks using a biomass-augmented catch-MSY approach. Page 24. Western Pacific Regional Fishery Management Council, Honolulu, HI.
- Seki, M. P.  
1986. Carangidae. Pages 86–87 *Fishery atlas of the Northwestern Hawaiian Islands*. NOAA Tech. Rep. NMFS 38.
- Smith, A., and P. Dalzell.  
1993. Fisheries resources and management investigations in Woleai Atoll, Yap State, Federated States of Micronesia. Inshore Fish. Res. Proj., Tech. Doc., South Pacific Commission. Noumea, New Caledonia.
- Smith, S. G., J. S. Ault, J. A. Bohnsack, D. E. Harper, J. Luo, and D. B. McClellan.  
2011. Multispecies survey design for assessing reef-fish stocks, spatially explicit management performance, and ecosystem condition. *Fisheries Research* **109**:25–41.

- Sudekum, A. E., J. D. Parrish, R. L. Radtke, and S. Ralston.  
1991. Life history and ecology of large jacks in undisturbed, shallow, oceanic communities. *Fishery Bulletin* **89**:493–513.
- Tagawa, A., and T. Clayward.  
2006. Hawaii's ulua and papio tagging project 2000 to 2004. 06-01, DAR Technical Report. Honolulu, HI.
- Then, A. Y., J. M. Hoenig, N. G. Hall, and D. A. Hewitt.  
2015. Evaluating the predictive performance of empirical estimators of natural mortality rate using information on over 200 fish species. *ICES Journal of Marine Science* **72**:82–92.
- Toonen, R. J. et al.  
2011. Defining boundaries for ecosystem-based management: a multispecies case study of marine connectivity across the Hawaiian Archipelago. *Journal of Marine Biology* **2011**:1–13.
- Williams, I. D., B. L. Richards, S. A. Sandin, J. K. Baum, R. E. Schroeder, M. O. Nadon, B. Zgliczynski, P. Craig, J. L. McIlwain, and R. E. Brainard.  
2011. Differences in reef fish assemblages between populated and remote reefs spanning multiple archipelagos across the central and western Pacific. *Journal of Marine Biology* **2011**:1–14.
- Williams, I., and H. Ma.  
2013. Estimating catch weight of reef fish species using estimation and intercept data from the Hawaii Marine Recreational Fishing Survey. Page 61. Honolulu: NOAA Pacific Islands Fisheries Science Center Administrative Report H-13-04.
- Williamson, D. H. et al.  
2016. Large-scale, multidirectional larval connectivity among coral reef fish populations in the Great Barrier Reef Marine Park. *Molecular Ecology* **25**: 6039-6054.
- Wren, J. L. K., D. R. Kobayashi, Y. Jia, and R. J. Toonen.  
2016. Modeled Population Connectivity across the Hawaiian Archipelago. *PLOS ONE* **11**:e0167626.
- Zeller, D., S. Booth, and D. Pauly.  
2005. Reconstruction of coral reef fisheries catches for US associated islands in the Western Pacific Region, 1950 to 2002. Western Pacific Regional Fishery Management Council report, Honolulu.

## TABLES AND FIGURES

Table 1.--List of parameters used in the document.

Parameter	Definition
$\alpha, \beta$	Parameters of the length-weight relationship
$a_\lambda$	Oldest recorded age (i.e., longevity)
$a_0$	Theoretical age at which length equals zero from the von Bertalanffy growth curve
$B$	Total population biomass
$C_{30}$	Catch limit resulting in $SPR = 0.3$
$F$	Instantaneous annual fishing mortality rate
$F_{30}$	Instantaneous annual fishing mortality rate resulting in $SPR = 0.3$
$K$	Brody growth coefficient of the von Bertalanffy growth curve
$L_{\text{bar}}$ or $\bar{L}$	Average length in the exploited phase of a stock
$L_{C_{30}}$	Size at first capture limit resulting in $SPR = 0.3$
$L_{\text{inf}}$ or $L_\infty$	Expected length at infinite age from the von Bertalanffy growth curve
$L_\lambda$	Expected length at the oldest recorded age
$L_{\text{mat}}$	Length at which 50% of females reach maturity
$L_{\text{max}}$	Longest length in a growth study or 99 <sup>th</sup> percentile of lengths in a population survey
$L_{S50}$	Length at 50% selectivity
$L_{S95}$	Length at 95% selectivity
$M$	Instantaneous annual natural mortality rate
$OFL$	Overfishing limit, defined as the median of the $C_{30}$ distribution
$S$	Survivorship at maximum recorded age
$SPR$	Spawning potential ratio
$Z$	Instantaneous annual total mortality rate

Table 2.-- Information summary of the two principal regions of the Hawaiian Islands, including the four subregions of the main Hawaiian Islands. Reef area is to 30-m depth and excludes soft bottom habitat. Source: CREP and Hawaii Mapping Research Group bathymetric synthesis data set.

Region	Human population (2010)	Reef area (km <sup>2</sup> )	Prop. of total reef in region	Pop. per reef area (# km <sup>-2</sup> )	Channel width (km) <sup>a</sup>
Main Hawaiian Is.					
Hawaii	185,079	168	0.18	1099	48
Maui Nui	154,950	269	0.28	577	48-42
Oahu	953,000	251	0.26	3794	42-116
Kauai-Niihau	65,819	274	0.28	240	116-220

<sup>a</sup> Channel widths from east to west.

Table 3.-- Area of sea floor in the MHI by depth zones, in state and federal waters (soft and hard bottom). Depth range extends to 250 m which is close to the maximum recorded depths for the species included in this report. Source: CREP and Hawaii Mapping Research Group bathymetric synthesis data set.

Depth (m)		Hectares of sea floor			Cumulative percent in fed. Waters
Min	Max	State waters	Federal waters	Total	
0	30	126,930	0	126,930	0
30	40	37,670	57	37,727	< 1
40	50	32,960	23,097	56,057	10
50	60	32,770	27,893	60,663	18
60	70	31,725	19,711	51,436	21
70	80	31,667	9,151	40,818	21
80	90	27,698	6,343	34,041	21
90	100	25,343	7,560	32,903	21
100	150	88,823	32,365	121,188	22
150	200	44,440	24,036	68,476	24
200	250	39,120	32,261	71,381	26

Table 4.--Coefficient of variation of 4 life history parameters at various sample sizes.  $L_{\text{inf}}$ ,  $K$ , and  $a_{\text{max}}$  from Kritzer et al. (2001), and  $L_{\text{mat}}$  from Nadon (unpubl.).

Sample size	CV $L_{\text{inf}}$	CV $K$	CV $L_{\text{mat}}$	CV $a_{\text{max}}$
25	0.167	0.568	0.070	0.190
50	0.111	0.299	0.048	0.168
75	0.078	0.250	0.036	0.147
100	0.060	0.222	0.031	0.129
125	0.050	0.190	0.027	0.118
150	0.045	0.172	0.025	0.113
200	0.040	0.142	0.021	0.092
300	0.030	0.120	0.017	0.074
500	0.021	0.095	0.013	0.056

Table 5.--Species summary of selected stock metrics. Bold text indicates stocks considered overfished /overfishing according to the  $SPR = 30\%$  based biological reference point. Overfishing is defined as  $F/F_{30} > 1$  and overfished is defined as  $SPR < 0.30$ .

Species	Man. unit	Group	LH source	Max depth (m)	Percent in fed. waters	$F/F_{30}$	$SPR$	Pop. from catch (kg)	Pop. from survey (kg)
<b>Acanthuridae</b>									
<i>Acanthurus blochii</i>	C	Ac	E	80	21	<b>2.3</b>	<b>0.12</b>	-	365,000
<i>Acanthurus dussumieri</i>	C	Ac	E	131	22	0.8	0.36	356,000	719,000
<i>Naso brevirostris</i>	C	Ac	E	122	22	<b>1.9</b>	<b>0.14</b>	-	132,000
<i>Naso hexacanthus</i>	C	Ac	E	124	22	<b>2.0</b>	<b>0.13</b>	30,000	-
<i>Naso lituratus</i>	C	Ac	E	93	21	<b>1.3</b>	<b>0.25</b>	30,000	452,000
<i>Naso unicornis</i>	C	Ac	L	120	22	<b>6.0</b>	<b>0.03</b>	-	364,000
<b>Carangidae</b>									
<i>Carangoides orthogrammus</i>	C	Ca	S	235	26	0.7	0.41	123,000	-
<i>Caranx ignobilis</i>	B	ND7	L	228	26	<b>1.1</b>	<b>0.28</b>	1,070,000	-
<i>Caranx melampygus</i>	C	Ca	L	230	26	0.7	0.40	811,000	-
<b>Lutjanidae</b>									
<i>Aprion virescens</i>	B	ND7	L	203	24	0.9	0.33	758,000	434,000
<i>Lutjanus fulvus</i>	C	Lu	S	128	22	0.9	0.33	48,000	180,000
<i>Lutjanus kasmira</i>	B	ND7	L	265	26	0.3	0.62	181,000	496,000
<b>Mullidae</b>									
<i>Mulloidichthys flavolineatus</i>	C	Mu	L	97	21	0.5	0.49	307,000	42,000
<i>Mulloidichthys pfluerigi</i>	C	Mu	S	242	26	0.7	0.41	21,000	-
<i>Mulloidichthys vanicolensis</i>	C	Mu	L	132	22	0.4	0.55	139,000	35,000
<i>Parupeneus cyclostomus</i>	C	Mu	S	113	21	<b>1.3</b>	<b>0.24</b>	12,000	77,000
<i>Parupeneus insularis</i>	C	Mu	S	90	21	0.4	0.57	5,000	42,000
<i>Parupeneus porphyreus</i>	C	Mu	L	140	22	<b>1.9</b>	<b>0.15</b>	15,000	14,000
<b>Scaridae</b>									
<i>Calotomus carolinus</i>	C	Sc	S	71	21	<b>2.2</b>	<b>0.13</b>	-	38,000
<i>Chlorurus perspicillatus</i>	C	Sc	S	80	21	0.5	0.54	-	79,000
<i>Chlorurus spilurus</i>	C	Sc	S	34	0	<b>1.4</b>	<b>0.23</b>	-	139,000
<i>Scarus dubius</i>	C	Sc	S	80	21	0.6	0.45	-	33,000
<i>Scarus psittacus</i>	C	Sc	S	48	10	0.7	0.41	-	130,000
<i>Scarus rubroviolaceus</i>	C	Sc	L	68	21	<b>1.2</b>	<b>0.26</b>	-	624,000
<b>Other families</b>									
<i>Cephalopholis argus</i>	C	Se	L	80	21	0.1	0.80	232,000	777,000
<i>Monotaxis grandoculis</i>	C	Le	S	101	22	0.8	0.38	29,000	232,000
<i>Myripristis berndti</i>	C	Ho	L	159	22	0.4	0.59	-	260,000

Management units: C = Coral Reef Management Unit Species (CREMUS), B = Bottomfish Management Unit Species (BMUS)  
 Grouping: Ac = Acanthuridae, Ca = Carangidae, Ho = Holocentridae, Le = Lethrinidae, Lu = Lutjanidae, Mu = Mullidae, ND7 = Non-deep 7 bottomfish, Se = Serranidae, Sc = Scaridae.

Life history source: E = external (different geographic location), L = local (from Hawaii), S = stepwise approach (Nadon & Ault, 2016).




Figure 1.--Map of the Hawaiian Islands (including Northwestern Hawaiian Islands in inset), with the four subregions. Figure from Nadon et al. (2015).






Figure 2.--Map of the 8 main Hawaiian Islands with deep water depth zones (0-m to 250-m depths). Black contour lines represent the 3 nautical mile State waters limit. Islands are not to scale. Data source: CREP and Hawaii Mapping Research Group.




Figure 3.--Cumulative sea floor area in hectares (top panel) and percentage of total (bottom panel) from 0 m to 250 m. Blue area is federal waters and red area is state waters. Soft and hard bottom included.




Figure 4.--Overall approach and decision points used to calculate stock status and obtain overfishing limits (OFL).




Figure 5.--Survey sectors of the MHI (islands at different scales). These sectors were used when running bootstrap analyses for biomass density data from diver surveys. There are 3 habitat categories (simple, complex, and coral-dominated) and 3 fishing intensity category (green = low, yellow = moderate, and red = high). Habitat categories were derived from CREP habitat survey data and the fishing categories are based on accessibility and human population density.



## **SPECIES REPORTS**

# *Acanthurus blochii*

Ringtail surgeonfish, *pualu*  
Acanthuridae (surgeonfishes)



## Life history and other input parameters

Parameter	Value	SD	Unit	n	Source
$L_{inf}$	363	61	mm	24	Mean: Choat & Robertson (2002), SD: Kritzer (2001)
$K$	0.25	0.14	yr <sup>-1</sup>		
$a_0$	-0.38	-	yr		
$L_{mat}$	276	19	mm	24	Mean: 76% of $L_{inf}$ , SD: Nadon (unpublished)
Longevity	35	6.7	yr	24	Mean: Choat & Robertson (2002), SD: Kritzer (2001)
L-W $\alpha$	1.87e-5	-	-	-	Kulbicki (2005)
L-W $\beta$	3.03	-	-	-	
$L_{S50}$	225	-	mm	-	Estimated.
$L_{S95}$	250				
$\bar{L}$ diver survey	299	3	mm	672	NOAA-CREP diver survey
$\bar{L}$ commercial	-	-	mm	-	-
$\bar{L}$ combined	-	-	mm	-	-
Max. depth	80	-	m	-	Pyle et al. (2016)
Federal waters	21	-	%	-	-

## Stock status and other output parameters

Parameter	Median	SD	Unit
$M$	0.09	0.02	yr <sup>-1</sup>
$F$	0.27	0.21	yr <sup>-1</sup>
$F_{30}$	0.12	0.03	yr <sup>-1</sup>
$F/F_{30}$	2.3	1.7	-
$SPR$	0.12	0.17	-
$SPR < 0.30$ iterations	84	-	%
$L_{C30}$	290	-	mm

Parameter	Median	SD	Unit
$B$ from catch	-	-	-
$B$ from survey	365,392	48,886	kg
Commercial catch	3,604	1,362	kg
Recreational catch	1,394	4,005	kg
Total catch	5,668	3,042	kg
$C_{30}$ from catch	-	-	-
$C_{30}$ from survey	38,300	9,000	kg


## General comments

The DAR commercial reporting system combines *A. xanthopterus* with this species. The commercial catch for these two species is listed above, but as a reference only. In the HMRFS data set, the *A. xanthopterus* catch is 4× larger than *A. blochii*.

Population abundance has been slowly rising since 2007 while  $L_{bar}$  has remained fairly stable at around 30cm, suggesting a population near equilibrium. Selectivity had to be estimated from *A. dussumieri*, a similar species, given the absence of fishery data. A sensitivity run with  $L_{S50}$  at 250 and  $L_{S95}$  at 280 had little impact on the results ( $F$ : 0.28,  $SPR$ : 0.11). There were some concerns with the LH parameters, which came from an Australian study with a low sample size and we therefore ran extra analyses using the stepwise approach ( $L_{max}$ : 420 mm gave  $L_{inf}$ : 390 mm,  $K$ :0.42,  $M$ : 0.1). The higher  $L_{inf}$  value resulted in a much higher  $F$  (0.61) and lower  $SPR$  (0.03). However, the  $C_{30}$  was mostly unchanged (40,300 kg).

The annual total catch estimate (5,668 kg) is fairly small compared to the *OFL* estimate (38,300 kg), even though it includes *A. xanthopterus* catches. This amount of catch is too small to explain the level of fishing mortality estimated from  $L_{bar}$ . However, the diver survey data for this species is fairly reliable (i.e., easily identifiable, common species, non-cryptic) and therefore the  $C_{30}$  estimates from these surveys should be reliable.


*Acanthurus blochii*





**Life history parameter distributions.**



*Acanthurus blochii*




**Abundance index from UVS (blue circles,  $\pm$ SE).**




**Size structure and average length time series from UVS ( $\pm$ SE).**

*Acanthurus blochii*




**Stock status parameter distributions (*SPR*: small bar shows 0.30 level).**



**$C_{30}$  (left) and population size (right) distributions.**

*Acanthurus blochii*



**Overfishing probability for a range of  $C_{30}$  levels (UVS – blue dotted line). *OFL* is represented by a small vertical bar.**

**Probability of overfishing for various  $C_{30}$  levels.**

Overfishing probability	$C_{30}$ from survey (1000 kg)	Overfishing probability	$C_{30}$ from survey (1000 kg)
0.10	28.4	0.31	34.2
0.11	28.8	0.32	34.4
0.12	29.1	0.33	34.7
0.13	29.5	0.34	34.9
0.14	29.8	0.35	35.1
0.15	30.1	0.36	35.4
0.16	30.4	0.37	35.5
0.17	30.7	0.38	35.7
0.18	31.0	0.39	35.9
0.19	31.4	0.40	36.2
0.20	31.6	0.41	36.4
0.21	31.9	0.42	36.7
0.22	32.2	0.43	36.9
0.23	32.5	0.44	37.1
0.24	32.7	0.45	37.3
0.25	32.9	0.46	37.5
0.26	33.1	0.47	37.8
0.27	33.4	0.48	38.0
0.28	33.6	0.49	38.2
0.29	33.9	0.50	38.3
0.30	34.0		

*Acanthurus blochii*

**Probability of overfishing at various minimum sizes.**

Overfishing probability	$L_{C_{30}}$ (mm)	Overfishing probability	$L_{C_{30}}$ (mm)
0.10	364	0.31	324
0.11	362	0.32	322
0.12	358	0.33	320
0.13	356	0.34	317
0.14	353	0.35	315
0.15	351	0.36	315
0.16	349	0.37	313
0.17	349	0.38	310
0.18	346	0.39	308
0.19	344	0.40	306
0.20	342	0.41	306
0.21	340	0.42	304
0.22	338	0.43	302
0.23	335	0.44	299
0.24	335	0.45	299
0.25	333	0.46	297
0.26	331	0.47	295
0.27	331	0.48	295
0.28	328	0.49	292
0.29	326	0.50	290
0.30	324		

# *Acanthurus dussumieri*

Eyestripe surgeonfish, *palani*  
Acanthuridae (surgeonfishes)



## Life history and other input parameters

Parameter	Value	SD	Unit	n	Source
$L_{inf}$	371	41	mm	43	Mean: Choat & Robertson (2002), SD: Kritzer (2001)
$K$	0.296	0.089	yr <sup>-1</sup>		
$a_0$	-0.29	-	yr		
$L_{mat}$	282	14	mm	50	Mean: Choat & Robertson (2002), SD: Nadon (unpublish.)
Longevity	28	4.7	yr	43	Mean: Choat & Robertson (2002), SD: Kritzer (2001)
L-W $\alpha$	2.33e-5	-	-	-	Kulbicki (2005)
L-W $\beta$	3.03	-	-	-	
$L_{S50}$	230	-	mm	-	DAR commercial data
$L_{S95}$	260				
$\bar{L}$ diver survey	323	3	mm	1198	NOAA-CREP diver survey
$\bar{L}$ commercial	334	1	mm	6403	DAR commercial data
$\bar{L}$ combined	333	2	mm	-	-
Max. depth	131	-	m	-	Pyle et al. (2016)
Federal waters	22	-	%	-	-

## Stock status and other output parameters

Parameter	Median	SD	Unit
$M$	0.11	0.02	yr <sup>-1</sup>
$F$	0.12	0.08	yr <sup>-1</sup>
$F_{30}$	0.14	0.02	yr <sup>-1</sup>
$F/F_{30}$	0.8	0.6	-
$SPR$	0.36	0.21	-
$SPR < 0.30$ iterations	37	-	%
$L_{C30}$	129	-	mm


Parameter	Median	SD	Unit
$B$ from catch	356,000	938,000	kg
$B$ from survey	718,622	106,385	kg
Commercial catch	13,370	2,294	kg
Recreational catch	21,529	12,334	kg
Total catch	35,959	11,712	kg
$C_{30}$ from catch	44,900	121,000	kg
$C_{30}$ from survey	90,100	18,600	kg

## General comments

Both population abundance and  $L_{bar}$  (UVS and commercial) time series were relatively stable suggesting a population mostly at equilibrium. Recreational catch did seem to be increasing, but the yearly estimates are fairly variable and this trend may have been spurious.  $L_{bar}$  estimates from the UVS and commercial data sets were almost identical. The LH parameters were obtained from an Australian study with a limited sample size. A sensitivity run using the stepwise approach gave the following results:  $L_{max}$ : 428 mm,  $L_{inf}$ : 399 mm,  $K$ : 0.40,  $M$ : 0.1,  $F_{30}$ : 0.13. The higher  $L_{inf}$  value resulted in a higher  $F$  (0.25) and lower  $SPR$  (0.14). The diver-survey  $C_{30}$  was mostly unchanged (81,000 kg), but the  $C_{30}$  from catch was reduced to 24,000 kg due to the higher  $F$ .


The population estimates derived from the catch was about half the size from the diver surveys. The lower population estimate from the catch data results in a lower catch-derived  $OFL$  estimate (44,900 kg vs. 90,100 kg). Given the quality of the diver data (i.e., high observation count, appropriate sampling design) and the similarity in  $F_{30}$  estimates between the external LH parameters and the stepwise-derived LH parameters, the survey-derived  $C_{30}$  is likely more reliable. The true population size may be even larger than the diver estimate (718,622 kg) given that this species' habitat extends far beyond diver survey depth (131 m vs. 30 m).

*Acanthurus dussumieri*




**Life history parameter distributions.**

*Acanthurus dussumieri*




**Abundance index from UVS (blue circles,  $\pm$ SE) and total catch time series from recreational (green squares) and commercial (orange triangles) sectors.**



**Size structure from commercial catch (top left) and UVS (top right). Average length time series (blue circles – UVS, orange triangles – commercial data,  $\pm$ SE).**


*Acanthurus dussumieri*



**Stock status parameter distributions (*SPR*: small bar shows 0.30 level).**




*Acanthurus dussumieri*



$C_{30}$  and current total catch (left) and population size (right) distributions.

*Acanthurus dussumieri*



**Overfishing probability for a range of  $C_{30}$  levels (catch - orange dashed line, UVS – blue dotted line).  $OFLs$  are represented by small vertical bars.**

**Probability of overfishing for various  $C_{30}$  levels.**

Overfish. probability	$C_{30}$ from catch (1000 kg)	$C_{30}$ from survey (1000 kg)	Overfish. probability	$C_{30}$ from catch (1000 kg)	$C_{30}$ from survey (1000 kg)
0.10	20.4	68.4	0.31	32.2	81.2
0.11	21.1	69.2	0.32	32.8	81.7
0.12	21.8	70.1	0.33	33.4	82.3
0.13	22.4	71.0	0.34	33.9	82.7
0.14	23.0	71.7	0.35	34.6	83.1
0.15	23.6	72.4	0.36	35.1	83.6
0.16	24.2	73.2	0.37	35.7	84.0
0.17	24.8	73.9	0.38	36.3	84.6
0.18	25.3	74.5	0.39	36.9	85.1
0.19	25.8	75.0	0.40	37.6	85.6
0.20	26.3	75.6	0.41	38.3	86.0
0.21	26.9	76.1	0.42	38.9	86.5
0.22	27.4	76.7	0.43	39.7	86.9
0.23	28.0	77.3	0.44	40.2	87.3
0.24	28.6	77.8	0.45	41.0	87.7
0.25	29.1	78.4	0.46	41.7	88.2
0.26	29.7	78.9	0.47	42.6	88.7
0.27	30.2	79.3	0.48	43.3	89.1
0.28	30.7	79.7	0.49	44.1	89.6
0.29	31.2	80.3	0.50	44.9	90.1
0.30	31.7	80.8			

*Acanthurus dussumieri*

**Probability of overfishing at various minimum sizes.**

Overfishing probability	$L_{C_{30}}$ (mm)	Overfishing probability	$L_{C_{30}}$ (mm)
0.10	315	0.31	246
0.11	310	0.32	242
0.12	308	0.33	237
0.13	304	0.34	232
0.14	301	0.35	230
0.15	299	0.36	225
0.16	294	0.37	218
0.17	292	0.38	214
0.18	287	0.39	207
0.19	285	0.40	202
0.20	283	0.41	196
0.21	278	0.42	191
0.22	276	0.43	186
0.23	274	0.44	177
0.24	271	0.45	170
0.25	267	0.46	163
0.26	264	0.47	156
0.27	260	0.48	147
0.28	255	0.49	138
0.29	253	0.50	129
0.30	248		

# *Naso brevirostris*

Paletail unicornfish, *kala lolo*

Acanthuridae (surgeonfishes)



## Life history and other input parameters

Parameter	Value	SD	Unit	n	Source
$L_{inf}$	327	16	mm	120	Mean: Choat & Robertson (2002), SD: Kritzer (2001)
$K$	0.402	0.076	yr <sup>-1</sup>		
$a_0$	-0.21	-	yr		
$L_{mat}$	269	7	mm	120	Mean: Choat & Robertson (2002), SD: Nadon (unpublish.)
Longevity	25	3.0	yr	120	Mean: Choat & Robertson (2002), SD: Kritzer (2001)
L-W $\alpha$	6.09e-6	-	-	-	Kulbicki (2005)
L-W $\beta$	3.24				
$L_{S50}$	200	-	mm	-	Best estimate based on similar surgeonfishes.
$L_{S95}$	220				
$\bar{L}$ diver survey	271	7	mm	561	NOAA-CREP diver survey
$\bar{L}$ commercial	-	-	mm	-	No species-specific catch data.
$\bar{L}$ combined	-	-	mm	-	-
Max. depth	122	-	m	-	Pyle et al. (2016)
Federal waters	22	-	%	-	-

## Stock status and other output parameters

Parameter	Median	SD	Unit
$M$	0.13	0.02	yr <sup>-1</sup>
$F$	0.30	0.16	yr <sup>-1</sup>
$F_{30}$	0.16	0.02	yr <sup>-1</sup>
$F/F_{30}$	1.9	1.0	-
$SPR$	0.14	0.12	-
$SPR < 0.30$ iterations	88	-	%
$L_{C30}$	264	-	mm

Parameter	Median	SD	Unit
$B$ from catch	-	-	kg
$B$ from survey	132,456	31,857	kg
Commercial catch	-	-	kg
Recreational catch	-	-	kg
Total catch	-	-	kg
$C_{30}$ from catch	-	-	kg
$C_{30}$ from survey	17,800	4,680	kg


## General comments

The commercial catch data set for this species includes both *N. unicornis* and *N. annulatus*, and therefore could not be used to estimate  $L_{bar}$ . The HMRFS data was also fairly unreliable given that this species only had 1 year with a recreational catch estimate.

Population abundance and  $L_{bar}$  appeared fairly stable, suggesting a population near equilibrium. Size selectivity had to be estimated, based on other surgeonfishes. A sensitivity run with  $L_{S50}$  at 225 and  $L_{S95}$  at 250 had some moderate impact on the results ( $F = 0.23$ ,  $SPR = 0.22$ ), but did not change overall conclusions. The life history parameters are from a study in Australia, with a decent sample size. Using the stepwise approach to generate LH parameters, we obtained the following estimates:  $L_{max}$ : 416 mm,  $L_{inf}$ : 388 mm,  $K$ : 0.39,  $M$ : 0.10,  $F_{30}$ : 0.12. Similarly to previous surgeonfishes, the  $L_{inf}$  was much higher than the Australian study value. This led to a higher  $F$  (0.71) and lower  $SPR$  (0.02).  $F_{30}$  (0.12) and  $C_{30}$  (13,745 kg) were only slightly lower given the similarity in  $M$  estimates.


Given the quality of the diver data (i.e., high observation count, appropriate sampling design) and the similarity in  $F_{30}$  estimates between the external LH parameters and the stepwise-derived LH parameters, the survey-derived  $C_{30}$  is likely reasonable. The true population size may be even larger than the diver estimate (132,456 kg) given that this species' habitat extends beyond diver survey depth (122 m vs. 30 m).

*Naso brevirostris*





**Life history parameter distributions.**

*Naso brevirostris*




**Abundance index from UVS (blue circles,  $\pm$ SE).**




**Size structure and average length time series from UVS ( $\pm$ SE).**

*Naso brevirostris*




Stock status parameter distributions ( $SPR$ : small bar shows 0.30 level).



$C_{30}$  (left) and population size (right) distributions.

*Naso brevirostris*



**Overfishing probability for a range of  $C_{30}$  levels (UVS – blue dotted line). *OFL* is represented by a small vertical bar.**

**Probability of overfishing for various  $C_{30}$  levels.**

Overfishing probability	$C_{30}$ from survey (1000 kg)	Overfishing probability	$C_{30}$ from survey (1000 kg)
0.10	12.0	0.31	15.6
0.11	12.3	0.32	15.7
0.12	12.5	0.33	15.9
0.13	12.7	0.34	16.0
0.14	12.9	0.35	16.1
0.15	13.1	0.36	16.2
0.16	13.3	0.37	16.3
0.17	13.5	0.38	16.4
0.18	13.7	0.39	16.6
0.19	13.9	0.40	16.7
0.20	14.1	0.41	16.8
0.21	14.2	0.42	16.9
0.22	14.4	0.43	17.0
0.23	14.5	0.44	17.2
0.24	14.7	0.45	17.3
0.25	14.8	0.46	17.4
0.26	14.9	0.47	17.5
0.27	15.1	0.48	17.6
0.28	15.2	0.49	17.7
0.29	15.3	0.50	17.8
0.30	15.5		



*Naso brevirostris*

**Probability of overfishing at various minimum sizes.**

Overfishing probability	$L_{C_{30}}$ (mm)	Overfishing probability	$L_{C_{30}}$ (mm)
0.10	302	0.31	282
0.11	300	0.32	282
0.12	300	0.33	280
0.13	300	0.34	280
0.14	298	0.35	280
0.15	296	0.36	278
0.16	296	0.37	278
0.17	294	0.38	276
0.18	294	0.39	276
0.19	292	0.40	274
0.20	292	0.41	274
0.21	292	0.42	272
0.22	290	0.43	272
0.23	290	0.44	270
0.24	288	0.45	270
0.25	288	0.46	268
0.26	286	0.47	268
0.27	286	0.48	266
0.28	286	0.49	266
0.29	284	0.50	264
0.30	284		

# *Naso hexacanthus*

Sleek unicornfish, *kala lolo*  
Acanthuridae (surgeonfishes)



## Life history and other input parameters

Parameter	Value	SD	Unit	n	Source
$L_{inf}$	599	57	mm	59	Mean: Choat & Robertson (2002), SD: Kritzer (2001)
$K$	0.221	0.061	yr <sup>-1</sup>		
$a_0$	-0.22	-	yr		
$L_{mat}$	511	22	mm	59	Mean: Choat & Robertson (2002), SD: Nadon (unpub.)
Longevity	44	6.9	yr	59	Mean: Choat & Robertson (2002), SD: Kritzer (2001)
L-W $\alpha$	4.12e-5	-	-	-	Choat and Axe (1996)
L-W $\beta$	2.85	-	-	-	
$L_{S50}$	350	-	mm	-	DAR commercial data
$L_{S95}$	410				
$\bar{L}$ diver survey	-	-	mm	-	NOAA-CREP diver survey
$\bar{L}$ commercial	507	2	mm	1249	DAR commercial data
$\bar{L}$ combined	-	-	mm	-	-
Max. depth	124	-	m	-	BRFA BotCam project
Federal waters	22	-	%	-	-

## Stock status and other output parameters

Parameter	Median	SD	Unit
$M$	0.07	0.01	yr <sup>-1</sup>
$F$	0.16	0.08	yr <sup>-1</sup>
$F_{30}$	0.08	0.01	yr <sup>-1</sup>
$F/F_{30}$	2.0	0.9	-
$SPR$	0.13	0.14	-
$SPR < 0.30$ iterations	88	-	%
$L_{C_{30}}$	497	-	mm


Parameter	Median	SD	Unit
$B$ from catch	29,900	75,600	kg
$B$ from survey	-	-	kg
Commercial catch	1,361	1,434	kg
Recreational catch	2,279	3,884	kg
Total catch	4,185	3,664	kg
$C_{30}$ from catch	2,260	5,170	kg
$C_{30}$ from survey	-	-	kg

## General comments

The number of diver survey observations for this species was insufficient to generate reliable  $L_{bar}$  or population biomass estimates. The commercial catch increased from 2007 to 2013, but dropped back down to 2003-2006 levels in the last few years. The recreational catch estimates were fairly variable, making it hard to discern any patterns.  $L_{bar}$  have been stable since 2003, which indicated a population near equilibrium.


The life history parameters came from an Australian study with low sample size. We obtained the following parameters from the stepwise approach, which are fairly similar:  $L_{max}$ : 642 mm,  $L_{inf}$ : 614 mm,  $K$ : 0.28,  $M$ : 0.10,  $F_{30}$ : 0.11.  $F$  (0.20),  $SPR$  (0.15), and  $C_{30}$  (2,578 kg) were also close to the original results. We also tried reducing the survivorship estimate for the  $M$  calculation from 0.04 to 0.01. This increased  $M$  from 0.07 to 0.10 and did not change the results significantly ( $F$ : 0.21,  $SPR$ : 0.15).

*Naso hexacanthus*





**Life history parameter distributions.**

*Naso hexacanthus*




**Total catch time series from recreational (green squares) and commercial (orange triangles) sectors.**




**Size structure and average length time series from commercial data ( $\pm$ SE)**

*Naso hexacanthus*




Stock status parameter distributions (*SPR*: small bar shows 0.30 level).

*Naso hexacanthus*



$C_{30}$  and current total catch (left) and population size (right) distributions.

*Naso hexacanthus*



**Overfishing probability for a range of  $C_{30}$  levels (commercial data – orange dashed line). *OFL* is represented by a small vertical bar.**

**Probability of overfishing for various  $C_{30}$  levels.**

Overfishing probability	$C_{30}$ from catch (1000 kg)	Overfishing probability	$C_{30}$ from catch (1000 kg)
0.10	0.86	0.31	1.53
0.11	0.89	0.32	1.57
0.12	0.93	0.33	1.60
0.13	0.96	0.34	1.63
0.14	1.00	0.35	1.66
0.15	1.02	0.36	1.70
0.16	1.05	0.37	1.74
0.17	1.09	0.38	1.77
0.18	1.12	0.39	1.81
0.19	1.14	0.40	1.86
0.20	1.18	0.41	1.90
0.21	1.22	0.42	1.93
0.22	1.25	0.43	1.98
0.23	1.27	0.44	2.02
0.24	1.30	0.45	2.07
0.25	1.34	0.46	2.11
0.26	1.38	0.47	2.15
0.27	1.41	0.48	2.18
0.28	1.45	0.49	2.22
0.29	1.47	0.50	2.26
0.30	1.49		

*Naso hexacanthus*

**Probability of overfishing at various minimum sizes.**

Overfishing probability	$L_{C_{30}}$ (mm)	Overfishing probability	$L_{C_{30}}$ (mm)
0.10	578	0.31	536
0.11	574	0.32	532
0.12	572	0.33	532
0.13	570	0.34	528
0.14	567	0.35	528
0.15	567	0.36	525
0.16	564	0.37	525
0.17	564	0.38	522
0.18	560	0.39	522
0.19	556	0.40	518
0.20	556	0.41	514
0.21	556	0.42	514
0.22	553	0.43	511
0.23	550	0.44	511
0.24	550	0.45	508
0.25	546	0.46	508
0.26	546	0.47	504
0.27	542	0.48	500
0.28	539	0.49	500
0.29	539	0.50	497
0.30	536		



# *Naso lituratus*

Orangespine unicornfish, *umaumalei*  
Acanthuridae (surgeonfishes)



## Life history and other input parameters

Parameter	Value	SD	Unit	n	Source
$L_{inf}$	322	36	mm	58	Mean: Nadon (unpublished), SD: Kritzer (2001)
$K$	0.341	0.10	yr <sup>-1</sup>		
$a_0$	-0.66	-	yr		
$L_{mat}$	250	37	mm	66	Mean: Nadon (unpublished), SD: Nadon (unpublished)
Longevity	25	4.2	yr	58	Mean: Nadon (unpublished), SD: Kritzer (2001)
L-W $\alpha$	7.20e-5	-	-	-	Smith and Dalzell (1993)
L-W $\beta$	2.84	-	-	-	
$L_{S50}$	215	-	mm	-	DAR commercial data
$L_{S95}$	230				
$\bar{L}$ diver survey	276	1	mm	1563	NOAA-CREP diver-survey data set
$\bar{L}$ commercial	287	2	mm	1098	DAR commercial data set
$\bar{L}$ combined	280	2	mm	-	-
Max. depth	93	-	m	-	Pyle et al. (2016)
Federal waters	21	-	%	-	-

## Stock status and other output parameters

Parameter	Median	SD	Unit
$M$	0.13	0.02	yr <sup>-1</sup>
$F$	0.21	0.13	yr <sup>-1</sup>
$F_{30}$	0.17	0.04	yr <sup>-1</sup>
$F/F_{30}$	1.3	0.9	-
$SPR$	0.25	0.20	-
$SPR < 0.30$ iterations	62	-	%
$L_{C30}$	215	-	mm

Parameter	Median	SD	Unit
$B$ from catch	30,000	205,000	kg
$B$ from survey	451,619	37,121	kg
Commercial catch	544	1,934	kg
Recreational catch	3,730	7,888	kg
Total catch	4,924	7,317	kg
$C_{30}$ from catch	4,390	27,000	kg
$C_{30}$ from survey	66,200	15,400	kg


## General comments

Population abundance appeared to be relatively stable, with a sharp increase in 2016 which may be spurious.  $L_{bar}$  from both data sets were similar and stable from 2003 to 2016, suggesting a population near equilibrium. The commercial catch increased slightly from 2008 to 2012, but appear to be declining in recent years. The recreational catch was too variable to infer on any temporal patterns.

The life history parameters for this species came from specimens collected in the Mariana islands by PIFSC, but they appeared appropriate for the Hawaii population. A sensitivity run using the stepwise approach gave nearly identical parameters:  $L_{max}$ : 350 mm,  $L_{inf}$ : 321 mm,  $K$ : 0.44,  $M$ : 0.1,  $F_{30}$ : 0.15, and  $F$ : 0.25.


There was a strong discrepancy between the population size estimates from the catch vs. those from diver surveys (30,000 kg vs. 451,619 kg). The catch estimates from both the HMRFS and commercial data sets seemed unrealistically low. The diver-derived population estimate should be fairly reliable given that this is a commonly encountered species that is easily identifiable and is not particularly afraid of divers. Therefore, the  $C_{30}$  from the survey data is likely more reliable than the  $C_{30}$  derived from the catch. The population estimate was likely negatively biased given that the population extends to 93 m, which is beyond the diver survey depth (30 m) from which the population size is derived.

*Naso lituratus*




**Life history parameter distributions.**

*Naso lituratus*




Abundance index from UVS (blue circles, ±SE) and total catch time series from recreational (green squares) and commercial (orange triangles) sectors.




Size structure from commercial catch (top left) and UVS (top right). Average length time series (blue circles – UVS, orange triangles – commercial data, ±SE).

*Naso lituratus*




**Stock status parameter distributions (*SPR*: small bar shows 0.30 level).**

*Naso lituratus*



$C_{30}$  and current total catch (left) and population size (right) distributions.

*Naso lituratus*



**Overfishing probability for a range of  $C_{30}$  levels (catch - orange dashed line, UVS – blue dotted line). *OFLs* are represented by small vertical bars.**

**Probability of overfishing for various  $C_{30}$  levels.**

Overfish. probability	$C_{30}$ from catch (1000 kg)	$C_{30}$ from survey (1000 kg)	Overfish. probability	$C_{30}$ from catch (1000 kg)	$C_{30}$ from survey (1000 kg)
0.10	1.18	48.4	0.31	2.62	59.1
0.11	1.24	49.1	0.32	2.70	59.4
0.12	1.30	49.8	0.33	2.78	59.8
0.13	1.36	50.4	0.34	2.85	60.2
0.14	1.43	51.0	0.35	2.92	60.6
0.15	1.50	51.7	0.36	3.01	61.0
0.16	1.58	52.3	0.37	3.08	61.4
0.17	1.65	52.8	0.38	3.17	61.7
0.18	1.72	53.4	0.39	3.25	62.2
0.19	1.78	53.9	0.40	3.35	62.6
0.20	1.86	54.5	0.41	3.44	62.9
0.21	1.93	54.9	0.42	3.51	63.4
0.22	1.99	55.4	0.43	3.63	63.7
0.23	2.06	55.8	0.44	3.73	64.1
0.24	2.13	56.3	0.45	3.82	64.5
0.25	2.19	56.8	0.46	3.92	64.8
0.26	2.26	57.1	0.47	4.02	65.1
0.27	2.32	57.5	0.48	4.14	65.5
0.28	2.40	57.9	0.49	4.27	65.9
0.29	2.48	58.2	0.50	4.39	66.2
0.30	2.55	58.7			

*Naso lituratus*

**Probability of overfishing at various minimum sizes.**

Overfishing probability	$L_{C_{30}}$ (mm)	Overfishing probability	$L_{C_{30}}$ (mm)
0.10	295	0.31	254
0.11	292	0.32	254
0.12	290	0.33	252
0.13	288	0.34	247
0.14	286	0.35	247
0.15	284	0.36	245
0.16	282	0.37	243
0.17	279	0.38	241
0.18	277	0.39	239
0.19	275	0.40	236
0.20	275	0.41	234
0.21	273	0.42	232
0.22	271	0.43	230
0.23	269	0.44	228
0.24	267	0.45	226
0.25	264	0.46	224
0.26	264	0.47	221
0.27	262	0.48	219
0.28	260	0.49	217
0.29	258	0.50	215
0.30	256		

# *Naso unicornis*

Bluespine unicornfish, *kala*  
Acanthuridae (surgeonfishes)



## Life history and other input parameters

Parameter	Value	SD	Unit	n	Source
$L_{inf}$	480	4	mm	534	Mean and SD: Andrews et al. (2016)
$K$	0.44	0.02	yr <sup>-1</sup>		
$a_0$	-0.12	-	yr		
$L_{mat}$	355	10	mm	295	Mean and SD: Eble (2009)
Longevity	50	5.4	yr	534	Mean and SD: Andrews et al. (2016)
L-W $\alpha$	1.65e-5	-	-	-	Kulbicki (2005)
L-W $\beta$	3.035	-	-	-	
$L_{S50}$	230	-	mm	-	Estimated from HMRFS data and other surgeonfishes.
$L_{S95}$	260				
$\bar{L}$ diver survey	348	5	mm	522	NOAA-CREP diver survey
$\bar{L}$ commercial	-	-	mm	-	DAR commercial data
$\bar{L}$ combined	-	-	mm	-	-
Max. depth	120	-	m	-	BRFA BotCam project
Federal waters	22	-	%	-	-

## Stock status and other output parameters

Parameter	Median	SD	Unit
$M$	0.06	0.01	yr <sup>-1</sup>
$F$	0.60	0.07	yr <sup>-1</sup>
$F_{30}$	0.10	0.01	yr <sup>-1</sup>
$F/F_{30}$	6.0	0.9	-
$SPR$	0.03	0.01	-
$SPR < 0.30$ iterations	100	-	%
$L_{C30}$	467	-	mm

Parameter	Median	SD	Unit
$B$ from catch	-	-	kg
$B$ from survey	364,114	74,939	kg
Commercial catch	11,282	3,350	kg
Recreational catch	8,169	4,829	kg
Total catch	20,197	5,743	kg
$C_{30}$ from catch	-	-	kg
$C_{30}$ from survey	33,200	7,270	kg

## General comments

The commercial data set could not be used to generate  $L_{bar}$  estimates given that *N. annulatus* and *N. brevirostris* are included with *N. unicornis* in the reporting system. The commercial catch for these 3 species is reported in the table above and in the graph further below as reference only.


Population abundance and  $L_{bar}$  were stable from 2005 to 2016, likely indicating a population near equilibrium. The recreational catch was fairly variable, but overall appeared stable as well. The LH parameters came from an in-depth study conducted locally and is likely highly reliable. There was some HMRFS data ( $n = 18$ ) that could be used to help infer the selectivity parameters. A sensitivity run with  $L_{S50}$  at 200 and  $L_{S95}$  at 230 had little impact on the results ( $M$ : 0.11,  $F = 0.54$ ,  $SPR = 0.05$ ).

The elevated  $F/F_{30}$  ratio and correspondingly low  $SPR$  estimate could not be explained by the relatively modest catch estimate, which was below the  $C_{30}$  estimate (20,197 kg vs. 33,100 kg). The reason for this may be the unusually high maximum age recorded for this species (50 year, from a bomb radio-carbon validated aged Oahu specimen). This high longevity leads to an extremely low  $M$  estimate, which in turn leads to a high  $F$  estimate and low  $SPR$ . The 50-year max age could be from an outlier individual, not representative of a 4% cohort survival rate (see equation 7). To test this scenario, we also ran the analyses using  $S = 1\%$  and longevity at 40 years and obtained the following estimates:  $F$ : 0.54,  $SPR$ : 0.05.

Despite this discrepancy, the population estimate from diver surveys should be reliable given that this is a common species that is easily identifiable. The population estimate is likely biased downward given that this species has been recorded to 120-m depths and diver surveys only reach 30-m depths.




*Naso unicornis*




**Life history parameter distributions.**

*Naso unicornis*




**Abundance index from UVS (blue circles,  $\pm$ SE) and total catch time series from recreational (green squares) and commercial (orange triangles) sectors (presented as a reference, see comments section).**




**Size structure and average length time series from UVS ( $\pm$ SE).**

*Naso unicornis*




**Stock status parameter distributions (SPR: small bar shows 0.30 level).**



**$C_{30}$  (left) and population size (right) distributions.**

*Naso unicornis*



**Overfishing probability for a range of  $C_{30}$  levels (UVS – blue dotted line).  
**OFL is represented by a small vertical bar.****

**Probability of overfishing for various  $C_{30}$  levels.**

Overfishing probability	$C_{30}$ from survey (1000 kg)	Overfishing probability	$C_{30}$ from survey (1000 kg)
0.10	24.2	0.31	29.6
0.11	24.5	0.32	29.8
0.12	24.9	0.33	29.9
0.13	25.2	0.34	30.2
0.14	25.5	0.35	30.3
0.15	25.8	0.36	30.5
0.16	26.1	0.37	30.7
0.17	26.4	0.38	30.9
0.18	26.6	0.39	31.1
0.19	26.8	0.40	31.3
0.20	27.1	0.41	31.4
0.21	27.4	0.42	31.7
0.22	27.7	0.43	31.8
0.23	27.9	0.44	32.0
0.24	28.1	0.45	32.2
0.25	28.3	0.46	32.4
0.26	28.5	0.47	32.6
0.27	28.7	0.48	32.8
0.28	28.9	0.49	32.9
0.29	29.1	0.50	33.2
0.30	29.4		

*Naso unicornis*

**Probability of overfishing at various minimum sizes.**

Overfishing probability	$L_{C_{30}}$ (mm)	Overfishing probability	$L_{C_{30}}$ (mm)
0.10	474	0.31	469
0.11	474	0.32	469
0.12	474	0.33	469
0.13	474	0.34	469
0.14	471	0.35	469
0.15	471	0.36	469
0.16	471	0.37	469
0.17	471	0.38	469
0.18	471	0.39	469
0.19	471	0.40	469
0.20	471	0.41	469
0.21	471	0.42	469
0.22	471	0.43	467
0.23	471	0.44	467
0.24	471	0.45	467
0.25	471	0.46	467
0.26	471	0.47	467
0.27	469	0.48	467
0.28	469	0.49	467
0.29	469	0.50	467
0.30	469		

# Carangoides orthogrammus

Island jack, *ulua*  
Carangidae (jacks)



## Life history and other input parameters

Parameter	Value	SD	Unit	n	Source
$L_{inf}$	773	81	mm	-	Mean and SD: Nadon & Ault (2016) $L_{max}$ : 685 (3) from DAR commercial data
$K$	0.290	0.101	yr <sup>-1</sup>		
$a_0$	-0.6	-	yr		
$L_{mat}$	454	58	mm	-	Mean and SD: Nadon & Ault (2016)
Longevity	11	3.4	yr	-	Mean and SD: Nadon & Ault (2016)
L-W $\alpha$	1.29e-5	-	-	-	Kulbicki (2005)
L-W $\beta$	2.994				
$L_{S50}$	325	-	mm	-	DAR commercial data
$L_{S95}$	350				
$\bar{L}$ diver survey	-	-	mm	-	NOAA-CREP diver survey
$\bar{L}$ commercial	502	3	mm	3128	DAR commercial data
$\bar{L}$ combined	-	-	mm	-	-
Max. depth	235	-	m	-	BRFA BotCam project
Federal waters	26	-	%	-	-

## Stock status and other output parameters

Parameter	Median	SD	Unit
$M$	0.27	0.08	yr <sup>-1</sup>
$F$	0.18	0.19	yr <sup>-1</sup>
$F_{30}$	0.26	0.07	yr <sup>-1</sup>
$F/F_{30}$	0.7	0.8	-
$SPR$	0.41	0.25	-
$SPR < 0.30$ iterations	36	-	%
$L_{C30}$	0	-	mm

Parameter	Median	SD	Unit
$B$ from catch	123,000	609,000	kg
$B$ from survey	-	-	kg
Commercial catch	2,686	1,098	kg
Recreational catch	13,162	16,036	kg
Total catch	16,714	14,139	kg
$C_{30}$ from catch	24,300	134,000	kg
$C_{30}$ from survey	-	-	kg


## General comments

There were insufficient UVS observations to generate population or  $L_{bar}$  estimates for this species. Commercial and recreational catches appeared relatively stable from 2003 to 2016, although the recreational catch estimates were highly variable.  $L_{bar}$  estimates from the commercial data set appeared to be slowly rising in the early year of the time series before slowly declining in recent years.

No life history parameters exist for this species and we therefore used the stepwise approach to generate estimates. We used the  $L_{max}$  estimate from the commercial data since the diver estimate was deemed unreliable due to a low observation number. The estimate of  $L_{max}$  for the NWHI from diver surveys was 700 mm (from 95 UVS observations). A sensitivity run using an even larger  $L_{max}$  (720 mm) generated the following LH parameters estimates:  $L_{inf}$ : 798 mm,  $K$ : 0.27,  $M$ : 0.27,  $F$ : 0.18,  $F_{30}$ : 0.25,  $SPR$ : 0.38,  $C_{30}$ : 22,358 kg). These results were fairly similar to the original ones.


Note that the estimated  $L_{inf}$  parameter is higher than the  $L_{max}$  due to the indeterminate growth curve typical of jacks (high  $M/K$  ratio).

*Carangoides orthogrammus*




**Life history parameter distributions.**

*Carangoides orthogrammus*




**Total catch time series from recreational (green squares) and commercial (orange triangles) sectors.**




**Size structure and average length time series from commercial data ( $\pm$ SE)**



*Carangoides orthogrammus*




**Stock status parameter distributions ( $SPR$ : small bar shows 0.30 level).**



**$C_{30}$  and current total catch (left) and population size (right) distributions.**

*Carangoides orthogrammus*



**Overfishing probability for a range of  $C_{30}$  levels (commercial data – orange dashed line). *OFL* is represented by a small vertical bar.**

**Probability of overfishing for various  $C_{30}$  levels.**

Overfishing probability	$C_{30}$ from catch (1000 kg)	Overfishing probability	$C_{30}$ from catch (1000 kg)
0.10	7.0	0.31	14.5
0.11	7.4	0.32	14.8
0.12	7.8	0.33	15.1
0.13	8.1	0.34	15.5
0.14	8.4	0.35	16.0
0.15	8.8	0.36	16.4
0.16	9.1	0.37	16.9
0.17	9.4	0.38	17.5
0.18	9.8	0.39	17.9
0.19	10.2	0.40	18.3
0.20	10.5	0.41	18.9
0.21	10.9	0.42	19.3
0.22	11.2	0.43	19.9
0.23	11.6	0.44	20.4
0.24	11.9	0.45	21.0
0.25	12.2	0.46	21.6
0.26	12.4	0.47	22.3
0.27	12.9	0.48	22.9
0.28	13.2	0.49	23.6
0.29	13.7	0.50	24.3
0.30	14.1		

*Carangoides orthogrammus*

**Probability of overfishing at various minimum sizes.**

Overfishing probability	$L_{C_{30}}$ (mm)	Overfishing probability	$L_{C_{30}}$ (mm)
0.10	539	0.31	358
0.11	530	0.32	348
0.12	520	0.33	338
0.13	510	0.34	325
0.14	500	0.35	318
0.15	494	0.36	306
0.16	484	0.37	292
0.17	478	0.38	276
0.18	471	0.39	266
0.19	462	0.40	250
0.20	452	0.41	240
0.21	442	0.42	224
0.22	436	0.43	205
0.23	426	0.44	188
0.24	419	0.45	169
0.25	413	0.46	150
0.26	406	0.47	118
0.27	393	0.48	0
0.28	387	0.49	0
0.29	377	0.50	0
0.30	367		

# *Caranx ignobilis*

Giant trevally, *ulua aukea*

Carangidae (jacks)



## Life history and other input parameters

Parameter	Value	SD	Unit	n	Source
$L_{inf}$	2170	310	mm	10	Mean and SD: Sudekum (1991)
$K$	0.111	0.02	yr <sup>-1</sup>		
$a_0$	0.097	-	yr		
$L_{mat}$	839	31	mm	10	Mean and SD: Sudekum (1991)
Longevity	11	3	yr	10	Mean and SD: Sudekum (1991)
L-W $\alpha$	2.22e-5	-	-	-	Seki (1986)
L-W $\beta$	2.913	-	-	-	
$L_{S50}$	350	-	mm	-	DAR commercial data
$L_{S95}$	430				
$\bar{L}$ diver survey	-	-	mm	-	-
$\bar{L}$ commercial	761	6	mm	5372	DAR commercial data
$\bar{L}$ combined	-	-	mm	-	-
Max. depth	228	-	m	-	BRFA BotCam project
Federal waters	26	-	%	-	-

## Stock status and other output parameters

Parameter	Median	SD	Unit
$M$	0.26	0.05	yr <sup>-1</sup>
$F$	0.18	0.10	yr <sup>-1</sup>
$F_{30}$	0.17	0.03	yr <sup>-1</sup>
$F/F_{30}$	1.1	0.7	-
$SPR$	0.28	0.22	-
$SPR < 0.30$ iterations	54	-	%
$L_{C30}$	430	-	mm


Parameter	Median	SD	Unit
$B$ from catch	1,070,000	3,930,000	kg
$B$ from survey	-	-	kg
Commercial catch	6,689	2,079	kg
Recreational catch	133,921	117,016	kg
Total catch	142,429	113,530	kg
$C_{30}$ from catch	147,000	564,000	kg
$C_{30}$ from survey	-	-	kg

## General comments

There were insufficient diver observations in the MHI to generate a reliable  $L_{bar}$  estimate. The catch was dominated by the recreational sector and, although fairly variable from year to year, it appear to be mostly constant.  $L_{bar}$  from the commercial catch were also stable throughout the period under consideration, suggesting equilibrium conditions.


The life history parameters for this species came from a local study with a very limited sample size. We also ran the analyses using the stepwise approach and generated the following values:  $L_{max}$ : 1351 mm,  $L_{inf}$ : 1679 mm,  $K$ : 0.17,  $M$ : 0.21,  $F_{30}$ : 0.15,  $F$ : 0.21,  $SPR$ : 0.19. These are fairly similar to the original analyses and did not change the conclusions.

*Caranx ignobilis*





**Life history parameter distributions.**

*Caranx ignobilis*




**Total catch time series from recreational (green squares) and commercial (orange triangles) sectors.**




**Size structure and average length time series from commercial data ( $\pm$ SE)**

*Caranx ignobilis*



**Stock status parameter distributions ( $SPR$ : small bar shows 0.30 level).**


*Caranx ignobilis*



$C_{30}$  and current total catch (left) and population size (right) distributions.



*Caranx ignobilis*



**Overfishing probability for a range of  $C_{30}$  levels (commercial data – orange dashed line). *OFL* is represented by a small vertical bar.**

**Probability of overfishing for various  $C_{30}$  levels.**

Overfishing probability	$C_{30}$ from catch (1000 kg)	Overfishing probability	$C_{30}$ from catch (1000 kg)
0.10	51.8	0.31	96.9
0.11	54.0	0.32	99.2
0.12	56.3	0.33	102.0
0.13	59.3	0.34	104.1
0.14	61.6	0.35	106.4
0.15	63.8	0.36	108.8
0.16	65.7	0.37	111.3
0.17	67.8	0.38	113.9
0.18	70.2	0.39	116.6
0.19	71.8	0.40	119.4
0.20	73.9	0.41	121.9
0.21	76.2	0.42	124.2
0.22	77.8	0.43	126.5
0.23	80.0	0.44	129.1
0.24	82.3	0.45	132.7
0.25	84.6	0.46	135.2
0.26	86.6	0.47	138.4
0.27	88.7	0.48	141.5
0.28	90.4	0.49	144.1
0.29	92.6	0.50	147.0
0.30	94.9		

*Caranx ignobilis*

**Probability of overfishing at various minimum sizes.**

Overfishing probability	$L_{C_{30}}$ (mm)	Overfishing probability	$L_{C_{30}}$ (mm)
0.10	959	0.31	710
0.11	945	0.32	696
0.12	931	0.33	686
0.13	917	0.34	679
0.14	906	0.35	668
0.15	892	0.36	654
0.16	878	0.37	640
0.17	864	0.38	626
0.18	858	0.39	616
0.19	844	0.40	602
0.20	833	0.41	584
0.21	819	0.42	570
0.22	808	0.43	556
0.23	798	0.44	536
0.24	788	0.45	518
0.25	777	0.46	504
0.26	763	0.47	486
0.27	752	0.48	466
0.28	742	0.49	452
0.29	732	0.50	430
0.30	721		

# Caranx melampygu

Bluefin trevally, 'omilu

Carangidae (jacks)



## Life history and other input parameters

Parameter	Value	SD	Unit	n	Source
$L_{inf}$	1041	174	mm	14	Mean: Sudekum (1991), SD: Kritzer (2001)
$K$	0.233	0.13	yr <sup>-1</sup>		
$a_0$	-0.044	-	yr		
$L_{mat}$	476	33	mm	14	Mean: Sudekum (1991), SD: Nadon (unpublished)
Longevity	7	1.3	yr	14	Mean: Sudekum (1991), SD: Kritzer (2001)
L-W $\alpha$	2.38e-5	-	-	-	Seki (1986)
L-W $\beta$	2.94	-	-	-	
$L_{S50}$	325	-	mm	-	DAR commercial data
$L_{S95}$	370				
$\bar{L}$ diver survey	465	9	mm	169	NOAA-CREP diver survey
$\bar{L}$ commercial	557	3	mm	4501	DAR commercial data
$\bar{L}$ combined	550	3	mm	-	-
Max. depth	230	-	m	-	Pyle et al. (2016)
Federal waters	26	-	%	-	-

## Stock status and other output parameters

Parameter	Median	SD	Unit
$M$	0.44	0.07	yr <sup>-1</sup>
$F$	0.27	0.21	yr <sup>-1</sup>
$F_{30}$	0.37	0.06	yr <sup>-1</sup>
$F/F_{30}$	0.7	0.6	-
$SPR$	0.40	0.23	-
$SPR < 0.30$ iterations	34	-	%
$L_{C30}$	88	-	mm


Parameter	Median	SD	Unit
$B$ from catch	811,000	3,500,000	kg
$B$ from survey	-	-	kg
Commercial catch	3,147	718	kg
Recreational catch	144,775	50,248	kg
Total catch	148,127	50,035	kg
$C_{30}$ from catch	205,000	911,000	kg
$C_{30}$ from survey	-	-	kg

## General comments

Population abundance appeared to be slowly increasing in recent years. There were enough diver observations to obtain an  $L_{bar}$  estimate, but the fast swimming behavior of this species made estimating population size unreliable.  $L_{bar}$  from the UVS were not abundant enough on a yearly basis to generate a long term pattern.  $L_{bar}$  from the commercial data appeared fairly stable, which suggested that the population was mostly at equilibrium. The recreational sector dominated the catch and appeared fairly constant, although variable from year to year.


As for *C. ignobilis*, the life history parameters for this species came from a local study with a limited sample size. We also ran the analyses using the stepwise approach and generated the following values:  $L_{max}$ : 843 mm,  $L_{inf}$ : 921 mm,  $K$ : 0.25,  $M$ : 0.26,  $F_{30}$ : 0.22,  $F$ : 0.22,  $SPR$ : 0.32,  $C_{30}$ : 141,420 kg. These were reasonably similar to the original analyses and did not change the conclusions.

*Caranx melampygius*




**Life history parameter distributions.**

*Caranx melampygus*




**Abundance index from UVS (blue circles,  $\pm$ SE) and total catch time series from recreational (green squares) and commercial (orange triangles) sectors.**




**Size structure from commercial catch (top left) and UVS (top right). Average length time series (blue circles – UVS, orange triangles – commercial data,  $\pm$ SE).**

*Caranx melampygus*




**Stock status parameter distributions (*SPR*: small bar shows 0.30 level).**

*Caranx melampyus*



$C_{30}$  and current total catch (left) and population size (right) distributions.

*Caranx melampygus*



**Overfishing probability for a range of  $C_{30}$  levels (commercial data – orange dashed line). *OFL* is represented by a small vertical bar.**

**Probability of overfishing for various  $C_{30}$  levels.**

Overfishing probability	$C_{30}$ from catch (1000 kg)	Overfishing probability	$C_{30}$ from catch (1000 kg)
0.10	92.1	0.31	143.7
0.11	95.6	0.32	146.4
0.12	98.2	0.33	149.0
0.13	100.4	0.34	151.5
0.14	102.5	0.35	153.8
0.15	104.9	0.36	156.7
0.16	107.5	0.37	159.6
0.17	110.4	0.38	162.7
0.18	112.7	0.39	165.0
0.19	115.4	0.40	167.7
0.20	117.5	0.41	170.9
0.21	119.4	0.42	174.2
0.22	121.6	0.43	177.2
0.23	123.6	0.44	181.3
0.24	126.0	0.45	185.4
0.25	128.4	0.46	188.6
0.26	130.7	0.47	192.7
0.27	132.7	0.48	196.5
0.28	135.6	0.49	200.0
0.29	138.1	0.50	205.3
0.30	141.4		



*Caranx melampyus*

**Probability of overfishing at various minimum sizes.**

Overfishing probability	$L_{C_{30}}$ (mm)	Overfishing probability	$L_{C_{30}}$ (mm)
0.10	536	0.31	354
0.11	526	0.32	344
0.12	517	0.33	332
0.13	507	0.34	325
0.14	500	0.35	315
0.15	491	0.36	302
0.16	484	0.37	292
0.17	474	0.38	280
0.18	465	0.39	267
0.19	458	0.40	254
0.20	452	0.41	240
0.21	445	0.42	224
0.22	436	0.43	211
0.23	426	0.44	198
0.24	416	0.45	182
0.25	410	0.46	166
0.26	400	0.47	146
0.27	393	0.48	130
0.28	384	0.49	107
0.29	374	0.50	88
0.30	364		

# Aprion virescens

Green jobfish, *uku*

Lutjanidae (snappers)



## Life history and other input parameters

Parameter	Value	SD	Unit	n	Source
$L_{inf}$	771	50	mm	379	Mean and SD: O'Malley (unpubl.)
$K$	0.372	0.021	yr <sup>-1</sup>		
$a_0$	-0.51	-	yr		
$L_{mat}$	489	16	mm	103	Mean: Everson (1989), SD: Nadon (unpublished)
Longevity	31	2.3	yr	379	Mean and SD: O'Malley (unpubl.)
L-W $\alpha$	6.44e-4	-	-	-	HMFRS data
L-W $\beta$	2.404	-	-	-	
$L_{S50}$	425	-	mm	-	DAR commercial data
$L_{S95}$	475				
$\bar{L}$ diver survey	613	10	mm	278	NOAA-CREP diver survey
$\bar{L}$ commercial	665	1	mm	17634	DAR commercial data
$\bar{L}$ combined	664	1	mm	-	-
Max. depth	203	-	m	-	BRFA BotCam project
Federal waters	24	-	%	-	-

## Stock status and other output parameters

Parameter	Median	SD	Unit
$M$	0.10	0.01	yr <sup>-1</sup>
$F$	0.15	0.07	yr <sup>-1</sup>
$F_{30}$	0.16	0.01	yr <sup>-1</sup>
$F/F_{30}$	0.9	0.5	-
$SPR$	0.33	0.16	-
$SPR < 0.30$ iterations	42	-	%
$L_{C30}$	349	-	mm

Parameter	Median	SD	Unit
$B$ from catch	758,000	1,590,000	kg
$B$ from survey	434,419	83,661	kg
Commercial catch	36,867	11,226	kg
Recreational catch	51,001	38,316	kg
Total catch	92,248	36,805	kg
$C_{30}$ from catch	104,000	226,000	kg
$C_{30}$ from survey	60,000	12,100	kg

## General comments


Population abundance appeared to be increasing from 2003 to 2016. The commercial catch also appeared to be increasing although it has been flat in the last 2 years. Recreational catch, although fairly variable, appeared to be relatively stable.  $L_{bar}$  from both the commercial and UVS data sets were similar and mostly stable, suggesting that the population was mostly at equilibrium.

The life history parameters for this species came from a local study conducted by the PIFSC life history group (J. O'Malley) with a fairly large sample size. Length at maturity came from a local study as well. Given the reliability of these sources, we did not deem it necessary to run the stepwise analyses for this species.

The diver  $L_{bar}$  was lower than the commercial one. A sensitivity run using this  $L_{bar}$  generated the following results:  $F$ : 0.33,  $F/F_{30}$ : 2.1,  $SPR$ : 0.15. Another sensitivity run using selectivity parameters of 475 mm and 525 mm resulted in the following values:  $F$ : 0.18,  $F/F_{30}$ : 1.1,  $SPR$ : 0.30.


There were also sufficient diver observations to obtain a population size estimate (434,419 kg) which was relatively close to the one derived from the total catch estimate (758,000 kg). The lower population estimate from diver surveys could be related to the depth limitation of these surveys compared to the full population range for this species (30 m vs. 203 m).

*Aprion virescens*




**Life history parameter distributions.**

*Aprion virescens*




**Abundance index from UVS (blue circles,  $\pm$ SE) and total catch time series from recreational (green squares) and commercial (orange triangles) sectors.**




**Size structure from commercial catch (top left) and UVS (top right). Average length time series (blue circles – UVS, orange triangles – commercial data,  $\pm$ SE).**

*Aprion virescens*




**Stock status parameter distributions (*SPR*: small bar shows 0.30 level).**

*Aprion virescens*



$C_{30}$  and current total catch (left) and population size (right) distributions.

*Aprion virescens*



**Overfishing probability for a range of  $C_{30}$  levels (catch - orange dashed line, UVS – blue dotted line). *OFLs* are represented by small vertical bars.**

**Probability of overfishing for various  $C_{30}$  levels.**

Overfish. probability	$C_{30}$ from catch (1000 kg)	$C_{30}$ from survey (1000 kg)	Overfish. probability	$C_{30}$ from catch (1000 kg)	$C_{30}$ from survey (1000 kg)
0.10	51.3	44.6	0.31	78.3	54.0
0.11	52.9	45.4	0.32	79.1	54.4
0.12	54.0	46.1	0.33	80.2	54.7
0.13	55.0	46.7	0.34	81.4	55.1
0.14	56.4	47.3	0.35	82.6	55.5
0.15	57.8	47.7	0.36	84.0	55.7
0.16	59.1	48.2	0.37	85.3	56.1
0.17	60.5	48.6	0.38	86.4	56.4
0.18	61.8	49.0	0.39	87.7	56.6
0.19	63.2	49.4	0.40	89.3	57.0
0.20	64.8	49.8	0.41	90.5	57.4
0.21	66.1	50.2	0.42	91.8	57.7
0.22	67.3	50.7	0.43	93.3	58.0
0.23	68.4	51.2	0.44	94.9	58.4
0.24	69.5	51.5	0.45	96.3	58.6
0.25	70.6	51.9	0.46	97.9	58.9
0.26	71.7	52.3	0.47	99.4	59.2
0.27	73.0	52.6	0.48	100.9	59.5
0.28	74.4	53.0	0.49	102.4	59.8
0.29	75.8	53.3	0.50	104.3	60.0
0.30	76.9	53.7			

*Aprion virescens*

**Probability of overfishing at various minimum sizes.**

Overfishing probability	$L_{C_{30}}$ (mm)	Overfishing probability	$L_{C_{30}}$ (mm)
0.10	625	0.31	502
0.11	616	0.32	497
0.12	608	0.33	489
0.13	604	0.34	480
0.14	595	0.35	476
0.15	591	0.36	468
0.16	586	0.37	463
0.17	582	0.38	455
0.18	574	0.39	446
0.19	570	0.40	438
0.20	565	0.41	429
0.21	557	0.42	425
0.22	552	0.43	421
0.23	548	0.44	408
0.24	540	0.45	404
0.25	536	0.46	391
0.26	531	0.47	378
0.27	523	0.48	370
0.28	518	0.49	357
0.29	514	0.50	348
0.30	506		



# *Lutjanus fulvus*

Blacktail snapper, *to'au*

Lutjanidae (snappers)



## Life history and other input parameters

Parameter	Value	SD	Unit	n	Source
$L_{inf}$	342	19	mm	-	Mean and SD: Nadon & Ault (2016) $L_{max}$ : 372 (7) from MHI diver data
$K$	0.43	0.17	yr <sup>-1</sup>		
$a_0$	-0.6	-	yr		
$L_{mat}$	240	37	mm	-	Mean and SD: Nadon & Ault (2016)
Longevity	19	7	yr	-	Mean and SD: Nadon & Ault (2016)
L-W $\alpha$	2.04e-5	-	-	-	Kulbicki (2005)
L-W $\beta$	2.97				
$L_{S50}$	210	-	mm	-	DAR commercial data
$L_{S95}$	220				
$\bar{L}$ diver survey	264	3	mm	471	NOAA-CREP diver survey
$\bar{L}$ commercial	295	2	mm	4771	DAR commercial data
$\bar{L}$ combined	283	2	mm	-	-
Max. depth	128	-	m	-	Pyle et al. (2016)
Federal waters	22	-	%	-	-

## Stock status and other output parameters

Parameter	Median	SD	Unit
$M$	0.17	0.07	yr <sup>-1</sup>
$F$	0.20	0.18	yr <sup>-1</sup>
$F_{30}$	0.22	0.09	yr <sup>-1</sup>
$F/F_{30}$	0.9	0.9	-
$SPR$	0.33	0.23	-
$SPR < 0.30$ iterations	45	-	%
$L_{C30}$	132	-	mm

Parameter	Median	SD	Unit
B from catch	47,900	161,000	kg
B from survey	180,433	29,498	kg
Commercial catch	1,606	832	kg
Recreational catch	6,682	2,358	kg
Total catch	8,488	2,488	kg
$C_{30}$ from catch	8,670	36,000	kg
$C_{30}$ from survey	33,100	11,400	kg

## General comments


Note: this species is non-native and considered invasive. It was introduced in 1956 from Tahiti.

Population abundance for this species has been stable. The commercial  $L_{bar}$  have been stable as well, while the recreational  $L_{bar}$  have declined a bit from the early years. The recreational catch was much larger than the commercial one and both have been steady in recent years (with the recreational catch declining slightly in the last 2 years).

There is no published life history parameters for this species. We therefore used the stepwise approach to generate estimates, using an  $L_{max}$  value of 372 mm from diver survey data (from the MHI, since this species is not present in the NWHI). A sensitivity run with a higher  $L_{max}$  (400 mm), generated the following estimates:  $L_{inf}$ : 364 mm,  $K$ : 0.40,  $M$ : 0.17,  $F_{30}$ : 0.21,  $F$ : 0.32,  $C_{30}$  survey: 32,359 kg).


The population size estimate derived from the catch was lower than the estimate from diver survey. This could be expected given the uncertainty in both the total catch estimate and the life history parameters. The population size estimate derived from the diver survey is likely more reliable given that this is a common species that is easily identifiable. Consequently, the  $C_{30}$  derived from the diver survey population estimate is likely more reliable.

*Lutjanus fulvus*




**Life history parameter distributions.**

*Lutjanus fulvus*




**Abundance index from UVS (blue circles,  $\pm$ SE) and total catch time series from recreational (green squares) and commercial (orange triangles) sectors.**




**Size structure from commercial catch (top left) and UVS (top right). Average length time series (blue circles – UVS, orange triangles – commercial data,  $\pm$ SE).**

*Lutjanus fulvus*




**Stock status parameter distributions (*SPR*: small bar shows 0.30 level).**

*Lutjanus fulvus*



$C_{30}$  and current total catch (left) and population size (right) distributions.

*Lutjanus fulvus*



**Overfishing probability for a range of  $C_{30}$  levels (catch - orange dashed line, UVS – blue dotted line). *OFLs* are represented by small vertical bars.**

**Probability of overfishing for various  $C_{30}$  levels.**

Overfish. probability	$C_{30}$ from catch (1000 kg)	$C_{30}$ from survey (1000 kg)	Overfish. probability	$C_{30}$ from catch (1000 kg)	$C_{30}$ from survey (1000 kg)
0.10	3.5	21.3	0.31	5.9	28.0
0.11	3.6	21.7	0.32	6.0	28.2
0.12	3.7	22.1	0.33	6.1	28.5
0.13	3.8	22.5	0.34	6.2	28.8
0.14	3.9	22.8	0.35	6.4	29.0
0.15	4.0	23.2	0.36	6.5	29.3
0.16	4.1	23.6	0.37	6.6	29.6
0.17	4.2	24.0	0.38	6.8	29.8
0.18	4.3	24.3	0.39	6.9	30.2
0.19	4.4	24.6	0.40	7.1	30.5
0.20	4.5	24.9	0.41	7.3	30.8
0.21	4.6	25.2	0.42	7.4	31.1
0.22	4.8	25.5	0.43	7.5	31.3
0.23	4.9	25.8	0.44	7.7	31.6
0.24	5.0	26.0	0.45	7.8	31.9
0.25	5.2	26.3	0.46	8.0	32.2
0.26	5.3	26.5	0.47	8.1	32.5
0.27	5.4	26.8	0.48	8.3	32.8
0.28	5.5	27.0	0.49	8.5	33.0
0.29	5.6	27.4	0.50	8.7	33.3
0.30	5.7	27.7			

*Lutjanus fulvus*

**Probability of overfishing at various minimum sizes.**

Overfishing probability	$L_{C_{30}}$ (mm)	Overfishing probability	$L_{C_{30}}$ (mm)
0.10	294	0.31	231
0.11	290	0.32	229
0.12	288	0.33	225
0.13	284	0.34	220
0.14	281	0.35	218
0.15	279	0.36	216
0.16	275	0.37	210
0.17	273	0.38	208
0.18	271	0.39	202
0.19	268	0.40	197
0.20	265	0.41	193
0.21	263	0.42	187
0.22	260	0.43	181
0.23	258	0.44	174
0.24	254	0.45	168
0.25	250	0.46	162
0.26	248	0.47	156
0.27	246	0.48	147
0.28	242	0.49	141
0.29	239	0.50	132
0.30	235		

# *Lutjanus kasmira*

Bluestripe snapper, *ta'ape*

Lutjanidae (snappers)



## Life history and other input parameters

Parameter	Value	SD	Unit	n	Source
$L_{inf}$	340	15	mm	171	Mean: Morales-Nin and Ralston (1990), SD: Kritzer (2001)
$K$	0.29	0.05	yr <sup>-1</sup>		
$a_0$	-1.37	-	yr		
$L_{mat}$	200	5	mm	100?	Mean: Allen (1985), SD: Nadon (unpublished)
Longevity	8	1	yr	171	Mean: Loubens (1980), SD: Kritzer (2001)
L-W $\alpha$	4.30e-6	-	-	-	Kulbicki (2001)
L-W $\beta$	3.25	-	-	-	
$L_{S50}$	220	-	mm	-	DAR commercial data
$L_{S95}$	240				
$\bar{L}$ diver survey	274	4	mm	1402	NOAA-CREP diver survey
$\bar{L}$ commercial	271	1	mm	7460	DAR commercial data
$\bar{L}$ combined	271	1	mm	-	-
Max. depth	265	-	m	-	Pyle et al. (2016)
Federal waters	26	-	%	-	-

## Stock status and other output parameters

Parameter	Median	SD	Unit
$M$	0.39	0.04	yr <sup>-1</sup>
$F$	0.21	0.13	yr <sup>-1</sup>
$F_{30}$	0.75	0.14	yr <sup>-1</sup>
$F/F_{30}$	0.3	0.2	-
$SPR$	0.62	0.16	-
$SPR < 0.30$ iterations	0	-	%
$L_{C30}$	0	-	mm

Parameter	Median	SD	Unit
$B$ from catch	181,000	576,000	kg
$B$ from survey	496,235	98,182	kg
Commercial catch	16,165	2,813	kg
Recreational catch	10,734	7,534	kg
Total catch	27,968	7,429	kg
$C_{30}$ from catch	80,100	288,000	kg
$C_{30}$ from survey	221,000	50,000	kg

## General comments

Note: this species is non-native and considered invasive. It was introduced in 1956 from Tahiti.


Population abundance appears to have declined from a peak in 2005. However, the survey effort in this early year had a limited sample size and this may simply be an outlier. Abundance appeared to be fairly stable for the rest of the time series.  $L_{bar}$  from both data sets were very similar and constant throughout the years, which suggest a population near equilibrium. The commercial catch has been declining in recent years while the recreational catch has remained highly variable but stable.

The growth parameters for this species came from a local study with a proper sample size. However, the maturity parameter source was not ideal (i.e. a fish ID guide). The stepwise approach for this species provided the following results:  $L_{max}$ : 340 mm,  $L_{inf}$ : 316 mm,  $L_{mat}$ : 224,  $K$ : 0.45,  $M$ : 0.20,  $F$ : 0.37,  $F_{30}$ : 0.30,  $SPR$ : 0.23,  $C_{30}$  survey: 118,185 kg. The  $L_{mat}$  value obtained from the stepwise approach was close to the original one. The rest of the parameters were similar except for a higher longevity and thus higher  $M$ . The max age for this species is estimated at 8 years while snapper typically live longer (16 years). This led to the higher  $F$  value and low  $SPR$  value.

The population size estimate derived from the catch was less than half the estimate from diver surveys. This can be expected given the uncertainty in the total catch estimate. The population size estimate derived from the diver survey is likely more reliable given that this is a common species that is easily identifiable. The  $OFL$  derived from the diver survey population estimate is therefore likely more reliable.




*Lutjanus kasmira*




**Life history parameter distributions.**

*Lutjanus kasmira*




**Abundance index from UVS (blue circles,  $\pm$ SE) and total catch time series from recreational (green squares) and commercial (orange triangles) sectors.**



**Size structure from commercial catch (top left) and UVS (top right). Average length time series (blue circles – UVS, orange triangles – commercial data,  $\pm$ SE).**


*Lutjanus kasmira*




**Stock status parameter distributions (*SPR*: small bar shows 0.30 level).**

.

*Lutjanus kasmira*



$C_{30}$  and current total catch (left) and population size (right) distributions.



**Overfishing probability for a range of  $C_{30}$  levels (catch - orange dashed line, UVS – blue dotted line).  $OFLs$  are represented by small vertical bars.**

**Probability of overfishing for various  $C_{30}$  levels.**

Overfish. probability	$C_{30}$ from catch (1000 kg)	$C_{30}$ from survey (1000 kg)	Overfish. probability	$C_{30}$ from catch (1000 kg)	$C_{30}$ from survey (1000 kg)
0.10	36.6	161.4	0.31	57.2	197.0
0.11	37.7	163.5	0.32	58.1	198.0
0.12	38.8	165.5	0.33	59.1	199.6
0.13	39.9	167.3	0.34	60.0	200.8
0.14	41.1	169.1	0.35	61.1	202.2
0.15	42.3	171.2	0.36	62.1	203.4
0.16	43.3	173.7	0.37	63.0	204.5
0.17	44.1	175.5	0.38	64.1	205.6
0.18	45.0	177.2	0.39	65.0	206.8
0.19	45.9	179.1	0.40	66.2	208.5
0.20	46.7	180.9	0.41	67.3	209.7
0.21	47.6	182.5	0.42	68.4	210.9
0.22	48.4	184.0	0.43	69.8	212.1
0.23	49.5	185.8	0.44	71.1	213.4
0.24	50.6	187.2	0.45	72.3	214.6
0.25	51.5	188.7	0.46	73.7	215.7
0.26	52.4	190.2	0.47	75.4	216.9
0.27	53.4	191.5	0.48	77.1	218.2
0.28	54.3	193.0	0.49	78.6	219.4
0.29	55.2	194.2	0.50	80.1	220.5
0.30	56.3	195.7			

*Lutjanus kasmira*

**Probability of overfishing at various minimum sizes.**

Overfishing probability	$L_{C_{30}}$ (mm)	Overfishing probability	$L_{C_{30}}$ (mm)
0.10	172	0.31	0
0.11	169	0.32	0
0.12	167	0.33	0
0.13	163	0.34	0
0.14	161	0.35	0
0.15	158	0.36	0
0.16	156	0.37	0
0.17	152	0.38	0
0.18	150	0.39	0
0.19	146	0.40	0
0.20	143	0.41	0
0.21	141	0.42	0
0.22	136	0.43	0
0.23	134	0.44	0
0.24	130	0.45	0
0.25	128	0.46	0
0.26	123	0.47	0
0.27	119	0.48	0
0.28	112	0.49	0
0.29	0	0.50	0
0.30	0		

# Mulloidichthys flavolineatus

Yellowstripe goatfish, *weke'a*

Mullidae (goatfishes)



## Life history and other input parameters

Parameter	Value	SD	Unit	n	Source
$L_{inf}$	371	41	mm	50	Mean: Holland (1993), SD: Kritzer (2001)
$K$	0.564	0.170	yr <sup>-1</sup>		
$a_0$	-0.36	-	yr		
$L_{mat}$	199	7	mm	95	Mean: Cole (2009), SD: Nadon (unpublished)
Longevity	6	2	yr	-	Mean: Estimated, SD: estimated
L-W $\alpha$	4.12e-6	-	-	-	Holland (1993)
L-W $\beta$	3.21				
$L_{S50}$	215	-	mm	-	DAR commercial data
$L_{S95}$	230				
$\bar{L}$ diver survey	281	10	mm	414	NOAA-CREP diver survey
$\bar{L}$ commercial	293	5	mm	2035	DAR commercial data
$\bar{L}$ combined	291	6	mm	-	-
Max. depth	97	-	m	-	Pyle et al. (2016)
Federal waters	21	-	%	-	-

## Stock status and other output parameters

Parameter	Median	SD	Unit
$M$	0.46	0.12	yr <sup>-1</sup>
$F$	0.33	0.26	yr <sup>-1</sup>
$F_{30}$	0.64	0.18	yr <sup>-1</sup>
$F/F_{30}$	0.5	0.5	-
$SPR$	0.49	0.22	-
$SPR < 0.30$ iterations	18	-	%
$L_{C30}$	0	-	mm

Parameter	Median	SD	Unit
$B$ from catch	307,000	1,840,000	kg
$B$ from survey	42,455	14,919	kg
Commercial catch	2,152	843	kg
Recreational catch	61,857	68,173	kg
Total catch	64,806	65,901	kg
$C_{30}$ from catch	118,000	787,000	kg
$C_{30}$ from survey	16,300	6,260	kg

## General comments


Population abundance from UVS was fairly variable, but overall appeared constant. The commercial catch was relatively small and stable. The recreational catch was very high but this is likely an artefact of the HMRFS sampling protocol (see discussion below).  $L_{bar}$  from both data sets were similar and appear to be slightly increasing in recent years.

The growth parameters for this species came from a local mark-recapture study. There is, however, no estimate of maximum age. Longevity for this species is estimated at around 6 years, which is the average for the goatfish family. A sensitivity run with longevity set at the highest value recorded for goatfishes (11 years) gave the following estimates:  $M$ : 0.29,  $F_{30}$ : 0.41,  $SPR$ : 0.30,  $C_{30}$  survey: 12,487 kg.

The recreational catch for this species is 25× larger than the commercial catch, which is highly unusual (i.e., the next highest catch ratio for goatfishes is 5× for *P. cyclostomus*). There is a fishery for the juvenile of this species (called oama). It is very likely that HMRFS surveyors do not often measure lengths or weights when encountering oama (Hongguang Ma, pers. comm.), but still record counts. This likely biases the average weight upward. Since average weight multiplied by number recorded is used to infer total catch by weight, the total recreational catch estimate is very likely positively biased. If we multiply the commercial catch by the (continued on next page)

(continued from previous page)


average recreational to commercial catch ratio for goatfishes (4), we get an estimated recreational catch of 8,608 kg, which, combined with the commercial catch, provides a population biomass estimate of 47,164 kg. This value is close to the population biomass estimate from diver surveys (42,455 kg). Given this issue, the diver survey  $C_{30}$  estimates should be used for this species. The “corrected” total catch estimate of 10,760 kg is significantly below the  $OFL$  value of 16,300 kg which is concordant with the  $SPR$  value of 0.49.




**Life history parameter distributions.**



*Mulloidichthys flavolineatus*




**Abundance index from UVS (blue circles,  $\pm$ SE) and total catch time series from recreational (green squares) and commercial (orange triangles) sectors.**




**Size structure from commercial catch (top left) and UVS (top right). Average length time series (blue circles – UVS, orange triangles – commercial data,  $\pm$ SE).**

*Mulloidichthys flavolineatus*




**Stock status parameter distributions (*SPR*: small bar shows 0.30 level).**

*Mulloidichthys flavolineatus*



$C_{30}$  and current total catch (left) and population size (right) distributions.

*Mulloidichthys flavolineatus*



**Overfishing probability for a range of  $C_{30}$  levels (catch - orange dashed line, UVS – blue dotted line). *OFLs* are represented by small vertical bars.**

**Probability of overfishing for various  $C_{30}$  levels.**

Overfish. probability	$C_{30}$ from catch (1000 kg)	$C_{30}$ from survey (1000 kg)	Overfish. probability	$C_{30}$ from catch (1000 kg)	$C_{30}$ from survey (1000 kg)
0.10	34.9	8.8	0.31	70.5	13.2
0.11	36.6	9.1	0.32	72.3	13.4
0.12	38.2	9.3	0.33	74.3	13.6
0.13	39.6	9.6	0.34	76.2	13.8
0.14	40.9	9.9	0.35	78.5	14.0
0.15	42.4	10.1	0.36	80.7	14.2
0.16	44.3	10.3	0.37	83.1	14.3
0.17	45.8	10.5	0.38	85.2	14.5
0.18	47.3	10.7	0.39	87.6	14.7
0.19	48.9	10.9	0.40	89.7	14.8
0.20	50.3	11.2	0.41	92.3	15.0
0.21	52.0	11.4	0.42	95.0	15.1
0.22	54.0	11.6	0.43	98.4	15.3
0.23	55.5	11.8	0.44	100.6	15.4
0.24	57.4	12.0	0.45	103.8	15.6
0.25	59.7	12.2	0.46	106.9	15.7
0.26	61.4	12.4	0.47	109.8	15.9
0.27	62.7	12.6	0.48	112.6	16.0
0.28	64.8	12.8	0.49	115.2	16.1
0.29	66.6	12.9	0.50	118.3	16.3
0.30	68.2	13.1			

*Mulloidichthys flavolineatus*

**Probability of overfishing at various minimum sizes.**

Overfishing probability	$L_{C_{30}}$ (mm)	Overfishing probability	$L_{C_{30}}$ (mm)
0.10	245	0.31	161
0.11	241	0.32	157
0.12	234	0.33	153
0.13	230	0.34	148
0.14	226	0.35	142
0.15	224	0.36	138
0.16	219	0.37	131
0.17	215	0.38	127
0.18	211	0.39	118
0.19	209	0.40	114
0.20	204	0.41	108
0.21	200	0.42	101
0.22	198	0.43	92
0.23	194	0.44	84
0.24	189	0.45	64
0.25	185	0.46	0
0.26	181	0.47	0
0.27	176	0.48	0
0.28	174	0.49	0
0.29	170	0.50	0
0.30	166		

# *Mulloidichthys pfluegeri*

Pflueger's goatfish, *weke nono*

Mullidae (goatfishes)



## Life history and other input parameter

Parameter	Value	SD	Unit	n	Source
$L_{inf}$	557	26	mm	-	Mean and SD: Nadon & Ault (2016) $L_{max}$ : 484 (1) from DAR data
$K$	0.55	0.14	yr <sup>-1</sup>		
$a_0$	-0.6	-	yr		
$L_{mat}$	270	31	mm	-	Mean and SD: Nadon & Ault (2016)
Longevity	5.6	2.0	yr	-	Mean and SD: Nadon & Ault (2016)
L-W $\alpha$	2.87e-6	-	-	-	Kulbicki (2005)
L-W $\beta$	3.29				
$L_{S50}$	300	-	mm	-	DAR commercial data
$L_{S95}$	330				
$\bar{L}$ diver survey	-	-	mm	-	-
$\bar{L}$ commercial	401	1	mm	3541	DAR commercial data
$\bar{L}$ combined	-	-	mm	-	-
Max. depth	242	-	m	-	BRFA BotCam project
Federal waters	26	-	%	-	-

## Stock status and other output parameters

Parameter	Median	SD	Unit
$M$	0.57	0.19	yr <sup>-1</sup>
$F$	0.53	0.37	yr <sup>-1</sup>
$F_{30}$	0.81	0.33	yr <sup>-1</sup>
$F/F_{30}$	0.7	0.7	-
$SPR$	0.41	0.23	-
$SPR < 0.30$ iterations	31	-	%
$L_{C30}$	168	-	mm


Parameter	Median	SD	Unit
$B$ from catch	21,100	167,000	kg
$B$ from survey	-	-	kg
Commercial catch	2,195	668	kg
Recreational catch	3,036	9,128	kg
Total catch	6,042	5,736	kg
$C_{30}$ from catch	9,010	89,100	kg
$C_{30}$ from survey	-	-	kg

## General comments

This species mostly occurs in deep water and therefore was not encountered during the diver surveys. The recreational and commercial catch were similar and relatively constant (except for a spike in 2004 from the recreational catch, which is likely spurious). The  $L_{bar}$  from the commercial data has been constant since 2003 reflecting a population likely near equilibrium conditions.


There were no published life history parameters for this species and we therefore used the stepwise approach to generate estimates from an  $L_{max}$  value of 484 mm. This  $L_{max}$  value was close to the maximum length value reported on fishbase.org (480 mm) and no specimen were reported above 500 mm in the commercial data set. A sensitivity run with an  $L_{max}$  of 520 mm generated the following estimates:  $L_{inf}$ : 520 mm,  $K$ : 0.52,  $M$ : 0.61,  $F_{30}$ : 0.82,  $F$ : 0.73,  $SPR$ : 0.34,  $C_{30}$  catch: 6,923 kg. These values were reasonably similar to the original results and did not change the conclusion of our analyses for this species.

*Mulloidichthys pfluegeri*





**Life history parameter distributions.**

*Mulloidichthys pfluegeri*




**Total catch time series from recreational (green squares) and commercial (orange triangles) sectors.**



**Size structure and average length time series from commercial data ( $\pm$ SE).**




*Mulloidichthys pfluegeri*




**Stock status parameter distributions (*SPR*: small bar shows 0.30 level).**

*Mulloidichthys pfluegeri*



**C<sub>30</sub> and current total catch (left) and population size (right) distributions.**

*Mulloidichthys pfluegeri*



**Overfishing probability for a range of  $C_{30}$  levels (commercial data – orange dashed line). *OFL* is represented by a small vertical bar.**

**Probability of overfishing for various  $C_{30}$  levels.**

Overfishing probability	$C_{30}$ from catch (1000 kg)	Overfishing probability	$C_{30}$ from catch (1000 kg)
0.10	2.80	0.31	5.64
0.11	2.94	0.32	5.80
0.12	3.07	0.33	5.95
0.13	3.18	0.34	6.12
0.14	3.31	0.35	6.27
0.15	3.42	0.36	6.40
0.16	3.55	0.37	6.55
0.17	3.67	0.38	6.70
0.18	3.82	0.39	6.87
0.19	3.95	0.40	7.05
0.20	4.08	0.41	7.24
0.21	4.20	0.42	7.42
0.22	4.30	0.43	7.59
0.23	4.42	0.44	7.74
0.24	4.56	0.45	7.91
0.25	4.72	0.46	8.13
0.26	4.88	0.47	8.31
0.27	5.04	0.48	8.54
0.28	5.20	0.49	8.78
0.29	5.35	0.50	9.01
0.30	5.50		

*Mulloidichthys pfluegeri*

**Probability of overfishing at various minimum sizes.**

Overfishing probability	$L_{C_{30}}$ (mm)	Overfishing probability	$L_{C_{30}}$ (mm)
0.10	372	0.31	279
0.11	366	0.32	276
0.12	360	0.33	270
0.13	354	0.34	267
0.14	351	0.35	261
0.15	345	0.36	258
0.16	342	0.37	252
0.17	336	0.38	249
0.18	333	0.39	243
0.19	330	0.40	237
0.20	324	0.41	231
0.21	321	0.42	225
0.22	315	0.43	222
0.23	312	0.44	215
0.24	306	0.45	207
0.25	303	0.46	201
0.26	297	0.47	192
0.27	294	0.48	186
0.28	291	0.49	180
0.29	288	0.50	168
0.30	282		

# Mulloidichthys vanicolensis

Yellowtail goatfish, *weke 'ula*

Mullidae (goatfishes)



## Life history and other input parameters

Parameter	Value	SD	Unit	n	Source
$L_{inf}$	267	30	mm	50	Mean: Cole (2009), SD: Kritzer (2001)
$K$	1.3	0.39	yr <sup>-1</sup>		
$a_0$	-1.1	-	yr		
$L_{mat}$	206	6	mm	118	Mean: Cole (2009), SD: Nadon (unpublished)
Longevity	5	1	yr	50	Mean: Cole (2009), SD: Kritzer (2001)
L-W $\alpha$	1.83e-5	-	-	-	Jehangeer (2003)
L-W $\beta$	2.96	-	-	-	
$L_{S50}$	200	-	mm	-	DAR commercial data
$L_{S95}$	210				
$\bar{L}$ diver survey	256	8	mm	595	NOAA-CREP diver survey
$\bar{L}$ commercial	278	2	mm	2175	DAR commercial data
$\bar{L}$ combined	276	3	mm	-	-
Max. depth	132	-	m	-	BRFA BotCam project
Federal waters	22	-	%	-	-

## Stock status and other output parameters

Parameter	Median	SD	Unit
$M$	0.61	0.12	yr <sup>-1</sup>
$F$	0.35	0.40	yr <sup>-1</sup>
$F_{30}$	0.83	0.22	yr <sup>-1</sup>
$F/F_{30}$	0.4	0.5	-
$SPR$	0.55	0.23	-
$SPR < 0.30$ iterations	15	-	%
$LC_{30}$	-	-	mm

Parameter	Median	SD	Unit
$B$ from catch	139,000	897,000	kg
$B$ from survey	34,896	17,672	kg
Commercial catch	14,267	4,772	kg
Recreational catch	9,875	38,536	kg
Total catch	28,511	22,058	kg
$C_{30}$ from catch	61,600	402,000	kg
$C_{30}$ from survey	15,500	7,710	kg


## General comments

Population abundance appeared to be declining in the last year of the UVS surveys. Further surveys should clarify if this is a clear pattern or caused by an outlier data point. The catch from both sectors were similar and constant, except for an anomalous jump in the recreational catch in 2015. This jump could be due to the same juvenile goatfish reporting issue as for *M. flavolineatus* (see discussion for this species). It is unlikely to reflect a real increase in recreational catch for this species. Commercial and recreational  $L_{bar}$  appeared relatively constant for this species, suggesting a population near equilibrium. However, the commercial  $L_{bar}$  was higher than  $L_{inf}$  for this species, which is problematic given the low  $M/K$  ratio. We thus ran the analyses with the diver survey  $L_{bar}$ , given the high observation count.

The life history parameters for this species came from a local study with a limited sample size. Using the stepwise approach, we obtained the following results:  $L_{max}$ : 325 mm,  $L_{inf}$ : 377 mm,  $K$ : 0.59,  $M$ : 0.61,  $F_{30}$ : 0.80,  $F$ : 0.70,  $SPR$ : 0.34,  $C_{30}$  survey: 13,890 kg. Note that goatfishes usually have indeterminate growth curves, resulting in a fairly high  $L_{inf}$  value for the stepwise approach in this example.


Similarly to *M. flavolineatus*, there was a discrepancy between the population sizes estimated from the catch vs. diver surveys (139,000 kg vs. 34,896 kg). This led to significant difference in  $C_{30}$  estimates from the catch vs. diver surveys as well. The smaller population estimate could be due to the depth limit associated with diver surveys (30 m) vs. the actual depth range of this species (132 m). Note that the catch-derived population size from the sensitivity run was much smaller (80,965 kg) given the much higher  $F$  value (0.70 vs. 0.35). The  $C_{30}$  from the diver survey is likely more reliable than the catch-derived estimates.

*Mulloidichthys vanicolensis*




**Life history parameter distributions.**

*Mulloidichthys vanicolensis*




**Abundance index from UVS (blue circles,  $\pm$ SE) and total catch time series from recreational (green squares) and commercial (orange triangles) sectors.**



**Size structure from commercial catch (top left) and UVS (top right). Average length time series (blue circles – UVS, orange triangles – commercial data,  $\pm$ SE).**


*Mulloidichthys vanicolensis*



**Stock status parameter distributions (*SPR*: small bar shows 0.30 level).**




*Mulloidichthys vanicolensis*



$C_{30}$  and current total catch (left) and population size (right) distributions.

*Mulloidichthys vanicolensis*



**Overfishing probability for a range of  $C_{30}$  levels (catch - orange dashed line, UVS – blue dotted line).  $OFLs$  are represented by small vertical bars.**

**Probability of overfishing for various  $C_{30}$  levels.**

Overfish. probability	$C_{30}$ from catch (1000 kg)	$C_{30}$ from survey (1000 kg)	Overfish. probability	$C_{30}$ from catch (1000 kg)	$C_{30}$ from survey (1000 kg)
0.10	18.7	6.4	0.31	37.0	11.7
0.11	19.6	6.7	0.32	38.2	11.9
0.12	20.4	7.0	0.33	39.2	12.1
0.13	21.3	7.3	0.34	40.3	12.3
0.14	22.0	7.6	0.35	41.3	12.5
0.15	22.9	7.9	0.36	42.3	12.8
0.16	23.6	8.2	0.37	43.5	13.0
0.17	24.4	8.4	0.38	44.7	13.2
0.18	25.5	8.7	0.39	45.9	13.4
0.19	26.4	9.0	0.40	46.9	13.6
0.20	27.2	9.2	0.41	48.0	13.8
0.21	28.1	9.5	0.42	49.5	14.0
0.22	28.9	9.7	0.43	50.8	14.1
0.23	29.6	10.0	0.44	52.3	14.3
0.24	30.3	10.2	0.45	53.7	14.5
0.25	31.2	10.4	0.46	55.3	14.7
0.26	31.8	10.6	0.47	56.5	14.9
0.27	33.1	10.9	0.48	58.1	15.1
0.28	34.0	11.1	0.49	60.1	15.2
0.29	35.0	11.3	0.50	61.6	15.5
0.30	35.9	11.5			

# *Parupeneus cyclostomus*

Blue goatfish, *moano hulu*

Mullidae (goatfishes)



## Life history and other input parameters

Parameter	Value	SD	Unit	n	Source
$L_{inf}$	565	30	mm	-	Mean and SD: Nadon & Ault (2016) $L_{max}$ : 494 (10) from NWHI diver survey
$K$	0.512	0.146	yr <sup>-1</sup>		
$a_0$	-0.6	-	yr		
$L_{mat}$	269	33	mm	-	Mean and SD: Nadon & Ault (2016)
Longevity	5.4	1.9	yr	-	Mean and SD: Nadon & Ault (2016)
L-W $\alpha$	8.28e-6	-	-	-	Kulbicki (2001)
L-W $\beta$	3.13				
$L_{S50}$	225	-	mm	-	DAR commercial data
$L_{S95}$	250				
$\bar{L}$ diver survey	318	4	mm	362	NOAA-CREP diver survey
$\bar{L}$ commercial	342	2	mm	3070	DAR commercial data
$\bar{L}$ combined	336	3	mm	-	-
Max. depth	113	-	m	-	Pyle et al. (2016)
Federal waters	21	-	%	-	-

## Stock status and other output parameters

Parameter	Median	SD	Unit
$M$	0.60	0.18	yr <sup>-1</sup>
$F$	0.73	0.46	yr <sup>-1</sup>
$F_{30}$	0.59	0.17	yr <sup>-1</sup>
$F/F_{30}$	1.3	1.0	-
$SPR$	0.24	0.22	-
$SPR < 0.30$ iterations	61	-	%
$L_{C30}$	263	-	mm

Parameter	Median	SD	Unit
$B$ from catch	12,000	66,300	kg
$B$ from survey	77,428	30,321	kg
Commercial catch	700	257	kg
Recreational catch	3,684	1,871	kg
Total catch	4,474	1,829	kg
$C_{30}$ from catch	4,000	25,200	kg
$C_{30}$ from survey	25,600	11,300	kg


## General comments

Population abundance jumped unexpectedly in 2012 with no easy explanation for this pattern. Future surveys should reveal if this is a persistent pattern or simply an anomaly. The commercial catch appeared to be rising slightly from 2003 to 2014 but fell in 2016. The recreational catch was fairly variable but also appear to be rising slightly. The commercial and recreational  $L_{bar}$  were reasonably similar and mostly stable in recent years.

There were no published life history parameters for this species and the stepwise approach was used to generate estimates. The  $L_{max}$  used for this analysis was obtained from the pristine NWHI and was likely a fairly reasonable estimate. A sensitivity run with a larger  $L_{max}$  (530 mm) generated the following estimates:  $L_{inf}$ : 608 mm,  $K$ : 0.50,  $M$ : 0.63,  $F_{30}$ : 0.57,  $F$ : 1.0,  $SPR$ : 0.16,  $C_{30}$  survey: 26,100 kg.


The catch data for this species seemed unusually small (4,474 kg) compared to other species with similar abundance (*M. vanicolensis*, *M. flavolineatus*). There is little reason to doubt the diver surveys, given that this is a fairly common and easily identifiable species. The diver survey  $C_{30}$  is likely more reliable than the catch-derived  $C_{30}$ .

*Parupeneus cyclostomus*




**Life history parameter distributions.**

*Parupeneus cyclostomus*




**Abundance index from UVS (blue circles,  $\pm$ SE) and total catch time series from recreational (green squares) and commercial (orange triangles) sectors.**




**Size structure from commercial catch (top left) and UVS (top right). Average length time series (blue circles – UVS, orange triangles – commercial data,  $\pm$ SE).**

*Parupeneus cyclostomus*




**Stock status parameter distributions (*SPR*: small bar shows 0.30 level).**

*Parupeneus cyclostomus*



$C_{30}$  and current total catch (left) and population size (right) distributions.

*Parupeneus cyclostomus*



**Overfishing probability for a range of  $C_{30}$  levels (catch - orange dashed line, UVS – blue dotted line). *OFLs* are represented by small vertical bars.**

**Probability of overfishing for various  $C_{30}$  levels.**

Overfish. probability	$C_{30}$ from catch (1000 kg)	$C_{30}$ from survey (1000 kg)	Overfish. probability	$C_{30}$ from catch (1000 kg)	$C_{30}$ from survey (1000 kg)
0.10	1.92	12.3	0.31	2.93	20.3
0.11	1.99	12.9	0.32	2.99	20.6
0.12	2.03	13.4	0.33	3.03	20.9
0.13	2.09	14.0	0.34	3.08	21.2
0.14	2.14	14.4	0.35	3.13	21.5
0.15	2.19	15.0	0.36	3.17	21.8
0.16	2.24	15.3	0.37	3.22	22.0
0.17	2.28	15.8	0.38	3.28	22.3
0.18	2.33	16.1	0.39	3.34	22.6
0.19	2.37	16.5	0.40	3.39	22.8
0.20	2.41	16.9	0.41	3.45	23.1
0.21	2.46	17.3	0.42	3.52	23.4
0.22	2.51	17.6	0.43	3.57	23.7
0.23	2.56	18.0	0.44	3.63	23.9
0.24	2.61	18.2	0.45	3.69	24.2
0.25	2.65	18.5	0.46	3.76	24.5
0.26	2.70	18.8	0.47	3.81	24.7
0.27	2.75	19.1	0.48	3.87	25.0
0.28	2.79	19.4	0.49	3.94	25.3
0.29	2.84	19.7	0.50	4.00	25.6
0.30	2.89	20.1			



*Parupeneus cyclostomus*

**Probability of overfishing at various minimum sizes.**

Overfishing probability	$L_{C_{30}}$ (mm)	Overfishing probability	$L_{C_{30}}$ (mm)
0.10	403	0.31	326
0.11	398	0.32	324
0.12	394	0.33	322
0.13	389	0.34	317
0.14	385	0.35	315
0.15	382	0.36	310
0.16	378	0.37	308
0.17	376	0.38	306
0.18	371	0.39	304
0.19	367	0.40	299
0.20	364	0.41	297
0.21	360	0.42	295
0.22	358	0.43	290
0.23	353	0.44	286
0.24	351	0.45	281
0.25	349	0.46	279
0.26	344	0.47	274
0.27	340	0.48	270
0.28	338	0.49	266
0.29	333	0.50	263
0.30	331		

# *Parupeneus insularis*

Island goatfish, *munu*

Mullidae (goatfishes)



## Life history and other input parameters

Parameter	Value	SD	Unit	n	Source
$L_{inf}$	376	17	mm	-	Mean and SD: Nadon & Ault (2016) $L_{max}$ : 324 (4) from NWHI diver survey
$K$	0.60	0.14	yr <sup>-1</sup>		
$a_0$	-0.6	-	yr		
$L_{mat}$	203	25	mm	-	Mean and SD: Nadon & Ault (2016)
Longevity	6.4	2.2	yr	-	Mean and SD: Nadon & Ault (2016)
L-W $\alpha$	9.15e-6				
L-W $\beta$	3.13				
$L_{S50}$	220	-	mm	-	DAR commercial data
$L_{S95}$	240				
$\bar{L}$ diver survey	278	3	mm	179	NOAA-CREP diver survey
$\bar{L}$ commercial	305	2	mm	1598	DAR commercial data
$\bar{L}$ combined	296	2	mm	-	-
Max. depth	90	-	m	-	Pyle et al. (2016)
Federal waters	21	-	%	-	-

## Stock status and other output parameters

Parameter	Median	SD	Unit
$M$	0.50	0.16	yr <sup>-1</sup>
$F$	0.28	0.24	yr <sup>-1</sup>
$F_{30}$	0.72	0.25	yr <sup>-1</sup>
$F/F_{30}$	0.4	0.4	-
$SPR$	0.57	0.22	-
$SPR < 0.30$ iterations	12	-	%
$L_{C30}$	0	-	mm

Parameter	Median	SD	Unit
$B$ from catch	4,740	37,500	kg
$B$ from survey	41,983	7,557	kg
Commercial catch	176	153	kg
Recreational catch	496	1,358	kg
Total catch	776	1,011	kg
$C_{30}$ from catch	1,960	16,900	kg
$C_{30}$ from survey	17,100	4,300	kg


## General comments

Population abundance was stable from 2005 to 2016. The commercial catch increased in the early year before going down from 2013 to 2016. The recreational catch was elevated in the early years and lower in recent years, although it is not clear if this is a real pattern given the variability associated with this data set.  $L_{bar}$  from both data set appeared to be declining slightly.

There are no published life history parameters for this species and the stepwise approach was used to generate estimates using an  $L_{max}$  from NWHI surveys. A sensitivity run with a  $L_{max}$  of 360 mm generated the following results:  $L_{inf}$ : 412 mm,  $K$ : 0.56,  $M$ : 0.57,  $F_{30}$ : 0.73,  $F$ : 0.48,  $SPR$ : 0.42,  $C_{30}$  survey: 16,783 kg.


Similarly to *P. cyclostomus*, The catch data for this species seemed unusually small (776 kg) compared to other species with similar abundance (*M. vanicolensis*, *M. flavolineatus*). There is no clear explanation for this observation. However, the diver survey data is likely more reliable to estimate population size and *OFLs*.

*Parupeneus insularis*




**Life history parameter distributions.**

*Parupeneus insularis*




**Abundance index from UVS (blue circles,  $\pm$ SE) and total catch time series from recreational (green squares) and commercial (orange triangles) sectors.**




**Size structure from commercial catch (top left) and UVS (top right). Average length time series (blue circles – UVS, orange triangles – commercial data,  $\pm$ SE).**

*Parupeneus insularis*




**Stock status parameter distributions (*SPR*: small bar shows 0.30 level).**

*Parupeneus insularis*



$C_{30}$  and current total catch (left) and population size (right) distributions.

*Parupeneus insularis*



**Overfishing probability for a range of  $C_{30}$  levels (catch - orange dashed line, UVS – blue dotted line). *OFLs* are represented by small vertical bars.**

**Probability of overfishing for various  $C_{30}$  levels.**

Overfish. probability	$C_{30}$ from catch (1000 kg)	$C_{30}$ from survey (1000 kg)	Overfish. probability	$C_{30}$ from catch (1000 kg)	$C_{30}$ from survey (1000 kg)
0.10	0.49	12.1	0.31	1.09	15.1
0.11	0.52	12.3	0.32	1.13	15.1
0.12	0.55	12.5	0.33	1.17	15.2
0.13	0.58	12.6	0.34	1.20	15.4
0.14	0.60	12.8	0.35	1.24	15.5
0.15	0.63	13.0	0.36	1.28	15.6
0.16	0.65	13.1	0.37	1.31	15.7
0.17	0.67	13.3	0.38	1.35	15.8
0.18	0.70	13.4	0.39	1.40	15.9
0.19	0.72	13.6	0.40	1.45	16.1
0.20	0.75	13.7	0.41	1.50	16.2
0.21	0.78	13.8	0.42	1.55	16.3
0.22	0.82	14.0	0.43	1.58	16.4
0.23	0.84	14.1	0.44	1.63	16.5
0.24	0.87	14.2	0.45	1.69	16.6
0.25	0.90	14.3	0.46	1.74	16.7
0.26	0.93	14.4	0.47	1.79	16.8
0.27	0.96	14.5	0.48	1.83	16.9
0.28	0.99	14.7	0.49	1.90	17.0
0.29	1.02	14.8	0.50	1.96	17.1
0.30	1.06	14.9			

*Parupeneus insularis*

**Probability of overfishing at various minimum sizes.**

Overfishing probability	$L_{C_{30}}$ (mm)	Overfishing probability	$L_{C_{30}}$ (mm)
0.10	216	0.31	0
0.11	209	0.32	0
0.12	205	0.33	0
0.13	200	0.34	0
0.14	196	0.35	0
0.15	189	0.36	0
0.16	183	0.37	0
0.17	178	0.38	0
0.18	172	0.39	0
0.19	165	0.40	0
0.20	158	0.41	0
0.21	152	0.42	0
0.22	145	0.43	0
0.23	136	0.44	0
0.24	128	0.45	0
0.25	117	0.46	0
0.26	91	0.47	0
0.27	0	0.48	0
0.28	0	0.49	0
0.29	0	0.50	0
0.30	0		



# *Parupeneus porphyreus*

White-saddle goatfish, *kumu*

Mullidae (goatfishes)



## Life history and other input parameters

Parameter	Value	SD	Unit	n	Source
$L_{inf}$	547	76	mm	36	Mean: Moffitt (1979), SD: Kritzer (2001)
$K$	0.538	0.231	yr <sup>-1</sup>		
$a_0$	-0.446	-	yr		
$L_{mat}$	264	13	mm	36	Mean: Moffitt (1979), SD: Nadon (unpublished)
Longevity	6	1.1	yr	36	Mean: Moffitt (1979), SD: Kritzer (2001)
L-W $\alpha$	6.99e-6	-	-	-	Kulbicki (2005) - <i>P. multifasciatus</i>
L-W $\beta$	3.211	-	-	-	
$L_{S50}$	225	-	mm	-	DAR commercial data
$L_{S95}$	250	-	mm	-	
$\bar{L}$ diver survey	299	8	mm	179	NOAA-CREP diver survey
$\bar{L}$ commercial	323	1	mm	5074	DAR commercial data
$\bar{L}$ combined	323	2	mm	-	-
Max. depth	140	-	m	-	Pyle et al. (2016)
Federal waters	22	-	%	-	-

## Stock status and other output parameters

Parameter	Median	SD	Unit
$M$	0.53	0.11	yr <sup>-1</sup>
$F$	1.00	0.58	yr <sup>-1</sup>
$F_{30}$	0.54	0.11	yr <sup>-1</sup>
$F/F_{30}$	1.9	1.1	-
$SPR$	0.15	0.17	-
$SPR < 0.30$ iterations	80	-	%
$L_{C30}$	317	-	mm

Parameter	Median	SD	Unit
$B$ from catch	14,600	73,000	kg
$B$ from survey	13,810	4,225	kg
Commercial catch	1,717	1,507	kg
Recreational catch	5,045	2,266	kg
Total catch	7,130	2,704	kg
$C_{30}$ from catch	4,860	25,000	kg
$C_{30}$ from survey	4,580	1,480	kg


## General comments

Population abundance was highly variable due to the number of observations in individual years for this species being low. The commercial catch increased from 2008 to 2011 but has been going down since then. The recreational catch also appeared to be declining. The commercial  $L_{bar}$  has been steady (there were not enough observations per year to generate a  $L_{bar}$  time series from UVS).

The life history parameters for this species came from a local study with a limited sample size. The stepwise approach applied to this species generated the following estimates:  $L_{max}$ : 435 mm,  $L_{inf}$ : 496 mm,  $K$ : 0.54,  $M$ : 0.59,  $F_{30}$ : 0.63,  $F$ : 0.60,  $SPR$ : 0.32,  $C_{30}$  survey: 5,157 kg,  $C_{30}$  catch: 7,931 kg. The lower  $L_{inf}$  estimate lead to a lower  $F$  and higher  $SPR$ .


There was a good agreement between catch-derived and survey-derived population and  $C_{30}$  estimates. The survey-derived  $C_{30}$  had lower variability and is usually more reliable than catch-derived  $C_{30}$ .

*Parupeneus porphyreus*




**Life history parameter distributions.**

*Parupeneus porphyreus*




**Abundance index from UVS (blue circles, ±SE) and total catch time series from recreational (green squares) and commercial (orange triangles) sectors.**




**Size structure from commercial catch (top left) and UVS (top right). Average length time series (blue circles – UVS, orange triangles – commercial data, ±SE).**

*Parupeneus porphyreus*




**Stock status parameter distributions ( $SPR$ : small bar shows 0.30 level).**

*Parupeneus porphyreus*



$C_{30}$  and current total catch (left) and population size (right) distributions.

*Parupeneus porphyreus*



**Overfishing probability for a range of  $C_{30}$  levels (catch - orange dashed line, UVS – blue dotted line). *OFLs* are represented by small vertical bars.**

**Probability of overfishing for various  $C_{30}$  levels.**

Overfish. probability	$C_{30}$ from catch (1000 kg)	$C_{30}$ from survey (1000 kg)	Overfish. probability	$C_{30}$ from catch (1000 kg)	$C_{30}$ from survey (1000 kg)
0.10	2.64	2.73	0.31	3.80	3.84
0.11	2.70	2.80	0.32	3.86	3.88
0.12	2.77	2.86	0.33	3.90	3.92
0.13	2.83	2.93	0.34	3.95	3.96
0.14	2.89	3.00	0.35	4.00	4.00
0.15	2.95	3.05	0.36	4.06	4.05
0.16	3.01	3.11	0.37	4.10	4.09
0.17	3.07	3.17	0.38	4.16	4.14
0.18	3.12	3.23	0.39	4.22	4.18
0.19	3.18	3.30	0.40	4.27	4.22
0.20	3.23	3.34	0.41	4.33	4.25
0.21	3.28	3.39	0.42	4.39	4.28
0.22	3.33	3.44	0.43	4.44	4.32
0.23	3.39	3.49	0.44	4.49	4.36
0.24	3.44	3.54	0.45	4.54	4.39
0.25	3.50	3.59	0.46	4.61	4.42
0.26	3.56	3.64	0.47	4.67	4.47
0.27	3.60	3.68	0.48	4.73	4.50
0.28	3.66	3.72	0.49	4.80	4.54
0.29	3.70	3.77	0.50	4.86	4.58
0.30	3.75	3.80			

*Parupeneus porphyreus*

**Probability of overfishing at various minimum sizes.**

Overfishing probability	$L_{C30}$ (mm)	Overfishing probability	$L_{C30}$ (mm)
0.10	418	0.31	360
0.11	414	0.32	358
0.12	412	0.33	356
0.13	407	0.34	353
0.14	405	0.35	351
0.15	400	0.36	349
0.16	398	0.37	346
0.17	396	0.38	344
0.18	394	0.39	342
0.19	389	0.40	340
0.20	387	0.41	338
0.21	385	0.42	335
0.22	382	0.43	333
0.23	380	0.44	331
0.24	376	0.45	328
0.25	374	0.46	326
0.26	374	0.47	324
0.27	369	0.48	322
0.28	367	0.49	320
0.29	364	0.50	317
0.30	362		

# *Calotomus carolinus*

Stareye parrotfish, *ponuhunuhu*

Scaridae (parrotfishes)



## Life history and other input parameters

Parameter	Value	SD	Unit	n	Source
$L_{inf}$	366	25	mm	-	Mean and SD: Nadon & Ault (2016) $L_{max}$ : 430 (11) from NWHI diver survey
$K$	0.534	0.168	yr <sup>-1</sup>		
$a_0$	-0.6	-	yr		
$L_{mat}$	253	29	mm	-	Mean and SD: Nadon & Ault (2016)
Longevity	13	4.8	yr	-	Mean and SD: Nadon & Ault (2016)
L-W $\alpha$	8.31e-6	-	-	-	Smith & Dalzell (1993)
L-W $\beta$	3.17				
$L_{S50}$	200	-	mm	-	Estimated.
$L_{S95}$	220				
$\bar{L}$ diver survey	275	4	mm	141	NOAA-CREP diver survey
$\bar{L}$ commercial	-	-	mm	-	-
$\bar{L}$ combined	-	-	mm	-	-
Max. depth	71	-	m	-	Pyle et al. (2016)
Federal waters	21	-	%	-	-

## Stock status and other output parameters

Parameter	Median	SD	Unit
$M$	0.24	0.08	yr <sup>-1</sup>
$F$	0.61	0.33	yr <sup>-1</sup>
$F_{30}$	0.29	0.10	yr <sup>-1</sup>
$F/F_{30}$	2.2	1.2	-
$SPR$	0.13	0.14	-
$SPR < 0.30$ iterations	87	-	%
$L_{C30}$	276	-	mm

Parameter	Median	SD	Unit
$B$ from catch	-	-	kg
$B$ from survey	38,102	9,192	kg
Commercial catch	-	-	kg
Recreational catch	-	-	kg
Total catch	-	-	kg
$C_{30}$ from catch	-	-	kg
$C_{30}$ from survey	8,430	3,010	kg

## General comments

Parrotfish catches are grouped at the family level and therefore there is no species-level commercial data.


Population abundance for this species appeared to be increasing significantly. There were not enough observation in individual years to generate a  $L_{bar}$  time series.

There are currently no published life history parameters for this species. We used the stepwise approach to generate LH parameters, using a  $L_{max}$  value of 430 mm from the pristine NWHI. Analyses using the lower  $L_{max}$  value found in the MHI (410 mm) generated the following results:  $L_{inf}$ : 349 mm,  $K$ : 0.57,  $M$ : 0.25,  $F_{30}$ : 0.32,  $F$ : 0.46,  $SPR$ : 0.20,  $C_{30}$  survey: 8,879 kg. Furthermore, selectivity for parrotfishes had to be estimated, given the absence of catch data. A sensitivity run with  $L_{S50}$  at 230 mm and  $L_{S95}$  at 260 mm had little impact on the results ( $F$ : 0.62,  $SPR$ : 0.16).

The population biomass estimate may be biased downward given that this species' range extends to 68 m depth, which is beyond the depth of the diver surveys (30 m).




*Calotomus carolinus*




**Life history parameter distributions.**

*Calotomus carolinus*




**Abundance index from UVS (blue circles, ±SE).**




**Size structure and average length time series from UVS (±SE).**

*Calotomus carolinus*




**Stock status parameter distributions ( $SPR$ : small bar shows 0.30 level).**



$C_{30}$  (left) and population size (right) distributions.

*Calotomus carolinus*



**Overfishing probability for a range of  $C_{30}$  levels (UVS – blue dotted line). *OFL* is represented by a small vertical bar.**

**Probability of overfishing for various  $C_{30}$  levels.**

Overfishing probability	$C_{30}$ from survey (1000 kg)	Overfishing probability	$C_{30}$ from survey (1000 kg)
0.10	5.12	0.31	6.99
0.11	5.24	0.32	7.07
0.12	5.34	0.33	7.15
0.13	5.46	0.34	7.23
0.14	5.58	0.35	7.32
0.15	5.66	0.36	7.39
0.16	5.74	0.37	7.46
0.17	5.85	0.38	7.51
0.18	5.94	0.39	7.58
0.19	6.03	0.40	7.64
0.20	6.13	0.41	7.73
0.21	6.22	0.42	7.81
0.22	6.29	0.43	7.90
0.23	6.37	0.44	7.97
0.24	6.45	0.45	8.04
0.25	6.53	0.46	8.13
0.26	6.63	0.47	8.22
0.27	6.70	0.48	8.29
0.28	6.77	0.49	8.35
0.29	6.84	0.50	8.43
0.30	6.92		

*Calotomus carolinus*

**Probability of overfishing at various minimum sizes.**

Overfishing probability	$L_{C_{30}}$ (mm)	Overfishing probability	$L_{C_{30}}$ (mm)
0.10	334	0.31	302
0.11	332	0.32	302
0.12	330	0.33	300
0.13	328	0.34	298
0.14	326	0.35	296
0.15	324	0.36	296
0.16	324	0.37	294
0.17	322	0.38	292
0.18	320	0.39	292
0.19	320	0.40	290
0.20	318	0.41	288
0.21	316	0.42	288
0.22	314	0.43	286
0.23	312	0.44	284
0.24	312	0.45	282
0.25	310	0.46	282
0.26	310	0.47	280
0.27	308	0.48	278
0.28	306	0.49	278
0.29	306	0.50	276
0.30	304		

# *Chlorurus perspicillatus*

Spectacled parrotfish, *uhu uliuli*

Scaridae (parrotfishes)



## Life history and other input parameters

Parameter	Value	SD	Unit	N	Source
$L_{inf}$	500	22	mm	-	Mean and SD: Nadon & Ault (2016) $L_{max}$ : 574 (4) from NWHI diver survey
$K$	0.377	0.135	yr <sup>-1</sup>		
$a_0$	-0.6	-	yr		
$L_{mat}$	348	28	mm	-	Mean and SD: Nadon & Ault (2016)
Longevity	19	6.5	yr	-	Mean and SD: Nadon & Ault (2016)
L-W $\alpha$	1.06e-5	-	-	-	Smith & Dalzell (1993) taken for <i>S. rubroviolaceus</i>
L-W $\beta$	3.11				
$L_{S50}$	240	-	mm	-	Estimated.
$L_{S95}$	260				
$\bar{L}$ diver survey	404	13	mm	123	NOAA-CREP diver survey
$\bar{L}$ commercial	-	-	mm	-	-
$\bar{L}$ combined	-	-	mm	-	-
Max. depth	80	-	m	-	Pyle et al. (2016)
Federal waters	21	-	%	-	-

## Stock status and other output parameters

Parameter	Median	SD	Unit
$M$	0.17	0.06	yr <sup>-1</sup>
$F$	0.09	0.08	yr <sup>-1</sup>
$F_{30}$	0.19	0.06	yr <sup>-1</sup>
$F/F_{30}$	0.5	0.4	-
$SPR$	0.54	0.21	-
$SPR < 0.30$ iterations	13	-	%
$L_{C30}$	0	-	mm

Parameter	Median	SD	Unit
$B$ from catch	-	-	kg
$B$ from survey	78,752	22,829	kg
Commercial catch	-	-	kg
Recreational catch	-	-	kg
Total catch	-	-	kg
$C_{30}$ from catch	-	-	kg
$C_{30}$ from survey	12,400	5,050	kg

## General comments


Parrotfish catches are grouped at the family level and therefore there is no species-level commercial data.

Population abundance was fairly variable from year to year for this species, likely due to the relatively low observation counts. Similarly, there were not enough yearly observations to generate an  $L_{bar}$  time series.

There are currently no published life history parameters for this species. We used the stepwise approach to generate LH parameters using an  $L_{max}$  value (574 mm) from the NWHI, where this species is fairly abundant. Analyses using a higher  $L_{max}$  estimate from the MHI (610 mm) generated the following results:  $L_{inf}$ : 526 mm,  $K$ : 0.33,  $M$ : 0.17,  $F_{30}$ : 0.17,  $F$ : 0.11,  $SPR$ : 0.45,  $C_{30}$  survey: 11,609 kg. Furthermore, selectivity for parrotfishes had to be estimated, given the absence of catch data. A sensitivity run with  $L_{S50}$  at 260 mm and  $L_{S95}$  at 300 mm had little impact on the results ( $F$ : 0.09,  $SPR$ : 0.52).


The population biomass estimate may be biased downward given that this species' range extends to 68 m depth, which is beyond the depth of the diver surveys (30 m).

*Chlorurus perspicillatus*




**Life history parameter distributions.**

*Chlorurus perspicillatus*




Abundance index from UVS (blue circles,  $\pm$ SE).




Size structure and average length time series from UVS ( $\pm$ SE).



*Chlorurus perspicillatus*




**Stock status parameter distributions ( $SPR$ : small bar shows 0.30 level).**



**$C_{30}$  (left) and population size (right) distributions.**

*Chlorurus perspicillatus*



**Overfishing probability for a range of  $C_{30}$  levels (UVS – blue dotted line). *OFL* is represented by a small vertical bar.**

**Probability of overfishing for various  $C_{30}$  levels.**

Overfishing probability	$C_{30}$ from survey (1000 kg)	Overfishing probability	$C_{30}$ from survey (1000 kg)
0.10	6.9	0.31	10.1
0.11	7.1	0.32	10.2
0.12	7.3	0.33	10.3
0.13	7.5	0.34	10.4
0.14	7.6	0.35	10.6
0.15	7.8	0.36	10.7
0.16	8.0	0.37	10.8
0.17	8.1	0.38	10.9
0.18	8.4	0.39	11.1
0.19	8.5	0.40	11.2
0.20	8.6	0.41	11.3
0.21	8.8	0.42	11.4
0.22	8.9	0.43	11.6
0.23	9.1	0.44	11.7
0.24	9.2	0.45	11.8
0.25	9.3	0.46	11.9
0.26	9.4	0.47	12.0
0.27	9.6	0.48	12.1
0.28	9.7	0.49	12.2
0.29	9.8	0.50	12.4
0.30	9.9		

*Chlorurus perspicillatus*

**Probability of overfishing at various minimum sizes.**

Overfishing probability	$L_{C_{30}}$ (mm)	Overfishing probability	$L_{C_{30}}$ (mm)
0.10	278	0.31	0
0.11	266	0.32	0
0.12	257	0.33	0
0.13	240	0.34	0
0.14	228	0.35	0
0.15	214	0.36	0
0.16	194	0.37	0
0.17	173	0.38	0
0.18	158	0.39	0
0.19	137	0.40	0
0.20	108	0.41	0
0.21	0	0.42	0
0.22	0	0.43	0
0.23	0	0.44	0
0.24	0	0.45	0
0.25	0	0.46	0
0.26	0	0.47	0
0.27	0	0.48	0
0.28	0	0.49	0
0.29	0	0.50	0
0.30	0		

# *Chlorurus spilurus*

Bullethead parrotfish, *uhu*

Scaridae (parrotfishes)



## Life history and other input parameters

Parameter	Value	SD	Unit	n	Source
$L_{inf}$	341	21	mm	-	Mean and SD: Nadon & Ault (2016) $L_{max}$ : 397 (7) from NWHI diver survey
$K$	0.59	0.22	yr <sup>-1</sup>		
$a_0$	-0.6	-	yr		
$L_{mat}$	235	37	mm	-	Mean and SD: Nadon & Ault (2016)
Longevity	13	4.2	yr	-	Mean and SD: Nadon & Ault (2016)
L-W $\alpha$	2.61e-5	-	-	-	Kulbicki (2005)
L-W $\beta$	2.97				
$L_{S50}$	220	-	mm	-	Estimated.
$L_{S95}$	240				
$\bar{L}$ diver survey	283	3	mm	435	NOAA-CREP diver survey
$\bar{L}$ commercial	-	-	mm	-	-
$\bar{L}$ combined	-	-	mm	-	-
Max. depth	34	-	m	-	NOAA-CREP BRUV survey
Federal waters	0	-	%	-	-

## Stock status and other output parameters

Parameter	Median	SD	Unit
$M$	0.26	0.09	yr <sup>-1</sup>
$F$	0.50	0.36	yr <sup>-1</sup>
$F_{30}$	0.37	0.15	yr <sup>-1</sup>
$F/F_{30}$	1.4	1.1	-
$SPR$	0.23	0.20	-
$SPR < 0.30$ iterations	64	-	%
$L_{C30}$	222	-	mm

Parameter	Median	SD	Unit
$B$ from catch	-	-	kg
$B$ from survey	139,438	17,478	kg
Commercial catch	-	-	kg
Recreational catch	-	-	kg
Total catch	-	-	kg
$C_{30}$ from catch	-	-	kg
$C_{30}$ from survey	38,400	11,000	kg

## General comments

Parrotfish catches are grouped at the family level and therefore there is no species-level commercial data.


Population abundance has been rising steadily since 2008 while  $L_{bar}$  seemed to be slightly declining. Selectivity had to be estimated given the absence of fishery data. A sensitivity run with  $L_{S50}$  at 190 mm and  $L_{S95}$  at 220 mm had little impact on the results ( $F$ : 0.57,  $SPR$ : 0.16).

Life history parameters were available for this species from a study in American Samoa but the  $L_{inf}$  parameter was too small (289 mm vs. an  $L_{max}$  of 380 mm in the MHI). We therefore used the stepwise approach for this species. The American Samoa LH parameters generated the following estimates:  $F$ : 0.03,  $SPR$ : 0.88. The much lower  $F$  value (and higher  $SPR$  value) were expected given the unrealistically low  $L_{inf}$  estimate from this source.

Note that the longevity estimate obtained through the stepwise approach was similar to the maximum recorded age for this species (13 yr vs. 10 yr). The NWHI  $L_{max}$  is identical to the MHI estimate (397 mm), which suggested this number was appropriate.


The population biomass estimate should be reasonably accurate given that this species extends only slightly beyond the maximum diver survey depth (30 m).

*Chlorurus spilurus*





**Life history parameter distributions.**

*Chlorurus spilurus*




**Abundance index from UVS (blue circles,  $\pm$ SE).**




**Size structure and average length time series from UVS ( $\pm$ SE).**

*Chlorurus spilurus*




**Stock status parameter distributions ( $SPR$ : small bar shows 0.30 level).**



**$C_{30}$  (left) and population size (right) distributions.**

*Chlorurus spilurus*



**Overfishing probability for a range of  $C_{30}$  levels (UVS – blue dotted line). *OFL* is represented by a small vertical bar.**

**Probability of overfishing for various  $C_{30}$  levels.**

Overfishing probability	$C_{30}$ from survey (1000 kg)	Overfishing probability	$C_{30}$ from survey (1000 kg)
0.10	26.6	0.31	33.3
0.11	26.9	0.32	33.7
0.12	27.4	0.33	33.9
0.13	27.8	0.34	34.2
0.14	28.2	0.35	34.4
0.15	28.6	0.36	34.7
0.16	29.0	0.37	35.0
0.17	29.3	0.38	35.1
0.18	29.6	0.39	35.4
0.19	29.9	0.40	35.8
0.20	30.3	0.41	36.1
0.21	30.5	0.42	36.3
0.22	30.9	0.43	36.5
0.23	31.1	0.44	36.8
0.24	31.4	0.45	37.1
0.25	31.7	0.46	37.3
0.26	32.0	0.47	37.6
0.27	32.3	0.48	37.8
0.28	32.6	0.49	38.1
0.29	32.8	0.50	38.4
0.30	33.1		



*Chlorurus spilurus*

**Probability of overfishing at various minimum sizes.**

Overfishing probability	$L_{C_{30}}$ (mm)	Overfishing probability	$L_{C_{30}}$ (mm)
0.10	301	0.31	264
0.11	299	0.32	262
0.12	297	0.33	260
0.13	295	0.34	257
0.14	293	0.35	257
0.15	290	0.36	255
0.16	288	0.37	253
0.17	286	0.38	251
0.18	284	0.39	249
0.19	284	0.40	246
0.20	282	0.41	244
0.21	279	0.42	242
0.22	277	0.43	240
0.23	275	0.44	238
0.24	273	0.45	235
0.25	273	0.46	233
0.26	271	0.47	231
0.27	268	0.48	229
0.28	266	0.49	227
0.29	264	0.50	224
0.30	301		

# Scarus dubius

Regal parrotfish, *lauia*

Scaridae (parrotfishes)



## Life history and other input parameters

Parameter	Value	SD	Unit	n	Source
$L_{inf}$	335	17	mm	-	Mean and SD: Nadon & Ault (2016) $L_{max}$ : 375 (2) from NWHI diver survey
$K$	0.63	0.22	yr <sup>-1</sup>		
$a_0$	-0.6	-	yr		
$L_{mat}$	232	26	mm	-	Mean and SD: Nadon & Ault (2016)
Longevity	13	4.5	yr	-	Mean and SD: Nadon & Ault (2016)
L-W $\alpha$	3.86e-7	-	-	-	Froese (1998)
L-W $\beta$	3.75				
$L_{S50}$	190	-	mm	-	Estimated.
$L_{S95}$	220				
$\bar{L}$ diver survey	287	5	mm	121	NOAA-CREP diver survey
$\bar{L}$ commercial	-	-	mm	-	-
$\bar{L}$ combined	-	-	mm	-	-
Max. depth	80	-	m	-	Pyle et al. (2016)
Federal waters	21	-	%	-	-

## Stock status and other output parameters

Parameter	Median	SD	Unit
$M$	0.26	0.09	yr <sup>-1</sup>
$F$	0.18	0.18	yr <sup>-1</sup>
$F_{30}$	0.31	0.10	yr <sup>-1</sup>
$F/F_{30}$	0.6	0.6	-
$SPR$	0.45	0.23	-
$SPR < 0.30$ iterations	26	-	%
$L_{C30}$	0	-	mm

Parameter	Median	SD	Unit
$B$ from catch	-	-	kg
$B$ from survey	32,840	7,868	kg
Commercial catch	-	-	kg
Recreational catch	-	-	kg
Total catch	-	-	kg
$C_{30}$ from catch	-	-	kg
$C_{30}$ from survey	7,690	2,660	kg

## General comments


Parrotfish catches are grouped at the family level and therefore there is no species-level commercial data.

Population abundance was fairly variable from year to year for this species, but appeared to be increasing from 2003 to 2015 before falling slightly in 2016. There was not enough yearly observation to generate a proper  $L_{bar}$  time series.

There are currently no published life history parameters for this species. We used the stepwise approach to generate these parameters using an  $L_{max}$  value (375 mm) from the NWHI. The  $L_{max}$  estimate from the MHI was nearly identical (378 mm). Furthermore, selectivity for parrotfishes had to be estimated, given the absence of catch data. A sensitivity run with  $L_{S50}$  at 230 mm and  $L_{S95}$  at 250 mm had little impact on the results ( $F$ : 0.21,  $SPR$ : 0.43).


The population biomass estimate may be biased downward given that this species' range extends to 68 m depth, which is beyond the depth of the diver surveys (30 m).

*Scarus dubius*





**Life history parameter distributions.**

*Scarus dubius*




**Abundance index from UVS (blue circles).**




**Size structure and average length time series from UVS ( $\pm$ SE).**

*Scarus dubius*




**Stock status parameter distributions ( $SPR$ : small bar shows 0.30 level).**



**$C_{30}$  (left) and population size (right) distributions.**

*Scarus dubius*



**Overfishing probability for a range of  $C_{30}$  levels (UVS – blue dotted line). *OFL* is represented by a small vertical bar.**

**Probability of overfishing for various  $C_{30}$  levels.**

Overfishing probability	$C_{30}$ from survey (1000 kg)	Overfishing probability	$C_{30}$ from survey (1000 kg)
0.10	4.7	0.31	6.4
0.11	4.8	0.32	6.5
0.12	4.9	0.33	6.5
0.13	5.0	0.34	6.6
0.14	5.1	0.35	6.7
0.15	5.2	0.36	6.7
0.16	5.3	0.37	6.8
0.17	5.4	0.38	6.9
0.18	5.4	0.39	7.0
0.19	5.5	0.40	7.0
0.20	5.6	0.41	7.1
0.21	5.7	0.42	7.2
0.22	5.8	0.43	7.2
0.23	5.8	0.44	7.3
0.24	5.9	0.45	7.4
0.25	6.0	0.46	7.4
0.26	6.0	0.47	7.5
0.27	6.1	0.48	7.6
0.28	6.2	0.49	7.6
0.29	6.3	0.50	7.7
0.30	6.3		

*Scarus dubius*

**Probability of overfishing at various minimum sizes.**

Overfishing probability	$L_{C_{30}}$ (mm)	Overfishing probability	$L_{C_{30}}$ (mm)
0.10	260	0.31	136
0.11	256	0.32	125
0.12	253	0.33	108
0.13	249	0.34	0
0.14	243	0.35	0
0.15	239	0.36	0
0.16	236	0.37	0
0.17	232	0.38	0
0.18	226	0.39	0
0.19	222	0.40	0
0.20	217	0.41	0
0.21	211	0.42	0
0.22	203	0.43	0
0.23	198	0.44	0
0.24	190	0.45	0
0.25	185	0.46	0
0.26	179	0.47	0
0.27	173	0.48	0
0.28	165	0.49	0
0.29	158	0.50	0
0.30	148		

# Scarus psittacus

Palenose parrotfish, *uhu*

Scaridae (parrotfishes)



## Life history and other input parameters

Parameter	Value	SD	Unit	n	Source
$L_{inf}$	291	16	mm	-	Mean and SD: Nadon & Ault (2016) $L_{max}$ : 326 (2) from NWHI diver survey
$K$	0.72	0.25	yr <sup>-1</sup>		
$a_0$	-0.6	-	yr		
$L_{mat}$	201	25	mm	-	Mean and SD: Nadon & Ault (2016)
Longevity	11	3	yr	-	Mean and SD: Nadon & Ault (2016)
L-W $\alpha$	5.02e-6	-	-	-	Kulbicki (2005)
L-W $\beta$	3.32				
$L_{S50}$	200	-	mm	-	Estimated
$L_{S95}$	210				
$\bar{L}$ diver survey	254	2	mm	344	NOAA-CREP diver survey
$\bar{L}$ commercial	-	-	mm	-	-
$\bar{L}$ combined	-	-	mm	-	-
Max. depth	48	-	m	-	NOAA-CREP BRUV survey
Federal waters	10	-	%	-	-

## Stock status and other output parameters

Parameter	Median	SD	Unit
$M$	0.29	0.10	yr <sup>-1</sup>
$F$	0.31	0.29	yr <sup>-1</sup>
$F_{30}$	0.45	0.16	yr <sup>-1</sup>
$F/F_{30}$	0.7	0.7	-
$SPR$	0.41	0.23	-
$SPR < 0.30$ iterations	32	-	%
$L_{C30}$	0	-	Mm

Parameter	Median	SD	Unit
$B$ from catch	-	-	kg
$B$ from survey	130,295	41,926	kg
Commercial catch	-	-	kg
Recreational catch	-	-	kg
Total catch	-	-	kg
$C_{30}$ from catch	-	-	kg
$C_{30}$ from survey	39,900	16,300	kg

## General comments

Parrotfish catches are grouped at the family level and therefore there is no species-level commercial data.

Population abundance has been increasing since 2003, with a potential small decrease in 2016.  $L_{bar}$  has remained stable from 2008 to 2016. Selectivity had to be estimated given the lack of catch data.


The life history parameters were available from a study conducted in American Samoa, but the  $L_{inf}$  for this study was deemed too small compared to  $L_{max}$  values in the NWHI (278 mm vs. 326 mm). This study also estimated a  $K$  parameter that seemed extreme for parrotfishes (1.65). The American Samoa LH parameters would have generated the following estimates:  $L_{inf}$ : 278 mm,  $K$ : 1.65,  $L_{mat}$ : 196 mm,  $M$ : 0.50,  $F_{30}$ : 0.82,  $F$ : 0.87,  $SPR$ : 0.30,  $C_{30}$  survey: 57,913 kg.

A sensitivity run with  $L_{S50}$  at 220 mm and  $L_{S95}$  at 240 mm had a slight impact on the results ( $F$ : 0.51,  $SPR$ : 0.31).

The population biomass estimate should be reasonably accurate given that this species extends only slightly beyond the maximum diver survey depth (30 m).




*Scarus psittacus*





**Life history parameter distributions.**

*Scarus psittacus*




**Abundance index from UVS (blue circles).**




**Size structure and average length time series from UVS ( $\pm$ SE).**

*Scarus psittacus*




**Stock status parameter distributions (*SPR*: small bar shows 0.30 level).**



**$C_{30}$  (left) and population size (right) distributions.**

*Scarus psittacus*



**Overfishing probability for a range of  $C_{30}$  levels (UVS – blue dotted line). *OFL* is represented by a small vertical bar.**

**Probability of overfishing for various  $C_{30}$  levels.**

Overfishing probability	$C_{30}$ from survey (1000 kg)	Overfishing probability	$C_{30}$ from survey (1000 kg)
0.10	22.0	0.31	32.4
0.11	22.7	0.32	32.8
0.12	23.4	0.33	33.2
0.13	23.9	0.34	33.6
0.14	24.4	0.35	34.0
0.15	24.9	0.36	34.4
0.16	25.4	0.37	34.8
0.17	26.0	0.38	35.2
0.18	26.5	0.39	35.7
0.19	27.0	0.40	36.0
0.20	27.5	0.41	36.3
0.21	28.0	0.42	36.7
0.22	28.4	0.43	37.0
0.23	28.9	0.44	37.4
0.24	29.5	0.45	37.8
0.25	29.9	0.46	38.3
0.26	30.4	0.47	38.7
0.27	30.8	0.48	39.2
0.28	31.1	0.49	39.5
0.29	31.6	0.50	39.9
0.30	31.9		

*Scarus psittacus*

**Probability of overfishing at various minimum sizes.**

Overfishing probability	$L_{c30}$ (mm)	Overfishing probability	$L_{c30}$ (mm)
0.10	240	0.31	180
0.11	236	0.32	174
0.12	234	0.33	170
0.13	230	0.34	168
0.14	228	0.35	162
0.15	226	0.36	158
0.16	222	0.37	154
0.17	220	0.38	146
0.18	218	0.39	140
0.19	214	0.40	132
0.20	212	0.41	124
0.21	208	0.42	118
0.22	206	0.43	106
0.23	204	0.44	88
0.24	200	0.45	0
0.25	198	0.46	0
0.26	196	0.47	0
0.27	194	0.48	0
0.28	190	0.49	0
0.29	186	0.50	0
0.30	182		

# Scarus rubroviolaceus

Redlip parrotfish, *uhu* 'ele'ele

Scaridae (parrotfishes)



## Life history and other input parameters

Parameter	Value	SD	Unit	n	Source
$L_{inf}$	563	23	mm	182	Mean and SD: Howard (2008)
$K$	0.288	0.041	yr <sup>-1</sup>		
$a_0$	-0.809	-	yr		
$L_{mat}$	374	9	mm	182	Mean and SD: Howard (2008)
Longevity	22	2	yr	182	Mean and SD: Howard (2008)
L-W $\alpha$	7.89e-6	-	-	-	Smith & Dalzell (1993)
L-W $\beta$	3.11				
$L_{S50}$	240	-	mm	-	Estimated.
$L_{S95}$	260				
$\bar{L}$ diver survey	401	8	mm	1054	NOAA-CREP diver survey
$\bar{L}$ commercial	-	-	mm	-	-
$\bar{L}$ combined	-	-	mm	-	-
Max. depth	68	-	m	-	Pyle et al. (2016)
Federal waters	21	-	%	-	-

## Stock status and other output parameters

Parameter	Median	SD	Unit
$M$	0.15	0.01	yr <sup>-1</sup>
$F$	0.18	0.04	yr <sup>-1</sup>
$F_{30}$	0.15	0.01	yr <sup>-1</sup>
$F/F_{30}$	1.2	0.3	-
$SPR$	0.26	0.08	-
$SPR < 0.30$ iterations	68	-	%
$LC_{30}$	302	-	mm

Parameter	Median	SD	Unit
$B$ from catch	-	-	kg
$B$ from survey	623,532	80,385	kg
Commercial catch	-	-	kg
Recreational catch	-	-	kg
Total catch	-	-	kg
$C_{30}$ from catch	-	-	kg
$C_{30}$ from survey	82,500	11,900	kg

## General comments


Parrotfish catches are grouped at the family level and therefore there is no species-level commercial data.

Population abundance has been fluctuating between 2005 and 2016, with no clear temporal pattern.  $L_{bar}$  from UVS has also been fluctuating but has remained stable overall. As with other parrotfishes, selectivity had to be estimated given the lack of catch data. A sensitivity run with  $L_{S50}$  at 260 mm and  $L_{S95}$  at 300 mm had little impact on the results ( $F$ : 0.19,  $SPR$ : 0.25).

The life history parameters for this species came from an in-depth local study and there was little reasons to doubt the validity of these parameters. As an exercise, the stepwise approach was used to generate alternate numbers (using an  $L_{max}$  of 681 mm from the NWHI) and generated the following values:  $L_{inf}$ : 580 mm,  $K$ : 0.26,  $M$ : 0.15,  $L_{mat}$ : 403 mm,  $F$ : 0.17,  $SPR$ : 0.24.  $C_{30}$  survey: 75,920 kg. These values are fairly close to the original analyses.


The population biomass estimate may be biased downward given that this species' range extends to 68 m depth, which is beyond the depth of the diver surveys (30 m).

*Scarus rubroviolaceus*





**Life history parameter distributions.**

*Scarus rubroviolaceus*




**Abundance index from UVS (blue circles,  $\pm$ SE).**




**Size structure and average length time series from UVS ( $\pm$ SE).**



*Scarus rubroviolaceus*




Stock status parameter distributions ( $SPR$ : small bar shows 0.30 level).



$C_{30}$  (left) and population size (right) distributions.

*Scarus rubroviolaceus*



**Overfishing probability for a range of  $C_{30}$  levels (UVS – blue dotted line). *OFL* is represented by a small vertical bar.**

**Probability of overfishing for various  $C_{30}$  levels.**

Overfishing probability	$C_{30}$ from survey (1000 kg)	Overfishing probability	$C_{30}$ from survey (1000 kg)
0.10	67.9	0.31	76.7
0.11	68.5	0.32	77.0
0.12	68.9	0.33	77.3
0.13	69.5	0.34	77.6
0.14	70.1	0.35	77.9
0.15	70.6	0.36	78.3
0.16	71.2	0.37	78.6
0.17	71.6	0.38	78.9
0.18	72.0	0.39	79.1
0.19	72.3	0.40	79.4
0.20	72.7	0.41	79.7
0.21	73.1	0.42	80.0
0.22	73.6	0.43	80.3
0.23	73.9	0.44	80.7
0.24	74.4	0.45	81.0
0.25	74.7	0.46	81.3
0.26	75.0	0.47	81.5
0.27	75.4	0.48	81.8
0.28	75.7	0.49	82.2
0.29	76.1	0.50	82.5
0.30	76.4		

*Scarus rubroviolaceus*

**Probability of overfishing at various minimum sizes.**

Overfishing probability	$L_{C_{30}}$ (mm)	Overfishing probability	$L_{C_{30}}$ (mm)
0.10	386	0.31	355
0.11	384	0.32	353
0.12	382	0.33	353
0.13	379	0.34	350
0.14	379	0.35	348
0.15	377	0.36	348
0.16	374	0.37	346
0.17	372	0.38	343
0.18	370	0.39	341
0.19	370	0.40	338
0.20	367	0.41	338
0.21	365	0.42	336
0.22	362	0.43	334
0.23	360	0.44	334
0.24	360	0.45	331
0.25	358	0.46	329
0.26	355	0.47	326
0.27	353	0.48	324
0.28	350	0.49	324
0.29	350	0.50	322
0.30	348		

# Cephalopholis argus

Peacock grouper, *roi*  
Serranidae (groupers)



## Life history and other input parameters

Parameter	Value	SD	Unit	N	Source
$L_{inf}$	506	11	mm	590	Mean: Donovan (2013), SD: Kritzer (2001)
$K$	0.075	0.007	yr <sup>-1</sup>		
$a_0$	-6.5	-	yr		
$L_{mat}$	268	8	mm	100?	Mean: Myers (1999), SD: Nadon (unpublished)
Longevity	25	1.4	yr	590	Mean: Donovan (2013), SD: Kritzer (2001)
L-W $\alpha$	2.05e-5	-	-	-	Kulbicki (2005)
L-W $\beta$	2.99	-	-	-	
$L_{S50}$	270	-	mm	-	DAR commercial data
$L_{S95}$	310				
$\bar{L}$ diver survey	373	4	mm	614	NOAA-CREP diver survey
$\bar{L}$ commercial	374	4	mm	2234	DAR commercial data
$\bar{L}$ combined	374	4	mm	-	-
Max. depth	80	-	m	-	Pyle et al. (2016)
Federal waters	21	-	%	-	-

## Stock status and other output parameters

Parameter	Median	SD	Unit
$M$	0.13	0.01	yr <sup>-1</sup>
$F$	0.02	0.02	yr <sup>-1</sup>
$F_{30}$	0.16	0.01	yr <sup>-1</sup>
$F/F_{30}$	0.1	0.1	-
$SPR$	0.80	0.11	-
$SPR < 0.30$ iterations	0	-	%
$L_{C30}$	-	-	mm

Parameter	Median	SD	Unit
$B$ from catch	232,000	494,000	kg
$B$ from survey	777,397	56,961	kg
Commercial catch	830	980	kg
Recreational catch	3,192	6,708	kg
Total catch	4,552	5,594	kg
$C_{30}$ from catch	33,300	72,200	kg
$C_{30}$ from survey	111,000	11,300	kg

## General comments


Note: this species is non-native and considered invasive. It was introduced in 1956 from Tahiti.

Population abundance has been stable for the last 10 years, except for a higher estimate in the first year. The commercial catch increased steadily from 2003 to 2011 before going back down. The recreational catch was too variable to infer on any temporal trends. The  $L_{bar}$  from the commercial and recreational sector were identical and stable.

The growth parameters for this species came from an in-depth local study. However, the maturity parameter came from a less reliable source. Furthermore, the growth curve did not have juvenile age estimates which explains the highly negative  $a_0$  and the resulting low  $K$  value. We could not run the stepwise approach for this species given that the grouper family is not currently available for this method.


Population estimate from the catch was about a third the size of the population estimate from diver surveys. This is not entirely surprising given that the catch estimate seemed low. For example, the average weight of this species is around 1 kg, which would suggest that only 3000 individuals are caught recreationally every year. This would mean only 8 individuals are caught on a daily basis across the entire island chain, which seems unlikely. The diver survey population should be fairly representative given that this is a commonly encountered species that is easily identified, although it is likely biased downward given that this species' range extends beyond diver depths, to 80 m. Note: no  $L_{C30}$  could be generated given the very low fishing mortality rates.

*Cephalopholis argus*




**Life history parameter distributions.**

*Cephalopholis argus*




**Abundance index from UVS (blue circles,  $\pm$ SE) and total catch time series from recreational (green squares) and commercial (orange triangles) sectors.**




**Size structure from commercial catch (top left) and UVS (top right). Average length time series (blue circles – UVS, orange triangles – commercial data,  $\pm$ SE).**

*Cephalopholis argus*



**Stock status parameter distributions (*SPR*: small bar shows 0.30 level).**


*Cephalopholis argus*



$C_{30}$  and current total catch (left) and population size (right) distributions.



*Cephalopholis argus*



**Overfishing probability for a range of  $C_{30}$  levels (catch - orange dashed line, UVS – blue dotted line). *OFLs* are represented by small vertical bars.**

**Probability of overfishing for various  $C_{30}$  levels.**

Overfish. probability	$C_{30}$ from catch (1000 kg)	$C_{30}$ from survey (1000 kg)	Overfish. probability	$C_{30}$ from catch (1000 kg)	$C_{30}$ from survey (1000 kg)
0.10	8.9	97.1	0.31	19.8	105.4
0.11	9.3	97.6	0.32	20.4	105.7
0.12	9.7	98.1	0.33	21.0	106.0
0.13	10.3	98.6	0.34	21.7	106.2
0.14	10.8	99.2	0.35	22.3	106.5
0.15	11.3	99.7	0.36	22.9	106.8
0.16	11.8	100.1	0.37	23.5	107.1
0.17	12.2	100.4	0.38	24.2	107.4
0.18	12.7	100.9	0.39	24.8	107.7
0.19	13.2	101.2	0.40	25.5	108.0
0.20	13.6	101.6	0.41	26.2	108.3
0.21	14.1	101.9	0.42	26.9	108.6
0.22	14.5	102.4	0.43	27.8	108.9
0.23	15.2	102.7	0.44	28.6	109.2
0.24	15.7	103.2	0.45	29.3	109.4
0.25	16.3	103.5	0.46	30.0	109.7
0.26	16.8	103.9	0.47	30.7	110.0
0.27	17.4	104.1	0.48	31.4	110.4
0.28	17.9	104.5	0.49	32.3	110.7
0.29	18.6	104.8	0.50	33.3	110.9
0.30	19.0	105.1			

# Monotaxis grandoculis

Bigeye bream, *mu*

Lethrinidae (emperors)



## Life history and other input parameters

Parameter	Value	SD	Unit	n	Source
$L_{inf}$	527	22	mm	-	Mean and SD: Nadon & Ault (2016) $L_{max}$ : 597 (19) from NWHI diver survey
$K$	0.37	0.18	yr <sup>-1</sup>		
$a_0$	-0.5	-	yr		
$L_{mat}$	389	28	mm	-	Mean and SD: Nadon & Ault (2016)
Longevity	21	9	yr	-	Mean and SD: Nadon & Ault (2016)
L-W $\alpha$	1.93e-5	-	-	-	Smith & Dalzell (1993)
L-W $\beta$	3.02				
$L_{S50}$	250	-	mm	-	DAR commercial data
$L_{S95}$	300				
$\bar{L}$ diver survey	359	7	mm	278	NOAA-CREP diver survey
$\bar{L}$ commercial	448	4	mm	1951	DAR commercial data
$\bar{L}$ combined	425	5	mm	-	-
Max. depth	101	-	m	-	Pyle et al. (2016)
Federal waters	22	-	%	-	-

## Stock status and other output parameters

Parameter	Median	SD	Unit
$M$	0.15	0.06	yr <sup>-1</sup>
$F$	0.12	0.12	yr <sup>-1</sup>
$F_{30}$	0.16	0.07	yr <sup>-1</sup>
$F/F_{30}$	0.8	0.6	-
$SPR$	0.38	0.22	-
$SPR < 0.30$ iterations	36	-	%
$L_{C30}$	0	-	Mm

## Management information

Parameter	Median	SD	Unit
$B$ from catch	28,600	82,200	kg
$B$ from survey	231,797	49,031	kg
Commercial catch	1,346	1,270	kg
Recreational catch	1,381	1,129	kg
Total catch	2,998	1,668	kg
$C_{30}$ from catch	3,950	11,700	kg
$C_{30}$ from survey	31,700	12,800	kg


## General comments

The population abundance has been stable except for a drastic jump in the last survey year which was likely a statistical outlier. Commercial  $L_{bar}$  have been generally steady and higher than the UVS  $L_{bar}$ . It is not entirely clear why there is such a discrepancy between  $L_{bar}$  sources. The commercial catch increased drastically between 2008 and 2011 before falling quickly again to its original level. The recreational catch was too variable to infer on temporal trends.

There are currently no published life history parameters for this species. We used the stepwise approach to generate LH parameters, using an  $L_{max}$  value from NWHI diver surveys. A sensitivity run using an alternate  $L_{max}$  value from the MHI UVS (556 mm) generated the following results:  $L_{inf}$ : 493 mm,  $K$ : 0.45,  $M$ : 0.16,  $F_{30}$ : 0.18,  $F$ : 0.07,  $SPR$ : 0.59,  $C_{30}$  catch: 6,828 kg,  $C_{30}$  survey: 35,538 kg.


The total catch estimate seemed fairly low for a relatively common and prized species. The population biomass estimate from diver surveys was much high than the catch-derived estimate. The diver survey biomass estimate, and the  $C_{30}$  generated from it, is likely more reliable than the catch data for this species.

*Monotaxis grandoculis*




**Life history parameter distributions.**

*Monotaxis grandoculis*




**Abundance index from UVS (blue circles, ±SE) and total catch time series from recreational (green squares) and commercial (orange triangles) sectors.**




**Size structure from commercial catch (top left) and UVS (top right). Average length time series (blue circles – UVS, orange triangles – commercial data, ±SE).**

*Monotaxis grandoculis*




**Stock status parameter distributions (*SPR*: small bar shows 0.30 level).**

*Monotaxis grandoculis*



$C_{30}$  and current total catch (left) and population size (right) distributions.

*Monotaxis grandoculis*



**Overfishing probability for a range of  $C_{30}$  levels (catch - orange dashed line, UVS – blue dotted line). *OFLs* are represented by small vertical bars.**

**Probability of overfishing for various  $C_{30}$  levels.**

Overfish. probability	$C_{30}$ from catch (1000 kg)	$C_{30}$ from survey (1000 kg)	Overfish. probability	$C_{30}$ from catch (1000 kg)	$C_{30}$ from survey (1000 kg)
0.10	1.50	18.7	0.31	2.62	26.0
0.11	1.55	19.2	0.32	2.67	26.3
0.12	1.60	19.7	0.33	2.72	26.6
0.13	1.65	20.2	0.34	2.79	26.9
0.14	1.71	20.5	0.35	2.85	27.2
0.15	1.76	20.9	0.36	2.90	27.5
0.16	1.82	21.3	0.37	2.98	27.8
0.17	1.87	21.6	0.38	3.05	28.2
0.18	1.92	21.9	0.39	3.11	28.5
0.19	1.97	22.2	0.40	3.16	28.7
0.20	2.04	22.6	0.41	3.23	29.1
0.21	2.09	22.9	0.42	3.32	29.4
0.22	2.14	23.3	0.43	3.39	29.7
0.23	2.18	23.6	0.44	3.49	30.0
0.24	2.24	23.9	0.45	3.58	30.2
0.25	2.28	24.2	0.46	3.64	30.5
0.26	2.34	24.5	0.47	3.72	30.8
0.27	2.39	24.9	0.48	3.79	31.0
0.28	2.45	25.1	0.49	3.86	31.3
0.29	2.51	25.4	0.50	3.95	31.7
0.30	2.56	25.7			

*Monotaxis grandoculis*

**Probability of overfishing at various minimum sizes.**

Overfishing probability	$L_{C_{30}}$ (mm)	Overfishing probability	$L_{C_{30}}$ (mm)
0.10	415	0.31	288
0.11	410	0.32	278
0.12	405	0.33	265
0.13	400	0.34	255
0.14	398	0.35	250
0.15	390	0.36	238
0.16	385	0.37	225
0.17	380	0.38	215
0.18	372	0.39	205
0.19	368	0.40	190
0.20	360	0.41	175
0.21	355	0.42	160
0.22	350	0.43	140
0.23	345	0.44	122
0.24	340	0.45	85
0.25	332	0.46	0
0.26	325	0.47	0
0.27	320	0.48	0
0.28	312	0.49	0
0.29	302	0.50	0
0.30	295		



# *Myripristis berndti*

Bigscale soldierfish, 'u'u  
Holocenridae (soldierfishes)



## Life history and other input parameters

Parameter	Value	SD	Unit	n	Source
$L_{inf}$	271	14	mm	126	Mean: Craig & Franklin (2008), SD: Kritzer (2001)
$K$	0.148	0.028	yr <sup>-1</sup>		
$a_0$	-4.48	-	yr		
$L_{mat}$	175	5	mm	100?	Mean: Murty (2002) , SD: Nadon (unpublished)
Longevity	27	3.2	yr	126	Mean: Craig & Franklin (2008), SD: Kritzer (2001)
L-W $\alpha$	2.14e-5	-	-	-	Kulbicki (2005)
L-W $\beta$	3.00	-	-	-	
$L_{S50}$	170	-	mm	-	Estimated with some HMRFS data.
$L_{S95}$	180				
$\bar{L}$ diver survey	224	4	mm	824	NOAA-CREP diver survey
$\bar{L}$ commercial	-	-	mm	-	-
$\bar{L}$ combined	-	-	mm	-	-
Max. depth	159	-	m	-	Pyle et al. (2016)
Federal waters	22	-	%	-	-

## Stock status and other output parameters

Parameter	Median	SD	Unit
$M$	0.12	0.01	yr <sup>-1</sup>
$F$	0.06	0.04	yr <sup>-1</sup>
$F_{30}$	0.16	0.02	yr <sup>-1</sup>
$F/F_{30}$	0.4	0.3	-
$SPR$	0.59	0.17	-
$SPR < 0.30$ iterations	3	-	%
$L_{C30}$	-	-	mm

Parameter	Median	SD	Unit
$B$ from catch	-	-	kg
$B$ from survey	260,111	47,848	kg
Commercial catch	-	-	kg
Recreational catch	-	-	kg
Total catch	-	-	kg
$C_{30}$ from catch	-	-	kg
$C_{30}$ from survey	36,100	7,700	kg

## General comments


This species is reported with other soldierfishes in the commercial data set, preventing the use of catch data for the current analyses.

Population abundance for this species appear to have been relatively stable.  $L_{bar}$  were higher in the earlier survey years (2005–2008) but appear to have declined in 2009, staying relatively stable since then. Selectivity had to be estimated given the lack of catch data. However, there were some length estimates in the HMRFS recreational data set ( $n = 15$ ) which provided some basis for our estimate. A sensitivity run with  $L_{S50}$  at 200 mm and  $L_{S95}$  at 220 mm generated similar values ( $F$ : 0.07 and  $SPR$ : 0.63).

The growth parameters came from a local study. However, the maturity parameter came from a study in India for a different species (*M. murdjan*). Further, the growth study lacked age estimates for juveniles resulting in a fairly negative  $a_0$  parameter and low  $K$  estimate. We could not run the stepwise approach for this species given that the soldierfish family is not currently available for this method.


This species occurs at depths much greater than the maximum diver survey depth (159 m vs. 30 m) and it is likely that the population size estimate is bias downward.

*Myripristis berndti*





**Life history parameter distributions.**

*Myripristis berndti*




**Abundance index from UVS (blue circles,  $\pm$ SE).**




**Size structure and average length time series from UVS ( $\pm$ SE).**

*Myripristis berndti*




**Stock status parameter distributions ( $SPR$ : small bar shows 0.30 level).**



**$C_{30}$  (left) and population size (right) distributions.**

*Myripristis berndti*



**Overfishing probability for a range of  $C_{30}$  levels (UVS – blue dotted line). *OFL* is represented by a small vertical bar.**

**Probability of overfishing for various  $C_{30}$  levels.**

Overfishing probability	$C_{30}$ from survey (1000 kg)	Overfishing probability	$C_{30}$ from survey (1000 kg)
0.10	26.7	0.31	32.4
0.11	27.1	0.32	32.6
0.12	27.5	0.33	32.8
0.13	27.9	0.34	33.0
0.14	28.2	0.35	33.2
0.15	28.5	0.36	33.4
0.16	28.8	0.37	33.6
0.17	29.1	0.38	33.8
0.18	29.4	0.39	34.0
0.19	29.7	0.40	34.2
0.20	29.9	0.41	34.4
0.21	30.2	0.42	34.6
0.22	30.4	0.43	34.8
0.23	30.6	0.44	35.1
0.24	30.8	0.45	35.2
0.25	31.1	0.46	35.4
0.26	31.3	0.47	35.5
0.27	31.6	0.48	35.8
0.28	31.8	0.49	35.9
0.29	32.0	0.50	36.1
0.30	32.2		