

NOAA Technical Memorandum NWS WR-225

CLIMATE OF SANTA BARBARA, CALIFORNIA

Gary Ryan Weather Service Office Santa Maria, California

December 1994

U.S. DEPARTMENT OF COMMERCE

National Oceanic and Atmospheric Administration National Weather Service



NOAA TECHNICAL MEMORANDA National Weather Service, Western Region Subseries

The National Weather Service (NWS) Western Region (WR) Subseries provides an informal medium for the documentation and quick dissemination of results not appropriate, or not yet ready, for formal publication. The series is used to report on work in progress, to describe technical procedures and practices, or to relate progress to a limited audience. These Technical Memoranda will report on investigations devoted primarily to regional and local problems of interest mainly to personnel, and hence will not be widely distributed.

Papers 1 to 25 are in the former series, ESSA Technical Memoranda, Western Region Technical Memoranda (WRTM); papers 24 to 59 are in the former series, ESSA Technical Memoranda, Weather Bureau Technical Memoranda (WBTM). Beginning with 60, the papers are part of the series, NOAA Technical Memoranda NWS. Out-of-print memoranda are not listed.

Papers 2 to 22, except for 5 (revised edition), are available from the National Weather Service Western Region, Scientific Services Division, P.O. Box 11188, Federal Building, 125 South State Street, Salt Lake City, Utah 84147. Paper 5 (revised edition), and all others beginning with 25 are available from the National Technical Information Service, U.S. Department of Commerce, Sills Building, 5285 Port Royal Road, Springfield, Virginia 22161. Prices vary for all paper copies; microfiche are \$3.50. Order by accession number shown in parentheses at end of each entry.

ESSA Technical Memoranda (WRTM)

- Climatological Precipitation Probabilities. Compiled by Lucianne Miller, December 1965. Western Region Pre- and Post-FP-3 Program, December 1, 1965, to February 20, 1966. Edward D. Diemer, March 1966.
- Edward D. Diemer, March 1966. Station Descriptions of Local Effects on Synoptic Weather Patterns. Philip Williams, Jr., April 1966 (Revised November 1967, October 1969). (PB-17800) Interpreting the RAREP. Herbert P. Benner, May 1966 (Revised January 1967). Some Electrical Processes in the Atmosphere. J. Latham, June 1966. A Digitalized Summary of Radar Echoes within 100 Miles of Sacramento, California. J. A. Youngberg and L. B. Overaas, December 1966. An Objective Aid for Forecasting the End of East Winds in the Columbia Gorge, July through October. D. John Coparanis, April 1967. Derivation of Radar Horizons in Mountainous Terrain. Roger G. Pappas, April 1967.

ESSA Technical Memoranda, Weather Bureau Technical Memoranda (WBTM)

- Verification of Operation Probability of Precipitation Forecasts, April 1966-March 1967.
 W. Dickey, October 1967. (PB-176240)
 A Study of Winds in the Lake Mead Recreation Area. R. P. Augulis, January 1968. (PB-177830)
- 177830) Weather Extremes. R. J. Schmidli, April 1968 (Revised March 1986). (PB86 177672/AS). (Revised October 1991 PB92-115062/AS) Small-Scale Analysis and Prediction. Philip Williams, Jr., May 1968. (PB178425) Numerical Weather Prediction and Synoptic Meteorology. CPT Thomas D. Murphy, USAF, March 1065. (AD 627365)
- May 1968. (AD 673365)
- (AD 01305)
 (cipitation Detection Probabilities by Salt Lake ARTC Radars. Robert K. Belesky, July
 (PB 179084) 1968. Probability Forecasting-A Problem Analysis with Reference to the Portland Fire Weather District. Harold S. Ayer, July 1968. (PB 179289)
- Temperature Trends in Sacramento-Another Heat Island. Anthony D. Lentini, February 1969. (PB 183055) Disposal of Logging Residues Without Damage to Air Quality. Owen P. Cramer, March 1959. (PB 183057)
- per-Air Lows Over Northwestern United States. A.L. Jacobson, April 1969. PB 184296) e Man-Machine Mix in Applied Weather Forecasting in the 1970s. L.W. Snellman, August 40 Ť 1969. (PB 185068)
- Forecasting Maximum Temperatures at Helena, Montana. David E. Olsen, October 1969. (PB 185762) Estimated Return Periods for Short-Duration Precipitation in Arizona. Paul C. Kangieser,
- October 1969. (PB 187763) Applications of the Net Radiometer to Short-Range Fog and Stratus Forecasting at Eugene, Oregon. L. Yee and E. Bates. December 1969. (PB 190476) Statistical Analysis as a Flood Routing Tool. Robert J.C. Burnash, December 1969. (PB
- 188744)
- 188/44) Tsunami. Richard P. Augulis, February 1970. (PB 190157) Predicting Precipitation Type. Robert J.C. Burnash and Floyd E. Hug, March 1970. (PB
- 190962) Statistical Report on Aeroallergens (Pollens and Molds) Fort Huschuca, Arizona, 1969. Wayne S. Johnson, April 1970. (PB 191743) Western Region Sea State and Surf Forecaster's Manual. Gordon C. Shields and Gerald B. Burdwell, July 1970. (PB 193102) Sacramento Weather Radar Climatology. R.G. Pappas and C. M. Veliquette, July 1970. (PB 190240)
- 193347)

- 193347) A Refinement of the Vorticity Field to Delineate Areas of Significant Precipitation. Barry B. Aronovitch, August 1970. Application of the SSARR Model to a Basin without Discharge Record. Vail Schermerhorn and Donal W. Kuehl, August 1970. (PB 194394) Areal Coverage of Precipitation in Northwestern Utah. Philip Williams, Jr., and Werner J. Heck, September 1970. (PB 194389) Preliminary Report on Agricultural Field Burning vs. Atmospheric Visibility in the Willamette Valley of Oregon. Earl M. Bates and David O. Chilcote, September 1970. (PB 194210)
- 194710)Air Pollution by Jet Aircraft at Seattle-Tacoma Airport. Wallace R. Donaldson, October
- Air Pollution by Jet Aircraft at Seattle-Tacoma Airport. Wallace R. Donauson, October 1970. (COM 71 60017) Application of PE Model Forecast Parameters to Local-Area Forecasting. Leonard W. Snellman, October 1970. (COM 71 00016) An Aid for Forecasting the Minimum Temperature at Medford, Oregon, Arthur W. Fritz, October 1970. (COM 71 00120) 700-mb Warm Air Advection as a Forecasting Tool for Montana and Northern Idaho. Norris E. Woerner, February 1971. (COM 71 00349) Wind and Weather Regimes at Great Falls, Montana. Warren B. Price, March 1971. Climate of Sacramento, California. Tony Martini, April 1990. (Fifth Revision) (PB89 207781/AS)

- A Preliminary Report on Correlation of ARTCC Radar Echoes and Precipitation. Wilbur K. Hall, June 1971. (COM 71 00829) National Weather Service Support to Soaring Activities. Ellis Burton, August 1971. (COM
- 71 00956)
- Western Region Synoptic Analysis-Problems and Methods. Philip Williams, Jr., February 1972. (COM 72 10433) Thunderstorms and Hail Days Probabilities in Nevada. Clarence M. Sakamoto, April 1972. (COM 72 10554)

- A Study of the Low Level Jet Stream of the San Joaquin Valley. Ronald A. Willis and Philip Williams, Jr., May 1972. (COM 72 10707) Monthly Climatological Charts of the Behavior of Fog and Low Stratus at Los Angeles International Airport. Donald M. Gales, July 1972. (COM 72 11140) A Study of Radar Echo Distribution in Arizona During July and August. John E. Hales, Jr., July 1972. (COM 72 11126)

- International Airport. Donald M. Gales, July 1972. (COM 72 11140) A Study of Radar Echo Distribution in Arizona During July and August. John E. Hales, Jr., July 1972. (COM 72 11136) Forecasting Precipitation at Bakersfield, California, Using Pressure Gradient Vectors. Earl T. Riddiough, July 1972. (COM 72 11146) Climate of Stockton, California. Robert C. Nelson, July 1972. (COM 72 10920) Estimation of Number of Days Above or Below Selected Temperatures. Clarence M. Sakamoto, October 1972. (COM 72 10021) An Aid for Forecasting Summer Maximum Temperatures at Seattle, Washington. Edgar G. Johnson, November 1972. (COM 73 10150) Flash Flood Forecasting and Warning Program in the Western Region. Philip Williams, Jr., Chester L. Glenn, and Roland L. Raetz, December 1972, (Revised March 1978). (COM 73 10251) 10251)
- Comparison of Manual and Semiautomatic Methods of Digitizing Analog Wind Records.
 Glenn E. Rasch, March 1973. (COM 73 10669)
 Conditional Probabilities for Sequences of Wet Days at Phoenix, Arizona. Paul C. Kangieser, June 1973. (COM 73 11264)
 A Refinement of the Use of K-Values in Forecasting Thunderstorms in Washington and Oregon. Robert Y.G. Lee, June 1973. (COM 73 11276)
 Objective Forecast Precipitation Over the Western Region of the United States. Julia N. Paegle and Larry P. Kierulff, September 1973. (COM 73 11946/3AS)
 Arizona 'Eddy' Tornadoes. Robert S. Ingram, October 1973. (COM 73 10465)
 Smoke Management in the Willamette Valley. Earl M. Bates, May 1974. (COM 74 11277/AS)
- 92

- 11277/AS) An Operational Evaluation of 500-mb Type Regression Equations. Alexander E. MacDonald, June 1974. (COM 74 11407/AS) Conditional Probability of Visibility Less than One-Half Mile in Radiation Fog at Fresno, California. John D. Thomas, August 1974. (COM 74 11555/AS) Climate of Flagstaff, Arizona. Paul W. Sorenson, and updated by Reginald W. Preston, January 1987. (PB87 143160/AS) Map type Precipitation Probabilities for the Western Region. Glenn E. Rasch and Alexander E. MacDonald, February 1975. (COM 75 10428/AS) Eastern Pacific Cut-Off Low of April 21-28, 1974. William J. Alder and George R. Miller, January 1976. (PB 250 711/AS)

- January 1976. (PB 250 711/AS) Study on a Significant Precipitation Episode in Western United States. Ira S. Brenner, April 1976. (COM 75 10719/AS) A Study of Flash Flood Susceptibility-A Basin in Southern Arizona. Gerald Williams, August 1975. (COM 75 11360/AS) A Set of Rules for Forecasting Temperatures in Napa and Sonoma Counties. Wesley L. Tuft, October 1975. (PB 246 902/AS) Application of the National Weather Service Flash-Flood Program in the Western Region. Gerald Williams, January 1976. (PB 250 053/AS) Objective Aids for Forecasting Minimum Temperatures at Reno. Nevada, During the Summer Months. Christopher D. Hill, January 1976. (PB 252 866/AS) Forecasting the Mono Wind. Charles P. Ruscha, Jr., February 1976. (PB 254 650) Use of MOS Forecast Parameters in Temperature Forecasting. John C. Plankinton, Jr., March 1976. (PB 256 49) Map Types as Aids in Using MOS PoPs in Western United States. Ira S. Brenner, August 1976. (PB 259 594) Other Kinds of Wind Shear. Christopher D. Hill, August 1976. (PB 260 437/AS) Forecasting North Winds in the Upper Sacramento Valley and Adjoining Foreate

- Other Kinds of Wind Shear. Christopher D. Hill, August 1976. (PB 260 437/AS) Forecasting North Winds in the Upper Sacramento Valley and Adjoining Forests. Christopher E. Fontana, September 1976. (PB 273 677/AS) Cool Inflow as a Weakening Influence on Eastern Pacific Tropical Cyclones. William J. Denney, November 1976. (PB 284 655/AS) The MAN/MOS Program. Alexander E. MacDonald, February 1977. (PB 265 941/AS) Winter Sesson Minimum Temperature Formula for Bakersfield, California, Using Multiple Regression. Michael J. Oard, February 1977. (PB 273 694/AS) A Study of Wind Gusts on Lake Mead. Bradley Colman, April 1977. (PB 268 847) The Relative Frequency of Cumulonimbus Clouds at the Nevada Test Site as a Function of K-Value. R.F. Quiring, April 1977. (PB 272 831) Moisture Distribution Modification by Upward Vertical Motion. Ira S. Brenner, April 1977.

- MOSTURE INSTITUTION INCOMPANY OF A STATE OF

- (PB 271 299/AS) Climatological Prediction of Cumulonimbus Clouds in the Vicinity of the Yucca Flat Weather Station. R.F. Quiring, June 1977. (PB 271 704/AS) A Method for Transforming Temperature Distribution to Normality. Morris S. Webb, Jr., June 1977. (PB 271 742/AS) Statistical Guidance for Prediction of Eastern North Pacific Tropical Cyclone Motion Part I. Charles J. Neumann and Preston W. Leftwich, August 1977. (PB 272 661) Statistical Guidance on the Prediction of Eastern North Pacific Tropical Cyclone Motion -Part II. Preston W. Leftwich and Charles J. Neumann, August 1977. (PB 273 155/AS) Climate of San Francisco. E. Jan Null, February 1978. Revised by George T. Pericht, April 1988. (PB88 208624/AS)
- Development of a Probability Equation for Winter-Type Precipitation Patterns in Great Falls, Montana. Kenneth B. Mielke, February 1978. (PB 281 387/AS) Hand Calculator Program to Compute Parcel Thermal Dynamics. Dan Gudgel, April 1978. (PB 283 08/AS)

- Hand Calculator Frogram to Compute Farter Internal Dynamics. Dat Gouger, 1991 2010.
 (PB 283 080/AS)
 Fire whirls. David W. Goens, May 1978. (PB 283 866/AS)
 Flash-Flood Procedure. Ralph C. Hatch and Gerald Williams, May 1978. (PB 286 014/AS)
 Automated Fire-Weather Forecasts. Mark A. Mollner and David E. Olsen, September 1978. 131
- (FD 229 510/AS) Estimates of the Effects of Terrain Blocking on the Los Angeles WSR-74C Weather Radar. R.G. Pappas, R.Y. Lee, B.W. Finke, October 1978. (PB 289767/AS) Spectral Techniques in Ocean Wave Forecasting. John A. Jannuzzi, October 1978.
- (PB291317/AS)

- (PB291317/AS) Solar Radiation. John A. Jannuzzi, November 1978. (PB29195/AS) Application of a Spectrum Analyzer in Forecasting Ocean Swell in Southern California Coastal Waters. Lawrence P. Kierulff, January 1979. (PB292716/AS) Basic Hydrologic Principles. Thomas L. Dietrich, January 1979. (PB292247/AS) LFM 24-Hour Prediction of Eastern Pacific Cyclones Refined by Satellite Images. John R. Zimmerman and Charles P. Ruscha, Jr., January 1979. (PB294224/AS) A Simple Analysis/Diagnosis System for Real Time Evaluation of Vertical Motion. Scott Heflick and James R. Fors, February 1979. (PB294224/AS) A Simple Analysis/Diagnosis System for Real Time Evaluation of Vertical Motion. Scott Heflick and James R. Fors, February 1979. (PB294216/AS) Aids for Forecasting Minimum Temperature in the Wenatchee Frost District. Robert S. Robinson, April 1979. (PB298339/AS) Influence of Cloudiness on Summertime Temperatures in the Eastern Washington Fire Weather district. James Holcomb, April 1979. (PB298674/AS) Comparison of LFM and MFM Precipitation Guidance for Nevada During Doreen. Christopher Hill, April 1979. (PB298613/AS)

NOAA Technical Memorandum NWS WR-225

CLIMATE OF SANTA BARBARA, CALIFORNIA

Gary Ryan Weather Service Office Santa Maria, California

December 1994

UNITED STATES DEPARTMENT OF COMMERCE Ronald H. Brown, Secretary National Oceanic and Atmospheric Administration D. James Baker, Under Secretary and Administrator National Weather Service Elbert W. Friday, Jr., Assistant Administrator for Weather Services



This publication has been reviewed and is approved for publication by Scientific Services Division, Western Region

26 miller

Kenneth B. Mielke, Chief Scientific Services Division Salt Lake City, Utah

TABLE OF CONTENTS

Ι.	INTRODUCTION
п.	THE CITY OF SANTA BARBARA 2
III .	HISTORY
IV.	SANTA BARBARA WEATHER RECORDS
V .	CHARACTERISTICS OF SANTA BARBARA CLIMATE
VI.	TEMPERATURE 9
VII.	PRECIPITATION
VIII.	FLOODING
IX.	WIND
Х.	DOWNSLOPE "SUNDOWNER" WINDS
XI.	MISCELLANY
XII.	REFERENCES AND ACKNOWLEDGMENTS
XIII.	APPENDIX

LISTS OF TABLES AND FIGURES

<u>LIST (</u>	DF TABLES	AGE
1.	DAILY TEMPERATURE MEANS BY MONTH	13
2.	AN EARLY SANTA BARBARA TEMPERATURE RECORD	14
3.	RAINFALL STATISTICS BY MONTH	18
4.	MONTHLY AND SEASONAL PRECIPITATION	
_	1867-1993	19
5.	ANNUAL RAINFALL: CUMULATIVE FREQUENCY	
_	DISTRIBUTION	29
6.	MAXIMUM RAINFALL FOR CONSECUTIVE MINUTES	30
7.	MAXIMUM RAINFALL BY MONTH	
8.	SANTA BARBARA WEATHER RECORDS, MEANS AND	
	EXTREMES	49
9.	SANTA BARBARA DAILY AND MONTHLY WEATHER	
	NORMALS	50
10.	SANTA BARBARA DAILY RECORD HIGH/LOW	
	TEMPERATURES	62
11.	AIRPORT DAILY RECORD HIGH/LOW	
	TEMPERATURES	74
12.	SUNRISE AND SUNSET TABLE	86

LIST OF FIGURES

4

-3

 $\left(\right)$

PAGE

1.	MAP OF SOUTHEASTERN SANTA BARBARA	
	COUNTY	
2.	MAP OF THE CITY OF SANTA BARBARA 4	
3.	SANTA BARBARA PRECIPITATION 1868-1990	
4a.	SMOOTHED RAINFALL AVERAGES 1867-1992	
	(ROSSMAN)	,
4b.	SMOOTHED RAINFALL AVERAGES 1868-1993	
	(GOODRIDGE)	
5.	PRECIPITATION DEPTH/DURATION/FREQUENCY	
	CURVE	,
6.	SANTA BARBARA STREAM GROUP FLOOD PLAIN	
	MAP	
7.	STAGE HYDROGRAPH: ARROYO BURRO CREEK	
8.	PAST FLOOD PICTURES 33	,
9.	WIND ROSES IN THE SANTA BARBARA AREA	,
10.	NEWSPAPER ACCOUNT OF 1917 DOWNSLOPE	
	WINDSTORM	ł
11.	GEORGE W. RUSSELL'S WEATHER LOG,	
	JUNE 1917)

12.	DOWNSLOPE WINDS WHIP SYCAMORE FIRE,	
	JULY 1977	41
13.	WINDS BLOW FIRE ACROSS SANTA BARBARA,	
	JUNE 1990	42

THE CLIMATE OF SANTA BARBARA, CALIFORNIA

Gary Ryan National Weather Service Santa Maria, California

ABSTRACT

The city of Santa Barbara, California has a moderate Mediterranean climate regime in general, with some surprising and unique features. The United States Weather Bureau last issued a climatology of Santa Barbara in 1962. This paper represents an effort, to update that study and assemble the latest relevant climatological data for the city of Santa Barbara. Within this paper, long-term temperature records are analyzed, precipitation trends are fully discussed, and research into the causes and effects of downslope wind events are outlined. Tables of daily temperature and precipitation values (averages and extremes) are also included.

I. INTRODUCTION

November 1962: In San Francisco, C. Robert Elford, the United States Weather Bureau's State Climatologist, completed a climate summary entitled "The Climate of Santa Barbara." Mr. Elford, with his assistant, Max R. McDonough, wrote a three page climate analysis for the city of Santa Barbara, along with a more extensive 30-page report on Santa Barbara County This 1962 Santa Barbara climate. climatology, until now the most recent U.S. Weather Service survey of the subject, was based on two sources: (1) the official downtown weather station record from 1931 to 1960 and (2) a relatively short record of aviation observations from Santa Barbara Airport in Goleta.

There have been sweeping changes in the monitoring of weather parameters in the Santa Barbara area since 1962. Today, several departments maintain weather observing equipment within and just outside of the city. These agencies include the National Weather Service, Federal Aviation Administration, U.S. Forest Service, State of California Department of Water Resources, County of Santa Barbara Flood Control and Water Conservation District, and County of Santa Barbara Air Pollution Control Corporations, educational District. institutions, and private citizens are also engaged in weather and climate monitoring, for a wide variety of reasons. As a result of this activity, the climatological information database for the Santa Barbara area has been greatly expanded.

This paper represents an effort to assemble the latest relevant climatological data for the city of Santa Barbara from as many sources as possible. The result is designed to be

an overview of the local climate. The reader is invited to contact the individual agencies noted in this report for more detailed information.

The city of Santa Barbara has a longterm official weather record dating from just after the Civil War. The general weather regime described by that record is a tranquil one, as befits a southern California coastal community. But in its extremes, Santa Barbara weather is unique, surprising and, at times, even awesome.

II. THE CITY OF SANTA BARBARA

On the southern California coast, 90 miles northwest of Los Angeles, beneath the ridges and canyons of the Santa Ynez Mountains, lies the city of Santa Barbara (Figs. 1 and 2). The city is located on an east-west oriented section of coastline facing the Santa Barbara Channel and the Channel Islands to the south. The 1993 population of the city was 87,500; the population of the city and its environs was 191,367. Goleta, Santa Barbara's sister city and the site of Santa Barbara Airport, is located eight miles west of Santa Barbara.


Santa Barbara and Goleta are built on a narrow, one-to-five mile wide coastal plain which rises precipitously to the Santa Ynez Mountain ridgeline. Elevations along the ridgeline, which extends to within five miles of the city of Santa Barbara, range from 2224 feet at San Marcos Pass, 10 miles northwest of the city, to 4299 feet at Santa Ynez Peak, 19 miles west northwest of downtown. La Cumbre Peak, at 3985 feet, is only six miles north of Santa Barbara.


Gibraltar Reservoir, in the upper Santa Ynez River Valley, lies eight miles north of Santa Barbara. To the east of the city are the communities of Montecito and Summerland. Carpinteria, at the Ventura County line, is 14 miles east. Located to the west are El Capitan Beach (20 miles), Gaviota Pass (34 miles) and Point Conception (46 miles).

III. HISTORY

Chumash Indians occupied the area near Santa Barbara as early as 10,000 years ago. This Native American culture was flourishing and relatively advanced when Juan Rodrigues Cabrillo claimed the California coast for Spain in 1542. Sebastian Vizcaino anchored near Santa Barbara on December 4, 1602, the feast day of Saint Barbara, and named the channel and shoreline in honor of the patron saint of mariners.

In April 1782, Father Junipero Serra conducted ceremonies establishing the Santa Barbara Presidio Real (Royal Fortress). Jose Francisco Ortega was appointed commandante of the first permanent outpost in the district. By the late 1780s, the population of the settlement stood at 4,200, of which 4,000 were Native Americans.





On December 21, 1812, the worst California earthquake in history destroyed most of the settlement at Santa Barbara. In 1820, the Santa Barbara Mission (which stands today) was completed to replace the facility destroyed in the earthquake. In 1822, the Mexican Revolution initiated a period of instability resulting in the establishment of military rule, which lasted until 1826. In the War with Mexico, U.S. forces seized Santa Barbara and in 1850 established civilian jurisdiction.

The late nineteenth century was an age of rapid change for the city. Stearns Wharf was completed in 1872, changing the character of the community into a port city. The Arlington Hotel opened in 1876, bringing tourists into the area. Electric street lights were installed and railroad service was established in 1877. The following year, State Street was paved. By the turn of the century the city population had reached 6,587.

The foundation for Gibraltar Dam was laid in 1913. A severe earthquake occurred on June 29, 1925, killing 11 and damaging many buildings. A new courthouse was finished in 1929 at a cost of \$1.5 million. By 1950, the population of Santa Barbara had reached 44,759. The dam at Cachuma Lake, holding the current water supply for the city, was completed in 1953 and took five years to spill. La Cumbre Plaza was finished in 1967. In 1969, a giant oil slick caused environmental damage to Santa Barbara Channel.

Santa Barbara today is, arguably, one of the most beautiful cities in the United States. The community has attempted to maintain a Spanish colonial image and merge that with modern American lifestyle.

IV. SANTA BARBARA WEATHER RECORDS

Outside of early settlers' general remarks and notes about weather and climate, the first weather record in the Santa Barbara area dates from June 1859. In the winter of 1861-62, journals recorded severe flooding in the district, followed by the worst drought in historic times.

Regularly scheduled weather observations began in Santa Barbara on July 1, 1867, with the establishment of a daily precipitation record. This was more than three years before the United States government first organized a national meteorological service.

A station history was compiled circa 1962 by Max R. McDonough, Assistant State Climatologist. An unedited version of the original handwritten document is presented herewith:

SANTA BARBARA STATION HISTORY

Precipitation record at Santa Barbara date [sic] back to July 1, 1867. A standard rain gage was located at the residence of Prof. Hugh D. Vail, 1325 Chapala St., 1/2 mile west of the present Post Office. The elevation was 92 feet.

On January 1, 1897, temperature record became of the observation part program with the addition of a Cotton Region Shelter and max & min thermometers. Professor Vail remained the observer until July 1, 1900. On this date the station was relocated 4/10 of a mile northwest to the home of James A. Dodge. The Dodge residence was at 116 W. Islay St., 9/10 mile WNW of the Post Office at an elevation of 128 ft.

Mr. continued Dodge as observer until November 1, 1906 when the station was again moved 4/10 of a mile west to the residence of Mr. W. Russell. 2135George Castillo St: 1.3 miles WNW of Post Office, elevation 121 ft. The climate station remained this location at through October 31, 1931. Mr. Russell was succeeded by his wife Mrs. Ella M. Russell on May 1, 1925, who carried on the observational work until the appointment of Mrs. Edvthe On November 1. Hensel. 1931, the station was relocated approximately sixtv vards the southwest to Hensel Residence, at 427 Los Olivos St., elevation now 118 ft.

In May 1940, a [recording rain gage] was added, to remain part of the observational program to the present date. Mrs. Hensel retired in February 1948. On February 5, 1948, Mr. August F. Grimm was appointed, and the station was moved to the Rose Garden Nursery, 606 W. Pueblo St., two tenths (2/10) mile west, 1-1/2 mile WNW of Post Office, elevation 120 ft. Mr. Grimm's association was terminated in December 1954 when the nursery property was purchased by the State Division of Highways. On December 22, the move was to the Santa Barbara Fire Dept. residential substation (Firehouse no. 3) at 415 E. Sola St. New location one and four tenths (1.4) miles ENE of the Post Office, elevation 100 feet. Observation duties were performed by onduty personnel at the fire station. Thus, continuing a long and homogeneous weather record from the city of Santa Barbara.

As with those who preceded them, the continuous published record they have furnished and the original forms they have prepared stand as a monument to their contribution to the climatological and hydrologic [sic] data of this community and state. This is that effort that makes this publication possible.

The McDonough record is fine in its broad outline through 1954. but requires some correction and augmentation. It is not certain who started the original precipitation record at Santa Barbara in 1867. Historical documents suggest that it was the work of Dr. J. B. Shaw. Dr. Shaw utilized a Smithsonian model rain gage. His

logbooks contain original precipitation records, and it is likely that the records he maintained were copied by Professor Vail and others in the 1880's.

Temperature data were collected by Dr. L. N. Dimmick beginning in 1875 and continuing through 1879. Dr. Dimmick establish temperature wished to averages for Santa Barbara. In the fashion of the day, he diligently read the thermometer three times a day (at 7 a.m., 2 p.m. and 9 p.m.) and averaged the three readings. This average was considered at the time to be quite scientific; the average annual temperature computed from this method was roughly two degrees F. higher than the modern calculation.

Dr. Shaw continued this temperature analysis from 1879 to 1883. In January 1883, William Field obtained "maximum and minimum self-registering thermometers" and recorded official high and low temperatures at the Arlington Hotel until 1885. At that time, Professor Vail took responsibility for the weather observation program.

The Vail record continued through July 31, 1900. James A. Dodge's name appears on Santa Barbara records beginning in August 1900. In January 1901, the meteorological forms used by the Department of Agriculture (Weather Bureau) underwent significant revision.

The weather records of George W. Russell and his wife Ella spanned 25 years from 1906 to 1931. Their records were complete and were accompanied by substantial annotation --- frequently regarding agriculture. One of George Russell's records appears as Table 2.

The Weather Bureau operated the official substation at the fire department at 415 E. Sola Street beginning on December 22, 1954. It was located 1.5 miles inland from the harbor and 0.6 miles north-northwest of the post office, not 1.4 miles eastnortheast of the post office, as in the McDonough report. The site was judged excellent for precipitation only "fair" measurement but for temperature shelter exposure.

In 1964, the Fire Department decided to blacktop the backyard of Fire Station No. 3. The Weather Bureau judged change in this exposure to be unsatisfactory, and moved the substation once again. The substation was relocated to its present site at the Santa Barbara (El Estero) Water Treatment Plant at 402 E. Mason Street on March 25, 1964. This location is 0.9 miles east-southeast of the post office, close to the 101 Freeway and the ocean, at an elevation of five feet MSL.

Upon moving the observing site to El Estero in 1964, a Weather Bureau report cautioned "...it is thought that maximum temperatures may be somewhat lower at the new location due to its proximity to the Pacific Ocean." Santa Barbara maximum (and minimum) temperatures have indeed exhibited а pronounced oceaninfluenced moderation since 1964, which will be addressed in this report within the section concerning temperature analysis.

The El Estero station, number 04-7902-6, is located at 34° 25' N and 119° 41' W. The observation site is equipped with a Fischer & Porter rain gage, an eight-inch standard rain gage, and a cotton region shelter enclosing a standard maximum and minimum thermometer. Observations are taken daily at midnight.

On March 17, 1938, the Federal Aviation Administration established an airport weather station (station number 04-7905, WBAN number 23190) in an operations building at Santa Barbara Airport in Goleta. The same site is still in use today. The station is located one mile from the ocean and eight miles west of the Santa Barbara city center, at an elevation of nine feet MSL. Map coordinates are 34 degrees 26 minutes north latitude and 119 degrees 50 minutes west longitude.

Daily temperature records from Santa Barbara Airport began January 1, 1941. On September 1, 1942, a standard eight-inch rain gage was added to station equipment. Maximum and minimum thermometers were installed on May 9, 1947. An HO83 electronic hygrothermometer was commissioned in March 1986, resulting in some (undetermined) dampening of high and low temperature data.

Flight Service Station personnel provided hourly and special aviation weather observations through the years at Santa Barbara Airport. Then, beginning in March 1993, the station was staffed with private contractor weather observers, operating on a 24hour basis. Observations are currently taken on standard federal forms, which are checked for accuracy by officials at the National Climatic Data Center in Asheville, North Carolina.

V. CHARACTERISTICS OF SANTA BARBARA CLIMATE

Santa Barbara is noted for its moderate climate. Under the modified Koppen climate classification system, the Santa Barbara climate is categorized as "Csb," or Mediterranean, with cool, dry summers (Trewartha and Horn, 1980). There are pronounced seasonal changes in rainfall, but relatively modest seasonal transitions in temperature.

The mean annual temperature at Santa Barbara is 60°F. The average daily high temperature is 71°F and the average low is 49°F.

Rainfall is concentrated during the winter season, primarily from November through April. Significant precipitation events from May through October are unusual. The average annual precipitation is 17.91 inches.

The eastern Pacific high pressure area, a semi-permanent feature of the general atmospheric circulation pattern in the Northern Hemisphere, controls weather along the California coast for much of the year. Prevailing wind along the state's coastline is generally from the northwest or west, although surface wind patterns show some complexity due to terrain. In the Santa Barbara area, the northwest wind flow is normally blocked by the Santa Ynez Mountains, creating daytime sea-breeze conditions with winds blowing from the south or west quadrant. Wind speeds average four to five mph annually, with peak winds on most days of about 10 to 15 mph.

A pronounced marine layer exists along the coast in the summer months through mid-October. In this season, low clouds and fog frequently occur from late at night to about midday, then "burn off" to a clear but somewhat hazy sky. Dense fog occurs from time to time in the late summer and early fall, but is not as frequent at Santa Barbara as it is along the coastline north of Point Conception.

Severe weather is uncommon. Winter storms occasionally bring gusty southeast winds to about 40 mph to the district with locally heavy rain. These storms can cause flooding within the city of Santa Barbara. Generally, however, winter storms are migratory; so wet and dry periods tend to alternate, albeit with considerable irregularity in timing and duration.

Northwesterly downslope winds, known locally as "sundowners," occur within the city on an average of two or three times each year. Sundowner winds, usually warm and dry, can locally reach speeds in excess of 40 miles per hour and cause a variety of public safety hazards.

VI. TEMPERATURE

Historical temperature data at Santa Barbara present problems for climatologists. The official Santa

Barbara observation site has been in at least seven different locations since it was established in 1867. Until 1964, the point of observation was near the city center at elevations between 82 feet and 130 feet MSL. Temperature records from those locations may be considered homogeneous and reliable. In March 1964, the official observation site was moved to the El Estero Water Treatment Plant near the ocean and at elevation of five MSL. an feet Temperature averages 1964since manifest a predictable maritimeinfluenced moderation. The average daily maximum temperature decreased 1.1°F in the post-1964 period, while the average daily minimum temperature increased 2.2°F.

High and low daily temperature records at Santa Barbara were set at the rate of 1.63 per month during the 1940s, but that frequency dropped to 0.26 per month during the 1980s. Airport records indicate 1.80 daily record temperature extremes registered per month during the 1940s and 1.29 records per month in the 1980s. Such changes in comparative data suggest an inhomogeneity in Santa Barbara city records. El Estero temperature records should be employed with caution and probably should not be used in computing long-term temperature averages for the city.

For the purposes of statistical analysis, Santa Barbara city temperature data are utilized only through 1960, and data from El Estero are not considered. For all-time daily record computations, the entire existing continuous record from January 1883 to November 1993

addition was used, in to some fragmentary records beginning in 1875. The complete city temperature record was utilized, including El Estero, because there was no clear alternative. It has been noted that it has been demonstrably more difficult to set daily temperature records at Santa Barbara since 1964. Therefore, for the sake of comparison, Santa Barbara Airport alltime daily temperature records are included within this report.

Because of inhomogeneities within the historic temperature record at Santa Barbara, the station data support no conclusions in regard to long-term temperature trends. However, regional trends have been described by Fritts (1991) and others engaged in dendroclimatologic research, which indicate that warming has been occurring across the Santa Barbara area during the twentieth century. Temperatures during this period have averaged slightly warmer than the 30-year normal ending in 1900.

The average annual temperature at the city of Santa Barbara, based on data from 1901-1960, is 60.1°F. This compares with 66°F at Los Angeles and 57°F at the San Francisco Airport. Santa Barbara's average daily high temperature is 71.4°F and the average daily low is 48.8°F. Average daily maximum temperatures peak in August and September at 78°F, dropping to 64°F in late December and early January. Average daily minimum temperatures peak at 57°F in July and August and drop to 40°F for a short time around New Year's Day. At the Santa Barbara Airport, daily maximum temperatures average 2.1°F cooler and daily minimum temperatures average 0.2°F warmer than Santa Barbara city readings.

transitions Temperature between seasons is gradual and smooth, as reflected in daily normal temperature tables. The most significant warming occurs during a sixty day period in May and June, when average daily temperatures rise by 6°F. Cooling in the fall occurs more sharply; average temperatures drop by 6°F in a 35-day period between late October and late November.

Although moderate temperatures are the rule at Santa Barbara due to the marine influence. temperature extremes can and do occur. The official high temperature record of 115°F occurred during a desiccating downslope windstorm on June 17, 1917. The coldest temperature reported in the city, 20°F, occurred twice during the record cold month of January 1949. The highest temperature recorded unofficially at Santa Barbara was an incredible 136°F on Friday, June 17, 1859. A description of that event is included in the chapter on downslope "sundowner" winds.

Hot Temperatures

Santa Barbara has a southerly exposure at the base of the Santa Ynez Range, and is subject to warming downslope winds which cross the crest of the Santa Ynez Range from the north. Therefore, the city can become quite warm at any time of year. Maximum temperatures have been recorded in excess of 90°F in every month of the year. The temperature reaches 90°F or higher on an average of 4.5 times per year. Temperatures of 100°F or more have occurred historically in the months from April through October. Maxima of 100°F or higher occur in about one year in three. The average annual maximum temperature recorded at Santa Barbara is 97°F. The month in which the annual maximum temperature is most likely to occur is September.

Very high temperatures are sometimes caused by and accompanied by gusty downslope winds, which are known locally as "sundowners." These winds can reach gale force within the city limits, but such conditions are rare. Hot temperatures are not always associated with downslope windstorms, however. Sometimes the heat occurs with light winds, when a strong high pressure ridge aloft phases with surface high pressure to the east of the district, creating a light offshore wind pattern. In this situation temperatures along the Santa Barbara littoral can reach to from 100-110°F.

Cold Temperatures

Cold weather outbreaks have historically dropped temperatures into the 30s°F at Santa Barbara in all months except June, July, and August. These incidents of cold temperatures are usually caused when strong high pressure areas push northward from the eastern Pacific into the Gulf of Alaska, which results in cold air from the Canadian Arctic moving southward into the western United States. When offshore pressure gradients are light and low-level winds are quiescent, temperatures can cool rapidly at Santa Barbara in the relatively dry air and clear skies.

The winter of 1948-49, which featured below normal temperatures throughout the western United States, marked the century's benchmark cold outbreak at Santa Barbara. From September low through February, 50 daily temperature records were established; these records have endured for almost a half century. Sixteen daily low temperature records were set in January 1949 alone. The average daily low temperature for that January was 30.5°F, 10.9°F below normal, and the only time on record when a monthly minimum temperature averaged below freezing. The all-time record low of 20°F was set on January 4 and equalled on January 10. ų. Lais

Freezing temperatures are uncommon at Santa Barbara, occurring on an average of four times per season, twice in December and twice in January. In 38 percent of the winter seasons, the lowest observed temperature is above freezing --- 33°F or higher. In 53 percent of winter seasons, the lowest temperature ranges from 28-32°F, and in nine percent of winter seasons, the lowest temperatures are colder than 28°F. Freezing temperatures have occurred as early as November 12 and as late as April 28, based on the 110year continuous temperature record.

Elford (1962), in a study prepared with 1931-1960 data, indicated an average first freeze date (32°F) at Santa Barbara of December 31 and a last freeze date of January 23, with a resultant growing season of 342 days. This study is based on data reflecting a high degree of homogeneity, and is valid for the area near the city center. An analysis of Santa Barbara Airport temperatures (1951-1974) yields an average first freeze date of December 20, a last freeze date of January 20, and a growing season of 334 days. The heating degree day seasonal average (base $65^{\circ}F$) for the city for the 1931-1960 period stands at 2086. Even during California's coldest overall synoptic patterns, daily high temperatures at Santa Barbara usually climb into the 50s and 60s°F. The coolest daytime high temperature ever recorded in the city occurred on January 6, 1913, when the mercury managed to rise to only 44°F. TABLE 1.

DAILY TEMPERATURE MEANS BY MONTH AT SANTA BARBARA, CA

MONTH	MAXIMUM	MINIMUM	MEAN	DEGREE <u>HEAT</u>	DAYS ING/COOLING
JANUARY	64.7	41.4	53.1	384	0
FEBRUARY	65.5	43.2	54.3	308	0
MARCH	67.3	44.8	56.0	282	0
APRIL	69.1	47.6	58.3	198	0
MAY	70.5	49.9	60.2	140	0
JUNE	73.1	52.8	63.0	75	24
JULY	76.9	56.4	66.7	28	55
AUGUST	77.5	56.7	67.1	22	91
SEPTEMBER	77.5	54.8	66.2	36	95
OCTOBER	75.3	50.7	63.0	84	24
NOVEMBER	72.3	45.1	58.7	210	0
DECEMBER	67.0	42.3	54.7	319	0
ANNUAL	71.4	48.8	60.1	2086	289
Sources: 1	lemperature Heating deg	data, Sant ree days, Sa	a Barbara c anta Barbara anta Barbara	ity 1901-1 city 1931	.960. 1960.

13

Form No. 1078 A-Met'l. U. S. DEPARTMENT OF AGRICULTURE, WEATHER BUREAU.

Stati	Con.C									_				aj	1
Lati								···· ون						, 130	feei
				De	ata, ζ	M	an (Ten	her	atur	l				-
Year.	Jan.	Feb.	March.	April,	May.	June.	July.	August.	Sept.	Oct.	Nov.	Dec.	Annual.	Mar	ority. Min
1885	53.2	56.7	59.1	60.7	60.0	62.4	61.1	68.0	66.9	63.0	58.9	57.2	60.6		
6	55. o	59.6	53.1	55.7	60.5	62.0	66.3	68.2	63.8	<u> (62.3)</u>	<u> (57.4</u>)	55.8	(6 <u>0.0)</u>		•
7	547	50.4	57.0	58.4	60.1	63.7	64.6	64.8	66.0	65.0	58.9	(54.8)	(v9.9)		· · · · · · · · · · · · · · · · · · ·
8	49.0	53.8	53.0	59.9	62.8	64.4	67.0	66.3	67.9	63.5	59.1	55.8	60.2		
9	53.0	55.4	59.0	59.9	60.0	62.5	62.2	67.3	68.8	63.9	59.6	54.2	60.5		
1890	48.4	52.6	55.6	56.6	60.0	62.4	67.3	67.9	66.5	64.0	63.3	58.4	60.2		
	<i>54, 4</i>	52.6	56.6	56.3	59.0	62.4	67.2	69.1	69,3	63,3	58.8	51.9	60.1	96.	33
2	54.5	53.7	55.3	57.8	60.3	61.0	63,5	66.3	64.2	62.0	60.3	54.4	59.4	98	38
3	55.7	J3.4	53.1	57.0	59.2	62.2	64.6	66.1	62.7	62.0	57.0	18.0	59.2	88	38
4	49.7	50.9	53.4	57.1	58.0	61.4	62.8	65.9	65.9	62.6	57.4	54,1	18.3	94	33
5	51.7	55.8	53.6	57.5	60.5	62.5	65.0	65.0	65.6	63,4	19.9	55.1	59.6	91	38
6	56.4	61.2	57.6	55.3	61.4	64.3	66.9	67.5	64.7	63.3	58.7	58.1	61.3	98	36
	54.6	52.8	52.7	59.6	59.5	63.0	65.3	71.9	67.0	61.0	58.6	544	59.6	93	32
8	50.7	55.9	54.4	60.2	18.3	64.0	65.5	68.6	66.0	61.8	19.1	55.6	60.0	95	34
9	54.8	54.2	56.6	56.8	56.4	61.4	64.6	6516	66.0	62.4	60.4	57.1	19.7	93	29
1900	57.5	58.0	58.3	56.0	60.8	63.8	65.9	65.0	65.2	62.8	64.3	59.8	61.4	96.	38
01	546	56.4	59.0	55.0	57.8	61.6	65.3	67.2	61.6	63.6	59.2	v7.4	59.9	96	34
02	54.8	54.8	54.0	57.2	59.0	62.8	65.3	65.4	66.0	61.8	57.6	54.2	59.4	90	35
03	56.4	51.1	54.6	57.0	58.4	63.4	65.2	66.1	66.1	64.8	62.1	57.6	60.2	95	32
04	SS. 0	54.6	56.0	58.4	59.1	64.4	64.8	69.0	69.2	65.0	63.1	58.4	61.4	95	33
05	56.6	56.8	59.1	58.2	58.5	61.0	65.6	66.2	66.5	644	57.8	55.8	60.7	96	35
06	548	58.4	56.4	57.8	58.2	65.1	67.9	66.4	67.3	65.6	56.4	<i>55.</i>]	60.8	98	34
07	51.0	59.0	54.0	58.2	60.2	62.8	67.8	65.5	64.1	63.1	58.6	56.0	60.0	108	34
08	54.6	52.2	55.9	59.0	58.2	60.2	65.8	66.5	64.9	59.6	56.2	51.5	18.7	91	32
														100	
1	•												1 I		······
Means	53.8	SUG	11.1-	5.2			<u>l. 1. 1.</u>		′ <u>,</u>	1	100	<u> </u>	61.0		1

REMARKS.

TABLE 2.

AN EARLY SANTA BARBARA TEMPERATURE RECORD

Courtesy: Jim Goodridge

.

VII. PRECIPITATION

City of Santa Barbara precipitation data published by the National Weather Service and by Santa Barbara County Flood Control and Water Conservation District were checked against the original historical record. The data were remarkably consistent from 1867 through 1977. Then, beginning in 1978, significant divergence occurred in the reprinting of the El Estero data.

Precipitation data published in this report represent, as far as can be determined, official figures noted in original source material; the entire corrected monthly city precipitation record is published herein for the first time (Table 4). Statistical analysis of the Santa Barbara precipitation record is based on data from July 1867 (the Shaw record) through June 1993, inclusive, except as noted.

Since rainfall in the Santa Barbara area is generally a winter season phenomenon occurring from November through April, statistics are frequently presented on a "water year" basis, as opposed to a calendar year format. The water year displays a more coherent picture of rainfall data. The National Weather Service computes the water year from July 1 through June 30. Santa Barbara Flood Control and Water Conservation District reckons water year boundaries from September 1 through August 31. This bookkeeping difference is unimportant because noteworthy rainfall rarely falls at Santa Barbara during July and August. In fact, some of the earliest weather observers in Santa Barbara did not include any July or August rainfall data at all as part of the annual totals; a tenmonth water year was started on September 1 and ended on June 30! This may account for what appears to be an unusual string of no-precipitation totals during the months of July and August from 1867 to 1894.

The annual average precipitation at Santa Barbara, based on a continuous and homogeneous 126-year record from 1867-1993, is 17.91 inches. The airport average (1961-1990) is 16.25 inches. The monthly average at Santa Barbara for January is 3.92 inches. By midsummer, rainfall is scanty. The average July precipitation is only 0.03 inches, and what moisture does fall is most often associated with a deep layer of marine air, up to about 4,000 feet thick, which causes widespread drizzle and local areas of fog.

Most of the season's rains occur during the months from November through April. Only eight percent of total seasonal rainfall normally occurs from May through October and less than one percent of the total occurs during the three-month period from June through August.

Heaviest rains are associated with storms approaching California from the west or southwest which frequently tap a moisture supply from the subtropics. These storms often produce south or southeast low-level winds, which provide pronounced orographic lifting (and thus, rain enhancement) in the Santa Barbara area. Extremely heavy rains, up to 15 inches in one day, have been recorded near the crest of the Santa Ynez Range just a few miles north of the city. Within the city, the heaviest daily rainfall on record is 6.95 inches, which occurred on January 25, 1914. Probable maximum precipitation at Santa Barbara is plotted in graph form (Fig. 5); a rainfall depth/duration/frequency analysis is found in Tables 6 and 7.

Thunderstorms are unusual in Santa Barbara. Sometimes accompanied by small hail and gusty winds, they occur on an average of 2.4 days per year. Thunderstorms can develop during the winter or early spring, usually spawned by or accompanying rapidly moving cold fronts that swing southeastward across California from the Gulf of Alaska. Summer thunderstorms have ิล different origin; they tend to drift into the Santa Barbara area from the south. This happens when the so-called southwest monsoon brings moist subtropical air into southern California.

Snowfall is extremely rare. The Weather Bureau historically included occurrences of hail and ice pellets with snowfall data, so that statistics relating "snow" to may not describe Using the inclusive "snowflakes." Weather Bureau definition of snow, some form of frozen precipitation (presumably mostly ice pellets) has fallen in Santa Barbara during the months from December through April. Snowfall depths have never reached more than trace levels within the city -that is, any snow or ice that has fallen has never accumulated to a depth of a A documented half-inch or more. snowfall occurred at Santa Barbara in 1882. An account of the snowfall was

given by Dr. J. B. Shaw in his weather logbook as follows:

January 12, 1882. This morning all the hills, down to the houses in town [are] being covered with snow. The streets in Los Angeles, Lompoc, and Los Alamos [are] being covered with snow: such an event was never known before.

And from <u>The Daily Press</u> of the same date:

Flakes of snow fell in the streets of Santa Barbara today. We believe this to be totally unprecedented. The boys of Santa Barbara were wild with excitement this morning, and many of them made a beeline for the top of the foothill, in search of a genuine snowball. Some of them, California-born, have never before had the opportunity to roll this delightfully chilly article into a missile and grasped the chance and the snow with avidity.

Trends and Fluctuations in the Rainfall Record

Attempts have been made to find evidence of periodicity in historic rainfall patterns in the Santa Barbara area. A computer analysis by Rossman (Fig. 4a) graphed 10-year rolling precipitation means at 5-year intervals for the entire historic record beginning in 1867. Solid vertical lines represent 30-year intervals beginning in 1870. Dashed lines mark low points in the rainfall record. This smoothing revealed an apparent dry period of about 30 years. This periodicity had been noted by several others working with this and other rainfall data from the area.

This pattern was still extant in another graphic study, completed in December 1993 by Jim Goodridge, using the same base data (Fig. 4b). Goodridge combined continuous nine-year rolling means with an individual data plot. He considered \mathbf{the} resultant pattern "remarkable," but concluded that the dry period is <u>not</u> cyclic because of the random wavelength displayed in the raw data plot. Moreover, statistical analysis of such a relatively short (by planetary time standards) rainfall record cannot be taken as conclusive evidence of cycling. In a well-done tree ring study of southern California climate, Michaelsen et al., (1990) reconstructed annual streamflow (and by extension, patterns precipitation patterns) from the year 1460 to the present. He found that the patterns are approximately "random and stationary" and "do not show longterm trends." The flow of the nearby Santa Ynez River has averaged near the 500-year normal since 1900. Michaelsen noted very wet and very dry climatic cycles before 1800, which have not been matched in the record since 1800.

This is not to say that dramatic changes in precipitation frequency have not occurred in historic times. The floods of 1861-62 were followed immediately by the worst drought the Santa Barbara area has known. O. H. O'Neill (1939) reported that after a "50inch" rainfall during the winter of 1861-62...

The Dry Spell began in the fall of 1862 and lasted to the winter of 1864-65. The rainfall for the season of 1862-63 did not exceed four inches. In the fall of 1863 a few showers fell, but not enough to start the grass.

The severe drought that hit the county in the early 1860s wiped out the local cattle industry and closed the Matanza (slaughterhouse), which had been established in the city. Lacking the infrastructure and emergency water supply of the modern era, the early cattle owners had no choice but to watch their stock perish. One account mentioned that "there was no available grass within 400 miles of Santa Barbara." Reliable reports state that of 300,000 head of cattle in Santa Barbara County in 1862, less than 500 were alive after the winter of 1864.

	STATISTICS BY MONTH	FOR TH	HE CITY OF	SANTA	BARBARA, CA
month	AVERAGE	MAXIM	 <u>M (YEAR)</u>	 Max D.	AILY (DATE)
JANUARY	3.92	17.22	(1916)	6.95	(1914/25)
FEBRUARY MARCH	3.77 2.95	17.33 12.38	•	4.25 3.80	(1914/18) (1952/15)
APRIL	1.22	6.55	(1965)	3.14	(1926/ 4)
MAY	0.36	2.96	(1977)	1.60	(1949/18)
JUNE	0.09	1.62	(1884)	1.13	(1963/11)
JULY	0.03	0.81	(1950)	0.63	(1950/ 9)
AUGUST	0.04	1.48	(1983)	0.78	(1983/19)
SEPTEMBER	0.33	7.15	(1904)	4.50	(1904/25)
OCTOBER	0.60	8.65	(1889)	2.45	(1907/22)
NOVEMBER	1.57	9.84	(1885)	3.90	(1902/10)
DECEMBER	3.03	12.67	(1867)	5.32	(1977/28)
ANNUAL	17.91	45.21	(1940-41)	6.95	(1/25/1914)
Sources:	Santa Barbara city				
Data:	from 1 July 1867 th Rainfall in inches reading at time of through the histor:	. Dail daily ic peri	ly records observatio lod.	reflec on, whi	t gage ch varies
===========				======	

à.

1

CITY OF SANTA BARBARA, CALIFORNIA MONTHLY AND SEASONAL PRECIPITATION

YEAR	JUL	AUG	SEP	OCT	NOV	DEC	JAN	FEB	MAR	APR	MAY	JUN	SEASON
1867-68	0.00	0.00	0.00	0.00	2.31	12.67	3.97	2.00	1.08	2.44	0.72	0.00	25.19
1868-69	0.00	0.00	0.00	0.00	1.25	4.26	3.26	2.12	4.22	0.46	0.20	0.00	15.77
1869-70	0.00	0.00	0.00	0.30	0.65	0.57	0.25	5.87	0.83	0.99	0.74	0.07	10.27
1870-71	0.00	0.00	0.00	1.04	0.27	1.41	0.86	2.92	0.02	2.02	0.37	0.00	8.91
1871-72	0.00	0.00	0.00	0.09	1.83	6.56	2.53	1.81	0.18	1.80	0.00	0.14	14.94
1872-73	0.00	0.00	0.05	0.00	0.00	4.34	0.58	5.48	0.05	0.00	0.00	0.00	10.50
1873-74	0.00	0.02	0.00	0.00	0.27	5.26	4.54	3.17	0.78	0.28	0.14	0.00	14.46
1874-75	0.00	0.00	0.00	1.91	1.30	0.00	14.84	0.18	0.38	0.10	0.00	0.00	18.71
1875-76	0.00	0.00	0.00	0.00	6.53	0.31	7.56	5.67	2.73	0.27	0.00	0.00	23.07
1876-77	0.00	0.00	0.00	0.32	0.00	0.00	2.72	0.00	0.82	0.13	0.45	0.00	4.49
1877-78	0.00	0.00	0.00	0.00	1.32	3.12	7.17	11.73	2.47	3.34	0.29	0.07	29.51
1878-79	0.00	0.00	0.00	0.32	0.00	5.16	5.24	0.71	0.34	1.60	0.21	0.00	13.58
1879-80	0.00	0.00	0.00	0.41	1.62	4.57	1.30	10.86	1.15	5.73	0.00	0.00	25.64
1880-81	0.00	0.00	0.00	0.25	0.28	9.73	2.83	0.30	1.25	0.59	0.00	0.00	15.23
1881-82	0.00	0.00	0.44	1.47	0.33	0.95	1.13	2.38	5.74	1.63	0.00	0.20	14.27
1882-83	0.00	0.00	0.00	0.37	0.77	0.10	2.18	2.92	3.64	0.29	2.79	0.35	13.41
1883-84	0.00	0.00	0.00	1.32	0.00	2.76	6.33	9.68	9.77	2.60	0.39	1.62	34.47
1884-85	0.00	0.00	0.00	1.02	0.79	6.62	1.23	0.07	0.35	3.00	0.00	0.00	13.08
1885-86	0.00	0.00	0.00	0.19	9.84	2.47	5.12	1.19	2.03	3.40	0.00	0.00	24.24
1886-87	0.00	0.00	0.00	0.39	0.87	0.86	0.31	8.64	0.13	1.43	0.33	0.03	12.99
1887-88	0.00	0.00	0.38	0.31	1.10	4.43	10.15	1.30	3.86	0.16	0.02	0.00	21.71
1888-89	0.00	0.00	0.03	0.07	5.62	5.59	0.29	1.29	7.31	0.49	0.76	0.13	21.58
1889-90	0.00	0.00	0.00	8.65	3.21	10.64	5.32	2.96	1.10	0.31	0.18	0.00	32.37
1890-91	0.00	0.00	1.50	0.05	0.48	3.53	0.45	7.92	1.56	1.57	0.30	0.00	17.36
1891-92	0.00	0.00	0.15	0.00	0.00	2.43	1.10	2.55	2.95	0.46	1.12	0.00	10.76
1892-93	0.00	0.00	0.00	0.26	4.27	6.66	4.41	3.10	7.80	0.38	0.09	0.00	26.97
1893-94	0.00	0.00	0.00	0.82	0.07	2.94	0.99	0.76	0.29	0.24	0.91	0.00	7.02
1894-95	0.12	0.00	1.36	0.68	0.07	4.67	6.25	0.67	1.99	0.46	0.02	0.05	16.34
1895-96	0.00	0.00	0.00	0.55	0.77	0.93	6.84	0.00	2.37	1.78	0.08	0.05	13.37
1896-97	0.40	0.00	0.00	0.92	3.51	2.92	4.35	3.65	2.73	0.02	0.00	0.00	18.50
1897-98	0.00	0.00	0.00	1.44	0.00	0.00	0.63	1.39	0.28	0.00	1.25	0.00	4.99
1898-99	0.00	0.00	3.17	0.14	0.00	0.36	4.48	0.02	2.78	0.64	0.00	0.78	12.37
1899-00	0.00	0.00	0.00	2.06	1.97	2.35	2.32	0.05	1.58	0.42	1.90	0.01	12.66

ŝ.

19

.

TABLE 4.

CITY OF SANTA BARBARA, CALIFORNIA MONTHLY AND SEASONAL PRECIPITATION

YEAR	JUL	AUG	SEP	OCT	NOV	DEC	JAN	FEB	MAR	APR	MAY	JUN	SEASON
1900-01	0.02	0.00	0.04	0.15	3.99	0.02	4.86	3.65	0.16	2.07	0.34	0.10	15.40
1901-02	0.06	0.09	0.36	2.42	1.16	0.00	1.36	4.40	2.89	1.40	0.07	0.00	14.21
1902-03	0.00	0.00	0.00	1.48	4.01	2.24	2.06	1.63	6.12	2.91	0.27	0.02	20.74
1903-04	0.00	0.00	0.00	0.00	0.05	0.00	0.46	4.69	4.40	1.89	0.09	0.00	11.58
1904-05	0.00	0.10	7.15	0.51	0.00	1.53	3.73	8.22	6.40	0.51	1.44	0.05	29.64
1905-06	0.18	0.00	0.03	0.16	1.14	0.07	4.26	3.67	9.96	0.83	2.40	0.00	22.70
1906-07	0.00	0.04	0.00	0.00	0.35	6.46	12.46	2.34	5.64	0.27	0.00	0.16	27.72
1907-08	0.00	0.03	0.00	6.23	0.00	1.80	4.29	5.96	0.21	0.49	0.20	0.00	19.21
1908-09	0.00	0.00	1.16	0.20	1.84	2.48	15.67	7.92	6.91	0.00	0.03	0.08	36.29
1909–10	0.00	0.01	0.17	0.57	2.34	9.53	2.91	0.08	3.62	0.39	0.00	0.00	19.62
1910-11	0.02	0.00	2.56	0.29	0.33	0.75	14.21	4.92	7.76	1.02	0.03	0.05	31.94
1911-12	0.00	0.00	0.12	0.28	0.02	2.33	0.42	0.00	9.48	2.12	1.58	0.00	16.35
1912-13	0.00	0.00	0.00	0.28	0.21	0.00	3.14	6.58	0.64	1.04	0.19	0.50	12.58
1913-14	0.09	0.07	0.17	0.00	3.43	2.71	15.91	7.30	0.95	0.70	0.03	0.16	31.52
1914-15	0.05	0.00	0.00	0.12	0.04	4.38	4.94	8.03	1.15	0.97	1.57	0.00	21.25
1915-16	0.00	0.00	0.05	0.00	0.65	4.06	17.22	1.89	1.71	0.30	0.00	0.00	25.88
1916-17	0.00	0.11	1.90	2.82	0.10	6.12	3.25	7.61	0.28	0.28	0.09	0.00	22.56
1917-18	0.00	0.03	0.05	0.00	0.17	0.03	0.51	10.47	10.37	0.05	0.00	0.00	21.68
1918-19	0.25	0.58	2.13	0.02	3.64	0.83	1.20	1.95	2.62	0.17	1.07	0.00	14.46
1919-20	0.00	0.00	0.84	0.27	0.23	2.11	0.33	5.81	4.20	0.81	0.00	0.08	14.68
1920-21	0.00	0.00	0.00	0.40	0.56	1.51	5.32	1.58	1.77	0.38	2.69	0.10	14.31
1921-22	0.00	0.00	0.24	0.32	0.02	7.25	4.64	3.48	2.73	0.09	0.45	0.00	19.22
1922-23	0.00	0.00	0.00	0.37	1.98	8.70	1.96	0.91	0.00	3.29	0.00	0.03	17.24
1923-24	0.00	0.03	0.13	0.25	0.00	0.08	1.63	0.06	3.56	0.62	0.00	0.00	6.36
1924-25	0.00	0.00	0.00	0.85	1.20	1.20	0.60	1.45	2.79	1.89	2.23	0.05	12.26
1925-26	0.05	0.00	0.00	0.71	0.80	2.57	2.08	4.28	0.25	6.13	0.00	0.00	16.87
1926-27	0.00	0.00	0.00	0.36	6.84	0.62	1.94	9.86	2.28	0.78	0.00	0.00	22.68
1927-28	0.00	0.05	0.11	3.48	1.49	3.28	0.00	1.95	2.46	0.17	0.50	0.05	13.54
1928-29	0.00	0.00	0.00	0.10	2.46	4.41	1.53	2.28	2.39	1.17	0.00	0.20	14.54
1929-30	0.00	0.00	0.05	0.00	0.00	0.00	5.82	1.21	4.93	0.95	0.63	0.12	13.71

TABLE 4.

1

CITY OF SANTA BARBARA, CALIFORNIA MONTHLY AND SEASONAL PRECIPITATION

YEAR	JUL	AUG	SEP	OCT	NOV	DEC	JAN	FEB	MAR	APR	MAY	JUN	SEASON
1930-31	0.00	0.00	0.01	0.04	2.64	0.00	4.25	4.07	0.00	1.43	2.11	0.00	14.55
1931-32	0.00	0.23	0.01	0.00	2.81	9.84	2.40	6.17	0.23	0.35	0.09	0.00	22.13
1932-33	0.00	0.00	0.11	0.10	0.00	0.67	6.42	0.00	0.30	0.20	0.11	0.75	8.66
1933-34	0.00	0.00	0.00	0.88	0.11	6.28	1.49	3.67	0.00	0.00	0.00	1.00	13.43
1934-35	0.00	0.00	0.04	1.89	3.48	3.63	4.10	1.58	3.16	3.32	0.00	0.00	21.20
1935-36	0.07	0.27	0.02	0.78	0.71	1.46	0.73	10.49	1.97	0.65	0.01	0.01	17.17
1936-37	0.01	0.70	0.00	1.86	0.00	6.93	3.09	7.99	4.79	0.03	0.11	0.00	25.51
1937-38	0.00	0.00	0.00	0.16	0.09	4.40	1.90	8.20	10.26	1.09	0.00	0.00	26.10
1938-39	0.00	0.00	0.19	0.14	0.08	4.94	2.84	1.27	3.62	0.17	0.10	0.00	13.35
1939-40	0.00	0.00	0.26	0.09	0.02	1.41	6.39	4.87	0.82	1.06	0.02	0.00	14.94
1940-41	0.00	0.00	0.00	0.75	0.43	8.92	9.68	8.21	11.71	5.50	0.01	0.00	45.21
1941-42	0.03	0.01	0.00	0.89	0.44	5.00	0.80	0.75	1.76	3.19	0.00	0.00	12.87
1942-43	0.00	0.00	0.03	1.44	0.62	1.36	12.84	4.21	2.92	0.92	0.03	0.00	24.37
1943-44	0.00	0.00	0.00	0.39	0.12	5.57	1.44	7.05	1.74	1.57	0.01	0.06	17.95
1944-45	0.00	0.00	0.00	0.00	2.66	1.23	0.60	5.87	4.87	0.00	0.00	0.00	15.23
1945-46	0.00	0.00	0.06	0.73	0.37	6.35	0.40	0.72	2.69	0.00	0.05	0.02	11.39
1946-47	0.00	0.00	0.00	0.89	5.95	3.17	0.60	0.76	1.80	0.10	0.08	0.00	13.35
1947-48	0.00	0.01	0.05	0.22	0.00	0.37	0.00	1.71	4.29	2.01	0.43	0.17	9.26
1948-49	0.00	0.00	0.00	0.08	0.00	2.64	1.40	1.35	2.78	0.24	2.43	0.03	10.95
1949-50	0.00	0.00	0.00	0.02	1.72	4.16	2.54	2.76	1.29	0.61	0.05	0.01	13.16
1950-51	0.81	0.02	0.41	1.21	1.88	0.50	2.53	1.21	1.20	1.45	0.01	0.01	11.24
1951-52	0.00	0.06	0.00	0.49	2.04	4.80	13.89	0.71	7.37	1.79	0.00	0.08	31.23
1952-53	0.03	0.01	0.04	0.10	3.60	5.26	1.78	0.03	0.71	1.42	0.17	0.29	13.44
1953-54	0.00	0.00	0.01	0.00	2.08	0.09	5.98	2.95	3.81	0.44	0.06	0.02	15.44
1954-55	0.00	0.02	0.00	0.03	2.03	3.60	4.39	2.29	0.70	3.45	0.40	0.01	16.92
1955-56	0.00	0.01	0.00	0.00	1.36	6.07	7.19	1.15	0.00	2.42	1.64	0.00	19.84
1956-57	0.00	0.00	0.00	0.11	0.00	0.14	5.39	3.74	0.54	2.31	1.57	0.06	13.86
1957-58	0.00	0.00	0.00	1.41	0.51	4.51	3.71	9.84	6.20	5.43	0.33	0.00	31.94
1958-59	0.00	0.00	0.27	0.00	0.11	0.04	2.68	5.05	0.00	0.89	0.02	0.00	9.06
1959-60	0.00	0.00	0.01	0.01	0.00	1.01	3.12	3.39	0.63	2.64	0.00	0.01	10.82

i.

TABLE 4.

CITY OF SANTA BARBARA, CALIFORNIA MONTHLY AND SEASONAL PRECIPITATION

YEAR	JUL	AUG	SEP	OCT	NOV	DEC	JAN	FEB	MAR	APR	MAY	JUN	SEASON
1960-61	0.00	0.00	0.00	0.09	6.57	0.41	1.81	0.02	0.80	0.20	0.09	0.00	9.99
1961-62	0.00	0.01	0.04	0.00	3.74	1.47	2.18	17.33	1.41	0.00	0.00	0.00	26.18
1962-63	0.00	0.00	0.00	0.42	0.00	0.12	1.79	5.39	4.09	2.42	0.29	1.21	15.73
1963-64	0.00	0.07	0.91	0.88	3.56	0.00	1.45	0.00	2.33	0.84	0.10	0.05	10.19
1964-65	0.00	0.00	0.01	0.80	2.59	4.94	0.76	0.46	2.33	6.55	0.02	0.00	18.46
1965-66	0.00	0.00	0.05	0.00	8.26	3.53	1.51	0.78	0.06	0.02	0.12	0.01	14.34
1966-67	0.01	0.00	0.04	0.04	3.31	5.63	7.61	0.50	2.57	5.18	0.00	0.00	24.89
1967-68	0.00	0.00	0.23	0.00	4.05	1.09	1.44	2.02	4.22	0.62	0.00	0.00	13.67
1968-69	0.00	0.02	0.00	1.03	0.65	1.81	15.55	8.35	1.00	1.92	0.06	0.08	30.47
1969-70	0.02	0.00	0.05	0.07	2.03	0.20	3.23	3.80	2.48	0.00	0.00	0.00	11.88
1970-71	0.02	0.00	0.00	0.05	4.54	4.67	1.21	0.88	0.82	0.73	1.10	0.00	14.02
1971-72	0.00	0.00	0.00	0.00	0.48 [.]	7.33	0.12	0.53	0.00	0.15	0.02	0.01	8.64
1972-73	0.01	0.00	0.01	0.13	5.47	0.84	6.37	7.38	3.01	0.05	0.03	0.03	23.33
1973-74	0.00	0.03	0.01	0.70	1.75	1.54	8.04	0.00	4.93	0.28	0.00	0.05	17.33
1974-75	0.00	0.00	0.05	0.90	0.09	7.21	0.00	5.27	3.86	0.80	0.00	0.00	18.18
1975-76	0.00	0.03	0.00	0.03	0.20	0.07	0.00	5.61	1.25	0.79	0.01	0.20	8.19
1976-77	0.00	0.00	4.01	0.12	1.06	0.94	4.01	0.19	1.59	0.00	2.96	0.10	14.98
1977-78	0.00	0.50	0.00	0.00	0.00	7.04	9.35	9.62	12.38	1.98	0.00	0.05	40.92
1978-79	0.00	0.00	1.22	0.00	3.18	1.70	4.30	4.82	6.47	0.05	0.00	0.00	21.74
1979-80	0.00	0.00	1.07	0.45	0.20	1.21	6.71	11.11	3.05	0.56	0.23	0.00	24.59
1980-81	0.36	0.03	0.02	0.00	0.00	1.70	3.60	2.67	6.00	0.65	0.00	0.00	15.03
1981-82	0.00	0.00	0.00	0.50	2.10	0.69	2.94	0.47	5.79	2.57	0.01	0.04	15.11
1982-83	0.00	0.00	2.04	0.62	4.64	2.49	10.45	8.67	4.68	4.45	0.18	0.03	38.25
1983-84	0.00	1.48	3.66	0.83	3.07	4.66	0.21	0.12	0.54	0.12	0.00	0.01	14.70
1984-85	0.01	0.16	0.67	0.35	1.99	4.17	1.57	2.19	1.87	0.02	0.00	0.00	13.00
1985-86	0.01	0.00	0.09	0.58	3.50	0.88	2.07	7.94	6.20	0.87	0.00	0.00	22.14
1986-87	0.00	0.00	1.45	0.00	1.35	0.12	1.56	3.00	3.66	0.16	0.00	0.17	11.47
1987-88	0.13	0.00	0.00	2.45	1.08	3.12	2.43	2.19	0.02	3.97	0.04	0.02	15.45
1988-89	0.02	0.00	0.17	0.00	1.01	3.64	0.35	2.53	0.90	0.34	0.39	0.05	9.40
1989-90	0.00	0.00	0.08	0.55	0.63	0.00	2.01	2.36	0.00	0.09	0.80	0.00	6.52
		0.00	0.00	0.00	0.15	0.05	0.04	0 55	11 05	0.00	0.00	0.11	17 50
1990-91	0.00	0.00	0.06	0.00	0.15	0.05	2.26	3.55	11.05	0.03	0.00	0.44	17.59
1991-92	0.00	0.18	0.02	0.50	0.09	3.24	2.38	8.74	4.16	0.02	0.27	0.00	19.61
1992-93	0.51	0.01	0.00	0.87	0.00	4.97	9.29	7.45	3.05	0.00	0.05	0.71	26.90

.

SANTA BARBARA PRECIPITATION 1868-1990




Figure 3. Source: Santa Barbara County Flood Control and Water Conservation District





Figure 4a. PRECIPITATION TRENDS AT SANTA BARBARA, CA (1867-1992)


Courtesy: Corey Rossman



Inches

25

Courtesy: Jim Goodridge



D W R. No 535

VIII. FLOODING

The city of Santa Barbara lies within Santa Barbara County's south coast watershed. which encompasses a drainage basin of 289 square miles. Streams flowing through the watershed generally originate on the orographically favored steep southfacing slopes of the Santa Ynez Mountain Range. The streams, coursing southward through gravelly alluvial fans, are relatively short in and respond swiftly length and dramatically to heavy rainfall.

The U.S. Army Corps of Engineers has identified four creeks that subject the city of Santa Barbara to flooding: Sycamore, Mission, Arroyo Burro, and San Roque (Fig. 6). These streams drain an area of 29.7 square miles; Mission Creek having the largest drainage area with 11.5 square miles. Mission Creek parallels the 101 Freeway from Oak Park through the central business district, crossing under the freeway near the Amtrak station. There is an additional section, the Central Drainage Area, covering 2.5 square miles north and east of lower State Street, which is inadequately drained and is subject to frequent urban and street flooding.

Streamflow within the city of Santa Barbara is negligible, except during and immediately following precipitation. Neither climate nor watershed type is supportive of continuous runoff within the district.

Heavy rains can occur locally in combination with areas of compacted or

impervious soils, steep channel gradients, burned-over sections, and man-made projects and obstructions. Downpours can then translate rapidly into intense, debris-laden floods within the city. The U.S. Army Corps of Engineers (1975) has cautioned that "even worse flooding" is possible in the future.

There is evidence that flooding has occurred periodically in the Santa Barbara area far back into prehistoric times. The Santa Barbara littoral is itself constructed of alluvium deposited from the Santa Ynez Mountains.

Michaelsen and Haston (1988) studied tree ring growth and reconstructed river flow data for the nearby Santa Ynez River watershed. This record indicates that remarkable flooding likely occurred in Santa Barbara County in 1565 and 1568. The 1568 event was extreme throughout southern California and may have signalled monthly rainfall rates approaching 10,000-year return period projections. The very heavy streamflows in the late sixteenth century predate European settlement in the area; thus no written record is available, but confidence is high that such events did occur. Excessive streamflow values in 1868 that had been projected by Michaelsen and Haston correlate well with the modern record December rainfall, 12.67 inches, observed by Dr. Shaw at Santa Barbara in 1867.

Flooding was mentioned by Spanish missionaries in the eighteenth and early nineteenth centuries. In the modern era, flooding was reported at Santa Barbara in 1862, 1867, 1875, 1877, 1888, 1907, 1909, 1911, 1914, 1918, 1938, 1941, 1943, 1952, 1964, 1967, 1969, 1971, 1978, and 1983.

Beginning on December 24, 1861 and continuing through the end of January 1862, a five week period of heavy rains caused some of the largest flood discharges ever experienced in California. Measurable rain fell in nearby Ventura for 60 consecutive days. O'Neill (1939) stated that "immense slides of earth and rocks took place in the mountains [of Santa Barbara resulting in considerable County], change in the appearance of the country." The silting of the relatively shallow harbor at Goleta effectively ended its usefulness as a port.

On January 21, 1909, a record daily rainfall of 6.40 inches was recorded by observer George W. Russell, which flooded some of the Oak Park neighborhood where Mr. Russell lived, and also flooded lower State Street to a depth of three feet. The flooding at Oak Park was exacerbated by street improvements that were underway in the area at that time.

In January 1914, a 48-hour rainfall total of 8.20 inches, part of a heavy monthly rainfall of 15.91 inches, caused damage or destruction to a dozen houses, several commercial establishments, and six bridges in the Mission Creek area. In his official Weather Bureau record, George W. Russell wrote that "the enormous rainfall of 6.95 [inches] in 24 hours on the 25th followed by heavy damage to entire lower end of county." Newspaper accounts told of four and a half inches of rain in four hours within the city on the afternoon of January 25. If correct, these reports would document a rainfall with a return record of 500 years. During the event, the official daily rainfall record established on that day, 6.95 inches, has stood through the 1992-93 water year.

In 1940-41, the wettest water year on record at Santa Barbara, 45.21 inches of rain was recorded. There were reports of "flooding and silt deposition" within the city, but the rainfall was spread relatively evenly from December into April, mitigating damage.

In January 1952, Mission Creek flooded on two separate occasions. Many homes were damaged, and water one or two feet in depth flooded the lower State Street area (Fig. 8). In January 1967 and again in January 1969, heavy mudslides. caused debris rains deposition, and erosion. Within the city, the 1967 event caused considerable property damage and stream overflows. The 1969 flood, while generally much worse in the county, caused only relatively minor damage in the Oak Park area.

In February and March 1978, flooding occurred once again in the city as 22 inches of precipitation fell in the twomonth period. Mudslides did millions of dollars of damage to residences, as the heavy rains occurred after the destructive Sycamore Fire, which had occurred in the summer of 1977. Quasi-stationary storms, such as the January 1969 rainstorm that inundated the adjacent communities of Montecito and Carpinteria, have the potential to wreak havoc in Santa Barbara. Strong orographic uplifting along the precipitous slopes of the mountains just north of the city helps create a serious short-fused flood danger. High velocity water movements can move large quantities of debris through the city, blocking bridges and clogging channels.

A 100-year flood (the 1914 event approaches a 100-year storm) would be capable of creating such high stream velocities, resultant severe erosion and, in overflow areas, heavy deposition of debris and mud. Stream velocities of 3 to 15 feet per second (2 to 10 mph) would occur in main channels, except in lined sections of Mission Creek, where velocities could reach 33 feet per second. After the onset of heavy rains, flood stage is reached within the Santa Barbara Stream Group in 1.5 hours, and extreme flood peaks occur in about 3.5 hours. (Stage hydrograph, page 32. Note that an intermediate regional flood approximates a 100-year flood.) This allows for only a very short-fused warning time: Flood control and warning agencies must be extremely vigilant preceding and during heavy precipitation events.

A fire, such as the 1990 Painted Cave Fire, creates an additional flood hazard. With protective vegetation cover eliminated, ash and fine silt can combine to create a clay-like surface that promotes almost immediate runoff during moderate or heavy precipitation Such flooding occurred in episodes. November 1964, following the Coyote Fire. In that event, a wall of mud and debris up to 25 feet high was reported to have rushed through San Antonio Creek, just west of the city of Santa Barbara.

TABLE 5. CUMULATIVE FREQUENCY DISTRIBUTION OF ANNUAL PRECIPITATION FOR SANTA BARBARA, CA
Probability of receiving less than the value indicated, values in inches. (based on data from 1931-1960)
Probability %

	5	10	25	33	50	67	75 _	90	95
	6.9	8.7	12.0	13.5	16.6	20.0	22.0	28.3	32.4

TABLE 6. RAINFALL MAXIMUM	DEPTH/DU	•				-		
Return <u>Period (</u>		15min	30min	1hr	3hr	12hr	24hr	W/Y


RP2	0.15	0.31	0.48	0.68	1.18	2.27	2.79	16.72
RP5	0.21	0.44	0.69	0.97	1.70	3.27	4.02	24.05
RP10	0.26	0.54	0.85	1.20	2.10	4.03	4.95	28.86
RP25	0.31	0.64	1.00	1.42	2.47	4.76	5.84	34.77
RP50	0.34	0.72	1.13	1.59	2.79	5.36	6.58	39.03
RP100	0.38	0.80	1.25	1.77	3.09	5.94	7.29	43.13
RP200	0.42	0.88	1.37	1.93	3.38	6.51	7.99	47.15
RP500	0.47	0.98	1.53	2.15	3.77	7.25	8.90	52.35
RP1000	0.50	1.05	1.65	2.32	4.06	7.81	9.60	56.21
RP10000	0.62	1.29	2.03	2.86	5.00	9.61	11.81	68.83
========		========		========	========		=======	=====
Source:	Jim Good	dridge						
	_				-			

Based on rainfall statistics from 1940-1992 Data: Precipitation in inches; RP = return period in years

TABLE 7.


RAINFALL DEPTH/DURATION/FREQUENCY AT SANTA BARBARA, CA MAXIMUM RAINFALL FOR INDICATED MONTH

	<u>RP2</u>	RP10	RP50	RP100	RP500	RP1000	RP10000
JANUARY	3.02	8.59	13.69	15.83	21.37	22.80	29.62
FEBRUARY	3.11	8.79	13.43	15.30	19.11	21.23	26.35
MARCH	2.49	6.37	9.54	10.83	13.38	14.87	18.32
			2.01				20.02
APRIL	0.86	3.01	5.02	5.87	8.39	8.65	11.78
MAY	0.12	1.19	2.41	2.95	4.55	4.81	6.76
JUNE	0.00	0.36	0.91	1.17	1.84	2.29	2.84
0 0 1 1 2				/			
JULY	0.00	0.15	0.46	0.61	0.90	1.15	1.50
AUGUST	0.00	0.18	0.59	0.79	1.45	1.53	
SEPTEMBER	0.00	1.27	3.25	4.20	7.18	7.54	11.10
OCTOBER	0.38	1.62	2.79	3.29	4.65	4.94	6.59
NOVEMBER	1.28	3.74	5.76		8.40	9.14	
DECEMBER	2.61	6.45	9.51	10.74	13.47	14.59	18.20
5202.152.1							20120
SEASON	16.50	28.53	38.36	42.33	51.11	54.87	66.90
	======		=======			=========	======
Source: Jim	Goodri	dge an	d Santa	Barbara	County	Flood Co	ntrol
				stics fro			
						iod in y	ears



Engineers


31



PAST FLOODS



The 1952 Flood on Mission Creek perils this house near Gutierrez Street...



 \ldots and the house is finally destroyed by the rampaging floodwaters.

Figure 8.

Source: US Army Corps of Engineers

IX. WIND

Reliable, long-term wind data for the city of Santa Barbara do not exist. However, statistical data are available for Santa Barbara Airport, and some very good automated stations have been installed close to Santa Barbara since the mid-1980s.


Elford (1962) utilized an FAA study of surface winds at Santa Barbara Airport conducted between August 1936 and August 1940. These data showed that light wind (three mph or less) occurred 37 percent of the time. Wind speeds of 4-15 mph occurred 60 percent of the time and speeds greater than 15 mph were noted three percent of the time. The predominant wind direction, when winds measured four mph or more, was from the south or southwest. But when winds were stronger, 16 mph or more, the predominant direction was from the west or northwest.

Wind roses from Santa Barbara County Air Pollution Control District sites at Goleta, about nine miles west of the city of Santa Barbara, are included in this report (Fig. 9). The Exxon 10 site is at the ocean while the Goleta sensors are two miles inland. Both locations manifest a light nighttime drainage wind from the north-northeast.

Recent (1988-1991) data yield an average annual wind speed at Santa Barbara Airport of 6.1 mph. Concurrent data from the California Department of Water Resources suggest an average annual wind speed of 4.5 mph for the city of Santa Barbara. Highest average daily wind speeds, 6-8 mph, tend to be noted during the time from March into early June. Winds are lighter than average from July through October.

Strong winds, 40 mph or more, occur at Santa Barbara from three sources: (1) significant winter storms that approach the district from the Pacific Ocean, causing gusty winds to blow from the southeast across the Santa Barbara coastal strip; (2) the eastern Pacific high pressure area, as it pushes up against relatively low pressure over the Colorado River Basin and sets up a strong west wind in the Santa Barbara area; and (3) downslope, so-called "sundowner", winds, from the north or northwest, which can strongly affect sections of the city.

Elford (1962) estimated that a peak wind of 60 mph at Santa Barbara might be experienced "as often as once in 50 years" and that a peak wind of 80 mph might occur "once in 100 years." If and when such extreme winds are measured in Santa Barbara, they would most likely fit into the category of downslope winds.



Wind Rose at Goleta and Exxon 10 - West Campus

. 2

X. DOWNSLOPE WINDS

The Santa Barbara area's unique downslope winds, somewhat analogous to the Santa Ana winds that occur near Los Angeles, are known locally as "sundowners." Sundowners often occur during evening hours, hence the name.

Sundowners, like most downslope winds, occur in various degrees of severity. Light sundowner conditions create irregular rises in temperature at Santa Barbara with gentle offshore breezes. Stronger episodes, occurring about two or three times per year, result in sharp temperature increases and localized gusty winds. Rarely. perhaps only a few times in a century, a severe sundowner episode occurs, resulting in very strong and hot winds descending along the south-facing slopes of the Santa Ynez Range into the city of Santa Barbara. In these events, winds can reach gale force or higher within the city and downslope heating can make temperatures soar to well over 100°F.

The most phenomenal sundowner struck the city on Friday, June 17, 1859. Very hot north to northwest winds caused a severe dust storm that frightened the inhabitants of Santa Barbara. A United States Survey ship was anchored in Goleta Harbor that day and measured the temperature at 133°F at 2 p.m. The survey report, written in 1869 and quoted by Tompkins, stated:

> No human being could withstand such heat out of doors. All betook themselves

to their dwellings and closed every door and window, the thick adobe walls giving admirable protection. Calves, rabbits and cattle died on their feet.

Another report on the 1859 sundowner comes from an historical text on the History and Resources of Santa Barbara published in 1876 and quoted later in the Santa Barbara Morning Press:

On the 17th day of June, 1859, a hot wind, like a sirocco, visited the city of Santa Barbara. The wind was from the northwest and blew furiously, with a dense cloud of dust. The temperature rose to 136 Fahrenheit's scale. It commenced blowing about noon and continued until about half past three in the afternoon. Birds, rabbits, and tender lambs were killed; the leaves on the side toward the wind were scorched and died, and some fruit was blasted.

The temperature reading of 136°F (57.8° C) at Santa Barbara, if it were validated, would tie the record high temperature for the earth, set in Libya, North Africa, in 1922. Because of unique atmospheric mountain wave activity that occurs during downslope wind events near Santa Barbara, it is indeed possible that temperatures in the 130s°F could be reached within the city. Consistency within the historical source material inspires further confidence in the 1859 observations. However, the temperature measurements taken in that exceptional event are well outside of the official Santa Barbara temperature record, both statistically and temporally. Also, weather observation techniques were not standardized by the U.S. Government until the 1870's. Therefore, the observed 136°F and 133°F readings are considered to be possibly accurate, but not official.

Not quite as powerful as the 1859 downslope windstorm, another severe sundowner struck Santa Barbara on Saturday night and Sunday, June 16-17, 1917. Fires burned throughout the district, whipped by strong winds. The situation was especially serious at Carpinteria, where residents carried their property to the beaches to avoid an advancing canyon fire. In Santa Barbara, strong winds created a dust storm (Fig. 10) "such as had never been known in this city before." The winds blew at gale force from 6 p.m. on Saturday until 2 a.m. Sunday. The winds returned at 2 p.m. Sunday, when the temperature reached its all-time modern record of 115°F and the fourday heat wave peaked. George W. Russell wrote in the official log (Fig. 11):

> The hottest day ever recorded on the 17th. A hot wind during night 16th with velocity of 35 to 45 miles per hour, damaged fruit & other trees. Nuts are damaged, beans seem to have escaped.

Another spectacular sundowner windstorm occurred on June 27, 1990. Winds gusting at 40 to 70 mph caused the Painted Cave Fire to race from near the ridgeline of the Santa Ynez Mountains to across the 101 Freeway within the city limits of Santa Barbara. Temperatures reached 112°F in the area during the holocaust, as 5,000 acres and over 500 structures were burned (Fig. 13).

Although local meteorologists have long appreciated the forces at work in the downslope winds that affect Santa Barbara, many advances in \mathbf{the} understanding atmospheric involved dynamics in sundowner activity have been made only in the past few years. As late as 1982, an Independent Insurance Agent's report on the devastating Sycamore Fire (another wind-driven blaze) made the following observations:

The sundowner is of a much more local nature [than the Santa Ana wind]. Not much is known about sundowners at present. It is thought that in the Santa Barbara area they normally start over the ocean west of the Lompoc-Guadaloupe area and move until they are the behind Santa Ynez Mountains. When it drops to the ground, [the air] heats up and the humidity is very low. As it rushes down the mountainsides and through the canyons in the late afternoons and evenings it creates the winds known as "sundowners," lasting only a few hours. Wind speeds may be 40 to 50 mph but can be 80 mph or more [Fig. 12].

ŝ

At this writing, the dynamics of Santa Barbara downslope winds are under continuing investigation. It is believed that these conditions are caused by a combination of factors including (1) differences in atmospheric pressures between Santa Barbara and the portion of the county north of the Santa Ynez Mountains and (2) the formation of an atmospheric wave on the leeward side of the Santa Ynez Mountains. It is during the months of June and July that Santa Barbara is most vulnerable to the combination of severe downslope winds and heating.

The forecasting of sundowner winds at Santa Barbara has been enhanced in the past few years by research into these events, although some forecast techniques have been used for decades. The National Weather Service issues local forecasts for the city of Santa Barbara and environs, and routinely checks the potential for downslope windstorms. Using present technology, sundowners may be predicted as much as several days in advance.

XI. MISCELLANY

Relative Humidity.

Humidity is an indicator of the amount of water vapor in the air. Relative humidity is a ratio between the amount of water vapor actually in the air at a certain temperature and the amount of water vapor present when the air is saturated at the same temperature. This is expressed as a percentage. Relative humidity averages 67 percent at Santa Barbara on an annual basis. The average daily maximum relative humidity is 82 percent, generally occurring in the early morning hours around sunrise. The average daily minimum relative humidity is 55 percent, usually occurring in the early afternoon.

Humidities are higher than the annual average, by 10 to 20 percent, during the time from June through early October. Lowest levels of the atmosphere tend to be drier in the Santa Barbara area during the winter season; relative humidities are generally 10 to 20 percent below the annual average from November through April. Moist conditions are common in Santa Barbara. Relative humidities are frequently near 100 percent during late night and morning hours in the summer "marine laver" season.

Dry air with very low relative humidity is not frequently observed. Relative humidities of less than 20 percent occur with warming downslope (offshore) winds, when the district is under the influence of a large high pressure area. These conditions are most likely to be found from November through May, and probably occur on an average of from 10 to 15 days per year. Relative humidities as low as five percent have measured with been reliable instrumentation at Santa Barbara.

Evapotranspiration.

The term evapotranspiration refers to the total transfer of moisture from the soil to the air in a field growing a well-



and Jack Bailard were work > road from Shepard's In he coast highway was fille mobiles and wagons carry as from the probable put

heat of this district wa led by the heat on the Ven road from Foster Park of stricken town. Scores s drove over this highwa id to the persons still a rying residents to the coast illo this side of Ojal a free broken out at four o'clock y-up call had been sent t r aid. Ventura's resource e rapidly being used up fo calls for men also for worl on and for a blaze south o

Ojai in Ruine

noted resort town, was lesolution yesterday. Th duck with the burnt stub handsome residences wer ruins. Here and there b of good luck a home ha fire but half the residence were gone.

he women had left Oja e night before when th the greatest. All thes re sent over the Sant as the Ventura road ha the order of Sheri

roadside was burne faces between the tw ot enough to cause muc assing autos, it appeare But the action of the she for the autos leaving Oj with women and childre om the vigil and wit nsion as a result of th ips they had undergone. something-mighty-inspi way the Ojal people tod There wasn't a long fa s of Ojal yesterday afte the water supply hi the cheerful people to grace. The men we ruised with their batt nes. There were only illage for most of the ting fresh fires <u>a</u>! spread of the older con-At five o'clock yesterday flames which had apingcontrol a few hours inned by new gusts of sent to Ventura

Figure 10.

Weather Observer Says Relief Will Come Soon; Many Go to Beach.

Never in the history of Santa Barbara has there been such heat as that prevailing since the Ojai and Carpinteria fires started. Weather Observer George W. Russell's reports show that the nearest heat record to that which yesterday set a new top notch at 115 degrees, was in September, 1914, when the mercury climbed to 108. That heat, also, was occasioned by the presence of a fire, the Hope Park district then being in the grasp of the flames.

While a part of the heat was due to desert winds, it is stated that the highrecord was forced through the heat from the forest fires. The severity of the winds, also, are attributed to the fires, and normal weather conditions are expected when the fires which now are raging in the Ojal and above Carpinteria, have been extinguished.

In Santa Barbara a dust storm prevalled Saturday night, such as has never been known in this city before. The winds whipped papers from bill boards, lashed numerous American flags, left flying over the stores, into shreds, and drove the finest dust into dwellings and stores, no place being too securely closed to prevent the particles from sifting through.

The heat through the evening sent thousands to the beach. Last night the number was materially diminished, as a breeze from the ocean set in just before midnight, and gave a cool night to the city.

Never before have such numbers of people sought the surf to cool off, hundreds of surf bathers spent most of the Sunday either in the water or lolling on the beach, taking an occasional plunge to keep cool. The bath house yesterday served 750 bathers, and there were many others who used their automobiles in which to don their bathing suits, it being estimated that during the day, from Graham's on the east, to far beyond Castle Rock, over 1000 people took to the water during the day.

at High School

low-

man

lohn

A musical program, to which the public has been invited, will be given at the High School tonight Three throughout the ented late Carol Mar-

Musical Program

SANTA BARBARA MORNING PRESS FEATURE Monday, 18 June 1917 "HEAT RECORDS SET FOR CITY Headline: BY FIRES AND WIND" Mary Compton Courtesy: no novert Foxen Miss Gertrude Masters

of Incomparable

Every woman should be in creations of Richard Hudnut, 1 Perfumer.

The more exacting the requi tain that a Hudnut article will Violet Bec. Tollet Water

75a and \$1.40 bottle Violet Sec. Bath Salts 750 and \$1.50 bottle Violet Sec. Scap 25º Cake



HAPPY HOME C "NO PLACE LIKE

Monday, June 18, 1917.)

Steadfast, unmovable-for as : your labor is not in vain in the Lore

THE COMMONPI

God bless the commonplace! V Through wearisome and unpro Striving to carve new destinie

A trail through unaccustomed la The feverish years possess us, In our tense seeking for untroc The common heritage, nor care Altars to dear familiar things. When shadows lengthen and the Of life falls faintly on the half-he With vision dimmed and feeble

Back to the homely joys of bygor Love and a hearthstone and a d And through our tears we bless

NEW FOOTHILL SU

THE LATEST AND THE BEST-BUILDING

SURE THE VALUE OF YOUR LOT_ON

- JUST OPENED -

Contraction of the second	ALON	th of . ·	1	yan.	ne.	ENN C	, 191	Stati	on, <u>42</u> 	nd Pa	Ho	L.; County, ur of Observ	Sarta Ba	7 Am	Nean maximum, 77.8
	S	-		sl.	_	- Latit				ido, 117	; Tin	no used on t	vation, 64m	}	Mean minimum, 53 22
	PATE	<u>.</u>		ERATUR			1	BECIPITAT	<u> </u>	Darrs or Brow	PREVAILURE	OF PAY.	i Mincellan Prenomei		Nas, 65.5
		И.11- лон.	Marata Marata	Bares	Kat.	Tist or Beels sys.	TINS of RESIDE.	t Amount.	BROWFALL, 28 LINNAR,	64 GANEER AT TIME OF OMER- VALUES.	WIND Disseries,	. OF PAT.	PRENOLEI	34. · · · · · · · · · · · · · · · · · · ·	Marimum, // 5; date,/ 5
		70	4	82	2			•			h	ober	Tra to	10	Minimum 44 45 date, 75 of
	8	70			7				1		n	4		9	Greatest daily range, 3-3- 17
	8	70		21	\$	1					n	11		£	PRECIPITATION.
	4.	67	1		<u></u>			•			hr.			10	
	s	1.6.7	1	5 2						·[m.	*		10	TetalOinches
	6	65			4					· · · · · · · · · · · · · · · · · · ·	h	······			Greatest in 24 hours (date,
	Ţ	6.	<u> </u> 5 5					.		· • · · · · · · · · · · · · · · · · · ·	hr				
	8	22	S		S				<u> </u>	.					SNOW.
	9	11	S		3	•				-	•	- •			Total fall, inches; on ground 15th,
	11	20	1		4			-			•	1	•	·//	lackes; at end of month, inches,
	18	78	£ 4	53								. ,	1	9	
	13	80		<u>8.3</u>	3		.]	<u>.</u>			<u> </u>		··· .		
	16	9.		18 4	λ		•	•			·		Tota .		With .01 inch or more precipitation,
	15		5	4.4	ਸ <u>਼</u>	•	·[······;·				<u>A</u> C		Hels Ary		Clear, J.A .; partly cloudy, 0 ;; cloudy, 0
	16	107			X] N	·	······	·			3.0	·····	dia por	mall in	
		1/15	6		M M			•		1	2		Ton care		DATES OF-
	18	5			L		•				m	••	andtos	.	Killing frost,
	***	75	+ 6		st						n	••			Thunderstorms,
95. 	\$1	2			5						r	4	. 4	5-0.	Hall
	##	<u>. 7</u>		9.1	- [_n		· · ·		
	13	<u> ZS</u>		1 .			-				<u>P</u>	4	······	2,30	SHEEL
	¥	1.	5		<u>9</u>		+			· [· · · · · · · · · · · · · · · · · ·	hr in				Âuroras,
	95			31	 J					+			· · · ·	1	REMARKS.
	H	-4.6		3	⊊ 2		+		1			·	4	- ty	the hoterst-day ever und
	======================================	84	5		7 1					1	<u> </u>	~		••••••••••••••••••••••••••••••••••••••	
		84			2				1		Ĺ	~			tatte 17th a hot much dure
	**		- S ,	5 3							3			······	Might 16 the with Kelnety 5
	31								h	l					35- 15- miles for from dam
5/ 1 S		7 224	155	73 73	1	1	1			43310	1 14	1 .	1	1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1	fiel home flam

COPY JUNE 1917

~

י די 17






Figure 13. Santa Barbara resident waters roof with garden hose as downslope winds gusting to 70 mph spread flames across city, June 1990. The home was engulfed by flames just after this picture was taken.

Courtesy: Santa Barbara News-Press

established crop. Some of the water loss is by evaporation from the surface of the soil while other moisture is carried upward and transpired from the leaves and other surfaces of the plants. At the State of California Department of Water Resources site at Santa Barbara, evapotranspiration is measured for a grassy surface, which has good coverage and is well irrigated.

At Santa Barbara, the average annual evapotranspiration is 45.3 inches. The average daily evapotranspiration rate is 0.124 inches, with a range from 0.07 inches per day in January to 0.18 inches per day in May and June.

The estimated average date of exhausting soil moisture reserves after the wet season at Santa Barbara is June 5. This is the "dry date," the date when range grasses stop growing.

Average annual evaporation, as measured from a standard four-foot pan, is 69.0 inches per year. The monthly average ranges from 3.7 inches in February to 8.1 inches in July. Approximately two-thirds of the total annual evaporation occurs in the months from May through October.

Sky Cover, Clouds, and Fog.

Nineteenth century data that purported to measure cloud cover and fog was based on observations that were too imprecise to be of any scientific value. In fact, the early weather journals frequently reflected a desire on the part of the authors to prove that Santa Barbara's climate was superior to that of Boston or Philadelphia or Oakland.

An attempt was made between 1936 and 1938 to study the incidence of clouds and fog at the airport in Goleta. The study was of too short a duration to provide reliable climatological data.

Long-term aviation weather observations at Santa Barbara Airport, and the addition of solar radiation measurements at California Department of Water Resources sites, have helped to clarify sky cover patterns at Santa Barbara.

Dense fog, visibilities of 1/4 mile or less. occurs on an average of approximately 14 days per year. Dense fog resulting from airmass advection is typically observed from late August through the middle of October. September averages 2.9 days with dense fog and October averages 3.3 A typical statistical analysis days. indicates that February also has a significant number of dense fog days (2.8), but this anomaly is the result of including precipitation and fog together as an obstruction of vision due to fog. This yields a fog/day result that is excessive and unrepresentative.

1000

Santa Barbara is a mostly sunny place. The city averages 255 clear days per year (with zero to three tenths sky covered by clouds). There are an average of 50 partly cloudy days (four to seven tenths sky cover) and 60 cloudy days (eight tenths or more sky cover). Elford (1962) discussed general cloud cover patterns at Santa Barbara:

Cloudiness associated with migrating storms is minimal in Santa Barbara County, but there is a considerable amount of stratus cloudiness along the immediate coast. Sunshine averages around 60% to 70% of possible at Santa Barbara, increasing to near 80% during the fall period of the year.

More precise statistics on measured solar radiation are now available. Solar radiation has been monitored at Santa Barbara Airport since March 1988 by the Department of Water Resources at CIMIS station #67. This station received an annual average of 56.8 percent of the extra terrestrial radiation (ETR). The average daily solar radiation at Santa Barbara Airport is 412 Langleys per day (Ly/dy) (calories per square centimeter) for the four years from 1989 through 1992. The monthly average solar radiation ranges from 226 Ly/dy in December to 578 Ly/dy in June.

Solar radiation in California varies from a low of 45 percent of ETR at Caspar Creek on the North Coast to a high of 70 percent of ETR at Lancaster in the high desert of Los Angeles County.

XII. REFERENCES AND ACKNOWLEDGMENTS

AUTHORS:

- Elford, C. Robert, 1962: The Climate of Santa Barbara. Original typed manuscript.
- Elford, C. Robert, et al., 1965: *The Climate of Santa Barbara County*. University of California Agricultural Extension Service, Santa Barbara.
- Fritts, Harold C., 1991: Reconstructing Large-scale Climatic Patterns from Tree-Ring Data. University of Arizona Press, Tucson.
- Kelly, Robert, 1985: A Century of Flooding on Mission Creek, Santa Barbara County 1870's - 1970's: Two Histories in Conflict. University of California, Santa Barbara.
- Mason, J. D., 1883: *History of Santa Barbara County, California*. Thompson and West, Oakland.
- Michaelsen Joel and Haston Laura, 1988: Reconstruction of the Flows of the Santa Ynez River at Bradbury Dam, Santa Barbara County, California 1537-1964. University of California, Santa Barbara.
- Michaelsen, Joel et al., 1990: Estimating Drought Probabilities in California Using Tree Rings. University of California, Santa Barbara.
- O'Neill, O. H., 1939: *History of Santa Barbara County*. Union Printing Company, Santa Barbara.
- Tompkins, Walker A., Undated: Santa Barbara's Incredible Hottest Day (based on Tompkins' historical text, Goleta: The Good Land) Santa Barbara National Bank.
- Trewartha, Glenn and Horn, Lyle, 1980: An Introduction to Climate. McGraw-Hill Book Company, New York.

PUBLICATIONS:

County of Santa Barbara Flood Control and Water Conservation District Precipitation Data Report (1990).

- Flood Plain Information Santa Barbara Stream Group, U.S. Army Corps of Engineers, Los Angeles District (1975).
- Monthly Station Normals of Temperature, Precipitation, and Heating and Cooling Degree Days, 1961-90: California, National Climatic Data Center, Asheville, NC (1992)
- The Lessons of the Sycamore Fire, Santa Barbara, California, Independent Insurance Agents of Santa Barbara (1982)

ACKNOWLEDGMENTS

The following persons and agencies contributed time, data, and ideas to the preparation of this report. Their assistance is gratefully acknowledged.

Paul Angel Jim Ashby Linda S. Brainard Sally Cappon Mary Compton Peter C. Felsch Dennis R. Gibbs James D. Goodridge **Richard Hallerman** Phil Mann Kenneth B. Mielke Keith Meier Elaine Robinson Chris Parker Michael Redmon Tom Ross Corey Rossman

National Weather Service Western Regional Climate Center California Department of Water Resources **Free-Lance Writer KEYT** Television National Weather Service Santa Barbara County Flood Control State Climatologist (retired) Santa Barbara County Air Pollution Control **KEYT** Television National Weather Service, Western Region Headquarters National Weather Service, Western Region Headquarters National Weather Service, Western Region Headquarters Flight Instructor Santa Barbara Historical Society Museum National Climatic Data Center California State University, Chico

APPENDIX

ITEM	

<u>PAGE</u>

TABLE	8	WEATHER RECORDS, MEANS AND EXTREMES 49
TABLE	9	SANTA BARBARA DAILY AND MONTHLY NORMALS 50
TABLE	10	SANTA BARBARA DAILY RECORD HIGH/LOW TEMPS 62
TABLE	11	SANTA BARBARA AIRPORT RECORD TEMPS
TABLE	12	SUNRISE AND SUNSET TABLE

.

TABLE 8.

SANTA BARBARA, CALIFORNIA WEATHER RECORDS, MEANS AND EXRTREMES 1867-1993

34 deg 25' N Lat, 119 deg 42' W Long, elevation 130 feet

TEMPERATURE (deg F)	
Mean Annual Temperature	_60.1
Average Daily High	71.4
Average Daily Low	48.8
January Average Daily High	_64.7
January Average Daily Low	
August Average Daily High	77.5
August Average Daily Low	56.7
Highest Temperature All-Time	
Lowest Temperature All-Time	20 on 4 and 10 January 1949
Average Annual Heating Degree Days	2086
Average Annual Cooling Degree Days	289
Average Annual Heating Degree Days Average Annual Cooling Degree Days Average Growing Season (Above 32 deg)	342 days
PRECIPITATION (inches)	
Average annual	17.91
Average January	3.91
Average July	0.03
Maximum Precipitation in: 15 minutes	1.18 (1978)
t have a	1 70 (1000 00)
1 day	6.95 (25 Jan 1914)
1 month	17.33 (Feb 1962)
1 season	45.21 (1940-41)
calendar yr	_1.70 (1982-83) _6.95 (25 Jan 1914) _17.33 (Feb 1962) _45.21 (1940-41) _41.48 (1941)
Least Precipitation in: 1 season	4.49 (1876-77)
calendar yr	3.99 (1947)
Ave no of days/year rainfall 0.10 or mo	
Ave no of days/year rainfall 1.00 or mo	
THUNDERSTORMS	
Ave no of days/year with thunder	2.4
WIND (miles per hour)	
Average annual wind speed	4.5
Prevailing direction	from southwest
Maximum wind speed	data unavailable
	··
RELATIVE HUMIDITY (percent)	
Averge annual67	
Average daily maximum82	
Average daily minimum55	
BAROMETRIC PRESSURE	
	inches (1014.8 mb)

TABLE 9.

CITY OF SANTA BARBARA, CALIFORNIA DAILY AND MONTHLY WEATHER NORMALS

MONTH OF JANUARY

DAY	<u>HIGH</u>	LOW	PCPN	SEASON/DATE
01	64	40	.12	5.72
02	64	40		
03		40		
04	_64	40		
05	64	41		
06	_64	41	.12	6.32
07	64	41	.12	6.44
0880	64	41		
09	64	41		
10	65	_41		
11	65	_41	.12	6.92
12	65	41	. 12	7.04
13	65		.12	7.16
14	65	41	.13	7.29
15	65	41	.13	_7.42
16	65	42	.13	7.55
17	65	_42		
18	65	_42	.13	7.81
19	65	42		
20	65	42		8.07
21	_65	42	.13	8.20
22	65	42		
23	65	42		8.46
24	65	42		
25	65	_42	.13	8.72
26	65	42	.13	8.85
27	65	42		
28	65	42		
29	65	42		
30	65		.14	
31	65	42	.14	9.52

AVE DAILY MAX TEMP: 64.7 AVE DAILY MIN TEMP: 41.4 MEAN MONTHLY TEMP: 53.1

AVE DAILY MAX TEMP: 64.7 AVE MONTHLY PRECIP: 3.92 IN.

MONTH OF FEBRUARY

DAY	HIGH	LOW	PCPN	SEASON/DATE
01	65	42	. 14	9.66
02				9.80
03	65	42		9.94
04				10.08
05			.14	
06	65	42	.14	10.36
07			. 14	10.50
08	65		. 14	10.64
09	65		.14	
10	65			10.92
11	65	43	. 14	11.06
12	65		. 14	
13	65		. 14	
14	66			11.48
15	66	43	. 13	11.61
16	66	43	.13	11.74
17			.13	
18			.13	
19		44	.13	
20	66		. 13	
21	66	44	.13	12.39
22	66	44	.13	12.52
23	66	44	.13	12.65
24	66	44	.13	12.78
25	66	44	.13	
26	66	44	.13	13.04
27			.13	
28	66	44	.12	

AVE DAILY MAX TEMP: 65.5 AVE DAILY MIN TEMP: 43.2 MEAN MONTHLY TEMP: 54.3 AVE MONTHLY PRECIP: 3.77 IN.

TEMPERATURES BASED ON 1901-1960 DATA PRECIPITATION BASED ON 1867-1993 DATA

NOTE: NO NORMAL VALUES FOR FEBRUARY 29TH ARE INCLUDED; IN COMMON PRACTICE THE NORMAL VALUES FOR THE 28TH ARE USED FOR THE 29TH IN EACH LEAP YEAR. FEBRUARY TEMPERATURE AND PRECIPITATION MONTHLY VALUES ARE NOT ADJUSTED FOR LEAP YEARS.

MONTH OF MARCH

.

DAY	HIGH	LOW	PCPN	SEASON/DATE
01	67	44	.12	13.41
02		44	.12	13.53
03	67	44	.12	13.65
04	67		.12	
05	67		.11	
06	67	44	.11	13.99
07		44	.11	14.10
08	67	44	.11	14.21
09	67	44	.11	14.32
10	67			14.43
11	67	45	.11	14.54
12	67	45		14.64
13	67	45	.10	14.74
14	67	45	.10	14.84
15	67		. 10	
16	67	45	.10	15.04
17	67	45		15.14
18		45	.09	15.23
19			.09	15.32
20	67		. 09	
21	67	45	.09	15.50
22		45		15.58
23	~ ~	45	.08	15.66
24	<u> </u>	45	.08	15.74
25	68	45	.08	15.82
26	68	45	.07	15.89
27		45	. 07	15.96
28		45	.07	16.03
29		46	.07	16.10
30	·····	46	.07	16.17
31				16.24

AVE DAILY MAX TEMP: 67.3 AVE DAILY MIN TEMP: 44.8 MEAN MONTHLY TEMP: 56.0

AVE MONTHLY PRECIP: 2.95 IN.

MONTH OF APRIL

٤,

DAY	HIGH	LOW	PCPN	SEASON/DATE
01	68	46	.06	16.30
02	68	46	.06	16.36
03	68		.06	
04	68	47	.06	16.48
05	68		.06	
06	68	47	.06	16.60
07	69		.05	
08	69		.05	
09	69	47	05	
10	69			16.80
11	69	47	05	16.85
12			.05	
13			.05	
14			.04	
15	69		.04	
16	69	48	04	17.07
17	69	48		17.11
18	69			
19	69			17.19
20	69	48	.03	17.22
21	70		.03	
22	70	48	.03	17.28
23	70	48	.03	17.31
24	70	48	03	17.34
25	70	48	.02	17.36
26	70	49	.02	17.38
27	70	49		17.40
28	70			17.42
29	70	49	.02	17.44
30	70	49	.02	17.46
•••		· =		

AVE DAILY MAX TEMP: 69.1 AVE DAILY MIN TEMP: 47.6 MEAN MONTHLY TEMP: 58.3 AVE MONTHLY PRECIP: 1.22 IN.

MONTH OF MAY

DAY	HIGH	LOW	PCPN	SEASON/DATE
01	70	49	. 02	17.48
02	70	49	.02	17.50
03	70	49	.02	17.52
04		49	.02	17.54
05	70	49	.02	17.56
06	70	49	.01	17.57
07	70	49	.01	17.58
08	70	49	.01	17.59
09	70	49	.01	17.60
10	70	49	.01	17.61
11	70	49	.01	17.62
12	70	49		17.63
13	70			17.64
14	70	49	.01	17.65
14 15	70	49	.01	17.66
16	70	50	.01	17.67
17		50	.01	17.68
18	71	50	.01	17.69
19	71	50		17.70
20		50		17.71
21	71	50	.01	17.72
22	71	50	.01	17.73
23	71	50	.01	17.74
24	71	51		17.75
25	71	51	.01	17.76
26	71	51	.01	17.77
27	71	51		
28	71	51		17.79
29	71	52		
30	71		.01	17.81
31	72	52	.01	17.82

MEAN MONTHLY TEMP: 60.2

AVE DAILY MAX TEMP: 70.5 AVE MONTHLY PRECIP: 0.36 IN. AVE DAILY MIN TEMP: 49.9

_

MONTH OF JUNE

DAY	HIGH	LOW	PCPN	SEASON/DATE
01	72	52	.00	17.82
02	72	52	.00	
03	72	52	.00	
04	72	52	.00	
05	72			17.82
06	72	52	.00	17.82
07	72	52	.00	
08	72	52	.00	
09	72	52	.00	
10	72	52	.01	17.83
11	72	52	.01	17.84
12	72		.01	
13			.01	
14	73	52	.01	17.87
15	73	53	.01	17.88
16	73	53	.01	17.89
17	73	53	.01	17.90
			.01	
19	73	53	.00	17.91
20	73	53	.00	17.91
21	73	53	.00	17.91
22	74	53	.00	
23	74	54	.00	
24	74	54	.00	
25	74	54	. 00	
26	75	54	.00	17.91
27	75	54	.00	17.91
28	75	04	. 00	17.91
29	76	54	.00	17.91
30	76	55	.00	17.91

AVE DAILY MAX TEMP: 73.1 AVE DAILY MIN TEMP: 52.8 MEAN MONTHLY TEMP: 63.0

AVE MONTHLY PRECIP: 0.09 IN.

MONTH OF JULY

DAY	HIGH	LOW	PCPN	SEASON/DATE
01	76	55	.01	0.01
02	76	55	.01	0.02
03	76	55	01	0.03
04	77	55	.00	0.03
05	77		.00	0.03
06	_77	56	.00	0.03
07	77	56	00	0.03
08	77	56	.00	0.03
09	77	56		0.03
10	77			0.03
11	77	56	.00	0.03
12	77	56	.00	0.03
13	77	56	.00	0.03
14	77	56	.00	0.03
15	77	56	.00	0.03
16	77	57	.00	0.03
17	77	57		0.03
18	77	57	.00	0.03
19	77	57		0.03
20	77		.00	
21	77	57	.00	0.03
22	77	57		0.03
23	77	57		0.03
24	77	57		0.03
25	77		.00	
26	77	57	.00	0.03
27	77	57		0.03
28	77	57		0.03
29	77	57		0.03
30		57	.00	
31	77	57	00	0.03

.

AVE DAILY MAX TEMP: 76.9 AVE MONTHLY PRECIP: 0.03 IN. AVE DAILY MIN TEMP: 56.4 MEAN MONTHLY TEMP: 66.7

MONTH OF AUGUST

DAY	HIGH	LOW	PCPN	SEASON/DATE
01	77	57	.00	_0.03
02				0.03
03	77	_57		
04	77	57		
05	_77	57	.00	0.03
06	77	57	.00	_0.03
07	77	57		
08	77	_57		
09	77	57		0.03
10	77	_57		
11	77	57	.00	0.03
12	77	57	.00	0.03
13	77	57		
14	_77			
15	77			
16	77	57	.00	0.03
17	78	57	.00	0.03
18	78	57		0.03
19	78	57		0.03
20	78			_0.03
21	78	57	.00	0.03
22	78	57	.00	0.03
23	78	56	.00	_0.03
24	78	56	.00	0.03
25	78	56		_0.03
26	78	56	.00	0.03
27	78	56	.00	0.03
28	78			
29	78		.01	0.05
30	78	56	01	_0.06
31	78	56	.01	0.07

AVE DAILY MAX TEMP: 77.5 AVE DAILY MIN TEMP: 56.7 MEAN MONTHLY TEMP: 67.1 AVE MONTHLY PRECIP: 0.04 IN.

MONTH OF SEPTEMBER

DAY	HIGH	LOW	PCPN	SEASON/DATE
01	78	56		0.08
02	78	56		0.09
03	78	56	.01	0.10
04		56		0.11
05		56		0.12
06	78			
07	78		.01	0.14
08	78	56		0.15
09	78	56		0.16
10	78	56	.01	0.17
11	78		.01	
12	78		01	
13	78		.01	
14	78			0.21
15	78	55	.02	0.23
16	77	55	.02	0.25
17	77		02	
18	77		01	
19	77		.01	
20	77	54	.01	0.30
21	77			
22	77	54	.01	0.32
23	77	54	.01	0.33
24	77	54	.01	0.34
25	77	54	01	
26	77	53		0.36
27	77		.01	
28	77	53	.01	0.38
29	77	53	.01	0.39
30	77	53	.01	0.40

AVE DAILY MAX TEMP: 77.5 AVE DAILY MIN TEMP: 54.8 MEAN MONTHLY TEMP: 66.2 AVE MONTHLY PRECIP: 0.33 IN.

MONTH OF OCTOBER

DAY	HIGH	LOW	PCPN	SEASON/DATE
01	77	53	.01	0.41
02	77			
03	77	53	.01	0.43
04	76			_0.44
05	76	53	.01	0.45
06	76	53	01	0.46
07	76			0.47
08	76			0.48
09	76		.01	_0.49
10	76	52		
11	76	52	.01	0.51
12	76	52		
13	76	52	.01	0.53
14	76	51	.01	0.54
15	76		.01	_0.55
16	75	51	.01	0.56
17	75	51	.02	0.58
18	75	51	.02	0.60
19	75	_50		
20	75			
21	75	50	. 02	0.66
22	75	50	.02	0.68
23	75	49	.03	0.71
24	74	49	.03	0.74
25	74	49		
26	74	49	03	0.80
27	74	48	.04	0.84
28	74	48	.04	_0.88
29	74	48	04	_0.92
30	74	47	04	0.96
31	74	47	04	1.00

AVE DAILY MAX TEMP: 75.3 AVE DAILY MIN TEMP: 50.7 MEAN MONTHLY TEMP: 63.0 AVE MONTHLY PRECIP: 0.60 IN.

MONTH OF NOVEMBER

DAY	HIGH	LOW	PCPN	SEASON/DATE
01	74	47	.04	1.04
02	74	47		1.08
03	74		.04	
04	74	47	.05	1.17
05	74	47	.05	1.22
06	74		.05	
07	74	47	.05	1.32
08	74	46	.05	1.37
09	74	46	.05	1.42
10	73	46	.05	1.47
11	73	46	.05	1.52
12	73	46	.05	1.57
13	73	46	.05	1.62
14	73	45	.05	1.67
15	73	45	.05	1.72
16	72	45	.05	1.77
17	72	45	.06	
18	72		.06	
19	72	45	.06	
20	72	44	.06	2.01
21	71	44	.06	2.07
22	71	44	.06	
23	71	44	.06	2.19
24	71	43	.05	2.24
25	71	43	.05	2.29
26	71	43	.05	2.34
27	70		.05	
28	70	43		2.45
29	70	43	.06	2.51
30	70		.06	

AVE DAILY MAX TEMP: 72.3 AVE DAILY MIN TEMP: 45.1 MEAN MONTHLY TEMP: 58.7 AVE MONTHLY PRECIP: 1.57 IN.

MONTH OF DECEMBER

DAY	HIGH	LOW	PCPN	SEASON/DATE
01	70	43	.07	2.64
02		43		
03		43	.07	2.78
04		43		
05	69		.08	
06	69	43	.08	3.01
07	69	43	.08	3.09
08	69			3.18
09		43		
10	68	43		
11	68	43	.09	3.45
12	68	43		3.54
13	67	43		
14	67			3.74
15	67	43		
16	67	43	. 10	3.94
17	67	43	. 10	4.04
18	67	43	. 10	4.14
19	66	43	.10	4.24
20	66	42		
21	66	42	- 10	4.44
22	00 66	42		4.55
23	66	42	. 1 1	4.66
24	65	42		4.77
25	65	41	11	4.88
26	65	41	. 12	5.00
07	05 65	41	12	5.12
28	65	41		
29	05 65	40		5.36
30	64	40	.12	5.48
		<u> </u>		
31	64	40	12	5.60

AVE DAILY MAX TEMP: 67.0 AVE DAILY MIN TEMP: 42.3 MEAN MONTHLY TEMP: 54.7 AVE MONTHLY PRECIP: 3.03 IN.

TABLE 10.

CITY OF SANTA BARBARA, CALIFORNIA RECORD HIGH AND LOW TEMPERATURES

MONTH OF JANUARY

DAY	HIGHEST	YEAR	LOWEST	YEAR
01	78	1898	27	1919
02	78 1	893/1930	25	1924
03	;0;	1893	23	19/9
0.0			20	1949
05	0 +	1924	20 ^,1	1949
00	0J	1 324	25	1949
06	8019	02/03/62	25	1949
07	89	1962	29	1950
08	86	1923	26	1949
09	91 X	1923	25	1937
10	80	1908/61	20 X,Y	1949
11	83	1961	27	1949
12	85	_1928	25	1949
13	82	1948	29	1950/63
14	83	1975	29	1932
15	83	1975	30	1947
16	86	1920	27	1949
17	86	1920 1920/61	25	1949
10	85	1975	31	1937/43/47
19	85	1971	31	1910/43/47
20	87	1975	29	1922/37
21	80	1976	26	1937
22	81	1950	26	1937
23	80	1968	25	1937
24	81	1951	27	1949
25	84	1951	32	1932
26	70	1931/47	27	1949
20		1928	21	1949
28	02	1931	28	1945
29	_04	1931	29	1937
30	02 85	1883	28 29	1949
30	05		23	1949
31	_82	1935/53	29	1937/49
			~ * ~	
		14 ON 06 JAN 1		
-WARMEST	MINIMUM: 6	65 ON 30 JAN 1	883	
X - RECOR	D FOR MONT	гн ү	- RECORD A	LL TIME

CITY OF SANTA BARBARA, CALIFORNIA RECORD HIGH AND LOW TEMPERATURES

MONTH OF FEBRUARY

DAY	HIGHEST	YEAR	LOWEST	YEAR
01	85	1935	29	1919/49
		1963		
04	83	1963	29	1948/49
05	81	1963	29	1949
06	87	1953	29	1899
07		1954	31	1883
08	85	1907	27 X	1949
09	82	1907	27 X	1929
10	82	1 907 1 907	30	1929/49
11	85	1886	32	1929
12	86	1924	27 X	1949
13	90	1924	28	1949
		1943		
		1916		
16	87	1930	30	1990
17		1930		
18	87	1924	33	1949
19	82	1896/1928	33	1945
20		1961	34	1951
21	82	1961	35	1953
22	84	1961	33	1897
23	85	1954	35	1897/1939
24	90	1921	33	1942
25	94 X	1921	33	1942
26	85	1926/32	33	1935
27	89	1926	34	1935
28	87	1926	32	1951/62
29	80	1924/36	35	1948
30				
21				

31____

-COOLEST MAXIMUM: 46 ON 23 FEB 1913 -WARMEST MINIMUM: 64 ON 22 FEB 1900

X - RECORD FOR MONTH

Y - RECORD ALL TIME

CITY OF SANTA BARBARA, CALIFORNIA RECORD HIGH AND LOW TEMPERATURES

MONTH OF MARCH

.

DAY	HIGHEST	YEAR	LOWEST	YEAR
01	87	_1963	_34	1948/71
02	86	1936	32 19	48/53/87
03	86	1936 1931	30 X	1953
04		1959/68	_00 / <u></u>	1987
04	_02	1972	33	1040
03	_91	_1972		1340
06	_8518	399/1914	36	_1939
07	92	1904	_34	_1945
08	_821906/1	4/16/46	_35	1925
09	87	1916	_32	_1935
10	89	1934	37	1935
				_
11	91	1916	_33	_1935
12	86	1926	33	1893
13	89	1926	_34	_1954
14	88	1926	33	1944
15	88	1915	32	1917
16	90	1915	35	1927
17	88	1914	35	1898
18	87	1914	34	1898
19	90	1925	35	1886
20	92	1931	35	1935
	<u></u>			_
21	90	_1931	_3519	35/48/52
22		1926	32	1935
23	95	1926	34	1936
24	86 193	30/44/88	34	1929/33
25	94	1988	33	1948
26	96 X	1988	35	1948
27	87	1923/32	37	1942/48
28		1956	36 1	884/1945
29	89	1879	36	1884
30		1918		1938
31	87	1989	_33	_1977
		5 ON 23 MAR 193		
-WARMEST	MINIMUM: 60) ON 23 MAR 193	1 (ALSO 25	MAR 1988)
X - RECOR	D FOR MONTH	H Y -	RECORD ALL	TIME

CITY OF SANTA BARBARA, CALIFORNIA RECORD HIGH AND LOW TEMPERATURES

MONTH OF APRIL

DAY	HIGHEST	YEAR	LOWEST	YEAR
01	90	1985	36	1936/38
		1950		
03	92	1961	36	1945
	96	1989	37	1945/48
		1989		1928/46/75
06	101 X	1989	36	1921
07	89	1989	34	1929
08	85	1932	37	1929
09	89	1932		1922
		1968		
11	92	1940	38	1927/45
12		1940		
13	95	1888/98	37	1945
14	98	1925	39	1913/22
15		1966	38	1921/45
16	92	1931	36	1917/22
17	88	1889	34	1917/44
18	90	1924	38	1933
19	93	1924	38	1927
20	94	1958	38	1896
21	94	1958	37	1896
22	96	1910	39	_1904/32/44
23	97	1910	38	1932/61
24	90	1898 _1898/1965	39	1883
25	90	_1898/1965	38	1944
26		1965	40	1900/52
27	93	1921	39	1904
28	97	1921	30 X	1970
29	90	1959	37	1948
30	85	1916/29	40	1937/62
31			·	
-COOLES	ST MAXIMUM:	53 ON 11 APR	1912	

-COOLEST MAXIMUM: 53 ON 11 APR 1912 -WARMEST MINIMUM: 67 ON 15 APR 1914 (ALSO 21 APR 1958)

X - RECORD FOR MONTH Y - RECORD ALL TIME

MONTH OF MAY

DAY	HIGHEST	YEAR	LOWEST	YEAR		
01	88	1916	40 19	20/22/76		
02	88	1929	41	1991		
03	92	1984	40	1991		
04	90	1948		_1920/91		
05	92	1928/53	38	1991		
06	94	1941	38	1933		
07	98	1941	39	1921		
08	98	1941	39	_1930		
09	101 X	1923		_1930		
10	92	1934	44	1948		
11	92	1926	41	1922		
12	88	1926	42	1944		
13	95	1979	42			
14	97	1979		1908		
15	94	1956	39	1883		
16	99	1970	40	1908		
17	97	1967		1991		
18	98	1892		_1991		
19	89	1942	39			
20	94	1883		1991		
21	90	1911	42 19	23/39/48		
22	85	1930		_1908		
23	90	1930		1908		
24	92	1927		1908		
25	98	1896	40	1921		
				—		
26	101 X	1968	40	1953		
27	96	1960	_4519	17/24/29		
28	91	1960	_45	1917		
29	_87	1973	_44	1953		
30	92	_1909	_41	1988		
31	94	_1909	40	1923		
				—		
-COOLEST MAXIMUM: 55 ON 03 MAY 1909						
-WARMEST MINIMUM: 71 ON 02 MAY 1987						
X - RECORD FOR MONTH Y - RECORD ALL TIME						

MONTH OF JUNE

02 03	95 90	1904	40 X	1916
02 03	95 90	1904	4.0	
03	90	1000	42	1923
	97	1090	42	1933
04			43	1908
05	98	1883		1939
06	93	1890	41	1922
07	92	1890	44	1923/45
08	90	1906	43	1908
09	91	1894	44	1909
	102			1924
11	102	1877	45	1913
			45	
			42	
14	95	1917	44 188	7/1943/44
15	105	1917	45	1944/91
16	107	1917	_45	1962/91
17	115 X,Y	1917	451	
	100			1916
19	100	1929	46	1923/40
20	98	1929	45	1923
21	102	1973	48	1943
22	92	1929	44	1923
23		1931	45	1912
24	98	1976	46	1912/15
	100	1976	45	1912
26	90	1990	46	1943
27	103	1990	46	1923
28	96	1976		
29	92	1976	491935	5/45/47/49
30	101	1937	491	912/38/55
31				

-COOLEST MAXIMUM: 59 ON 13 JUN 1983 -WARMEST MINIMUM: 67 ON 25 JUN 1931

X - RECORD FOR MONTH Y - RECORD ALL TIME

MONTH OF JULY

DAY	HIGHEST	YEAR	LOWEST	YEAR
01	108 X	1937	49	1936/38
02	105	1985		
03	108 X	1907	47	
04	96	1907	4819	48/51/55
05	96	1907		1948
06	96	_1954	44 X	1948
07	101		49	1908/48
08		1954	47	1948
09	_89	1959	46	
10	98	1959	47	1944
				—
11	92	1953	48	1991
12	100			
	104		47	1932
14	97		50	1914
15	97			
16	105	1978	49	1940
17	98		49	
18	95		45	
19	96		45	
20		1936	45	
21	93	1960	47	_1943
22	91		50	1928/34
23	96		50	1928/47
24		1948	46	1948
25	97			1913/24
26	921	891/1943	48	_1913
27	107	1889	51	
28	90	_1931	50	
29	100		511	
30	99	_1930		
31	_93	_1930	45	1956
		2 ON 06 JUL 19		
-WARMEST	MINIMUM: 8	2 ON 19 JUL 19	92	
X - RECOR	D FOR MONT	Н Ү-	RECORD ALL	. TIME

MONTH OF AUGUST

DAY	HIGHEST	YEAR	LOWEST	YEAR
01	90	1946	50	1911
02	93	1918	49	1944
03		1920/29	49	1944/53
04	90	1917	50	1935
05	90	1983	49	1935/50
05	90	1363	43	1935750
06	96	1983	45 X	1923
07	91	1983	50	1923
08	91	1936	48	1950
09	90	1965	49	1939
10	99 X	1978	49	1948/50
		······································		
11	90	1922	501892	2/1919/32
12	90	1898		
13	91	1965	49	1917
14	91	1929		1924
15	91	1889	49	1934
, .				
16	93			918/24/49
17	94	1890	50 1910	/13/18/40
18	98	1890	46	
19	91	1986		1924
20	90	1930	48	1947
21	89	1891	49	1947
22	91	1925/59	48	1947
23	93	1968	49	1944/47
24	95	1931	49	1944
25	93	1931	501	944/47/49
26	94	1931		1940/53
27	98	1883	50	1947
28	97	1883	48	1908
29	96	1905		910/12/42
30	_98	1915	48	1945/46
31	98	1955	49	1947
		63 ON 31 AUG 1	91/	
		69 ON 20 AUG 1		0 4116 1984
		AND 19 AUG	•	U AUG 1304
X - RECOF	RD FOR MON	тн ү	- RECORD AL	L TIME
PERIOD OF	RECORD: (CONTINUOUS FRO	M JANUARY 1	883

MONTH OF SEPTEMBER

DAY	HIGHEST	YEAR	LOWEST	YEAR
01	103	1955	50	1928/51
		1955		
03	94	1945	48	1914/42
04	99	1961	49	1921/33
05	93	1961	46	1933
06	92	1928	46	1933
07	95	1944	49	1911/21
08	97	1944	46	1946
09	97	1984	48	1953
10	90	1983	47	1943
11	95	1983	44	1946
12	91	1959	46	1952
13	98	1971	46	1915
14	97	1909	46	1915/52
15	100	1909	47	1921
16	103	1913	49	1907/08/36
17	108 X	1913	46	1908
18	102	1939	45	1947
19	102	1912	44	1947
20	104	1939	45	1944
21	104	_1885/1939	43	1944
22	103	1939/43	45	
23	102	1939/44	45	1923/24
24	102	1978	45	1923/48
25	105	1978	43	1948
26	105	1963	38 X	1948
27	103	1963	41	1948
28	101	1963	43	1948
29	98	1924	42	1948
30	93	1945	42	1950
31	· · · · · · · · · · · · · · · · · · ·			
	ST MAYTMUM	60 ON 08 SEP	1914	

-COOLEST MAXIMUM: 60 ON 08 SEP 1914 -WARMEST MINIMUM: 74 ON 29 SEP 1917

X - RECORD FOR MONTH Y - RECORD ALL TIME

MONTH OF OCTOBER

DAY	HIGHEST	YEAR	LOWEST	YEAR	
01	98	1931	42	1950	
02	98	1945	42	1950	
	101	1958	41	1983	
04	01	1933	43		
05	101	1953	44	1908	
03	101	1999		1000	
06	100	1930	41	1937	
07	100	1971	42	1916	
08	99	1971	43	1937	
09	98	1909	42	1941	
10	94	1909/88	40	1916	
	0.0	1070	07	1004	
11		1976	37		
12	95	1940	36	1924	
13	97	1939	40	1924	
14	103 X	1950	42	1923/28	
15	102	1961	43	1934/48	
16	102	1961	40	1921/84	
17	95	1958/67	40	1938	
18	99	1933		1938/49	
19	100		00 	1932	
20	95			1920	
20	00		0`		
21	98	1929	37	1920	
22	98	1929/39	39	1920	
23	99	1959	40	1935	
24	95	1965	37	1935	
25	94	1968	38	1935	
0.6	91	1000	38	1954	
26 27	91	1983	38	1920/49	
		1935	41	1920/49 1919	
28	94	1931	40	1046/56	
29	97 96	1931		1946/56 1971	
30	90	1931/39	36	1971	
31	92	1918	34 X	1935	
-COOLEST MAXIMUM: 55 ON 28 OCT 1942 -WARMEST MINIMUM: 67 ON 08 OCT 1905					
-WARME	SI MINIMUM:	DI UN US UCI	1905		
X - RECORD FOR MONTH Y - RECORD ALL TIME					

MONTH OF NOVEMBER

DAY	HIGHEST	YEAR	LOWEST	YEAR
01	94	_1966	34	1971
02	91	1926		
03	94	1921	35	1935
04	95	1921	34	1935
05	91	1941	35	_1935
06	_93	_1956	35	_1935
07	_94	_1956	36	_1938
08	_98 X	_1914	_36	_1936/38
09	_97	_1956	_35	1948
10	_97	_1956	34	_1948
11	88	_1930/42	35	_1948
12		1900		
13	_89	_1974	_29	_1938
14	89	_1949	32	_1938
15		_1932	_34	_1938
16	90	_1919	_34	_1938
17	_89	_1936	_32	_1958
18	_92	_1932/36	36	_1941/58
19	_881	895/1932	_35	_1941/48
20	_92	_1917	_34	_1941
21	88	1924	32	_1947
22	89	1924	33	1948
23	90	1933	32	_1931
24	_92	1933	_28 X	_1931
25	_88	1956	_32	_1931
26	88	1956	35	1906
		1903		
28	_85	_1977	_30	_1919
29	88	_1977	31	_1948
30	87	1956	32	1948
31				

-COOLEST MAXIMUM: 55 ON 30 NOV 1975 (ALSO EARLIER DATES) -WARMEST MINIMUM: 74 ON 02 NOV 1992

X - RECORD FOR MONTH

Y - RECORD ALL TIME

MONTH OF DECEMBER

DAY	HIGHEST	YEAR	LOWEST	YEAR	
01	89	1959	32	1948	
02		1958	34	1913/36	
03		_1958		1948	
04	85	_1939		1909	
05	83 87			1948	
00		_1340	2J	_1340	
06		/1900/38	34	1920/42	
07	_90	1938	32	1948	
08	_91	1938	34	1960/71	
09		_1957	29	1951	
10	86	1921	31	1951	
11	_84	1921/58	32	1923/47	
12	88	1956	29	1949	
13	85	1953	32	1931/67	
14	82	1953	31	1945	
15	82	1942	26	1948	
16	82	1904	31	1948	
17		1929	32		
18	86		_27	1924	
19	84	1929	27	1924	
20	81 19	04/53/60	30	1948	
· · · · · · · · · · · · · · · · · · ·					
21	80	1906	29	1968	
22	80	1960	28	1990	
23	81	_1960	28	1990	
24	83	1929	26	1948	
25	85	1925	23 X	1924	
26	8219	25/47/56	24	1924	
27	84		27	1924	
28	86	1956	1	926/48/84	
29	86	_1956	28	1987	
30	85	_1917	27	1987	
31	82	_1980	26	1987	
		5 ON 26 DEC 19			
-WARMEST MINIMUM: 68 ON 05 DEC 1989					
X - RECORD FOR MONTH Y - RECORD ALL TIME					

TABLE 11.

RECORD HIGH AND LOW TEMPERATURES SANTA BARBARA, CALIFORNIA (AIRPORT)

MONTH OF JANUARY

DAY	HIGHEST	YEAR	LOWEST	YEAR
01	73	1969	25 X	1976
02	79		26	
03			28	
04			28	
04	75		27	
05	/ 5	_1954/58	27	1949
06	78	1964	27	1961
07	86 X	1962	29	1950
08	82	1941	28	1989
09			30	
10			26	
• -				
11			30	
12	79	_1986	29	1989
13	86 X	1991	26	1963
14		1991		1962/89
15	81	1991		
16	81	1976	29	1987
17			26	
18		1994	28	1987
19	79	1965	30	1943
20		1944	30	1987
21	86 X	1942	311	963/76/90
22	83	1950	31	1991
23	78	1953	32	1945
24	79	1951	31	1949
25		1951		1950
26	82	1947	31	1950/89
27	78	1986		1950
28	81	1976	30	1957/75
29	82	1953		1975
	81	1962	31	1968
31	82	1953	31	1946/85
		5 ON 11 JAN		
-WARMEST	MINIMUM: 5	9 ON 25 JAN	1969 (ALSO 1	1 JAN 1980)
X - RECORD FOR MONTH Y - RECORD ALL TIME				
PERIOD OF RECORD: CONTINUOUS FROM 01 JANUARY 1941 DATE OF COMPUTATION: FEBRUARY 1994				

.

MONTH OF FEBRUARY

DAY	HIGHEST	YEAR	LOWEST	YEAR
01	79	1954/76	32	1946
02		1954	33	1946/51
03		1963	30	1972
04		1954		1948/88
05	80	1953	30	1988
06	85	1953	25 X	1989
07	83	1954	29	1989
08	84	1954	33	
09	76	1988	33	
10	82	1988	33	1965
11	80	1971	33	1966
	83	1943	32	1944/48
13	84	1943		1942/48
		1943	31	1949
15	86 X	1977	26	
16	79	1977	32	1944/90
		1953		
		1981		
		1982		
		1982		
21	78	1948	32	1987
22		1992		
23		1968		
24		1986		
25		1992		
26	79	1992	33	1987
27		1976		
28		1976		
29		1968/72		
30	·			
31				
-COOLE	ST MAXIMUM:	49 ON 23 FEB	1951	
		58 ON 09 FEB		
X – RE	CORD FOR MON	ТН Ү	- RECORD AL	L TIME
DEDTOD				V 1011
		CONTINUOUS FR N: OCTOBER 1		(† 1941
		75		

,

MONTH OF MARCH

DAY	HIGHEST	YEAR	LOWEST	YEAR
01	87	1963	34	1962
02	86	1972	32	1971
03			30 X	1966
04		1972		1966/76
05	78	1993	35	1948/67
05		1993	30	1946/07
06	80	1979/93	35	1971/82
07	78	1955	33	1961
08	_80		33	1961
09	81	1944	33	1961
10	84	1984		951/58/60
	<u></u>			
11		1947		1951/88
12		1993	34	1950
13		1989	34	1954
14	84	1951	31	1954
15	84	1964	35	1944/73
16		1964	34	1991
17	87	1947	36	1963
18		1950	37	1954
19		1989		1982
20	_80	1988/90	34	1982
• •				
21	78			1942
22	78	1950		1944
23		1988	37	1952
24		1988		1957
25	87	1988	381	942/48/67
06	00	1000	26	1061
26		1988	36	
27	85	1988	33	
28		1989		
29	77		35	1961
30	80	1944	37	1949
31	_90	1989	37	1951
		51 ON 02 MAR 19 52 ON 20 MAR 19		7 MAR 1982)
X - RECORD FOR MONTH Y - RECORD ALL TIME				
PERIOD OF RECORD: CONTINUOUS FROM 01 JANUARY 1941				

MONTH OF APRIL

DAY	HIGHEST	YEAR	LOWEST	YEAR
01	87	1959	38	1975
		1950		
04	94	1989	39	1945
05	96 X	1989	37	1976
06	96 X	1989	39	1955/75
07	91			1955/75
08	81	1987/89	411967	/72/78/91
09	81	1980	34	1945
10	84	1955/68	40	1976
11	87	1947	39	1945
12	88	1947	39	1976
13	90	1947	37	1961
14	88	1964	40	1972
15	95	1966	40	1970
16	81	1958		
17	79	1948/92	35	1944
18	79	1992	37	1968
19	86	1958	39	1972
20	94	1958	37	1961
21		1987		
22	86	1987	38	1968/70
23	82	1966	37	1961
24	87	1985	36	1960
25	88	1948	401	960/64/67
26	88	1993	39	1989
27	85	1991	38	1984
28	85		41	1944
29	83	1959/81	39	1948
30	80	1993	40	1990
31				

-COOLEST MAXIMUM: 52 ON 01 APR 1967 -WARMEST MINIMUM: 67 ON 15 APR 1966

4

()

X - RECORD FOR MONTH Y - RECORD ALL TIME

MONTH OF MAY

DAY	HIGHEST	YEAR	LOWEST	YEAR					
01	83	1952/61	43 1950/	54/55/67/88/90					
02	85	1970	_38 X						
03	88	1984	41	1942					
04	90	1948	40	_1965					
05	89	1953	39	1975					
			····· ··· ····························						
06	86	1991	40	1975					
07	81	1991	40	1964					
08	_86	1984	41	_1988					
09	_89	_1943	43	1957/64					
10	_80	_1950/53	41	1982					
11	78	_1979/88	_441942/	82/85/68					
12	92	_1988	42_1944/	67/70/85/89					
13	91	_1979		_1989					
14	_90	_1970	42	1961					
15	_91	1967/70	_43	1955/68					
-									
16	89	1956	_4419	53/60/86					
17	87	1955/71	42	1953					
18	85	1978	40	1991					
19	87	1959	39	1944					
20	86	1942	41	1948					
,									
21	78	1989	42	_1948					
22	77	_1949	_42	1961					
23	_82	1949	_39	1960					
24	89	1989	_42	_1957					
25	_90	1968	_42	_1953/89					
26	101 X	_1968	39	_1953					
27	95	_1960	44	_1980/87					
28		1978	_44	_1991					
29	81	_1989	41	_1953					
30	82	1943	_42	1944/61					
31	84	1943	_44	_1988					
-COOLEST MAXIMUM: 56 ON 18 MAY 1949 (ALSO ON 19 MAY 1972) -WARMEST MINIMUM: 62 ON 11 MAY 1992									
	· · ·								
X - RECOR	D FOR MONT	H Y -	RECORD ALL	TIME					

MONTH OF JUNE

DAY	HIGHEST	YEAR	LOWEST	YEAR
01	87	1949	41 X	1955
02	84	1951	43	1955/67
03	83	1990	44	1953/80
04	84		45	1942
05	_86	1946	41 X	1943
06		1954/85	45	1943/80
07	82		42	
08	_84	1984	44	1988
			44	1952
10	96	_1979	45	1952/64
11	_88	1946	46	1953/54
		_1981	44	1977
13	92	1956	45	1943
14	92	_1981	44	1943
15	95	1981	46	1955/62
16	100	1981	45	1962
17	96		46	1950
18	95	1957	48	1978
19	86	1971	48	1978
20	96	1973	46	1975
21		1973	47	1943
22	_96		47	
23	_97	1976	47	1991
24	95	1976	48	1991
25	_98	1962	46	1943
	103	1990	45	
	109 X,Y		47	
28	83	1992	47	1963
29	95	1960		1949/63/64
30	87		48	1943/55
31		<u> </u>		

-COOLEST MAXIMUM: 61 ON 03 JUN 1973 -WARMEST MINIMUM: 67 ON 09 JUN 1945

X - RECORD FOR MONTH Y - RECORD ALL TIME

MONTH OF JULY

DAY	HIGHEST	YEAR	LOWEST	YEAR							
01	97	1985	49	1955							
02	109 X,Y			1963							
03	102		50 1955/5								
04		_1983 1957	501955/56/78/87 461955								
05	92	1990	_40 47	1955							
05	92	_1990	_4 /	1900							
06	99	1954	49	1948/55							
07	101	1954	49	1966							
08	_89	1992	50	1944							
09		1959/92	47								
10	95	1971		1944							
11		_1964	_50	1952							
12	85	1953/90		1978							
13	86	_1990	_48	1941							
14		1978		1941							
15	105	1978	_49	1941							
16	82	_1946/92	_50	1941/43							
17	84	_1992	_50	1943/66							
18	_84	1992	45 X	1944							
19		1992									
		1992									
21		_1960									
22		_1960									
23	85	1943	_51	1987							
24	82 194	43/59/60	49	1948							
25		1943/77									
26		_1977	_50								
27	97	_1947	_50	1965							
28	86	1947	49	1949							
29	81 194	47/54/60	51	1987							
30		1947		8/49/70							
31	93	_1977	_50	1942/70							
-0001 557			O (ALCO ON	00 88 4070)							
		4 ON 01 JUL 196									
-WARMES	MINIMUM: 68	B ON 12 JUL 199	Z (ALSU UN	13 JUL 1992)							
X - RECOR	RD FOR MONTH	H Y - I	RECORD ALL	TIME							
	REPTOD OF RECORD, CONTINUOUS FROM OF LANUARY 1941										

MONTH OF AUGUST

DAY	HIGHEST	YEAR	LOWEST	YEAR					
01	87	1947	48	1944					
02	82	1943/45		942/44/56					
03		1945		1953					
04		1976		1956					
05		1961		1950					
•••	00								
06	91	1983	50	1951					
07	86	1983		1944					
08				1950					
09				1941/50					
10			48	1944					
11	85	1965	51	1949					
12									
13									
14		1992	49						
15	85	1992		1949/88					
<u></u>									
16	84	1962	50	1949/86					
17			511						
18		1945							
19	89	1986	50						
20	84	1941	50	1944					
21	85	1972	43 X	1941					
22	105 X	1972	49	1987					
23	90	1968	49	1944/87					
24	85		48	1944					
25	85	1985	48	1944					
26	86	1977		1944					
27	98	1971		1975					
28	92	1971	49	1942					
29	96	1971	49	1946					
30	101	1984	49	1941/57					
31	94	1955	48	1941					
	T MAXIMUM: (T MINIMUM: (G 1965 (ALSO C	DN 23 AUG 1972					
X - REC	ORD FOR MON	гн	Y - RECORD AL	L TIME					
PERIOD OF RECORD: CONTINUOUS FROM 01 JANUARY 1941 DATE OF COMPUTATION: OCTOBER 1993									

81

MONTH OF SEPTEMBER

DAY	HIGHEST	YEAR	LOWEST	YEAR
01	99	1955	48	1941
		1955		
		1982		
		1988		
05	_89	1961	_49	_1970
06	_97	1955	_46	_1973
07	_93	_1955		
08	_92	_1984	_46	
09	95	1984	47	
10	_94	_1956	_45	_1943
11	89	_1959/83	45	1946
12	88	1971	47	1985
13	93	_1963	47	
14	86	_1959	46	1952/66
15	_93	_1979	_45	_1970
16	90	1979	_48	1970/86
		1979		
18	104 X			
19	_95	_1984	_45	_1947
20	_95	_1984	_45	_1986
21	85	_1942	44	1944
		1987		
		1944		
		1978		
25	_98	1978	_47	
26	103	_1963	43	1948
27	99	1963	46	1970
		_1963		
29	91	_1945	44	1955
30	90	1980	_42 X	1950
31				

-COOLEST MAXIMUM: 62 ON 23 SEP 1955 -WARMEST MINIMUM: 81 ON 18 SEP 1979

X - RECORD FOR MONTH

Y - RECORD ALL TIME

MONTH OF OCTOBER

DAY	HIGHEST	YEAR	LOWEST	YEAR						
01	92	1965	42	1950						
02	91	1945/91	42	1950						
03	98		42	1950						
04	103 X	1987	42	1989						
05	98	1971	45	1955/57						
			· • • _							
06	97	1971	42	1941						
07	93	1951	42	1954						
08	92	1976	42	1970						
09		1988	40	1941						
10	93	1969	42	1953						
11		1983/91	42	1990						
12	83	1950	43	1960						
13		1950		1981						
14		1961								
15	98	1961	40	1941/60/66						
16	92	1971	39	1960						
17	92	1971		1966/71						
18		1970		1971						
19		1964		1949						
20	94 94	1964 1964		1949						
20	0+	100+	07							
21	93	1942	38	1949						
22		1965	40	1949						
23	97	1959	40	1953/75						
24	95	1965	36	1975						
25	90	1965	37	1975						
0.0	00	1000	20	1054						
26	90 86	1993	38 39	1954						
27		1993 1979	39 35	1970/89 1970						
28 29										
		1958/73	31 X							
30	85	1962/80	34	1971						
31	88	1949	37	1971						
	-COOLEST MAXIMUM: 59 ON 28 OCT 1942 -WARMEST MINIMUM: 64 ON 01 OCT 1963 (ALSO 03 OCT 1963 08 OCT 1972, 14 OCT 1976 AND 24 OCT 1982)									
X - RECORD FOR MONTH Y - RECORD ALL TIME										

PERIOD OF RECORD: CONTINUOUS FROM 01 JANUARY 1941 DATE OF COMPUTATION: NOVEMBER 1993

83

MONTH OF NOVEMBER

DAY	HIGHEST	YEAR	LOWEST	YEAR
01	91	_1966 _1992	36	_1971
02	_97 X	1992	_3819	43/71/89
03	_90194	45/50/93	_37	1943/89
04	93	_1976	35	1943
05	_87	_1976	_37	_1981
06	_83	_1941	_37	_1973
07	90	1956	36 19	47/61/90
08	_93	1956	_37	1990
09	_96	_1956	_34	_1943
10	_92	_1990	_35	_1943
11	86	1942	34	1943
		1986		
13	84	1949	35	
14	86	1949	34	1943
15	_85	_1949	_35	_1956/85
16	79	_1977/89	34	1958/64
		1948		
18	85	1949	34	1964
		1992		
20	_82	1989	_30 X	_1988
21	84	_1954/59	31	1979
22	83	1954	34	1988
23	90	1950	32	1941
24	82	1956/79	31	1941
25		1953/77		
26	89	_1977	34	1980/84
27	90	1977		
28	90	1977	32	1989
29	82	1988	30 x	1989
30	87	1977	32	_1975
31	<u> </u>			_

-COOLEST MAXIMUM: 53 ON 11 NOV 1978 -WARMEST MINIMUM: 62 ON 09 NOV 1958

X - RECORD FOR MONTH

Y - RECORD ALL TIME

MONTH OF DECEMBER

DAY	HIGHEST	YEAR	LOWEST	YEAR
01	85	1958	31	1991
02	82	1959	33	
03	89_X	1958	33	1957/90
04	78	1977	00 33	1989
04 05	82	1962	32	1968
05	02	1902	32	1908
06	83	1990	32	1959/78
07	82	1989	29	1978
08	80	1975	29	1960/78
09	82	1957		1956
10	81	1958	28	1951
11		1958		1976/80
12	86	1956	31	1949/85
13	82	1953	25	1967
14	79	1953/83	25	1967
15	82	1942	27	1967
16		1980	28	1975
17	76	1973	29	1990
18		1950	31	1975
19	79	1953	32	1975/92
20	80	1953	32	1967/91
21	80	1953	28	1968
22	76	1989	20 X,Y	
23	70 77	1950	20 ,,,21	
23	80	1989	21	1990
24	80	1989 1947	25	1990
29	01	1347	25	1990
26	82	1947	27	1987/90
27	83	1975	26	1988
28	81	1956	28	1990
29		945/56/80		1962/69
30		1980	29	1969
31			28	1990
		47 ON 18 DEC 60 ON 21 DEC AND 27 DE	: 1943 (ALSO 2	24 DEC 1964
X - REC	ORD FOR MON	тн	Y - RECORD AL	L TIME
PERIOD	OF RECORD:	CONTINUOUS F	ROM 01 JANUAR	RY 1941

DATE OF COMPUTATION: OCTOBER 1993 85 TABLE 12.

SUNRISE AND SUNSET AT SANTA MARIA, CALIFORNIA PACIFIC STANDARD TIME

NO. 1046

D 8 W	JAN.		JAN. FF		FEB. MAR.		MAR. APR.		MAY		JUNE		JU	LY	AUG.		SEPT.		OCT.		NOV.		DEC.	
DAY	Rise A.M.	Set P.M.	Rise A.M.	Set P.M.	Rise A.M.	Set P.M.	Rise A.M.	Set P.M.	Rise A.M.	Set P.M.	Rise A.M.	Set P.M.	Rise A.M.	Set P.M.	Rise Å.M.	Set P.M.	Rise A.M.	Set P.M.	Rise A.M.	Set P.M.	Rise A.M.	Set P.M.	Rise A.M.	Set P.M.
1 2 3 4 5	7 10 7 10 7 10 7 10 7 10 7 10	5 01 5 02 5 03 5 04 5 05	7 01 7 00 6 59 6 58 6 58	5 31 5 32 5 33 5 34 5 35	6 32 6 30 6 29 6 28 6 26	5 57 5 58 5 59 6 00 6 01	5 49 5 48 5 46 5 45 5 44	6 24 6 24	5 11 5 10 5 09 5 08 5 07	6 47 6 48 6 48 6 49 6 50	4 49 4 49 4 49 4 48 4 48	7 10 7 11 7 11 7 12 7 12 7 12	4 52 4 52 4 52 4 53 4 53	7 19 7 19 7 19	5 11 5 12 5 13 5 13 5 13 5 14	7 05 7 04 7 03 7 02 7 01	5 34 5 35 5 36 5 36 5 37	6 29 6 27 6 26 6 25 6 23	5 56 5 57 5 58 5 59 5 59 5 59	5 46 5 45 5 43 5 42 5 41	6 22 6 23 6 24 6 25 6 26	5 08 5 07 5 06 5 05 5 04	6 51 6 52 6 53 6 54 6 55	4 50 4 50 4 50
6 7 8 9 10	7 10 7 10 7 10 7 10 7 10 7 10	5 05 5 06 5 07 5 08 5 09	6 57 6 56 6 55 6 54 6 53	5 36 5 37 5 38 5 39 5 40	6 25 6 24 6 22 6 21 6 20	6 02 6 03 6 03 6 04 6 05	5 42 5 41 5 40 5 38 5 37	6 27 6 28 6 28 6 29 6 30	5 06 5 05 5 04 5 04 5 03	6 51 6 52 6 52 6 53 6 54	4 48 4 48 4 48 4 48 4 48 4 48	7 13 7 13 7 14 7 14 7 15	4 54 4 54 4 55 4 55 4 55	7 19 7 19 7 18 7 18 7 18 7 18	5 15 5 16 5 16 5 17 5 18	7 00 6 59 6 58 6 57 6 56	5 38 5 39 5 39 5 40 5 41	6 22 6 20 6 19 6 18 6 16	6 00 6 01 6 02 6 03 6 03	5 39 5 38 5 36 5 35 5 34	6 27 6 28 6 29 6 30 6 31	5 03 5 02 5 02 5 01 5 01	6 55 6 56 6 57 6 58 6 59	4 50 4 50
11 12 13 14 15	7 10 7 10 7 10 7 10 7 10 7 09	5 10 5 11 5 12 5 13 5 14	6 52 6 51 6 50 6 49 6 48	5 41 5 42 5 43 5 44 5 45	6 18 6 17 6 16 6 14 6 13	6 06 6 07 6 08 6 08 6 09	5 35 5 34 5 33 5 32 5 30	6 32 6 32 6 33	5 02 5 01 5 00 4 59 4 59	6 55 6 56 6 56 6 57 6 58	4 47 4 47 4 47 4 47 4 47 4 47	7 15 7 16 7 16 7 17 7 17 7 17	4 57 4 57 4 58 4 58 4 59	7 17 7 17 7 17 7 16 7 16 7 16	5 19 5 19 5 20 5 21 5 22	6 55 6 54 6 53 6 51 6 50	5 42 5 42 5 43 5 44 5 44	6 15 6 13 6 12 6 10 6 09	6 04 6 05 6 06 6 07 6 07	5 32 5 31 5 30 5 29 5 27	6 32 6 33 6 34 6 35 6 36	4 59 4 59 4 58 4 57 4 57	6 59 7 00 7 01 7 01 7 02	4 51 4 51 4 52
16 17 18 19 20	7 09 7 09 7 08 7 08 7 08 7 08	5 15 5 16 5 17 5 17 5 18	6 47 6 46 6 45 6 44 6 42	5 45 5 46 5 47 5 48 5 49	6 11 6 10 6 09 6 07 6 06	6 10 6 11 6 12 6 13 6 13	5 29 5 28 5 26 5 25 5 24	6 35 6 36 6 36 6 37 6 38	4 58 4 57 4 56 4 56 4 55	6 59 7 00 7 00 7 01 7 02	4 48 4 48 4 48 4 48 4 48 4 48	7 17 7 18 7 18 7 18 7 18 7 18	5 00 5 00 5 01 5 02 5 02	7 15 7 15 7 14 7 14 7 13	5 22 5 23 5 24 5 25 5 25	6 49 6 48 6 47 6 46 6 44	5 45 5 46 5 47 5 47 5 48	6 08 6 06 6 05 6 03 6 02	6 08 6 09 6 10 6 11 6 12	5 26 5 25 5 24 5 22 5 21	6 37 6 38 6 39 6 40 6 41	4 56 4 55 4 55 4 54 4 54	7 03 7 03 7 04 7 05 7 05	4 52 4 53
21 22 23 24 25	7 07 7 07 7 06 7 06 7 05	5 19 5 20 5 21 5 23 5 24	6 41 6 40 6 39 6 38 6 37	5 50 5 51 5 52 5 53 5 54	6 04 6 03 6 02 6 00 5 59	6 14 6 15 6 16 6 17 6 17	5 23 5 22 5 20 5 19 5 18	6 39 6 40 6 40 6 41 6 42	4 54 4 54 4 53 4 53 4 52	7 03 7 03 7 04 7 05 7 05	4 48 4 49 4 49 4 49 4 49 4 49	7 19 7 19 7 19 7 19 7 19 7 19	5 03 5 04 5 05 5 05 5 06	7 13 7 12 7 11 7 11 7 11 7 10	5 26 5 27 5 28 5 28 5 29	6 43 6 42 6 41 6 39 6 38	5 49 5 50 5 50 5 51 5 52	6 00 5 59 5 58 5 56 5 55	6 13 6 13 6 14 6 15 6 16	5 20 5 19 5 18 5 16 5 15	6 42 6 43 6 44 6 45 6 46	4 53 4 53 4 53 4 52 4 52	7 06 7 06 7 07 7 07 7 07 7 07	
26 27 28 29 30 31	7 05 7 04 7 03 7 03 7 02 7 01	5 29	6 35 6 34 6 33 6 33	5 55 5 56 5 57 5 57	5 57 5 56 5 55 5 53 5 52 5 50	6 18 6 19 6 20 6 21 6 21 6 22	5 17 5 16 5 15 5 14 5 12	6 43 6 44 6 44 6 45 6 46	4 52 4 51 4 51 4 50 4 50 4 50	7 06 7 07 7 07 7 08 7 09 7 09	4 50 4 50 4 50 4 51 4 51	7 19 7 19 7 19 7 19 7 19 7 19	5 07 5 07 5 08 5 09 5 10 5 10	7 09 7 09 7 08 7 07 7 06 7 05	5 30 5 31 5 31 5 32 5 33 5 33	6 37 6 35 6 34 6 33 6 31 6 30	5 53 5 53 5 54 5 55 5 56	5 53 5 52 5 50 5 49 5 48	6 17 6 18 6 19 6 20 6 21 6 22	5 14 5 13 5 12 5 11 5 10 5 09	6 46 6 47 6 48 6 49 6 50	4 52 4 51 4 51 4 51 4 51 4 51	7 08 7 08 7 09 7 09 7 09 7 09 7 09	4 58 4 58 4 59

Add one hour for Daylight Saving Time if and when in use.

E. W. WOOLARD Director Nautical Almanac U. S. Naval Observatory

I certify that the above data are the result of an accurate and true computation by the Nautical Almanac Office, United States Naval Observatory, an agency charged by Federal Statute (9 Stat. L 374, 375) with the duty of making such computations and publishing the results.

C. G. CHRISTIE Captain, USN Superintendent U.S. Naval Observatory

- The Usefulness of Data from Mountaintop Fire Lookout Stations in Determining Atmospheric Stability. Jonathan W. Corey, April 1979. (PE298899/AS) The Depth of the Marine Layer at San Diego as Related to Subsequent Cool Season Precipitation Episodes in Arizona. Ira S. Brenner, May 1979. (PE298817/AS) Arizona Cool Season Climatological Surface Wind and Pressure Gradient Study. Ira S. Brenner, May 1979. (PE298900/AS) The BART Experiment. Morris S. Webb, October 1979. (PB80 155112) Occurrence and Distribution of Flash Floods in the Western Region. Thomas L. Dietrich, December 1979. (PB80 160344) Misinterpretations of Precipitation Probability Forecasts. Allan H. Murphy, Sarah Lichtenstein, Baruch Fischhoff, and Robert L. Winkler, February 1980. (PB80 174576) Annual Date and Verification Tabulation Eastern and Central North Pacific Tropical Storms and Hurricanes 1979. Emil B. Guuther and Staff. EPHC, April 1980. (PB80 220486) NMC Model Performance in the Northeast Pacific. James E. Overland, PMEL-ERL, April 1980. (PB80 196033) 1980. (PB80 196033)

- 1980. (PB80 196033) Climate of Salt Lake City, Utah. Wilbur E. Figgins (Retired) and Alexander R. Smith. Fith Revision, July 1992. (PB92 220177) An Automatic Lightning Detection System in Northern California. James E. Rea and Chris E. Fontana, June 1980. (PB80 225592) Regression Equation for the Peak Wind Gust 6 to 12 Hours in Advance at Great Falls During Strong Downslope Wind Storms. Michael J. Oard, July 1980. (PB91 108367) A Raininess Index for the Arizona Monsoon. John H. Ten Harkel, July 1980. (PB81 106404)

- Diving Strong Downsor white Softials, Michael J. Oki (1997) 105001 (PB81) 106494)
 A Raininess Index for the Arizona Monsoon. John H. Ten Harkel, July 1980. (PB81) 106494)
 The Effects of Terrain Distribution on Summer Thunderstorm Activity at Reno, Nevada. Christopher Dean Hill, July 1980. (PB81 102501)
 An Operational Evaluation of the Scofield/Oliver Technique for Estimating Precipitation Rates from Satellite Imagery. Richard Ochoa, August 1980. (PB81 108227)
 Hydrology Practicum. Thomas Dietrich, September 1980. (PB81 134033)
 Tropical Cyclone Effects on California. Arnold Court, October 1980. (PB81 133779)
 Eastern North Pacific Tropical Cyclone Occurrences During Intraseasonal Periods. Preston W. Leftwich and Gail M. Brown, February 1981. (PB81 205494)
 Solar Radiation as a Sole Source of Energy for Photovoltaics in Las Vegas, Nevada, for July and December. Darryl Randerson, April 1981. (PB81 224503)
 A Systems Approach to Real-Time Runoff Analysis with a Deterministic Rainfall-Runoff Model. Robert J.C. Burnash and R. Larry Ferral, April 1981. (PB81 224495)
 A Comparison of Two Methods for Forecasting Thunderstorms at Luke Air Force Base, Arizona. LTC Keith R. Cooley, April 1981. (PB81 225393)
 An Objective Aid for Forecasting Afternoon Relative Humidity Along the Washington Cascade East Slopes. Robert S. Robinscon, April 1981. (PB82 20336)
 Preliminary Estimates of Wind Power Potential at the Nevada Test Site. Howard G. Booth, June 1981. (PB82 127036)
 ARAP User's Guide. Mark Mathewson, July 1981, Revised September 1981. (PB82 196783)
 Forecasting the Onset of Coastal Gales Off Washington-Oregon. John R. Zimmerman and William D. Burton, August 1981. (PB82 157051)
 A Statistical-Dynamical Model for Prediction of Tropical Cyclone Motion in the Eastern North Pacific Ocean. Preston W. Leftwich, Jr., October 1981. (PB82195298)

- Forecasting Heavy Snow at Wenatchee, Washington. James W. Holcomb, December 1981. (PB82 177783) Central San Joaquin Valley Type Maps. Thomas R. Crossan, December 1981. (PB82
- 196064) ARAP Test Results. Mark A. Mathewson, December 1981. (PB82 198103) Approximations to the Peak Surface Wind Gusts from Desert Thunderstorms. Darryl Randerson, June 1982. (PB82 253089) Climate of Phoenix, Arizona. Robert J. Schmidli, April 1969 (Revised December 1986). (DB27 140082(AS)

- Climate of Phoenix, Arizona. Robert J. Schmidli, April 1969 (Revised December 1986). (PB87 142063/AS) Annual Data and Verification Tabulation, Eastern North Pacific Tropical Storms and Hurricanes 1982. E.B. Gunther, June 1983. (PB85 106078) Stratified Maximum Temperature Relationships Between Sixteen Zone Stations in Arizona and Respective Key Stations. Ira S. Brenner, June 1983. (PB83 249904) Standard Hydrologic Exchange Format (SHEF) Version I. Phillip A. Pasteris, Vernon C. Bissel, David G. Bennett, August 1983. (PB85 106052) Quantitative and Spacial Distribution of Winter Precipitation along Utah's Wasatch Front. Lawrence B. Dunn, August 1983. (PB85 106052) 500 Millibar Sign Frequency Teleconnection Charts Winter. Lawrence B. Dunn, December 1983. (PB85 111367) Collection and Use of Lightring Strike Data in the Western US During Summer 1000

- Collection and Use of Lightning Strike Data in the Western U.S. During Summer 1983.
 Glenn Rasch and Mark Mathewson, February 1984. (PB85 110534)
 S00 Millibar Sign Frequency Teleconnection Charts Summer. Lawrence B. Dunn, March
- 1884. (PB85 111359) Annual Data and Verification Tabulation eastern North Pacific Tropical Storms and Hurricanes 1983. E.B. Gunther, March 1984, (PB85 109635) 500 Millibar Sign Frequency Teleconnection Charts Fall. Lawrence B. Dunn, May 1984. (PB85 110930)
- The Use and Interpretation of Isentropic Analyses. Jeffrey L. Anderson, October 1984. (PB85 132694)
- (FDSS 132094) Annual Data & Verification Tabulation Eastern North Pacific Tropical Storms and Hurricanes 1984. E.B. Gunther and R.L. Cross, April 1985. (PB85 1878887AS) Great Salt Lake Effect Snowfall: Some Notes and An Example. David M. Carpenter, October 1985. (PB86 119158/AS)
- Great Salt Lake Effect Snowfall: Some Notes and An Example. David M. Carpenter, October 1985. (PB86 119153/AS) Large Scale Patterns Associated with Major Freeze Episodes in the Agricultural Southwest. Ronald S. Hamilton and Glenn R. Lussky, December 1985. (PB86 144474AS) NWR Voice Synthesis Project: Phase I. Glen W. Sampson, January 1986. (PB86

- NWR Voice Synthesis Project: Phase I. Glen W. Sampson, January 1986. (PB86 145604/AS)
 The MCC An Overview and Case Study on Its Impact in the Western United States. Glenn R. Lussky, March 1986. (PB86 170651/AS)
 Annual Data and Verification Tabulation Eastern North Pacific Tropical Storms and Hurricanes 1985. E.B. Gunther and R.L. Cross, March 1986. (PB86 170641/AS)
 Radid Interpretation Guidelines. Roger G. Pappas, March 1986. (PB86 17064)/AS)
 A Mesoscale Convective Complex Type Storm over the Desert Southwest. Darryl Randerson, April 1986. (PB86 (19098/AS))
 The Effects of Eastern North Pacific Tropical Oyclones on the Southwestern United States. Walter Smith, August 1986. (PB87 106258AS)
 Preliminary Lightning Climatology Studies for Idaho. Christopher D. Hill, Carl J. Gorski, and Michael C. Conger, April 1987. (PB87 180196/AS)
 Heavy Rains and Flooding in Montana: A Case for Slantwise Convection. Glenn R. Lussky, April 1987. (PB87 185229/AS)

- Annual Data and Verification Tabulation Eastern North Pacific Tropical Storms and Hurricanes 1986. Roger L. Cross and Kenneth B. Mielke, September 1987. (PB88 110895/AS)
 An Inexpensive Solution for the Mass Distribution of Satellite Images. Glen W. Sampson and George Clark, September 1987. (PB88 114038/AS)
 Annual Data and Verification Tabulation Eastern North Pacific Tropical Storms and Weight Structure 1997. (PB88 114038/AS)
- Hurricanes 1987. Roger L. Cross and Kenneth B. Mielke, September 1988. (PB88 101935/AS)
- An Investigation of the 24 September 1986 "Cold Sector" Tornado Outbreak in Northern California. John P. Monteverdi and Scott A. Braun. October 1988. (PB89 121297/AS)
- California. John P. Monteverdi and Scott A. Braun. October 1988. (PB89 121297/AS) Preliminary Analysis of Cloud-To-Ground Lightning in the Vicinity of the Nevada Test Site. Carven Scott, November 1988. (PB89 128649/AS) Forecast Guidelines For Fire Weather and Forecasters How Nighttime Humidity Affects Wildland Fuels. David W. Goens, February 1989. (PB89 162549/AS) A Collection of Papers Related to Heavy Precipitation Forecasting. Western Region Headquarters, Scientific Services Division, August 1989. (PB89 230833/AS) The Las Vegas McCarran International Airport Microburst of August 8, 1989. Carven A. Scott, Une 1000 (PB00 240289)
- June 1990. (PB90-240268) Meteorological Factors Contributing to the Canyon Creek Fire Blowup, September 6 and 7, 1988. David W. Goens, June 1990. (PB90-245085)
- Stratus Surge Prediction Along the Central California Coast. Peter Felsch and Woodrow Whitlatch, December 1990. (PB91-129239)

- Hydrotools. Tom Egger. January 1991. (PB91-151787/AS) A Northern Utah Soaker. Mark E. Struthwolf, February 1991. (PB91-168716) Preliminary Analysis of the San Francisco Rainfall Record: 1849-1990. Jan Null, May 1991. (PB91-208439) Idaho Zone Preformat, Temperature Guidance, and Verification. Mark A. Mollner, July 1991.
- (PB91-227405/AS) Emergency Operational Meteorological Considerations During an Accidental Release of
- Hazzrdous Chemicals. Peter Mueller and Jerry Galt, August 1991. (PB91-235424) Hazzrdous Chemicals. Peter Mueller and Jerry Galt, August 1991. (PB91-235424) WeatherTools. Tom Egger, October 1991. (PB93-184950) Creating MOS Equations for RAWS Stations Using Digital Model Data. Dennis D. Gettman, December 1991. (PB92-131473/AS)
- Forecasting Heavy Snow Events in Missoula, Montana. Mike Richmond, May 1992. (PB92-196104)
- NWS Winter Weather Workshop in Portland, Oregon. Various Authors, December 1992. (PB93-146785)
- A Case Study of the Operational Usefulness of the Sharp Workstat on in Forecasting a Mesocyclone-Induced Cold Sector Tornado Event in California. John P. Monteverdi, March 1993. (PB93-178697)
- Climate of Pendleton, Oregon. Claudia Bell, August 1993. (PB93-227536)
- Utilization of the Bulk Richardson Number, Helicity and Sounding Modification in the Assessment of the Severe Convective Storms of 3 August 1992. Eric C. Evenson, September 1993. (PB94-131943)
- Convective and Rotational Parameters Associated with Three Tornado Episodes in Northern and Central California. John P. Monteverdi and John Quadros, September 1993. (PB94-131943)
- Climate of San Luis Obispo, California. Gary Ryan, February 1994. (PB94-162062) Climate of Wenatchee, Washington. Michael W. McFarland, Roger G. Buckman, and Gregory E. Matzen, March 1994. PB94-164308

NOAA SCIENTIFIC AND TECHNICAL PUBLICATIONS

The National Oceanic and Atmospheric Administration was established as part of the Department of Commerce on October 3, 1970. The mission responsibilities of NOAA are to assess the socioeconomic impact of natural and technological changes in the environment and to monitor and predict the state of the solid Earth, the oceans and their living resources, the atmosphere, and the space environment of the Earth.

The major components of NOAA regularly produce various types of scientific and technical information in the following kinds of publications.

PROFESSIONAL PAPERS--Important definitive research results, major techniques, and special investigations.

CONTRACT AND GRANT REPORTS-Reports prepared by contractors or grantees under NOAA sponsorship.

ATLAS--Presentation of analyzed data generally in the form of maps showing distribution of rainfall, chemical and physical conditions of oceans and atmosphere, distribution of fishes and marine mammals, ionospheric conditions, etc. TECHNICAL SERVICE PUBLICATIONS--Reports containing data, observations, instructions, etc. A partial listing includes data serials; prediction and outlook periodicals; technical manuals, training papers, planning reports, and information serials; and miscellaneous technical publications.

TECHNICAL REPORTS--Journal quality with extensive details, mathematical developments, or data listings.

TECHNICAL MEMORANDUMS--Reports of preliminary, partial, or negative research or technology results, interim instructions, and the like.



Information on availability of NOAA publications can be obtained from:

NATIONAL TECHNICAL INFORMATION SERVICE

U. S. DEPARTMENT OF COMMERCE

5285 PORT ROYAL ROAD

SPRINGFIELD, VA 22161