

NOAA Technical Memorandum NWS WR-231

WASHINGTON STATE TORNADOES

Treste' Huse Colorado Basin River Forecast Center Salt Lake City, Utah

July 1995

U.S. DEPARTMENT OF COMMERCE

National Oceanic and Atmospheric Administration National Weather Service



NOAA TECHNICAL MEMORANDA National Weather Service, Western Region Subseries

The National Weather Service (NWS) Western Region (WR) Subseries provides an informal medium for the documentation and quick dissemination of results not appropriate, or not yet menum for the documentation and quick dissemination of results has applying the of hor yet ready, for formal publication. The series is used to report on work in progress, to describe technical procedures and practices, or to relate progress to a limited audience. These Technical Memoranda will report on investigations devoted primarily to regional and local problems of interest mainly to personnel, and hence will not be widely distributed.

Papers 1 to 25 are in the former series, ESSA Technical Memoranda, Western Region Technical Memoranda (WRTM); papers 24 to 59 are in the former series, ESSA Technical Memoranda, Weather Bureau Technical Memoranda (WBTM). Beginning with 60, the papers are part of the series, NOAA Technical Memoranda NWS. Our-of-print memoranda are not listed.

Papers 2 to 22, except for 5 (revised edition), are available from the National Weather Service Western Region, Scientific Services Division, P.O. Box 11188, Federal Building, 125 South State Street, Salt Lake City, Utah 84147. Paper 5 (revised edition), and all others beginning with 25 are available from the National Technical Information Service, U.S. Department of Commerce, Sills Building, 5285 Port Royal Road, Springfield, Virginia 22161. Prices vary for all paper copies; microfiche are \$3.50. Order by accession number shown in parentheses at end of each entry.

ESSA Technical Memoranda (WRTM)

- Climatological Precipitation Probabilities. Compiled by Lucianne Miller, December 1965. Climatological Precipitation Probabilities. Compiled by Lucianne Miller, December 1965. Western Region Pre- and Post-FP-3 Program, December 1, 1965, to February 20, 1966. Edward D. Diemer, March 1966. Station Descriptions of Local Effects on Synoptic Weather Patterns. Philip Williams, Jr., April 1966 (Revised November 1967, October 1969). (PB-17800) Interpreting the RAREP. Herbert P. Benner, May 1966 (Revised January 1967). Some Electrical Processes in the Atmosphere. J. Latham, June 1966. A Digitalized Summary of Radar Echces within 100 Miles of Sacramento, California. J. A. Youngberg and L. B. Overaas, December 1966. An Objective Aid for Forecasting the End of East Winds in the Columbia Gorge, July through October. D. John Coparanis, April 1967. Derivation of Radar Horizons in Mountainous Terrain. Roger G. Pappas, April 1967. 3

ESSA Technical Memoranda, Weather Bureau Technical Memoranda (WBTM)

- Verification of Operation Probability of Precipitation Forecasts, April 1966-March 1967.
 W. Dickey, October 1967. (PB-176240)
 A Study of Winds in the Lake Mead Recreation Area. R. P. Augulis, January 1968. (PB-1797020)
- 177830)
- 177330) Weather Extremes. R. J. Schmidli, April 1968 (Revised March 1986). (PB86 177672/AS). (Revised October 1991 PB92-115062/AS) Small-Scale Analysis and Prediction. Philip Williams, Jr., May 1968. (PB178425) Numerical Weather Prediction and Synoptic Meteorology. CPT Thomas D. Murphy, USAF, May 1968. (AD 673365) Precipitation Detection Probabilities by Salt Lake ARTC Radars. Robert K. Belesky, July
- Temperature Trends in Sacramento-Another Heat Island. Anthony D. Lentini, February
- Temperature Trends in Sacramento-Another Heat Island. Anthony D. Lentini, February 1969. (PB 183055) Disposal of Logging Residues Without Damage to Air Quality. Owen P. Cramer, March 1969. (PB 183057)
- Upper-Air Lows Over Northwestern United States. ALL Jacobson, April 1969. PB 184296) The Man-Machine Mix in Applied Weather Forecasting in the 1970s. L.W. Snellman, August
- Forecasting Maximum Temperatures at Helena, Montana. David E. Olsen, October 1969. (PB 188762) 1969. (PB 185068)
- Estimated Return Periods for Short-Duration Precipitation in Arizona. Paul C. Kangieser,
- Estimated Return Periods for Subriduation Programmed in Fischer 1969. (PB 187763) October 1969. (PB 187763) Applications of the Net Radiometer to Short-Range Fog and Stratus Forecasting at Eugene, Oregon. L. Yee and E. Bates, December 1969. (PB 190476) Statistical Analysis as a Flood Routing Tool. Robert J.C. Burnash, December 1969. (PB 188744)
- Tsunami. Richard P. Augulis, February 1970. (PB 190157) Predicting Precipitation Type. Robert J.C. Burnash and Floyd E. Hug, March 1970. (PB 10/062)
- 190962) Statistical Report on Aeroallergens (Pollens and Molds) Fort Huachuca, Arizona, 1969. Wayne S. Johnson, April 1970. (PB 191743) Western Region Sea State and Surf Forecaster's Manual. Gordon C. Shields and Gerald B. Burdwell, July 1970. (PB 193102) Sacramento Weather Radar Climatology. R.G. Pappas and C. M. Veliquette, July 1970. (PB
- 193347)

- 193347) A Refinement of the Vorticity Field to Delineate Areas of Significant Precipitation. Barry B. Aronovitch, August 1970. Application of the SSARR Model to a Basin without Discharge Record. Vail Schermerhorn and Donal W. Kuehl, August 1970. (PB 194394) Areal Coverage of Precipitation in Northwestern Utah. Philip Williams, Jr., and Werner J. Heck, September 1970. (PB 194389) Preliminary Report on Agricultural Field Burning vs. Atmospheric Visibility in the Willamette Valley of Oregon. Earl M. Bates and David O. Chilcote, September 1970. (PB 194710)
- 194710)
 Air Pollution by Jet Aircraft at Seattle-Tacoma Airport. Wallace R. Donaldson, October 1970. (COM 71 00017)
 Application of PE Model Forecast Parameters to Local-Area Forecasting. Leonard W. Sneliman, October 1970. (COM 71 00016)
 An Aid for Forecasting the Minimum Temperature at Medford, Oregon, Arthur W. Fritz, October 1970. (COM 71 00120)
 700-mb Warm Air Advection as a Forecasting Tool for Montana and Northern Idaho. Norris E. Woerner, February 1971. (COM 71 00349)
 Wind and Weather Regimes at Great Falls, Montana. Warren B. Price, March 1971. Climate of Sacramento, California. Tony Martini, April 1990. (Fifth Revision) (PB89 207781/AS)

- 207781/AS) A Preliminary Report on Correlation of ARTCC Radar Echoes and Precipitation. Wilbur K. Hall, June 1971. (COM 71 00829)
- National Weather Service Support to Soaring Activities. Ellis Burton, August 1971. (COM
- Valoual vesteel burne cupper to boaring Activities. Inthe burnes, August 1972. (COM Western Region Synoptic Analysis-Problems and Methods. Philip Williams, Jr., February 1972. (COM 72 10433)
- Thunderstorms and Hail Days Probabilities in Nevada. Clarence M. Sakamoto, April 1972. (COM 72 10554)

- A Study of the Low Level Jet Stream of the San Joaquin Valley. Ronald A. Willis and Philip Williams, Jr., May 1972. (COM 72 10707) Monthly Climatological Charts of the Behavior of Fog and Low Stratus at Los Angeles International Airport. Donald M. Gales, July 1972. (COM 72 11140) A Study of Radar Echo Distribution in Arizona During July and August. John E. Hales, Jr., July 1972. (COM 72 11136)
- July 1972. (COM 72 11136) Forecasting Precipitation at Bakersfield, California, Using Pressure Gradient Vectors. Earl

- Forecasting Precipitation at Bakersfield, California, Using Pressure Gradient Vectors. Earl T. Riddiough, July 1972. (COM 72 11146) Climate of Stockton, California. Robert C. Nelson, July 1972. (COM 72 10920) Estimation of Number of Days Above or Below Selected Temperatures. Clarence M. Sakamoto, October 1972. (COM 72 10021) An Aid for Forecasting Summer Maximum Temperatures at Seattle, Washington. Edgar G. Johnson, November 1972. (COM 73 10150) Flash Flood Forecasting and Warning Program in the Western Region. Philip Williams, Jr., Chester L. Glenn, and Roland L. Raetz, December 1972, (Revised March 1978). (COM 73 10251) 10251)
- 10251) A comparison of Manual and Semiautomatic Methods of Digitizing Analog Wind Records. Glenn E. Rasch, March 1973. (COM 73 10669) Conditional Probabilities for Sequences of Wet Days at Phoenix, Arizona. Paul C. Kangieser, June 1973. (COM 73 11264) A Refinement of the Use of K.Values in Forecasting Thunderstorms in Washington and Oregon. Robert Y.G. Lee, June 1973. (COM 73 11276) Objective Forecast Precipitation Over the Western Region of the United States. Julia N. Pargie and Larry P. Kierulff, September 1973. (COM 73 11946/3AS) Arizona "Eddy Tornadoes. Robert S. Ingram, October 1973. (COM 73 10465) Smoke Management in the Willsmette Valley. Earl M. Bates, May 1974. (COM 74 11277/AS)

- 11277/AS)
- 11277/AS) An Operational Evaluation of 500-mb Type Regression Equations. Alexander E. MacDonald, June 1974. (COM 74 11407/AS) Conditional Probability of Visibility Less than One-Half Mile in Radiation Fog at Fresno, California. John D. Thomas, August 1974. (COM 74 11555/AS) Climate of Flagstaff, Arizona. Paul W. Sorenson, and updated by Reginald W. Preston, January 1987. (PB87 143160/AS) Map type Precipitation Probabilities for the Western Region. Glenn E. Rasch and Alexander E. MacDonald, February 1975. (COM 75 10428/AS) Eastern Parcific Cut-Off Law of April 21-28, 1974. William J. Alder and George R. Miller, January 1976. (PB 250 711/AS)

- on a Significant Precipitation Episode in Western United States. Ira S. Brenner, April (COM 75 10719/AS) Study

- 1976. (COM 75 10719/AS) A Study of Flash Flood Susceptibility-A Basin in Southern Arizona. Gerald Williams, August 1975. (COM 75 11360/AS) A Set of Rules for Forecasting Temperatures in Napa and Sonoma Counties. Wesley L. Tuft, October 1975. (PB 246 902/AS) Application of the National Weather Service Flash-Flood Program in the Western Region. Gerald Williams, January 1976. (PB 253 053/AS) Objective dids for Forecasting Minimum Temperatures at Reno, Nevada, During the Summer Months. Christopher D. Hill, January 1976. (PB 252 866/AS) Forecasting the Mono Wind. Charles P. Ruscha, Jr., February 1976. (PB 254 650) Use of MOS Forecast Parameters in Temperature Forecasting. John C. Plankinton, Jr., Mar Types as Aids in Using MOS PoPa in Western United States. Irs S. Brenner, August

- March 1976. (FD 259 594) Map Types as Aids in Using MOS PoPs in Western United States. Ira S. Brenner, August 1976. (FB 259 594) Other Kinds of Wind Shear. Christopher D. Hill, August 1976. (FB 260 437/AS)
- Other Kinds of Wind Shear. Christopher D. Hill, August 1976. (PS 2601 437/A53) Forecasting North Winds in the Upper Sacramento Valley and Adjoining Forests. Christopher E. Fontana. September 1976. (PB 273 677/AS) Cool Inflow as a Weakening Influence on Eastern Pacific Tropical Cyclones. William J. Denney, November 1976. (PB 264 655/AS) The MAN/MOS Program. Alexander E. MacDonald, February 1977. (PB 265 941/AS) Winter Season Minimum Temperature Formula for Bakersfield, California, Using Multiple Regression. Michael J. Oard, February 1977. (PB 273 676/AS) A Study of Wind Gusts on Lake Mead. Bradley Colman, April 1977. (PE 268 847) The Relative Frequency of Cumulonimbus Clouds at the Nevada Test Site as a Function of K-Value. R.F. Quiring, April 1977. (PB 272 831) Moisture Distribution Modification by Upward Vertical Motion. Ira S. Brenner, April 1977.

- (PB 268 740) Relative Frequency of Occurrence of Warm Season Echo Activity as a Function of Stability Indices Computed from the Yucca Flat, Nevada, Rawinsonde. Darryl Randerson, June 1977. (PB 271 290/AS)
- (PB 271 290/AS) Climatological Prediction of Cumulonimbus Clouds in the Vicinity of the Yucca Flat Weather Station. R.F. Quiring, June 1977. (PB 271 704/AS) A Method for Transforming Temperature Distribution to Normality. Morris S. Webb, Jr., June 1977. (PB 271 742/AS)

- June 1977. (PB 271 742/AS) Statistical Guidance for Prediction of Eastern North Pacific Tropical Cyclone Motion Part I. Charles J. Neumann and Preston W. Leftwich, August 1977. (PB 272 661) Statistical Guidance on the Prediction of Eastern North Pacific Tropical Cyclone Motion -Part II. Preston W. Leftwich and Charles J. Neumann, August 1977. (PB 273 155/AS) Climate of San Francisco. E. Jan Null, February 1978. Revised by George T. Pericht, April 1988. (PB88 208624/AS) Development of a Probability Equation for Winter-Type Precipitation Patterns in Great Falls, Montana. Kenneth B. Mielke, February 1978. (PB 281 387/AS) Hand Calculator Program to Compute Parcel Thermal Dynamics. Dan Gudgel, April 1978.

- Hand Calculator Program to compute Autor. (PB 283 866/AS) (PB 283 080/AS) Fire whirls. David W. Goens, May 1978. (PB 283 866/AS) Flash-Flood Procedure. Ralph C. Hatch and Gerald Williams, May 1978. (PB 286 014/AS) Automated Fire-Weather Forecasts. Mark A. Mollner and David E. Olsen, September 1978.
- Automated Fire-Weather Forecasts. Mark A. Mollner and David E. Olsen, September 1978. (PB 289 916/AS) Estimates of the Effects of Terrain Blocking on the Los Angeles WSR-74C Weather Radar. R.G. Pappas, R.Y. Lee, B.W. Finke, October 1978. (PB 289767/AS) Spectral Techniques in Ocean Wave Forecasting. John A. Jannuzzi, October 1978. (PB291317/AS)
- Solar Radiation. John A. Jannuzzi, November 1978. (PB291195/AS)

- Solar Radiation. John A. Jannuzzi, November 1978. (PE291195/AS) Application of a Spectrum Analyzer in Forecasting Ocean Swell in Southern California Coastal Waters. Lawrence P. Klerulft, January 1979. (PE292716/AS) Basic Hydrologic Principles. Thomas L. Dietrich, January 1979. (PE292247/AS) LFM 24+Hour Prediction of Eastern Pacific Cyclones Refined by Satellite Images. John R. Zimmerman and Charles P. Ruscha, Jr., January 1979. (PE294324/AS) A Simple Analysis/Diagnosis System for Real Time Evaluation of Vertical Motion. Scott Heflick and James R. Fors, February 1979. (PE294216/AS) Aids for Forecasting Minimum Temperature in the Wenatchee Frost District. Robert S. Robinson, April 1979. (PE29339/AS) Influence of Cloudiness on Summertime Temperatures in the Eastern Washington Fire Weather district. James Holcomb, April 1979. (PE29674/AS) Comparison of LFM and MFM Precipitation Guidance for Nevada During Doreen. Christopher Hill, April 1979. (PE2983613/AS)

Û

NOAA Technical Memorandum NWS WR-231

WASHINGTON STATE TORNADOES

Treste' Huse Colorado Basin River Forecast Center Salt Lake City, Utah

July 1995

UNITED STATES DEPARTMENT OF COMMERCE Ronald H. Brown, Secretary National Oceanic and Atmospheric Administration D. James Baker, Under Secretary and Administrator National Weather Service Elbert W. Friday, Jr., Assistant Administrator for Weather Services



This publication has been reviewed and is approved for publication by Scientific Services Division, Western Region

Jhhh

Delain A. Edman Scientific Services Division Salt Lake City, Utah

TABLE OF CONTENTS

÷

Ι.	INTRODUCTION 1
Ш.	DATA COLLECTION 1
111.	CLIMATOLOGY 1
IV.	SYNOPTIC AND LOCAL ENVIRONMENTS 2
V.	EVENT DESCRIPTION 4
VI.	DISCUSSION 4
VII.	ACKNOWLEDGEMENTS 5
VIII.	REFERENCES 5

TABLES AND FIGURES

r,

-.

Figure 1.	Geographical location and corresponding F-scale intensity of Washington tornadoes from 1950 through 1994.			
Figure 2.	Population density map of Washington based on 1990 U.S. Bureau of Census data.			
Figure 3.	Monthly and Hourly distributions of tornadoes in (a) the United States, (b) western Washington and (c) eastern Washington			
Figure 4.	(a) (b)	Western Washington 850 mb Composite Eastern Washington 850 mb Composite		
Figure 5.	(a) (b)	Western Washington 500 mb Composite Eastern Washington 500 mb Composite		
Figure 6.	(a) (b)	SW flow cases for western Washington at 850 mb SW flow cases for western Washington at 500 mb		
Figure 7.	(a) (b)	NW flow cases for western Washington at 850 mb NW flow cases for western Washington at 500 mb		
Figure 8.	(a) (b)	SW flow cases for eastern Washington at 850 mb SW flow cases for eastern Washington at 500 mb		
Figure 9.	(a) (b)	NW flow cases for eastern Washington at 850 mb NW flow cases for eastern Washington at 500 mb		
Figure 10.	-	vses of a) 850 mb and b) 500 mb heights for 1200 UTC 5 1972		
Figure 11.	Hodo	graph for Spokane 1200 UTC 5 April 1972 sounding		

Table 1.Average values of CAPE and S.R. Helicity for tornadic events
using soundings from Quillayute and Spokane

WASHINGTON STATE TORNADOES

Treste' Huse Colorado Basin River Forecast Center Salt Lake City, UT (Former affiliation NWSFO Seattle)

ABSTRACT

This study researches past tornado occurrences in Washington to identify elements conducive for tornado potential. Tornadic events were examined through composites and case study analysis. The composites revealed either southwest or northwest mid- and upper-level flow regimes. Environments more favorable for classic supercell development were implied from the southwest flow cases.

I. INTRODUCTION

Tornadoes in Washington, while rare compared to those east of the Rockies, should not be dismissed as insignificant features of the climate of the state. Tornadoes have been reported in 28 of the last 44 years in Washington; with a total of 53 tornados from 1950 through 1994. Although Washington tornadoes are generally weak, several F3 tornadoes have occurred. The most destructive tornado killed 6 people and injured 300 others in April 1972. This study examines the climatology of tornadoes in Washington and the meteorological conditions associated with these notable events.

II. DATA COLLECTION

NSSFC tornado statistics for Washington provided precise times and locations, path dimensions and severity information for tornadic events. To further assess damage, *Storm Data* were retrieved. Additionally, the surface and upper-level analyses with nearest synoptic times to the events were obtained from a microfilm library at the University of Washington. To gain more

insight into similarities of the analyses and their deviation from climatology, composite height fields were produced for 850, 700, 500 and 200 mb; composite sea-level pressure analyses were also generated. All composites were created using the NMC Grid Point Data Set on Compact Disc. These data are available twice daily over a hemispheric domain with a grid resolution of 380 km. Soundings from Quillayute for western Washington and Spokane for eastern Washington were taken from the Radiosonde Data of North America Compact Disc (NOAA/FSL 1993). These soundings were examined and, in certain cases, modified using the interactive Skew T/Hodograph Analysis and Research Program (SHARP). These modifications will be described later.

III. CLIMATOLOGY

Figure 1 illustrates the geographic locations and intensities of tornadoes in Washington. Tornadoes appear to be concentrated in or near Seattle and Spokane (Fig. 2). This apportionment reflects a bias to population centers. The tornadoes in western Washington were mainly observed in the central and southern interior. Thirty four (64 percent) of the tornadoes were reported east of the Cascades even though the population density is less than in western Washington. The frequency of tornadoes was less along the east slopes of the Cascades, increasing significantly over the eastern third of the state. A number of these reports were from areas with population densities of fewer than four persons per square mile. No tornadoes were reported over the Cascade Mountains.

There was a preponderance of F0 and F1 tornadoes in the state with 73 percent of the events in these categories. A higher proportion of Washington tornadoes were classified as F0 and F1 than the 63 percent for the United States as a whole (Fujita 1987). Eleven F2 and three F3 tornadoes have been reported in Washington since The average path length for 1950. tornadoes in Washington was determined to be 1.8 miles. To compare these statistics to path lengths in other areas of the Unites States. Fujita's averages for path lengths by F-scale were obtained. These data show that tornado mean path lengths increase with higher F-scale classifications. A population weighted mean was then constructed using the Fscale classification of the 53 Washington tornadoes and these Fujita averages. This technique revealed an expected mean path length of 3.0 miles, considerably greater than observed. This result indicates that the typical path length for tornadoes in Washington is shorter than for the nation as a whole.

Histograms were constructed showing the frequency of tornadoes by month and hour (Fig. 3). Histograms for eastern Washington were generally similar to national statistics with April and May the most common months for tornado activity. A secondary maximum was indicated July through August. While tornadoes were more evenly distributed by month west of the Cascades, the fewest instances occurred in mid-summer and late winter. This suggests a greater likelihood of tornadoes during transitional seasons in western Washington. However, this is clearly speculation due to the small sample size. The greatest diurnal frequency west of the Cascades is indicated during the late morning and afternoon. The connection between tornado events and solar heating is more evident east of the Cascades where tornadoes tend to peak in the afternoon and early evening.

IV. SYNOPTIC AND LOCAL ENVIRONMENTS

To define the range of synoptic conditions associated with tornadoes in Washington, constant pressure fields were analyzed and composited. The protective barrier afforded by the Cascade Mountains bars the inland penetration of cool maritime air from the Pacific, which promotes stronger diurnal heating as well as a drier climate east of the Cascades. Because of the differing climates, eastern and western Washington were composited separately.

The western Washington composite at 850 mb (Fig. 4a) shows a trough positioned at 130°W with consolidated westerly flow over the state. In contrast, the 850 mb height field for the eastern Washington events (Fig. 4b) indicates a flat ridge extending from the central Pacific into Washington. Both composites at 500 mb (Fig. 5) are characterized by a trough from the southern British Columbia coast to the northern California coast. However, a deeper trough with stronger southwest flow into Washington prevails on the western Washington composites. The standard deviations of the composite values were quite large (not shown) over the eastern Pacific. Due to these significant height variations among the analyses, it was deemed necessary to further classify the tornado events.

A survey of the collected fields revealed that there were two broadly similar synoptic patterns for both eastern and western Washington. The events generally exhibited either northwest or southwest flow at 500 mb. Therefore, the events were subdivided into four classifications based on location and synoptic flow This process, to some degree, regimes. reduced the standard deviation in the near field in the composites. It should be noted that seven cases did not fit in either synoptic flow pattern and these cases were not composited.

The southwest flow cases for western Washington (Fig. 6) are characterized by a deep long-wave trough over the eastern Pacific with strong southwest flow over the state. Difluence is indicated downstream from the upper-level trough axis inferring upward motion over Washington. The synoptic-scale northwest flow regime in western Washington (Fig. 7) consists of a trough extending from British Columbia southwest across Vancouver Island. Northwest winds over Washington are implied to be quite strong at upper levels with significant difluence over Washington at 500 mb.

The southwest flow cases for eastern Washington (Fig. 8) delineate a negatively tilted trough off the Washington coast, typifying short-wave troughs moving northeast. Very strong flow is evident on the backside of the trough. Analogous to the western Washington southwest flow cases, the long-wave pattern places the state in the preferred region of ascent downstream from the trough axis. Finally, for the northwest flow scenarios in eastern Washington (Fig. 9) an 850 mb positively tilted ridge extends from the eastern Pacific northeast across Washington. At 500 mb, a ridge is positioned over the eastern Pacific with a trough centered over Idaho. Northerlies at 850 mb and northwest flow at 500 mb are indicated over the state.

To explore local environments conducive to tornado formation, soundings from Quillayute (for western Washington) and Spokane (for eastern Washington) were examined using the SHARP software package. Due to diurnal heating variations, the soundings were not always indicative of the tornadic environment. This was especially true for western Washington cases given the location of Quillayute on the extreme northwest coast. The afternoon temperatures at Quillayute were usually considerably lower than those across the interior. To provide more representative surface conditions, high temperatures were gathered from a proximity observing site for the day of the tornadic event. If \mathbf{the} maximum temperature at the observing site deviated significantly from the initial sounding, the surface temperature of the sounding was modified. Of the 33 soundings available, 22 were modified.

To determine the available buoyant energy in the actual and modified soundings, Convective Available Potential Energy (CAPE) was calculated. CAPE represents the vertically integrated positive area of a parcel rising adiabatically within the ambient environment. Larger temperature differences between the warmer parcel and cooler environment will lead to greater CAPE, updraft strength, and essentially more vigorous convection. Storm-relative helicity was also examined to estimate rotation potential. Storm-relative helicity measures the thunderstorm's potential to

develop a rotating updraft as it moves through a vertically sheared environment. It combines the effects of storm motion. inflow strength, and horizontal vorticity. The results are presented in Table 1. The majority of soundings had CAPE values that are considered marginally unstable (between 0 and 1000 J Kg⁻¹). However, several of the modified soundings on both sides of the Cascades were moderately unstable (between 1000 and 2000 J Kg⁻¹). All soundings, with the exception of one, exhibited storm-relative helicity values less than 150 $m^2 s^2$; which is the approximate threshold indicated for supercell development. Values of CAPE and stormrelative helicity were similar for eastern and western Washington. Interestingly, CAPE and helicity were considerably higher for the southwest flow cases than the northwest cases. This suggests the southwest flow cases provide a more environment for supercell favorable development. However, the classifications consisted of relatively small numbers of soundings, with only four available for the western Washington southwest flow scenario.

V. EVENT DESCRIPTION

The most severe tornadic outbreak occurred during the afternoon of 5 April 1972, with four tornadoes observed in Washington. Two were classified as F3 tornadoes. The most damaging tornado created a nine mile path of destruction in and near Vancouver in extreme southwest Washington. The tornado killed 6 people and injured 300 others, the majority in a shopping center. The other three tornadoes occurred in eastern Washington and touched down in more remote areas demolishing numerous farm buildings.

The general synoptic situation for the event depicted a deep trough located over

the eastern Pacific. This placed Washington under a strong, consolidated southwest flow in the mid- and uppertroposphere (Fig. 10), consistent with the southwest flow composites presented earlier.

According to reports, a rapidly moving squall line advanced across Washington during the late morning and afternoon hours of 5 April; the tornadoes were attendant to this squall line. The first tornado initially touched down in north Portland, Oregon, lifted to a funnel cloud as it crossed the Columbia River and then touched down again near Vancouver at 1245 PM PST (2045 UTC). The last tornado was reported in northeast Washington at 6 PM PST (0200 UTC 6 April).

The only radiosonde information available for the event was the 1200 UTC sounding at Spokane. Interestingly, this sounding exhibited an impressive storm-relative helicity value of 281 m² s⁻²; by far the highest of any of the events studied. The hodograph displayed a strong veering wind profile with large wind shear through the lower levels (Fig. 11). The modified CAPE value was 35 J Kg⁻¹. This case study is consistent with the previous suggestion that southwest flow cases are closer to a classic supercell environment.

VI. DISCUSSION

This study examined tornado events in Washington for the past 44 years. The data indicated tornadoes in Washington were less damaging than in other areas of the United States, but Washington's sample size is quite limited. Washington tornadoes also exhibited a shorter path length than those of corresponding intensity across the United States. There was a greater occurrence of tornadoes in eastern Washington than western Washington. The seasonal and diurnal distributions of tornadoes in eastern Washington were similar to national averages, with occurrences highest in the spring. The diurnal maximum was found to be late in the day. Meanwhile, western Washington events tended to peak during transitional seasons and around midday.

Composite maps for selected constant pressure surfaces display either southwest or northwest mid- and upper-level flow regimes for tornadic events. A trough pattern favorable for large-scale ascent was implied from the southwest flow cases. Furthermore, southwest flow is typically characterized by warm advection (although not indicated by the geostrophic shear in the composite) and would generally be considered more prone to thunderstorm Higher CAPE and helicity outbreaks. values for the southwest flow cases also suggest more favorable local environments for classic tornadic activity. The northwest flow cases are indicative of a colder environment and cold advection; conditions not considered conducive for severe storm or supercell development. However, this type of environment is sometimes capable of producing weak tornadoes known as cold-air funnels (Bluestein 1993). In addition, local terrain-induced circulations may contribute to tornado formation in the Puget Sound region. West-northwest flow is the necessary direction for the Puget Sound Convergence Zone to develop (Whitney 1993). This would be a favored region for the generation of antecedent vorticity due to the interaction of low-level wind with topography (Colman 1992).

Several synoptic patterns conducive to tornado formation have been detailed in this study. The patterns found are not

dramatically different from typical weather patterns over Washington. However, it is important to note that there are broadly different patterns that can be conducive to tornadic formation across this region. An awareness of the local climate in combination with synoptic pattern recognition can help alert the forecaster to the potential for tornadic activity and lead to a better understanding of these events. With an approaching vigorous short wave strong instability; operational and forecasters need to be aware of the potential for tornadic events. With the installation of WSR-88D radars in the Pacific Northwest, examination of reflectivity and velocity data will provide more insight into mesoscale boundaries more directly associated with tornado These will be extremely occurrences. useful data for better understanding and predicting tornadoes in Washington.

VII. ACKNOWLEDGMENTS

Special appreciation is extended to Brad Colman, Science and Operations Officer at NWSFO Seattle for his support and professional guidance during the development of this memorandum. I would also like to thank Area Manager Chris Hill and the Western Region Scientific Services Division for their helpful comments and reviews, and Dennis Cain, forecaster at NWSO Hanford, for his aid in preparing graphical displays.

VIII. REFERENCES

Bluestein, H.B., 1993: Synoptic Dynamic Meteorology in Midlatitudes Volume II: Observations and Theory of Weather Systems., Oxford University Press., 594 pp. Colman, B.R., 1992: The Operational Considerations of Non-Supercell Tornadoes Particularly Those Where Local Terrain Is of Foremost Importance to Tornadogenesis. Proceedings Meteorology, Canadian AES/CMOS, Whistler, B.C., Canada, September.

Forecast Systems Laboratory and National Climatic Data Center, 1993: *Radiosonde Data of North America 1946-1992*. CDROM, Version 1.0.

Fujita, T.T., 1987: U.S. Tornadoes Part I: 70-year Statistics., Satellite and Mesometeorology Research Project, Paper 218., 121 pp.

Storm Data, Vol. 1, No. 1 through Vol. 36, No. 4.

Washington., World Book Encyclopedia. 1994. Vol 21.

Whitney, W.M., R.L. Doherty, and B.R. Colman, 1993: A Methodology for Predicting the Puget Sound Convergence Zone and Its Associated Weather. *Wea. Forecasting.*, 8, 214-222.





Fig 1. Geographical location and corresponding F-scale intensity of Washington tornadoes from 1950 through 1994.



Persons per sq. mi. More than 100 25 to 100 10 to 25 Less than 10

Fig. 2. Population density map of Washington based on 1990 U.S. Bureau of Census data (World Book Encyclopedia, 1994).






Fig 3. Monthly and Hourly distributions of tornadoes in (a) the United States, (b) western Washington and (c) eastern Washington.





___ _




Fig 6. SW flow cases for western Washington at a) 850 mb and b) 500 mb.




Fig 7. NW flow cases for western Washington at a) 850 mb and b) 500 mb.




Fig 8. SW flow cases for eastern Washington at a) 850 mb and b) 500 mb.




Fig 9. NW flow cases for eastern Washington at a) 850 mb and b) 500 mb.




Fig. 10. Analyses of a) 850 mb and b) 500 mb heights for 1200 UTC 5 April 1972.




Fig. 11. Hodograph for Spokane 1200 UTC 5 April 1972 sounding (heights are labeled in km).

	Initial CAPE	Modified CAPE	SR Helicity
UIL - all cases	163	780	39
UIL - SW flow cases	104	946	53
UIL - NW flow cases	192	696	32
GEG - all cases	458	790	35
GEG - SW flow cases	616	927	62
GEG - NW flow cases	276	614	7

Table 1. Average values of CAPE and S.R. Helicity for tornadic events using soundings from Quillayute (for western Washington) and Spokane (for eastern Washington).

- The Usefulness of Data from Mountaintop Fire Lookout Stations in Determining Atmospheric Stability. Jonathan W. Corey, April 1979. (PB298899/AS) The Depth of the Marine Layer at San Diego as Related to Subsequent Cool Season Precipitation Episodes in Arizona Ira S. Brenner, May 1979. (PB298817/AS) Arizona Cool Season Climatological Surface Wind and Pressure Gradient Study. Ira S. Brenner, May 1979. (PB298800/AS) 142 143
- 146
- December 1979. (PB80 160344) Occurrence and Distribution of Flash Floods in the Western Region. Thomas L. Dietrich, December 1979. (PB80 160344) 147
- 149 150
- December 1979. (FBS0 160344) Misinterpretations of Precipitation Probability Forecasts. Allan H. Murphy, Sarah Lichtenstein, Baruch Fischhoff, and Robert L. Winkler, February 1980. (PB80 174576) Annual Data and Verification Tabulation Eastern and Central North Pacific Tropical Storms and Hurricanes 1979. Emil B. Gunther and Staff, EPHC, April 1980. (PB80 220486) NMC Model Performance in the Northeast Pacific. James E. Overland, PMEL-ERL, April 2020. 151
- 152
- NMC Model Performance in the Northeast Pacific. James E. Overland, PMEL-EKL, April 1980. (PB80 196033)
 Climate of Salt Lake City, Utah. Wilbur E. Figgins (Retired) and Alexander R. Smith. Fifth Revision, July 1992. (PB92 220177)
 An Automatic Lightning Detection System in Northern California. James E. Rea and Chris E. Fontana, June 1980. (PB80 225592)
 Regression Equation for the Peak Wind Gust 6 to 12 Hours in Advance at Great Falls During Strong Downslope Wind Storms. Michael J. Oard, July 1980. (PB91 108367)
 A Rainliness Index for the Arizona Monsoon. John H. Ten Harkel, July 1980. (PB81 106264) 153
- 154 155
- 156
- The Effects of Terrain Distribution on Summer Thunderstorm Activity at Reno, Nevada. Christopher Dean Hill, July 1980. (PB81 102501) An Operational Evaluation of the Scofield/Oliver Technique for Estimating Precipitation 157
- 158
- 159
- 160
- 161 162
- 163
- 164
- 165
- Christopher Dean Hill, July 1980. (PB81 102501) An Operational Evaluation of the Scofield/Oliver Technique for Estimating Precipitation Rates from Satellite Imagery. Richard Ochoa, August 1980. (PB81 108227) Hydrology Practicum. Thomas Dietrich, September 1980. (PB81 134033) Tropical Cyclone Effects on California. Arnold Court, October 1980. (PB81 133779) Eastern North Pacific Tropical Cyclone Occurrences During Intrascessonal Periods. Preston W. Leftwich and Gail M. Brown, February 1981. (PB81 2063494) Solar Radiation as a Sole Source of Energy for Photovoltaics in Las Vegas, Nevada, for July and December. Darryl Randerson, April 1981. (PB81 2063494) Solar Radiation as a Sole Source of Energy for Photovoltaics in Las Vegas, Nevada, for July and December. Darryl Randerson, April 1981. (PB81 2063494) Solar Radiation as a Sole Source of Energy for Photovoltaics in Las Vegas, Nevada, for July and December. Darryl Randerson, April 1981. (PB81 2063494) A Systems Approach to Real-Time Hunoff Analysis with a Deterministic Rainfall-Runoff Model. Robert J.C. Burnash and R. Larry Ferral, April 1981. (PB81 224495) A Comparison of Two Methods for Forecasting Thunderstorms at Luke Air Force Base, Arizona. LTC Keith R. Cooley, April 1981. (PB81 23078) Annual Data and Verification Tabulation, Eastern North Pacific Tropical Storms and Hurricance 1980. Emil B. Gunther and Staff, May 1981. (PB81 23078) Annual Data and Verification Tabulation, Eastern North Pacific Tropical Storms and Hurricance 1980. Emil B. Gunther and Staff, May 1981. (PB82 12036) Preliminary Estimates of Wind Power Potential at the Nevada Test Site. Howard G. Booth, June 1981. (PB82 127051) A Statistical-Dynamical Model for Prediction of Tropical Cyclone Motion in the Eastern North Pacific Ocean. Jerston W. Leftwich, J., October 1981. (PB82195298) An Enhanced Plotter for Surface Airways Observations. Andrew J. Spry and Jeffrey L. Anderson, October 1981. (PB82 153883) Verification of 72-Hour 50-0MB Map-Type Predictions. RF. Quiring, November 1981. (PB82 158098 166
- 167 168
- 169
- 170
- 171 (PB82 158098)
- 172 Forecasting Heavy Snow at Wenatchee, Washington. James W. Holcomb, December 1981. (PB82 177783)
- 173 Central San Joaquin Valley Type Maps. Thomas R. Crossan, December 1981. (PB82 196064
- 174 176
- 177
- 178
- 179
- 180
- Central San Josquin Valley 19pe Maps. 100mas R. Crossan, December 1901. (PBo2 196064) ARAP Test Results. Mark A. Mathewson, December 1961. (PB82 198103) Approximations to the Peak Surface Wind Gusts from Desert Thunderstorms. Darryl Randerson, June 1982. (PB82 253089) Climate of Phoenix, Arizona. Robert J. Schmidli, April 1969 (Revised December 1986). (PB87 142063/AS) Annual Data and Verification Tabulation, Eastern North Pacific Tropical Storms and Hurricanes 1982. E.B. Gunther, June 1983. (PB85 106078) Stratified Maximum Temperature Relationships Between Sixteen Zone Stations in Arizona and Respective Key Stations. Ira S. Brenner, June 1983. (PB83 249904) Standard Hydrologic Exchange Format (SHEF) Version I. Phillip A. Pasteris, Vernon C. Bissel, David G. Bennett, August 1983. (PB85 106052) Guantitative and Spacial Distribution of Winter Precipitation along Utah's Wasatch Front. Lawrence B. Dunn, August 1983. (PB85 106012) 500 Millibar Sign Frequency Teleconnection Charts Winter. Lawrence B. Dunn, December 1983. (PB83 106276) 181 182
- 183
- 1900 Million Sign Frequency Teleconnection Charts Spring, Lawrence B. Dunn, January 1984. (PB85 111367) 184
- 1984. (PB85 111367)
 Collection and Use of Lightning Strike Data in the Western U.S. During Summer 1983.
 Glenn Rasch and Mark Mathewson, February 1984. (PB85 110534)
 500 Millibar Sign Frequency Teleconnection Charts Summer. Lawrence B. Dunn, March
 Annual Data and Verification Tabulation eastern North Pacific Tropical Storms and
 Hurricanes 1983. E.B. Gunther, March 1984. (PB85 109635)
 500 Millibar Sign Frequency Teleconnection Charts Fall. Lawrence B. Dunn, May 1984. 185
- 186
- 187
- The Use and Interpretation of Isentropic Analyses. Jeffrey L. Anderson, October 1984. 188

- 188 The Use and Interpretation of Isentropic Analyses. Jeffrey L. Anderson, October 1984. (PB85 132664)
 189 Annual Data & Verification Tabulation Eastern North Pacific Tropical Storms and Hurricanes 1984. E.B. Gunther and R.L. Cross. April 1985. (PB85 1878887AS)
 190 Great Salt Lake Effect Snowfall: Some Notes and An Example. David M. Carpenter, October 1985. (PB86 119153/AS)
 191 Large Scale Patterns Associated with Major Freeze Episodes in the Agricultural Southwest. Ronald S. Hamilton and Glenn R. Lussky, December 1985. (PB86 144474AS)
 192 NWR Voice Synthesis Project: Phase I. Glen W. Sampson, January 1986. (PB86 145664/AS)
 193 The MCC An Overview and Case Study on Its Impact in the Western United States. Glenn R. Lussky, March 1986. (PB86 170651/AS)
 194 Annual Data and Verification Tabulation Eastern North Pacific Tropical Storms and Hurricanes 1985. E.B. Gunther and R.L. Cross, March 1986. (PB86 17064/AS)
 195 Radid Interpretation Guidelines. Roger G. Pappas, March 1986. (PB86 17060/AS)
 196 Amesocale Convective Complex Type Storm over the Desert Southwest. Darryl Randerson, April 1986. (PB86 19098/AS)
 197 The Effects of Eastern North Pacific Tropical States. Walter Smith, August 1986. (PB86 (PB86 107650/AS)
 198 Preliminary Lightning Climatology Studies for Idaho. Christopher D. Hill, Carl J. Gorski, and March 2000.

- Preliminary, Lightin 1900. (FBS (10253AS) Preliminary Lighting Climatology Studies for Idaho. Christopher D. Hill, Carl J. Gorski, and Michael C. Conger, April 1987. (PB87 180196/AS) Heavy Rains and Flooding in Montans: A Case for Slantwise Convection. Glenn R. Lussky, April 1987. (PB87 185229/AS) 199

- Annual Data and Verification Tabulation Eastern North Pacific Tropical Storms and 200 Hurricanes 1986. Roger L. Cross and Kenneth B. Mielke. September 1987. (PB88 110895/AS) 201 An Inexpensive Solution for the Mass Distribution of Satellite Images. Gien W. Sampson and
- An mexpensive Source for the Wass Distribution of Satellite images. Gien W. Sampson and George Clark, September 1987. (PB88 114038/AS) Annual Data and Verification Tabulation Eastern North Pacific Tropical Storms and 202
- Hurricanes 1987. Roger L. Cross and Kenneth B. Mielke, September 1988. (PB88 101935/AS)
- An Investigation of the 24 September 1986 "Cold Sector" Tornado Outbreak in Northern 203 An investigation of the properties in the second second research of the second research in the second research is a second research in the second research in the second research is a second research in the second research in the second research is a second research in the second research in the second research is a second research in the second research in the second research is a second research in the second research in the second research in the second research in the second research is a second research in the second research in the second research is a second research in the second re
- 204 Carven Scott, November 1988. (PB89 128649/AS) Forecast Guidelines For Fire Weather and Forecasters -- How Nighttime Humidity Affects
- 205 Wildland Fuels. David W. Goens, February 1989. (PB89 162549/AS) A Collection of Papers Related to Heavy Precipitation Forecasting. Western Region 206
- Headquarters, Scientific Services Division, August 1989. (PB89 230833/AS) The Las Vegas McCarran International Airport Microburst of August 8, 1989. Carven A. Scott, 207
- June 1990. (PB90-240268) 208
- Meteorological Factors Contributing to the Canyon Creek Fire Blowup, September 6 and 7, 1988. David W. Goens, June 1990. (PB90-245085) Stratus Surge Prediction Along the Central California Coast. Peter Felsch and Woodrow 209
- Whitlatch, December 1990. (PE91-129239) Hydrotools. Tom Egger. January 1991. (PE91-151787/AS) A Northern Utah Soaker. Mark E. Struthwolf. February 1991. (PE91-168716) Preliminary Analysis of the San Francisco Rainfall Record: 1849-1990. Jan Null, May 1991. 210
- 911
- 212
- (PB91-208439) Idaho Zone Preformat, Temperature Guidance, and Verification. Mark A. Mollner, July 1991. 213
- (PB91-227405/AS) Emergency Operational Meteorological Considerations During an Accidental Release of 214
- Hazardous Chemicals. Peter Mueller and Jerry Galt, August 1991. (PB91-235424) WeatherTools. Tom Egger, October 1991. (PB93-184950)
- Creating MOS Equations for RAWS Stations Using Digital Model Data. Dennis D. Gettman, December 1991. (PB92-131473/AS) 216
- Forecasting Heavy Snow Events in Missoula, Montana. Mike Richmond, May 1992. (PB92-217 196104)
- 218 NWS Winter Weather Workshop in Portland, Oregon. Various Authors, December 1992. (PB93-146785)
- A Case Study of the Operational Usefulness of the Sharp Workstation in Forecasting a Mesocycione-Induced Cold Sector Tornado Event in California. John P. Monteverdi, March 1993. (PB93-178697) 219
- Climate of Pendleton, Oregon. Claudia Bell, August 1993. (PB93-227536)
- Utilization of the Bulk Richardson Number, Helicity and Sounding Modification in the Assessment of the Severe Convective Storms of 3 August 1992. Eric C. Evenson, September 221 1993. (PB94-131943)
- Convective and Rotational Parameters Associated with Three Tornado Episodes in Northern and Central California. John P. Monteverdi and John Quadros, September 1993. 222 (PB94-131943)
- (PB94-D1940) Climate of San Luis Obispo, California. Gary Ryan, February 1994. (PB94-162062) Climate of Wenatchee, Washington. Michael W. McFarland, Roger G. Buckman, and Gregory E. Matzen, March 1994. (PB94-164308) 224
- 225 Climate of Santa Barbara, California. Gary Ryan, December 1994. (PB95-173720) Climate of Yakima, Washington. Greg DeVoir, David Hogan, and Jay Neher, December 1994.
- 226 (PB95-173688)
- 997
- Climate of Kalispell, Montana. Chris Maier, December 1994. (PB95-169488) Forecasting Minimum Temperatures in the Santa Maria Agricultural District. Wilfred Pi and 228
- Poterseing imminimi reinpracties in eite band anna reinverteita Disarte von et i auto Peter Felsch, December 1994. (PB95-171088)
 The 10 February 1994 Oroville Tornado-A Case Study. Mike Staudenmaier, Jr., April 1995.
 Santa Ana Winds and the Fire Outbreak of Fall 1993. Ivory Small, June 1995.

NOAA SCIENTIFIC AND TECHNICAL PUBLICATIONS

The National Oceanic and Atmospheric Administration was established as part of the Department of Commerce on October 3, 1970. The mission responsibilities of NOAA are to assess the socioeconomic impact of natural and technological changes in the environment and to monitor and predict the state of the solid Earth, the oceans and their living resources, the atmosphere, and the space environment of the Earth.

The major components of NOAA regularly produce various types of scientific and technical information in the following kinds of publications.

PROFESSIONAL PAPERS--Important definitive research results, major techniques, and special investigations.

CONTRACT AND GRANT REPORTS--Reports prepared by contractors or grantees under NOAA sponsorship.

ATLAS--Presentation of analyzed data generally in the form of maps showing distribution of rainfall, chemical and physical conditions of oceans and atmosphere, distribution of fishes and marine mammals, ionospheric conditions, etc. TECHNICAL SERVICE PUBLICATIONS--Reports containing data, observations, instructions, etc. A partial listing includes data serials; prediction and outlook periodicals; technical manuals, training papers, planning reports, and information serials; and miscellaneous technical publications.

TECHNICAL REPORTS-Journal quality with extensive details, mathematical developments, or data listings.

TECHNICAL MEMORANDUMS--Reports of preliminary, partial, or negative research or technology results, interim instructions, and the like.

 \hat{V}

10



Information on availability of NOAA publications can be obtained from:

NATIONAL TECHNICAL INFORMATION SERVICE

U. S. DEPARTMENT OF COMMERCE

5285 PORT ROYAL ROAD

SPRINGFIELD, VA 22161