NOAA Technical Report NMFS SSRF-714



4

Wind Stress and Wind Stress Curl Over the California Current

Craig S. Nelson

August 1977

U.S. DEPARTMENT OF COMMERCE National Oceanic and Atmospheric Administration National Marine Fisheries Service

NOAA TECHNICAL REPORTS

National Marine Fisheries Service, Special Scientific Report-Fisheries

The major responsibilities of the National Marine Fisheries Service (NMFS) are to monitor and assess the abundance and geographic distribution of fishery resources, to understand and predict fluctuations in the quantity and distribution of these resources, and to establish levels for optimum use of the resources. NMFS is also charged with the development and implementation of policies for managing national fishing grounds, development and enforcement of domestic fisheries regulations, surveillance of foreign fishing off United States coastal waters, and the development and enforcement of international fishery agreements and policies. NMFS also assists the fishing industry through marketing service and economic analysis programs, and mortgage insurance and vessel construction subsidies. It collects, analyzes, and publishes statistics on various phases of the industry.

The Special Scientific Report—Fisheries series was established in 1949. The series carries reports on scientific investigations that document long-term continuing programs of NMFS, or intensive scientific reports on studies of restricted scope. The reports may deal with applied fishery problems. The series is also used as a medium for the publication of bibliographies of a specialized scientific nature.

NOAA Technical Reports NMFS SSRF are available free in limited numbers to governmental agencies, both Federal and State. They are also available in exchange for other scientific and technical publications in the marine sciences. Individual copies may be obtained (unless otherwise noted) from D825, Technical Information Division, Environmental Science Information Center, NOAA, Washington, D.C. 20235. Recent SSRFs are:

649. Distribution of forage of skipjack tuna (*Euthynnus pelamis*) in the eastern tropical Pacific. By Maurice Blackburn and Michael Laurs. January 1972, iii + 16 p., 7 figs., 3 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

650. Effects of some antioxidants and EDTA on the development of rancidity in Spanish mackerel (*Scomberomorus maculatus*) during frozen storage. By Robert N. Farragut. February 1972, iv + 12 p., 6 figs., 12 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

651. The effect of premortem stress, holding temperatures, and freezing on the biochemistry and quality of skipjack tuna. By Ladell Crawford. April 1972, iii + 23 p., 3 figs., 4 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

653. The use of electricity in conjunction with a 12.5-meter (Headrope) Gulf-of-Mexico shrimp trawl in Lake Michigan. By James E. Ellis. March 1972, iv + 10 p., 11 figs., 4 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

654. An electric detector system for recovering internally tagged menhaden, genus *Brevoortia*. By R. O. Parker, Jr. February 1972, iii + 7 p., 3 figs., 1 app. table. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

655. Immobilization of fingerling salmon and trout by decompression. By Doyle F. Sutherland. March 1972, iii + 7 p., 3 figs., 2 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

656. The calico scallop, Argopecten gibbus. By Donald M. Allen and T. J. Costello. May 1972, iii + 19 p., 9 figs., 1 table. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

657. Making fish protein concentrates by enzymatic hydrolysis. A status report on research and some processes and products studied by NMFS. By Malcolm B. Hale. November 1972, v + 32 p., 15 figs., 17 tables, 1 app. table. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

658. List of fishes of Alaska and adjacent waters with a guide to some of their literature. By Jay C. Quast and Elizabeth L. Hall. July 1972, iv + 47 p. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

659. The Southeast Fisheries Center bionumeric code. Part I: Fishes. By Harvey R. Bullis, Jr., Richard B. Roe, and Judith C. Gatlin. July 1972, xl + 95 p., 2 figs. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

660. A freshwater fish electro-motivator (FFEM)-its characteristics and operation. By James E. Ellis and Charles C. Hoopes. November 1972, iii + 11 p., 2 figs.

661. A review of the literature on the development of skipjack tuna fisheries in the central and western Pacific Ocean. By Frank J. Hester and Tamio Otsu. January 1973, iii + 13 p., 1 fig. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

662. Seasonal distribution of tunas and billfishes in the Atlantic. By John P. Wise and Charles W. Davis. January 1973, iv + 24 p., 13 figs., 4 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

663. Fish larvae collected from the northeastern Pacific Ocean and Puget Sound during April and May 1967. By Kenneth D. Waldron. December 1972, iii + 16 p., 2 figs., 1 table, 4 app. tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

664. Tagging and tag-recovery experiments with Atlantic menhaden, *Brevoortia tyrannus*. By Richard L. Kroger and Robert L. Dryfoos. December 1972, iv + 11 p., 4 figs., 12 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

665. Larval fish survey of Humbolt Bay, California. By Maxwell B. Eldrige and Charles F. Bryan. December 1972, iii + 8 p., 8 figs., 1 table. For sale by the Superintendent of Documents, U.S. Government Printing, Office, Washington, D.C. 20402.

666. Distribution and relative abundance of fishes in Newport River, North Carolina. By William R. Turner and George N. Johnson. September 1973, iv + 23 p., 1 fig., 13 tables. For sale by the Superinten dent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

667. An analysis of the commercial lobster (*Homarus americanus* fishery along the coast of Maine, August 1966 through December 1970. By James C. Thomas. June 1973, v + 57 p., 18 figs., 11 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

668. An annotated bibliography of the cunner, *Tautogolabrus adspersus* (Wilbaum). By Fredric M. Serchuk and David W. Frame. May 1973, ii + 43 p. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

669. Subpoint prediction for direct readout meterological satellites. By L. E. Eber. August 1973, iii + 7 p., 2 figs., 1 table. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

670. Unharvested fishes in the U.S. commercial fishery of western Lake Erie in 1969. By Harry D. Van Meter. July 1973, iii + 11 p., 6 figs., 6 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

671. Coastal upwelling indices, west coast of North America, 1946-71. By Andrew Bakun. June 1973, iv + 103 p., 6 figs., 3 tables, 45 app. figs. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

NOAA Technical Report NMFS SSRF-714



Wind Stress and Wind Stress Curl Over the California Current

Craig S. Nelson

August 1977

U.S. DEPARTMENT OF COMMERCE Juanita M. Kreps, Secretary National Oceanic and Atmospheric Administration Robert M. White, Administrator

National Marine Fisheries Service Robert W. Schoning, Director

For Sale by the Superintendent of Documents, U.S. Government Printing Office Washington D.C. 20402 Stock No. 003-020-00139-4 The National Marine Fisheries Service (NMFS) does not approve, recommend or endorse any proprietary product or proprietary material mentioned in this publication. No reference shall be made to NMFS, or to this publication furnished by NMFS, in any advertising or sales promotion which would indicate or imply that NMFS approves, recommends or endorses any proprietary product or proprietary material mentioned herein, or which has as its purpose an intent to cause directly or indirectly the advertised product to be used or purchased because of this NMFS publication.

CONTENTS

T			
-1-2	-	-	~
1	-	0.1	പ
	a	\sim	6
		<u> </u>	

Introduction	1
Data reduction	2
Dependence of stress estimates on C_D	6
Effects of averaging	6
Effect of atmospheric stability	6
Wind stress distributions	9
Spatial and temporal variability	12
Seasonal cycle at the coast	17
Wind stress curl	19
Monthly curl distributions	20
Coastal time series	22
Discussion	22
Physical implications	23
Biological implications	24
Conclusions	25
Acknowledgments	25
Literature cited	25
Appendix I Monthly surface wind stress distributions	27
Charts 1-19 Resultant surface wind stress vectors January through December	28
Charts 13-24 East component surface wind stress January through December	40
Charts 25-26. North component surface wind stress, January through December	52
Appendix II. Standard errors of the means	64
Appendix II. Monthly wind stress our distributions	75
Charte 27.49 Wind stress curl distributions	10
Unarts 57-46. Wind stress curi, January through December	10

Table

1. W	Vind speed	equivalents of	Beaufort	force estimates	in knots	and meters	per second		5
------	------------	----------------	----------	-----------------	----------	------------	------------	--	---

Figures

1.	Chart of the west coast of the United States showing the grid of 1-degree squares used for summaries of		
	wind observations		3
2.	Distribution of observations per 1-degree square		4
3.	Equivalent neutral drag coefficient for June		7
4.	Equivalent neutral drag coefficient for December		8
5.	The effect of atmospheric stability on surface wind stress for June		10
6.	The effect of atmospheric stability on surface wind stress for December		11
7.	Wind stress constancy for June		13
8.	Wind stress constancy for December		14
9.	Relative frequency surfaces and frequency histograms for June		15
10.	Relative frequency surfaces and frequency histograms for December		16
11.	Relative frequency surfaces and frequency histograms for July		17
12.	Relative frequency surfaces and frequency histograms for January		18
13.	Seasonal cycle of alongshore surface wind stress near the coast		19
14.	Seasonal cycle of onshore surface wind stress near the coast		20
15.	A conceptual diagram of the relationship of wind stress curl to the divergence and convergence of sur-	·	20
	face Ekman transport offshore of the primary coastal unwelling zone	•	20
16.	Discretization grid used in calculating the wind stress curl		21
17.	Seasonal cycle of wind stress curl near the coast	•	22
18	Distributions of: A wind stress curl B surface currents and C anchovy subnonulations	•	24
	Distributions of, M. white stress curr, D. surface currents, and C. anchovy subpopulations		41

Wind Stress and Wind Stress Curl Over the California Current

CRAIG S. NELSON¹

ABSTRACT

Historical surface marine wind observations are summarized by 1-degree square areas and months to describe the seasonal distribution of wind stress over the California Current. Off the coasts of southern California and Baja California, an alongshore equatorward component of surface wind stress is present throughout the year. The distributions of wind stress north of Cape Mendocino are characterized by marked changes in direction and magnitude between summer and winter. The predominant wind stress maximum shifts northward coherently from off Point Conception in March to south of Cape Blanco in September, and extends approximately 500 km in the offshore direction and 1,000 km in the alongshore direction. Maximum values of surface wind stress occur during July near Cape Mendocino.

The wind stress curl is positive near the coast and negative in the region offshore. A line of zero wind stress curl parallels the coast 200 to 300 km offshore, except off central Baja California. The patterns of wind stress curl are consistent with the existence of an equatorward Sverdrup transport offshore and a poleward transport near the coast.

INTRODUCTION

Spatial variation of surface wind stress has long been recognized as a fundamental quantity in discussions of the wind-driven ocean circulation. Sverdrup's (1947) transport balance relates the vertically integrated meridional mass transport in the interior ocean to the open ocean wind stress curl. Munk (1950) extended the theory to a closed basin and estimated the mass transports for several oceanic circulations from a knowledge of the wind stress alone. Recent theoretical developments have attemped to explain the well-known westward intensification, but have largely ignored the wind-driven circulation in eastern boundary currents, such as the California Current.

The California Current flows southeastward along the west coast of the United States as one branch of the large anticyclonic gyre in the North Pacific Ocean. The predominant surface current occurs between a cell of high atmospheric pressure to the west and a continental thermal low situated over California. Seasonal variations in the direction and strength of the surface wind are related to shifts in location of the high pressure system and intensification of the semipermanent thermal low.

The basic flow of the California Current system may be described in terms of the local wind stress (Reid et al. 1958). The equatorward surface current and poleward undercurrent are typical of eastern boundary currents (Wooster and Reid 1963). Munk (1950) has suggested that the average wind distribution off California is consistent with the existence of an equatorward surface current offshore and a poleward current inshore. However,

Pacific Environmental Group, National Marine Fisheries Service, NOAA, Monterey, CA 93940. lack of an adequate data base has precluded a detailed analysis of the relationships among the distributions of surface wind stress, the southward flowing California Current, and the northward flowing countercurrent which appears seasonally off the coasts of California, Oregon, and Washington.

A related phenomenon in eastern boundary currents is the process of coastal upwelling. The role of upwelling in bringing nutrients into the surface layers where they are available for organic production is widely recognized. According to the simplified model (Sverdrup 1938), equatorward wind stress parallel to the coast induces flow in the surface Ekman layer of the ocean which is deflected offshore by the earth's rotation (Ekman 1905). When this occurs over an expanse of coast where horizontal surface flow cannot compensate for that driven offshore, the balance is maintained by upwelling of subsurface water. Smith (1968) suggested that wind stress, being the driving force in this mechanism, may be the most important single variable in coastal upwelling.

Previous estimates of the surface wind stress over the oceans have been based primarily on wind data taken from the U.S. Hydrographic Office Pilot Chart wind roses (Hidaka 1958; Hellerman 1967). The large-scale features evident in these distributions are consistent with the occurrence of the major cyclonic and anticyclonic circulations in the oceans. However, the coarse spatial and temporal resolution of these data (5-degree quadrilateral space average and 3-mo time average) are not adequate to describe the locally driven processes evident in the California Current system.

Wooster and Reid (1963) used Hidaka's estimates of surface wind stress to calculate the offshore Ekman transport and described the locations and seasonal timing of the major coastal upwelling regimes in terms of these "upwelling indices." Bakun (1973) extended the concept of the upwelling index by computing estimates of Ekman transport from routinely available analyzed surface atmospheric pressure fields. Temporal fluctuations are well described by these data. However, both the derived quantities based on Hidaka's wind stress data and Bakun's upwelling indices suffer from an inability to characterize small-scale features of the Ekman transport field, particularly near the coast. For example, where these data may indicate surface divergence on the large scale, smaller scale surface convergence might exist, with associated effects on distributions of organisms within the coastal upwelling zone.

This report provides more detailed descriptions of the surface wind stress distributions over the California Current. The study differs from previous work by calculating monthly mean values on a 1-degree square area basis. Surface marine wind observations have been utilized in the computations. Roden (1974) evaluated the surface wind stress on a 1-degree latitude-longitude grid. However, Roden's distributions were derived from monthly mean surface atmospheric pressure analyses based on a 5-degree latitude-longitude grid. Thus, any information concerning space scales smaller than 5 degrees is due to the particular interpolation scheme used to refine the data to a 1-degree grid, rather than due to observed data.

The monthly mean data described in this report adequately resolve the seasonal cycle, which is the dominant time scale for coastal upwelling (Mooers et al. 1976). The high resolution in space and time may provide the observational background for more detailed investigations of the relationships among the local wind stress distributions, the equatorward surface current offshore, the poleward surface and subsurface flow inshore, and distributions of certain species of fishes, such as the northern anchovy, *Engraulis mordax*, and the Pacific mackerel, *Scomber japonicus*.

DATA REDUCTION

The monthly mean distributions of surface wind stress presented in Appendix I are based on summaries of data contained in the National Climatic Center's file of surface marine observations (Tape Data Family-11). The total file contains approximately 40 million individual ship reports dating from the mid-19th century. Over 1 million of the reports are within the area of the California Current system.

I have compiled long-term composite monthly fields of surface wind stress on a 1-degree square area basis within the geographical area outlined in Figure 1. The data grid extends from lat. 20°N to 50°N and parallels the coastline configuration, extending 10 degrees of longitude in the offshore direction. Each 1-degree quadrilateral is centered on a whole degree of latitude and longitude. Approximately 25% of the total available reports contain positions recorded to the nearest whole degree of latitude and longitude. The grid orientation used in this study thus minimizes spatial bias which might be introduced by summarizing the data according to the Marsden square numbering system.

The historical data contain errors in position measurement, and processing. A single pass editor was used to remove gross errors in the data, including duplicate reports (0.5%), position errors (0.1%), and measurement errors (0.5%). Erroneous wind directions and wind speeds greater than 100 m s⁻¹ were removed Reports of variable winds were treated as calms.

An estimate of the surface wind stress was calculated for each wind velocity report according to the bulk formula:

$$(\tau_x, \tau_y) = \rho_a C_D (|\overline{W}_{10}| U_{10}, |\overline{W}_{10}| V_{10})$$
(1)

where τ_x and τ_y denote the eastward and northward components of stress, ρ_a is the density of air which wa considered to have a constant value of 0.00122 g cm⁻³ $|\vec{W}_{10}|$ is the observed wind speed, and U_{10} and V_{10} are the eastward and northward components of the wind ve locity measured at a height of 10 m. The empirical drag coefficient C_D , referred to the 10-m level, was given a constant value of 0.0013 (Kraus 1972).

The resultant long-term monthly mean wind stres vectors were computed as the arithmetic means by eas and north components of all available reports from 1850 to 1972 within a 1-degree square area. The appropriate average is defined in Equation (2):

$$(\tilde{\tau}_x, \tilde{\tau}_y) = \frac{1}{N} \sum_{i=1}^{N} (\tau_x, \tau_y)_i$$
⁽²⁾

where N is the total number of reports within a 1-degre square area and month. The values $(\tau_x, \tau_y)_i$ wer evaluated according to Equation (1). A mean value for each long-term month and square is therefore forme from a data set which is independent of all other month and squares. The monthly fields of surface wind stress ar displayed in Appendix I as vector quantities (Charts 1 t 12) and as east (Charts 13 to 24) and north (Charts 25 t 36) components. No attempt has been made to smoot the fields, either by removing data which do not appea to fit the distributions or by applying objectiv smoothing procedures. The mean values were contoured by computer and "bull's-eyes" in the contours, even where they possibly reflect erroneous data, were left in the charts as indications of the general degree of consis tency in the composite distributions. The bull's-eye may reflect either a paucity of ship observations, or ex treme variability associated with inadequate sampling o strong winds.

The spatial distribution of observations is biased in that "ship of opportunity" reports are generally confined to coastwise shipping lanes. Contours indicating the total numbers of observations per 1-degree square area are shown in Figure 2. The highest density of reports is found within 300 km of the coast, exceeding 40,000 observations per 1-degree square in the area south of Point Conception. The number of reports per 1-degree square



Figure 1.—Chart of the west coast of the United States showing the grid of 1-degree squares used for summaries of wind observations. Large dots indicate squares for which frequency diagrams are displayed in Figures 9-12.



Figure 2.-Distribution of observations per 1-degree square. The contour interval is 2,500 observations. Values greater than 5,000 are shaded.

per month varies from less than 20 (in January off Vancouver Island) to more than 3,800 (in May off Los Angeles). A significant increase in the number of observations is evident along the shipping route between San Francisco and Hawaii. Less than 20% of the vector means are based on fewer than 50 observations per 1-degree square per month. Some temporal bias may also exist, since approximately 50% of the total reports have been taken since 1945. However, the general coherence of the resulting vector fields indicates that the composite wind stress distributions can be used to describe the dominant seasonal cycle in the California Current system.

The historical surface marine observations used in this study have been obtained primarily from ship logs, ship weather reporting forms, and published ship observations. The reports differ markedly in method and precision of measurement, ranging from observations made aboard 19th century vessels, to those taken aboard modern oceanographic research ships. Possible errors in wind measurement have been summarized by Han⁺el (1970). These include influences of anemometer height, atmospheric stability, wind gusts, and duration of the observation. Sources of error in wind estimates include the wind effects observed, error in determining the true wind from the observed apparent wind, and the unavoidable subjectivity of the observer.

A substantial portion of the historical data consists of wind reports based on the antiquated Beaufort wind force scale (Kinsman 1968). Table 1 lists the conversion

Table 1Wine	d speed equi	val	ents of
Beaufort force	e estimates	in	knots
and meters pe	r second.		

Regufort	Knots	-1	
Deautore	TEHOUS		
0	0	0.0	
1	2	0.9	
2	5	2.4	
3	9	4.4	
4	13	6.7	
5	18	9.3	
6	24	12.3	
7	30	15.5	
8	37	18.9	
9	44	22.6	
10	52	26.4	
11	60	30.5	
12	68	34.8	

scale between the Beaufort number and the equivalent wind speed in knots and meters per second. These estimates are equivalent to a wind speed measurement at a height of 10 m.^2

Based on the documentation for the National Climatic Center's data file (TDF-11),³ a determination was made of the number of wind observations estimated and those actually measured by an anemometer. Approximately 35% of the reports in this study consisted of Beaufort wind force estimates. An additional 53% of the reports were estimated wind speeds which did not correspond to the Beaufort wind force scale. Less than 12% of the total reports consisted of measured quantities.

In addition to the errors introduced by the necessary calculation of true wind from the measured apparent wind, the reported directions varied in precision. Resolution varied from $\pm 11.25^{\circ}$ to $\pm 5^{\circ}$ corresponding to observations based on 16 points of a 32-point compass and a 36-point compass, respectively.

The above considerations lead to the conclusion that random observational errors may be as large as real nonseasonal fluctuations in the distributions of surface wind stress. The problem may be formalized by expressing the individual stress estimates τ_x and τ_y as the sums of monthly mean values $\tilde{\tau}_x$ and $\tilde{\tau}_y$, deviations from the monthly means τ'_x and τ'_y , and error terms τ^{ϵ}_x and τ^{ϵ}_y . The second and higher moment statistics will consist of the nonzero correlations between the deviations from the monthly means and the error terms. If the observational errors are greater than the real fluctuations, then the standard deviations of the monthly means will reflect observational noise rather than actual nonseasonal variability. However, provided that these errors are not systematic, the resultant first moment statistics will be the appropriate estimates of the monthly mean wind stress.

The standard error of the mean provides a more appropriate quantitative measure of the relation between the standard deviation of a set of measurements and the precision of the mean value of the data set (Young 1962). The standard errors of the means are defined by:

$$(S_{\overline{\tau}_{x}}, S_{\overline{\tau}_{y}}) = (S_{\tau_{x}} / N, S_{\tau_{y}} / N)$$
(3)

where $S_{\overline{\tau}_x}$ and $S_{\overline{\tau}_y}$ are the standard errors of the means of the eastward and northward components, respectively, S_{τ_x} and S_{τ_y} are the standard deviations, and N is the number of observations. Large values of N and small values of S_{τ_x} and S_{τ_y} correspond to mean values $\overline{\tau}_x$ and $\overline{\tau}_y$ which closely approximate the population means. The computed standard errors of the means and num-

bers of observations of each 1-degree square area and long-term month are tabulated in Appendix II. Values less than 0.1 dyne cm⁻² occur throughout a large portion of the summary area, although spatial and seasonal dependence is evident. Standard errors greater than 0.1 dyne cm⁻² occur over large areas north of Cape Mendocino and between 500 and 1,000 km off the coast of Baja California. High values are generally associated with regions of sparse date, although inadequate sampling of intense storms may also lead to large values. Typical ratios of the standard error to the mean stress $(S_{\overline{\tau}}/\overline{\tau})$ range from 0.01 off southern California to 1.0 offshore from Cape Blanco. Near Cape Mendocino, this ratio varies between 0.02 and 0.10. Minimum values adjacent to the coast south of lat. 40°N and along the shipping lane between San Francisco and Hawaii are well correlated with the distribution of observations shown in Figure 2.

²Resolution 9, International Meteorological Committee, Paris, 1946. ³National Climatic Center, Tape Data Family 11, NOAA/EDS/NCC, Asheville, N.C.

DEPENDENCE OF STRESS ESTIMATES ON C_D

A constant drag coefficient was used to compute the surface wind stress data displayed in Appendix I. Stress calculations were made for each wind report and then averaged to form the resultant monthly mean vectors. Different results might have been obtained if the stress were calculated from monthly mean wind data, or if effects of atmospheric stability were considered. These aspects are discussed below.

Errors associated with the observational data used in this study place certain limitations on the form of the bulk exchange formula expressed in Equation (1). If measurement errors are large, there may be little value in refining the parameterization by replacing C_D by a variable drag coefficient which is a function of stability or wind speed. Even with a constant C_D , the bulk formula contains nonlinearities. Therefore, the drag coefficient must be appropriate for the particular timeaveraged winds used in the calculations. That is, a drag coefficient appropriate for synoptic wind observations would not be appropriate for winds averaged over a month.

Effects of Averaging

The empirically derived transfer coefficient C_D , determined by eddy correlation and profile methods, is based on wind measurements averaged over periods of 30 to 60 min. Pond (1975) indicated that the appropriate values of $|\overline{W}_{10}|$, U_{10} , and V_{10} should be obtained over a period of a few minutes to a few hours at most. If the surface stress were calculated using winds averaged over much longer periods, a higher drag coefficient would be required. The surface stress calculated from a monthly mean surface wind field will be significantly less than the surface stress field calculated as the mean of the individual stress estimates, if the same drag coefficient is used and all other parameters are held constant.

The difference described above has been determined for the data used in this study. For each 1-degree square area and long-term month, the following ratio has been calculated:

$$(C_D)_{\text{equ}} = \left| \overrightarrow{\tau} \right| / \left[\rho_a (\left| \overrightarrow{W_{10}} \right|)^2 \right]$$
(4)

where $\lceil \tau \rceil_{15}$ the magnitude of the monthly mean surface stress, $\lceil \overline{W}_{10} \rceil$ is the magnitude of the monthly mean surface wind, and $(C_D)_{equ}$ is an equivalent drag coefficient. This is the value which would be necessary to make the stress values calculated from the monthly mean wind data agree with the values calculated as the means of individual stress estimates.

Values of (C_D) equ for the months of June and December are shown in Figures 3 and 4. The values are significantly different from the constant value of $C_D = 0.0013$ employed in this study. In all cases, (C_D) equ is greater than C_D . Spatial and seasonal

variability is marked. There is a tendency for much higher values in the northern section of the grid than in the southern area. Large values of $(C_D)_{eou}$ are more evident in December than in June. Geographical and seasonal variations in the quantity $(C_D)_{equ}$ agree with the general geographical and seasonal fluctuations in meteorological conditions over the northeast Pacific Ocean. This would indicate that a large part of the monthly variance of the wind stress data is due to actual intramonth fluctuations and not due to observational errors. If a resultant surface wind stress were calculated from the monthly mean wind field, these calculations imply that the wind stress estimates would be underestimated, on average, by as much as 50% to 100%. The departures would be even larger off Oregon and Washington.

Effect of Atmospheric Stability

Investigations on the functional form of the drag coefficient have alternately suggested constant values, or a dependence on wind speed. Wilson (1960) gave a detailed review of the available data and adopted values of $C_D = 0.0024 \pm 0.005$ for strong winds and $C_D = 0.0015 \pm 0.0008$ for light winds. Smith (1970) proposed a constant value of $C_D = (1.35 \pm 0.34) \times 10^{-3}$. Wu (1969) adopted the form $C_D = (0.5 \times \overline{U}_{10}^{-1/2}) \times 10^{-3}$. Considering the large scatter in the open ocean determinations of C_D , the lack of conclusive data at wind speeds greater than 15 ms⁻¹, and the lack of agreement among individual observers, a constant value for the neutral drag coefficient was used in this study.

Recent investigations by Davidson (1974) and Denman and Miyake (1973) have demonstrated the dependence of the drag coefficient on atmospheric stability and the spectral shape of the ocean wave field. A generally accepted formulation of the dependence of C_D on wave properties does not seem well enough established to be incorporated in the calculations of surface wind stress based on historical ship observations. However, effects of atmospheric stability on the momentum exchange may be significant in regions where seasonal upwelling typically produces a stable boundary layer.

The specific effects of atmospheric stability on the exchange of momentum have not been completely determined and are still under investigation. However, for a given wind speed, the effect of stable (unstable) stratification is to decrease (increase) the magnitude of the momentum exchange (surface wind stress) across the atmosphere-ocean interface.

In order to investigate the effects of atmospheric stability, the monthly mean fields of surface wind stress were recomputed, replacing the constant value C_D in Equation (1) by a coefficient varying with atmospheric stability. Deardorff (1968) defined the bulk Richardson number as an appropriate dimensionless measure of atmospheric stability. Deardorff's method was adopted to parameterize the dependence of the drag coefficient on stability, while neutral stability was assumed when the absolute value of the air-sea temperature difference was

NCC - TDF-11 - DNE DEGREE SUMMARIZATION



Figure 3.-Equivalent neutral drag coefficient for June. The plotted value is the drag coefficient (× 10') appropriate for wind observations averaged over a month. The contour interval is 2.5.

NCC	- TDF-	-11 -	DNE	DEGRE	E SUN	1MAR I	ZATIC	N										-110		1.1.77	110	115			0.11	2 1	
137	136	135	134	133	132	131	130	129	128	127	126	125	124	123	122	121	120	119	118	117	116	115	112	1 11	3 11	2 1	
						-		1	G	- mar	3																
50	4.21	4.96	3.19	4.08	5.36	4.60	3.83	7.14	4.95	3.29	VAN	COLVER	15LAN	-													Si
	-	4 40	A 74	A 177	4.56	3.95	4.41	7.00	5.82	5.90	6.93	22	~	K			[ND	AA					45
49		4.40	4.14		A				5			X	2	23	5			NAT	PACIF	ICE	INE F	ISHE	TAL	GRDU	INICE		63
48			4.23	3.33	6.53	3.11	5,401	4.73	4.39	5.74	6.25	4.95	1	4	1				М	ONTE	RΕΥ,	CALI	FDRN	IA			
47				3.08	3,36	4.16	4.37	4.05	6.45	5.71	4.57	4.35	4.03							EQI	JIV	ALE	ENT	Г			4
46				3.30	3.11	2.71	3,92	3.78	4.59	5.26	3,86	4.34	4.85	- COLU	MBLA A	IVER		[RA	G	COE	FF	ICI	EN	IT		4
45				3.42	3.72	3.43	4.11	3,39	4.03	6,05	7:53	5.89	5.28														4
				2.86	3, 15	3.54	4.51	3.82	4.91	5.51	5.20	5.15	3.91														4
							10.00	700	2 60		6 M	E 20	1							LO	NG TE	RM M	EAN				4
43				4.19	3.//	4.03	19.01	14 50	12.00				f }	CHPS	: BLHNC					1	FCE	MBF	R				
42				3.63	3.53	3.88	4.68	3.51	6.09	4.31	7.76	7.69	5.52								.B6Ø ·	- 197	2				4
41				3.37	3.39	3.59	4.88	3.79	4,56	9.35	0.53	10.99	11.50	0000	HEND	100					-						4
40	1			-5.0t	4.98	4.60	6.36	8.23	7,20	5.16	7 62	9.17	10.08	L.P.F.L	. HEAD	AC LING											4
39)	 	4,99	4.38	4,64	7.59	6.60	8.63	8.63	7,28	8.00	14.83														3
38					5.10	6.76	8.69	6.57	10.94	13.83	7,51	8.98	9.47	13.38	2												3
					12	Lon	1	E 04	1.00			4.05	6 38	E AR	A												-
37		ļ				4.Y		0,04	4900			0	0.00		3												
36						-4-81	6.38	6.87	6,36	6.68	5.78	5.33	7.38	4.55	4.12	1											F
35							7.91	5,86	5.38	6.05	4991	6.15	4.15	3,34	3.79	3.97	POI	NT CON	CEPTIC	N							9
34		1					-		5.14	4.07	3,15	3.87	3.58	3,65	3.00	3,30	4. 39	8.89	2								1.1
33							1			3.66	3.58	3.58	3.60	2.87	2.79	2.61	3.30	3.56	4.67	>							12
32							J				4.72	4.46	3.16	2.73	2.52	2.58	2.58	2.66	4.06	3.53							
		;;		 							3 03	1 3 10	3 17	3 39	3 34	2 74	4.15	2.85	2.73	3.11	1		~	2			
31							j				0.00	1	5.11	3,30	5.54	1		2.00	2.75		7	1	1		(
30												3,43	2.08	2.64	2.64	2.33	3.11	Z.42	21.52	2.63	2.5	5	1	-	1		F
29]	<u> </u>]						2.45	3.05	2.41	2.49	2.35	1.84	2.28	2.43	2.35	2.5	2	10	0	R	1
28						10.00						-	3.06	2:49	2.29	2.41	2.44	-2952	2.33	2.07	2.25	2.3	-	PUNTA	EUGEN	IA	4
27	1				1	1	1					-		3.07	3.20	2.14	1.97	1.88	2.33	1.84	2.00	2.3	3 2.	12	1	2	4
26)						2.89	2.07	2.66	1.86	2.03	4.13	3.42	2.2	8 2.	58 2	35	1	
25															2 00	2.14	2.07	2.32	1.02	2 5	123	12.	 a 2	27 2	42		2
								1							2.03	2114	2.07	LIDE							6	2	1
24				1 	1 1 1	1 1 	- 					1 1 					2.04	1.93	2,08	2.00	2.1	2.2	9 2.	23 2	45 2	.75	2.54
23	1		1	1				1									1.90	2.03	2.01	1.79	1.7	3 2.4	2.	71 3	. 10 2	.54	2.72
22				-									1				1.67	1.72	1.87	1.77	2.2	2.8	9 1.	85 1	.99 Z	. 86	2.73
21		1	1	1	1		1	1	1			1	1	1			1.63	1.75	1.72	1.75	1.5	1.7	4 2.	15 2	.28 2	.32	2.18
13	7 120	1.25	124	199	100	1.01	120	120	1.00	107	1.20	100	104	100	122	101	100	110	110	117	110				10.1	10.1	

Figure 4.—Equivalent neutral drag coefficient for December. The plotted value is the drag coefficient (\times 10³) appropriate for wind observations averaged over a month. The contour interval is 2.5. less than 1°C. Details of the calculations will be reported elsewhere.

The effect of atmospheric stability was computed as a percentage increase (decrease) in the magnitude of the monthly mean surface wind stress above (below) the appropriate neutral values. The following ratio was calculated:

$$R = \left(\frac{\left| \overrightarrow{\tau} \right|_{\text{var}}}{\left| \overrightarrow{\tau} \right|_{\text{neu}}} - 1 \right) \times 100$$
 5

where $|\vec{\tau}|_{var}$ is the magnitude of the monthly mean surface wind stress calculated using a drag coefficient which varies with stability, and $|\vec{\tau}|_{neu}$ is the magnitude of the monthly mean surface wind stress computed with a constant (neutral) coefficient. A value of R = 0.0 corresponds to neutrally stable conditions.

The ratio defined by Equation (5) is displayed in Figures 5 and 6 for the months of June and December, respectively. The average percentage difference is between 2 and 5%. Values greater than 10% are rare. In December, the values are positive, indicating unstable conditions. Negative values occur in June. These are related to the stable stratification induced by positive air-sea temperature differences within the coastal upwelling zone. The effects of atmospheric stability determined in this study are less than the values of 6 to 15% reported by Husby and Seckel (1975) for Ocean Weather Station "Victor," but consistent with the values of less than 5% reported by Dorman et al. (1974) for Ocean Weather Station "November."

WIND STRESS DISTRIBUTIONS

The long-term composite monthly mean fields of surface wind stress are displayed in Appendix I as resultant vectors (Charts 1 to 12), east components (Charts 13 to 24), and north components (Charts 25 to 36). These data indicate two characteristic features. The mean stress tends to be directed equatorward along the coast from Cape Mendocino to Baja California throughout the year. Off the coasts of Oregon and Washington, the wind stress exhibits marked seasonal variability in both magnitude and direction.

The distributions of surface wind stress off Baja California have been previously discussed by Bakun and Nelson (1975).⁴ They concluded that most of the seasonal variability is in magnitude rather than direction. The mean surface wind stress tends to have an alongshore, equatorward component during all months, implying conditions generally favorable for coastal upwelling throughout the year. This is contrasted with the situation to the north. Off the coasts of Oregon and Washington, patterns of surface wind stress change seasonally. From December to February, the stress impinges on the coast at Cape Blanco, forming the southern boundary of the low pressure system which develops in the Gulf of Alaska during the winter. From May through September, the surface wind stress veers toward the south. The components are directed onshore and equatorward. The months of March-April and October-November appear as transition periods. The surface wind stress is directed primarily onshore off Washington and Oregon during these months.

The most evident feature in these distributions is the position and extent of the predominant wind stress maximum. For purposes of discussion, this maximum is defined by the 1.0 dyne cm⁻² contour and is highlighted by light shading in Charts 1 to 12. Large values of stress occurring between lat. 45° N and 50° N during winter are probably associated with either high winds or sparse data, and will be ignored in this discussion.

The maximum is first evident in March, south of Point Conception. During succeeding months, the maximum strengthens, increases in size, and shifts northward. In April, values exceeding 1.0 dyne cm⁻² cover an area from Point Conception to San Francisco. Winds off Oregon have veered, suggesting an alongshore component and implying conditions generally favorable for coastal upwelling. The pattern of surface wind stress is repeated in May. The northern boundary of the central maximum reaches the coast south of Cape Blanco.

The mean wind stress reaches maximum intensity during June and July off Cape Mendocino, where characteristic values exceed 1.5 dynes cm⁻². The large-scale maximum extends approximately 500 km in the offshore direction and 1,000 km in the longshore direction. A more intense coastal maximum ($|\vec{\tau}| \ge 1.5$ dynes cm⁻²) is coherent over an area approximately one-fifth this size. Maximum values of mean surface wind stress typically occur 200 to 300 km off the coast, rather than adjacent to the coast. This feature was previously observed by Munk (1950) and Reid et al. (1958). Mean distributions for August and September suggest relaxing conditions; the maximum is reduced in extent and intensity. These months still exhibit a degree of equatorward stress along the coasts of Oregon and Washington, but in October, the direction backs to the north. The poleward components correspond to onshore Ekman transport and subsequent downwelling at the coast. The distributions for November, December, January, and February indicate characteristic winter conditions. Mean values are typically less than 0.5 dyne cm⁻². High winds associated with intense storm activity in the North Pacific Ocean result in regions of mean surface wind stress exceeding 1.0 dyne cm⁻² off the coasts of Oregon and Washington.

Seasonality of the surface wind stress and its association with coastal upwelling are easily observed along the northern section of the grid. Large changes in mean direction are apparent. Maximum magnitudes occur during the winter. Winds favorable for coastal upwelling occur from April to September. During the remaining months, the direction of the mean surface wind stress corresponds to downwelling at the coast.

[']Bakun, A. and C. S. Nelson, 1975. Climatology of upwelling-related processes off Baja California. Unpubl. manuscr., 41 p. On file at Pacific Environmental Group, National Marine Fisheries Service, NOAA, Monterey, CA 93940.



Figure 5.—The effect of atmospheric stability on surface wind stress for June. Plotted values are the percentage increase (decrease) in the surface wind stress above (below) neutrally stable conditions. The contour interval is 10.0. Positive values correspond to unstable conditions. Negative values are shaded.

NCC - TDF-11 ----ONE DEGREE SUMMARIZATION



Figure 6.- The effect of atmospheric stability on surface wind stress for December. Plotted values are the percentage increase (decrease) in the surface wind stress above (below) neutrally stable conditions. The contour interval is 10.0. Positive values correspond to unstable conditions. Negative values are shaded.

The coast along southern California and Baja California is characterized by winds favorable for upwelling throughout the year. Peak values of surface wind stress occur in April and May. Values exceeding 0.5 dyne cm⁻² are evident from February to July. A local maximum immediately north of Punta Eugenia is easily observed in May. This feature is consistent throughout the year. A region of local wind stress minima is indicated along the coast south of Point Conception. This feature corresponds in location with the semipermanent cyclonic eddy which dominates the ocean surface circulation in the Southern California Bight (Reid et al. 1958).

Spatial and Temporal Variability

Spatial and seasonal variability of the monthly distributions will be described in terms of standard errors of the means (Equation (3)), constancy of the wind stress, and frequency diagrams for selected 1-degree square areas and months. In general, these data indicate greater variability in magnitude and direction in the region north of Cape Mendocino than in the area to the south. Summer distributions are characterized by well-defined mean directions and magnitudes. Broad-banded frequency histograms are typical of winter months.

The degree of variability in direction and magnitude of the surface wind stress is reflected in the spatial distributions of the standard errors of the means. In regions outside of the primary shipping lanes (see Fig. 2), monthly mean distributions are based on approximately the same number of observations per 1-degree square area. If one assumes that there are no systematic sampling errors in these regions, then the magnitudes of the computed standard errors of the means should be a function of the wind stress variability. Accordingly, standard errors to the north of Cape Mendocino are larger by a factor of 2 to 3 (Appendix II) than those to the south of Cape Mendocino. Thus, the data imply a greater degree of variation in the direction and magnitude of the surface wind stress off Oregon and Washington than off California.

A contrast between winter and summer conditions is also evident. South of Cape Mendocino, magnitudes of the standard errors of the means remain nearly constant throughout the year. Off Oregon and Washington, the computed values decrease to a minimum during the summer, and increase to a maximum during the winter. Typical values range from 0.10 dyne cm⁻² in June to 0.30 dyne cm⁻² in December.

Similar features of the large-scale temporal and spatial variations are evident in monthly distributions of constancy of the wind stress as defined by the ratio of the magnitude of the average stress to the average magnitude of the stress. Figures 7 and 8 show the patterns for June and December. During December, values greater than 0.5 occur south of Point Conception. To the north, wind stress constancy decreases to relatively small values, implying a high degree of directional variability. An increase in directional constancy is indicated during June. Values greater than 0.50 extend from Baja California to Vancouver Island. Values less than 0.50 occur only in the northwest section of the grid. South of Cape Mendocino, the ratios approach a value of 1.0, implying very little variation in wind stress direction.

The patterns of wind stress constancy indicate possible error in estimates of the total mechnical energy transfer at the air-sea interface. In the region off California, the magnitude of the mean vector wind stress appears to be a good estimator of the total stress acting on the sea surface. However, in the northern regions of the data grid, these data show that the magnitude of the mean vector wind stress underestimates the total stress acting on the sea surface. This has important implications for mixed-layer modelling, in which the input of turbulent energy depends on the total stress acting on the sea surface (Denman 1973), the direction of the surface stress, and the rate at which the stress is applied to the sea surface (Niiler 1975).

The large-scale spatial and temporal variations in surface wind stress distributions are finally described in terms of selected frequency diagrams. Data for the 10 squares indicated in Figure 1 are displayed in Figures 9 to 12. Figures 9 and 10 show data for June and December taken from the five squares indicated in the inset. Similar data at five different locations are presented for July and January in Figures 11 and 12.⁶

Available wind stress data have been classed by direction and magnitude for each 1-degree square area and month. Relative frequencies have been determined for 16 direction bands and for 11 magnitude bands. A category for calm winds also includes variable winds. The direction bands are 22.5° wide and the magnitude bands correspond to equivalent wind speed intervals of 2.0 m s⁻¹.

The relative frequency surfaces shown in these figures display contours of percentage of total reports falling within a given direction and magnitude band. The 2.5% and 7.5% contours are indicated by a solid line. Dashed line contours indicate relative frequencies of 5.0% and 10.0%. The mean vector magnitude and direction is indicated by an arrow. Note that directions are defined with the oceanographic convention (i.e., the direction toward which the wind blows).

The histograms shown to the right of each frequency surface display relative frequencies for magnitude at the top, and direction at the bottom. Relative percent is labelled along the ordinate. Midpoints of the magnitude and direction class intervals are labelled on the abscissa.

The contrast between winter and summer distributions in the southern section of the grid appears as a change in the character of the frequency surfaces (Figs. 9, 10). In general, the number of contours indicated for June is greater than the number of contours appearing in December. Evidently, the wind stress is relatively constant in magnitude and direction during the summer

⁵Wind roses for all 1-degree squares within the area north of lat. 34°N may be found in *Climatic study of the near coastal zone; West Coast of the United States*, published by the Director, Naval Oceanography and Meteorology, June 1976, 133 p.





Figure 7.—Wind stress constancy for June. The plotted values are defined as the ratios of the vector means to the scalar means. The contour interval is 0.25.



Figure 8.—Wind stress constancy for December. The plotted values are defined as the ratios of the vector means to the scalar means. The contour interval is 0.25.



Figure 9.—Relative frequency surfaces and frequency histograms for June. Data are shown for the 1-degree squares labelled 1 through 5 in the upper left inset. Contours of relative frequency are drawn at intervals of 2.5%. Mean vector wind stress is indicated by an arrow.

months. A slight shift in direction and magnitude of the wind stress is indicated between summer and winter. The mean stress in December is directed more toward the south, and the magnitude has decreased.

The frequency histograms also show these features. During June, the direction histogram is characteristically narrow-banded. Over 70% of the observations may be concentrated within three direction intervals. In December, the observations tend to be spread over a wider range of directions. The histograms are broader and the peaks in direction are less well-defined. Peak magnitudes in June are generally one class interval larger than peak magnitudes in December. However, the winter distributions are characterized by an increase in relative frequency at high values of wind stress magnitude.

A similar pattern of contrasts between summer and winter distributions is apparent in the northern section of the grid. As shown in Figures 11 and 12, the direction histograms are generally well-defined during the summer, although bimodal distributions are evident. During winter, observations are nearly uniformly spread among all directions. There is a lack of consistency in the frequency surfaces in Figure 12. Direction histograms for



Figure 10.—Relative frequency surfaces and frequency histograms for December. Data are shown for the 1-degree squares labelled 1 through 5 in the upper left inset. Contours of relative frequency are drawn at intervals of 2.5%. Mean vector wind stress is indicated by an arrow.

January are broad and flat and the peaks which generally characterize the summer distributions are missing. Mean directions shift from equatorward to poleward between summer and winter. A complete reversal in the mean direction occurs at point 7 (Figs. 11, 12). A shift in magnitude is equally pronounced. Peak magnitudes are higher in January than in July indicating a greater contribution of high wind speeds in January.

The above discussion adds a new dimension to the seasonal descriptions of the surface wind stress distributions over the California Current. Pronounced seasonal variations in magnitude and direction of the monthly mean wind stress are indicated along the entire west coast of the United States. These changes are most evident off the coasts of Oregon and Washington. In addition, these data suggest month to month changes in the large-scale spatial variability. Frequency histograms for direction and magnitude are broad and flat during the winter months, and along the northern coast. Welldefined peaks in magnitude and direction characterize the distributions during the summer and along the southern coast.



Figure 11.—Relative frequency surfaces and frequency histograms for July. Data are shown for the 1-degree squares labelled 6 through 10 in the upper left inset. Contours of relative frequency are drawn at intervals of 2.5%. Mean vector wind stress is indicated by an arrow.

Seasonal Cycle at the Coast

Time series of the surface wind stress within the 1degree square areas immediately adjacent to the coast are displayed in Figure 13 as the alongshore component, and in Figure 14 as the onshore component. For these displays, the vector means have been resolved into components parallel and perpendicular to the coast. The coastline angles were determined by visually fitting a line to the dominant trend of the coast within each 1degree square. The months are indicated along the tops of the figures. The latitudes of the 1-degree squares are indicated on the right and left sides of the figures. Negative values are shaded and indicate equatorward alongshore stress in Figure 13 and offshore stress in Figure 14.

Several characteristic features are apparent in these figures. South of lat. 40°N there is an equatorward component throughout the year, implying conditions favorable for coastal upwelling in all months. Off the coasts of Oregon and Washington, the data suggest that upwelling occurs seasonally between the months of April and September.

Three relative maxima occur in the time/space domain. Off the coast of Baja California, maximum values



Figure 12.—Relative frequency surfaces and frequency histograms for January. Data are shown for the 1-degree squares labelled 6 through 10 in the upper left inset. Contours of relative frequency are drawn at intervals of 2.5%. Mean vector wind stress is indicated by an arrow.

of wind stress are evident near Punta Eugenia in May. A large maximum occurs just south of Cape Mendocino between May and August. A smaller, local maximum occurs in May and June at lat. 36°N.

Figure 13 is similar in appearance to a time series of offshore Ekman transport shown by Bakun et al. (1974). They showed a good correlation between the occurrence of maximum offshore transport at lat. 39°N and a suppression of seasonal warming in the adjacent coastal waters during early summer, which is indicative of upwelling. The timing of the central wind stress maximum off the coast from Cape Mendocino agrees with the description of the mean yearly cycle of indicated upwelling given by Bakun (1973). However, Bakun's data are spatially distorted, and indicate maximum values at lat. 33°N, long. 119°W in the middle of the Southern California Bight. The spatial distortion is primarily caused by development of an intense thermal low over southern California during the summer. The influence of this low pressure system, and the effects of coastal mountain ranges distort the analyzed pressure fields used in Bakun's computations.

The time series of alongshore surface wind stress (Fig. 13) suggests a slight tilt, with time and space, to the region of maximum values. This corresponds to a northward shift in the intensity of the surface wind stress from



April and May off the coast of Baja California to June, July, and August off Cape Mendocino and Cape Blanco.

Figure 14 indicates a tendency for the surface wind stress to be directed onshore throughout the year north of lat. 30°N. Off the coast of Baja California, surface stress is characterized by offshore components, except between the months of April and October. Offshore components are also apparent in the vicinity of Cape Mendocino and Point Conception. Near these points, abrupt changes in coastline orientation may influence the direction and magnitude of the surface wind stress.

WIND STRESS CURL

The surface wind stress curl is the forcing function for the vertically integrated mass transport of the winddriven ocean circulation. Under linear, steady-state conditions on the " β -plane," in the absence of friction and interactions with bottom topography, the meridional component of mass transport (M_y) is directly proportional to the vertical component of the curl of the wind stress as expressed in Equation (6):

$$M_{y} = \frac{\overrightarrow{k} \cdot (\overrightarrow{\nabla} \times \overrightarrow{\tau})}{\beta} \tag{6}$$

where M_y is the meridional component of the vertically integrated mass transport, β is the meridional derivative of the Coriolis parameter f, and $\vec{k} \cdot (\nabla \times \vec{\tau})$ is the vertical component of the wind stress curl. According to the simplified model, positive (negative) wind stress curl is associated with northward (southward) meridional transport. Surface Ekman divergence (convergence) corresponding to positive (negative) wind stress curl is balanced by geostrophic convergence (divergence) in the northward (southward) meridional flow.

Coastal upwelling occurs only at the ocean boundary. However, wind induced upwelling will occur whenever divergence in the surface wind drift is not balanced by other modes of horizontal surface flow. Figure 15 shows a mechanism by which the wind stress curl determines the divergent or convergent nature of the surface wind drift offshore of the primary coastal upwelling zone. A seaward increase in the equatorward wind stress parallel to the coast (Fig. 15A) is characterized by positive wind stress curl. In this situation, the offshore component of Ekman transport increases in the offshore direction, resulting in continued surface divergence. Upwelling is required to maintain the mass balance. If the equatorward longshore surface wind stress decreases in the offshore direction (Fig. 15B), the wind stress curl is negative. Convergence in the surface wind drift will result. Frontal formation and downwelling may occur just offshore of the primary coastal upwelling zone.

Figure 13.—Seasonal cycle of alongshore surface wind stress near the coast. Means of the alongshore components of wind stress were computed by month for the 1-degree squares immediately adjacent to the coast. Units are dyne cm⁻². The contour interval is 0.5 dyne cm⁻². Equatorward alongshore stress is shaded. Numbers along ordinate are degrees north latitude.





Figure 15.—A conceptual diagram of the relationship of wind stress curl to the divergence and convergence of surface Ekman transport offshore of the primary coastal upwelling zone.

Direct measurements of wind stress curl are not available; however, the curl of the monthly mean surface wind stress fields has been computed. The vertical component of the curl of the wind stress in spherical coordinates is defined by:

$$\vec{k} \cdot (\nabla \times \vec{\tau}) = \frac{1}{R \cos \varphi} \left[\frac{\partial \vec{\tau}_y}{\partial \lambda} - \frac{\partial}{\partial \varphi} (\vec{\tau}_x \cos \varphi) \right] (7)$$

where R is the radius of the earth, φ and λ denote geographic latitude and longitude, respectively, and $\overline{\tau}_x$ and $\overline{\tau}_y$ denote the eastward and northward components of the mean surface wind stress. A finite difference equation approximating the curl at the grid point (i, j) in Figure 16 is given by:

$$\vec{k} \cdot (\nabla \times \vec{\overline{\tau}}_{i,j}) = \frac{1}{R \cos \varphi_{i,j}} \left[\frac{\vec{\tau}_{y_{i+1,j}} - \vec{\tau}_{y_{i-1,j}}}{2\Delta \lambda} - \left(\frac{(\vec{\tau}_x \cos \varphi)_{i,j+1} - (\vec{\tau}_x \cos \varphi)_{i,j-1}}{2\Delta \varphi} \right) \right] (8)$$

where $\Delta \varphi = \Delta \lambda = 1^{\circ}$. The method is similar to calculations described by Hantel (1970). Centered differences were utilized throughout the interior of the data grid. Forward and backward differences were required along the boundaries. As a result, certain artificial features may have been introduced at these boundaries.

Monthly Curl Distributions

Monthly fields of surface wind stress curl are displayed in Appendix III (Charts 37 to 48). These distributions correspond to the monthly mean wind stress distributions shown in Appendix I. Values are plotted in units of dyne cm⁻² per 100 km. A value of 1 dyne cm⁻²

Figure 14.—Seasonal cycle of onshore surface wind stress near the coast. Means of the onshore components of wind stress were computed by month for the 1-degree squares immediately adjacent to the coast. Units are dyne cm⁻². The contour interval is 0.5 dyne cm⁻². Offshore stress is shaded. Numbers along ordinate are degrees north latitude.



Figure 16.—Discretization grid used in calculating the wind stress curl.

per 100 km is equivalent to 1×10^{-7} dyne cm⁻³. Negative values of wind stress curl are shaded.

Small scale features evident in these distributions should be viewed with caution. Detail within a single 1degree square area which is not supported by similar values in surrounding squares probably reflects "noise" in the monthly means of surface wind stress. Objective smoothing procedures applied to the monthly mean wind stress fields would result in more homogeneous distributions of wind stress curl. A particular method has been described by Evenson and Veronis (1975).

Characteristic absolute magnitudes of the spatially averaged wind stress curl are 1×10^{-8} dyne cm⁻³. This value is approximately a factor of two less than the summertime mean reported by Halpern (1976) for an upwelling region near the Oregon coast. Considering the time and space averages used in the present study, this difference appears to be reasonable. However, the values are consistent with wind stress curl calculations reported by Saunders (1976). The probable errors associated with these estimates of wind stress curl may be calculated from a knowledge of the spatial distributions of standard errors of the (wind stress) component means (see Appendix II). On average, the expected error is 1×10^{-8} dyne cm^{-3} , with maximum and minimum errors of 4×10^{-8} dyne cm⁻³ and 1 \times 10⁻¹⁰ dyne cm⁻³, respectively. Of course, these values apply to particular individual 1degree squares. Large values generally correspond to "holes" in the distributions, which are easily seen in the monthly charts. For larger space scales, errors associated with the gradients of the mean wind stress components would tend to cancel. Thus, greater confidence may be expected for the patterns of wind stress curl which are consistent over several degrees of latitude and longitude.

The large-scale features of positive and negative curl along the coast are significant. Important details to note are the sign of the wind stress curl at the coast, and the position of the line of zero wind stress curl. A general feature common to all months is the occurrence, on average, of positive wind stress curl near the coast, and negative curl at some distance offshore. This feature is well developed from May to September. Greater spatial variability is evident in winter distributions.

Existence of an offshore wind stress maximum results in a line of zero wind stress curl approximately parallel to the coast. Positive curl occurs inshore of the maximum wind stress. Negative curl in the offshore region is associated with the anticyclonic atmospheric circulation over the interior ocean. The positive curl near the coast is related to topography and to local features in the surface wind stress distributions.

The distributions from December to March are characterized by positive wind stress curl near the coast from San Francisco to northern Baja California, and south of Punta Eugenia. Large areas of associated Ekman divergence extend several hundred kilometers off the coast. The patterns of wind stress curl are less well behaved near the coasts of Oregon and Washington. However, negative wind stress curl near this coastal region appears to be typical of the distributions during the winter.

During spring and summer upwelling seasons, the dominant patterns of surface wind stress curl are easily recognized. The line of zero wind stress curl parallels the coast approximately 200 to 300 km offshore, along the entire boundary from northern Baja California to Vancouver Island, and from June through September. Yoshida and Mao (1957) placed this line at approximately 500 km from the coast. Considering the coarse resolution (5-degree squares of latitude and longitude) of their data, this disparity is not surprising.

The months of April-May and October-November appear as transitional periods. During transition from spring to summer, the negative curl along the coasts of Oregon and Washington shifts to positive curl. The off-shore distribution takes on a more uniform character. Scattered regions of positive and negative curl are replaced by a large area of negative curl. The late fall transition is marked by a total breakdown in the curl distributions within the northern sector of the grid.

Several local (positive) curl maxima are associated with major topographic changes in the coastline configuration. Large values of positive wind stress curl are found near Cape Blanco, Cape Mendocino, San Francisco, and Point Conception. These features may be real, or they may be artifices of the finite difference calculations or the data distributions. However, one does note that near Point Conception, the values of positive wind stress curl are consistent with a decrease in the magnitude of the surface wind stress in the lee of the point. Where areas of positive wind stress curl extend offshore of capes, there would tend to be a continued, although much reduced, level of upwelling outside of the primary coastal upwelling regime. A region of negative wind stress curl (Ekman convergence) reaches the coast of Baja California between Punta Eugenia and Punta Baja to the north. This feature appears consistently throughout the year. A partial breakdown in this system occurs in August, October, and November. However, considering the probable uncertainties in these derived data, one might reasonably conclude that the coastal region near Punta Eugenia can be characterized in the mean by convergence in the surface wind drift. The distributions of wind stress curl in this area imply favorable conditions for formation of fronts and convergent patches of recently upwelled water.

Coastal Time Series

The mean annual cycle of wind stress curl near the coast is shown in Figure 17. Positive wind stress curl occurs along most of the coast throughout the year. Exceptions to this generalization are found in the region south of lat. 30°N and between lat. 40°N and 47°N. The temporal persistence of the negative wind stress curl near Punta Eugenia (lat. 29°N) is clearly seen in this time series. A similar feature occurs near the tip of Baja California. Seasonal variation between positive and negative curl is apparent from Cape Mendocino to the Columbia River.

Yoshida and Mao (1957) presented evidence indicating that open-ocean upwelling is related to wind stress curl. This process is distinct from coastal upwelling which is primarily a boundary phenomenon. However, the two mechanisms are not totally independent. In regions where positive wind stress curl occurs, such as between lat. 30°N and 40°N throughout the year, and near the coasts of Oregon and Washington during the summer, ascending motion related to the wind-induced divergence offshore enhances the upwelling associated with the more dominant coastal divergence.

DISCUSSION

Monthly distributions of surface wind stress show an offshore maximum which progressively develops over a large extent of the California Current and shifts northward and intensifies from winter to summer. This process is a primary forcing function for coastal upwelling. Off Baja California, wind-induced surface divergence occurs on average throughout the year. Downwelling occurs near the coast of Vancouver Island, except during the restricted period between May and August when a small equatorward component of surface wind stress is observed.

The major seasonal variations in the magnitude and direction of the surface wind stress can be described

Figure 17.—Seasonal cycle of wind stress curl near the coast. The wind stress curl is shown by month for the 1-degree squares immediately adjacent to the coast. The calculations were based on the monthly mean surface wind stress distributions. Units are dyne cm⁻² per 100 km. The contour interval is 0.25 dyne cm⁻² per 100 km. Negative values are shaded. Numbers along ordinate are degrees north latitude.



simply in terms of the general atmospheric circulation over the northeast Pacific Ocean. Monthly mean surface pressure charts typically show two well-developed pressure cells. A high pressure system over the ocean shifts northward and increases in strength from winter to summer. The center of the cell moves in a northwestward direction. This shift in the large-scale anticyclonic circulation results in the observed winter to summer reversal in the alongshore component of wind off the coasts of Oregon and Washington. A low pressure system is situated over the southwestern United States. This semipermanent thermal low is fully developed over the Central Valley in California during the summer. Cyclonic circulation associated with the low leads to equatorward surface wind stress parallel to the coast. The amplitude of the annual cycle is large. During winter, both of these pressure cells weaken. The high pressure system moves southward and the coasts of Oregon and Washington come under the influence of the intense low pressure system in the Gulf of Alaska.

The primary mechanism controlling the location and strength of the wind stress maximum and the associated coastal upwelling is described by seasonal variations in the gradient between the two pressure cells. During the winter, this gradient is weak. Strong heating over the continent during summer deepens the low and increases the amplitude of the onshore-offshore pressure gradient. As a result of the northward shift and strengthening of the high, and deepening of the low, the region of maximum wind stress moves from the area south of Point Conception to the vicinity of Cape Mendocino. These variations are, of course, a function of differential oceancontinent heating related to the annual cycle of solar radiation.

Physical Implications

Climate of the adjacent coastal regions is influenced by upwelling. During summer, the dome of high pressure which develops over the North Pacific Ocean favors large-scale subsidence and a strong temperature inversion over the west coast of the United States. This suppresses deep cloud formation and greatly inhibits precipitation. Unpublished distributions of cloud cover,⁶ summarized from ship observations, show a large onshore-offshore gradient in the total cloud amount. Minima occur at the coast. The effect of the large-scale subsidence is noted in the true desert climate of Baja California, and in the almost complete lack of rainfall along the coasts of California, Oregon, and Washington during the summer. Coastal upwelling primarily influences the local climate of the nearshore zone within 10 to 20 km of the coast and contributes to the formation of low stratus clouds and fog typical of much of the coast along California and Oregon.

A secondary mechanism may account for local intensification of surface wind stress and persistence of coastal upwelling over periods ranging from several weeks to a few months (Bakun 1974). In a simplified positive feedback model, wind stress parallel to the coast brings cold water to the surface and cools the adjacent air. The resulting temperature contrast between the continent and the ocean increases the local pressure gradient. Alongshore surface winds are increased and upwelling is enhanced (Ramage 1971). This mechanism may be slightly modified by the effect of atmospheric stability. In the summertime coastal upwelling zone, the air-sea temperature difference is usually positive. This stable stratification decreases the magnitude of the surface wind stress. The resulting negative feedback may partially offset the increase in surface winds associated with the described changes in the local pressure gradient.

The existence of maximum wind stress some 200 to 300 km from the coast is an interesting feature. Of course, a maximum in the onshore-offshore pressure gradient offshore may explain this phenomenon. Positive feedback associated with wind-induced upwelling extending hundreds of kilometers off the coast may act to intensify the alongshore winds. However, this feature also suggests a coastal boundary layer which acts to frictionally retard the winds near the coast, leading to a positive wind stress curl in the nearshore region.

A characteristic feature of the wind stress curl distributions is the occurrence of a line of zero curl at some distance from the coast. Observations show positive curl inshore of this line and negative curl in the offshore region. A theoretical analysis suggests that a poleward undercurrent along an eastern boundary is favored by positive wind stress curl along the coast and a poleward decrease in surface heating (Pedlosky 1974a). The monthly distributions of wind stress curl presented in this study are generally consistent with an equatorward Sverdrup flow offshore and a poleward Sverdrup flow near the coast, except in the region from Punta Eugenia to Punta Baja, where the wind stress curl is negative. The general pattern of positive wind stress curl along the coast and observations of the California Countercurrent (Wooster and Jones 1970; Wickham 1975) are consistent with Pedlosky's theory.

The Sverdrup transport balance expressed in Equation (6) may provide a simple and reasonable explanation for the existence of the current-countercurrent system observed along the west coast of the United States. Transport calculations, based on the July wind stress curl data for the line of 1-degree squares extending offshore of Cape Blanco (lat. 43°N), show a net southward transport of $3.5 \times 10^{12} \text{ g s}^{-1}$. Within 300 km of the coast, an integrated northward transport of $2.2 \times 10^{12} \text{ g s}^{-1}$ is required. Negative wind stress curl between long. 127°W and 133°W is associated with a southward vertically integrated mass transport of $5.7 \times 10^{12} \text{ g s}^{-1}$. These values underestimate, by a factor of two, the total volume transport of 10 sv (1 sverdrup = $10^6 \text{ m}^3 \text{ s}^{-1}$) for the California Current suggested by Sverdrup et al. (1942:724).

⁶Distributions of monthly mean cloud cover over the California Current are on file at Pacific Environmental Group, National Marine Fisheries Service, NOAA, Monterey, CA 93940.

RAMAGE, C. S.

1971. Monsoon meteorology. Academic Press, N.Y., 296 p.

REID, J. L., JR., G. I. RODEN, and J. G. WYLLIE. 1958. Studies of the California Current system. Calif. Coop. Oceanic Fish. Invest., Prog. Rep., 1 July 1956 to 1 January 1958, p. 27-57.

RODEN, G. I.

1974. Thermohaline structure, fronts, and sea-air energy exchange of the trade wind region east of Hawaii. J. Phys. Oceanogr. 4:168-182.

SAUNDERS, P. M.

1976. On the uncertainty of wind stress curl calculations. J. Mar. Res. 34:155-160.

SMITH, R. L.

1968. Upwelling. Oceanogr. Mar. Biol. Annu. Rev. 6:11-46. SMITH, S. D.

1970. Thrust-anemometer measurements of wind turbulence, Reynolds stress, and drag coefficient over the sea. J. Geophys. Res. 75:6758-6770.

SVERDRUP, H. U.

1938. On the process of upwelling. J. Mar. Res. 1:155-164.

1947. Wind-driven currents in a baroclinic ocean; with application to the equatorial currents of the eastern Pacific. Proc. Natl. Acad. Sci. 33:318-326.

SVERDRUP, H. U., M. W. JOHNSON, and R. H. FLEMING.

1942. The oceans. Their physics, chemistry, and general biology. Prentice Hall, Englewood Cliffs, N.J., 1087 p. VROOMAN, A. M., and P. E. SMITH.

1971. Biomass of the subpopulations of northern anchovy, Engraulis mordax Girard. Calif. Coop. Oceanic Fish. Invest. Rep. 15: 49-51.

1975. Observations of the California countercurrent. J. Mar. Res. 33:325-340.

WILSON, B. W.

1960. Note on surface wind stress over water at low and high wind speeds. J. Geophys. Res. 65:3377-3382.

WOOSTER, W. S., and J. H. JONES.

1970. California undercurrent off northern Baja California. J. Mar. Res. 28:235-250.

WOOSTER, W. S., and J. L. REID, JR.

1963. Eastern boundary currents. In M. N. Hill (editor), The sea, 2:253-280. Interscience Publ., N.Y.

WU, J.

1969. Wind stress and surface roughness at air-sea interface. J. Geophys. Res. 74:444-455.

WYRTKI, K.

1975. El Niño—The dynamic response of the equatorial Pacific Ocean to atmospheric forcing. J. Phys. Oceanogr. 5:572-584.

YOSHIDA, K., and H.-L. MAO.

1957. A theory of upwelling of large horizontal extent. J. Mar. Res. 16:40-54.

YOUNG, H. D.

1962. Statistical treatment of experimental data. McGraw Hill, N.Y., 172 p.

WICKHAM, J. B.

APPENDIX I

Monthly Surface Wind Stress Distributions

The long term composite monthly mean surface wind stress distributions are displayed in Charts 1 to 12 as resultant vectors, in Charts 13 to 24 as the eastward components, and in Charts 25 to 36 as the northward components. The values are plotted in units of dyne cm⁻². The month is indicated in the figure legend in the upper right corner of the charts. The two years displayed below the month, for example 1858-1972, correspond to the year of the earliest report and the year of the most recent report, respectively. The coastline configuration is superimposed on the grid as a visual aid and does not represent a conformal mapping.

In Charts 1 to 12, the vectors are plotted according to the scale shown below the figure legend. The mean vector magnitudes are contoured at intervals of 0.5 dyne $\rm cm^{-2}$. Magnitudes greater than 1.0 dyne $\rm cm^{-2}$ are shaded.

Charts 13 to 24 display contoured values of the eastward component of the resultant surface wind stress. The contour interval is 0.5 dyne cm⁻². Positive values correspond to eastward components. Negative values are shaded.

The northward component of the resultant stress is plotted in Charts 25 to 36. The contour interval is 0.5 dyne cm⁻². Positive values correspond to northward components. Negative values are shaded.



Chart 1.—Resultant surface wind stress vectors for January.



Chart 2.-Resultant surface wind stress vectors for February.



Chart 3.-Resultant surface wind stress vectors for March.



Chart 4.-Resultant surface wind stress vectors for April.



Chart 5.-Resultant surface wind stress vectors for May.



Chart 6.-Resultant surface wind stress vectors for June.


Chart 7.-Resultant surface wind stress vectors for July.



Chart 8.-Resultant surface wind stress vectors for August.



Chart 9.-Resultant surface wind stress vectors for September.



Chart 10.-Resultant surface wind stress vectors for October.



Chart 11.-Resultant surface wind stress vectors for November.



Chart 12.-Resultant surface wind stress vectors for December.

NCC -	TDF-	-11 -	ONE	DEGRE	E SUM	1MARI	ZHIIO	IN			1.00	1.05	101	122	122	121	1.20	110 7118 117 116 115	11/ 113 112 111
137	136	135	134	133	132	131	130	129	128	127	126	125	124	123	122	121	12.0	113 110 11/ 110 115	114 115 112 111
									G	5	2	-~-							
50	Ø.71	Ø.74	0.91	0.56	0.44	Ø.33	0.69	0.21	-0,03-	Ø.58	VANO	COUVER	1SLAND	-1					
19		0.70	0.74	0.60	Ø.74	Ø. 13	0.12	0.27	Ø. 16	0.04	0.02	-2	7	×				NDAR	PIES SERVICE
48			Ø.68	0,55	0.20	0.29	0.48	1.57	Ø.68	10.40	6.1	0.18	1	2	5			PACIFIC ENVIRONMEN MONTEREY, CALI	TAL GROUP
47			-	0.31	0.74	0.22	0.43	0.65	0.30	Ø.62	Ø.26	0.10	-0.26					EAST COMP	ONENT
46				Ø.88	p.39	Ø.69	Ø.67	0.54	Ø. 38	0.59	Ø.3Ø	Ø.26	0.60	> colu	IMBIA F	RIVER		SURFACE S	TRESS
45				Ø.53	0.61	0.58	0.43	0.49	0.54	0.76	Ø.28	Ø.32	0.06					C DYNE CM-	2)
44				Ø.65	Ø949	Ø.67	0.45	0.39	0.82	0.12	Ø.32	0.40	0.03					1000 7504	501
43				Ø.56	Ø.66	19,54	Ø.22	Q.71	0.62	0.20	Ø.33	Ø.31	0.14	CAPE	BLAN	00		FOR	ICHN
42) 			Ø.83	Ø.73	Ø.36	Ø.15	0.23	Ø.82	0.71	0,07	0.20	0.14					JANUAR	
41				Ø.35	Ø.37	0.62	0.88	0,50	0.72	10.41	Ø.47	0.07	0.11					1857 - 19	2
40	J 1 1 1			0.51	Ø. 47	0.35	Ø.36	0.36	Ø.35	Ø.19	Ø.23	0.08	-0.26	CAPE	. MENU	UCINU			
39	 			0.55	0.30	6.)57	0.01	Ø.14	Ø.24	Ø.10	Ø.28	0.24	0.14						
38				-	0.36	Ø.30	Ø.21	Ø.36	Ø.16	Ø.20	Ø.16	Ø.15	Ø. 19	0.02	8				
37						Ø.38	Ø.34	0.20	Ø.36	0.25	Ø.32	Ø.28	Ø.28	Ø.13	2.09				
36						Ø.36	0.27	Ø.31	Ø.29	Ø.28	Ø. 32	0.30	Ø.15	0.21	Ø. 19	1			
35							Ø. 32	Ø.26	Ø.31	Ø.31	Ø.29	Ø.34	Ø.23	Ø.22	Ø.27	0.17	POI	NT CONCEPTION	
34									Ø.28	Ø.18	0.30	Ø.15	Ø.24	Ø.29	Ø.22	Ø.28	0.21	0.23	
33										Ø.19	0.21	Ø.16	0.24	Ø.31	Ø.28	Ø.35	Ø.38	0.36 0.17	
32											0.03	Ø.13	Ø.13	Ø. 15	Ø.26	Ø.34	Ø.33	0.33 0.22 0.11	\sim
31											0.02	Ø.14	0.09	Ø.Ø8	0.28	Ø.29	Ø.26	0.24 0.27 0.18	57
30						- - 						Ø.Ø8	Ø.13	Ø.15	0.06	Ø.15	Ø. 19	0.24 0.26 0.21 0.24	6.1
29			1										-0.17	9.21	0.02	2.03	Ø.16	0.19 0.14 0.24 0.20 0.2	163
28												1	-0.23	-0.10	-3.08	-0.08	Ø.29	0.10 0.09 0.25 0.18 0.2	PUNTA EUGENIA
27											 	1		-0.07	-Ø.22	-Ø.11	10:02	-0.04 0.13 0.14 0.21 0.2	8 8.18 h
26									· · · · · · · ·		 	 			-0.20	-0.23	-0.04	-0.09 0.09 0.07 0.15 0.2	5 0.16 0.20
25						1 1 1 1 1 1					1 				-Ø.18	-Ø.33	-0.22	-0.09-0.15-0.05 0.18 0.5	4 0.15 0.14
24																	-0.27	-0.12-0.02-0.08-0.04 0.0	0 0.16 0.09 0.13 0.09
23					- - 	ļ							 				-Ø. 18	-9.15-9.18-9.07 0.99-9.1	2-0.81 8.84 8.84 8.88
22											 						-Ø.31	-0.32-0.02-0.13-0.03-0.1	3 0,00-0.04-0.02 0.06
21	7 100	1	1	1 122	1 1 2 2	101	100	100	100	107	100	105	104	100	100	101	-0.50	-0.40-0.29-0.23-0.35-0.3	6 0.04-0.01-0.02 0.04
1 13,	100	1 1 3 3	134	+ 100	132	101	130	123	120	121	120	120	124	123	ILL	141	120	113 110 11/ 110 113	114 115 112 111

Chart 13.-East component surface wind stress for January.

137	136	135	134	133	132	131	130	129	128	127	126	125	124	123	122	121	120	1 119 118 117 116 115 114 113 112 111
	10.47	0 68	6 13	0 40	0 36	0 08	a 20	0 01	Chill and	-0 38	VAN		ISLAM	ND				
	10.41		0.15		0.00		0.20	0.01	0.00		2	2	~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~	2				NDBA
49		0.45	0.34	0.17	0.19	10.23	0.15	0.19	0.14	0.13	-0.11	2	-	2	ζ			NATIONAL MARINE FISHERIES SERVICE
48	¦		0.68	0.36	Q.59	2.20	0.17	0.52	0.31	0.06	-0.28	-0.03	1	8	2			MONTEREY, CALIFORNIA
47				Ø.38	0.20	Ø.39	0.68	-(0.39	-0.25	Ø.28	-0701	0.14	-0.26					EAST COMPONENT
46				Ø. 18	Ø.50	Ø. 12	10.59	Ø.28	0.44	Ø.43	Ø.13	0.33	0.07	COL	UMBIA	RIVER		SURFACE STRESS
15				Ø.83	0.32	0.40	0.28	Ø.36	0.51	Ø.33	Ø.27	0.22	0.24	1 7 1				(DYNE CM-2)
14				0.58	8.48	Ø.36	0.41	0.29	0.12	Ø.15	Ø. 27	Ø.29	0.09					
13				0.49	Ø.83	0.39	Ø. ⁶ 51	0.45	0.89	0.56	0.44	Ø. 19	0.04	CAP	E BLAN	NCO		FOR
12				0.35	Ø.74	14.36	9.33	0,49	0.37	0.29	Ø.28	Ø. 17	0.06					FEBRUARY
41				Ø.36	0,48	0.97	0,45	Ø.56	Ø.25	Ø.4Ø	Ø.29	0.08	0.07	5 5 5 6 1				1858 - 1972
40				0.24	Ø. 38	0.68	0.34	Ø.34	Ø.54	Ø.32	Ø.28	Ø. 18	-0.02	CAP	E MEND	DOCINO		in area a sea to a location of
39	j			0.29	Ø. 17	0.38	12.54	Ø.35	0.32	0.32	0.21	Ø.13	Ø. 16					
38					8.49	0.17	0.24	0.20	0.30	0.27	0.25	0.21	0.30	Ø.29	52			
						9.15	Ø. 26	Ø. 26	Ø.33	Ø. 22	0.24	0.30	10.41	0.33	A-21			
			}			0.24	1 22	0.15	0 20	0.22	7 20	0.00	0 17	0 22	5			
						0.34	0.00	0.15	0.20	0.00	0.23	0.22	D. 17	0.22	0.33	1		
							12.36	10.30	0.3/	10.20	0.25	12.19	0.29	0.31	10.21	0.20	PO	INT CONCEPTION
34									0.27	0.29	0.26	Ø.34	0.37	0.42	Ø.41	0.38	0.4	4 0.10
33										0.24	Ø.19	0.20	0.31	Ø.35	0.40	0.47	Ø. 59	3 0.42 0.28
32											Ø.19	Ø. 18	Ø.23	Ø.33	Ø.43	Ø.46	0.40	0 0.50 0.28 0.14
31											-0.00	0.08	Ø.12	0.22	0.36	0.36	Ø.39	9 2.33 0.40 0.26
30												0.20	0.25	Ø. 47	-0,52	Ø.33	0.24	4 0.19 0.27 0.27 0.32
29							11.1						0.10	0.00	0.10	0.05	Ø.11	1 0.21 0.22 0.30 0.38 0.49
28													0.09	Ø.27	0.04	0,01	0.6	2 0.17 0.26 0.13 0.29 0.29 PUNTA EUGENIA
27														Ø.11	0.10	-ø. jz	0.31	1 0.29 0.23 0.11 0.23 0.30 0.26
26				4											-Ø.Ø8	6.05	0.04	4 0.15 0.03 0.11 0.11 0.21 0.24 0.24
25	J														-0.09	-Ø.18	-Ø. 19	9-0:01 0.03 0.05 0.14 0.11 0.21 0.25
24																	-Ø, 15	5-8.17-8.13 0.00-8.00 0.09 0.07 0.21 0.21 0.2
23																	-0.17	7-0, 11-0, 24-0, 08-0, 04 0, 01 0, 00 0, 04 0, 17 0, 11
																	-10.3b	5-3.25-79.33-79.24-74.85-73.82-79.81 0.190 0.06 0.1
107	1 100	105	104	100	100	121	100	100	120	107	100	105	101	100	100	101	-0.35	9-0.32-0.26-0.19-0.06-0.28-0.20-0.20-0.01 0.0

Chart 14.-East component surface wind stress for February.

NCC -	TDF-	-11 -	DNE	DEGRE	E SUI	MARI	ZATIC	IN	100	107	120	1.95	124	122	122	121	120 1	19 118 117	116	115	114 1	113	112	111
137	136	135	134	133	132	131	130	129	128	12/	126	125	124	123	122	121	120 1	115 110 117	110	115	114 1	115	1.1.6.	
									5	5	2	-20	101 011											
ø	Ø. 59	Ø.77	0.34	10.58	B-49	0.43	Ø.41	0.04	0.30	0.01	VANU	COUVER		1			Г						_	
9		Ø. 56	0.73	Ø.66	Ø., 59	9.45	Ø.21	Ø.35	Ø.28	0.16	2.29	R	2	3	-			NATIONAL MA	RINE F	HA ISHER	IES S	ERVI	CE	
.8		1	Ø.64	0.60	0.46	0.50	0.55	0.42	0.30	Ø. 32	0.03	0.04	1	2	3			PACIFIC MONT	ENVIRO	CALIFO	AL GR	OUP		
.7				0.58	0.53	0.38	0.81	0.56	0.41	0.41	Ø.35	0.04	0.12					EAST	CO	MPO	NEN	IT	-	
.6				Ø.67	0.61	0.65	Ø.82	0.62	0.48	0.64	Ø.34	Ø. 38	0.10	COLU	IMBIA R	IVER		SURF	ACE	ST	RES	S		
5				1.10	Ø.42	0.45	0.56	Ø.52	0.66	Ø.29	0.42	0.23	0.34					(DYNE	CM-2	1			
4				Ø.69	0.70	0.61	0.31	Ø.79	0.90	9.43	Ø.31	Ø.33	0.38						ONG TE	RM ME	AN			
3				0.62	0.72	5. 44	Ø.98	Ø. 79	0.37	0.41	Ø.22	0.34	0.33	CAPE	BLANC	:0			FC	R				
2			1	1.00	0.72	6.46	50.55	10.44	Ø.75	0.71)0.41	Ø.23	0.13						MAR	CH				
1			1	0.84	Ø.58	Ø.64	Ø. 55	-\$)12	1102	0.15	Ø.63	Ø.13	0.17	CODE	MEND	00.1100	L	10121	1057	15/2				
.0	1		1	Ø.58	Ø.52	0.52	Ø.75	0.37	0.46	Ø.12	6.6)	Ø.23	-0.12	CHI	- FRENDE	JE LINU								
39				Ø. 65	0.57	0,45	0.42	Ø., 32	Ø.36	0.24	0.34	0.29	Ø.32	-										
8]			0.45	Ø.38	Ø.44	Ø.43	Ø.36	Ø.46	0.42	0.47	0.55	0.49	3									
7					1 1 1 1	DA. 52	Ø.36	Ø.25	Ø.38	Ø.36	Ø.41	Ø.46	0.49	Ø.36	0.33									
36]	9.42	Ø.31	Ø.38	Ø.31	0.37	0.37	Ø.34	Ø.29	Ø.41	0.46	1								
35		1					0.31	Ø.41	Ø.29	0.33	Ø.32	Ø.36	Ø.34	0.40	0.44	0.49	PDIN							
34									Ø.32	Ø.32	0.34	Ø.34	Ø.37	0.44	8.48	Ø.66	0.61	0.20						
33	1 1 1 1]								0.20	0.19	Ø.29	Ø.31	0.58	Ø.65	0.75	0.82	0.63 0.37						
32											Ø. 19	Ø.26	0.20	0.49	0.47	0.61	0.68	0.60 0.39 0.	1	r	~			
31]			<u>.</u>						Ø. 14	Ø. 16	0.26	0,26	Ø.29	Ø.34	0.39	0.56 0.49 0.	38	1		2		
3Ø	1]]			-0.07	6.00	Ø.36	Ø.25	9.32	Ø.45	Ø.33 E.39 Ø.	39 0.42	~	5	, /	1	
29													Ø.12	Ø.23	0.24	0.03	0.27	0.30 0.35 0.	40 0.40	0.69	1	3	a	
28						1]]					-0.09	20.00	0.07	0.07	0.03	0.24 0.19 0.	25 Ø.36	0.41	PUN		ENIR	1
27								1						0.05	40.06	-Ø.Ø9	0,01	0.09 0.42 0.	39 0.27	Ø. 42	0.41	2	r	(
26]					1				1 	1 1 1 1	1			-0.13	-Ø.07	-0.02	0.02-0.02-0.	15 0.26	Ø.34	0.40	0.42	1	1
25			1			1]						-Ø.15	-0.03	0.00	-0.04-0.03-0.	83 0.22	Ø.39	0.48	0.37	4	
24				1	1	1 1 1 1			- - 		1						-0.06-	-0.11 0.02 0-	20 0.07	Ø.16	0.24	0.28	0.37	0.3
23										1							-0.15-	-0.09-0.06-0.	29-0.0	Ø.Ø8	Ø. 15	Ø. 18	0.26	0.3
22					1							1					-Ø. 30-	-0.21-2.15-0.	20-0.61	0.05	0.12	0.14	Ø. 18	0.1
21													1				-0.34-	-0.30-0.13-0.	16-0.18	-0.14	0.00	-0.05	0.03	0.3
137	7 136	6 135	5 134	4 133	3 132	131	130	129	128	127	126	125	124	123	122	121	120	119 118 11	7 116	115	114	113	112	11

Chart 15.-East component surface wind stress for March.



13	7 136	135	134	133	132	131	130	129	128	127	126	125	124	123	122	121	120	119 118 117 116 115 114 113 112 111
									Ś	1	2							
50	Ø.78	Ø.59	0.43	0.44	Ø.27	0.30	Ø.42	Ø.41	0.12	-0.53	VAN	COUVER	15LAN	3				58
49		Ø.62	Ø.63	0.39	0.37	Ø. 17	0.43	0.20	0.29	Ø. 14	0.30	2 cc	2	7	7			NORA NATIONAL MARINE FISHERIES SERVICE
48			0.51	0.60	9.36	Ø. 46	Ø.43	Ø.38	Ø.28	0.33	Ø.16	0.29		2	2			PACIFIC ENVIRONMENTAL GROUP MONTEREY, CALIFORNIA
47				0.66	0.45	Ø.45	Ø. 45	Ø. 15	Ø.5Ø	Ø.26	0,27	Ø.31	0.26					EAST COMPONENT
46				0,83	0.40	0.45	0.55	18,49	0.34	Ø.26	0.22	0.26	0.34	- COL	UMBIA	RIVER		SURFACE STRESS
45				0.43	0.50	Ø.55	Ø.57	0.50	0.51	0.51	0.40	Ø.19	0.25					(DYNE CM-2) 45
44				0.59	Ø.36	0.33	0.42	8.52	0.58	0.58	-0.21	Ø.23	0.08					44
43				Ø. 39	0.21	0.71	Ø.28	0.24	0.53	0.42	0.21	Ø.18	0.05	CAP	E BLAN	со		FOR 43
42				Ø. 36	Ø. 48	Ø.41	Ø.25	0.52	0.46	Ø.26	Ø.38	0.22	0.13					APRIL 42
41				Ø.28	Ø.37	Ø.43	0.21	0.33	0.30	0.50	0.40	Ø.19	0.13					1858 - 1972 41
40				Ø.34	Ø.22	Ø.29	0.30	Ø.43	0.28	Ø.21	Ø.35	0.27	0.13	CRP	E MEND	OCINO		42
39				Ø. 38	Ø.3Ø	Ø.23	0.21	Ø.38	Ø.36	Ø.32	0.31	Ø.38	0.44					39
38					Ø.28	Ø.35	Ø.36	Ø.39	Ø.28	Ø.34	0,43	0.49	0.58	0.74	3			38
37			1.			Ø. 16	Ø.29	Ø.28	0.32	0.40	0.47	0.56	0.70	0.50	0.43			37
36						Ø.31	Ø.37	0.32	Ø.35	Ø.4Ø	0.44	0.14	0.43	10.53	0.55	1		36
35							Ø.29	0.38	Ø.37	Ø. 37	Ø.33	0.40	Ø.33	0.49	Ø.54	0.63	PO	INT CONCEPTION 35
34									0.25	0.30	Ø.24	Ø.46	Ø. 44	8.51	0.56	0.72	0.75	8.20
33										0.27	0.25	Ø.35	0.45	0.57	0.57	Ø.74	Ø.83	3 8.66 8.48
32										ane.	Ø. 25	Ø.33	Ø.33	0.42	Ø.56	0.60	0.78	5 0.59 0.42 0.20 x 32
31							1				0.04	Ø.21	0.27	0.30	0.66	0.51	0.68	5 0.57 0.50 0.39 31
3Ø										-		Ø. 12	Ø.13	0.22	0.40	0.47	0.43	3 8.38 8.49 8.47 8.45
29								1					0.08	Ø. 19	0.29	Ø.23	Ø.43	3 8.38 8.39 8.43 8.69 8.50 29
28													-0.03	0.04	Ø.20	0.22	Ø. 19	0 8.37 8.42 8.33 8.45 8.48 PUNTA EUGENTA 28
27														-0.08	0.04	0.08	0,07	7 0.32 0.25 0.58 0.37 0.49 0.49 2.49
26															-0.09	-0,01	0.27	7 0.07 0.26 0.20 0.44 0.37 0.49 0.52 26
25							1								-0.15	0,00	-0.0	8 8.11 8.18 8.06 8.14 8.27 8.39 8.47
24																	0.01	-8.84 0,08 0.17 8.87 8.25 8.33 8.42 8.43 8.4824
23																	-0.14	-8.10-8.84 8-08 8.85 8.89 8.25 8.19 8.48 8.4323
22							1										-0.22	2-8.83-8.18-8.1 8.18 8.84 8.12 8.25 8.16 8.3522
21									 								-0.26	-0.34-0.10-0.13-0.22-0.02 0.02 0.02 0.02 0.23 0.2621
13	7 136	135	134	133	132	131	130	129	128	127	126	125	124	123	122	121	120	119 118 117 116 115 114 113 112 111

Chart 16.-East component surface wind stress for April.

NCC	- TDF-	-11 -	DNE	DEGRE	E SUN	1MARI	ZATIO	IN	100	107	1.20	105	101	100	122	101	120	110	110	117	116	115	114	113	110	2 111	-
137	136	135	134	133	132	131	130	129	128	127	126	125	124	123	122	121	1210	113	110	11/	110	115	1.14	11.	<u>, 116</u>		1
									à	Z	2	~															l
50	-0.50	Ø.37	Ø. 43	0.30	Ø.32	Ø.21	0.24	0.25	0.25	0.21	VAN	COLVER	ISLAN	4											_	_	N.
49		Ø.31	0.27	0.32	Ø.38	0.29	0.23	0.21	Ø. 17	Ø.33	0.25	s.	1	K				NOT	TONOL	MOR	NDE	RA	RIES	SER	VICE		4
48			Ø.16	Ø.29	Ø.13	Ø.29	Ø.31	Ø.33	Ø.35	0.65	0.35	0.12	1	2	5			net	PACIF	IC E	NVIRO	NMEN	TAL	GROU	P		4
47				Ø. 45	Ø.19	Ø.13	Ø.23	Ø. 15	Ø.24	0.22	Ø.19	Ø.27	Ø. 17						EA	ST	CO	MPC	NE	NT			4
46				0.44	Ø.22	Ø.21	Ø.19	Ø.18	0.25	Ø. 16	Ø.26	Ø.23	0.39	COLI	IMBIA I	RIVER			SU	RFA	ACE	ST	RE	SS			4
45				a) 51	Ø.16	Ø.12	Ø,26	Ø.23	Ø. 38	0.25	Ø.19	Ø.19	0.13							([DYNE	CM-2	2.)				4
44				Ø.35	Ø.26	Ø.21	0.42	Ø.39	Ø.36	0.20	0.21	Ø.21	Ø.15							1.01	-		-				4
43	1			0.37	0.6	Ø.31	Ø.31	0.47	Ø.3Ø	Ø. 37	Ø.16	Ø.13	0.13	CAPI	E BLAN	co				Lur	FO	R	HN				4
42	1			Ø. 32	Ø.36	0.32	Ø. 3Ø	0.04	Ø. 29	Ø.33	Ø.26	0.35	Ø. 19								MA	Y	2				4
41	1			0.20	Ø.22	0.27	0.21	Ø.36	0.03	0.22	Ø.31	Ø.28	0.31		6.6					1	057 -	197,	٢				4
40				0.25	Ø.25	Ø. 17	Ø.32	Ø.44	0.20	Ø.32	0.47	0.44	0.09	CAP	e menu	UCINU											4
39				Ø.19	0.29	Ø. 12	Ø.35	Ø.19	Ø.21	0,30	0.40	0/58	0.78	\													60
38		1			0.22	Ø.23	Ø.28	0.28	Ø.32	0.31	10.44	0.60	Ø. 78	0.71	8												00
37					8 6 6 7	Ø.27	Ø.25	Ø. 38	Ø.31	Ø, 35	0.48	0.59	Ø.75	0.73	0.51												101
36]				0.21	Ø.24	0.24	Ø.28	Ø.37	0.40	0.58	-ð.50	0.55	Ø.69	1											6.0
35			 				Ø.17	Ø.18	0.32	Ø.31	0.40	Ø.46	0.39	9/50	0.56	0.67	POI	NT CON	CEPTIO	N							(0)
34]							Ø.24	Ø.36	Ø.48	6.51	Ø.64	0.60	0.58	0.77	0.81	0.29	5								101
33										Ø. 17	Ø. 19	Ø.32	Ø.34	2,52	0.53	0.86	Ø.97	0.69	0.37	1							111
32											Ø. 19	0.24	Ø.32	0.43	Ø.71	0.63	Ø. 68	0.63	0.45	0.28		Y	~	2			(1)
31											0.11	Ø.14	0.25	Ø.34	0.49	0.60	0,50	0.52	0.51	8.42	7	(١	7			102
30												Ø.13	0.20	0.30	Ø.36	Ø.37	0.44	0.63	0.51	0.53	0.49	~	7		3		(1)
29						1 1 1 1					1 		0.03	0.21	Ø.37	Ø.29	Ø.33	0.35	0.38	0.43	0,51	0.64	1	S	2	5	2
28		1						Y 1 1 1					0.05	0.09	0.18	Ø.13	Ø.27	Ø.31	0.33	Ø.34	0.43	10.58	~	UNTA I	EUGENI	A	2
27						k 1 1 1 1 1 1			1 	 				0.01	0.11	Ø.14	Ø.14	Ø.12	0.20	0.34	Ø. 46	10.52	0.5	er.	1. 4	h	12
26															-0.01	2.03	0.07	0.03	Ø.22	0.09	Ø.28	0.54	0.5	54 Ø.	54	5	AN.
25					ļ							1 1 1 1			-0.05	-0.08	0.01	0.12	Ø. 17	Ø.13	Ø,25	0.20	ø) s	il Br	50 2	2	Y
24			1 1 1 1 					1 1 1 1			1 1 1 1						-Ø.10	0.06	0.00	0.09	0.22	0.18	0.3	30 0.	42 0.4	8 0.5	44
23			1 						1 1 1 1 1 1	1 	1 1 1 1 1 1 1 1						-Ø. 12	2-0.17	-0.24	0.04	0.09	Ø. 10	0.2	2 Ø.	27 0.4	17-0.5	12
22				- - 							 						-0.11	-Ø.22	-0.08	-Ø.04	-0.10	0.09	0.2	29 Ø.	20 0.3	34 Ø.3	82
21		-	-	1	1	1	1	1	1	1		1					-0.34	-0.14	-0.09	-Ø.05	-Ø.11	-0:01	0.1	4 2.	11 0.3	24 0.2	82
13	7 136	5 135	134	133	132	131	130	129	128	127	126	125	124	123	122	121	120	119	118	117	116	115	114	4 11	3 112	2 11	

Chart 17.-East component surface wind stress for May.

NCC - TDF-11 - ONE DEGREE SUMMARIZATION

137	136	135	134	133	132	131	130	129	128	127	126	125	124	123	122	121	120	119	118	117	116	115	11	4 11	3 11	2 11	1
									G	2	2																
50	Ø =51	Ø.31	Ø.35	Ø.35	Ø.31	0.30	0.22	0.40	ø. 16	0.13	VAN	COUVER									1.25		-			_	50
49		0.21	Ø.38	Ø.33	Ø.35	Ø.27	0.32	Ø. 38	Ø.39	Ø.41	0.40	- Ce	2	57	ξ			NR	TIONE	L MAR	NC RINE I	FISHE	RIE	S SE	RVICE		49
48			Ø.39	Ø.46	0.24	0.31	0.79	2.37	0.39	0.41	0.41	Ø.15	1	3	4				FACI	IDNTE	REY,	CALI	FDRM	NIA	UF		48
47				Ø.34	Ø.44	Ø.3Ø	Ø.34	0.42	0.54	Ø.66	0.43	Ø.22	0.12						EF	IST	CO	MP	INC	ENT	-		47
46				Ø.36	Ø.27	0.23	Ø.41	Ø.36	0.43	0.49	0.43	Ø.22	0.22	COL	UMBIA	RIVER			SL	JRF	ACE	S	TRI	ESS	5		46
45]			Ø.29	0.35	Ø.28	Ø.28	Ø. 38	Ø.28	Ø.27	Ø.17	0.20	Ø. 15	, , ,						ſ	DYNE	CM-	2)				45
44]			Ø.33	Ø.25	Ø.31	0.09	0.21	0.05	Ø.28	Ø.33	Ø.13	0.09								NO TO		CON				44
43				0.40	0.29	Ø.28	Ø.22	0.01	0.21	Ø. 15	0.06	Ø. 15	0.06	CAP	E BLAN	ICO				Lu	NG TE	DR	EHN				43
42				Ø. 19	Ø.28	Ø.16	0.24	Ø.27	0.31	Ø. 12	Ø. 19	Ø.23	Ø. 16								JL	INE					42
41				Ø.28	Ø.18	Ø. 19	0.20	0.20	Ø.17	Ø.23	0.28	Ø.23	0.25								1858	- 197	2				41
40				Ø. 16	Ø. 16	Ø.17	Ø.18	Ø.23	Ø.26	0.27	Ø.36	Ø.38	0.23	CAP	E MEND	001100		64 J.	His,							_	40
39	1			0.14	Ø.16	Ø.16	Ø.15	Ø.20	0.30	Ø.31	0.43	0.59	Ø.68														39
38					Ø.13	0.20	Ø.23	Ø.26	Ø.29	Ø.38	0.56	0.67	Ø.77	Ø.77	2												38
37						Ø.13	0.24	Ø.23	Ø.28	0.41	10.49	0.61	0.74	Ø.73	0.49												37
36						0.04	0.06	Ø. 16	0.20	Ø.34	0.43	0.49	2.55	Ø.64	0.68												36
35							0.12	Ø.14	Ø.14	0.27	Ø.42	Ø.42	0.56	Ø.64	Ø. 55	0.65	PO	NT CO	NCEPTI	ИС							35
34									0.07	0.25	Ø.31	Ø.45	0.51	Ø.61	0.72	0.71	0.6	10.20	5								34
33										Ø. 17	0.17	Ø.32	0.41	0.56	0.70	Ø.66	Ø.78	0.55	Ø.25	5							33
32											Ø.16	Ø.26	Ø.31	0.49	0.59	Ø.67	0.59	0.50	Ø.34	0.21							32
31											0.03	Ø. 14	Ø.25	Ø. 26	0.39	0.57	Ø.57	0.58	2.48	Ø. 36	1	{	~	3			31
30												0.11	Ø.13	Ø. 19	0.32	0.34	Ø. 48	Ø.45	Ø.45	Ø. 42	0.4		1		1		30
29													0.02	Ø.14	Ø. 16	Ø.17	0.23	Ø.33	Ø.36	Ø.46	Ø.44	6.54	2	2	5		29
28					1								0.00	0,00	Ø.Ø8	Ø.19	Ø. 10	Ø.32	0.21	0.29	0.40	0.43	2	PUNTO	FLIGEN	1	28
27														-ø.ø1	Ø.1Ø	0.26	Ø.Ø2	Ø.18	Ø.18	0.26	Ø.43	0.52	Nø.	51	1	L	27
26															-0.01	0.01	0.05	Ø.11	Ø.16	0.11	Ø. 33	0.45	ð Ø.	52 10.	49	2	26
25															-0.10	-0.24	-0.01	0.03	Ø.13	0.23	Ø.21	Ø. 38	a los	B1 0.	44	,	25
24																	-0.0	0.04	0.04	Ø.11	0.20	0.29	Ø.:	35 Ø.	42 0.	42 0.	4924
23																	0.00	0200	0.09	0.06	0.04	0.09	Ø.,	25 Ø.	35 Ø.	40 0.	4523
22						 											0.05	-0.09	0.00	0.09	Ø.13	Ø.11	Ø.	17 0.	23 0.	30 0.	4522
21																	-0.20	-0.21	-0.17	-0.07	10.00	0.07	Ø.	11 0.	19 0.	15 0.	2421
13	7 136	135	134	133	132	131	130	129	128	127	126	125	124	123	122	121	120	119	118	117	116	115	11	4 11	3 11	2 11	1

Chart 18.-East component surface wind stress for June.

10	7 120	195	134	133	132	131	130	129	128	127	126	125	124	123	122	121	120	119	118	117	11	6	115	114	113	3 11	2 11	1
13	1.30	100	1.54	100	- OL	101	1	1	4	-	2																	
	tan	1 9 22	0.75	9. 27	R 34	0.48	PL 41	0.41	a.K	0.18	VRINI	COUVER	ISLAN	٥														
	10.3	0.33	0.25	0.3/	0.54	0.40	0.91				2	2	~	2			[NCA	A				٦	
49		12.37	0.30	0.32	0.33	0.45	0,28	0.41	0.45	10.50	0.45	- Ce	-	2 2	7			NRI	TONA	L MA	RIN	E FI	ISHE	RIES	SER	VICE		
48			0.23	0.25	0.31	Ø.35	Ø.34	0.26	0.44	0.59	0.24	0.26	1	6	1				1	IONTI	EREY	· . C	ALIF	ORNI	A			
47			 	Ø.43	Ø.36	Ø.39	0.22	0.36	0.43	Ø.34	0.40	10.23	0.12						EF	IST	C	10:	1PC	NE	NT			
46	-			0.21	0.43	Ø,31	Ø. 3Ø	Ø.26	0.34	0.44	Ø.38	0.20	0.27	- COLI	IMBIA F	RIVER			SL	IRF	AC	E	ST	RE	SS			
45		-		Ø.13	Ø.23	Ø. 12	0.21	Ø. 3Ø	Ø.24	Ø.38	Ø.28	Ø. 12	0.09							(DYN	Æ	CM-2	2)				
44		1		0.22	Ø.19	0,20	Ø.13	Ø.23	Ø.14	0.13	Ø.17	Ø.13	0.26										-	-011				
43				Ø. 18	0.25	Ø.15	Ø.13	0.09	Ø.16	0.08	0.02	0.05	0.08	CAPI	E BLAN	00				U	UNG	FOR	en ne R	HN				
42		1		Ø.19	Ø.18	Ø.14	Ø.13	Ø. 17	0.02	0.08	2.17	0.25	0.32									JUL	_Y					
41		J		0,20	0.09	Ø. 19	0.14	Ø. 10	0.14	Ø.13	40.05	Ø.36	Ø.42					5.4			185	8 -	197	2				
10		1		0.07	0.07	2,08	0.07	Ø. 14	0,21	10,27	0,42	0.45	0.22	CAP	E MENDI	DCINO												
				0.00	01-010		2 29	Ø. 13	0.24	11.28	0.37	6.55	0.69															
				0.00	0.00	0.01	0.0	D. 13	0.10	0.24		0 70	0 79	25	1>													
38			ļ		10.106	print 1	0.04	10.13	0.15	0.34	D.49	0.13	0.78		10													
37				 	i 	►Ø. Ø5	E7/08	Ø.11	Ø. 19	0.26	0.40	12.52	0.65	10.54	0.39													
36						-Ø. Ø1	-Ø.Ø3	0.02	Ø.11	Ø.25	Ø.32	Ø. 48	0.54	0.57	0.54	1												
35							-2.12	-10.102	0.07	Ø. 16	Ø.31	Ø. 32	0.38	0.54	Ø.57	0.51	POI	NT COM	ICEPTI	ON								
34									-0.02	0.18	Ø.18	0.35	0,53	0.59	0.64	0.67	0.54	Ø. 19	2									
33				1						20001	0.10	0.25	Ø.32	0.49	0.63	0.60	10.59	1.47	0.29	1								
32		1		1			1	1	1	1	0.08	Ø. 17	0.21	Ø.32	0.17	1.52	Ø.57	0.51	Ø.38	0.2	5		4					
31											-0.03	10-00	Ø. 11	Ø.18	Ø.26	0.41	10.50	0.42	0.3	0.3	12		(5			
30		1	1						J			-0.02	10.01	0.10	Ø. 12	0.23	Ø. 33	Ø. 16	Ø. 38	B Ø.3	36 0	5		1		1		
20					J = = = = = .				 				l øl øg	0.13	0.20	0,13	0.19	Ø.16	0.3	2 0.3	36 Ø	.36	0.47	1	18	2		
20) J					1		0.03	Ø. 14	0.17	0.19	Ø-1	3 12.3	25 0	.34	0.34	3	1	5	2	
20				į) J = = = = -		j						10.00	Lan	0.05	0.04	0.07	0.10	0 1	2 12 1		24	5	PPP P	UNTA	EUGEN	IA	
21														-10×103	0.01	0.04	10.07	0.06	20.10	0.0		.24	0.37	0.3	5	1	4	
26				1											-0.03	-Ø.14	10.106	-12.191	0.0	9 0.1	18 10	. 15	0.28	0.3	6 Ø.	31	1	5
25										1 1 1 1 1					-0.13	-Ø. Ø4	0.00	@¥00	0.0	3 Ø.1	15 Ø	. 17	0.23	0.3	2 0.	32	5	-
24						ļ	1				· · · · · ·	ļ					-0.12	-0.07	-0.0	1 0.0	05 Ø	. 10	0.23	0.2	1 Ø.	28 Ø	27 0	2
23				-	-	-	-	-				1					-0.15	0.00	-0.0	5-0.1	H Ø	.Ø4	0.09	Ø.1	6 Ø.	15 Ø.	19 0	2
22				8 8 9 9			-		1		-	-					-0.10	-Ø.Ø8	-Ø. 1	8-Ø.	19-10	.02	0.05	0.0	4 0.	Ø4 Ø.	Ø9 Ø	1
21	4 4 4 4 4		1		1			1 0 1		1	1	1	1	1 1 1			-0,29	-Ø.28	-0.2	2-10.1	23-0	.23	-Ø. 18	-0.0	18 0	00 0	00-0	2
1	37 13	6 135	134	133	132	131	130	129	128	127	126	125	124	123	122	121	120	119	118	11	7 1	16	115	114	1 11	3 1	12 1	1

Chart 19.-East component surface wind stress for July.

137	136	135	134	133	132	131	130	129	128	127	126	125	124	123	122	121	120	119 118 117 116 115 114 113 112 111
									2		2							
50	Ø.33	0.44	Ø.48	Ø.46	Ø. 49	10.44	Ø.36	Ø.31	0.25	0.08	VRI	ICOUVE	ISLA					
49		0.43	Ø. 38	Ø. 42	0.40	2.47	0.40	Ø.35	Ø. 43	0.21	Ø. 18	22	1	j k	5			NDRA NATIONAL MARINE FISHERIES SERVICE PACIFIC ENVIRONMENTAL GROUP
40			0.43	0.00	2.00	0.00	.41	2.33	0.00	0.00	0.10		1	6	~			MONTEREY, CALIFORNIA
47				0.33	10.42	12.30	0.22	0.38	0.30	0.34	Ø.32	0.08	0.11		WOLD I			EAST COMPONENT
46				0.48	Ø.35	Ø.35	0.36	Ø.35	Ø.36	0.30	0.34	0.14	Ø. 16		UMBIH	KIVER		SURFACE STRESS
45				Ø.46	Ø.33	Ø.33	0.49	Ø.27	0.27	0.27	0.27	0.09	0.06					(DYNE CM-2)
44				Ø.22	Ø. 29	0.23	Ø.30	0.30	0.29	Ø.15	0.08	0.02	0.04	ļ				LONG TERM MEON
43				Ø.37	0.21	Ø. 19	0.15	Ø.18	0.29	Ø.Ø8	0.08	0.07	0.12	CAP	E BLAN	СО		FOR
42				Ø.19	Ø. 19	Ø.21	Ø . 19	Ø. 13	Ø.1Ø	Ø. 12	Ø. 10	0.21	Ø. 16					AUGUST
41				0.10	Ø. 1Ø	Ø. 15	0.12	Ø. 17	Ø.14	Ø. 18	0.29	0.27	Ø. 12					1857 - 1972
40				Ø. 15	0.03	Ø.14	0.06	Ø.13	Ø. 17	Ø. 18	0.20	0.33	0.19	CAP	e mendi	OCINO		
39				0.03	0.03	Ø.Ø3	0.05	Ø . 12	0.21	Ø.26	0.30	1 19.52	0.59					
38					0.04	0.06	Ø.12	0.11	0.20	Ø.35	Ø.45	0.56	0.56	0.59	2			
37		4				0.02	0.02	Ø.11	Ø.23	0.26	Ø.35	0.52	0.63	9.54	0.37			
36						Ø.Ø6	-0.02	0.01	0.04	0.21	0.33	B.44	0.42	0.54	0.50			
35			 	 			-0.05	0.01	0.06	Ø.15	0.22	Ø.27	Ø.41	0.51	Ø.52	0.49	POI	
34			4						0.07	0.12	10.13	Ø.38	Ø.36	Ø.51	0.51	0.61	Ø.67	1 8.15
33										2.01	0.07	Ø.21	2.29	0.42	0.46	Ø.58	0.74	6 8.45 E.30
32											0.05	17.12	0.26	0.31	RAI	1.53	0.65	954 934 921
31											-0 -07	0 019	01 11	id 15	0 17	9 38		
20											D.D.	0.00	0.11	0.10	G 14	0.00	0. 20	
		}										10107	0.03	0.03	0.14	0.21	0.00	
													-0.92	0.23	6.05	0.15	10.15	5 8.16 9.21 0.30 0.35 0.36 C
)) 						-0.21	-Ø.13	10.04	0.06	0.07	0.20 0.18 0.25 0.29 0.32 PUNTA EUGENIA
27														-Ø. 12	-Ø. 13-	-12.123	0.11	0.13 0.06 0.14 0.22 0.35 0.34
26															-Ø.13-	-Ø.05	-0.03	5 0.01 0.08 0.06 0.12 0.25 0.33 0.36
25															-Ø. 12-	-Ø.23	2.92	- 8.00 C.02-2.02 0.12 0.16 0.26 0.30
24																	-Ø. 40	9-10.21-10.102 10.06 10.09 10.04 10.11 10.24 10.25 10.32
23						 											-Ø. 33	3-0.07-0.27-0.17-0.06 0.18 0.12 0.16 0.09 0.17
22																	-Ø.06	-0.32-0.12-9.12-0.50-0.25-0.82 0.08 0.13 0.13
21																	-2.06	-0.32-0.25-0.09-0.06 0.00-0.17-0.25 0.08 0.11
137	136	135	134	133	132	131	130	129	128	127	126	125	124	123	122	121	120	119 118 117 116 115 114 113 112 111

Chart 20.-East component surface wind stress for August.

	137	136	1.35	134	133	132	131	130	129	128	127	126	125	124	123	122	121	120	119*	118	117	116	115	11/	4 11	3 1	12	111	1
								1	1	9	-	3	~																l
50		Ø.55	9.48	Ø. 53	0.44	0.42	Ø.36	Ø.33	Ø.11	0.05	0.03	VAN	COUVER	ISLAN	0 ~												_	5	Ø
49			0.72	0.62	0.56	0,50	Ø.36	Ø.31	0.34	Ø.30	Ø. 19	0.00	Re all	2	Je -	5			NAT	TONAL	MAR	ND INE F	AA FISHE DNMEN	RIE	s sei Gro	RVIC	E	4	9 8
48				0.59	0.00	10.44	0.32	0.30	-0-	0.00	0.10			1	6	~				м	IDNTE	REY,	CALI	FORM	IIA				7
47					0.38	Ø.29	Ø.29	Ø.39	Ø.49	0.39	0.53	0.37	0.15	0.00		WOTO I				EA	ST	CO	MP	ONE	ENT			ľ	1
46]		Ø. 46	Ø.42	0.47	0.42	Ø.26	0.40	Ø.18	0.64	Ø.19	0.05	CUL	JUBIH I	NIVEN			SU	RF	ACE	S	TRE	ESS	ò		4	0
45					0.31	Ø.47	0.21	Ø.29	2,56	Ø.37	Ø.32	Ø.34	0.07	0.10							(DYNE	CM-	2)				4	- Un
44					Ø.28	0.33	Ø.17	Ø.16	Ø. 19	Ø.19	Ø.39	Ø.23	0.03	0.07							10	NC TE	0N M	FON				4	4
43	1	1			0.33	Ø.18	0.06	Ø.11	Ø.24	0.27	2.00	0.02	0.00	0.01	CRPI	E BLAN	60					F	R	LINA				4	3
42					Ø.19	Ø.23	Ø.11	Ø.29	Ø.29	0.00	0.12	0.05	Ø.11	-0.04							SE	EPTE	MBE	R				4	2
41					0.05	0.06	0.09	0.10	Ø.18	Ø.16	-0.10	0.14	Ø.13	0.09								1854	- 197	2				4	11
40					0.04	0.06	70701	0.22	10.14	0.03	0.11	Ø.13	0.28	0.17	CAPI	E MENDO	OCINO											4	Ø
39					-Ø.12	-0.02	0.00	0,00	0.02	0.10	0.10	Ø.15	Ø.38	0.3														0	19
38					<u>Landa</u>	0.02	0.00	0.03	0.09	0.06	Ø.17	0.24	Ø.39	0.47	0.31	R												(1)	18
37							10.00	0_01	0.06	Ø. 12	0.22	0.28	Ø.39	Ø.46	2.41	Ø. 33												10	57
36							-e. 02	-0.10	0.03	Ø.Ø6	Ø.19	Ø.23	0.27	0.29	Ø.38	0.41												101	36
35						 		1-0.07	10.04	0.06	Ø. 11	Ø.17	0.30	Ø.27	0.35	Ø. 42	0.39	P01	NT. CON	CEPTIC	IN							9	S
34									_	0.03	0.09	0.25	10.22	0.35	0.39	0.44	Ø.59	0.51	0.20									102	34
30											-0.03	0.10	Ø. 18	1.22	1.32	0.40	0.66	Ø. 66	1.45	8.29	1								33
20			 		 							0 03	0 14	G 19	13.32		a 18	0.54	7 49	0.35	R 19								2
34												0.05	0.14	0,10	0.52	0.00	0.40	0.04	0.40	0.00	0.77	1	1	~	2				54
31						ļ						10.01	0.05	10 . 10	10.20	10.29	10.43	0.41	10.31	0.3/	10, 32	7		1		1			1.
30													0.05	10.104	0.21	0.22	0.27	0.32	0.34	0.35	10.34	9.34	~	7	10	1		1	80
29														0.01	0.01	0.15	0.09	Ø, 16	0.27	Ø.16	0.31	0.36	0.5	1	2	5	Z	. 2	27
28						1 1 1 1 1 1								-9.95	0.04	0.05	Ø. 12	0.14	0.27	0.22	Ø. 17	Ø.28	0.3	Z	PUNTA	EUGE	NIA	4	8
27						1 1 1 1									-10.09	-0.14	Ø.13	Ø.11	Ø. 15	0.21	Ø.16	Ø.24	0.3	7 0.	33	2	P		7
26								 								-0.06	9-02	0.04	Ø.10	Ø.18	0.15	8.22	Ø.2	2 Ø.:	34 Ø.	32	1	1	6
25																-Ø. 14	-Ø.24	0.400	0.00	0.10	0.25	0.37	Ø.1	7 Ø.	26 Ø	28	L	1	S
24				1														-0.10	0.00	0.08	Ø. 12	9.10	0.2	Ø Ø.:	29 Ø	28	3.23	0.26	4
23																		-Ø.18	-Ø.16	-0.09	Ø.22	0.06	Ø. 1	5 Ø.	14 0.	26	0.25	0.172	20 -
22																		-0.05	0. 28	0.00	-0.25	0.39	Ø.3	3 0.1	Ø6 Ø.	. 19	8.22	0.142	2
21								1					1					-0.03	-0.03	-0.64	0.18	0.07	0.0	B Ø.	40 3	16	8.30	0.002	1
L	137	136	135	134	133	132	131	130	129	128	127	126	125	124	123	122	121	120	119	118	117	116	115	11	4 1	13 1	12	111	

Chart 21.-East component surface wind stress for September.



137	136	135	134	133	132	131	130	129	128	127	126	125	124	123	122	121	120	119	118	117	116	115	5 114	1 113	3 112	111
	-								G		3															
50	0.70	0.60	Ø.67	0.71	0.57	Ø.37	0.22	Ø.13	0.06	Ø. 39	VAN	COUVER	ISLA													S
49		0.39	Ø.58	Ø . 51	0.52	Ø.34	Ø.27	Ø. 19	0.28	-0.14	-0.20	es.	2	J'Z	7			NF	TIONF	AL MA	N RINE	DAR FISH	ERIES	SER	/ICE	4
48			1.1	0.81	Ø.75	10.41	0.45	0.28	Ø.38	Ø . 13	Ø.18	-0.10	5	2	2				PACI	IFIC MONTE	ENVIF	CALI	NTAL FORN	GROUP	2	4
47				0.67	0.94	0.78	1.)6	0.42	Ø.45	Ø.28	0.20	0.07	-Ø. 13						EF	AST	CC	MP	ONE	NT		4
46				Ø.75	Ø.56	Ø.92	Ø.79	0.53	0.33	0.54	0.28	0.23	-0.05	> col	UMBIA	RIVER		200	SL	JRF	ACE	S	TRE	SS		4
45				0.44	0.62	10.44	0.63	0.72	0.57	0.60	0.46	Ø.12	0.06							ſ	DYNE	CM-	2]			4
44				Ø.73	0.45	0.59	Ø.39	Ø.34	Ø.45	Ø.15	0.09	0.08	0.05										EON			4
43				Ø.63	0.77	0.59	Ø.33	Ø.29	0.30	0.21	Ø.13	(-Ø. 13	0.00	CAF	PE BLAN	IC0				u	F	OR	IEHN			4
42				Ø. 42	0.32	0.20	Ø.38	Øg51	0.21	Ø.19	0.07	0.05	0.02								OCTI	DBEF	3			4
41				0.21	Ø. 1Ø	Ø.27	0.26	Ø.39	Ø.17	Ø.11	Ø. 15	0.00	0.09								1854	- 197	12			4
40				Ø.14	Ø.44	Ø.26	Ø . 15	Ø . 16	Ø.23	0.20	0.21	0.10	6.03		'E MENU	IULINU										4
39				0.20	Ø.21	0.22	Ø.24	0.20	Ø.Ø8	Ø. 17	<i>u</i> .Ø4	0.22	Ø.31													3
38					Ø.37	0.27	Ø.24	0.09	Ø.17	Ø.19	Ø. 19	0.31	Ø.33	0.26	3											3
37		1				Ø. 18	0.04	Ø.16	Ø.14	Ø.21	0.23	Ø.35	0.34	Ø.38	0.22											3
36						0.06	70.02	0.05	Ø.Ø8	Ø.17	Ø. 19	Ø. 19	0.22	Ø.3Ø	10.32											3
35		1					0.01	Ø.11	0.03	Ø.12	0.09	Ø.16	0.27	0.24	Ø. 38	0.34	PO	INT CO	NCEPTI	DN						3
34									Ø.14	0.03	Ø.13	Ø. 17	0.24	0.34	0.54	0.51	Ø.39	0.11	2							3
33		1								Ø.11	Ø.11	0.21	0.23	0.26	2.42	0.45	0.47	Ø.40	0.23	5						3.
32		1									Ø.Ø6	Ø.14	Ø. 16	Ø.32	Ø.31	Ø.39	0-51	Ø. 42	Ø.37	0.22		5	-			3
31											0.01	0.08	Ø. 17	0.26	0.23	0.31	Ø.31	0.39	Ø.37	Ø.26	1	1	1	7		3
30									-			0.03	Ø. 10	0.10	Ø.25	Ø. 25	Ø.27	0.31	0.27	Ø.30	0.3	Ĺ	1		5	30
29													0.07	0.07	Ø. 12	Ø.13	Ø. 28	0.17	Ø.28	Ø.31	Ø.34	0.3	1	2	à	29
28													0.03	-0.03	Ø. 14	0.07	Ø. 10	0.11	Ø.10	0.24	Ø.29	0.32	PL	INTA EL	JGENIA	28
27														-0.03	Ø. 62	0.10	0.03	Ø. 10	0.09	-0.2)	Ø.29	0.35	0.3	in	7	2
26															-0.07	-0.03	10.01	Ø.12	0.07	0.11	Ø. 19	0.28	Ø.30	6 Ø.3	7	2 26
25															-Ø.13	-Ø.12	0.04	0-00	0.05	Ø. 12	0.09	0.22	0.43	3 Ø.30	1	25
24		1															-Ø.15	-Ø.05	-0.01	0.02	Ø.Ø8	Ø. 12	0.1	0.23	Ø.26	10.2824
23				-						1							-0.23	0.00	-0.04	0.00	-0.9	Ø. 12	0.17	0.14	Ø.22	Ø.2423
22																	-Ø. 17	-Ø.11	-0.14	0.00	0.03	-8.02	0.00	5 Ø. 10	0.09	Ø. 1922
21		1								1							-Ø.25	-Ø.20	-0.03	-Ø.Ø2	-0.08	0.04	0.05	0.07	0.07	-0.0321
137	136	135	134	133	132	131	130	129	128	127	126	125	124	123	122	121	120	119	118	117	116	115	114	113	112	111

Chart 22.-East component surface wind stress for October.

NC	- 32	TDF-	11 -	DNE	DEGRE	E SUP	1714141.	120	120	128	127	126	125	124	123	122	121	120 119 118 117 116 115 114 113 112 111
1	.37	136	135	134	133	132	131	130	129	120	161	120	125	164	120	h fai ha	1 60 1	
-			1							S	5	Za						
50		Ø. 73	0.66	Ø.71	0.72	0.60	Ø.37	Ø.24	0.28	9.05	3.06	VANC	OUVER	1SLAND	1			50
19			0.81	0.73	0.57	0.48	0.40	Ø.34	Ø.28	0.01	0.00	0.00	5	1	×			NOAA 49
-									0 20	0 20	0 43	a anst	0 16	-	5	}		PACIFIC ENVIRONMENTAL CROUP 48
48 -				N. 80	0.75	Ø.55	0.13	640	0.20	0.23	0.45		1	1000	6	~		MONTEREY, CALIFORNIA
47	1				0.96	0.52	0.70	Ø.95	\$.4Y	0.63	0.23	0.35	0.13-	0.27				EAST COMPONENT
46	1				10.98	Ø.48	0.80	0.40	0.32	Ø.73	0.49	1010	0.17	0.19	COLU	MBIR F	RIVER	SURFACE STRESS 46
45					Ø.79	0.56	0.30	0.54	0.55	0.55	0.64	10.33	Ø.22	0.05				(DYNE CM-2) 45
44					0.65	0.76	p. 1	Ø. 92	Ø.92	0.54	0.45	Ø.29	Ø.10	0.01				4.4
43		i			0.83	0.59	0.57	0.67	0.57	0.65	0.01	0.23	Ø.13	0.09	CRPE	BLAN	00	LONG TERM MEAN FOR 43
-					0 55	0.67	Ø.58	0.73	9 AB	R. 27	0.35	0.21	-0(01	0.10				NOVEMBER 42
42 -					a 22			0.52		G 14	a051	11 22	alad	1 00				1854 - 1972
41 -					10.72	10.73	0.65	0.5/	967.0	0.14	0151	0.22		3	CAPE	MEND	OK I NO	
40 -					Ø.58	0.58	Ø.36	0.48	0.99	0.21	0.05	0.26	0.03	10.15				
39	1				0.74	0,51	2 51	Ø.41	Ø.26	0.25	0.42	Ø.14	Ø.15	0.03	1			39
38						Ø. 39	0.40	Ø.39	Ø.31	0.34	0.33	Ø.36	Ø.18	Ø.13	0.11	3		38
37							Ø.28	Ø.39	Ø.29	0.28	Ø.26	Ø.32	0.26	Ø.27	0.20	0.12		37
36							0.53	0,32	Ø.29	Ø.25	0.25	Ø.21	0.23	Ø. 15	0.25	0.20	1	36
35					1		1	0.22	Ø.25	0.21	0.23	Ø.19	Ø.27	Ø.18	0.20	0.27	0.19	POINT CONCEPTION 35
34										Ø.14	0.27	Ø.19	0.26	Ø.26	Ø. 3Ø	0.30	Ø.39	9 8.33 8.88
33				1 1 1 1	1	1 1 1		1			0.14	Ø.13	Ø.14	0.22	Ø.31	0.32	Ø. 47	7 8.45 8.32 8.21
32							 					0.09	Ø.17	Ø.18	0.32	0.28	Ø.26	6 8.34 8.36 8.26 8.1
31												0.08	0.09	0.06	0.13	Ø.33	0.21	1 0.33 0.38 0.30 0.21
30									l				Ø. 16	0.02	0.06	0.20	0.10	8 8.34 8.26 8.25 8.27 8.25
29							·							-0.04	0.02	.03	0:52	2 8.87 8.13 8.21 8.22 8.25 8.18 18 28
28				1 2										-9.97	0.05	10.03	0-00	8 8.85 8.89 8.17 8.18 8.24 8.25
20				j											Lad		-01 010	
21															0.0(2		
26																10.108		
25													 			-Ø. 13	-0.15	5-0.09-0.08 8.02-0.04 0.09 0.00 0.18 0.20
24																		-0.12-0.17-0.15 0.01-0.00 0.00 0.01 0.16 0.19 0.1924
23																		-0.39-0.29-0.08-0.22-0.05-0.03 0.09 0.03 0.13 0.1523
22]	1							-0.33-0.41-0.07-0.20-0.13-0.09-0.84 0.05 0.09 0.0722
21			1				-	-	-	1								-0.42-0.23-0.20-0.09-0.12-0.02-0.06-0.05-0.05-0.06-0.0921
L	137	136	135	134	133	132	131	130	129	128	127	126	125	124	123	122	121	120 119 118 117 116 115 114 113 112 111

Chart 23.-East component surface wind stress for November.



Chart 24.-East component surface wind stress for December.



Chart 25.-North component surface wind stress for January.



Chart 26.-North component surface wind stress for February.

	137	136	135 1	34	133	132	131	130	129	128	127	126	125	124	123	122 121	0 119 118 117 116 115 1	14 113 112 111
		6 1- 1-	1	1						5	5	2						
50	F	0.20	0.05-0	.04	0.15	0.02	0.01	Ø.24	Øv53	0.06	0.35	VRN	COUVER	ISLAN	0			
49			Ø.05 Ø	. 10	Ø. 19	0.07	Ø.36	Ø.4Ø	0.14	0.20	Ø.33	0.40	Te	2	2	7	NDAA NATIONAL MARINE FISHER POCIFIC ENVIRONMENT	IES SERVICE
48			-0	1.12-	0.05	Ø.15	0.22	Ø.19	Ø.19	Ø. 16	Ø. 19	Ø.26	0.55	1	Z	1	MONTEREY, CALIFO	RNIA
47]			0.09	Ø. 11	0.05	Ø.47	0.10	Ø.26	Ø.38	0.50	Ø.28	0.14			NORTH COMP	DNENT
46					Ø.16	Ø. 15	13.07	-0.04	-0,03	0.04	Ø.13	Ø.22	0.49	Ø.75	> COLI	IMBIA RIVER	SURFACE STI	RESS ⁴⁶
45	1				0.03	0.05	-0.00	10.02	Ø.18	Ø.36	Ø. 38	0.43	0.70	0.34			(DYNE CM-2) 45
4.4		1			Ø. 18	-2.39	-0,20	0.09	Ø.19	Ø. 16	-2.27	0,02	Ø.36	0.45			LOND TEDM MED	44
43					Ø. 10	-Ø.25	-Ø.26	0.20	Ø. 31	Ø. 17	0.05	Ø.14	Ø.35	0.07	CAPI	BLANCO	FOR	43
42					Ø. 17	-Ø.20	-Ø.ØG	-0.08	-0.41	-0.32	0.32	Ø.44	-0.10	-0.04			MARCH	42
41					Ø.17	-Ø.14	-Ø.12	-Ø.14	-0 53	-10.71	ø.ei	0.20	-Ø. 02	-Ø.11			1857 - 1972	41
40					Ø.37	-0.20	-0.30	0.21)ø. 14	-0.48	-Ø.26	-0.55	-Ø.19	-0.20	CAP	E MENDOCINO		42
39		L			Ø.41	-Ø.38	-Ø.24	-0.26	-Ø.39	-0-51	-Ø.56	-0.55	0.47	=0.48				39
38						-0.13	-0,45	-Ø.15	-Ø.18	-Ø.27	-Ø. 46	-0.30	10.62	-Ø.67	-0.33	8		38
37							-Ø, 43	-0.33	9-0.27	-0,51	-Ø.55	-0.53	-Ø.57	-0.57	E. 44	-0.32		37
36							Ø. 38	-Ø.40	Í-Ø₊44	-0.98	-0.53	40.47	-0 .50	-ø. 3}	-Ø.68	-0.65		36
35								-0.38	9-0.34	1-10.57	-Ø.36	-0.93	-Ø.41	-0.53	-0.39	-2.69-2.56	POINT CONCEPTION	35
34								1	1	-0.52	-g_48	9-0.39	-0.52	-0.48	0.61	-0.74-0.81	48-0.07	34
33											-Ø.64	-Ø.62	-Ø.69	-Ø.76	-1681	-0.89-0.94	.70-0.35-0.14	33
32				1			 					-0.60	-Ø.62	-0.70	-Ø.88	-0.96-0.97	.78-0.54-0.25-0.18	32
31												-0.51	D. 48	-Ø.56	-3:49	-0.63-0.79	.60-0.67-0.53-0.40	31
30								-				}	-Ø.56	-Ø.59	-Ø.87	-0.84-0-52	.53-0.62-0.55-0.50-0.53	30
29								1					1	-0.61	10 745	-0.86(-0.1	.77-0.93-0.50-0.59-0.59-0.69	10 2 20
28														-0.71	-Ø.61	-0.69-0.6	.61-0.57-0.53-0.51-0.58-0.60	PUNTA EUGENIA
27				1						, , , , , ,					-Ø.63	-0.52-0.75	.55-0.58-0.92-0.58-0.63-0.59-	0.42 h 2
26	5															-0.49-0.6	A9-0.55-0.64-0.79-0.81-0.62-	0.55-0 50 20
25	5															-0.35-0.4	. 41-0 49-0. 79-0. 56-0. 75-0. 58-	0.64-0.49 2 25
24																	. 48-0. 46-0. 50-0. 63-0. 70-0. 61-	0.56-0.48-0.43-0.48
23	3										1						.31-0,49-0.40-0.48-0.55-0.55-	0.58-0.52-0.40-0.442
22		 				1				1	 			 	 		.55-0.55-0.31-0.58-0.51-0.56-	0.64-0.53-0.54-0.3822
21									1								. 40-0. 49-0. 33-0. 52-0. 67-0. 48-	0.50-0.55-0.46-0.4021
	137	136	135	134	133	132	131	130	129	128	127	126	125	124	123	122 121	20 119 118 117 116 115	114 113 112 111

Chart 27.-North component surface wind stress for March.



Chart 28.-North component surface wind stress for April.

NLL 13	7 136	135	134	133 1	132 1	31	130	129 1	128 1	27 1	126	125	124	123	122 1	21	120	119 118 117 116 115 114 113 112 11	5
	1 130	100	104	100	1		1		3	e	34								
50	10.68	0.23	0.02	0.17	0.08-	0.01-	0.02-	Ø. Ø9-	0.26-1	0.02	VAND	OUVER	ISLAN	1					50
49	Dies	Ø. 16	Ø.13 Ø.Ø7	Ø. 03	0.02 0.03	0.03	0.12-	Ø.05 Ø.08	0.17) 0.04-1	0.10 0.49-	0.00	2	100	N.S.	5			NDAA NATIONAL MARINE FISHERIES SERVICE PACIFIC ENVIRONMENTAL CROUP MONTEREY, CALIFORNIA	49 48
47				0.60-	0.10	0,01	B, Ø1-	Ø. 11-	Ø.11	0.00	a. 63-	Ø. 22	0.03					NORTH COMPONENT	47
46				+Ø. 16-	Ø. 18	0,00	0.01	0.06	0.01-	Ø. 11-	0.32	-Ø.27	-0.12	► COLL	MBIA R	IVER		SUBFACE STRESS	46
45				Ø.Ø8	0.03	9.00	-Ø. 12-	0.14	Ø. 16)	Ø.1Ø-	0.21-	-Ø. 38	-0.20					(DYNE CM-2)	45
11				-0.04	0-00-	Ø.11-	-2.19-	Ø. 12-	0.27-	Ø.22-	0.27	10.00	-0.15						44
13				a.a.	Ø.25-	Ø. 39 ⁻	-0.24-	-0.23-	0.42-	0.26-	0.14	-0.87	6.29	CAPS	FIL FILL	0		LONG TERM MEAN	43
40				a 12	0.24-	0 22	7 59	a 82	0 28-	0 45	D. F.A	A	-0.73	Grute				MAY	42
42				-Øx 12-	0.22	a 20	2.30	DE CZ	0.20	a 70	3.74	-0 92	-0 84					1857 - 1972	41
41				-10×21-	10.22-	0.20	-10.24	2 24	0.02	0.70	1.14	1.14	6	CAPE	MENDO	CINO			10
40				-Ø.16-	0.20-	Ø. 47	-0.43	10.31	0.64	Øx 76-	1,00	-1.14	3						20
39				-0,26-	0.39-	0.22	-0.28	-0/49-	-Ø, 78-	0.90-	1100	-1.19	-1.29	5	_				00
38					Ø. 43-	0.37	-0.48	-Ø.66-	0.61+	0.86-	1.05	-1,06	-1. [1	-0.57	8				30
37						0.42	-ø.si	-Ø. 73-	0.80-	0.92-	0.98	-1.06	-1.13	-0.88	-0.59				37
36						Ø _x 57	Ø.61	-Ø. 69-	0.68	Ø.85	0.86	-1/01	-Ø.79	-0.88	0.92	1			36
35							-0.55	Ø.64	-Ø.70-	Ø.91-	Ø. 99	-0.90	-0.80	Ø.89	-0.80-	0.77	PO	INT CONCEPTION	35
34									-0.79-	0.63	0.94	-15201	-0.97	-1,99	-Ø.85-	V. 88-	-0.56	5-0.00	34
33									ŀ	0.90	Ø.71	-Ø.89	-0.96	-1.05	-0.89-	0.95	-0.77	18.34-0.10	33
32											-Ø. 86	-Ø.95	-0.93	-1.04	1.28-	Ø.77-	-0.71	1-0 49-0.24-0.15	32
31	1				1						-0.70	-Ø.79	-2.84	-0.85	-0.76-	Ø.83	-2.69	9-8.58-9.45-8.48	31
30												-Ø.81	-0.96	-Ø.82	-0.70	Ø.74	-0.76	6-0.70-0.58-0.57-0.56	30
29													-0.72	-Ø.73	-0.62	Ø.61	-0.60	8-9.70-0.58-0.62-0.63-0.63	29
28													-0.62	-Ø.67	-Ø.83-	Ø.54	-0.74	4-0.55-0.58-0.58-0.57-0.50 PUNTA EUGENIA	-28
27		1	d	4 4										-0.60	-Ø.66-	Ø. 68	-Ø.72	2-0.56 049-0.70-0.77-0.51-0.59	27
26			J												-0.70-	0.63	-0.70	8-8.69-8.68-8.49-8.78-8.88-8.58-8.51	28
25					 										-0.46-	Ø.58	-0.58	8-0.72-0.78-0.57-0.67-0.30-0.68-0.51	125
24			J												<u> </u>		-0251	0-0.71-0.69-0.58-0.89-0.65-0.51-0.56-0.49-0.	4424
23												 					-0. 4	5-20, 51-0, 60-0, 77-0, 69-0, 82-0, 63-0, 59-0, 58-0.	472
22																	-9 4	8-9.72-9.47-9.69-9.69-9.71-9.74-9.61-9.69-9	5022
21			1		1		 										-0.40		45 21
	37 136	135	134	133	132	131	130	129	128	127	126	125	124	123	122	121	120		1

Chart 29.—North component surface wind stress for May.

137	136 135	134	133 1	132	131	130	129	128	127	126	125	124	123	122 12	21 12	0 119 118 117 116 115 114 113 112
157	1 1 1	104	100 1	102	101	150	120	120		3	12.5	124	120	166 16		<u>0 110 110 117 110 110 114 110 112</u>
	0 00 0 03	G 10	7 00	0 01		0.00	-0 25	5	0 20	VAN	COUVE	R ISLA	ND			
	0.00 0.05		0.05		0.00					2	5	~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~	2			
8	Ø. 13	0.06 0.05	Ø. Ø8-	0.06 0.10	0.02	0.07	-Ø.23 -Ø.29	-Ø.29 -Ø.14	-Ø.31	-9.19 -0.23	0.01	-	3	5		NUHH NATIONAL MARINE FISHERIES SERVICE PACIFIC ENVIRONMENTAL GRDUP MONTEREY, CALIFORNIA
7			0.07	1.60	0.08	0,04	-Ø.26	-Ø.37	-6.54	-0.21	-Ø.11	-Ø.05				NORTH COMPONENT
6			Ø.05-I	Ø.14-	Ø.05-	0.20	-Ø. 13	-Ø.33	-Ø.36	-1254	-Ø.23	-0.00	COL	JMBIA RIV	/ER	SURFACE STRESS
5		ŀ	@.@4~	Ø. 17-	Ø.27	Ø\$Ø2	-0.30	Ø.43	-Ø.29	-0.23	-Ø.37	-0.24	}			(DYNE CM-2)
1			Ø. 17-1	0.31-	0.25	0.77	70.19	Ø.15	-Ø.33	- 0-4 9	-10-56	-0.24				
3			Ø. 19-1	Ø.43-	Ø.34-	0.2	-0.81	-0.61	-0.63	-Ø.78	-0.73	-0.27	CAP	E BLANCO		LONG TERM MEAN FDR
2			-Ø.28-1	0.26-	0.42-	0.68	-Ø.64	-Ø.89	1.02	-0.75	-1.21	-0.55				JUNE
1			0.24-1	0.26-	0.43	2 ,49	-Ø.61	-0.70	1.17	-1.07	-1.18	-0.74				1858 - 1972
8			Ø.31-i	Ø.32-	0.43-	9.83	-Ø.62	-1.03	() 96	-(.62	×1.34	6.97	CRP	E MENDOCI	(NO	
			-Ø. 41-I	Ø.51-	Ø.50-	0.53	-Ø. 79	-0.91	-1.19	-1.30	1.59	-1.18				
				0.54	-01-77-	0.71	-Ø.88	-0.94	-1 - 13	-1.23	-1.37	-1.23	-9.67	~		
					-Ø 66-	-07.71	-0.80	-1 03	-1 013	-1 15	-1 25	-1 15	-01 87	A 53		
					-01 158-		-0.87	1 10	-1 11	-1 15	-1 10	เ่–ตานต	-1 86	L IN		
]					1	0.74		4.00		4 4 19			1.60		3	
						10.74	-0*09	1.05	70.00	-1.1/	1.106	-1.14	-1.180	-0.00-0.	"L	OINT CONCEPTION
+								-0.93	1	-Topi	-1.04	-1.62	** L = 1 L	-1.03-0.	85-0.	20-10-10-20
5									-0.87	-0.91	1.05	-1.02	-1.01	-1.07-0.	.85-0.1	54-0.27-0.05
										-0.87	-1-22	-0.94	-1.100	-0.93-0.	80-0.6	52-0,48-0,15-0,00
										-0.81	-0.85	i-1.02	-0.93	-Ø.80-0.	82-0.	71-9.61-90.46-90.35
ð 											-Ø.79	-Ø.70	-Ø.85	-0.70-0.	.57-0.0	55-0.66-0.60-0.47-0.48
												-0.60	-Ø.81	-Ø.59-Ø.	61,-0.6	52-10.62-10.62-10.55-10.60 B
3	ļ											-Ø. 53	-0.57	-Ø.66-Ø.	59-0.8	51-0.58-0.53-0.44-0.54-0.53 PUNTA EUGENTA
7	ļ												-Ø.65	-0.58-0.	45-0.3	37-0.53-0.58-0.55-0.69-0.50-0.50
6														-0.42-0.	50-0.	43-9.51-0.58-0.55-0.67-0.69-0.53-0.46
5														-0.47-0.	47-0.	18-0.54-0.44-0.56-0.52-0.54-0.60-0.43
4														1	-Ø.	18-0.55-0.39-0.61-0.53-0.53-0.53-0.47-0.38-1
3															-Ø.3	34-0.54-0.62-0.55-0.40-0.44-0.48-0.50-0.50-
2												1 1 1			-Ø.3	39-0.40-0.44-0.41-0.46-0.58-0.52-0.37-0.44-1
	JJ														H:+:H	

Chart 30.-North component surface wind stress for June.

13	7 136 135 134 133 132 131 130 129 128 127 126 125 124 123 122 121 120 119 118 117 116 115 114 113 112 111
	and a second sec
0	
9	-0.01-0.05-0.01-0.13-0.01-0.19-0.25-0.35-0.29-0.20-0.20-0.20-0.20-0.20-0.20-0.20
8	-6.04-6.03-6.17-6.24-6.17-9.30 PLST 9.45 0.11 PLS
7	-0, 16-0, 35-0, 23-0, 46-0, 18-0, 37-0, 38-0, 36-0, 24-0, 05 NORTH COMPONENT
6	-0.11-0.48-0.19-0.32-0.38-0.38-0.62-0.62-0.41-0.27 SURFACE STRESS
5	-0.26-0.26-0.49-0.44-0.51-0.56-0.54-0.67-0.53-0.23 (DYNE CM-2)
4	-0.21-0.37-0.44-0.59-0.44-0.58-0.81-0.72-0.59-0.35
3	-Ø.18-0.23-0.47-0.81-0.69-0.89-0.90-1.18-0.81-0.67 CAPE BLANCO FDR
2	-8.39-0.51-0.57-0.51-0.95-1.09-0.95-1.10-1.53-1.19 JULY
1	0.08-0.58-0.68-0.74-0.74-1.22-1.31-1.39-1.42-1.28
0	-0.44-0.49-0.63(1.22)0.65-1.01-1.29-1.28-1.49-1.42
9	<u>a.49-0.53-0.57-0.34-0.93</u> -1.07-1.31-1.33-1.48-1.33
	-0.66-0.78-0.85-1.09-1.24-1.30-1.47-1.25-0.43
	-0.85-0.93-1.07-1.12-1.18-1.13-1.08-0.73-0.34
	-0.55-0.84-0.95-1.15-1.14-1.19-0.9657.03-2.69
	-0.85-0.90-1.04-1.03-1.18-1.14-0.95-0.96-0.65 POINT CONCEPTION
	-a. 85-a. 97-1. 01-1. 11-1. 03-1. 03-0. 95-0. 34-0. 04
3	-0.98-1.02-0.58-0.99-0.79-0.96-0.78-0.51-0.22-0.07
2	-0.93-9.88-0.78-0.78-0.86-0.73-0.53-0.42-0.22-0.10
1	-0.87-0.70-0.67-0.77-0.78-0.73-0.69-0.49-0.36-0.33
10	-0.54-0.79-0.57-0.68-0.59-0.58-0.12-0.45-0.42-0.38
	-0.85-0.75-0.71-0.52-0.42-0.45-0.47-0.43-0.51
	-8.63-9.53-9.52-9.52-9.52-9.53-9.41-9.49-9.44-9.39
	-0. 58-0. 53-0. 45-0. 45-0. 46-0. 44-0. 44-0. 44-0. 44-0. 34
	- a 13-0 57-0 52-0 50-0 35-0 42-0 37-0 45-0 34-0 29
25	D 49-0 40-0 39-0 46-0 35-0 47-0 39-0 41-0 37-0 25
21	
-	
-	10. 32-01. 45-10. 45-10. 45-10. 25-10. 25-10. 25-10. 25-10. 25-10. 25-10. 21-10. 1
	-0.43-0.54-0.33-0.00-0.19-0.00-0.53-0.31-0.17-0.1

Chart 31.-North component surface wind stress for July.

137 136 135	- UNE DEGREE SUMPHRIZHIUN 5 134 133 132 131 130 129 128 127 126 125 124 123 122 121 120	119 118 117 116 115 114 113 112 1
0.07 0.0	6-0.03-0.16 0.15-0.10-0.21-0.10-0.36-0.01 VANCOUVER ISLAND	
-0.2	13 0.00-0.05-0.15-0.13-0.20-0.33-0.40-0.15-0.09 -0.13 0.02-0.05-0.06-0.28-0.26-0.35-0.39-0.06 0.00	NDAA NATIONAL MARINE FISHERIES SERVICE PACIFIC ENVIRONMENTAL GROUP MONTEREY, CALIFORNIA
	-Ø. 13-0. 17-0. 09-0. 27-0. 48-0. 31-0. 42-0. 30-0. 10-0. 12	NORTH COMPONENT
	-0.15-0.21-0.11-0.31-0.31-0.38-0.40-0.52-0.37-0.25 COLUMBIA RIVER	SURFACE STRESS
	-0.01-0.22-0.26-0.40-0.26 0.64-0.45-0.39-0.41-0.40	(DYNE CM-2)
	0.39-0.35-0.15-0.35-0.43-0.50-0.53-0.60-0.60-0.17	LONG TERM MEAN
	-0.33-0.29-0.37-0.40-0.62-0.47-0.83-0.91-0.69-0.49 CAPE BLANCO	FOR
	-0.26-0.35-0.40-0.22-0.37-0.77-0.63-0.87-1.00-0.63	
	-0.23-0.51-0.47-0.70-0.67-0.89-0.99-1.07-0.99-0.36	1972 - 7001
	-0.41-0.51-0.48-(1.22-0.90-0.95-1.17-1.01-1.42-1.23	
	0,49-0.57-0.52-0.72-0.90-1.03-1.03-1.19-1.30-1.10	
	-0.51-0,50-0.68-0.91-0.84-1.05-1.00-1.12-1.06-0140	
	0.52-0.74-0.76-0.98-1.02-0.97-1.02-0.97-0.74-0.42	
	-0.51-0.54-0.74-0.91-0.93-0.92-0.97-0.80-0.89-0.7	
	-0.59-0.69-0.92-0.95-0.78-0.38-0.36-0.96-0.77-0.67	INT CONCEPTION
	-0.72-0.88-0.77-1.04-0.95-0.89-0.73-0.79-0.4	8-0.04
	-Ø.74-0.92-0.85-0.84-0.74-0.78-0.78-0.8	1-0.21-0.08
	-0.76-0.76-0.92-0.97-0.76-0.6	3-0.42-0.21-0.12
	-0.59-0.87-0.75-0.75-0.75-0.55-0.5	3-0.49-2.36-0.33
	-0.71-0.54-0.74-0.69-0.53-0.5	-0.40-0.37-0.36-0.39
	-0.51-0.50-0.46-0.52-0(4	9-0.43-0.31-0.43-0.42-0.4 C
	-0. 18-9. 57-9. 57-9. 57-8. 5	2-0.57-0.39-0051-0.40-0.40 PUNTA EUGENIA
	-0.51-0.60-0.43-0.3	2-0.33-0.47-0.41-0.36-0.41-0.34
	-0.51-0.57-0.4	5-0.44-0.43-0.29-0.37-0.41-0.31-0.28
	-B. 42-B. 58-0. 4	3-0.42-0.48-0.39-0.31-0.33-0.35-0.26
	-9.6	9-0.38-0.31-0.39-0.40-0.25-0.28-0.28-0.23-0
	-0.3	4-0.50-0.31-0.17-0.28-0.36-0.29-0.26-0.21-0
	-0.4	1-0.34-0.35-0.40 0,00 0.00-0.34-0.20-0.25-0
	-0.3	9-0.34-0.23-0.15 0.02-0.23-0.23 0.26-0.22-0
137 136 13	5 134 133 132 131 130 129 128 127 126 125 124 123 122 121 120	119 118 117 116 115 114 113 112 1

Chart 32.-North component surface wind stress for August.

137	136 1	35	134	133	132	131	130	129	128	127	126	125	124	123	122	121	120 119 118 117 116 115 114 113 112
1.57	130 1	1	1.54	100	1	1	1		2	~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~	3						
	a 20 a		G (30)	01 (15)	C 13	0 02	0.08	-0.14	0.55	0.03	VAI	ICOUVE	A ISLA	10			
	0.25			0.05		W.					2	2	~	2			NDRA
	6	3.24	0.08	0.06	0.07	-Ø. 04-	(-)	-10.108	-0.15	-10, 14	-0.0	2	~	5	ζ		NATIONAL MARINE FISHERIES SERVICE
3	· · · · · ·		0.31	0.09	0.01-	-Ø. 19	12-103	-Ø.32	-Ø.44	-Ø.15	-0.9	0.09	5	2	2		MONTEREY, CALIFORNIA
7				0.04	Ø. 33	-Ø. 25-	2,55	-Ø.32	-Ø. 48	-0.25	-0.0	j-Ø. 1	-0.06				NORTH COMPONENT
				-Ø×Ø4	Ø.26	-Ø. 19	Ø.19	-Ø.44	-0.39	-Ø. 48	-Ø. 42	-0.2	0.09	COL	UMBIA	RIVER	SURFACE STRESS
				Ø.13	Ø. 49	-Ø. 37	0.20	-Ø.32	-0.63	-Ø.22	-0.3	-Ø.2	-ø.øs				(DYNE CM-2)
				Ø.13	0.31	-Ø.36	0.52	-0.58	-0.48	-0.41	-2.1	-ø. 3	-0.13				and a set a set of the set of the
				0.20	-@_55	-0.39-	-Ø.91	-Ø.61	-0.78	-Ø.85	-lord	-0.6	0.26	CAP	E BLAN	CO	LONG TERM MEAN FOR
				-0.28	-0.35	-10.52-	-0.47	-0.49	-0.63	-0.99	-1.2	6.8) -Ø.56				SEPTEMBER
				ato	TA DE		-17 70	201 10	-0 51	1 21	-1 4	i la p	1-0.44				1854 - 1972
				0.00	0.00	1	0.10	R. 98	0.01	A.C.	114		K	CRF	E MEND	OC IND	
				10.65	-0.54	10.54	-10. 78	-10. 78	10.03	10 . / 4	0.0	()	0.00				
				-Ø.54	-0.63	-Ø.65	-0.73	-Ø.78	-1.05	-0.82	-0.9	-1,18	3-0.73	>			
					18.)48	-Ø.67	-0.65	-Ø.73	0.90	-0.80	0.9	7-0.8	9-0.81	-0.34	3		
						-0.55	-0.70	-0.80	0.95	-0.84	-Ø.8	7-0.9	2-0.85	-0.62	0.38		
5						-Ø.65	-0.84	-Ø.7	Ø.81	-0.88	0.8	3-0.7	4-0.65	-Ø.75	-0.63	1	
5							-0.75	-0.86	-Ø. 87	-0.75	0.7	3-0.7	2-0.71	-Ø.59	-0.68	-0.51	POINT CONCEPTION
-			1	1		1			-0.67	-Ø.80	0.6	ð-Ø.7	1-0.86	-0.70	-0.73	-0.77	1-8.35-8.89
										-0.7	-0.7	3-0.7	2-0.70	-0.69	-0.80	-0.83	-0.02-0.23-0.11
2		!							1		-0.6	3-0.5	7-0.77	-0.70	-0.78	-0.73	-8.66-9.45-8.26-8.15
									1		-0.5	5-Ø.6	8-0.73	-0.75	-0.84	-Ø.91	-8.65-0.53 8.47-0.39
3											-	-ø.5	8-0.57	-0.62	-0.72	-Ø.61	8.48-2.59-2.47-3.65-2.44
3										1	1	-	-0.53	-0.55	-0.57	-2.52	-0.41-0.53-0.37-0. 5-0.52-0.53 20 A
													-0.60	-0.45	-0.38	-Ø.61	2. 16-8. 63-0.49-8. 34-8. 43-8. 47
													-	10.43	1.54	-0.33	-9, 52-9, 55-9, 77-9, 49-9, 54-9, 45-9, 30
									J					2	-0.52	-9.72	-9.54-9. A)-9.57-9.55-9.55-9.42-9.38-9.72
																-0.54	
												J			E. 98	-10.64	1 0. 45 0. 05 0. 50 0. 14 0. 45 0. 50 0. 4C 0. 25
									j								-0.55-0.56-0.66-0.44-0.59-0.52-0.39-0.31-0.22
								1 									-0.75-0.70-0.62-0.61-0.45-0.51-0.34-0.40-0.27
																	-0.14-0.45-0.23-0.33-0.30-0.53-0.30-0.30-0.23
					1			1		1. 1. 1.		1					-0.45-0.64-0.52-0.42-0.30-0.44 0.20-0.30-0.20

Chart 33.-North component surface wind stress for September.





Chart 34.-North component surface wind stress for October.





Chart 35.-North component surface wind stress for November.



Chart 36.-North component surface wind stress for December.

APPENDIX II

Standard Errors of the Means

The following tables list the computed standard errors of the monthly means by 1-degree square area for the eastward and northward components of the resultant surface wind stress. The standard errors of the means are defined by Equation (3). Each page contains values for 30 1-degree squares blocked in groups of 10. Within each group of 10, all 1-degree squares are defined by a common latitude. The first set of values corresponds to the 1degree square immediately adjacent to the coast. The final tabulations refer to the 1-degree square farthest from the coast.

The standard error of the mean is tabulated for each month and square. The latitude and longitude of the center of the 1-degree square is listed at the left. The first two numbers below each month are the standard errors of the means for the eastward and northward components, respectively. The units are dyne cm⁻². The last value is the number of observations.

LATILON	JAN	÷EB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
21 N 111 W	• 04 • 05 74	• 03 • 03 71	• 0 3 • 0 4 95	• 03 • 05 75	• 03 • 04 88	• 0 4 • 0 4 87	.06 .04 101	.04 .04 85	.08 .06 107	• 06 • 04 95	•] 8 • 05 81	• 02 • 03 94
21 N 112 W	•03 •05 51	.02 .03 68	•02 •07 •52	•03 •07 48	.04 .06 61	• 03 • 05 66	•08 •06 57	•11 •07 35	•08 •08 57	•04 •06 45	•04 •04 46	• 0 4 • 0 4 5 7
21 N 113 W	• 0 4 • 0 6 3 5	•11 •08 50	•04 •10 38	•04 •09 39	• 05 • 08 3 9	• 04 • 07 39	•05 •06 40	•15 •16 25	.08 .16 33	•04 •10 38	•08 •10 37	• 05 • 05 58
21 N 114 W	•04 •08 41	• 10 • 07 39	•04 •08 33	• 07 • 17 22	.05 .11 24	•08 •10 21	•08 •24 32	•15 •16 18	•35 •28 23	•07 •04 28	•03 •07 32	• 05 • 08 42
21 N 115 W	•16 •09 29	• 09 • 10 26	•08 •07 35	.08 .13 13	• 05 • 06 23	•06 •08 23	.03 .02 31	•05 •08 17	.08 .11 19	.07 .11 20	•03 •08 27	•04 •06 40
21 N 116 W	•15 •12 26	• 0 4 • 0 6 2 8	• 07 • 09 32	•09 •14 15	•10 •14 13	.07 .07 14	•08 •12 26	•14 •16 15	•20 •15 18	•08 •06 18	.04 .09 18	• 0 3 • 0 6 2 5
21 N 117 W	• 06 • 09 • 38	• 03 • 06 26	•07 •09 25	.11 .09 19	.05 .10 12	• 05 • 09 22	•16 •08 15	•13 •19 21	•14 •10 23	•09 •09 22	.02 .06 24	• 04 • 07 26
21 N 118 W	.09 .08 .34	•10 •05 21	• 06 • 07 • 24	•14 •14 18	.08 .12 19	•08 •06 19	•12 •09 22	•19 •24 18	•11 •15 21	•03 •03 38	•05 •08 23	• 05 • 08 34
21 N 119 W	• 09 • 08 • 24	• 07 • 08 26	• 06 • 09 30	•11 •11 27	.04 .10 19	•09 •06 27	•16 •13 21	•11 •06 22	.08 .18 30	• 0 4 • 0 5 3 3	•04 •06 43	• 0 7 • 0 7 • 3 7
21 N 120 W	•09 •10 24	• 05 • 06 34	• 06 • 05 30	•05 •09 28	• 06 • 07 38	•06 •05 31	•12 •07 39	•05 •09 34	•07 •07 37	•09 •13 40	•05 •05 40	• 0 8 • 06 27
22 N 111 W	.02 .04 128	.03 .03 101	•02 •03 164	•04 •05 122	•02 •03 169	•03 •03 150	•03 •02 143	•03 •03 150	•07 •07 161	•03 •03 160	•02 •02 150	•03 •05 120
22 N 112 W	• 02 • 04 73	• 02 • 04 6 0	•03 •05 78	•06 •05 80	.03 .05 .99	.03 .04 117	.02 .03 105	•15 •08 116	.04 .04 110	.04 .04 100	•02 •03 90	• 03 • 05 69
22 N 113 W	.03 .04 51	•04 •11 47	• 03 • 06 67	•05 •12 57	• 03 • 07 67	•03 •05 75	.03 .03 84	•04 •05 68	.09 .11 70	•03 •04 73	•03 •05 77	• 06 • 04 57
22 N 114 W	.03 .06 34	•04 •09 42	•04 •09 45	•02 •07 52	• 05 • 08 53	•03 •07 50	.04 .04 61	•09 •07 35	•11 •10 58	.03 .08 45	04 05 38	• 03 • 04 66
22 N 115 W	•07 •09 37	•10 •10 30	•06 •09 54	•04 •10 38	• 05 • 09 36	•03 •09 37	•05 •07 32	•19 •18 34	•09 •07 31	•07 •07 34	•04 •07 22	•07 •06 30
22 N 116 W	.06 .10 24	.05 .10 30	•04 •12 32	•11 •19 15	• 05 • 11 32	•08 •09 30	.03 .07 26	•32 •29 17	*26 *29 24	.08 .06 21	•04 •07 32	07 06 33
22 N 117 W	•05 •11 33	• 05 • 09 29	•08 •10 34	•05 •11 23	.07 .14 19	.08 .12 15	.07 .10 21	*13 *27 17	•12 •10 17	•05 •06 25	•06 •07 26	• 05 • 05 27
22 N 118 W	•06 •09 28	•11 •11 27	•04 •06 33	•10 •13 21	.08 .11 17	.06 .12 18	•15 •14 15	.05 .12 17	•21 •17 17	.05 .07 19	•06 •09 23	•05 •07 34
22 N 119 W	.07 .08 31	• 0.7 • 27 24	•08 •08 24	•07 •05 25	•10 •10 19	•04 •06 24	•06 •12 19	•12 •10 26	•19 •18 21	•10 •07 21	•12 •07 26	• 05 • 09 32
22 N 120 W	• 07 • 08 • 33	•08 •10 26	•07 •06 31	•09 •08 23	• 04 • 07 38	•17 •14 17	•08 •10 20	•14 •13 19	•11 •14 22	•07 •07 23	•06 •06 34	• 07 • 08 20
23 N 111 W	.01 .01 1059	.01 .01 1030	.01 .01 1125	.01 .01 1159	.01 .01 1241	.01 .01 1128	.01 0.00 1178	•02 •03 1152	•01 •01 1132	.01 .01 1199	.01 .01 1083	•03 •01 986
23 N 112 W	• 02 • 02 155	.02 .03 124	•02 •03 146	•03 •04 135	•03 •03 192	• 02 • 03 245	.02 .01 203	.04 .03 181	•03 •03 206	.02 .02 187	.02 .03 154	03 05 140
23 N 113 W	• 0 3 • 0 9 • 47	• 0 3 • 0 4 4 8	• 06 • 06 • 47	•03 •06 54	•03 •05 78	• 05 • 05 78	•02 •03 76	•04 •05 72	•04 •10 79	.03 .04 75	•02 •04 70	• 0 6 • 1 1 47
23 N 114 W	•04 •04 45	• 02 • 05 • 44	•04 •08 49	• 05 • 13 59	•03 •06 79	.03 .05 71	.02 .03 84	•04 •04 96	•05 •06 74	•04 •06 73	•03 •05 59	•06 •09 47
23 N 115 W	• 11 • 0 4 42	• 0 3 • 0 7 4 5	•04 •06 55	•03 •06 67	•03 •09 65	.02 .05 71	•03 •04 62	• 05 • 05 5 2	•04 •06 71	•04 •04 68	• 05 • 05 55	• 05 • 07 57
23 N 116 W	• 04 • 07 42	• 07 • 09 37	• 04 • 06 48	•05 •07 46	• 03 • 07 63	•05 •06 46	03 05 62	•07 •04 50	•05 •08 51	•04 •06 47	•05 •06 45	* 0 6 * 0 5 5 4
23 N 117 W	• 09 • 10 30	•04 •11 34	• 0 6 • 0 7 6 0	•05 •12 38	• 06 • 09 48	• 0 4 • 0 7 • 46	•04 •05 43	•16 •19 28	•09 •09 40	•06 •08 38	.09 .10 33	• 05 • 04 52
23 N 118 W	• 05 • 08 34	•10 •10 29	• 03 • 08 37	• 05 • 10 33	•06 •08 36	•06 •10 27	•05 •06 32	•16 •16 22	•11 •11 29	•06 •06 32	•06 •06 26	•04 •05 38
23 N 119 W	•06 •08 35	• 05 • 09 38	•03 •09 36	• 05 • 09 33	05 07 38	•06 •08 39	•06 •08 30	.07 .12 16	•10 •11 24	•07 •09 26	•17 •10 24	*10 *08 37
23 N 120 W	• 10 • 10 37	.05 .06 30	.06 .10 21	•11 •14 17	• 04 • 06 22	• 05 • 09 23	.08 .17 17	•08 •09 •22	•07 •10 24	.08 .10 18	•14 •08 32	• 06 • 08 42

LATILON	JAN	ΞEB	MAR	APR	MAY	JUN	JUL	AUG ,	SEP	OCT	NOV	DEC
24 N 111 W	• 01 • 01 572	• 01 • 01 494	.01 .01 590	•01 •01 587	.02 .01 591	•01 •01 603	•01 •01 596	•01 •01 602	•03 •06 566	.01 .01 535	•01 •01 594	•01 •01 504
24 N 112 W	0.00 .01 1226	.01 .01 1209	.02 .01 1375	.01 .01 1380	•01 •01 1458	.01 .01 1398	.01 0.00 1453	.01 .01 1324	•01 •01 1382	0.00 .01 1379	0.00 .01 1336	.01 .01 1208
24 N 113 W	.02 .04 143	• 03 • 04 142	•03 •03 195	•03 •05 150	• 02 • 03 176	•02 •03 210	.02 .02 193	•02 •02 223	.03 .03 214	•03 •03 177	•02 •02 172	• 02 • 03 141
24 N 114 W	.07 .06 42	• 06 • 09 39	•05 •08 45	• 05 • 07 38	• 04 • 05 58	• 0 4 • 0 5 5 4	.03 .05 54	•04 •06 70	•05 •06 78	•04 •05 62	•05 •08 44	•14 •18 25
24 N 115 W	.07	• 05 • 08 31	• 05 • 08 39	• 07 • 15 46	•04 •07 59	.07 .10 49	•03 •06 87	•04 •05 51	•05 •06 68	•05 •05 56	•03 •07 44	• 07 • 10 22
24 N 115 W	• 05 • 07 35	.03 .06 43	• 08 • 09 37	•04 •08 55	• 04 • 08 67	• 05 • 06 68	•02 •04 75	•06 •05 73	•04 •08 56	•03 •05 59	•02 •06 56	•04 •10 39
24 N 117 W	• 05 • 07 43	• 06 • 15 29	• 0 4 • 0 9 3 3	•05 •18 44	• 04 • 08 62	•03 •07 71	.03 .06 71	•04 •05 62	•04 •06 60	•03 •05 56	•04 •05 51	• 04 • 05 57
24 N 118 W	• 05 • 06 35	• 06 • 08 • 47	•04 •05 49	• 05 • 09 40	• 05 • 08 52	• 03 • 06 52	•02 •06 47	•09 •06 44	•05 •08 50	•04 •05 57	•04 •06 48	• 07 • 06 39
24 N 119 W	.03	:11 :13 41	•08 •07 56	• 05 • 10 44	• 84 • 07 57	• 03 • 09 39	•06 •08 53	•13 •08 38	.05 .09 36	•04 •07 40	•08 •08 36	•04 •04 50
24 N 120 W	.07 .08 47	• 0 4 • 0 7 5 5	• 0 3 • 0 8 57	• 05 • 08 45	.05 .06 41	• 05 • 09 30	•08 •05 38	•15 •10 29	•08 •11 35	•07 •05 38	•05 •06 46	• 05 • 05 48
25 N 113 W	0.00 .01 1297	.01 1257	•01 •01 1441	•01 •01 1409	.01 .01 1509	•01 •01 1467	0.00 1508	0.00 .01 1431	.01 .01 1431	0.00 .01 1426	0.00 .01 1408	01 01 1259
25 N 114 W	.02 .04 147	• 0 2 • 0 4 1 2 8	•04 •05 139	•03 •05 150	•03 •04 176	•03 •03 206	.04 .04 177	•02 •02 225	•03 •04 200	.13 .06 185	03 04 149	.03 .03 125
25 N 115 W	•56 •65 31	•04 •09 32	• 05 • 08 42	• 03 • 08 59	• 03 • 07 86	•05 •08 66	.03 .04 81	•04 •04 73	.03 .06 .57	•02 •04 66	•05 •05 46	•08 •09 30
25 N 116 W	.10 .17 15	• 0 8 • 0 6 2 2	•12 •13 27	•05 •12 23	•08 •08 35	•06 •09 31	.03 .05 46	.02 .03 53	•14 •08 44	•04 •11 25	•05 •06 37	•10 •16 14
25 N 117 W	• 0 4 • 0 7 2 3	• 11 • 10 27	•10 •12 30	• 0 4 • 0 9 3 5	• 04 • 06 52	•05 •06 40	•03 •05 51	•06 •06 59	•06 •08 38	•05 •07 45	•05 •08 34	•11 •16 22
25 N 118 W	• 06 • 06 33	• 04 • 10 29	•06 •10 34	•05 •23 41	•04 •08 61	•04 •06 58	.02 .05 67	•05 •07 57	.06 .10 48	.02 .05 61	•05 •08 42	• 0 3 • 0 8 5 0
25 N 119 W	• 06 • 08 52	• 06 • 10 34	• 05 • 07 5 0	•05 •14 42	.05 .08 54	• 0 4 • 0 7 7 0	•03 •07 58	•03 •05 59	•04 •09 56	•03 •05 58	•05 •11 49	• 05 • 06 48
25 N 120 W	.07 .09 37	.07 .12 36	•04 •06 49	•04 •08 52	.04 .07 61	.03 .05 44	.02 .04 55	•04 •06 51	•04 •07 41	•04 •06 45	•04 •09 54	• 07 • 08 36
25 N 121 W	• 07 • 06 46	.05 .10 42	•03 •06 53	• 05 • 07 38	•04 •06 49	.04 .06 41	•03 •06 55	•08 •08 46	•08 •08 41	.05 .05 60	•09 •15 33	• 07 • 07 37
25 N 122 W	• 05 • 08 5 0	• 07 • 08 52	•04 •06 56	•05 •08 59	• 02 • 05 68	04 05 36	•04 •06 44	.06 .06 45	• 06 • 09 48	•05 •06 54	•10 •06 49	•] 9 •] 6 56
25 N 113 W	• 01 • 02 372	•01 •02 399	• 0 2 • 0 2 • 2 2	• 02 • 02 397	• 02 • 02 444	• 01 • 02 452	•01 •06 398	•01 •01 408	• 0 2 • 0 2 42 8	•01 •02 427	•01 •02 439	• 02 • 02 389
25 N 114 W	.01 .01 1199	.01 .01 1201	•01 •01 1438	.01 1327	.01 .01 1367	.01 .01 1334	.01 .01 1360	.01 .01 1346	.01 .01 1331	.02 1333	•01 •01 1253	.01 .01 1165
25 N 115 W	.04 .04 120	.03 .04 115	•04 •05 126	•03 •05 117	•04 •04 135	•02 •04 158	.02 .03 148	•02 •03 150	•03 •05 147	•03 •04 129	•02 •04 129	• 03 • 05 96
26 N 115 W	• 08 • 09 30	• 67 • 14 24	•08 •10 34	•06 •10 45	•04 •07 65	.04 .07 68	•03 •04 68	•03 •05 59	•04 •07 66	• 03 • 05 55	•05 •18 60	•15 •80 30
25 N 117 W	.07 .16 16	•12 •18 15	•24 •28 16	•04 •09 33	• 0 3 • 07 30	• 03 • 09 37	•04 •05 59	•05 •08 48	.04 .09 .42	•03 •06 35	•04 •05 30	•20 •19 16
25 N 118 W	•04 •09 32	•06 •09 24	•12 •12 32	•04 •15 26	• 04 • 08 35	• 04 • 07 34	•04 •05 35	• 05 • 06 43	•04 •07 52	•04 •07 39	•05 •10 25	•10 •14 13
25 N 119 W	• 05 • 09 33	•10 •11 29	• 0 6 • 0 9 2 6	•05 •14 33	•04 •08 41	• 04 • 08 36	•03 •06 43	•02 •08 38	•06 •10 24	•05 •07 30	• 0 4 • 0 7 • 3 7	• 0 4
26 N 120 W	• 1 0 • 07 3 0	• 10 • 15 24	•09 •15 34	•11 •09 38	• 03 • 08 57	• 02 • 05 59	•05 •08 50	• 03 • 06 45	•06 •09 48	•03 •06 57	• 06 • 06 43	• 05 • 09 38
125 N 121 W	• 05 • 07 59	•10 •12 26	• 0 4 • 1 0 4 0	•04 •09 54	• 0 4 • 0 9 5 5	•03 •06 69	•08 •08 43	• 0 3 • 0 5 6 6	•04 •09 47	•04 •05 51	•05 •14 42	• 07 • 08 46
26 N 122 W	• 05 • 06 37	• 04 • 06 52	.06 .06 .61	•03 •09 54	• 03 • 07 55	• 02 • 04 57	•03 •05 60	• 0 3 • 0 5 5 7	•03 •06 60	• 0 3 • 0 4 6 3	•03 •05 70	• 0 8 • 0 7 5 0

LATILON	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
27 N 114 W	.02 .02 468	.02 .02 458	•02 •02 542	.02 .02 517	•02 •02 548	•02 •02 488	•01 •01 527	.01 .01 452	•02 •02 549	.01 .01 524	.02 .02 511	• 02 • 02 521
27 N 115 W	.01 .01 1113	.01 .01 1085	.01 .01 1245	.01 .01 1204	.01 .01 1250	.01 .01 1194	.01 .01 1209	.01 .01 1188	.01 .01 1169	.01 .01 1183	.01 .01 1122	•01 •02 973
27 N 116 W	• 06 • 07 76	.04 .06 102	.02 .10 88	.03 .04 108	.04 .05 101	•04 •05 110	•02 •04 99	.02 .03 120	•03 •06 130	•03 •06 84	•03 •05 86	• 0 4 • 0 7 6 2
27 N 117 W	.08 .12 29	.05 .07 .34	.08 .10 43	.05 .11 48	.05 .08 51	•03 •06 64	.03 .06 71	.03 .06 71	•03 •05 76	• 43 • 73 55	.06 .07 51	• 06 • 10 25
27 N 118 W	.06 .10 28	.07 .16 20	•10 •14 23	.05 .11 .32	•04 •08 36	• 03 • 06 4 3	•02 •05 70	•03 •04 53	• 0 4 • 0 8 4 9	.03 .04 31	•05 •09 37	•24 •10 19
27 N 119 W	.33 .09 19	:10 :32 12	•11 •08 16	•05 •09 34	•04 •10 32	• 0 \ • 10 32	•03 •10 29	•04 •05 32	•04 •07 37	•03 •06 37	•04 •05 26	• 0 4 • 22 2 0
27 N 120 W	• 07 • 11 34	:11 :27 19	•08 •11 23	.05 .11 20	• 03 • 14 30	• 03 • 08 33	•04 •07 29	•04 •07 38	•08 •09 36	•04 •09 26	•09 •14 17	.06 .08 15
27 N 121 W	•13 •09 40	• 06 • 05 25	•11 •12 29	•05 •12 42	•04 •08 46	•05 •10 18	•04 •12 31	•03 •05 49	•06 •13 24	•04 •07 49	•07 •07 33	.08 .14 17
27 N 122 W	• 07 • 06 36	.09 .12 33	.05 .11 51	•03 •10 53	• 05 • 08 58	•05 •06 64	•05 •09 54	•04 •08 62	•06 •09 48	•02 •05 75	•07 •08 53	• 07 • 05 62
27 N 123 W	. 04 . 07 69	•08 •09 52	• 0 6 • 0 9 6 3	•03 •07 75	• 02 • 05 93	•03 •06 90	•05 •08 50	•03 •05 80	•03 •04 91	•02 •04 105	•06 •08 77	•06 •09 51
20 11	17		24	0.2		6.4	0.4					
115 W	• 06 588	.02 517	.02 742	•02 •03 720	.02 723	.01	•01 632	.01 685	•02 709	•01 695	•02 676	• 02 • 02 624
28 N 115 W	.01 .02 810	.01 .02 747	•01 •02 865	.01 .02 843	.01 .01 874	•01 •01 903	.01 .01 831	.01 .01 840	.01 835	.01 875	.01 .01 821	•02 735
28 N 117 W	• 06 • 05 75	.05 .07 72	.05 .06 65	.03 .06 69	•04 •06 59	.03 .04 69	•04 •05 53	03 07 66	05 75	04 05 71	•04 •08 55	•09 •12 56
28 N 118 W	.05 .06 34	.07 .11 24	• 05 • 09 25	.07 .15 30	• 05 • 06 41	• 0 3 • 0 7 4 7	•02 •06 56	•04 •04 63	•04 •07 54	•02 •06 43	•05 •10 33	• 05 • 14 22
28 N 119 W	• 06 • 06 38	.04 .11 30	•05 •08 33	•05 •09 43	• 04 • 05 53	•04 •07 72	•03 •07 67	•03 •07 55	• 14 • 11 39	•03 •06 39	•06 •10 33	•13 •14 19
28 N 120 W	•20 •15 17	·29 ·36 18	.08 .09 36	.06 .11 43	• 06 • 11 33	.03 .08 39	•06 •06 36	•04 •07 41	•03 •06 45	•05 •08 22	•05 •12 31	:10 :20 20
28 N 121 W	• 06 • 09 35	.08 .25 14	.08 .11 28	.05 .10 41	.04 .09 31	.05 .10 41	•05 •09 29	•02 •05 48	•04 •09 32	•04 •07 39	.03 .08 31	•06 •10 23
28 N 122 W	.06 .11 56	.04 .10 34	.07 .12 45	.05 .10 32	•05 •13 44	.05 .08 44	•03 •08 55	•05 •08 58	.03 .05 43	.03 .08 54	•08 •16 27	• 05 • 11 35
28 N 123 W	•10 •12 •47	.08 .10 50	• 06 • 12 44	.03 .10 68	• 05 • 06 7 0	•05 •11 37	.03 .10 41	•07 •07 65	•04 •05 65	•03 •04 80	•07 •09 75	•06 •06 58
28 N 124 W	•06 •08 48	.05 .07 46	•05 •12 43	.05 .09 44	•03 •07 65	• 06 • 06 54	•06 •09 46	•08 •07 47	.09 .08 38	02 04 88	•06 •07 67	.06 .09 51
29 N 115 W	• 07 • 09 62	.07 .08 88	•11 •09 90	•07 •09 80	•09 •10 71	• 05 • 06 • 66	•07 •07 61	•07 •04 63	•05 •07 •59	•05 •05 58	•12 •07 73	• 08 • 09 67
29 N 116 W	.02 .01 1030	.01 .01 1)49	.01 .01 1209	.01 .01 1155	.01 .01 1246	.01 .01 1080	.01 .01 1081	.01 .01 1115	.01 .01 1109	.01 .01 1112	.01 .01 1051	•01 •01 983
29 N 117 W	•03 •03 252	• 02 • 03 249	• 02 • 03 336	• 02 • 03 308	• 02 • 03 250	• 02 294	•02 •02 267	*01 •02 280	•02 •02 273	.02 .02 265	•02 •03 261	• 0 3 • 0 3 229
29 N 118 W	.03 .06 108	•08 •10 58	.07 .08 71	.07 .07 65	• 05 • 07 • 59	•04 •07 64	•05 •06 49	.03 .05 56	•08 •06 54	•04 •05 55	•05 •05 56	• 0 6 • 0 9 37
29 N 119 W	•07 •09 27	• 07 • 09 21	•10 •22 34	• 05 • 08 35	• 05 • 10 43	•04 •09 45	•03 •07 43	•05 •10 33	•04 •09 42	• 0 2 • 0 5 5 4	•10 •09 33	• 05 • 07 35
23 N 120 W	.06 .11 29	.07 .09 28	•10 •15 25	• 05 • 09 56	• 04 • 07 48	• 0 4 • 0 8 • 4 4	•05 •07 •52	•03 •05 68	•04 •08 55	•06 •09 36	• 06 • 09 27	• 06 • 11 36
29 N 121 W	• 05 • 08 5 9	• 11 • 12 29	•09 •12 42	• 05 • 08 5 0	• 05 • 08 51	• 03 • 07 6 9	•04 •08 59	•03 •07 54	•07 •07 45	• 0 4 • 0 8 • 4 4	1.27	• 0 8 • 0 9 3 4
29 N 122 W	• 05 • 10 73	.09 .13 48	•07 •13 51	•04 •10 56	• 05 • 06 81	• 0 3 • 0 6 8 4	•04	•03 •06 86	• 0 5 • 0 6 • 4 4	•03 •06 82	• 07 • 07 62	• 0 8 • 0 8 3 9
29 N 123 W	•08 •12 63	• 05 • 13 39	• 0 9 • 11 3 9	• 05 • 12 45	• 04 • 07 65	• 03 • 08 61	•03 •06 65	•16 •07 61	• 0 2 • 0 5 7 3	• 0 3 • 0 6 6 8	• 07 • 09 54	• 06 • 09 58
29 N 124 W	.06 .11 51	• 07 • 07 • 50	•07 •14 36	• 06 • 08 5 0	• 03 • 07 76	• 0 4 • 0 7 5 4	•04 •10 54	•02 •05 57	•06 •05 53	•04 •06 80	• 07 • 08 89	.08 .11 31

LATILON	JAN	= E B	MAR	APR	MAY	ЛПИ	JUL	AUG	SEP	OCT	NON	DEC
30 N 116 H	. 32	.02	• 0 2 • 0 3 6 3 7	•01 •02 581	•01 •02 624	•01 •02 593	.01 .01 547	•01 •01 537	•01 •02 553	•01 •02 557	•01 •02 545	• 01 • 02 481
30 N 117 W	•01 •02 504	• 01 • 02 530	•01 •02 698	• 01 • 02 568	• 05 • 04 658	•01 •01 654	.01 .01 666	.01 .01 693	•01 •02 651	•01 •01 672	•03 •04 645	• 02 • 02 591
30 N 118 W	.04 .05 194	• 0 2 • 0 3 2 0 9	.03 .04 187	.04 .04 201	.03 .04 193	•03 •04 190	•02 •03 195	.02 .02 188	.02 .03 180	•02 •02 203	•03 •03 182	•04 •05 165
30 N 119 W	• 07 • 08 76	• 0.6 • 0.9 65	• 07 • 08 60	• 0 4 • 0 5 6 5	•10 •10 38	•04 •06 73	•02 •03 139	•03 •06 52	•05 •08 60	•06 •05 91	•05 •10 48	•08 •10 42
30 N 120 W	•05 •09 63	.05 .08 113	•08 •10 58	• 05 • 07 99	. 113 . 05 86	•05 •06 91	•04 •06 67	•05 •06 70	•04 •05 56	•04 •04 95	•07 •07 •37	•06 •10 52
30 N 121 W	• 37 • 08 71	.06 .07 81	•07 •11 58	-04 -08 107	.03 .07 91	•04 •05 86	•03 •05 84	.03 .05 102	•04 •06 70	•03 •06 89	•04 •08 35	•10 •11 53
30 N 122 W	• 05 • 07 92	•12 •16 60	• 0 9 • 12 47	•08 •11 51	.04 .08 100	.03 .08 104	•03 •06 77	.04 .06 88	•04 •07 73	• 05 • 06 74	.05 .08 51	• 05 • 09 49
30 N 123 W	• 05 • 08 78	•13 •13 50	•12 •15 48	•07 •14 54	• 04 • 06 87	• 03 • 07 8 0	•03 •06 72	•03 •12 73	•06 •09 69	.03 .09 82	•04 •10 69	•03 •05 96
30 N 124 W	•11 •11 66	•09 •11 80	• 0 4 • 0 7 55	•03 •07 75	.04 .07 110	.02 .05 108	•04 •08 60	•05 •07 83	.03 .07 66	.03 .07 113	•04 •09 87	• 05 • 09 63
31 N 125 W	• 8 • 12 78	•10 •15 66	•06 •10 57	•05 •10 66	. 07 - 08 55	• 0 4 • 0 8 7 8	•04 •08 64	•04 •08 78	•04 •06 70	•03 •07 86	•06 •09 72	• 0 6 • 0 8 6 5
31 N 117 W	.01 .01 776	.01 .02 793	•01 •01 922	.01 .01 880	.01 .01 916	•01 •01 866	•01 •01 826	•01 •01 837	.01 .01 861	•01 •01 821	•01 •02 793	• 01 • 02 706
31 N 118 W	•02 •03 394	•03 •03 402	•02 •03 446	• 0 2 • 0 2 4 4 2	.02 .03 441	.02 .02 384	•01 •02 360	.01 .01 443	•02 •02 499	.02 .02 441	• 02 • 02 • 059	• 0 3 • 0 3 372
31 N 119 W	.02 .03 294	• 0 2 • 0 4 2 9 4	•04 •06 236	•04 •04 286	• 03 • 03 253	• 02 • 03 296	•02 •03 255	•02 •03 214	.02 .03 267	.02 .02 345	•04 •04 222	•04 •05 154
31 N 125 W	• 05 • 07 169	•04 •06 143	.05 .07 131	•05 •06 149	•04 •06 144	•03 •05 124	.03 .05 158	•03 •04 126	•04 •06 98	•03 •03 212	.04 .07 122	•03 •06 84
31 N 121 W	•05 •08 103	•04 •08 145	•05 •10 101	•03 •06 131	.06 .06 119	•05 •07 105	.03 .04 132	•04 •05 95	•04 •08 78	•04 •05 131	•04 •06 101	•06 •08 78
31 N 122 W	.04 .06 115	.05 .10 98	•05 •10 97	•08 •11 7]	.06 .07 104	.03 .07 104	.03 .05 106	•04 •06 87	•04 •09 79	•03 •08 127	.05 .10 107	• 05 • 08 87
31 N 123 W	.03 .06 171	.06 .10 110	•06 •08 84	.04 .10 91	•04 •06 109	•03 •67 123	•02 •05 170	.03 .05 142	.03 .06 133	•03 •07 153	.04 .06 111	•05 •10 79
31 N 124 W	.07 .09 112	.07 .12 86	•05 •10 105	•04 •07 173	• 02 • 04 2 07	.03 .06 169	•02 •05 166	.02 .06 102	.03 .06 133	•03 •05 146	•05 •08 119	• 06 • 10 107
31 N 125 W	•06 •08 134	•05 •09 121	•04 •07 139	•05 •08 144	•03 •06 126	•03 •06 150	•02 •04 160	.03 .07 125	.03 .06 124	.03 .08 166	•05 •09 122	• 05 • 11 92
31 N 126 W	.05 .08 157	•06 •10 137	•06 •07 152	•04 •07 145	•03 •05 149	•03 •05 155	•03 •06 158	.02 .05 145	.02 .04 144	.03 .06 180	•05 •06 148	•05 •09 131
32 N 117 W	.01 .01 1.46	.01 .01 912	•01 •01 1)55	•01 •01 967	.01 .01 1095	0.00	0.00 0.00 931	0.00 0.00 1097	.01 .01 1041	.01 .01 1062	•01 •01 887	•01 •01 714
32 N 118 W	.01 2163	.01 .01 1933	.01 .01 2268	.01 .01 2283	•01 0.00 2375	0.00 0.00 2221	.01 0.00 1934	0.00 0.00 2512	.01 0.00 2363	.01 .01 2284	.01 .31 1882	.01 .01 1133
32 N 119 W	.01 .02 1108	.02 .02 .02 .081	.02 .02 1244	.01 .02 1105	.01 .01 1140	.01 .01 1000	.01 .01 1054	.02 .01 1128	.01 .01 1220	.01 .01 1399	.02 1017	• 0 2 • 0 2 7 6 8
32 N 120 W	•02 •03 451	• 0 3 • 0 5 3 5 0	•03 •03 491	•03 •04 479	• 03 • 03 465	•03 •04 354	•02 •02 500	• 0 2 • 0 2 4 3 3	•02 •02 573	•02 •03 492	•02 •03 419	• 04 • 05 262
32 N 121 W	.03 .06 241	• 0 4 • 0 6 2 5 2	•04 •07 235	•04 •05 249	• 03 • 05 254	.04 .05 196	•02 •04 272	• 0 3 • 0 4 225	.03 .04 245	•02 •04 293	•05 •06 233	•05 •07 189
32 N 122 W	•04 •07 198	•05 •08 214	.04 .08 212	•05 •08 214	.04 .06 188	•03 •06 190	•03 •05 229	.04 .06 171	•03 •05 191	•03 •04 302	•03 •05 208	• 05 • 08 176
32 N 123 W	• 0 3 • 0 6 2 87	.05 .08 197	.04 .07 218	.05 .08 198	•04 •07 190	.03 .07 172	.02 .04 203	•03 •05 188	.03 .05 181	•03 •06 253	•03 •06 232	•04 •08 184
32 N 124 W	·01 •04 771	• 0 2 • 0 4 5 3 1	.02 .03 630	• 03 • 05 432	• 01 • 04 569	.01 .03 551	.01 .02 662	.01 .02 711	•01 •02 662	•01 •04 828	• 02 • 03 623	• 01 • 03 637
32 N 125 W	• 03 • 05 302	• 0 3 • 0 6 2 7 3	.05 .06 289	•04 •05 267	• 04 • 05 273	• 02 • 05 263	•02 •05 268	•02 •04 284	•02 •04 237	• 02 • 05 353	•02 •04 323	•03 •06 363
32 N 126 W	• 03 • 05 289	• 05 • 07 252	• 03 • 05 307	•03 •05 300	• 03 • 05 278	•02 •04 257	•02 •05 267	•02 •04 280	•01 •04 247	• 0 2 • 0 5 3 2 0	•03 •04 352	•03 •06 311

LATILON	JAN	= EB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NDV	DEC
33 N 118 W	0.00 0.00 3538	.01 .01 3329	.01 0.00 3831	•01 0.00 3775	0.00 0.00 3805	0.00 3305	0.00 0.00 3107	0.00 0.00 3712	0.00 0.00 3483	0.00 0.00 3798	0.00 0.00 3536	•01 •01 2251
33 N 119 W	.01 .01 1351	.02 .02 1121	•02 •91 1557	.02 .01 1412	.02 .01 1492	.01 1320	.01 .01 1079	.01 0.00 1595	.01 .01 1476	.01 .01 1686	.01 .01 1461	.02 .01 1023
33 N 120 W	• 02 • 02 966	.03 .04 510	• 02 • 02 930	• 0 3 • 0 2 9 3 5	.02 .02 1064	.02 .02 1124	•02 •02 823	•01 •02 967	.01 .02 1000	•01 •02 975	•02 •02 934	•03 •03 699
33 N 121 W	• 0 2 • 0 4 473	• 03 • 06 289	•04 •06 392	•04 •05 333	•05 •06 410	.03 .04 415	•03 •05 303	•02 •03 341	•03 •04 384	.03 .04 310	•03 •05 341	• 03 • 05 293
33 N 122 W	• 03 • 06 345	.07	•04 •06 372	• 04 • 05 350	•03 •04 332	•04 •06 276	•04 •05 306	•02 •04 269	•02 •04 298	•03 •06 297	•03 •05 347	• 03 • 05 332
33 N 123 W	.03 .06 363	.03 .06 313	•04 •06 336	•02 •04 575	• 02 • 05 462	•03 •05 329	•02 •03 411	•02 •04 286	.02 .04 308	.02 .04 310	•03 •05 345	• 03 • 05 362
33 N 124 W	.02	•03 •06 286	• 0 3 • 0 5 3 9 5	.04 .05 313	• 02 • 05 389	•03 •05 319	•02 •05 314	•01 •03 378	•02 •04 287	•02 •05 406	•03 •05 377	• 0 3 • 0 5 377
33 N 125 W	• 03 • 05 356	.03 .06 318	• 0 3 • 0 6 3 3 0	•03 •06 272	• 02 • 05 324	• 03 • 05 285	•02 •04 273	•01 •04 306	•02 •05 238	•02 •05 329	•03 •04 339	• 0 3 • 0 6 3 5 6
33 N 125 W	.06 .07 244	•04 •08 187	•04 •08 217	•04 •08 185	• 02 • 05 217	•03 •06 164	•02 •06 204	•03 •06 187	•02 •95 182	.02 .06 216	•03 •06 221	• 03 • 06 256
33 N 127 W	.35 .38 196	.05 .09 178	.04 .07 199	.07 .08 154	•03 •07 164	.04 .06 152	•03 •06 178	.03 .06 137	.03 .07 143	.03 .06 182	•05 •06 183	•06 •11 199
34 N 119 W	.01 .01 1505	.01 .01 1537	•01 0•00 1774	.01 0.03 1990	.01 .01 1924	0.00 0.00 1615	0.00 0.00 1623	0.00 D.00 1420	.01 0.00 1448	•01 0•00 1472	•01 •01 1261	•02 •01 995
34 N 120 W	.02 .02 599	•03 •03 588	.03 .02 743	•03 •03 705	•04 •03 643	.02 .02 721	•02 •02 775	.03 .02 650	.02 .02 656	.02 .02 640	•02 •02 652	• D 2 • D 2 7 5 9
34 N 121 W	• 02 • 03 576	•03 •04 506	•03 •04 594	•03 •04 535	•03 •04 543	• 02 • 03 676	•02 •03 603	•02 •02 650	•02 •03 577	.02 .03 567	•02 •03 558	.03 .04 495
34 N 122 W	• 03 • 05 276	.04 .07 261	•04 •07 274	•04 •07 239	•04 •06 258	•07 •09 265	•03 •04 284	•03 •04 232	•03 •05 246	•04 •07 228	•04 •08 272	•03 •05 256
34 N 123 W	•04 •07 230	• 05 • 08 228	•03 •08 258	•04 •07 185	.05 .08 189	.05 .08 190	•04 •06 197	.04 .06 156	.04 .05 172	•03 •07 263	•04 •06 207	•03 •07 212
34 N 124 H	•03 •05 265	• 0 5 • 0 7 2 0 4	•04 •07 229	•05 •07 218	• 06 • 08 147	•04 •07 166	•04 •08 152	•04 •06 146	.03 .13 188	•03 •06 273	•04 •07 206	•04 •07 241
34 N 125 W	• 04 • 06 229	•05 •08 199	•04 •07 240	• 05 • 07 204	•05 •08 152	•04 •07 143	•04 •07 149	.07 .07 137	.03 .06 175	•02 •07 234	•03 •05 217	• 06 • 09 234
34 N 125 W	•07 •08 206	.05	•05 •07 204	•05 •09 196	•05 •09 149	•04 •08 132	•03 •07 141	.04 .07 101	.04 .06 142	•03 •06 189	•05 •08 184	• 0 4 • 17 232
34 N 127 W	• 05 • 08 2 08	• 66 • 07 182	.06 .07 212	•05 •08 185	•05 •08 138	•04 •07 134	•04 •08 122	•03 •08 96	.03 .08 145	•03 •06 164	•05 •08 187	• 05 • 06 218
34 N 128 W	•05 •08 230	•06 •08 184	•06 •07 224	.06 .09 182	•04 •07 179	•04 •07 148	.03 .06 127	•04 •06 129	.03 .07 130	•05 •06 169	.05 .08 186	•05 •07 215
35 N 121 W	• 02 • 03 525	• 0 3 • 0 4 477	.03 .04 613	• 03 • 04 572	•03 •03 586	.03 .04 651	•02 •02 631	•02 •03 564	•02 •02 536	•02 •03 658	•02 •03 559	• 0 3 • 0 4 4 4 7
35 N 122 W	•03 •05 296	•03 •05 310	•04 •05 376	•03 •05 388	•03 •04 432	• 02 • 04 480	•03 •04 441	•02 •03 445	.02 .93 418	•03 •04 422	•02 •04 423	• 02 • 05 274
35 N 123 W	•04 •07 243	•04 •08 218	•04 •07 224	•03 •05 284	•03 •06 276	• 04 • 06 296	•03 •04 293	•02 •06 325	•02 •04 306	•02 •04 262	•03 •06 283	•04 •08 211
35 N 124 W	•04 •08 222	• 05 • 09 202	•04 •08 185	•05 •08 189	•03 •06 243	•04 •06 229	•03 •06 225	•03 •05 243	•02 •04 204	•02 •05 279	•03 •06 237	•05 •09 202
35 N 125 W	• 07 • 12 207	.04 .07 178	.05 .08 171	.05 .09 165	.05 .09 161	•04 •07 200	•03 •06 177	.03 .06 174	•04 •07 174	•03 •07 187	•03 •06 207	•06 •10 169
35 N 126 W	•06 •12 189	.07 .10 147	•05 •12 157	.04 .10 183	.06 .08 141	•04 •08 134	•04 •08 138	•04 •06 147	.04 .07 139	•04 •08 174	.04 .08 179	06 08 184
35 N 127 W	.05 .09 193	.05 .08 181	•06 •08 198	•05 •08 171	•04 •09 153	•05 •07 154	•04 •08 129	•03 •06 150	.03 .07 111	•05 •08 206	.05 .09 191	•05 •09 201
35 N 128 W	•04 •06 366	.05 .07 317	• 04 • 06 373	•06 •05 259	• 03 • 05 270	•02 •06 243	•02 •05 313	.02 .04 316	•03 •06 273	•02 •05 356	•03 •05 370	• 05 • 06 368
35 N 129 W	•04 •06 400	.05 .06 394	•03 •05 479	• 04 • 05 379	• 82 • 84 352	•02 •04 337	•02 •04 312	•02 •03 326	•02 •05 324	•03 •05 404	•03 •05 373	•04 •05 411
35 N 130 W	•04 •05 568	• 04 • 05 523	• 03 • 05 577	• 0 4 • 0 4 4 8 7	• 02 • 04 492	• 02 • 03 442	• 02 • 04 464	•01 •03 446	•02 •04 476	•02 •04 592	•03 •04 554	• 03 • 05 562
LATILON	JAN	÷ЕВ	MAR	APR	MAY	ЛЛИ	JUL	AUG	SEP	OCT	ИОМ	DEC
---------------	-------------------------	-------------------------	-------------------------	-----------------------	-----------------------	-------------------------	-------------------------	-------------------------	-------------------------	------------------------	-------------------------	-------------------------
35 N 122 W	• 02 • 04 552	.03 .05 465	• 03 • 04 592	.02 .04 621	• 03 • 04 631	• C 2 • C 3 7 5 9	•02 •03 767	•02 •03 561	•02 •03 604	•02 •04 553	•02 •03 561	•03 •04 411
35 N 123 W	•03 •06 286	• 0 4 • 0 6 3 4 2	•03 •06 298	•03 •07 364	•02 •04 468	.02 .04 591	•02 •04 577	•02 •03 525	•02 •03 478	•02 •05 418	•03 •05 380	• 0 4 • 0 9 2 4 7
35 N 124 W	• 05 • 08 240	• 0 3 • 0 8 2 2 8	.04 .06 215	• 03 • 07 279	.03 .05 360	•03 •05 351	•03 •05 339	•02 •04 360	•03 •05 286	•02 •06 269	•03 •06 257	•04 •06 246
35 N 125 W	• 05 • 38 296	•04 •09 206	• 0 4 • 17 225	•04 •08 271	•84 •06 281	•03 •06 298	•03 •06 297	•03 •05 291	•02 •07 258	•03 •07 250	•03 •06 270	• 0 4 • 0 7 2 5 5
35 N 125 W	.04 .08 446	•03 •06 414	.03 .06 449	•03 •06 368	• 03 • 05 371	• C 3 • 06 357	• 0 2 • 0 5 3 5 8	•02 •04 382	•02 •06 352	•03 •06 398	•02 •05 530	• 0 4 • 0 7 3 9 7
35 N 127 W	• 0 2 • 0 4 8 9 9	•03 •05 748	•03 •04 694	• 02 • 04 624	.02 .04 651	•02 •04 655	•02 •04 584	•01 •03 714	•02 •04 646	•02 •04 699	•02 •04 660	• 0 2 • 0 5 7 3 4
36 N 128 W	• 62 • 04 1060	03 04 375	• 0 2 • 0 3 96 3	• 02 • 03 855	•01 •03 769	•01 •03 868	•01 •03 861	•01 •02 952	•01 •04 929	•02 •03 913	.02 .03 1168	• 0 2 • 0 3 9 9 7
36 N 129 W	.04 .05 519	•04 •05 511	•03 •04 682	.03 .04 616	• 02 • 04 566	.02 .04 581	•01 •03 552	•01 •03 585	•01 •03 637	•03 •04 612	•03 •04 598	• 0 4 • 0 6 6 2 0
35 N 130 W	•06 •08 303	•06 •06 315	•04 •06 382	•05 •06 331	•03 •06 274	• 0 2 • 0 5 2 9 8	•02 •05 249	•02 •04 259	•04 •05 299	• 0 3 • 0 4 37 3	•04 •05 404	• 06 • 07 322
35 N 131 W	•08 •09 235	•08 •08 266	•05 •08 272	• 05 • 07 238	•04 •06 212	•04 •05 209	•03 •05 138	•03 •05 185	.04 .06 191	•04 •06 250	•06 •08 251	•07 •08 234
37 N 122 ₩	•03 •05 412	• 03 • 05 337	•03 •04 343	•03 •04 378	•03 •04 432	•03 •04 391	•01 •02 581	•02 •03 354	•02 •03 400	•02 •03 433	•02 •04 376	• 0 3 • 0 5 3 2 5
37 N 123 W	• 03 • 05 5 88	.03 .06 591	• 9 4 • 0 5 6 8 8	• 03 • 05 672	•03 •04 738	•02 •03 998	•02 •02 881	•01 •03 836	•01 •02 882	•02 •04 797	.03 .04 687	• 0 4 • 0 6 6 3 7
37 N 124 W	• 84 • 7 578	•04 •06 572	•03 •05 605	.03 .05 514	•04 •06 504	• 03 • 05 570	•03 •04 523	•02 •04 537	•02 •04 533	•03 •05 602	•02 •04 645	• 03 • 05 534
37 N 125 W	• J 2 • 05 731	. U 3 . C 5 586	.02 .04 789	.02 .05 719	• 02 • 04 748	•02 •04 696	•02 •03 706	.02 .03 681	•02 •04 664	•02 •04 755	•02 •04 883	• D 2 • D 5 6 8 8
37 N 126 W	• 03 • 05 633	.02 .05 545	• 02 • 05 696	• 02 • 05 7 03	• 02 • 05 685	•02 •04 720	•02 •04 684	•01 •03 603	•02 •04 602	•01 •04 651	•02 •04 722	• 02 • 05 617
37 N 127 W	•04 •06 435	.04 .08 391	• 03 • 06 424	• 0 3 • 0 5 473	• 02 • 05 5 3 2	• 02 • 06 447	•02 •05 438	•02 •05 406	•02 •05 453	• 02 • 05 528	•04 •05 489	• 0 3 • 0 7 4 3 0
37 N 128 W	•06 •08 313	• 0.6 • 0.8 299	.04 .07 352	•03 •06 343	.02 .05 369	•02 •05 361	•03 •06 286	.02 .06 274	.02 .06 405	.02 .05 410	•04 •06 357	•06 •07 337
37 N 129 W	*06 *08 263	.06 .08 236	•06 •09 279	•04 •05 322	•04 •06 306	•03 •06 331	•02 •06 253	•02 •05 283	• 02 • 05 333	•03 •05 387	.04 .06 351	• 06 • 09 271
37 N 130 W	• 06 • 09 233	•06 •08 230	• 05 • 08 243	•04 •07 312	.04 .05 281	•02 •05 306	•02 •05 271	•03 •05 261	•03 •05 345	•03 •05 338	•05 •06 298	.06 .10 257
37 N 131 W	• 07 • 09 223	•07 •10 201	.08 .10 209	•05 •06 266	•04 •05 229	•04 •06 239	•02 •05 234	•02 •04 242	•03 •05 262	.04 .05 291	•05 •07 266	• 07 • 08 254
38 N 123 W	•02 •03 956	•02 •03 919	•03 •03 937	•03 •03 1139	•02 •03 1121	•03 •03 1034	.01 .02 1132	•02 •02 931	•01 •02 1025	•01 •03 986	• 0 2 • 0 3 8 2 7	•03 •03 791
38 N 124 W	.04 .09 319	•04 •08 332	•04 •06 351	•03 •05 464	• 04 • 06 459	•03 •05 568	•03 •05 560	•02 •04 563	•02 •04 533	• 0 3 • 0 4 42 3	•03 •05 530	•04 •07 453
38 N 125 W	• 05 • 89 208	•05 •10 222	•05 •12 216	•04 •08 296	•04 •09 274	•04 •07 315	•03 •06 338	•03 •05 370	•03 •07 327	• 0 3 • 0 7 287	•03 •06 289	•05 •09 222
38 N 126 W	•08 •14 162	.04 .11 229	•05 •09 191	•04 •10 215	•03 •08 259	• 0 3 • 0 8 2 4 8	•04 •06 292	•03 •05 340	• 0 3 • 0 6 3 0 0	• 0 3 • 0 9 272	•06 •08 239	.07 .12 211
38 N 127 W	• 07 • 17 175	•07 •11 176	• 05 • 08 233	•04 •08 243	•02 •07 318	•03 •06 338	•03 •06 384	.03 .06 349	.03 .06 318	•03 •08 285	•04 •06 266	.08 .11 214
38 N 128 W	• 06 • 13 206	.09 .12 193	•04 •07 261	•04 •07 277	• 03 • 05 383	.02 .06 381	•02 •05 449	•02 •04 435	.02 .05 416	•04 •07 300	•05 •07 282	•07 •09 239
38 N 129 W	• 08 • 13 177	•06 •10 206	•06 •09 214	• 05 • 06 247	•03 •05 307	•02 •09 330	•02 •05 355	• 0 2 • 0 5 3 3 7	•03 •05 320	•03 •07 249	•05 •06 253	•08 •09 217
38 N 130 W	.09 .15 140	•12 •11 172	.07 .10 170	•05 •08 180	• 03 • 06 247	•03 •05 258	•03 •05 273	.02 .04 281	•03 •05 272	•05 •08 192	•08 •09 206	•10 •10 196
38 N 131 W	.89 .13 160	•08 •12 146	•07 •12 151	.07 .09 151	•03 •05 248	.04 .08 201	•03 •05 232	.02 .04 291	.04 .06 221	•06 •08 198	.07 .08 174	• 08 • 12 190
38 N 132 W	* 11 * 17 126	•13 •12 139	•08 •08 166	•07 •09 145	•05 •07 189	•04 •06 181	•03 •05 208	•06 •05 225	•04 •04 238	•08 •08 193	•07 •08 178	•11 •12 157

LATILON	JAN	= E B	MAR	APR	MAY	JUN	JUL	AUG	SEP	007	NOV	DEC
33 N 124 W	• 0 4 • 0 8 3 3 7	.04 .07 393	• 0 3 • 0 6 3 9 2	•04 •05 437	• 05 • 08 423	•04 •06 496	•03 •07 465	.03 .05 415	•03 •05 420	•02 •05 406	•03 •05 422	•09 •12 344
39 N 125 W	•10 •12 181	.06 .10 171	.08 .15 162	•05 •10 202	.05 .10 229	• 05 • 10 273	.04 .08 311	•04 •08 267	•09 242	.04 .08 218	• 05 • 12 157	•05 •12 158
39 N 126 W	.07 .17 172	.09 .11 160	.07 .11 165	•04 •09 231	• 04 • 09 2 32	.03 .07 301	.03 .06 331	•03 •07 318	•03 •07 269	•07 •17 220	•06 •09 206	•08 •13 155
39 N 127 W	.09 .17 134	•06 •09 173	•07 •13 167	.05 .10 213	.05 .10 173	.05 .10 182	•04 •08 222	•03 •07 244	•04 •09 196	•05 •12 197	.08 .10 174	• 08 • 13 141
39 N 128 W	• 37 • 14 138	.10 .11 135	•07 •13 169	.05 .11 164	.04 .10 161	•04 •09 167	•03 •07 233	•03 •07 219	•03 •07 227	•05 •10 171	•07 •20 163	•08 •11 136
39 N 129 W	.07 .11 133	.10 .10 162	.07 .10 202	•05 •09 229	.04 .06 249	.03 .06 254	•02 •06 288	•03 •06 248	•03 •07 238	•06 •09 178	•06 •09 184	•07 •09 172
39 N 13. W	.07 .08 304	.07 .07 337	• 05 • 07 248	•03 •07 349	•03 •04 451	• 02 • 04 339	•02 •05 356	•02 •05 324	.03 .06 251	•06 •10 239	.07 .09 187	• 05 • 07 335
39 N 131 W	•13 •13 138	.11 .13 149	.06 .08 192	.04 .05 215	• 04 • 04 347	• 03 • 05 264	•02 •04 400	.02 .03 357	.04 .07 284	•06 •08 187	•09 •09 198	•08 •10 174
39 N 132 W	• 10 • 12 135	•13 •15 101	•10 •12 157	•06 •08 163	•05 •07 185	.04 .05 251	•02 •04 269	•03 •06 219	•04 •06 235	.04 .07 159	•09 •10 172	•13 •14 148
39 N 133 W	•12 •18 132	•10 •21 112	.08 .11 145	.05 .11 149	•05 •08 162	•03 •05 206	•03 •04 240	•03 •05 186	•04 •04 246	•05 •06 316	•09 •08 220	•08 •11 150
40 N 124 W	• 03 • 09 727	•03 •06 710	• 0 3 • 0 7 7 4 7	• 02 • 05 736	• 02 • 05 853	.02 .04 605	•03 •05 658	•05 •06 517	•02 •04 478	•02 •06 490	•02 •07 525	• 0 4 • 0 8 4 5 4
40 N 125 W	.07 .16 178	• 07 • 14 168	•05 •11 184	•04 •11 210	• 05 • 12 167	• 0 4 • 0 9 2 2 3	•05 •09 212	•04 •09 233	•04 •11 193	•03 •12 191	.05 .11 213	•07 •15 151
40 N 126 W	.09 .18	• 0 6 • 28 75	•11 •19 80	•05 •13 102	•10 •14 124	•06 •17 105	•05 •12 127	•04 •11 127	• 05 • 13 95	•06 •13 114	•07 •17 128	• 0 9 • 2 3 1 0 1
40 N 127 W	.09	•09 •13	•13 •16 115	.05 .11 165	•04 •09 167	• 03 • 08 196	•04 •09 204	• 0 3 • 0 6 27 0	•04 •09 162	•05 •12 169	•07 •16 122	•08 •13 123
40 N 128 W	.06 .12 287	.09	•10 •11 153	• 05 • 08 212	• 03 • 08 201	• 03 • 21 248	•02 •07 301	•03 •06 247	•04 •18 205	• 05 • 89 190	•08 •12 136	• 0 8 • 3 9 225
40 N 129 W	.07 .10	.06	• 05 • 07 276	• 05 • 07 305	• 04 • 05 279	• 03 • 06 230	•03 •05 268	•04 •06 249	• 05 • 08 243	•04 •08 253	•08 •09 218	• 07 • 11 158
40 N 130 W	.07	• 0 6 • 0 7 2 9 5	• 05 • 05 568	• 0 4 • 0 5 47 9	.03 .04 519	•02 •04 519	• 0 2 • 0 5 499	•02 •06 393	•04 •06 359	•04 •05 472	•05 •06 396	• 05 • 08 197
40 N 131 W	•10 •15 108	• 11 • 12 156	•08 •10 145	• 07 • 08 168	• 05 • 15 191	• 03 • 06 185	•04 •05 216	• 0 2 • 0 4 2 4 5	• 05 • 06 170	•07 •07 147	•08 •10 153	•11 •12 129
40 N 132 W	•11 •10 165	•10 •09 162	•08 •10 182	• 05 • 07 183	• 05 • 06 232	• 02 • 04 261	•02 •04 304	•02 •03 271	•04 •05 233	•13 •08 213	•08 •09 177	•10 •13 173
40 N 133 W	•09 •11 172	•17 •09 166	•08 •08 225	•04 •05 291	•04 •05 268	• 0 2 • 0 3 3 8 4	•02 •03 375	.02 .04 341	•03 •04 429	•11 •06 362	•08 •07 235	•09 •11 203
41 N 124 W	•08 •17 102	.04 .10 112	•05 •12 101	• 05 •21 114	• 05 • 12 125	•04 •11 120	•08 •14 114	.02 .06 180	•02 •06 208	•03 •06 212	•04 •08 209	•07 •07 178
41 N 125 W	.06 .16 204	• 05 • 14 227	•03 •09 256	•03 •11 225	•03 •10 243	•03 •09 258	•04 •09 271	•03 •08 236	•03 •10 236	•04 •11 271	•03 •11 246	•11 •17 192
41 N 126 W	• 17 • 16 77	•09 •19 77	•12 •29 82	• 08 • 15 93	•04 •16 102	• 05 • 14 87	•12 •29 84	.06 .14 87	•04 •18 91	•08 •27 118	•05 •15 101	•10 •19 98
41 N 127 W	• 12 • 22 88	•16 •17 84	•16 •26 79	•12 •17 117	•04 •10 105	•08 •18 66	•07 •14 64	•06 •12 85	•09 •20 60	•07 •15 82	•11 •19 94	•14 •21 84
41 N 128 W	·29 ·25 77	•13 •14 60	•20 •36 65	•06 •09 101	• 08 • 13 69	• 06 • 13 76	•04 •13 83	•04 •14 77	• 05 • 12 65	.08 .12 89	•16 •24 83	•14 •14 86
41 N 129 W	•13 •18 103	•12 •18 97	•50 •43 108	•05 •08 120	.06 .09 119	•03 •06 154	•03 •07 137	•05 •07 122	•04 •09 118	•09 •11 111	•10 •12 107	•20 •14 113
41 N 130 W	•18 •14 113	• 11 • 16 156	•11 •09 164	•04 •05 216	•04 •06 218	• 03 • 05 274	•02 •05 253	•02 •05 295	•05 •06 199	•08 •08 206	•08 •10 154	•10 •15 122
41 N 131 W	• 89 • 12 175	•12 •16 202	.09 .10 221	•05 •06 289	•04 •06 268	•02 •04 270	•05 •11 344	•03 •03 341	•04 •05 247	• 05 • 08 227	•08 •12 195	.09 .11 151
41 N 132 W	•10 •12 116	•14 •18 113	•10 •12 115	•08 •09 155	•08 •09 161	.05 .05 161	•03 •05 171	•04 •07 152	•06 •07 148	•09 •08 162	•12 •11 139	•12 •17 131
41 N 133 W	•12 •16 117	•12 •18 99	•13 •16 115	•07 •09 133	.08 .08 118	•04 •05 157	•04 •26 152	•03 •05 168	.05 .07 141	•08 •11 137	•15 •12 113	•10 •13 123

LATILON	JAN	÷ E B	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
42 N 124 W	.08 .15 103	.06 .14 107	• 06 • 12 74	•05 •15 92	•05 •13 96	.05 .09 112	•05 •13 128	•05 •13 90	.03 .10 112	.02 .09 112	•07 •17 88	- 16
42 N 125 W	•06 •12 221	• 06 • 11 241	•04 •08 273	.03 .11 258	•04 •14 214	.03 .10 303	•03 •09 296	•03 •09 238	•03 •10 243	•03 •08 268	•03 •10 272	•06 •15 196
42 N 126 W	•17 •28 73	·15 ·25 60	•15 •21 77	-08 -14 80	• 08 • 15 75	*06 *13 61	•07 •17 63	•04 •12 76	•08 •21 •50	·13 ·23	•09 •15 77	•17 •23
42 N 127 W	*18 *21 86	• 0 9 • 17 82	•18 •23 95	•10 •21 89	• 05 • 09 83	.07 .11 73	•08 •12 72	.06 .11 64	-08 -18 -59	•11 •15 94	•12 •15 114	·14 ·16 102
42 N 128 W	• 24 • 20 • 94	• 10 • 18 64	*23 *24 87	•10 •13 68	•07 •14 70	*10 *25 67	•06 •15 67	.07 .18 61	•09 •11 53	.08 .13 82	*14 *25 86	•16 •20 110
42 N 129 W	•12 •16 212	.10 .13 166	•06 •09 260	•05 •07 244	•06 •07 241	- 04 - 07 157	•04 •06 143	-02 -05 264	:11 :13 48	·14 •19 61	•11 •13 152	·10 ·11 257
42 N 130 W	•10 •14 146	•14 •17 142	•12 •12 124	.07 .09 93	.05 .09 100	.04 .07 139	-04 -09 128	·32 ·39 100	•21 •09	•12 •12 76	•12 •10 137	•14 •19 114
42 N 131 W	.07 .09 187	•14 •13 158	.07 .10 199	.07 .05 216	.05 .07 160	.04 .05 183	•03 •04 280	.02 .04 288	•05 •07 206	•10 •09 166	•10 •13 158	·13 ·12 138
42 N 132 M	•13 •17 121	*11 18 120	*15 *13 123	•08 •09 185	•05 •05 194	• 04 • 05 228	•03 •05 249	•04 •05 234	.04 .06 190	•08 •09 159	•12 •19 116	·12 ·21 126
42 N 133 W	• 15 • 15 144	•16 •18 110	*13 *10 167	.07 .09 181	• 06 • 07 256	• 05 • 08 257	•03 •04 233	.03 .03 259	.04 .05 219	•08 •07 200	•15 •17 119	·12 ·15 139
43 N 124 W	• 11 • 14 95	.09 .11 100	*06 *12 80	.03 .12 84	*06 *10 87	.03 .07 104	.03 .10 92	•03 •09 76	•02 •06 123	.02 .08 103	*08 *21 72	.07 .13 104
43 N 125 W	•06 •13 254	•04 •13 272	•06 •10 255	.03 .09 241	•03 •09 241	20. 70. 280	• 0 2 • 0 6 30 4	.02 .06 213	.02 .09 281	·10 •07 456	•04 •09 423	*06 *15 224
43 N 125 W	•13 •19 71	•13 •22 58	*15 *14 61	•13 •12 59	.06 .14 59	.06 .17 41	•07 •16 42	•03 •17 52	•07 •21 47	•07 •12 87	•13 •15 77	*12 *23 68
43 N 127 W	*24 *33 71	•13 •22 64	•12 •20 84	•10 •18 64	•06 •09 64	.05 .13 48	-04 -10 58	•11 •17 •54	•07 •14 53	.09 .13 105	*11 *15 110	• 20 • 25 • 83
43 N 128 W	*18 *21 85	*23 *31 78	* 30 * 22 75	•11 •14 68	*06 *11 69	•04 •08 64	.09 .11 64	•05 •08 54	•07 •12 71	+09 +17 88	•14 •15 104	*15 *22 105
43 N 129 W	*14 *17 136	*16 *21 98	*14 *15 123	•09 •13 93	*12 *13 97	*09 *23 87	•05 •08 76	.05 .10 86	•07 •08 77	*12 *17 82	*15 *18 105	•13 •23 107
43 N 130 W	• 08 • 10 330	•08 •12 156	•08 •10 306	.05 .08 317	.03 .05 346	.02 .05 369	•02 •05 320	.02 .04 330	•03 •05 288	.09 .10 173	•12 •12 118	.07 .10 332
43 N 131 W	• 12 • 15 119	.10 .14 149	*11 *11 139	+11 +14 117	.05 .10 133	.04 .05 150	•04 •06 150	.02 .05 169	.05 .09 126	.10 .09 130	•11 •12 116	•19 •22 103
43 N 132 W	* 15 * 19 99	·12 .26 101	•13 •15 114	*11 *15 95	•10 •09 85	.07 .08 83	•05 •07 94	.05 .08 84	.08 .10 75	•14 •12 81	•19 •20 78	•17 •22 79
43 N 133 W	• 14 • 13 127	•12 •17 112	.09 .10 117	.09 .09 147	80. 90. 122	.04 .06 128	.04 .05 115	.06 .05 129	.07 .07 100	.09 .12 149	*13 *12 143	.20 .47 121
44 N 124 W	• 09 • 17 126	• 08 • 10 133	•09 •16 118	•04 •08 136	•03 •06 162	.01 .07 133	.02 .05 161	•01 •05 95	•03 •05 138	.03 .08 156	•04 •09 174	•11 •16 94
44 N 125 W	• 07 • 14 229	• 05 • 10 252	•05 •09 305	•03 •08 237	• 06 • 07 259	• 02 • 06 271	•01 •05 341	.01 .05 286	.01 .04 540	.02 .07 492	•03 •06 437	.08 .18 221
44 N 126 W	• 17 • 23 84	·14 •25 45	•09 •16 68	•18 •65 52	.06 .10 73	.05 .12 63	•05 •17 61	:02 :09 76	*10 *24 39	•16 •19 80	•09 •13 76	·17 ·32 65
44 N 127 W	• 20 • 28 75	•20 •23 •62	*13 *21 77	*14 *17 55	• 08 • 12 77	•08 •11 55	•05 •15 42	:04 :09 64	*10 *12 61	•13 •16 77	.09 .14 103	:17 :30 81
44 N 128 W	• 19 • 22 99	• 16 • 23 • 124	•15 •17 103	•1) •10 90	.06 .11 109	•27 •47 83	•05 •11 79	.05 .08 74	.05 .10 124	•05 •07 288	.07 .12 270	·25 ·22 103
44 N 129 W	• 15 • 18 99	•14 •17 94	*16 *17 115	•10 •13 125	.07 .06 102	.05 .11 107	.04 .07 104	.04 .08 94	.05 .08 119	.07 .11 168	•16 •12 145	•18 •22 139
44 N 130 W	• 10 • 14 134	• 11 • 15 • 116	*16 *16 126	•10 •15 102	.09 .14 110	.07 .20 110	•04 •09 110	•06 •09 91	•05 •09 113	•09 •14 125	•13 •20 117	·16 ·19 117
44 N 131 W	• 10 • 12 231	·09 ·10 194	.07 .07 318	• 05 • 06 256	• 04 • 06 288	.02 .04 417	•02 •04 384	•05 •03 385	.03 .05 381	•04 •07 372	•06 •08 273	·13 ·14 238
44 N 132 W	•12 •14 170	·11 ·17 150	*11 *11 167	•07 •07 181	•06 •08 177	• 05 • 06 169	•03 •05 175	•03 •05 149	•05 •08 143	.08 .10 173	•13 •12 151	•11 •17 177
44 N 133 W	• 14 • 16 137	• • 15 • 16 133	•13 •12 118	•12 •11 101	.07 .07 105	•08 •07 110	•06 •06 108	•04 •07 109	.07 .10 100	•11 •10 137	•13 •14 116	•11 •15 135

LATILON	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
45 N 124 W	•11 •12 175	•06 •08 188	.05 .09 164	.04 .09 150	•04 •06 219	•01 •04 230	•01 •05 204	.01 .04 182	.02 .07 161	.04 .07 211	•10 •12 175	•11 •14 141
45 N 125 W	.06 .10 277	•04 •08 338	.06 .10 329	•04 •06 277	• 03 • 05 344	•01 •05 349	.01 .04 368	.01 .04 238	.01 .03 422	•03 •06 458	•05 •10 355	•10 •14 233
45 N 126 W	•13 •24 109	•08 •12 116	•14 •18 83	.07 .15 87	.05 .08 103	.03 .08 129	.05 .13 87	•05 •08 96	.09 .12 99	•14 •10 115	•11 •16 113	•18 •27 101
45 N 127 W	•19 •26 98	•14 •17 101	•12 •17 115	.09 .18 82	.06 .11 104	•05 •08 75	.06 .11 81	.04 .07 81	•06 •12 72	•12 •12 110	•11 •15 132	•20 •29 85
45 N 128 W	• 11 • 20 102	.13 .14 99	•10 •15 113	.09 .21 105	• 04 • 04 297	•05 •07 100	•05 •08 79	.03 .07 147	.07 .11 71	•17 •15 113	•14 •13 121	•29 •27 85
45 N 129 W	.14 .19 119	•13 •13 97	•11 •16 144	.08 .10 109	.06 .09 100	.05 .07 115	•05 •08 97	•04 •07 95	•08 •08 123	•19 •14 117	•12 •18 131	•11 •16 109
45 N 130 H	• 06 • 07 5 07	• 06 • 08 500	• 05 • 05 539	•04 •05 554	•03 •05 386	•03 •03 420	•01 •04 559	•03 •03 307	.03 .03 471	•06 •06 379	•07 •09 482	•08 •10 425
45 N 131 W	.08 .10 257	•09 •10 273	•07 •07 284	• 05 • 07 223	•04 •06 260	.03 .04 251	•09 •04 274	•03 •05 222	•05 •07 209	.07 .08 216	•10 •12 199	•11 •12 255
45 N 132 W	•12 •17 166	•12 •12 143	•11 •10 163	•08 •08 192	.07 .07 170	•04 •06 167	•05 •06 129	.05 .06 141	•06 •08 145	•10 •11 160	•09 •13 132	•18 •16 127
45 N 133 W	·23 ·29 61	• 29 • 25 5 3	•23 •22 59	•15 •16 59	•16 •12 55	.07 .09 68	•08 •08 56	•09 •12 45	.07 .09 56	•13 •15 93	*16 *18 79	• 25 • 34 75
45 N	. 04	.04	.03	• 03	.03	.02	.01	.01	.02	.03	.04	•06
124 M	907 .09	.07	.08	•05 694 •05	709	.02	·02 •02	.01	.04	.05 818 .07	719 • 07	.13
125 W	·10 220 •13	·10 ·13	269 •18	204	246	291	314 • 04	255	269 •19	285	258 •24	•14 154 •27
125 W 45 N	.19 100 .14	•19 93 •11	•17 84 •11	•13 74 •08	•10 82 •05	•13 78 •07	•08 95 •05	•08 87 •03	•16 88 •06	•27 89 •15	•27 115 •12	•21 76 •18
127 W	•16 126	13 120	14 140	•11 108 •08	.05	•10 96	.09 162	•06 98	•11 95	.15 105 .11	.14 181 .14	15 114
128 W	.17 139	.15 148	.11 164	10 118	.08 109	.08 106	.08 113	.06 110	.07 118	•12 139	•17 145 •18	•13 145
123 W	.17 145	16 136	·11 154	·11 129	.09 116	.06 115	.06 107	•05 120	.09 113	14 133	•12 136	.18 145
137 W	14 131	.16 130	12 152	12 123	.06 120	08 140	07	98	.12 102	.11 123	12	16 130
45 N 131 N	•14 •15 141	•12 •16 107	•11 •15 112	.10 158	• 05 • 08 157	.05 142	.07 123	•05 •05 144	.08 .08 141	•13 •13 125	•15 •17 154	•12 •12 148
45 N 132 W	•10 •10 240	•09 •09 182	•09 •08 254	•08 •08 295	•03 •05 229	•06 •06 145	•14 •12 138	.06 .08 121	.07 .08 151	•09 •12 121	•13 •14 114	•16 •17 134
46 N 133 W	•15 •14 106	•17 •20 92	•15 •15 83	•12 •12 95	•14 •14 92	•07 •08 77	•06 •09 74	•05 •06 86	•10 •13 90	•13 •12 105	•17 •18 91	•24 •23 108
47 N 124 W	•18 •15	.10 .11	.07 .10 81	•08 •10 83	•04 •09	• 02 • 03	•02 •03	•02 •04	•03 •04	•10 •06	•07 •09 173	•18 •19 74
47 N 125 W	•10 •12 228	.08 .10	• 06 • 08 252	•07 •09 195	•04 •05	• 02 • 03 356	.01 .02 450	•01 •02 436	.03 .03 274	• 0 4 • 17 357	• 07 • 09 304	• 11 • 15 155
47 N 125 W	•14 •16	•12 •13	• 10 • 11 177	•09 •10 185	• 05 • 08 174	•06 •08 152	•03 •06 143	•03 •06 174	•09 •10 157	•09 •11 192	•10 •11 223	•12 •14 138
47 N 127 W	• 15 • 13 171	· 10 · 12 177	•11 •09 215	.08 .12 171	• 05 • 08 177	•08 •11 175	•03 •06 149	•05 •06 151	•08 •10 181	•07 •10 170	•11 •13 186	•14 •13 159
47 N 128 W	•15 •12	•16 •19 127	•09 •12	•10 •13 130	• 06 • 06 128	•09 •12 119	•08 •06 108	•04 •08 116	.06 .10 183	•08 •09 293	•10 •19 154	•15 •17 125
47 N 129 W	•19 •18 88	·21 ·20	•13 •16 83	.09 .13 94	• 06 • 06 99	•06 •08 95	•64 •07 94	• 05 • 08 86	•08 •09 107	•09 •10 198	•11 •16 115	•16 •18 98
47 N 130 W	•10 •12 252	•10 •12 283	•10 •09 299	• 05 • 06 275	• 05 • 05 213	• 03 • 06 247	•02 •03 270	• 02 • 05 242	.07 .06 257	•13 •14 148	•19 •21 78	• 0 9 • 0 8 2 8 2
47 N 131 W	• 08 • 13 237	• 12 • 16 120	•08 •09 180	• 05 • 08 200	•04 •07 198	• 0 3 • 0 4 267	•03 •04 237	•03 •05 316	•04 •06 326	•11 •11 202	•10 •14 185	•10 •12 262
47 N 132 W	•11 •16 159	• 08 • 11 206	•09 •10 171	•13 •09 157	• 04 • 06 265	• 0 3 • 0 3 374	•03 •04 292	•04 •05 259	•04 •06 232	•08 •10 257	•08 •10 248	• 11 • 17 196
47 N 133 W	•16 •17	•18 •19	•19 •15	•11 •17	•10 •14 75	•08 •11 75	•07 •08	•06 •10 85	•08 •15 96	•11 •12 114	•20 •22	•18 •17 91

LATION	JAN	= E B	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
48 N 125 W	•06 •05 930	• 06 • 06 739	•04 •06 848	• 0 3 • 0 4 77 0	•02 •02 767	•01 763	.01 .01 1213	.01 .02 1138	•02 •03 754	•03 •05 795	• 0 5 • 0 5 82 3	•07 •08 629
48 N 125 W	•12 •12 304	·11 ·09 209	•10 •08 252	•07 •07 198	.05 .09 185	•05 •05 197	• 0 2 • 0 3 32 4	•02 •02 327	•03 •04 298	.07 .05 400	•09 •09 331	•12 •12 201
48 N 127 ₩	• 12 • 12 143	• 12 • 13 131	.10 .12 130	.09 .14 101	.08 .10 151	• 06 • 07 140	•06 •07 137	.04 .06 134	•04 •06 206	•09 •11 136	•09 •11 247	•17 •15 137
48 N 128 W	•12 •14 114	•11 •16 131	•09 •09 139	•06 •08 147	•06 •08 148	.05 .06 133	.05 .08 118	•05 •07 115	.08 .14 130	•13 •14 134	•14 •16 142	•15 •16 136
48 N 129 W	•12 •17 111	•14 •12 125	•10 •11 126	.07 .10 110	•06 •08 147	.05 .08 113	•05 •08 135	•04 •08 131	-08 -14 141	•11 •15 117	•13 •16 114	•13 •15 122
48 N 130 W	•14 •15 132	.11 .17 108	.09 .10 136	•13 •11 123	.07 .09 131	.45 .18 138	•06 •09 120	.04 .06 148	-08 -09 179	•10 •15 131	•14 •16 134	•15 •16 141
48 N 131 W	• 19 • 18 143	• 11 • 13 126	• 11 • 12 134	• 05 • 07 167	.05 .06 207	.04 .05 168	.03 .04 173	.04 .06 161	.07 .09 178	•10 •14 169	•11 •15 162	•10 •15 184
48 N 132 W	• 0.8 • 0.8 322	.07 .07 302	•06 •06 335	• 05 • 07 321	.03 .04 380	.03	.04 .06 192	.04 .05 192	•07 •08 209	•13 •13 161	•11 •12 197	•08 •10 295
48 N 133 W	• 12 • 13 124	•14 •16 95	•13 •15 116	•08 •12 134	•08 •08 136	.07 .07 133	•04 •05 137	.06 .09 107	.10 .09 147	•13 •12 143	•12 •16 147	•15 •15 153
48 N 134 W	•16 •17 82	•18 •26 107	•15 •14 97	•13 •13 104	.08 .07 118	.06 .08 118	•05 •05 94	•08 •07 91	•10 •13 96	•22 •34 114	•13 •14 132	·28 ·25 114
49 N	.20	.09	.20	.08	.05	.06	.03	.02	.05	.07	.09	.15
126 W	•13 98 •12	.09 140 .09	.13 124 .10	.08 141 .08	.05 149 .06	145	406	378 .03	235	207	122 .26	·16 106 •12
127 W	•12 152 •13	.10 159 .08	•12 170 •10	.10 193 .06	.06 213 .04	209	.03 331 .04	-03 283 -04	.06 268	.08 223 .10	·16 212 •10	.11 169 .14
128 W	.13 185	.12 175	.09 234	.09 197	+05 453	273	325 04	322 0.3	272	207	10 267	.13 184
129 W	14 184	.09 190	.09 212	245	289	.05	272	:05 251	249	233	11 246	14
49 N 130 W	•13 •12 308	.19 .11 316	.07 314	*06 *07 316	•04 •05 298	•03 •04 298	•03 •05 303	•03 •05 325	•05 •06 345	•07 •09 280	•07 •08 332	•11 •12 304
43 N 131 W	• 09 • 09 430	.09 .09 388	.05 .08 411	• 05 • 08 452	•04 •05 330	• 02 • 03 520	•04 •05 458	•03 •04 330	•05 •06 420	•05 •06 362	•07 •08 368	.07 .11 406
43 N 132 W	*15 *14 182	.10 .11 197	.10 .11 189	•09 •09 191	• 05 • 05 240	.06 .05 211	.03 .04 235	•04 •05 220	•06 •08 225	•09 •10 227	•11 •10 235	•12 •11 217
43 N 133 W	•11 •14 196	•10 •12 185	•11 •08 225	•07 •09 250	*06 *06 235	•04 •05 244	•03 •04 220	•04 •05 272	.06 .07 251	•11 •10 233	*13 *13 238	•11 •13 190
49 N 134 W	•14 •13 200	•12 •11 179	.10 .09 201	•09 •12 210	•05 •06 249	• 05 • 05 240	•04 •04 227	•04 •05 252	•07 •07 277	•11 •12 257	*11 *12 262	•11 •12 211
49 N 135 W	•12 •11 183	•11 •11 165	•10 •11 199	•09 •12 217	•08 •07 213	•08 •05 200	.05 .05 184	•05 •06 211	•06 •07 263	•08 •11 243	.09 .11 306	•11 •10 241
5] N 127 W	• 38 • 26 19	• 17 • 16 30	• 28 • 33 21	•55	• 09 • 02 5 9	•09 •10	•05 •05	•04 •01	•03 •02	·10 ·07	2.75	.54
50 N 128 W	• 18 • 17 • 68	• 33 • 35 48	•19 •13	•11 •17 63	•13 •13 •62	.08	.09	.06	•09 •12	•10 •10	•12 •17	• 32
50 N 129 W	· 17 · 20 93	•18 •16 78	•13 •17 89	•11 •15	• 06 • 07 • 136	.08	.05	•05 •07	.07	•13 •14	.20	• 22
53 N 130 W	•19 •14 124	•13 •13 •13	•15 •14 121	.07 .10	• 05 • 06	• 06	•04	.05	.05	•11 •11	•13 •20	•11
50 N 131 W	•13 •17	• 12 • 13	•10 •11	.07	• 05	.05	.04	.04	.07	•10	•11 •12	•12
50 N 132 W	• 12 • 16	.12	•12	• 08 • 08	• 05	.04	.04	.09	-06 -08	•12 •16	187 •13 •12	154 •14 •23
50 N 133 W	•14 •16	•14	•15	• 06	.06	• 06 • 07	.05	•04 •05	.07	148 •14 •12	170 •16 •15	125 •16 •17
50 N 134 W	•12 •11 727	• 0 8 • 1 0	.07	• 06	• 04 • 05	• 03 • 04	• 0 3 • 0 4	230 •04	-07 -08	166 •11 •11	147 •11 •10	144 • 10 • 12
50 N 135 W	•14	·13	•11	•08 •08	• 05 • 05	• 04	.03 .04	.05 .06	• 05 • 06	163 •10 •11	284 •11 •13	289 •15 •15
50 N 136 W	.08	.09	• 05	• 04 • 05	- 04 - 04	263 •04 •04	237 •03 •03	237 .04 .04	48 % • 0 4 • 0 6	228 .07 .08	222 .08 .07	188 •09 •09
	485	463	649	681	595	537	384	406	446	499	536	566

APPENDIX III

Monthly Wind Stress Curl Distributions

The monthly mean wind stress curl distributions are displayed in Charts 37 to 48. The plotted values are estimates of the vertical component of the wind stress curl, calculated by applying a finite difference, spherical coordinate curl operator to the fields of monthly mean surface wind stress. The contoured values are plotted in units of dyne cm⁻²per 100 km. A value of 1.0 dyne cm⁻² per 100 km is equivalent to 1.0×10^{-7} dyne cm⁻³. The contour interval is 0.25 dyne cm⁻²per 100 km. Positive wind stress curl is associated with surface Ekman divergence (upwelling). Negative values are shaded. These correspond to surface Ekman convergence (downwelling).

In the following charts, the month is indicated in the figure legend in the upper right corner of the charts. The coastline configuration is superimposed on the grid as a visual aid and does not represent a conformal mapping.

137	136 135 134 133 132 131 130 129 128 127 126 125 124 123 122 121 120 119 118 117 116 115 114 113 112 11
	and a second sec
3	9,68 0,03 0.11-0.05 0.24-0.11-0.12 0.19 0.04 0.60 VANCOUVER ISLAND
3	0.12 0.05 0.20 0.05 0.15 0.05 0.45 0.04 0.37
3	0.73 8.46 8.20 5.14-0.16 0.39 0.30 0.30 0.30 0.30 0.30 MONTEREY, CALIFORNIA
7	0.39 0.35 0.73 0.09(0.45 0.01 0.44 0.35 0.17 0.02 WIND STRESS
6	0.76 0.44 0.22 0.16 0.20 0.35 0.04 0.11 0.62 0.19 CUUMBIN RIVER CURL
	0.36 0.31-0.08-0.16 0.02 0.45-0.21-5.04 0.12-0.13 (DYNE CM-2/100KM)
4	0.10 0.17-0.06-0.93 0.32 0.07-0.03 0.31-0.04-0.37
3	0.35 0.45-0.40-0(57-0)55 0.56 0.30-0.02=0.01-0.26 CAPE BLANCO FOR
2	0.11 0,0470.43-0,33, 0.21 0.53-0.29 0.47 0-38-0.35 JANUARY
1	-0.15 0.00-9.11 D.D9-0.08-0.48 0.04 0.23 0.22 0.39
	8.21 0.01 9.19-0.42-0.38 0.14-0.41 0.05 0.73 0.64
	a 36 a 24 a 32 a 34 a 37 a 34 - 3 36 - 3 24 0 - 37 a 30
/	0.12 0/00-0.13 0.06 0.05-0.23 0.11 0.02 0.14 0.23
6	0.14 0.03 0.02-0.06 0.09 0.07 0.06-0.16-0.09 0.14
5	0.01-0.11-0.00 0.09 0.09-0.13-0.25-0.07 0.00 0.23 POINT CONCEPTION
4	-0.02-0.11-0.05-0.11-0.07 (0.02-0.12 0.12 0.34 0.38
3	0.13 0.07-0.12-0.20-0.10-0.01 0.07 0.24 0.34 0.24
2	-0.07 - 9.04 - 0.07 - 0.12 - 0.02 0.05 0.01 0.17 0.19
1	-0.11-0.10 0.05 0.03-0.15-0.15 0.03 0.09 0.19
0	-0.24-0.33 0.12-0.18-0.22 0.02 0.26-0.02 0.04-0.01
.9	-0.45-9.20 9.00- 0.01 0.17 0.10-0.19-0.03-0.07-0.05 D
.8	6.23-0.14/0.07-0.05-0.32-0.01-0.06-0.12-0.03 0.02 PINTA FIRENTA
	0.05 9.00-0.15-0.26-0.09-0.04-0.16 9.03 0.07 0.02
26	-0.05-0.16-0.14-0.12 0.00 p. 30 0.03-0.07
	a 12 a 11-a az a de a 12-a 16-a 12 data zo a de
24	
	10.28 0.12-0.04-3,05 6.13 0.06-0.13-0.14-0.01-0.
22	-8.13-8.05-0.09-0.14-0.29 0.00 0.13 0.01-3.07-0.
1	-0.35-0.17-0.13-0.07-0.31-0.16 0.02 0.07 0.01-0.

Chart 37.-Wind stress curl for January.

NCC - TDF-11 - ONE DEGREE SUMMARIZATION



Chart 38.-Wind stress curl for February.

13	7 136 135 134 133	132 131 130 129 128	3 127 126 125 124 123 122 121 120 119 118 117 116 115 114 113 112 111
		G	Surger and a second sec
50	0.2070.05 0.59 0.12	0.00 0.18 0.18 0.15-0.	S 0.54 VANCOUVER ISLAND SE
49	8.10 0.23 0.00	0.11 0.26-0.07 0.03 0.1	NATIONAL MARINE FISHERIES SERVICE
48	C. 02 0. 15	0.16 0.00 0.16 0.08 0.0	ACT R. 18 G. 39 MONTEREY, CALIFORNIA
47	ØØ6	5.05 0.3P 0.16 0.05 0	27 0.31 0.16 0.08 0.20 WIND STRESS 4
46	>0.23	0. 19-0.08 0.01 3.00 0.2	22 0.12 0.27 0.43 0.43 COLUMBIR RIVER CURL 4
45	-Ø=04	0.03-0.02-0.07 0.21 0.2	32-8.84 0.19-0.07-0.32 (DYNE CM-2/100KM) 4
44	Dec	2. 08 10. 30 0. 44 0. 17 70.	41-0/02 0.30 0.32 0.11
43	-D. 19	-0.07 0.27 0.42-0.16-0.	21 0.11 0.23 0.46-0.62 CAPE BLANCO FOR 4
42	0.07	0.01 0.17-1.39-0.54 0.	74 9.35-6.19-9.38 0.60 MARCH 4
41	-Ø _× 14	-0.02 0.02-0.13-0.36 0.	19 0.26 0.07-0.16-0.21
40	Ø. 12	0.04 0.16 0.04-0.20-0.	35 0,00 0,403 6,28 0.05 CAPE MENDOCINO
39	11.0	0.07 0.01-0.21-0.11-0.	13 0.13-0.02 0.15 0.23
38		-0.45 0.01 0.1370.09-0.	14 8.04+0.05-0.18 0.24 0.22
37		0.12 0.03-0.11-0.	17-0.04-0.02-0.07-0.03 0.11 0.25
36		-0.10-0.05 0.03-0.	08 0130-0.01 0.01-0.15-0.13 0.00
35		0.04-0.07 8	20 0.00-2.03 0.06 0.05-2.06-0.09 0.32 POINT CONCEPTION
34		Ø.	07 0.01-0.07-0.05-0.05 9.00 0.28 0.59 0.78
33			-0.88-0.05-0.10-0.24-0.04 0.03 0.08 0.35 0.45 9-24
32			0.02-0.07-0.15-0.25-0.04-0.21 0.21 0.24 0.20
31			-9.01-0.17-9.09-0.09-0.23-0.11-0.05-0.08 0.14 0.20
30			0.23-0.22-0.14 0.16 0.02-0.10-0.12 0.00 0.02-0.04
29			0. 2-0228 0. 10-0. 06-0.51 0. 10 0. 08-0. 10-0. 07-0. 39
28			-0.08-0.06-0.14-0.01-0.08-0.05 0.06-0.02-0.10-0.13 PUNTA EUGENIA
27			0.15-0.15-0.07 0.06-0.28-0.14-0.03 0.00 0.04 0.09
26			-0.15 0.02 0.02-0.13 0.32 0.27 0.06 0.11 0.05 0.00
25			-0.07 0.00-0.05-0.24-0.05 0.00-0.04-0.01-0.02 0.08
24			-0.04-0.03-0.09-0.12-0.10-0.06-0.04-0.01-0.05 0.03
23			0,28-3.09-0.07-3.16-0.06-3.06-0.83 0.02-0.94-0.11
22			-0.06 0.02-0.04-0.12-0.16 0.00-0.05-0.02 0.07
21			-0.12-0.04 0.00-0.12-0.13-0.08-0.14-0.15-0.05 0.04
1	37 136 135 134 133	3 132 131 130 129 12	28 127 126 125 124 123 122 121 120 119 118 117 116 115 114 113 112 111

Chart 39.-Wind stress curl for March.

NCC - TDF-11 - 1	UNE DEGREE SUMMENTIZATION 134 133 132 131 130 129 128 127 126 125 124 123 122 121 120	119 118 117 116 115 114 113 112 111
0 0.39 2.29-	0.02-0.10 0.22-0.16-0.02-0.21 0.32 0.88 VANCOUVER ISLAND	
9 0.28	D.27 0.86 8.19 8.09-0.12-0.84-8.84 8.29-0.33 2	NDAA NATIONAL MARINE FISHERIES SERVICE
8	0.10-0.03 0.18 0.07 0.03 0.14-0.05 0.10 0.52	MONTEREY, CALIFORNIA
7	0.17 0.11-2,24 0.20 0.06-2,02 0.09 0.04 0.01 0.02	WIND STRESS
6	3.00 0.02 0.15 0.01 0.18 0.14 0.13 0.00 0.16 0.36 COLUMBIA RIVER	CURL
5	0.15-0.01-0.05-0.05 0.16 0.24-0.05 0.40 0.04 0.10	(DYNE CM-2/100KM)
4	-0.17-0.19 0:26-0.06-0.19 0.18 0.75 0.31 20 37-0.10	
3	-0.17 0.15-0428-0.16 0.32-0.13-0.18 0.27 0.00 0.03 CAPE BLANCO	LONG TERM MEAN FOR
2	0.01 0.07-0.14 0.29-0/01-0255 0.15-0.07-0.17 0.37	APRIL
1	-0.01-0526-0.16 0.12-0.08-0.11-0.27-0.24 0.05-0.11	
.0	-0.67-0.02-0.16-0.10-0.07-0.16-0.06-0.24-0.18 0.22	
9	0.09 0.11 0.11-0.15 0.50-0.05-0.03-0.18 0.09 0.4	
8	0.04 0.02-0.01-0.17-0.11-0.24-0.03 0.16 0.29 0.26	
7	-0.12-0.01-0.13-0.08-0.01-0.03-0.01 0.09 0.15 0.38	
16	0.05-0.04 0.04 0.04-0.13-0.14-0.03-0.19-0.01 0.28	
35	-0.15 0.04-0.12 0.35 0.21 0.22 0.03 0.03 0.07 0.13	INT CONCEPTION
34	-0.12-0.12-0.05 0.10 0.08-0.02 0.05 0.21 0.5	rie.97
13	-8.94-0.91-9.16-9.12-0.95 6.05 0.14 8.2	0.49 0.33
12	-0, 10-0, 15-0, 14-0, 16, 0, 05-2022, 0, 17	0.26 0.19 0.16
31	- 9, 38-9, 22-9, 13-9, 15 p. 00 p. 08-9, 07	10-00-00 0 0 19 1 0 00 00 00 00
	10.10 m 1.10 m. 17 m. 12 15 m	
	<u>16. 17-10. 19-10. 08-10. 07-10. 19</u>	PUNTA EUGENIA
	<u>-0.2</u> -0.22-0.03 0.10	9-9.11-0.03 8-93 D.06-10.02 0.10
·D	-0.15-0.10-0.1	1-0.02 0.06-0528-0.09 0.07 0.03 0.00
	-0.11-0.05-0.23	20.22 0.01 0.20-0.16-0.06 0.00 0.03
-4	-6.9	5-8.16-0.22 0-01-0.97-0.02 0.00-0.04 0.00-0.0
23	-0, 93	2 0.02-0.05-0.09 0.06-0.20-3.05 0.06fa.01 0.0
22	8.11	4-0.12-9.14-0.23-0.11 0.13-0.16-0.08 0.04-9.0
21	-8.29	8-0.32 6.10-0.04-0.40-0.06-0.04-0.12 0.06-0.0
137 136 135	134 133 132 131 130 129 128 127 126 125 124 123 122 121 120	119 118 117 116 115 114 113 112 111

Chart 40.-Wind stress curl for April.



Chart 41.-Wind stress curl for May.

NCC - TDF-11 - DNE DEGREE SUMMARIZATION



Chart 42.-Wind stress curl for June.



Chart 43.-Wind stress curl for July.

NCC - TDF-11 - ONE DEGREE SUMMARIZATION



Chart 44 .- Wind stress curl for August.

DI TOTTON

N	C - 10F-11 - UNE DEGREE SUMMERIZATION	
Г	37 136 135 134 133 132 131 130 129 128 127 126 125 124 125 122 121 120 115 116 117 116 115 116 116	1
	A PART A PART A PART A PART A PART A PART VANCOUVER ISLAND	50
510	NDRA	49
49	RETIONAL MARINE FISHERIES SERVICE	10
48	-0.31-0.27-0.01-0.04-0.24 0.15 0.42 0.33 0.24 MONTEREY, CALIFORNIA	40
47	-0.47-0.14-0.07-0.01-0.05 0.05 0.29 0.31 0.03 0.03 WIND STRESS	47
46	-0.31 0.00 0.01-0.28-0.08 0.02-0.18 0.08 0.23 0.33 COLUMBIA RIVER CURL	46
45	-0.53-0.18 0.05-0.08-0.30-0.02 028+0.18 0.06 0.18 (DYNE CM-2/100KM)	45
44	0.21-3.26-0.19-0.21-0.11 0.06 0.05-0.07 9 00 0.17	44
43	-0.45-0.15-0.24-0.07 0.17-0.23 0.02 0.04 0.16 0.43 CRPE BLANCO FOR	43
42	-0.20-0.19-0.05 0.01-0.12-0.37-0.39 0.76 0.44 0.33 SEPTEMBER	42
41	-0.12-0.03-0.14-0.01 0.04-0.45-0.53 0.25 0.65 0.66	41
10	0, 05 0, 02-0, 18-0, 18-0, 09 0, 02-0, 10 0, 12 0, 21	40
20		39
00		38
35		37
37	-0.17-0.19-0.16-0.82 0.03-0.10-0.10 0.03 6.2	26
36	-0.22-0.08 0,00-0.10-0.11 0.03 0.11+0.08-0.04 0.11	50
35	-0.02-0.05 5.01 0.03 0.02 0.00 0.02 0.05 (0.37 POINT CONCEPTION	35
34	0.16-0.02 0.01-0.19-0.01 0.06-0.03 0.33 0.53 0.53	34
33	0.86-0.07-0.01-0.06-0.08-0.12 0.05 0.34 0.49 8.18	33
32	-2.08-0.13-0.10-0.05-0.07-0.03 0.24 0.17 0.19 0.23	32
31	-9.11-0.12-0.09-0.13 0(30 0.16 0.03 0.07 0.15	31
30	0.01=0.01=0.15=0.05=0.02=0.03=0.03=0.02 9.01 0.06	30
29	-0.06-0.09-0.25 0.00-0.13 0.00 0.91-0.15-0.04-0.10 1 B	29
28	0.11 0.09-0.19-0.02-0.03-0.06 0.17-0.03-0.12-0.09 PUNTA EUGENIA	28
27	8.22 0.62-8.83-0.15-0.28 8.91 0.16 0.08 0.83 0.89	27
26	-0.10-0.17 (0.18)0.06 0.17 0.03 0.18 0.06 0.04 0.06	26
25	-0.11-0.17-0.25-0.26 0.01 0.05-0.06 0.00 0.11 0.11	25
21	0.04-10.02-10.02 0/00-10.17 0.09 0.05 0.07 0.18 0.01	224
25	a as a as a and a la a 12 h at a la a	123
20		122
24		22
21	137 136 135 134 133 132 131 130 129 128 127 126 125 124 123 122 121 120 119 118 117 116 115 114 113 112 111	Je 1

Chart 45.-Wind stress curl for September.

NCC - TDF-11 - DNE DEGREE SUMMARIZATION



Chart 46.-Wind stress curl for October.

NCC - TDF-11 - DNE DEGREE SUMMARIZATION

NL	37 136 135 134 133 132 131 130 129 128 127 126 125 124 123 122 121 120	119 118 117 116 115 114 113 112 111
	Eng	
50	0.00-0.12-0.03 0.17-0.23 0.18 0.36-0.03-0.81 2.41 VANCOUVER ISLAND	50
49	0.03 0.06 0.10-0.01 0.10 0.03 0.30-010 51 0.43	NDAA NATIONAL MARINE FISHERIES SERVICE
48	-0.11 0.28 0.22 0.24 0.23 0.11 0.05-0.10 0.58 0.73	PACIFIC ENVIRONMENTAL GROUP 48 MONTEREY, CALIFORNIA
47	0.21 0.19 0.17-0.21 0.24 0.65 0.17 0.25 G.00 0.01	WIND STRESS
46	0.22 0.03-0.32-0.04 0.17 0.15 0.45-0.05-0.02 0.19 COLUMBIA RIVER	CURL
45	0.34 0.30 0.24 0.10 0.16 0.01 0.10 0627 6.12 0.49	(DYNE CH-2/100KM) 45
44	0.30 0.10 0.03-0.03-0.13 0.21-0.20-0.12 0.23 0.33	LONG TERM MEAN
43	0.12-0.12 0.11 0.18-0.18-0.15 0.08 0.13 0.06 0.01 CAPE BLANCO	FDR 43
42	0.37-0.02-0.04-0.07 0.35-0.16-0.23-0.04 0.21 0.10	NUVEMBER 42
41	9-27 0.12-0.19-0.25 0.01 0.10-0.22-0.11 0.08-0.16	41
40	-0-52-8.16-0.23-0.05 0.02 0.08-0.33-0.15 0.68 0.63	40
39	-0.09-0.11 0.03-0.11 0.13 0.16-0.40-0.00 0.30 0.23	39
38	-8.04-0.13-0.10-0.05 0.08 0.03 0.04 0.20 0.18	30
37	0.15 0.02-0.06-0.08-0.10 0.01-0.02 0.22 0.29	3:
36	0.00-0.18-0.07-0.03 0.00-0.02-0.03-0.20 0.01 0.20	30
35	0-01-0.03-0.09-0.07 0.02 0.00-0.10 0.04 0.21 0.42 pr	DINT CONCEPTION 35
34	0.14-0.01-0.12-0.14-0.09-0.05 0.05 0.37 0.3	3
33	-0.37-0.12-0.07-0.08-0.04-0.08 0.03 0.3	3 0.33 0.17
32	-0.11-0.06-0.13-0.09 0.08 0.05 0.0	3
31	-9.91-9.02-0.94-9.05 0.03-0.25-0.0	17-8) 82 (0. 99 (0. 18) 3
30	-9.05-9.05 0.01-9.49-1	7-0.22 0/00 0-00 0.03
29	-0.03 9.00 0.32 p/or - 200	10 0. 15-0. 05-0. 06-0. D1 0. 07 2 02 2
28	0.18 0.02 0.03 0.19 0.0	0.23 0.28 0.44 0.07 0.04 PUNTA EUGENIA
27	0.19 0.08-0.07-0.2	13-0.21-0.12 0.00-0.03-0.03 0.06 h
26	-0.37 0 01 0.2	38) Ø. 06-9. 11-9. 08-9. 92-9. 05 0. 00-9. 01
25	-0.28-0.13-0.2	36-0.06-0.02 0.03 0.03-0.12-0.08 0.06
24	-0.2	28-0.15-0.03-0.08-0.03-0.02 0.00 0.01-0.05-0.072
23	-0.	35-0.13 001-0.15-0.01-0.04 0.06 0.00-0.09-0.042
22	0.2	15-0.05-0.08 0.07-0.04 0.04-0.09 0.00-0.03-0.132
21	P. 0	0. 20-20. 06 0. 12-0. 01 0. 02-0. 08-0. 04 0. 01-0. 192
	13/ 130 135 134 133 132 131 130 129 128 127 126 125 124 123 122 121 120	

Chart 47.-Wind stress curl for November.

NCC - TDF-11 - ONE DEGREE SUMMARIZATION



Chart 48.-Wind stress curl for December.