



NOAA Technical Memorandum NMFS

AUGUST 2018

UPDATED ESTIMATES OF THE PROBABILITY OF STRIKING A WESTERN NORTH PACIFIC GRAY WHALE DURING THE PROPOSED MAKAH HUNT

Jeffrey E. Moore and David W. Weller

Marine Mammal and Turtle Division, Southwest Fisheries Science Center
National Marine Fisheries Service, National Oceanic and Atmospheric Administration
8901 La Jolla Shores Drive, La Jolla, CA 92037 USA

NOAA-TM-NMFS-SWFSC-605

U.S. DEPARTMENT OF COMMERCE
National Oceanic and Atmospheric Administration
National Marine Fisheries Service
Southwest Fisheries Science Center

About the NOAA Technical Memorandum series

The National Oceanic and Atmospheric Administration (NOAA), organized in 1970, has evolved into an agency which establishes national policies and manages and conserves our oceanic, coastal, and atmospheric resources. An organizational element within NOAA, the Office of Fisheries is responsible for fisheries policy and the direction of the National Marine Fisheries Service (NMFS).

In addition to its formal publications, the NMFS uses the NOAA Technical Memorandum series to issue informal scientific and technical publications when complete formal review and editorial processing are not appropriate or feasible. Documents within this series, however, reflect sound professional work and may be referenced in the formal scientific and technical literature.

SWFSC Technical Memorandums are available online at the following websites:

SWFSC: <https://swfsc.noaa.gov>

NOAA Repository: <https://repository.library.noaa.gov/>

NTIS National Technical Reports Library: <https://ntrl.ntis.gov/NTRL/>

Accessibility information

NOAA Fisheries Southwest Fisheries Science Center (SWFSC) is committed to making our publications and supporting electronic documents accessible to individuals of all abilities. The complexity of some of SWFSC's publications, information, data, and products may make access difficult for some. If you encounter material in this document that you cannot access or use, please contact us so that we may assist you.
Phone: 858-546-7000

Recommended citation

Moore, Jeffrey E., and David W. Weller. 2018. Updated estimates of the probability of striking a western North Pacific gray whale during the proposed Makah hunt. U.S. Department of Commerce, NOAA Technical Memorandum NMFS-SWFSC-605. 8 p. <https://doi.org/10.25923/je72-t228>

ACKNOWLEDGEMENTS

We extend our gratitude to the NOAA Fisheries West Coast Regional Office and Southwest Fisheries Science Center for their contributions to, and support of, this work.

EXECUTIVE SUMMARY

Observations of gray whales (*Eschrichtius robustus*) from the western North Pacific (WNP) migrating to areas off the coast of North America (Alaska to Mexico) raised concerns that this small population could be encountered during a hunt of eastern North Pacific (ENP) gray whales proposed by the Makah Indian Tribe in northern Washington, USA. In 2013, an analysis was conducted to estimate the probability of striking (i.e. killing or seriously injuring) a WNP whale under the Makah Tribe's hunt proposal (Moore and Weller 2013). NOAA Fisheries is considering a draft proposal that would govern ENP gray whale hunts by the Makah for up to 10 years. Under the draft proposal, hunting seasons would alternate between winter-spring hunts in even-numbered years and summer hunts during odd-numbered years. It is presumed that only in even-numbered years (thus, for 5 of the 10 years) would WNP whales potentially be encountered during the hunt. In each of these years, the draft proposal would allow for up to 3 gray whales to be struck. Based on this alternative hunting scheme and the availability of updated gray whale data, this report re-estimates the probability of striking a WNP whale reported earlier (Moore and Weller 2013). One of the models from the 2013 analysis (Model 2A) was used to generate new estimates. We estimate that for an individual strike on a gray whale, the expected probability of it being a WNP whale is 0.004 (95% CRI: 0.002 – 0.007). For a single year's hunt (3 strikes), the expected probability of striking ≥ 1 WNP whale would be 0.012 (0.006 – 0.019). Across the 10-year hunt period (15 strikes), the probability of striking ≥ 1 WNP whale would be 0.058 (0.030 – 0.093).

INTRODUCTION

Two gray whale (*Eschrichtius robustus*) populations are recognized in the North Pacific Ocean. Significant mitochondrial and nuclear genetic differences have been found between whales in the western North Pacific (WNP) and those in the eastern North Pacific (ENP) (LeDuc *et al.*, 2002, Lang *et al.* 2010, Lang *et al.*, 2011). The ENP population ranges from wintering areas in Baja California, Mexico, to feeding areas in the Bering, Beaufort, and Chukchi Seas (Fig. 1). An exception to this generality is the relatively small number (100s) of whales that summer and feed along the Pacific coast between Kodiak Island, Alaska, and northern California (Weller *et al.*, 2013). These whales are collectively called the Pacific Coast Feeding Group (PCFG). The International Whaling Commission (IWC) has defined PCFG whales as individuals observed between 1 June and 30 November from 41°N to 52°N in two or more years (IWC, 2012), and NOAA Fisheries has adopted this definition in recent assessments (Weller *et al.*, 2013). The usual and accustomed (U&A) fishing grounds of the Makah Indian Tribe are off the coast of northern Washington, USA, and overlap with a portion of the PCFG summering area (Fig. 1).




Figure 1. Areas in the western and eastern North Pacific mentioned in the report.

The WNP population feeds in the Okhotsk Sea off Sakhalin Island, Russia (Weller *et al.*, 1999; Weller *et al.* 2012), and in nearshore waters of the southwestern Bering Sea off the southeastern Kamchatka Peninsula (Tyurneva *et al.*, 2010). The historical distribution of gray whales in the Okhotsk Sea greatly exceeded what is found today (Reeves *et al.*, 2008). Whales associated with the Sakhalin feeding area can be absent for all or part of a given feeding season (Bradford *et al.*, 2008), indicating they use other areas during the summer and fall feeding period. Some of the whales identified feeding in the coastal waters off Sakhalin, including reproductive females and calves, have been documented off the southern and eastern coast of Kamchatka (Tyurneva *et al.*, 2010). A small number of whales observed off Sakhalin have also been sighted off the northern Kuril Islands in the eastern Okhotsk Sea and Bering Island in the western Bering Sea (Weller *et al.*, 2003).

Mixing of whales identified in the WNP and ENP has been observed (Weller *et al.*, 2012). Lang (2010) reported that two adult individuals from the WNP, sampled off Sakhalin in 1998 and 2004, matched the microsatellite genotypes, mtDNA haplotypes, and sexes (one male, one female) of two whales sampled off Santa Barbara, California in March 1995. Between 2010 and 2012 three whales outfitted with satellite transmitters were tracked moving from Sakhalin in the WNP to the ENP (Mate *et al.*, 2015). Finally, photographic matches between the WNP and ENP, including matches between Sakhalin, Vancouver Island and Laguna San Ignacio (Fig. 1), have further confirmed use of areas in the ENP by whales identified in the WNP (Weller *et al.*, 2012, Urbán *et al.*, 2012). Despite this level of mixing, significant mtDNA and nuclear genetic differences between whales in the WNP and ENP have been found (LeDuc *et al.* 2002, Lang *et al.*, 2011).

In 1995, following the 1994 delisting of ENP gray whales under the U.S. Endangered Species Act, the Makah Indian Tribe notified NOAA Fisheries of its interest in re-establishing limited ceremonial and subsistence whale hunting. The decision-making history on this issue is complex and not described here except to note that in 2005, the Makah Tribe submitted a detailed proposal for hunting ENP gray whales in the coastal portion of its U&A off northern Washington, USA, as part of a request for a waiver of the U.S. Marine Mammal Protection Act's (MMPA) take moratorium (16 USC 1371(a)(3)(A)). Subsequently, observations of WNP gray whales migrating through areas off the coast of North America (Alaska to Mexico) emphasized the need to evaluate the probability of a WNP gray whale being encountered in aboriginal hunts for ENP gray whales (IWC, 2012). Following recommendations of the Scientific Committee of the International Whaling Commission (IWC), analyses were conducted to estimate such probability in the context of the Makah Tribe's hunt proposal (Moore and Weller, 2013). These analyses informed a draft Environmental Impact Statement (DEIS), completed in 2015 (NMFS, 2015), pertaining to the Makah Tribe's MMPA waiver request.

NOAA Fisheries is presently considering a MMPA waiver and associated draft proposal that would govern a modified version of the Tribe's hunt proposal. The objective of the analysis reported here was to provide updated estimates of the probability that one or more WNP whales might be subjected to strikes¹, unsuccessful strike attempts (i.e., harpoon throws that do not penetrate), and vessel approaches during hunts and hunt training exercises considered in the draft proposal. This report is based on the methods used by Moore and Weller (2013) and incorporates updated information about the population sizes of ENP and WNP gray whales and their occurrence within the proposed hunt area.

METHODS

Hunt proposal

NOAA Fisheries' draft proposal would govern a Makah Tribe hunt of ENP gray whales in the coastal portion of the U&A (i.e., the "hunt area") over a 10-year hunt period. In odd-numbered years, the hunt would take place from 1 July through 31 October, a period when no sightings of WNP whales have been recorded in the ENP, and when gray whales generally (apart from PCFG

¹ As described in NOAA Fisheries' DEIS (NMFS, 2015), the term "strike" is interpreted to be consistent with the IWC Schedule definition as meaning "to penetrate with a weapon used for whaling."

animals) are in northern feeding areas. Thus, hunted animals in these odd-numbered years would presumably belong to the PCFG and it is assumed that WNP whales would not be at risk from proposed hunt operations. In even-numbered years, the hunt would take place from 1 December through 31 May. This period coincides with both the southward (December to mid-February) and northward (mid-February to late May) migration of ENP whales and overlaps with the time when WNP gray whales have been sighted in the ENP. Thus, in even-numbered years there is a potential risk to WNP whales from proposed hunt operations. In each of the even-numbered years, a maximum of 3 gray whales per year could be struck (including “struck and lost” animals). Over the 10-year period of the proposed hunt, a maximum of 15 whales could be struck (in even-numbered years) that would have some probability of being WNP whales. We therefore evaluate the probability of striking at least one WNP whale per even-numbered year (out of 3 strikes) and for the 10-year period (out of 15 strikes). We also evaluate associated rates of WNP whales being subjected to aforementioned “unsuccessful strike attempts” (i.e., harpoon throws that do not penetrate) and “approaches” (i.e., whales approached by vessels during hunts and hunt training exercises).

Data

Abundance estimates - The most recent ENP abundance estimate (for 2015/2016) is 26,960 (CV = 0.05) (Durban *et al.*, 2017). The most recent WNP abundance estimate (for 2015) is 200 (CV = 0.03) for the 1+ population (i.e., excluding calves) (Cooke 2018). We then multiplied the WNP estimate by 1.099 to account for calves. This multiplier is based on the ratio of the population size with and without calves in 2012 (IUCN, 2012).

Mixing proportions based on sightings in the Makah Hunt Area - During spring surveys (March to May) in 1996-2012 there were 181 observed whale-days in the Makah hunt area (Calambokidis *et al.*, 2014). To clarify the term “whale-day” – all sightings of an individual on a particular day collectively count as 1 whale-day (e.g., multiple sightings of the same individual on the same day count as just 1 whale-day, but the same individual seen the next day would count as a second whale-day). None of the 181 whale-days observed included WNP whales²; 73 (40.3%) were considered PCFG whales; and the rest (108, or 59.7%) were assumed to be migrating ENP whales.

However, rather than use 40.3% as the expected PCFG proportion in the hunt area during an even-year hunt, we use 28% for this mixing proportion (i.e. 72% of animals encountered during an even-year hunt are likely to be non-PCFG animals). This value is based on analyses summarized in a 2018 IWC workshop (IWC, 2018).

Proportion of WNP whales migrating with ENP whales - The proportion of the WNP population that migrates along the North American coast is unknown but estimated to be at least 0.37 based on analysis by Cooke (2015) and reported to a 2015 IWC workshop on gray whale population structure (IWC, 2016).

² Although not in the Makah hunt area, Weller *et al.* (2012) report observing three WNP whales on 2 May 2004 and three more on 25 April 2008 near Barkley Sound off the west coast of southern Vancouver Island, British Columbia, Canada.

Model

Moore and Weller (2013) considered four models in their analysis but they based final inferences on what they termed Model 2B. Here, we use Model 2A instead. Models 2A and 2B are similar. The difference is that for Model 2A, the conditional probability of a non-PCFG whale being a WNP (rather than ENP) whale is simply based on the ratio of WNP:ENP population size. This is an intuitive estimator, though it does rely on the assumption that WNP and ENP animals migrating together are using the same migration corridors and behaving similarly. For Model 2B, this assumption is relaxed and we allow for broader uncertainty by stating that the conditional probability varies uniformly from zero (if the WNP whales do not migrate through the Makah area at all) to some maximum value that is based on (but not equivalent to) the ratio of WNP:ENP population size. However, it is difficult to define that maximum value, and allowing a lower probability of zero is not precautionary and arguably should not be considered without supporting evidence.

Model 2 (A and B) makes use of the mixing proportion/sightings data for the Makah hunt area, as well as WNP and ENP abundance estimates. WNP whales are assumed to be moving with the ENP migrants, so that the marginal probability of a WNP whale being struck is the probability that the struck whale is a migrant, P_{mig} (i.e., probability of not being a PCFG whale), multiplied by the conditional probability of being a WNP whale given that it is a migrant ($P_{\text{WNP|mig}}$). Thus, $P_{\text{WNP}} = P_{\text{mig}}P_{\text{WNP|mig}}$.

P_{mig} is defined as $1 - P_{\text{PCFG}}$, where P_{PCFG} is given by an informative prior: $P_{\text{PCFG}} \sim \text{Beta}(5.3648, 13.7952)$ which has a mean of 0.28 and SD of 0.1 (IWC 2018).

We assume that the per-capita likelihood of a migrating (non-PCFG) whale in the hunt area being a WNP whale (i.e., $P_{\text{WNP|mig}}$) is simply given by the proportion of the migrating population made up of WNP whales. This proportion depends on what fraction of the WNP population migrates along the U.S. West Coast, which we call m , and the relative size of the WNP to the ENP population. Thus, $P_{\text{WNP|mig}} = mN_{\text{WNP}} / (mN_{\text{WNP}} + N_{\text{ENP}})$. Let $m \sim \text{Uniform}(0.37, 1)$, based on Cooke *et al.* (2015). N_{WNP} and N_{ENP} are treated as lognormally distributed variables with means and CVs as given above.

Estimation

Earlier analyses (Moore and Weller, 2013) used Bayesian estimation. In the current exercise, analysis was conducted using OpenBUGS software, but estimation is not strictly Bayesian because there are no new data updating the informative prior inputs. Rather, the present analysis is essentially a Monte Carlo procedure, with distributions for the parameters of interest (e.g., probability of striking a WNP whale) being derived from random draws from informed prior distributions for the input parameters. Derived parameter distributions are summarized from two MCMC chains, each 25,000 samples in length (50,000 samples total).

Derived parameters

The key parameter of interest is the per-strike probability of striking a WNP whale. Derived from this parameter are the probabilities of striking at least one WNP out of 3 gray whale strikes (i.e., the annual probability of striking a WNP whale, for the even-numbered years) or out of 15 gray

whale strikes (i.e., probability for the whole 10-year period). These are calculated as $P(x > 0) = 1 - (1 - P_{WNP})^X$, where X is 3 or 15. Additionally, we can derive the expected number of WNP strikes as $E(x) = P_{WNP}X$. Using data collected during previous hunts (NMFS, 2015), the following two assumptions were used to calculate analogous estimates for vessel approaches and unsuccessful strike attempts: (1) there will be 353 vessel approaches per year (3530 across all 10 years)³, and (2) there will be 6 unsuccessful strike attempts for every strike in an even-year hunt⁴.

RESULTS

Parameter estimates

Estimated parameters from all model sets are in Table 1. Figure 2 shows the distribution for P_{WNP} . It is straightforward to integrate across the uncertainty in P_{WNP} to obtain a single probability estimate. We did this for the probability of striking ≥ 1 WNP whale over the entire 10-year hunt period (i.e., out of 15 strikes). This probability was 0.058.

Table 1. Distribution summaries for key model parameters. “Prob(WNP)” is the probability of at least 1 WNP animal being struck or subjected to unsuccessful strike attempts or vessel approaches given the specified number of events.

Parameter	Posterior mean	2.5% CRI	Posterior median	97.5% CRI
Prob(WNP) for a single interaction, i.e., P_{WNP}	0.004	0.002	0.004	0.007
Prob(WNP 3 strikes in 1 yr)	0.012	0.006	0.012	0.019
Prob(WNP 15 strikes in 10 yrs)	0.058	0.030	0.057	0.093
Prob(WNP 18 unsuccessful strike attempts in 1 yr)	0.070	0.036	0.069	0.110
Prob(WNP 90 unsuccessful strike attempts in 10 yrs)	0.299	0.167	0.298	0.442
Prob(WNP 353 approaches in 1 yr)	0.735	0.511	0.751	0.899
Prob(WNP 3530 approaches in 10 yrs)	~ 1.0	0.999	~ 1.0	~ 1.0
Expected WNP 3 strikes in 1 yr	0.012	0.006	0.012	0.019
Expected WNP 15 strikes in 10 yrs	0.060	0.030	0.059	0.097
Expected WNP 18 unsuccessful strike attempts in 1 yr	0.072	0.036	0.071	0.116
Expected WNP 90 unsuccessful strike attempts in 10 yrs	0.361	0.182	0.353	0.582
Expected WNP 353 approaches in 1 yr	1.416	0.714	1.386	2.283
Expected WNP 3530 approaches in 10 yrs	14.160	7.141	13.860	22.830

³ This number is conservative because it assumes that all approaches (hunting and training) in both even and odd years occur during the winter/spring period when WNP whales may be present. Realistically we would expect a substantial number of approaches to occur outside this period, i.e., during the summer when ocean conditions are more favorable and, in odd years, when hunting approaches are restricted to July - October.

⁴ We expect zero in odd years because the draft proposal limits training strikes (which count as unsuccessful strike attempts) to the summer-fall hunting season, when WNP whales are not expected to be present.




Figure 2. Posterior distribution for probability that any given strike is a WNP whale.

DISCUSSION

Estimates from our analysis may be precautionary since they assume that the Makah hunt will achieve proposed maximum strike limits, and because the assumption of Model 2A is that WNP whales are homogeneously mixed with ENP whales. The likelihood of striking a WNP whale is overestimated if fewer total animals are struck or if in reality the WNP animals use a different migration corridor and are less likely to travel through the Makah hunt area. Given uncertainties associated with the model and scenario assumptions, these results serve as a rough approximation of the potential for WNP gray whales to be subjected to strikes, unsuccessful strike attempts and vessel approaches during a Makah hunt operating under a draft proposal currently being considered by NOAA Fisheries.

REFERENCES

- Bradford, A.L., Weller, D.W., Wade, P.R., Burdin, A.M. and Brownell, R.L., Jr. 2008. Population abundance and growth rate of western gray whales *Eschrichtius robustus*. *Endangered Species Research* 6(1):1-14.
- Calambokidis, J., Laake, J., and A. Perez. 2014. Updated analysis of abundance and population structure of seasonal gray whales in the Pacific Northwest, 1996-2012. Final Report to National Marine Mammal Laboratory, Seattle, WA. [Available at: <http://www.afsc.noaa.gov/nmml/>]
- Cooke, J.G. 2015. Implications of observed whale movements on the relationship between the Sakhalin gray whale feeding aggregation and putative breeding stocks of the gray whale. Paper SC/A15/GW02 presented to the Second Workshop on the Rangewide Review of the

- Population Structure and Status of North Pacific Gray Whales, 1-3 April 2015, La Jolla, CA, USA. [Available at: <http://www.iwcoffice.org/>]
- Cooke, J.G., Weller, D.W., Bradford, A.L., Sychenko, O., Burdin, A.M., Lang, A.R. and Brownell, R.L. 2015. Updated population assessment of the Sakhalin gray whale aggregations based on the Russia-US photo-identification study at Piltun, Sakhalin, 1994-2014. Paper WGWP 16/17 presented at the 16th meeting of the Western Gray Whale Advisory Panel. [Available at: http://cmsdata.iucn.org/downloads/wgwap_16_17_cooke_et_al.pdf]
- Cooke, J.G. 2018. Abundance estimates for western North Pacific gray whales for use with stock structure hypotheses of the Range-wide Review of the Population Structure and Status of North Pacific gray whales. Paper SC/67B/ASI/02 presented to the International Whaling Commission Scientific Committee. [Available at: <http://www.iwcoffice.org/>]
- Durban, J.W., Weller, D.W. and Perryman, W.L. 2017. Gray whale abundance estimates from shore-based counts off California in 2014/15 and 2015/16. Paper SC/A17/GW/06 presented to the International Whaling Commission Scientific Committee. [Available at: <http://www.iwcoffice.org/>]
- IUCN 2012. Report of the 11th Meeting of the Western Gray Whale Advisory Panel. Geneva, Switzerland. [Available at: http://www.iucn.org/wgwap/publications_and_reports/]
- IWC 2012. Report of the Scientific Committee. Annex E. Report of the standing working group on an aboriginal subsistence whaling management procedure. Journal of Cetacean Research and Management (Supplement) 13:130-153. [Available at: <http://www.iwcoffice.org/>]
- IWC 2016. Report of the Scientific Committee. Report of the 2nd workshop on the rangewide review of the population structure and status of North Pacific gray whales, 1-3 April 2015, La Jolla, CA, USA. Journal of Cetacean Research and Management (Supplement) 17:567-581. [Available at: <http://www.iwcoffice.org/>]
- IWC. 2018. Report of the Fifth Rangewide Workshop on the Status of North Pacific Gray Whales. Paper SC/67B/REP/07 Rev 1 presented to the International Whaling Commission Scientific Committee. [Available at: <http://www.iwcoffice.org/>]
- Lang, A.R. 2010. The population genetics of gray whales (*Eschrichtius robustus*) in the North Pacific. Ph.D. dissertation, University of California, San Diego, CA 222 pp.
- Lang, A.R., Weller, D.W., LeDuc, R., Burdin, A.M., Pease, V.L., Litovka, D., Burkanov, V. and Brownell, R.L. 2011. Genetic analysis of stock structure and movements of gray whales in the eastern and western North Pacific. Paper SC/63/BRG10 presented to the International Whaling Commission Scientific Committee. [Available at: <http://www.iwcoffice.org/>]
- LeDuc, R.G., Weller, D.W., Hyde, J., Burdin, A.M., Rosel, P.E., Brownell, R.L., Würsig, B. and Dizon, A.E., 2002. Genetic differences between western and eastern gray whales (*Eschrichtius robustus*). Journal of Cetacean Research and Management 4(1):1-5.
- Mate, B.R., Ilyashenko, V.Y., Bradford, A.L., Vertyankin, V.V., Tsidulko, G.A., Rozhnov, V.V. and Irvine, L.M. 2015. Critically endangered western gray whales migrate to the eastern North Pacific. Biology Letters 11:20150071.
- Moore, J.E. and Weller, D.W. 2013. Probability of taking a western North Pacific gray whale during the proposed Makah hunt. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-SWFSC-506. [Available at: <https://swfsc.noaa.gov/publications/TM/SWFSC/NOAA-TM-NMFS-SWFSC-506.pdf>]

- NMFS 2015. Draft Environmental Impact Statement on the Makah Tribe Request to Hunt Gray Whales. February 2015. 1,230 pp. [Available at: http://www.westcoast.fisheries.noaa.gov/protected_species/marine_mammals/cetaceans/makah_tribal_whale_hunt.html]
- Reeves, R.R., Smith, T.D. and Josephson, E.A. 2008. Observations of western gray whales by ship-based whalers in the 19th century. *Journal of Cetacean Research and Management* 10(3):247-256.
- Tyurneva, O. Yu, Yakovlev, Yu M., Vertyankin, V.V. and Selin, N.I. 2010. The peculiarities of foraging migrations of the Korean-Okhotsk gray whale (*Eschrichtius robustus*) population in Russian waters of the Far Eastern seas. *Russian Journal of Marine Biology* 36(2):117-124.
- Urbán R., J., Weller, D.W., Tyurneva, O., Swartz, S., Bradford, A., Yakovlev, Y., Sychenko, O., Rosales N., H., Martínez A., S., Burdin, A. and Gómez-Gallardo U., A. 2012. Report on the photographic comparison of the western and Mexican gray whale catalogues. Paper SC/64/BRG13 presented to the International Whaling Commission Scientific Committee. [Available at: <http://www.iwcoffice.org/>]
- Weller, D.W., Burdin, A.M., Ivashchenko, Y.V., Tsidulko, G.A. and Brownell, R.L. 2003. Summer sightings of western gray whales in the Okhotsk and western Bering Seas. Paper SC/55/BRG9 submitted to the International Whaling Commission Scientific Committee. [Available at: <http://www.iwcoffice.org/>]
- Weller, D.W., Würsig, B., Bradford, A.L., Burdin, A.M., Blokhin, S.A., Minakuchi, H. and Brownell, R.L. 1999. Gray whales (*Eschrichtius robustus*) off Sakhalin Island, Russia: seasonal and annual patterns of occurrence. *Marine Mammal Science* 15:1208-1227.
- Weller, D.W., Klimek, A., Bradford, A.L., Calambokidis, J., Lang, A.R., Gisborne, B., Burdin, A.M., Szaniszló, W., Urbán, J., Gomez-Gallardo Unzueta, A., Swartz, S. and Brownell, R.L. 2012. Movements of gray whales between the western and eastern North Pacific. *Endangered Species Research* 18:193-199
- Weller, D.W., Bettridge, S., Brownell, R.L., Laake, J.L., Moore, J.E., Rosel, P.E., Taylor, B.L and Wade, P.R. 2013. Report of the National Marine Fisheries Service gray whale stock identification workshop. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-SWFSC-507. [Available at: <https://swfsc.noaa.gov/publications/TM/SWFSC/NOAA-TM-NMFS-SWFSC-507.pdf>]