

Chapter 7

Assessment of the Kamchatka Flounder stock in the Bering Sea and Aleutian Islands

By

Thomas K. Wilderbuer, James Ianelli,
Daniel G. Nichol and Robert Lauth

Alaska Fisheries Science Center
NMFS/NOAA 7600 Sand Point Way NE
Seattle WA 98115

Executive Summary

This assessment is a full update of the 2014 stock assessment. The 2015 assessment was an off-cycle assessment that did not re-run an updated assessment model; instead the projection model was run with updated catch information only to provide estimates of 2016 and 2017 ABC and OFL without re-estimating the stock assessment model parameters and biological reference points. The stock is estimated to be 13% above $B_{40\%}$ and has been harvested at about half the ABC level the past 3 years.

Summary of changes in assessment input

- 1) Estimate of catch for 2015 and 2016. The estimated 2016 catch of 4,530 t was used as the catch value for the 2017 and 2018 ABC and OFL projections.
- 2) 2016 slope survey biomass and standard error estimates.
- 3) 2015 and 2016 shelf survey length composition
- 4) 2016 and 2016 shelf survey biomass and standard error estimates.
- 5) 2015 Aleutian Islands survey biomass and standard error.
- 6) 2016 Aleutian Islands survey length composition.
- 7) 2016 slope survey length composition.

No changes were made to the assessment methodology.

Summary of Results

| | Tier 3 assessment model | | | |
|--------------------------------------|-------------------------------------|---------|-------------------------------------|---------|
| Quantity | As estimated last year for | | As estimated this year for | |
| | 2016 | 2017 | 2017 | 2018 |
| M (natural mortality rate) | 0.11 | 0.11 | 0.11 | 0.11 |
| Tier | 3 | 3 | 3 | 3 |
| Projected total (age 2+) biomass (t) | 182,300 | 189,100 | 170,300 | 181,000 |
| Projected female spawning biomass | | | | |
| Projected | 61,700 | 63,800 | 60,300 | 62,200 |
| $B_{100\%}$ | 132,500 | 132,500 | 127,000 | 127,000 |
| $B_{40\%}$ | 53,000 | 53,000 | 50,800 | 50,800 |
| $B_{35\%}$ | 46,400 | 46,400 | 44,400 | 44,400 |
| F_{OFL} | 0.076 | 0.076 | 0.078 | 0.078 |
| $maxF_{ABC}$ | 0.065 | 0.065 | 0.066 | 0.066 |
| F_{ABC} | 0.065 | 0.065 | 0.066 | 0.066 |
| OFL (t) | 11,100 | 11,700 | 10,360 | 10,700 |
| maxABC (t) | 9,500 | 10,000 | 8,880 | 9,200 |
| ABC (t) | 9,500 | 10,000 | 8,880 | 9,200 |
| Status | As determined <i>last</i> year for: | | As determined <i>this</i> year for: | |
| | 2014 | 2015 | 2015 | 2016 |
| Overfishing | no | n/a | no | n/a |
| Overfished | n/a | no | n/a | no |
| Approaching overfished | n/a | no | n/a | no |

Responses to SSC and Plan Team Comments on Assessments in General

1. Assessment authors should routinely do retrospective analyses extending back 10 years, plot spawning biomass estimates and error bars, plot relative differences, and report Mohn's rho (revised).
2. If a model exhibits a retrospective pattern, try to investigate possible causes.
3. Communicate the uncertainty implied by retrospective variability in biomass estimates.
4. For the time being, do not disqualify a model on the grounds of poor retrospective performance alone.

This comment was from 2014 and was addressed for the first time in this assessment (off-cycle year for full assessment was in 2015).

Responses to SSC and Plan Team Comments Specific to this Assessment

No comments were provided specific to this assessment from 2015.

Introduction

In 2013 a Tier 3 approach was used to describe the stock status of Kamchatka flounder using survey and fishery age and length structured modeling. The assessment previously used Tier 5 methodology reliant upon trawl survey biomass from the Bering Sea shelf, slope and the Aleutian Islands and an estimate of natural mortality. ABC and OFL were determined from a 7-year averaging technique of survey biomass.

The Kamchatka flounder (*Atheresthes evermanni*) is a relatively large flatfish which is distributed from Northern Japan through the Sea of Okhotsk to the Western Bering Sea north to Anadyr Gulf (Wilimovsky et al. 1967) and east to the eastern Bering Sea shelf and south of the Alaska Peninsula (there is also a catch record from California). In U.S. waters they are found in commercial concentrations in the Aleutian Islands where they generally decrease in abundance from west to east (Zimmerman and Goddard 1996). They are also present in Bering Sea slope waters but are absent in survey catches east of Chirikof Island.

In the eastern part of their range, Kamchatka flounder overlap with arrowtooth flounder (*Atheresthes stomias*), a species that is very similar in appearance and was not routinely distinguished in the commercial catches until 2007. Until about 1991, these species were also not consistently separated in trawl survey catches (Fig. 7-1) and were combined in the arrowtooth flounder stock assessment (Wilderbuer et al. 2009). However, managing the two species as a complex became undesirable in 2010 due to the emergence of a directed fishery for Kamchatka flounder in the BSAI management area. Since the ABC was determined by the large amount of arrowtooth flounder relative to Kamchatka flounder (the complex was about 93% arrowtooth flounder), the possibility arose of an overharvest of Kamchatka flounder as the *Atheresthes* sp. ABC exceeded the Kamchatka flounder biomass. Arrowtooth and Kamchatka flounder have been managed separately since 2011. There is no evidence of stock structure.

Fishery

Catch History

Historical Kamchatka flounder catch is combined in catch records for arrowtooth flounder and Greenland turbot from the 1960s. The fisheries for Greenland turbot intensified during the 1970s and the bycatch of arrowtooth flounder and Kamchatka flounder is assumed to have also increased. Catches of these species decreased after implementation of the MFCMA and the Kamchatka flounder resource remained lightly exploited with the combined catches with arrowtooth flounder averaging 12,831 t from 1977-2008 (Table 7-1). It is estimated that only a small fraction (<10%) of this catch was Kamchatka flounder. This decline resulted from catch restrictions placed on the fishery for Greenland turbot and phasing out of the foreign fishery in the U.S. EEZ. Catches in Table 7-1 through 2006 are for arrowtooth flounder and Kamchatka flounder combined, catches thereafter are those for Kamchatka flounder only. The total combined catch for arrowtooth and Kamchatka flounder reported by the Alaska Regional Office (catches were not differentiated by species until 2011) is a blend of vessel reported catch and observer at-sea sampling of the catch. However, observers have separately identified the two species from catches aboard trawl vessels since 2007 and their sampling has indicated that the proportion of Kamchatka flounder in the combined catch has steadily increased from 10% in 2007 to 55% in 2010.

| year | Percent of combined catch |
|------|---------------------------|
| 2007 | 10% |
| 2008 | 31% |
| 2009 | 45% |
| 2010 | 55% |

The increased harvest was the result of a recently developed foreign market for Kamchatka flounder, which has now become a fishery target. Based on the above observer-derived percentages, the 2010 estimated catch of Kamchatka flounder was 21,153 t, taken primarily in area 514 and to a lesser extent in area 518. The 2011 catch of 9,935 is less than half of the 2010 combined total (TAC and ABC = 17,700 t, OFL = 23,600 t) and was evenly split between area 541 in the central Aleutian Islands (51%) and area 524 in the northern Bering Sea (34%). Based on this result in 2011, area apportionment has not been pursued in the assessment. The 2012 catch of 9,514 t was about the same as the 2011 and has declined since then to 4,994 t in 2015 and 4,533 t in 2016 (through October 15). The 2016 catch is 48% of the ABC of 9,500 t (Table 7-1). The Kamchatka catch by week in 2016 (Fig. 7-1, mostly trawl catch) indicates that targeting for Kamchatka flounder began May 1 and continued throughout the summer. Peak weekly amounts occurred in May and to a lesser extent in August. Catches were lower the rest of the year.

Data

The data used in this assessment includes estimates of total fishery catch, bottom trawl survey biomass estimates and length composition from the Bering Sea shelf, slope and Aleutian Islands surveys. Age data are available from the 2010 Aleutian Islands survey and from the 2002 and 2012 slope surveys. All survey length-weight observations were included.

Fishery catch

Fishery catch from 2007-2016 were included in the model as listed above. Catches from 1991-2006, years when Kamchatka and arrowtooth flounder were not identified to species were calculated by assuming that Kamchatka flounder comprised 10% of the catch during that time period. The 2016 catch used the current estimate from the Alaska Regional Office at the time of the assessment (October 15), assuming the fishery is nearly finished for 2016, while also acknowledging that a small amount of catch will still be attained.

Absolute Abundance from Trawl Surveys

Biomass estimates (t) for Kamchatka flounder from the standard shelf survey area in the eastern Bering Sea, slope surveys and the Aleutian Islands region are shown in Table 7-2. Reliable estimates of Kamchatka flounder became available in 1991 and they were estimated at an average biomass of 45,500 t through 1994 on the Bering Sea shelf (Fig. 7-2). During the following 10 years the biomass was estimated at a lower level (25,000 t average) before increasing to high and stable levels the past 10 years (52,700 t average). On the continental shelf they are usually found in highest concentrations at depths greater than 200 meters around the Pribilof Islands and also in the large shelf area west of St. Matthew Island (Fig. 7-3) and were present in 244 of the 376 stations in the standard survey area in 2016 (Table 7.3, Fig. 7-2). Trends of abundance from the slope and Aleutian Islands surveys indicate a stable resource at a higher level than 10 years ago. They are common in the deeper waters of the slope area (500 to 800 meters, Zimmerman and Goddard 1996) in both the Aleutian Islands and the eastern Bering Sea slope (Figs. 7-3 and 7-4).

Population length composition estimates for the three trawl surveys are shown by year and sex in Figure 7-5 and Tables 7.4, 7.6.

Data

The data available for Kamchatka flounder are:

| | |
|---|-----------|
| Fishery catch | 1991-2016 |
| Shelf survey biomass estimates and standard error | 1982-2016 |

| | |
|---|--|
| Slope survey biomass estimates and standard error | 2002, 2004, 2008, 2010, 2012, 2016 |
| Aleutian Islands survey biomass and S.E. | 1991, 1994, 1997, 2000, 2002, 2004, 2006, 2010, 2012, 2014, 2016 |
| Shelf survey length composition | 1991-2016 |
| Slope survey length composition | 2004, 2008, 2010, 2016 |
| Aleutian Islands survey length composition | 1991, 1994, 1997, 2000, 2002, 2004, 2006, 2012, 2014, 2016 |
| Fishery length data | 2008 - 2013 |
| Slope survey age data | 2002, 2012 |
| Aleutian Islands survey age data | 2010 |

Analytic Approach

Model Structure

This stock assessment utilizes the AD Model Builder software to model the population dynamics of Bering Sea and Aleutian Islands Kamchatka flounder starting in 1991. The model is a sex-specific length and age-based approach where survey and fishery length composition observations are used to calculate estimates of population numbers-at-age by the use of a length-age (growth) matrix. The model simulates the dynamics of the population as well as the surveys and fisheries, and compares the expected values of for the survey and fishery quantities to those observed from surveys and fishery sampling programs. This is accomplished by the simultaneous estimation of the parameters in the model using the maximum likelihood estimation procedure. The fit of the simulation values to the observed values is optimized by maximizing the log(likelihood) function given the following distributional assumptions about the observed data (see Tables 7-8 and 7-9).

The suite of parameters estimated by the base model are classified by the following likelihood components:

| Data Component | Distribution assumption |
|--|-------------------------|
| Trawl fishery size composition | Multinomial |
| Shelf survey population size composition | Multinomial |
| Slope survey population size composition | Multinomial |
| Slope survey age composition (2002 and 2012) | Multinomial |
| Aleutian Islands survey size composition | Multinomial |
| Aleutian Islands age composition (2010) | Multinomial |
| Trawl survey biomass estimates and S.E. | Log normal |
| Slope survey biomass estimates and S.E. | Log normal |
| Aleutian Islands biomass estimates and S.E. | Log normal |

The total log likelihood is the sum of the likelihoods for each data component. The model allows for the individual likelihood components to be weighted by an emphasis factor. Equal emphasis was placed on fitting all data components for this assessment with the exception that a large emphasis was placed on fitting the fishery catch. The number of parameters estimated for the base configuration of the model are presented below:

| Fishing mortality | Selectivity | Shelf and Aleutian survey q | Year class strength | Total |
|-------------------|-------------|-------------------------------|---------------------|-------|
| 27 | 16 | 2 | 50 | 95 |

The recruitment parameters are comprised of the 24 initial ages in 1991 (ages 2-25), the 24 subsequent recruitment deviation estimates from 1992-2014 and the mean log of the initial recruitment and the log of all recruitment. Fishing mortality (F) parameters include the log of average F and the 26 annual fishing mortality deviations. Selectivity parameters are from the logistic model for 3 surveys and a single fishery, for each sex. It was assumed that the shelf, slope and Aleutian Islands surveys measure non-overlapping segments of the Kamchatka flounder stock. Biomass was apportioned between the three areas by calculating the average of the annual proportions estimated from the trawl surveys (Fig 7-8). The resulting proportions are (53% shelf, 20% slope and 26% in the Aleutian Islands) were used as starting values in the area-specific catchability estimation. The length-age conversion matrices (sex-specific) were constructed using fitted von Bertalanffy growth curves to the available age data. The variability in length at age was estimated to reflect a CV of about 8% (in cm). This provided the variance in growth for the length-age conversions.

In addition, two more parameters can be estimated in a later stage to estimate the annual relationship between bottom water temperature and shelf survey catchability and bottom water temperature and the overall value of catchability which relates to the capture process and availability of the stock (discussed in the next section).

Parameters Estimated Outside of the Assessment Model

Length-weight, length and weight at age, maturity and natural mortality

All length-weight measurements collected during RACE surveys (1,074 total, 483 males and 591 females) were used to describe the Kamchatka flounder length (cm)-weight (g) relationship (Fig 7.6) by the equation:

$$\text{Males: } W = 4.73 \times 10^{-6} L^{3.757}$$

$$\text{Females } W = 2.08 \times 10^{-3} L^{3.393}$$

Length at age calculations from the ageing of 450 otoliths from the 2010 Aleutian Islands survey were fit to a von Bertalanffy growth model to obtain male and female length at age. These data were then multiplied by the sex-specific length-weight data to obtain estimates of weight-at-age for the assessment model. Weight-at-age data indicate that females and males grow at a similar rate until about the age of maturation, after which females continue to grow to a larger size (Fig 7.7). Maturity was determined in a study by Stark (2011) from a histological examination of ovary samples collected in the Bering Sea (Table 7.7).

Both sexes have been found in relatively equal numbers and the oldest fish have been aged as old as 49 years, indicating that Kamchatka flounder are similar in life history to other Bering Sea flatfish. The assessment model was used to explore estimates of natural mortality.

Parameters Estimated Inside the Assessment Model

Catchability

Examination of Bering Sea shelf survey biomass estimates indicate that some of the annual variability seemed to positively co-vary with bottom water temperature. Variations in shelf survey biomass were particularly evident during the coldest year (1999) and the warm trend that occurred from 2001-2005.

The relationship between average annual bottom water temperature collected during the survey and annual survey biomass estimates can be better understood by modeling survey catchability as:

$$q = e^{-\alpha + \beta T}$$

where q is catchability, α and β are parameters estimated by the model, and T_t is the average annual bottom water temperature for year t . The catchability equation has two parts. The e^α term is a constant or time-independent estimate of q . The second term, $e^{\beta T}$ is a time-varying (annual) q which relates to the metabolic aspect of herding or distribution (availability) which can vary annually with bottom water temperature.

Year class strengths

The population simulation specifies the numbers-at-age in the beginning year of the simulation, the number of recruits in subsequent years as deviations from overall mean log recruitment, and the survival rate for each cohort as it moves through the population calculated from the population dynamics equations (see Table 7-8 and Table 7-9).

Fishing Mortality

The fishing mortality rates for each age and year are calculated to approximate the catch weight by solving for F while still allowing for observation error in catch measurement. A large emphasis (300) was placed on the catch likelihood component to closely match the observed catches.

Selectivity

Survey results indicate that fish less than about 4 years old (< 30 cm) are found mostly on the Bering Sea shelf and to a lesser extent in the Aleutian Islands. Males and females from 30-50 cm are found on the shelf and in deeper waters of the Aleutian Islands and Bering Sea slope waters, and males and females > 50 cm are mainly found at depths below 200 meters. Sex specific "domed-shaped" selectivity was freely estimated for males and females in the shelf survey due to the lack of larger fish there. We assumed an asymptotic selectivity pattern for both sexes in the slope surveys and the Aleutian Islands surveys. Selectivity was assumed constant over all survey years.

Up to the present, the low sampling intensity for length measurements from the fishery may not provide sufficient information for the model to reliably estimate fishery selectivity. The input sample size for fitting this data was set at a low level (25) and may be overemphasized. This results in sample size problems which make estimates of fishery selectivity unreliable. The shape of the selectivity curve was fixed asymptotic for older fish in the fishery since the directed fishery for Kamchatka flounder presumably targets the larger fish.

Results

Model Evaluation

- 1) Started with q 's (catchability) apportioned by their relative survey biomass estimates for the three survey areas.
- 2) Examination of the results from the initial model run indicated that fishery selectivity is poorly determined (presumably due to the low sample sizes,) and that there are males present in the length records that are larger than those observed in any survey data. It is suspected that this is the result of some mis-sexing of Kamchatka flounder in the commercial fishery sampling. This was resolved by fixing the slope of the logistic curve (age at 50% selection is still estimated for each sex) which produced more sensible results (Fig. 7-9) and estimated reference F values similar to other Bering Sea flatfish species.

- 3) Based on selectivity patterns, the shelf survey showed big differences in the ages of fish available to these different surveys (Fig. 7-10). The slope survey selectivity estimates seemed most stable hence: Alternative values of q were fixed for the slope survey and q values for the shelf and Aleutian Islands surveys were freely estimated.
- 4) Since q is confounded with natural mortality, M was estimated as a free parameter but the model would not converge even though q was fixed for the slope survey. Profiling over M with catchability fixed for only one of the three surveys gave a value of 0.09. It was decided to continue the use of $M=0.11$ for this assessment as in the past but need to explore estimating M (profiling) with q fixed for all three surveys.

Estimates of catchability from the slope survey profile and the associated likelihood indicated that slope q is less than 0.3, but flat from about 0.2-0.5. Estimates of female spawning biomass derived from slope $q = 0.1$ and $q = 0.18$ are shown in figure 7-11. The difference in total likelihood between these models was only 1.95, with the $q=0.1$ model being favored (in terms of total log likelihood) since the best fit based on the overall likelihood is a low slope q (Fig. 7-12). Since the likelihood surface was so flat between $q=0.1$ and 0.18, the fixed value of 0.18 was retained for slope q . With the model configured in this way (slope survey $q=0.18$, $M = 0.11$ and fishery selectivity logistic slope fixed) the model was run to estimate the status and the population dynamics of the Kamchatka flounder stock over the period 1991-2014.

Time Series Results

Model results estimate that the total biomass of Kamchatka flounder steadily increased from 1991 to 2009 to about 170,000 t, then declined by 17,000 t in 2011 before increasing to 171,000 t in 2016 (Fig. 7-13). The female spawning biomass trend mirrors the total biomass with a parallel trend that peaks at 61,000 t in 2009 and has remained stable through 2016 (Fig. 7-13, Fig. 7.14 and Table 7.10). The model estimates of shelf, slope and Aleutian Islands surveys fit the trends estimated by those data sources reasonably well (Fig. 7-15). Selectivity, as previously discussed, was constrained for the fishery and was freely estimated for the surveys. It is clear that the shelf survey samples a younger portion of the population than those surveys conducted on the Bering Sea slope and in the Aleutian Islands (Fig. 7-10). Model estimates of male and female numbers-at-age are shown in Table 7.11 and estimated female spawning biomass-at-age in Table 7.12.

Model estimates of fishing mortality indicate that the stock was lightly harvested from 1991 to 2007, with an average annual full selection F of 0.019 (Fig 7-16). As the fishery developed for Kamchatka flounder in 2008 the fishing mortality was much higher peaking at 0.155 in 2010. For the last 4 years fishing mortality has averaged 0.045. The average annual F for the past 5 years is 0.044, well-below the $F_{40\%}$ value of 0.066.

Examination of the model fit to the survey length composition data was made by comparing the average observed proportion at length from the time-series to the average predicted proportion at length from the model (Fig. 7-17). Overall the model fits the general shape of the length compositions but has some residual trends for large fish on the slope and the Aleutian Islands. Fits to the individual annual length compositions, by sex, are shown in Figure 7.18.

Projections and Harvest Recommendations

The reference fishing mortality rate for Kamchatka flounder is determined by the amount of reliable population information available (Amendment 56 of the Fishery Management Plan for the groundfish fishery of the Bering Sea/Aleutian Islands). Estimates of $B_{40\%}$, $F_{40\%}$, and $SPR_{40\%}$ were obtained from a spawner-per-recruit analysis. Assuming that the average recruitment from 1989-2011 year classes estimated in this assessment represents a reliable estimate of equilibrium recruitment, then an estimate of

$B_{40\%}$ is calculated as the product of $SPR_{40\%}$ * equilibrium recruits (=50,800 t). The 2017 spawning biomass is estimated at 60,300 t. Since reliable estimates of 2017 spawning biomass (B), $B_{40\%}$, $F_{40\%}$, and $F_{35\%}$ exist and $B > B_{40\%}$ (60,300 t > 50,800 t shown in Fig. 7.14 and Fig. 7.19), The reference fishing mortality for Kamchatka flounder is defined in tier 3a of Amendment 56. For this tier, F_{ABC} is constrained to be $\leq F_{40\%}$, and F_{OFL} is defined as $F_{35\%}$. The values of these quantities are:

| | | |
|---------------------------|---|----------|
| 2017 SSB estimate (B) | = | 60,300 t |
| $B_{40\%}$ | = | 50,800 t |
| $F_{40\%}$ | = | 0.066 |
| F_{ABC} | = | 0.066 |
| $F_{35\%}$ | = | 0.078 |
| F_{OFL} | = | 0.078 |

The estimated catch level for year 2017 associated with the overfishing level of $F = 0.078$ is 10,360 t. **The 2017 recommended ABC associated with F_{ABC} of 0.066 is 8,880 t.** Projections of Kamchatka flounder female spawning biomass (described below) at a harvest rate equal to the average fishing mortality rate of the past five years indicate that the stock could increase to a stable level of over 70,000 t from 2022-2029 (Fig. 7.14).

A standard set of projections is required for each stock managed under Tiers 1, 2, or 3 of Amendment 56. This set of projections encompasses seven harvest scenarios designed to satisfy the requirements of Amendment 56, the National Environmental Policy Act, and the Magnuson-Stevens Fishery Conservation and Management Act (MSFCMA).

For each scenario, the projections begin with the vector of 2016 numbers at age estimated in the assessment. This vector is then projected forward to the beginning of 2017 using the schedules of natural mortality and selectivity described in the assessment and the best available estimate of total (year-end) catch for 2016. In each subsequent year, the fishing mortality rate is prescribed on the basis of the spawning biomass in that year and the respective harvest scenario. In each year, recruitment is drawn from an inverse Gaussian distribution whose parameters consist of maximum likelihood estimates determined from recruitments estimated in the assessment. Spawning biomass is computed in each year based on the time of peak spawning and the maturity and weight schedules described in the assessment. Total catch is assumed to equal the catch associated with the respective harvest scenario in all years. This projection scheme is run 1000 times to obtain distributions of possible future stock sizes, fishing mortality rates, and catches.

Five of the seven standard scenarios will be used in an Environmental Assessment prepared in conjunction with the final SAFE. These five scenarios, which are designed to provide a range of harvest alternatives that are likely to bracket the final TAC for 2017, are as follows (“ $max F_{ABC}$ ” refers to the maximum permissible value of F_{ABC} under Amendment 56):

Scenario 1: In all future years, F is set equal to $max F_{ABC}$. (Rationale: Historically, TAC has been constrained by ABC, so this scenario provides a likely upper limit on future TACs.)

Scenario 2: In all future years, F is set equal to a constant fraction of $max F_{ABC}$, where this fraction is equal to the ratio of the F_{ABC} value for 2017 recommended in the assessment to the $max F_{ABC}$ for 2017. (Rationale: When F_{ABC} is set at a value below $max F_{ABC}$, it is often set at the value recommended in the stock assessment.)

Scenario 3: In all future years, F is set equal to 50% of $max F_{ABC}$. (Rationale: This scenario provides a likely lower bound on F_{ABC} that still allows future harvest rates to be adjusted downward when stocks fall below reference levels.)

Scenario 4: In all future years, F is set equal to the 2012-2016 average F . (Rationale: For some stocks, TAC can be well below ABC, and recent average F may provide a better indicator of F_{TAC} than F_{ABC} .)

Scenario 5: In all future years, F is set equal to zero. (Rationale: In extreme cases, TAC may be set at a level close to zero.)

The recommended F_{ABC} and the maximum F_{ABC} are equivalent in this assessment, and five-year projections of the mean Kamchatka flounder harvest and spawning stock biomass for the remaining four scenarios are shown in Table 10.11.

Two other scenarios are needed to satisfy the MSFCMA’s requirement to determine whether the Alaska plaice stock is currently in an overfished condition or is approaching an overfished condition. These two scenarios are as follows (for Tier 3 stocks, the MSY level is defined as $B_{35\%}$):

Scenario 6: In all future years, F is set equal to F_{OFL} . (Rationale: This scenario determines whether a stock is overfished. If the stock is expected to be above its MSY level in 2017 under this scenario, then the stock is not overfished.)

Scenario 7: In 2017 and 2018, F is set equal to $max F_{ABC}$, and in all subsequent years, F is set equal to F_{OFL} . (Rationale: This scenario determines whether a stock is approaching an overfished condition. If the stock is expected to be above its MSY level in 2029 under this scenario, then the stock is not approaching an overfished condition.)

The results of these two scenarios indicate that Kamchatka flounder are neither overfished nor approaching an overfished condition. With regard to assessing the current stock level, the expected stock size in the year 2016 of scenario 6 is well above its $B_{35\%}$ value of 44,400 t. With regard to whether the stock is likely to be in an overfished condition in the near future, the expected stock size in the year 2029 of scenario 7 is also greater than its $B_{35\%}$ value. Figure 7.19 shows the relationship between the estimated time-series of female spawning biomass and fishing mortality and the tier 3 control rule for Kamchatka flounder. The simulation results for the 7 harvest scenarios are shown in Table 7.13.

Scenario Projections and Two-Year Ahead Overfishing Level

In addition to the seven standard harvest scenarios, Amendments 48/48 to the BSAI and GOA Groundfish Fishery Management Plans require projections of the likely OFL two years into the future. While Scenario 6 gives the best estimate of OFL for 2017, it does not provide the best estimate of OFL for 2018, because the mean 2018 catch under Scenario 6 is predicated on the 2017 catch being equal to the 2017 OFL, whereas the actual 2017 catch will likely be less than the 2017 ABC. Therefore, the projection model was re-run with the 2018 catch fixed at the 2017 level.

| Year | Catch | ABC | OFL |
|------|-------|-------|--------|
| 2017 | 4,530 | 8,880 | 10,360 |
| 2018 | 4,530 | 9,200 | 10,700 |

Retrospective analysis

Retrospective patterns of female spawning biomass estimates were investigated from stock assessment model estimates for 2007-2016 (Figure 7.20). Between year model estimates exhibited jumps in FSB between groups of years relative to the 2016 reference year estimates as the population increased. Groups in 2007-2009, 2010-2013 and 2014-2016 represent different levels of increasing biomass being fit by the assessment model from the three surveys. Differences of 20% were seen between the reference model estimate and the terminal year estimates from 2007-2010. Survey biomass increases are likely affecting the retrospective trends. Mohn's statistic was computed at 0.12 for Kamchatka flounder.

Ecosystem Considerations

Predators of Kamchatka flounder

Kamchatka flounder have rarely been found in the stomachs of other groundfish species in samples collected by the Alaska Fisheries Science Center. Their presence has only been documented in 17 stomach samples from the BSAI where the predators included Pacific cod, pollock, Pacific halibut, arrowtooth flounder and two sculpin species.

Kamchatka flounder predation

The prey of Kamchatka flounder can be discerned from 152 stomachs collected in 1983 (Yang and Livingston 1986). The principle diet was composed of walleye pollock, shrimp (mostly Crangonidae) and euphausiids. Pollock was the most important prey item for all sizes of fish, ranging from 56 to 86% of the total stomach content weight. An examination of diet overlap with arrowtooth flounder indicated that these two congeneric species basically consume the same resources. Therefore the following sections are from the arrowtooth flounder assessment but pertain to Kamchatka flounder.

Ecosystem Effects on the stock

Prey availability/abundance trends

Arrowtooth flounder diet varies by life stage as indicated in the previous section. Regarding juvenile prey and its associated habitat, information is not available to assess the abundance trends of the benthic infauna of the Bering Sea shelf. The original description of infaunal distribution and abundance by Haflinger (1981) resulted from sampling conducted in 1975 and 1976 and has not been re-sampled since. Information on pollock abundance is available in Chapter 1 of this SAFE report. It has been hypothesized that predators on pollock, such as adult arrowtooth flounder, may be important species which control (with other factors) the variation in year-class strength of juvenile pollock (Hunt et al. 2002). The populations of arrowtooth flounder which have occupied the outer shelf and slope areas of the Bering Sea over the past twenty years for summertime feeding do not appear food-limited. These populations have fluctuated due to the variability in recruitment success which suggests that the primary infaunal food source has been at an adequate level to sustain the arrowtooth flounder resource.

Predator population trends

As juveniles, it is well-documented from studies in other parts of the world that flatfish are prey for shrimp species in near shore areas. This has not been reported for Bering Sea arrowtooth flounder due to a lack of juvenile sampling and collections in near shore areas, but is thought to occur. As late juveniles they are found in stomachs of pollock and Pacific cod, mostly on small arrowtooth flounder ranging from 5 to 15 cm standard length..

Past, present and projected future population trends of these predator species can be found in their respective SAFE chapters in this volume. Encounters between arrowtooth flounder and their predators may be limited as their distributions do not completely overlap in space and time.

Changes in habitat quality

Changes in the physical environment which may affect Kamchatka flounder distribution patterns, recruitment success, migration timing and patterns are catalogued in the Ecosystem Considerations Appendix of this SAFE report. Habitat quality may be enhanced during years and warmer bottom water temperatures with reduced ice cover (higher metabolism with more active feeding). Environmental factors important to juvenile survival are presently not well known.

Fishery Effects on the Ecosystem

Ecosystem effects on Kamchatka flounder

| Indicator | Observation | Interpretation | Evaluation |
|--|--|--|----------------------------------|
| <i>Prey availability or abundance trends</i> | | | |
| Benthic infauna | Stomach contents | Stable, data limited | Unknown |
| <i>Predator population trends</i> | | | |
| Fish (Pollock, Pacific cod) | Stable | Possible increases to Kamchatka mortality | |
| <i>Changes in habitat quality</i> | | | |
| Temperature regime | Cold years Kamchatka catchability and herding may decrease | Deeper water species so less likely to affect surveyed stock | No concern (dealt with in model) |
| Winter-spring environmental conditions | Affects pre-recruit survival | Probably a number of factors | Causes natural variability |

Arrowtooth flounder effects on ecosystem

| Indicator | Observation | Interpretation | Evaluation |
|--|---|---|------------------|
| <i>Fishery contribution to bycatch</i> | | | |
| Prohibited species | Stable, heavily monitored | Minor contribution to mortality | No concern |
| Forage (including Pollock, shrimp and euphausiids) | Stable, heavily monitored | Bycatch levels small relative to forage biomass | No concern |
| HAPC biota | Low bycatch levels of (spp) | Bycatch levels small relative to HAPC biota | No concern |
| Marine mammals and birds | Very minor direct-take | Safe | No concern |
| Sensitive non-target species | Likely minor impact | Data limited, likely to be safe | No concern |
| <i>Fishery concentration in space and time</i> | | | |
| | Recent high exploitation rate | Little detrimental effect | No concern |
| <i>Fishery effects on amount of large size target fish</i> | | | |
| | Recent high exploitation rate, but unknown effect | Natural fluctuation | No concern |
| <i>Fishery contribution to discards and offal production</i> | | | |
| | Stable trend | Improving, but data limited | Possible concern |
| <i>Fishery effects on age-at-maturity and fecundity</i> | | | |
| | Unknown | NA | Possible concern |

Data Gaps and Research Priorities

A significant improvement in the estimate of fishery selectivity would likely result from an increase in the amount of Kamchatka flounder length data collected when Kamchatka flounder are targeted in the commercial fishery.

References

- Stark, J. W. 2011. Female maturity, reproductive potential, relative distribution, and growth compared between arrowtooth flounder (*Atheresthes stomias*) and Kamchatka flounder (*A. evermanni*) indicating concerns for management. *J. Appl. Ichthyol.* (2011) 1-5.
- Wilderbuer, T. K., D. G. Nichol, and K. Aydin 2009. Arrowtooth flounder. In Stock Assessment and Fishery Evaluation Document for Groundfish Resources in the Bering Sea/Aleutian Islands Region as Projected for 2010, Chapter 6. North Pacific Fishery Management Council, P.O. Box 103136, Anchorage Alaska 99510.
- Wilimovsky, N. J., A. Peden, and J. Peppar. 1967. Systematics of six demersal fishes of the north Pacific Ocean. *Fish. Res. Board Can., Tech. Rep.* 34, 52 p.
- Yang, M. S. and P. A. Livingston. 1986. Food habits and diet overlap of two congeneric species, *Atheresthes stomias* and *Atheresthes evermanni*, in the eastern Bering Sea. *Fish. Bull.* Vol. 82 (8)615-623.
- Zimmermann, Mark, and Pamela Goddard 1996. Biology and distribution of arrowtooth (*Atheresthes stomias*) and Kamachatka (*A. evermanni*) flounders in Alaskan waters. *Fish. Bull* 94:358-370.

Table 7-1. Total combined catch (t) of arrowtooth and Kamchatka flounder in the eastern Bering Sea and Aleutian Islands region, 2001-2006. Catches from 2007 to present, when the two species were differentiated in commercial catches, are reported for Kamchatka flounder only in this table.

| year | catch | TAC | ABC | OFL |
|------|--------|--------|--------|--------|
| 1991 | 22,052 | | | |
| 1992 | 10,382 | | | |
| 1993 | 9,338 | | | |
| 1994 | 14,366 | | | |
| 1995 | 9,280 | | | |
| 1996 | 14,652 | | | |
| 1997 | 10,054 | | | |
| 1998 | 15,241 | | | |
| 1999 | 10,573 | | | |
| 2000 | 12,929 | | | |
| 2001 | 13,908 | | | |
| 2002 | 11,540 | | | |
| 2003 | 12,834 | | | |
| 2004 | 17,809 | | | |
| 2005 | 13,685 | | | |
| 2006 | 13,309 | | | |
| 2007 | 1,183 | | | |
| 2008 | 6,819 | | | |
| 2009 | 12,802 | | | |
| 2010 | 21,153 | | | |
| 2011 | 9,935 | 17,700 | 17,700 | 23,600 |
| 2012 | 9,514 | 17,700 | 18,600 | 24,800 |
| 2013 | 7,772 | 10,000 | 12,200 | 16,300 |
| 2014 | 6,220 | 7,100 | 7,100 | 8,270 |
| 2015 | 4,994 | 6,500 | 9,000 | 10,500 |
| 2016 | 4,533 | 6,500 | 9,500 | 11,000 |

Table 7-2. Estimated Kamchatka flounder biomass and coefficient of variation (CV) from the three BSAI bottom trawl surveys (shelf, slope, and Aleutian Islands). Reliable estimates of Kamchatka flounder biomass are only available after 1991 when Kamchatka and arrowtooth flounder were differentiated.

| | shelf | shelf CV | slope | slope CV | Aleutian Islands | Aleutian Islands CV |
|------|--------|-------------|--------|-------------|------------------|------------------------|
| 1982 | 0 | | | | | |
| 1983 | 17,299 | | | | 1,034 | |
| 1984 | 20,695 | | | | | |
| 1985 | 31 | | | | | |
| 1986 | 0 | | | | 565 | |
| 1987 | 40 | | | | | |
| 1988 | 13,723 | | | | | |
| 1989 | 17,108 | | | | | |
| 1990 | 32,799 | | | | | |
| 1991 | 37,152 | 0.11 | | | 16,255 | 0.27 |
| 1992 | 50,081 | 0.11 | | | | |
| 1993 | 38,376 | 0.09 | | | | |
| 1994 | 56,268 | 0.12 | | | 49,156 | 0.38 |
| 1995 | 28,393 | 0.10 | | | | |
| 1996 | 24,196 | 0.10 | | | | |
| 1997 | 18,282 | 0.10 | | | 37,664 | 0.25 |
| 1998 | 23,474 | 0.09 | | | | |
| 1999 | 18,974 | 0.14 | | | | |
| 2000 | 21,551 | 0.11 | | | 28,535 | 0.23 |
| 2001 | 31,120 | 0.09 | | | | |
| 2002 | 25,213 | 0.12 | 18,631 | 0.11 | 49,035 | 0.28 |
| 2003 | 27,531 | 0.11 | | | | |
| 2004 | 29,663 | 0.09 | 14,740 | 0.10 | 39,219 | 0.24 |
| 2005 | 46,084 | 0.07 | | | | |
| 2006 | 61,644 | 0.08 | | | 45,369 | 0.24 |
| 2007 | 65,191 | 0.08 | | | | |
| 2008 | 53,967 | 0.10 | 24,822 | 0.19 | | |
| 2009 | 47,252 | 0.11 | | | | |
| 2010 | 51,927 | 0.08 | 27,856 | 0.10 | 49,069 | 0.38 |
| 2011 | 46,094 | 0.09 | | | | |
| 2012 | 40,951 | 0.08 | 32,685 | 0.22 | 35,100 | 0.40 |
| 2013 | 46,380 | 0.08 | | | | |
| 2014 | 58,036 | 0.02 | | | 45,157 | 0.37 |
| 2015 | 60,331 | 0.06 | | | | |
| 2016 | 55,324 | 0.057 | 21,369 | 0.097 | 27,968 | 0.23 |

Table 7.3. Kamchatka flounder sample sizes from the Eastern Bering Sea shelf survey. The hauls columns refer to the number of hauls from which either lengths or otoliths were obtained.

| Year | Total Hauls | Hauls w/Lengths | Number lengths | Hauls w/otoliths | Hauls w/ages | Number otoliths | Number ages |
|------|-------------|-----------------|----------------|------------------|--------------|-----------------|-------------|
| 1982 | 334 | | | | | | |
| 1983 | 353 | 13 | 692 | | | | |
| 1984 | 355 | 27 | 741 | | | | |
| 1985 | 357 | | | | | | |
| 1986 | 354 | | | | | | |
| 1987 | 357 | 1 | 5 | | | | |
| 1988 | 373 | 18 | 142 | | | | |
| 1989 | 374 | 33 | 424 | | | | |
| 1990 | 371 | 51 | 643 | | | | |
| 1991 | 372 | 92 | 1056 | | | | |
| 1992 | 356 | 98 | 1039 | 20 | | 165 | |
| 1993 | 375 | 146 | 1117 | 15 | | 148 | |
| 1994 | 375 | 122 | 1241 | | | | |
| 1995 | 376 | 100 | 816 | 7 | | 74 | |
| 1996 | 375 | 136 | 826 | 9 | | 103 | |
| 1997 | 376 | 100 | 698 | 2 | | 31 | |
| 1998 | 375 | 138 | 1099 | | | | |
| 1999 | 373 | 94 | 805 | | | | |
| 2000 | 372 | 124 | 1054 | | | | |
| 2001 | 375 | 127 | 1111 | | | | |
| 2002 | 375 | 118 | 1053 | | | | |
| 2003 | 376 | 158 | 1530 | | | | |
| 2004 | 375 | 165 | 3034 | | | | |
| 2005 | 373 | 182 | 3582 | | | | |
| 2006 | 376 | 141 | 4126 | | | | |
| 2007 | 376 | 132 | 2954 | | | | |
| 2008 | 375 | 154 | 2724 | | | | |
| 2009 | 376 | 132 | 2074 | | | | |
| 2010 | 376 | 160 | 3219 | | | | |
| 2011 | 376 | 189 | 2130 | | | | |
| 2012 | 376 | 136 | 2953 | | | | |
| 2013 | 376 | 151 | 2954 | 62 | | 519 | |
| 2014 | 376 | 185 | 2490 | 31 | | 314 | |
| 2015 | 376 | 195 | 2953 | 65 | | 453 | |
| 2016 | 376 | 244 | 3081 | 74 | | 524 | |

Table 7-4. Bering Sea shelf survey female size composition estimates (1,000s of fish).

| year/size (cm) | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22 | 23 | 24 | 25 | 26 | 27 | 28 | 29 | 30 | 31 | 32 | 33 |
|----------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
| 1991 | 68 | 203 | 725 | 354 | 80 | 155 | 0 | 226 | 158 | 745 | 718 | 624 | 1837 | 642 | 1030 | 569 | 1842 | 3414 | 2868 |
| 1992 | 256 | 148 | 1318 | 221 | 354 | 0 | 50 | 318 | 568 | 494 | 663 | 146 | 267 | 715 | 232 | 2052 | 1190 | 1069 | 1221 |
| 1993 | 239 | 166 | 463 | 884 | 151 | 274 | 116 | 284 | 735 | 1047 | 436 | 620 | 274 | 239 | 327 | 301 | 1115 | 199 | 533 |
| 1994 | 74 | 219 | 221 | 168 | 40 | 211 | 271 | 566 | 417 | 597 | 770 | 809 | 898 | 692 | 962 | 901 | 976 | 1163 | 821 |
| 1995 | 0 | 56 | 55 | 0 | 115 | 59 | 27 | 53 | 199 | 56 | 107 | 113 | 63 | 215 | 190 | 498 | 575 | 978 | 471 |
| 1996 | 0 | 123 | 48 | 185 | 63 | 90 | 23 | 66 | 126 | 173 | 224 | 184 | 249 | 220 | 127 | 154 | 152 | 224 | 248 |
| 1997 | 100 | 402 | 200 | 570 | 190 | 239 | 338 | 172 | 142 | 308 | 149 | 149 | 243 | 58 | 81 | 58 | 59 | 99 | 60 |
| 1998 | 292 | 314 | 352 | 345 | 299 | 214 | 380 | 250 | 585 | 1012 | 713 | 675 | 583 | 739 | 392 | 419 | 275 | 354 | 169 |
| 1999 | 195 | 220 | 273 | 192 | 146 | 110 | 145 | 163 | 167 | 405 | 636 | 904 | 674 | 518 | 705 | 312 | 1164 | 653 | 701 |
| 2000 | 141 | 201 | 0 | 66 | 0 | 92 | 215 | 376 | 732 | 1052 | 740 | 689 | 410 | 200 | 558 | 612 | 504 | 882 | 699 |
| 2001 | 189 | 64 | 205 | 318 | 306 | 422 | 222 | 309 | 789 | 451 | 240 | 137 | 163 | 454 | 936 | 1478 | 1792 | 1250 | 1489 |
| 2002 | 551 | 886 | 901 | 455 | 383 | 170 | 171 | 226 | 426 | 764 | 364 | 193 | 447 | 394 | 537 | 620 | 347 | 471 | 254 |
| 2003 | 86 | 146 | 318 | 686 | 795 | 681 | 814 | 743 | 1254 | 1263 | 1166 | 985 | 663 | 506 | 348 | 347 | 387 | 403 | 520 |
| 2004 | 830 | 1984 | 4070 | 7817 | 7175 | 6595 | 4630 | 3755 | 2096 | 1210 | 1420 | 2549 | 1392 | 2147 | 1386 | 1495 | 1608 | 1823 | 1522 |
| 2005 | 1237 | 1273 | 2253 | 2161 | 1729 | 2570 | 2342 | 4279 | 2417 | 4199 | 5980 | 8191 | 8525 | 6565 | 6816 | 5023 | 3579 | 2822 | 2749 |
| 2006 | 282 | 425 | 313 | 428 | 368 | 953 | 547 | 526 | 1016 | 1756 | 2723 | 4636 | 4468 | 5828 | 8832 | 6877 | 5699 | 7239 | 5673 |
| 2007 | 351 | 1233 | 1528 | 1456 | 566 | 213 | 134 | 265 | 529 | 640 | 363 | 768 | 733 | 1158 | 1976 | 2840 | 2891 | 4988 | 4905 |
| 2008 | 214 | 513 | 714 | 627 | 586 | 872 | 920 | 1874 | 1356 | 706 | 465 | 314 | 700 | 852 | 651 | 1136 | 1314 | 1240 | 1458 |
| 2009 | 141 | 449 | 705 | 617 | 138 | 218 | 346 | 464 | 630 | 484 | 871 | 956 | 777 | 874 | 1031 | 883 | 1393 | 623 | 611 |
| 2010 | 1018 | 3640 | 3874 | 3340 | 1154 | 1279 | 1557 | 2453 | 2387 | 1294 | 1168 | 877 | 866 | 1064 | 1016 | 941 | 1298 | 1261 | 1221 |
| 2011 | 450 | 687 | 840 | 543 | 574 | 343 | 1300 | 1119 | 2382 | 2352 | 2168 | 1270 | 2209 | 1047 | 1419 | 1225 | 1172 | 996 | 724 |
| 2012 | 1542 | 1332 | 1371 | 2258 | 1382 | 1685 | 1331 | 1917 | 1704 | 2523 | 1457 | 1958 | 2477 | 2608 | 2369 | 2777 | 1704 | 1764 | 1001 |
| 2013 | 393 | 792 | 379 | 1301 | 332 | 395 | 752 | 2027 | 2519 | 2904 | 3408 | 3037 | 2136 | 2164 | 1966 | 2067 | 2035 | 3718 | 2863 |
| 2014 | 867 | 1246 | 1027 | 2452 | 1037 | 763 | 705 | 636 | 326 | 1393 | 1048 | 1298 | 1020 | 1404 | 2450 | 3534 | 4021 | 4350 | 3435 |
| 2015 | 1230 | 1169 | 1264 | 856 | 888 | 685 | 881 | 528 | 517 | 826 | 1086 | 1541 | 2468 | 1535 | 780 | 1393 | 836 | 1402 | 1240 |
| 2016 | 710 | 848 | 1289 | 1360 | 1346 | 2207 | 1634 | 1431 | 1694 | 1758 | 1379 | 1599 | 2212 | 859 | 1465 | 1385 | 2116 | 2217 | 1593 |

Table 7-4. Continued.

| year/size (cm) | 34 | 35 | 36 | 37 | 38 | 39 | 40 | 41 | 42 | 43 | 44 | 45 | 46 | 47 | 48 | 49 | 50 | 51 | 52 | 53 | 54 |
|----------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|-----|
| 1991 | 1873 | 2059 | 2215 | 1964 | 2991 | 1933 | 780 | 1052 | 1002 | 543 | 486 | 339 | 1502 | 606 | 175 | 384 | 416 | 682 | 425 | 202 | 45 |
| 1992 | 1462 | 1334 | 730 | 1315 | 2028 | 2309 | 2075 | 2487 | 2236 | 2491 | 1476 | 1320 | 1467 | 514 | 799 | 211 | 669 | 568 | 315 | 1158 | 94 |
| 1993 | 471 | 606 | 983 | 639 | 844 | 424 | 1464 | 913 | 1258 | 1241 | 2060 | 1471 | 1065 | 915 | 1136 | 798 | 636 | 1011 | 254 | 468 | 407 |
| 1994 | 501 | 686 | 721 | 547 | 760 | 883 | 743 | 1349 | 877 | 1082 | 1180 | 2089 | 1239 | 1231 | 1041 | 1185 | 955 | 1585 | 1068 | 544 | 712 |
| 1995 | 380 | 553 | 857 | 317 | 654 | 427 | 254 | 490 | 601 | 394 | 221 | 643 | 399 | 658 | 562 | 580 | 614 | 448 | 430 | 463 | 344 |
| 1996 | 469 | 469 | 272 | 402 | 851 | 730 | 289 | 404 | 590 | 312 | 495 | 389 | 219 | 363 | 347 | 321 | 421 | 408 | 479 | 544 | 744 |
| 1997 | 173 | 276 | 59 | 88 | 115 | 59 | 267 | 413 | 522 | 473 | 642 | 574 | 607 | 334 | 208 | 116 | 212 | 61 | 275 | 191 | 404 |
| 1998 | 295 | 319 | 254 | 123 | 96 | 473 | 144 | 273 | 454 | 717 | 458 | 472 | 392 | 524 | 249 | 308 | 601 | 139 | 338 | 254 | 133 |
| 1999 | 442 | 173 | 583 | 226 | 327 | 406 | 413 | 180 | 192 | 127 | 419 | 85 | 128 | 126 | 114 | 320 | 258 | 257 | 213 | 408 | 364 |
| 2000 | 1152 | 558 | 420 | 455 | 659 | 255 | 315 | 617 | 143 | 63 | 152 | 31 | 129 | 209 | 98 | 172 | 203 | 203 | 77 | 185 | 368 |
| 2001 | 529 | 618 | 697 | 1148 | 988 | 1074 | 878 | 586 | 537 | 806 | 498 | 382 | 457 | 279 | 264 | 475 | 251 | 176 | 57 | 285 | 46 |
| 2002 | 414 | 554 | 1041 | 1002 | 841 | 294 | 367 | 418 | 414 | 491 | 807 | 620 | 781 | 774 | 447 | 151 | 545 | 91 | 255 | 154 | 109 |
| 2003 | 564 | 638 | 505 | 332 | 518 | 581 | 286 | 1111 | 474 | 227 | 608 | 268 | 377 | 320 | 321 | 233 | 308 | 194 | 236 | 29 | 48 |
| 2004 | 1220 | 736 | 979 | 604 | 473 | 849 | 189 | 1031 | 298 | 287 | 231 | 264 | 338 | 219 | 267 | 176 | 240 | 212 | 326 | 250 | 110 |
| 2005 | 2551 | 2146 | 2598 | 2160 | 781 | 1673 | 760 | 498 | 349 | 229 | 284 | 131 | 394 | 92 | 513 | 137 | 491 | 151 | 193 | 172 | 149 |
| 2006 | 6141 | 5321 | 4762 | 3063 | 3697 | 2665 | 1184 | 1465 | 633 | 557 | 390 | 515 | 229 | 173 | 247 | 272 | 27 | 29 | 88 | 210 | |
| 2007 | 5214 | 4622 | 5165 | 6456 | 5419 | 4120 | 3755 | 2137 | 2344 | 879 | 1062 | 518 | 310 | 707 | 59 | 372 | 226 | 524 | 166 | 90 | 62 |
| 2008 | 2962 | 2428 | 2548 | 2968 | 4291 | 3005 | 4431 | 3825 | 3956 | 2345 | 1492 | 953 | 964 | 303 | 939 | 145 | 369 | 430 | 62 | 166 | 130 |
| 2009 | 524 | 1053 | 857 | 1462 | 1762 | 2310 | 2399 | 2053 | 2331 | 2612 | 2974 | 2340 | 1667 | 1450 | 1426 | 438 | 329 | 288 | 159 | 78 | 113 |
| 2010 | 2271 | 1945 | 828 | 972 | 363 | 1513 | 913 | 901 | 1775 | 1373 | 1527 | 1892 | 2244 | 1869 | 2115 | 1699 | 1605 | 508 | 557 | 551 | 192 |
| 2011 | 736 | 968 | 736 | 705 | 813 | 759 | 401 | 830 | 518 | 479 | 661 | 745 | 1241 | 1120 | 1675 | 1199 | 1328 | 1035 | 1283 | 917 | 569 |
| 2012 | 945 | 598 | 837 | 297 | 935 | 697 | 818 | 559 | 794 | 468 | 757 | 425 | 726 | 575 | 605 | 619 | 1171 | 734 | 757 | 840 | 904 |
| 2013 | 3022 | 2257 | 1661 | 1544 | 974 | 1028 | 828 | 494 | 766 | 483 | 904 | 734 | 427 | 455 | 478 | 388 | 499 | 434 | 643 | 880 | 401 |
| 2014 | 2388 | 2071 | 1534 | 2045 | 1723 | 3054 | 1173 | 2385 | 983 | 958 | 605 | 1575 | 309 | 448 | 653 | 472 | 181 | 325 | 519 | 482 | 436 |
| 2015 | 1831 | 2599 | 2459 | 2650 | 3109 | 3007 | 2447 | 3043 | 2921 | 2273 | 2257 | 1590 | 1424 | 1251 | 486 | 338 | 700 | 354 | 382 | 282 | 610 |
| 2016 | 2106 | 1160 | 1407 | 1659 | 1597 | 1625 | 2576 | 1646 | 2307 | 2215 | 2417 | 1330 | 1179 | 616 | 982 | 1019 | 1207 | 347 | 294 | 334 | 528 |

Table 7-4. Continued.

| year/size (cm) | 55 | 56 | 57 | 58 | 59 | 60 | 61 | 62 | 63 | 64 | 65 | 66 | 67 | 68 | 69 | 70+ |
|----------------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| 1991 | 103 | 205 | 136 | 33 | 45 | 0 | 0 | 206 | 178 | 37 | 33 | 0 | 43 | 0 | 0 | 0 |
| 1992 | 613 | 222 | 228 | 384 | 288 | 282 | 0 | 68 | 123 | 0 | 35 | 0 | 0 | 0 | 0 | 37 |
| 1993 | 435 | 306 | 416 | 400 | 299 | 150 | 0 | 94 | 136 | 0 | 57 | 28 | 116 | 28 | 164 | 116 |
| 1994 | 679 | 118 | 454 | 376 | 595 | 146 | 185 | 527 | 190 | 299 | 35 | 88 | 0 | 0 | 52 | 123 |
| 1995 | 469 | 611 | 353 | 50 | 30 | 140 | 236 | 66 | 25 | 0 | 27 | 66 | 0 | 62 | 0 | 0 |
| 1996 | 332 | 212 | 418 | 331 | 103 | 159 | 143 | 32 | 174 | 189 | 150 | 0 | 65 | 0 | 0 | 148 |
| 1997 | 135 | 180 | 165 | 232 | 179 | 77 | 27 | 0 | 98 | 32 | 58 | 89 | 57 | 0 | 29 | 98 |
| 1998 | 90 | 288 | 328 | 168 | 140 | 239 | 153 | 75 | 184 | 148 | 135 | 86 | 131 | 0 | 0 | 288 |
| 1999 | 114 | 139 | 67 | 97 | 0 | 0 | 128 | 328 | 213 | 62 | 32 | 33 | 0 | 84 | 150 | 334 |
| 2000 | 179 | 174 | 307 | 29 | 228 | 57 | 49 | 82 | 0 | 43 | 167 | 271 | 57 | 58 | 229 | 612 |
| 2001 | 257 | 120 | 193 | 167 | 213 | 176 | 124 | 23 | 374 | 172 | 29 | 256 | 731 | 73 | 143 | 293 |
| 2002 | 105 | 209 | 469 | 82 | 337 | 78 | 112 | 0 | 166 | 0 | 0 | 161 | 247 | 164 | 75 | 175 |
| 2003 | 237 | 62 | 0 | 114 | 327 | 104 | 200 | 56 | 58 | 223 | 0 | 81 | 280 | 110 | 127 | 394 |
| 2004 | 188 | 46 | 179 | 53 | 82 | 0 | 46 | 66 | 169 | 53 | 169 | 47 | 102 | 28 | 18 | 582 |
| 2005 | 295 | 0 | 305 | 0 | 146 | 108 | 35 | 107 | 149 | 92 | 128 | 0 | 25 | 0 | 45 | 611 |
| 2006 | 54 | 55 | 29 | 27 | 28 | 144 | 123 | 29 | 27 | 0 | 53 | 0 | 103 | 28 | 84 | 110 |
| 2007 | 123 | 30 | 592 | 106 | 479 | 120 | 131 | 57 | 29 | 222 | 178 | 286 | 234 | 0 | 145 | 192 |
| 2008 | 167 | | | | | | | | | | | | | | | |

Table 7-4. Bering Sea shelf survey male size composition estimates (1,000s of fish).

| year/size (cm) | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22 | 23 | 24 | 25 | 26 | 27 | 28 | 29 | 30 | 31 | 32 | 33 |
|----------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
| 1991 | 75 | 217 | 160 | 15 | 168 | 0 | 28 | 267 | 291 | 670 | 253 | 1175 | 277 | 759 | 777 | 976 | 2044 | 2690 | 2244 |
| 1992 | 156 | 770 | 1094 | 1225 | 217 | 41 | 144 | 490 | 220 | 710 | 577 | 294 | 169 | 399 | 511 | 898 | 1132 | 863 | 456 |
| 1993 | 98 | 934 | 790 | 1246 | 384 | 92 | 28 | 402 | 764 | 475 | 435 | 213 | 369 | 243 | 586 | 713 | 623 | 313 | 418 |
| 1994 | 28 | 314 | 86 | 179 | 369 | 164 | 389 | 212 | 403 | 875 | 1013 | 1008 | 1259 | 925 | 1595 | 1025 | 1420 | 1193 | 1070 |
| 1995 | 46 | 181 | 83 | 90 | 82 | 135 | 142 | 269 | 89 | 109 | 304 | 140 | 168 | 246 | 653 | 435 | 1219 | 530 | 1011 |
| 1996 | 75 | 64 | 61 | 269 | 92 | 100 | 61 | 133 | 131 | 356 | 260 | 410 | 186 | 188 | 96 | 335 | 328 | 391 | 265 |
| 1997 | 28 | 821 | 582 | 789 | 619 | 459 | 578 | 229 | 576 | 309 | 420 | 211 | 313 | 173 | 106 | 31 | 102 | 120 | 124 |
| 1998 | 168 | 591 | 470 | 628 | 582 | 166 | 330 | 459 | 385 | 662 | 875 | 1180 | 1114 | 558 | 500 | 530 | 393 | 535 | 528 |
| 1999 | 151 | 500 | 1716 | 732 | 954 | 58 | 134 | 333 | 300 | 596 | 1566 | 1224 | 801 | 615 | 407 | 945 | 767 | 1441 | 1061 |
| 2000 | 204 | 186 | 58 | 95 | 0 | 213 | 345 | 659 | 887 | 1502 | 1331 | 984 | 630 | 349 | 421 | 1330 | 1197 | 1578 | 632 |
| 2001 | 70 | 23 | 36 | 183 | 59 | 169 | 137 | 279 | 207 | 550 | 693 | 521 | 157 | 473 | 1051 | 1627 | 1745 | 1574 | 940 |
| 2002 | 732 | 724 | 859 | 531 | 859 | 254 | 105 | 194 | 268 | 404 | 633 | 498 | 631 | 328 | 467 | 604 | 442 | 715 | 636 |
| 2003 | 32 | 202 | 470 | 640 | 972 | 872 | 673 | 858 | 1220 | 1568 | 1338 | 1205 | 1291 | 1220 | 843 | 586 | 296 | 898 | 570 |
| 2004 | 967 | 2808 | 4598 | 8840 | 9036 | 8040 | 5753 | 2798 | 1910 | 1566 | 1474 | 1883 | 1099 | 1950 | 1092 | 1508 | 1054 | 1433 | 1569 |
| 2005 | 1456 | 1538 | 1849 | 1846 | 2349 | 2219 | 3493 | 3751 | 2522 | 5835 | 6106 | 6404 | 7012 | 6700 | 4950 | 3194 | 2697 | 1851 | 2985 |
| 2006 | 111 | 302 | 364 | 694 | 1043 | 667 | 866 | 978 | 1549 | 2161 | 4462 | 5132 | 5562 | 8590 | 7349 | 5828 | 7433 | 8061 | 6573 |
| 2007 | 671 | 2556 | 2003 | 1155 | 570 | 95 | 187 | 368 | 335 | 537 | 656 | 1033 | 625 | 991 | 2013 | 2506 | 3539 | 4826 | 5279 |
| 2008 | 314 | 689 | 964 | 704 | 890 | 1109 | 1470 | 1964 | 1414 | 862 | 493 | 392 | 692 | 881 | 826 | 1247 | 1188 | 1083 | 1654 |
| 2009 | 199 | 410 | 631 | 644 | 357 | 364 | 662 | 277 | 961 | 616 | 1195 | 486 | 1157 | 1122 | 1386 | 934 | 656 | 523 | 785 |
| 2010 | 1668 | 4969 | 4829 | 4354 | 2098 | 1877 | 3207 | 3932 | 2393 | 1736 | 794 | 723 | 569 | 1440 | 1011 | 1201 | 1996 | 1068 | 1718 |
| 2011 | 385 | 512 | 1312 | 434 | 953 | 487 | 686 | 941 | 3030 | 1808 | 2669 | 2035 | 1621 | 1637 | 2059 | 750 | 509 | 485 | 711 |
| 2012 | 1142 | 1635 | 1831 | 2085 | 1452 | 1001 | 1644 | 1775 | 2176 | 1819 | 1753 | 2269 | 2086 | 3244 | 2672 | 2970 | 2021 | 1828 | 1383 |
| 2013 | 340 | 772 | 663 | 408 | 306 | 372 | 489 | 1265 | 1688 | 2434 | 3224 | 2903 | 2542 | 2144 | 2052 | 2933 | 2114 | 3047 | 3150 |
| 2014 | 1303 | 1428 | 1756 | 1738 | 1300 | 1000 | 1043 | 975 | 1529 | 1380 | 1423 | 1763 | 1528 | 1979 | 2835 | 5131 | 4430 | 5074 | 3785 |
| 2015 | 1599 | 1661 | 989 | 969 | 999 | 829 | 763 | 530 | 992 | 1062 | 1490 | 2084 | 2817 | 2310 | 1440 | 1794 | 1722 | 1572 | 2139 |
| 2016 | 669 | 1063 | 1503 | 1152 | 1515 | 2791 | 2383 | 2934 | 1994 | 2216 | 1581 | 1587 | 1760 | 1948 | 1288 | 1832 | 2186 | 2268 | 1440 |

Table 7-4. Continued.

| year/size (cm) | 34 | 35 | 36 | 37 | 38 | 39 | 40 | 41 | 42 | 43 | 44 | 45 | 46 | 47 | 48 | 49 | 50 | 51 | 52 | 53 | 54 |
|----------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|-----|------|-----|-----|
| 1991 | 2676 | 3162 | 1969 | 2161 | 1788 | 1714 | 718 | 661 | 889 | 1258 | 301 | 778 | 513 | 389 | 156 | 414 | 268 | 177 | 115 | 92 | 0 |
| 1992 | 1177 | 2682 | 1988 | 1676 | 1836 | 2425 | 1827 | 1696 | 1627 | 1491 | 938 | 2131 | 456 | 831 | 820 | 478 | 1001 | 904 | 79 | 108 | 87 |
| 1993 | 367 | 841 | 1156 | 1130 | 1031 | 985 | 1899 | 1817 | 1530 | 1428 | 1147 | 1171 | 1260 | 600 | 579 | 437 | 246 | 548 | 67 | 260 | 175 |
| 1994 | 589 | 291 | 507 | 589 | 1294 | 704 | 1373 | 1508 | 1889 | 1599 | 1117 | 1395 | 2074 | 1974 | 866 | 1186 | 688 | 879 | 167 | 597 | 259 |
| 1995 | 855 | 768 | 924 | 550 | 246 | 392 | 235 | 252 | 458 | 709 | 407 | 254 | 902 | 865 | 630 | 633 | 618 | 840 | 1068 | 326 | 297 |
| 1996 | 119 | 406 | 451 | 441 | 846 | 571 | 741 | 771 | 628 | 466 | 535 | 273 | 632 | 244 | 487 | 504 | 457 | 591 | 210 | 169 | 0 |
| 1997 | 198 | 86 | 60 | 222 | 204 | 309 | 962 | 415 | 791 | 831 | 592 | 517 | 703 | 360 | 388 | 476 | 214 | 170 | 156 | 83 | 115 |
| 1998 | 608 | 170 | 102 | 120 | 216 | 261 | 305 | 359 | 612 | 510 | 365 | 801 | 486 | 291 | 410 | 213 | 534 | 337 | 258 | 155 | 287 |
| 1999 | 441 | 406 | 226 | 140 | 264 | 124 | 203 | 47 | 250 | 80 | 149 | 260 | 390 | 466 | 122 | 340 | 239 | 0 | 213 | 181 | 56 |
| 2000 | 795 | 343 | 1100 | 695 | 847 | 704 | 490 | 664 | 214 | 387 | 204 | 185 | 181 | 203 | 259 | 221 | 192 | 68 | 51 | 120 | 233 |
| 2001 | 521 | 1418 | 906 | 1411 | 818 | 1272 | 908 | 1213 | 833 | 574 | 532 | 436 | 340 | 270 | 130 | 82 | 310 | 282 | 168 | 208 | 174 |
| 2002 | 869 | 790 | 1039 | 1255 | 431 | 277 | 606 | 589 | 525 | 563 | 206 | 282 | 468 | 308 | 602 | 16 | 301 | 199 | 109 | 0 | 420 |
| 2003 | 1016 | 465 | 527 | 592 | 1085 | 801 | 854 | 807 | 760 | 586 | 570 | 669 | 446 | 252 | 337 | 202 | 240 | 337 | 100 | 226 | 341 |
| 2004 | 1089 | 756 | 478 | 735 | 343 | 439 | 457 | 218 | 242 | 310 | 334 | 335 | 278 | 239 | 389 | 138 | 272 | 53 | 27 | 189 | 63 |
| 2005 | 2264 | 1736 | 1205 | 973 | 818 | 696 | 157 | 536 | 186 | 132 | 296 | 237 | 69 | 165 | 130 | 213 | 121 | 28 | 274 | 0 | 97 |
| 2006 | 6496 | 5580 | 4950 | 2522 | 1745 | 1192 | 1297 | 708 | 877 | 440 | 438 | 173 | 420 | 111 | 195 | 191 | 56 | 240 | 75 | 226 | 28 |
| 2007 | 5668 | 4680 | 6341 | 5592 | 4836 | 3292 | 3396 | 652 | 1550 | 351 | 632 | 661 | 131 | 389 | 64 | 240 | 133 | 519 | 36 | 176 | 0 |
| 2008 | 3380 | 4088 | 4862 | 3586 | 4822 | 4049 | 4799 | 3247 | 3200 | 1230 | 1307 | 1016 | 1059 | 116 | 544 | 132 | 189 | 129 | 153 | 341 | 156 |
| 2009 | 557 | 577 | 1152 | 2148 | 1405 | 1820 | 1549 | 2505 | 2335 | 1659 | 1147 | 911 | 720 | 451 | 150 | 226 | 0 | 51 | 56 | 0 | 0 |
| 2010 | 1558 | 1183 | 1110 | 641 | 824 | 498 | 1063 | 1597 | 2493 | 2714 | 2839 | 2082 | 2469 | 717 | 1362 | 344 | 329 | 120 | 212 | 20 | 0 |
| 2011 | 626 | 1320 | 1232 | 1047 | 529 | 840 | 798 | 575 | 653 | 820 | 1138 | 1209 | 1817 | 1288 | 1426 | 894 | 464 | 138 | 144 | 181 | 65 |
| 2012 | 953 | 1278 | 1122 | 593 | 1223 | 795 | 1040 | 253 | 701 | 454 | 951 | 479 | 1091 | 520 | 779 | 687 | 467 | 233 | 123 | 185 | 125 |
| 2013 | 3567 | 2246 | 2071 | 1544 | 1088 | 522 | 650 | 851 | 662 | 815 | 510 | 890 | 455 | 702 | 348 | 481 | 447 | 203 | 196 | 80 | 46 |
| 2014 | 3081 | 2679 | 1737 | 1810 | 2367 | 2262 | 2163 | 1581 | 550 | 771 | 421 | 1677 | 348 | 633 | 196 | 936 | 787 | 153 | 583 | 126 | 78 |
| 2015 | 1743 | 2283 | 3867 | 3393 | 3622 | 2230 | 1967 | 2214 | 1875 | 1742 | 1822 | 1141 | 949 | 460 | 603 | 471 | 317 | 226 | 357 | 269 | 203 |
| 2016 | 2068 | 1920 | 1663 | 1744 | 1768 | 1693 | 2373 | 3198 | 2109 | 1745 | 1593 | 1222 | 1349 | 661 | 681 | 393 | 374 | 583 | 470 | 185 | 108 |

Table 7-4. Continued.

| year/size (cm) | 55 | 56 | 57 | 58 | 59 | 60 | 61 | 62 | 63 | 64 | 65 | 66 | 67 | 68 | 69 | 70+ |
|----------------|-----|-----|-----|-----|-----|-----|-----|----|----|----|----|-----|----|----|----|-----|
| 1991 | 66 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1992 | 94 | 0 | 123 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1993 | 24 | 24 | 0 | 24 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1994 | 106 | 52 | 283 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1995 | 177 | 80 | 210 | 68 | 0 | 0 | 133 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1996 | 30 | 42 | 89 | 30 | 90 | 90 | 38 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1997 | 20 | 102 | 0 | 29 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 102 | 0 | 0 | 0 | 0 |
| 1998 | 51 | 28 | 37 | 0 | 74 | 0 | 64 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1999 | 93 | 0 | 0 | 64 | 0 | 28 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2000 | 208 | 17 | 21 | 0 | 0 | 26 | 0 | 26 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2001 | 338 | 0 | 42 | 69 | 27 | 0 | 0 | 27 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2002 | 256 | 76 | 0 | 27 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2003 | 195 | 114 | 157 | 130 | 41 | 32 | 29 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2004 | 75 | 239 | 76 | 20 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2005 | 325 | 25 | 0 | 0 | 65 | 159 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2006 | 56 | 27 | 47 | 0 | 75 | 0 | 19 | 28 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2007 | 57 | 118 | 85 | 28 | 0 | 50 | 36 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2008 | 29 | 256 | 29 | 109 | 0 | 35 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2009 | 84 | 55 | 0 | 0 | 134 | 0 | 55 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2010 | 78 | 69 | 103 | 99 | 0 | 0 | 0</ | | | | | | | | | |

Table 7-5. Bering Sea slope survey female size composition estimates (1,000s of fish).

| year/size (cm) | 21 | 22 | 23 | 24 | 25 | 26 | 27 | 28 | 29 | 30 | 31 | 32 | 33 | 34 | 35 | 36 | 37 | 38 | 39 | 40 |
|----------------|----|----|----|----|----|----|----|----|----|----|----|----|----|-----|-----|-----|-----|-----|-----|-----|
| 2002 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 5 | 0 | 0 | 0 | 6 | 19 | 37 | 35 | 12 | 83 | 78 |
| 2004 | 5 | 0 | 9 | 4 | 3 | 12 | 15 | 5 | 29 | 30 | 36 | 45 | 34 | 35 | 30 | 80 | 48 | 68 | 60 | 64 |
| 2008 | 0 | 8 | 0 | 0 | 0 | 4 | 0 | 4 | 0 | 4 | 0 | 15 | 13 | 71 | 102 | 118 | 199 | 216 | 202 | 412 |
| 2010 | 0 | 0 | 0 | 5 | 0 | 4 | 4 | 0 | 0 | 0 | 4 | 11 | 0 | 21 | 21 | 14 | 21 | 54 | 40 | 144 |
| 2012 | 0 | 0 | 10 | 0 | 0 | 7 | 24 | 0 | 15 | 38 | 11 | 5 | 5 | 0 | 0 | 8 | 4 | 39 | 29 | 5 |
| 2016 | 0 | 0 | 0 | 41 | 52 | 14 | 12 | 42 | 13 | 27 | 30 | 88 | 57 | 104 | 99 | 158 | 169 | 236 | 146 | 247 |

Table 7-5. Continued.

| year/size (cm) | 41 | 42 | 43 | 44 | 45 | 46 | 47 | 48 | 49 | 50 | 51 | 52 | 53 | 54 | 55 | 56 | 57 | 58 | 59 | 60 |
|----------------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| 2002 | 78 | 105 | 111 | 194 | 196 | 152 | 213 | 122 | 152 | 145 | 81 | 58 | 58 | 172 | 158 | 141 | 120 | 153 | 196 | 199 |
| 2004 | 47 | 55 | 69 | 93 | 92 | 75 | 83 | 73 | 103 | 100 | 80 | 68 | 97 | 124 | 100 | 103 | 91 | 68 | 78 | 77 |
| 2008 | 454 | 497 | 473 | 337 | 307 | 284 | 279 | 191 | 189 | 191 | 145 | 194 | 118 | 144 | 154 | 98 | 169 | 127 | 178 | 94 |
| 2010 | 168 | 249 | 314 | 395 | 428 | 790 | 698 | 846 | 598 | 461 | 498 | 393 | 352 | 310 | 313 | 152 | 178 | 213 | 117 | 170 |
| 2012 | 16 | 0 | 55 | 93 | 0 | 16 | 32 | 66 | 64 | 140 | 59 | 294 | 32 | 64 | 122 | 209 | 70 | 79 | 18 | 23 |
| 2016 | 278 | 295 | 200 | 336 | 296 | 419 | 288 | 368 | 233 | 270 | 263 | 203 | 290 | 311 | 414 | 468 | 614 | 402 | 468 | 545 |

Table 7-5. Continued.

| year/size (cm) | 61 | 62 | 63 | 64 | 65 | 66 | 67 | 68 | 69 | 70 | 71 | 72 | 73 | 74 | 75 | 76 | 77 | 78 | 79 | 80 |
|----------------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|----|----|----|----|-----|
| 2002 | 163 | 164 | 153 | 155 | 201 | 159 | 145 | 177 | 98 | 159 | 96 | 156 | 91 | 67 | 59 | 39 | 26 | 52 | 38 | 12 |
| 2004 | 74 | 64 | 66 | 92 | 88 | 102 | 81 | 106 | 79 | 56 | 70 | 43 | 54 | 55 | 50 | 40 | 49 | 33 | 23 | 47 |
| 2008 | 122 | 92 | 110 | 65 | 97 | 126 | 79 | 130 | 107 | 103 | 124 | 90 | 74 | 72 | 63 | 61 | 69 | 28 | 52 | 16 |
| 2010 | 118 | 92 | 95 | 118 | 58 | 90 | 81 | 103 | 67 | 104 | 82 | 34 | 52 | 31 | 59 | 51 | 48 | 57 | 33 | 34 |
| 2012 | 22 | 10 | 12 | 19 | 10 | 9 | 4 | 13 | 0 | 45 | 9 | 0 | 9 | 33 | 14 | 11 | 0 | 0 | 0 | 4 |
| 2016 | 446 | 426 | 236 | 221 | 135 | 225 | 204 | 131 | 140 | 107 | 82 | 106 | 132 | 123 | 133 | 95 | 95 | 65 | 73 | 115 |

Table 7-5. Bering Sea slope survey male size composition estimates (1,000s of fish).

| year/size (cm) | Male size composition estimates from the slope surveys (1,000s of fish) | | | | | | | | | | | | | | | | | | | |
|----------------|---|----|----|----|----|----|----|----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|------|
| | 21 | 22 | 23 | 24 | 25 | 26 | 27 | 28 | 29 | 30 | 31 | 32 | 33 | 34 | 35 | 36 | 37 | 38 | 39 | 40 |
| 2002 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 6 | 38 | 42 | 68 | 51 | 77 | 186 | 169 |
| 2004 | 4 | 0 | 3 | 16 | 12 | 33 | 21 | 58 | 72 | 109 | 120 | 114 | 85 | 138 | 135 | 181 | 228 | 144 | 112 | 92 |
| 2008 | 4 | 0 | 0 | 0 | 0 | 0 | 4 | 0 | 16 | 25 | 31 | 68 | 115 | 220 | 314 | 333 | 641 | 619 | 767 | 1001 |
| 2010 | 6 | 0 | 0 | 4 | 4 | 0 | 5 | 16 | 10 | 20 | 18 | 57 | 36 | 64 | 58 | 87 | 130 | 190 | 214 | 390 |
| 2012 | 0 | 0 | 5 | 0 | 14 | 20 | 5 | 0 | 39 | 52 | 41 | 26 | 31 | 5 | 26 | 76 | 0 | 221 | 21 | 31 |
| 2016 | 14 | 0 | 14 | 14 | 0 | 55 | 36 | 71 | 138 | 99 | 76 | 307 | 158 | 221 | 247 | 379 | 779 | 612 | 809 | 949 |

Table 7-5. Continued.

| year/size (cm) | 41 | 42 | 43 | 44 | 45 | 46 | 47 | 48 | 49 | 50 | 51 | 52 | 53 | 54 | 55 | 56 | 57 | 58 | 59 | 60 |
|----------------|------|------|------|------|------|------|-----|-----|------|-----|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| 2002 | 269 | 295 | 285 | 230 | 216 | 155 | 133 | 125 | 299 | 222 | 246 | 350 | 290 | 305 | 323 | 238 | 143 | 174 | 78 | 72 |
| 2004 | 158 | 229 | 196 | 229 | 217 | 270 | 224 | 372 | 363 | 352 | 403 | 278 | 318 | 283 | 267 | 231 | 164 | 101 | 81 | 87 |
| 2008 | 1171 | 1100 | 767 | 506 | 539 | 451 | 537 | 342 | 473 | 487 | 524 | 384 | 468 | 234 | 285 | 198 | 220 | 211 | 119 | 87 |
| 2010 | 576 | 805 | 1130 | 1058 | 1182 | 1083 | 817 | 679 | 479 | 539 | 330 | 339 | 283 | 291 | 224 | 319 | 195 | 257 | 130 | 123 |
| 2012 | 10 | 14 | 39 | 94 | 0 | 49 | 139 | 78 | 92 | 85 | 9 | 46 | 18 | 9 | 16 | 158 | 27 | 37 | 26 | 0 |
| 2016 | 667 | 768 | 872 | 853 | 920 | 761 | 527 | 986 | 1072 | 897 | 1065 | 638 | 636 | 759 | 446 | 500 | 463 | 369 | 273 | 300 |

Table 7-5. Continued.

| year/size (cm) | 61 | 62 | 63 | 64 | 65 | 66 | 67 | 68 | 69 | 70 | 71 | 72 | 73 | 74 | 75 | 76 | 77 | 78 | 79 | 80 |
|----------------|-----|-----|-----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|
| 2002 | 66 | 27 | 0 | 12 | 21 | 0 | 0 | 6 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2004 | 28 | 27 | 7 | 8 | 3 | 6 | 4 | 3 | 0 | 0 | 3 | 0 | 0 | 0 | 0 | 5 | 0 | 0 | 0 | 0 |
| 2008 | 59 | 62 | 30 | 7 | 13 | 4 | 0 | 0 | 0 | 0 | 0 | 4 | 4 | 4 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2010 | 101 | 75 | 39 | 18 | 9 | 12 | 19 | 5 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2012 | 13 | 10 | 4 | 0 | 0 | 0 | 4 | 7 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2016 | 138 | 181 | 108 | 76 | 7 | 7 | 9 | 5 | 9 | 0 | 0 | 16 | 0 | 0 | 0 | 16 | 0 | 0 | 0 | 0 |

Table 7-6. Aleutian Islands survey female size composition estimates (1,000s of fish).

| year/size (cm) | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22 | 23 | 24 | 25 | 26 | 27 | 28 | 29 | 30 | 31 | 32 | 33 |
|----------------|-----|-----|-----|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
| 1991 | 17 | 75 | 242 | 330 | 393 | 308 | 436 | 576 | 628 | 363 | 430 | 325 | 511 | 581 | 631 | 603 | 746 | 719 | 662 |
| 1994 | 18 | 111 | 287 | 330 | 530 | 539 | 619 | 616 | 652 | 803 | 627 | 809 | 653 | 766 | 890 | 895 | 1025 | 1098 | 1137 |
| 1997 | 157 | 295 | 630 | 868 | 1068 | 1099 | 924 | 1177 | 986 | 963 | 1033 | 1093 | 1523 | 2045 | 1550 | 2611 | 1530 | 1477 | 1855 |
| 2000 | 112 | 273 | 608 | 1229 | 1821 | 1281 | 1140 | 1199 | 1362 | 1203 | 2140 | 2515 | 2166 | 3029 | 2782 | 2198 | 2004 | 2220 | 1464 |
| 2002 | 41 | 162 | 483 | 728 | 833 | 1067 | 681 | 1476 | 1672 | 2264 | 2482 | 2570 | 2140 | 2201 | 2213 | 2059 | 2282 | 2847 | 2698 |
| 2004 | 254 | 258 | 264 | 854 | 907 | 924 | 791 | 789 | 973 | 1072 | 1367 | 1166 | 1264 | 1670 | 1661 | 1700 | 1825 | 2124 | 1910 |
| 2006 | 40 | 159 | 182 | 658 | 1045 | 1340 | 1037 | 972 | 847 | 1416 | 1146 | 1716 | 1558 | 1442 | 1298 | 1542 | 1117 | 1308 | 1615 |
| 2010 | 34 | 96 | 320 | 673 | 714 | 745 | 716 | 920 | 959 | 1044 | 1015 | 1057 | 1499 | 1441 | 1494 | 1582 | 1619 | 1417 | 1786 |
| 2012 | 8 | 14 | 71 | 92 | 116 | 138 | 95 | 198 | 118 | 180 | 177 | 300 | 278 | 259 | 182 | 319 | 298 | 230 | 224 |
| 2014 | 256 | 500 | 634 | 1123 | 1546 | 1464 | 1321 | 1333 | 1127 | 1258 | 1495 | 1484 | 1688 | 1777 | 1812 | 2032 | 1856 | 1710 | 2054 |
| 2016 | 26 | 69 | 179 | 87 | 312 | 284 | 368 | 579 | 375 | 301 | 169 | 153 | 163 | 208 | 190 | 189 | 93 | 149 | 344 |

Table 7-6. Continued.

| year/size (cm) | 34 | 35 | 36 | 37 | 38 | 39 | 40 | 41 | 42 | 43 | 44 | 45 | 46 | 47 | 48 | 49 | 50 | 51 | 52 | 53 | 54 |
|----------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
| 1991 | 724 | 767 | 830 | 773 | 890 | 918 | 847 | 784 | 503 | 527 | 452 | 605 | 398 | 584 | 390 | 498 | 551 | 406 | 262 | 250 | 245 |
| 1994 | 1163 | 1327 | 1296 | 1255 | 1228 | 1358 | 1331 | 1567 | 1452 | 1365 | 1528 | 1414 | 1763 | 1492 | 1676 | 1154 | 1170 | 1220 | 1216 | 1195 | 980 |
| 1997 | 1707 | 1508 | 1496 | 1200 | 1383 | 1321 | 1395 | 1296 | 1340 | 1384 | 1583 | 1579 | 1661 | 1356 | 1348 | 1084 | 1060 | 1545 | 1312 | 1121 | 1358 |
| 2000 | 1822 | 1234 | 1479 | 1182 | 1618 | 1675 | 1383 | 1860 | 1424 | 1342 | 1516 | 1155 | 979 | 1055 | 962 | 702 | 771 | 916 | 821 | 1087 | 804 |
| 2002 | 2777 | 2356 | 2430 | 1944 | 2464 | 1900 | 1945 | 2152 | 1587 | 1587 | 1607 | 1586 | 1558 | 1265 | 1478 | 1183 | 1415 | 1014 | 1287 | 1072 | 1240 |
| 2004 | 1866 | 1790 | 2105 | 2266 | 2327 | 2784 | 2267 | 2773 | 2317 | 2070 | 2371 | 2140 | 2120 | 1634 | 1505 | 1532 | 1334 | 1384 | 1157 | 1528 | 1444 |
| 2006 | 1476 | 1512 | 1631 | 1875 | 2046 | 2791 | 2248 | 2263 | 2132 | 2595 | 2491 | 3067 | 3492 | 2900 | 3421 | 4398 | 5088 | 3884 | 3293 | 3524 | 2920 |
| 2010 | 1650 | 2091 | 1801 | 1808 | 1591 | 1807 | 1618 | 1640 | 1322 | 1280 | 1007 | 1242 | 754 | 974 | 1293 | 1160 | 1494 | 1355 | 1500 | 1502 | 1579 |
| 2012 | 261 | 227 | 189 | 237 | 141 | 146 | 119 | 170 | 89 | 96 | 62 | 82 | 49 | 105 | 36 | 74 | 89 | 87 | 139 | 87 | 110 |
| 2014 | 1778 | 1528 | 1628 | 1588 | 1365 | 1565 | 1286 | 1368 | 1193 | 1456 | 1347 | 1288 | 1332 | 1431 | 1345 | 995 | 1203 | 1116 | 1386 | 1382 | 1710 |
| 2016 | 85 | 155 | 101 | 77 | 81 | 107 | 103 | 98 | 98 | 151 | 138 | 59 | 67 | 90 | 30 | 148 | 106 | 42 | 57 | 112 | 26 |

Table 7-6. Continued.

| year/size (cm) | 55 | 56 | 57 | 58 | 59 | 60 | 61 | 62 | 63 | 64 | 65 | 66 | 67 | 68 | 69 | 70+ |
|----------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|-------|
| 1991 | 120 | 198 | 194 | 168 | 167 | 160 | 187 | 193 | 129 | 171 | 175 | 154 | 72 | 102 | 80 | 1239 |
| 1994 | 1066 | 1018 | 1029 | 1095 | 1211 | 1038 | 992 | 1108 | 603 | 605 | 577 | 500 | 416 | 360 | 238 | 1844 |
| 1997 | 1500 | 1284 | 1002 | 1149 | 973 | 1024 | 936 | 833 | 914 | 780 | 766 | 515 | 543 | 640 | 445 | 2384 |
| 2000 | 1157 | 1025 | 1006 | 708 | 876 | 807 | 567 | 616 | 566 | 634 | 487 | 450 | 398 | 369 | 338 | 3726 |
| 2002 | 1333 | 1180 | 1526 | 1341 | 1221 | 1512 | 930 | 1102 | 1218 | 1228 | 1334 | 1017 | 956 | 982 | 1149 | 5780 |
| 2004 | 1198 | 1142 | 1496 | 1041 | 1518 | 1286 | 1198 | 1131 | 775 | 937 | 787 | 800 | 787 | 660 | 493 | 4314 |
| 2006 | 3558 | 3470 | 3384 | 4034 | 2788 | 3106 | 3064 | 3064 | 3245 | 2451 | 2740 | 1590 | 1796 | 1288 | 1005 | 5896 |
| 2010 | 1094 | 1374 | 914 | 844 | 690 | 982 | 769 | 752 | 707 | 677 | 420 | 756 | 527 | 554 | 415 | 4188 |
| 2012 | 153 | 108 | 158 | 72 | 174 | 120 | 36 | 92 | 79 | 82 | 113 | 61 | 54 | 88 | 74 | 1416 |
| 2014 | 1309 | 1326 | 1441 | 2032 | 1770 | 2063 | 1554 | 2085 | 1691 | 2053 | 1930 | 1946 | 1634 | 1861 | 1503 | 14208 |
| 2016 | 97 | 29 | 62 | 200 | 114 | 137 | 123 | 93 | 149 | 34 | 174 | 78 | 26 | 61 | 77 | 925 |

Table 7-6. Aleutian Islands survey male size composition estimates (1,000s of fish).

| year/size (cm) | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22 | 23 | 24 | 25 | 26 | 27 | 28 | 29 | 30 | 31 | 32 | 33 |
|----------------|-----|-----|-----|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
| 1991 | 95 | 115 | 182 | 237 | 207 | 336 | 298 | 297 | 312 | 188 | 346 | 561 | 508 | 422 | 640 | 585 | 1092 | 785 | 1097 |
| 1994 | 82 | 181 | 357 | 548 | 423 | 533 | 493 | 477 | 428 | 464 | 527 | 468 | 488 | 737 | 824 | 1010 | 1197 | 1195 | 1247 |
| 1997 | 205 | 589 | 792 | 894 | 1045 | 1120 | 907 | 536 | 893 | 1231 | 1130 | 1633 | 1625 | 1344 | 1295 | 1279 | 1419 | 1358 | 1665 |
| 2000 | 165 | 302 | 467 | 1454 | 1345 | 1090 | 1107 | 910 | 1015 | 1377 | 1853 | 2022 | 2346 | 2575 | 2244 | 2581 | 2413 | 1853 | 1824 |
| 2002 | 91 | 316 | 617 | 626 | 535 | 516 | 675 | 612 | 999 | 1432 | 1836 | 1475 | 1344 | 1071 | 1761 | 1869 | 2211 | 1760 | 1523 |
| 2004 | 73 | 205 | 510 | 880 | 1171 | 630 | 851 | 838 | 790 | 889 | 969 | 1105 | 1142 | 1321 | 1389 | 1368 | 1722 | 1669 | 1226 |
| 2006 | 96 | 379 | 738 | 995 | 958 | 658 | 623 | 911 | 1018 | 1120 | 1278 | 1255 | 1477 | 1441 | 1150 | 1389 | 891 | 1080 | 995 |
| 2010 | 22 | 165 | 249 | 893 | 882 | 1185 | 867 | 754 | 705 | 727 | 1067 | 1317 | 1590 | 1831 | 1830 | 2271 | 2302 | 2481 | 2681 |
| 2012 | 13 | 20 | 3 | 58 | 144 | 153 | 161 | 170 | 226 | 260 | 183 | 194 | 168 | 327 | 473 | 402 | 430 | 338 | 321 |
| 2014 | 379 | 640 | 957 | 1758 | 1726 | 2050 | 1567 | 1514 | 1507 | 1697 | 1576 | 1770 | 2257 | 2526 | 2793 | 3069 | 3098 | 2602 | 2489 |
| 2016 | 9 | 99 | 177 | 241 | 557 | 428 | 800 | 605 | 492 | 381 | 321 | 252 | 361 | 287 | 235 | 302 | 320 | 396 | 240 |

Table 7-6. Continued.

| year/size (cm) | 34 | 35 | 36 | 37 | 38 | 39 | 40 | 41 | 42 | 43 | 44 | 45 | 46 | 47 | 48 | 49 | 50 | 51 | 52 | 53 | 54 |
|----------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
| 1991 | 1054 | 1021 | 731 | 928 | 837 | 577 | 731 | 401 | 436 | 304 | 346 | 369 | 159 | 235 | 133 | 184 | 157 | 204 | 186 | 154 | 175 |
| 1994 | 1350 | 1165 | 1560 | 1524 | 1419 | 1429 | 1751 | 1799 | 1611 | 1492 | 1966 | 1575 | 1138 | 1177 | 1066 | 871 | 987 | 723 | 1024 | 1174 | 1513 |
| 1997 | 1136 | 1248 | 934 | 1341 | 1134 | 1293 | 1299 | 1250 | 1676 | 1389 | 1975 | 1790 | 1728 | 1549 | 1217 | 1378 | 953 | 819 | 827 | 636 | 872 |
| 2000 | 1967 | 1664 | 1729 | 1850 | 1987 | 1489 | 1190 | 1010 | 1179 | 1013 | 926 | 1302 | 1551 | 1298 | 1354 | 893 | 802 | 608 | 609 | 635 | 463 |
| 2002 | 1308 | 1463 | 1800 | 1832 | 1560 | 1617 | 1415 | 1464 | 1173 | 1767 | 1912 | 1704 | 1635 | 1186 | 1172 | 1229 | 1233 | 1043 | 855 | 667 | 447 |
| 2004 | 1180 | 1332 | 1432 | 1404 | 1578 | 1464 | 1453 | 1559 | 1849 | 1506 | 1508 | 1680 | 1150 | 1208 | 1311 | 1182 | 1128 | 1092 | 806 | 684 | 773 |
| 2006 | 1223 | 1219 | 1218 | 1490 | 1069 | 1374 | 1858 | 2562 | 3559 | 3416 | 3518 | 3212 | 2615 | 2952 | 2233 | 1913 | 1230 | 1416 | 1340 | 738 | 1092 |
| 2010 | 2051 | 3100 | 2751 | 2615 | 2442 | 1974 | 2010 | 3485 | 2277 | 2516 | 2244 | 2400 | 2041 | 1448 | 1083 | 1081 | 818 | 718 | 749 | 618 | 869 |
| 2012 | 341 | 370 | 350 | 309 | 362 | 231 | 177 | 115 | 161 | 130 | 128 | 101 | 181 | 175 | 227 | 276 | 272 | 404 | 278 | 171 | 247 |
| 2014 | 2447 | 2525 | 2407 | 2855 | 2463 | 2324 | 2539 | 2655 | 2629 | 2729 | 2561 | 3010 | 2570 | 2905 | 3006 | 3490 | 3642 | 4016 | 4272 | 4063 | 5569 |
| 2016 | 339 | 221 | 237 | 317 | 150 | 265 | 192 | 180 | 211 | 247 | 358 | 231 | 210 | 213 | 142 | 62 | 46 | 223 | 67 | 99 | 266 |

Table 7-6. Continued.

| year/size (cm) | 55 | 56 | 57 | 58 | 59 | 60 | 61 | 62 | 63 | 64 | 65 | 66 | 67 | 68 | 69 | 70+ |
|----------------|------|------|------|------|------|------|------|------|------|------|------|------|------|-----|-----|-----|
| 1991 | 100 | 169 | 122 | 183 | 111 | 145 | 71 | 83 | 94 | 43 | 7 | 4 | 15 | 4 | 0 | 5 |
| 1994 | 1488 | 1278 | 1745 | 1030 | 1037 | 729 | 703 | 552 | 327 | 122 | 51 | 96 | 3 | 11 | 0 | 14 |
| 1997 | 908 | 873 | 699 | 788 | 675 | 789 | 563 | 402 | 251 | 225 | 175 | 78 | 30 | 14 | 16 | 4 |
| 2000 | 579 | 467 | 601 | 566 | 586 | 626 | 536 | 395 | 372 | 200 | 242 | 86 | 23 | 69 | 65 | 338 |
| 2002 | 416 | 416 | 438 | 336 | 462 | 451 | 451 | 473 | 384 | 501 | 498 | 132 | 157 | 101 | 55 | 521 |
| 2004 | 875 | 726 | 790 | 757 | 917 | 797 | 1009 | 660 | 672 | 423 | 282 | 186 | 126 | 81 | 85 | 323 |
| 2006 | 750 | 597 | 804 | 818 | 882 | 834 | 794 | 614 | 728 | 524 | 515 | 198 | 210 | 116 | 105 | 951 |
| 2010 | 498 | 816 | 1052 | 1109 | 911 | 1179 | 1554 | 1354 | 1031 | 974 | 552 | 540 | 216 | 302 | 51 | 491 |
| 2012 | 471 | 453 | 298 | 293 | 537 | 572 | 563 | 804 | 599 | 774 | 521 | 359 | 259 | 226 | 109 | 31 |
| 2014 | 5307 | 5252 | 6169 | 5641 | 6177 | 6546 | 6501 | 5541 | 4950 | 4324 | 2840 | 1932 | 1152 | 868 | 547 | 163 |
| 2016 | 307 | 318 | 182 | 239 | 289 | 395 | 274 | 395 | 344 | 477 | 474 | 392 | 296 | 264 | 184 | 245 |

Table 7.7. Estimated maturity at age for female Kamchatka flounder (Stark 2011).

| age | proportion mature |
|-----|----------------------|
| 2 | 0.00 |
| 3 | 0.01 |
| 4 | 0.01 |
| 5 | 0.02 |
| 6 | 0.05 |
| 7 | 0.10 |
| 8 | 0.18 |
| 9 | 0.31 |
| 10 | 0.48 |
| 11 | 0.66 |
| 12 | 0.80 |
| 13 | 0.89 |
| 14 | 0.94 |
| 15 | 0.97 |
| 16 | 0.99 |
| 17 | 0.99 |
| 18 | 1.00 |
| 19 | 1.00 |
| 20 | 1.00 |
| 21 | 1.00 |
| 22 | 1.00 |
| 23 | 1.00 |
| 24 | 1.00 |
| 25 | 1.00 |

Table 7-8. Key equations used in the population dynamics model.

| | |
|---|--|
| $N_{t,1} = R_t = R_0 e^{\tau}, \tau_t \sim N(0, \delta_R^2)$ | Recruitment $t=1969-1990$ |
| $N_{t,1} = R_t = R_y e^{\tau}, \tau_t \sim N(0, \delta_R^2)$ | Recruitment $t=1991-2012$ |
| $C_{t,a} = \frac{F_{t,a}}{Z_{t,a}} (1 - e^{-z_{t,a}}) N_{t,a}$ | Catch in year t for age a fish |
| $N_{t+1,a+1} = N_{t,a} e^{-z_{t,a}}$ | Numbers of fish in year $t+1$ at age a |
| $N_{t+1,A} = N_{t,A-1} e^{-z_{t,A-1}} + N_{t,A} e^{-z_{t,A}}$ | Numbers of fish in the “plus group” |
| $S_t = \sum N_{t,a} W_{t,a} \phi_a$ | Spawning biomass |
| $Z_{t,a} = F_{t,a} + M$ | Total mortality in year t at age a |
| $F_{t,a} = s_a \mu^F \exp^{\varepsilon^F_t}, \varepsilon^F_t \sim N(0, \sigma^{2F})$ | Fishing mortality |
| $s_a = \frac{1}{1 + (e^{-\alpha + \beta a})}$ | Age-specific fishing selectivity |
| $C_t = \sum C_{t,a}$ | Total catch in numbers |
| $P_{t,a} = C_{t,a} / C_t$ | Proportion at age in catch |
| $SurB_t = q \sum N_{t,a} W_{t,a} v_a$ | Survey biomass |
| $reclike = \lambda \left(\sum_{i=1965}^{endyear} \bar{R} - R_i \right)^2 + \sum_{a=1}^{20} \left(\bar{R}_{init} - R_{init,a} \right)^2$ | Recruitment likelihood |
| $catchlike = \lambda \sum_{i=startyear}^{endyear} (\ln C_{obs,i} - \ln C_{est,i})^2$ | catch likelihood |
| $surveylike = \lambda \frac{(\ln B - \ln \hat{B})^2}{2\sigma^2}$ | survey biomass likelihood |
| $SurvAgelike = \sum_{t,a} n_t P_{t,a} (\ln \hat{P}_{t,a} + 0.001) - \sum_{t,a} n_t P_{t,a} (\ln P_{t,a} + 0.001)$ | survey age comp likelihood |
| $SurvLengthlike = \sum_{t,a} n_t P_{t,a} (\ln \hat{P}_{t,a} + 0.001) - \sum_{t,a} n_t P_{t,a} (\ln P_{t,a} + 0.001)$ | survey length comp likelihood |

Table 7-9. Variables used in the population dynamics model.

| Variables | |
|-------------------|---|
| R_t | Age 1 recruitment in year t |
| R_0 | Geometric mean value of age 1 recruitment, 1956-75 |
| R_γ | Geometric mean value of age 1 recruitment, 1976-96 |
| τ_t | Recruitment deviation in year t |
| $N_{t,a}$ | Number of fish in year t at age a |
| $C_{t,a}$ | Catch numbers of fish in year t at age a |
| $P_{t,a}$ | Proportion of the numbers of fish age a in year t |
| C_t | Total catch numbers in year t |
| $W_{t,a}$ | Mean body weight (kg) of fish age a in year t |
| ϕ_a | Proportion of mature females at age a |
| $F_{t,a}$ | Instantaneous annual fishing mortality of age a fish in year t |
| M | Instantaneous natural mortality, assumed constant over all ages and years |
| $Z_{t,a}$ | Instantaneous total mortality for age a fish in year t |
| s_a | Age-specific fishing gear selectivity |
| μ^F | Median year-effect of fishing mortality |
| ε_t^F | The residual year-effect of fishing mortality |
| v_a | Age-specific survey selectivity |
| α | Slope parameter in the logistic selectivity equation |
| β | Age at 50% selectivity parameter in the logistic selectivity equation |
| σ | Standard error of the survey biomass in year t |

Table 7.10. Estimated total biomass (ages 2+), female spawning biomass, and recruitment (age 2 fish).

| | female spawning biomass | | | total biomass | | | age 2 recruitment |
|------|-------------------------|-------------|-------------|---------------|-------------|-------------|-------------------|
| | estimate | lower bound | upper bound | estimate | lower bound | upper bound | |
| 1991 | 21,330 | 17,105 | 25,555 | 67,314 | 59,492 | 75,136 | 10313 |
| 1992 | 23,004 | 18,905 | 27,103 | 71,923 | 64,038 | 79,808 | 14951 |
| 1993 | 25,641 | 21,601 | 29,681 | 77,444 | 69,463 | 85,425 | 9143 |
| 1994 | 28,901 | 24,869 | 32,933 | 82,610 | 74,508 | 90,712 | 6067 |
| 1995 | 32,459 | 28,389 | 36,529 | 86,711 | 78,478 | 94,944 | 8784 |
| 1996 | 36,466 | 32,303 | 40,629 | 90,890 | 82,531 | 99,249 | 13889 |
| 1997 | 40,024 | 35,746 | 44,302 | 94,286 | 85,798 | 102,774 | 22026 |
| 1998 | 43,284 | 38,887 | 47,681 | 98,068 | 89,453 | 106,683 | 19512 |
| 1999 | 45,657 | 41,161 | 50,153 | 101,370 | 92,620 | 110,120 | 20540 |
| 2000 | 47,713 | 43,141 | 52,285 | 105,070 | 96,181 | 113,959 | 11760 |
| 2001 | 49,213 | 44,587 | 53,839 | 108,570 | 99,534 | 117,606 | 16018 |
| 2002 | 50,362 | 45,701 | 55,023 | 112,400 | 103,186 | 121,614 | 30305 |
| 2003 | 51,545 | 46,855 | 56,235 | 117,370 | 107,930 | 126,810 | 44360 |
| 2004 | 52,818 | 48,093 | 57,543 | 124,050 | 114,282 | 133,818 | 67651 |
| 2005 | 54,167 | 49,386 | 58,948 | 131,100 | 120,939 | 141,261 | 22602 |
| 2006 | 56,113 | 51,243 | 60,983 | 139,310 | 128,689 | 149,931 | 13168 |
| 2007 | 58,367 | 53,385 | 63,349 | 147,890 | 136,762 | 159,018 | 18040 |
| 2008 | 61,025 | 55,908 | 66,142 | 156,570 | 144,912 | 168,228 | 19434 |
| 2009 | 61,684 | 56,411 | 66,957 | 159,060 | 146,878 | 171,242 | 21500 |
| 2010 | 60,470 | 54,999 | 65,941 | 155,800 | 143,077 | 168,523 | 57972 |
| 2011 | 56,029 | 50,306 | 61,752 | 143,640 | 130,416 | 156,864 | 33685 |
| 2012 | 56,550 | 50,457 | 62,643 | 143,460 | 129,613 | 157,307 | 41843 |
| 2013 | 56,913 | 50,425 | 63,401 | 143,720 | 129,220 | 158,220 | 20135 |
| 2014 | 57,596 | 50,743 | 64,449 | 146,450 | 131,232 | 161,668 | 32076 |
| 2015 | 58,485 | 51,312 | 65,658 | 151,390 | 135,328 | 167,452 | 50341 |
| 2016 | 60,038 | 52,570 | 67,506 | 159,030 | 141,920 | 176,140 | 62092 |

Table 7.11. Estimated numbers at age (thousands) from the stock assessment model for ages 2-24.

| | females | | | | | | | | | | | |
|------|---------|-------|-------|-------|-------|-------|-------|-------|------|------|------|------|
| | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 |
| 1991 | 5157 | 4523 | 5744 | 10233 | 10172 | 5524 | 3487 | 3361 | 2760 | 820 | 2121 | 1454 |
| 1992 | 7476 | 4611 | 4033 | 5092 | 9000 | 8882 | 4802 | 3025 | 2914 | 2392 | 711 | 1838 |
| 1993 | 4571 | 6691 | 4122 | 3597 | 4526 | 7975 | 7856 | 4244 | 2673 | 2574 | 2113 | 628 |
| 1994 | 3033 | 4093 | 5984 | 3679 | 3201 | 4018 | 7070 | 6959 | 3758 | 2367 | 2279 | 1871 |
| 1995 | 4392 | 2715 | 3657 | 5333 | 3265 | 2831 | 3546 | 6232 | 6132 | 3311 | 2085 | 2008 |
| 1996 | 6944 | 3932 | 2428 | 3266 | 4750 | 2902 | 2513 | 3145 | 5526 | 5437 | 2936 | 1848 |
| 1997 | 11013 | 6215 | 3515 | 2165 | 2900 | 4204 | 2563 | 2217 | 2774 | 4874 | 4795 | 2589 |
| 1998 | 9756 | 9860 | 5559 | 3138 | 1928 | 2577 | 3731 | 2273 | 1966 | 2460 | 4321 | 4251 |
| 1999 | 10270 | 8732 | 8813 | 4956 | 2787 | 1707 | 2277 | 3293 | 2006 | 1734 | 2169 | 3811 |
| 2000 | 5880 | 9195 | 7811 | 7869 | 4414 | 2477 | 1514 | 2019 | 2919 | 1778 | 1537 | 1923 |
| 2001 | 8009 | 5264 | 8222 | 6969 | 6999 | 3915 | 2193 | 1340 | 1786 | 2582 | 1572 | 1359 |
| 2002 | 15152 | 7169 | 4706 | 7334 | 6196 | 6205 | 3465 | 1939 | 1184 | 1578 | 2282 | 1390 |
| 2003 | 22180 | 13566 | 6412 | 4202 | 6531 | 5504 | 5504 | 3071 | 1719 | 1050 | 1399 | 2022 |
| 2004 | 33826 | 19856 | 12132 | 5723 | 3739 | 5797 | 4879 | 4875 | 2720 | 1522 | 929 | 1238 |
| 2005 | 11301 | 30274 | 17747 | 10814 | 5082 | 3309 | 5119 | 4304 | 4299 | 2398 | 1342 | 819 |
| 2006 | 6584 | 10117 | 27075 | 15840 | 9624 | 4511 | 2933 | 4534 | 3811 | 3806 | 2123 | 1188 |
| 2007 | 9020 | 5894 | 9049 | 24171 | 14104 | 8549 | 4002 | 2600 | 4018 | 3377 | 3373 | 1881 |
| 2008 | 9717 | 8076 | 5274 | 8083 | 21545 | 12547 | 7596 | 3554 | 2308 | 3567 | 2998 | 2994 |
| 2009 | 10750 | 8680 | 7184 | 4652 | 7047 | 18584 | 10753 | 6491 | 3033 | 1969 | 3043 | 2557 |
| 2010 | 28986 | 9580 | 7676 | 6257 | 3965 | 5890 | 15349 | 8833 | 5321 | 2484 | 1612 | 2491 |
| 2011 | 16843 | 25735 | 8394 | 6552 | 5145 | 3152 | 4589 | 11848 | 6793 | 4086 | 1907 | 1237 |
| 2012 | 20922 | 15020 | 22799 | 7339 | 5623 | 4342 | 2633 | 3815 | 9832 | 5633 | 3387 | 1580 |
| 2013 | 10067 | 18660 | 13311 | 19946 | 6305 | 4751 | 3633 | 2193 | 3172 | 8168 | 4678 | 2813 |
| 2014 | 16038 | 8986 | 16569 | 11695 | 17266 | 5384 | 4025 | 3066 | 1848 | 2671 | 6877 | 3939 |
| 2015 | 25171 | 14327 | 7994 | 14617 | 10196 | 14893 | 4615 | 3439 | 2617 | 1577 | 2278 | 5866 |
| 2016 | 31046 | 22500 | 12765 | 7077 | 12824 | 8872 | 12896 | 3987 | 2968 | 2258 | 1360 | 1965 |

Table 7.11. (continued).

| | females continued | | | | | | | | | | | | | | |
|------|-------------------|------|------|------|------|------|------|------|------|------|------|------|--|--|--|
| | 14 | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22 | 23 | 24 | 25 | | | |
| 1991 | 597 | 473 | 479 | 368 | 314 | 296 | 339 | 295 | 324 | 274 | 239 | 790 | | | |
| 1992 | 1260 | 517 | 410 | 415 | 319 | 272 | 257 | 294 | 255 | 281 | 238 | 892 | | | |
| 1993 | 1623 | 1113 | 457 | 362 | 367 | 282 | 240 | 227 | 259 | 225 | 248 | 998 | | | |
| 1994 | 556 | 1437 | 986 | 404 | 320 | 325 | 249 | 213 | 201 | 230 | 200 | 1103 | | | |
| 1995 | 1648 | 490 | 1266 | 868 | 356 | 282 | 286 | 220 | 188 | 177 | 202 | 1147 | | | |
| 1996 | 1780 | 1461 | 434 | 1123 | 770 | 316 | 250 | 253 | 195 | 166 | 157 | 1197 | | | |
| 1997 | 1630 | 1570 | 1289 | 383 | 990 | 679 | 279 | 221 | 224 | 172 | 147 | 1194 | | | |
| 1998 | 2295 | 1445 | 1392 | 1143 | 339 | 878 | 602 | 247 | 196 | 198 | 152 | 1188 | | | |
| 1999 | 3749 | 2024 | 1275 | 1227 | 1008 | 299 | 774 | 531 | 218 | 173 | 175 | 1182 | | | |
| 2000 | 3378 | 3323 | 1794 | 1130 | 1088 | 893 | 265 | 686 | 470 | 193 | 153 | 1203 | | | |
| 2001 | 1700 | 2987 | 2939 | 1587 | 999 | 962 | 790 | 235 | 607 | 416 | 171 | 1199 | | | |
| 2002 | 1201 | 1503 | 2640 | 2597 | 1402 | 883 | 850 | 698 | 207 | 536 | 368 | 1211 | | | |
| 2003 | 1231 | 1064 | 1332 | 2339 | 2301 | 1243 | 782 | 753 | 619 | 184 | 475 | 1398 | | | |
| 2004 | 1790 | 1090 | 942 | 1179 | 2071 | 2037 | 1100 | 693 | 667 | 548 | 163 | 1659 | | | |
| 2005 | 1092 | 1578 | 961 | 831 | 1039 | 1826 | 1796 | 970 | 611 | 588 | 483 | 1606 | | | |
| 2006 | 725 | 966 | 1397 | 851 | 736 | 920 | 1617 | 1590 | 859 | 541 | 521 | 1849 | | | |
| 2007 | 1052 | 643 | 856 | 1238 | 754 | 652 | 815 | 1432 | 1409 | 761 | 479 | 2100 | | | |
| 2008 | 1670 | 934 | 570 | 760 | 1099 | 669 | 579 | 724 | 1272 | 1251 | 675 | 2289 | | | |
| 2009 | 2554 | 1424 | 797 | 487 | 648 | 937 | 571 | 493 | 617 | 1084 | 1067 | 2529 | | | |
| 2010 | 2093 | 2090 | 1166 | 652 | 398 | 531 | 767 | 467 | 404 | 505 | 888 | 2943 | | | |
| 2011 | 1911 | 1606 | 1604 | 895 | 500 | 306 | 407 | 589 | 358 | 310 | 388 | 2939 | | | |
| 2012 | 1026 | 1584 | 1331 | 1329 | 741 | 415 | 253 | 338 | 488 | 297 | 257 | 2758 | | | |
| 2013 | 1312 | 852 | 1316 | 1106 | 1104 | 616 | 344 | 210 | 280 | 405 | 247 | 2504 | | | |
| 2014 | 2368 | 1105 | 717 | 1108 | 931 | 929 | 518 | 290 | 177 | 236 | 341 | 2315 | | | |
| 2015 | 3360 | 2020 | 942 | 612 | 945 | 794 | 793 | 442 | 247 | 151 | 201 | 2266 | | | |
| 2016 | 5060 | 2898 | 1743 | 813 | 528 | 815 | 685 | 684 | 381 | 213 | 130 | 2128 | | | |

Table 7.11. (continued).

| | males | | | | | | | | | | | |
|------|-------|-------|-------|-------|-------|-------|-------|-------|-------|------|------|------|
| | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 |
| 1991 | 5157 | 4523 | 5744 | 10233 | 10172 | 5524 | 3487 | 3361 | 2760 | 820 | 2121 | 1454 |
| 1992 | 7476 | 4617 | 4046 | 5127 | 9093 | 8972 | 4834 | 3035 | 2918 | 2394 | 711 | 1838 |
| 1993 | 4571 | 6695 | 4134 | 3620 | 4577 | 8092 | 7958 | 4278 | 2683 | 2578 | 2114 | 628 |
| 1994 | 3033 | 4095 | 5995 | 3699 | 3233 | 4078 | 7190 | 7057 | 3790 | 2376 | 2283 | 1872 |
| 1995 | 4392 | 2717 | 3665 | 5361 | 3300 | 2874 | 3611 | 6348 | 6222 | 3340 | 2093 | 2011 |
| 1996 | 6944 | 3934 | 2433 | 3280 | 4791 | 2942 | 2556 | 3206 | 5632 | 5518 | 2961 | 1856 |
| 1997 | 11013 | 6219 | 3522 | 2176 | 2927 | 4260 | 2607 | 2259 | 2830 | 4968 | 4867 | 2612 |
| 1998 | 9756 | 9864 | 5569 | 3152 | 1944 | 2610 | 3789 | 2314 | 2004 | 2509 | 4405 | 4315 |
| 1999 | 10270 | 8738 | 8831 | 4981 | 2813 | 1729 | 2312 | 3348 | 2043 | 1768 | 2213 | 3885 |
| 2000 | 5880 | 9199 | 7824 | 7902 | 4450 | 2507 | 1537 | 2052 | 2969 | 1811 | 1567 | 1962 |
| 2001 | 8009 | 5266 | 8236 | 7000 | 7057 | 3963 | 2226 | 1362 | 1816 | 2627 | 1602 | 1386 |
| 2002 | 15152 | 7173 | 4715 | 7368 | 6250 | 6282 | 3517 | 1971 | 1204 | 1606 | 2322 | 1416 |
| 2003 | 22180 | 13572 | 6423 | 4219 | 6583 | 5570 | 5584 | 3121 | 1747 | 1068 | 1423 | 2057 |
| 2004 | 33826 | 19866 | 12152 | 5747 | 3769 | 5865 | 4949 | 4952 | 2764 | 1547 | 945 | 1260 |
| 2005 | 11301 | 30295 | 17785 | 10868 | 5128 | 3351 | 5195 | 4373 | 4370 | 2438 | 1364 | 833 |
| 2006 | 6584 | 10122 | 27125 | 15912 | 9708 | 4569 | 2977 | 4606 | 3873 | 3869 | 2159 | 1208 |
| 2007 | 9020 | 5897 | 9063 | 24272 | 14217 | 8652 | 4062 | 2642 | 4084 | 3433 | 3429 | 1913 |
| 2008 | 9717 | 8079 | 5281 | 8111 | 21695 | 12682 | 7701 | 3610 | 2346 | 3626 | 3048 | 3044 |
| 2009 | 10750 | 8698 | 7223 | 4706 | 7180 | 18995 | 10975 | 6612 | 3087 | 2003 | 3094 | 2600 |
| 2010 | 28986 | 9617 | 7763 | 6409 | 4125 | 6168 | 15971 | 9094 | 5439 | 2532 | 1641 | 2533 |
| 2011 | 16843 | 25904 | 8560 | 6841 | 5529 | 3437 | 4954 | 12511 | 7036 | 4186 | 1945 | 1260 |
| 2012 | 20922 | 15070 | 23130 | 7605 | 6014 | 4777 | 2916 | 4150 | 10414 | 5841 | 3472 | 1612 |
| 2013 | 10067 | 18720 | 13457 | 20556 | 6688 | 5200 | 4057 | 2446 | 3460 | 8661 | 4853 | 2884 |
| 2014 | 16038 | 9010 | 16727 | 11977 | 18138 | 5821 | 4459 | 3444 | 2066 | 2917 | 7295 | 4087 |
| 2015 | 25171 | 14356 | 8055 | 14907 | 10602 | 15881 | 5037 | 3829 | 2945 | 1764 | 2489 | 6223 |
| 2016 | 31046 | 22535 | 12841 | 7187 | 13232 | 9332 | 13854 | 4368 | 3310 | 2543 | 1522 | 2147 |

Table 7.11. (continued). Males

| | males continued | | | | | | | | | | | |
|------|-----------------|------|------|------|------|------|------|------|------|------|------|------|
| | 14 | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22 | 23 | 24 | 25 |
| 1991 | 597 | 473 | 479 | 368 | 314 | 296 | 339 | 295 | 324 | 274 | 239 | 790 |
| 1992 | 1260 | 517 | 410 | 415 | 319 | 272 | 257 | 294 | 255 | 281 | 238 | 892 |
| 1993 | 1623 | 1113 | 457 | 362 | 367 | 282 | 240 | 227 | 259 | 225 | 248 | 998 |
| 1994 | 556 | 1437 | 986 | 404 | 320 | 325 | 249 | 213 | 201 | 230 | 200 | 1103 |
| 1995 | 1649 | 490 | 1266 | 868 | 356 | 282 | 286 | 220 | 188 | 177 | 202 | 1147 |
| 1996 | 1783 | 1462 | 434 | 1123 | 770 | 316 | 250 | 253 | 195 | 166 | 157 | 1197 |
| 1997 | 1637 | 1572 | 1290 | 383 | 990 | 679 | 279 | 221 | 224 | 172 | 147 | 1194 |
| 1998 | 2315 | 1451 | 1394 | 1143 | 339 | 878 | 602 | 247 | 196 | 198 | 152 | 1188 |
| 1999 | 3806 | 2042 | 1280 | 1229 | 1008 | 299 | 774 | 531 | 218 | 173 | 175 | 1182 |
| 2000 | 3443 | 3373 | 1810 | 1134 | 1090 | 894 | 265 | 686 | 470 | 193 | 153 | 1203 |
| 2001 | 1735 | 3045 | 2983 | 1601 | 1003 | 964 | 790 | 235 | 607 | 416 | 171 | 1199 |
| 2002 | 1225 | 1533 | 2691 | 2636 | 1415 | 887 | 852 | 699 | 207 | 536 | 368 | 1211 |
| 2003 | 1254 | 1085 | 1359 | 2385 | 2336 | 1254 | 786 | 755 | 619 | 184 | 475 | 1399 |
| 2004 | 1821 | 1110 | 961 | 1203 | 2111 | 2068 | 1110 | 695 | 668 | 548 | 163 | 1659 |
| 2005 | 1111 | 1606 | 979 | 847 | 1060 | 1861 | 1823 | 978 | 613 | 589 | 483 | 1606 |
| 2006 | 738 | 983 | 1422 | 867 | 750 | 939 | 1648 | 1614 | 866 | 543 | 522 | 1849 |
| 2007 | 1070 | 654 | 871 | 1260 | 768 | 664 | 832 | 1460 | 1430 | 768 | 481 | 2101 |
| 2008 | 1698 | 950 | 580 | 773 | 1118 | 682 | 590 | 738 | 1296 | 1270 | 681 | 2292 |
| 2009 | 2596 | 1448 | 810 | 495 | 660 | 954 | 581 | 503 | 630 | 1106 | 1083 | 2536 |
| 2010 | 2128 | 2125 | 1186 | 663 | 405 | 540 | 781 | 476 | 412 | 516 | 905 | 2962 |
| 2011 | 1944 | 1633 | 1631 | 910 | 509 | 311 | 414 | 599 | 365 | 316 | 396 | 2967 |
| 2012 | 1044 | 1611 | 1354 | 1352 | 754 | 422 | 258 | 343 | 497 | 303 | 262 | 2787 |
| 2013 | 1339 | 867 | 1338 | 1124 | 1122 | 626 | 350 | 214 | 285 | 412 | 251 | 2532 |
| 2014 | 2428 | 1127 | 730 | 1127 | 946 | 945 | 527 | 295 | 180 | 240 | 347 | 2343 |
| 2015 | 3486 | 2071 | 962 | 623 | 961 | 807 | 806 | 450 | 252 | 154 | 205 | 2295 |
| 2016 | 5369 | 3007 | 1787 | 830 | 537 | 829 | 696 | 695 | 388 | 217 | 133 | 2156 |

Table 7.12. Estimate of the spawning biomass at age (t) from the stock assessment model.

| | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 |
|------|---|----|----|-----|-----|------|------|------|------|------|------|-------|-------|
| 1991 | 1 | 2 | 12 | 73 | 219 | 325 | 507 | 1069 | 1664 | 802 | 2926 | 2554 | 1247 |
| 1992 | 1 | 2 | 9 | 36 | 194 | 523 | 698 | 962 | 1757 | 2339 | 981 | 3228 | 2633 |
| 1993 | 1 | 4 | 9 | 26 | 98 | 469 | 1141 | 1350 | 1611 | 2516 | 2916 | 1102 | 3391 |
| 1994 | 0 | 2 | 13 | 26 | 69 | 236 | 1027 | 2214 | 2266 | 2314 | 3145 | 3286 | 1161 |
| 1995 | 0 | 1 | 8 | 38 | 70 | 167 | 515 | 1983 | 3696 | 3237 | 2877 | 3526 | 3444 |
| 1996 | 1 | 2 | 5 | 23 | 102 | 171 | 365 | 1001 | 3331 | 5316 | 4051 | 3247 | 3719 |
| 1997 | 1 | 3 | 7 | 15 | 63 | 247 | 372 | 705 | 1672 | 4765 | 6616 | 4547 | 3406 |
| 1998 | 1 | 5 | 12 | 22 | 42 | 152 | 542 | 723 | 1185 | 2405 | 5962 | 7466 | 4795 |
| 1999 | 1 | 5 | 19 | 35 | 60 | 100 | 331 | 1048 | 1209 | 1695 | 2993 | 6693 | 7833 |
| 2000 | 1 | 5 | 17 | 56 | 95 | 146 | 220 | 642 | 1760 | 1738 | 2121 | 3377 | 7057 |
| 2001 | 1 | 3 | 17 | 50 | 151 | 230 | 319 | 426 | 1076 | 2524 | 2169 | 2387 | 3553 |
| 2002 | 2 | 4 | 10 | 53 | 134 | 365 | 503 | 617 | 714 | 1543 | 3148 | 2440 | 2510 |
| 2003 | 3 | 7 | 14 | 30 | 141 | 324 | 800 | 977 | 1036 | 1026 | 1930 | 3551 | 2572 |
| 2004 | 4 | 11 | 26 | 41 | 81 | 341 | 709 | 1551 | 1639 | 1488 | 1282 | 2175 | 3740 |
| 2005 | 1 | 16 | 38 | 77 | 110 | 195 | 744 | 1369 | 2592 | 2344 | 1851 | 1439 | 2281 |
| 2006 | 1 | 5 | 57 | 113 | 207 | 266 | 426 | 1442 | 2297 | 3721 | 2929 | 2086 | 1515 |
| 2007 | 1 | 3 | 19 | 173 | 304 | 503 | 581 | 827 | 2422 | 3301 | 4654 | 3304 | 2199 |
| 2008 | 1 | 4 | 11 | 58 | 464 | 738 | 1104 | 1131 | 1391 | 3487 | 4137 | 5258 | 3489 |
| 2009 | 1 | 5 | 15 | 33 | 152 | 1094 | 1562 | 2065 | 1828 | 1925 | 4198 | 4491 | 5335 |
| 2010 | 3 | 5 | 16 | 45 | 85 | 347 | 2230 | 2810 | 3208 | 2429 | 2225 | 4375 | 4373 |
| 2011 | 2 | 14 | 18 | 47 | 111 | 186 | 667 | 3769 | 4095 | 3995 | 2631 | 2173 | 3993 |
| 2012 | 2 | 8 | 48 | 53 | 121 | 256 | 383 | 1214 | 5927 | 5507 | 4674 | 2776 | 2143 |
| 2013 | 1 | 10 | 28 | 143 | 136 | 280 | 528 | 698 | 1912 | 7985 | 6455 | 4941 | 2742 |
| 2014 | 2 | 5 | 35 | 84 | 372 | 317 | 585 | 975 | 1114 | 2611 | 9489 | 6918 | 4948 |
| 2015 | 3 | 8 | 17 | 105 | 220 | 876 | 670 | 1094 | 1577 | 1541 | 3144 | 10303 | 7019 |
| 2016 | 4 | 12 | 27 | 51 | 276 | 522 | 1873 | 1268 | 1789 | 2207 | 1877 | 3452 | 10572 |

Table 7.12. (continued).

| | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22 | 23 | 24 | 25 |
|------|------|------|------|------|------|------|------|------|------|------|-------|
| 1991 | 1126 | 1268 | 1064 | 979 | 987 | 1197 | 1096 | 1265 | 1116 | 1009 | 3458 |
| 1992 | 1232 | 1085 | 1200 | 995 | 907 | 907 | 1093 | 996 | 1143 | 1005 | 3901 |
| 1993 | 2653 | 1209 | 1046 | 1143 | 939 | 849 | 844 | 1012 | 918 | 1049 | 4364 |
| 1994 | 3425 | 2610 | 1169 | 999 | 1081 | 881 | 792 | 783 | 935 | 844 | 4825 |
| 1995 | 1167 | 3352 | 2510 | 1111 | 940 | 1010 | 818 | 732 | 720 | 855 | 5020 |
| 1996 | 3483 | 1149 | 3245 | 2400 | 1052 | 884 | 944 | 760 | 677 | 663 | 5236 |
| 1997 | 3741 | 3412 | 1107 | 3087 | 2262 | 984 | 821 | 872 | 699 | 620 | 5222 |
| 1998 | 3444 | 3685 | 3303 | 1058 | 2924 | 2126 | 919 | 763 | 807 | 644 | 5198 |
| 1999 | 4824 | 3375 | 3549 | 3142 | 997 | 2734 | 1976 | 850 | 702 | 739 | 5172 |
| 2000 | 7919 | 4751 | 3266 | 3392 | 2975 | 937 | 2554 | 1836 | 786 | 646 | 5262 |
| 2001 | 7119 | 7781 | 4588 | 3115 | 3205 | 2790 | 873 | 2367 | 1693 | 722 | 5246 |
| 2002 | 3582 | 6990 | 7509 | 4373 | 2942 | 3004 | 2599 | 809 | 2182 | 1555 | 5296 |
| 2003 | 2537 | 3526 | 6763 | 7176 | 4140 | 2764 | 2805 | 2414 | 748 | 2009 | 6118 |
| 2004 | 2598 | 2495 | 3408 | 6458 | 6788 | 3886 | 2579 | 2603 | 2229 | 688 | 7257 |
| 2005 | 3761 | 2545 | 2402 | 3241 | 6083 | 6346 | 3611 | 2383 | 2394 | 2041 | 7026 |
| 2006 | 2303 | 3699 | 2460 | 2293 | 3066 | 5711 | 5920 | 3351 | 2200 | 2201 | 8090 |
| 2007 | 1531 | 2267 | 3579 | 2350 | 2171 | 2880 | 5332 | 5498 | 3097 | 2025 | 9186 |
| 2008 | 2226 | 1510 | 2198 | 3426 | 2229 | 2044 | 2694 | 4961 | 5091 | 2855 | 10015 |
| 2009 | 3394 | 2110 | 1407 | 2022 | 3122 | 2016 | 1837 | 2409 | 4414 | 4510 | 11062 |
| 2010 | 4982 | 3087 | 1886 | 1242 | 1768 | 2710 | 1739 | 1576 | 2057 | 3753 | 12877 |
| 2011 | 3827 | 4247 | 2586 | 1560 | 1018 | 1439 | 2191 | 1399 | 1261 | 1639 | 12860 |
| 2012 | 3776 | 3525 | 3844 | 2312 | 1382 | 895 | 1257 | 1904 | 1209 | 1086 | 12066 |
| 2013 | 2029 | 3483 | 3196 | 3442 | 2051 | 1217 | 783 | 1094 | 1649 | 1043 | 10953 |
| 2014 | 2633 | 1898 | 3202 | 2902 | 3096 | 1831 | 1080 | 691 | 960 | 1442 | 10130 |
| 2015 | 4814 | 2495 | 1768 | 2946 | 2645 | 2801 | 1646 | 965 | 615 | 851 | 9913 |
| 2016 | 6907 | 4614 | 2350 | 1645 | 2715 | 2419 | 2546 | 1488 | 868 | 551 | 9311 |

Table 7.13. Projections of spawning biomass (1,000s t), catch (1,000s t), and fishing mortality rate for each of the seven scenarios. The value of $B_{40\%}$ and $B_{35\%}$ are 50,800 t and 44,400 t, respectively.

| Scenarios 1 and 2 | | | | Scenario 3 | | | |
|--|------------|--------------|----------|--------------------------------|------------|--------------|----------|
| Maximum ABC harvest permissible | | | | 1/2 maximum ABC harvest | | | |
| Year | FSB | catch | F | Year | FSB | catch | F |
| 2016 | 59,321 | 4,533 | 0.034 | 2016 | 59,321 | 4,533 | 0.034 |
| 2017 | 60,172 | 8,879 | 0.066 | 2017 | 60,340 | 4,440 | 0.033 |
| 2018 | 60,210 | 9,197 | 0.066 | 2018 | 62,245 | 4,798 | 0.033 |
| 2019 | 64,105 | 9,514 | 0.066 | 2019 | 66,424 | 5,157 | 0.034 |
| 2020 | 64,376 | 9,610 | 0.066 | 2020 | 68,799 | 5,351 | 0.034 |
| 2021 | 64,573 | 9,691 | 0.066 | 2021 | 71,124 | 5,536 | 0.034 |
| 2022 | 64,785 | 9,715 | 0.066 | 2022 | 73,465 | 5,687 | 0.034 |
| 2023 | 65,162 | 9,686 | 0.066 | 2023 | 75,974 | 5,802 | 0.034 |
| 2024 | 65,682 | 9,624 | 0.066 | 2024 | 78,627 | 5,891 | 0.034 |
| 2025 | 66,087 | 9,544 | 0.066 | 2025 | 81,115 | 5,960 | 0.034 |
| 2026 | 66,133 | 9,450 | 0.066 | 2026 | 83,111 | 6,010 | 0.034 |
| 2027 | 65,788 | 9,348 | 0.066 | 2027 | 84,521 | 6,044 | 0.034 |
| 2028 | 65,143 | 9,240 | 0.066 | 2028 | 85,407 | 6,064 | 0.034 |
| 2029 | 64,393 | 9,137 | 0.066 | 2029 | 85,996 | 6,076 | 0.034 |

| Scenario 4 | | | | Scenario 5 | | | |
|--|------------|--------------|----------|-------------------|------------|--------------|----------|
| Harvest at avg F over the past five years | | | | No fishing | | | |
| Year | FSB | catch | F | Year | FSB | catch | F |
| 2016 | 59,321 | 4,533 | 0.034 | 2016 | 59,321 | 4,533 | 0.05 |
| 2017 | 60,336 | 4,533 | 0.033 | 2017 | 60,408 | 0 | 0 |
| 2018 | 62,203 | 5,573 | 0.039 | 2018 | 64,295 | 0 | 0 |
| 2019 | 66,324 | 6,613 | 0.044 | 2019 | 69,044 | 0 | 0 |
| 2020 | 68,025 | 6,801 | 0.044 | 2020 | 75,065 | 0 | 0 |
| 2021 | 69,646 | 6,975 | 0.044 | 2021 | 82,233 | 0 | 0 |
| 2022 | 71,258 | 7,106 | 0.044 | 2022 | 89,964 | 0 | 0 |
| 2023 | 73,016 | 7,193 | 0.044 | 2023 | 97,586 | 0 | 0 |
| 2024 | 74,897 | 7,250 | 0.044 | 2024 | 104,671 | 0 | 0 |
| 2025 | 76,612 | 7,284 | 0.044 | 2025 | 111,008 | 0 | 0 |
| 2026 | 77,866 | 7,299 | 0.044 | 2026 | 116,546 | 0 | 0 |
| 2027 | 78,588 | 7,297 | 0.044 | 2027 | 121,297 | 0 | 0 |
| 2028 | 78,856 | 7,283 | 0.044 | 2028 | 125,333 | 0 | 0 |
| 2029 | 78,890 | 7,263 | 0.044 | 2029 | 128,745 | 0 | 0 |

Table 7.13. (continued).

| Scenario 6 | | | | Scenario 7 | | | |
|-------------------------------------|------------|--------------|----------|---|------------|--------------|----------|
| Determination of overfishing | | | | Determination of approaching overfishing | | | |
| Year | FSB | catch | F | Year | FSB | catch | F |
| 2016 | 59,321 | 4,533 | 0.034 | 2016 | 59,321 | 4,533 | 0.034 |
| 2017 | 60,114 | 10,361 | 0.078 | 2017 | 60,172 | 8,881 | 0.066 |
| 2018 | 59,149 | 10,260 | 0.078 | 2018 | 59,879 | 8,887 | 0.066 |
| 2019 | 58,648 | 10,239 | 0.078 | 2019 | 59,984 | 10,449 | 0.078 |
| 2020 | 58,298 | 10,287 | 0.078 | 2020 | 59,605 | 10,483 | 0.078 |
| 2021 | 57,938 | 10,330 | 0.078 | 2021 | 59,202 | 10,511 | 0.078 |
| 2022 | 57,669 | 10,317 | 0.078 | 2022 | 58,874 | 10,482 | 0.078 |
| 2023 | 57,637 | 10,253 | 0.078 | 2023 | 58,770 | 10,402 | 0.078 |
| 2024 | 57,814 | 10,162 | 0.078 | 2024 | 58,864 | 10,294 | 0.078 |
| 2025 | 57,947 | 10,056 | 0.078 | 2025 | 58,905 | 10,173 | 0.078 |
| 2026 | 57,807 | 9,940 | 0.078 | 2026 | 58,670 | 10,043 | 0.078 |
| 2027 | 57,361 | 9,820 | 0.078 | 2027 | 58,127 | 9,909 | 0.078 |
| 2028 | 56,691 | 9,684 | 0.078 | 2028 | 57,363 | 9,769 | 0.078 |
| 2029 | 55,968 | 9,531 | 0.077 | 2029 | 56,548 | 9,612 | 0.077 |

Table 7.14. Selected parameter estimates and standard deviations for the Kamchatka flounder stock assessment model (2016 assessment).

| details | parameter name | value | std dev | details | parameter name | value | std dev |
|------------|----------------------------|--------|---------|---------|-------------------------|--------|---------|
| | shelf survey q | 1.05 | 0.06 | 2007 | total_biomass | 147890 | 5677 |
| | Aleutian Islands survey q | 0.31 | 0.03 | 2008 | total_biomass | 156570 | 5948 |
| | mean_log_recruitment | 8.40 | 0.06 | 2009 | total_biomass | 159060 | 6215 |
| | fishery_inflection_females | 4.25 | 0.30 | 2010 | total_biomass | 155800 | 6492 |
| | fishery_inflection_males | 4.84 | 0.42 | 2011 | total_biomass | 143640 | 6747 |
| ascending | survey1_slope_females | 0.94 | 0.14 | 2012 | total_biomass | 143460 | 7065 |
| ascending | survey1_inflection_females | 1.87 | 0.24 | 2013 | total_biomass | 143720 | 7398 |
| descending | survey1_slope_females | 0.33 | 0.03 | 2014 | total_biomass | 146450 | 7764 |
| descending | survey1_inflection_females | 7.42 | 0.75 | 2015 | total_biomass | 151390 | 8195 |
| ascending | survey1_slope_males | 0.82 | 0.21 | 2016 | total_biomass | 159030 | 8730 |
| ascending | survey1_inflection_males | 0.93 | 0.28 | 1991 | female spawning biomass | 21330 | 2156 |
| descending | survey1_slope_males | 0.54 | 0.06 | 1992 | female spawning biomass | 23004 | 2092 |
| descending | survey1_inflection_males | 7.75 | 0.50 | 1993 | female spawning biomass | 25641 | 2061 |
| ascending | survey2_slope_females | 1.10 | 0.14 | 1994 | female spawning biomass | 28901 | 2057 |
| ascending | survey2_inflection_females | 5.80 | 0.22 | 1995 | female spawning biomass | 32459 | 2077 |
| ascending | survey2_slope_males | 1.78 | 0.25 | 1996 | female spawning biomass | 36466 | 2124 |
| ascending | survey2_inflection_males | 4.23 | 0.16 | 1997 | female spawning biomass | 40024 | 2183 |
| ascending | survey3_slope_females | 1.54 | 0.49 | 1998 | female spawning biomass | 43284 | 2244 |
| ascending | survey3_inflection_females | 1.61 | 0.17 | 1999 | female spawning biomass | 45657 | 2294 |
| ascending | survey3_slope_males | 0.94 | 0.29 | 2000 | female spawning biomass | 47713 | 2333 |
| ascending | survey3_inflection_males | 1.47 | 0.20 | 2001 | female spawning biomass | 49213 | 2360 |
| 1991 | total_biomass | 67314 | 3991 | 2002 | female spawning biomass | 50362 | 2378 |
| 1992 | total_biomass | 71923 | 4023 | 2003 | female spawning biomass | 51545 | 2393 |
| 1993 | total_biomass | 77444 | 4072 | 2004 | female spawning biomass | 52818 | 2411 |
| 1994 | total_biomass | 82610 | 4134 | 2005 | female spawning biomass | 54167 | 2439 |
| 1995 | total_biomass | 86711 | 4200 | 2006 | female spawning biomass | 56113 | 2485 |
| 1996 | total_biomass | 90890 | 4265 | 2007 | female spawning biomass | 58367 | 2542 |
| 1997 | total_biomass | 94286 | 4330 | 2008 | female spawning biomass | 61025 | 2611 |
| 1998 | total_biomass | 98068 | 4396 | 2009 | female spawning biomass | 61684 | 2690 |
| 1999 | total_biomass | 101370 | 4465 | 2010 | female spawning biomass | 60470 | 2791 |
| 2000 | total_biomass | 105070 | 4535 | 2011 | female spawning biomass | 56029 | 2920 |
| 2001 | total_biomass | 108570 | 4610 | 2012 | female spawning biomass | 56550 | 3109 |
| 2002 | total_biomass | 112400 | 4701 | 2013 | female spawning biomass | 56913 | 3310 |
| 2003 | total_biomass | 117370 | 4816 | 2014 | female spawning biomass | 57596 | 3496 |
| 2004 | total_biomass | 124050 | 4984 | 2015 | female spawning biomass | 58485 | 3660 |
| 2005 | total_biomass | 131100 | 5184 | 2016 | female spawning biomass | 60038 | 3810 |
| 2006 | total_biomass | 139310 | 5419 | | | | |

Table 7.15. Non-target catch (t) when Kamchatka flounder were fishery targets, 2011-2016.

| Row Labels | 2011 | 2012 | 2013 | 2014 | 2015 | 2016 | Grand Total |
|--------------------------------|--------------|-------------|-------------|-------------|-------------|-------------|--------------|
| Benthic urochordata | 0.0 | | | 0.0 | 0.0 | 0.0 | 0.0 |
| Bivalves | 0.0 | | | 0.0 | 0.0 | 0.0 | 0.0 |
| Brittle star unidentified | 0.8 | 0.0 | | 0.0 | 0.0 | 0.0 | 0.9 |
| Corals Bryozoans | 0.1 | 0.0 | 0.0 | 0.8 | 0.3 | 1.3 | 2.6 |
| Deep sea smelts (bathylagidae) | | | | | 0.0 | 0.0 | 0.0 |
| Eelpouts | 14.6 | 19.3 | 8.4 | 3.2 | 0.5 | 1.7 | 47.7 |
| Eulachon | 0.0 | | 0.0 | | | | 0.0 |
| Giant Grenadier | 241.0 | 4.6 | 11.8 | 10.1 | 1.2 | 15.7 | 284.4 |
| Grenadier | 0.1 | 0.0 | | | 0.4 | 2.1 | 2.7 |
| Hermit crab unidentified | | 0.0 | | | 0.0 | | 0.0 |
| Invertebrate unidentified | 5.6 | | | | 0.1 | 0.0 | 5.8 |
| Lanternfishes (myctophidae) | 0.0 | | 0.0 | 0.0 | 0.3 | 0.1 | 0.3 |
| Large Sculpins | 3.0 | 0.6 | 0.1 | 3.1 | 11.2 | 6.2 | 24.2 |
| Misc crabs | | | 0.0 | 0.0 | 0.0 | 0.1 | 0.1 |
| Misc crustaceans | | | | | 0.3 | | 0.3 |
| Misc deep fish | | | | 0.0 | 0.1 | 0.0 | 0.1 |
| Misc fish | 0.5 | 0.0 | 0.0 | 0.1 | 0.1 | 1.1 | 1.8 |
| Misc inverts (worms etc) | 0.0 | 0.0 | | 0.0 | 0.0 | 0.0 | 0.0 |
| Other Sculpins | 26.5 | 0.1 | 0.4 | 11.1 | 15.6 | 10.0 | 63.8 |
| Pandalid shrimp | 0.1 | 0.0 | 0.0 | 0.1 | 0.1 | 0.2 | 0.5 |
| Polychaete unidentified | 0.0 | 0.0 | | | 0.0 | 0.0 | 0.0 |
| Scypho jellies | 0.7 | 0.0 | 0.0 | 0.2 | 0.1 | 0.4 | 1.4 |
| Sea anemone unidentified | 1.2 | 0.6 | 0.2 | 0.1 | 0.1 | 0.0 | 2.2 |
| Sea pens whips | 0.0 | | | | 0.0 | | 0.0 |
| Sea star | 2.6 | 0.6 | 0.3 | 0.5 | 1.6 | 0.7 | 6.3 |
| Snails | 0.1 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 |
| Sponge unidentified | 16.0 | 0.0 | 0.2 | 1.5 | 11.1 | 6.2 | 35.0 |
| Stichaeidae | | | | | 0.0 | | 0.0 |
| urchins dollars cucumbers | 0.4 | 0.5 | 0.0 | 0.3 | 0.1 | 0.0 | 1.4 |
| Grand Total | 313.4 | 26.4 | 21.5 | 31.1 | 43.3 | 46.1 | 481.8 |




Figure 7-1. 2016 Kamchatka flounder catch (t) by week from Alaska Regional Office catch reports.




Figure 7.2—Number of hauls where arrowtooth flounder and Kamchatka flounder were identified during the annual Bering Sea shelf surveys, 1982-2016.





Figure 7-3. Distribution and relative of abundance of Kamchatka flounder from the three surveys conducted in 2016 with catch > 5 kg/ha. (top panel) and from the 2010 slope survey (bottom panel). Plus signs indicates stations with no catch of Kamchatka flounder.




Figure 7-4. Distribution and relative abundance of Kamchatka flounder from the 2006 Aleutian Islands survey.




Figure 7-4 (continued).




Figure 7-4 (continued).




Figure 7.5. Kamchatka flounder population length composition estimates from the shelf, slope and Aleutian Islands survey for males and females.




Figure 7.5. (continued).





Figure 7-6 Kamchatka flounder length-weight plots for male and females.




Figure 7-7 Estimated weight-at-age for male and female Kamchatka flounder from a 2010 age sample from the Aleutian Islands.




Figure 7-8 Initial area-specific catchability values were assigned in the assessment model according to the proportion of the average biomass from the time-series of each trawl survey (shelf, slope and Aleutian Islands) in years when all three surveys were conducted (2002, 2010, 2012, 2016).




Figure 7-9 Estimated fishery selectivity from two model runs, unconstrained (left panel) and estimated with slope parameter fixed (right panel). Maturity curve is also plotted.




Figure 7-10 Model estimates of survey selectivity, males and females, for the shelf, slope and Aleutian Islands surveys.




Figure 7-11 Comparison of spawning biomass estimates with slope survey catchability fixed at 0.18 (solid line) and 0.1 (dotted line). The difference in total likelihood between these models was 1.95 (with the higher biomass model being favored).




Figure 7-12 Plot of $-\log(\text{likelihood})$ values for model components when profiling over values of slope survey q ranging from 0.05 to 0.3.




Figure 7-13 Assessment model estimate of female spawning biomass and total Kamchatka flounder total biomass (t) and 95% (?) confidence bounds from 1991-2016.




Figure 7-14 Assessment model estimate of female spawning biomass (t) and 95% (?) confidence bounds relative to $B_{40\%}$.




Figure 7-15 Assessment model fit (yellow line) to the shelf, slope and Aleutian Islands surveys (shown with 95% confidence intervals).




Figure 7-16. Assessment model estimates of full selection F_s for 1991-2016.




Fig. 7-17 Comparison of the average observed (open circles) proportion at length from the time-series to the average predicted (solid line) proportion at length from the model for the fishery, and the three surveys on the Bering Sea shelf, slope and the Aleutian Islands.




Figure 7-18 Assessment model fit (black dotted line) to the shelf, slope and Aleutian Islands survey size compositions (red solid line).




Figure 7-18 continued.




Figure 7-18 continued.




Figure 7.18 (continued).




Figure 7.18 (continued).




Figure 7.19. Phase plane figure of Kamchatka flounder female spawning biomass (t) and annual fishing mortality rate.




Figure 7.20. Restrospective plot of Kamchatka flounder female spawning biomass for 2007-2016.