## OKEANOS EXPLORER ROV DIVE SUMMARY

|  |   |  | 1  |             |                   |
|--|---|--|--|-------------|-------------------|
| Site Name  |   | East of Necke  | r  |             |                   |
| ROV<br>Lead/Expediti<br>on<br>Coordinator              | Karl I  | Mcletchie/ Brian R   | C Kennedy  | 1 th        | EX1603 Dive 1     |
| Science Team<br>Leads                                  | Daniel Wagner and Jonathan Tree   |  |  | Necker Is   |                   |
| General Area<br>Descriptor                             | Papahanaumokuakea Marine National Monument  |  |  |             |                   |
| ROV Dive   | Cruise  | e Season   | Leg  |             | Dive Number       |
| Name   | EX1603  |  | 1  |             | DIVE01            |
| Equipment  | ROV:  |  | Deep Discoverer  |             |                   |
| Deployed   | Camera Platform:  |  | Seirios  |             |                   |
|  | 🛛 D2 CTD  |  | Depth  |             | Altitude          |
| ROV  | Scanning S  | onar   | USBL Position  |             | Heading           |
| Measurement  | Pitch   |  | 🛛 Roll   |             | HD Camera 1       |
| S  | HD Camera 2   |  | 🛛 ROV HD 2   |             | Seirios CTD       |
|  | Temperature Probe   |  | 🛛 D2 DO Sensor   |             | Seirios DO sensor |
| Equipment<br>Malfunctions                              | The Kraft manipulator arm malfunctioned on the pre-dive. After an hour of troubleshooting failed to resolve the issue we proceeded with the dive. |  |  |             |                   |
| ROV Dive<br>Summary<br>(From<br>processed<br>ROV data) |   | 23°, 34.3<br>2016-02<br>23°, 35.4<br>2016-02<br>23°, 34.4<br>2016-02 | -27T19:40:59.542000<br>305' N ; 164°, 01.568' W<br>-28T02:47:11.227000<br>172' N ; 164°, 02.549' W<br>-28T00:17:45.380000<br>486' N ; 164°, 01.839' W<br>-27T21:57:22.670000<br>409' N ; 164°, 01.644' W |             |                   |
| Special Notes  |   |  |  |             |                   |
| Scientists   | Name  | Affiliation  |  | Email Addı  | ress              |
| Involved   | Steve   |  |  |             | cavitch@temple.e  |
| (please  | Auscavitch  | Temple University  |  | du          |                   |
| provide name<br>/ location /<br>affiliation /          | Barry<br>Eakins   | NOAA NCEI  |  | barry.eakir | ns@noaa.gov       |
| affiliation /<br>email)                                | Scott<br>France   | University of Louisiana at Lafayette                                 |  |             | uisiana.edu       |

|  | David<br>Jourdan     | Nauticos  | dave@nauticos.com                           |
|--|----------------------|---|---|
|  | Chris Kelley         | University of Hawaii                                      | ckelley@hawaii.edu                          |
|  | Spencer<br>King      | Nauticos LLC  | spencerking@tampabay.rr.co<br>m             |
|  | Tina<br>Molodtsova   | P.P. Shirshov Institute of Oceanology<br>RAS              | tina@ocean.ru;<br>tina.molodtsova@gmail.com |
|  | Nicole<br>Morgan     | Florida State University                                  | nmorgan@fsu.edu                             |
|  | Jonathan<br>Parshall | Nauticos  | jonp@combinedfleet.com                      |
|  | Brendan<br>Roark     | Texas A&M University                                      | broark@geos.tamu.edu                        |
|  | Jonathan<br>Tree     | University of Hawaii Geology and<br>Geophysics Department | jtree@hawaii.edu                            |
|  | Michael<br>Vecchione | NMFS Systematics Lab                                      | vecchiom@si.edu                             |
|  | Katharine<br>Woodard | NOAA/NCEI   | katharine.woodard@noaa.go<br>v              |

## Purpose of the Dive

This dive, located on the northeast side of Necker Island (Mokumanamana), was carried out at the request of the Extended Continental Shelf (ECS) Program in order to obtain information that might be able to show a connection between Necker Island and Necker Ridge, a narrow feature that extends over 400 miles and protrudes past the current exclusive economic zone (EEZ) of the United States. The primary objective was to collect two geological samples on the ridge in order to determine whether these have the same composition as specimens that have previously been collected near Necker Island. Additionally, the ROV would survey biological communities in the area as time permits. The target start point of the dive was a flat surface located at a depth of 4302 m, which transitioned into a steep slope towards the northwest. The plan of the dive was for the ROV to survey up the steep slope moving northwest to a target depth of approximately 3900 m, collecting two geological samples along the way, as well as opportunistically documenting the biological community in the area.

## Description of the Dive:


The ROV landed on flat, heavily-sedimented surface at 4291 m with weak or nor current. The bottom had coalescent sediment cover with scattered Mn-encrusted boulders and pebbles. As the ROV moved northwestward towards the slope of the ridge, a few animals were observed including shrimps, sponges, sea anemones, sea cucumbers and two fish. At ~4280 m, the heavily-sedimented surface gave way to a more sloped surface with hardpan and isolated pillow lavas. Upon traversing up the slope near a depth of 4250 m, lobes of intact flows of pillow lavas were observed. A loose manganese-crusted volcanic rock was collected at 4245 m near the base of this flow. As the ROV moved up the slope, a couple of bamboo corals and anemones were observed. Sediment cover continued to decrease upslope to a surface composed of more Mn-encrusted pillow lavas with pockets of sediment. The Mn crusts appeared to be thin relative to the thicker crusts present on older (Cretaceous age) flows. This observation suggests that these flows are younger in age and were emplaced during


Hawaiian volcanic growth ~10 Myrs ago. A second manganese-crusted volcanic rock was collected at the based of another pillow lava flow very close to the end of the dive at 4222 m. Based on observation of the Mn crust thickness, both of the geological samples collected are hypothesized to be Hawaiian in origin and age. This suggests that the Necker Ridge continuation was not observed on the NE flank of the Necker complex. However, further analysis will be needed reveal the absolute age and composition of these samples to confirm this notion. The ROV left the bottom after a total bottom time of 2:18h.

## Animals observed during dive

| Phylum      | Group           | Species                    |
|-------------|-----------------|----------------------------|
| Anellida    | Polychaetes     | Polychaete? (on sponge)    |
| Arthropods  | Shrimp          | Unidentified red shrimp    |
| Arthropods  | Shrimp          | Aristopenaeus? sp.         |
| Cnidarians  | Actiniarians    | Unidentifed anemone        |
| Cnidarians  | Gorgonians      | Bathygorgia? sp.           |
| Cnidarians  | Gorgonians      | Chrysogorgia sp. planar    |
| Cnidarians  | Gorgonians      | Isidella? sp.              |
| Echinoderms | Holothurians    | Psychropotes? sp.          |
| Echinoderms | Holothurians    | Elpidiidae                 |
| Fishes      | Argentiniformes | Alepocephalidae?           |
| Fishes      | Aulopiformes    | Bathysaurus mollis?        |
| Mollusks    | Cephalopod      | Pink Octopus               |
| Sponges     | Hexactinellids  | Hyalonema (Corynonema) sp. |
|             |                 |                            |
|             |                 |                            |

Map of ROV Dive Area





| Comments                    | subangular |   |
|-----------------------------|------------|---|
| Please direct inquiries to: |            | NOAA Office of Ocean Exploration & Research<br>1315 East-West Highway (SSMC3 10 <sup>th</sup> Floor)<br>Silver Spring, MD 20910<br>(301) 734-1014 |