



---

VERIFICATION OF SEVERE LOCAL STORMS FORECASTS ISSUED BY THE  
NATIONAL SEVERE STORMS FORECAST CENTER: 1993

Richard W. Anthony  
National Severe Storms Forecast Center  
Kansas City, Missouri 64106-2877

June 1994

---

U.S. DEPARTMENT OF  
COMMERCE

/ National Oceanic and  
Atmospheric Administration

/ National Weather  
Service

National Weather Service  
National Severe Storms Forecast Center

The National Severe Storms Forecast Center (NSSFC) has the responsibility for the issuance of tornado and severe thunderstorm watches for the contiguous 48 states. Watches are issued for those areas where thunderstorms are forecast to produce one or more of the following: (1) hailstones of 3/4 inch diameter or greater, (2) surface wind gusts of 50 knots or greater, or (3) tornadoes.

NOAA Technical Memoranda in the NWS, NSSFC subseries are produced under the technical guidance of the NSSFC, Techniques Development Unit. They facilitate rapid dissemination of material of general interest in the field of severe storm meteorology. These papers may be preliminary in nature, and may be formally published elsewhere at a later date.

These papers are available from the National Technical Information Service (NTIS), U.S. Department of Commerce, Sills Building, 5285 Port Royal Road, Springfield, Virginia 22161. Price varies, \$3.50 for microfiche.

Previous issued in this series:

- No. 1      New Severe Thunderstorm Radar Identification Techniques and Warning Criteria: A Preliminary Report. Leslie R. Lemon, July 1977, 60 p., (PB 273049).
- No. 2      A Subjective Assessment of Model Initial Conditions Using Satellite Imagery. John E. Hales, Jr., November 1978, 19 p., (PB 291593).
- No. 3      Severe Thunderstorm Radar Identification Techniques and Warning Criteria. Leslie R. Lemon, April 1980, 60 p., (PB 231409).
- No. 4      The Enhanced-V, A Satellite Observable Severe Storm Signature. Donald W. McCann, March 1981, 31 p., (PB 230336).
- No. 5      The Operational Meteorology of Convective Weather Volume I: Operational Mesoanalysis. Charles A. Doswell III, November 1982, 160 p., (PB83 162321).
- No. 6      Severe Local Storm Warning and Event Summaries Available in AFOS. Preston W. Leftwich, Jr. and Lawrence C. Lee, January 1984, 10 p., (PB84 150291).
- No. 7      Severe Thunderstorm Cases of 1984. John E. Hales, Jr. and Hugh G. Crowther, May 1985, 88 p., (PB85 210748/AS).

VERIFICATION OF SEVERE LOCAL STORMS FORECASTS ISSUED BY THE  
NATIONAL SEVERE STORMS FORECAST CENTER: 1993

Richard W. Anthony  
National Severe Storms Forecast Center  
Kansas City, Missouri 64106-2877

June 1994

UNITED STATES  
DEPARTMENT OF COMMERCE  
Robert A. Mosbacher  
Secretary

National Oceanic and  
Atmospheric Administration  
John A. Knauss  
Under Secretary

National Weather  
Service  
Elbert W. Friday, Jr.  
Assistant Administrator



## TABLE OF CONTENTS

	Page No.
Abstract . . . . .	1
1. Introduction . . . . .	1
2. Verification Procedures . . . . .	2
3. Convective Outlooks . . . . .	3
4. Severe Local Storm Watches . . . . .	4
5. Summary . . . . .	8
6. Acknowledgements . . . . .	8
7. References . . . . .	9

VERIFICATION OF SEVERE LOCAL STORM FORECASTS ISSUED BY THE  
NATIONAL SEVERE STORMS FORECAST CENTER: 1993

by

Richard W. Anthony  
National Severe Storms Forecast Center  
Kansas City, Missouri 64106

ABSTRACT. The SELS Unit of the National Severe Storms Forecast Center routinely issues convective outlooks and severe local storm watches to delineate areas that are favorable for development of severe local storms. This report summarizes verification of those forecasts that were issued during 1993.

The threat of severe local storms was identified well via the second-day outlooks. Prediction of the location of subsequent events improved in the first-day outlooks. During 1993 the SELS Unit issued 957 severe local storm watches. Of these, 350 were tornado watches. The total of 957 was the most watches issued in one year, exceeding the previous high of 940 in 1992. There were 13986 severe local storm events during 1993. Of these, 1173 were tornadoes. In terms of the total number of events, this was the most active season on record.

## 1. INTRODUCTION

The Severe Local Storms (SELS) Unit of the National Severe Storms Forecast Center (NSSFC) has responsibility for issuing convective outlooks, tornado watches and severe thunderstorm watches for the contiguous United States. Convective outlooks, which depict expected areas and densities of severe local storms in a preliminary sense, are issued daily at 0700, 1500, and 1900 UTC. Forecast periods begin at 1200, 1500, and 1900 UTC, respectively, and continue until 1200 UTC the following day. Although convective outlooks include forecasts for all thunderstorms, only forecasts of severe local storms are addressed in this report. An initial second-day severe local storm outlook is issued at 0800 UTC and updated at 1800 UTC each day. Both outlooks are valid for the 24-hr period beginning at 1200 UTC the next day. Tornado and severe thunderstorm watches are issued, as needed, to delineate areas in which conditions will become favorable for severe local storms to occur within the next few hours.

Various forms of verification of products issued by the SELS Unit have long been an important aspect of operations. Such data aid the assessment of forecast quality, identify possible areas of improvement, and provide helpful feedback concerning the progression of events during specific severe local storm episodes.

Results of these earlier verification efforts have been published by Galway (1967), Galway (1975) and Pearson and Weiss (1979) and Anthony and Leftwich (1992a). In 1982, the National Weather Service (NWS) formulated a National Verification Plan (NWS 1982) to provide guidelines for verification of the various products. As an integral part of the national program, the NSSFC publishes verification summaries each year. For example, Anthony (1993) summarized verification of watches and outlooks issued by the NSSFC during 1992. This report documents verification of severe local storm watches and convective outlooks issued during 1993.

## 2. VERIFICATION PROCEDURES

The first step in verification of any forecast is collection of both the issued forecast messages and reports of events that occurred during the forecast period. Collection of watch and outlook messages is accomplished in real-time as they are disseminated at the NSSFC. They are automatically encoded for processing via electronic computer. To qualify as a valid severe local storm event that is used in watch and convective outlook verification, reports must satisfy one of the criteria listed in Table 1. Although severe local storm reports are collected at the NSSFC as a part of operational procedures, reports of tornadoes, large hail and thunderstorm wind events are not retained for use in official (final) verification. All other reports are obtained via "Storm Data and Unusual Weather Phenomena". This monthly report, also known as Form F-8, is prepared for each state by the various NWS Forecast Offices. Further restrictions are applied before any event is included in the verification procedure. Multiple reports of hail and/or wind occurring within both 10 statute miles and 15 minutes of each other, and in the same county, are considered as one event. However, all tornado reports appearing in the F-8 summary are included as separate events. Additional information regarding processing of severe local storm reports is given by Grenier and Halmstad (1986).

Table 1  
Criteria for severe local storm events

- 
- a. Tornado - a rotating circulation touching the ground and associated with a convective cloud.
  - b. Hail equal to or greater than 3/4 inch (1.9 cm) in diameter.
  - c. Thunderstorm wind gust of at least 50 knots (93 km/h).
  - d. Thunderstorm wind damage.
  - e. Extreme turbulence (reported by aircraft) associated with a thunderstorm.
- 

Once data have been compiled, various verification statistics are computed. Primary statistics are Percent Verified (PV), Probability of Detection (POD), False Alarm Ratio (FAR) and Critical Success Index (CSI). The latter three statistics were adapted from those described by Donaldson et al. (1975). Adaptations were necessary because the statistics described by Donaldson et al. considered point forecasts, whereas both watches and outlooks are area forecasts. Modifications and the computational procedures that are currently followed are discussed in detail by Weiss et al. (1980).

### 3. CONVECTIVE OUTLOOKS

During 1993 a total of 719 convective outlooks specified a threat of severe local storms for the current, or first, day. Figure 1 shows overall convective outlook verification statistics for the ten-year period 1984-1993. Verification statistics for 1993, stratified by the three issuance times, are given in Section A of Table 2. The two later outlooks, which are based on additional diagnostic analysis and later National Meteorological Center (NMC) numerical model guidance, contained progressively higher percentages of reports (higher POD). Increases in CSI were also observed for these later outlooks.




Figure 1. Convective outlook verification for the years 1984-1993.

There were 221 initial (issued at 0800 UTC) second-day outlooks that specified a threat of severe local storms for the following day. A threat of severe local storms was initiated or retained in 230 updated outlooks issued at 1800 UTC. Verification of these outlooks is given in Section B of Table 2. Updates improved the initial outlooks in terms of POD.

Comparison of second-day outlooks with first-day outlooks shows noteworthy improvement in values of POD with little change in FAR. This results in higher values of CSI for successive outlooks. Improvement in POD and little change in FAR results from second-day outlooks being, on the average, smaller in area than first-day outlooks. Second-day outlooks are based almost entirely on indications derived from numerical model guidance, which emphasizes synoptic-scale features. For the first-day outlooks important information is obtained from mesoscale analysis as well as later runs of numerical guidance. Generally, a larger area of threat is evident in mesoscale analyses than in numerical guidance alone. Also, such subsynoptic-scale analysis leads to better placement of outlook areas. Increased area and better placement of outlook areas combine to produce similar values of FAR and higher values of POD and CSI.



Table 2

Verification scores for severe local storm outlooks during 1993

## A. First-Day

Issue Time (UTC)	Number Issued	POD	FAR	CSI
0700	238	.65	.55	.35
1500	242	.71	.53	.38
1900	239	.73	.51	.41
All	719	.69	.53	.38

## B. Second-Day

Issue Time (UTC)	Number Issued	POD	FAR	CSI
0800	221	.49	.66	.25
1800	230	.57	.69	.25
All	451	.53	.68	.25

## 4. SEVERE LOCAL STORM WATCHES

During 1993 the SELS Unit issued 957 severe local storm watches. Of these, 350 were tornado watches and 607 were severe thunderstorm watches. The total of 957 was the most watches issued in one year, exceeding the previous high of 940 in 1992. There were 13986 severe local storm events during 1993. Of these, 1173 were tornadoes. In terms of the total number of events, this was the most active season on record (Grenier and Halmstad 1992) and the second most active for total number of tornadoes. Also, the mean area contained in severe local storm watches in 1993 was 22856 square nautical miles, the smallest on record.

Issuance of either a tornado watch or a severe thunderstorm watch denotes the threat of severe local storms. Analysis of the overall forecasting of severe local storms is obtained via verification of all watches and all severe local storm events considered together. Trends in some of these statistics for the period 1984-1993 are shown in Figure 2.

Issuance of tornado watches emphasizes the threat of tornadoes. To assess this aspect of tornado watches separately, additional verification of tornado watches is done considering tornadoes only. That is, a tornado watch is considered to be verified only when a tornado occurs during the valid time period and within the watch area. Statistics computed in this manner are shown in Figure 3 for the period 1984-1993.

As shown by Table 3, skill was shown during 1993 in distinguishing severe local storm situations which have a greater threat of tornadoes from situations when tornado development is not likely. While at least one tornado occurred in 50% of the tornado watches, only 19% of the severe thunderstorm watches contained tornadoes. Tornado watches also reflected an increased threat of severe local storms as only 13% of them did not contain any reported events.




Figure 2. Verification for all severe local storm watches issued during the period 1984-1993.

Values of POD for various intensities of tornadoes during 1993 relative to valid severe local storm watches are given in Table 4. A tornado is considered to have occurred within a valid watch if any portion of its track occurred within a valid watch area. Intensities of tornadoes are indicated by F-scale (Fujita 1981) values ranging from "0" (weakest) to "5" (most violent). During 1993, 60% of the weak (F0-F1) tornadoes occurred in valid watches, while 83% of the violent (F4-F5) tornadoes occurred in valid watches. For strong and violent tornadoes, which caused 79% of the tornado-related fatalities during 1993, the Probability of Detection was 0.75. The POD for all tornadoes was 0.62.




Figure 3. Verification of tornado watches via tornadoes only for the period 1984-1993.

Table 3

Frequencies of severe local storms in valid watches, 1993

<u>Watch Type</u>	<u>Number Issued</u>	<u>Observed Tornado</u>	<u>Only Other Severe Types</u>	<u>No Severe</u>
Tornado	350	50%	37%	13%
Severe Thunderstorm	607	19%	61%	20%

Table 4

Probability of Detection (POD) for various intensities (F-scale) of tornadoes relative to valid severe local storm watches during 1993

	<u>Weak (F0-F1)</u>	<u>Strong (F2-F3)</u>	<u>Violent (F4-F5)</u>	<u>Strong/Violent (F2-F5)</u>	<u>All (F0-F5)</u>
Total	1058	109	6	115	1173
In Watch	638	81	5	86	724
POD	0.60	0.74	0.83	0.75	0.62

Strong and violent (F2-F5) tornadoes consistently pose an increased threat to life and property. Values of POD for all, weak (F0-F1) and strong/violent (F2-F5) tornadoes for the period 1984-1993 are shown in Figure 4. The closeness of the POD for all tornadoes to the POD for weak tornadoes reflects the high percentage of weak tornadoes. POD values for strong/violent tornadoes are consistently higher than values for weak tornadoes. Results for the past several years have been consistent with the earlier findings of Ostby and Higginbotham (1982).




Figure 4. Annual values of Probability of Detection (POD) for weak, strong/violent, and all tornadoes, respectively, for the years 1984-1993.

During 1993, there were 16 tornadoes that caused at least one fatality. Twelve of the 16 "killer" tornadoes occurred within valid severe local storm watches. Further statistics concerning severe local storm events resulting in fatalities and injuries are given in Table 5. During 1993, tornadoes caused 33 fatalities in the United States. Severe thunderstorm winds caused 22 additional fatalities. Twenty-two (67%) tornado related fatalities occurred within valid watches. Twelve (55%) of the 22 fatalities resulting from thunderstorm wind gusts occurred within valid watches. Valid severe local storm watches contained 60% of tornado-related injuries and 76% of injuries attributable to other severe thunderstorms.

Table 5

## Severe local storm-related fatalities and injuries in 1993

	<u>Tornado</u>		<u>Severe Thunderstorm</u>	
	<u>Fatalities</u>	<u>Injuries</u>	<u>Fatalities</u>	<u>Injuries</u>
Total	33	848	22	479
Occurring within valid watches	22	508	12	365
% Within Watches	67	60	55	76

## 5. SUMMARY

Severe local storm outlooks and watches are issued by the SELS Unit of the NSSFC. These products identify areas in which development of severe thunderstorms and/or tornadoes is likely. Verification of these products is done routinely for purposes of feedback to the forecasters and quality control of issued products.

In 1993 there was some improvement in FAR between second-day and first-day severe local storm outlooks. However, the location of severe local storm occurrence (as reflected in the POD) was better predicted by the first-day outlooks.

For severe local storm watches, ability was exhibited in distinguishing the added threat of tornadoes from that of hail or thunderstorm wind gusts. During 1993, 67% of the tornado-related fatalities and 60% of the tornado-related injuries occurred within valid severe local storm watches.

## 6. ACKNOWLEDGEMENTS

The author thanks Janalee Snyder for her technical assistance in preparation of this report.

## 7. REFERENCES

- Anthony, R. W., 1993: Verification of Severe Local Storm Forecasts Issued by the National Severe Storms Forecast Center: 1992. NOAA Technical Memorandum NWS NSSFC-35, National Severe Storms Forecast Center, Kansas City, Missouri, 11 pp.
- Anthony, R. W. and P. W. Leftwich, 1992a: Trends in Severe Local Storm Watch Verification at the National Severe Storms Forecast Center. Wea. Forecasting, 7, 613-622.
- Donaldson, R.J., R.M. Dyer and M.J. Kraus, 1975: An objective evaluator of techniques for predicting severe weather events. Preprints, Ninth Conference on Severe Local Storms (Norman, OK), Amer. Meteor. Soc., Boston, MA, 395-402.
- Fujita, T.T., 1981: Tornadoes and downbursts in the context of generalized planetary scales. J. Atmos. Sci., 38, 1511-1534.
- Galway, J.G., 1967: SELS forecast verification 1952-1966. Preprints, Fifth Conference on Severe Local Storms (St. Louis, MO), Amer. Meteor. Soc., Boston, MA., 140-145.
- \_\_\_\_\_, 1975: Relationship of tornado deaths to severe weather watch areas. Mon. Wea. Rev., 103, 737-741.
- Grenier, L.A. and J.T. Halmstad, 1986: Severe Local Storm Warning Verification Preliminary Procedures. NOAA Technical Memorandum NWS NSSFC-12, National Severe Storms Forecast Center, Kansas City, Missouri, 10 pp.
- \_\_\_\_\_ and \_\_\_\_\_, 1992: Severe Local Storm Warning Verification: 1991. NOAA Technical Memorandum NWS NSSFC-33, National Severe Storms Forecast Center Kansas City, Missouri, 28 pp.
- National Weather Service, 1982: National Verification Plan. U.S. Dept. of Commerce, NOAA, 81 pp.
- Ostby, F. P., and D.M. Higginbotham, 1982: Tornado predictability as a function of geography and intensity. Preprints, Twelfth Conference on Severe Local Storms (San Antonio, TX), Amer. Meteor. Soc., Boston, MA., 176-179.
- Pearson, A.D. and S.J. Weiss, 1979: Some trends in forecast skill at the National Severe Storms Forecast Center. Bull. Amer. Meteor. Soc., 60, 319-326.

Weiss, S.J., D.L. Kelly and J.T. Schaefer, 1980: New objective verification techniques at the National Severe Storms Forecast Center. Preprints, Eighth Conference on Weather Forecasting and Analysis (Denver, CO), Amer. Meteor. Soc., Boston, MA., 140-145.

- No. 8      A Minimum Assumption Tornado Hazard Probability Model. Joseph T. Schaefer, Donald L. Kelly, and Robert F. Abbey, May 1985, 30 p., (PB85 20692/AS).
- No. 9      Verification of Severe Local Storm Forecasts Issued by the National Severe Storms Forecast Center: 1984. Preston W. Leftwich, Jr., November 1985, 23 p., (PB86 128105/AS).
- No. 10     Severe Local Storm Warning Verification: 1984. Preston W. Leftwich, Jr. and Leo A. Grenier, December 1985, 14 p., (PB86 148244).
- No. 11     Severe Thunderstorm Cases of 1985. John E. Hales, Jr. and Hugh G. Crowther, February 1986, 51 p., (PB86 164340/AS).
- No. 12     Severe Local Storm Warning Verification Preliminary Procedures. Leo A. Grenier and John T. Halmstad, April 1986, 16 p., (PB86 194362).
- No. 13     Verification of Severe Local Storm Forecasts Issued by the National Severe Storms Forecast Center: 1985. Preston W. Leftwich, Jr., November 1986, 9 p., (PB87 137139/AS).
- No. 14     Severe Local Storm Warning Verification: 1985. Preston W. Leftwich, Jr. and Leo A. Grenier, December 1986, 16 p., (PB87 137147/AS).
- No. 15     An Examination of the National Weather Service Severe Local Storm Warning Program and Proposed Improvements. John E. Hales, Jr., January 1987, 32 p., (PB87 147948/AS).
- No. 16     Severe Thunderstorm Cases of July 1985 through June 1986. John E. Hales, Jr. and Hugh G. Crowther, February 1987, 72 p., (PB87 163911/AS).
- No. 17     Severe Local Storm Warning Verification: 1986. Leo A. Grenier, John T. Halmstad and Preston W. Leftwich, Jr., June 1987, 19 p., (PB87 195939).
- No. 18     Verification of Severe Local Storm Forecasts Issued by the National Severe Storms Forecast Center: 1986. Preston W. Leftwich, Jr., September 1987, 9 p., (PB88 101407).
- No. 19     Severe Thunderstorm Cases of July 1986 through June 1987. John E. Hales, Jr. and Hugh G. Crowther, April 1988, 83 p., (PB88 214085).



- No. 20      Severe Local Storm Warning Verification: 1987. Leo A. Grenier, John T. Halmstad and Preston W. Leftwich, Jr., June 1988, 19 p., (PB88 241393).
- No. 21      Verification of Severe Local Storm Forecasts Issued by the National Severe Storms Forecast Center: 1987. Preston W. Leftwich, Jr., December 1988, 11 p., (PB89 159719/AS).
- No. 22      Severe Thunderstorm Cases of July 1987 thru June 1988. John E. Hales, Jr. and Hugh G. Crowther, April 1989, 92 p., (PB89 206411/AS)
- No. 23      Severe Local Storm Warning Verification: 1988. Leo A. Grenier, John T. Halmstad and Preston W. Leftwich, Jr., June 1989, 26 p., (PB89 226310/AS).
- No. 24      Verification of Severe Local Storm Forecasts Issued by the National Severe Storms Forecast Center: 1988. Preston W. Leftwich, Jr., September 1989, 18 p., (PB90-140211/AS).
- No. 25      A Dyad of Papers Concerning Joint Verification of Severe Local Storm Watches and Warnings During Tornado Events: Preston W. Leftwich, Jr. and John E. Hales, Jr., January 1990, 36 p., (PB90-219387/AS).
- No. 26      Severe Thunderstorm Cases of July 1988 thru June 1989: John E. Hales, Jr. and Hugh G. Crowther, April 1990, 106 p., (PB90-226424/AS)
- No. 27      Severe Local Storm Warning Verification: 1989. Leo A. Grenier, John T. Halmstad and Preston W. Leftwich, Jr., May 1990, 25 p., (PB90-248592)
- No. 28      Verification of Severe Storms Forecasts Issued by the National Severe Storms Forecast Center: 1989. Preston W. Leftwich, Jr. and Richard W. Anthony, September 1990, 17 p., (PB91-132472/AS)
- No. 29      Severe Thunderstorm Cases of July 1989 thru June 1990: John E. Hales, Jr. and Hugh G. Crowther, April 1991, 131 p., (PB91-201392)
- No. 30      Severe Local Storm Warning Verification: 1990. Leo A. Grenier, John T. Halmstad, May 1991, 32 p., (PB91-227520)
- No. 31      Verification of Severe Local Storm Forecasts Issued By the National Severe Storms Forecast Center: 1990. Preston W. Leftwich, Jr. and Richard W. Anthony, October 1991, 12 p., (PB92-124452/AS)

- No. 32      Severe Thunderstorm Cases of July 1990 thru June 1991:  
John E. Hales, Jr. and Hugh G. Crowther, February 1992,  
159 p., (PB92-158336/AS)
- No. 33      Severe Local Storm Warning Verification: 1991. Leo A.  
Grenier, John T. Halmstad, April 1992, 28 p., (PB92-  
185982/AS)
- No. 34      Verification of Severe Local Storm Forecasts Issued By  
the National Severe Storms Forecast Center: 1991.  
Richard W. Anthony and Preston W. Leftwich, Jr., May  
1992, 12 p., (PB92-186063)
- No. 35      Verification of Severe Local Storms Forecasts Issued By  
the National Severe Storms Forecast Center: 1992.  
Richard W. Anthony, June 1993, 14 p., (PB93-216844)
- No. 36      Severe Thunderstorm Cases of July 1991 thru June 1992:  
John E. Hales, Jr. and Kevin L. Polston, June 1993,  
200 p., (PB93-227460)
- No. 37      Severe Local Storm Warning Verification: 1992. Hugh  
C. Crowther, and John T. Halmstad, May 1993, 27 p.,  
(PB93-227452)