

2019 UPDATE TO “AN IMPACT ASSESSMENT OF GREAT LAKES AQUATIC NONINDIGENOUS SPECIES”

El Lower¹, Nicholas Boucher², Rochelle Sturtevant¹, Ashley Elgin³

¹ Michigan Sea Grant

² Cooperative Institute for Great Lake Lakes Research, University of Michigan

³ NOAA Great Lakes Environmental Research Laboratory

NOAA Great Lakes Environmental Research Laboratory
4840 S. State Road, Ann Arbor, Michigan

Wednesday, July 08, 2020



UNITED STATES
DEPARTMENT OF COMMERCE

Wilbur L. Ross, Jr.
Secretary

NATIONAL OCEANIC AND
ATMOSPHERIC ADMINISTRATION

Dr. Neil Jacobs,
Acting Administrator

NOTICE

Mention of a commercial company or product does not constitute an endorsement by NOAA. Use of information from this publication concerning proprietary products or the tests of such products for publicity or advertising purposes is not authorized. This is GLERL Contribution No. xxxx. Funding was awarded to Cooperative Institute for Great Lakes Research (CI GLR) through the NOAA Cooperative Agreement with the University of Michigan (NA17OAR4320152). This CI GLR contribution number is xxxx. This publication is available as a PDF file and can be downloaded from GLERL's web site: www.glerl.noaa.gov or by emailing GLERL Information Services at oar.pubs.glerl@noaa.gov.

TABLE OF CONTENTS

1.0 Summary	1
2.0 Addenda	4
3.0 Risk Assessments	5
Scientific Name: <i>Cercopagis pengoi</i>	5
Organism Impact Assessment	5
Scientific Name: <i>Ctenopharyngodon idella</i>	7
Organism Impact Assessment	7
Scientific Name: <i>Daphnia galeata galeata</i>	13
Organism Impact Assessment	13
Scientific Name: <i>Hemimysis anomala</i>	16
Organism Impact Assessment	16
Scientific Name: <i>Lepomis humilis</i>	19
Organism Impact Assessment	19
Scientific Name: <i>Oncorhynchus gorbuscha</i>	21
Organism Impact Assessment	21
Scientific Name: <i>Oncorhynchus tshawytscha</i>	24
Organism Impact Assessment	24
Scientific Name: <i>Schizopera borutzkyi</i>	27
Organism Impact Assessment	27
4.0 References	30

LIST OF TABLES

Table 1. Summary of impact assessment results by taxonomic group. For each impact category (i.e. environmental, socio-economic, beneficial), the number of species whose impact was assessed as high (H), moderate (M), low (L), or unknown (U) is given. Note: “Arthropods” refers to non-crustacean arthropods. Relative to Sturtevant et al. (2014), “+” indicates an increase in the number of species in the category, while “-” indicates a decrease.	3
Table 2. New species and major changes to the assessments, etc. originally published in Sturtevant et al. (2014).	4
Table 3. Changes and additions to Tables 2-11 in Sturtevant et al. (2014).	4

NOAA TECHNICAL MEMORANDUM GLERL-161C

2019 UPDATE TO “AN IMPACT ASSESSMENT OF GREAT LAKES AQUATIC NONINDIGENOUS SPECIES”

El Lower, Nicholas Boucher, Rochelle Sturtevant, Ashley Elgin

1.0 SUMMARY

This report includes all major updates to the earlier Risk Assessments on nonindigenous species conducted by the GLANSIS project during the 2019 calendar year. All new assessments were conducted following the same methods outlined in the original technical memorandum, [NOAA Technical Memorandum GLERL-161](#) “An impact assessment of Great Lakes aquatic nonindigenous species” (Sturtevant et al, 2014). All re-assessments are based on new literature surveys using the original as a baseline and conducted to the same methods. All assessments were reviewed by members of the GLANSIS Team (according to expertise) and by select external reviewers. Results of each risk assessment are incorporated into the species profiles found on the GLANSIS website (www.glerl.noaa.gov/glansis/).

To be included in the GLANSIS nonindigenous list, the species in question must meet a particular set of criteria:

1. Records of the species appeared suddenly and had not been recorded in the basin previously;
2. It subsequently spreads within the basin.
3. Its distribution in the basin is restricted compared with native species.
4. Its global distribution is anomalously disjunct (meaning it contains widely scattered and isolated populations).
5. Its global distribution is associated with human vectors of dispersal.
6. The basin is isolated from regions possessing the most genetically and morphologically similar species.

Additionally, to be listed on the nonindigenous list rather than the GLANSIS Watchlist, a species must have a reproducing population within the basin that is capable of overwintering, as inferred from multiple discoveries of adult and juvenile life stages over at least two consecutive years.

A total of 82 species were reassessed in 2019, with 10 of those species undergoing changes detailed in the following sections of this document.

Grass carp (*Ctenopharyngodon idella*) was listed in [NOAA Technical Memorandum GLERL -169b](#) (Lower et al, 2019) as a watchlist species, but has since been moved to the nonindigenous list. Grass carp have been periodically found in the Great Lakes since the 1970s as a result of stocking triploids for control of aquatic plants, but the first diploid was not confirmed until 2011. Populations of diploid grass carp have been found in the Maumee and Sandusky Rivers, and are both overwintering and reproducing (producing viable eggs) in these waterways, meeting the criteria for inclusion on our list of nonindigenous species present within the Great Lakes basin. As such, *Ctenopharyngodon idella* was re-assessed using the methods for this list, and its new organism impact assessment score is included here in full.

Northern bur-reed (*Sparganium glomeratum*) was listed in Sturtevant et al. (2014) as a nonindigenous species and is now considered to be native to portions of Minnesota proximate to the Great Lakes basin, based on the most up-to-date species distribution maps. As such, this species is now considered a range expander rather than nonindigenous. It is retained in the GLANSIS system, but designated as a range expander and will no longer be included in updates to this tech memo series.

Sweetscent (*Pluchea odorata odorata* and *Pluchea odorata succulent*) were previously listed separately in GLANSIS and had separate assessments in Sturtevant et al. (2014) but are now listed together under *Pluchea odorata*. These two subspecies are both nonindigenous to the Great Lakes, but do not differ significantly in their ecology, habitat, niche, or impact within the Great Lakes, so they have been combined as a single species in this assessment.

The impact assessment scores for the other seven species were updated upon reassessment based on review of the more recent literature. Five species previously designated as ‘current research is inadequate to support proper assessment’ for one or more portions of the impact assessment were able to be scored in the reanalysis. The environmental impact score for chinook salmon (*Oncorhynchus tshawytscha*) was changed from moderate to high based on additional literature highlighting its impacts on native species, including direct competition with lake trout and other native species and top-down effects on forage fish (Bunnell et al. 2014), as well as having served as a vector for the disease *Renibacterium salmoninarum*, which is now considered to be widespread in the Great Lakes (GLFHC 2015). The socioeconomic impact score for fishhook waterflea (*Cercopagis pengoi*) was changed from low to high based on literature detailing the annual economic cost of damages caused by and mitigation of *C. pengoi* at \$5 million USD (Pimentel et al., 2005).

Two species were re-named in this assessment. The quagga mussel (*Dreissena bugensis*) was renamed *Dreissena rostriformis bugensis*, which is the preferred scientific name according to the CABI Invasive Species Compendium. Spotted ladysthumb (*Polygonum persicaria*) is now listed as its alternate synonym *Persicaria maculosa*.

Seventy-two additional species were re-assessed during this period, but the qualitative impact assessment categories (high, medium, low, or unknown for environmental, socio-economic, or beneficial impacts) did not change for any of these additional species.

An updated version of Table 1 from Sturtevant et al. (2019) is presented on the next page. General changes to the table include a net overall loss of one species from the nonindigenous list based on addition of one fish species (*Ctenopharyngodon idella*), the reclassification of one plant as a range expander (*Sparganium glomeratum*), and the collapse of the two *Pluchea odorata* subspecies into a single entry.

Table 1. Summary of impact assessment results by taxonomic group. For each impact category (i.e. environmental, socio-economic, beneficial), the number of species whose impact was assessed as high (H), moderate (M), low (L), or unknown (U) is given. Note: “Arthropods” refers to non-crustacean arthropods. Relative to Sturtevant et al. (2014), “+” indicates an increase in the number of species in the category, while “-” indicates a decrease.

Taxon	Environmental				Socio-Economic				Beneficial			
	H	M	L	U	H	M	L	U	H	M	L	U
Fishes (n=28) (+1)	8	5	0(-)	14	3	1	19(-)	4	8	2	12(-)	5
Annelids (n=6)	0	0	0	6	0	0	6	0	0	0	5	1
Arthropods (n=2)	0	0	0	2	0	0	2	0	0	1	1	0
Bryozoans (n=1)	0	0	0	1	0	0	0	1	0	0	1	0
Coelenterates (n=2)	0	0	0	2	0	0	1	1	0	0	2	0
Crustaceans (n=24)	2	2(+)	0	20(+)	0	1(+)	21(+)	2	0	1(+)	20(+)	3(+)
Mollusks (n=18)	3	2	1	12	2	2	11	3	0	0	16	2
Plants (n=55) (-2)	6(+)	20(+)	3	28	4	9(+)	41(+)	3	4	15(+)	33	5
Algae (n=27)	0	4	20	3	1	3	23	0	0	1	26	0
Amoebae (n=3)	0	0	0	3	0	0	3	0	0	0	3	0
Parasites and Diseases (n=20)	7	1	12	0	2	0	18	0	0	0	20	0
Total (n=186) (-1)	26(+)	34(+)	36(-)	91(+)	12	16(+)	145(+)	14	12	20(+)	139(+)	16(+)

In addition, none of the summary statements in the original Sturtevant et al. (2014) have substantively changed, and they are as follows:

1. Additional research is still needed to understand the environmental impacts of nonindigenous species. The state of knowledge is inadequate to assess the environmental impact for nearly half (now 48% instead of 49%) of the established species.
2. At least 32% (previously 31%) of the nonindigenous species found in the Great Lakes have significant (moderate to high) environmental impact. If the 91 species for which the state of scientific knowledge is insufficient to complete the assessment of environmental impact follow the trends of the assessed species this number will be closer to 50%. References in the literature (e.g., Williamson and Fitter 1996) and popular media of approximately 10% of non-native species becoming invasive is a severe underestimate for the Great Lakes.
3. We estimate between 14 and 16% of the nonindigenous species found in the Great Lakes have moderate to high socioeconomic impact.

Of the 32 species assessed as having significant (moderate to high) benefits, only one – *Puccinellia distans* – is still assessed as having low environmental and socioeconomic impact.

2.0 ADDENDA

Table 2. New species and major changes to the assessments, etc. originally published in Sturtevant et al. (2014).

Species	Addenda	Author, date added
<i>Cercopagis pengoi</i>	Socio-economic impacts changed from low to high.	Boucher, 2019
<i>Ctenopharyngodon idella</i>	Status changed from watchlist to nonindigenous list.	Boucher, 2019
<i>Daphnia galeata galeata</i>	Environmental impacts changed from unknown to high.	Boucher, 2019
<i>Dreissena rostriformis bugensis</i>	Species name change, scores unchanged from TM-161.	Sturtevant and Elgin, 2019
<i>Hemimysis anomala</i>	Environmental impacts changed from unknown to moderate	Boucher, 2019
<i>Lepomis humilis</i>	Beneficial impacts changed from unknown to low	Boucher, 2019
<i>Oncorhynchus gorbuscha</i>	Environmental impact changed from unknown to moderate.	Boucher and Lower, 2019
<i>Oncorhynchus tshawytscha</i>	Environmental impact changed from moderate to high.	Boucher and Lower, 2019
<i>Pluchea odorata</i>	Newly combined profile, scores unchanged.	Lower, 2019
<i>Pluchea odorata odorata</i>	**subspecies combined in one profile (<i>Pluchea odorata</i>)	Lower, 2019
<i>Pluchea odorata succulenta</i>	**subspecies combined in one profile (<i>Pluchea odorata</i>)	Lower, 2019
<i>Persicaria maculosa</i>	Species name changed from <i>Polygonum persicaria</i> , scores unchanged.	Lower, 2019
<i>Schizopera borutzkyi</i>	Environmental impacts changed from unknown to low.	Boucher, 2019

Table 3. Changes and additions to Tables 2-11 in Sturtevant et al. (2014).

Scientific Name	Common Name	Family	Environmental Impact		Socio-Economic Impact		Beneficial Impact	
			Score	# Unknown	Score	# Unknown	Score	# Unknown
<i>Cercopagis pengoi</i>	Fishhook waterflea	Cercopagidae	7	2	7	0	0	1
			High		High		Low	
<i>Ctenopharyngodon idella</i>	Grass carp	Cyprinidae	20	0	5	0	1	0
			High		Moderate		Low	
<i>Daphnia galeata galeata</i>	A waterflea	Daphniidae	7	0	0	0	0	0
			High		Low		Low	
<i>Hemimysis anomala</i>	Bloody red shrimp	Mysidae	3	2	0	1	1	0
			Moderate		Low		Low	
<i>Lepomis humilis</i>	Orange-spotted sunfish	Centrarchidae	0	2	0	1	0	0
			Unknown		Low		Low	
<i>Oncorhynchus gorbuscha</i>	Pink salmon	Salmonidae	3	2	1	0	2	0
			Moderate		Low		Moderate	
<i>Oncorhynchus tshawytscha</i>	Chinook salmon	Salmonidae	13	1	0	0	13	1
			High		Low		High	
<i>Pluchea odorata</i>	Sweetscent	Asteraceae	1	4	0	0	1	0
			Unknown		Low		Low	
<i>Schizopera borutzkyi</i>	An oarsman	Diosaccidae	0	1	0	0	0	1
			Low		Low		Low	

3.0 RISK ASSESSMENTS

Scientific Name: *Cercopagis pengoi*

Common Name: Fishhook waterflea

Organism Impact Assessment

IMPACT RESULTS

Environmental: High

Socio-Economic: High

Beneficial: Low

Comments: Socioeconomic impact score changed from low to high. Other scores unchanged.

SOCIO-ECONOMIC IMPACT

NOTE: In this section, a “Not significantly” response should be selected if there have been no reports of a particular impact. An “Unknown” response is appropriate if the potential for a particular impact might be inferred from a significant environmental impact but has not been explicitly reported or if there is an unresolved debate about a particular impact.

Does the species pose some hazard or threat to human health (e.g., it magnifies toxin levels, is poisonous, a virus, bacteria, parasite, or a vector of one)

Yes, significant effects on human health have already been observed	6
Yes, but negative consequences have not been widespread, long lasting, or severe AND/OR It has significantly affected human health in past invasions outside of the Great Lakes	1
Not significantly	0 ✓
Unknown	U

- *Not reported.*

Does it cause damage to infrastructure (such as water intakes, pipes, or any other industrial or recreational infrastructure)?

Yes, it is known to cause significant damage	6
Yes, but the costs have been small and are largely repairable or preventable AND/OR It has a history of causing significant infrastructural damage in past invasions outside of the Great Lakes	1
Not significantly	0 ✓
Unknown	U

- *Not reported.*

Does it negatively affect water quality?

Yes, it has significantly affected water quality, and is costly or difficult to reverse	6
Yes, but the effects are negligible and/or easily reversed AND/OR It has a history of significantly affecting water quality in past invasions outside of the Great Lakes	1
Not significantly	0✓
Unknown	U

- *Not reported.*

Does it harm any markets or economic sectors (e.g., commercial fisheries, aquaculture, agriculture)?

Yes, it has caused significant damage to one or more markets or economic sectors	6✓
Yes, some damage to markets or sectors has been observed, but negative consequences have been small AND/OR It has a history of harming markets or economic sectors in past invasions outside of the Great Lakes	1
Not significantly	0
Unknown	U

- *Costs in damages and control associated with C. pengoi in the U.S. are currently estimated at about five million US dollars annually (Pimentel et al., 2005).*

Does it inhibit recreational activities and/or associated tourism (e.g., through frequent water closures, equipment damage, decline of recreational species)?

Yes, it has caused widespread, frequent, or otherwise expensive inhibition of recreation and tourism	6
Yes, but negative consequences have been small	1✓
Not significantly	0
Unknown	U

- *Cercopagis pengoi fouls fishing lines, which acts both as a nuisance and as a possible mechanism of its dispersal and expansion. In a study by Jacobs and MacIsaac (2007), fouling was found to be most intense with longer lines and larger trolling distances; accumulation of C. pengoi on a single fishing line towed 1 km in Lake Ontario was as high as 1,024 individuals and 106 diapausing eggs. Lines specially designed to reduce waterflea fouling experienced diminished C. pengoi accumulation (Jacobs and MacIsaac 2007).*

Does it diminish the perceived aesthetic or natural value of the areas it inhabits?

Yes, the species has received significant attention from the media/public, significantly diminished the natural or cultural character of the area, or significantly reduced the area's value for future generations	6
Yes, but negative consequences have been small	1
Not significantly	0✓
Unknown	U

- *Not reported.*

Socio-Economic Impact Total	7
Total Unknowns (U)	0

Scientific Name: *Ctenopharyngodon idella*

Common Name: Grass carp

Organism Impact Assessment

IMPACT RESULTS

Environmental: High

Socio-Economic: Moderate

Beneficial: Low

Comments: This species was moved from the Watchlist to the Nonindigenous list in 2019.

ENVIRONMENTAL IMPACT

Does the species pose some hazard or threat to the health of native species (e.g., it magnifies toxin levels; is poisonous; is a pathogen, parasite, or a vector of either)? ✓

Yes, and it has impacted threatened/endangered species, resulted in the reduction or extinction of one or more native populations, affects multiple species, or is a reportable disease	6
Yes, but negative consequences have been small (e.g., limited number of infected individuals, limited pathogen transmissibility, mild effects on populations and ecosystems)	1 ✓
Not significantly	0
Unknown	U

- *Grass carp are known to be carriers of numerous parasitic organisms. Shireman and Smith (1983) thoroughly list a wide array of organisms, from viruses to protozoans to trematodes, which are parasites of Grass carp. Worth noting is Bothriocephalus acheilognathi, the Asian tapeworm. This parasite has been introduced by grass carp, to every continent except Antarctica (Bain 1993, Salgado-Maldonado and Pineda-Lopez 2003). Additionally, grass carp are the source of Ergasilus spp. in United Kingdom waters (Cowx 1997). However, disease and parasitism are not as prevalent in wild populations as in fish culture (Shireman and Smith 1983).*

Does it out-compete native species for available resources (e.g., habitat, food, nutrients, light)?

Yes, and it has resulted in significant adverse effects (e.g., impacted threatened/endangered species or caused critical reduction, extinction, behavioral changes including modified spawning behavior) on one or more native populations	6
Yes, and it has caused some noticeable stress to (e.g., decrease in growth, survival, fecundity) or decline of at least one native population	1 ✓
Not significantly	0
Unknown	U

- *Grass carp is known to out-compete native species for both food and habitat. Research in small closed systems has demonstrated that due to grass carp's preference for native aquatic plants over watermilfoil (*Myriophyllum spp.*); these fish compete with waterfowl, which feed on these plants as well (Fowler and Robson, 1978; McKnight and Hepp, 1995; Pine et al., 1990; Pine and Anderson, 1991).*

- *Furthermore, direct competition for plant material may also occur between grass carp and other native fishes that include macrophytes in their diet, such as gizzard shad (*Dorosoma cepedianum*), lake sturgeon (*Acipenser fulvescens*), as well as several species of buffalo (*Ictobius spp.*)(Cudmore and Mandrak, 2004; Coker et al., 2001).*
- *Grass carp may compete with planktonic and benthic species, including catfishes and hybrid sunfishes for aquatic plants (Shireman and Smith, 1983), especially during grass carp juvenile stages and at lower water temperatures (Fedorenko and Fraser, 1978).*
- *Direct competition for habitat has been found to occur between grass carp and other fish species, particularly bluegill. With their schooling habit, grass carp invade and disturb bluegill spawning areas, greatly reducing bluegill weight and numbers (Forester and Lawrence, 1978).*

Does it alter predator-prey relationships?

Yes, and it has resulted in significant adverse effects (e.g., impacted threatened/endangered species, caused significant reduction or extinction of one or more native populations, creation of a dead end or any other significant alteration in the food web)	6 [✓]
Yes, and it has resulted in some noticeable stress to (e.g., decrease in growth, survival, fecundity) or decline of at least one native population AND/OR Yes, and it has resulted in some alteration of the food web structure or processes, the effects of which have not been widespread or severe	1
Not significantly	0
Unknown	U

- *Grass carp have environmental impacts on the ecosystems they have been introduced. For instance, grass carp is known to be the source of major alterations to the trophic structure and food chains of aquatic systems. Many of these changes in plant, invertebrate, and fish communities are largely secondary consequences of reductions in the density and composition of aquatic plant communities (Bain 1993, Cudmore and Mandrak 2004). When stocked at high densities, grass carp can eliminate all vegetation in even large aquatic systems (e.g., 8100-ha Lake Conroe, Klussman et al., 1988). Declines have occurred in the diversity and density of organisms that are dependent on structured littoral habitats and food chains based on plant detritus, macrophytes, and attached algae as a consequence of reduced plant surface habitat, increased invertebrate food supplies (i.e. plant detritus), altered substrate conditions, and increased dissolved oxygen conditions (Bain 1993, Martin and Shireman 1976, Vinogradov and Zolotova 1974).*

Has it affected any native populations genetically (e.g., through hybridization, selective pressure, introgression)?

Yes, and it has caused a loss or alteration of genes that may be irreversible or has led to the decline of one or more native species (or added pressure to threatened/endangered species)	6
Yes, some genetic effects have been observed, but consequences have been limited to the individual level	1
Not significantly	0 [✓]
Unknown	U

- *No reports of affecting native fish genetically.*

Does it negatively affect water quality (e.g., increased turbidity or clarity, altered nutrient, oxygen, or other chemical levels/cycles)?

Yes, and it has had a widespread, long-term, or severe negative effect on water quality AND/OR Yes, and it has resulted in significant negative consequences for at least one native species	6✓
Yes, it has affected water quality to some extent, but the alterations and resulting adverse effects have been limited or inconsistent (as compared with above statement)	1
Not significantly	0
Unknown	U

- *Grazing by grass carp has been associated with alterations of water quality. The decay of these large volumes of dead aquatic plants due to grass carp's grazing and waste production elevate nutrient levels in water, induce phytoplankton blooms, reduce water clarity, and decrease oxygen levels (Bain 1993, Boyd 1971, Vinogradov and Zolotova 1974).*

Does it alter physical components of the ecosystem in some way (e.g., facilitated erosion/siltation, altered hydrology, altered macrophyte/phytoplankton communities, physical or chemical changes to substrate)?

Yes, and it has had a widespread, long term, or severe negative effect on the physical ecosystem AND/OR Yes, and it has resulted in significant negative consequences for at least one native species	6✓
Yes, it has affected the physical ecosystem to some extent, but the alterations and resulting adverse effects have been mild	1
Not significantly	0
Unknown	U

- *The herbivorous grass carp has a significant impact on macrophyte communities through intense grazing pressure (Bain 1993).*

Environmental Impact Total	20
Total Unknowns (U)	0

SOCIO-ECONOMIC IMPACT

NOTE: In this section, a "Not significantly" response should be selected if there have been no reports of a particular impact. An "Unknown" response is appropriate if the potential for a particular impact might be inferred from a significant environmental impact but has not been explicitly reported or if there is an unresolved debate about a particular impact.

Does the species pose some hazard or threat to human health (e.g., it magnifies toxin levels, is poisonous, a virus, bacteria, parasite, or a vector of one)?

Yes, significant effects on human health have already been observed	6
Yes, but negative consequences have not been widespread, long lasting, or severe	1✓
Not significantly	0
Unknown	U

- *Their contribution to increased algae blooms may affect drinking water quality.*

Does it cause damage to infrastructure (e.g., water intakes, pipes, or any other industrial or recreational infrastructure)?

Yes, it is known to cause significant damage	6
--	---

Yes, but the costs have been small and are largely reparable or preventable	1
Not significantly	0 ✓
Unknown	U

- *Has not been reported to damage infrastructure.*

Does it negatively affect water quality (i.e. in terms of being less suitable for human use)?

Yes, it has significantly affected water quality, and is costly or difficult to reverse	6
Yes, but the effects are negligible and/or easily reversed	1 ✓
Not significantly	0
Unknown	U

- *One of the undesirable consequences of stocking grass carp is increased turbidity, either algal or abiotic (Bonar et al. 2002, Lembi et al. 1978, Maceina et al. 1992, Water Environmental Services Incorporated 1994).*

Does it negatively affect any markets or economic sectors (e.g., commercial fisheries, aquaculture, agriculture)?

Yes, it has caused significant damage to one or more markets or economic sectors	6
Some damage to markets or sectors has been observed, but negative consequences have been small	1 ✓
Not significantly	0
Unknown	U

- *When in excessive numbers it destroys existing food chain relationships (Petr and Mitrofanov 1998).*
- *May threaten yellow perch (Perca flavescens) spawning by consuming emerging macrophytes where Yellow perch lay their eggs (Kocovsky pers. com., 2019)*

Does it inhibit recreational activities and/or associated tourism (e.g., through frequent water closures, equipment damage, decline of recreational species)?

Yes, it has caused widespread, frequent, or otherwise expensive inhibition of recreation and tourism	6
Yes, but negative consequences have been small	1 ✓
Not significantly	0
Unknown	U

- *May cause decline in recreational species through habitat modification, direct competition, or competition with species that act as forage fishes.*

Does it diminish the perceived aesthetic or natural value of the areas it inhabits?

Yes, the species has received significant attention from the media/public, significantly diminished the natural or cultural character of the area, or significantly reduced the area's value for future generations	6
Yes, but negative consequences have been small	1 ✓
Not significantly	0
Unknown	U

- *Grazing can promote algal blooms and increased turbidity (Bonar et al., 2002; Lembi et al., 1978; Maceina et al., 1992; Water Environmental Services Incorporated 1994).*

Socio-Economic Impact Total	5
Total Unknowns (U)	0

BENEFICIAL EFFECT

NOTE: In this section, a “Not significantly” response should be selected if there have been no reports of a particular effect. An “Unknown” response is appropriate if the potential for a particular effect might be inferred but has not been explicitly reported or if there is an unresolved debate about a particular effect.

Does it act as a biological control agent for aquatic weeds or other harmful nonindigenous organisms?

Yes, it has succeeded significantly as a control agent	6
Yes, it has had some success as a control agent, but may be inconsistent or lack a desired level of effectiveness	1 ✓
Not significantly	0
Unknown	U

- *Grass carp is being widely introduced throughout the United States to control aquatic vegetation in lakes and ponds (Chilton and Muoneke 1992, Page and Burr 1991). Grass carp can effectively control and eliminate aquatic plants in a variety of situations. Private fish farms have been producing large numbers of sterile, triploid grass carp as interest in stocking open systems increases (Bain 1993). Grass carp also are now routinely stocked in irrigation canals of the western United States (Bain 1993) and in Saskatchewan, Canada (Cudmore and Mandrak 2004).*
- *However, grass carp populations are below the necessary threshold to have an effect on submerged aquatic vegetation in the Great Lakes, and at the population level necessary to control SAV they could possibly contribute to eutrophication by releasing nutrients sequestered in wetlands (Cudmore and Mandrak, 2004).*

Is it commercially valuable (e.g., for fisheries, aquaculture, agriculture, bait, ornamental trade)?

Yes, it is economically important to at least one of these industries	6
Yes, but its economic contribution is small	1
Not significantly	0 ✓
Unknown	U

- *Sometimes captured for food in its native range, however, they rarely comprise a large proportion of the catch and are taken incidentally in common or silver carp fisheries in the Amur basin (Shireman and Smith 1983)*
- *Not currently accepted as a food fish in the United States.*

Is it recreationally valuable (e.g., for sport or leisurely fishing, as a pet, or for any other personal activity)?

Yes, it is commonly employed recreationally and has some perceived value for local communities and/or tourism	6
It is sometimes employed recreationally, but adds little value to local communities or tourism	1
Not significantly	0 ✓
Unknown	U

- *Not reported.*

Does the species have some medicinal or research value (i.e. outside of research geared towards its control)?

Yes, it has significant medicinal or research value	6
It has some medicinal or research value, but is not of high priority OR It is potentially important to medicine or research and is currently being or scheduled to be studied	1
Not significantly	0✓
Unknown	U

- *No reported medicinal or research value.*

Does the species remove toxins or pollutants from the water or otherwise increase water quality?

Yes, it reduces water treatment costs or has a significant positive impact for the health of humans and/or native species	6
Yes, but positive impact for humans or native species is considered negligible	1
Not significantly	0✓
Unknown	U

- *Not reported to remove toxins or pollutants from the water.*

Does the species have a positive ecological impact outside of biological control (e.g., increases the growth or reproduction rates of other species, fills an important gap in the food web, supports the survival of a species that is threatened, endangered species, or commercially valuable)?

Yes, it significantly contributes to the ecosystem in one or more of these ways	6
Yes, it provides some positive contribution to the ecosystem, but is not vital	1
Not significantly	0✓
Unknown	U

- *No other positive ecological effects reported.*

Beneficial Effect Total	1
Total Unknowns (U)	0

Scientific Name: *Daphnia galeata galeata*

Common Name: A waterflea

Organism Impact Assessment

IMPACT RESULTS

Environmental: High

Socio-Economic: Low

Beneficial: Low

Comments: Environmental impacts changed from unknown to high.

ENVIRONMENTAL IMPACT

Does the species pose some hazard or threat to the health of native species (e.g., it magnifies toxin levels, is poisonous, a virus, bacteria, parasite, or a vector of one)?

Yes, and it has resulted in the reduction or extinction of one or more native species populations, affects multiple species, or is a reportable disease	6
Yes, but negative consequences have been small (e.g., limited number of infected individuals, limited pathogen transmissibility, mild effects on populations and ecosystems) AND/OR It has significantly affected similar species in past invasions outside of the Great Lakes	1
Not significantly	0 ✓
Unknown	U

- *Not reported to threaten the health of native species.*

Does it out-compete native species for available resources (e.g., habitat, food, nutrients, light)?

Yes, and it has resulted in significant adverse effects (e.g., critical reduction, extinction, behavioral changes) on one or more native species populations	6
Yes, and it has caused some noticeable stress to or decline of at least one native species population	1 ✓
Not significantly	0
Unknown	U

- *Hybrids appear much more successful than parents, as hybrids represented 90-100% of swarms in four 'mixed' lakes (Taylor and Hebert 1993).*

Does it alter predator-prey relationships?

Yes, and it has resulted in significant adverse effects (e.g., added pressure to threatened/endangered species, significant reduction or extinction of any native species populations, creation of a dead end or any other significant alteration in the food web)	6
--	---

Yes, and it has resulted in some noticeable stress to or decline of at least one native species population AND/OR Yes, and it has resulted in some alteration of the food web structure or processes, the effects of which have not been widespread or severe	1
Not significantly	0✓
Unknown	U

- *Not reported to alter predator-prey relationships.*

Has it affected any native populations genetically (e.g., through hybridization, selective pressure, introgression)?

Yes, and it has caused a loss or alteration of genes which may be irreversible or has led to the decline or extinction of one or more native species	6✓
Yes, some genetic effects have been observed, but consequences have been limited to the individual level AND/OR It has genetically affected the same or similar species in past invasions outside of the Great Lakes	1
Not significantly	0
Unknown	U

- *The North American and European subspecies have hybridized in the Great Lakes basin. Morphologies and genetics of hybrid D. g. galeata x D. g. mendotae tend more toward the European D. g. galeata form in Lake Erie and more towards the North American D. g. mendotae form in the Lake Ontario drainage. Before, these populations were genetically distinct and mated non-randomly under sympatry. Hybrids appear much more successful than parents, as hybrids represented 90-100% of swarms in four 'mixed' lakes (Taylor and Hebert 1993).*

Does it negatively affect water quality (e.g., increased turbidity or clarity, altered nutrient, oxygen, or other chemical levels/cycles)?

Yes, and it has had a widespread, long term, or severe negative effect on water quality AND/OR Yes, and it has resulted in significant negative consequences for at least one native species	6
Yes, it has affected water quality to some extent, but the alterations and resulting adverse effects have been mild AND/OR It has significantly affected water quality in past invasions outside of the Great Lakes	1
Not significantly	0✓
Unknown	U

- *No reported effects on water quality.*

Does it alter the physical ecosystem in some way (e.g., facilitated erosion/siltation, altered hydrology, altered macrophyte/phytoplankton communities, changes to substrate (physical or chemical))?

Yes, and it has had a widespread, long term, or severe negative effect on the physical ecosystem AND/OR Yes, and it has resulted in significant negative consequences for at least one native species	6
Yes, it has affected the physical ecosystem to some extent, but the alterations and resulting adverse effects have been mild	1

AND/OR	
It has significantly altered physical ecosystems in past invasions outside of the Great Lakes	
Not significantly	0✓
Unknown	U

Environmental Impact Total	7
Total Unknowns (U)	0

Scientific Name: *Hemimysis anomala*

Common Name: Bloody red shrimp

Organism Impact Assessment

IMPACT POTENTIAL RESULTS

Environmental: Moderate

Socio-Economic: Low

Beneficial: Low

Comments: Environmental impacts changed from unknown to moderate.

POTENTIAL ENVIRONMENTAL IMPACT

Does the species pose some hazard or threat to the health of native species (e.g., it magnifies toxin levels; is poisonous; is a pathogen, parasite, or a vector of either)?

Yes, and it has impacted threatened/endangered species, resulted in the reduction or extinction of one or more native populations, affects multiple species, or is a reportable disease	6
Yes, but negative consequences have been small (e.g., limited number of infected individuals, limited pathogen transmissibility, mild effects on populations and ecosystems)	1
Not significantly	0
Unknown	U✓

- *A mysid introduction can increase the biomagnification of contaminants in piscivores through a lengthening of the food chain; for example, concentrations of polychlorinated biphenyls and mercury in fishes have been shown to be higher in lakes containing mysids than in mysid-free lakes (Cabana et al., 1994; cf. Rasmussen et al., 1990).*
- *Through direct transmission and indirect effects on the food web, introduced mysids may cause increased parasitism by nematodes, cestodes, and acanthocephalans in fishes (Lasenby et al., 1986; Northcote 1991).*

Does it out-compete native species for available resources (e.g., habitat, food, nutrients, light)?

Yes, and it has resulted in significant adverse effects (e.g., impacted threatened/endangered species or caused critical reduction, extinction, behavioral changes including modified spawning behavior) on one or more native populations	6
Yes, and it has caused some noticeable stress to (e.g., decrease in growth, survival, fecundity) or decline of at least one native population	1✓
Not significantly	0
Unknown	U

- *Hemimysis anomala may compete with, or prey upon, other invertebrate predators, such as Bythotrephes longimanus and Leptodora kindti. Its omnivory may also reduce local phytoplankton if small-sized juvenile mysids are abundant (Ketelaars et al., 1999).*
- *Found to consume a broader diet than native zooplankton, including the ability to prey on larger diet items as juveniles (Evans et al., 2018).*
- *Displays high diet plasticity in the St. Lawrence River (Marty et al., 2012).*

Does it alter predator-prey relationships?

Yes, and it has resulted in significant adverse effects (e.g., impacted threatened/endangered species, caused significant reduction or extinction of one or more native populations, creation of a dead end or any other significant alteration in the food web)	6
Yes, and it has resulted in some noticeable stress to (e.g., decrease in growth, survival, fecundity) or decline of at least one native population AND/OR Yes, and it has resulted in some alteration of the food web structure or processes, the effects of which have not been widespread or severe	1 ✓
Not significantly	0
Unknown	U

- *Based on its impacts in some European reservoirs (Ketelaars et al. 1999), H. anomala may reduce zooplankton biomass and diversity in invaded areas, with cladocerans, rotifers, and ostracods being most affected.*
- *Hemimysis anomala may compete with, or prey upon, other invertebrate predators, such as Bythotrephes longimanus and Leptodora kindti. Its omnivory may also reduce local phytoplankton if small-sized juvenile mysids are abundant (Ketelaars et al., 1999)*
- *Predatory functional responses were generally higher than those of the comparator native species (Mysis salemaai and M. diluviana). Had similar or higher attack rates, consistently lower prey handling times and higher maximum feeding rates compared to those of the two Mysis species, formerly known as 'Mysis relicta', which itself has an extensive history of food web disruption in lakes to which it has been introduced. (Dick et al., 2012).*
- *In Europe, Hemimysis invasions have been associated with a decline in phytoplankton and zooplankton populations (Ketelaars et al., 1999), which in turn, may potentially lead to a reduction in energy flow to higher trophic levels of the food web (Marty et al., 2012).*
- *Sinclair et al., found Hemimysis strongly favors cladocerans over copepods due to copepods' stronger ability to avoid predation.*

Has it affected any native populations genetically (e.g., through hybridization, selective pressure, introgression)?

Yes, and it has caused a loss or alteration of genes that may be irreversible or has led to the decline of one or more native species (or added pressure to threatened/endangered species)	6
Yes, some genetic effects have been observed, but consequences have been limited to the individual level	1 ✓
Not significantly	0
Unknown	U

- *Sinclair et al., (2016) demonstrated in a mesocosm experiment that Hemimysis predation selected for larger Daphnia spp.*

Does it negatively affect water quality (e.g., increased turbidity or clarity, altered nutrient, oxygen, or other chemical levels/cycles)?

Yes, and it has had a widespread, long-term, or severe negative effect on water quality AND/OR Yes, and it has resulted in significant negative consequences for at least one native species	6
Yes, it has affected water quality to some extent, but the alterations and resulting adverse effects have been limited or inconsistent (as compared with above statement)	1

Not significantly	0
Unknown	U✓

- *Hemimysis feeds rapidly, even at low prey densities, and its fecal pellets may alter the local physico-chemical environment (Ketelaars et al., 1999; Olenin and Leppäkoski 1999; Pienimäki and Leppäkoski 2004).*

Does it alter physical components of the ecosystem in some way (e.g., facilitated erosion/siltation, altered hydrology, altered macrophyte/phytoplankton communities, physical or chemical changes to substrate)?

Yes, and it has had a widespread, long term, or severe negative effect on the physical ecosystem AND/OR Yes, and it has resulted in significant negative consequences for at least one native species	6
Yes, it has affected the physical ecosystem to some extent, but the alterations and resulting adverse effects have been mild	1
Not significantly	0✓
Unknown	U

Environmental Impact Total	3
Total Unknowns (U)	2

Scientific Name: *Lepomis humilis*

Common Name: Orangespotted sunfish

Organism Impact Assessment

IMPACT POTENTIAL RESULTS

Environmental: Unknown

Socio-Economic: Low

Beneficial: Low

Comments: Beneficial impacts changed from unknown to low.

BENEFICIAL EFFECT

NOTE: In this section, a “Not significantly” response should be selected if there have been no reports of a particular effect. An “Unknown” response is appropriate if the potential for a particular effect might be inferred but has not been explicitly reported or if there is an unresolved debate about a particular effect.

Does it act as a biological control agent for aquatic weeds or other harmful nonindigenous organisms?

Yes, it has succeeded significantly as a control agent	6
Yes, it has had some success as a control agent, but may be inconsistent or lack a desired level of effectiveness	1
Not significantly	0✓
Unknown	U

- *Consume mosquito larvae (Barney and Anson, 1923).*

Is it commercially valuable (e.g., for fisheries, aquaculture, agriculture, bait, ornamental trade)?

Yes, it is economically important to at least one of these industries	6
Yes, but its economic contribution is small	1
Not significantly	0✓
Unknown	U

- *Too small for commercial fishing, not a popular aquarium species.*

Is it recreationally valuable (e.g., for sport or leisurely fishing, as a pet, or for any other personal activity)?

Yes, it is commonly employed recreationally and has some perceived value for local communities and/or tourism	6
It is sometimes employed recreationally, but adds little value to local communities or tourism	1
Not significantly	0✓
Unknown	U

- *Smaller species than other centrarchids, not popular for fishing.*

Does the species have some medicinal or research value (i.e. outside of research geared towards its control)?

Yes, it has significant medicinal or research value	6
It has some medicinal or research value, but is not of high priority OR It is potentially important to medicine or research and is currently being or scheduled to be studied	1
Not significantly	0✓
Unknown	U

- *No reported medicinal or research value.*

Does the species remove toxins or pollutants from the water or otherwise increase water quality?

Yes, it reduces water treatment costs or has a significant positive impact for the health of humans and/or native species	6
Yes, but positive impact for humans or native species is considered negligible	1
Not significantly	0✓
Unknown	U

- *Not reported.*

Does the species have a positive ecological impact outside of biological control (e.g., increases the growth or reproduction rates of other species, fills an important gap in the food web, supports the survival of a species that is threatened, endangered species, or commercially valuable)?

Yes, it significantly contributes to the ecosystem in one or more of these ways	6
Yes, it provides some positive contribution to the ecosystem, but is not vital	1
Not significantly	0✓
Unknown	U

- *Not reported.*

Beneficial Effect Total	0
Total Unknowns (U)	0

Scientific Name: *Oncorhynchus gorbuscha*

Common Name: Pink salmon

Organism Impact Assessment

IMPACT RESULTS

Environmental: Moderate

Socio-Economic: Low

Beneficial: Moderate

Comments: Environmental impact changed from unknown to moderate.

ENVIRONMENTAL IMPACT

Does the species pose some hazard or threat to the health of native species (e.g., it magnifies toxin levels, is poisonous, a virus, bacteria, parasite, or a vector of one)?

Yes, and it has resulted in the reduction or extinction of one or more native species populations, affects multiple species, or is a reportable disease	6
Yes, but negative consequences have been small (e.g., limited number of infected individuals, limited pathogen transmissibility, mild effects on populations and ecosystems) AND/OR It has significantly affected similar species in past invasions outside of the Great Lakes	1
Not significantly	0 ✓
Unknown	U

- *Not reported.*

Does it out-compete native species for available resources (e.g., habitat, food, nutrients, light)?

Yes, and it has resulted in significant adverse effects (e.g., critical reduction, extinction, behavioral changes) on one or more native species populations	6
Yes, and it has caused some noticeable stress to or decline of at least one native species population	1 ✓
Not significantly	0
Unknown	U

- *Pink salmon may displace native chubs by way of food competition and may also compete with native cisco (Coregonus artedii) (Becker 1983).*
- *Pink salmon has also been identified as utilizing spawning habitats similar to those used by brook trout (Salvelinus fontinalis), potentially providing another mechanism of competition (Kocik and Jones 1999).*

Does it alter predator-prey relationships?

Yes, and it has resulted in significant adverse effects	6
---	---

(e.g., added pressure to threatened/endangered species, significant reduction or extinction of any native species populations, creation of a dead end or any other significant alteration in the food web)	
Yes, and it has resulted in some noticeable stress to or decline of at least one native species population AND/OR Yes, and it has resulted in some alteration of the food web structure or processes, the effects of which have not been widespread or severe	1 ✓
Not significantly	0
Unknown	U

- *Individuals over one year old feed heavily on rainbow smelt (Osmerus mordax) and alewife (Alosa pseudoharengus), which are important components of the diets of other Great Lakes salmonids (Diana 1990; Kocik and Taylor 1987; Kocik et al., 1991).*

Has it affected any native populations genetically (e.g., through hybridization, selective pressure, introgression)?

Yes, and it has caused a loss or alteration of genes which may be irreversible or has led to the decline or extinction of one or more native species	6
Yes, some genetic effects have been observed, but consequences have been limited to the individual level AND/OR It has genetically affected the same or similar species in past invasions outside of the Great Lakes	1 ✓
Not significantly	0
Unknown	U

- *Genetic analysis of populations in the St. Marys River, Mich. indicates that pink salmon is capable of hybridizing with recreationally important chinook salmon (Oncorhynchus tshawytscha) (Kirkpatrick et al., 2007). Hybridization has the potential to create further competition for the parental species, especially since the hybrid appears to have growth rates that exceed those of pink and chinook salmon.*

Does it negatively affect water quality (e.g., increased turbidity or clarity, altered nutrient, oxygen, or other chemical levels/cycles)?

Yes, and it has had a widespread, long term, or severe negative effect on water quality AND/OR Yes, and it has resulted in significant negative consequences for at least one native species	6
Yes, it has affected water quality to some extent, but the alterations and resulting adverse effects have been mild AND/OR It has significantly affected water quality in past invasions outside of the Great Lakes	1
Not significantly	0
Unknown	U ✓

- *Unknown.*

Does it alter the physical ecosystem in some way (e.g., facilitated erosion/siltation, altered hydrology, altered macrophyte/phytoplankton communities, changes to substrate (physical or chemical))?

Yes, and it has had a widespread, long term, or severe negative effect on the physical ecosystem AND/OR	6
--	---

Yes, and it has resulted in significant negative consequences for at least one native species	
Yes, it has affected the physical ecosystem to some extent, but the alterations and resulting adverse effects have been mild AND/OR It has significantly altered physical ecosystems in past invasions outside of the Great Lakes	1
Not significantly	0
Unknown	U✓

- *Unknown.*

Environmental Impact Total	3
Total Unknowns (U)	2

Scientific Name: *Oncorhynchus tshawytscha*

Common Name: Chinook salmon

Organism Impact Assessment

IMPACT RESULTS

Environmental: High

Socio-Economic: Low

Beneficial: High

Comments: Environmental impact changed from moderate to high.

ENVIRONMENTAL IMPACT

Does the species pose some hazard or threat to the health of native species (e.g., it magnifies toxin levels, is poisonous, a virus, bacteria, parasite, or a vector of one)?

Yes, and it has resulted in the reduction or extinction of one or more native species populations, affects multiple species, or is a reportable disease	6✓
Yes, but negative consequences have been small (e.g., limited number of infected individuals, limited pathogen transmissibility, mild effects on populations and ecosystems) AND/OR It has significantly affected similar species in past invasions outside of the Great Lakes	1
Not significantly	0
Unknown	U

- *Renibacterium salmoninarum* is believed to have been introduced to Lake Michigan when Chinook salmon (*Oncorhynchus tshawytscha*) stocking began in 1967 (Holey et al. 1998). The province of Ontario considers this species to be endemic - present in most samples at 1-33% without signs of clinical disease or gross lesions (GLFHC 2015).
- Great Lakes native species found harboring *R. salmoninarum* include lake whitefish (*Coregonus clupeaformis*), bloater (*C. hoyi*), lake herring (*C. artedii*), mottled scuplin (*Cottus bairdi*), white sucker (*Catostomus commersonii*), muskellunge (*Esox masquinongy*), channel catfish (*Ictalurus punctatus*), lake sturgeon (*Acipenser fulvescens*), and walleye (*Sander vitreus*) (COSEWIC 2005; GLFHC 2006; GLFHC 2012; Hay 2003; Jonas et al., 2002; Nuhfer et al., 2005; Starliper et al., 1997).

Does it out-compete native species for available resources (e.g., habitat, food, nutrients, light)?

Yes, and it has resulted in significant adverse effects (e.g., critical reduction, extinction, behavioral changes) on one or more native species populations	6
Yes, and it has caused some noticeable stress to or decline of at least one native species population	1✓
Not significantly	0
Unknown	U

- *In the Great Lakes, Chinook salmon competes with native lake trout (Salvelinus namaycush)(Page and Laird 1993).*
- *Scott et al. (2003) found that the presence of Chinook salmon causes delayed nesting and reduced survival of Atlantic salmon (Salmo salar) during spawning in Lake Ontario.*

Does it alter predator-prey relationships?

Yes, and it has resulted in significant adverse effects (e.g., added pressure to threatened/endangered species, significant reduction or extinction of any native species populations, creation of a dead end or any other significant alteration in the food web)	6✓
Yes, and it has resulted in some noticeable stress to or decline of at least one native species population AND/OR Yes, and it has resulted in some alteration of the food web structure or processes, the effects of which have not been widespread or severe	1
Not significantly	0
Unknown	U

- *Chinook salmon is a predatory fish and may impact populations of smaller fish. Jones et al.(1993) predicted that maintaining high levels of predator demand by stocking Chinook and other top predators at the current rate would eventually lead to an alewife collapse, possibly followed by the further collapse of other small forage fish population.*
- *Bunnell et al., (2014) found that predation by Chinook salmon has top-down effects on forage fish in the Great Lakes which causes resource limitation for native piscivores.*

Has it affected any native populations genetically (e.g., through hybridization, selective pressure, introgression)?

Yes, and it has caused a loss or alteration of genes which may be irreversible or has led to the decline or extinction of one or more native species	6
Yes, some genetic effects have been observed, but consequences have been limited to the individual level AND/OR It has genetically affected the same or similar species in past invasions outside of the Great Lakes	1
Not significantly	0✓
Unknown	U

- *Not reported.*

Does it negatively affect water quality (e.g., increased turbidity or clarity, altered nutrient, oxygen, or other chemical levels/cycles)?

Yes, and it has had a widespread, long term, or severe negative effect on water quality AND/OR Yes, and it has resulted in significant negative consequences for at least one native species	6
Yes, it has affected water quality to some extent, but the alterations and resulting adverse effects have been mild AND/OR It has significantly affected water quality in past invasions outside of the Great Lakes	1
Not significantly	0✓
Unknown	U

- *Not reported.*

Does it alter the physical ecosystem in some way (e.g., facilitated erosion/siltation, altered hydrology, altered macrophyte/phytoplankton communities, changes to substrate (physical or chemical))?

Yes, and it has had a widespread, long term, or severe negative effect on the physical ecosystem AND/OR Yes, and it has resulted in significant negative consequences for at least one native species	6
Yes, it has affected the physical ecosystem to some extent, but the alterations and resulting adverse effects have been mild AND/OR It has significantly altered physical ecosystems in past invasions outside of the Great Lakes	1
Not significantly	0
Unknown	U✓

- *Crawford et al. (2001) pointed out that salmonids have the potential to alter the energy and nutrient cycles of the Great Lakes system through increased energy transfer between open water and streams/tributaries. This energy transfer includes the addition of nitrogen and phosphorous to tributaries through decaying salmonine carcasses, as well as the addition of salmon eggs and dead fish as a food source in streams (Ivan et al., 2011; Parmenter and Lamarra 1991; Rand et al., 1992).*
- *The presence of live salmonids may have an even greater effect on nutrients in streams through the excretion of ammonium and soluble reactive phosphorus and their mechanical disturbance of the stream bottom (Ivan et al. 2011; Tiegs et al., 2009).*

Environmental Impact Total	13
Total Unknowns (U)	1

Scientific Name: *Schizopera borutzkyi*

Common Name: An oarsman

Organism Impact Assessment

IMPACT RESULTS

Environmental: Low

Socio-Economic: Low

Beneficial: Low

Comments: Environmental impact changed from unknown to low.

ENVIRONMENTAL IMPACT

Does the species pose some hazard or threat to the health of native species (e.g., it magnifies toxin levels, is poisonous, a virus, bacteria, parasite, or a vector of one)?

Yes, and it has resulted in the reduction or extinction of one or more native species populations, affects multiple species, or is a reportable disease	6
Yes, but negative consequences have been small (e.g., limited number of infected individuals, limited pathogen transmissibility, mild effects on populations and ecosystems) AND/OR It has significantly affected similar species in past invasions outside of the Great Lakes	1
Not significantly	0✓
Unknown	U

- *Not reported.*

Does it out-compete native species for available resources (e.g., habitat, food, nutrients, light)?

Yes, and it has resulted in significant adverse effects (e.g., critical reduction, extinction, behavioral changes) on one or more native species populations	6
Yes, and it has caused some noticeable stress to or decline of at least one native species population	1
Not significantly	0✓
Unknown	U

- *Schizopera borutzkyi has altered the species composition of nearshore harpacticoid communities, comprising up to 75% of the community at deep sites (15 m) in Lake Michigan. Impact on the food web in these communities is unknown, but it is likely that S. borutzkyi is competing with native species for similar resources or has the ability to exploit previously unused resources (Horvath et al., 2001).*
- *Dominant harpacticoid in Lake Michigan following introduction; no evidence that S. borutzkyi has altered food webs or ecosystem level processes where it has established in the Great Lakes; low consequences of establishment (Grippo et al., 2017).*

Does it alter predator-prey relationships?

Yes, and it has resulted in significant adverse effects (e.g., added pressure to threatened/endangered species, significant reduction or extinction of any native species populations, creation of a dead end or any other significant alteration in the food web)	6
Yes, and it has resulted in some noticeable stress to or decline of at least one native species population AND/OR Yes, and it has resulted in some alteration of the food web structure or processes, the effects of which have not been widespread or severe	1
Not significantly	0
Unknown	U [✓]

- *Unknown.*

Has it affected any native populations genetically (e.g., through hybridization, selective pressure, introgression)?

Yes, and it has caused a loss or alteration of genes which may be irreversible or has led to the decline or extinction of one or more native species	6
Yes, some genetic effects have been observed, but consequences have been limited to the individual level AND/OR It has genetically affected the same or similar species in past invasions outside of the Great Lakes	1
Not significantly	0 [✓]
Unknown	U

- *Not reported.*

5) Does it negatively affect water quality (e.g., increased turbidity or clarity, altered nutrient, oxygen, or other chemical levels/cycles)?

Yes, and it has had a widespread, long term, or severe negative effect on water quality AND/OR Yes, and it has resulted in significant negative consequences for at least one native species	6
Yes, it has affected water quality to some extent, but the alterations and resulting adverse effects have been mild AND/OR It has significantly affected water quality in past invasions outside of the Great Lakes	1
Not significantly	0 [✓]
Unknown	U

- *Not reported.*

Does it alter the physical ecosystem in some way (e.g., facilitated erosion/siltation, altered hydrology, altered macrophyte/phytoplankton communities, changes to substrate (physical or chemical))?

Yes, and it has had a widespread, long term, or severe negative effect on the physical ecosystem AND/OR Yes, and it has resulted in significant negative consequences for at least one native species	6
Yes, it has affected the physical ecosystem to some extent, but the alterations and resulting adverse effects have been mild AND/OR	1

It has significantly altered physical ecosystems in past invasions outside of the Great Lakes	
Not significantly	0✓
Unknown	U

Environmental Impact Total	0
Total Unknowns (U)	1

4.0 REFERENCES

- Bailey, R.M., and G.R. Smith. 1992. Names of Michigan Fishes. Michigan Department of Natural Resources 7 pp.
- Bailey, R.M., and M.O. Allum. 1962. Fishes of South Dakota. Miscellaneous Publications of the Museum of Zoology, University of Michigan, Ann Arbor, MI 119:1-131.
- Bain, M. 1993. Assessing impacts of introduced aquatic species: Grass Carp in large systems. *Environmental Management* 17:211-224.
- Bain, M.B., D.H. Webb, M.D. Tangedal, and L.N. Mangum. 1990. Movements and habitat use by Grass Carp in a large mainstream reservoir. *Transactions of the American Fisheries Society* 119:553-561.
- Barney, R.L., and B.J. Anson. 1923. Life History and Ecology of the Orange-Spotted Sunfish *Lepomis humilis*. Bureau of Fisheries. 16 pp.
- Bauers, Sandy. 1995. Debate on Fish that Can't Scale Back. The Philadelphia Inquirer, City & Region Section. Published on 4 June, 1995.
- Becker, G.C. 1983. Fishes of Wisconsin. University of Wisconsin Press, Madison, WI. 1052 pp. Available: <http://digital.library.wisc.edu/1711.dl/EcoNatRes.FishesWI>
- Bence, J.R., and K.D. Smith. 1999. An overview of recreational fisheries of the Great Lakes. In Taylor, W.W., and C.P. Ferreri (Eds.), Great Lakes Fisheries Policy and Management: A Binational Perspective. Michigan State University Press, East Lansing, MI. pp. 259-306.
- Benoit, H.P., O.E. Johannsson, D.M. Warner, W.G. Sprules, and L.G. Rudstam. 2002. Assessing the impact of a recent predatory invader: the population dynamics, vertical distribution, and potential prey of *Cercopagis pengoi* in Lake Ontario. *Limnology and Oceanography* 47(3):626-635.
- Berg, L.S. 1949. Freshwater fishes of the USSR and adjacent countries. Izdatel'stvo Akademii Nauk SSSR, Moscow.
- Bij de Vaate, A., K. Jazdzewski, H.A.M. Ketelaars, S. Gollasch, and G. Van der Velde. 2002. Geographical patterns in range extension of Ponto-Caspian macroinvertebrate species in Europe. *Canadian Journal of Fisheries and Aquatic Sciences* 59: 1159-1174.
- Boersma, M. 1995. The allocation of resources to reproduction in *Daphnia galeata*: against the odds? *Ecology* 76(4): 1251-1261.
- Bondarenko, M.V., and E.A. Yablonskaya. 1979. The rate of oxygen consumption by the mysid *Paramysis lacustris* (Czern.) from the northern Caspian Sea. In Effect of water management on the biological productivity of the Azov and Caspian seas. VNIRO, Moscow. pp. 65-69. (In Russian with English summary.)
- Borcherding, J., S. Murawski, and H. Arndt. 2006. Population ecology, vertical migration and feeding of the Ponto-Caspian invader *Hemimysis anomala* in a gravel-pit lake connected to the River Rhine. *Freshwater Biology* 51: 2376-2387.

- Boschung, H.T. 1992. Catalogue of freshwater and marine fishes of Alabama. *Alabama Museum of Natural History Bulletin* 14:1-266.
- Brooking, T.E., L.G. Rudstam, S.D. Krueger, J.R. Jackson, A.B. Welsh, and W.W. Fetzer. 2010. First occurrence of the mysid *Hemimysis anomala* in an inland lake in North America, Oneida Lake. *Journal of Great Lakes Research* 36(3): 577-581.
- Brooks, J.L. 1957. The systematics of North American *Daphnia*. *Memoirs of the Connecticut Academy of Arts and Sciences* 13: 1-180.
- Brown, M., J. Roth, B. Smith, B. Boscarino. 2017. The Light at the end of the funnel?: Using light-based traps for the detection and collection of a nearshore aquatic, invasive invertebrate, *Hemimysis anomala*. *Journal of Great Lakes Research* 43: 717-727.
- Bubinas, A.D. 1980. Formation of benthic fauna as a food base for fish in the reservoir of the Kaunas hydro electric power plant, Lithuanian-SSR USSR. *Lietuvos TSR Mokslu Akademijos Darbai Serija C Biologijos Mokslai* 4: 91-96.
- Bushnoe, T.M., D.M. Warner, L.G. Rudstam, and E.L. Mills. 2003. *Cercopagis pengoi* as a new prey item for alewife (*Alosa pseudoharengus*) and rainbow smelt (*Osmerus mordax*) in Lake Ontario. *Journal of Great Lakes Research* 29(2):205-212.
- Cabana, G., A. Tremblay, J. Kalff, and J.B. Rasmussen. 1994. Pelagic food chain structure in Ontario lakes: a determinant of mercury levels in lake trout (*Salvelinus namaycush*). *Canadian Journal of Fisheries and Aquatic Sciences* 51: 381-389.
- CABI. Invasive Species Compendium. *Dreissena rostriformis bugensis*. CAB International. [Accessed Mar 9, 2020]. <https://www.cabi.org/isc/datasheet/107770>
- Carl, L.M. 1982. Natural reproduction of coho salmon and chinook salmon in some Michigan streams. *North American Journal of Fisheries Management* 4:375-380.
- Cavaletto, J., H. Vanderploeg, R. Pichlová-Ptácníková, S. Pothoven, J. Liebig, and G.L. Fahnenstiel. 2010. Temporal and spatial separation allow coexistence of predatory cladocerans: *Leptodora kindtii*, *Bythotrephes longimanus*, and *Cercopagis pengoi*, in southeastern Lake Michigan. *Journal of Great Lakes Research* 36(SP3):65-73.
- Chapman D.C., Davis J.J., Jenkins J.A., Kocovsky P.M., Miner J.G., Farver J., and Jackson P.R. First evidence of Grass Carp recruitment in the Great Lakes Basin. *Journal of Great Lakes Research* 39 4 547-554
- Charlebois, P.M., M.J. Raffenberg, and J.M. Dettmers. 2001. First occurrence of *Cercopagis pengoi* in Lake Michigan. *Journal of Great Lakes Research* 27(2):258-261.
- Chilton, II, E.W., and M.I. Muoneke Biology and management of Grass Carp (*Ctenopharyngodon idella*, Cyprinidae) for vegetation control: a North American perspective. *Reviews in Fish Biology and Fisheries* 2: 283-320
- Cisneros, R., E. Hooker, and L. E. Velasquez. 1991b. Natural diet of herbivorous zooplankton in Lake Xolotlán (Managua). *Hydrobiological Bulletin* 25(2):163-167.

- Clearwater, S.J., C.W. Hickey, and M.L. Martin. 2008. Overview of potential piscicides and molluscicides for controlling aquatic pest species in New Zealand. Science & Technical Publishing, New Zealand Department of Conservation, Wellington, New Zealand.
- Colorado Division of Wildlife. 2010. Fishery Survey Summaries - Union Reservoir.
- COSEWIC. 2008. Committee on the Status of Endangered Wildlife in Canada Annual Report.
- Courtenay, W. R., Jr., D. A. Hensley, J. N. Taylor, and J. A. McCann. 1984. Distribution of exotic fishes in the continental United States. Pages 41-77 in W. R. Courtenay, Jr., and J. R. Stauffer, Jr., editors. Distribution, biology and management of exotic fishes. Johns Hopkins University Press, Baltimore, MD.
- Crawford, H.M., D.A. Jensen, B. Peichel, P.M. Charlebois, B.A. Doll, S.H. Kay, V.A. Ramey, and M.B. O'Leary. 2001. Sea grant and invasive aquatic plants: a national outreach initiative. *Journal of Aquatic Plant Management* 39(1):8-11.
- Crawford, S.S. 2001. Salmonine introductions to the Laurentian Great Lakes: an historical review and evaluation of ecological effects. *Canadian Special Publication of Fisheries and Aquatic Sciences* No. 132. 205 pp.
- Cross, D.G. 1970. The tolerance of Grass Carp *Ctenopharyngodon idella* (Val.) to seawater. *Journal of Fish Biology* 2:231-233.
- Cross, F.B. 1967. Handbook of Fishes of Kansas. University of Kansas Museum of Natural History Miscellaneous Publication No.45. University of Kansas, Topeka, KS. 357 pp.
- Cross, F.B., and J.T. Collins. 1995. Fishes in Kansas. University of Kansas Natural History Museum. Lawrence, KS.
- Cudmore-Vokey, B., and E.J. Crossman. 2000. Checklists of the fish fauna of the Laurentian Great Lakes and their connecting channels. *Canadian Manuscript Report of Fisheries and Aquatic Sciences* 2500: v + 39 pp.
- Dahlberg, M.D., and D.C. Scott. 1971b. Introductions of freshwater fishes in Georgia. *Bulletin of the Georgia Academy of Science* 29:245-252
- DeVaney, S.C., K.M. McNyset, J.B. Williams, A.T. Peterson, and E.O. Wiley. 2009. A tale of four "carp": invasion potential and ecological niche modeling. *PLoS ONE* 4(5): e5451.
- Diana, J.S. 1990. Food habits of angler-caught salmonines in western Lake Huron. *Journal of Great Lakes Research* 16(2):271-278.
- Dill, W.A., and A.J. Cordone. 1997. History and status of introduced fishes in California, 1871-1996. *California Department of Fish and Game Fish Bulletin*, volume 178.
- Dodson, S.I. 1988. Cyclomorphosis in *Daphnia galeata mendotae* Birge and *D. retrocurva* Forbes as a predator-induced response. *Freshwater Biology* 19: 109-114.
- Dumont, H.J., and S. Negrea. 1996. A conspectus of the Cladocera of the subterranean waters of the world. *Hydrobiologia* 325: 1-30.

- Dumont, S. 2006. Notes and News. A new invasive species in the north-east of France, *Hemimysis anomala* G. O. Sars, 1907 (Mysidacea). *Crustaceana* 79: 1269-1274.
- Dumont, S., and C.D. Muller. 2009. Distribution, ecology and impact of a small invasive shellfish, *Hemimysis anomala* in Alsatian water. *Biological Invasions* 12(3): 495-500.
- Ebert, D. 2005. Chapter 9, Population Dynamics and Community Ecology in Ecology, Epidemiology, and Evolution of Parasitism in *Daphnia*. National Center for Biotechnology Information. Bethesda, MD.
- Ebner, M.P. [Ed.]. 1995. The State of Lake Huron in 1992. Great Lakes Fishery Commission Special Publication. 95-2. 140 pp.
- Ellis, S. and H.J. MacIsaac. 2009. Salinity tolerance of Great Lakes invaders. *Freshwater Biology* 54(1):77-89.
- Embke, H.S., P.M. Kocovsky, C.A. Richter, J.J. Pritt, C.M. Mayer, and S.S. Qian. 2016. First direct confirmation of grass carp spawning in a Great Lakes tributary. *Journal of Great Lakes Research* 42 (4), 899-903.
- Emery, L. 1985. Review of fish introduced into the Great Lakes, 1819-1974. Great Lakes Fishery Commission Technical Report 45:1-31.
- Etnier, D.A., and W.C. Starnes. 1993. The Fishes of Tennessee. The University of Tennessee Press, Knoxville, TN. 681 pp.
- Evans, T.M., R. Naddafi, B. Weidel, B. Lantry, M. Walsh, B. Boscarino, O. Johannsson, L. Rudstam. 2018. Stomach contents and stable isotopes analysis indicate *Hemimysis anomala* in Lake Ontario are Broadly Omnivorous. *Journal of Great Lakes Research* 44:467-475.
- Embke, H.S. et al. 2016. First direct confirmation of Grass Carp spawning in a Great Lakes tributary. *Journal of Great Lakes Research* 42(4):899-903
- Fisheries and Oceans Canada. 2008. Survey of recreational fishing in Canada: Selected results for the Great Lakes fishery, 2005. Catalogue No. Fs23-522/2005-1E. Fisheries and Oceans Canada, Ottawa, Ontario. Available: <https://waves-vagues.dfo-mpo.gc.ca/Library/346362.pdf>
- Flobner, D., and K. Kraus. 1986. On the taxonomy of the *Daphnia hyalina-galeata* complex (Crustacea: Cladocera). *Hydrobiologia* 137: 97-115.
- Freidhoff, J. pers. comm. Buffalo State College, Buffalo, NY
- Fuller, P.L. 2003. Freshwater aquatic vertebrate introductions in the United States: patterns and pathways. In G.M. Ruiz and J.T. Carlton (eds.), *Invasive Species: Vectors and Management Strategies* (pp. 123-151). Island Press.
- Garza, E.L., and R. L. Whitman. 2004. The nearshore benthic invertebrate community of southern Lake Michigan and its response to beach nourishment. *Journal of Great Lakes Research* 30(1): 114-122.

- Gharrett, A. J. and M. A. Thomason. 1987. *Canadian Journal of Fisheries and Aquatic Sciences*, 44(4): 787-792.
- Gherardi, F., S. Gollasch, D. Minchin, S. Olenin, and V.E. Panov. 2009. Alien invertebrates and fish in European inland waters. In J.A. Drake (ed.), *DAISIE Handbook of Alien Species in Europe* (pp. 81-92). Dordrecht: Springer.
- Glagolev, S.M. 1986. Species composition of *Daphnia* in Lake Glubokoe with notes on the taxonomy and geographical distribution of some species. *Hydrobiologia* 141: 55-82.
- GLMRIS. 2012. Appendix C: Inventory of Available Controls for Aquatic Nuisance Species of Concern, Chicago Area Waterway System. U.S. Army Corps of Engineers.
- Global Invasive Species Database. IUCN-World Conservation Union—Invasive Species Specialist Group. 27 January 2011. <http://www.issg.org/database/>
- Gorokhova, E., T. Fagerberg, S. Hansson. 2004. Predation by herring (*Clupea harengus*) and sprat (*Sprattus sprattus*) on *Cercopagis pengoi* in a western Baltic Sea bay. *ICES Journal of Marine Science* 61(6):959-965.
- Grippio, M.A., I. Hlohowskyj, L. Fox, B. Herman, J. Pothoff, C. Yoe, J. Hayse. 2017. Aquatic Nuisance Species in the Great Lakes and Mississippi River Basin--A Risk Assessment in Support of GLMRIS. *Environmental Management* 59(1): 154-173.
- Guillory, V. and R.D. Gasaway. 1978. Zoogeography of the Grass Carp in the United States. *Transactions of the American Fisheries Society* 107:105-112
- Halpern, T., N. Paulson, and J.T. Hatch. 2002. Pink salmon. Minnesota Department of Natural Resources. http://academics.cehd.umn.edu/hatch/research/fish/fishes/pink_salmon.html
- Hansen, M.J. [Ed.] 1994 The State of Lake Superior in 1992. Great Lakes Fishery Commission Special Publication 94-1.
- Hassan-Williams, C., and T.H. Bonner. n.d. *Lepomis humilis*: orangespotted sunfish. Texas State University, San Marcos-Biology Department/Aquatic Station.
- Hegrenes, S. 2001. Diet-induced phenotypic plasticity of feeding morphology in the orangespotted sunfish, *Lepomis humilis*. *Ecology of Freshwater Fish* 10:35-42.
- Heisey, D., and K.G. Porter. 1977. The effect of ambient oxygen concentration on filtering and respiration rates of *Daphnia galeata mendotae* and *Daphnia magna*. *Limnology and Oceanography* 22(5): 839-845.
- Herborg, L.-M., N.E. Mandrak, B.C. Cudmore, and H.J. MacIsaac. 2007. Comparative distribution and invasion risk of snakehead (Channidae) and Asian carp (Cyprinidae) species in North America. *Canadian Journal of Fisheries and Aquatic Sciences* 64: 1723-1735.
- Hocutt, C. H., R. E. Jenkins, and J.R. Stauffer, Jr. 1986. Zoogeography of the fishes of the central Appalachians and central Atlantic Coastal Plain. Pages 161-212 in C.H. Hocutt, and E.O. Wiley, editors. *The Zoogeography of North American Freshwater Fishes*. John Wiley and Sons, New York, NY.

- Holdich, D., S. Gallagher, L. Rippon, P. Harding, and R. Stubbington. 2006. The invasive Ponto-Caspian mysid, *Hemimysis anomala*, reaches the UK. *Aquatic Invasions* 1(1): 4-6.
- Horvath, T.G., R.L. Whitman, and L. L. Last. 2001. Establishment of two invasive crustaceans (Copepoda: Harpacticoida) in the nearshore sands of Lake Michigan. *Canadian Journal of Fisheries and Aquatic Sciences* 58: 1261-1264.
- Hubbs, C., R.J. Edwards, and G.P. Garrett. 1991. An annotated checklist of freshwater fishes of Texas, with key to identification of species. *Texas Journal of Science, Supplement* 43(4):1-56.
- Hulsmann, S., and T. Mehner. 1997. Predation by underyearling perch (*Perca fluviatilis*) on a *Daphnia galeata* population in a short-term enclosure experiment. *Freshwater Biology* 38(1):209-219.
- International Union for Conservation of Nature (IUCN). 2010. *Cercopagis pengoi*. Global Invasive Species Database. <http://www.issg.org/database/species/ecology.asp?fr=1&si=118>
- Ivan, L.N., E.S. Rutherford, and T.H. Johengen. 2011. Impacts of adfluvial fish on the ecology of two Great Lakes tributaries. *Transactions of the American Fisheries Society* 140:1670-1682.
- Ives, J.T., J. Marty, Y. DeLafontaine, T.B. Johnson, M.A. Koops, and M. Power. 2013. Spatial variability in trophic offset and food sources of *Hemimysis anomala* in lentic and lotic ecosystems within the Great Lakes basin. *Journal of Plankton Research* 35(4): 772-784.
- Jacobs, M.J., and H.J. MacIsaac. 2007. Fouling of fishing line by the waterflea *Cercopagis pengoi*: a mechanism of human-mediated dispersal of zooplankton? *Hydrobiologia* 583(1):119-126.
- Janas, U., and P. Wysocki. 2005. *Hemimysis anomala* G.O. Sars, 1907 (Crustacea, Mysidacea) – first record in the Gulf of Gdansk. *Oceanologia* 47: 405-408.
- Jazdzewski, K., A. Konopacka, and M. Grabowski. 2005. Native and alien malacostracan crustacean along the Polish Baltic Sea coast in the twentieth century. *Oceanological and Hydrobiological Studies XXXIV*(suppl. 1): 175-193.
- Jenkins, R.E., and N.M. Burkhead. 1994. *Freshwater Fishes of Virginia*. American Fisheries Society, Bethesda, MD.
- Jones, M.L., J.F. Koonce, and R. O'Gorman. 1993. Sustainability of hatchery-dependent salmonine fisheries in Lake Ontario: the conflict between predator demand and prey supply. *Transactions of the American Fisheries Society* 122:1002-1018.
- Kerr, S.J., and G.C. LeTendre. 1991. The State of the Lake Ontario Fish Community in 1989. Great Lakes Fishery Commission Special Publication 91-3. 38 pp.
- Kestrup, Å.M. and A. Ricciardi. 2008. Occurrence of the Ponto-Caspian mysid shrimp *Hemimysis anomala* (Crustacea, Mysida) in the St. Lawrence River. *Aquatic Invasions* 3(4): 461-464.
- Kestrup, Å.M., J.T.A. Dick, and A. Ricciardi. 2011. Interactions between invasive and native crustaceans: differential functional responses of intraguild predators towards juvenile hetero-specifics. *Biological Invasions* 13(3):731-737.

- Ketelaars, H.A.M., F.E. Lambrechts-van de Clundert, C.J. Carpentier, A.J. Waqenvoort, and W. Hooqenboezem. 1999. Ecological effects of the mass occurrence of the Ponto-Caspian invader, *Hemimysis anomala* G.O. Sars, 1907 (Crustacea: Mysidacea), in a freshwater storage reservoir in the Netherlands, with notes on its autecology and new records. *Hydrobiologia* 394: 233-248.
- Kilambi, R.V., and A. Zdinak. 1980. The effects of acclimation on the salinity tolerance of Grass Carp. *Journal of Fish Biology* 16:171-175.
- Kilby, J.D., E. Crittenden, and L.E. Williams. 1959. Several fishes new to Florida freshwaters. *Copeia* 1959(1):77-78.
- Kirkpatrick, N.S., D.W. Everitt, and B.I. Evans. 2007. Asymmetric hybridization of pink (*Oncorhynchus gorbuscha*) and Chinook (*O. tshawytscha*) salmon in the St. Marys River, Michigan. *Journal of Great Lakes Research* 33(2):358-365.
- Kocik, J.F., and M.L. Jones. 1999. Pacific salmonines in the Great Lakes basin. In Taylor, W.W. and C.P. Ferreri, (Eds.). *Great Lakes Fisheries Policy and Management: A Binational Perspective*. Michigan State University Press, East Lansing, MI, pp. 455-488.
- Kocik, J.F., and W.W. Taylor. 1987. Diet and movements of age-1+ pink salmon in western Lake Huron. *Transactions of the American Fisheries Society* 116(4):628-633.
- Kocik, J.F., W.W. Taylor, and W.C. Wagner. 1991. Abundance, size, and recruitment of pink salmon (*Oncorhynchus gorbuscha*) in selected Michigan tributaries of the upper Great Lakes, 1984-1988. *Journal of Great Lakes Research* 17(2):203-213.
- Kocovsky, P.M., D.C. Chapman, and J.E. McKenna
- Kolar, C. S., D. C. Chapman, W. R. Courtenay Jr., C. M. Housel, and J. D. Williams. 2005. Asian Carps of the Genus *Hypophthalmichthys* (Pisces, Cyprinidae) ? A Biological Synopsis and Environmental Risk Assessment. U.S. Fish and Wildlife Service.
- Komarova, T.I. 1989. Ecological and faunistic review of Mysidacea (Crustacea: Mysidae) in the Sea of Azov, USSR. *Vestnik Zoologii* 4: 3-7.
- Kwain, W., and G.A. Rose. 1986. Spawning migration of Great Lakes pink salmon (*Oncorhynchus gorbuscha*): size and sex distributions, river entrance and exit. *Journal of Great Lakes Research* 12(2):101-108.
- Langeland, A., J.I. Koksvisk, and J. Nydal. 1991. Impact of the introduction of *Mysis relicta* on the zooplankton and fish populations in a Norwegian lake. *American Fisheries Society Symposium* 9: 98-114.
- Lantry, B.F., M.G. Walsh, J.H. Johnson, and J.E. McKenna. 2010. Occurrence of the Great Lake's most recent invader, *Hemimysis anomala*, in the diet of fishes in southeastern Lake Ontario. *Journal of Great Lakes Research* 36(1): 179-183.
- Lasenby, D.C., T.G. Northcote, and M. Fürst. 1986. Theory, practice, and effects of *Mysis* introductions to North American and Scandinavian lakes. *Canadian Journal of Fisheries and Aquatic Sciences* 43: 1277-1284.

Latta, W.C. – Michigan Department of Natural Resources (d. 2008).

Laxson, C.L., K.N. McPhedran, J.C. Makarewicz, I.V. Telesh, and H.J. MacIsaac. 2003. Effects of the non-indigenous cladoceran *Cercopagis pengoi* on the lower food web of Lake Ontario. *Freshwater Biology* 48(12):2094-2106.

Lee, D.S., C.R. Gilbert, C.H. Hocutt, R.E. Jenkins, D.E. McAllister, and J.R. Stauffer, Jr. 1980. Atlas of North American freshwater fishes. North Carolina State Museum of Natural History, Raleigh, NC.

Leibold, M., and A.J. Tessier. 1991. Contrasting patterns of body size of *Daphnia* species that segregate by habitat. *Oecologia* 86: 324-348.

Lesko, L.T., P.L. Hudson, J.W. Reid, and M.A. Chriscinske. 2003. Harpacticoid copepods of the Laurentian Great Lakes. Great Lakes Science Center, Ann Arbor, Michigan.
<http://www.glsc.usgs.gov/greatlakescopepods/Key.asp?GROUP=Harpacticoid>

Litvinchuk, L.F., and I.V. Telesh. 2006. Distribution, population structure, and ecosystem effects of the invader *Cercopagis pengoi* (Polyphemoidea, Cladocera) in the Gulf of Finland and the open Baltic Sea. *Oceanologia* 48 (S):243-257.

López, C., L. M. Soto, L. Dávalos-Lind, and O. Lind. 2008. Occurrence of *Diaphanosoma fluviatile* Hansen 1899 (Cladocera: Sisidae) in two reservoirs in central Texas. *The Southwestern Naturalist* 53(3):412-414.

Lundberg, S., and J.E. Svensson. 2004. The mysid shrimp *Hemimysis anomala* G. O. Sars documented in Sweden. *Fauna och Flora (Stockholm)* 99(1): 38-40.

Lynch, M., L.J. Weider, and W. Lampert. 1986. Measurement of the carbon balance in *Daphnia*. *Limnology and Oceanography* 31(1): 17-33.

Lysebo, E.M. 1995. Behavioural and morphological changes in polymorphic *Daphnia* related to different predation regimes. *Hydrobiologia* 307: 185-191.

Maceina, M.J., and J.V. Shireman. 1979. Grass Carp: effects of salinity on survival, weight loss, and muscle tissue water content. *Progressive Fish Culturist* 41:69-72

Makarewicz, J.C., I.A. Grigorovich, E. Mills, E. Damaske, M.E. Cristescu, W. Pearsall, M.J. LaVoie, R. Keats, L. Rudstam, P. Hebert, H. Halbritter, T. Kelly, C. Matkovich, and H.J. MacIsaac. 2001. Distribution, fecundity, and genetics of *Cercopagis pengoi* (Ostroumov) (Crustacea, Cladocera) in Lake Ontario. *Journal of Great Lakes Research* 27(1):19-32.

Mandrak, N. E. 1989. Potential Invasion of the Great-Lakes by Fish Species Associated with Climatic Warming. *Journal of Great Lakes Research* 15:306-316.

Mandrak, N. E., and B. Cudmore. 2005. Risk assessment for Asian carps in Canada. Research Document 2004/103. Fisheries and Oceans Canada.

Mandrak, N.E, and E.J. Crossman. 1992. A checklist of Ontario freshwater fishes. Royal Ontario Museum, Toronto.

- Marrone, G. 1996. Chinook salmon. South Dakota Department of Game, Fish and Parks, Division of Wildlife, Pierre, SD. <http://www3.northern.edu/natsource/FISH/Chinoo1.htm> (accessed 3/4/2011).
- Marty, J., J. Ives, Y. DeLafontaine, S. Despatie, M. Koops, M. Power. 2012. Evaluation of carbon pathways supporting the diet of invasive *Hemimysis anomala* in a large river. *Journal of Great Lakes Research* 38: 45-51.
- Marty, J., K. Bowen, M.A. Koops, and M. Power. 2010. Distribution and ecology of *Hemimysis anomala*, the latest invader of the Great Lakes basin. *Hydrobiologia* 647(1):71-80.
- McAfee, W. R. 1966. Landlocked king salmon. Pages 294-295 in A. Calhoun, editor. *Inland Fisheries Management*. California Department of Fish and Game.
- Menhinick, E. F. 1991. The freshwater fishes of North Carolina. North Carolina Wildlife Resources Commission. 227 pp.
- Mettee, M.F., P.E. O'Neil, and J.M. Pierson. 1996. Fishes of Alabama and the Mobile Basin. Oxmoor House, Inc., Birmingham, AL.
- Michigan Department of Natural Resources (MIDNR). 2003. Pink salmon (*Oncorhynchus gorbuscha*). https://www.michigan.gov/dnr/0,4570,7-350-79135_79218_79614_82590---,00.html. (Accessed 6/24/2020).
- Michigan Department of Natural Resources and Environment (MIDNRE). 2011. Chinook salmon studies. https://www.michigan.gov/dnr/0,4570,7-350-79135_79218_79614_82588---,00.html (Accessed 6/24/2020).
- Mills, E.L., and J.L. Forney. 1983. Impact on *Daphnia pulex* of predation by young yellow perch in Oneida Lake, New York. *Transactions of the American Fisheries Society* 112:151-161.
- Mills, E.L., J.M. Casselman, R. Dermott, J.D. Fitzsimmons, G. Gal, K.T. Holeck, J.A. Hoyle, O.E. Hohannsson, B.F. Lantry, J.C. Makearewicz, E.S. Millard, I.F. Munawar, M. Munawar, R. O'Gorman, R.W. Owens, L.G. Rudstam, T. Schaner, and T.J. Stewart. 2003. Lake Ontario: food web dynamics in a changing ecosystem (1970-2000). *Canadian Journal of Fisheries and Aquatic Sciences* 60: 471-490.
- Mitzner, L. 1978. Evaluation of biological control of nuisance aquatic vegetation by Grass Carp. *Transactions of the American Fisheries Society* 107:135-145.
- Monchenko, V.I. 1967. Beitrag zur Kenntnis der Gattung Schizopera (Crustacea, Harpacticoida) im Schwarzen Meer. *Zoologischer Anzeiger* 178: 367-374.
- Monchenko, V.I. 1974. New data on the distribution of rare harpacticoids off the northwestern shores of the Black Sea. *Hydrobiological Journal* 10(5): 78-80.
- Monchenko, V.I. 1995. Harpacticoida of River Dnieper (results of faunistic studies). *Gidrobiologicheskii Zhurnal* 31(5): 33-37.
- Mordukhai-Boltovskoi, F.D. 1979. Composition and distribution of Caspian fauna in light of modern data. *Internationale Revue der gesamten Hydrobiologie* 64: 1-38.

- Murphy, E.A., and Jackson, P.R., 2013, Hydraulic and water-quality data collection for the investigation of Great Lakes tributaries for Asian carp spawning and egg-transport suitability: U.S. Geological Survey Scientific Investigations Report 2013–5106, 30 p., <http://pubs.usgs.gov/sir/2013/5106/>.
- Negus, M.T. 1995. Bioenergetics modeling as a salmonine management tool applied to Minnesota Waters of Lake Superior. *North American Journal of Fisheries Management* 15:60-78.
- Nelson, J.S., and S.D. Gerking. 1968. Annotated key to the fishes of Indiana. Project 342-303-815. Department of Zoology, Indiana Aquatic Research Unit, Indiana State University, Bloomington, IN.
- Nixon, D.E., and R.L. Miller. 1978. Movements of Grass Carp, *Ctenopharyngodon idella*, in an open reservoir system as described by radiotelemetry. *Transactions of the American Fisheries Society* 107:146-148.
- Northcote, T.G. 1991. Success, problems, and control of introduced mysid populations in lakes and reservoirs. *American Fisheries Society Symposium* 9: 5-16.
- Ojaveer, H., L.A. Kuhns, R.P Barbiero, and M.L. Tuchman. 2001. Distribution and population characteristics of *Cercopagis pengoi* in Lake Ontario. *Journal of Great Lakes Research* 27(1):10-18.
- Olenin, S., and E. Leppäkoski. 1999. Non-native animals in the Baltic Sea: alteration of benthic habitats in coastal inlets and lagoons. *Hydrobiologia* 393: 233-243.
- Ontario Federation of Anglers and Hunters. 2009. OFAH Database - 2009 Download.
- Ontario's Invading Species Awareness Program. Bloody Red Shrimp *Hemimysis anomala*. <http://www.invadingspecies.com/bloody-red-shrimp/>. Accessed on 06/24/2020.
- Ontario's Invading Species Awareness Program. Spiny and Fishhook Waterfleas. <http://www.invadingspecies.com/-and-fishhook-waterfleas/>. Accessed on 06/24/2020.
- Owen, J.B., D.S. Elsen, and G.W. Russell. 1981. Distribution of Fishes in North and South Dakota Basins Affected by the Garrison Diversion Unit. Fisheries Research Unit, University of North Dakota, Grand Forks, ND.
- Page, L.M. and B.M. Burr. 2011. A field guide to freshwater fishes of North America north of Mexico. Boston: Houghton Mifflin Harcourt, 663 pp.
- Page, L.M., and B.M. Burr. 1991. A field guide to freshwater fishes of North America north of Mexico. The Peterson Guide Series, vol. 42. Houghton Mifflin Company, Boston, MA.
- Page, L.M., and C.A. Laird. 1993. The identification of the nonnative fishes inhabiting Illinois waters. Report prepared by Center for Biodiversity, Illinois Natural History Survey, Champaign, for Illinois Department of Conservation, Springfield. Center for Biodiversity Technical Report 1993(4). 39 pp.
- Panov, V.E., P.I. Krylov, and N. Riccardi. 2004. Role of diapause in dispersal and invasion success by aquatic invertebrates. *Journal of Limnology* 63(Suppl. 1): 56-69.
- Parmenter, R.R., and V.A. Lamarra. 1991. Nutrient cycling in a freshwater marsh: The decomposition of fish and waterfowl carrion. *Limnology and Oceanography* 36(5):976-987.

- Peck, J.W., T.S. Jones, W.R. MacCallum, and S.T. Schram. 1999. Contribution of hatchery-reared fish to Chinook salmon population and sport fisheries in Lake Superior. *North American Journal of Fisheries Management* 19:155-164.
- Petr, T. and V. P. Mitrofanov. 1998. The impact on fish stocks of river regulation in Central Asia and Kazakhstan. *Lakes & Reservoirs: Research & Management* 3:143-164.
- Pflieger, W. 1997. The fishes of Missouri. Missouri Department of Environmental Conservation, Jefferson City, MO.
- Pichlová, R., and J. Vijverberg. 2001. A laboratory study of functional response of *Leptodora kindtii* to some cladoceran species and copepod nauplii. *Archiv für Hydrobiologie* 150(4):529-544.
- Pichlová-Ptácníková, R., and H.A. Vanderploeg. 2009. The invasive cladoceran *Cercopagis pengoi* is a generalist predator capable of feeding on a variety of prey species of different sizes and escape abilities. *Fundamental and Applied Limnology* 173 (4): 267-279.
- Pienimäki, M.E., and E. Leppäkoski. 2004. Invasion pressure on the Finnish Lake District: invasion corridors and barriers. 2004. *Biological Invasions* 6: 331-346.
- Pimentel, D. 2005. Aquatic nuisance species in the New York State Canal and Hudson River systems and the Great Lakes basin: an economic and environmental assessment. *Environmental Management* 35(5):692-701.
- Pligin, Y.V., and L.V. Yemel'yanova. 1989. Acclimatization of Caspian invertebrates in Dnieper reservoirs. *Hydrobiological Journal* 25: 1-9.
- Post, J.R., and D.J. McQueen. 1987. The impact of planktivorous fish on the structure of a plankton community. *Freshwater Biology* 17(1):79-89.
- Pothoven, S. A., H. A. Vanderploeg, S. A. Ludsin, T. O. Hook and S. B. Brandt. 2009. Feeding Ecology of emerald shiners and rainbow smelt in Central Lake Erie. *Journal of Great Lakes Research* 35(2):190-198.
- Pothoven, S.A., H.A. Vanderploeg, J.F. Cavaletto, D.M. Krueger, D.M. Mason, and S.B. Brandt. 2007. Alewife planktivory controls the abundance of two invasive predatory cladocerans in Lake Michigan. *Freshwater Biology* 52(3):561-573.
- Pothoven, S.A., I.A. Grigorovich, G.L. Fahnenstiel, and M.D. Balcer. 2007. Introduction of the Ponto-Caspian bloody-red mysid *Hemimysis anomala* into the Lake Michigan basin. *Journal of Great Lakes Research*. 33: 285-292.
- Quinn, T. P. 2005. The Behavior and Ecology of Pacific Salmon and Trout: Ist edition. American Fisheries Society and University of Washington Press.
- Rand, P.S., and D.J. Stewart. 1998. Prey fish exploitation, salmonine production, and pelagic food web efficiency in Lake Ontario. *Canadian Journal of Fisheries and Aquatic Sciences* 55 (February, 1998):318-327.

- Rand, P.S., C.A.S. Hall, W.H. McDowell, N.H. Ringler, and J.G. Kennen. 1992. Factors limiting primary productivity in Lake Ontario tributaries receiving salmon migrations. *Canadian Journal of Fisheries and Aquatic Sciences* 49(11):2377-2385.
- Rasmussen, J.B., Rowan, D.J., Lean, D.R.S., and Carey, J.H. 1990. Food chain structure in Ontario lakes determines PCB levels in lake trout (*Salvelinus namaycush*) and other pelagic fish. *Canadian Journal of Fisheries and Aquatic Sciences* 47: 2030-2038.
- Repka, S. 1997. Effects of food type on the life history of *Daphnia* clones from lakes differing in trophic state. I. *Daphnia galeata* feeding on *Scenedesmus* and *Oscillatoria*. *Freshwater Biology* 37: 675-683.
- Ricciardi, A., and J. B. Rasmussen. 1998. Predicting the identity and impact of future biological invaders: a priority for aquatic resource management. *Canadian Journal of Fisheries and Aquatic Sciences* 55: 1759-1765.
- Ricciardi, A., S. Avlijas, and J. Marty. 2011. Forecasting the ecological impacts of the *Hemimysis anomala* invasion in North America: Lessons from other freshwater mysid introductions. *Journal of Great Lakes Research* 38(2): 7-13.
- Rixon, C.A.M., I.C. Duggan, N.M.N. Bergeron, A. Ricciardi, and H.J. MacIsaac. 2005. Invasion risks posed by the aquarium trade and live fish markets on the Laurentian Great Lakes. *Biodiversity and Conservation* 14: 1365-1381.
- Robison, H.W., and T.M. Buchanan. 1998. *Fishes of Arkansas*. University of Arkansas Press, Fayetteville, AR.
- Rutherford, E.S. 1997. Evaluation of natural reproduction, stocking rates, and fishing regulations for Steelhead *Oncorhynchus mykiss*, Chinook salmon *O. tshawytscha*, and coho salmon *O. kisutch* in Lake Michigan. Federal Aid in Sport Fish Restoration, Project F-35-R-22, Final Report, Fisheries Division, Michigan Department of Natural Resources, Ann Arbor, Mich.
- Salemaa, H., and V. Hietalahti. 1993. *Hemimysis anomala* G. O. Sars (Crustacea: Mysidacea) – immigration of a Pontocaspian mysid into the Baltic Sea. *Annales Zoologici Fennici* 30(4): 271-276.
- Sanders, S., C. Castiglione, and M. H. Hoff. 2018. Risk Assessment Mapping Program: RAMP. U.S. Fish and Wildlife Service. USGS (U.S. Geological Survey). 2018. *Diaphanosoma fluviatile*. USGS Nonindigenous Aquatic Species Database, Gainesville, Florida. Available: <https://nas.er.usgs.gov/viewer/omap.aspx?SpeciesID=2648>. (July 2018).
- Schofield, P. J., J. D. Williams, L. G. Nico, P. Fuller, and M. R. Thomas. 2005. Foreign Nonindigenous Carps and Minnows (Cyprinidae) in the United States-A guide to their Identification, Distribution, and Biology. US Geological Survey Scientific Investigations Report.
- Schreiner, D.R. (Ed.). 1995. Fisheries management plan for the Minnesota waters of Lake Superior. Special Publication #149. Minnesota Department of Natural Resources, Division of Fish and Wildlife, Duluth, MN, 87 pp.
- Schwenk, K., D. Posada, and P.D.N. Hebert. 2000. Molecular systematics of European *Hyalodaphnia*: the role of contemporary hybridization in ancient species. *Proceedings: Biological Sciences* 267(1455): 1833-1842.

- Scott, R.J., D.L.G. Noakes, F.W.H. Meamish, and L.M. Carl. 2003. Chinook salmon impede Atlantic salmon conservation in Lake Ontario. *Ecology of Freshwater Fish* 12:66-73.
- Scott, W. B., and E. J. Crossman. 1973. Freshwater fishes of Canada. *Fisheries Research Board of Canada*, Bulletin 184. 966 pp.
- Shireman, J.V., and C.R. Smith. 1983. Synopsis of biological data on the Grass Carp, *Ctenopharyngodon idella* (Cuvier and Valenciennes, 1884). *FAO Fisheries Synopsis* 135.
- Sinclair, J. S., Arnott, S. E., and A. Cox. 2016. The quick and the dead: copepods dominate as cladocerans decline following invasion by *Hemimysis anomala*. *Canadian Journal of Fisheries and Aquatic Sciences* 73:793-803.
- Spencer, C.N., B.R. McClelland, and J.A. Stanford. 1991. Shrimp stocking, salmon collapse, and eagle displacement. *BioScience* 44: 14-21.
- Stanley, J. G. 1976. Reproduction of the Grass Carp (*Ctenopharyngodon idella*) outside its native range. *Fisheries* 1:7-10.
- Stanley, J.G., W.W. Miley, and D.L. Sutton. 1978. Reproductive requirements and likelihood for naturalization of escaped Grass Carp in the United States. *Transactions of the American Fisheries Society* 103:587-592.
- Starnes, W.C., J. Odenkirk, and M.J. Ashton. 2011. Update and analysis of fish occurrences in the lower Potomac River drainage in the vicinity of Plummers Island, Maryland—Contribution XXXI to the natural history of Plummers Island, Maryland. *Proceedings of the Biological Society of Washington* 124(4):280-309.
- Stauffer, J.R., Jr., J.M. Boltz, and L.R. White. 1995. The Fishes of West Virginia. West Virginia Department of Natural Resources. Academy of Natural Sciences of Philadelphia, Philadelphia, PA. 389 pp.
- Stewart, D. J., and M. Ibarra. 1991. Predation and production by salmonine fishes in Lake Michigan, 1978-88. *Canadian Journal of Fisheries and Aquatic Science* 48: 909-922.
- Stewart, T.J., O.E. Johannsson, K. Holeck , W.G. Sprules, and R. O’Gorman. 2010. The Lake Ontario zooplankton community before (1987-1991) and after (2001-2005) invasion-induced ecosystem change. *Journal of Great Lakes Research* 36(4):596-605.
- Stewart, T.J., W.G. Sprules, and R. O’Gorman. 2009. Shifts in the diet of Lake Ontario alewife in response to ecosystem change. *Journal of Great Lakes Research* 35(2):241-249.
- Stollewerk, A. 2010. The water flea *Daphnia* - a ‘new’ model system for ecology and evolution? *Journal of Biology*. 9(2): 21.
- Storch, A., K. Schulz, C. Cáceres, P. Smyntek, J. Dettmers, and M. Teece. 2007. Consumption of two exotic zooplankton by alewife (*Alosa pseudoharengus*) and rainbow smelt (*Osmerus mordax*) in three Laurentian Great Lakes. *Canadian Journal of Fisheries and Aquatic Science* 64(10):1314-1328.
- Sublette, J.E., M.D. Hatch, and M. Sublette. 1990. The fishes of New Mexico. University of New Mexico Press, Albuquerque, NM.

Swift, C.C., C.R. Gilbert, S.A. Bortone, G.H. Burgess, and R.W. Yerger. 1986. Zoogeography of the Fishes of the Southeastern United States: Savannah River to Lake Pontchartrain. In C.H. Hocutt and E.O. Wiley, eds. *The Zoogeography of North American Freshwater Fishes*. John Wiley and Sons, New York, NY. 213-266.

Tanner, H. A., and W. H. Tody. 2002. History of the Great Lakes Salmon fishery: a Michigan perspective. Pages 139-154 in K. D. Lynch, M. L. Jones, and W. W. Taylor, editors. *Sustaining North American Salmon: perspectives across regions and disciplines*, American Fisheries Society, Bethesda, Maryland.

Taylor, D.J., and P.D.N. Hebert. 1993. Cryptic intercontinental hybridization in *Daphnia* (Crustacea): the ghost of introductions past. *Proceedings of the Royal Society of London Series B* 254: 163-168.

Taylor, D.J., P.D.N. Hebert, and J.K. Colbourne. 1996. Phylogenetics and evolution of the *Daphnia longispina* group (Crustacea) based on 12S rDNA sequence and allozyme variation. *Molecular Phylogenetics and Evolution* 5(3): 495-510.

Thermal and hydrologic suitability of Lake Erie and its major tributaries for spawning of Asian carps. *Journal of Great Lakes Research* 38 1 159-166

Therriault, T.W., I.A. Grigorovich, D.D. Kane, E.M. Haas, D.A. Culver, and H.J. MacIsaac. 2002. Range expansion of the exotic zooplankter *Cercopagis pengoi* (Ostroumov) into western Lake Erie and Muskegon Lake. *Journal of Great Lakes Research* 28(4):698-701.

Thompson, E., J.C. Makarewicz, and T.W. Lewis. 2005. Additional link in the food web does not biomagnify a persistent contaminant in Lake Ontario: the case of *Cercopagis pengoi*. *Journal of Great Lakes Research* 31(2):210-218.

Tiegs, S.D., E.Y. Campbell, P.S. Levi, J. Rüegg, M.E. Benbow, D.T. Chaloner, R.W. Merritt, J.L. Tank, and G.A. Lamberti. 2009. Separating physical disturbance and nutrient enrichment caused by Pacific salmon in stream ecosystems. *Freshwater Biology* 54(9): 1864-1857.

Tilmant, J.T. 1999. Management of nonindigenous aquatic fish in the U.S. National Park System. National Park Service. 50 pp.

Tilmant, J.T. 1999. Management of nonindigenous aquatic fish in the U.S. National Park System. National Park Service. 50 pp.

Trautman, M.B. 1981. *The Fishes of Ohio*. Ohio State University Press, Columbus, OH.

Tsehay, I., M. L. Jones, J. R. Bence, T. O. Brenden C. P. Madenjian, and D.M. Warner. 2014. A multispecies statistical age-structured model to assess predator-prey balance: application to an intensively managed Lake Michigan pelagic fish community. *Canadian Journal of Fisheries and Aquatic Sciences* 71: 627-644.

U.S. Environmental Protection Agency (USEPA). 2008. EPA Monitoring Data. EPA Great Lakes National Program Office. Available <http://www.epa.gov/grtlakes/monitoring/biology/exotics/cercopagis.html>

U.S. Environmental Protection Agency. 2008a. Effects of climate change on aquatic invasive species and implications for management and research [electronic resource]. Washington, D.C.: National

Center for Environmental Assessment, Office of Research and Development, U.S. Environmental Protection Agency, 2008.

U.S. Environmental Protection Agency. 2008b. Predicting future introductions of nonindigenous species to the Great Lakes [electronic resource] / National Center for Environmental Assessment, Office of Research and Development, U.S. Environmental Protection Agency. Washington, DC: National Center for Environmental Assessment, 2008.

U.S. Fish and Wildlife Service, Region 3 Fisheries Program, and Great Lakes Fishery Commission (USFWS/GLFC). 2010. Great Lakes Fish Stocking database. Available: <http://www.glfc.org/fishstocking/index.htm>

Vanderploeg, H.A., T.F. Nalepa, D.J. Jude, E.L. Mills, K.T. Holeck, J.R. Leibig, I.A. Grigorovich, and H. Ojaveer. 2002. Dispersal and emerging ecological impacts of Ponto-Caspian species in the Laurentian Great Lakes. *Canadian Journal of Fisheries and Aquatic Sciences* 59(7):1209-1228.

Vinyard, G.L. 2001. Fish Species Recorded from Nevada. Biological Resources Research Center. University of Nevada Reno. 5 pp.

W. Stone, pers. comm. - New York Dept. Environmental Conservation, Delmar, NY.

Walsh, M.G., B.F. Lantry, B. Boscarino, K. Bowen, J. Gerlofsma, T. Schaner, R. Back, J. Questel, G.A. Smythe, R. Cap, M. Goehle, B. Young, M. Chalupnicki, J.H. Johnson, and J.E. McKenna. 2010. The early observations on an emerging Great Lakes invader *Hemimysis anomala* in Lake Ontario. *Journal of Great Lakes Research* 36(3) 499-504.

Warner, D., L.G. Rudstam, H. Benoit, E.L. Mills, and O. Johannsson. 2006. Changes in seasonal nearshore zooplankton abundance patterns in Lake Ontario following establishment of the exotic predator *Cercopagis pengoi*. *Journal of Great Lakes Research* 32(3):531-542.

Wieringa J.G., S.J. Herbst, and A.R. Mahon. 2017. The reproductive viability of Grass Carp (*Ctenopharyngodon idella*) in the western basin of Lake Erie. *Journal of Great Lakes Research* 43(2):405-09.

Williamson, M. and Fitter, A., 1996. The varying success of invaders. *Ecology*, 77(6), pp.1661-1666.

Witt, A.M., J.M. Dettmers, and C.E. Cáceres. 2005. *Cercopagis pengoi* in southwestern Lake Michigan in four years following invasion. *Journal of Great Lakes Research* 31(3):245-252.

Wittman M.E., C.L. Jerde, J.G. Howeth, S.P. Maher, A.M. Deines, and J.A. Jenkins. 2014. Grass Carp in the great lakes region: establishment potential, expert perceptions, and re-evaluation of experimental evidence of ecological impact. *Canadian Journal of Fisheries and Aquatic Sciences* 71(7):992-1002.

Wittman, K.J. 2006. Distribution and invasion potential of the Ponto-Caspian Mysidae (Mysidacea: Crustacea Malacostraca: Peracarida: Mysida). In *Neobiota: From Ecology to Conservation* (W.F. Rabitsch, F. Klingenstein and F. Essl, editors). Federal Agency for Nature Conservation, Vienna.

Wurster, C., W. Patterson, D. Stewart, J. Bowlby, and T. Stewart. 2005. Thermal histories, stress, and metabolic rates of Chinook salmon (*Oncorhynchus tshawytscha*) in Lake Ontario: evidence from

intra-otolith stable isotope analyses. *Canadian Journal of Fisheries and Aquatic Sciences* 62 (March, 2005): 700-713.

Wyda, J. - The Ecosystem Center, the Marine Biological Laboratory, Woods Hole, MA.

Yerger, R.W. 1977. Fishes of the Apalachicola River. *Florida Marine Research Publications* 26:22-33.

Young, B.A., T.L. Welker, M.L. Wildhaber, C.R. Berry, and D. Scarnecchia, editors. 1997. Population structure and habitat use of benthic fishes along the Missouri and lower Yellowstone rivers. Annual Report of Missouri River Benthic Fish Study PD-95-5832. U.S. Army Corps of Engineers and the U.S. Bureau of Reclamation. 207 pp.

Yuille, M.J., T.B. Johnson, S.E. Arnott, and L.M. Campbell. 2012. *Hemimysis anomala* in Lake Ontario food webs: stable isotope analysis of nearshore communities. *Journal of Great Lakes Research*. 38(Supplement 2):86-92.

Zhang, L. N., X. Li, W. Lu, H. X. Shen, and Y. K. Luo. 2011. Quality predictive models of Grass Carp (*Ctenopharyngodon idellus*) at different temperatures during storage. *Food Control* 22:1197-1202.

Zhao, Z. G., S. L. Dong, F. Wang, X. L. Tian, and Q. F. Gao. 2011. Respiratory response of Grass Carp (*Ctenopharyngodon idellus*) to temperature changes. *Aquaculture* 322:128-133.

Znachor, P., J. Hejzlar, J. Vrba, J. Nedoma, J. Seda, K. Šimek, J. Komárková, J. Kopáček, J. Šorf, J. Kubecka, J. Matena, M. Říha, J. Peterka, M. Čech, and M. Vašek. 2016. Brief history of long-term ecological research into aquatic ecosystems and their catchments in the Czech Republic. Part I: Manmade reservoirs. Volume 1. 1st edition. Institute of Hydrobiology, Biology Centre CAS, České Budějovice.