

TOPICAL REVIEW • OPEN ACCESS

Essential gaps and uncertainties in the understanding of the roles and functions of Arctic sea ice

To cite this article: Sebastian Gerland *et al* 2019 *Environ. Res. Lett.* **14** 043002

View the [article online](#) for updates and enhancements.

Recent citations

- [Recent Arctic Ocean Surface Air Temperatures in Atmospheric Reanalyses and Numerical Simulations](#)
Allison B. Marquardt Collow *et al*
- [Polar Code application areas in the Arctic](#)
Meric Karahalil *et al*
- [Quantification of the Arctic Sea Ice Driven Atmospheric Circulation Variability in Coordinated Large Ensemble Simulations](#)
YuChiao Liang *et al*

Environmental Research Letters



TOPICAL REVIEW

OPEN ACCESS

RECEIVED
31 August 2018

REVISED
5 February 2019

ACCEPTED FOR PUBLICATION
22 February 2019

PUBLISHED
4 April 2019

Original content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](#).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



Essential gaps and uncertainties in the understanding of the roles and functions of Arctic sea ice

Sebastian Gerland¹ , David Barber², Walt Meier³ , Christopher J Mundy², Marika Holland⁴, Stefan Kern⁵, Zhijun Li⁶, Christine Michel⁷, Donald K Perovich⁸ and Takeshi Tamura⁹

¹ Norwegian Polar Institute, Fram Centre, Tromsø, Norway

² Centre for Earth Observation Science, University of Manitoba, Winnipeg, Canada

³ National Snow and Ice Data Center, University of Colorado, Boulder, United States of America

⁴ National Center for Atmospheric Research, Boulder, CO, United States of America

⁵ Integrated Climate Data Center, CEN, University of Hamburg, Hamburg, Germany

⁶ State Key Laboratory of Coastal and Offshore Engineering, Dalian University of Technology, Dalian, People's Republic of China

⁷ Fisheries and Oceans Canada, Winnipeg, Manitoba, Canada

⁸ Thayer School of Engineering, Dartmouth College, Hanover, NH, United States of America

⁹ National Institute of Polar Research, Tokyo, Japan

E-mail: gerland@npolar.no

Keywords: sea ice, arctic, climate, observation gaps, knowledge gaps

Abstract

While Arctic sea ice is changing, new observation methods are developed and process understanding improves, whereas gaps in observations and understanding evolve. Some previous gaps are filled, while others remain, or come up new. Knowing about the status of observation and knowledge gaps is important for interpreting observation and research results, interpretation and use of key climate indicators, and for research and observation planning. This paper deals with identifying some of the important current gaps connected to Arctic sea ice and related climate indicators, including their role and functions in the sea ice and climate systems. Subtopics that are discussed here include Arctic sea-ice extent, concentration, and thickness, sea-ice thermodynamics, age and dynamic processes, and biological implications of changing sea ice. Among crucial gaps are few *in situ* observations during the winter season, limited observational data on snow and ice thickness from the Arctic Basin, and wide gaps in biological rate measurements in or under sea ice. There is a need to develop or improve analyzes and products of remote sensing, especially for new sensors and technology such as remotely operated vehicles. Potential gaps in observations are inevitably associated with interruptions in long-term observational time series due to sensor failure or cuts in observation programmes.

1. Introduction and background

The identification of gaps in understanding the roles and functions of Arctic sea ice is important for assessing data and new findings of Arctic sea ice changes, and to develop research and monitoring strategies. The goal of this paper is to summarize and detail important current gaps of knowledge, and observation gaps related to Arctic sea ice, and, to a limited extent, to discuss the recent development of detecting gaps and ways and strategies for filling them. The paper includes discussing gaps in connection with key sea ice climate indicators.

On the first view, it might be easier to detect observation gaps rather than knowledge gaps. But the two

things go hand in hand; more knowledge, i.e. an improved understanding of the system is also necessary to make better decisions on where to do which type of observations.

Our discussion of gaps is not comprehensive. Sea ice is a broad and interdisciplinary scientific subject with numerous sub-disciplines and processes at play, and we by no means claim to present here a complete overview on Arctic sea ice gaps, rather a selection of some of the important ones. Our focus is on the observational record and methods used to observe a variety of sea ice conditions.

The starting point for this work is based on the sea ice chapter of the recent Arctic Marine Assessment Program (AMAP) Snow, Water, Ice and Permafrost

Assessment (SWIPA) 2017 report (AMAP 2017), where knowledge and observation gaps were addressed regarding (i) sea-ice extent, concentration, and thickness, (ii) sea-ice thermodynamics, age and dynamic processes, and (iii) biological implications of changing sea ice, respectively (Barber *et al* 2017). Of those topics, especially ice extent, thickness and age are often used as indicators of climate change, which makes existence or potential existence of gaps even more relevant to be discussed.

Gaps that are addressed in the SWIPA 2017 report are, for example, the fact that there are very few *in situ* observations in winter; limited data available on snow thickness from the Arctic Basin; and that there is the need to develop or improve analyzes and products of remote sensing, including for example new sensors or satellites.

How have the gaps discussed here changed over time? On one hand, the number of gaps is gradually becoming less since some scientific questions are answered or observation systems put in place. On the other hand, more sophisticated methods and models may also need better input data, which do not always exist, and they might identify new processes for consideration, meaning that new gaps will arise. Changes in Arctic sea ice can lead to new or different sea ice processes becoming dominant, which may lead to new knowledge requirements, and by extension new methods that will need to be tested and validated.

Information on how gaps have changed over time will be supported by comparing statements and sub-chapters on knowledge gaps from the SWIPA 2017 report (AMAP 2017) and recent literature with statements in ACIA (2005), SWIPA 2011 (AMAP 2011) and other earlier assessments.

Uncertainties of observational methods are not explicitly gaps of knowledge, but they can affect our ability to gain new insights on processes of interest. For example, if a method is too coarse to reveal heterogeneity at some spatial scales, it is difficult to assess the role of that heterogeneity on relevant processes. Therefore, we also include to some extent uncertainties in the discussion. Under uncertainties we understand the sum of several contributions: (i) the difference between a true quantity and its observed value—bias or accuracy; (ii) the repeatability with which a quantity is observed by the same method—precision; (iii) the degree with which the spatio-temporal distribution of a quantity is observed by the same method—representativity error; and (iv) the degree with which the spatio-temporal distribution of a quantity is observed by the same method with unchanged contributions (i)–(iii)—stability. When discussing the uncertainties of different methods, reasons for uncertainties are often related to the physics of sensors, assumptions/simplifications of the methods, the sea ice system, or economic limits. While some uncertainties can be reduced through ideas and action, others cannot be changed. Methods that have low

accuracy or coarse spatial or temporal resolution often have other advantages that makes them attractive for applications where the accuracy is not the one and only criterion for the usability/usefulness of the method but the high stability is more relevant.


Although we focus here on gaps related to Arctic sea ice, several of the points raised apply equally for Antarctic sea ice, especially in recent years as the Arctic sea ice has transitioned to a state with a larger portion of seasonal sea ice at the cost of older, multiyear ice.

2. Existing gaps and uncertainties

2.1. Gaps in knowledge for sea ice concentration, extent and thickness

The sea ice concentration and extent record from passive microwave instruments provides a nearly complete and consistent record since late 1978. On one hand, this almost 40 year long time series is a very important tool for climate studies, and it is used widely as a climate indicator to quantify and illustrate Arctic sea ice change. On the other hand, the fact that the time series is limited to four decades, that it has limited spatial resolution, some varying data quality, and uncertainty depending on season and changing satellite sensors over time all limit the applicability of the time series. Intercalibration between sensors has not always been optimal (Eisenman *et al* 2014). This could be improved via use of longer overlaps period that are now available, employing better calibrated source data and, as is now being done in some products (Comiso *et al* 2017, Lavergne *et al* 2019), implementing adaptive time-varying algorithm coefficients that adjust for changing surface conditions. Nevertheless, there is a looming gap due to the potential loss of satellite coverage. Passive microwave sensors have been routinely launched since 1987 and for many years multiple sensors have been in orbit. However, recent failures have reduced the number currently operating sensors and have significantly increased the risk of a gap in coverage in the next years. AMSR-E failed in 2011, and the DMSP F-19 SSMIS failed in early 2016 after only two years of operation. In April 2016, the F-17 SSMIS started behaving anomalously and its data quality has been reduced. The suite of DMSP SSMI/SSMIS instruments has been the workhorse of passive microwave observations, but all currently operating (as of June 2018) SSMIS instruments (F-16, F-17, F-18) have been in orbit for at least 8.5 years, well beyond their design lifetime of three years (figure 1). The JAXA AMSR2 sensor, operating for a little over six years (as of June 2018) is the youngest passive microwave sensor, but it is now also past its design lifetime (five years). The only remaining sensor potentially ready to be launched in the near future (the DMSP F-20 SSMIS) has been cancelled.

While discussions concerning follow-on sensors are ongoing (including an AMSR follow-on, a



EUMETSAT instrument on their METOP constellation, and possible new US radiometers) these are all likely to be a minimum of five years from launch. At that point, the youngest sensor commonly used for climate monitoring (AMSR2), if still operating, would be nine years old. There is a passive microwave sensor (MWRI) on the Chinese FengYung-3B and -3C platforms. This has not been employed as part of the long-term passive microwave sea ice climate record, but has potential to fill the looming gap due to the loss of current capabilities. First intercalibration efforts of the FY-3C satellite data with other passive microwave satellite data have been made (Li *et al* 2016, Wang *et al* 2018, Wu and Liu 2018).

Another important, but more difficult to measure sea ice climate indicator is sea ice thickness. Significant development has been made to quantify sea-ice thickness and its seasonal and interannual change. However, there are still a number of significant gaps in knowledge about sea-ice thickness and the methods for measuring it. One method to estimate sea-ice thickness is to measure freeboard from aircraft or satellite and calculate ice thickness from that. Here, there is a gap in interpreting the radar and laser returns to yield accurate freeboard measurements (specifically for radar altimetry; see recent work by Ricker *et al* 2017, King *et al* 2018). Second, better snow and ice density information is needed to convert freeboard estimates into ice thickness. The most significant gap in this context, however, is the lack of reliable basin-scale snow depth observations (see e.g. Webster *et al* 2018). The snow weighs down the ice, changing the

freeboard to thickness ratio. For laser altimeters, which reflect off the top of the snow surface, the information about snow depth is only required for the freeboard-to-thickness conversion. For radar altimeters, which penetrate the snow depending on its properties, the snow depth is required for both, accurate freeboard retrieval and freeboard-to-thickness conversion. The comparably poor spatiotemporal coverage with altimeter measurements and changing satellite sensors limit the present-day maturity of sea-ice thickness products further. Other methods that do not have the same problem are *in situ* surveys with classical thickness measurements from drillings, and those using electromagnetics (from air and ground), thermal methods (e.g. Mäkynen and Karvonen 2017) or microwave radiometry like from SMOS (e.g. Kaleschke *et al* 2016). However, these types of measurements have a number of limitations. Drillings are only possible in a few places of the Arctic at a given time. Electromagnetic measurements do not distinguish between snow and sea ice, and wrong information of snow thickness can lead to over- or underestimation of the ice thickness. Thermal measurements are only useful and provide accurate thickness information for relatively thin sea ice, i.e. below ~ 0.5 m, and the same applies for the usage of microwave radiometry.

It is also important to mention that CryoSat-2 is beyond its life time, and that CryoSat-2 is the only currently flown radar altimeter providing data north of 81.5° N, covering the Arctic Ocean up to 88° N. The laser altimeter on ICESat-2, launched in September

2018, covers latitudes up to 86° N. In any case there remains an observation gap for the central Arctic Ocean.

The combination of ice extent, ice concentration and ice thickness information to ice volume estimates is another important climate indicator, and has been used in local (Gerland and Renner 2007) and panarctic contexts (Laxon *et al* 2013). Sea ice volume estimate is a powerful climate indicator, because it takes into account dynamic effects, which can lead to apparent reductions or increases of sea ice amounts due to for example ridging, rafting and spreading existing sea ice into larger areas. Complete sea ice volume datasets are rare since they require both ice extent, concentration and thickness data with sufficient spatial and temporal resolution (e.g. Kwok and Cunningham 2015, Lindsay and Schweiger 2015, Tilling *et al* 2018). The uncertainty of such sea-ice volume estimates naturally depends on the uncertainties of the input data which we discuss in the following sections. Alternatively, coupled ice-ocean models into which sea-ice information is assimilated, as e.g. PIOMAS (Zhang and Rothrock 2003), provide a reasonable measure of the Arctic sea-ice volume, which can be applied for further studies, such as on sea ice volume fluxes (e.g. Schweiger *et al* 2011, Zhang *et al* 2012, Zhang *et al* 2017). However, evaluating the quality of such results in the context of their value as an independent powerful climate indicator remains a challenge and requires careful evaluation taking into account the model's capability to properly resolve physical processes at the grid resolution used.

2.2. Sea Ice concentration and extent uncertainty

Estimates of uncertainty from passive microwave sea-ice concentration (SIC) products is a challenge owing to the large scale, varying surface properties, and limited availability of comparison data; summer is especially challenging due to the effect of melt water on the passive microwave signal (e.g. Kern *et al* 2016). Evaluation of SIC products is an ongoing effort—because new algorithms are still being developed (Kongoli *et al* 2011, Tikhonov *et al* 2015), because new satellite sensors are becoming available (e.g. AMSR2, Beitsch *et al* 2014), and because new independent data sets are emerging which help to better quantify SIC uncertainties. SIC derived from clear-sky satellite optical imagery such as MODIS for summer Arctic sea ice (Rösel *et al* 2012) can help quantify uncertainties in satellite microwave radiometry-based SIC during summer melt (Kern *et al* 2016), but such sensors depend on clear sky and daylight conditions. Thin ice thickness distribution derived, under freezing conditions and for close to 100% sea ice, from the ESA SMOS sensor (Kaleschke *et al* 2012, Huntemann *et al* 2014, Tian-Kunze *et al* 2014, Kaleschke *et al* 2016) can be used to infer biases in SIC over thin ice (Heygster *et al* 2014).

More effort has been dedicated during recent years to assess the retrieval uncertainty of SIC or at least providing SIC data sets with an uncertainty estimate—which is an important pre-requisite for assimilating SIC data into numerical models. A few approaches have been developed. One is to apply Gaussian error propagation to the equations used to estimate SIC and to calculate SIC uncertainty as a function of algorithm tie point, brightness temperature, and gridding uncertainties (Kern 2004, Spreen *et al* 2008). This is used in the EUMETSAT OSI-SAF SIC data set (Tonboe *et al* 2016, Lavergne *et al* 2019) and the ESA-CCI SIC data set (<http://esa-cci.nersc.no>, Lavergne *et al* in 2019). A second approach is to take the standard deviation of the SIC within a certain area as a measure for the quality of the SIC estimate (Peng *et al* 2013, Meier *et al* 2014) based on the fact that regions of higher SIC variability will have higher uncertainty because of the low sensor spatial resolution. The Enhanced NASA Team ('NASA Team 2') algorithm takes advantage of the iterative nature of the algorithm and estimates the uncertainty from the variability of the concentration as the iteration converges to a minimum cost function (Brucker *et al* 2014). These approaches provide a measure of the precision only.

More difficult is a quantitative estimate of the uncertainty of the sea-ice area or extent, which goes beyond a simple computation of the variation in these quantities over time. Factors contributing to uncertainty in sea-ice area and extent are (i) the choice of the threshold ice concentration used, (ii) filters used to flag spurious weather-influence induced SIC over open water (see e.g. Lavergne *et al* 2019), (iii) the accuracy of the SIC itself (e.g. low biases during melt conditions), (iv) filters used to flag land-spillover effects, and (v) use of different land masks.

Meier and Stewart (2019) presented an approach to quantify the variability in extent based on the above five factors as a proxy for an uncertainty estimate. They compared sea ice total extent from several different products and found a variability range of 500 000 square kilometers or more, depending on season. This generally corresponds to relative biases in ice edge location of 25–50 km and is due the factors mentioned above. The differences in extent, while varying seasonally, are consistent over the years for similar times of years. This means that trends and variability between the different products are more consistent than the absolute extents. In another part of the Meier and Stewart (2019) study, they found that using a consistent algorithm and processing yielded very stable results with standard deviations estimates at ~20 000 square kilometers. However, estimating the accuracy of sea ice area remains challenging because of potential biases in the SIC due to snow and sea-ice surface processes and weather effects not accounted for in the algorithms.

Another aspect is the uncertainties in trends. Trend uncertainties are often given as the uncertainty

(standard deviation) of the linear fit. However, trend uncertainties are affected by the limited consistency of the satellite record. This issue has not been thoroughly investigated, but Eisenman *et al* (2014) found that trend uncertainties may be higher than the quoted standard deviation of the trend. Another issue when examining trends and anomalies is the common assumption that while absolute extent or area values may be biased due to surface melt, snow, and ice conditions, confidence in trends would still be high because these surface effects are consistent over the years (i.e. every summer, surface melt causes an underestimation in concentration and area). This assumption, however, might not be valid anymore because of the observed earlier melt onset and the occurrence of more highly degraded or 'rotten' ice and resulting underestimation of the ice cover by passive microwave instruments (Barber *et al* 2009, Meier *et al* 2015). Specifically, for certain times of the year negative trends in sea-ice extent could appear larger than they are.

2.3. Sea ice thickness uncertainty

Regarding ice thickness from altimetry, Zygmunowska *et al* (2014) reported that when uncertainties in the input parameters for the freeboard-to-thickness conversion are considered appropriately, the changes in sea-ice volume between the ICESat and CryoSat-2 periods are less dramatic than was reported in the literature at that time; this view was later shared by Kwok and Cunningham (2015). Uncertainty sources in the freeboard-to-thickness conversion and the freeboard retrieval itself are manifold. Various publications have shown that both sea ice and snow density play a major role in the uncertainty budget of the sea-ice thickness derived from satellite radar altimetry (e.g. Alexandrov *et al* 2010, Forsström *et al* 2011, Kern *et al* 2015).

The validation of these data is in its infancy. Many uncertainties still exist in the derivation of the basic parameter, the freeboard (Kwok 2014, Ricker *et al* 2015). In addition, data used for the validation might also not be free of potential biases or have unknown uncertainty (e.g. Kurtz *et al* 2013, Lindsay and Schweiger 2015), which complicates validation procedures.

Laxon *et al* (2013) used additional data to distinguish first-year and multi-year ice and applied different sea ice densities and snow depth values for the freeboard-to-thickness conversion. Zygmunowska *et al* (2013) suggested an approach that would allow this discrimination by using CryoSat-2 data itself—without the need for additional data and with the potential of improved freeboard retrieval.


Several publications (Ricker *et al* 2014, 2017, Kurtz *et al* 2014, Tilling *et al* 2015, 2018) address enhanced solutions for the freeboard retrieval and freeboard-to-thickness conversion using CryoSat-2 data. The limited range resolution of CryoSat-2, the unknown tracking point of the impinging radar wave in the ice-

snow system, the change in radar wave propagation speed in a snow layer of unknown thickness, and the presence of saline snow on seasonal sea ice can complicate the freeboard retrieval (Kwok 2014, Nandan *et al* 2017). One of the main sources of uncertainty is the snow depth on sea ice (Forsström *et al* 2011, Kwok 2014, Ricker *et al* 2015). Its retrieval remains a challenge.

The laser altimetry satellite ICESat ceased operation in 2009, but is still useful for extending the CryoSat-2 thickness time series backwards. However, stitching together these products is difficult considering the still limited knowledge regarding uncertainties. For example, Bi *et al* (2014) analyzed ICESat data and found large estimation errors over thick ice. Radar altimeter data were also provided by the ERS1/2 satellites (Laxon *et al* 2003) and the Envisat satellite (Giles *et al* 2008, Schwegmann *et al* 2016, Paul *et al* 2018). One of the challenges here, which is only solved partly (e.g. Paul *et al* 2018), is to produce a consistent data record of freeboard and sea-ice thickness extending from the ERS1/2 era, starting in 1993, through today (CryoSat-2 and SARAL-AIiKa). The ICESat-2 satellite will continue the sea ice altimeter record and its new technologies will help refine estimates of thickness and potentially also of the snow depth on sea ice (Lawrence *et al* 2018), particularly if the radar altimeter on CryoSat-2 continues to operate beyond the start of the operative phase of the laser altimeter on ICESat-2.

2.4. Gaps in knowledge for sea-ice thermodynamics, age and dynamic processes

Information on snow thickness over Arctic sea ice is crucial, since snow strongly affects thermodynamic ice growth, melt pond formation, and radiative transfer. Information on snow thickness on Arctic sea ice is still sparse, although the amount of data has increased recently due to new *in situ*, airborne and satellite surveys (e.g. Laxon *et al* 2013, Renner *et al* 2014, Haas *et al* 2017, Rösel *et al* 2018). Reanalysis data can fill in where observations are lacking, but intercomparisons with *in situ* measurements have shown that they do not always match well (Boisvert *et al* 2018). Autonomous buoys with snow thickness sensors (ultrasonic sensors or active thermistor chains as used in IMBs) give quantitative time series data with a higher temporal resolution, and usually with a smaller footprint than airborne and satellite surveys (e.g. Brucker and Markus 2013, Maaß *et al* 2013, 2015). These instruments can directly measure changes in ice thickness and delineate surface and bottom melt (Perovich and Richter-Menge 2015). However, to collect representative data for a region requires many buoys. So far, the total amount of buoys at a given time in the Arctic is still rather limited, and the distribution often not very regular, so that vast areas might be without buoys at a given time (figure 2). Initiatives such as the International Arctic Buoy Programme



(Rigor *et al* 2016, 2000) help coordinate deployments of sea ice buoys. Sophisticated buoy setups that combine measurements related to surface energy and mass balance have been developed recently (e.g. Jackson *et al* 2013, Wang *et al* 2014, 2016, Hoppmann *et al* 2015), as well as buoys that float and have the potential to ‘survive’ a regional ice-free summer (see e.g. Polashenski *et al* 2011). Such new developments will contribute to denser and accurate autonomously collected sea ice datasets. Understandably, relatively cheap drifter buoys with GPS but not much other sensors are the most frequent buoys used in the Arctic, while more sophisticated and multi-sensor setups are relatively rare. However, for collecting information on sea ice thermodynamics more ice mass balance buoys (IMBs) would be necessary, along with global data archives where all respective data would be accessible for the scientific community.

Recent Arctic expeditions have made increased use of new technology with mobile sensor platforms. These include autonomous underwater vehicle, unmanned airborne vehicle, remotely operated vehicle and instrument sledges. Some of these systems are still at the development stage, but with new systems becoming standard tools, it will be possible to obtain high quality sea ice data over larger spatial scales in the near future. There are still fewer *in situ* sea ice surveys undertaken in winter months than summer months, but recent (Granskog *et al* 2016, 2018, Assmy *et al* 2017, Rösel *et al* 2018) and planned interdisciplinary expeditions (e.g. the Multidisciplinary drifting Observatory for the Study of Arctic Climate (MOSAIC), www.mosaicobservatory.org) are adding to the

existing winter data and providing insights into the changing Arctic sea ice system.

Ice age, also often used as a climate indicator (e.g. Perovich *et al* 2018, Stroeve and Notz 2018) is now less correlated with ice thickness than it used to be (Tschudi *et al* 2016a), since the older ice classes have become thinner. First-year ice with little snow cover can reach a thickness similar to second-year ice (Notz 2009), which has also been found in a recent field experiment in the Arctic Ocean (Granskog *et al* 2016). This can be a challenge for deriving ice classes from ice thickness data (Hansen *et al* 2014). Distinguishing first- and second-year ice by means of its thickness was probably also not always trivial previously, and is complicated using satellite microwave radiometry as well (Comiso 2006), but in total the range of thicknesses has become narrower, with the thickest ice classes becoming rare, making this more difficult. Assessing the age of ice can be supported by information on its physical properties, for example via their signature in satellite microwave scatterometry (Belmonte Rivas *et al* 2018), or by tracking ice movement over time (Tschudi *et al* 2016a). The Lagrangian tracking of ice parcels used here is subject to errors in the motion tracking, especially for regions with high spatial variability in ice dynamics, types and age within the tracked ice parcels which triggered attempts for a revision of the sea-ice age retrieval method (Korosov *et al* 2018). But the general good agreement between the ice age fields and ice thickness and multi-year ice extent estimates (Maslanik *et al* 2011) make this pan-Arctic dataset still a valuable tool to assess sea ice age on large spatial scales.

The Lagrangian ice age fields are derived from sea ice motion products (e.g. Lavergne *et al* 2010, Tschudi *et al* 2016b), which use feature tracking algorithms to estimate sea ice displacement in satellite imagery. The most commonly used source is passive microwave because of its all-sky capability. However, the low spatial resolution is a significant limitation on the precision and accuracy of the estimates and important small-scale features such as lead formation and ridging cannot be detected. The growing catalogue of SAR sensors has the potential to routinely provide daily, near-Arctic wide coverage at much higher spatial resolutions (e.g. RADARSAT Constellation), though the passive microwave record will still be valuable for long-term context.

Scientists are monitoring and studying sea ice as it drifts through Fram Strait, the area between Greenland and Svalbard, a major gate and only deepwater connection connecting the Arctic Ocean to the other world oceans (Kwok *et al* 2004, Spreen *et al* 2009, Smedsrud *et al* 2011, Hansen *et al* 2013, Renner *et al* 2014, Smedsrud *et al* 2017). Through this gate, most of the sea ice that is exported from the Arctic Ocean passes through here. Findings from here can give information on a larger area 'upstream' (Kwok 2009, Hansen *et al* 2013, Krumpen *et al* 2016). With current technology and infrastructure, continuous monitoring at such gates is only possible with moored instruments (e.g. upward looking sonars for measuring sea ice draft), satellite remote sensing, and local meteorological information. Crucial gaps in this context are or can be that there are only few or no observation sites at a gate, and potential data gaps due to instrument failures. Additional limitations arise from current remote sensing capabilities for studying the complex dynamics at such gates and a mis-match in the spatial resolution of the applied data sources (e.g. Ricker *et al* 2018). Given the large across-strait variability in sea-ice type, thickness and motion, SAR-based estimates of the sea-ice motion are possibly the best way to proceed (Smedsrud *et al* 2017) provided enough *in situ* data, e.g. from buoys, allow continued evaluation.

2.5. Sea-ice thermodynamics, age and dynamic processes uncertainties

Information about ice thermodynamics with high temporal resolution is often retrieved from IMBs and *in situ* measurements. Uncertainties are related to the question how homogenous the ice cover is, or in other words how representative a IMB site is for the region it is placed in. Depending what technology in IMBs is used, detection of snow ice and superimposed ice might be not trivial, and this can lead to an overestimation of the snow thickness and an underestimation of the ice thickness.

Drifter buoys with iridium transmitted GPS positions have a better accuracy than Argos-based buoys without GPS earlier, but ice motion can be

underestimated due to motion with changing drift direction between individual position recordings. For avoiding that, we recommend position recordings at least once an hour. Sophisticated methods have been developed that use iridium transmission GPS beacons to examine sea ice motion following methodology developed for atmospheric dynamics (e.g. Lukovich *et al* 2015). Methods of floe tracking from satellite products are well suited to local and regional scale motion tracking (e.g. Komarov and Barber 2014, Itkin *et al* 2018) and detection and modelling of sea ice hazards to marine navigation (e.g. Barber *et al* 2014). A key gap is to develop an improved understanding of sea ice dynamic growth as a function of sea ice ridging and rubbing. Essential for this is also improved information on the spatial distribution and variability of surface roughness. The process of dynamic growth is expected to result locally and regionally in thicker ice classes even though thermodynamic growth will be reduced (see case study in Itkin *et al* 2018).

2.6. Biological implications of changing sea ice; uncertainties and gaps in knowledge

Knowledge and data gaps continue to limit the certainty of predictions concerning how the ice-associated ecosystem and its coupling to the pelagic and benthic food webs will respond under a changing climate. The relative ease of accessing coastal landfast ice versus the central Arctic ice-pack has resulted in biogeochemical observations being focused in key locations around the periphery of the Arctic Ocean, leaving large data gaps for vast areas of the shelves and central basins (e.g. Leu *et al* 2015). Similarly, logistical considerations, as well as a focus on the biologically productive period, have skewed the seasonal observational base towards spring-summer when environmental conditions are more favorable to field studies. The few autumn-winter studies that do exist (e.g. Niemi *et al* 2011, Niemi and Michel 2015, Assmy *et al* 2017, Melnikov *et al* 2016, Olsen *et al* 2017) reveal interesting ecological features of the ice-associated ecosystem, calling for a better characterization of biogeochemical processes during this time of year. Little remains known about the distribution and bloom development of sub- and under-ice primary producers and there are few within-ice primary production estimates.

Baseline information on multi-year sea ice communities encompassing biomass, productivity and biodiversity, and role in the cycling of carbon and other elements, is also an important knowledge gap which requires urgent attention given the rapid decline in Arctic multiyear ice. With multiyear ice potentially being more productive than traditionally assumed (Lange *et al* 2017), there is a critical need for rate measurements in multiyear ice. The likelihood of changing ice and ocean conditions enhancing the potential for toxin-producing algal blooms in the

Arctic highlights a need for a better understanding of the presence, distribution, and recurrence of these species and their influence on Arctic food webs, as well as a need to improve the reporting of toxin occurrences across the Arctic. Repeat observations of higher trophic levels are also required, including fishes and marine mammals, in order to constrain their ecological responses and distributional changes in relation to the changing sea-ice cover.

Owing to the interacting effects of light and nutrients as the main factors controlling primary production at the sea ice interface and under the ice, uncertainties associated with observations and projections of the physical variables by which they are controlled, confound the capacity to predict the future structure and function of the sea-ice ecosystem. On Arctic shelves, sea ice and sea ice export contribute significantly to structure spatially diverse productivity regimes, also influenced by nutrient inventories and dynamics (Michel *et al* 2015, Tremblay *et al* 2015). As these different productivity regimes respond differently to on-going and future Arctic change (Ardyna *et al* 2011), a key challenge remains in resolving meso- to large-scale heterogeneity across the shelves and basins, for primary production estimates. In addition, poorly constrained estimates of the contribution of ice algae and phytoplankton to primary production and transfers to the food web, including harvest resources, require further attention. Furthermore, these uncertainties increase when predicting the responses at higher trophic levels, due to the added complexity of interactions at the species, population, community and ecosystem level.

Beyond gaps connected with sea ice ecosystem observations, it is important to mention that there is also a rather limited amount of modelling studies that address both the complexity of the physical system, and the wide range of components of the ecosystem (e.g. Duarte *et al* 2015, 2017).

3. Changes of gaps over time

In table 1, we give an overview on Arctic sea ice observation gaps. Some gaps disappear, while new ones emerge. This is expected and connected to both the change in sea ice conditions and processes, and in the research and monitoring activities scientists develop. New methods, designed to fill gaps, require proper validation and development in order to produce accurate data with high temporal and spatial resolution. Changing sea ice conditions result in changing knowledge needs. For example, certain processes can become less important while new, not yet fully understood processes can become relevant. Additionally, ice conditions can go beyond known examples, such as when new record low ice extents are registered, and regions become ice free, or change from perennial to seasonal ice cover. Because of this, it

is critical to avoid potential gaps, for example in the passive microwave satellite record, to continue research to enhance knowledge and understanding, and to extend observations to better characterize the spatial heterogeneity of conditions. Redundancy in observation systems is also desirable. Previous assessment reports treated gaps of knowledge with varying levels of depth.

In the ACIA report (2005), sea ice related gaps of knowledge were not addressed specifically. However, there were more general statements made that more observations, process studies and monitoring would be necessary; sea ice freezing and melting was mentioned. Recommendations included more use of autonomous platforms, improved physics-ecosystem modelling, and establishing of data bases. Parts of these recommendations were followed up, but for most of them there is still a way to go. Several international collaborations, for example those that were a part of the International Polar Year 2007–2009 led to more observations, interdisciplinary approaches, and modern technological solutions.

The fifth assessment report of the IPCC (Vaughan *et al* 2013) discussed gaps in the observation record of sea ice extent, and the approach to use climatology data to fill gaps in the time series for times prior to the passive microwave satellite era (prior to 1979). This gives insights into a longer time series, but it also increases the uncertainties and size of error bars of these earlier periods 1870–1953 (twice) and 1954–1978 (factor 1.5), relative to the period from 1979 onwards. The frequent use of sea ice extent data from passive microwave satellite observations as an indicator of climate change illustrates the need to improve these datasets, to better characterize the uncertainties, and to ensure continuation of the observation time series.

The first AMAP SWIPA report from 2011 (AMAP 2011) addresses several of the gaps that were also a subject of the most recent SWIPA report (AMAP 2017). This is not particularly surprising since there was only about six years between the two reports. The lack of basin-wide snow thickness measurements was highlighted already. Ice thickness data from satellite sensors was still in development; at the time of SWIPA 2011, the methods were not yet ready for routine use. As detailed above, there is still some way to go, but the situation has improved. Several newer publications deal with use of satellite-based altimetry data (e.g. Laxon *et al* 2013, Kurtz *et al* 2014, Ricker *et al* 2017, Tilling *et al* 2018, Paul *et al* 2018), and one has more insights into the nature of the data received. In September 2018, the NASA laser altimetry satellite ICESat-2 was launched, likely leading to better insights into sea ice thickness distributions and changes. SWIPA 2011 also mentioned shortcomings in process understanding regarding ecosystem processes, and few long-term biological sampling initiatives/programmes. Much progress has been made but

Table 1. Overview on observation gaps for Arctic sea ice.

Parameter	Method	Gap
		(1) No data prior to 1978
Ice extent	Passive microwave satellite sensors	(1) Lacking validation data, resulting in unclear uncertainty (1) Potential of future data gaps because of few satellites and long running times
Ice extent (high spatial resolution)	SAR satellite sensors	No Arctic-wide coverage as for passive microwave
Sea ice concentration	SAR and passive microwave, optical satellite RS	Spatial resolution, data processing algorithms, changing sensors, unclear uncertainty (1) Detailed snow thickness data and density data for conversion of freeboard to thickness is lacking
Sea ice thickness	Satellite-based altimeters	(1) Only few <i>in situ</i> and airborne validation datasets (1) Lack in interpretation knowledge of radar signals to derive correct freeboard (1) Gap of observation north of 88° N
Snow thickness	Satellites, airborne, <i>in situ</i>	Generally too few data
Sea ice mass balance	Autonomous buoys	Too few buoys running at a given time, large gaps with regions without buoys
Sea ice temperature	Autonomous buoys	Too few buoys running at a given time, large gaps with regions without buoys
Ice age	Ice thickness (all methods)	Ice age can be less easy derived from ice thickness, because ice thicknesses for different ice ages can be more similar now than earlier
Ice drift speed	Floe tracking, buoys	Data might be biased towards underestimating speeds; few buoys in parts of the Arctic
Biogeochemical and ecosystem properties and projections	Methods that depend on sea ice physics data and biogeochemical observations	(1) Large spatial gaps of observation; (2) Gaps in physics data result in gaps in ecosystem data where those are needed as input
Sea ice pollution	<i>In Situ</i> /sampling and remote sensing	Large spatial gaps of observation, and methods developed parallel to new pollutants coming up
Various parameters	Autonomous mobile and remote sensing based platforms	New platforms give new datasets, but calibration and validation is needed
Various physics and ecosystem parameters	Various methods	Lack of winter <i>in situ</i> data

establishing long-term time-series continues to be a challenge. Some examples of current efforts include the Joint Ocean-Ice Studies in the Beaufort Sea (<http://dfo-mpo.gc.ca/science/collaboration/jois-eng.html>), ArcticNet in the coastal Canadian Arctic (<http://www.arcticnet.ulaval.ca/>), the Distributed Biological Observatory in the Pacific sector of the Arctic (<https://www.pmel.noaa.gov/dbo/>), MOSJ Environmental Monitoring of Svalbard and Jan Mayen (<http://www.mosj.no/en/>), and the Nansen Legacy in the northern Barents Sea and adjacent Arctic Basin (<http://www.nansenlegacy.org>). Notably, some predictions from the SWIPA 2011 report were directly addressed by the scientific community, e.g. the predicted occurrence of dual algal blooms associated with longer open water seasons, studied in Ardyna *et al* (2011). The

upcoming MOSAiC project with fieldwork in 2019–20 will address gaps about ecosystem process understanding, and the need of more community-based observations. Community-based observing programs provide a mechanism for two-way knowledge transfers. This point was also highlighted in SWIPA 2017.

Limitations in sea ice observations present challenges for our ability to integrate observations with models to enhance understanding. The presence of short and/or inconsistent records, inadequate spatial and temporal sampling, and challenges in estimating observational uncertainty are all problematic for the optimal use of observations to inform models. This difficulty spans across possible model applications. For comparisons to climate simulations of change, the

influence of internal variability needs to be thoughtfully considered. Indeed, even when considering relatively long timeseries, such as that of the nearly 40 year passive microwave ice concentration, internal variability can play an important role (e.g. Ding *et al* 2017) and complicate the comparison of model simulations and observations. The challenge of quantifying observational uncertainty has implications for the useful assimilation of observations for model forecasting. The lack of adequate spatial and temporal sampling of variables like snow thickness, under-ice ecosystems, and processes that drive their variability makes it difficult to use these observations to improve the model representation of relevant processes. In general, given that gaps and limitations in observations will always be present, there is a need for better methods to integrate observations and models, which explicitly consider the limitations inherent in both the observations and the models.

Traditional knowledge, for example from Inuit, can provide important value-added content to data products and serves to make the data more relevant to northern users. The Inuit have for millennia used the sea ice as a travel corridor, a platform for resource harvesting and a key cultural icon for their way of life. Closer ties are beginning to form, where the western science approach is integrated with Inuit knowledge (e.g. Krupnik *et al* 2010), beginning at the start of the research process and working hand-in hand through to conclusion and reporting (e.g. Barber and Barber 2009).

The fact that gaps of knowledge are more explicitly discussed recently might also reflect that there is more awareness about the relevance and importance of topics and aspects that are not fully covered and solved within the current status of knowledge.

4. Future, perspectives and strategies to reduce gaps

Some options to reduce gaps are discussed in the previous section. Although knowledge gaps will always exist, a better understanding of the Arctic sea ice system will provide a better knowledge base for decision makers to choose options for sufficient protection and sustainable management.

Uncertainties arise in the development of new methods. However, once these methods are fully developed, they help to close knowledge and observation gaps, as it is the strategy with new satellites (e.g. ESA Sentinels, NASA ICESat-2). More and more available SAR satellite data (e.g. through Sentinel) with higher resolution than passive microwave and different polarisations can help to create and support almost daily improved Arctic wide ice extent datasets in the near future, to be useful for operational purposes, validation and climate research.

In some cases, rapid progress can be made to address knowledge gaps. For example, the recent awareness about microplastic pollution in sea ice has led to upcoming research. And while insights into processes, amounts and distribution are currently rather limited, first studies do indicate substantial levels and motivate future work (Obbard *et al* 2014, Peeken *et al* 2018).

Large international efforts and projects such as MOSAiC will provide more interdisciplinary datasets in current sea ice regimes especially for seasons with few existing observations (autumn, winter). Better access to the Arctic can be a result of both new and more platforms and ships and also changing ice conditions. In general, more information and data about Arctic sea ice is essential to close gaps: this concerns new data to be collected, also through more community-based observations including automatic sensors on commercial ships and aircrafts, but also better access to existing data via data portals and data bases. The value of the data also depends on consistency of datasets collected by different groups and in different regions. Efforts to coordinate future expeditions and data collection in a more sophisticated way will help to increase the possibilities of data use and intercomparability. This includes also considering possible future datasets, to be collected with methods that do not yet exist. Archiving sea ice, snow and water samples can allow for later analysis with improved or refined methods future technologies that do not exist yet.

Communication between scientists through publications, modern social media, international projects and symposia is and will be important. Building integrated, interdisciplinary communities of modelers, field observers, and remote sensing research is a critical component of future efforts. Models can be used to inform observational strategies. The observations can then be used to improve parameterizations used in models. This approach is necessary in order to develop a new generation of creative sea ice scientists, developing projects to address crucial gaps of knowledge.

Acknowledgments

We thank five anonymous reviewers for their valuable and constructive comments on an earlier version of this paper. We are grateful for the efforts of the contributing authors to sea ice chapter in the Snow, Water, Ice and Permafrost in the Arctic (SWIPA) 2017 report, and we thank the Arctic Monitoring and Assessment Programme (AMAP) and its secretariat for coordinating the work for that report, from which the sea ice chapter represents the base for this publication. We thank Anders Skoglund, Norwegian Polar Institute, for support producing the map in figure 2. The project ID Arctic, funded by the Norwegian Ministries of Foreign Affairs and Climate and Environment (programme Arktis 2030)

supported the collaboration of authors from Norway, Canada and USA. Zhijun Li's contribution was supported by the Global Change Research Program of China (2015CB953901). Marika Holland's contribution was supported by the National Center for Atmospheric Research, which is a major facility sponsored by the U.S. National Science Foundation under Cooperative Agreement No. 1852977.

ORCID iDs

Sebastian Gerland  <https://orcid.org/0000-0002-2295-9867>

Walt Meier  <https://orcid.org/0000-0003-2857-0550>

References

- ACIA 2005 *Arctic Climate Impact Assessment: Scientific Report*. (Cambridge: Cambridge University Press) p 1042
- Alexandrov V, Sandven S, Wahlin J and Johannessen O M 2010 The relation between sea ice thickness and freeboard in the Arctic *Cryosphere* **4** 373–80
- AMAP 2011 Snow, water, ice and permafrost in the arctic (SWIPA): climate change and the cryosphere *Arctic Monitoring and Assessment Programme (AMAP)* (Oslo, Norway) pp 538
- AMAP 2017 Snow, water, ice and permafrost in the arctic (SWIPA) 2017 *Arctic Monitoring and Assessment Programme (AMAP)* (Oslo, Norway) pp 269
- Ardyna M, Gosselin M, Michel C, Poulin M and Tremblay J-É 2011 Environmental forcing of phytoplankton community structure and function in the Canadian high Arctic: contrasting oligotrophic and eutrophic regions *Mar. Ecol. Prog. Ser.* **442** 37–47
- Assmy P et al 2017 Leads in Arctic pack ice enable early phytoplankton blooms below snow-covered sea ice *Sci. Rep.* **7** 40850
- Barber D G and Barber D 2009 *Two Ways of Knowing: Merging Science and Traditional Knowledge During the Fourth International Polar Year*. (Winnipeg, Canada: University of Manitoba Press) p 287
- Barber D G, Galley R, Asplin M G, De Abreu R, Warner K-A, Pucko M, Gupta M, Prinsenberg S and Julien S 2009 Perennial pack ice in the southern Beaufort Sea was not as it appeared in the summer of 2009 *Geophys. Res. Lett.* **36** L24501
- Barber D G, McCullough G, Babb D G, Komarov A S, Candlish L M, Lukovich J V, Asplin M, Prinsenberg S, Dmitrenko I and Rysgaard S 2014 Climate change and ice hazards in the beaufort sea *Elem. Sci. Anth.* **2** 000025
- Barber D G et al 2017 *Arctic sea ice. Chapter 5 of Snow, Water, Ice and Permafrost in the Arctic (SWIPA)*. (Oslo: AMAP) 103–36
- Beitsch A, Kaleschke L and Kern S 2014 Investigating high-resolution AMSR2 sea ice concentrations during the february 2013 fracture event in the Beaufort Sea *Remote Sens.* **6** 3841–56
- Belmonte Rivas M, Otosaka I, Stoffelen A and Verhoef A 2018 A scatterometer record of sea ice extents and backscatter: 1992–2016 *Cryosphere* **12** 2941–53
- Bi H, Huang H, Su Q, Yan L, Liu Y and Xu X 2014 An Arctic sea ice thickness variability revealed from satellite altimetric measurements *Acta Oceanol. Sin.* **33** 134–40
- Boisvert L N, Webster M A, Petty A A, Markus T, Bromwich D H and Cullather R I 2018 Intercomparison of precipitation estimates over the Arctic Ocean and its peripheral seas from reanalyses *J. Clim.* **31** 8441–8462
- Brodzik M J and Stewart J S 2016 *Near-Real-Time SSM/I-SSMIS EASE-Grid Daily Global Ice Concentration and Snow Extent, Version 5. [Sea ice concentration]*. (Boulder, Colorado USA: NASA National Snow and Ice Data Center Distributed Active Archive Center) (Accessed: 17 August 2018) (<https://doi.org/10.5067/3KB2JPLFPK3R>.)
- Brucker L, Cavalieri D J, Markus T and Ivanov A 2014 NASA Team 2 sea ice concentration algorithm retrieval uncertainty *IEEE Trans. Geosci. Remote Sens.* **52** 7336–52
- Brucker L and Markus T 2013 Arctic-scale assessment of satellite passive microwave-derived snow depth on sea ice using operation icebridge airborne data *J. Geophys. Res. Oceans* **118** 2892–905
- Comiso J C 2006 Impacts of the variability of second-year ice types on the decline of the Arctic perennial sea-ice cover *Ann. Glaciol.* **44** 375–82
- Comiso J C, Gersten R A, Stock L V, Turner J, Perez G J and Cho K 2017 Positive trend in the Antarctic sea ice cover and associated changes in surface temperature *J. Clim.* **30** 2251–67
- Ding Q et al 2017 Influence of high-latitude atmospheric circulation changes on summertime Arctic sea ice *Nat. Clim. Change* **7** 289–95
- Duarte P, Assmy P, Hop H, Spreen G, Gerland S and Hudson S R 2015 Modeling potentials and limitations for estimating production of ice algae in the Arctic Ocean *J. Mar. Syst.* **145** 69–90
- Duarte P et al 2017 Sea-ice thermohaline-dynamics and biogeochemistry in the Arctic Ocean: empirical and model results *J. Geophys. Res.-Biogeosci.* **122** 1632–54
- Eisenman I, Meier W N and Norris J R 2014 A spurious jump in the satellite record: has Antarctic sea ice expansion been overestimated? *Cryosphere* **8** 1289–96
- Forsström S, Gerland S and Pedersen C A 2011 Thickness and density of snow-covered sea ice and hydrostatic equilibrium assumption from in situ measurements in Fram Strait, the Barents Sea and the Svalbard coast *Ann. Glaciol.* **57** 261–70
- Gerland S and Renner A H H 2007 Sea ice mass balance in an Arctic fjord *Ann. Glaciol.* **46** 435–42
- Giles K A, Laxon S W and Ridout A L 2008 Circumpolar thinning of Arctic sea ice following the 2007 record ice extent minimum *Geophys. Res. Lett.* **35** L22502
- Granskog M A, Assmy P, Gerland S, Spreen G, Steen H and Smedsrud L H 2016 Arctic research on thin ice: consequences of Arctic sea ice loss *Eos Trans. AGU* **97** 22–6
- Granskog M A, Fer I, Rinke A and Steen H H 2018 Atmosphere-iceocean-ecosystem processes in a thinner Arctic sea ice regime: the Norwegian young sea ICE (N-ICE2015) expedition *J. Geophys. Res. Oceans* **123** 1586–1594
- Haas C, Beckers J, King J, Silis A, Stroeve J, Wilkinson J, Notenboom B, Schweiger A and Hendricks S 2017 Ice and snow thickness variability and change in the high Arctic Ocean observed by in situ measurements *Geophys. Res. Lett.* **44** 10462–69
- Hansen E, Gerland S, Granskog M A, Pavlova O, Renner A H H, Haapala J, Løyning T B and Tschudi M 2013 Thinning of Arctic sea ice observed in Fram Strait: 1990–2011 *J. Geophys. Res.—Oceans* **118** 5202–5221
- Hansen E, Ekeberg O-C, Gerland S, Pavlova O, Spreen G and Tschudi M 2014 Variability in categories of Arctic sea ice in Fram Strait *J. Geophys. Res. Oceans* **119** 7175–89
- Heygster G, Huntemann M, Ivanova N, Saldo R and Pedersen L T 2014 Response of passive microwave sea ice concentration algorithms to thin ice 2014 *IEEE Int. Proc. Geoscience and Remote Sensing Symp. (IGARSS) (Quebec City, 13–18 July)* pp 3618–21
- Hoppmann M, Nicolaus M, Hunkeler P A, Heil P, Behrens L-K, König-Langlo G and Gerdes R 2015 Seasonal evolution of an ice-shelf influenced fast-ice regime, derived from an autonomous thermistor chain *J. Geophys. Res. Oceans* **120** 1703–24
- Huntemann M, Heygster G, Kaleschke L, Krumpfen T, Mäkynen M and Drusch M 2014 Empirical sea ice thickness retrieval during the freeze up period from SMOS high incident angle observations *Cryosphere* **8** 439–51
- Itkin P, Spreen G, Hvidegaard S M, Skourup H, Wilkinson J, Gerland S and Granskog M A 2018 Contribution of

- deformation to sea ice mass balance: a case study from an N-ICE2015 storm *Geophys. Res. Lett.* **45** 789–796
- Jackson K, Wilkinson J, Maksym T, Meldrum D, Beckers J, Haas C and Mackenzie D 2013 A novel and low-cost sea ice mass balance buoy *J. Atmos. Oceanic Technol.* **30** 2676–88
- Kaleschke L, Tian-Kunze X, Maaß N, Mäkynen M and Drusch M 2012 Sea ice thickness retrieval from SMOS brightness temperatures during the Arctic freeze-up period *Geophys. Res. Lett.* **39** L05501
- Kaleschke L et al 2016 SMOS sea ice product: operational application and validation in the Barents Sea marginal ice zone *Remote Sens. Environ.* **180** 264–73
- Kern S 2004 A new method for medium-resolution sea ice analysis using weather-influence corrected special sensor microwave/imager 85 GHz data *Int. J. Remote Sens.* **25** 4555–82
- Kern S, Khvorostovsky K, Skourup H, Rinne E, Parsakhoo Z S, Djepa V, Wadhams P and Sandven S 2015 The impact of snow depth, snow density and ice density on sea ice thickness retrieval from satellite radar altimetry: results from the ESA-CCI sea ice ECV project round robin exercise *Cryosphere* **9** 37–52
- Kern S, Rösel A, Pedersen L T, Ivanova N, Saldo R and Tonboe R T 2016 The impact of melt ponds on summertime microwave brightness temperatures and sea-ice concentrations *Cryosphere* **10** 2217–39
- King J, Skourup H, Hvidegaard S M, Rösel A, Gerland S, Spreen G and Liston G E 2018 Comparison of freeboard retrieval and ice thickness calculation from ALS, ASIRAS, and CryoSat-2 in the Norwegian Arctic to field measurements made during the N-ICE2015 expedition *J. Geophys. Res. Oceans* **123** 1123–1141
- Komarov A and Barber D G 2014 Sea Ice motion tracking from sequential dual-polarized Radarsat-2 images *IEEE Trans. Geosci. Remote Sens.* **52** 121–36
- Kongoli C, Boukabara S A, Yan B, Weng F and Ferraro R 2011 A new sea ice concentration algorithm based on microwave surface emissivities—application to AMSU measurements *IEEE Trans. Geosci. Remote Sens.* **49** 175–89
- Korosov A A, Rampal P, Toudal Pedersen L, Saldo R, Ye Y, Heygster G, Lavergne T, Aaboe S and Girard-Ardhuin F 2018 A new tracking algorithm for sea ice age distribution estimation *Cryosphere* **12** 2073–85
- Kruppen T, Gerdes R, Haas C, Hendricks S, Herber A, Selyuzhenok V, Smedsrud L H and Spreen G 2016 Recent summer sea ice thickness surveys in Fram Strait and associated ice volume fluxes *Cryosphere* **10** 523–34
- Krupnik I, Aporta C, Gearheard S, Laidler G J and Kielsen Holm L (ed) 2010 *SIKU: Knowing Our Ice*. (Netherlands: Springer) p 501
- Kurtz N T, Farrell S L, Studinger M, Galin N, Harbeck J P, Lindsay R, Onana V D, Panzer B and Sonntag J G 2013 Sea ice thickness, freeboard, and snow depth products from Operation IceBridge airborne data *Cryosphere* **7** 1035–56
- Kurtz N T, Galin N and Studinger M 2014 An improved CryoSat-2 sea ice freeboard retrieval algorithm through the use of waveform fitting *Cryosphere* **8** 1217–37
- Kwok R 2009 Outflow of Arctic Ocean sea ice into the Greenland and Barents Seas: 1979–2007 *J. Clim.* **22** 2438–57
- Kwok R 2014 Simulated effects of a snow layer on retrieval of CryoSat-2 sea ice freeboard *Geophys. Res. Lett.* **41** 5014–20
- Kwok R and Cunningham G F 2015 Variability of Arctic sea ice thickness and volume from CryoSat-2 *Phil. Trans. R. Soc. A* **373** 20140157
- Kwok R, Cunningham G F and Pang S S 2004 Fram Strait sea ice outflow *J. Geophys. Res.* **109** C01009
- Lange B A et al 2017 Pan-Arctic sea ice-algal chl a biomass and suitable habitat are largely underestimated for multi-year ice *Glob. Change Biol.* **23** 4581–97
- Lavergne T, Eastwood S, Teffah Z, Schyberg H and Breivik L -A 2010 Sea ice motion from low-resolution satellite sensors: An alternative method and its validation in the Arctic *J. Geophys. Res.* **115** C10032
- Lavergne T et al 2019 Version 2 of the EUMETSAT OSI SAF and ESA CCI sea ice concentration climate data records *Cryosphere* **13** 49–78
- Lawrence I R, Tsamados M C, Stroeve J C, Armitage T W K and Ridout A L 2018 Estimating snow depth over Arctic sea ice from calibrated dual-frequency radar freeboards *Cryosphere* **12** 3551–64
- Laxon S, Peacock H and Smith D 2003 High interannual variability of sea ice thickness in the Arctic region *Nature* **425** 947–950
- Laxon S W et al 2013 CryoSat-2 estimates of Arctic sea ice thickness and volume *Geophys. Res. Lett.* **40** 732–7
- Leu E, Mundy C J, Assmy P, Campbell K, Gabrielsen T M, Gosselin M, Juul-Pedersen T and Gradinger R 2015 Arctic spring awakening—steering principles behind the phenology of vernal ice algae blooms *Prog. Oceanogr.* **139** 151–70
- Li L, Chen H and Guan L 2016 Study on the retrieval of snow depth from FY3B/MWRI in the Arctic *Int. Arch. Photogramm., Remote Sens. Spat. Inf. Sci. ISPRS Arch.* **V41** 513–20
- Lindsay R and Schweiger A 2015 Arctic sea ice thickness loss determined using subsurface, aircraft, and satellite observations *Cryosphere* **9** 269–83
- Lukovich J V, Hutchings J K and Barber D G 2015 On sea-ice dynamical regimes in the Arctic Ocean *Ann. Glaciol.* **56** 69
- Maaß N, Kaleschke L, Tian-Kunze X and Drusch M 2013 Snow thickness retrieval over thick Arctic sea ice using SMOS satellite data *Cryosphere* **7** 1971–89
- Maaß N, Kaleschke L, Tian-Kunze X and Tonboe R 2015 Snow thickness retrieval from L-band brightness temperatures: a model comparison *Ann. Glaciol.* **56** 9–17
- Maslanik J, Stroeve J, Fowler C and Emery W 2011 Distribution and trends in Arctic sea ice age through spring 2011 *Geophys. Res. Lett.* **38** L13502
- Mäkynen M and Karvonen J 2017 MODIS sea ice thickness and open water—sea ice charts over the Barents and Kara Seas for development and validation of sea ice products from microwave sensor data *Remote Sens.* **9** 1324
- Meier W N, Peng G, Scott D J and Savoie M H 2014 Verification of a new NOAA/NSIDC passive microwave sea-ice concentration climate record *Polar Res.* **33** 21004
- Meier W N, Fetterer F, Stewart J S and Helfrich S 2015 How do sea ice concentrations from operational data compare with passive microwave estimates? Implications for improved model evaluations and forecasting *Ann. Glaciol.* **56** 332–40
- Meier W N and Stewart J S 2019 Assessing uncertainties in sea ice extent climate indicators *Environ. Res. Lett.* **14** 035005
- Melnikov I A, Zhitina L S and Semenova T N 2016 Recent condition of the sea ice biodiversity within the North Pole region *Probl. Arktiki I Antarkt.* **4** 104–10 (in Russian) (http://aari.ru/misc/publicat/paa_arj_jour_en.php?idnum=477)
- Michel C, Hamilton J, Hansen E, Barber D, Reigstad M, Iacozza J, Seuthe L and Niemi A 2015 Arctic Ocean outflow shelves in the changing Arctic: a review and perspectives *Prog. Oceanogr.* **139** 66–88
- Nandan V, Geldsetzer T, Yackel J, Mahmud M, Scharien R, Howell S, King J, Ricker R and Else B 2017 Effect of snow salinity on CryoSat-2 Arctic first-year sea ice freeboard measurements *Geophys. Res. Lett.* **44** 10419–26
- Niemi A and Michel C 2015 Temporal and spatial variability in sea-ice carbon:nitrogen ratios on Canadian Arctic shelves *Elem.: Sci. Anth.* **3** 000078
- Niemi A, Michel C, Hille K and Poulin M 2011 Protist assemblages in winter sea ice: setting the stage for the spring ice algal bloom *Polar Biol.* **34** 1803–17
- Notz D 2009 The future of ice sheets and sea ice: between reversible retreat and unstoppable loss *Proc. Natl Acad. Sci.* **106** 20590–5
- Obbard R W, Sadri S, Wong Y Q, Khitun A A, Baker I and Thompson R C 2014 Global warming releases microplastic legacy frozen in Arctic Sea ice *Earth's Future* **2** 315–20
- Olsen L M et al 2017 The seeding of ice algal blooms in Arctic pack ice: The multiyear ice seed repository hypothesis *J. Geophys. Res. Biogeosciences* **122** 1529–1548

- Paul S, Hendricks S, Ricker R, Kern S and Rinne E 2018 Empirical parametrization of Envisat freeboard retrieval of Arctic and Antarctic sea ice based on CryoSat-2: progress in the ESA climate change initiative *Cryosphere* **12** 2437–60
- Peeken I, Primpke S, Beyer B, Gütermann J, Katlein C, Krumpfen T, Bergmann M, Hehemann L and Gerdt G 2018 Arctic sea ice is an important temporal sink and means of transport for microplastic *Nat. Commun.* **9** 1505
- Peng G, Meier W N, Scott D J and Savoie M H 2013 A long-term and reproducible passive microwave sea ice concentration data record for climate studies and monitoring *Earth Syst. Sci. Data* **5** 311–8
- Perovich D K and Richter-Menge J A 2015 Regional variability in sea ice melt in a changing Arctic *Proc. R. Soc.* **373** 20140165
- Perovich D et al 2018 Sea ice cover. Section 5d in state of the climate in 2017 *Bull. Am. Meteorol. Soc.* **99** S147–52
- Polashenski C, Perovich D, Richter-Menge J and Elder B 2011 Seasonal ice mass-balance buoys: adapting tools to the changing Arctic *Ann. Glaciol.* **52** 18–26
- Renner A H H, Gerland S, Haas C, Spreen G, Beckers J F, Hansen E, Nicolaus M and Goodwin H 2014 Evidence of Arctic sea ice thinning from direct observations *Geophys. Res. Lett.* **41** 5029–36
- Ricker R, Girard-Ardhuin F, Krumpfen T and Lique C 2018 Satellite derived sea ice export and its impact on Arctic ice mass balance *Cryosphere* **12** 3017–32
- Ricker R, Hendricks S, Helm V, Skourup H and Davidson M 2014 Sensitivity of CryoSat-2 Arctic sea-ice freeboard and thickness on radar-waveform interpretation *Cryosphere* **8** 1607–22
- Ricker R, Hendricks S, Kaleschke L, Tian-Kunze X, King J and Haas C 2017 A weekly Arctic sea-ice thickness data record from merged CryoSat-2 and SMOS satellite data *Cryosphere* **11** 1607–23
- Ricker R, Hendricks S, Perovich D K, Helm V and Gerdes R 2015 Impact of snow accumulation on CryoSat-2 range retrievals over Arctic sea ice: an observational approach with buoy data *Geophys. Res. Lett.* **42** 4447–55
- Rigor I G, Colony R L and Martin S 2000 Variations in surface air temperature observations in the Arctic, 1979–97 *J. Clim.* **13** 896–914
- Rigor I G, Clemente-Colón P, Reinking C, Keith D and Reams M 2016 *The International Arctic Buoy Programme (IABP)—A Model for Sustaining Arctic Observing Networks White paper Arctic Observing Summit 2016 (Fairbanks Alaska, USA)* p 9
- Rösel A, Itkin P, King J, Divine D, Wang C, Granskog M A, Krumpfen T and Gerland S 2018 Winter and spring development of sea-ice and snow thickness distributions north of Svalbard observed during N-ICE2015 *J. Geophys. Res.—Oceans* **123** 1156–1176
- Rösel A, Kaleschke L and Birnbaum G 2012 Melt ponds on Arctic sea ice determined from MODIS satellite data using an artificial neural network *Cryosphere* **6** 431–46
- Schwegmann S, Rinne E, Ricker R, Hendricks S and Helm V 2016 About the consistency between Envisat and CryoSat-2 radar freeboard retrieval over Antarctic sea ice *Cryosphere* **10** 1415–25
- Schweiger A, Lindsay R, Zhang J, Steele M, Stern H and Kwok R 2011 Uncertainty in modeled Arctic sea ice volume *J. Geophys. Res.* **116** C00D06
- Smedsrud L H, Halvorsen M H, Stroeve J C, Zhang R and Kloster K 2017 Fram Strait sea ice export variability and September Arctic sea ice extent over the last 80 years *Cryosphere* **11** 65–79
- Smedsrud L H, Sirevaag A, Kloster K, Sorteberg A and Sandven S 2011 Recent wind driven high sea ice area export in the Fram Strait contributes to Arctic sea ice decline *Cryosphere* **5** 821–9
- Spreen G, Kaleschke L and Heygster G 2008 Sea ice remote sensing using AMSR-E 89-GHz channels *J. Geophys. Res.* **113** C02S03
- Spreen G, Kern S, Stammer D and Hansen E 2009 Fram Strait sea ice volume export estimated between 2003 and 2008 from satellite data *Geophys. Res. Lett.* **36** L19502
- Stroeve J and Notz D 2018 Changing state of Arctic sea ice across all seasons *Environ. Res. Lett.* **13** (10) 103001
- Tian-Kunze X, Kaleschke L, Maaß N, Mäkynen M, Serra N, Drusch M and Krumpfen T 2014 SMOS-derived thin sea ice thickness: algorithm baseline, product specifications and initial verification *Cryosphere* **8** 997–1018
- Tikhonov V V, Repina I A, Raev M D, Sharkov E A, Ivanov V V, Boyarskii D A, Alexeeva T A and Komarova N Y 2015 A physical algorithm to measure sea ice concentration from passive microwave remote sensing data *Adv. Space Res.* **56** 1578–89
- Tilling R L, Ridout A, Shepherd A and Wingham D J 2015 Increased Arctic sea ice volume after anomalously low melting in 2013 *Nat. Geosci.* **8** 643–6
- Tilling R L, Ridout A and Shepherd A 2018 Estimating Arctic sea ice thickness and volume using CryoSat-2 radar altimeter data *Adv. Space Res.* **62** 1203–25
- Tomboe R T, Eastwood S, Lavergne T, Sørensen A M, Rathmann N, Dybkjær G, Pedersen L T, Høyer J L and Kern S 2016 The EUMETSAT sea ice concentration climate data record *Cryosphere* **10** 2275–90
- Tremblay J-É, Anderson L G, Matrai P, Coupel P, Bélanger S, Michel C and Reigstad M 2015 Global and local drivers of nutrient supply, primary production and CO₂ drawdown in the changing Arctic ocean *Prog. Oceanogr.* (<https://doi.org/10.1016/j.pocean.2015.08.009>)
- Tschudi M A, Stroeve J C and Stewart J S 2016a Relating the age of Arctic sea ice to its thickness, as measured during NASA's ICESat and IceBridge campaigns *Remote Sens.* **8** 457
- Tschudi M, Fowler C, Maslanik J, Stewart J S and Meier W 2016b *Polar Pathfinder Daily 25 km EASE-Grid Sea Ice Motion Vectors, Version 3*. (Boulder, Colorado USA: NASA National Snow and Ice Data Center Distributed Active Archive Center) (<https://doi.org/10.5067/O57VAIT2AYYY>)
- Vaughan D G et al 2013 Observations: cryosphere *climate change 2013 The Physical Science Basis. Contribution of Working Group I to the 5th Assessment Report of the Intergovernmental Panel on Climate Change* ed et al (Cambridge: Cambridge University Press) pp 319–82
- Wang C, Granskog M, Gerland S, Hudson S R, Perovich D K, Nicolaus M, Karlsen T-I, Fossan K and Bratrein M 2014 Autonomous observations of solar energy partitioning in first-year sea ice in the Arctic Basin *J. Geophys. Res. Oceans* **119** 2066–80
- Wang C, Granskog M, Hudson S R, Gerland S, Pavlov A K, Perovich D K and Nicolaus M 2016 Atmospheric conditions in the central Arctic Ocean through the melt seasons of 2012 and 2013: impact on surface conditions and solar energy deposition into the ice-ocean system *J. Geophys. Res. Atmos.* **21** 1043–58
- Wang X Y, Guan L and Li L L 2018 Comparison and validation of sea ice concentration from FY-3B/MWRI and Aqua/AMSR-E observations *J. Remote Sens.* **22** 723–36 (in Chinese)
- Webster M, Gerland S, Holland M, Hunke E, Kwok R, Lecomte O, Massom R, Perovich D and Sturm M 2018 Snow in the changing sea-ice systems *Nat. Clim. Change* **8** 946–53
- Wu S and Liu J 2018 Comparison of Arctic sea ice concentration datasets *Haiyang Xuebao* **40** 64–72 (in Chinese)
- Zhang J, Lindsay R, Schweiger A and Rigor I 2012 Recent changes in the dynamic properties of declining Arctic sea ice: a model study *Geophys. Res. Lett.* **39** L20503
- Zhang J and Rothrock D 2003 Modeling global sea ice with a thickness and enthalpy distribution model in generalized curvilinear coordinates *Mon. Weather Rev.* **31** 845–61
- Zhang Z, Bi H, Sun K, Huang H, Liu Y and Yan L 2017 Arctic sea ice volume export through the Fram Strait from combined satellite and model data: 1979–2012 *Acta Oceanol. Sin.* **36** 44–55
- Zygmuntowska M, Khvorostovsky K, Helm V and Sandven S 2013 Waveform classification of airborne synthetic aperture radar altimeter over Arctic sea ice *Cryosphere* **7** 1315–24
- Zygmuntowska M, Rampal P, Ivanova N and Smedsrud L H 2014 Uncertainties in Arctic sea ice thickness and volume: new estimates and implications for trends *Cryosphere* **8** 705–20