





Draft Genome Sequence of the Fish Pathogen *Flavobacterium columnare* Strain MS-FC-4

 Ryan P. Bartelme,^a Paul Barbier,^b Ryan S. Lipscomb,^c Scott E. LaPatra,^d Ryan J. Newton,^a Jason P. Evenhuis,^c
 Mark J. McBride^b

^aSchool of Freshwater Sciences, University of Wisconsin—Milwaukee, Milwaukee, Wisconsin, USA

^bDepartment of Biological Sciences, University of Wisconsin—Milwaukee, Milwaukee, Wisconsin, USA

^cU.S. Department of Agriculture, National Center for Cool and Cold Water Aquaculture, Agriculture Research Service, Kearneysville, West Virginia, USA

^dClear Springs Foods, Inc., Buhl, Idaho, USA

ABSTRACT *Flavobacterium columnare* MS-FC-4 is a highly virulent genetic group 1 (formerly genomovar I) strain isolated from rainbow trout (*Oncorhynchus mykiss*). The draft genome consists of three contigs totaling 3,449,277 bp with 2,811 predicted open reading frames. *F. columnare* MS-FC-4 is a model strain for functional genomic analyses.

Flavobacterium columnare causes columnaris disease in wild and cultured fish (1, 2). It is a common pathogen of farmed freshwater fish and causes major economic losses worldwide. *F. columnare* isolates are genetically diverse and were classified historically into multiple genomovars (3, 4). Recently, the genomovars were described as four distinct genetic groups (5). Almost all isolates recovered from salmonids belong to genetic group 1 (formerly genomovar 1). *F. columnare* MS-FC-4 is a highly virulent strain that was isolated from rainbow trout (*Oncorhynchus mykiss*) (6). It is amenable to genetic manipulation, making it attractive for functional genomic studies of virulence.

F. columnare MS-FC-4 was cultured for 24 h in tryptone yeast extract salt medium at 30°C and 150 rpm. Genomic DNA (gDNA) was isolated using a cetyltrimethylammonium bromide (CTAB)/phenol-chloroform/isoamyl alcohol protocol (7) optimized for *F. columnare*.

For long-read sequencing, a gDNA library was prepared according to the standard Pacific Biosciences RS II large-insert library protocol. Two flow cells (C4 chemistry) were each loaded with 0.06 nM of a BluePippin (Sage Science, Beverly, MA, USA) size-selected (>10-kb) SMRTbell gDNA library. One microgram of MS-FC-4 gDNA was also sequenced on an Illumina HiSeq 2000 platform with 100-bp paired-end reads. PacBio reads were assembled using Canu version 1.6 with the following options: genome size 3.3 Mb, corrected error rate 0.035, and corMaxEvidenceErate 0.15 (8). Subread coverage of the genome was ~865×. The Illumina HiSeq paired-end reads were mapped to the PacBio assembly using Bowtie 2 version 2.3.4 (9) with the no-unaligned parameter to discard unaligned reads. HiSeq coverage of the contigs was ~360×. The output SAM file was converted to a BAM file using SAMtools version 1.4 (10). Using the Canu assembly and the BAM file as input, Pilon version 1.22 (11) corrected two insertions, seven deletions, and one substitution. The Pilon corrections did not close the gaps in the Canu PacBio assembly, resulting in three contigs of 2,709,164, 716,735, and 23,378 bp. The corrected MS-FC-4 genome assembly was annotated using the NCBI Prokaryotic Genome Annotation Pipeline (12).

The draft *F. columnare* MS-FC-4 genome has a cumulative total size of 3,449,277 bp with a G+C content of 31.9%. The annotated genome had 2,811 coding genes, 19 complete rRNA operons, 95 tRNAs, and 2 clustered regularly interspaced short palin-

Received 16 April 2018 Accepted 17 April 2018 Published 17 May 2018

Citation Bartelme RP, Barbier P, Lipscomb RS, LaPatra SE, Newton RJ, Evenhuis JP, McBride MJ. 2018. Draft genome sequence of the fish pathogen *Flavobacterium columnare* strain MS-FC-4. Genome Announc 6:e00429-18. <https://doi.org/10.1128/genomeA.00429-18>.

This is a work of the U.S. Government and is not subject to copyright protection in the United States. Foreign copyrights may apply.

Address correspondence to Jason P. Evenhuis, jason.evenhuis@ars.usda.gov, or Mark J. McBride, mcbride@uwm.edu.

R.P.B. and P.B. contributed equally to this work.

dromic repeat (CRISPR) arrays. Two incomplete prophage regions of 12.4 and 25.9 kb were identified using PHASTER (13). The two-way average nucleotide identities (ANIs) (14) between the MS-FC-4 genome sequence and those of strains ATCC 49512 (15), Pf1 (16), and CSF 298-10 (17) were >99%, confirming the *F. columnare* genetic group 1 (genomovar I) classification (5, 6). The MS-FC-4 genome contains the core genes of the *Bacteroidetes*-specific type IX secretion system linked to gliding motility and virulence (18, 19). The availability of the MS-FC-4 genome sequence should facilitate construction of targeted gene deletions to identify critical *F. columnare* virulence factors.

Accession number(s). This whole-genome shotgun project has been deposited at DDBJ/ENA/GenBank under the accession number [PVLU00000000](https://www.ncbi.nlm.nih.gov/nuclseq/PVLU00000000). The version described in this paper is the first version, PVLU01000000.

ACKNOWLEDGMENTS

This project used the University of Wisconsin—Milwaukee Great Lakes Genomics Center DNA sequencing and bioinformatics services. Illumina HiSeq sequencing was done by Arizona Research Labs (Tucson, AZ, USA).

This research was supported by funds from the USDA-ARS (project numbers 5090-31320-004-03S and 8082-32000-006) and by funds from the University of Wisconsin Sea Grant Institute under grants from the National Sea Grant College Program, National Oceanic and Atmospheric Administration, U.S. Department of Commerce, and the State of Wisconsin, federal grant NA14OAR4170092, project R/SFA-11. R.P.B. was supported by a fellowship from the University of Wisconsin—Milwaukee. The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication. The USDA is an Equal Opportunity Employer.

REFERENCES

1. Declercq AM, Haesebrouck F, Van den Broeck W, Bossier P, Decostere A. 2013. Columnaris disease in fish: a review with emphasis on bacterium-host interactions. *Vet Res* 44:27. <https://doi.org/10.1186/1297-9716-44-27>.
2. Wahli T, Madsen L. 2018. Flavobacteria, a never ending threat for fish: a review. *Curr Clin Microbiol Rep* 5:26–37. <https://doi.org/10.1007/s40588-018-0086-x>.
3. Triyanto, Wakabayashi H. 1999. Genotypic diversity of strains of *Flavobacterium columnare* from diseased fishes. *Fish Pathol* 34:65–71. <https://doi.org/10.3147/jsfp.34.65>.
4. LaFrentz BR, García JC, Dong HT, Waldbieser GC, Rodkhum C, Wong FS, Chang SF. 2017. Optimized reverse primer for 16S-RFLP analysis and genomovar assignment of *Flavobacterium columnare*. *J Fish Dis* 40: 1103–1108. <https://doi.org/10.1111/jfd.12583>.
5. LaFrentz BR, García JC, Waldbieser GC, Evenhuis JP, Loch TP, Liles MR, Wong FS, Chang SF. 2018. Identification of four distinct phylogenetic groups in *Flavobacterium columnare* with fish host associations. *Front Microbiol* 9:452. <https://doi.org/10.3389/fmicb.2018.00452>.
6. Evenhuis JP, Mohammed H, Lapatra SE, Welch TJ, Arias CR. 2016. Virulence and molecular variation of *Flavobacterium columnare* affecting rainbow trout in Idaho, USA. *Aquaculture* 464:106–110. <https://doi.org/10.1016/j.aquaculture.2016.06.017>.
7. Wilson K. 2001. Preparation of genomic DNA from bacteria, p 2.4.1–2.4.5. *In* Current protocols in molecular biology. John Wiley & Sons, Inc., Hoboken, NJ.
8. Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM. 2017. Canu: scalable and accurate long-read assembly via adaptive *k*-mer weighting and repeat separation. *Genome Res* 27:722–736. <https://doi.org/10.1101/gr.215087.116>.
9. Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nat Methods* 9:357–359. <https://doi.org/10.1038/nmeth.1923>.
10. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R. 2009. The sequence alignment/map format and SAMtools. *Bioinformatics* 25:2078–2079. <https://doi.org/10.1093/bioinformatics/btp352>.
11. Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, Cuomo CA, Zeng Q, Wortman J, Young SK, Earl AM. 2014. Pilon: an integrated tool for comprehensive microbial variant detection and genome assembly improvement. *PLoS One* 9:e112963. <https://doi.org/10.1371/journal.pone.0112963>.
12. Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI prokaryotic genome annotation pipeline. *Nucleic Acids Res* 44:6614–6624. <https://doi.org/10.1093/nar/gkw569>.
13. Arndt D, Grant JR, Marcu A, Sajed T, Pon A, Liang Y, Wishart DS. 2016. PHASTER: a better, faster version of the PHAST phage search tool. *Nucleic Acids Res* 44:W16–W21. <https://doi.org/10.1093/nar/gkw387>.
14. Goris J, Konstantinidis KT, Klappenbach JA, Coenye T, Vandamme P, Tiedje JM. 2007. DNA–DNA hybridization values and their relationship to whole-genome sequence similarities. *Int J Syst Evol Microbiol* 57:81–91. <https://doi.org/10.1099/ijs.0.64483-0>.
15. Tekedar HC, Karsi A, Gillaspay AF, Dyer DW, Benton NR, Zaitshik J, Vamenta S, Banes MM, Gulsoy N, Aboko-Cole M, Waldbieser GC, Lawrence ML. 2012. Genome sequence of the fish pathogen *Flavobacterium columnare* ATCC 49512. *J Bacteriol* 194:2763–2764. <https://doi.org/10.1128/JB.00281-12>.
16. Zhang Y, Nie P, Lin L. 2016. Complete genome sequence of the fish pathogen *Flavobacterium columnare* Pf1. *Genome Announc* 4(5):e00900-16. <https://doi.org/10.1128/genomeA.00900-16>.
17. Evenhuis JP, LaPatra SE, Graf J. 2017. Draft genome sequence of the fish pathogen *Flavobacterium columnare* strain CSF-298-10. *Genome Announc* 5(15):e00173-17. <https://doi.org/10.1128/genomeA.00173-17>.
18. Sato K, Naito M, Yukitake H, Hirakawa H, Shoji M, McBride MJ, Rhodes RG, Nakayama K. 2010. A protein secretion system linked to bacteroidete gliding motility and pathogenesis. *Proc Natl Acad Sci U S A* 107:276–281. <https://doi.org/10.1073/pnas.0912010107>.
19. Li N, Zhu Y, LaFrentz BR, Evenhuis JP, Hunnicutt DW, Conrad RA, Barbier P, Gullstrand CW, Roets JE, Powers JL, Kulkarni SS, Erbes DH, Garcia JC, Nie P, McBride MJ. 2017. The type IX secretion system is required for virulence of the fish pathogen *Flavobacterium columnare*. *Appl Environ Microbiol* 83:e01769-17. <https://doi.org/10.1128/AEM.01769-17>.