



Mitochondrial DNA Part B Resources

ISSN: (Print) 2380-2359 (Online) Journal homepage: <https://www.tandfonline.com/loi/tmdn20>

Complete mitochondrial genome of *Carijoa riisei* (Duchassaing & Michelotti, 1860) (Octocorallia: Alcyonacea: Stolonifera: Clavulariidae)

Erin E. Easton & David Hicks

To cite this article: Erin E. Easton & David Hicks (2020) Complete mitochondrial genome of *Carijoa riisei* (Duchassaing & Michelotti, 1860) (Octocorallia: Alcyonacea: Stolonifera: Clavulariidae), Mitochondrial DNA Part B, 5:2, 1826-1827, DOI: [10.1080/23802359.2020.1750998](https://doi.org/10.1080/23802359.2020.1750998)

To link to this article: <https://doi.org/10.1080/23802359.2020.1750998>



© 2020 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



[View supplementary material](#)



Published online: 13 Apr 2020.



[Submit your article to this journal](#)



Article views: 169



[View related articles](#)



[View Crossmark data](#)

MITOGENOME ANNOUNCEMENT



Complete mitochondrial genome of *Carijoa riisei* (Duchassaing & Michelotti, 1860) (Octocorallia: Alcyonacea: Stolonifera: Clavulariidae)

Erin E. Easton  and David Hicks

School of Earth, Environmental and Marine Sciences, University of Texas Rio Grande Valley, Brownsville, TX, USA

ABSTRACT

We report the first complete Stolonifera mitochondrial genome. *Carijoa riisei* (Duchassaing & Michelotti, 1860) isolate CLP2_A03 was collected by scuba at 32 m on the USTS *Texas Clipper* (27° 53.7827'N, 93° 36.2702'W). The complete mitogenome has the ancestral octocoral gene order for its 14 protein-coding genes, two rRNA genes, and one tRNA gene. It is 18,714 bp (30.7% A, 15.8% C, 18.8% G, and 34.7% T). Of the Alcyonacea mitogenomes published to date, it is most genetically similar (94% uncorrected) to *Sinularia ceramensis* Verseveldt, 1977 (NC_044122).

ARTICLE HISTORY

Received 20 March 2020
Accepted 27 March 2020

KEYWORDS

Artificial reef; Cnidaria; Gulf of Mexico; mesophotic

Carijoa Müller, 1867 consists of three accepted species of which two, *C. riisei* and *C. operculata*, are reported in the Gulf of Mexico. These species are distinguishable by anthocodial spiculation, which forms an operculum in the latter species (Bayer 1961). Although *C. riisei* was described from the Atlantic-Caribbean, it is widely distributed in the Atlantic, Pacific, and Indian Oceans and has a likely origin in the Indo-Pacific, whereas, the Atlantic and Caribbean populations are inferred to be the youngest (Concepcion et al. 2010). Colonies can reach sexual maturity in a few months, can maintain growth rates ~ 0.5 cm week⁻¹ for several months, undergo vegetative propagation, and are gonochoric (rarely hermaphroditic) with continuous and asynchronous gamete release (Kahng et al. 2008; Barbosa et al. 2014). These characteristics enable *C. riisei* to proliferate rapidly and to form dense aggregations and may contribute to its colonization of artificial structures (Kahng et al. 2008).

The specimen was collected by scuba at 32 m on the USTS *Texas Clipper* (27° 53.7827'N, 93° 36.2702'W) on 22 August 2017. This vessel was sunk as part of the Texas Parks and Wildlife Department's Artificial Reef Program on 17 November 2007 (Curley 2011). DNA was extracted with GeneJET Genomic DNA Purification Kit (ThermoFisher Scientific, Waltham, MA) per manufacture's protocol and submitted to Biopolymers Facility at Harvard Medical School for library preparation (Illumina Nextera XT2) and next-generation sequencing (NextSeq 500). Trimmed reads (Trimmomatic-0.32, Bolger et al. 2014) were assembled de novo with SPAdes (Bankevich et al. 2012) on the University of New Hampshire server. Trimmed reads (BBduk v. 37.25) were mapped to the SPAdes contig to generate a consensus sequence in Geneious Prime 2020.0.5 (<https://www.geneious.com>).

Genes were annotated by manually adjusting *Muricea crassa* Verrill, 1869 (NC029697) annotations mapped to the consensus sequence in Geneious. The *C. riisei* mitogenome was deposited in GenBank (MT161608) and the specimen and sclerite SEM plate (see Supplemental Material) were deposited in the Smithsonian National Museum of Natural History (USNM1616994). The complete mitogenome was aligned with default MUSCLE (Edgar 2004) parameters in Geneious to 26 representative species for which the complete mitochondrial genomes with the ancestral gene order were available in GenBank. A maximum-likelihood, phylogenetic tree, rooted with the Pennatulacea clade, was constructed with RaxML 8.2.11 (Stamatakis 2014) (Figure 1): 100 bootstrap replicates (rapid bootstrapping with search for best-scoring ML tree), no outgroup, and nucleotide model = GTR CAT I. Extended methods and alignments are available in Supplemental Material.

The complete mitogenome is 18,714 bp (30.7% A, 15.8% C, 18.8% G, and 34.7% T), has the ancestral octocoral gene order, and has 14 protein-coding genes, two rRNA genes, and one tRNA gene. This mitogenome report is the first for suborder Stolonifera. *C. riisei* is sister to the two species in Alcyoniidae and $\sim 94\%$ similar (uncorrected) to *Sinularia ceramensis* Verseveldt, 1977 (Alcyoniina: Alcyoniidae) (Figure 1). As found in previous studies (Figueroa and Baco 2015; Polisenio et al. 2017), phylogenetic reconstructions result in polyphyly of octocoral taxa, including orders, such as Alcyonacea (Figure 1).

Disclosure statement

The authors report no conflicts of interest. The authors alone are responsible for the content and writing of this article.

CONTACT Erin E. Easton  erin.easton@utrgv.edu  UTRGV-SEEMS, 33363 Marine Lab Dr., South Padre Island, TX 78597, USA

 Supplemental data for this article is available on line at <https://doi.org/10.1080/23802359.2020.1750998>.

© 2020 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.




Figure 1. Maximum-likelihood, phylogenetic tree of the complete mitochondrial genomes of *Carijoa riisei* (*this study, GenBank accession number, species name, field sample ID) and 26 representative octocorals (GenBank accession number, species name). In Geneious Prime 20.0.5, complete mitochondrial genomes were aligned with default MUSCLE parameters; the resulting alignment was used to construct the phylogenetic tree with RaxML 8.2.11 plugin with the following changes to the default settings: bootstrap replicates = 100, algorithm = rapid bootstrapping and search for best-scoring ML tree, nucleotide model = GTR CAT I. Bootstrap values >50 are report at the nodes. See Supplemental Material for methods details.

Funding

This publication was made possible by the National Oceanic and Atmospheric Administration, Office of Education Educational Partnership Program award NA16SEC4810009. Its contents are solely the responsibility of the award recipient and do not necessarily represent the official views of the U.S. Department of Commerce, National Oceanic and Atmospheric Administration. Research cruise funding provided by Texas Parks and Wildlife Department—Artificial Reef Program [Grant No. 475342, 2016–2018] and research expenses supported by award NA16SEC4810009 and New Hampshire-INBRE through an Institutional Development Award (IDeA), P20GM103506, from the National Institute of General Medical Sciences of the NIH. Sequencing costs supported by an Institutional Grant [NA14OAR4170102] to the Texas Sea Grant College Program from the National Sea Grant Office, National Oceanic and Atmospheric Administration, U.S. Department of Commerce. Funds for contracted labor (SEM imaging) were provided by Southeast Deep Coral Initiative to CSS, Inc, led by NOAA National Centers for Coastal Ocean Science with support from NOAA Deep Sea Coral Research and Technology Program.

ORCID

Erin E. Easton  <http://orcid.org/0000-0003-4853-6026>

References

- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, et al. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol.* 19(5):455–477.
- Barbosa TM, Gomes PB, Bergeron A-S, Santos AM, Chagas C, Freitas EM, Perez CD. 2014. Comparisons of sexual reproduction in *Carijoa riisei* (Cnidaria, Alcyonacea) in South Atlantic, Caribbean, and Pacific areas. *Hydrobiologia.* 734(1):201–212.
- Bayer FM. 1961. The shallow-water Octocorallia of the West Indian region. *Stud Fauna Curaçao Other Caribbean Islands.* 12(1):1–373.
- Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for illumina sequence data. *Bioinformatics.* 30(15):2114–2120.
- Concepcion G, Kahng S, Crepeau M, Franklin E, Coles S, Toonen R. 2010. Resolving natural ranges and marine invasions in a globally distributed octocoral (genus *Carijoa*). *Mar Ecol Prog Ser.* 401:113–127.
- Curley SJ. 2011. The ship that would not die: USS Queens, SS Excambion, and USTS Texas Clipper. Centennial Series of the Association of Former Students, Texas A&M University (Book 117). College Station: Texas A&M University Press. p. 256.
- Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res.* 32(5):1792–1797.
- Figueroa DF, Baco AR. 2015. Octocoral mitochondrial genomes provide insights into the phylogenetic history of gene order rearrangements, order reversals, and cnidarian phylogenetics. *Genome Biol E.* 7(1): 391–409.
- Kahng SE, Benayahu Y, Wagner D, Rothe N. 2008. Sexual reproduction in the invasive octocoral *Carijoa riisei* in Hawaii. *Bull Mar Sci.* 82(1): 1–17.
- Poliseno A, Feregrino C, Sartoretto S, Aurelle D, Wörheide G, McFadden CS, Vargas S. 2017. Comparative mitogenomics, phylogeny and evolutionary history of *Leptogorgia* (Gorgoniidae). *Mol Phylogenet Evol.* 115:181–189.
- Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics.* 30(9): 1312–1313.