

**NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION AND THE UNIVERSITY OF  
MARYLAND CENTER FOR ENVIRONMENTAL SCIENCE**

# **2018 Status Report Scoring Methodology for Pacific Jurisdictions**

**Indicator calculations and scoring for coral  
reef status reports of the Pacific  
jurisdictions: American Samoa, Hawaiian  
Archipelago, Pacific Remote Islands, Guam,  
and Northern Mariana Islands**

Caroline Donovan, Justine Kimball, Dione Swanson, Courtney Couch, Bernardo Vargas-Angel, Ivor Williams, Adel Heenan, Thomas Oliver, Rusty Brainard, Matt Gorstein, Arielle Levine, Peter Edwards, Jarrod Loerzel, Mark Eakin, Erick Geiger, Heath Kelsey, Alexandra Fries

## **Table of Contents**

- I. Background**
- II. Pacific jurisdictions with reef areas**
- III. Indicator Development**
- IV. Indicators and scoring process**
- V. References**
- VI. Indicator data citations**

## I. Background

### **The motivation behind status and trends reporting**

The National Oceanic and Atmospheric Administration (NOAA) Coral Reef Conservation Program (CRCP) invests significant funds to support a National Coral Reef Monitoring Program (NCRMP) throughout the U.S. Pacific, Atlantic, Gulf of Mexico, and Caribbean coral reef areas. A key component of this program is a periodic, national-level assessment on the status and trends of U.S. coral reef areas. To develop and implement this report framework, CRCP partnered with the Integration and Application Network (IAN) at the University of Maryland Center for Environmental Science (UMCES). The framework, termed herein a status report, is based on the timely and transparent assessment of biophysical and human dimension indicators against reference and assessment points, which are synthesized into overall condition scores for each jurisdiction. The primary purpose of the CRCP status report products is to communicate the status and trends of U.S. coral reefs to Congress, NOAA leadership, and the interested public. The primary purpose of this document is to describe the scoring process used for all the Pacific jurisdictions: American Samoa, Hawaiian Archipelago, Pacific Remote Islands, Guam, and Northern Mariana Islands.

Ecosystem condition assessments are a common approach to synthesizing a large amount of ecosystem monitoring data into a public-friendly report that can be understood by decision makers, managers, and scientists alike. Fundamentally, status reports help answer the question “How is the ecosystem doing?” The goals of a status report are to: provide a broad-level assessment, communicate complex information, use real data, and engage communities. These assessments are produced by a variety of groups from small, community-based organizations to regional management agencies, to large international partnerships. To advance this effort, CRCP and IAN-UMCES brought together science experts, natural resource managers, and other stakeholders from NOAA and local jurisdictions in order to develop biological, climatological, and human connection indicators and reference/assessment points for coral reefs.


In the case of the CRCP and IAN-UMCES partnership to develop a coral reef status report, there was an identified need to summarize and communicate coral reef monitoring in the U.S. jurisdictions and create a framework to support continued reporting. These assessments provide the status of U.S. coral reef areas in order to track change over time and are evaluating ecosystem condition, not management efforts or restoration success. The goals stated above are accomplished by producing a simple and concise product that tells the story of coral reefs using effective visual and narrative elements. All jurisdictions were assessed on coral & algae, fish, climate, and human connections indicators. To allow for regionally and context specific baselines, the scoring of indicators is unique to each jurisdiction. Reference and assessment points against which data is evaluated are developed based on literature review, regulatory guidelines, institutional goals, biological limits, reference conditions, historical benchmarks, and expert judgement. These reference and assessment points are determined by a group of experts—scientists and managers—from NOAA and local jurisdictional agencies. By developing jurisdiction-specific reference and assessment points, indicator scores are representative of that system. By using the same indicators across jurisdictions, indicator scores are comparable.

The data used in the scoring process is restricted to NCRMP data. Localized data, such as those collected by jurisdictional agencies, are not included in the scoring of indicators. This is because blending data that are collected under different sampling designs and/or different sampling methodologies is difficult. In the absence of a targeted calibration exercise that would allow for integrating disparate datasets, we focus here on the NCRMP data that is designed to monitor coral reefs at a jurisdictional scale. It is a goal of future efforts to include this other data in indicator scoring. However, local long-term data products are included when possible as a time-series or highlight story. Highlight stories are meant to message locally relevant information.

## Create a conceptual framework

The first step in developing a status report is to conceptualize the ecosystem and determine the threats to that system and the parts of the system that people value. A two-day stakeholder engagement workshop, which includes not only decision makers and managers of the systems, but also the scientists that gather and analyze data in the region, is critical to creating buy-in and support for the assessment process and final product. Workshops were held for each jurisdiction. These workshops brought together scientists, managers, and decision makers to determine the conceptual framework, potential indicators and thresholds, and draft layout of the report.

*This 5-Step process was used to create the status reports:*



Create a framework defining goals and major aspects of each goal that should be evaluated over time.



Select indicators that convey meaningful information and can be reliably measured.



Define status categories, reporting regions, and method of measuring threshold attainment.



| Region      | Year | MeanSIZE |
|-------------|------|----------|
| Ofu&Olosega | 2014 | 1.02     |
| Rose        | 2014 | 1.08     |
| Swains      | 2014 | 1.07     |
| Ta'u        | 2014 | 1.06     |
| Tutuila N   | 2014 | 1.01     |
| Tutuila S   | 2014 | 0.94     |

Calculate indicator scores and combine into index grades.



Communicate results using visual elements such as photos, maps, and conceptual diagrams.

## II. Pacific jurisdictions with reef areas

### American Samoa

American Samoa is an unincorporated United States Territory in the South Pacific. The Territory consists of five volcanic islands and two atolls, all of which are surrounded by fringing coral reefs. American Samoa was divided into six regions based on geographic location and data availability. The six regions are North Tutuila, South Tutuila, Ta'u, Swains Island, Ofu & Olosega, and Muliava/Rose Atoll.

*American Samoa Status report regions and areas. Note that the total area for each reporting region is the hardbottom forereef habitat less than 30 meters in depth:*

| Region               | Area (km <sup>2</sup> ) |
|----------------------|-------------------------|
| North Tutuila        | 19.23                   |
| South Tutuila        | 21.61                   |
| Ta'u                 | 9.04                    |
| Ofu & Olosega        | 7.93                    |
| Swains               | 2.81                    |
| Muliava (Rose Atoll) | 1.20                    |

### Hawaiian Archipelago

The Hawaiian Archipelago includes the state of Hawai'i and the Northwestern Hawaiian Islands. The archipelago consists of volcanic islands, atolls, and seamounts that stretch over 1500 miles from southeast to northwest. For the status report, there are two main regions, the Main Hawaiian Islands (the State of Hawai'i) and the Northwestern Hawaiian Islands. The Main Hawaiian Islands were further sub-divided into four regions based on geographic location and data availability. The four regions are O'ahu, Hawai'i, Maui Nui, and Kaua'i + Ni'ihau.

*Hawaiian Archipelago Status report Main Hawaiian Islands regions and areas. Note that the total area for each reporting region is the hardbottom forereef habitat less than 30 meters in depth:*

| Region             | Area (km <sup>2</sup> ) |
|--------------------|-------------------------|
| O'ahu              | 251.19                  |
| Hawai'i            | 168.40                  |
| Maui Nui           | 280.56                  |
| Kaua'i and Ni'ihau | 273.93                  |

*Hawaiian Archipelago Status report Northwestern Hawaiian Islands regions and areas. Note that the total area for each reporting region is the hardbottom forereef habitat less than 30 meters in depth:*

| Region         | Area (km <sup>2</sup> ) |
|----------------|-------------------------|
| French Frigate | 169.02                  |
| Kure           | 24.39                   |
| Laysan         | 33.996                  |
| Lisianski      | 309.55                  |
| Maro           | 256.07                  |
| Midway         | 32.94                   |
| Pearl & Holmes | 84.98                   |

## Pacific Remote Islands

*Pacific Remote Islands Status report regions and areas. Note that the total area for each reporting region is the hardbottom forereef habitat less than 30 meters in depth:*

| Region          | Area (km <sup>2</sup> ) |
|-----------------|-------------------------|
| Johnston        | 65.74                   |
| Kingman Palmyra | 49.78                   |
| Howland Baker   | 5.63                    |
| Jarvis          | 3.66                    |
| Wake            | 2.80                    |

## Guam

*Guam Status report regions and areas. Note that the total area for each reporting region is the hardbottom forereef habitat less than 30 meters in depth:*

| Region                 | Area (km <sup>2</sup> ) |
|------------------------|-------------------------|
| Western Guam           | 22.30                   |
| Eastern Guam           | 21.13                   |
| Marine Protected Areas | 7.70                    |

## Northern Mariana Islands

Located just north of Guam in the Western Pacific, the Northern Mariana Islands is a three-hundred-mile archipelago consisting of 14 islands. The Northern Mariana Islands were divided into four sub-regions, Saipan, Tinian, and Aguijan, Northern Islands, Rota, and National Monument.

*Northern Mariana Islands Status report regions and areas. Note that the total area for each reporting region is the hardbottom foreereef habitat less than 30 meters in depth:*

| Region                  | Area (km <sup>2</sup> ) |
|-------------------------|-------------------------|
| Saipan, Tinian, Aguijan | 53.59                   |
| Northern Islands        | 31.07                   |
| Rota                    | 13.31                   |
| National Monument       | 7.01                    |

### III. Indicator Development

NOAA's National Coral Reef Monitoring Program defines its four main monitoring data themes in its monitoring plan (NOAA, 2014). The four themes are fish, benthos, climate, and human connections monitoring with associated indicators within each theme. During the initial workshop, presentations of available data were given by experts followed by breakout sessions to determine appropriate indicators for this product within each theme (fish, benthos, climate, and human connections). The criteria which experts used to choose indicators were: 1) data availability, 2) sufficient understanding of reference conditions, and 3) importance to overall ecosystem health. These indicators were refined over months of discussion between different groups, jurisdictions, and NOAA headquarters.

*Coral reef status report indicators, indicator categories, and scoring system.*

| Indicators                     | Indicator categories |                                                                                     | Scoring system for all indicators                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
|--------------------------------|----------------------|-------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Coral reef cover               | Coral and algae      |  | <h3>What do the scores mean?</h3> <div><div>90–100% <b>Very good</b></div><p>All or almost all indicators meet reference values. Conditions in these locations are unimpacted, or minimally impacted or have not declined. *Human connections are very high.</p><div>80–89% <b>Good</b></div><p>Most indicators meet reference values. Conditions in these locations are lightly impacted or have lightly declined. *Human connections are high.</p><div>70–79% <b>Fair</b></div><p>Some indicators meet reference values. Conditions in these locations are moderately impacted or have declined moderately. *Human connections are moderate.</p><div>60–69% <b>Impaired</b></div><p>Few indicators meet reference values. Conditions in these locations are very impacted or have declined considerably. *Human connections are lacking.</p><div>0–59% <b>Critical</b></div><p>Very few or no indicators meet reference values. Conditions in these locations are severely impacted or have declined substantially. *Human connections are severely lacking.</p><div>Insufficient data, not scored</div></div> |
| Coral populations              |                      |                                                                                     |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| Herbivory                      |                      |                                                                                     |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| Mortality                      |                      |                                                                                     |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| Diversity (not scored)         |                      |                                                                                     |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| Reef fish                      | Fish                 |  |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| Sustainability                 |                      |                                                                                     |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| Sharks and other predators     |                      |                                                                                     |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| Diversity (not scored)         |                      |                                                                                     |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| Temperature stress             | Climate              |  |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| Ocean acidification            |                      |                                                                                     |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| Reef material growth           |                      |                                                                                     |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| Awareness                      | Human Connections    |  |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| Support for management actions |                      |                                                                                     |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| Pro-environmental behavior     |                      |                                                                                     |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |

## Define reference and assessment points

The reference point (or baseline) is the value against which the current status is evaluated. The reference points for all fish and climate indicators were chosen to represent an historical or pre-human impact condition. For benthic indicators, the reference points were either a combination of expert opinion, published literature, or the best available data. Human Connection reference and assessment points were chosen differently – please refer to that section for more information. Assessment points are the breakpoints used to determine different scoring bins. Reference and assessment points can be determined in several ways, including using regulatory criteria assessment, established management goals, literature reviews of best practices, and expert opinions. At each of the workshops, breakout groups proposed potential ideas for assessment of specific indicators. Most of the reference and assessment points were determined through a series of exploratory data analyses with input from a variety of stakeholders in each jurisdiction.

## IV. Indicators and scoring process

The following sections detail the process by which individual indicators for each Pacific jurisdiction were scored. The sections are organized by theme – coral and algae (benthos), fish, climate, and human connections. Scores are calculated on a 0-100% scale, with descriptive words and narrative text accompanying each score.

### Corals and Algae (Benthos)

Indicators in this theme include: Coral reef cover (coral cover, algal cover and CCA cover), coral populations (juvenile and adult coral density), partial mortality of adult corals, herbivory, and diversity.

Coral cover, crustose coralline algae cover, and macroalgae cover are averaged and reported as coral reef cover. Juvenile density and adult density scores are averaged, but are shown as individual scores on the status report. The overall coral and algae score consists of the average of coral reef cover, coral populations (density), partial mortality, and herbivory.

The goal was to include a limited number of metrics that provide stable indicators of coral population and benthic community status, and can be tracked over time with NCRMP data. Surveys were located using a stratified-random survey design with the survey domain being all hard bottom forereef habitat from 0-30 m.

### *Coral reef cover indicator (benthic cover)*

Raw benthic cover data are derived from the analysis of benthic photos collected at all coral and fish survey sites. Coral reef cover includes three sub-indicators: percent coral cover, percent crustose coralline algae (CCA) cover, and percent macroalgae cover. For some islands lacking analyzed benthic imagery, the benthic cover indicator metric was derived from visual estimates collected at the fish survey sites.

### *Adult and juvenile coral density*

Juvenile and adult coral densities are measures of the population's ability to reproduce and sustain itself. Juvenile coral density was included in this status report to provide additional information about the potential outlook for these coral populations – Are there babies on the reef that will potentially grow into adults?

Raw coral colony density data is derived from benthic demographic surveys conducted at each site (see table below) within two, 18-m belt transects. Adult coral colonies ( $\geq 5$  cm) were surveyed within 4, 1 x 2.5 m<sup>2</sup> segments (total of 10 m<sup>2</sup>) on each transect. Adult colonies were identified to genus or species and measured (maximum diameter) to the nearest cm. Juvenile coral colonies ( $< 5$  cm) were surveyed within 3, 1 m<sup>2</sup> segments (total 3 m<sup>2</sup>) on each transect and identified in the field by a distinct features that distinguished them from asexual fragments of larger adult colonies. Each juvenile colony was identified to the lowest taxonomic level possible (genus or species) and measured (both the maximum and perpendicular diameter to the nearest 2 mm).

### *Partial mortality*

Partial mortality is a measure of cumulative mortality and may be a proxy for loss of reproductive biomass within coral populations. The adult coral partial mortality indicator was calculated as mean 'old dead' percent. Old dead mortality is defined as the non-living portion of a colony where the skeletal structures are either eroded or covered over by organisms that are not easily removed. This portion of the colony is where the live tissue is presumed to have died within the last few months to years or longer and has a low probability of tissue recovery.

Raw coral partial mortality data is derived from the same surveys conducted to assess adult coral colony densities (as above), whereby for each adult colony assessed partial colony mortality is visually quantified as the percent of dead tissue.

### *Herbivory*

This herbivory indicator is a measure of the level of feeding pressure by fish on corals and algae. While the herbivory indicator is included as a benthic indicator, the sampling data, methodologies and scoring follow the same protocols as the reef fish biomass indicator (see that section). The status of each region was scored based on a study published in 2015 that used data from these islands and >30 others in the U.S. Pacific, and which assessed the importance of a range of human, oceanographic (e.g. temperature, oceanic productivity) and habitat drivers (e.g. coral cover, habitat structural complexity). That study estimated what 'baseline biomass' would be at each location in the absence of humans – i.e. what would be natural for each location, based entirely on its oceanographic and habitat setting (Williams et al PLoS One, 2015). For the herbivory indicator, local and baseline biomass are for those species classified as 'primary consumers', which include parrotfishes, chub, most surgeonfishes, as well as a number of other mostly small fishes such as some damselfishes.

## Diversity

Biodiversity is the complex variety of life within our ecosystems that forms the foundation for the vast array of ecosystem services that contribute to human well-being. Each of the many 1000s of species occupying coral reefs provide essential ecological functions that maintain ecosystem health. Since different related species often play similar and redundant functional roles but have levels of sensitivity to certain stressors, reefs with higher levels of biodiversity tend to have increased functional redundancy and likelihood of reef survival during episodic disturbance events, such as disease outbreaks or coral bleaching. Hence, biodiversity serves as a useful indicator of ecological resilience and the ability to survive or recover from disturbance events.

At the time of the status report process, we were not yet able to come to consensus on appropriate biodiversity thresholds that relate meaningfully to scores for ecosystem health. In future status reports, however, biodiversity will hopefully be developed into a scored indicator.

## American Samoa

Data collected in 2015 were used for the current status report.

### *Coral reef cover indicator (benthic cover)*

Coral reef cover includes three sub-indicators: percent coral cover, percent crustose coralline algae (CCA) cover, and percent macroalgae cover. Each of these sub-indicators followed a similar scoring process. Benthic cover estimates were calculated for each site. The site-specific percent cover estimates were pooled by sector and area weighted to provide an island-wide or sector-wide mean. These area-weighted benthic cover estimates were scored using criteria below. Island/sector- wide means were area-weighted by percent forereef habitat when rolled up to the jurisdiction-level.

For American Samoa, reference and assessment points were chosen using expert judgement by the NOAA Pacific Islands Fisheries Science Center, Ecosystem Sciences Division benthic team (Bernardo Vargas-Ángel and Dione Swanson) and reviewed by a larger expert group of partners and collaborators.

*Number of sampling sites for all benthic cover indicator data (coral and fish survey sites).*

| Island or sector     | # of sites |
|----------------------|------------|
| South Tutuila        | 142        |
| North Tutuila        | 104        |
| Ofu & Olosega        | 82         |
| Muliava (Rose Atoll) | 76         |
| Ta'u                 | 67         |
| Swains               | 50         |

### *Coral cover*

For corals, a maximum of 50% and above was considered very good based on expert opinion, while <10% was considered a critical condition.

*Coral cover assessment points and associated scoring. The score is determined using an equation that relates coral cover to a percent score.*

| Coral cover (%) | Score (%) | Equation |
|-----------------|-----------|----------|
| ≥50             | 100       | $Y=100$  |
| 40-<50          | 90-<100   | $Y=x+50$ |
| 30-<40          | 80-<90    | $Y=x+50$ |
| 20-<30          | 70-<80    | $Y=x+50$ |
| 10-<20          | 60-<70    | $Y=x+50$ |
| 0-<10           | 0-<60     | $Y=6x$   |

### *CCA cover*

For CCA, a maximum of 30% and above was considered very good based on expert opinion, while <2% was considered severely deteriorated (critical).

*Crustose coralline algae assessment points and associated scoring. The score is determined using an equation that relates crustose coralline algae cover to a percent score.*

| CCA cover (%) | Score (%) | Equation        |
|---------------|-----------|-----------------|
| ≥30           | 100       | $Y=100$         |
| 20-<30        | 90-<100   | $Y=x+70$        |
| 10-<20        | 80-<90    | $Y=x+70$        |
| 5-<10         | 70-<80    | $Y=2x+60$       |
| 2-<5          | 60-<70    | $Y=3.33x+53.33$ |
| 0-<2          | 0-<60     | $Y=30x$         |

### *Macroalgal cover*

For macroalgae a reverse scale was set, whereby lower macroalgal levels scored higher. As such, macroalgal cover of 2.5% or less was considered very good while 30% and above was considered a critical reef condition.

*Macroalgae assessment points and associated scoring. The score is determined using an equation that relates macroalgal cover to a percent score. For macroalgae, this is a negative relationship (more macroalgae receives a worse score).*

| Macroalgae (%) | Score (%) | Equation    |
|----------------|-----------|-------------|
| ≤2.5           | 100       | $Y=100$     |
| >2.5-5         | 90-<100   | $Y=-4x+110$ |
| >5-10          | 80-<90    | $Y=-2x+100$ |
| >10-20         | 70-<80    | $Y=-x+90$   |
| >20-30         | 60-<70    | $Y=-x+90$   |
| >30            | 0-<60     | $Y=-6x+240$ |

### *Adult and juvenile coral density*

The juvenile and adult coral density indicators were generated from a select list of the most abundant and ecologically relevant taxa represented across all islands within the region. The use of the selected taxa (genera and species) provides information about existing coral populations and incorporates a mechanism to determine how coral populations are changing over time (fluctuations in one or more of the most abundant or important taxa). The selected 10 genera and 4 species that were used to generate estimates for the coral population indicators are listed in the table below.

*Number of sampling sites for all benthic population indicator data.*

| Island or sector     | # of sites |
|----------------------|------------|
| South Tutuila        | 48         |
| North Tutuila        | 41         |
| Ofu & Olosega        | 31         |
| Muliava (Rose Atoll) | 11         |
| Ta'u                 | 21         |
| Swains               | 18         |

The juvenile and adult coral density indicators were generated from a select list of abundant and important coral genera and species represented across all islands within the region. The use of the selected taxa (genera and species) provides information about existing coral populations and incorporates a mechanism to determine how coral populations are changing over time (fluctuations in one or more of the most abundant or important taxa). The following 10 genera and 4 species were used to generate estimates for the coral population indicators:

| Genera      | Species                         |
|-------------|---------------------------------|
| Acropora    | <i>Astreopora myriophthalma</i> |
| Astreopora  | <i>Galaxea fascicularis</i>     |
| Galaxea     | <i>Astrea curta</i>             |
| Goniastrea  | <i>Pavona varians</i>           |
| Isopora     |                                 |
| Montipora   |                                 |
| Pavona      |                                 |
| Pocillopora |                                 |
| Porites     |                                 |
| Psammocora  |                                 |

Throughout the literature, there are not clear benchmarks to establish reference and assessment points for density of juvenile and adult corals due to the natural variability in abundance across habitat space. Island and sector level scores for adults and juveniles were calculated similar to percent cover in American Samoa using the following steps.

1. Reef area-weighted island and sector density estimates were generated from strata means for each of the 10 genera and 4 species listed above for 2015 when the stratified random surveys were conducted.
2. The score was calculated as a percent of the maximum island or sector density as follows:
  - a. Taxon density score (%) = (taxon A density estimate for island A / maximum island or sector density estimate for taxon A) \* 100.
  - b. The scores were average across all taxa for a given island or sector.
  - c. The scores were then converted to a 0-100% scale.

*Coral population density assessment points and associated scoring. The score is determined using an equation that relates the percent of maximum density to a percent score.*

| Coral population (colonies per square meter) | Score (%) | Equation     |
|----------------------------------------------|-----------|--------------|
| 60-100                                       | 90-100    | $Y=0.25x+75$ |
| 40-<60                                       | 80-<90    | $Y=0.50x+60$ |
| 20-<40                                       | 70-<80    | $Y=0.50x+60$ |
| 10-<20                                       | 60-<70    | $Y=x+50$     |
| 0-<10                                        | 0-<60     | $Y=6x$       |

The bins were truncated at the lower range as a conservative approach to account for habitat specific difference in density for adults and juveniles. For example, some habitats have naturally low coral density and low density of juveniles can co-occur with high adult density because the amount of available space for settlement is low.

### *Partial mortality*

The partial mortality indicator was generated from the same select list of abundant and important coral species and genera used for juvenile and adult density.

Similar to colony density, there are not clear benchmarks for thresholds of partial mortality estimates other than low values are good and higher values may be bad. In the process of threshold development, we evaluated the range in island estimates in American Samoa as well as other regions in the Pacific and Atlantic. The maximum of the range was set at 30% for all genera and species. This represents a conservative approach but hopefully can indicate significant shifts in partial mortality.

Mean partial mortality scores were generated for each genus and species using the following formula for each taxon and each island or sector:

Taxon partial mortality score (%) = (taxon A partial mortality estimate for island A (%)) / 30 % \* 100.

Mean scores were then calculated from the 10 genera and 4 species for each island and sector.

*Coral mortality assessment points and associated scoring. The score is determined using an equation that relates the percent of maximum mortality to a percent score. For mortality, this is a negative relationship (more mortality receives a worse score).*

| Coral mortality (%) | Score (%) | Equation          |
|---------------------|-----------|-------------------|
| 0-20                | >90-100   | $Y = -0.5x + 100$ |
| 20-40               | 80-90     | $Y = -0.5x + 100$ |
| 40-60               | 70-80     | $Y = -0.5x + 100$ |
| 60-80               | 60-70     | $Y = -0.5x + 100$ |
| 80-100              | 0-60      | $Y = -3x + 300$   |

## Hawaiian Archipelago

In the Main Hawaiian Islands, data collected in 2016 were used for the current status report. For the Northwestern Hawaiian Islands, the data collected from a combination of survey efforts was used for the status report to provide the most recent estimates for each island across the Northwestern portion of the archipelago. These survey years include 2010 through 2016.

To account for differences in how benthic communities form across a range of habitats (e.g. coral-rich, pavement, rock and boulder), the benthic maps in the main Hawaiian Islands are divided into sectors of broad reef structure which include coral-rich (e.g. aggregate reef, spur and groove), complex (e.g. rock and boulder), and simple (e.g. pavement). Sites within each sector and island were classified by these habitat types.

In the Northwestern Hawaiian Islands, mapped benthic habitats are currently unreliable. Instead we used diver visual habitat classifications between 2010 and 2016 to calculate the percent of sites for a given island that fell within certain habitat structure types. We then assigned each island a habitat structure type if greater than 50% of the sites were classified of that habitat structure. Habitat structure types consisted of coral-rich (e.g. aggregate reef, spur and groove) and simple (e.g. pavement) for the islands included in this status report.

### *Coral reef cover indicator (benthic cover)*

Coral reef cover includes three sub-indicators: percent coral cover, percent crustose coralline algae (CCA) cover, and percent macroalgae cover. Each of these sub-indicators followed a similar scoring process. Average benthic cover estimates were calculated for each site. The site-specific percent cover estimates were pooled by depth stratum, averaged, and stratum area weighted to provide an island-wide or sector-wide mean. These area-weighted benthic cover estimates were scored using criteria below. Island/sector- wide means were area-weighted by percent forereef habitat when rolled up to Main Hawaiian Island –wide level and Northwestern Hawaiian Island –wide level.

For each sub-indicator, the most recent percent cover values for a given island or sector were compared to the maximum island or sector-level value reported between 2013 and 2016 for the Main Hawaiian Islands and between 2010 and 2016 for the Northwestern Hawaiian Islands. These were categorized for a given habitat structure to calculate the percent of maximum. In other words, islands dominated by coral-rich habitats were only compared to other coral-rich islands. The percent of maximum values were then converted to a 0–100% scale using the equations listed in the tables below. For macroalgae, this is a negative relationship (more macroalgae receives a worse score). To calculate scores for macroalgae the 60–100% scores were inverted so that 0% max = 100%.

*Number of sampling sites for all benthic cover indicator data (coral and fish survey sites). The number of sites are listed for Main Hawaiian Islands (MHI) and Northwestern Hawaiian islands (NWHI). CR= Coral Rich, CM= Complex, SI= Simple*

| OBS_YEAR | REGION | ISLAND  | Habitat | n  |
|----------|--------|---------|---------|----|
| 2013     | MHI    | Hawaii  | CM      | 58 |
| 2013     | MHI    | Hawaii  | CR      | 22 |
| 2013     | MHI    | Kauai   | SI      | 50 |
| 2013     | MHI    | Lanai   | CM      | 21 |
| 2013     | MHI    | Lanai   | CR      | 22 |
| 2013     | MHI    | Maui    | CM      | 28 |
| 2013     | MHI    | Maui    | SI      | 12 |
| 2013     | MHI    | Molokai | CM      | 21 |
| 2013     | MHI    | Molokai | CR      | 17 |
| 2013     | MHI    | Molokai | SI      | 21 |
| 2013     | MHI    | Niihau  | CM      | 7  |
| 2013     | MHI    | Niihau  | SI      | 36 |
| 2013     | MHI    | Oahu    | SI      | 22 |
| 2013     | MHI    | Oahu    | SI      | 74 |
| 2015     | MHI    | Hawaii  | CM      | 32 |
| 2015     | MHI    | Hawaii  | CR      | 65 |
| 2015     | MHI    | Kauai   | SI      | 20 |
| 2015     | MHI    | Lanai   | CM      | 15 |
| 2015     | MHI    | Maui    | CM      | 28 |
| 2015     | MHI    | Maui    | SI      | 2  |
| 2015     | MHI    | Molokai | CR      | 32 |
| 2015     | MHI    | Molokai | SI      | 16 |
| 2015     | MHI    | Niihau  | CM      | 15 |
| 2015     | MHI    | Niihau  | SI      | 33 |
| 2015     | MHI    | Oahu    | SI      | 1  |
| 2015     | MHI    | Oahu    | SI      | 34 |
| 2016     | MHI    | Hawaii  | CM      | 36 |
| 2016     | MHI    | Hawaii  | CR      | 54 |
| 2016     | MHI    | Kauai   | SI      | 47 |
| 2016     | MHI    | Lanai   | CM      | 29 |
| 2016     | MHI    | Lanai   | CR      | 12 |
| 2016     | MHI    | Maui    | CM      | 34 |
| 2016     | MHI    | Maui    | SI      | 10 |
| 2016     | MHI    | Molokai | CM      | 9  |
| 2016     | MHI    | Molokai | CR      | 10 |
| 2016     | MHI    | Molokai | SI      | 15 |
| 2016     | MHI    | Niihau  | CM      | 2  |
| 2016     | MHI    | Niihau  | SI      | 15 |

|           |      |                |    |    |
|-----------|------|----------------|----|----|
| 2016      | MHI  | Oahu           | SI | 23 |
| 2016      | MHI  | Oahu           | SI | 63 |
| 2016      | MHI  | Kahoolawe      | CR | 35 |
| 2010-2012 | NWHI | French Frigate | CR | 35 |
| 2010-2012 | NWHI | Kure           | CR | 30 |
| 2010-2012 | NWHI | Laysan         | S  | 23 |
| 2010-2012 | NWHI | Lisianski      | CR | 59 |
| 2010-2012 |      | Midway         | S  | 17 |
| 2010-2012 | NWHI | Pearl & Hermes | CR | 48 |
| 2013-2015 | NWHI | French Frigate | CR | 31 |
| 2013-2015 | NWHI | Kure           | CR | 8  |
| 2013-2015 | NWHI | Laysan         | S  | 8  |
| 2013-2015 | NWHI | Lisianski      | CR | 46 |
| 2013-2015 |      | Midway         | S  | 42 |
| 2013-2015 | NWHI | Pearl & Hermes | CR | 21 |
| 2016      | NWHI | French Frigate | CR | 73 |
| 2016      | NWHI | Kure           | CR | 58 |
| 2016      | NWHI | Laysan         | S  | 11 |
| 2016      | NWHI | Lisianski      | CR | 56 |
| 2016      | NWHI | Midway         | S  | 10 |
| 2016      | NWHI | Pearl & Hermes | CR | 75 |

### *Coral and CCA cover*

*Coral and CCA cover assessment points and associated scoring. The score is determined using an equation that relates the percent maximum coral and CCA cover to a percent score.*

| <b>Coral and CCA cover (% of max)</b> | <b>Score (%)</b> | <b>Equation</b> |
|---------------------------------------|------------------|-----------------|
| 80-100                                | >90-100          | $Y=0.5x+50$     |
| 60-<80                                | >80-<90          | $Y=0.5x+50$     |
| 40-<60                                | >70-<80          | $Y=0.5x+50$     |
| 20-<40                                | >60-<70          | $Y=0.5x+50$     |
| 0-<20                                 | 0-<60            | $Y=3x$          |

### *Macroalgal cover*

*Macroalgae assessment points and associated scoring. The score is determined using an equation that relates the percent of maximum macroalgal cover to a percent score. For macroalgae, this is a negative relationship (more macroalgae receives a worse score).*

| Macroalgae (%) | Score (%) | Equation          |
|----------------|-----------|-------------------|
| 0-<20          | >90-100   | $Y = -0.5x + 100$ |
| 20-<40         | >80-90    | $Y = -0.5x + 100$ |
| 40-<60         | >70-80    | $Y = -0.5x + 100$ |
| 60-<80         | >60-70    | $Y = -0.5x + 100$ |
| 80-100         | 0-60      | $Y = -3x + 300$   |

### *Adult and juvenile coral density*

The juvenile and adult coral density indicators were generated from a select list of the most abundant and important coral genera and species represented across all islands within a given habitat. The use of the selected taxa (genera and species) provides information about existing coral populations and incorporates a mechanism to determine how coral populations are changing over time (fluctuations in one or more of the most abundant or important taxa). The selected coral taxa for each habitat structure type within the Main Hawaiian Islands and Northwestern Hawaiian Islands that were used to generate estimates for the coral population indicators are listed in the table below.

*Number of sampling sites for all benthic population indicator data in the (A) Main Hawaiian Islands and (B) Northwestern Hawaiian Islands. The dashes indicate the habitat structure type was not found and the zeros indicate not surveyed.*

#### Main Hawaiian Islands

| Island    | Coral-Rich | 2013 Complex | Simple | Coral-Rich | 2016 Complex | Simple |
|-----------|------------|--------------|--------|------------|--------------|--------|
| Hawaii    | 6          | 18           | -      | 18         | 13           | -      |
| Kahoolawe | 0          | -            | -      | 11         | -            | -      |
| Kauai     | -          | -            | 14     | -          | -            | 17     |
| Lanai     | 7          | 7            | -      | 4          | 11           | -      |
| Maui      | -          | 6            | 2      | -          | 11           | 4      |
| Molokai   | 6          | 8            | 7      | 3          | 3            | 5      |
| Niihau    | -          | 6            | 11     | -          | 0            | 5      |
| Oahu      | -          | 6            | 35     | -          | 12           | 20     |

#### Northwest Hawaiian Islands

| Island | n (2011-2012) | n (2014-2015) | n (2016) | Habitat type |
|--------|---------------|---------------|----------|--------------|
| FFS    | 4             | 16            | 26       | Coral-rich   |
| KUR    | 9             | 6             | 19       | Coral-rich   |
| LAY    | 9             | 5             | 0        | Simple       |
| LIS    | 17            | 21            | 16       | Coral-rich   |
| MAR    | 10            | 10            | 0        | Coral-rich   |
| MID    | 6             | 19            | 0        | Simple       |
| PHR    | 13            | 14            | 20       | Coral-rich   |

*The selected taxa of scleractinian coral used to generate all coral population indicators for the Main Hawaiian Islands and Northwestern Hawaiian Islands by habitat structure type.*

| <b>Main Hawaiian Islands</b>        | <b>Northwestern Hawaiian Islands</b> |
|-------------------------------------|--------------------------------------|
| <b>Coral-rich habitat structure</b> |                                      |
| <i>Montipora capitata</i>           | <i>Acropora cytherea</i>             |
| <i>Montipora patula</i>             | <i>Cyphastrea ocellina</i>           |
| <i>Porites compressa</i>            | <i>Montipora capitata</i>            |
| <i>Porites lobata</i>               | <i>Montipora patula</i>              |
| <i>Pocillopora meandrina</i>        | <i>Porites compressa</i>             |
|                                     | <i>Porites lobata</i>                |
|                                     |                                      |
| <b>Complex habitat structure</b>    |                                      |
| <i>Montipora capitata</i>           |                                      |
| <i>Montipora patula</i>             |                                      |
| <i>Porites compressa</i>            |                                      |
| <i>Porites lobata</i>               |                                      |
| <i>Pocillopora meandrina</i>        |                                      |
|                                     |                                      |
| <b>Simple habitat structure</b>     |                                      |
| <i>Montipora capitata</i>           | <i>Montipora capitata</i>            |
| <i>Montipora patula</i>             | <i>Porites compressa</i>             |
| <i>Porites lobata</i>               | <i>Pavona duerdeni</i>               |
| <i>Pocillopora meandrina</i>        | <i>Porites lobata</i>                |
|                                     | <i>Pocillopora meandrina</i>         |

Throughout the literature, there are not clear benchmarks to establish reference and assessment points for density for juvenile and adult corals due to the natural variability in abundance across habitat space. The island level scores for adults and juveniles we calculated similar to the percent cover using the following steps:

1. Reef area-weighted island and sector density estimates were generated from strata means for each of the species listed above for 2013 through 2016 (MHI) and 2011 through 2016 (NWHI) when stratified random surveys were conducted.
2. The score was calculated as a percent of the maximum island or sector density as follows:
  - a.  $\text{Taxon density \% max score} = (\text{taxon A density estimate for island or sector A} / \text{maximum island or sector density estimate for habitat A reported for all years for taxon A}) * 100.$
  - b. The scores were averaged across all taxa for a given island or sector.
  - c. The scores were then converted to a 0-100% scale using equations in the table below.
  - d. These scores were generated separately for juveniles and adults.

*Coral population density assessment points and associated scoring. The score is determined using an equation that relates the percent of maximum density to a percent score.*

| % of Maximum | Score (%) | Equation        |
|--------------|-----------|-----------------|
| 80-100       | 90-100    | $y = 0.5x + 50$ |
| 60-80        | 80-89     | $y = 0.5x + 50$ |
| 40-60        | 70-79     | $y = 0.5x + 50$ |
| 20-40        | 60-69     | $y = 0.5x + 50$ |
| 0-20         | 0-59      | $y = 3x$        |

### *Partial mortality*

The partial mortality indicator was generated from the same select list of abundant and important coral species used for juvenile and adult density.

Similar to density, there are not clear benchmarks to establish reference and assessment points for partial mortality due to the natural variability in abundance across habitat space. The island level scores for partial mortality we calculated similar to the other benthic metrics as follows:

1. Reef area-weighted island and sector density estimates were generated from strata means for each of the taxa listed above for 2013 through 2016 (MHI) and 2011 through 2016 (NWHI) when stratified random surveys were conducted.
2. The score was calculated as a percent of the maximum island or sector density as follows:
  - a. Taxon density % max score = (taxon A density estimate for island or sector A / maximum island or sector density estimate for habitat A reported for all years for taxon A) \* 100.
  - b. The scores were averaged across all taxa for a given island or sector.
  - c. The scores were then converted to a 0-100% on an inverted scale using equations in table below.

*Coral mortality assessment points and associated scoring. The score is determined using an equation that relates the percent of maximum mortality to a percent score. For mortality, this is a negative relationship (more mortality receives a worse score).*

| % of Maximum | Score (%) | Equation          |
|--------------|-----------|-------------------|
| 0-20         | 90-100    | $y = -0.5x + 100$ |
| 20-40        | 80-89     | $y = -0.5x + 100$ |
| 40-60        | 70-79     | $y = -0.5x + 100$ |
| 60-80        | 60-69     | $y = -0.5x + 100$ |
| 80-100       | 0-59      | $y = -3x + 300$   |

### **Pacific Remote Islands**

Data collected in 2015 were used for the current status report for all islands except Jarvis Island and Wake Atoll. Recent surveys conducted during 2017 were used these two locations. The habitat structure types were determined to be similar among all island and atolls in the Pacific Remote jurisdiction.

### *Coral reef cover indicator (benthic cover)*

Coral reef cover includes three sub-indicators: percent coral cover, percent crustose coralline algae (CCA) cover, and percent macroalgae cover. Each of these sub-indicators followed a similar scoring process. Average benthic cover estimates were calculated for each site. The site-specific percent cover estimates were pooled by stratum, averaged, and stratum area weighted to provide an Island-wide or sector-wide mean. These area weighted benthic cover estimates were scored using criteria below. Island/sector-wide means were area weighted by percent forereef habitat when rolled up to Pacific Remote-wide level.

For each sub-indicator, the most recent percent cover values for a given island were compared to the maximum island-level value reported between 2014 and 2017 to calculate the percent of maximum. The percent of maximum values were then converted to a 60-100% scale using the equations listed in the table below. To calculate scores for macroalgae the 60–100% scores were inverted so that 0% max = 100%.

*Number of sampling sites for all benthic cover indicator data (coral and fish survey sites). CR = coral-rich*

| Year | Island   | Habitat | n   |
|------|----------|---------|-----|
| 2014 | Wake     | CR      | 65  |
| 2015 | Baker    | CR      | 51  |
| 2015 | Howland  | CR      | 56  |
| 2015 | Jarvis   | CR      | 102 |
| 2015 | Johnston | CR      | 46  |
| 2015 | Kingman  | CR      | 73  |
| 2015 | Palmyra  | CR      | 114 |
| 2017 | Wake     | CR      | 81  |
| 2017 | Baker    | CR      | 23  |
| 2017 | Howland  | CR      | 20  |
| 2017 | Jarvis   | CR      | 60  |

### *Coral and CCA cover*

*Coral and CCA cover assessment points and associated scoring. The score is determined using an equation that relates coral and CCA cover to a percent score. The score relates to narrative text describing coral condition.*

| % of maximum | Score (%) | Equation    |
|--------------|-----------|-------------|
| 80-100       | 90-100    | $Y=0.5x+50$ |
| 60-<80       | 80-<90    | $Y=0.5x+50$ |
| 40-<60       | 70-<80    | $Y=0.5x+50$ |
| 20-<40       | 60-<70    | $Y=0.5x+50$ |
| 0-<20        | 0-<60     | $Y=3x$      |

### *Macroalgal cover*

*Macroalgae assessment points and associated scoring. The score is determined using an equation that relates the percent of maximum macroalgal cover to a percent score. For macroalgae, this is a negative relationship (more macroalgae receives a worse score).*

| % of maximum | Score (%) | Equation      |
|--------------|-----------|---------------|
| 0-<20        | >90-100   | $Y=-0.5x+100$ |
| 20-<40       | >80-90    | $Y=-0.5x+100$ |
| 40-<60       | >70-80    | $Y=-0.5x+100$ |
| 60-<80       | >60-70    | $Y=-0.5x+100$ |
| 80-100       | 0-60      | $Y=-3x+300$   |

### *Adult and juvenile coral density*

The juvenile and adult coral density indicators were generated from a select list of the most abundant and important coral genera and species unique to each island or atoll given the physical distance between them. The use of the selected taxa (genera and species) provides information about existing coral populations and incorporates a mechanism to determine how coral populations are changing over time (fluctuations in one or more of the most abundant or important taxa). The selected coral taxa for each island or atoll in the Pacific Remote jurisdiction that were used to generate estimates for the coral population indicators are listed in the table below.

*Number of sampling sites for all benthic population indicator data.*

| Island/year | n  |
|-------------|----|
| 2014-2015   |    |
| Baker       | 15 |
| Howland     | 21 |
| Jarvis      | 41 |
| Johnston    | 10 |
| Kingman     | 21 |
| Palmyra     | 39 |
| Wake        | 20 |
|             |    |
| 2016        |    |
| Jarvis      | 30 |
|             |    |
| 2017        |    |
| Jarvis      | 32 |
| Wake        | 28 |

The selected taxa of scleractinian coral used to generate all coral population indicators for each island or atoll in the Pacific Remote Island Areas jurisdiction.

|                        |                     |
|------------------------|---------------------|
| <b>Baker</b>           | <b>Kingman</b>      |
| Genera                 | Genera              |
| <i>Acropora</i>        | <i>Acropora</i>     |
| <i>Montipora</i>       | <i>Favia</i>        |
| <i>Pocillopora</i>     | <i>Fungia</i>       |
| <i>Porities</i>        | <i>Montipora</i>    |
| Species                | <i>Pavona</i>       |
| <i>Favia mattahaii</i> | <i>Pocillopora</i>  |
|                        | <i>Porities</i>     |
|                        | Species             |
|                        | <i>Astrea curta</i> |
|                        |                     |
| <b>Howland</b>         | <b>Palmyra</b>      |
| Genera                 | Genera              |
| <i>Acropora</i>        | <i>Favia</i>        |
| <i>Montipora</i>       | <i>Montipora</i>    |
| <i>Pavona</i>          | <i>Pavona</i>       |
| <i>Pocillopora</i>     | <i>Pocillopora</i>  |
| <i>Porites</i>         | <i>Porites</i>      |
|                        | Species             |
|                        | <i>Astrea curta</i> |
|                        |                     |
| <b>Jarvis</b>          | <b>Wake</b>         |
| Genera                 | Genera              |
| <i>Fungia</i>          | <i>Acanthastrea</i> |
| <i>Montipora</i>       | <i>Astreopora</i>   |
| <i>Pavona</i>          | <i>Favia</i>        |
| <i>Pocillopora</i>     | <i>Goniastrea</i>   |
| <i>Porites</i>         | <i>Montipora</i>    |
|                        | <i>Pocillopora</i>  |
|                        | <i>Porites</i>      |
|                        |                     |
| <b>Johnston</b>        |                     |
| Genera                 |                     |
| <i>Acropora</i>        |                     |
| <i>Montipora</i>       |                     |
| <i>Pavona</i>          |                     |
| <i>Pocillopora</i>     |                     |
| <i>Porites</i>         |                     |

Throughout the literature, there are no clear benchmarks to establish reference and assessment points for density for juvenile and adult corals due to the natural variability in abundance across habitat space. The island level scores for adults and juveniles we calculated similar to the percent cover using the following steps:

1. Reef area-weighted island and sector density estimates were generated from strata means for each of the coral taxon listed above for the most recent data (2015 or 2017) when stratified random surveys were conducted.
2. The score was calculated as a percent of the maximum island or sector density as follows:
  - a. Taxon density % max score = (taxon A density estimate for island A / maximum island or sector density estimate reported between 2014 and 2017 for taxon A) \* 100.
  - b. The scores were averaged across all taxa for a given island or sector.
  - c. The scores were then converted to a 60-100% scale using the equations below.
  - d. These scores were generated separately for juveniles and adults.

*Coral population density assessment points and associated scoring. The score is determined using an equation that relates the percent of maximum density to a percent score.*

| % of maximum | Score (%) | Equation     |
|--------------|-----------|--------------|
| 60-100       | 90-100    | $Y=0.25x+75$ |
| 40-<60       | 80-<90    | $Y=0.50x+60$ |
| 20-<40       | 70-<80    | $Y=0.50x+60$ |
| 10-<20       | 60-<70    | $Y=x+50$     |
| 0-<10        | 0-<60     | $Y=6x$       |

### *Partial mortality*

The partial mortality indicator was generated from the same select list of abundant and important coral species and genera used for juvenile and adult density.

Similar to density, there are no clear benchmarks to establish reference and assessment points for partial mortality due to the natural variability in abundance across habitat space. The island level scores for partial mortality we calculated similar to the other benthic metrics as follows:

1. Reef area-weighted island and sector density estimates were generated from strata means for each of the coral taxon listed above for 2015 and 2017 when stratified random surveys were conducted.
2. The score was calculated as a percent of the maximum island or sector density as follows:
  - a. Taxon density % max score = (taxon A density estimate for island A / maximum island or sector density estimate reported between 2014 and 2017 for taxon A) \* 100.
  - b. The scores were averaged across all taxa for a given island or sector.
  - c. The scores were then converted to a 60-100% on an inverted scale.

*Coral mortality assessment points and associated scoring. The score is determined using an equation that relates the percent of maximum mortality to a percent score. For mortality, this is a negative relationship (more mortality receives a worse score).*

| % of maximum | Score (%) | Equation          |
|--------------|-----------|-------------------|
| 0-20         | 90-100    | $Y = -0.5x + 100$ |
| 20-<40       | 80-<90    | $Y = -0.5x + 100$ |
| 40-<60       | 70-<80    | $Y = -0.5x + 100$ |
| 60-<80       | 60-<70    | $Y = -0.5x + 100$ |
| 80-100       | 0-<60     | $Y = -3x + 300$   |

## Guam

Habitat structure types around Guam consisted of coral-rich habitat structure (e.g. aggregate reef, spur and groove). Benthic indicator estimates are scaled as described below by estimates generated throughout Guam and the Northern Mariana Islands. Habitat structure types throughout the Northern Mariana Islands included the habitat structure types coral-rich, mixed habitat structures (no predominant habitat types), and complex (e.g. rock and boulder).

### *Coral reef cover indicator (benthic cover)*

Coral reef cover includes three sub-indicators: percent coral cover, percent crustose coralline algae (CCA) cover, and percent macroalgae cover. Each of these sub-indicators followed a similar scoring process. Average benthic cover estimates were calculated for each site. The site-specific percent cover estimates were pooled by stratum, averaged, and stratum area weighted to provide an Island-wide or sector-wide mean. These area weighted benthic cover estimates were scored using criteria below. Sector-wide means were area weighted by percent for reef habitat when rolled up to Guam-wide level.

*Number of sampling sites for all benthic cover indicator data (coral and fish survey sites). CR = Coral Rich.*

| Year | Habitat | n   |
|------|---------|-----|
| 2014 | CR      | 108 |
| 2017 | CR      | 97  |

For each sub-indicator, the most recent percent cover values for a given island were compared to the maximum island-level value reported between 2014 and 2017 for a given habitat structure type to calculate the percent of maximum. In other words, islands dominated by coral-rich habitats were only compared to other coral rich islands. The percent of maximum values were then converted to a 60-100% scale using the equations listed below. To calculate scores for macroalgae the 60-100% scores were inverted so that 0% max = 100%.

### *Coral and CCA cover*

*Coral and CCA cover assessment points and associated scoring. The score is determined using an equation that relates the percent of maximum coral and CCA cover to a percent score.*

| % of maximum | Score (%) | Equation    |
|--------------|-----------|-------------|
| 80-100       | 90-100    | $Y=0.5x+50$ |
| 60-<80       | 80-<90    | $Y=0.5x+50$ |
| 40-<60       | 70-<80    | $Y=0.5x+50$ |
| 20-<40       | 60-<70    | $Y=0.5x+50$ |
| 0-<20        | 0-<60     | $Y=3x$      |

### *Macroalgal cover*

*Macroalgae assessment points and associated scoring. The score is determined using an equation that relates the percent of maximum macroalgal cover to a percent score. For macroalgae, this is a negative relationship (more macroalgae receives a worse score).*

| % of maximum | Score (%) | Equation      |
|--------------|-----------|---------------|
| 0-<20        | >90-100   | $Y=-0.5x+100$ |
| 20-<40       | >80-90    | $Y=-0.5x+100$ |
| 40-<60       | >70-80    | $Y=-0.5x+100$ |
| 60-<80       | >60-70    | $Y=-0.5x+100$ |
| 80-100       | 0-60      | $Y=-3x+300$   |

### *Adult and juvenile coral density*

The juvenile and adult coral density indicators were generated from a select list of the most abundant and important coral genera and species represented across all islands within Guam and Northern Marianas Islands for a given habitat structure type. The use of the selected taxa provides information about existing coral populations and incorporates a mechanism to determine how coral populations are changing over time (fluctuations in one or more of the most abundant or important taxa). The selected coral taxa for the coral-rich habitat structure type, which include east and west sectors of Guam (open areas outside of Marine Protected Areas) that were used to generate estimates for the coral population indicators, are listed in the table below.

*Number of sampling sites for all benthic population indicator data for Guam sectors.*

| Guam sectors | 2014 (n) | 2017 (n) |
|--------------|----------|----------|
| East Open    | 13       | 12       |
| West Open    | 11       | 11       |

*The selected taxa of scleractinian coral used to generate all coral population indicators for Guam.*

| Coral-rich habitat structure |
|------------------------------|
| Genera                       |
| <i>Acropora</i>              |

|                    |
|--------------------|
| <i>Astreopora</i>  |
| <i>Cyphastrea</i>  |
| <i>Favia</i>       |
| <i>Goniastrea</i>  |
| <i>Leptastrea</i>  |
| <i>Montipora</i>   |
| <i>Pavona</i>      |
| <i>Pocillopora</i> |
| <i>Porites</i>     |

Throughout the literature, there are no clear benchmarks to establish reference and assessment points for density for juvenile and adult corals due to the natural variability in abundance across habitat space. The island level scores for adults and juveniles we calculated similar to the percent cover using the following steps:

1. Reef area-weighted island and sector density estimates were generated from strata means for each of the coral taxon listed above when stratified random surveys were conducted.
2. The score was calculated as a percent of the maximum island or sector density as follows:
  - a. Taxon density % max score = (taxon A density estimate for sector A / maximum island or sector density estimate for habitat A reported for all years for taxon A) \* 100.
  - b. The scores were averaged across all taxa for a given island or sector.
  - c. The scores were then converted to a 0-100% scale using equations in the table below.
  - d. These scores were generated separately for juveniles and adults.

*Coral population density assessment points and associated scoring. The score is determined using an equation that relates the percent of maximum density to a percent score*

| % of maximum | Score (%) | Equation     |
|--------------|-----------|--------------|
| 60-100       | 90-100    | $Y=0.25x+75$ |
| 40-<60       | 80-<90    | $Y=0.50x+60$ |
| 20-<40       | 70-<80    | $Y=0.50x+60$ |
| 10-<20       | 60-<70    | $Y=x+50$     |
| 0-<10        | 0-<60     | $Y=6x$       |

### *Partial mortality*

The partial mortality indicator was generated from the same select list of abundant and important coral species and genera used for juvenile and adult density.

Throughout the literature, there are not clear benchmarks to establish reference and assessment points for density for partial mortality due to the natural variability in abundance across habitat space. The island level scores for partial mortality we calculated similar to the other benthic metrics as follows:

1. Reef area-weighted island and sector density estimates were generated from strata means for each of the coral taxon listed above when stratified random surveys were conducted.
2. The score was calculated as a percent of the maximum island or sector density as follows:
  - a. Taxon density % max score = (taxon A density estimate for sector A / maximum island or sector density estimate for habitat A reported for all years for taxon A) \* 100.
  - b. The scores were averaged across all taxa for a given island or sector.
  - c. The scores were then converted to a 60-100% on an inverted scale using equations below.

*Coral mortality assessment points and associated scoring. The score is determined using an equation that relates the percent of maximum mortality to a percent score. For mortality, this is a negative relationship (more mortality receives a worse score).*

| % of maximum | Score (%) | Equation          |
|--------------|-----------|-------------------|
| 0-20         | 90-100    | $Y = -0.5x + 100$ |
| 20-<40       | 80-<90    | $Y = -0.5x + 100$ |
| 40-<60       | 70-<80    | $Y = -0.5x + 100$ |
| 60-<80       | 60-<70    | $Y = -0.5x + 100$ |
| 80-100       | 0-<60     | $Y = -3x + 300$   |

## Northern Mariana Islands

To account for differences in how benthic communities form in across a range of habitats (e.g. coral-rich, pavement, rock and bolder), we used diver visual habitat classifications between 2011 and 2017 to calculate the percent of sites for a given island that fell within certain habitat types. We then assigned each island a habitat type if greater than 50% of the sites were classified that that habitat type. Habitat structure types throughout the Northern Marianas Islands included the habitat structure types coral-rich, mixed habitat structures (no predominant habitat types), and complex (e.g. rock and boulder).

### *Coral reef cover indicator (benthic cover)*

Coral reef cover includes three sub-indicators: percent coral cover, percent crustose coralline algae (CCA) cover, and percent macroalgae cover. Each of these sub-indicators followed a similar scoring process. Average benthic cover estimates were calculated for each site. The site-specific percent cover estimates were pooled by stratum, averaged, and stratum area weighted to provide an Island-wide or sector-wide mean. These area weighted benthic cover estimates were scored using criteria below. Island/sector-wide means were area weighted by percent forereef habitat when rolled up to Northern Marianas Islands –wide level.

*Number of sampling sites for all benthic cover indicator data (coral and fish survey sites). MX = mixed, CR = coral-rich.*

| Year | Island   | Habitat | n  |
|------|----------|---------|----|
| 2014 | Aguijan  | MX      | 16 |
| 2014 | Alamagan | MX      | 17 |

|      |                     |    |    |
|------|---------------------|----|----|
| 2014 | Asuncion            | CM | 33 |
| 2014 | Farallon de Pajaros | CM | 18 |
| 2014 | Guguan              | MX | 16 |
| 2014 | Maug                | CR | 60 |
| 2014 | Pagan               | MX | 62 |
| 2014 | Rota                | CR | 38 |
| 2014 | Saipan              | CR | 70 |
| 2014 | Sarigan             | CM | 16 |
| 2014 | Tinian              | CR | 26 |
| 2017 | Agrihan             | MX | 26 |
| 2017 | Aguijan             | CR | 27 |
| 2017 | Alamagan            | MX | 13 |
| 2017 | Asuncion            | CM | 31 |
| 2017 | Farallon de Pajaros | CM | 28 |
| 2017 | Guguan              | MX | 12 |
| 2017 | Maug                | CR | 65 |
| 2017 | Pagan               | MX | 59 |
| 2017 | Rota                | CR | 41 |
| 2017 | Saipan              | CR | 58 |
| 2017 | Sarigan             | CM | 14 |
| 2017 | Tinian              | CR | 38 |

For each sub-indicator, the most recent percent cover values for a given island were compared to the maximum island-level value reported between 2014 and 2017 for a given habitat type to calculate the percent of maximum. In other words, islands dominated by coral-rich habitats were only compared to other coral rich islands. The percent of maximum values were then converted to a 60-100% scale using the equations listed below. To calculate scores for macroalgae the 60-100% scores were inverted so that 0% max = 100%.

### *Coral and CCA cover*

*Coral and CCA cover assessment points and associated scoring. The score is determined using an equation that relates percent of maximum coral and CCA cover to a percent score.*

| % of maximum | Score (%) | Equation    |
|--------------|-----------|-------------|
| 80-100       | 90-100    | $Y=0.5x+50$ |
| 60-<80       | 80-<90    | $Y=0.5x+50$ |
| 40-<60       | 70-<80    | $Y=0.5x+50$ |
| 20-<40       | 60-<70    | $Y=0.5x+50$ |
| 0-<20        | 0-<60     | $Y=3x$      |

### *Macroalgal cover*

*Macroalgae assessment points and associated scoring. The score is determined using an equation that relates the percent of maximum macroalgal cover to a percent score. For macroalgae, this is a negative relationship (more macroalgae receives a worse score). The score relates to narrative text describing macroalgal condition.*

| % of maximum | Score (%) | Equation          |
|--------------|-----------|-------------------|
| 0-<20        | >90-100   | $Y = -0.5x + 100$ |
| 20-<40       | >80-90    | $Y = -0.5x + 100$ |
| 40-<60       | >70-80    | $Y = -0.5x + 100$ |
| 60-<80       | >60-70    | $Y = -0.5x + 100$ |
| 80-100       | 0-60      | $Y = -3x + 300$   |

### *Adult and juvenile coral density*

The juvenile and adult coral density indicators were generated from a select list of the most abundant and important coral genera and species represented across all islands within a given habitat. The use of the selected taxa (genera and species) provides information about existing coral populations and incorporates a mechanism to determine how coral populations are changing over time (fluctuations in one or more of the most abundant or important taxa). The selected coral taxa for each habitat structure type for the Mariana Islands that were used to generate estimates for the coral population indicators are listed in the table below.

*Number of sampling sites for all benthic population indicator data. Habitat structure types are noted for each island.*

| Island              | 2014 (n) | 2017 (n) | Habitat type |
|---------------------|----------|----------|--------------|
| Farallon de Pajaros | 7        | 12       | Complex      |
| Maug                | 22       | 27       | Coral-rich   |
| Asuncion            | 12       | 12       | Complex      |
| Agrihan             | 0        | 7        | Mixed        |
| Pagan               | 19       | 19       | Mixed        |
| Alamagan            | 6        | 4        | Mixed        |
| Guguan              | 5        | 3        | Mixed        |
| Sarigan             | 5        | 5        | Complex      |
| Saipan              | 22       | 22       | Coral-rich   |
| Tinian              | 7        | 14       | Coral-rich   |
| Aguijan             | 6        | 10       | Coral-rich   |
| Rota                | 10       | 13       | Coral-rich   |

*The selected taxa of scleractinian coral used to generate all coral population indicators for the Northern Mariana Islands.*

|                                     |
|-------------------------------------|
| <b>Coral-rich habitat structure</b> |
| Genera                              |
| <i>Acropora</i>                     |
| <i>Astreopora</i>                   |
| <i>Cyphastrea</i>                   |
| <i>Favia</i>                        |
| <i>Goniastrea</i>                   |
| <i>Leptastrea</i>                   |
| <i>Montipora</i>                    |
| <i>Pavona</i>                       |
| <i>Pocillopora</i>                  |
| <i>Porites</i>                      |
|                                     |
| <b>Mixed habitat structure</b>      |
| Genera                              |
| <i>Astreopora</i>                   |
| <i>Cyphastrea</i>                   |
| <i>Favia</i>                        |
| <i>Galaxea</i>                      |
| <i>Goniastrea</i>                   |
| <i>Leptastrea</i>                   |
| <i>Montipora</i>                    |
| <i>Pavona</i>                       |
| <i>Pocillopora</i>                  |
| <i>Porites</i>                      |
|                                     |
| <b>Complex habitat structure</b>    |
| Genera                              |
| <i>Astreopora</i>                   |
| <i>Cyphastrea</i>                   |
| <i>Favia</i>                        |
| <i>Goniastrea</i>                   |
| <i>Leptastrea</i>                   |
| <i>Montipora</i>                    |
| <i>Pavona</i>                       |
| <i>Pocillopora</i>                  |
| <i>Porites</i>                      |

Throughout the literature, there are no clear benchmarks to establish reference and assessment points for density for juvenile and adult corals due to the natural variability in abundance across habitat space. The island level scores for adults and juveniles we calculated similar to the percent cover using the following steps:

1. Reef area-weighted island and sector density estimates were generated from strata means for each of the coral taxa listed above when stratified random surveys were conducted.
2. The score was calculated as a percent of the maximum island or sector density as follows:
  - a. Taxon density % max score = (taxon A density estimate for island A / maximum island or sector density estimate for habitat A reported for all years for taxon A) \* 100.
  - b. The scores were averaged across all taxa for a given island or sector.
  - c. The scores were then converted to a 0-100% scale using equations in the table below.
  - d. These scores were generated separately for juveniles and adults.

*Coral population density assessment points and associated scoring. The score is determined using an equation that relates the percent of maximum density to a percent score.*

| <b>Coral population (colonies per square meter)</b> | <b>Score (%)</b> | <b>Equation</b> |
|-----------------------------------------------------|------------------|-----------------|
| 60-100                                              | 90-100           | $Y=0.25x+75$    |
| 40-<60                                              | 80-<90           | $Y=0.50x+60$    |
| 20-<40                                              | 70-<80           | $Y=0.50x+60$    |
| 10-<20                                              | 60-<70           | $Y=x+50$        |
| 0-<10                                               | 0-<60            | $Y=6x$          |

### *Partial mortality*

The partial mortality indicator was generated from the same select list of abundant and important coral species and genera used for juvenile and adult density.

Throughout the literature, there are not clear benchmarks to establish reference and assessment points for density for partial mortality due to the natural variability in abundance across habitat space. The island level scores for partial mortality we calculated similar to the other benthic metrics as follows:

1. Reef area-weighted island and sector density estimates were generated from strata means for each of coral taxa listed above when stratified random surveys were conducted.
2. The score was calculated as a percent of the maximum island or sector density as follows:
  - a. Taxon density % max score = (taxon A density estimate for island A / maximum island or sector density estimate for habitat A reported for all years for taxon A) \* 100.
  - b. The scores were averaged across all taxa for a given island or sector.
  - c. The scores were then converted to a 60-100% on an inverted scale using equations below.

*Coral mortality assessment points and associated scoring. The score is determined using an equation that relates the percent of maximum mortality to a percent score. For mortality, this is a negative relationship (more mortality receives a worse score).*


| % of maximum | Score (%) | Equation          |
|--------------|-----------|-------------------|
| 0-20         | 90-100    | $Y = -0.5x + 100$ |
| 20-<40       | 80-<90    | $Y = -0.5x + 100$ |
| 40-<60       | 70-<80    | $Y = -0.5x + 100$ |
| 60-<80       | 60-<70    | $Y = -0.5x + 100$ |
| 80-100       | 0-<60     | $Y = -3x + 300$   |

## Fish indicators


The indicators for the fish theme are reef fish, predators, sustainability, and diversity. Field sampling included stationary point counts (SPC) and towed diver surveys. SPC surveys were located using a stratified-random survey design with the survey domain being all hardbottom forereef habitat 0-30 m. Towed diver surveys were conducted in ~10-20m deep forereef areas, with tows located haphazardly around islands with a goal of broadly spreading them around the island circumference. All fish sampling sites were co-located with benthic cover sampling.

Sharks and other predators are averaged together before averaged with other fish indicators. The overall fish score consists of the average of reef fish biomass, average of sharks + other predators, and sustainability.

*Stationary point count collected data are used in the reef fish biomass, sustainability and herbivory indicators:*




Towed diver survey data are used in the predator indicator:



## Reef fish biomass

The Reef fish indicator is a measure of the amount of fish present, specifically the weight of fish per unit area (i.e. ‘biomass’). Total reef fish biomass is probably the most widely used measure for reef fish condition by research and monitoring scientists, and has been repeatedly shown to be responsive to human impacts such as fishing and protection. For the status report, the status of each region was scored based on a study published in 2015 that used data from these islands and >30 others in the US Pacific, and which assessed the importance of a range of human, oceanographic (e.g. temperature, oceanic productivity) and habitat drivers (e.g. coral cover, habitat structural complexity). That study estimated what ‘baseline biomass’ would be at each location in the absence of humans – i.e. what would be natural for each location, based entirely on its oceanographic and habitat setting (Williams *et al* PLoS One, 2015). Some results from that study are shown in the figure below.

### Modeled Biomass Baselines



Reef fish biomass from surveys and modeled ‘baseline biomass’ for each of 40 islands surveyed by Pacific National Coral Reef Monitoring Program between 2010 and 2013. Percentages over American Samoa islands represent the difference between survey mean and modeled baseline biomass at each island in that group. This figure and results taken from Williams *et al* PLoS ONE 2015.


The indicator is measured as the total instantaneous reef fish biomass from the SPC surveys as a proportion of modeled 'baseline' biomass per island (from *Williams et al PLoS One 2015*). In American Samoa those ranged from 44% at Tutuila, i.e. that actual biomass of reef fishes around Tutuila was ~44% of what it would be in the absence of human impacts, to 80% at Swains. Sharks and jacks were excluded for the biomass estimates and the baseline modeling exercise because their biomass is generally not well sampled by small-scale surveys such as the SPC. As a general rule, small-area surveys tend to overestimate densities of such highly mobile species and are prone to bias due to different behavioral responses of those species to divers in different locations (Ward et al 2010, Parrish et al. 2000).

Reef fish biomass as a proportion of baseline biomass was converted to an indicator score on a 0-100% scale, with a conversion function and break points determined by expert opinion. It is becoming widely recognized that even small human populations tend to have quite significant impacts on local reef fish biomass, and that fish biomass around human-populated islands is typically <50% and not uncommonly <25% of pristine biomass (Williams et al. 2015, MacNeil et al. 2015). Thus, the scoring system was created to reflect that understanding – specifically that (i) reef fish biomass > 80% of pristine biomass was about the highest that would be likely to occur around any populated island, and that (ii) even biomass of around 1/3<sup>rd</sup> of baseline is not uncommon for populated islands. Specifically, reef fish biomass > 80% of baseline was considered 'very good' (score 90%+), with the expectation that would only occur at uninhabited island, and a reef fish biomass level of 33% of baseline would be considered 'fair' (score=70%). Most populated islands in the Pacific would score 'poor' (score=60-70%) or 'fair' in this system, but American Samoa tends to score higher than islands in other populated regions.

## *Predators*

The predators indicator is a measure of the status of upper trophic level fishes – i.e. fishes that primarily eat other fishes. Predators are considered both ecologically important, and also a suitable indicator of human impacts, because large predators tend to be preferentially targeted and highly vulnerable to human impacts. The predators' indicator was separated into two sub-indicators: shark abundance and biomass of other predators. Those were broken out separately for 2 main reasons: (i) they are better surveyed by different methods – sharks by towed diver surveys, 'other predators' by SPC; and (ii) because reef sharks recently became protected in American Samoa, and as such tracking the status of reef shark populations separately is of management interest. Drawbacks to using shark abundance as an indicator include their tendency to avoid divers around populated areas and that, generally, large portions of their populations are in waters that are deeper than can be surveyed by divers on SCUBA. Nevertheless towed diver abundance is a useful measure of relative abundance, and is one that we expect to respond well to changes in their populations.

As with reef fish, sharks abundance is scored by first comparing the density from surveys with a model generated estimate of baseline abundance in the absence of humans (Nadon et al 2012). And the converting proportion of baseline values into scores using a scoring function.



For this status report, we modified the approach used by Nadon et al (2012) by including only sharks estimated to be 1m or larger in counts, and adjusting baseline estimates to remove the average number of sharks smaller than that size in the complete data set (i.e. 12% of all encounters). Small (< 1m long) sharks were excluded because divers occasionally encounter large schools of juvenile gray reef sharks (as occurred at Swains Island in 2012). Not only do those encounters introduce a lot of variability into the raw count data, they are also likely not to be representative of the long-term shark populations.

'Other predator' biomass was generated from biomass of all predators other than sharks and jacks in SPC surveys, and compared against baseline biomass estimates generated in the same way as for the reef fish indicator, and documented in Williams et al (2015). Biomass of 'other predators' primarily comes from snapper, grouper, and barracuda species, but includes other predatory emperors, moray eels, goatfish, wrasse, mackerel and others.

## Sustainability

Sustainability indicates whether the targeted fish stocks have large numbers of mature individuals and particularly of large mature individuals. Therefore this indicator is intended to represent a measure of local stocks' ability to reproduce themselves. This indicator is sensitive to fishing pressure, as size and number of large individuals both respond to human impacts such as fishing, and management e.g. fishery limits or closed areas. Additionally, the maintenance of sufficient breeding stock is a core fisheries and conservation goal.

The following methods used for Sustainability were developed after the completion of the American Samoa status report. These methods are used for the Hawaiian Archipelago, Pacific Remote Islands, Guam, and the Northern Mariana Islands status reports produced and released in 2018. For methods for American Samoa, see below. In the future, all iterations of all status report will use these methods, including American Samoa.

## *Defining ‘Generally Targeted Species’*

In order to develop a method that was suitable across all regions, it was necessary to come up with broad definitions of potential target species. Specifically, not all species are present in all regions, and even where they are present, there are differences in fishing preference among regions – in fact, for the Pacific Remote Islands, there is little or no fishing of reef fishes. Therefore, for the purposes of this indicator, target species were defined as all large-bodied species (maximum length greater than 40cm) of the following, commonly-targeted, families: Jack, Surgeon, Parrot, Emperor, Grouper, Snapper, Goatfishes, and Squirrel/Soldierfish (Carangidae, Acanthuridae, Scaridae, Lethrinidae, Serranidae, Lutjanidae, Mullidae, and Holocentridae). Following discussions with local resource managers, a number of those species were excluded, because they are not generally preferred targets or because visual survey data are likely to poorly represent their real size distributions. Those species were: the jacks *Trachinotus* spp, *Seriola* spp, and *Elagatis bipinnulata*, the surgeonfish *Naso brevirostris*, *N. tonganus*, and *N. vlamingii*; and the snappers *Lutjanus bohar* and *L. fulvus*.

### Calculating regionally-appropriate sizes-at-maturity per species

Given the lack of information on size at maturity for most species at most regions, we used a standardized approach to generate size at maturity (L50) per species per region – L50 being the size at which 50% of individuals are expected to be sexually mature. First, we determined the maximum observed size (from survey data) by species within each region (*regional LMax*). Because the Pacific Remote Islands are an administrative and not a geographic region, Wake was pooled with the Marianas and Johnson with Hawai’i for this assessment. The remaining Pacific Remote Islands make up the US Line & Phoenix Islands, which was considered a reasonable biogeographic grouping. Species with fewer than 20 observations within a region were dropped from that region on the basis that the *regional LMax* could not be reliably estimated. The *regional L50* was derived from a published empirical relationship for teleost fishes:

$\log_{10}(L50) = -0.1189 + 0.9157 * \log_{10}(Lmax)$  [Binholland & Froese J Appl Ichthyology 2009].


### Calculating interim sustainability metrics by species, family, and status report unit

To generate interim sustainability metrics per species, we first calculated the mean size of fishes of that species observed in each region; first dropping fishes smaller than 40% of their *regional LMax* – those were considered likely to be juvenile fishes. That size cut off is intended to ensure that the measured stocks are of fishable size classes and to prevent large recruitment pulses from bringing down the mean size (e.g., when there are many young-of-year fishes). Species for which fewer than 5 individuals were observed in a reporting unit were dropped on the basis that a reliable mean size could not be generated for that location. For each species and reporting unit, mean-size was divided by the *regional L50* for that species, to generate a metric that represented mean size relative to size at maturity for that species in that location. Those values were averaged within families – so that each family would be weighted equally – and the overall sustainability metric for each location was generated by averaging the metrics for each of the families at that location. Finally, we cubed those values in order to convert what was essentially a length-based metric to a volume-based value. Fishes are 3 dimensional objects – thus, for

example, doubling their size results in an approximately 8-fold increase in volume and therefore biomass, and likely a similar increase in their reproductive output.

### Converting sustainability metric per location into a 'sustainability score'

The sustainability metric described above is essentially a derived measure of mean volume of a large number of fish species at a location relative to their sizes at maturity. High scores indicate that mean sizes of those fishes are generally large relative to the size at which they will start contributing to the regional breeding pool – and higher values therefore represent higher likely population fecundity. To convert that to a score that is applicable across all locations and comparable with other status report scores (which are scored as percentages), we first developed a standard for what we would expect from largely un-fished stocks. Specifically, we calculated the average of the sustainability metrics from NCRMP locations where harvesting of reef fishes is likely to be negligible, either because of their remoteness or well enforced prohibition of fishing: The Northwestern Hawaiian Islands, Jarvis, Kingman & Palmyra, Howland & Baker, Mariana Trench Marine National Monument, and Northern Mariana Unpopulated Islands. We considered that value as a benchmark for relatively unfished reef fish stocks, and generated the sustainability score for each region by dividing the sustainability metric by that remote-location benchmark. Those scores were capped at 100%.



## Diversity

Biodiversity is the complex variety of life within our ecosystems that forms the foundation for the vast array of ecosystem services that contribute to human well-being. Each of the many 1000s of species occupying coral reefs provide essential ecological functions that maintain ecosystem health. Since different related species often play similar and redundant functional roles but have levels of sensitivity to certain stressors, reefs with higher levels of biodiversity tend to have increased functional redundancy and likelihood of reef survival during episodic disturbance events, such as disease outbreaks or coral bleaching. Hence, biodiversity serves as a useful indicator of ecological resilience and the ability to survive or recover from disturbance events. At the time of the status report completion, we were not yet able to come to consensus on appropriate biodiversity thresholds that relate meaningfully to scores for ecosystem health. In future status reports, however, biodiversity will be an indicator that is scored and incorporated into the overall scores for each jurisdiction.

## American Samoa

Field sampling included stationary point counts (SPC) at 765 sites and 226 towed diver surveys. For this status report, data are from 2010 to 2015.


*# of SPC sites and towed-diver surveys per Reporting Group*

| Reporting Unit | # SPC Sites | # Fish Towed Diver Surveys |
|----------------|-------------|----------------------------|
| Ofu & Olosega  | 112         | 34                         |
| Rose           | 94          | 23                         |
| Swains         | 94          | 25                         |
| Tau            | 92          | 39                         |
| Tutuila North  | 166         | 51                         |
| Tutuila South  | 207         | 54                         |

*Reef fish biomass assessment points and associated scoring. The score is determined using an equation that relates reef fish biomass percent of modeled baseline to a percent score. The graph below is a visual representation of the equations used to generate the scores.*

| % of Modeled Biomass | Score (%)     | Equation                            |
|----------------------|---------------|-------------------------------------|
| < 33.33%             | 0 – 60%       | $y = x * (70/33.33)$                |
| > 33.33%             | 60 - 100% (1) | $y = 70 + ((x-33.33) * (80-33.33))$ |


Notes: (1) Scores capped at 100%



*Other predators assessment points and associated scoring. The score is determined using an equation that relates predator percent of modeled baseline to a percent score. The graph below is a visual representation of the equations used to generate the scores.*

| % of Modeled Biomass | Score (%)     | Equation                        |
|----------------------|---------------|---------------------------------|
| < 10 %               | 0 – 60%       | $y = 6 * x$                     |
| > 10%                | 60 - 100% (1) | $y = 60 + ((x - 10) * (40/80))$ |


Notes: (1) Scores capped at 100%



*Shark abundance assessment points and associated scoring. The score is determined using an equation that relates shark abundance percent of modeled baseline to a percent score. The graph below is a visual representation of the equations used to generate the scores. Shark abundance or 'other predator' biomass from surveys are represented as a proportion of baseline abundance or biomass for each island ( $x$ =axis), and those are converted into a score for this indicator ( $y$ -axis) using the functions shown by the red dashed line.*

| % of Modeled Density | Score (%)     | Equation                            |
|----------------------|---------------|-------------------------------------|
| < 2 %                | 40 – 60% (1)  | $y = 30 \cdot x$                    |
| 2 – 10 %             | 60 - 70%      | $y = 60 + ((x - 2) \cdot (10/8))$   |
| > 10%                | 70 - 100% (2) | $y = 70 + ((x - 10) \cdot (30/50))$ |

Notes: (1) Scores minimum set to 40% in recognition of fact that sharks are present in all locations, but difficult to survey by divers when they are not abundant and because large portions of their populations may be in waters deeper than can readily be surveyed by divers on SCUBA. (2) Scores capped at 100%



### *Sustainability*

While there are different potential methods for evaluating sustainability, the method applied in American Samoa was the best available to date at the time of completion. Specifically, for each of the 25 most commonly captured reef species recorded by the Tutuila biosampling program, we estimate the mean size of that species of all fishes 15cm and above, that were larger than 30% of the species' maximum size. The size cut off is intended to ensure that the measured stocks are of fishable size classes and to prevent large recruitment pulses from bringing down the mean size (e.g., when there are many young of the year). The mean size of each species was then divided by the size at maturity of that species (L50), i.e., the size at which 50% of individuals are expected to be sexually mature.

L50 was derived from a published empirical relationships for teleost fishes:  $\log_{10}(L50) = -0.1189 + 0.9157 * \log_{10}(Lmax)$  [Binholand & Froese J Appl Ichthyology 2009]. That value for each species (mean size divided by L50) was cubed to represent biomass rather than size, as biomass better represent fecundity. Finally, the value was averaged across all 25 species for each reporting unit.

*Top 25 species in catch recorded by American Samoa Biosampling Program. Note that all samples are of fishes caught around Tutuila:*

|                              |
|------------------------------|
| <b>Surgeonfishes</b>         |
| <i>Acanthurus guttatus</i>   |
| <i>Acanthurus lineatus</i>   |
| <i>Acanthurus nigricans</i>  |
| <i>Ctenochaetus striatus</i> |
| <i>Naso lituratus</i>        |
| <i>Naso unicornis</i>        |

|                              |
|------------------------------|
| <b>Parrotfishes</b>          |
| <i>Chlorurus japanensis</i>  |
| <i>Scarus frenatus</i>       |
| <i>Scarus globiceps</i>      |
| <i>Scarus oviceps</i>        |
| <i>Scarus rubroviolaceus</i> |

|                         |
|-------------------------|
| <b>Snapper</b>          |
| <i>Lutjanus gibbus</i>  |
| <i>Lutjanus kasmira</i> |

|                                   |
|-----------------------------------|
| <b>Emperor</b>                    |
| <i>Lethrinus rubrioperculatus</i> |
| <i>Lethrinus xanthochilus</i>     |

|                                 |
|---------------------------------|
| <b>Grouper</b>                  |
| <i>Cephalopholis argus</i>      |
| <i>Epinephelus melanostigma</i> |
| <i>Variola albimarginata</i>    |


|                             |
|-----------------------------|
| <b>Goatfishes</b>           |
| <i>Parupeneus insularis</i> |

A Pacific-wide comparison was not used to generate the scoring function because different regions have species assemblages and different target species preferences. Therefore, the scoring function used for American Samoa was generated subjectively.

*Sustainability assessment points and associated scoring. The score is determined using an equation that relates sustainability to a percent score. The graph below is a visual representation of the equations used to generate the scores. The x-axis represents a measure of mean size relative to L50 for 25 targeted species in Samoa. This is converted into a score for this indicator (y-axis) using the function shown by the blue line above. American Samoan islands are shown as black circles.*

| % of Modeled Density | Score (%)     | Equation                        |
|----------------------|---------------|---------------------------------|
| > 10%                | 65 - 100% (1) | $y = 65 + ((x - 70) * (35/80))$ |

Notes: (1) Scores set to range between 65% and 100%




### *Hawaiian Archipelago*

Field sampling included stationary point counts (SPC) at 1,850 sites and 334 towed-diver surveys for Hawaii. For this status report, fish data from 2010 to 2017 was used.



*# of SPC sites and towed-diver surveys per Reporting Group*

| Reporting Unit                | # SPC Sites | # Fish Towed Diver Surveys |
|-------------------------------|-------------|----------------------------|
| Northwestern Hawaiian Islands | 669         | 134                        |
| Hawaii                        | 257         | 58                         |
| Kauai-Niihau                  | 216         | 46                         |
| Maui-nui                      | 481         | 82                         |
| Oahu                          | 227         | 14                         |


## Reef fish biomass



## Predators



## Sustainability




## Pacific Remote Islands

For this status report, fish data from 2010 to 2017 was used.



# of SPC sites and towed-diver surveys per Reporting Group.

| Reporting Unit    | # SPC Sites | # Fish Towed Diver Surveys |
|-------------------|-------------|----------------------------|
| Howland & Baker   | 171         | 51                         |
| Jarvis            | 192         | 25                         |
| Johnston          | 63          | 40                         |
| Kingman & Palmyra | 237         | 108                        |
| Wake              | 128         | 24                         |


## Reef fish biomass



## Predators



## Sustainability




## Guam & Northern Mariana Islands

For these status reports, fish data from 2010 to 2017 was used.



Table # of SPC sites and towed-diver surveys per Reporting Group.

| Reporting Unit                      | # SPC Sites | # Fish Towed Diver Surveys |
|-------------------------------------|-------------|----------------------------|
| Mariana Trench MNM                  | 207         | 34                         |
| NORTHERN MARIANA ISLANDSUnpopulated | 235         | 56                         |
| Saipan-Tinian-Aguian                | 217         | 58                         |
| Rota                                | 80          | 19                         |
| Guam East                           | 79          | 24                         |
| Guam West                           | 106         | 25                         |
| Guam Marine Preserves               | 119         | 12                         |


## Reef fish biomass



## Predators



## Sustainability



## Climate

The indicators for the climate theme are reef material growth, ocean acidification, and temperature stress. Each indicator has its own field methodology, which will be described in each section.

### Reef material growth

The reef growth indicator directly measures the rate that calcium carbonate, i.e. the skeleton of corals and other reef organisms, accumulates in a given environment (Vargas-Angel et al. 2015). Data were collected by using Calcification Accretion Units (CAUS), a set of 5 stacks of 2 PVC plates per site out-planted for 3 years on a given reef. Units were recovered and the net weight of carbonate that accreted over the deployment period was recorded. Data presented here are from fore-reef sites. For jurisdictions with two years of data, reef material growth values for one year were scored (see threshold table below) and then averaged into an annual reef material growth score. Reef material growth values for the second year were scored (see threshold table below) and then averaged into an annual reef material growth score. The two scores were then averaged into an overall reef material growth score for the jurisdiction. For example, in American Samoa, reef material growth values for 41 sites in 2012 were

scored and then averaged into an annual 2012 reef material growth score. Reef material growth values for 39 sites in 2015 were scored and then averaged into an annual 2015 reef material growth score. The 2012 and 2015 scores were then averaged into an overall American Samoa reef material growth score.

In American Samoa, 41 and 39 sites were averaged from 2012 and 2015, respectively. In the Main Hawaiian Islands and Northwestern Hawaiian Islands, 29 and 19 sites, respectively, were averaged from 2016. For the Northern Mariana Islands, Guam, and Wake Island, 37 and 41 sites were averaged from 2014 and 2017, respectively. For the Pacific Remote Islands, with the exception of Wake Island, 35 sites were averaged from 2015.

*Years of collection and number of sites for each jurisdiction for reef material growth.*

| Jurisdiction                                    | Years      | # of sites |
|-------------------------------------------------|------------|------------|
| American Samoa                                  | 2012, 2015 | 41, 39     |
| Main Hawaiian Islands                           | 2016       | 29         |
| Northwestern Hawaiian Islands                   | 2016       | 19         |
| Northern Mariana Islands, Guam, and Wake Island | 2014, 2017 | 37, 41     |
| Pacific Remote Islands except Wake Island       | 2015       | 35         |

Thresholds were guided by CAU data from throughout all Pacific jurisdictions, with specific thresholds coming from the log-transformed quantiles of Pacific CAU accretion rates. Confidence in using this as an indicator of climate effects on coral reefs is moderate, as we have a broad spatial database of similar metrics. However, this metric does not directly target coral accretion rates, but rather the more OA sensitive crustose coralline algae.

*Reef material growth assessment points and associated scoring. The score is determined using an equation that relates reef material growth to a percent score.*

| RMG_Rate   | Score | Justification                                                                                                                                |
|------------|-------|----------------------------------------------------------------------------------------------------------------------------------------------|
| 0.22615648 | 100   | These breaks are based on the log distribution of RMG_Rate.<br>$\text{Mean}(\text{Log}(\text{RMG}) + 2 * \text{sd}(\text{Log}(\text{RMG})))$ |
| 0.10991417 | 90    | $\text{Mean}(\text{Log}(\text{RMG}) + \text{sd}(\text{Log}(\text{RMG})))$                                                                    |
| 0.05341932 | 80    | $\text{Mean}(\text{Log}(\text{RMG}))$                                                                                                        |
| 0.02596229 | 70    | $\text{Mean}(\text{Log}(\text{RMG}) - \text{sd}(\text{Log}(\text{RMG})))$                                                                    |
| 0.01261792 | 60    | $\text{Mean}(\text{Log}(\text{RMG}) - 2 * \text{sd}(\text{Log}(\text{RMG})))$                                                                |
| 0.0001     | 1     | 0; Less than 0.012 = functional zero.                                                                                                        |

## Ocean acidification

The ocean acidification indicator measures if the water chemistry is suitable for the growth of corals and other reef calcifiers. Aragonite is a form of calcium carbonate that is used by corals to build the hard

parts of the reef. The aragonite saturation state, which is unitless, measures the carbonate ion concentration in the water column. Data were collected by standard seawater sampling for carbonate measurements, using diver deployed Niskin bottle sampling at the surface immediately above focal coral reefs. For jurisdictions with two years of data, ocean acidification values for one year were scored (see threshold table below) and then averaged into an annual ocean acidification score. Ocean acidification values for the second year were scored (see threshold table below) and then averaged into an annual ocean acidification score. The two scores were then averaged into an overall ocean acidification score for that jurisdiction. For the current status reports, this only applies to American Samoa, where there is data from 2012 and 2015.

In American Samoa, 45 and 85 sites were averaged from 2012 and 2015, respectively. In the Main Hawaiian Islands and Northwestern Hawaiian Islands, 87 and 55 sites, respectively, were averaged from 2016. For the Northern Mariana Islands, Guam, and Wake Island, 106 sites were averaged from 2014. For the Pacific Remote Islands, with the exception of Wake Island, 73 sites were averaged from 2015.

*Years of collection and number of sites for each jurisdiction for ocean acidification.*

| Jurisdiction                                    | Years      | # of sites |
|-------------------------------------------------|------------|------------|
| American Samoa                                  | 2012, 2015 | 45, 85     |
| Main Hawaiian Islands                           | 2016       | 87         |
| Northwestern Hawaiian Islands                   | 2016       | 55         |
| Northern Mariana Islands, Guam, and Wake Island | 2014       | 106        |
| Pacific Remote Islands except Wake Island       | 2015       | 73         |

The thresholds for this indicator were determined by using ecologically and climatologically relevant thresholds (see table below) and were the same for both pilot areas. Past and future conditions  $\Omega_{\text{arag}}$  estimated by keeping total alkalinity, salinity, and water temperature constant while altering the partial pressure of carbon dioxide. One annual mean Aragonite saturation value in each year was determined for each region from multiple sample values. That one mean Aragonite saturation value was compared against the threshold and scored. The scores for multiple years were then averaged for a regional score.

*Ocean acidification assessment points and associated scoring. The score is determined using an equation that relates ocean acidification to a percent score.*

| Aragonite saturation | Narrative             | Score | Equation       |
|----------------------|-----------------------|-------|----------------|
| $\geq 4.6$           | Pre-industrial        | 100   | $Y=100$        |
| $4.28 - <4.6$        |                       | 90    | $Y=30.8x-41.5$ |
| $3.95 - <4.28$       |                       | 80    | $Y=30.8x-41.5$ |
| $3.63 - <3.95$       |                       | 70    | $Y=30.8x-41.5$ |
| $3.3 - <3.63$        |                       | 60    | $Y=30.8x-41.5$ |
| $<3.3$               | Double pre-industrial | 0     | $Y=200x-600$   |

## Temperature stress

The temperature stress indicator grades corals' health based on the occurrence and severity of coral bleaching thermal stress they have experienced during the 4-year evaluation period. Mass coral bleaching due to anomalously warm water temperatures has occurred with increasing frequency and severity in recent decades and is now the most significant single contributor to the decline of coral reef ecosystems on a global scale. Coral mortality and disease outbreak often follow massive bleaching events, along with significantly reduced coral growth rate both during and after the bleaching and ability to fight off other stresses. NOAA Coral Reef Watch (CRW) has been using NOAA's operational near-real-time satellite sea surface temperature data to detect and monitor thermal stress conducive to mass coral bleaching globally since 1997 (Liu et al., 2013, 2014, 2017). Monitoring data for the status report target regions were extracted from CRW's global products and then statistical analysis was performed on the data to generate a grade. Data analysis follows these steps:

1. Determine the reporting period based on available satellite data. Scoring will be based on degree heating week event frequency and severity per 4-year period.
2. Define all the reef-containing data pixels for each area of interest and extract the daily time series of degree heating week values. For multi-pixel areas with 10 or more data pixels, the 90<sup>th</sup> percentile degree heating week (DHW) value is chosen for each time step in the series. For multi-pixel areas with less than 10 data pixels, the maximum degree heating week value is used for each time step.
3. Take the maximum degree heating week value for each year in the 4-year time series.
4. Use these 4 values along with the grading chart and find the corresponding score based on the frequency and severity values in the chart. The value resulting in the lowest score becomes the overall grade for that reporting period.

The scoring chart ranks thermal stress severity in 7 bins based on DHW ranges shown across the top of the chart. The frequency of events at these levels are shown below each bin and have varying distribution based on the relationship between DHW and coral bleaching and mortality. The corresponding score and an interpretation of what that score means are shown on the left hand side of the chart. Counting the number of times a DHW level is reached in a 4-year period and matching it to the proper severity column and frequency row will result in the corresponding score. The chart also takes into account consecutive years of high DHW events, which would have a greater impact on corals than non-consecutive events. These have their own frequency label of "2c". For example, during the period 2013-2016, if 2013 and 2015 saw DHW values of 9, the resulting grade would be 55%, but if 2013 and 2014 saw DHW values of 9 the resulting grade would be 45%.

**Coral Reef Watch NCRMP Status report Scoring Table for a Four-Year Evaluation Period**

| %Bleached | %Dead | Score | 0<N<4 | 4≤N<8 | 8≤N<12 | 12≤N<16 | 16≤N<20 | 20≤N<32 | 32≤N |
|-----------|-------|-------|-------|-------|--------|---------|---------|---------|------|
| <1%       |       | 100%  | 0     | 0     | 0      | 0       | 0       | 0       |      |
| 1%        |       | 95%   | 1-2   |       |        |         |         |         |      |
| 10%       |       | 85%   | 3-4   | 1     |        |         |         |         |      |
| 20%       |       | 75%   |       | 2     |        |         |         |         |      |
| 40%       |       | 65%   |       | 3     | 1      |         |         |         |      |
| 60%       | 10%   | 55%   |       | 4     | 2      | 1       |         |         |      |
| 80%       | 20%   | 45%   |       |       | 2c     | 2       | 1       |         |      |
| 90%       | 40%   | 35%   |       |       | 3      | 2c      | 2       |         |      |
| 100%      | 60%   | 25%   |       |       | 4      | 3       | 2c      | 1       |      |
|           | 80%   | 15%   |       |       |        | 4       | 3       |         |      |
|           | 90%   | 5%    |       |       |        |         | 4       | 2       |      |
|           | 100%  | 0%    |       |       |        |         |         | 3-4     | 1    |

**Key**



DHW Severity Ranges (N = DHW value)



Number of DHW occurrences in 4-Year period (2c = 2 years are consecutive)



Corresponding Score and Expected Impact on Corals

## Human connections

The human connections theme of the NCRMP gathers and monitors a collection of socioeconomic variables, such as demographics in coral reef areas, human use of coral reef resources, as well as knowledge, attitudes, and perceptions of coral reefs and coral reef management. The overall goal of the human connections monitoring component is to track relevant information regarding each jurisdiction's population, social and economic structure, and interactions between coral reef ecosystems and adjacent human communities. The selection of indicators was determined through workshops and consultations with partners (local jurisdictions as well as federal agencies). The human connections indicators are: awareness, participation in pro-environmental behaviors, and support for management actions. To operationalize these three indicators, secondary data collected from the jurisdiction and data collected from an NCRMP survey of jurisdictional residents are used. Threshold goals for these indicators were established through consultation with coral reef managers, environmental education and outreach coordinators, and relevant federal, state, and local agency staff.

Scoring scale for Human Connections indicators

| Percentage calculated by dividing indicator score by the established threshold | Score     |
|--------------------------------------------------------------------------------|-----------|
| 90%+                                                                           | Very good |
| 80%-89.99%                                                                     | Good      |
| 70%-79.99%                                                                     | Fair      |
| 60%-69.99%                                                                     | Impaired  |
| <60%                                                                           | Critical  |

## American Samoa


All indicators are evaluated at the jurisdictional level, not at the reporting region level. Due to resource constraints, the NCRMP team was not able to conduct a survey of residents of the less populated Manu'a Islands, although surveys in these locations are planned for future survey iterations. Thus, the indicators for awareness, support for management actions, and participation in pro-environmental behaviors reflect the most populated island of Tutuila. The score is the average of the three overall indicators – Awareness, Support for management actions, and Pro-environmental behavior.

### Awareness

The Awareness indicator is an indicator of residents' familiarity with threats to and the importance of reefs. Three awareness metrics, obtained from the NCRMP jurisdictional resident survey, were averaged into an overall Awareness indicator score. The three metrics for American Samoa are: familiarity with threats to coral reefs; familiarity with Marine Protected Areas (MPAs); and, the value or importance respondents place on coral reefs.

### *Familiarity with threats to coral reefs*

Survey respondents in American Samoa were asked to rate their familiarity with various threats posed to coral reefs on a scale of “very unfamiliar” to “very familiar.” Familiarity with threats indicates local awareness of the need for management action. The percentage of respondents that were at least “familiar” was calculated for each of the nine threats that were proposed in the American Samoa survey. A threshold of two-thirds was established (i.e., a goal that at least two-thirds of respondents were at least “familiar” with the threat). Coral reef managers in American Samoa confirmed that this goal was appropriate (i.e., professional judgement).




### *Familiarity with Marine Protected Areas*

Survey respondents in American Samoa were asked to rate their familiarity with MPAs on a scale of “very unfamiliar” to “very familiar.” Familiarity with MPAs indicates success of jurisdictional education and outreach campaigns and understanding of marine regulations. The percentage of respondents that were at least “familiar” with MPAs was calculated. A threshold of two-thirds was established (i.e. a goal that at least two-thirds of respondents were at least “familiar” with MPAs). Coral reef managers in American Samoa confirmed that this goal was appropriate (i.e., professional judgement).

### *Value or Importance of coral reefs*

The value or importance that respondents in American Samoa place on coral reefs was examined. This section of the survey contained four questions in which statements were proposed and respondents were asked to rate how much they “agree” with the statements on a scale of “strongly disagree” to “strongly agree.” Two of these statements were ambiguous in their interpretation (“Coral reefs are only important to fishermen, divers and snorkelers” and “Healthy coral reefs attract tourists to American Samoa”). Since agreement with these aforementioned statements can either be considered good or

bad, they were not analyzed in this section. However, the other two questions in this section were analyzed (“Coral reefs protect American Samoa from erosion and natural disasters” and “Coral reefs are important to American Samoan cultures”). Agreement with these statements can be interpreted as positive indicators. A threshold of two-thirds in agreement with the protection and culture statements was established after consultation with coral reef managers in American Samoa.




### *Support for management actions*

The support for management indicator measures the level of support that respondents to the NCRMP jurisdictional resident survey indicate for coral reef management activities. The American Samoa survey asks two sets of questions (agreement with various Marine Protected Area functions and support for coral reef management rules and regulations). After consultation with local partners, the threshold for both indicators was set at two-thirds of the respondents in agreement, or supportive.

### *Agreement with marine protected area functions*


The marine protected area question section contained nine statements, and respondents were asked to rate how much they “agree” with the statements on a scale of “strongly disagree” to “strongly agree.” Three of these statements were left out of this analysis due to the ambiguous nature of interpretation (“There should be fewer MPAs in American Samoa,” “Fishermen’s livelihoods have been negatively impacted from the establishment of MPAs in American Samoa,” and “MPAs help increase tourism in American Samoa”). The first two of these aforementioned statements can be considered as negative sentiments toward MPAs, and the third statement does not necessarily indicate agreement with MPA functions. The other statements in this section can all be considered agreement with the intended functions of MPAs (“MPAs protect coral reefs,” “MPAs increase the number of fish,” “There should be more MPAs in American Samoa,” “There has been economic benefit to American Samoa from the establishment of MPAs,” “I would support adding new MPAs in American Samoa if there is evidence that the ones we have are improving American Samoa’s marine resources,” and “I generally support the establishment of MPAs”). Therefore, only these “positive sentiment” statements are analyzed in this

section. A threshold of two-thirds agreement was established for the “positive” statements after consultation with coral reef managers in American Samoa to maintain consistency across the indicators.



### *Support for coral reef management rules and regulations*

Respondents were asked to rate their support for various management initiatives on a scale of “strongly oppose” to “strongly support.” The same threshold goal of two-thirds of survey respondents in support with the proposed management initiatives was set after consultation with coral reef managers.



### *Pro-environmental behavior*

The pro-environmental behavior indicator measures residents' (active) participation in activities, such as beach clean-ups, volunteering with an environmental group, recycling, etc. to protect the environment. NCRMP survey respondents in American Samoa were asked to rate their frequency of participation in any pro-environmental behaviors on a scale of "not at all" to "several times a month or more." Through discussion with coral reef managers, a goal of 70% participation was established (i.e., researchers wanted to observe that at least 70% of respondents participated in any form of pro-environmental behavior and at any frequency).

### **Hawaiian Archipelago**


All indicators are evaluated for the Main Hawaiian Islands, at the jurisdictional level, as well as the reporting region level. Human Connections data were not collected for the Northwest Hawaiian Islands, as they are not populated by people. Thus, the indicators for awareness, support for management actions, and participation in pro-environmental behaviors reflect the Main Hawaiian Islands as a whole, as well as Hawai'i Island, O'ahu, Kaua'i, and Maui Nui.

### *Awareness*

The Awareness indicator is an indicator of residents' familiarity with threats to and the importance of reefs. Three awareness metrics, were averaged into an overall Awareness indicator score. The three metrics for Hawai'i are: familiarity with threats to coral reefs; familiarity with MPAs; and, the value or importance respondents place on coral reefs.

### *Familiarity with threats to coral reefs*

Survey respondents in Hawai'i were asked to rate their familiarity with various threats posed to coral reefs on a scale of "very unfamiliar" to "very familiar." Familiarity with threats indicates local awareness of the need for management action. The percentage of respondents that were at least "familiar" was calculated for each of the ten threats that were proposed in the Hawai'i survey. A threshold of two-thirds was established (i.e., a goal that at least two-thirds of respondents were at least "familiar" with the threat). Coral reef managers in Hawai'i confirmed that this goal was appropriate (i.e., professional judgement).




### *Familiarity with Marine Protected Areas*

Survey respondents in Hawai'i were asked to rate their familiarity with MPAs on a scale of "very unfamiliar" to "very familiar." Familiarity with MPAs indicates success of jurisdictional education and outreach campaigns and understanding of marine regulations. The percentage of respondents that were at least "familiar" with MPAs was calculated. A threshold of two-thirds was established (i.e. a goal that at least two-thirds of respondents were at least "familiar" with MPAs). Coral reef managers in Hawai'i confirmed that this goal was appropriate (i.e., professional judgement).

### *Value or Importance of coral reefs*

The value or importance that respondents in Hawai'i place on coral reefs was examined. This section of the survey contained four questions in which statements were proposed and respondents were asked to rate how much they "agree" with the statements on a scale of "strongly disagree" to "strongly agree." Two of these statements were ambiguous in their interpretation ("Coral reefs are only important to fishermen, divers and snorkelers" and "Healthy coral reefs attract tourists to Hawai'i"). Since agreement with these aforementioned statements can either be considered good or bad, they were not analyzed in this section. However, the other two questions in this section were analyzed ("Coral reefs protect Hawai'i from erosion and natural disasters" and "Coral reefs are important to Hawaiian culture"). Agreement with these statements can be interpreted as positive indicators. A threshold of two-thirds in agreement with the protection and culture statements was established after consultation with coral reef managers in Hawai'i.




### *Support for management actions*

The support for management indicator measures the level of support that respondents to the NCRMP jurisdictional resident survey indicate for coral reef management activities. The Hawai'i survey asks one set of questions related to support for coral reef management rules and regulations. After consultation with local partners, the threshold for the indicator was set at two-thirds of the respondents in support.

### *Support for coral reef management rules and regulations*

Respondents were asked to rate their support for various management initiatives on a scale of "strongly oppose" to "strongly support." The same threshold goal of two-thirds of survey respondents in support with the proposed management initiatives was set after consultation with coral reef managers.



### *Pro-environmental behavior*

The pro-environmental behavior indicator measures residents' (active) participation in activities, such as beach clean-ups, volunteering with an environmental group, recycling, etc. to protect the environment. NCRMP survey respondents in Hawai'i were asked to rate their frequency of participation in any pro-environmental behaviors on a scale of "not at all" to "several times a month or more." Through discussion with coral reef managers, a goal of 70% participation was established (i.e., researchers wanted to observe that at least 70% of respondents participated in any form of pro-environmental behavior and at any frequency).

### **Guam**


All indicators are evaluated at the jurisdictional level, not at the reporting region level. Due to resource constraints, the NCRMP team was not able to collect representative samples from Eastern and Western Guam. Thus, the indicators for awareness, support for management actions, and participation in pro-environmental behaviors reflect the status of Guam as a whole.

### *Awareness*

The Awareness indicator is an indicator of residents' familiarity with threats to and the importance of reefs. Three awareness metrics, were averaged into an overall Awareness indicator score. The three metrics for Guam are: familiarity with threats to coral reefs; familiarity with MPAs; and, the value or importance respondents place on coral reefs.

### *Familiarity with threats to coral reefs*

Survey respondents in Guam were asked to rate their familiarity with various threats posed to coral reefs on a scale of “very unfamiliar” to “very familiar.” Familiarity with threats indicates local awareness of the need for management action. The percentage of respondents that were at least “familiar” was calculated for each of the eleven threats that were proposed in the Guam survey. A threshold of two-thirds was established (i.e., a goal that at least two-thirds of respondents were at least “familiar” with the threat). Coral reef managers in Guam confirmed that this goal was appropriate (i.e., professional judgement).




### *Familiarity with Marine Protected Areas*

Survey respondents in Guam were asked to rate their familiarity with MPAs on a scale of “very unfamiliar” to “very familiar.” Familiarity with MPAs indicates success of jurisdictional education and outreach campaigns and understanding of marine regulations. The percentage of respondents that were at least “familiar” with MPAs was calculated. A threshold of two-thirds was established (i.e. a goal that at least two-thirds of respondents were at least “familiar” with MPAs). Coral reef managers in Guam confirmed that this goal was appropriate (i.e., professional judgement).

### *Value or Importance of coral reefs*

The value or importance that respondents in Guam place on coral reefs was examined. This section of the survey contained four questions in which statements were proposed and respondents were asked to rate how much they “agree” with the statements on a scale of “strongly disagree” to “strongly agree.” Two of these statements were ambiguous in their interpretation (“Coral reefs are only important to fishermen, divers and snorkelers” and “Healthy coral reefs attract tourists to Guam”). Since agreement with these aforementioned statements can either be considered good or bad, they were not analyzed in

this section. However, the other two questions in this section were analyzed (“Coral reefs protect Guam from erosion and natural disasters” and “Coral reefs are important to Guam’s culture”). Agreement with these statements can be interpreted as positive indicators. A threshold of two-thirds in agreement with the protection and culture statements was established after consultation with coral reef managers in Guam.




### *Support for management actions*

The support for management indicator measures the level of support that respondents to the NCRMP jurisdictional resident survey indicate for coral reef management activities. The Guam survey asks two sets of questions (agreement with various Marine Protected Area functions and support for coral reef management rules and regulations). After consultation with local partners, the threshold for both indicators was set at two-thirds of the respondents in agreement, or supportive.

### *Agreement with marine protected area functions*


The marine protected area question section contained ten statements, and respondents were asked to rate how much they “agree” with the statements on a scale of “strongly disagree” to “strongly agree.” Four of these statements were left out of this analysis due to the ambiguous nature of interpretation (“There should be fewer MPAs in Guam,” “Fishermen’s livelihoods have been negatively impacted from the establishment of MPAs in Guam,” “The establishment of MPAs increases the likelihood that people will vacation in Guam,” and “MPAs help increase tourism in Guam”). The first two of these aforementioned statements can be considered as negative sentiments toward MPAs, and the last two statements do not necessarily indicate agreement with MPA functions. The other statements in this section can all be considered agreement with the intended functions of MPAs (“MPAs protect coral reefs,” “MPAs increase the number of fish,” “There should be more MPAs in Guam,” “There has been economic benefit to Guam from the establishment of MPAs,” “I would support adding new MPAs in Guam if there is evidence that the ones we have are improving Guam’s marine resources,” and “I generally support the establishment of MPAs”). Therefore, only these “positive sentiment” statements

are analyzed in this section. A threshold of two-thirds agreement was established for the “positive” statements after consultation with coral reef managers in Guam to maintain consistency across the indicators.



### *Support for coral reef management rules and regulations*

Respondents were asked to rate their support for various management initiatives on a scale of “strongly oppose” to “strongly support.” The same threshold goal of two-thirds of survey respondents in support with the proposed management initiatives was set after consultation with coral reef managers.



### *Pro-environmental behavior*

The pro-environmental behavior indicator measures residents' (active) participation in activities, such as beach clean-ups, volunteering with an environmental group, recycling, etc. to protect the environment. NCRMP survey respondents in Guam were asked to rate their frequency of participation in any pro-environmental behaviors on a scale of "not at all" to "several times a month or more." Through discussion with coral reef managers, a goal of two thirds participation was established (i.e., researchers wanted to observe that at least two thirds of respondents participated in any form of pro-environmental behavior and at any frequency).

### **Northern Mariana Islands**


All indicators are evaluated at the jurisdictional level, not at the reporting region level. Due to resource constraints, the NCRMP team was not able collect representative samples from Tinian and Rota, although responses from residents of those islands are still included in the overall figures for the Northern Mariana Islands. Thus, the indicators for awareness, support for management actions, and participation in pro-environmental behaviors reflect the status of the Northern Mariana Islands as a whole.

### *Awareness*

The Awareness indicator is an indicator of residents' familiarity with threats to and the importance of reefs. Three awareness metrics, were averaged into an overall Awareness indicator score. The three metrics for the Northern Mariana Islands are: familiarity with threats to coral reefs; familiarity with MPAs; and, the value or importance respondents place on coral reefs.

### *Familiarity with threats to coral reefs*

Survey respondents in Northern Mariana Islands were asked to rate their familiarity with various threats posed to coral reefs on a scale of “very unfamiliar” to “very familiar.” Familiarity with threats indicates local awareness of the need for management action. The percentage of respondents that were at least “familiar” was calculated for each of the ten threats that were proposed in the Northern Mariana Islands survey. A threshold of two-thirds was established (i.e., a goal that at least two-thirds of respondents were at least “familiar” with the threat). Coral reef managers in Northern Mariana Islands confirmed that this goal was appropriate (i.e., professional judgement).




### *Familiarity with Marine Protected Areas*

Survey respondents in Northern Mariana Islands were asked to rate their familiarity with MPAs on a scale of “very unfamiliar” to “very familiar.” Familiarity with MPAs indicates success of jurisdictional education and outreach campaigns and understanding of marine regulations. The percentage of respondents that were at least “familiar” with MPAs was calculated. A threshold of two-thirds was established (i.e. a goal that at least two-thirds of respondents were at least “familiar” with MPAs). Coral reef managers in Northern Mariana Islands confirmed that this goal was appropriate (i.e., professional judgement).

### *Value or Importance of coral reefs*

The value or importance that respondents in Northern Mariana Islands place on coral reefs was examined. This section of the survey contained four questions in which statements were proposed and respondents were asked to rate how much they “agree” with the statements on a scale of “strongly disagree” to “strongly agree.” One of these statements was negative in terms of its interpretation (“Coral reefs are only important to fishermen, divers and snorkelers”). Since agreement with this aforementioned statement can be considered good or bad, it was not analyzed in this section. However,

the other three questions in this section were analyzed (“Coral reefs protect Northern Mariana Islands from erosion and natural disasters,” “Coral reefs in good condition provide food for island communities to eat,” and “Coral reefs are important to Northern Mariana Island’s culture”). Agreement with these statements can be interpreted as positive indicators. A threshold of two-thirds in agreement with the protection, provisioning, and culture statements was established after consultation with coral reef managers in Northern Mariana Islands.




### *Support for management actions*

The support for management indicator measures the level of support that respondents to the NCRMP jurisdictional resident survey indicate for coral reef management activities. The Northern Mariana Islands survey asks two sets of questions (agreement with various Marine Protected Area functions and support for coral reef management rules and regulations). After consultation with local partners, the threshold for both indicators was set at two-thirds of the respondents in agreement, or supportive.

### *Agreement with marine protected area functions*


The marine protected area question section contained eleven statements, and respondents were asked to rate how much they “agree” with the statements on a scale of “strongly disagree” to “strongly agree.” Three of these statements were left out of this analysis due to the ambiguous nature of interpretation (“There should be fewer MPAs in Northern Mariana Islands,” “Fishermen’s livelihoods have been negatively impacted from the establishment of MPAs in Northern Mariana Islands,” “The establishment of MPAs increases the likelihood that people will vacation in Northern Mariana Islands,” and “MPAs help increase tourism in Northern Mariana Islands”). The first two of these aforementioned statements can be considered as negative sentiments toward MPAs, and the last two statements do not

necessarily indicate agreement with MPA functions. The other statements in this section can all be considered agreement with the intended functions of MPAs (“MPAs protect coral reefs,” “MPAs increase the number of fish,” “There should be more MPAs in Northern Mariana Islands,” “There has been economic benefit to Northern Mariana Islands from the establishment of MPAs,” “I would support adding new MPAs in Northern Mariana Islands if there is evidence that the ones we have are improving Northern Mariana Islands’s marine resources,” “I generally support the establishment of MPAs,” and “I generally support the establishment of the federally managed Marina Trench Marine National Monument”). Therefore, only these “positive sentiment” statements are analyzed in this section. A threshold of two-thirds agreement was established for the “positive” statements after consultation with coral reef managers in Northern Mariana Islands to maintain consistency across the indicators.



### *Support for coral reef management rules and regulations*

Respondents were asked to rate their support for various management initiatives on a scale of “strongly oppose” to “strongly support.” The same threshold goal of two-thirds of survey respondents in support with the proposed management initiatives was set after consultation with coral reef managers.



### *Pro-environmental behavior*

The pro-environmental behavior indicator measures residents' (active) participation in activities, such as beach clean-ups, volunteering with an environmental group, recycling, etc. to protect the environment. NCRMP survey respondents in Northern Mariana Islands were asked to rate their frequency of participation in any pro-environmental behaviors on a scale of "not at all" to "several times a month or more." Through discussion with coral reef managers, a goal of two thirds participation was established (i.e., researchers wanted to observe that at least two thirds of respondents participated in any form of pro-environmental behavior and at any frequency).

## Frequently asked questions

### **Quality assurance and quality control**

All NCRMP data goes through rigorous quality assurance and quality control protocols specific to its data type. References to specific standard operating procedures can be found in each indicator section.

### **Confidence**

While we can report confidence in our current data, it is not as straight-forward to calculate confidence after scores are calculated - therefore scores should be taken as an estimate rather than a precise value. The scores are reported as a percentage as a science communication tool, rather than expressing scientific precision.

## Why did you choose “pristine” or “pre-human” as a reference point/baseline and is it attainable?

Two options were seriously considered during discussions regarding reference/baseline values for each biophysical indicator: 1) pristine/pre-human and 2) a value reflecting an “attainable” state given human presence and impacts. We concluded that although an “attainable” value might be more practical and useful, it was too subjective at this time and the science doesn’t support what that value should be. We were also aware of the concept of shifting baselines and thought there could be value in not losing those historical values, even if it is unlikely to ever be attained again, at least in the foreseeable future.

## Local monitoring data

The data used in this status report effort is restricted to NCRMP data. Localized data, such as those collected by jurisdictional agencies, are not included in the scoring of indicators. This is because blending data that are collected under different sampling designs and/or different sampling methodologies is difficult. In the absence of a targeted calibration exercise that would allow for integrating disparate datasets, we focus here on the NCRMP data that is designed to monitor coral reefs at a jurisdictional scale. It is a goal of future efforts to include this other data in indicator scoring. However, local long-term data products are included when possible as a time-series or highlight story. Highlight stories are meant to message locally relevant information.

## Water quality

Water quality is critical to the condition and resiliency of reefs. While we would ideally like to include water quality indicators such as nutrients and sediments, NCRMP does not collect this information. There is also not spatially reliable water quality data from another source available. If spatially robust water quality data becomes available in the future, we can consider including it.

## V. References

Liu, G, Skirving WJ, Geiger EF, De La Cour JL, Marsh BL, Heron SF, Tirak KV, Strong AE, Eakin CM (2017) NOAA Coral Reef Watch's 5km Satellite Coral Bleaching Heat Stress Monitoring Product Suite Version 3 and Four-Month Outlook Version 4. Reef Encounter 45 32(1): 39-45.

Liu, G, Heron SF, Eakin CM, Muller-Karger FE, Vega-Rodriguez M, Guild LS, De La Cour JL, Geiger EF, Skirving WJ, Burgess TFR, Strong AE, Harris A, Maturi E, Ignatov A, Sapper J, Li J, Lynds S (2014). Reef-scale Thermal Stress Monitoring of Coral Ecosystems: New 5-km Global Products from NOAA Coral Reef Watch. Remote Sensing 6(11): 11579-11606, doi:10.3390/rs61111579.

Liu, G., J.L. Rauenzahn, S.F. Heron, C.M. Eakin, W.J. Skirving, T.R.L. Christensen, A.E. Strong, J. Li (2013). NOAA Coral Reef Watch 50 km Satellite Sea Surface Temperature-Based Decision Support System for Coral Bleaching Management. NOAA Technical Report NESDIS 143. NOAA/NESDIS. College Park, MD. 33pp..

MacNeil MA, NAJ Graham, JE Cinner, SK Wilson, ID Williams, J Maina, S Newman, AM Friedlander, S Jupiter, NVC Polunin, TR McClanahan. 2015. Recovery potential of the world's coral reef fishes. *Nature* 520:341–344.

Nadon MO, JK Baum, ID Williams, JM McPherson, BJ Zgliczynski, BL Richards, RE Schroeder, RE Brainard. 2012. Re-Creating Missing Population Baselines for Pacific Reef Sharks. *Conservation Biology* 26:493–503.

NOAA Coral Program. 2014. National Coral Reef Monitoring Plan. Silver Spring, MD, NOAA Coral Reef Conservation Program.

Parrish FA, MP Craig, TJ Ragen, GJ Marshall, BM Buhleier. 2000. Identifying diurnal foraging habitat of endangered Hawaiian monk seals using a seal-mounted video camera. *Marine Mammal Science* 16:392–412.

Ward-Paige C., J Mills Flemming, HK Lotze. 2010. Overestimating Fish Counts by Non-Instantaneous Visual Censuses: Consequences for Population and Community Descriptions. *PLoS ONE* 5:e11722.

Williams ID, JK Baum, A Heenan, KM Hanson, MO Nadon, RE Brainard. 2015. Human, Oceanographic and Habitat Drivers of Central and Western Pacific Coral Reef Fish Assemblages. *Plos One* 10:e0120516.

## VI. Indicator data citations

### *Coral Reef Cover*

NOAA Pacific Islands Fisheries Science Center, Ecosystem Sciences Division (2018). National Coral Reef Monitoring Program: Benthic cover derived from analysis of images collected during stratified random surveys (StRS) across American Samoa. NOAA National Centers for Environmental Information. Dataset. doi:10.7289/V52V2DFW

NOAA Pacific Islands Fisheries Science Center, Ecosystem Sciences Division (2018). National Coral Reef Monitoring Program: Benthic cover derived from analysis of images collected during stratified random surveys (StRS) of the Hawaiian Archipelago. NOAA National Centers for Environmental Information. Dataset. doi:10.7289/V5JS9NR4

Pacific Islands Fisheries Science Center, 2018: National Coral Reef Monitoring Program: Stratified Random Surveys (StRS) of Reef Fish, including Benthic Estimate Data of the Hawaiian Archipelago since 2013, <https://inport.nmfs.noaa.gov/inport/item/24447>.

Pacific Islands Fisheries Science Center, 2018: Pacific Reef Assessment and Monitoring Program: Stratified Random Surveys (StRS) of Reef Fish, including Benthic Estimate Data at Coral Reef Sites across the Pacific Ocean from 2008 to 2012, <https://inport.nmfs.noaa.gov/inport/item/34515>.

NOAA Pacific Islands Fisheries Science Center, Ecosystem Sciences Division (2018). National Coral Reef Monitoring Program: Benthic cover derived from analysis of images collected during stratified random surveys (StRS) across the Mariana Archipelago. NOAA National Centers for Environmental Information. Dataset. doi:10.7289/V57M0673

NOAA Pacific Islands Fisheries Science Center, Ecosystem Sciences Division (2018). National Coral Reef Monitoring Program: Benthic cover derived from analysis of images collected during stratified random surveys (StRS) across the Pacific Remote Island Areas. NOAA National Centers for Environmental Information. Dataset. doi:10.7289/V5154FBH

### *Coral Populations and Mortality*

NOAA Pacific Islands Fisheries Science Center, Ecosystem Sciences Division (2018). National Coral Reef Monitoring Program: Stratified random surveys (StRS) of coral demography (adult and juvenile corals) across American Samoa. NOAA National Centers for Environmental Information. Dataset. doi:10.7289/V579431K

NOAA Pacific Islands Fisheries Science Center, Ecosystem Sciences Division (2018). National Coral Reef Monitoring Program: Stratified random surveys (StRS) of coral demography (adult and juvenile corals) across the Hawaiian Archipelago. NOAA National Centers for Environmental Information. Dataset. doi:10.7289

Pacific Islands Fisheries Science Center, 2018: National Coral Reef Monitoring Program: Stratified Random Surveys (StRS) of Coral Demography (Adult and Juvenile Corals) across the Hawaiian Archipelago since 2013, <https://inport.nmfs.noaa.gov/inport/item/36164>.

Pacific Islands Fisheries Science Center, 2018: Pacific Reef Assessment and Monitoring Program: Belt Transect for Demographics and Condition of Corals at U.S. Pacific Reefs from 2007 to 2012, <https://inport.nmfs.noaa.gov/inport/item/25260>.

NOAA Pacific Islands Fisheries Science Center, Ecosystem Sciences Division (2018). National Coral Reef Monitoring Program: Stratified random surveys (StRS) of coral demography (adult and juvenile corals) across the Mariana Archipelago. NOAA National Centers for Environmental Information. Dataset. doi:10.7289/V53N21Q5

NOAA Pacific Islands Fisheries Science Center, Ecosystem Sciences Division (2018). National Coral Reef Monitoring Program: Stratified random surveys (StRS) of coral demography (adult and juvenile corals) across the Pacific Remote Island Areas. NOAA National Centers for Environmental Information. Dataset. doi:10.7289/V5ZW1J8B

### *Reef fish, Sustainability and Herbivory*

NOAA Pacific Islands Fisheries Science Center, Ecosystem Sciences Division (2018). National Coral Reef Monitoring Program: Stratified random surveys (StRS) of reef fish, including benthic estimate data of

American Samoa. NOAA National Centers for Environmental Information. Dataset.  
doi:10.7289/V5TB1564

NOAA Pacific Islands Fisheries Science Center, Ecosystem Sciences Division (2018). National Coral Reef Monitoring Program: Stratified random surveys (StRS) of reef fish, including benthic estimate data of the Hawaiian Archipelago. NOAA National Centers for Environmental Information. Dataset.  
doi:10.7289/V59C6VR5

Pacific Islands Fisheries Science Center, 2018: National Coral Reef Monitoring Program: Stratified Random Surveys (StRS) of Reef Fish, including Benthic Estimate Data of the Hawaiian Archipelago since 2013, <https://inport.nmfs.noaa.gov/inport/item/24447>.

Pacific Islands Fisheries Science Center, 2018: Pacific Reef Assessment and Monitoring Program: Stratified Random Surveys (StRS) of Reef Fish, including Benthic Estimate Data at Coral Reef Sites across the Pacific Ocean from 2008 to 2012, <https://inport.nmfs.noaa.gov/inport/item/34515>.

NOAA Pacific Islands Fisheries Science Center, Ecosystem Sciences Division (2018). National Coral Reef Monitoring Program: Stratified random surveys (StRS) of reef fish, including benthic estimate data of the Mariana Archipelago. NOAA National Centers for Environmental Information. Dataset.  
doi:10.7289/V5C827M0

NOAA Pacific Islands Fisheries Science Center, Ecosystem Sciences Division (2018). National Coral Reef Monitoring Program: Stratified random surveys (StRS) of reef fish, including benthic estimate data of the Pacific Remote Island Areas. NOAA National Centers for Environmental Information. Dataset.  
doi:10.7289/V58C9TKB

### *Sharks and other predators*

NOAA Pacific Islands Fisheries Science Center, Ecosystem Sciences Division (2018). National Coral Reef Monitoring Program: Towed-diver surveys of large-bodied fishes of American Samoa. NOAA National Centers for Environmental Information. Dataset. doi:10.7289/V5F769VR

NOAA Pacific Islands Fisheries Science Center, Ecosystem Sciences Division (2018). National Coral Reef Monitoring Program: Towed-diver surveys of large-bodied fishes of the Mariana Archipelago. NOAA National Centers for Environmental Information. Dataset. doi:10.7289/V5X34VR7

NOAA Pacific Islands Fisheries Science Center, Ecosystem Sciences Division (2018). National Coral Reef Monitoring Program: Towed-diver surveys of large-bodied fishes of the Hawaiian Archipelago. NOAA National Centers for Environmental Information. Dataset. doi:10.7289/V55M6417

NOAA Pacific Islands Fisheries Science Center, Ecosystem Sciences Division (2018). National Coral Reef Monitoring Program: Towed-diver surveys of large-bodied fishes of the Pacific Remote Island Areas. NOAA National Centers for Environmental Information. Dataset. doi:10.7289/V5QJ7FM7

Pacific Reef Assessment and Monitoring Program: Towed-diver Surveys of Large-bodied Fishes of the U.S. Pacific Reefs from 2000 to 2012 <https://inport.nmfs.noaa.gov/inport/item/34521>

### *Ocean Acidification*

NOAA Ecosystem Sciences Division, Pacific Islands Fisheries Science Center (2018). National Coral Reef Monitoring Program: Water chemistry of the coral reefs in the Pacific Ocean. NOAA National Centers for Environmental Information. Dataset. doi:10.25921/nnk7-5f86

### *Reef Material Growth*

NOAA Pacific Islands Fisheries Science Center, Ecosystem Sciences Division (2018). National Coral Reef Monitoring Program: Calcification rates of crustose coralline algae derived from calcification accretion units (CAUs) deployed across American Samoa. NOAA National Centers for Environmental Information. Dataset. doi:10.7289/V5222S2K

NOAA Pacific Islands Fisheries Science Center, Ecosystem Sciences Division (2018). National Coral Reef Monitoring Program: Calcification rates of crustose coralline algae derived from calcification accretion units (CAUs) deployed across the Hawaiian Archipelago. NOAA National Centers for Environmental Information. Dataset. doi:10.7289/V5PK0DFZ

NOAA Pacific Islands Fisheries Science Center, Ecosystem Sciences Division (2018). National Coral Reef Monitoring Program: Calcification rates of crustose coralline algae derived from calcification accretion Units (CAUs) deployed across the Mariana Archipelago. NOAA National Centers for Environmental Information. Dataset. doi:10.7289/V53T9FJ8

NOAA Pacific Islands Fisheries Science Center, Ecosystem Sciences Division (2018). National Coral Reef Monitoring Program: Calcification rates of crustose coralline algae derived from calcification accretion units (CAUs) deployed across the Pacific Remote Island Areas. NOAA National Centers for Environmental Information. Dataset. doi:10.7289/V5G15Z5H

Oliver, T; Vargas-Angel, B; Misa, P; Young, C; Clark, SJ; Pomeroy, N (2015). Calcification Rates of Crustose Coralline Algae derived from Calcification Accretion Units (CAUs) deployed across American Samoa and the Pacific Remote Island Areas in 2010 and recovered in 2012 (NCEI Accession 0137093). Version 2.2. NOAA National Centers for Environmental Information. Dataset.

### *Temperature stress indicator*

NOAA Coral Reef Watch. 2018. NOAA Coral Reef Watch Version 3.1 Daily Global 5-km Satellite Coral Bleaching Heat Stress Monitoring Products (1985-2018). College Park, Maryland, USA: NOAA Coral Reef Watch. Data last accessed 3-27-2018. at <https://coralreefwatch.noaa.gov/product/5km/index.php>.

### *Human Connections*

NOAA Coral Reef Conservation Program; NOAA National Centers for Coastal Ocean Science (2018). National Coral Reef Monitoring Program: Socioeconomic surveys of human use, knowledge, attitudes,

and perceptions in American Samoa. NOAA National Centers for Environmental Information. Dataset. doi:10.7289/V5K

NOAA Coral Reef Conservation Program; NOAA National Centers for Coastal Ocean Science (2018). National Coral Reef Monitoring Program: Socioeconomic surveys of human use, knowledge, attitudes, and perceptions in Hawaii. NOAA National Centers for Environmental Information. Dataset. doi:10.7289/V5G44NM8

NOAA Coral Reef Conservation Program; NOAA National Centers for Coastal Ocean Science (2018). National Coral Reef Monitoring Program: Socioeconomic surveys of human use, knowledge, attitudes, and perceptions in Guam. NOAA National Centers for Environmental Information. Dataset. doi:10.7289/V5QN653V

NOAA Coral Reef Conservation Program; NOAA National Centers for Coastal Ocean Science (2018). National Coral Reef Monitoring Program: Socioeconomic surveys of human use, knowledge, attitudes, and perceptions in the Commonwealth of Northern Mariana Islands (NORTHERN MARIANA ISLANDS). NOAA National Centers for Environmental Information. Dataset. doi:10.25921/1tkk-s064

## Figure data citations

### *Hawaiian Archipelago – Commercial Fishery Harvest Value Figure*

Hawai'i Division of Aquatic Resources. 2015. "State of Hawai'i Fisheries Statistics; Reported Landing Tables." Western Pacific Fisheries Information Network. NOAA Pacific Islands Fisheries Science Center. Honolulu, HI.

### *Hawaiian Archipelago – Coral Cover Decreasing Figure*

Data from Hawai'i Division of Aquatic Resources (DAR) and University of Hawaii Coral Reef Assessment and Monitoring Program (UH CRAMP). Personal communication - data for West Maui was provided by Hal Koike and for West Hawaii by Lindsey Kramer.

### *American Samoa – Average Coral Cover & Biomass of Large Fish Figures*

NOAA Pacific Islands Fisheries Science Center, Ecosystem Sciences Division (2018). National Coral Reef Monitoring Program: Towed-diver surveys of benthic habitats, key benthic species, and marine debris of American Samoa. NOAA National Centers for Environmental Information. Dataset. doi:10.7289/V59G5K30

Coral Reef Ecosystem Program; Pacific Islands Fisheries Science Center (2017). Pacific Reef Assessment and Monitoring Program: Towed-diver Surveys of Benthic Habitats, Key Benthic Species, and Marine Debris Sightings of the U.S. Pacific Reefs from 2000-09-09 to 2012-05-19 (NCEI Accession 0163745). Version 1.2. NOAA National Centers for Environmental Information. Dataset.

NOAA Pacific Islands Fisheries Science Center, Ecosystem Sciences Division (2018). National Coral Reef Monitoring Program: Towed-diver surveys of large-bodied fishes of American Samoa. NOAA National Centers for Environmental Information. Dataset. doi:10.7289/V5F769VR

Coral Reef Ecosystem Program; Pacific Islands Fisheries Science Center (2017). Pacific Reef Assessment and Monitoring Program: Towed-diver Surveys of Large-bodied Fishes of the U.S. Pacific Reefs from 2000-09-09 to 2012-05-19 (NCEI Accession 0163744). Version 1.2. NOAA National Centers for Environmental Information. Dataset.

### *American Samoa – Samoans support reef management figure*

NOAA Coral Reef Conservation Program; NOAA National Centers for Coastal Ocean Science (2018). National Coral Reef Monitoring Program: Socioeconomic surveys of human use, knowledge, attitudes, and perceptions in American Samoa. NOAA National Centers for Environmental Information. Dataset. doi:10.7289/V5K

### *Pacific Remote Islands – Jarvis Coral Cover Declined Figure*

NOAA Pacific Islands Fisheries Science Center, Ecosystem Sciences Division (2018). National Coral Reef Monitoring Program: Benthic cover derived from analysis of images collected during stratified random surveys (StRS) across the Pacific Remote Island Areas. NOAA National Centers for Environmental Information. Dataset. doi:10.7289/V5154FBH

Ecosystem Sciences Division, Pacific Islands Fisheries Science Center (2018). National Coral Reef Monitoring Program: Benthic Images Collected from Stratified Random Sites (StRS) across Jarvis Island of the Pacific Remote Island Areas in May 2016 (NCEI Accession 0176287).

Coral Reef Ecosystem Program; Pacific Islands Fisheries Science Center (2017). Pacific Reef Assessment and Monitoring Program: Towed-diver Surveys of Benthic Habitats, Key Benthic Species, and Marine Debris Sightings of the U.S. Pacific Reefs from 2000-09-09 to 2012-05-19 (NCEI Accession 0163745). Version 1.2. NOAA National Centers for Environmental Information. Dataset.