

NOAA COASTAL OCEAN PROGRAM
Decision Analysis Series No. 2



**TECHNOLOGY AND SUCCESS IN RESTORATION,
CREATION, AND ENHANCEMENT OF *SPARTINA*
ALTERNIFLORA MARSHES IN THE UNITED STATES**
Volume 1 -- Executive Summary and Annotated Bibliography

Geoffrey A. Matthews
Thomas J. Minello

August 1994



U.S. DEPARTMENT OF COMMERCE
National Oceanic and Atmospheric Administration
Coastal Ocean Office

The Decision Analysis Series has been established by NOAA's Coastal Ocean Program (COP) to present documents for coastal resource decision makers which contain analytical treatments of major issues or topics. The issues, topics, and principal investigators have been selected through an extensive peer review process. To learn more about the COP or the Decision Analysis Series, please write:

NOAA
Coastal Ocean Office
1315 East West Highway, Sta. 15140
Silver Spring, MD 20910

or phone 301-713-3338.

Science for Solutions

NOAA COASTAL OCEAN PROGRAM
Decision Analysis Series No. 2



**TECHNOLOGY AND SUCCESS IN RESTORATION,
CREATION, AND ENHANCEMENT OF *SPARTINA*
ALTERNIFLORA MARSHES IN THE UNITED STATES**
Volume 1 -- Executive Summary and Annotated Bibliography

Geoffrey A. Matthews
Thomas J. Minello

NOAA National Marine Fisheries Service
Southeast Fisheries Science Center
Galveston Laboratory
4700 Avenue U
Galveston, TX 77551

August 1994

U.S. DEPARTMENT OF COMMERCE
Ronald H. Brown, Secretary
National Oceanic and Atmospheric Administration
D. James Baker, Under Secretary
Coastal Ocean Office
Donald Scavia, Director

This publication should be cited as:

Matthews, Geoffrey A. and Thomas J. Minello. 1994. Technology and Success in Restoration, Creation, and Enhancement of *Spartina alterniflora* Marshes in the United States. Vol. 1 – Executive Summary and Annotated Bibliography. NOAA Coastal Ocean Program Decision Analysis Series No. 2. NOAA Coastal Ocean Office, Silver Spring, MD.

This publication does not constitute an endorsement of any commercial product or intend to be an opinion beyond scientific or other results obtained by the National Oceanic and Atmospheric Administration (NOAA). No reference shall be made to NOAA, or this publication furnished by NOAA, in any advertising or sales promotion which would indicate or imply that NOAA recommends or endorses any proprietary product mentioned herein, or which has as its purpose an interest to cause directly or indirectly the advertised product to be used or purchased because of this publication.


Note to Readers

The NOAA Coastal Ocean Program (COP) provides a focal point through which the agency, together with other organizations with responsibilities for the coastal environment and its resources, can make significant strides toward finding solutions to critical problems. By working together toward these solutions, we can ensure the sustainability of these coastal resources and allow for compatible economic development that will enhance the well-being of the Nation now and in future generations. The goals of the program parallel those of the NOAA Strategic Plan for 1995-2005.

A specific objective of COP is to provide the highest quality scientific information to coastal managers in time for critical decision making and in a format useful for these decisions. To help achieve this, COP inaugurated a program of developing documents that would synthesize information on issues that were of high priority to coastal managers. To develop such documents, a three-step process was used: 1) to compile a list of critical topics in the coastal ocean through a survey of coastal resource managers and to prioritize and select those suitable for the document series through the use of a panel of multidisciplinary technical experts; 2) to solicit proposals to do research on these topics and select principal investigators through a rigorous peer-review process; and 3) to develop peer-reviewed documents based on the winning proposals.

Seven topics and associated principal investigators were selected in the initial round. *Technology and Success in Restoration, Creation, and Enhancement of Spartina alterniflora Marshes in the United States* by Geoffrey A. Matthews and Thomas J. Minello of the NOAA National Marine Fisheries Service's Galveston Laboratory is the second document in this Decision Analysis Series to be published and is presented in two volumes. Information on Decision Analysis Series No. 1 is shown on the inside back cover. Other volumes will be published over the next two years on the following topics: seagrass restoration technology, coastal watershed restoration, restoring streams and anadromous fish habitat affected by logging, eutrophication and phytoplankton blooms, and management of cumulative coastal environmental impacts.

As with all of its products, COP is very interested in ascertaining the utility of the Decision Analysis Series particularly in regard to its application to the management decision process. Therefore, we encourage you to write, fax, call, or Internet us with your comments. Please be assured that we will appreciate these comments, either positive or negative, and that they will help us direct our future efforts. Our address and telephone and fax numbers are on the inside front cover. My Internet address is DSCAVIA@HQ.NOAA.GOV.



Donald Scavia
Director
NOAA Coastal Ocean Program

VOLUME 1

EXECUTIVE SUMMARY

SECTION I-- ANNOTATED BIBLIOGRAPHY OF SELECTED LITERATURE ON RESTORATION, CREATION, AND ENHANCEMENT OF *SPARTINA* *ALTERNIFLORA* MARSHES IN THE UNITED STATES

EXECUTIVE SUMMARY

Technology and Success in Restoration, Creation, and Enhancement of *Spartina alterniflora* Marshes in the United States

EXECUTIVE SUMMARY

Introduction

Extensive losses of coastal wetlands in the United States caused by sea-level rise, land subsidence, erosion, and coastal development have increased interest in the creation of salt marshes within estuaries. Smooth cordgrass *Spartina alterniflora* is the species utilized most for salt marsh creation and restoration throughout the Atlantic and Gulf coasts of the U.S., while *S. foliosa* and *Salicornia virginica* are often used in California. Salt marshes have many valuable functions such as protecting shorelines from erosion, stabilizing deposits of dredged material, dampening flood effects, trapping water-borne sediments, serving as nutrient reservoirs, acting as tertiary water treatment systems to rid coastal waters of contaminants, serving as nurseries for many juvenile fish and shellfish species, and serving as habitat for various wildlife species (Kusler and Kentula 1989). The establishment of vegetation in itself is generally sufficient to provide the functions of erosion control, substrate stabilization, and sediment trapping. The development of other salt marsh functions, however, is more difficult to assess. For example, natural estuarine salt marshes support a wide variety of fish and shellfish, and the abundance of coastal marshes has been correlated with fisheries landings (Turner 1977, Boesch and Turner 1984). Marshes function for aquatic species by providing breeding areas, refuges from predation, and rich feeding grounds (Zimmerman and Minello 1984, Boesch and Turner 1984, Kneib 1984, 1987, Minello and Zimmerman 1991). However, the relative value of created marshes versus that of natural marshes for estuarine animals has been questioned (Cammen 1976, Race and Christie 1982, Broome 1989, Pacific Estuarine Research Laboratory 1990, LaSalle et al. 1991, Minello and Zimmerman 1992, Zedler 1993). Restoration of all salt marsh functions is necessary to prevent habitat creation and restoration activities from having a negative impact on coastal ecosystems.

This project was undertaken to provide resource managers, habitat researchers, coastal planners, and the general public with an assessment of the technology and success in restoration, enhancement, and creation of salt marshes in the United States. The objective was to be accomplished through the development of three products: 1) an annotated bibliography of the pertinent literature, 2) an inventory of restored, enhanced, or created *Spartina alterniflora* marshes, and 3) a directory of people working in salt marsh creation and restoration. This executive summary describes these products and provides an overall assessment of our understanding regarding restoration, enhancement, and creation of salt marsh habitats. In particular, we have stressed *Spartina alterniflora* marshes and habitat functions related to the support of fishes, crustaceans, and other aquatic life.

Products

Section I. An Annotated Bibliography of Selected Literature on Restoration, Creation, and Enhancement of *Spartina alterniflora* Marshes in the United States.

What was included. This annotated bibliography summarizes the literature on created *Spartina alterniflora* salt marshes, particularly for the last decade. Literature on planting techniques and the establishment of vegetation is included, along with assessments of habitat value and functional equivalency. Publications that involve other species of *Spartina* such as *S. foliosa* have been included if they discuss marsh functions. A particular effort was made to include all studies involving nutrients, sediment organics, infaunal populations, and utilization of created *Spartina* marshes by aquatic and fisheries species.

The annotations were written to assist in determining whether a paper is pertinent to one's needs. Annotations were much like abstracts or summaries, but also included some interpretation on our part, particularly for articles that discussed the use of created salt marshes by fishery or related aquatic organisms.

Articles and project reports were requested from everyone we identified working in the field of salt marsh creation and restoration or conducting functional comparisons of created and natural marshes. Scientific literature was searched using Current Contents.

Lists of reports from various organizations and agencies were searched for articles concerning coastal marsh restoration activities. Articles were also obtained from proceedings of symposia on wetlands restorations, particularly those sponsored each spring by Hillsborough Community College in Florida, and those published by the Association of State Wetland Managers.

Endnote Plus, a computer reference program (Niles and Associates, Inc. Berkeley, CA), was used to create the directory. Entries are in alphabetical order according to the first author's last name. The directory is available in hard copy or as an electronic data file in Endnote Plus, Pro-Cite, Refer/BibIX, and text formats.

Limitations. A natural limitation of this annotated bibliography is that some important articles may have been missed, and new articles will have been published before the bibliography is printed. A few entries were included although their subject matter was tangential to the main focus. This distraction was considered more desirable than excluding the articles, because the entries showed some of the directions that salt marsh restoration activities were heading. Very few entries include quantitative comparisons of faunal utilization in created and natural salt marshes. This paucity of published research limits our ability to assess whether created marshes have attained functional equivalence; fortunately such studies are increasing in number.


Potential Benefits. This bibliography provides a comprehensive collation of literature on coastal salt marsh restoration, enhancement, and creation activities in the U.S. Although the focus was on evaluation of created salt marshes, the bibliography also covers construction techniques, mitigation requirements, sources of information about regional requirements of salt marshes, nutrient requirements and nutrient pools, and even a few articles on regional management practices for coastal zones. Thus, the bibliography should serve anyone looking for documentation on various aspects of coastal salt marsh restoration and coastal-zone wetlands restoration research needs.

Section II. An Inventory of Restored, Created and Enhanced *Spartina alterniflora* Marshes in the United States.

Development. This database contains information about restored, enhanced and created salt marshes in the coastal U.S. and was created using Microsoft Excel. Available

data on location, planting techniques, and success of each marsh was entered via 20 variables. A total of 787 marshes was located and entered in the database over our 1-year study period. Each marsh was assigned a unique inventory number and was also ranked in relation to the amount of available information on the marsh. These information rankings included 1) very little information available, a marsh was planned and permitted but may never have been constructed, 2) a marsh was planted but the location was not precisely identified, 3) a marsh was planted and the exact location is known, 4) a planted marsh was evaluated for successful establishment, 5) a marsh was successfully established and animal utilization was evaluated. The inventory also includes a few entries for marshes where no planting had been done, but restored tidal flow had allowed a marsh to develop (Information Ranking = RTE). Contractors throughout the U.S. provided information for the inventory on *Spartina alterniflora* marshes that they had designed or built. Site information on created marshes was also garnered from publications, reports, conference proceedings, and even assorted notes obtained from research ecologists working in this field. State environmental and fisheries agencies were consulted for their records about man-made salt marshes. Area and district offices of the U.S. Army Corps of Engineers (COE), Soil Conservation Service (SCS), U.S. Fish and Wildlife Service, and the Habitat Protection Branch of the National Marine Fisheries Service (NMFS) were also contacted for information about created *Spartina alterniflora* marshes in their respective coastal areas.

Limitations. The people we contacted in the field of restoration science were generally receptive toward our requests and provided available information. Frequently, however, data required to evaluate salt marsh construction and utilization by animals were never originally recorded, were incomplete, or were not readily available with a reasonable search effort. Thus, many of the marshes in the inventory have low information rankings. For example, some projects that were issued COE permits were not located, and it is unclear whether these marshes were failures or had never been constructed. In addition, we obtained some information from design or initial work plans for projects, and these plans may not have accurately described the actual planted marsh. In situations where we only suspected that a marsh had been created, the inventory entry was given an Information Rank of 1 (see frequency diagram below). For other marshes, only a general description of the marsh location was available (Information Rank 2).



(Rank categories are based on amount of information available [p.4., above], with 5 representing marshes for which the most information was available. RTE refers to marshes with restored tidal exchange.)

Information on created *Spartina alterniflora* marshes on the West Coast was not recorded. A few *Spartina alterniflora* marshes were planted in West Coast estuaries several decades ago, but this species is considered an exotic and is no longer used in that region.

Potential Benefits. Information on locations of created salt marshes is widely scattered, often making it difficult to study the progression of plant and animal populations and the functional development of these habitats. It may take many years for a created marsh to attain the functions of a natural marsh. After a few years, created marshes are often superficially identical to natural marshes, and precise location information is required to continue studying such sites for development of marsh functions. This computerized database is designed to help locate such marshes and to provide basic information about each marsh.

Despite data-collection limitations, the number of marshes in the inventory demonstrates the extent of marsh creation in the coastal areas of the United States. The inventory also demonstrates the amount of data lacking for most of these created marshes. In many cases, these data are relatively easy to record at marsh creation sites (for example the exact location of the marsh), and perhaps the existence of an inventory may stimulate better record keeping. Coastal managers should consider requiring such data records as part of the permitting process. Much of the missing data on marshes included in the inventory may be available somewhere, and we would like to solicit user's help in making the inventory as complete as possible. We anticipate updating the inventory periodically with the addition of new marshes and the inclusion of additional information on the marshes already entered.

Section III. A Directory of Human Resources Involved in the Restoration, Creation, and Enhancement of *Spartina alterniflora* Marshes.

Participants. Participation in the directory was strictly on a voluntary basis. Those included, therefore, have expressed their willingness to interact with the community of people interested in salt marsh restoration. Authors of pertinent research articles were solicited, and many notable estuarine ecologists are participating. The directory also includes staff of various federal permitting, regulatory, and research agencies including: U. S. Army Corps of Engineers, Waterways Experiment Station, National Marine Fisheries Service, U. S. Fish and Wildlife Service - National Biological Survey, and the U. S. Department of Agriculture - Soil Conservation Service. Staff of various state and municipal agencies involved in salt marsh restoration are included. In addition, the directory includes many contractors and others with hands-on knowledge of marsh design and construction. There are over 300 participants in the directory.

Directory Structure. An index lists individuals alphabetically by their last name, but the main contents are organized alphabetically by state. Individuals are listed first in each state, followed by companies and organizations. Each entry includes the person's name, firm/institute/agency, address, telephone number, FAX number, position and specialty, and a description of some of the participant's latest marsh related projects. The directory was created using Endnote Plus, and is available in hard copy or as an electronic data file in Endnote Plus, Pro-Cite, Refer/BibIX, and text formats.

Limitations. This directory is not a "complete" listing of all persons and organizations involved in preserving and restoring salt marshes and coastal wetlands. A large number of people were contacted and asked to participate. Not everyone chose to participate, and undoubtedly we failed to contact some important potential participants. The directory is dynamic and is intended to be updated periodically. We are still soliciting additional participants and updates to entries. Keywords about each individual were not included, but most reference programs can search all fields for descriptive key words; one might focus on the "specialty" or "projects" sections of each person's entry.

Potential Benefits. This directory provides a nucleus of concerned people involved in salt marsh restoration ecology. People from a variety of disciplines are included, and they can provide information about how to manage coastal salt marsh areas, how to create marshes, and how to assess whether the marshes are fulfilling their projected functions. The directory should facilitate information exchange among ecologists working in this field, lay people and managers concerned about their local estuary or marsh, coastal landowners interested in preserving their property from erosion, and companies needing assistance with habitat mitigation.

Conclusions and Recommendations

Smooth cordgrass *Spartina alterniflora* is the dominant salt marsh vegetation along shorelines of Atlantic and Gulf coasts of the United States. Techniques for seeding, planting, and transplanting *S. alterniflora* have been sufficiently developed so that the establishment of vegetation is highly likely when the prescribed conditions are met (Woodhouse 1979, Knutson et al. 1981, Webb and Newling 1985, Allen et al. 1986, Earhart and Garbisch 1983, Broome et al. 1988, Broome 1989, Nailon and Seidensticker 1991). The following key factors have been identified for successfully establishing *S. alterniflora*. 1) Young healthy plants should be used and should be obtained from as close to the planting area as possible. Plants from the project vicinity have the best chance for survival and good growth because they are adapted to the local conditions. 2) Planting should be conducted early in the growing season to provide an adequate length of time for establishment. 3) The soil must contain adequate nutrients. Graded down upland soils often need additions of fertilizer to supply sufficient nutrients, while dredged material or natural bay sediments usually have sufficient nutrients already. 4) Proper elevation

(0.2–0.5 m above Mean Low Water) at the site is critical. Reference should be made to the nearest flourishing natural marsh whenever possible. Smooth cordgrass will grow over a wider tidal range than where it grows best. Success has also been achieved when plants are placed at the higher end of the elevation range and allowed to grow into lower elevations on their own. 5) A gentle slope of 1-10% grade provides sufficient width and drainage for the marsh to develop. This slope should continue seaward from the planted area to reduce wave height and erosive forces. 6) Good water flow and tidal exchange are needed to supply nutrients and to prevent salt build-up in the sediment. 7) Protection from waves is particularly important for a new planting. The fetch should be less than 2 km; shorter fetches are required if the shoreline faces the direction of high winds during stormy weather. 8) Protection of the new plants from pests such as herbivorous fish, insects, small mammals, and man is often needed. Each marsh has a unique suite of potential pests. 9) Protection from activities on adjacent lands has become increasingly important as coastal development continues, and monitoring such activities can protect newly planted marshes from destructive abuse. Although following these nine guidelines and sound planting principles should result in the successful establishment of a *S. alterniflora* marsh, consultation with local experts will often provide technical refinements that will facilitate and insure a greater probability of success.

Above-ground biomass in created *Spartina alterniflora* marshes quickly reaches parity with natural marshes if these basic conditions for marsh establishment and survival are met (Cammen 1975, Webb et al. 1978, Seneca et al. 1985, Webb and Newling 1985, Broome et al. 1986, Broome 1989, and LaSalle et al. 1991). The presence of vegetation in itself is generally sufficient to provide the functions of erosion control, substrate stabilization, and sediment trapping (Knutson et al. 1982). The nutrient-rich sediments trapped by smooth cordgrass aid in maintaining and extending the vegetative zone for the species (Allen and Webb 1982). Apart from above-ground biomass, however, created salt marshes appear to differ from natural marshes in a number of characteristics. Created marshes generally have lower sediment organic content, below-ground biomass, densities of benthic infaunal prey organisms, and densities of nekton on the marsh surface. There is some evidence that these characteristics are linked, and that trophic support for nekton is relatively low in newly created salt marshes.

Below-ground biomass and sediment organic content appear to develop slowly in created salt marshes, and levels may take years to become comparable to natural marshes

(Cammen 1975, Webb et al. 1978, Lindau and Hossner 1981, Craft et al. 1988, Sacco 1989, Langis et al. 1991, LaSalle et al. 1991, Moy and Levin 1991, Minello and Zimmerman 1992). The origin of the sediment used to create the marsh is important, and organic content is generally lower in graded down uplands and sandy dredged material compared with fine-grained dredged material. Low initial levels usually increases the time to reach parity with natural sediments.

The density of benthic infauna such as annelid worms, insect larvae, and small crustaceans is generally lower in created marshes than in natural marshes. These prey organisms are important in supporting the food web of estuarine nekton. In Atlantic coast marshes Cammen (1976), Sacco (1989), and Moy and Levin (1991) found lower densities and generally different species composition for infaunal organisms in created salt marshes. In Texas, Minello and Zimmerman (1992) found that mean amphipod densities in transplanted marshes were only 20-40% of densities in natural marshes, and diversity of infaunal organisms was significantly lower in the transplanted marshes. Although polychaete densities in their study were not significantly different between the natural and transplanted marshes, there was some indication that the abundance of this group was related to the age of the marsh; differences from natural marshes were greatest for the youngest transplanted marsh (2 years old) and smallest for the oldest transplanted marsh (5 years old). LaSalle et al. (1991) also reported a positive relationship between the density of benthic infauna and marsh age. In most of the above studies, a positive relationship was observed between sediment organic content or macro-organic matter and the density of benthic infauna.

Quantitative sampling of nekton densities in salt marsh vegetation is difficult (Zimmerman et al. 1984, Kneib 1991, Rozas 1992), thus comparisons of surface utilization between created and natural marshes are limited. Relative abundance in pit traps or Breder traps is unreliable in making such comparisons, because the sampling area of these devices cannot be accurately defined. Minello and Zimmerman (1992) used a drop sampler to compare springtime densities in three created and three natural *S. alterniflora* marshes (3-5 years in age) on the Texas coast and found overall densities of large macrofauna and decapod crustaceans were significantly lower in the transplanted marshes. These differences were due mainly to daggerblade grass shrimp and young brown shrimp. Diversity of decapod crustaceans was also higher in the natural marshes. Densities of fish (mainly the darter goby and pinfish) as a group were not significantly different between natural and transplanted marshes. In addition, fish diversity was consistently higher in the transplanted marshes. In a

larger study of ten created salt marshes (3-15 years in age) in Galveston Bay, Texas, Minello and Webb (1993) also found reduced densities of commercially important crustaceans (brown shrimp, white shrimp, and blue crabs) when the marshes were compared with five natural marshes. In North Carolina, Meyer et al. (1993) used block nets to compare densities of fishes, shrimps, and crabs in a planted and natural *Spartina alterniflora* marsh. Two years following planting, overall mean densities of shrimp (mainly daggerblade grass shrimp and brown shrimp) were about three times higher in the natural reference marsh compared with the planted marsh, but differences were not statistically significant apparently due to high sample variability. Mean crab densities were two to three times higher in the created marsh, but again the differences were not significant. Fishes collected included spot, mummichog, pinfish, and pigfish, and the mean density for total fish was twice as high in the natural marsh as in the created marshes. This difference was statistically significant during two of the three sampling periods. These conflicting results in marsh utilization patterns may be the result of different sampling gears and techniques or may reflect regional differences in the way salt marshes function for nektonic organisms. The results may also be attributable to the natural variability in populations of organisms that occur from year to year and the natural variability in carrying capacity of habitats. Until a substantial number of such quantitative comparisons are available, it will be difficult to assess functional parity between created and natural salt marsh habitats.

Understanding the functional development of created marshes requires an understanding of how natural salt marsh systems function. There is evidence, for example, that salt marshes function differently for shrimp, crabs, and fishes in different coastal regions. Direct exploitation of the marsh surface is extensive in many marshes of the northern Gulf of Mexico (Zimmerman and Minello 1984, Thomas et al. 1990, Baltz et al. 1993, Rozas and Reed 1993, Peterson and Turner 1994), thus densities of decapods and fishes and their prey on the marsh surface should reflect habitat value for marshes in this region. Direct use of the marsh surface may be lower in marshes along the Atlantic coast, and nekton densities in these marshes appear to be substantially lower than in the Gulf (Hettler 1989, Mense and Wenner 1989, Kneib 1991, Fitz and Wiegert 1991). Atlantic coast marshes have relatively less edge, higher elevations, and different tidal inundation patterns, and all of these factors may affect utilization patterns of nekton (Rozas 1993). These regional differences increase the necessity for functional studies comparing natural and transplanted salt marshes in various parts of the country.

Indications of long-term retarded functional development in created salt marshes suggest that some habitat functions may never fully develop. However, under the assumption that most created marshes can eventually develop into fully functional habitats, there has been considerable interest in methods for stimulating marsh development rates. The correlative relationships identified between sediment organic content and infaunal populations have inspired work on the addition of organic amendments to sediments before planting. This work has shown promise for increasing the rate of marsh development, but success may be related to local and regional conditions (Currin, C.; Zedler, J.; Broome, S.; personal communications). The value of increasing sediment organic matter through the addition of soil amendments may depend upon the causal relationships between infaunal abundance and the development of marsh sediments. For example, if infaunal populations increase because sediment organic matter is used as food by deposit feeders, the relative amount of refractory organic matter may be important. However, sediment organics may also alter the physical structure of the sediment improving the habitat for infauna. Thus, factors such as porosity, bulk density, size of detrital particles, and inorganic grain size distribution may be important. In addition, the oxygenation of sediments by live plant roots may be a factor in increasing infaunal abundance. Controlling development rates of marsh sediments, therefore, may be difficult without a better understanding of sediment/infaunal relationships. Other research on improving functionality of created marshes indicates that the addition of edge through the construction of channels and creeks can dramatically increase use of the marsh surface by nekton (Minello et al. 1994). The value of adding edge, however, may also be a regional phenomena, and adding edge in Atlantic coast marshes may not be beneficial. Natural salt marshes in the northern Gulf of Mexico have relatively more edge and are apparently used more extensively by aquatic animals than marshes along the Atlantic coast (Rozas 1993).

Despite a great deal of rhetoric regarding functional equivalency in created marshes, quality research on the problem has been limited. Important questions that still must be addressed center on how natural salt marshes function for fishery and aquatic species and how these functions vary regionally. Without this basic understanding of salt marsh ecosystems, comparisons of natural and created systems will continue to be limited in scope. Comparisons of natural and created marsh use by fishery species and other aquatic nekton are needed, but these comparisons must be based on quantitative samples in marsh creeks and within the vegetation itself. The use of enclosure samplers is generally required for measuring animal densities in these vegetated habitats. A variety of such

devices have now been developed including throw traps and drop samplers (Kushlan 1981, Zimmerman et al. 1984), flume and block nets (McIvor and Odum 1986, Hettler 1989), and lift nets and flume weirs (Rozas 1992, Kneib 1991). Comparisons of functional equivalency in created and natural marshes may also require experimental measurements of animal growth and predation risk in these habitats.

A basic and pervasive problem in functional comparisons is a limited replication of study marshes. The development of our marsh inventory may be useful in this regard by helping to identify additional created marshes for analysis. Sound inferences concerning relative functioning of marshes require an estimate of variability among both natural marshes and created marshes. If for example, only one created marsh is examined in the study area, conclusions can only be made regarding that marsh and not created marshes in general. In a similar manner, variability among natural marshes is often great and should also be assessed. Data on spatial variability within marshes and seasonal and annual variability are also important (Zedler et al. 1986), but information on marsh to marsh variability will be most useful in assessing the overall value of created marshes compared with natural marshes.

Literature Cited

Allen, H. H., S. O. Shirley, and J. W. Webb Jr. "Vegetative stabilization of dredged material in moderate to high wave-energy environments for created wetlands." In: Proceedings of the thirteenth annual conference on wetland restoration and creation, in Hillsborough, FL, edited by F. J. Webb, Hillsborough, FL: Hillsborough Community College, 19-35, 1986.

Allen, H. H. and J. W. Webb Jr. "Influence of breakwaters on artificial salt marsh establishment on dredged material." In: Proceedings of the ninth annual conference on wetland restoration and creation, in Tampa, FL., edited by F. J. Webb, Hillsborough Community College, 18-35, 1982.

Baltz, D. M., C. Rakocinski, and J. W. Fleeger. "Microhabitat use by marsh-edge fishes in a Louisiana estuary." Environ. Biol. Fishes 36 (1993): 109-126.

Boesch, D. F. and R. E. Turner. "Dependence of fishery species on salt marshes: the role of food and refuge." Estuaries 7 (1984): 460-468.

Broome, S. W. "Creation and restoration of tidal wetlands of the southeastern United States." In: Wetland Creation and Restoration: The Status of the Science. Part 1, Regional Review., eds. J. A. Kusler and M. E. Kentula. 37-72. Washington, D.C.: Island Press, Inc., 1989.

Broome, S. W., E. D. Seneca, and W. W. Woodhouse Jr. "Long-term growth and development of transplants of the salt-marsh grass *Spartina alterniflora*." Estuaries 9 (1986): 63-74.

Broome, S. W., E. D. Seneca, and W. W. Woodhouse Jr. "Tidal salt marsh restoration." Aquat. Bot. 32 (1988): 1-22.

Cammen, L. M. "Accumulation rate and turnover time of organic carbon in a salt marsh sediment." Limnol. Oceanogr. 20 (1975): 1012-1015.

Cammen, L. M. "Abundance and production of macroinvertebrates from natural and artificially established salt marshes in North Carolina." American Midland Naturalist 96 (1976): 487-493.

Craft, C., S. Broome, and E. Seneca. "The role of transplanted marshes in processing nitrogen, phosphorus and organic carbon in estuarine waters." In: Increasing our wetland resources, proceedings of a conference. Natl. Wildl. Fed., eds. J. Zelazny and J. Feierabend. 327-332. Washington, DC.: 1988.

Earhart, H. G. and E. W. Garbisch Jr. "Habitat development utilizing dredged material at Barren Island, Dorchester County, Maryland." Wetlands 3 (1983): 108-119.

Earhart, H. G. and E. W. Garbisch Jr. "Beneficial uses of dredged materials at Barren Island, Dorchester County, Maryland." In: Proceedings of the thirteenth annual conference on wetlands restoration and creation. in Hillsborough, FL, edited by F. J. Webb Jr., Hillsborough, FL: Hillsborough Community College, 75-85, 1986.

Fitz, H. C. and R. G. Wiegert. "Utilization of the intertidal zone of a salt marsh by the blue crab *Callinectes sapidus* - Density, return frequency, and feeding habits." Mar. Ecol. Prog. Ser. 76 (1991): 249-260.

Hettler, W. F. "Nekton use of regularly-flooded salt marsh cordgrass habitat in North Carolina, USA." Mar. Ecol. Prog. Ser. 56 (1989): 111-118.

Kneib, R. T. "Patterns of invertebrate distribution and abundance in the intertidal salt marsh: causes and questions." Estuaries 7 (1984): 392-412.

Kneib, R. T. "Predation risk and use of intertidal habitats by young fishes and shrimp." Ecology 68 (1987): 379-86.

Kneib, R. T. "Flume weir for quantitative collection of nekton from vegetated intertidal habitats." Mar. Ecol. Prog. Ser. 75 (1991): 29-38.

Knutson, P. L., R. A. Brochu, W. N. Seelig, and M. Inskeep. "Wave damping in *Spartina alterniflora* marshes." Wetlands 2 (1982): 87-104.

Knutson, P. L., J. C. Ford, M. R. Inskeep, and J. Oyler. "National survey of planted salt marshes (vegetative stabilization and wave stress)." Wetlands 1 (1981): 129-157.

Kushlan, J. "Sampling characteristics of enclosure fish traps." Trans. Am. Fish. Soc. 110 (1981): 557-562.

Kusler, J. A. and M. E. Kentula, eds. Wetland Creation and Restoration. The Status of the Science. Washington, DC: Island Press, 1989.

Langis, R., M. Zalejko, and J. B. Zedler. "Nitrogen assessments in a constructed and a natural salt marsh of San Diego Bay." Ecol. Appl. 1 (1991): 40-51.

LaSalle, M. W., M. C. Landin, and J. G. Sims. "Evaluation of the flora and fauna of a *Spartina alterniflora* marsh established on dredged material in Winyah Bay, South Carolina." Wetlands 11 (1991): 191-208.

Lindau, C. W. and L. R. Hossner. "Substrate characterization of an experimental marsh and three natural marshes." Soil Sci. Soc. Amer. J. 45 (1981): 1171-76.

McIvor, C. C. and W. E. Odum. "The flume net: A quantitative method for sampling fishes and macrocrustaceans on tidal marsh surfaces." Estuaries 9(3) (1986): 219-224.

Mense, D. J. and E. L. Wenner. "Distribution and abundance of early life history stages of the blue crab, *Callinectes sapidus*, in tidal marsh creeks near Charleston, South Carolina." Estuaries 12 (1989): 157-68.

Meyer, D. L., M. S. Fonseca, D. R. Colby, W. J. Kenworthy, and G. W. Thayer. "An examination of created marsh and seagrass utilization by living marine resources." In: Coastal Zone '93, Volume 2. Proceedings of the 8th Symposium on Coastal and Ocean Management., eds. O. Magoon, W.S. Wilson, H. Converse, and L. T. Tobin. 1858-1863. New York.: American Society Of Civil Engineers, 1993.

Minello, T. J. and J. W. Webb Jr. "The development of fishery habitat value in created salt marshes." In: Coastal Zone '93, Volume 2. Proceedings of the 8th Symposium on Coastal and Ocean Management., eds. O. Magoon, W.S. Wilson, H. Converse, and L. T. Tobin. 1864-1865. New York: American Society Of Civil Engineers, 1993.

Minello, T. J. and R. J. Zimmerman. "The role of estuarine habitats in regulating growth and survival of juvenile penaeid shrimp." In: Frontiers in shrimp research., eds. P. DeLoach, W. J. Dougherty, and M. A. Davidson. 1-16. Amsterdam: Elsevier Sci. Publ., 1991.

Minello, T. J. and R. J. Zimmerman. "Utilization of natural and transplanted Texas salt marshes by fish and decapod crustaceans." Mar. Ecol. Prog. Ser. 90 (1992): 273-285.

Minello, T. J., R. J. Zimmerman, and R. Medina. "The importance of edge for natant macrofauna in a created salt marsh." Wetlands (in press) (1994):

Moy, L. D. and L. A. Levin. "Are *Spartina* marshes a replaceable resource? A functional approach to evaluation of marsh creation efforts." Estuaries 14 (1991): 1-16.

Nailon, R. W. and E. L. Seidensticker. "The effects of shoreline erosion in Galveston Bay, Texas." In: Coastal wetlands, ed. H. S. Bolton. 193-206. New York: Amer. Soc. Civil Engineers, 1991.

Pacific Estuarine Research Laboratory. A manual for assessing restored and natural coastal wetlands with examples from southern California. California Sea Grant Report No. T-CSGCP-021. La Jolla, California: 1990.

Peterson, G. W. and R. E. Turner. "The value of salt marsh edge vs interior as a habitat for fish and decapod crustaceans in a Louisiana tidal marsh." Estuaries 17 (1994): 235-262.

Race, M. S. and D. R. Christie. "Coastal zone development: mitigation, marsh creation, and decision-making." Environ. Manag. 6 (1982): 317-328.

Rozas, L. P. "Bottomless lift net for quantitatively sampling nekton in intertidal marshes." Mar. Ecol. Prog. Ser. 89 (1992): 287-292.

Rozas, L. P. "Nekton use of salt marshes of the Southeast region of the United States." In: Coastal Zone '93, Volume 2. Proceedings of the 8th Symposium on Coastal and Ocean Management., eds. O. Magoon, W.S. Wilson, H. Converse, and L. T. Tobin. 528-536. New York: American Society Of Civil Engineers, 1993.

Rozas, L. P. and D. J. Reed. "Nekton use of marsh-surface habitats in Louisiana (USA) deltaic salt marshes undergoing submergence." Mar. Ecol. Prog. Ser. 96 (1993): 147-157.

Sacco, J. Infaunal community development of artificially established salt marshes in North Carolina. Ph.D. Thesis, North Carolina State University. Raleigh: 1989.

Seneca, E. D., S. W. Broome, and W. W. Woodhouse Jr. "Comparison of *Spartina alterniflora* Loisel. transplants from different locations in a man-initiated marsh in North Carolina." Wetlands 5 (1985): 181-190.

Thomas, J. L., R. J. Zimmerman, and T. J. Minello. "Abundance patterns of juvenile blue crabs (*Callinectes sapidus*) in nursery habitats of two Texas bays." Bull. Mar. Sci. 46 (1990): 115-125.

Turner, R. E. "Intertidal vegetation and commercial yields of penaeid shrimp." Trans. Am. Fish. Soc. 106 (1977): 411-16.

Webb, J. W., J. D. Dodd, B. W. Cain, W. R. Leavens, L. R. Hossner, C. Lindau, R. R. Stickney, and H. Williamson. Habitat development field investigations, Bolivar Peninsula marsh and upland habitat development site, Galveston Bay, Texas, Appendix D: Propagation of vascular plants and postpropagation monitoring of botanical, soil, aquatic biota, and wildlife resources. Tech. Rept. D-78-15. Vicksburg, MS: U.S. Army Corps of Engineers, Waterways Experiment Station, 1978.

Webb, J. W., Jr. and C. J. Newling. "Comparison of natural and man-made salt marshes in Galveston Bay complex, Texas." Wetlands 4 (1985): 75-86.

Woodhouse, W. W., Jr. Building salt marshes along the coasts of the continental United States. Special Report No. 4. Fort Belvoir, VA: U.S. Army Corps Eng., Coastal Eng. Res. Cent., 1979.

Zedler, J. B. "Canopy architecture of natural and planted cordgrass marshes: selecting habitat evaluation criteria." Ecol. Appl. 3 (1993): 123-138.

Zedler, J. B., J. Covin, C. Nordby, P. Williams, and J. Boland. "Catastrophic events reveal the dynamic nature of salt-marsh vegetation in southern California." Estuaries 9 (1986): 75-80.

Zimmerman, R. J. and T. J. Minello. "Densities of *Penaeus aztecus*, *P. setiferus* and other natant macrofauna in a Texas salt marsh." Estuaries 7 (1984): 421-433.

Zimmerman, R. J., T. J. Minello, and G. Zamora. "Selection of vegetated habitat by brown shrimp, *Penaeus aztecus*, in a Galveston Bay salt marsh." Fish. Bull., U.S. 82 (1984): 325-336.

SECTION I

ANNOTATED BIBLIOGRAPHY OF SELECTED LITERATURE ON RESTORATION, CREATION, AND ENHANCEMENT OF *SPARTINA* *ALTERNIFLORA* MARSHES IN THE UNITED STATES

Foreword

Coastal development, sea level rise, and land subsidence have resulted in extensive losses of estuarine salt marsh habitat throughout much of the United States. Concomitant with this habitat loss is the loss of salt marsh functions. These functions include the stabilization of shorelines, prevention of erosion, flood control, sediment trapping, nutrient cycling, and removal of toxic wastes in watersheds. In addition, salt marshes provide valuable habitat for estuarine animals including economically important fishery species.

Efforts to implement a "no net wetland loss" policy in the United States will require a continuation and expansion of programs to restore and create salt marshes in regions of deteriorating coastal wetlands. These restoration projects frequently involve transplanting marsh vegetation on graded-down uplands or on dredged material. Although many of these efforts have failed, the establishment of vegetative growth in this manner often has been successful, and marsh planting techniques have been developed for a variety of coastal conditions.

Replacing or creating vegetative cover, however, is only the first step in creating a functional salt marsh. Habitat functions such as providing food, protection from predators, and reproductive sites for estuarine animals must also be replaced. The development of these habitat functions does not appear to parallel the growth of macrophytes, yet it is the growth of the plants that has been the usual measure of marsh creation success. This retarded functional development has raised serious questions about the relative value of created salt marshes. Few direct studies of animal growth, survival, or reproductive success have been conducted in created salt marshes. The abundance of animals in a salt marsh is generally used as an indicator of relative habitat value, although valuable habitats do not always support high densities of animals. Comparing animal abundance in natural and created salt marshes, however, is in itself a difficult problem, and the importance of collecting quantitative samples in assessing use of salt marshes by fishery and other aquatic organisms cannot be overemphasized. Sampling techniques often include the use of breeder traps, seines, trawls, or simple qualitative visual estimates. These techniques are generally not quantitative, and thus results based on their use must be weighed accordingly.

Resource managers, habitat researchers, and coastal planners need assistance in developing marsh restoration projects. The literature documenting efforts in this area is scattered and not readily available, and a synthesis of these data should be valuable for determining the most appropriate restoration and creation techniques applicable to different

coastal conditions. Of special interest is an analysis of the importance of regional differences in successful restoration techniques. In addition, an assessment of whether these projects have replaced salt marsh functions is of primary importance.

This annotated bibliography is intended to summarize the literature on created *Spartina alterniflora* salt marshes. Although papers that deal with planting techniques and the establishment of the vegetation itself are included in the collection, special emphasis is placed on publications that assess habitat value and investigate the replacement of natural salt marsh functions. Some papers that involve other species of *Spartina* such as *S. foliosa* have been included in the bibliography if the papers relate to functional values. We tried to include all studies involving nutrients, sediment organics, and infaunal populations and studies that examined utilization of created *Spartina* marshes by fishery species or other aquatic or terrestrial fauna. The annotations were written to assist the reader in determining whether a paper was pertinent for their needs; annotations were not designed to substitute for reading the papers themselves. In contrast to abstracts or summaries, the annotations include some interpretation on our part. This is especially true in relation to the use of created salt marshes by fishery or related aquatic organisms. In the annotations, we have adopted the practice of abbreviating elevation references using the capitalized first letter of each word, included are: MHW for mean high water, MHT for mean high tide, MSL for mean sea level, MLT for mean low tide, MLW for mean low water, MLLW for mean lower low water, MTL for mean tide level, and NGVD for national geodetic vertical datum. These abbreviations are defined in a glossary of terms in the Tide Tables 1994 High and Low Water Predictions, East Coast of North and South America Including Greenland (DOC/NOAA/NOS, Washington, DC).

This bibliography is not totally comprehensive, rather it contains a wide selection of papers that we thought were important in advancing the science of salt marsh restoration. Undoubtedly, we have missed some significant contributions to the literature. The reader may wish to consult annotated bibliographies by Wolf et al. (1986) and by the U.S. Fish and Wildlife Service (see Miller et al. 1991); these bibliographies are much broader in scope and cover most studies involving created wetlands.

Acknowledgments

Creating this bibliography was greatly facilitated by the many scientists who kindly sent us reprints of their articles and reports of their projects, and who discussed the projects with us when possible. We are very grateful for their aid and interest. Four regional assistants helped obtain much of the gray literature reviewed here, and we thank them for their work: Chris L. Sardella (Northeast Coast), Andrew L. Bunch (South Atlantic Coast), Joseph L. Staton (Gulf of Mexico), and Russell E. DiFiori (West Coast). Thanks are also extended to Mark E. Pattillo who also reviewed articles and wrote their annotations. This work was funded by the National Marine Fisheries Service, Southeast Fisheries Science Center, and by a Resource Information Delivery grant from NOAA Coastal Ocean Program.

Table of Contents

Section I -- Annotated Bibliography

FOREWORD	iii
ACKNOWLEDGMENTS	v
BIBLIOGRAPHY	1
AUTHOR INDEX	63
SUBJECT INDEX	67

Bibliography

Allen, H. "Biotechnical stabilization of dredged material shorelines." In: Beneficial Uses of Dredged Material, ed. M. C. Landin. 116-128. January 1988. Baltimore MD: US Army Corps of Engineers, 1988.

This review covers the efforts of the U.S. Army Corps of Engineer's Waterways Experiment Station in Vicksburg, Mississippi to develop biotechnical techniques for the stabilization of dredged material shorelines. Biotechnical stabilization combines the use of mechanical structures with biological elements (plants). Mechanical structures, including breakwaters and biodegradable mats, are useful in high energy (fetches over 9 km) environments for reducing erosion until plants such as *Spartina alterniflora* can become established.

Breakwaters are only necessary for 2-3 years until plants become established. The use of sand bags, floating tires, and tire-pole breakwaters is discussed. These techniques reduced shoreline energy under proper conditions. Other devices useful in the stabilization of plant stems include fibrous erosion control mats and burlap plant rolls. Techniques employing these devices are described and show promise in allowing the establishment of plants in high-energy areas. In general, the use of mats and plant rolls is more economical than breakwaters.

Allen, H. H., E. J. Clairain, R. J. Diaz, A. W. Ford, L. F. Junt, and B. R. Wells. "Habitat development field investigations--Bolivar Peninsula Marsh and upland habitat development site, Galveston, Texas: summary report." 73. City: U.S. Army Corps Eng. Waterways Experiment Station, Environmental Lab., 1978.

This report described an experiment to establish salt marsh and upland vegetation on a 2-yr old dredged material deposition site of about 7.3 ha on the Galveston Bay side of Bolivar Peninsula, Texas. The report also described the construction of a sandbag dike to reduce wave impacts, and a fence to keep large mammals out, including goats, raccoons and people out.

Most of the grading and sand moving on the site was accomplished with a small bulldozer and a rubber tire frontend loader. Intertidal plantings were in 3 elevation tiers. Winter, spring and summer plantings, various fertilizers and application methods, several plant species, and planting techniques were tried. Monthly samplings recorded changes in plant height, density, # stems/plant, # of stressed plants, # of stable plants, % foliar cover, vegetative reproduction, # of plants with flowers, seed heads and new growth, and above and below-ground biomass.

Spartina alterniflora grew best in 0.06-0.21 m above MSL. At this elevation, tidal inundation occurred 69-87% of the time from Feb.-Aug'77. *Spartina patens* grew best at 0.37 m above MSL; inundation was less than 30% of the time. Sprigging was much more successful than seeding, but seeding (more economical) was used successfully in the upper third of the intertidal zone, where inundation frequency was low but soil remained moist, and there was less washing out by even small waves. Fertilizers were not particularly effective. It was thought that the nutrient-rich fine sediments that became deposited in the marsh planting area promoted enough growth to mask the action of the fertilizers.

Sampling for fish and aquatic animals was conducted before and after marsh construction. No major changes were detected in fish diversity or abundance after marsh planting, however, no adequate sampling devices were available for sampling the marsh grass at that time. Increases were noted in the polychaetes and oligochaetes in the benthos in the marsh area protected by the dike. After marsh construction, some increases in bird and mammal use of the area were noted. More marsh birds were noted once the cordgrasses developed. Also, more rats and mice were noted when sufficient leaf growth was established to provide covered habitat. Marsh rabbits ate the new growth, but did little damage.

The overall results from this marsh establishment project were that a salt marsh could be created on dredged material, and that it could function like a natural marsh provided certain precautions were taken. Wave energy had to be greatly reduced to prevent erosion of plants, and new growth had to be protected from foraging mammals.

Allen, H. H., S. O. Shirley, and J. W. Webb, Jr. "Vegetative stabilization of dredged material in moderate to high wave-energy environments for created wetlands." In: Proceedings of the thirteenth annual conference on wetland restoration and creation. in Hillsborough, FL, edited by F. J. Webb, Jr. Hillsborough, FL: Hillsborough Community College, 19-35, 1986.

This paper reports the results of several attempts to stabilize shorelines using *Spartina alterniflora*. High wave energy erodes shoreline vegetation, and special devices to reduce waves must be used before cordgrass can be established to reduce erosion. Two breakwater designs and four planting enhancement treatments were tested using four test locations. The breakwaters were a 3-tier floating tire fence and a fixed tire-pole design that was two tires high by two tires wide. The enhanced treatments were: (1) multiple stemmed clumps, (2) roots and lower stems of clumps were wrapped in a piece of burlap, (3) a "plant roll" was made by rolling a 3.7 x 0.9 m burlap cloth around sandy soil with six clumps of *Spartina* spaced in it at 0.5-m intervals, and (4) single stems of *Spartina* sprigged in a woven natural fiber mat which was then anchored to the substrate. Results showed a breakwater was needed if a transplant project was to be successful in areas of high wave energy. Clumps proved as effective as the other more expensive treatments behind the breakwaters. Plant rolls functioned to protect single stems planted to landward of the rolls in areas of only moderate wave energy.

Allen, H. H. and J. W. Webb, Jr. "Influence of breakwaters on artificial salt marsh establishment on dredged material." In: Proceedings of the ninth annual conference on wetland restoration and creation. in Tampa, FL., edited by F. J. Webb, Jr. Tampa, FL.: Hillsborough Community College, 18-35, 1982.

Cordgrass was transplanted to a site with high wave energy to stabilize a dredged material dike and to provide marsh habitat. The site was located on the northwest side of a recently deposited dredged material island situated in the center of Mobile Bay, Alabama. Wind fetches were 4.8 to 6.4 km. An unprotected planting of 1.5 ha with *Spartina alterniflora* single stems on 1.0-m centers was a complete failure. A second planting was made in the same area, this time behind two types of wavebreaks. Two months after planting there was a 55% survival behind the floating-tire-breakwater, and a 24% survival behind a fixed-fence-breakwater. The fixed-fence-breakwater, however, was coming apart, and additional losses of transplants were expected due to washout. Use of a fertilizer during the planting did not appear to be of benefit. Spacing of transplants was also tested. Transplants set on 1.0-m centers yielded 34% survival versus 22% survival for those set on 0.5-m centers. A reason for the difference in survival was not offered.

In areas with significant wave energy, transplanting success will depend on a functional breakwater. The floating tire breakwater design appears to be successful though somewhat costly (about \$126 per linear meter). This breakwater can be partially disassembled, moved to another location and reassembled to protect another transplanting, thus making it a little more cost-effective.

Although the area receiving transplants on 1.0-m centers had slightly better survival, it did not have the coverage afforded by the area planted on 0.5-m centers. Coverage on the latter area was twice that of the 1.0-m centers. It depends on how fast one needs coverage as to which density of transplanting one should use. If you can wait three or four years for a stand to form, the 1.0-m center regimen would be less expensive and faster to plant.

Allen, H. H. and J. W. Webb, Jr. "Bioengineering methods to establish salt marsh on dredged material." In: Coastal Zone '93 in New Orleans, LA, edited by S. Laska and A. Puffer, New Orleans, LA: Amer. Soc. Civil Eng., 118-132, 1993.

This paper describes a test of low-cost wave stilling devices along with five transplanting treatments, all for use in establishing a fringe *Spartina alterniflora* marsh along a shoreline that is subject to moderate to high wave action (energy). The study was done on Bolivar Peninsula in the Galveston Bay area, Texas. Two breakwaters were used, one was a modified floating tire design (FTB) and the other was a fixed tire design. Transplanting treatments were: single stem, multi-stem clumps, multi-stem clumps wrapped in burlap, multi-stem clumps on 0.5-m intervals in a burlap roll with substrate between, and single stems planted through slits in an erosion control fiber mat that was anchored in the substrate. Only single stems were planted behind the breakwaters. Four test plots of each of the planting treatments were done outside the breakwaters.

Planting occurred in July 1984. In a January 1985 census only about 8% of the plants in the plots outside the breakwaters were surviving, and most was in two of the erosion control mat plots. However, 2.5 years later, there was at least a 25% cover in 3 mat plots, 2 multi-stem plots, and 1 burlap wrapped multi-stem plot. Growth continued in these plots, but the shoreline receded, leaving them as small islands out from shore. The fixed tire breakwater failed after about a year, and the 50% cover that had become established disappeared after about four years. The FTB worked well, and the stand of *Spartina*

expanded well beyond the initial plot by the end of five years, despite an initial sediment build-up behind the FTB which buried many transplants.

Athnos, D. L. "Compensatory mitigation in the Gulf coast states: Can we achieve "No net loss" of wetlands?" City: Inst. for Coastal and Estuarine Research, Univ. W. Florida, 1993.

This report summarizes wetland laws and regulations for each of the Gulf coast states, assesses the success of these regulations in achieving no net loss of wetlands through a review of recent literature, and presents recommendations which would help attain a "no net loss" status. Federal and state laws and regulations governing wetlands along the Gulf of Mexico coast were succinctly and clearly presented. Because the nation and many states do not have comprehensive laws protecting wetlands, wetlands are tentatively protected under the Clean Water Act, Section 404, which is under the administration by the U.S. Army Corps of Engineers and the Environmental Protection Agency.

The author concluded the system appeared to be failing and wetlands were being lost. A major portion of the loss was due to activities that are exempt for the 404 permitting process. But the process was not replacing the loss even when mitigation was available. The author offered 15 recommendations to assist in wetlands protection.

Baca, B. J. and T. W. Kana. "Methodology for restoring impounded coastal wetlands." In: Proceedings of the thirteenth annual conference on wetlands restoration and creation, in Hillsborough, FL, edited by F. J. Webb, Jr., Hillsborough, FL: Hillsborough Community College, 36-44, 1986.

This paper emphasizes the importance of water flows and tidal elevations at which various species of marsh plants are found, as the important factors to consider when restoring impounded intertidal areas. South Carolina has about 70,000 acres of coastal impoundments, and many of these could be reclaimed for tidal wetlands in the near future. Appropriate plants should be selected based on elevation and flushing patterns at a site.

Banner, A. "Revegetation and maturation of restored shoreline in Indian River, Florida." In: Proceedings of the fourth annual conference on restoration of coastal vegetation in Florida, in Tampa, FL, edited by R. R. Lewis, III and D. Cole, Tampa, FL: Hillsborough Community College, 13-42, 1977.

Restoration activities at a site just north of Vero Beach, Florida, are described. This was mitigation for destruction of natural waterfront and a slough. *Spartina alterniflora* (smooth cordgrass), *Rhizophora mangle* (red mangrove), *Laguncularia racemosa* (white mangrove), and *Avicennia germinans* (black mangrove) were transplanted to the site the same day they were dug from nearby donor sites along the Indian River. The substrate at the site was a coarse yellow sand that was mined locally, transported and dumped on the site, and smoothed to a slope of 7.5%. A breakwater was built to prevent erosion by storm waves. Plugs, 15 cm in diameter containing 30 cm tall cordgrass plants were planted on 0.5-m centers. Mangrove seedlings 40-90 cm tall were used. Planting was done in May, 1976.

Within two months after transplanting, about 75% of the red and black mangroves had lost their leaves. These plants never recovered. Mangrove plants continued to die through the remainder of the year. The four white mangrove seedlings that were

transplanted survived. On the other hand, *Spartina* transplants survived and grew, and by November many had set seed. In December, young *Spartina* seedlings were found on control (bare) areas and on areas where mangroves had been planted but had died. The seedlings were volunteers from the seed from nearby natural marshes. Although the combination planting of red mangroves with cordgrass did not help the red mangroves survive, the following year red mangrove propagules colonized the cordgrass stands.

Halodule and *Halophila* developed lush beds just subtidal to the mitigation site but on the coarse yellow sand. These were not planned, but offered a chance to observe bed development.

This simple experimental restoration project showed *Spartina alterniflora* was a valuable plant for stabilizing bare intertidal shoreline. Cordgrass grew rapidly and spread, forming stands that also assisted in the establishment of red mangroves by trapping and holding the propagules. Mangrove transplants were not as hardy nor as useful for shoreline stabilization.

Beeman, S. "Techniques for the creation and maintenance of intertidal saltmarsh wetlands for landscaping and shoreline protection." In: Proceedings of the tenth annual conference on wetlands restoration and creation, in Tampa, FL, edited by F. J. Webb, Jr. Tampa, FL: Hillsborough Community College, 33-43, 1983.

This paper describes techniques for establishing and maintaining an intertidal salt marsh along the Atlantic coast of Florida, based on literature and 10 years of experience (make that 20 years today) in the field of marsh creation. Factors considered important included: soil composition, elevation and slope of the shoreline, size of the plants used, planting density, time of year of installation, and proper choice of plant species.

Best results were achieved where a breakwater was constructed at the MLW seaward edge of the flattened slope, and a berm was constructed above the planting site to prevent erosion due to upland runoff. An intertidal marsh slope between 5° and 15° and a high marsh slope between 15° to 25° were recommended. Large plugs (rather than single sprigs) of *Spartina alterniflora* were planted on 1-ft centers between MLW and MHW. Plugs of *Spartina patens* were planted above MHW, and plugs of *Distichlis spicata* and *Sporobolus virginicus* were planted at even higher elevations. These elevations matched the zonation of natural marshes in the area. Planted areas filled in within 2-3 months, and were able to prevent erosion. Planting of mangroves was not necessary, because natural recruitment usually became established within three years.

Maintenance action was recommended to prevent the natural 3-4 yr cycle of growth and die-back. Actions consisted of removing flotsom wrack (prolonged shading by wrack killed cordgrasses), creeping vines, and dead tops after seed set.

Benner, C. S., P. L. Knutson, R. A. Brochu, and A. K. Hurme. "Vegetative erosion control in an oligohaline environment, Currituck Sound, North Carolina." Wetlands 2 (1982): 105-117.

Nine species of coastal marsh plants were tested for their use in stabilizing a shoreline in an oligohaline bay in North Carolina. Plants were sprigged (0.9-m interval) along transects from the shore-berm out 30.5 m into the intertidal zone. Two side-by-side

transects of each species were planted in four adjacent, replicating plots on a block of bare beach. A similar and adjacent bare area was used as a control.

Only four species survived (*Typha latifolia*, *Phragmites australis*, *Juncus roemerianus*, and *Spartina alterniflora*). Once they became fairly established, they were joined by 20 volunteer species that increased the vegetative cover substantially. As plant cover increased, erosion of the area decreased. Additional plant cover led to accretion of sediments and expansion of the marsh. A balance between erosion and accretion was finally established, and a year's net gain or loss of sediments then varied as weather and water conditions varied. The bare control area only experienced erosion through the eight years of the study.

Berger, J. J. ed. Ecological Restoration in the San Francisco Bay Area. Berkeley, CA: Restoring the Earth, 1990.

This is an excellent compendium of restoration projects of all types of habitats in the counties bordering San Francisco Bay. Plus, it contains a directory (Appendix A) of people, companies, and organizations (including governmental) that are involved in restoration activities and regulations.

Habitats covered include forests, grasslands, coastal dunes and prairies, mined lands, creeks, lakes, and marshes. Other restorations focused on wildlife such as fish, butterflies, birds, and mammals, and some focused on native plants. Each restoration project was summarized as to location, beginning date, current status, site characteristics, purpose of the project, procedures used, results to date (1990), monitoring activities, future plans, support, budget, contact people, and volunteer needs.

Appendix E explains how to create a restoration directory including a database of projects. This is a useful guide that can be used for many different types of directories.

Bernstein, P. G. and R. L. Zepp, Jr. "Evaluation of selected wetland creation projects authorized through the Corps of Engineers Section 404 program." 80. City: U.S. Fish and Wildlife Service, Permits/Licenses Branch, Annapolis, MD, 1990.

The goal of this study was to evaluate the success of mitigation for lost wetlands. Each of 66 randomly chosen mitigation projects located in the Baltimore, Norfolk, and Philadelphia Districts of the U.S. Army Corps of Engineers were reviewed, visited and evaluated. A short written report was included for each.

Four projects of the 66 permitted had not been done (no mitigation required), 44 of the 62 remaining were deemed failures according to the permit specifications, 9 were deemed successes, and 9 were still under construction. Among the failures were partial success--projects where wetlands were established but were inadequate to fulfill the permit requirements.

The authors concluded the permitting system was losing wetlands under its current system of operation. It needed modification if it was to preserve our wetlands. One recurring problem was the lack of thorough planning. Rarely was an approved design for the mitigated wetland (to be created or restored) submitted at the time of the permit request. Recommended improvements for the system were: (1) Standardize permitting requirements for all projects requiring compensation. (2) Implement a tracking system for all permits requiring mitigation. (3) Require a greater than 1:1 replacement ratio for some critical

types of wetland habitat. (4) Require routine follow-up investigations including comparisons of pre-construction and post-creation data. (5) Enforce permit requirements with prosecution.

Blair, C. "Successful tidal wetland mitigation in Norfolk, VA." In: Coastal Wetlands., ed. H. S. Bolton. 463-476. New York: American Society of Civil Engineers, 1991.

In 1984, a 2.8 ha man-made *Spartina alterniflora* marsh (Monkey Bottom marsh) was created on a scrape-down area connected to Willoughby Bay in Norfolk, VA. This mitigation area had four lateral ditches that drained into one main canal that connected with the bay via a culvert. Planting of *S. alterniflora* was done on 2-ft centers and at elevations that ranged above and below the expected marsh establishment zone.

The mitigation marsh was evaluated as to its sediments, vegetation, invertebrates, fish and birds four years after its construction. A comparison of Monkey Bottom marsh with two local natural *S. alterniflora* marshes found no significant differences in density, cover and standing crop of *S. alterniflora* among the three. Fish and crabs were found in abundance, though not quantitatively compared. *Mugil* and *Callinectes* accounted for 55% and 20% of the catch by numbers, respectively, at Monkey Bottom. *Brevoortia Fundulus* and *Menidia* were the dominants at the natural marsh, where mullet were seen but not captured. Hard, soft and razor clams were the three main benthic organisms at Monkey Bottom, but polychaetes were also common. The control marsh had many fiddler crabs and nematodes. Bird utilization was similar for the marshes, with the exception of more suburban species frequenting the control marsh which was surrounded by a long-established residential neighborhood. The overall conclusion was that the mitigation marsh was a success, and was functioning as a typical estuarine marsh.

Bolton, H. S., ed. Coastal Wetlands. Coastlines of the World. New York: American Society of Civil Engineers, 1991.

This is a collection of papers presented at Coastal Zone '91 held in Long Beach, CA. Topics run from political and legal considerations of coastal wetlands to environmental design and monitoring considerations.

Bontje, M. P. "The application of science and engineering to restore a salt marsh." In: Proceedings of the fifteenth annual conference on wetlands restoration and creation. in Tampa, FL, edited by F. J. Webb, Jr. Tampa, FL: Hillsborough Community College, 16-23, 1988.

The report compares the use of a 63-acre man-made *Spartina alterniflora* marsh by birds, mammals and fish, to that of a 113-acre neighboring, *Phragmites communis* (common reed) marsh. The report also briefly describes the creation of the *Spartina* marsh.

The cordgrass marsh was established by clearing the common reed by spraying with glyphosate, excavating and grading the area by digging wide, gently sloping, canals. Excavated material was piled into raised berms for shrubs and trees. Once the surface was at the proper slope and elevation (0.15-0.45 ft below mean high water), *Spartina alterniflora* was seeded into the area. The canals were connected with Mills Creek of the Hackensack River. Within a few growing seasons the cordgrass was well established.

Birds, mammals, fish and water were sampled for 11 months. Monthly bird counts were made by walking or boating through the entire site for 1-hr. Mammals were trapped with bait for 24-hr each month. Tracks and burrows were also recorded. Fish were collected bimonthly by pulling a 3-m seine once for 15 m.

Based on this sampling regime, results showed twice the diversity and seven times the abundance of birds were using the *Spartina* marsh as were using the larger *Phragmites* marsh. Muskrat burrows were twice as dense in the *Spartina* marsh as in the *Phragmites* marsh. *Fundulus heteroclitus* was the common fish in both areas. Fish diversity was low, reportedly due to poor water quality in the adjacent rivers and creeks where diversity was also low. Benthos was three times as plentiful and twice as diverse in the *Spartina* marsh as in the *Phragmites* marsh. Water quality was nearly identical for the two marshes.

The goal to improve wildlife habitat in the area was achieved by removing the common reed and establishing a cordgrass marsh. The cordgrass marsh apparently developed quite quickly in the area which had minimal wave stress, proper elevation, and a good inundation-drainage design.

Bontje, M. P. "A successful salt marsh restoration in the New Jersey meadowlands." In: Proceedings of the eighteenth annual conference on wetlands restoration and creation. in Tampa, FL, edited by F. J. Webb, Jr. Tampa, FL: Hillsborough Community College, 5-16, 1991.

This article provided a detailed account of a salt marsh restoration/creation project in Lyndhurst, NJ. An old dredged material deposition site had become overgrown with *Phragmites australis*. Wetland creation required killing the reed, decreasing the elevation of the site to match marsh levels, improving drainage across the site, and planting and establishment of *Spartina alterniflora*.

Innovative techniques were described that solved unforeseen problems that developed during the project. Rodeo was used to kill the *Phragmites*, via two aerial sprayings and one hand spraying of recruits. Earth moving was done by backhoes and dump trucks, saving money which would have been spent on specialized equipment.

This was a 14 acre site, in which about 9 acres of salt marsh, 2 acres of tidal channels, and 3 acres of upland berm were established. The salt marsh was established on a 1% slope that drained about 80% at low tide; only a few inches of water remained in the channels. At high tide only the dike and a central ridge were not submerged. *Spartina alterniflora* was planted as peat pots (3-4 stems/pot) on 3-ft centers. Fertilizer was added to the holes before the peat pots were inserted. No significant mortality occurred, and plant growth was good.

No scientific studies were done to test for use of the marsh by aquatic or land animals, but increased use of the area by birds was noted.

Boyd, M. J. "Salt marsh faunas: colonization and monitoring." In: Wetland restoration and enhancement in California. in Hayward, CA, ed. M. Josselyn, Hayward, CA: University of California Sea Grant College Program, 110, 1982.

This paper presents an overview of the marsh fauna found in California coastal marshes, and presents the needs for monitoring changes in the fauna at a restoration site.

Several species of invertebrates and vertebrates are listed. Detecting significant differences in population densities of various species was expected to require extensive sampling--well beyond what would be logistically or financially possible given current constraints. But some such assessment is needed if a restoration project is to be judged successful or not. A suggestion, by a participant in the discussion, was made to possibly use an indicator species.

Broome, S. W. "Creation and restoration of tidal wetlands of the southeastern United States." In: Wetland Creation and Restoration: The Status of the Science. Part 1, Regional Review., eds. J. A. Kusler and M. E. Kentula. 37-72. Washington, D.C.: Island Press, Inc., 1989.

This chapter includes descriptions and brief discussions of many facets of tidal salt marsh restorations and creations beginning with marsh functions, then project plans, and finally an evaluation of the success of a project. Smooth cordgrass, *Spartina alterniflora*, is one of the most important plants used in salt marsh restorations, particularly in the intertidal zone along the Atlantic and Gulf coasts of the U.S. The functions of these marshes are many, but the major ones eliciting restoration activities are shoreline erosion control, sediment stabilization, and fisheries and wildlife habitat development or replacement. Careful planning is required for success in such projects. Attention must be paid to elevation at the site, water circulation into and through the area, protection from erosive wave action, and adverse actions by pests, herbivores and man's activities in and around the site. Timing of the planting, health of the transplants, and fertility of the soil are also important factors. Monitoring is important for documenting the success of a technique. A photographic record can be valuable. Future research is needed on site selection, design and preparation, on plant propagation and culturing techniques, and on documentation of marsh development. Information and advice are offered on most of these subjects, and are based on the author's many years of first hand experience and reading.

Broome, S. W., S. M. Rogers, Jr., and E. D. Seneca. "Shoreline erosion control using marsh vegetation and low-cost structures." (1992).

This report describes the use of smooth cordgrass, *Spartina alterniflora*, marsh to control shoreline erosion. Suitable sites had a substantial tidal range and a regular diurnal tidal rhythm, and a gentle slope in the intertidal area. They were in estuaries where salinity ranged between 5 and 35 ‰, wave action was light or reducible at least while plants are becoming established. Planting in the spring to obtain a full growing cycle the first year was recommended, as was the use of a time-released fertilizer. Selection of other plant species was recommended for elevations above the normal tidal zone.

Broome, S. W., E. D. Seneca, and W. W. Woodhouse, Jr. "Establishing brackish marshes on graded upland sites in North Carolina." Wetlands 2 (1982): 152-178.

The objectives of this study were to determine: 1) the most suitable marsh plant species to use in a marsh creation project; 2) each specie's elevation requirement; 3) the effectiveness of several fertilizers; and 4) which methods were most effective when more than one process was possible. The study was done at a Texasgulf Chemical Co. phosphate

mining site adjacent to Bond Creek, a tributary of the Pamlico River estuary near Aurora, NC. The site was a series of borrow pits that were graded to suitable elevation (matching a local natural marsh) and slope, and connected to Bond Creek. Diurnal tidal flows were minimal, wind being a dominant force of water height change.

A great variety of plantings and tests were done over three years. Because of the lack of tidal fluctuation in water levels, plant species were limited to narrow elevation bands. *Spartina alterniflora* ranged from 0.06 to 0.43 m above mean sea level (MSL). *S. patens*, and *S. cynosuroides* were limited to 0.18 to 0.43 m above MSL. For transplanting, greenhouse grown seedlings of *S. patens* and *S. cynosuroides* grew better than field dug plants, but no such difference was found for *S. alterniflora*. Survival was 80-100% when plants were planted at the proper elevations. Direct seeding was only partially successful; irregular water levels during germination and seedling growth were damaging.

Some fertilizers were of benefit, some were not, and some were detrimental. The application of fertilizer directly to the planting hole caused root burn. Residual fertilizer benefits could be found even into the second growing season. When all was working well, it took 2-3 yr for the planted marshes to equal the above-ground production of the natural marshes.

Broome, S. W., E. D. Seneca, and W. W. Woodhouse, Jr. "The effects of source, rate and placement of nitrogen and phosphorous fertilizers on growth of *Spartina alterniflora* transplants in North Carolina." *Estuaries* 6 (1983): 212-226.

This study evaluated the effects of fertilizer rate, type of fertilizer material, and application methods on growth and survival of *Spartina alterniflora* transplants. Changes in erosion rates of planted and unplanted shoreline were also noted. The test plots were along the Neuse River near Oriental, NC. Soil tests of the area showed no organic matter in the compacted sandy clay loam, a pH of 5.17, and low nutrient concentrations. Tidal fluctuation was wind dominated, and salinity varied from 5 to 23 ‰. A previously attempted planting of *S. alterniflora* in the area showed fertilization was necessary for survival and growth, and that surface applications of fertilizers were ineffective. Subsurface application of fertilizers was tested in a randomized complete blocks design. Osmocote (3-4 month release), Mag Amp (medium texture), ammonium sulfate, urea, urea-formaldehyde, diammonium phosphate, concentrated superphosphate, and rock phosphate were used. Results showed the Osmocote fertilized plants survived significantly better than the others, and grew fastest. Plants with Mag Amp were slower to get started, but were doing well after 11 weeks. Test results also showed that both N and P were needed for satisfactory growth, and that K was not needed (probably enough was being supplied in the estuarine water). Phosphorous applied at a rate of 49 kg/ha produced maximum growth when adequate N was supplied. The maximum rate of N applied was 224 kg/ha, and growth continued to improve to this maximum. The most efficient and economical method of fertilizing was the time-released form of $\text{NH}_4\text{-N}$ combined with concentrated superphosphate, both applied underground but not necessarily in the planting hole. Some carry over of fertilizers was noted during the second year of growth. A top-dressing of N and P fertilizers after a stand has become established was said to be beneficial. Roots have developed which can take up the nutrients before they are washed away. With the

establishment of the plants, sediment accumulated along the landward margin of the stand. This build-up stimulated additional plant growth.

Broome, S. W., E. D. Seneca, and W. W. Woodhouse, Jr. "Long-term growth and development of transplants of the salt-marsh grass *Spartina alterniflora*." Estuaries 9 (1986): 63-74.

This paper reports about the effect of transplant spacing on growth and development of a *Spartina alterniflora* marsh, following the marsh development for 10 years, and comparing it with a nearby natural marsh. The test site was a Pine Knoll Shores, NC, on the barrier island of Bogue Banks. *Spartina alterniflora* plants of field (dug 11 km away) and greenhouse origins were planted on 45-, 60-, and 90-cm centers in three 15x12-m plots that were set in a randomized blocks design. Above-ground vegetative characteristics were measured from several 0.25 m² quadrats taken each October. Below-ground biomass was usually sampled by taking cores.

Results showed that 45- and 60-cm treatments were better for establishing a marsh in a marginal environment. Transplant survival rates at the end of the first growing season were 72%, 69% and 49% for the 45-, 60- and 90-cm treatments, respectively. Above-ground vegetation among the treatments was indistinguishable after two growing seasons, except that two of the 90-cm blocks had eroded away by the second year's monitoring. By the end of the fifth year, the planted marsh was as densely vegetated (above 600 stems/m²) as the natural control marsh that was 200 m to the west. Below-ground biomass increased fastest in the 45-cm treatment. It reached an equilibrium around 2 kg/m² by the end of the third year, and was matching the natural marsh.

The failure of the two 90-cm spacing blocks reaffirms the need to plan a marsh planting based on the harshness of the environment. A thicker planting will be more effective in breaking the erosive action of waves, but of course, the cost will be higher as many more plants will be required.

Broome, S. W., E. D. Seneca, and W. W. Woodhouse, Jr. "Tidal salt marsh restoration." Aquatic Botany 32 (1988): 1-22.

This article summarized the techniques to restore *Spartina alterniflora* salt marshes in the Southeastern U.S. Results showed that the site must be at the proper elevation--between MSL and MHW, and have a gentle slope--less than 10%, preferably less than 5%. Although salt marshes occurred in a variety of substrates, a sandy substrate was easiest for restoration planting; fertilizer was usually needed as sandy substrates were often nutrient poor.

Vegetation was usually restored by transplanting sprigs or plugs, or by seeding. Young plants were commonly available from nearby, healthy, natural donor marshes, or from nursery stocks. Labor costs ran about 100 man-hours per hectare for manual planting of sprigs on 1-m centers, and half of that for mechanical planting. Timing of planting was important, and was best scheduled for early in the growing season (April-June). Restoration of an oil-contaminated marsh should not proceed for at least six months even if the oil was removed.

Because salt marshes are very valuable habitats that serve many functions, the authors recommended that policies to protect marshes and restore damaged ones should be

strengthened. Restoration work should be guaranteed, and replanting should be required when deficiencies are detected during monitoring. Each project should be monitored for 3-5 years to determine its success. Documentation should be made of successes and failures, with reasons for failure being documented and discussed so future projects would not repeat mistakes.

Broome, S. W., W. W. Woodhouse, Jr., and E. D. Seneca. "The relationship of mineral nutrients to growth of *Spartina alterniflora* in North Carolina. II. The effects of N, P, and Fe fertilizers." Soil Science Society American Proceedings 39 (1975): 301-307.

Little is known about the effect of mineral nutrition on *Spartina alterniflora*. This study was initiated to evaluate the influence of N, P, and Fe on primary productivity by applying these nutrients to plots in natural marshes and on *S. alterniflora* seeded and transplanted on dredged material. In a marsh growing on a sandy substrate, additions of N alone increased yields of aboveground shoots significantly, and when P was also added, the yield increased about threefold. In a marsh growing on finer textured sediments, N fertilizer doubled the yield of short *S. alterniflora*, but there was no response to P. There was no growth response to applications of Fe to support previous speculation that iron nutrition might be a particularly important factor causing the chlorotic appearance of short *Spartina* and reducing its productivity. The results indicate that primary productivity of some *S. alterniflora* marshes is limited by the availability of N. When N is added, lack of P may become the factor limiting growth, particularly when the substrate is coarse in texture, indicating the importance of sediment as a factor in P supply to *S. alterniflora*. Lack of N is apparently one of a combination of factors which is responsible for producing the short form of *S. alterniflora*. The fact that N and P are the limiting factors in growth of *S. alterniflora* in some salt marshes has several ecological implications. Marshes may be acting as buffers for estuarine systems by providing sinks for excess nutrients from such sources as sewage and land runoff. The excess nutrients would produce increased growth of *S. alterniflora* thus providing and increased supply of food energy and nutrients to the detritus food chain of the estuary rather than altering energy pathways. The ability of salt marshes to utilize excess N and P may be important in managing estuarine systems. Nutrient-rich waste effluent dumped in marshes would have less impact on estuaries than that dumped in open estuarine waters.

Fertilization may be beneficial in propagating *S. alterniflora* on dredged material since establishing a full vegetative cover rapidly is important. Applications of N and P fertilizers enhanced growth of seedlings and transplants, but the response of *S. alterniflora* to fertilizer will depend on the inherent fertility of the substrate material.

Callaway, J. C. and M. N. Josselyn. "The introduction and spread of smooth cordgrass (*Spartina alterniflora*) in south San Francisco Bay." Estuaries 15 (1992): 218-226.

Spartina alterniflora was first introduced into south San Francisco Bay in the 1970's. Since that time it has spread to new areas within the south bay and is especially well established at four sites. The spread of this introduced species was evaluated by comparing its vegetative and reproductive characteristics to the native cordgrass, *Spartina foliosa*. The characters studied were intertidal distribution, phenology, aboveground and below-ground biomass, growth rates, seed production, and germination rates. *Spartina alterniflora* has a

wider intertidal distribution and was more productive than the native cordgrass in all aspects studied. These results indicate that the introduced species becomes established more readily in new areas than the native species, and once established, *S. alterniflora* spreads more rapidly vegetatively than *S. foliosa*, thus out-competing it. *S. alterniflora* is likely to continue to spread to new areas in the bay and displace the native species. This introduced species may also affect sedimentation dynamics, available detritus, benthic algal production, wrack deposition and disturbance, habitat structure for native wetland animals, benthic invertebrate populations, and shorebird and wading bird foraging areas due to its different physical and ecological characteristics. The evaluation of the extent of these impacts, however, can only be made after *S. alterniflora* has become widely established and very difficult to eradicate.

Cammen, L. M. "Accumulation rate and turnover time of organic carbon in a salt marsh sediment." Limnology and Oceanography 20 (1975): 1012-1015.

Organic carbon in the sediments was measured near Drum Inlet, North Carolina in a natural *Spartina alterniflora* salt marsh and on both unvegetated dredged material and dredged material planted with *S. alterniflora*. The planted marsh was about 1 year old at the time of sampling. Grain size analyses indicated that sediment texture was sandy and similar among the three sites. Above-ground *Spartina* biomass in the planted marsh was comparable to the natural marsh, but below-ground biomass in the planted marsh was between 19% (unfertilized) and 43% (fertilized) of the natural marsh. The annual accumulation rates of organic carbon (g/m²) in the upper 13 cm of sediment were 80.3 in the bare sediment, 87.0 in the planted marsh, and 96.8 in the planted and fertilized marsh. Organic carbon in the natural marsh sediment was 362.7 g/m². The data suggest that the dredged sediments would be comparable to the natural marsh sediment within 3-5 years, and that the presence of *Spartina alterniflora* did not appreciably increase the rate of organic matter accumulation. Bare sediment had accumulated almost as much organic carbon as planted dredged material, and the increases were attributed to detrital matter carried by tides and to benthic algae.

Cammen, L. M. "Abundance and production of macroinvertebrates from natural and artificially established salt marshes in North Carolina." American Midland Naturalist 96 (1976): 487-493.

Abundances of macro-infauna taken in core samples were studied from natural, man-made *Spartina alterniflora* marshes, and adjacent bare areas at two locations, one near Drum Inlet and one near Snow's Cut, North Carolina. At each location, a permanent set of sites were established along transects. The planted area transect ran from the upper extent of the cordgrass down the elevation gradient to the middle of the tidal creek at Drum Inlet, and to just above MLW at Snow's Cut. Transects through the bare areas ran parallel to the transect through the planted area. A similar transect crossed each natural marsh. Replicate samples, 1 m apart, were taken at each site with a piston corer that covered a surface area of 70.9 cm² and reached a depth of 13 cm.

Macro-infauna differed with the location. At Drum Inlet, insect larvae (Dolichopodidae) dominated the fauna in the bare and planted areas, while polychaetes (*Heteromastus filiformis* and *Capitella capitata*) dominated the natural marsh. At Snow's

Cut, polychaetes (*Laeonereis culveri*, *H. filiformis* and *C. capitata*) dominated the fauna in the bare area, while amphipods (*Lepidactylus dytiscus* and *Gammarus palustis*) and Dolichopodidae larvae dominated the fauna in the planted area. The natural marsh macro-infauna at Snow's Cut was dominated by polychaetes (*L. culveri* and *Nereis succinea*), an isopod (*Cyathura polita*), and a bivalve mollusk (*Arcuatula* (= *Modiolus*) *demissa*).

Annual macrofaunal production (excluding *Uca spp.*) was estimated to be equal to the maximum standing stock of the insect population, and twice the average standing stock of the rest of the macro-infauna. Production values (g dry wt./m²/yr) at Drum Inlet were 1.4 for the bare area, 1.2 for the transplanted area, 7.7 for the natural marsh, and 23-34 in the adjacent tidal creek bottom. At Snow's Cut production values were 13.8 (bare), 0.5 (transplanted), 5.7 (natural marsh), and 8-17 (tidal creek).

The low faunal diversity at the sites was partly attributed to infrequent sampling. It is also possible that the habitat was not developed enough at that time to support other less opportunistic species. Winter sampling was not conducted, and would have been useful for improving accuracy of annual production estimates.

Cammen, L. M. "Macroinvertebrate colonization of *Spartina* marshes artificially established on dredge spoil." *Estuarine and Coastal Marine Science* 4 (1976): 357-372.

This study investigated population dynamics of macro-infauna taken in core samples from natural and man-made *Spartina alterniflora* marshes, and adjacent bare areas. Two locations were studied, one near Drum Inlet and one near Snow's Cut, North Carolina. At each location, replicate core samples were taken at permanent sites that were established along transects that crossed the elevational plane from subtidal to the landward margin of the marsh.

Particle size analysis showed that the dredged material at the Drum Inlet was very similar to the natural marsh sediment. The dredged material at the bare and planted areas near Snow's Cut were similar to that at Drum Inlet, but differed from the natural marsh nearby. The natural marsh at Snow's Cut had much higher coarse-sand and silt-clay fractions.

The above-ground and below-ground biomass of *Spartina* differed for planted and natural marshes. The above-ground biomass values in g/m² were 709 (planted) and 1056 (natural) at Drum Inlet, and 984 (planted) and 637 (natural) at Snow's Cut. At each location, the planted marshes had only half the below-ground biomass that the natural marshes had, 605 and 2371 (planted) and 3169 and 4966 (natural) and the respective locations.

Macro-infauna differed between locations and among areas at the same location. At Drum Inlet, the macrofauna in the bare and planted areas had similar biomass, species composition, and numbers of taxa per sample. The bare area had higher densities of organisms during three of the five samplings. The natural marsh had a higher density, biomass, and diversity for two of the three dates when all three areas were sampled. At Snow's Cut, the macrofauna in the bare area consistently had greater biomass and density per date sampled. The natural marsh was only sampled once at Snow's Cut. The macro-infauna in the natural marsh had a greater density and diversity than the dredged material sites, but it had a biomass slightly less than that of the bare area and slightly greater than that of the planted area. Neither sediment characteristics nor the presence of *Spartina* were deemed

sufficient causes for these differences found in the fauna. Elevation levels, however, were different and were likely the cause of the infaunal differences.

Sufficient support for separating causes of differences in macro-infauna and in establishing statistically significant differences would have required an increase in sampling. Although the abundances of *Uca spp.* were not included, their presence could have impacted infauna abundances.

Cammen, L. M. "The macro-infauna of a North Carolina salt marsh." American Midland Naturalist 102 (1979): 244-253.

The community dynamics of the macro-infauna of a healthy *Spartina alterniflora* marsh near Beaufort, NC, were studied for one year. Monthly sampling involved taking at least 6 cores randomly in the marsh. These were processed through a 0.8 x 0.6-mm mesh sieve; the animals, macro-detritus and roots were caught and preserved in formalin. Greatest abundance was found during the late winter and early spring. Lowest abundance was during the summer and early autumn. Numbers of individuals ranged from 2200 to 15500 /m². There were 32 taxa identified in the samples, but only four dominated the community. These were: *Nereis succinea*, *Streblospio benedicti*, Capitellidae and Oligochaeta. They were present in over 93% of the cores. They accounted for 96% of the number of individuals caught, and averaged 85% of the monthly biomass. The low summer biomass was thought to be due to predation by juvenile fish and to mortality due to spring spawning. These taxa were also among the dominants in several other marshes in NC. The high numbers of infauna suggest that most of the marsh surface sediment and detritus is eaten and processed each year.

Chabreck, R. H. "Creation, restoration, and enhancement of marshes of the northcentral Gulf coast." In: Wetland creation and restoration: The status of the science. Volume I: Regional overviews. EPA/600/3-89/038., eds. J. A. Kusler and M. E. Kentula. 127-144. Corvallis, OR: U.S. Environmental Protection Agency, 1989.

Coastal marshes of the northcentral coast of the Gulf of Mexico encompass an estimated 1.2 million hectares. This is about half of all coastal marshes in the U.S., excluding Alaska. These coastal marshes are being threatened by subsidence, sea level rise, and erosion by wind generated waves. Efforts to protect and enhance the marshes in the northcentral Gulf involve use of dredged material and river water with its sediment. Weirs, dikes, and levees are also used around freshwater and brackish water marshes to protect them from saltwater intrusion. Information is given about things to consider when attempting to create or restore coastal marshes. Location, topography, hydrology, type of substrate, salinity, and wind and wave climates are all important factors to be considered in the planning process. Monitoring of a project is suggested to be a two level process. The first level is just a qualitative look to see if the project is developing as planned. The second level is a quantitative investigation of the project to assess the level of success attained.

Research is needed for developing marshes using diverted Mississippi River water. Opening channels in the banks of the larger delta channel of the river could allow mini-delta formation which could be planted or allowed to vegetate naturally. Dredged material use in marsh enhancement needs to be researched. Thin-layer deposition of dredged

material may improve marshes that are deteriorating because of subsidence, but methods for this need to be developed.

Cobb, R. A. "Mitigation evaluation study for the south Texas coast, 1975-1986." City: Corpus Christi State University and U.S. Fish and Wildlife Service, 1987.

This report presents an evaluation of the level of acceptance, implementation, and success of the U.S. Fish and Wildlife Service's recommendations made to preserve fish and wildlife habitat, or to mitigate its loss along the south Texas coast. For 59 permitted waterfront projects, the USFWS recommendations were unconditionally accepted in 78%, rejected in 5%, and modified in 16%; 1% were unresolved. However, most of the accepted recommendations were those to avoid impact to wetlands or to assure adequate water quality within excavated canals. Projects requiring habitat compensation had more modifications, and they had more non-compliance by the permittee. Non-compliance with permit conditions was found at 31% of the sites inspected--78% of this was unfulfilled mitigation requirements and 22% was additional work performed beyond that permitted.

In the proposed mitigation work involving marsh restoration or creation there was substantial variation in describing the substrate elevations where *Spartina alterniflora* was to be planted. Rarely were details of the planting (stems/m², as-planted maps, etc.) provided in a report. Monitoring was never done by the permittee, and only once by the Texas General Land Office and Corps of Engineers. Completion reports were rarely filed. Only one performance bond was required. Of the 15 projects investigated, 6 were deemed failures, 4 were partial successes, and 5 were fully successful according to the permit requirements.

Many of the usual reasons for failure of transplanted marshes were found among these failed projects including: (1) no soil conditioning, (2) incorrect elevation, (3) excessive slope, (4) excessive wave action and inadequate protection, (5) bad weather, (6) poor drainage and excessive salinity, (7) human disturbance, and (8) improper site preparation. In a few cases where failure was due to insufficient plant cover within the specified time (two years), the marshes have since developed sufficiently to be rated successful.

Although almost 90% of the recommendations proposed by the USFWS were incorporated in the permits, the overall final results after implementation yielded a net loss of wetlands--only 50% were recovered through mitigation. The report offers 21 recommendations to enhance the chances of successful marsh restoration. And 21 more are given for seagrass restoration projects.

Courtney, F. X., S. A. Peck, and M. O. Hall. "Post-harvest recovery of a donor *Spartina alterniflora* marsh." In: Proceedings of the eighteenth annual conference on wetlands restoration and creation in Tampa, FL, edited by F. J. Webb, Jr., Tampa, FL: Hillsborough Community College, 23-31, 1991.

A previously transplanted salt marsh at Spoil Island 2-D in Hillsborough Bay, Tampa Bay, FL, was used as a donor site for 300 m² of *Spartina alterniflora*. The harvest was made from 1/2 m or 1 m wide transects that were oriented either parallel or perpendicular to the shoreline. The cordgrass was hand pulled from the substrate. One half of the 24 harvest transects were fertilized shortly after harvest. After 17 months, mean culm

densities in donor transects were similar to those of the undisturbed marsh. Fertilizer made shoots more robust, but not more numerous. Limited harvest of planting material appears to cause only temporary damage to the vegetation of the donor marsh. No studies were done to assess changes in marsh utilization by aquatic fauna.

Covin, J. D. and J. B. Zedler. "Nitrogen effects on *Spartina foliosa* and *Salicornia virginica* in the salt marsh at Tijuana Estuary, California." Wetlands 8 (1988): 51-66.

Nitrogen effects were examined by experimentally enriching plots of pure *Spartina foliosa* and mixed *Spartina-Salicornia virginica* at Tijuana Estuary, California. Even with large inputs of organic nitrogen from sewage spills during the experimental period, plants responded to experimental urea enrichment. In pure plots, the addition of nitrogen increased *Spartina* growth (as measured by total stem length and August biomass) and foliar nitrogen (TKN) concentration. However, this effect is eliminated if increased foliar nitrogen stimulates insect predation resulting in heavy plant mortality. In mixed plots, enrichment had no apparent effect on *Spartina* but increased the growth of *Salicornia*. The lack of an enrichment effect on *Spartina* may be due to underground rhizomes and roots of *Salicornia* continuing to take up nitrogen through connections with plants outside the removal plots. The experimental removal of *Salicornia* from mixed stands increased *Spartina* production, but removal of *Spartina* did not affect *Salicornia*. Thus, there was a strong competitive effect. In mixed stands, urea additions are detrimental to *Spartina* because *Salicornia* has a greater response capacity. *Salicornia* is a superior competitor for nitrogen and checks the growth of *Spartina* in enriched and unenriched conditions. These experiments show that nitrogen, through its complex effects on growth and competition, is an important cause of spatial and temporal variability seen in long term observations of *Spartina*.

Craft, C. B., S. W. Broome, and E. D. Seneca. "Nitrogen, phosphorus and organic carbon pools in natural and transplanted marsh soils." Estuaries 11 (1988): 272-280.

The objectives of this study were to compare the nutrient and organic pools of transplanted and natural coastal saline marshes. Marsh sediments were studied in five created and nearby natural marshes in diverse locations in North Carolina. A couple of the marshes compared differed in their vegetation. The created marshes ranged in age from 1 to 15 years. Sediments were sampled using an 8.5 cm diameter corer. Ten to 20 cores, 30 cm deep, were taken randomly from each marsh. Macro organic matter (MOM) was separated from the sediments by sieving on a 2-mm mesh sieve. MOM and soil nutrient reservoirs were smaller in transplanted marshes than in the nearby natural marshes. About 12% and 20% of the net primary production of emergent vegetation was buried in sediments of the regularly flooded and irregularly flooded transplanted marshes, respectively. MOM pools developed rapidly in transplanted marshes, and were expected to match the pools in natural marshes in 15-30 years. Soil nutrient pools in transplanted marshes, however, developed slowly and were expected to take much longer to match those of natural marshes.

Craft, C. B., E. D. Seneca, and S. W. Broome. "Porewater Chemistry of Natural and Created Marsh Soils." Journal of Experimental Marine Biology and Ecology 152 (1991): 187-200.

Physical and chemical properties of soils and porewaters were compared from a natural marsh and a man-made marsh that was created from graded down upland. Porewaters were sampled monthly for a year from established wells set in the marshes. Water level, temperature, salinity, DO, pH, Eh, Fe, Mn, organic C, N, P, NH₄, NO₃, and PO₄ concentrations were monitored in the pore water.

The created marsh soil had about 1% organic matter compared with about 50% for the natural marsh. DO, Eh, Fe, Mn and NO₃-N concentrations were significantly higher in the created marsh pore waters, but the dissolved organic C and N, NH₄-N, PO₄-P, and pH concentrations were lower. Even five years after creation of a saline estuarine marsh, the soil characteristics remained typical of the upland soil from which it was made.

This study suggests that we should not expect mitigated wetlands created on upland soils to gain characteristics of a natural marsh within even a decade. The authors recommended avoiding damage to natural wetlands because wetland soils are not readily replaced.

Crewz, D. W. and R. R. Lewis, III. "An evaluation of historical attempts to establish emergent vegetation in marine wetlands in Florida." 113 pp. Florida Sea Grant, Univ. of Florida,, 1991.

The authors' objectives were: (1) to compile a database of past marine wetland projects, (2) to then survey the sites for elevation and plant cover based on a random selection of sites from the database, and (3) to provide guidelines based on the observations made at the sites. These guidelines should help increase the success rate of marine wetland creation projects where emergent vegetation is planted. Objectives 1 and 2 were dismissed due to lack of available information and lack of access.

For objective 3, 33 sites were visited where salt marsh or mangroves had been planted. The sites were chosen based on accessibility (physical and legal), accuracy of site location information, species planted, and geographic location. All sites were visited in 1986. Twenty sites had been planted with *Spartina alterniflora* along with other species. Plant densities were measured along transects that extended from the seaward edge to the upland edge of the marsh or mangrove area. Transects were spread about 25 m apart.

Results of the survey showed that marsh establishment was at least partially successful in 65% of the cases. Planted *Spartina alterniflora* survived from +0.2 - +0.6 m NGVD. Older natural marshes nearby ranged down to -0.1 m NGVD. Marsh failures were attributed to: (1) poor design and planning, (2) poor planting technique, (3) poor monitoring and remedial action, and (4) insufficient regulatory review. Critical factors found for establishing a salt marsh were: elevation, slope, drainage, substrate, plant selection, installation techniques, fetch, wave climate, marine connection, and elimination of pests from the site (including humans). Monitoring the development of a planted marsh was recommended because problems could be quickly identified and corrected.

The authors commented that despite advances in planting techniques, continued poor planning and planting led to failures of mitigated marshes. It was highly recommended that degradation of natural marshes should be avoided whenever possible. It was also

recommended that a strong monitoring program should be part of any wetland restoration or creation project.

Earhart, H. G. and E. W. Garbisch, Jr. "Habitat development utilizing dredged material at Barren Island, Dorchester County, Maryland." Wetlands 3 (1983): 108-119.

The objectives of this project were to stabilize a low island created by deposition of dredged material by planting *Spartina alterniflora* and to create a habitat for fish and wildlife. The project was done in the fall of 1981, in Chesapeake Bay just off the northeastern tip of Barren Island, Dorchester County, Maryland. About 135,831 m³ of fine-grained sediments were hydraulically deposited in a single unconfined location. The discharge pipe remained at a site until the water depth reached 1.8 m above MLW, then the outlet was moved to an adjacent lower portion of the site. The dredge material was allowed to consolidate for about five months (Nov. - April) before *Spartina alterniflora* was seeded over 8.4 ha at a rate of 96 seeds/m². The seeded area was fertilized a month later. Peat pots of *Spartina patens* were planted on a 0.6-m grid over about 2 ha at elevations above the *S. alterniflora*. During the winter about 460 m³ of oyster shell was placed in the center of the site to create nesting habitat for the least tern.

After eight months of growth, the shell island was nearly surrounded by 4.5 ha of flowering *Spartina alterniflora* at elevations ranging from 0.2 - 0.5 m above MLW, and 1.6 ha of *Spartina patens* at 0.4 - 0.7 m above MLW. A 4.1-ha pond, a 1.9-ha stretch of bare non-vegetated flatland, and a 0.1-ha area of oyster shell (tern nesting habitat) were also created.

This project required careful planning and diligence by the project team to keep the project on target. Constant monitoring by the project team was needed to keep the dredged material directed to the correct place. The project succeeded in creating *Spartina alterniflora* and *S. patens* marsh, plus some non-vegetated tern nesting habitat, but it did not create as much as was planned. When the project was completed, 69% of the designed marsh had become established. This paper provided a good comparison of project construction theory and construction reality, and it showed that even with a well planned design, circumstances during the construction phase can control the final success or failure of a project.

Earhart, H. G. and E. W. Garbisch, Jr. "Beneficial uses of dredged materials at Barren Island, Dorchester County, Maryland." In: Proceedings of the thirteenth annual conference on wetlands restoration and creation, in Hillsborough, FL, edited by F. J. Webb, Jr., Hillsborough, FL: Hillsborough Community College, 75-85, 1986.

Spartina alterniflora stems and clumps have been planted in dredged material to stabilize the material and establish salt marsh, but planting of large areas with sprigs of cordgrass can be very expensive. This paper describes a successful creation of 1.6 ha of *S. alterniflora* marsh using seeds. Judicious placement of the dredged material to specified elevations was required. This entailed constant monitoring of the additions of dredged material to the area. *Spartina alterniflora* marsh was made by broadcasting seed from an all-terrain-vehicle (ATV) during low tide over 5.3 ha. Broadcasting was followed by cultivating the substrate and seed by dragging a spiked metal mesh over it at low tide using the ATV. The site was fertilized four times during the summer. Seeding and fertilization

costs were only about one fifth those of single stem transplanting. *Ruppia maritima* voluntarily invaded the shallow waters between dredged material islands, establishing about 2.8 ha of *Ruppia maritima* beds. Natural beds were nearby.

Eleuterius, L. N. and J. I. Gill. "Long-term observations on seagrass beds and salt marsh established from transplants." In: Proceedings of the eighth annual conference on wetlands restoration and creation in Tampa, FL., edited by R. H. Stovall, Tampa, FL.: Hillsborough Community College, 74-86, 1981.

This paper reviews seagrass and saltmarsh transplant projects that were initiated five to ten years earlier in Mississippi. The projects involved transplanting a wide variety of seagrass and salt marsh species. In seagrass projects, *Halodule beaudettei*, *Thalassia testudinum* and *Cymodocea manatorum* were transplanted. Only about 30% of the transplants survived at the end of one year, but 80% of these survivors had begun spreading. *Halodule beaudettei* survived best, and spread rapidly. As the beds grew, they constantly changed shape. At these particular test sites the seagrass beds spread only westward, leaving their original planting sites empty as the eastern shoots died. This migration occurred within only a few growing seasons and was attributed to the predominant westward current in the area.

In salt marsh studies, growth and spread of transplanted *Panicum repens*, *Spartina alterniflora*, *Juncus roemerianus*, *Spartina cynosuroides*, *Spartina patens*, *Distichlis spicata*, and *Phragmites communis* were measured over various periods up to 10 years. Propagation trials showed that preliminary rooting of these marsh plants in peat pots was unnecessary. Individual shoots successfully grew into stands in 60 and 80% of the marshes transplanted from November to February. In random plot tests *Panicum repens*, *Spartina alterniflora*, *Spartina patens*, and *Distichlis spicata* formed closed stands within a year. *Spartina cynosuroides* and *Phragmites communis* survived but did not spread during the three years of the tests, apparently because they were out competed by other plants. *Juncus roemerianus* was a slow grower and spreader, but it continued to spread even when crowded by other species. *Juncus* formed closed stands in five years when planted on 1.2-m centers. Stands of each species were found to move due to competition with each other and to the local changes in environmental conditions. After about seven years the mosaic pattern of the natural saltmarshes in the area was attained at the test site.

In another study of transplanted *Spartina alterniflora*, all transplants in the lower elevations died due to wave action and sediment erosion. The transplants in the upper zone survived and flourished. Observations of the stand eight years later found *Spartina alterniflora* had invaded the lower zone and was now growing well, even at the lowest elevation that had originally been planted. This spread extended 15 m down the intertidal slope in about eight years. From this observation, a transplanting principle was formed: plant the upper portion of the intertidal zone and allow the cordgrass to invade the lower portion on its own.

Espey Huston & Associates Inc. "Monitoring of transplanted *Spartina alterniflora* on an unconfined dredged material disposal site, Chocolate Bay, Texas." Espey, Huston and Associates, Inc. DACW64-87-D-0002, 1988.

A *Spartina alterniflora* marsh was planted on 8.09 ha of unconfined dredged material in Chocolate Bay of the Galveston Bay System, Texas. Sprigs were obtained from a nearby marsh (control marsh) in Halls Lake, and planted on 1-m centers in June 1983.

Sampling to monitor marsh development was performed March - August and October 1984, April, June and October 1985, April, July and October 1986, and October 1987. Percent cover, stem density and height, and above-ground biomass were measured at 8 sites along each of 3 elevational transects through each marsh.

By 1986, the planted marsh had increased to an average of 79% cover and 382 stems/m² that averaged 29 cm in height. The control marsh had fallen to an average of 64% cover and 297 stems/m² that averaged 22 cm high. In October 1987, average cover was down slightly at 66% and 44% for the planted and control marshes respectively. Total number of stems averaged 436 and 348 stems/m², 37 and 34 cm high, respectively.

Surveys made for wildlife found periwinkles, fiddler crabs, oysters, and birds to be common in both marshes. Observations on marsh use, however, were very limited.

Espey Huston & Associates Inc. "Monitoring of transplanted *Spartina alterniflora* on an unconfined dredged material disposal site, Pelican Spit, Galveston Bay, Texas." Espey, Huston & Associates, Inc. DACW64-89-D-0003, 1990.

This report describes the results of a salt marsh creation. In April, 1987, about 2.6 ha of shoreline were planted with *Spartina alterniflora* on 1-m centers on recently deposited unconsolidated dredged material at Pelican Spit in Galveston Bay, Texas. Planting was done between +0.5 and +0.9 m MLW. Planted and nearby donor (control) marshes were monitored in Nov. 1987, Nov. 1988 and Oct. 1989 for vegetative growth and general marsh development.

The planted marsh increased to 91, 258 and 210 stems/m² compared with the 405, 145 and 150 stems/m² in the control marsh in 1987, 1988 and 1989, respectively. Marsh periwinkles (*Littorina irrorata*) were the only typical marsh invertebrates observed at the planted marsh, while fiddler crabs, blue crabs and periwinkles were found at the control marsh. Sampling for invertebrates was only qualitative.

Faber, P., A. Shepherd, and P. Williams. "Monitoring a tidal restoration site in San Francisco Bay- the Muzzi Marsh." In: Urban wetlands: proceedings of the national wetland symposium. in Oakland, CA, eds. J. A. Kusler, S. Daly, and G. Brooks, Oakland, CA: Assoc. State Wetland Managers, 331-335, 1988.

Observations on tidal channel formation, sedimentation, and revegetation are reported for the Muzzi Marsh at Corte Madera, Marin County, California. A 53-ha portion of the marsh was opened to tidal flows in 1976 by breaching the seaward dikes. Revegetation was slow, and was thought to be due to a lack of tidal flows to the landward portion of the marsh. In 1981 additional channels were dug to increase tidal flows to the interior and peripheral areas. Revegetation rates increased, and a substantial coverage of the flats by *Salicornia virginica* and of the channel banks by *Spartina foliosa* developed. Seed source was a nearby natural marsh.

Faber, P. M. "The Muzzi marsh, Corte Madera, California: long-term observations of a restored marsh in San Francisco Bay." In: Coastal Wetlands., ed. H. S. Bolton. 424-438. New York: American Society of Civil Engineers, 1991.

This article reports on observations made of drainage channel formation, sedimentation, and revegetation of the 53-ha portion of the Muzzi Marsh at Corte Madera, California, on San Francisco Bay. The Muzzi Marsh was a 81 ha natural coastal marsh that was diked in 1959. Subsequently, the marsh dried out, killing all the marsh vegetation. In 1976, 28 ha were further diked to retain dredged material, but another 53 ha were opened for tidal circulation by breaching the dikes in two areas. Periodic observations and studies showed the development and spread of drainage channels across the flat plain. The deposition of sediments was greatest just outside these channels. *Salicornia virginica* and *Spartina foliosa* became established fairly rapidly from the water born seeds that were generated by the plants growing on the bayside of the dikes and in the 27-ha natural marsh just to the north of the Muzzi marsh. There was constant competition between *Salicornia virginica* and *Spartina foliosa* along the MHW elevation. The competition was influenced by yearly rainfall. In wetter years *Spartina foliosa* became established at slightly higher elevations, but in drier years *Salicornia virginica* invaded lower elevations.

Florida Department of Environmental Regulation. "Report on the effectiveness of permitted mitigation." Florida Department of Environmental Regulation. 1991.

As of 1984, any alteration of a wetland area in Florida required a permit from the Florida Department of Environmental Regulation (FDER). Between January 1, 1985 and December 6, 1990, FDER issued 1,262 permits involving a loss of 3,305 acres and a preservation of 7,587 acres of natural wetlands. The permits also required creation of 3,345 acres and enhancement of 7,301 acres of wetlands. An evaluation was made of the success rate of the mitigation process.

A sample of the permits was drawn which included about 10% from each wetland category. Each project was examined for compliance with permit specifications, and for ecological success. Compliance was poor. Only 6% of 63 examined projects were in full compliance, and 34% of the mitigation projects had not even been attempted. Ecological success among the projects being constructed was projected to be about 27%. Success rates were lower (12%) for freshwater projects than for coastal marsh projects (45%).

As FDER concluded, the permitting policy needed revising if the wetlands were to be preserved, or "no net loss" were to be achieved. Permits that were granted needed to be monitored and enforced. Such changes would do much to safeguard wetlands.

Fowler, B. K., G. R. Hardaway, G. R. Thomas, C. L. Hill, J. E. Frye, and N. A. Ivison. "Vegetative growth patterns in planted marshes of the vegetative erosion control project." In: Proceedings of the twelfth annual conference on wetlands restoration and creation. in Tampa, FL, edited by F. J. Webb, Jr. Tampa, FL: Hillsborough Community College, 110-120, 1985.

This paper reports some of the results of the Vegetative Erosion Control Project, a study of 24 marsh plantings in the Chesapeake Bay system in Virginia. Sites were chosen to maximize diversity of conditions, mainly fetch and direction faced. Seven sites were classified as having low wave energy (average fetch exposure < 1.8 km), 10 sites were

classified as having medium wave energy (average fetch exposure from 1.8 to 9.2 km), and seven sites were classified as having high wave energy (average fetch exposure > 9.2 km). *Spartina alterniflora* was planted on 0.5-m centers from MHW to just below MSL. *Spartina patens* was planted above *S. alterniflora* at a few sites. About 30 ml of Osmocote fertilizer having a 14-5.2-11.6 slow release formula, was placed in each hole just before each culm was planted. Osmocote was scattered over the marsh surfaces one or twice each summer to encourage vigorous growth of vegetation.

Growth comparisons through time and with natural marshes nearby showed that marshes on low energy shorelines were more productive than those on high energy shorelines. *S. alterniflora* was more productive in the higher intertidal zone than the lower intertidal zone, particularly when marshes were getting started. Stem densities were slightly greater in marshes that faced south, i.e. the sunniest side and the leeward side (strongest winds usually came from the northeast).

Several recommendations were given to improve chances of successful establishment of fringe marshes. 1) A breakwater should be used to protect a planting if the fetch is greater than 6 km. 2) The fringe marsh should be made as wide as possible from MSL to MHW. 3) *S. patens* should be used on bare beach areas above the *S. alterniflora* zone. 4) Maintenance planting for a new marsh should be done early in each growing season. 5) Newly developing marshes should be fertilized twice each summer using a slow release type fertilizer like Osmocote.

Frenkel, R. E. and L. M. Kunze. "Introduction and spread of three *Spartina* species in the Pacific Northwest." In: Annual Meeting of the Association of American Geographers in Washington, DC, Washington, DC: Assoc. Amer. Geographers 1984.

This paper describes the introduction and spread of *Spartina alterniflora*, *S. patens*, and *S. anglica* in the Washington and Oregon estuaries. The potential spread of *Spartina* and the consequent loss of intertidal mudflats are also discussed. *Spartina alterniflora* appears to have been accidentally introduced into Willapa Bay prior to 1911, perhaps associated with some facet of the oyster industry that was established there in 1904. *S. alterniflora* was purposefully transplanted to enhance duck habitat in Thorndyke Bay, Gibson Spit, Kala Point, and Padilla Bay, Washington. These colonies are well established, but are spreading slowly.

Spartina patens appears to have been introduced at Cox Island, Oregon, probably in the early 1920's. First records were of a small patch at that time, but present records show it has expanded exponentially to cover about 3000 m². *S. patens* has since spread to the Dosewallips River in Washington.

Spartina anglica C.E. Hubbard, was planted on the eastern shore of Port Susan Bay near Stanwood, Washington. It has since spread to Livingston Bay, Iverson Spit, Triangle Cove, and Skagit Bay. *S. anglica* (sometimes call *S. townsendi*) is a fertile cross between *S. alterniflora* and *S. maritima* and is a vigorously spreading cordgrass. These three species are potential threats to native salt and brackish marsh plants in the Northwest, and to the bare mud and sand flats in the estuaries that are used by migratory birds.

Gallagher, J. L. "Salt marsh soil development." In: Rehabilitation and creation of selected coastal habitats: proceedings of a workshop. eds. J. C. Lewis and E. W. Bunce, U.S. Fish Wildl. Serv., 28-34, 1980.

This article reviews soil attributes that need be considered for successful planting of a salt marsh. The physical and chemical compositions of the soil are described in relation to five characteristics: stability, acidity, moisture, salinity and nutrients, with notes on interactions.

Stability is greater in sandy soils than in silt and clay oozes, but the latter can sometimes be improved by de-watering. Increasing stability increases the chance of plants staying in place and becoming established. Acidity was noted as a potential problem. Acidity is often caused by oxidation of the iron sulfides in dredged material when the material is pumped out and mixed with air and water. This is one reason for letting fresh dredged material sit and "age" for several months before planting; the acidity has a chance to be neutralized. Moisture content influences oxygenation of the soil. Soil, that was always saturated, became anaerobic. Plants capable of existing under anaerobic conditions (*Spartina alterniflora* being one) will grow in zones according to the lowest level of oxygen they can tolerate. Soil salinity influences plant establishment and zonation. Tidal inundation, salinity of the estuarine water, elevation, drainage, evapotranspiration and rainfall, all influence soil salinity. Nutrients are frequently limiting for plant growth in new marshes. Coarse sandy soils hold less nutrients than finer textured substrates, and fertilizers are often required to promote healthy plant growth when starting a marsh.

Gallagher, J. L., G. F. Somers, D. M. Grant, and D. M. Seliskar. "Persistent differences in two forms of *Spartina alterniflora*: A common garden experiment." Ecology 69(4) (1988): 1005-1008.

This paper provides a good review of the controversy around whether tall and short forms of *Spartina alterniflora* are genetically distinct, or whether these phenotypic differences are caused by environmental factors. Both tall and short form plants were transplanted from a Delaware salt marsh into common backyard garden plots also in Delaware. The plots were irrigated three times each week during the growing season with water (15-30 ‰) from a tidal creek. Initially the tall form was 30% to 100% taller than the short form. Differences between the height of the two forms persisted even after 9 years in the gardens. In addition, other morphological differences such as stem density, root-to-shoot ratios, and culm diameter varied between the two forms in a manner comparable to the differences in natural stands. Differences in productivity and underground reserves were also observed. The authors concluded that the phenotypic differences between tall and short form *S. alterniflora* are at least partially due to genetic differences between the forms. The possibility was also discussed that the forms are genetically similar, but certain genetic characters are turned on by environmental factors at the seedling stage, and that these differences persist despite long-term exposure to different environmental conditions.

Garbisch, E. W., Jr., P. B. Woller, and R. J. McCallum. "Salt marsh establishment and development." Fort Belvoir, VA.: U.S. Army Corps Eng., Coastal Eng. Research Center, Tech. Memo. 52, 114, 1975.

This study tested survival and growth of peat-potted *Spartina alterniflora*, *S. patens*, *S. cynosuroides*, *Distichlis spicata*, and *Ammophila breviligulata* seedlings that had been raised in a greenhouse. Seedlings were planted at inter-tidal and supra-tidal elevations. Grow-out sites were on natural beaches, sand and mud flats, and on dredged material beaches. Elevations, fertilizer applications, and planting dates were also investigated for influence on plant survival. Additional objectives were to determine shoreline stabilization potentials and sediment trapping potentials of the plants, and to determine the rate of macrobenthos colonization of the dredged material sites.

Seeds were harvested from natural stands near Assateague Island, VA during October, and grown out in sand filled peat pots the next spring. *Spartina alterniflora* was planted on 0.9-m centers; the others were planted on 0.6-m centers. All were in 10-cm diameter peat pots containing about five seedlings each. Plantings were made at four sites within a 40 km radius of St. Michaels, Maryland. All sites had gentle slopes of 1-6 degrees. Macrobenthos was sampled along four elevation contours at sites A, B, and #4. Elevations were +43 cm MLW (= MHW), +21 cm MLW (= MTL), MLW, and -15 cm MLW (= subtidal). Ten cores were taken at 0.5-m intervals along each elevation contour at each site. Cores were taken to a 15-cm depth using a corer with either a 21-cm² or a 46-cm² cross-sectional area. Organisms were separated from the substrate by washing the samples on 4.0, 1.0 and 0.5 mm sieves.

Results showed *Spartina patens*, *Distichlis spicata*, and *Ammophila breviligulata* all survived well at supratidal sites. *Spartina cynosuroides*, planted at MTL was completely killed within two years, most likely due to too much inundation. *Spartina alterniflora* suffered 60-90% mortality in areas from MHW to below MTL due to strong wave action. Those seedlings planted above MHW suffered about 10-40% mortality. Fertilizer applications to smooth cordgrass in moderate wave energy sandy areas increased net production 135-860%. Net production of smooth cordgrass in the unfertilized dredged material area was similar to that in the fertilized sandy areas.

Macrobenthos invaded the dredged material area rapidly. After only three months, intertidal macrobenthos at the transplant site was as dense as that in the nearby natural marsh. There were four dominant taxa which accounted for about 90% of the intertidal macrobenthos by number: *Laonereis culveri*, *Macoma balthica*, *Tubulanus pellucidus*, and Tubificid oligochaetes. Densities of macrobenthos tended to increase with time of inundation, i.e. from MHW to subtidal regions.

The use of peat pot plants in transplanting was recommended. This technique extends the planting season, reduces impacts on natural donor marshes, and allows transplant preparation. An important factor in transplant preparation is the acclimation of plants to site specific salinities prior to transplanting.

Hardaway, C. S., G. R. Thomas, B. K. Fowler, C. L. Hill, J. E. Frye, and N. A. Ibisson. "Results of the vegetative erosion control project in the Virginia Chesapeake Bay system." In: Proceedings of the twelfth annual conference on wetlands restoration and creation. in Tampa, FL, ed. F. J. Webb, Jr. Tampa, FL: Hillsborough Community College, 144-158, 1985.

This paper reports some of the results of the Vegetative Erosion Control Project conducted by the Virginia Institute of Marine Sciences. This was a study of 24

marsh plantings in the Chesapeake Bay system in Virginia. Sites were chosen to maximize diversity of conditions, mainly fetch and direction faced by the marsh. *Spartina alterniflora* was planted on 0.5-m centers from MHW to just below MSL. *Spartina patens* was planted above *S. alterniflora* at several sites. Osmocote fertilizer, a slow time-release type, was used; 30 ml was placed in the hole when each culm was planted. Osmocote was also broadcast into the marsh later when plants had started growing.

Results showed a fringe of marsh grass could be established with little or no maintenance on low energy shorelines, areas where the fetch was less than 1.8 km. Where wave energy was moderate (1.8-6.3 km fetch), the upper margin of the fringe marsh should be protected by establishing *S. patens* to trap sand and sediments, and preserve the back area from winter storm erosion. Areas with average fetches of 6.3-10 km need to be protected by a wave stilling device (breakwater). Fringe marshes should not be considered for beaches with average fetches greater than 10 km unless the beach is protected by a headland or spit. Also, maintenance planting of wash-out and die-back areas of the marshes should be required and anticipated.

Hartman, R. D., R. N. Reubsamen, P. M. Jones, and J. L. Koellen. "The National Marine Fisheries Service habitat conservation efforts in Louisiana, 1980 through 1990." *Mar. Fish. Rev.* 54 (1993): 11-20.

This article reviews the NMFS efforts to conserve habitat for fisheries and associated organisms in Louisiana from 1980 through 1990. During these years, NMFS reviewed 14,259 public notices to dredge, fill, or impound wetlands in Louisiana. The Habitat Conservation Division (HCD) of NMFS provided recommendations to the Corps of Engineers on 962 permit projects which would impact 240,000 ha of tidal influenced wetlands. Avoidance was recommended for about 113,000 ha, and mitigation was recommended for 63,5000 ha. Revisions or denials were recommended by HCD on only 12% of all proposed actions. The remaining permit applications were considered to have only minor impacts on marine fisheries. On a permit basis, 43% of HCD recommendations were accepted by the Corps, 34% were partially accepted, and 23% were rejected. Most of the permits involved oil and gas activities, followed by shoreline modifications, and then pipeline activities. A breakdown of permits and areas involved was given by drainage basins.

There is a need for greater awareness of coastal wetland loss through the Corp's Section 10/404 permitting program. An accurate continuing account of permitted wetland alterations and mitigation measures is needed to guide future decisions that will help preserve this important habitat.

Hoffman, W. E. and J. A. Rodgers. "A cost/benefit analysis of two large coastal plantings in Tampa Bay, Florida." In: Proceedings of the seventh annual conference on wetlands restoration and creation, in Tampa, FL, ed. D. P. Cole, Tampa, FL: Hillsborough Community College, 265-278, 1980.

This study compared costs of establishing a *Spartina alterniflora* marsh to those of establishing a mangrove stand. On the dredged material extension of Sunken Island in Hillsborough Bay of the Tampa Bay system, Florida, a 1.64 ha area was planted with smooth cordgrass using 12-cm diameter plugs planted in rows on 1.0-m centers with the

rows being 2 m apart. This effort required 995 man-hours/ha, and a total cost of about \$4,565/ha. About 93% survival was obtained, and after 14 months the plants had spread enough to almost hide the original planting layout. The mangrove planting was on a 0.52 ha plot on another dredged material island called CDA-D, also in Hillsborough Bay. Here *Avicennia germinans* (63%) and *Laguncularia racemosa* (37%) were planted on 2-m centers. Transplants were 30-190 cm tall. Survival was about 73% after 13 months. This planting required about 2541 man-hours/ha, and a total cost of about \$11,459/ha. Labor costs were about 70% of the total cost of each planting, but did not include planning and supervision.

An interesting point made in this study was that the *Spartina* donor marsh had fully recovered within 12 months. Plants had been removed from the donor marsh at about 1 plug/m². Another feature that was unusual was the clipping of the cordgrass to leave only 10 cm above the substrate once the stems had been transplanted. Clipping was done to reduce transpiration and possible shock due to root damage. Impacts on mangrove donor stands can be longer lasting, but no particular mention was made of this study's donor stand.

Josselyn, M. J., ed. Wetland restoration and enhancement in California. La Jolla, CA: California Sea Grant College Program, 1982.

This report includes eight presentations presented at a California wetlands restoration workshop held in February 1982 at the California State University, Hayward, CA. Seven of the presentations were followed by panel and audience discussions through which concerns about restoration gains and losses were voiced. Presentations reviewed past, current and future wetland restoration actions and potentials including development of regional wetland restoration goals and legal (and institutional) constraints. Design strategy, engineering features of circulation, sedimentation and water quality, techniques for restoring vegetation and for monitoring were also presented and discussed. Poster session abstracts and an ample bibliography were also valuable inclusions.

Many of the problems and concerns voiced at this workshop are still valid though a decade has passed. It appears there will always be confrontation between ecological action to preserve a clean environment and the lure of economic riches possible through residential and municipal development of wetlands.

Josselyn, M. N., J. B. Zedler, and T. Griswold. "Wetland mitigation along the Pacific coast of the United States." In: Wetland creation and restoration: The status of the Science. Part 1, Regional review., eds. J. A. Kusler and M. E. Kentula. 1-36. Washington, DC: Island Press, 1989.

This article reviews west coast wetland types, reviews considerations that aid in making restorations successful, and reviews the kinds of projects that have been done. Some critical features of restoration plans are described, including: site history, topography, water control structures, hydrology, flood events, sediment budget, edaphic characteristics, existing wetland characteristics (vegetation and wildlife), and adjacent site features and conditions. Two levels of monitoring are recommended: (1) the enforcement monitoring to make certain implementation is following permit requirements, and (2) environmental monitoring to see if the design was correct and the functions are being realized. Several restoration projects are profiled including successes and significant findings.

Kentula, M. E., R. P. Brooks, S. E. Gwin, C. C. Holland, A. D. Sherman, and J. C. Sifneos. An Approach to Improving Decision Making in Wetland Restoration and Creation. Corvallis, OR: U.S. Environmental Protection Agency, Environmental Research Laboratory, 1992.

The objective of this book is to provide a guide that assists people in conserving, restoring, and creating wetlands successfully. The book presents a technique for analyzing an area of the country to determine types and amounts of existing wetlands, and the types of wetlands needed. An approach for their successful restoration or creation of wetlands is recommended along with variables to be monitored to determine the least harmful environmental impacts and best chances for project success.

Restoration projects should include: (1) precise objectives, (2) detailed plans including scheduled actions, (3) detailed maps or diagrams of the site, and (4) a checklist of variables to be monitored. Variables that should be monitored relate to morphometry, hydrology, substrate, vegetation, fauna, and sometimes water quality.

The authors also recommended the establishment of a permanent database that contains the important information for evaluating the success of each project. Data suggested to be entered regarding each project included: permitting agency, permit number, date of permit, type of mitigation, location of the mitigation site (state, county, city and address), date construction began, date construction was completed, map showing the as-built conditions, name of the contractor/builder, name of the contracting company, and specific objectives of the project. Entries should be included for mid-course evaluation of the construction and for corrections made based on the evaluation. Entries should also be included that describe conditions found during a final evaluation after planting was completed. Additional data to add would be expected to come from monitoring reports that evaluate the development of the plant and animal communities in the wetland at various intervals (six months to a year) after planting was completed.

Kiraly, S. J., F. A. Cross, and J. D. Buffington. The federal effort to evaluate coastal wetland mitigation: A report by the National Ocean Pollution Policy board's habitat loss and modification working group. NOAA Technical Memorandum CS/NOPPO 91-2. 10 p. plus Appendices., 1991.

This report summarizes results of a workshop on wetland mitigation, held at San Diego State University in January 1991. Federal efforts to evaluate coastal wetland mitigation were assessed including analyses of functional values in created wetlands and follow-up studies for federally-permitted mitigation projects. Conclusions were that federal efforts in these areas should be improved. Additional research should be conducted on understanding how coastal wetland ecosystems function. A system should also be established for evaluating the success (including functional success) of permitted mitigation projects.

Knutson, P. L., R. A. Brochu, W. N. Seelig, and M. Inskeep. "Wave damping in *Spartina alterniflora* marshes." Wetlands 2 (1982): 87-104.

Field experiments were conducted to test a model of wave damping and to determine how well a marsh can dampen waves. The model has been used to explain how a marsh damps waves and reduces waves' erosional forces. Two *Spartina alterniflora* marshes in Chesapeake Bay in Virginia were chosen as test sites: Wescoat Cove north shore marsh

(the oldest known man-made cordgrass marsh, planted in 1928 by Mr. Wescoat) and Kings Creek north shore marsh. The marsh vegetation was characterized by clipping 0.25 m² areas of vegetation out of the marsh where each sensor was to be placed. Stem density, length and thickness ranged from about 180-350 stems/m², 20-30 cm (stem length was about half the plant's total height), and 5-6 mm in diameter, respectively. In each marsh, wave sensors were placed along a transect. Sensors were placed at the seaward edge and 2.5, 5, 10, 20 or 30 m in towards shore. The research vessel, Virginia Dare, generated waves for the tests; waves ranged in height from 0.06-0.30 m. Controls were run on an adjacent non-vegetated beach.

Analyses indicated the model worked well, only requiring minor changes in the coefficients to adjust for the plants. Results also showed the marshes significantly reduced wave height and erosional energy. Wave height was reduced by about 50% within the first 5 m of marsh, and by about 95% after crossing 30 m of marsh. Wave energy was reduced in these cases by 65% and nearly 100%, respectively.

In theory, marshes are most effective at damping waves when waves were less than plant height (the condition during the tests) but would not be during storm tides. Under storm conditions, when water levels rise and waves frequently exceed plant height, marshes would be less effective against erosion by waves.

Knutson, P. L., J. C. Ford, M. R. Inskeep, and J. Oyler. "National survey of planted salt marshes (vegetative stabilization and wave stress)." *Wetlands* 1 (1981): 129-157.

This study describes a technique for evaluating a coastal site's potential for vegetative stabilization based on the site's shoreline characteristics that relate to wave-climate severity. Results were based on the study of 104 salt marsh plantings in 12 coastal states. All marshes studied were exposed to wind waves, were located in brackish and salt water environments, were planted with *Spartina alterniflora* or *S. foliosa* at least one year prior to the survey, and had no rubble or man-made structures in the planted areas.

Results of correlation analyses indicated that sediment grain size in the swash zone, longest or average fetch, and shore configuration were useful indicators of a site's suitability for vegetative stabilization. There was an 80% success rate in establishing a fringe marsh when sediment grain size was 0.4 mm or less, and there was an 80% failure rate when sediment grain size was 0.8 mm or greater. Failures increased when fetches increased. Successes increased as protection increased, with the greatest success rate found in coves. Another recommendation was that the site should have at least 6 m of intertidal width and be planted over 60% of this area; this should cause sufficient wave dampening to prevent erosion during most of the year. On the basis of these observations, a site evaluation form was developed to predict the success of a *Spartina* planting to control an area's erosion. This form was named the Vegetative Stabilization Site Evaluation Form.

Kraus, D. B. and M. L. Kraus. "The establishment of a fiddler crab (*Uca minax*) colony on a manmade *Spartina* mitigation marsh, and its effect on invertebrate colonization." In: National wetland symposium: mitigation of impacts and losses in Berne, NY, eds. J. A. Kusler, M. L. Quammen, and G. Brooks, Berne, NY: Assoc. State Wetland Managers, 343-348, 1986.

The objective of this study was to establish fiddler crab populations in a man-made *Spartina alterniflora* marsh at the Mills Creek mitigation site, and to compare the macrobenthos at this marsh with that in a natural marsh area on Sawmill Creek, both in the Hackensack River basin in northeast New Jersey. Fiddler crabs (*Uca minax*) were collected from a colony at Moonachie Creek (also in the Hackensack River basin), transported to the test sites (7-m² sites in the developing marsh), and deposited one crab per artificial burrow during 29-30 May 1986. Artificial burrows were made about 50 cm deep on 25-cm centers using a broom handle. Censuses of the number of burrows, types of burrows, and crabs seen in each of the two test sites (= colonies) were made in June, July, August and September. Benthic macrofauna were sampled at each colony, two control sites (30 m to the side of each colony), and in a *Phragmites* marsh using a bulb corer (300 ml) to a depth of 10 cm. Replicates were taken in May (before the crabs), June, July and September. Animals were separated from the cores using a 1 mm mesh sieve.

Results of censuses showed that many of the crabs remained in each test site, forming two colonies. One month after transplanting the crabs, about 42% of the burrows were occupied. A month later a few additional burrows were noted, and the presence of small burrows indicated that recruitment may have occurred. By September, colony densities achieved those of natural colonies elsewhere in NJ. Benthic macroinvertebrates were five to ten times more abundant in the developing marsh and natural marsh than in the crab colonies or in the *Phragmites* marsh. Although sampling was not robust, fiddler crabs appeared to decrease the number of benthic invertebrates in the substrate of the colonies, perhaps through predation.

Kruczynski, W. L. "Salt marshes of the northeastern Gulf of Mexico." In: Creation and restoration of coastal plant communities, ed. R. R. Lewis, III. 71-88. Boca Raton, FL: CRC Press, Inc., 1982.

Salt marshes along the northeast coast of the Gulf of Mexico are generally similar to those along the Atlantic coast and the rest of the Gulf coast. However, *Juncus roemerianus* is more important along the northeast coast where it displaces much of the *Spartina alterniflora* normally found in the intermediate marsh. Descriptions of marshes in the northeast Gulf include marsh plant zonation and productivity, marsh animal communities, and marsh destruction. Salt marsh creation activities are summarized, as are the related uses of the various dominant plant species for restoration purposes. Species reviewed are: *Spartina alterniflora*, *Spartina patens*, *Spartina cynosuroides*, *J. roemerianus*, *Distichlis spicata*, *Phragmites communis*, *Panicum repens*, *Panicum amarum*, *Uniola paniculata*, and *Ammophila breviligulata*. Although *Juncus* is an important species in natural marshes of this area, it is not easily transplanted. Best success has been achieved by transplanting clumps of *Juncus*, rather than single sprigs.

Additional information was given on handling transplant materials, use of fertilizers, planting methods, and factors affecting successful establishment of a transplanted marsh. The most important factors to consider for success were erosion control (water and wind) and soil characteristics (soil water, soil salinity, and soil nutrients).

Kusler, J. A. and M. E. Kentula, eds. Wetland Creation and Restoration. The Status of the Science. Washington, DC: Island Press, 1989.

This book is a collection of chapters that summarize the status of the restoration science for various wetland types. The executive summary describes the adequacy of our scientific understanding concerning wetland restoration and creation, offers recommendations for needed scientific research to fill the information pages, and gives recommendations to wetland managers as to restoration and creation potential based on current scientific knowledge.

Scientific knowledge and data has been developing, but much is still unknown. Some blame should be placed on poorly specified goals and the lack of monitoring for many of the early, and even current, restoration/creation projects. Some wetlands appear to be easier to restore than others, as are some of the wetland functions. Rarely is a restoration a complete success, with all functions of a natural system being obtained, but partial successes that restore some of the wetland functions are beneficial and may lead to additional functional development in the future. The long term success of a wetland restoration is even more uncertain than the short term success. Both often depend upon our abilities to manipulate the site and its surrounding land, to maintain close supervision during all phases of a project, and to keep pests and intruders away from sensitive areas. Fourteen recommendations are offered to wetland managers to assist them in creating and maintaining wetlands:

1. Wetland restoration must be viewed with some cynicism, particularly where promises are made to create a natural system in exchange for a permit to destroy or degrade a natural system that already exists.
2. Multidisciplinary expertise in planning and project supervision is needed in all project phases.
3. Clear, site-specific project goals should be established first.
4. A detailed plan concerning all phases of a project should be prepared in advance to help regulatory agencies evaluate the probability and achievement of success.
5. Site-specific studies should be done for the original system prior to wetland alteration.
6. Careful attention to wetland hydrology is needed in the project design.
7. Wetlands should be designed to be self-sustaining systems and persistent features in the landscape.
8. Wetland design should consider relationships of the wetland to water sources, other wetlands in the watershed, and adjacent upland and deep water habitats.
9. Buffers, barriers, and other protective measures are often needed for a successful restoration or creation of a wetland.
10. Restoration should be favored over creation.
11. A project should incorporated monitoring and methods for mid-course corrections when needed.
12. Long term management is needed for some types of wetland systems.
13. Risks inherent in restoration and creation of wetlands should be carefully evaluated and be reflected in project design.
14. Restoration action for artificial or already altered systems requires special evaluation as to regional needs.

Landin, M. C. and J. W. Webb, Jr. "Wetland development and restoration as part of Corps of Engineer programs: Case studies." In: National Wetland Symposium: Mitigation of Impacts and Losses. in New Orleans, LA, eds. J. A. Kusler, M. Quammen, and G. Brooks, New Orleans, LA: Association of State Wetland Managers, 388-391, 1986.

This paper presents an overview of the Corps of Engineers' work to use dredged material constructively, generally to create, restore or enhance wetlands. It also presents short reviews of seven projects. Since about 1970, over 130 wetland sites have been constructed using dredged material. Many of the sites were in coastal saline and brackish waters. Sites ranged in size from 0.4 to thousands of hectares. Results showed that most man-made wetlands were at least partially successful. These wetlands required about 3-5 years to develop into habitats comparable to natural marshes. Although above-ground vegetation generally could be established in the marshes within a couple of growing seasons, sediment organics and root biomass required more time, perhaps 10 years or more, to approach the conditions found in neighboring natural marshes.

Langis, R., M. Zalejko, and J. B. Zedler. "Nitrogen assessments in a constructed and a natural salt marsh of San-Diego Bay." Ecological Applications 1 (1991): 40-51.

Differences in nitrogen content in soil, soil water, and plant stems and leaves were studied at a natural marsh (Paradise Creek) and a man-made marsh (Connector Marsh) in the San Diego Bay area, CA. Soil N pools and organic carbon were lower in the constructed marsh than in the adjacent natural marsh. Above-ground biomass and foliar N of *Spartina foliosa* were also lower in the constructed marsh. Rates of N fixation were lower in the surface (1-cm) sediments of the constructed marsh than the natural marsh, but not in the deeper sediments (down to 10 cm). Addition of organic matter to the soil increased N fixation rates in both marshes, more so when glucose was used than when ground-up dry *Spartina* plants were used. Nitrogen mineralization rates were high in both marshes. Results in general pointed to low import of nitrogen from tidal or stream flows, and high nitrogen demands by the marsh plants and ecosystem. Without a source of organics and nitrogen, constructed marshes will take a long time to develop production equivalent to natural marshes in the San Diego Bay area; longer perhaps than was estimated for Gulf and Atlantic coastal created marshes with *Spartina alterniflora* and *S. patens*.

LaPerriere, A. J. and M. M. Farmer. "Recent wetland restoration/creation actions in the New York District." In: Proceedings of the sixteenth annual conference on wetlands restoration and creation. in Plant City, FL., ed. F. J. Webb, Jr. Plant City, FL.: Hillsborough Community College, pp. 97-108, 1989.

Preliminary results of four New York District Corps of Engineers enforcement actions resulting in the restoration of one palustrine and three saltwater marshes are reported. The three saltmarsh restorations involved illegal fill being removed from sites in south central Long Island. At Site 1 (2.2 ha), the fill was removed within a few weeks of deposition, and was removed carefully so that about 25% of the original root mat was left intact and alive. At Site 2 (0.5 ha), the fill was removed a year after deposition, and all marsh plants and root matter had died. At Site 3 (0.04 ha), fill was removed two months after deposition, but conditions were such that the marsh was dead by the time removal was

complete. At all three sites, substrate elevations matching those present prior to filling were carefully restored and the areas were allowed to revegetate naturally.

Revegetation was estimated subjectively to be 50% coverage during the first growing season at Site 1. During the second growing season, transects with 1-m² plots established at 30.5-m intervals were established in each site. Percentage cover by each plant species was estimated and all were summed for a total coverage percentage. Percentages of total cover for Site 2 and Site 3 after one growing season were about 37%. For Site 1 after two growing seasons, total cover was about 67%.

This paper showed how enforcement of wetland regulations, coupled with planned restorative action, can be effective. It also showed that in some cases, expensive transplanting operations may not always be necessary.

LaSalle, M. W., M. C. Landin, and J. G. Sims. "Evaluation of the flora and fauna of a *Spartina alterniflora* marsh established on dredged material in Winyah Bay, South Carolina." *Wetlands* 11 (1991): 191-208.

The objectives of this study were to compare floral and faunal characteristics of 4 and 8 year old *Spartina alterniflora* marshes that developed naturally on unconfined dredged material deposited in Winyah Bay, SC. Both marshes were tall form *S. alterniflora*, and samples were collected in September of 1988. Most samples were collected at 10 randomly chosen sites along a 50-m transect in each marsh. Above-ground vegetative characteristics were assessed from a 0.25 m² quadrat at each site. Sediment, benthos, and below-ground biomass were sampled by coring at these sites. Large (1 to 2 cm) macrobenthos were collected from adjacent 1 m² quadrats. Fish, shrimp and crabs were collected only in the 4-yr old marsh with Breder traps and block nets.

Marsh sediments were similar, and substrata were mainly silts and clays. The percent organics in all sediments examined was about 11%. Percent cover by *Spartina alterniflora* was about 50% in both marshes. Stem density (257 vs. 199 stems/m²), below-ground biomass (3061 vs. 2204 g/m²) and total biomass (3692 vs. 3061 g/m²) were greater in the older marsh, but stem height (40 vs. 66 cm) and above-ground biomass (631 vs. 856 g/m²) were greater in the younger marsh.

Total density of benthic macrofauna from sediment cores was significantly greater in the 8-yr old marsh compared with the 4-yr old marsh, with mean values of 150 vs. 35 organisms/75 cm². Species composition in the two marshes was similar, and the infauna was dominated by oligochaetes. Differences in infaunal density between the two marshes were attributed to marsh age, although the authors acknowledged that other factors such as distance to open water may have affected the populations.

The fish and shellfish collected from the 4-yr old marsh in Breder traps were typical estuarine fauna reported by others for the natural marshes in Georgia and North Carolina. The mummichog, *Fundulus heteroclitus*, and the daggerblade grass shrimp, *Palaemonetes pugio*, were the dominants. Block net data were not presented, but apparently confirmed that large numbers of mummichogs and daggerblade grass shrimp were present in the marsh channels along with blue crabs, *Callinectes sapidus*. Gut contents from mummichogs indicated that *Uca* and insects were dominant prey items.

Lewis, R. R., III. "Creation and restoration of coastal plain wetlands in Florida." In: Wetland creation and restoration: The status of the Science. Part 1, Regional review., eds. J. A. Kusler and M. E. Kentula. 73-102. Washington, DC: Island Press, 1989.

This chapter reviews past activities related to salt marsh and mangrove restoration in Florida. Restoration and creation projects were generally initiated to mitigate destruction of natural wetlands, to enhance existing habitat, and to stabilize eroding shorelines. Despite hundreds of restoration and creation efforts, there are few reports from which to draw information to make restorations a science rather than an art. The more successful projects had paid attention to many factors including: location, wave climate, tidal range, salinity, shading, time of planting, substrate quality, condition of the transplants, buffer zones, and monitoring with mid-course corrections if needed. On the basis of critical review of projects and of the sparse literature, five factors appeared to be the most important to successful restorations: correct elevation, adequate drainage, protection from wave damage, appropriate plant material, and protection from human impacts. Future needs are seen as: a centralized data bank for restoration projects, research on natural recruitment versus transplanting, rates of faunal recruitment to restoration sites, and regional planning for restorations.

Lewis, R. R., III. "Wetlands restoration/creation/enhancement terminology: suggestions for standardization." In: Wetland creation and restoration: The status of the Science. Part 2. Perspectives., eds. J. A. Kusler and M. E. Kentula. 417-419. Washington, DC: Island Press, 1989.

This document provides a glossary of terms frequently used in restoration, creation, and enhancement research. The terms defined are: mitigation, mitigation banking, restoration, creation, enhancement, and success. Restoration is defined as "returned from a disturbed or totally altered condition to a previously existing natural, or altered condition by some action of man." Creation is defined as "the conversion of a persistent non-wetland area into a wetland through some activity of man." Enhancement is defined as "the increase in one or more values of all or a portion of an existing wetland by man's activities, often with the accompanying decline in other wetland values."

Lewis, R. R., III. "Coastal habitat restoration as a fishery management tool." In: Stemming the tide of coastal fish habitat loss., ed. R. H. Stroud. Savannah: National Coalition for Marine Conservation, Inc., 1992.

In response to declines in both the commercial and recreational harvests of fishery species, a number of fishery management tools have been proposed and implemented. In the past, management methods concentrated on increasing survival of late juveniles and adults to restore or increase egg production. In recent years, methods which increase the survival of larval and juvenile fishes came to be considered more important to a species' reproductive success than egg production. Coastal wetland restoration is one such method, particularly for those estuarine-dependent species whose life histories include a resident period in shallow low-salinity marine habitats. A summary of the use of coastal restoration in past studies shows both its importance as a management tool, and the paucity of published information concerning its use. This lack of information is another reason that wetlands

restoration is not generally listed or used as a fishery management tool. Another is the belief that restored wetlands can not reach a productive level equivalent to natural wetlands. Despite this generally negative attitude, restoration of coastal wetlands is generally acknowledged as being more predictable and assured of success if done correctly. Reviews of past projects show fish and wildlife populations closely approximating those found in natural wetlands, and suggest that this is an underutilized fishery management tool.

Lewis, R. R., III and C. S. Lewis. "Tidal marsh creation on dredged material in Tampa Bay, Florida." In: Proceedings of the fourth annual conference on restoration of coastal vegetation in Florida, in Tampa, FL, eds. R. R. Lewis, III and D. Cole, Tampa, FL: Hillsborough Community College, 45-67, 1977.

Three experimental plantings of smooth cordgrass, *Spartina alterniflora*, were done on a 12 year old dredged material island in Tampa Bay, Florida. The tests were to assist in developing marsh creation techniques.

In the first planting, 36 single stems were dug from an adjacent natural marsh and planted in six rows of six plants on 1.0-m centers. The substrate was uneven and the elevation of the planted plot ranged from 49 to 61 cm above MLW, or just a little lower than the natural marsh (58-64 cm above MLW). Planting was in September, 1976, and the test area was well protected from waves and had a gentle slope. About 91% of the transplants survived and increased so that nine months later there were 267 stems. The control area showed no volunteer establishment of *Spartina alterniflora* during the course of this study.

The second planting was adjacent to the first. It involved only 15 seedlings sent to Tampa. The seedlings had been grown in Maryland from seeds harvested near Assateague Island, Virginia. The seeds were germinated and raised by Environmental Concern, Inc. in Maryland. These seedlings were also planted in October, 1976, on 1.0-m centers. No details of survival percentages were given, but eight months later there were 331 stems.

The donor marsh for the first planting went into flower in November, and 237,000 spikelets were harvested and shipped to Environmental Concern, Inc. Eleven percent of the spikelets contained seeds, and 63% of the seeds germinated. Many of the seedlings produced were sent back to Tampa Bay for the third planting experiment. The 500 healthiest (=tallest) seedlings were planted on 1.0-m centers in another area adjacent to the previous planting. This planting was in May, 1977. One month later there appeared to be at least 90% survival.

The successful establishment of cordgrass on the island showed that a salt marsh could be established much more rapidly if assisted by transplanting. In addition the study showed that a northern variety of smooth cordgrass could be successfully transplanted in Florida. The experiments also showed planting could be successful even in the fall in Florida, probably because of the year-round warm weather. In photographs the plants appeared to be sparse and short, even in the natural marsh, but no mention was made of these features in the paper.

Lindall, W. N., Jr. and G. W. Thayer. "Quantification of National Marine Fisheries Service habitat conservation efforts in the southeast region of the United States." Mar. Fish. Rev. 44 (1982): 18-22.

The objectives of this study were to determine how many acres of coastal marsh were impacted by permitted alterations in the southeastern U.S. in a year, and to determine to what extent National Marine Fisheries Service's recommendations to protect this coastal habitat were being followed. From Oct. 1980 through Sept. 1981, there were 6,399 permit applications from the Corps of Engineers available for review by NMFS in the Southeast. The permits covered about 1,300 ha to be dredged, 2,590 ha to be filled, and 3,360 ha to be impounded. NMFS did not object to about 73% of the desired alteration after preliminary review showed they involved only minor alterations to wetlands. NMFS contracted 1,380 permit applications for thorough review, and did not assess 368 permit applications. Of the 7,272 ha in the 1,380 permits thoroughly reviewed, NMFS objected to alteration of 5,412 ha, but did not object to 1,860 ha being altered provided there was mitigation involving 1,012 ha of restoration and 324 ha of creation to reduce the loss of wetlands.

Almost all (98%) of the recommendations submitted by NMFS were incorporated in the permits by the Corps of Engineers, however, only 72% of these were complied with by permittees. Even though the study did not tell the percentage (by area) of wetlands that were preserved or improved, and did not evaluate the losses of wetlands from permitted activities versus non-permitted activities, it did show NMFS was effective in protecting a substantial amount of the country's coastal wetlands. The article also showed, however, that in one year in the southeast at least 524 ha of coastal wetlands were lost through the permitting process.

Lindau, C. W. and L. R. Hossner. "Substrate characterization of an experimental marsh and three natural marshes." Soil Science, American Journal 45 (1981): 1171-1176.

Changes were monitored in selected chemical and physical properties of dredged material used in the construction of a coastal marsh. Substrate properties were also compared with those of three natural marshes, all in the Galveston Bay area, Texas. The dredged material in all three elevational tiers at the transplant site contained about 97% sand, 2% clay, and 1% organic matter at the start of the study. Sixteen months later, the lowest tier had been covered by 2-30 cm of fine particles that had settled out of the water column due to a breakwater that had been constructed to protect the site from wave impacts. Baseline cores showed that organic matter, cation exchange capacity (CEC), total Kjeldahl nitrogen (TKN), and extractable phosphorus values were low, and nitrate and nitrite concentration were below detectable values. Ammonium was detected in half of the core samples, never exceeding 2.0 mg N/g. The clay content, CEC, TKN, extr-P, and sulfide values were highly variable due to the heterogeneity of the graded dredged material.

The intertidal and supratidal zones in the dredged material area were planted with *Spartina alterniflora*, *S. patens* and some other species. Trends of increasing concentrations of TKN, ammonium-N, organic matter, and extr-P, were found over the three post-planting samplings. These changes were mainly found in the lowest tier where most of the particulate matter accreted. Even with the increases found in the nutrients, their values were well below those for the natural marshes. Data suggest that concentrations of nutrients at the transplant site should approach those of the natural sites in a total of 2 to 5 years after

planting. The study also indicated that substrate concentrations of N and P did not show a response to the surface applications of fertilizer.

Lindau, C. W. and L. R. Hossner. "Sediment fractionation of Cu, Ni, Zn, Cr, Mn, and Fe in one experimental and three natural marshes." Journal of Environmental Quality 11 (1982): 540-545.

Clay mineralogical properties of a planted *Spartina alterniflora* marsh were compared with those of three natural marshes, all in the Galveston Bay area, Texas. At the transplanted marsh site the dredged material in all three elevational tiers sampled, contained about 97% sand, 2% clay, and 1% organic matter. A sequential chemical extraction procedure was used to obtain the concentration of the metals. Clay minerals found in the sediments of the experimental marsh were not significantly different from those in the natural marshes. Total elemental substrate concentrations of Cu, Ni, Cr, Zn, Mn and Fe averaged 7.9, 8.6, 25.5, 25.2, 123, and 12,200 ug/g, respectively, for the four marshes. About 30% of the total substrate Cu, Ni, and Zn was associated with the organic matter fraction in these marshes. About 53% of the experimental marsh Mn was associated with the easily reducible fraction, compared with only 11% in the natural marshes. Iron associated with the organic matter and sulfide fraction of the experimental marsh was about 10% lower than that for the natural marshes. The likelihood of heavy metals reaching toxic levels appears to be very low for these marshes.

Lyon, J. T., III. "A comparison of epiphytes on natural and planted *Spartina alterniflora* marshes." M.S. Thesis, North Carolina State University, 1975.

Epiphytic algae growing on stems of *Spartina alterniflora* were compared between two created salt marshes and nearby natural marshes at Beaufort and Snow's Cut, North Carolina. Comparisons were made through standing crop estimates and laboratory ¹⁴C uptake rates. Neither standing crop nor primary production varied consistently between transplanted and natural marshes. The age of stems and the marsh elevation appeared to be more important than whether the marsh was transplanted in determining epiphyte growth.

Mager, A. and G. W. Thayer. "NMFS habitat conservation efforts in the Southeast Region of the United States from 1981 through 1985." Marine Fisheries Review 48 (1986): 1-8.

The U.S. Army Corps of Engineers (COE) in the southeastern region of the United States regulates development activities affecting thousands of acres of wetlands every year. The National Marine Fisheries Service (NMFS) makes recommendations to the COE that are designed to minimize the effect of these projects on wetland resources. The NMFS Habitat Conservation Division's Southeast Region quantifies those COE projects relating to water development in the Southeast Region of the United States in a computerized system that tracks permit recommendations and proposed habitat alterations. This project was begun in late 1980, and the first five years of data are summarized. Such data are necessary in order to maintain a comprehensive view of wetlands modification to determine cumulative loss of habitat. This information is necessary to prevent avoidable damages to fisheries production and judge the effectiveness of the NMFS recommendations.

NMFS recommendations on permit applications are made by the Southeast Regional Office and its area offices, but are dependent on up-to-date research information provided by research laboratories of the Southeast Fisheries Center. The close links between these facilities and NMFS fisheries habitat conservation efforts are described.

Meyer, D. L., M. S. Fonseca, D. R. Colby, W. J. Kenworthy, and G. W. Thayer. "An examination of created marsh and seagrass utilization by living marine resources." In: Coastal Zone '93, Volume 2. Proceedings of the 8th Symposium on Coastal and Ocean Management., eds. O. Magoon, W. S. Wilson, H. Converse, and L. T. Tobin. 1858-1863. New York.: American Society Of Civil Engineers, 1993.

This paper summarizes the results of a study where fish, shrimp, and crab utilization of planted and natural *Spartina alterniflora* marshes and *Halodule wrightii* - *Zostera marina* seagrass beds was evaluated. Initially, *S. alterniflora* was planted in 1987 at three dredged-material sites in North Carolina in solid and reticulated (with access channels) patterns. By 1992, however, the marsh had coalesced into a solid vegetative stand. Habitat heterogeneity was then added by placing oyster cultch along certain areas of the marsh shoreline. Fishery utilization of the created marshes and nearby natural marsh was examined from 1987 through 1989 by the use of block and fyke nets that collected animals retreating off the marsh surface with the ebb tide.

Density data were presented from three collections of fisheries organisms made in the two years following transplanting. The overall mean density of shrimp (mainly daggerblade grass shrimp and brown shrimp) was 3.1 times as large in the natural reference marsh as in the planted marshes, but differences were not significant apparently due to high sample variability. Mean crab densities (mainly blue crabs) were twice as high in the planted marshes compared with the natural marsh, but again this difference was not statistically significant. Fishes collected included spot, mummichog, pinfish, and pigfish, and the mean density for total fish was twice as high in the natural marsh as in the created marshes. This difference was statistically significant during two of the three sampling periods. The results of this study highlight the difficulties encountered in collecting quantitative data from marsh surfaces in order to assess fishery utilization patterns in natural and created salt marshes.

The effect of depositing oyster cultch along the marsh shoreline was examined 3 months after cultch placement. Oysters, xanthid crabs, amphipods, and other reef organisms were found inhabiting this cultch. This addition appeared to increase animal diversity in the marsh because few of these organisms were collected in marsh areas where cultch was not added.

Miller, L., G. T. Auble, and K. A. Schneller-McDonald. "User's guide to the wetland creation/restoration data base, version 2." Wash. D.C.: U.S. Dept. Inter., Fish and Wildlife Service, Research and Development. 1991.

This is a guide to assist users in accessing a very large annotated bibliographical database about all types of wetland restorations and creations, and related topics. The guide facilitates finding articles about selected subjects in the database. The database (Wetland Creation/Restoration Data Base) contains about 500 entries referring to *Spartina alterniflora*, and should be of use to researchers.

Minello, T. J. and J. W. Webb, Jr. "The development of fishery habitat value in created salt marshes." In: Coastal Zone '93, Volume 2. Proceedings of the 8th Symposium on Coastal and Ocean Management., eds. O. Magoon, W. S. Wilson, H. Converse, and L. T. Tobin. 1864-1865. New York: American Society Of Civil Engineers, 1993.

This short paper describes preliminary results from a Coastal Ocean Program project in Galveston Bay, Texas, designed to compare ten created *Spartina alterniflora* marshes with five natural marshes. The created marshes were mainly transplanted on dredged material and ranged in age from 3 to 15 years at the time of sampling (fall 1990, spring 1991). Compared with natural marshes, above-ground plant biomass was equal or higher in most created marshes while below-ground biomass and sediment organic content was consistently lower in created marshes. Densities of juvenile fishery species within the marsh vegetation were estimated using a drop enclosure. Grass shrimp, commercial penaeid shrimp, blue crabs, pinfish, and gobies predominated. Created marshes generally supported lower numbers of natant macrofauna, especially juvenile brown shrimp, white shrimp, and blue crabs. Preliminary results from a caging study in the marshes indicated that growth rates of juvenile brown shrimp were similar in created and natural marshes, but survival in cages was significantly lower in created marshes than in natural marshes. A predator enclosure study in the marshes suggested production of benthic annelids was greater in natural marshes. There was little evidence in any of the results to indicate that utilization and value of the created marshes was increasing based solely on marsh age.

Minello, T. J. and R. J. Zimmerman. "Utilization of natural and transplanted Texas salt marshes by fish and decapod crustaceans." Marine Ecology Progress Series 90 (1992): 273-285.

The objective of this study was to compare three transplanted *Spartina alterniflora* salt marshes (2-5 years in age) with adjacent natural marshes on the Texas coast. Samples were collected only in the spring, limiting conclusions to this season, but the use of replicate marshes allowed a test of the null hypothesis that transplanted marshes on the Texas coast were equivalent to natural marshes. Quantitative drop enclosures (2.6 m² area) were used to collect juvenile fishes and crustaceans on the marsh surface. Above-ground density and biomass of macrophytes were also measured within these enclosures, and sediment cores were collected to examine sediment macro-organic matter (MOM) and benthic infaunal densities.

Mean values for stem density and above-ground biomass of *S. alterniflora* were consistently higher in the transplanted marshes, and the difference was statistically significant for stem density. Macro-organic matter in the upper 5 cm of sediment was significantly lower in the transplanted marshes. Densities of polychaetes and amphipods within transplanted marshes were positively correlated with this MOM. The transplanted marshes had significantly lower densities of decapod crustacea (primarily daggerblade grass shrimp, *Palaemonetes pugio*, and juvenile brown shrimp, *Penaeus aztecus*) compared with natural marshes. This reduced utilization may have been a response to low densities of benthic food organisms, and densities of decapods were positively correlated with densities of prey in sediment cores. In contrast to the utilization pattern of decapods, densities of fish (dominated by the darter goby, *Gobionellus boleosoma*, and pinfish, *Lagodon rhomboides*)

were similar between natural and transplanted marshes. These small fish may rely on salt marshes more for protective cover than for enhanced food resources, and above-ground structure in the transplanted marshes may have adequately provided this function.

The authors stressed that comparisons of functional equivalency between natural and transplanted salt marshes require adequate information on how salt marshes actually function for fish and decapod crustacea. For example, the use of prey density as an indicator of food value in a marsh can be misleading unless trophic pathways are well understood and access to the marsh surface is considered.

Minello, T. J., R. J. Zimmerman, and E. F. Klima. "Creation of fishery habitat in estuaries." In: Beneficial uses of dredged material; Proceedings of the first Interagency Workshop, 7-9 Oct. 1986, Pensacola, Florida., eds. M. C. Landin and H. K. Smith. 106-117. US Army COE, WES: Tech. Rep. D-87-1, 1987.

This paper discusses the importance of estuarine habitats, including salt marshes, in supporting the productivity of coastal fisheries. Extensive wetland losses have caused an increased willingness to create coastal salt marshes with dredged material. These marshes, however, are usually established over subtidal bay bottom. In order to properly assess the impacts of these habitat exchanges, the relative value of the involved habitats for fishery species must be known. Research conducted in fishery ecology and habitat functions at the Galveston Laboratory of the National Marine Fisheries Service is reviewed. Preliminary data is also reported indicating that transplanted *Spartina alterniflora* marshes support lower densities of nekton than natural marshes. Subsequently, these data have been analyzed fully by Minello and Zimmerman (1992). A cooperative program to create fishery habitat is also discussed; this program was in the initial stages of development between the NMFS and the U.S. Army Corps of Engineers.

Minello, T.J., R.J. Zimmerman, and R. Medina. "The importance of edge for natant macrofauna in a created salt marsh." *Wetlands* (in press, 1994).

The relationship between marsh edge and animal use was examined in a planted *Spartina alterniflora* marsh located in the Galveston Bay system of Texas. A completely randomized block experimental design was used with each of four blocks containing a control and experimental sector. Marsh edge was increased through the construction of channels in experimental sectors. Channel construction had no detectable effect on marsh surface elevation. Effects of these simulated tidal creeks on habitat use were examined by sampling nekton at high tide with drop enclosures both on the marsh surface and within the channels. Crustaceans dominated the nekton, and use of the marsh surface in experimental sectors was significantly higher than in controls; densities of brown shrimp *Penaeus aztecus*, white shrimp *P. setiferus*, and daggerblade grass shrimp *Palaemonetes pugio* were 4.6 to 13 times higher near the channels. Polychaete densities in marsh sediments were also significantly higher near channels, and densities of decapod predators were positively correlated with densities of these infaunal prey. Thus, channel effects on natant decapods may have been related to the distribution of prey organisms. However, increased densities of natant fauna along the channel edge may simply reflect a requirement for departure from the marsh surface at low tide. Marsh-surface densities of small bait fishes, bay anchovy *Anchoa mitchilli* and the inland silverside *Menidia beryllina*, also

increased near channels, but highest densities of these fishes were in the creeks themselves. The abundance and distribution of juvenile blue crabs *Callinectes sapidus* and gulf marsh fiddler crabs *Uca longisignalis* were not affected by the addition of experimental channels. Overall, the study results indicate that habitat value of created salt marshes can be enhanced by incorporating tidal creeks into the marsh design.

Morrison, J. and P. Williams. "Warm Springs marsh restoration." In: Urban wetlands: proceedings of the national wetland symposium, in Oakland, CA, eds. J. A. Kusler, S. Daly, and G. Brooks, Oakland, CA: Assoc. State Wetland Managers, 340-349, 1988.

This paper discusses the design and monitoring of the 81-ha tidal restoration portion of a previously diked wetland, the Warm Springs marsh area near Fremont, California. The area is at the southern end of San Francisco Bay. The design was made by a team of a landscape architect, a biologist, and a hydrologist. The objectives of the restoration were: 1. to establish an Alkali bulrush marsh as is found outside the diked area, 2. to enhance *Salicornia virginica* growth for habitat for the salt marsh harvest mouse, an endangered species, 3. to provide fill and flood protection for the industrial park, 4. to improve tidal circulation and water quality to the area, and 5. to improve public access to the area for fishing, hiking, picnicking, and bird watching.

The planned topography for the area included terraces and peninsulas along the embayment which would provide area for pickleweed growth. Elevated areas along the dikes were retained as refugia during extremely high water events. A main channel was established connecting the embayment with Coyote Slough at the south end, and a minor connection with Mud Slough at the north end. Channel erosion deepened and widened the south entrance during the first year as expected in the design. This channel stabilized within 15 months. The tidal inflow brought in seeds from adjacent marshes and *Salicornia virginica* became established along the higher terraces as planned. *Spartina sp.* started growing among the pickleweed at its lower elevations, but Alkali bulrush did not invade the restoration area as was expected it would. Increased use of the area by bird, fish and mammals use of the area was documented.

Although the Alkali bulrush did not become established, this carefully designed project is likely to be considered a success. Most of the goals of the restoration project are being met, including restoration of many natural marsh functions. The project is a good example of what careful and knowledgeable planning can do for wetland restoration.

Moy, L. D. and L. A. Levin. "Are *Spartina* marshes a replaceable resource? A functional approach to evaluation of marsh creation efforts." Estuaries 14 (1991): 1-16.

This study compared a 1-3 yr old man-made *Spartina alterniflora* salt marsh in Dills Creek, North Carolina with two adjacent natural marshes. Comparisons were made of sediment properties, infaunal composition, and *Fundulus heteroclitus* use of the marshes. Sediments and infauna were sampled in spring and fall of 1987 and spring and summer of 1988 along three isobath transects (8, 28 and 48 cm above MLW) in each marsh. On each transect sediment cores (3.2 to 4.7 cm diameter) were taken to a depth of 4 to 5 cm. During three of the above sampling periods, pit traps, consisting of plastic wash tubs buried in the

sediment, were used to collect juvenile *Fundulus* at five sites in each marsh. Gut analyses were also conducted on the collected fish.

Sediment grain-size was similar in the planted marsh and the natural marsh that shared the same drainage system (east marsh), but sediments were finer in the west marsh. Detritus and sediment organics were higher in both natural marshes than in the planted marsh. Sediment macrofauna in the natural marshes were dominated by the oligochaete *Monopylephorus evertus*, while the macrofauna of the planted marsh was dominated by polychaetes that occur nearer the sediment/water interface such as *Streblospio benedicti* and *Manayunkia aestuarina*. Meiofauna species composition was similar between east and planted marshes, the only two compared. Total mean density of meiofauna was about twice as high in the east marsh, but high variances prevented the detection of statistically significant differences.

Fundulus collected in pit traps on the natural marshes had mainly detritus and insects in their guts, while fundulids in the planted marsh had polychaetes and algae. These dietary differences reflected the available infaunal prey in the marshes. The number of fish caught in the pit traps was consistently higher in the natural marshes than in the planted marsh. The authors suggested that this increased catch indicated that the natural marshes supported larger populations of fish despite the observed dietary differences. Low *Spartina* stem densities in the planted marsh may have provided inadequate protection from predators or insufficient spawning sites. The different catches of *Fundulus* in pit traps, however, may also have been the result of variable catch efficiency in the marshes. The surface area of marsh sampled by the traps was unknown and may have varied among the marshes. Differences in the occurrence of depressions (natural 'pit traps') on the marsh surface may also have affected catches. This problem with pit traps, however, probably had little effect on the study conclusions that the planted salt marsh was ecologically different from the natural marshes in the area, and that salt marshes should not be treated as a replaceable resource.

Munro, J. W. "Wetland restoration in the mitigation context." Restoration & Management Notes 9 (1991): 80-86.

The author reviews some of the problems encountered in the permitting and mitigation process for wetland alteration. The process was characterized as costly, slow, and not well regulated (little standardization, and little checking of mitigation projects). Guidelines for planning a restoration project have not been formally established by responsible agencies, but several important variables to be considered in the planning, execution, and monitoring processes for a project's success were suggested, defined and briefly discussed. They included: sizing up the area, describing or defining the site, modeling the project, establishing constant reference points for monitoring, maintaining authenticity, defining the goals, developing a conceptual plan, specifying net gain, defining the pattern of the system, setting the ecological context of the project, defining the education potential, describing the area hydrology, planning for the long-term, scheduling, allowing for setbacks, having buffer zones, maintaining holism and continuity, using inoculants, making wildlife structures, preventing erosion, keeping some flexibility in the plan, using horticulture, using planting patterns, having sufficient sock for the project, accepting providence, protecting against exotics, developing biodiversity, ordering supplies, recycling

plants and soil if possible, planning for transportation, using soil banks, keeping accurate records, and publishing reports to help others.

The author concluded that without continuity of the designer overseeing the construction and monitoring its progress, projects could fall short of their goals. Timing and scheduling of planting was underscored as very important for obtaining that initial growth and plant establishment. Documentation of the entire project was deemed worthwhile even if the project failed--a lesson learned, and hopefully not repeated. Better area-wide planning of restoration activities could be done if a list of mitigation areas and projects were accessible to the public.

Nailon, R. W. and E. L. Seidensticker. "The effects of shoreline erosion in Galveston Bay, Texas." In: Coastal wetlands, ed. H S. Bolton. 193-206. New York: Amer. Soc. Civil Engineers, 1991.

The use of planted *Spartina alterniflora* is discussed for shoreline erosion control in Trinity Bay and East Galveston Bay, Texas. *S. alterniflora* was planted at four sites, and about 1825 meters of shoreline were vegetated. Before planting, erosion rates at the four sites ranged from 0.5 to 2.6 m per year, and this rate appeared related to fetch length. Christmas trees, plastic snow fence, and used cargo parachutes were installed to reduce shoreline energy while young transplants became established. Survival of transplants ranged from 60 to 70% after a year for the three sites protected from wave energy. At the one site with no protection, none of the transplants survived.

Fishery species were sampled with bag seines and cast nets in the waters just offshore from one successful transplant site and from the site with no surviving plants. The samples were collected in August and October of 1988, about 2 years after transplanting. White shrimp, Gulf menhaden, and striped mullet were most abundant in the catches and 87% of the finfish and crustaceans were collected adjacent to the successful transplant site. Although gear catch efficiencies may have varied among sites, and the samples were not collected within the vegetation, the large differences in observed catches suggested that the shoreline area with transplanted *Spartina alterniflora* was utilized more by these fishery organisms than was the bare shoreline.

National Research Council. Restoration of Aquatic Ecosystems: Science, Technology, and Public Policy. Washington D.C.: National Academy Press, 1992.

This book reviews historic degradation of aquatic ecosystems in the U.S. Of particular relevance were chapters 6 (Wetlands) and 8 (National restoration strategy: basic elements and related recommendations). The wetlands chapter included descriptions and discussion of the functions of wetlands, a brief history of the loss of wetlands in the U.S., a review of the potential for restoration actions, and a review of restoration research and technology. It also included a discussion of the difficulties that face wetland restorations, including ecological, biological and institutional constraints. Some discussion about what constitutes "successful" restoration was included, as was discussion about the research needs to assist wetland restoration. Among the most significant needs appeared to be the assessment of the restoration of functional equivalency, and what technology could be developed to accelerate the restoration process in each project.

Chapter 8 contains a proposal for a national restoration strategy. The elements come under four headings: (1) goal setting, (2) priority setting and decision-making principles, (3) redesign of federal policies and programs, and (4) innovation in financing.

This book provides an excellent base of information about the current status of restoration activities and thought in the U.S. It shows there is a large concern by ecologists, biologists, and naturalists that many important aquatic ecosystems are being destroyed with a consequent loss of associated fish and wildlife. It also shows how this loss of natural resources will continue if new policies are not established by state and federal agencies to offer better protection to aquatic ecosystems.

Newling, C. J., M. C. Landin, and S. D. Parris. "Long-term monitoring of the Apalachicola Bay wetland habitat development site." In: Proceedings of the tenth annual conference on wetland restoration and creation in Tampa, FL, ed. F. J. Webb, Jr. Tampa, FL: Hillsborough Community College, 164-186, 1983.

This paper describes changes in the man-made cordgrass marshes that occurred in the Apalachicola Bay wetland habitat development site, an area of dredged material deposition that was protected from wave action by dikes. The area, known as Drake Wilson Island, covers about 5 ha along the northern shore of Apalachicola Bay, FL. In 1976, *Spartina alterniflora* was planted in various configurations in the intertidal zone, and *Spartina patens* was planted in the supratidal zone. Based on randomly selected 0.5-m² quadrats, within two growing seasons all *S. alterniflora* plots that began with plants on 1-m centers or less had filled in for 100% coverage; those on larger centers had mostly been washed out, or were just surviving. Similar coverages were found for *S. patens* using 0.25-m² quadrats. By the second year, 42 species of plants had invaded the planted site; *Distichlis spicata* was the dominant invader in areas between the two cordgrasses.

By 1982, six years after the planting, several changes had occurred to the planted areas. *Scirpus robustus* was now found regularly interspersed with *S. alterniflora* along the landward margin. *S. patens* coverage had been greatly reduced by invading dune-type vegetation. Diversity in the dunes and marshes had increased to 97 species of plants by this time. The plant assemblage on the island was as diverse as those at the nearby natural marsh sites examined for comparison.

Based on visual observations, bird usage of the Drake Wilson Island was even greater than at the natural marshes. Wading birds were commonly found feeding at Drake Wilson Island, and included many clapper rails, plovers, herons, egrets and terns. No sampling was done for fisheries or fisheries food-chain organisms. Some concern was voiced about the eroding dikes and thus about the longevity of this marsh. Inferences were that repairs may become necessary to maintain the marsh.

Niering, W. A. "Vegetation dynamics in relation to wetland creation." In: Wetland creation and restoration: The status of the Science. Part 2. Perspectives., eds. J. A. Kusler and M. E. Kentula. 479-486. Washington, DC: Island Press, 1989.

Ecological concepts involved in the successional development of wetland vegetation are discussed. Discovering the hydrological basis of a wetland leads to a better understanding of how perturbations to water flow through a marsh will affect its vegetation. Disturbance is a natural part of coastal wetlands and depending on the severity of the

disturbance, the induced changes in wetland vegetation will vary from minor and short-termed to severe and persistent. Creating a wetland that will persist requires creating an area with an appropriate and stable topography and hydrology; once vegetation becomes established it should persist.

Pacific Estuarine Research Laboratory. A manual for assessing restored and natural coastal wetlands with examples from southern California. California Sea Grant Report No. T-CSGCP-021. La Jolla, California: 1990.

This manual discusses the problems encountered in assessing functional equivalency for created salt marshes. The basic premise is that man-made wetlands should be expected to replace natural wetland functions. Evaluation procedures are presented for wetland functions related to hydrology, water quality, sediment and nutrient dynamics, vegetation, and support of various animal populations. The main purpose of the manual is to standardize methods of assessing functions in created wetlands. Criteria for success need to be stated as testable goals that can be achieved through reasonable monitoring programs.

The assessment method recommended for fishery species was repetitive seining of enclosed areas in tidal creeks or ponds until populations decline. This seems extreme. Because salt marsh vegetation in southern California is infrequently flooded in comparison with other coastal regions, sampling of nonvegetated habitats within the marsh complex may be sufficient to assess fishery use. In other regions, however, where the vegetated areas in salt marshes are directly utilized by fishery species, animal densities on the marsh surface (within the vegetation) are important, and the quantitative sampling of this habitat requires different techniques.

Patience, N. and V. V. Klemas. "Wetland functional health assessment using remote sensing and other techniques: literature search." NOAA. Tech. Memo. NMFS-SEFSC-319, 114, 1993.

This report reviews techniques for determination of the functional health of wetlands using remote sensing and other techniques. It provides a concise review of concepts and technology, and points to areas of future research. Each section of the report is supported by an extensive review of pertinent literature. With improving technology and information transfer, system capabilities will make greater accuracy and timeliness possible for health assessments of many more wetlands and their functions. Remote sensing will also allow for monitoring of many more restored and created wetlands on an annual basis.

Reppert, R. "Wetlands mitigation banking concepts." U.S. Army Corps of Engineers. IWR Rep. 92-WMB-1, 33, 1992.

This report provides general and specific information about wetland mitigation banking. Mitigation banks that are functioning are reviewed and evaluated. The potential of mitigation banking for achieving "no net loss" of wetlands is also evaluated.

Wetland mitigation banks provide advanced compensation for unavoidable wetland losses by creation, restoration, enhancement or preservation of other wetlands of equivalent value. Such banks are usually large tracts whose tangible and intangible values are equated to credits. As wetlands are altered elsewhere, credits are withdrawn from the bank. When the credits are exhausted the bank is closed, and new projects must seek

mitigation credits elsewhere. Banks would appear to be very effective in mitigation efforts particularly when many small mitigation projects would not yield any one area of sufficient size to support some larger wetland animals that require a large area for a viable habitat.

Mitigation banks are sponsored by industries, highway departments, port authorities, federal projects, and commercial interests. Of the 37 banks identified in this report, 19 were sponsored by state highway departments, 8 by ports, and 7 by land developers. In 1988, there were only 12 banks, now there are 37, and there are plans for 65 new banks in the next couple of years. This shows a significant development towards banking for solutions to mitigation.

Banking has its good and bad features depending on your point of view, and such an increase in banks could be disturbing news. Some of the good features listed were:

1. A bank is generally a large block of land which can support more species, some requiring large areas as habitats, than small single mitigation projects.
2. A large unit in a bank should be more economical to manage than many small dispersed units.
3. A bank generally provides for mitigation before adverse impacts to wetlands occur, otherwise there could be loss of habitat and ecological functions for several years while a mitigation project develops to its full functioning state.
4. A bank can be more thoroughly planned, managed, and incorporated into regional wetland requirements because there is plenty of time for these actions.
5. A bank can serve more social value than small spread-out mitigation projects; elements for public appreciation and education can be offered in a larger area.
6. A bank can expedite conflict solution because it already exists and its costs are known.
7. A bank can provide a lifetime monitoring program to protect the habitat.

The establishment of a bank indicates permanence for the habitat as far as man's actions are concerned.

The main negative features of banks were listed as:

1. The planning may be less than adequate for the entire area and lifetime of the bank.
2. The existence of a bank may make it too easy to allow other wetlands to be destroyed where there may have been alternative actions that could have preserved them. Banks may act as short circuits to the regulatory process.
3. There is uncertainty as to the real value of a bank in terms of credits. Credit value has been difficult to establish for ecological functions of wetlands, and thus, crediting and debiting will always be controversial.
4. A bank may not provide for in-kind replacement, thereby potentially allowing for a particular habitat to be eliminated from a watershed.
5. There are uncertainties about management techniques for wetland habitats in general, and, when applied to a large bank, poor management could prove ecologically and economically costly.
6. The preservation of wetlands in the bank would only be beneficial if the lands were subject to loss to begin with; if not, there would be no replacement benefit by preserving the bank lands.

7. When all bank credits are exhausted, there is always the question of who is going to own the land and be responsible for its maintenance.

Mitigation banking is likely to increase as we press to use wetlands. To obtain the benefits offered by mitigation banking, we must support excellence in planning each bank and excellence in maintaining each bank. Yearly credits should be issued to each bank that would go to support the maintenance and monitoring. Certainly, the presence of mitigation banks should not be used as an excuse to destroy natural wetlands; avoidance of damage to natural systems should be emphasized.

Reubsamen, R. N. "National Marine Fisheries Service efforts in Texas to tailor mitigation to specific wetland types." In: Proceedings of the National Wetland Symposium: Mitigation of Impacts and Losses, in New Orleans, LA, New Orleans, LA: Assoc. State Wetl. Manag., 424-425, 1986.

This paper briefly summarizes the NMFS perspective toward mitigation for loss of wetlands in Texas. Avoidance of impacts on natural wetlands is the goal. When that is not possible, replacement by creation or restoration of in-kind wetland habitat of equal or greater size is sought. The replacement wetland should be located as close to the impacted wetland as practical (without itself becoming impacted). Increased manpower will be needed to review permit applications to alter wetlands (processed by the Corps of Engineers) and to monitor all the wetland mitigation efforts. Such monitoring is currently being accomplished for only a few projects.

Riggs, S. "Distribution of *Spartina alterniflora* in Padilla Bay, Washington, in 1991." Washington Dept. Ecology, Padilla Bay National Estuarine Research Reserve. Tech. Rep. 3, 63, 1992.

The objectives of this study were to document the distribution and spread of *Spartina alterniflora* in Padilla Bay, Washington. Baseline maps and data were available from a 1987 study by Wiggins and Binney (1987). Cordgrass stands that had been mapped in 1987 were mapped again in August and September, 1991. Comparing maps showed the cordgrass spread 1-2 m/yr. Area surveys also located several additional stands of *S. alterniflora*. The total area now covered was estimated at 4.8 ha compared with about 2.7 ha four years earlier. Only very minor die-back was found in central portions of some of the stands, and *Salicornia virginica* was found inhabiting those areas. There was still no evidence of flowering by *S. alterniflora* in Padilla Bay. (Note: Although not included in this report, S. Riggs informed us that 1992 was an extra warm summer for Padilla Bay, and that *S. alterniflora* did flower.)

The impact of *S. alterniflora* in the estuaries of Washington is considerable. Studies are needed to examine the function of cordgrass in the northwest.

Roberts, T. H.. "Habitat value of man-made coastal marshes in Florida." In: Sixteenth annual conference on wetlands restoration and creation, in Plant City, FL., ed. F.J. Webb, Jr., Plant City, FL.: Hillsborough Commun. College, 157-179, 1989.

This study was undertaken to determine the quality of man-made wetlands as habitat for fish and wildlife, and therefore, the effectiveness of marsh creation as mitigation for losses of coastal wetlands. Fish, bird, and mammal populations at 21 man-made,

established, coastal marshes of various ages and 6 natural marshes were sampled and compared for similarities. The majority of the sites were dominated by *Spartina alterniflora*. Created marshes were located throughout northern and central Florida and ranged in age from approximately 1 to 10 years old. Vegetation characteristics were highly variable, but man-made sites that were properly planned, constructed, and maintained, were considered to serve as viable habitat for animals normally associated with natural coastal marsh systems. Factors influencing site use by various animal groups, and suggestions for the design of future mitigation efforts are discussed.

A quantitative comparison of aquatic animal utilization in created vs. natural marshes was not possible with the data presented. Sampling methods used in the study were not quantitative, and variability in characteristics of the different marshes (size, configuration, location, etc.) may have influenced the results. However, results show qualitatively that established man-made wetlands have the ability to support the same animal species as natural wetlands.

Roberts, T. H. "Habitat value of man-made coastal marshes in Florida." Vicksburg, MS: U.S. Army Corps Eng., Waterways Experiment Station. Tech. Rep. WRP-RE-2, 42, 1991.

The objectives of the study were to determine the effectiveness of marsh creation as mitigation for losses of natural coastal marshes along Florida's Gulf and Atlantic coasts. Only 55% (21 of 38) of the man-made *Spartina alterniflora* marshes visited in Florida estuaries were identifiable or deemed successful enough to be sampled; 1 man-made *Juncus* marsh was also studied. The marshes ranged in size from 0.004 to 2.8 ha, and were compared with four *Spartina* and two *Juncus* natural marshes that ranged in size from 0.20 to 3.2 ha. The 22 man-made marshes ranged in age from 1 to 10 years old, but 7 were 1-2 yr old, 6 were 2-3 yr old, and 6 were 3-5 yr old. Data were gathered on soil characteristics such as substrate texture, particle size and organic content from 3 to 5 samples per site. Vegetation was sampled using stratified random transects with the point-intercept method, and data on species composition, percent cover, stem density, and height of *Spartina* plants were obtained. Below-ground biomass of *Spartina* was measured from 7-cm-diameter cores taken to a depth of 18 cm. Fish data were collected using fyke nets (2 per site), Breder traps, and a Wegener Ring net. The fyke nets were set at high tide, and fish and motile invertebrates were collected as they left the marsh with the ebbing water. The Wegener Ring net was used to collect fish in tidal channels in some marshes. Birds were surveyed in each marsh on three consecutive days, and bird calls were also identified and recorded. Mammals were trapped at each site using Sherman Box and Museum Special Snap traps, one each per station, 5-7 stations per marsh, and two nights in each marsh. Tracks and other signs of larger mammals were also noted.

Sufficiently large variability was found in the data for each variable sampled in each age group of man-made marshes that no differences could be detected among the age groups. The author attributed much of the variability to marsh size and shape. Organic matter averaged from 0.2 to 10.1 % in the man-made marshes, and showed no statistically significant increase with age ($r^2=0.04$). *Spartina alterniflora* cover in the man-made marshes ranged from 40% on a 2 yr old site to 80% on a 1 yr old site, and averaged about 60% for all. Many marshes had other species of plants growing in adjacent zones, and these added an

average of 10% to the total marsh cover for the sites. Below-ground biomass estimates ranged from 831-3,429 g/m² among man-made marshes and varied widely among sites. The minor correlation found between the age of a marsh and its below-ground biomass ($r^2=0.14$) did not appear to be significant. Only 50% of the man-made marshes supported marsh-dependent birds, and abundances were relatively low when birds were observed. Size and configuration of the man-made marshes were limiting, as were nearby development activities that appeared to be intolerable for the birds. Raccoons and marsh rabbits were found using about 30% of the man-made and natural marshes. Small mammals such as the cotton rat, rice rat, Norway rat, black rat, and house mouse were collected in man-made marshes. Their use of the marshes was probably influenced more by vegetation and nearby source of colonists than by age of the marsh.

Many of the fish species found in natural marshes were found in the man-made marshes. Again, no correlation was found between similarity index values (of the fish communities in a man-made marsh and the nearby natural marsh), and the ages of the man-made marshes ($r^2=0.003$). Most man-made marshes had three to five of the common eight species characteristic of natural marshes. Abundances in the man-made marshes were within the range of the natural marshes. Because the sampling was not quantitative in the sense that it could provide numbers of fish per m² of marsh, the abundance values were only taken as general indicators of marsh use. Other aquatic macroinvertebrates found in both man-made and natural marshes included *Littorina irrorata* and *Uca* spp. Again, there was great site-to-site variation in abundances of these animals.

This report points out the great variability in man-made and natural marshes. This variability makes it difficult to generalize from a study of only one or two marshes. The study also points out the difficulties in obtaining quantitative samples of marsh fauna; despite the use of multiple gears to sample fish and motile invertebrates, quantitative comparisons could not be made among marshes. Causes for differences among all the marshes were not obvious, though size, shape, and surroundings were likely sources for differences.

An important finding of this study was that established techniques to create or restore *Spartina alterniflora* marshes frequently appeared to be neglected, improper elevation of the substrate and a lack of protection from erosive waves were common in created marshes. Even when marsh creation was successful, other infringements and destructive acts were common when almost any other use could be made of the marsh or the adjacent land. Only by carefully incorporating species habitat requirements into project design, and locating wetlands where their habitat value can be realized, will mitigation efforts be fruitful and the loss of natural wetlands be offset.

Sacco, J. "Infaunal community development of artificially established salt marshes in North Carolina." Ph.D. Thesis, North Carolina State University., 1989.

Benthic infaunal populations were examined at six locations along the North Carolina coast; at each location, a transplanted *Spartina alterniflora* marsh was sampled along with a nearby natural marsh. The created marshes ranged from 1 to 17 years of age. The six locations differed in tidal range, salinity, and substrate composition. Infauna were dominated by annelid worms; polychaetes made up over 53% and oligochaetes 35% of the infauna in samples. Taxonomic composition and trophic diversity appeared similar among

the marshes, but infaunal densities were generally lower in the transplanted salt marshes. Sediment organic matter was also lower in created marshes. There appeared to be a relationship between high organic matter and high infaunal densities, but this relationship varied with local conditions. There was no apparent relationship between marsh age and the development of the infaunal community.

Sampling in natural marshes was conducted along the entire elevational range within the marsh, while sampling in created marshes was restricted to the original area of planting. Because elevation affects infaunal densities, this sampling regime may have affected results.

Seneca, E. D. and S. W. Broome. "Restoring tidal marshes in North Carolina and France." In: Restoring the nation's marine environment, ed. G. W. Thayer. 53-78. College Park: Maryland Sea Grant College, 1992.

This review article covers various techniques used to restore and create tidal marshes. The important factors discussed include site selection, elevation, slope, tidal range, wave climate, salinity, soil properties, cultural practices, sedimentation, wildlife impacts, post-planting maintenance, and traffic. The authors conclude that the primary goal of most wetland projects is to assist the system so that it can eventually attain functional equivalency with a natural wetland. The initial objective, therefore, is to restore the dominant emergent vegetation. These macrophytes then serve as substrata for other organisms, produce organic matter for the food web, provide habitat for organisms, and buffer shorelines from waves. Macrophytes also cycle nutrients and trap and stabilize sediments. Thus, establishment of a thriving macrophyte stand provides the basis for establishing fish and wildlife functions. These functions may or may not follow, but the establishment of the macrophytes is a key step in any restoration or creation project.

Seneca, E. D., S. W. Broome, and W. W. Woodhouse, Jr. "Comparison of *Spartina alterniflora* Loisel transplants from different locations in a man-initiated marsh in North Carolina." Wetlands 5 (1985): 181-190.

The objectives of this study were to determine how height form and latitude of origin might influence the variability of *Spartina alterniflora* when plants are transplanted to other sites as they are in restoration projects. Plants from four locations (short and tall forms from Beaufort) were transplanted on 0.9-m centers in a randomized split-plot design planting at the southern-most site. The grow-out site was a recently deposited (60 days prior to planting) area of dredged material that was 96% sand, 1% silt and 3% clay. Height, basal area, aboveground dry weight, culm density, and number of flower heads were measured periodically over five growing seasons.

Results showed that height and time of flowering of each location-type were maintained at the new site for several growing seasons. However, under the influence of elevation, the short form became taller when grown at lower elevations, and the tall form became shorter when grown at higher elevations. Within two growing seasons, the transplanted marsh had the standing crop equal to local natural marshes. Results also showed that yearly standing crop could vary as much as 45% in the transplanted marsh, which was the same as has been found for natural marshes. This variability should be considered in evaluating the success of a restored or created marsh.

Seneca, E. D., S. W. Broome, and W. W. Woodhouse, Jr. "The influence of duration-of-inundation on development of a man-initiated *Spartina alterniflora* Loisel marsh in North Carolina." *J. Exp. Mar. Biol. Ecol.* 94 (1985): 259-268.

This study was conducted in a *Spartina alterniflora* salt marsh that was planted in 1971 near Snow's Cut in North Carolina. Above-ground growth parameters measured included plant height, culm density, dry weight, basal area, and the number of flowers. Below-ground biomass was also measured. These growth parameters were measured at different elevations that corresponded to daily tidal flooding durations of 4, 7, 9, and 11 hours. The marsh was sampled during growing seasons 2, 3, 4, 5, and 12. During the second growing season, maximum above-ground growth occurred in the 7-h inundation zone. During growing seasons three through five, all growth variables generally increased with tidal inundation. By the twelfth growing season, *S. alterniflora* was completely displaced in the 4-h inundation zone by other marsh plants. *S. alterniflora* maintained dominance in the 9 and 11-h inundation zones and spread into areas of even lower elevation than originally planted.

Shisler, J. K. "Creation and restoration of the coastal wetlands of the northeastern United States." In: Wetland creation and restoration: The status of the Science. Volume I: Regional overviews. EPA/600/3-89/038., eds. J. A. Kusler and M. E. Kentula. 145-174. Corvallis: U.S. Environ. Prot. Agency, 1989.

This chapter characterizes the wetlands in the northeastern U.S., and describes the types of wetland restoration activities prevalent in that region. Most of the intertidal zone in salt and brackish marshes is dominated by smooth cordgrass, *Spartina alterniflora*. The area just above the intertidal zone is dominated by saltmarsh cordgrass, *Spartina patens*, mixed with some salt grass, *Distichlis spicata*. Several thousand hectares of this upper zone was diked off from tidal flows and the area used to grow salt hay (a combination of *Spartina patens*, *Distichlis spicata*, and *Juncus gerardii*). A couple of thousand hectares have recently been reopened to tidal circulation, but on a controlled-flow basis so the areas can be managed for waterfowl.

Factors to be considered in the design of creation or restoration projects are briefly described in this chapter. Some of the factors are: location of the project, hydrology at the site, topography of the site (elevation, slope, size), texture and quality of the substrate, species of plants to be used (matched to the environmental parameters expected from the design), adequacy of buffer zones, protection from pests, and degree of monitoring that will be required for success.

Research is needed to make restoration projects even more successful than they have been. Fourteen topics for research are given. Some concern plant species environmental requirements, tolerances of transplanting practices, and propagation potential. Other topics range from understanding pest control to simple developing a decent inventory of the wetlands.

Sinicrope, T. L., P. G. Hine, R. S. Warren, and W. A. Niering. "Restoration of an impounded salt marsh in New England." *Estuaries* 13 (1990): 25-30.

Vegetation changes of a tidal marsh in Stonington, CT, were documented for a 20 ha marsh 9 years after the reintroduction of tidal flushing. Vegetation was examined in aerial photography by comparing data from a study of the area by Hebard in 1976, with data obtained in 1986. Hebard's transects were then revisited in 1987 and 1988 to determine current coverage by species using the same line intercept method. Two extensions were added to transects T-3 and T-4 to cover what were previously mudflats.

Typha angustifolia decreased from 74% to 16% coverage, due to increased salinities over much of the area. *Phragmites australis* increased slightly as some of the *Typha* began dying out, but it too was exhibiting signs of stress in its stunted new growth. *Spartina alterniflora* increased from <1% coverage to 45%, invading much of the mudflats and intertidal area vacated by *Typha*. High marsh species such as *Distichlis spicata*, *Spartina patens*, *Juncus gerardi*, and *Salicornia europaea* also increased in areas upland from the *S. alterniflora* zone.

No systematic survey of the marsh and creek was made to document changes in utilization by aquatic species. However, increased use by ducks, shorebirds, and migratory birds was noted.

Soil Conservation Service. "Wetland restoration, enhancement, or creation." In: Engineering Field Handbook, ed. Soil Conservation Service. 79. 13. Washington DC: U.S. Department of Agriculture, 1992.

This is chapter 13 in the US DOA Soil Conservation Service's Engineering Field Handbook. It discusses wetland processes, functions and characteristics, and describes (with example data) the planning that is needed for a successful restoration/creation project. Consideration must be given to site selection, design, implementation, monitoring, and management. A wetland planning checklist is given in Appendix A, as is a monitoring site visit checklist in Appendix B.

Steinke, T. J. "Hydrologic manipulation and restoring wetland values: Pine Creek, Fairfield, Connecticut." In: National wetland symposium: mitigation of impacts and losses. in New Orleans, LA, eds. J. A. Kusler, M. L. Quammen, and G. Brooks, New Orleans, LA: Assoc. State Wetland Managers, 377-383, 1986.

This article presents a history of Pine Creek marsh in Fairfield, CT, and a description of the restoration activities associated with reestablishing tidal flows to this marsh. A detailed plan incorporated information about the area from hydrologists, biologists, public health officials, and civil engineers. The restoration began with 28 ha of an almost pure stand of 3-4 m tall *Phragmites australis* and has resulted in reestablishing the *Spartina patens* upper marsh and the *S. alterniflora* lower marsh after only five years of tidal flow renewal. Restoration was accomplished by building new dikes containing self-regulated tide gates before removing the old dikes. The restoration has also maintained flood protection and has eliminated 25-30 annual peat and *Phragmites* fires that threatened adjacent homes. A greater variety of birds has been noted in the marsh, and hunting and fishing has increased. Another 81 ha of marsh is under consideration for restoration action.

Stout, J. P. "Evaluation of coastal wetlands mitigation, Alabama." Alabama Dept. Environ. Management. Tech. Rep., 45, 1990.

The objectives of this study were to evaluate coastal mitigation work in Alabama. Recommendations were made that should improve success rates of coastal wetland restorations.

Of 12 wetland mitigation sites visited: no mitigation work had been started in three, work was incomplete in one, three were failures, two showed partial success, and three were deemed successful. The survey found very little monitoring of this mitigation work, and in most cases little was done to correct deficiencies when they were discovered during monitoring.

Problems with the mitigation system were varied. There was no standard format for a permit request that included all the needed information about a project to enable proper evaluation of the permit request. The limited documentation that was required was not routinely cataloged or available for examination and evaluation when needed. There was no central control or plan that could help direct mitigation efforts in Alabama. Such a plan or organization might act to preserve some of each kind of ecosystem in each watershed. A list of 25 factors to include in a plan was included.

Concern was expressed about a growing attitude that losses of natural wetlands could be mitigated by restoring or creating wetlands of equal value. The author suggested that we may be accepting a loss of valuable natural wetlands for equal areas of inferior quality created wetlands. Monitoring for 3-5 years after the restoration planting was recommended for marshes and submerged aquatic vegetation, and for 5-7 years for forested wetlands. There were 15 recommendations for monitoring which included photographic surveys from fixed markers and the use of aerial photos. Monitoring of vegetative, topographic, and hydrologic changes were also among the recommendations, along with utilization by waterfowl and aquatic fauna.

Because failures are largely ignored at present, changes in the system are sorely needed. A detailed mitigation design plan with clearly stated goals should be required for a permit to be granted. Concerned agencies must find staff to perform the monitoring, and performance of the work required by the permit must be enforced. Innovative approaches for mitigating loss of wetlands should be pursued. The process of mitigating an acre-planted for an acre-used is resulting in lost wetland functions as well as loss of fish and wildlife and their habitat.

Tanner, G. W. and J. D. Dodd. "Effects of phenological stage of *Spartina alterniflora* transplant culms on stand development." *Wetlands* 4 (1985): 57-74.

This study tested three phenological types of *Spartina alterniflora* for differences in survival and growth when used for marsh creation. Plants were dug from three natural donor marshes that had seedling, dwarf-form, and tall-form stands of *S. alterniflora*. Plantings were conducted at a dredged material site in Galveston Bay, Texas, and two substrates and three fertilizer treatments were also tested. Roots of all transplants were kept moist using wet burlap bags. Single culms were planted on 0.5-m centers in plots of 5 rows of 10 plants each. Treatments were configured in a randomized block design.

The seedling phenotype had a moderately superior initial survival and more rapid tiller production and biomass accumulation than the other two phenotypes. Seedling culms came from a newly established natural stand of plants that was in the process of rapid

expansion itself, while the other phenotypes came from mature stands that were in equilibrium.

The fine grain (silty) substrate supported more vegetative growth than the coarse grain (sandy) substrate, even though the sandy substrate had slightly better initial survival. Fertilizers had no apparent effects, but the root-stimulator treatment killed a significant number of transplants.

Thayer, G. W., ed. Restoring the Nation's Marine Environment. College Park, MD: University of Maryland Sea Grant College Program and DOC/NOAA/National Marine Fisheries Service, 1992.

This book brings together current philosophies, techniques, and recommendations for preserving, restoring, enhancing, and creating many kinds of habitats in the marine environment. Chapters on salt marsh restoration in southern California by Joy Zedler and in N. Carolina and France by Ernest Seneca and Stephen Broome emphasized the need to use appropriate techniques in the restoration and creation tidal marshes. Such techniques included detailed plans that considered many variables such as: site selection, elevation, slope, tidal range, wave climate, salinity, soil physical and chemical properties, sedimentation, variety of vegetation to be planted, and protective measures against herbivory and trampling.

Questions remained about how to restore all the functions associated with natural marshes. A particular concern was for restoration of faunal population dynamics, not just the habitat. Various studies indicated 2-10 years may be required for faunal dynamics of a created marsh to match those of a neighboring natural marsh. Long term marsh studies were expected to yield answers, and were recommended.

Underwood, S. G., G. D. Steyer, B. Good, and D. Chambers. "Bay bottom terracing and vegetative planting: an innovative approach for habitat and water quality enhancement." In: Proceedings of the eighteenth annual conference on wetland restoration and creation in Hillsborough, FL, ed. F. J. Webb, Jr. Hillsborough, FL: Hillsborough Community College, 164-173, 1991.

This paper presented the results of a project to prevent further shoreline and marsh erosion by waves, and to increase water quality in three ponds in a marsh along the north shore of Calcasieu Lake in the Sabine National Wildlife Refuge, Louisiana. There were 128 shallow bay terraces constructed and laid out in a checkerboard design in the ponds. *Spartina alterniflora* was planted on both sides of the terraces to stabilize them, trap sediment, and reduce the size of the wind generated waves. Preliminary results indicated reduced water turbidity in the terraced ponds, reduced size of wind generated waves in the ponds (greatly reduced fetch due to terraces), and good growth of cordgrass to stabilize the terraces.

The terracing technique may prove very useful in other areas of Louisiana for reestablishing marsh. Marshes that are subsiding may benefit greatly from terracing.

US Army Corps of Engineers. Engineering and Design. Environmental Engineering for Coastal Protection. Vol. EM 1110-2-1204. Washington DC: U.S. Army Corps of Engineers, 1989.

This manual provides guidance in environmental engineering for the protection of coastal areas. It summarizes information about various means of protecting shores, including the use of *Spartina alterniflora* marshes to reduce erosion. Sampling techniques are given for use in monitoring the effects of various projects.

Watzin, M. C. and J. G. Gosselink. "The fragile fringe: coastal wetlands of the continental United States." Louisiana Sea Grant College Program, Louisiana State University, Baton Rouge, LA; U.S. Fish and Wildlife Service, Washington, DC; and National Oceanic and Atmospheric Administration, Rockville, MD. 1992.

This report is an informative presentation of the nature and value of coastal wetlands in a popular format. It gives definitions, functions and needs, illustrated by examples that will help anyone to better understand the values of coastal wetlands and why their preservation is important.

Webb, J. W., Jr. "Establishment of vegetation for shoreline stabilization in Galveston Bay, Texas." Ph.D., Texas A&M Univ., College Station, TX, 1977.

The objectives of this research were to isolate and test plant species that would stabilize bay shorelines, to develop planting technology, to test wave-stilling devices, and to calculate a time-effort budget for a transplanting project. The study area was the north shoreline of East Bay, on the Anahuac National Wildlife Refuge, Chambers County, Texas. This area had an eroding bank that was receding at 1-2 m/yr. The soil was classified as loam or clay-loam texture, structurally unstable and subject to erosion. Thirteen plant species, two wave-stilling devices, and three planting techniques were tested.

Spartina alterniflora was the most successful species in the intertidal zone below MHW, but it required protection from wave impact. Wave-stilling fences of hay bales contained in chicken wire and in 14-gage welded mesh wire were unsuccessful. Tires on cables held in place by metal posts were somewhat successful, but lost effectiveness when they partially sank into the substrate. *Spartina spartinae* and *Spartina patens* were the most successful above MHW. Fertilizer was beneficial to plant growth in a natural marsh (control), but showed no significant benefit to the planted blocks.

Seeding an area failed to establish plants. Hand planting of tillers or culms was effective when protected by the wave stilling device (tire fence). Mechanical planting of tillers was also successful at low tide and behind a wave-stilling device.

Mechanical grading down of the banks to a smooth slope for establishing the planted marsh was not useful. Erosion of the loosened substrate occurred within several months.

Webb, J. W., Jr. "Salt marshes of the western Gulf of Mexico." In: Creation and restoration of coastal plant communities, ed. R. R. Lewis, III. 89-110. Boca Raton, FL: CRC Press, Inc., 1982.

This chapter reviews coastal salt marsh restoration and creation activities in the western Gulf of Mexico. There are six sections to the chapter which give excellent coverage to the topic. They include descriptions of: the plant community, the various levels of productivity of the salt marsh, the loss of this habitat through modification, specific projects, recommended techniques, and research needs.

The coastal marsh plant community in the western Gulf is quite similar in species composition to that along the Atlantic coast of the U.S. Smooth cordgrass, *Spartina alterniflora*, is the dominant species intertidally. It grows best from MLW to MHW. At a slightly higher elevation *Distichlis spicata* (salt grass) occurs. Overlapping *Distichlis* and moving slightly higher still, *Spartina patens* (saltmeadow cordgrass) is found, sometimes forming broad meadows. The higher margins of the salt meadows are shared with *Scirpus maritimus* (leafy threesquare), *Scirpus olneyi* (Olney bulrush), *Borrchia frutescens* (sea oxeye daisy), *Monanthochloe littoralis* (salt flat grass), *Limonium carolinanum* (sea lavender), *Batis maritima* (saltwort), *Salicornia bigelovii* (annual glasswort), *Salicornia virginica* (virginia or perennial glasswort), and *Iva frutescens* (bigleaf sumpweed or marsh elder) which often form clumps in this infrequently flooded area.

These salt marshes are very productive areas. Production of *Spartina alterniflora* can be 500 to 2800 g/m²/yr. Much of this production supplies the beginning material for a detritus based food web. The marshes act as feeding habitat that also offers protection for juvenile fish, shrimp and crabs of economic importance. Coastal marshes also serve as permanent habitat for otters, muskrats, nutrias, raccoons, and alligators, as well as winter habitat for thousands of ducks and geese.

Salt marshes are being lost every year through natural and man-induced causes. Subsidence and erosion are the two main natural causes, but man is contributing to these also. Dredging and filling operations are man's chief means of altering salt marshes. Now, sea level rise will add to the loss of salt marshes because much of the land on the higher side of the marshes is blocked. Thus, many of the marshes cannot migrate up the slope as the sea rises.

Eleven transplant projects are reviewed. Each gives insights and discoveries that led to improved restoration techniques. Tips on successful techniques are offered, among these were: 1. Select the proper species of plant for the elevation in the project area. 2. If possible, put transplants in the ground the same day as they are dug, and keep their roots moist at all times during the transfer. 3. Planting depth should be sufficient to allow the roots to extend normally. 4. Protect planting area from excessive wave energy by using a breakwater if necessary.

Several research needs were mentioned. Research is needed to develop ways to make transplanting less expensive. Combinations of transplanting and seeding might be developed for large areas to cut the costs. Research on "edge-effect" is needed. Does increasing the edge in a marsh increase production of fish, shrimp and crabs? Finally, research is needed to develop low-cost reusable wavebreaks.

In the ten years since this was written much research has been done, but many of the needs remain. The techniques given in the chapter are still used. Mitigation for loss of natural wetlands is not keeping up, and salt marsh is still being lost at a rapid rate in the western Gulf.

Webb, J. W., Jr. "Soil water salinity variations and their effects on *Spartina alterniflora*." Contributions in Marine Science 26 (1983): 1-13.

The effects of soil salinity on *Spartina alterniflora* were compared in a transplanted marsh and a natural marsh in the Galveston Bay area, Texas. soil salinity increased with elevation to a maximum at MHW, where it frequently reached 40 ‰. Highs

of 59 ‰ at the transplant site and 99 ‰ at the natural marsh site were found during the summer. The high salinity was damaging to *Spartina alterniflora*, and when combined with low soil moisture found above MHW was limiting to the cordgrass. High soil salinities were apparently caused by evaporation between tidal cycles.

Webb, J. W., Jr. and J. D. Dodd. "Wave-protected versus unprotected transplantings on a Texas bay shoreline." Journal of Soil & Water Conservation 38 (1983): 363-366.

Shoreline erosion is a dominant feature along many Texas bays. *Spartina alterniflora* has been planted along some stretches and effectively controlled further erosion. The objectives of this study were to see if a wave-stilling device would aid in establishment of transplanted salt marsh plants in an area where wave action was erosive, and to see at what elevations the various species of plants might become established.

The study site was along the north shoreline of East Bay in the Galveston Bay System, Texas. A 60-m stretch of shore bank was graded down to a 10% slope. The area was planted, but washed out within a few months. The area was re-graded to a 2% slope. A cable with tires threaded on it was suspended from poles driven into the bay bottom to act as a wave-stilling device. Plants were transplanted to the area, but within a few months the waves had washed them out again. The tires had sank partially, and waves were able to pass over them with minimum decrease in energy. A second string of tires was added, and the area replanted.

The two-tire cable was sufficient to break most of the wave action, and *Spartina alterniflora* survival was good (about 70% at the best elevation) after the first year. *Spartina spartinae*, *Spartina patens*, and *Distichlis spicata* also did well, surviving above mean high water much better than *S. alterniflora*.

After the tire line was removed there was some erosion of the *Spartina alterniflora*, but most of the marsh survived to prevent further erosion. The other species also acted to protect the shore at higher elevations, and were shown to be valuable in beach stabilization projects.

Webb, J. W., Jr. and J. D. Dodd. "*Spartina alterniflora* response to fertilizer, planting dates, and elevation in Galveston Bay, Texas." Wetlands 9 (1989): 61-72.

This study tested date, fertilizer rates, and elevation as they affected survival and growth of *Spartina alterniflora* transplants on a sandy dredged material deposition site on Bolivar Peninsula, Texas. Differences in seasonal tidal heights affected areas of best survival. Higher tides led to better survival at higher elevations (still below MHW). Better survival, tiller production, and growth was associated with the May planting versus the February planting. However, stem density was greater in the February planting area than in the May area after one full growing season despite the lower initial survival and tiller production. Fertilizer was of no apparent benefit. The use of a two-season planting was suggested where large expanses need to be covered. The lower elevations could be planted during the winter when water levels are usually lower, and the upper elevations could be planted during the spring when water levels are usually higher.

Webb, J. W., Jr., J. D. Dodd, A. T. Weichert, and B. H. Koerth. "*Spartina alterniflora* response to fertilizer rates, planting dates, and elevation in Galveston Bay, Texas." In: Estuaries. Second water quality and wetlands management conference. in New Orleans, LA, ed. N. V. Brodtmann, Jr., New Orleans, LA: , 379-399, 1985.

This study tested date, fertilizer rates, and elevation as they affected survival and growth of *Spartina alterniflora* transplants on a sandy dredged material deposition site on Bolivar Peninsula, Texas. Differences in seasonal tidal heights affected areas of best survival. Higher tides led to better survival at higher elevations--but below MHW. Better survival, tiller production, and growth was associated with the May planting versus the February planting. However, stem density was greater in the February planting area than in the May area after one full growing season despite the lower initial survival and tiller production. Fertilizer had no apparent effect.

Webb, J. W., Jr., M. C. Landin, and H. H. Allen. "Approaches and techniques for wetlands development and restoration of dredged material disposal sites." In: National wetland symposium: mitigation of impacts and losses. in New Orleans, LA, eds. J. A. Kusler, M. L. Quammen, and G. Brooks, New Orleans, LA: Assoc. State Wetland Managers, Inc., 132-134, 1986.

The possibilities of using dredged material and *Spartina alterniflora* to create salt marshes and stabilized shorelines are discussed. Techniques that have been tried and have shown promise are mentioned. Particular note is made of the erosion control mats with sprigs of *Spartina* planted through slits in the mat material. Breakwaters were found essential in areas of high wave energy.

Webb, J. W., Jr. and C. J. Newling. "Comparison of natural and man-made salt marshes in Galveston Bay complex, Texas." Wetlands 4 (1985): 75-86.

The vegetation of a man-made *Spartina alterniflora* marsh planted in 1976 on Bolivar Peninsula, Texas, was compared with three natural marshes in the Galveston Bay complex in 1978 and 1979. Samples for the analysis of above-ground biomass, live stem density, dead stem density, stem height, percent cover, and species composition were collected from 0.5 m² quadrats that were randomly placed along elevational transects. Below-ground biomass was collected in the same quadrats using about 8-10 cm diameter corers; core depths were 25 and 30 cm.

Spartina alterniflora dominated the sites below mean high water, but other plants (*Salicornia bigelovii*, *Spartina patens*, *Batis maritima*, *Sporobolus virginicus*, and *Distichlis spicata*) were more important above mean high water. The above-ground biomass of *S. alterniflora* and other species at Bolivar was within the variability shown among the three natural marshes. However, live biomass of *S. alterniflora* was significantly greater in 1978 in the man-made marsh than in the natural marshes. This difference was caused by greater stem heights in the created marsh which more than compensated for the generally lower stem densities.

At the lower elevations where *S. alterniflora* dominated, below-ground biomass was generally much lower in the man-made marsh than in the natural marshes. However, this biomass increased from 1978 to 1979. At the higher elevations, where other

species dominated, below-ground biomass at the created marsh was within the range of the natural marshes.

Overall, this study indicated that production of above-ground biomass in a 2-3 year old created salt marsh was comparable to production in nearby natural marshes. Below-ground biomass and species composition were changing in the created marsh, becoming more similar to the natural marshes of the area.

Wiggins, J. and E. Binney. "A baseline study of the distribution of *Spartina alterniflora* in Padilla Bay." Washington Dept. Ecology, Padilla Bay National Estuarine Research Reserve. Reprint Series 7, 28, 1987.

The objective of this study was to document the distribution of *Spartina alterniflora* in Padilla Bay, Washington. It is thought that *S. alterniflora* was transplanted into southern Padilla Bay sometime in the early 1960's. Stands of *Spartina* were located by walking the shoreline of all known salt marsh areas of the bay. Seven major stands were found in addition to the extensive stand on Dike Island, all still in southern Padilla Bay. The smaller stands were located south and east of Dike Island. All were mapped. There were also several small clumps (about 1 m² each) of *S. alterniflora* recorded in Telegraph Slough. The mapped area comprised over 2.4 ha. The elevation of the growing cordgrass ranged from 2.5 to 3.6 m above MLLW. *Salicornia virginica* intermixed with *S. alterniflora* at the higher elevation, and continued into the higher marsh where it was mixed with *Distichlis spicata*.

Based on comparisons of aerial photographs from 1968 and 1978, the spread of *Spartina alterniflora* in Padilla Bay appears to have been vegetative. It also appears that clumps may break off during storms, be transported onto other adjacent mudflats, and become established if the elevation is correct.

Williams, S. L. and J. B. Zedler. "Restoring sustainable coastal ecosystems on the Pacific coast: Establishing a research agenda." California Sea Grant College. Rep. T-CSGCP-026, 19, 1992.

This report lists prioritized research needs for Pacific coastal ecosystems, particularly estuarine wetlands. Priority weights were the averages of priorities determined by 15 scientists (most were from the West coast) who are familiar with coastal ecosystems and restoration potential. A prioritization was felt necessary to promote research in areas deemed critical for the perpetuation of coastal wetland ecosystems along the West coast. The West coast has probably lost 50% of its coastal estuarine wetlands already.

There were 25 research needs spread among four topics. Habitat specificity of organisms, habitat function determinants, and population dynamics were leaders in the conservation of biodiversity topic. Hydrology of the coastal wetlands was primary among physical processes research needs. Nutrient dynamics and establishing water quality criteria for vegetation were most important in water quality research. Habitat architecture, site selection criteria, and monitoring and evaluation of success were key needs under restoration research.

Wilsey, B. J., K. L. McKee, and I. A. Mendelsohn. "Effects of increased elevation and macro- and micronutrient additions on *Spartina alterniflora* transplant

success in salt-marsh dieback areas in Louisiana." Environmental Management 16 (1992): 505-511.

Extensive areas of salt marshes in coastal Louisiana are suffering dieback due to compaction, subsidence, and lack of sediment additions. The objectives of this study were to test elevation of the substrate and nutrient enhancement in an area of *Spartina alterniflora* degradation and dieback to see if the marsh could be restored by transplanting more *S. alterniflora* into the area given certain environmental modifications.

The test areas were dieback salt marshes in the lower Barataria Basin near Caminada Bay, Louisiana. Five sites, two elevations (0 and +30 cm), two macronutrient treatments (N, P, K; with and without), and two micronutrient treatments (Fe, Mn, Cu, Zn; with and without) were tested. One plant of *S. alterniflora* was used for each treatment at each site. All plants selected and dug from the nearby donor site were the same size.

After four months (i.e. on Nov. 17, 1989) the *S. alterniflora* in the elevated plots had twice the above-ground biomass and significantly more culms than the normal substrate level plots. The elevated plots were at the same height as the nearby, vigorously growing, *S. alterniflora* that lined the banks of the channels. Micronutrient addition appeared to inhibit growth, while macronutrient addition promoted growth only in the elevated plots. In these plots macronutrients increased culm density by about 61% over controls.

This paper reinforces the importance of planting at proper elevations in marsh restoration projects. Although elevations should generally match those of the nearest natural marsh, this study makes it clear that one should match elevations carefully, and should look to match the elevations of most vigorous growth in the natural marsh. The vigorous-growth area may be above the average elevation of the marsh.

Wolf, R. B., L. C. Lee, and R. R. Sharitz. "Wetland creation and restoration in the United States from 1970 to 1985: An annotated bibliography." Wetlands 6 (1986): 1-88.

This annotated bibliography covers studies concerning creation and restoration of salt and freshwater wetlands that were published between 1970 and 1985, and contains 304 citations accompanied by very brief annotations. Many of the cited studies are on the use of *Spartina alterniflora* in marsh creation or restoration activities. This bibliography will be useful in finding earlier articles about *S. alterniflora* restorations, or articles with a slightly different subject emphasis than our bibliography.

Woodhouse, W. W., Jr. "Building salt marshes along the coasts of the continental United States." Fort Belvoir, VA: U.S. Army Corps Eng., Coastal Eng. Res. Cent. Spec. Rep. 4, 96, 1979.

This report provides basic information about coastal marshes of the United States. Included are descriptions of the types of marshes, types of marsh plants, site requirements for establishing marshes, and plant propagation techniques. Also included are descriptions of planting and fertilizing techniques and their costs.

Woodhouse, W. W., Jr. and P. L. Knutson. "Atlantic coastal marshes." In: Creation and restoration of coastal plant communities, ed. R. R. Lewis, III. 45-70. Boca Raton, FL: CRC Press, Inc., 1982.

This chapter reviews various facets of salt marshes along the Atlantic coast of the U.S. There is a brief description of the loss of salt marshes and the need for their preservation. Much information is given to aid in creating or restoring salt marshes. Dominant coastal marsh plant species are discussed as to their shape, habitat, reproductive potential and timing, and as to techniques for harvesting, propagating, and culturing some of the species. The important species include: *Spartina alterniflora* (smooth cordgrass), *Spartina patens* (saltmeadow cordgrass), *Juncus roemerianus* (black needle rush), *Distichlis spicata* (saltgrass), *Spartina cynosuroides* (big cordgrass), and *Phragmites australis* (common reed). Planting procedures are described, and their costs estimated. A typical marsh creation project using *Spartina alterniflora* sprigs placed on 1-m centers requires about 10,000 sprigs and 100 man-hours to plant one hectare. This estimate does not include the planning and coordinating, obtaining of transplant material, nor the transport of material to the project site.

Recommendations are given for selecting a site, checking the soil, preparing the grade and elevation, selecting the planting time, and protecting the planted site. Protection must be established against grazing animals and trampling animals, including man and his off-road vehicles. Protection may also be necessary from erosive action by wind and wind and boat waves. The end product of a marsh creation project will be evaluated not only by the vegetation developed, but also by the wildlife and fisheries species that use the planted marsh as habitat.

Zedler, J. B. "Canopy architecture of natural and planted cordgrass marshes: selecting habitat evaluation criteria." Ecological Applications 3 (1993): 123-138.

Nesting requirements of the Light-footed Clapper Rail (*Rallus longirostris levipes*) were determined from its natural marsh habitat. Selected habitat criteria were identified for use in evaluating the value of created marshes as habitat for this bird. Height distribution and density of *Spartina foliosa* (California cordgrass) were preferred over percent cover for characterizing canopy architecture. Percent cover was found to be too subjective.

The bird appears to require a density of *Spartina foliosa* of at least 100 stems/m². Also, the required height frequency distribution of the stems has at least 90 stems over 60 cm tall, and of these at least 30 stems over 90 cm tall. Greater densities and taller plants make even better habitat. The size of the marsh is also important. The rail appears to need 0.8-1.6 ha of marsh with the specified architecture for its home range. There are additional requirements for the birds to use a marsh even if it appears to meet the vegetation requirements mentioned, but these other requirements have not yet been discovered.

Several associated recommendations were made based on work and observations made during this study. Destructive sampling of the cordgrass should be avoided; every stem is valuable. Densities and height frequency distributions to characterize a marsh should be based on 0.25 m² (or larger) quadrats. Long-term monitoring, 20 years, is recommended because establishment of planted marshes may meet the criteria for success for a year or two and fail thereafter.

Zedler, J. B., J. Covin, C. Nordby, P. Williams, and J. Boland. "Catastrophic events reveal the dynamic nature of salt-marsh vegetation in southern California." *Ecology* 9 (1986): 75-80.

Recent hydrological anomalies (flooding, dry-season stream flows, and droughts) were shown to greatly alter coastal wetland habitats in southern California in a six-year (1979-1984) study of the Tijuana Estuary. Permanent sampling sites (102) were set at 5-m intervals along 8 transects in lower intertidal marsh that was dominated by *Spartina foliosa*. During normally dry years (1979, 1981, 1982) interstitial soil salinities were between 35 and 45 ‰, but during years with wet winters and rare flooding (1980 and 1983) interstitial salinities dropped to between 15 and 35 ‰. During an extremely dry year (1984) the interstitial salinity reached 104‰ in September. *S. foliosa* reacted to these changes by increased growth during fresher years by mainly increased stem length when the freshwater inflow occurred early in the year (1980), and mainly by increased numbers of stems when freshwater inflow occurred later (1983). *S. foliosa* also responded quickly to the drought in 1984 with reduced stem density (62% from that found in 1983 or 28% from that found in the other years) and reduced stem heights (50% from 1983 and 30% from other years).

The study showed the importance of soil salinity in governing rates and types of growth of western cordgrass. Variations in tidal circulation and weather were also found to be important for marsh growth and establishment. Long-term dynamics of a salt marsh should be understood before the marsh is used as a reference site or as a site for restoration.

Zedler, J. B., M. Josselyn, and C. Onuf. "Restoration techniques, research, and monitoring: vegetation.", pp 63-72 In: Wetland Restoration and Enhancement in California. in Hayward, CA., ed. M. Josselyn, Hayward, CA.: California Sea Grant College Program, 1982.

This article summarizes the goals and techniques most commonly associated with California coastal marsh and seagrass restorations, and it suggests items and actions for a follow-up monitoring program for such projects. The goals are the same as for Atlantic coast restorations, perhaps with the decreased need for shoreline stabilization and increased concern for increasing habitat diversity for additional wildlife support. Restoration techniques included a basic sequence of actions: map the site, develop a conceptual plan, develop a site plan with engineering features including hydraulics, do some test planting, develop a comprehensive planting design and map including any needed protective devices, describe a monitoring plan and a management plan based on results from monitoring, and finally, develop a plan to share the information about the results of the restoration project with others (researchers, restorers, creators, and the public). Suggestions for future restoration research included determination of: (1) optimal habitat sizes and configurations for wildlife utilization by various species, (2) the tidal and freshwater flushing requirements of marsh vegetation, (3) the nutrient requirements of the vegetation and impacts of nutrient load in waste waters, (4) rates of marsh establishment--natural colonization versus transplants, (5) the relationship between plant density, flushing, and mosquito control.

Author Index

- Allen 1-3, 58
Athnos 4
Auble 38
Baca 4
Banner 4
Beeman 5
Benner 5
Berger 6
Bernstein 6
Binney 59
Blair 7
Boland 62
Bolton 7
Bontje 7-8
Boyd 8
Brochu 5, 28
Brooks 28
Broome 9-12, 17-18, 50- 51
Buffington 28
Callaway 12
Cammen 13-15
Chabreck 15
Chambers 54
Clairain 1
Cobb 16
Colby 38
Courtney 16
Covin 17, 62
Craft 17-18
Crewz 18
Cross 28
Diaz 1
Dodd 53, 57-58
Earhart 19
Eleuterius 20
Espey Huston & Associates,
Inc. 20-21
Faber 21-22
Farmer 32
Florida Department of
Environmental
Regulation 22
Fonseca 38
Ford 1, 29
Fowler 22, 25
Frenkel 23
Frye 22, 25
Gallagher 24
Garbisch 19, 24
Gill 20
Good 54
Gosselink 55
Grant 24
Griswold 27
Gwin 28
Hall 16
Hardaway 22, 25
Hartman 26
Hill 22, 25
Hine 51
Hoffman 26
Holland 28

- Hossner 36-37
 Hurme 5
 Ibison 22, 25
 Inskeep 28-29
 Jones 26
 Josselyn 12, 27, 62
 Junt 1
 Kana 4
 Kentula 28, 31
 Kenworthy 38
 Kiraly 28
 Klemas 45
 Klima 40
 Knutson 5, 28-29, 61
 Koellen 26
 Koerth 58
 Kraus 29
 Kruczynski 30
 Kunze 23
 Kusler 31
 Landin 32-33, 44, 58
 Langis 32
 LaPerriere 32
 LaSalle 33
 Lee 60
 Levin 41
 Lewis 18, 34-35
 Lindall 35
 Lindau 36-37
 Lyon 37
 Mager 37
 McCallum 24
 McKee 59
 Medina 40
 Mendelssohn 59
 Meyer 38
 Miller 38
 Minello 39-40
 Morrison 41
 Moy 41
 Munro 42
 Nailon 43
 National Research Council
 43
- Newling 44, 58
 Niering 44, 51
 Nordby 62
 Onuf 62
 Oyler 29
 Pacific Estuarine Research
 Laboratory 45
 Parris 44
 Patience 45
 Peck 16
 Reppert 45
 Reubsamen 26, 47
 Riggs 47
 Roberts 47-48
 Rodgers 26
 Rogers 9
 Sacco 49
 Schneller-McDonald 38
 Seelig 28
 Seidensticker 43
 Seliskar 24
 Seneca 9-12, 17-18, 50-51
 Sharitz 60
 Shepherd 21
 Sherman 28
 Shirley 2
 Shisler 51
 Sifneos 28
 Sims 33
 Sinicrope 51
 Soil Conservation Service 52
 Somers 24
 Steinke 52
 Steyer 54
 Stout 52
 Tanner 53
 Thayer 35, 37-38, 54
 Thomas 22, 25
 Underwood 54
 US Army Corps of Engineers
 54
 Warren 51
 Watzin 55
 Webb 2-3, 32, 39, 55-58

- Weichert 58
Wells 1
Wiggins 59
Williams 21, 41, 59, 62
Wilsey 59
Wolf 60
Woller 24
Woodhouse 9-12, 50-51,
60-61
Zalejko 32
Zedler 17, 27, 32, 59, 61-62
Zepp 6
Zimmerman 39-40

Subject Index

- Acidity 24
 Alabama 3, 53
 Alkali bulrush 41
Ammophila 25, 30
 Amphipods 14, 39
Anchoa 40
 Animal use 21, 30, 33, 38, 40, 48, 56
 Marsh rabbits 2
 Rats 2
 Annotated bibliography 38, 60
Arcuatula 14
 Atlantic coast 61
Avicennia germinans 4, 27
Batis 56, 58
 Benthos 2, 7, 15, 30, 33, 39
 macro-infauna 49
 Biomass 12-15, 21, 32, 33, 48, 51, 58
 below-ground 11, 33, 39
 above-ground 11, 32, 33, 39
 Birds 2, 7, 19, 21, 44, 47, 52
 Black mangrove 4, 27
 Block nets 33
 Blue crabs 39
Borrchia 56
 Breakwaters 1, 3-5, 26, 55, 57, 58
 Cargo parachutes 43
 Christmas trees 43
 Dike 3
 fixed tire fence 1-3
 floating tire 1
 floating tire fence 2, 3
 Plastic snow fence 43
 sand bag 1
 Terraces 54
 Breder traps 33, 48
Brevoortia 7
 Burlap plant rolls 1
 California 6, 8, 12, 17, 21, 22, 27, 32, 41, 45, 61, 62
Callinectes 7, 33, 39, 41
Capitella 13
 Created marsh requirements 50
 Clams 7
 Clipping of the cordgrass 27
 Clumps 19
 Coastal wetlands 55
 Competition 13, 20, 22
 Connecticut 52
 Cores 15, 25, 30, 33, 39, 41, 48, 49
 Cover 21
 Crabs 41
 Crustaceans 40
 Culturing 61
Cyathura 14
Cymodocea 20
 Definitions 34, 55
 Delaware 24
 Design 7, 11, 19, 27, 41, 42, 51, 52, 54, 55
 Dieback salt marshes 60
Distichlis 20, 25, 30, 44, 51, 56-59, 61
 Distribution and spread 47, 59
 Dolichopodidae 13
 Donor marsh 21, 27, 35
 Donor marsh recovery 16

- Dredged material 1, 3, 13, 19, 25,
 32, 33, 36, 37, 40, 50, 57, 58
 Chemical properties 36
 Physical properties 36
 Drop sampler 40
 Elevation 1, 5, 8-10, 13, 18-20,
 22-25, 37, 51, 57, 58, 60
 Enforcement 27, 32
 Epiphytic algae 37
 Erosion control 1, 26, 57
 Evaluation procedures 45
 Faunal diversity 14
 Fence, pest 1
 Fertilizer 1, 9, 10, 12, 17, 23, 25,
 26, 37, 55, 57, 58
 Organic nitrogen 17
 Osmocote 10, 26
 Fertilizers 30
 Fetch 22, 26, 43
 Fiber mat 2
 Fiddler crabs 7, 30, 41
 Fish 7, 33, 39, 40, 42, 47, 48
 Fish habitat 34
 Fisheries habitat conservation 38
 Fisheries organisms 38
 Fishery habitat 40
 Fishery species 43
 Florida 4, 5, 16, 19, 22, 26, 34,
 35, 44, 48
 Florida Department of Environmental
 Regulation (FDER) 22
 Functional equivalency 40, 45, 47, 55
 Functional health of wetlands 45
Fundulus 7, 33, 41
 Fyke nets 48
Gammarus 14
 Genetics 24
 Glossary of terms 34
Gobionellus 39
 Grading, bulldozer 1
 Growth rates 10, 12, 50-51, 54, 57,
 58, 60, 62
 Guide 55
 Habitat diversity 62
 Habitat functioning 40, 45, 47, 55
Halodule 5, 20, 38
Halophila 5
 Heavy metals 37
Heteromastus 13
 Hydrological anomalies 62
 Hydrology 41, 45
 Infauna 39, 41, 49
 Infaunal prey 40
 Invertebrates 7
 Iron sulfides 24
Iva 56
Juncus 6, 20, 30, 48, 51, 61
 Labor costs 11
Laeonereis 14, 25
Lagodon 39
Laguncularia racemosa 4, 27
Lepidactylus 14
 Light-footed Clapper Rail 61
Limonium 56
Littorina 21
 Louisiana 26, 54, 60
Macoma 25
 Macro-infauna 13-15
 Production 14
 Macrobenthos 25, 33
 Mag Amp, tests 10
 Mammals 7, 47-48
Manayunkia 42
 Mangrove propagules 5
 Marsh comparison
 created vs. natural 7, 10-14,
 17, 18, 21, 23, 32,
 37, 39, 41, 48-49, 50,
 56, 58
 Marsh creation 2, 4-11, 15, 21, 27,
 30, 32, 35, 43, 49, 50-
 51, 55, 57
 definitions 34, 55
 evaluation 26, 47, 53
 requirements 50, 56
 survey 18
 Marsh descriptions 60

- Marsh development 21, 32, 33, 35,
 41, 44, 51
 Marsh distribution 23
 Marsh edge 40
 Marsh evaluation 4, 18, 20, 30
 Marsh inventory 29
 Marsh restoration 34
 Marsh succession 52
 Maryland 19, 25
 Mats, erosion control 58
 Mechanical planting 55
 Meiofauna 42
Menidia 7, 40
 Mississippi 20
 Mitigation 16, 22, 28, 33, 36, 37,
 42, 46, 47, 49, 53
 evaluation 6, 22
 process 6
 Mitigation banking 45
 Mitigation policy 26, 42
 Model, wave dampening 28
Modiolus 14
 MOM 17, 39
Monanthochloe 56
 Monitoring 7, 8, 11, 15, 19, 27, 41,
 43, 52, 53, 55, 62
 Indicator species 9
Monopylephorus 42
Mugil 7
 Multiple species 6, 20, 55, 57
 National restoration strategy 43
 Nekton 40
 Nematodes 7
Nereis 14, 15
 Nesting requirements 61
 New Jersey 8, 30
 Hackensack River 7
 New York 32
 Nitrogen 32
 NMFS 26, 36, 37, 40, 47
 North Carolina 5, 10, 11, 13-15,
 17, 18, 37, 38, 41,
 49-51
 Northeastern U.S. 51
 Nutrient pools 17
 Nutrient requirements 12
 Nutrients 36, 60
 Oligochaetes 2, 15, 25, 49
 Oregon 23
 Organics 13, 17, 33, 42, 48
 sediments 49
 Osmocote 23, 26
 Oxidation 24
 Oyster cultch 38
Palaemonetes 33, 39, 40
Panicum 20, 30
 Peat pots 25
Penaeus 39, 40
 Permitting 16, 22, 26, 36, 37, 42,
 53
 Phenology 12, 24, 50, 53
 Philosophies 54
Phragmites 6-8, 20, 30, 52, 61
 Pit traps 42
 Planning 19, 27, 41, 52, 61
 Plant community 55
 Plant cover 18
 Plant growth 10, 11, 17, 23
 Plant roll 2
 Planting 1
 Planting dates 58
 Planting procedures 61
 Planting techniques 1-3, 5, 19, 30,
 35, 55
 Burlap roll 3
 Multi-stem clumps 3
 Multi-stem plugs 5
 Single stems 3
 Policy 53
 Polychaetes 2, 13, 15, 39, 49
 Pore water 18
 Potential spread 23
 Prey items 33
 Primary production 37
 Production, animal 14
 Productivity 55
 Propagating 61
 Quantitative drop enclosures 39

- Rallus* 61
 Reasons for created marsh failure
 16-18
 Recommendations 16, 18, 23, 27,
 31, 34, 43, 50, 53, 54,
 56, 61, 62
 Red mangrove 4
 Regional goals 27
 Regulations 4
 Remote sensing 45
 Research needs 15, 28, 34, 45, 51,
 54-56, 59, 62
 Restoration activities 6
 goals 41
 guide 38
 manual 45
 needs 55
 Restoration projects 32, 41
 Restoration techniques 50, 62
 Restored tidal exchange 4, 21, 22,
 41, 52
Rhizophora mangle 4
 Rodeo 8
 Root burn 10
Ruppia 20
Salicornia 17, 21, 22, 41, 52, 56,
 58, 59
 Sampling 61
 Sampling design, randomized block
 40
 Sampling sites 62
 Sampling strategy 45
Scirpus 44, 56
 Seagrasses 20
 Season 11, 58
 Sediment 33, 41
 Seed production 35
 Seeding 2, 10, 19
 Seedling 53
 Self-regulated tide gates 52
 Shoreline erosion 9
 Shoreline protection 55
 Shoreline stabilization 43
 Shrimp 39-40
 Slope 4, 5, 11, 25, 57
 Soil attributes 24
 Soil salinity 56, 62
 Soils 18
 South Carolina 4, 33
Spartina anglica 23
Spartina cynosuroides 10, 20, 25,
 30, 61
Spartina foliosa 12, 17, 21, 22, 32,
 61, 62
Spartina maritima 23
Spartina patens 2, 10, 19, 20, 23,
 25, 26, 30, 36, 44, 51, 52,
 55-58, 61
Spartina spartinae 55, 57
Spartina townsendi 23
Sporobolus 58
 Sprigging 2, 5, 58
 Stabilization 19
 Biotechnical 1
 Dredged material 1
 shoreline 1
Streblospio 15, 42
 Survey 53
 Techniques 54-56, 60, 61
 Terracing 54
 Texas 1, 3, 16, 21, 36, 37, 39, 40,
 43, 53, 55-58
Thalassia 20
 Time of inundation 25
 Topography 41, 45
 Transplant 50, 54
 Growth 25
 history 59
 Survival 25
 Transplant projects, evaluation 20
 Transplant spacing 11
 Transplanting 3, 4, 10, 11, 57
Tubulanus 25
Typha 6, 52
Uca 30
Uca longisignalis 41
Uniola 30

Vegetation

growth patterns 59

Vegetative Erosion Control

Project 22, 25

succession 45

Vegetative Stabilization Site

Evaluation Form 29

Virginia 7, 22, 25, 26, 28

Washington 23, 47, 59

Wave-climate severity 29

Waves' erosional forces 28

Wegener Ring net 48

West coast 59

Wetland Creation/Restoration Data

Base 38

Wetland functions 43

Wetland laws 4

decision making 28

design 28

planning 28

Wetland policy 4

Wetland restoration 31, 52, 53

fishery management tool 34

guide 31

Wetland types 27

White mangrove 4

Workshop

restoration 27

wetlands 28

Zonation

20, 22, 23, 30, 58

Zostera 38

**Other Titles in the
Decision Analysis Series**

No. 1. Able, Kenneth W. and Susan C. Kaiser. 1994 Synthesis of Summer Flounder Habitat Parameters.

Science for Solutions

NOAA COASTAL OCEAN PROGRAM
Decision Analysis Series No. 2



**TECHNOLOGY AND SUCCESS IN RESTORATION,
CREATION, AND ENHANCEMENT OF *SPARTINA
ALTERNIFLORA* MARSHES IN THE UNITED STATES**
Volume 2 -- Inventory and Human Resources Directory

Geoffrey A. Matthews
Thomas J. Minello

August 1994



U.S. DEPARTMENT OF COMMERCE
National Oceanic and Atmospheric Administration
Coastal Ocean Office

The Decision Analysis Series has been established by NOAA's Coastal Ocean Program (COP) to present documents for coastal resource decision makers which contain analytical treatments of major issues or topics. The issues, topics, and principal investigators have been selected through an extensive peer review process. To learn more about the COP or the Decision Analysis Series, please write:

NOAA
Coastal Ocean Office
1315 East West Highway, Sta. 15140
Silver Spring, MD 20910

phone: 301-713-3338
fax: 301-713-4044

Science for Solutions

NOAA COASTAL OCEAN PROGRAM

Decision Analysis Series No. 2



**TECHNOLOGY AND SUCCESS IN RESTORATION,
CREATION, AND ENHANCEMENT OF *SPARTINA*
ALTERNIFLORA MARSHES IN THE UNITED STATES
Volume 2 -- Inventory and Human Resources Directory**

Geoffrey A. Matthews

Thomas J. Minello

NOAA National Marine Fisheries Service

Southeast Fisheries Science Center

Galveston Laboratory

4700 Avenue U

Galveston, TX 77551

August 1994

U.S. DEPARTMENT OF COMMERCE

Ronald H. Brown, Secretary

National Oceanic and Atmospheric Administration

D. James Baker, Under Secretary

Coastal Ocean Office

Donald Scavia, Director

This publication should be cited as:

Matthews, Geoffrey A. and Thomas J. Minello. 1994. Technology and Success in Restoration, Creation, and Enhancement of *Spartina alterniflora* Marshes in the United States. Vol. 2 -- Inventory and Human Resources Directory. NOAA Coastal Ocean Program Decision Analysis Series No. 2. NOAA Coastal Ocean Office, Silver Spring, MD.

This publication does not constitute an endorsement of any commercial product or intend to be an opinion beyond scientific or other results obtained by the National Oceanic and Atmospheric Administration (NOAA). No reference shall be made to NOAA, or this publication furnished by NOAA, in any advertising or sales promotion which would indicate or imply that NOAA recommends or endorses any proprietary product mentioned herein, or which has as its purpose an interest to cause directly or indirectly the advertised product to be used or purchased because of this publication.


Note to Readers

The NOAA Coastal Ocean Program (COP) provides a focal point through which the agency, together with other organizations with responsibilities for the coastal environment and its resources, can make significant strides toward finding solutions to critical problems. By working together toward these solutions, we can ensure the sustainability of these coastal resources and allow for compatible economic development that will enhance the well-being of the Nation now and in future generations. The goals of the program parallel those of the NOAA Strategic Plan for 1995-2005.

A specific objective of COP is to provide the highest quality scientific information to coastal managers in time for critical decision making and in a format useful for these decisions. To help achieve this, COP inaugurated a program of developing documents that would synthesize information on issues that were of high priority to coastal managers. To develop such documents, a three-step process was used: 1) to compile a list of critical topics in the coastal ocean through a survey of coastal resource managers and to prioritize and select those suitable for the document series through the use of a panel of multidisciplinary technical experts; 2) to solicit proposals to do research on these topics and select principal investigators through a rigorous peer-review process; and 3) to develop peer-reviewed documents based on the winning proposals.

Seven topics and associated principal investigators were selected in the initial round. *Technology and Success in Restoration, Creation, and Enhancement of Spartina alterniflora Marshes in the United States* by Geoffrey A. Matthews and Thomas J. Minello of the NOAA National Marine Fisheries Service's Galveston Laboratory is the second document in this Decision Analysis Series to be published and is presented in two volumes. Information on Decision Analysis Series No. 1 is shown on the inside back cover. Other volumes will be published over the next two years on the following topics: seagrass restoration technology, coastal watershed restoration, restoring streams and anadromous fish habitat affected by logging, eutrophication and phytoplankton blooms, and management of cumulative coastal environmental impacts.

As with all of its products, COP is very interested in ascertaining the utility of the Decision Analysis Series particularly in regard to its application to the management decision process. Therefore, we encourage you to write, fax, call, or Internet us with your comments. Please be assured that we will appreciate these comments, either positive or negative, and that they will help us direct our future efforts. Our address and telephone and fax numbers are on the inside front cover. My Internet address is DSCAVIA@HQ.NOAA.GOV.



Donald Scavia
Director
NOAA Coastal Ocean Program

Foreword

Coastal wetlands are important habitats for many estuarine-dependent fishery species, and the productivity of these fisheries may well depend upon the health of wetland ecosystems. The low elevation wetlands on the Atlantic and Gulf of Mexico coasts of the United States are generally dominated by smooth cordgrass, *Spartina alterniflora*, and these salt marshes have been identified as valuable nursery habitats for many fishes and crustaceans. Coastal development, sea level rise, and land subsidence, however, have resulted in extensive losses of *S. alterniflora* habitat in the United States. Efforts to reduce the rate of wetland loss will likely require a continuation and expansion of programs to restore and create salt marshes in regions of deteriorating coastal wetlands. These restoration projects frequently involve establishment of marsh vegetation on graded-down uplands or on dredged material. Although many of the early efforts have failed, the technology for successfully establishing salt marsh vegetation by seeding and transplanting have now been developed for a variety of coastal conditions.

Salt marshes have long been built to reduce shoreline erosion and stabilize dredged material deposits. These projects have not inherently attempted to establish complete functional equivalence with natural marshes, and physically establishing the macrophytic vegetation has been the criteria used for determining success. When the goal of salt marsh creation becomes the replacement of a natural marsh, however, the entire suite of marsh functions must be considered. Replacing or creating vegetative cover is the first step in creating a functional salt marsh. Establishment of habitat functions such as providing a supportive food web, protection from predators, and reproductive sites for estuarine animals apparently requires more time than the establishment of macrophytes alone. Under some conditions, these habitat functions may never fully develop, and food webs of created salt marshes may never support fisheries organisms and other wildlife to the same extent as in natural marshes. The evidence for retarded functional development of created salt marshes has raised serious questions about the relative value of these habitats. Information on the establishment of salt marsh functions other than substrate stabilization is also more difficult to obtain and requires an understanding of how natural salt marsh systems function.

Our objectives in developing this database of created *Spartina alterniflora* marshes were: 1) to form a register of the marshes and estimate the number and

location of these marshes, 2) to determine how many of these marshes have been monitored for functional success and animal utilization, 3) to develop an inexpensive and easy-to-use data base and entry system to track future restoration/creation activities, 4) to promote via this database a greater awareness of the impacts of altering coastal wetlands and the need for full ecologically functional replacement.

Companion products from our Coastal Ocean Program project may also be of some value to users of this Inventory. These include a Directory of Human Resources in this volume and an Annotated Bibliography which appears in Volume 1. The information in these companion products was used extensively in creating this salt marsh inventory.

VOLUME 2

**SECTION II -- INVENTORY OF
RESTORED, CREATED, AND ENHANCED
SPARTINA ALTERNIFLORA MARSHES IN
THE UNITED STATES**

**SECTION III -- DIRECTORY OF HUMAN
RESOURCES**

SECTION II

AN INVENTORY OF RESTORED, CREATED, AND ENHANCED *SPARTINA ALTERNIFLORA* MARSHES IN THE UNITED STATES

Table of Contents

Section II -- Inventory

FOREWORD	iii
DATABASE	1
Data collection and limitations	1
Database structure	2
PRIMARY MARSHES	5
Connecticut	5
Delaware	10
Florida	17
Maine	104
Maryland	107
Massachusetts	127
New Jersey	128
New York	145
North Carolina	147
South Carolina	148
Texas	149
Virginia	229
MARSHES WITH MINIMAL INFORMATION	237
Alabama	237
Connecticut	238
Delaware	238
Florida	239
Georgia	247
Louisiana	248
Maine	257
Maryland	257
Massachusetts	283
Mississippi	283
New Hampshire	283
New Jersey	284
New York	285
North Carolina	286
Rhode Island	288
South Carolina	288

Texas	291
Virginia	297

Database

Data collection and limitations

In the development of this inventory, we utilized four major sources of information on restored and created *Spartina alterniflora* marshes, including 1) contractors who build the marshes, 2) research personnel who study marshes, 3) state agencies that monitor wetlands and marsh mitigation, and 4) federal agencies that monitor marshes or marsh organisms. We contacted contractors throughout the United States for information about restored, enhanced, or created *Spartina alterniflora* marshes that they had a role in building. We also reviewed information on created marshes from publications, reports, conference presentations, and even assorted notes obtained from consultants, contractors, and other research personnel working in this field. State environmental and fisheries agencies were consulted for their records about man-made salt marshes. We also contacted the offices of the U.S. Army Corps of Engineers (COE), Soil Conservation Service (SCS), U.S. Fish and Wildlife Service, and National Marine Fisheries Service (NMFS) for information about man-made *Spartina alterniflora* marshes in their respective areas of the coast.

For the most part, the people we contacted in the field of restoration science were receptive toward our requests and provided the available information. Often, however, data on salt marsh construction projects were never originally recorded, were incomplete or in some inappropriate format, were not readily available with a reasonable search effort, or were not entered into electronic files. The permitting process of the COE and the permit review records of the National Marine Fisheries Service Southeast Regional offices provided information on numerous marsh creation projects. Some projects that were issued COE permits, however, were not located, and it is unclear whether these marshes were failures or had never been constructed. In addition, we obtained some information from design or initial work plans for projects, and these plans may not always have been identical to the characteristics of the marsh actually planted.

Information on created *Spartina alterniflora* marshes on the West Coast was not recorded. A few *Spartina alterniflora* marshes were planted in West Coast estuaries several decades ago, but this species is no longer used in restoration/creation projects. *Spartina alterniflora* is considered an invasive exotic species on the West

coast, where it has spread rapidly, often out competing the native species, *Spartina foliosa*.

This inventory is intended to be dynamic. Over time, we anticipate that users of the inventory will identify new marshes and provide additional information on included marshes. We would like to encourage this type of input, and we hope that users will help make the inventory a more valuable resource for everyone working in the field of salt marsh creation and restoration. Therefore, corrections and additions are solicited and can be sent to the authors at the address on the cover page.

Database structure

During the course of our one year investigation we identified 787 marshes that were either planned, created, restored, or enhanced along coastal bays and estuaries from Maine to Texas. Each marsh was assigned a unique identification number when the data were entered into the electronic data file. The records are sorted alphabetically first by state, second by county, and third by bay in which the marshes were located. The database was constructed using Microsoft's Excel software and is also available on diskette as an Excel file. Excel was chosen because it was available for both IBM (and compatibles) and Macintosh computers and is fairly easy to use. All 787 marshes are included in the electronic data file.

Printed output was generated from the Excel database file and includes all marshes, but marsh information was presented in two forms to save space. Primary Marshes were identified on the basis of the information content in the database; the exact location of these marshes is known (INFO RANK of 3, 4, 5 and RTE, see below). All information available in the database is listed for each Primary Marsh. For marshes with minimal information (INFO RANK of 1 and 2), only a brief description is given.

The database has 22 variables to identify and characterize each marsh. These variables are:

<u>Variables</u>	<u>Specifications</u>
1) ID	A unique reference number to identify the marsh; given when the data were entered into the inventory database.
2) INFO RANK	Each planted marsh was ranked from 1-5 based on the amount of available information. 1=A COE permit was granted that required a marsh be created. It is unknown if the marsh was created. 2=A marsh was planted. The age and general location are known, but the exact location of the planting is unknown. 3=A marsh was planted and its specific location is known, but no assessments of its growth and functions were made. 4=A marsh was planted and was evaluated as to attainment of a 70% vegetative cover within two years after planting. 5=A marsh was planted, was successfully established, and was assessed for its utilization by animals. An RTE (restored tidal exchange) in this space indicates an area was opened to tidal exchange and a <i>Spartina alterniflora</i> marsh became established, though it was not planted.
3) STATE	State where the marsh is located.
4) COUNTY	County or parish where the marsh is located.
5) BAY	The body of water where the marsh is located; include bay, sub-bay, cove, river and creek when applicable.
6) ORIGINATOR	Who paid to have the marsh built. Includes name, address and telephone if available.
7) COE NO	U.S. Army Corps of Engineers permit number for a mitigation marsh.
8) PERMIT DATE	Month and year of the permit.
9) LOCATION	Directions to the actual site with identifying features of the marsh.

- 10) **LAT** Degree, minutes, and seconds where the marsh is located.
- 11) **LONG** Degree, minutes, and seconds where the marsh is located.
- 12) **DATE PLANTED** The month and year when the marsh planting was completed.
- 13) **ACRES** Area of the marsh.
- 14) **HECTARES** Area of the marsh.
- 15) **SOURCE PLANTING** Who supplied the information about the planting; includes name, address and telephone number when available, or a literature citation (intext style).
- 16) **SOURCE ASSESS** Name, address and telephone number of the person when available; or a literature citation (intext style).
- 17) **REFS** Literature citation(s) or person who knows about the marsh (address and telephone number if known).
- 18) **OTHER SPP PLANTED** List species that were planted along with the *Spartina alterniflora*..
- 19) **COVER** Visual estimation of the area of the marsh covered by *S. alterniflora*, and the month and year when the percentage was determined.
- 20) **ANIMAL USE** Yes or no. If yes, include reference. and which animals or animal groups were studied.
- 21) **COM1** Include any pertinent information about how this marsh was constructed, or how successful it was.
- 22) **COM2** Additional information that would not fit in the first comments box, or that you want separated. Marsh name generally included here.

Primary Marshes

Marsh Inventory No.: 661 **Information Rank (1-5):** RTE **State:** CT

Date Planted: 1979

Area Planted (1acre = 0.4047 ha): 52.611 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit:

Location of the marsh.

State: CT **County:** Fairfield **Latitude (N):** 41 07' **Longitude (W):** 73 15'

Bay and sub-bays: Long Island Sound; Pine Creek

Directions to the site: Exit I-95 and exit 20, take Post Rd. Go south on S. Pine Creek Rd. to Oldfield or Old Dam Rd, go east.

Source of planting information: Mr. Thomas J. Steinke, Conservation Commission - Town of Fairfield, Independence Hall, 725 Old Post Road, Fairfield, CT 06430, (203) 256-3071

Source of subsequent assessment information: John Trautman, Biol. Dept. Fairfield Univ. (203) 268-5103

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- No *Spartina alterniflora* planted, but the area was opened to more tidal exchange and *S. alterniflora* became established.

Marsh Inventory No.: 660 **Information Rank (1-5):** RTE **State:** CT

Date Planted: 1974

Area Planted (1acre = 0.4047 ha): 5.666 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit:

Location of the marsh.

State: CT **County:** **Latitude (N):** **Longitude (W):**

Bay and sub-bays: Ash Creek

Directions to the site:

Source of planting information: Mr. Thomas J. Steinke, Conservation Commission
- Town of Fairfield, Independence Hall, 725 Old Post Road, Fairfield, CT 06430, (203)
256-3071

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* : *Phragmites*

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- No *Spartina alterniflora* planted, but the area was opened to more tidal exchange and *S. alterniflora* became established.

Marsh Inventory No.: 663 **Information Rank (1-5):** RTE **State:** CT

Date Planted: 1990

Area Planted (1acre = 0.4047 ha): 60.705 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit:

Location of the marsh.

State: CT **County:** **Latitude (N):** **Longitude (W):**

Bay and sub-bays: Great Creek

Directions to the site: Milford, CT; in the state park.

Source of planting information: Mr. Thomas J. Steinke, Conservation Commission
- Town of Fairfield, Independence Hall, 725 Old Post Road, Fairfield, CT 06430, (203)
256-3071

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- No *Spartina alterniflora* planted, but the area was opened to more tidal exchange and *S. alterniflora* became established.

8

Marsh Inventory No.: 662 **Information Rank (1-5):** RTE **State:** CT

Date Planted: 1982

Area Planted (1acre = 0.4047 ha): 12.141 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit:

Location of the marsh.

State: CT **County:** **Latitude (N):** **Longitude (W):**

Bay and sub-bays: Sasco Creek

Directions to the site:

Source of planting information: Mr. Thomas J. Steinke, Conservation Commission
- Town of Fairfield, Independence Hall, 725 Old Post Road, Fairfield, CT 06430, (203)
256-3071

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

Marsh Inventory No.: 659 **Information Rank (1-5):** RTE **State:** CT

Date Planted: 1973

Area Planted (1acre = 0.4047 ha): 6.475 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit:

Location of the marsh.

State: CT **County:** **Latitude (N):** **Longitude (W):**

Bay and sub-bays: Turney Creek

Directions to the site:

Source of planting information: Mr. Thomas J. Steinke, Conservation Commission
- Town of Fairfield, Independence Hall, 725 Old Post Road, Fairfield, CT 06430, (203)
256-3071

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* : *Phragmites*

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- No *Spartina alterniflora* planted.

10

Marsh Inventory No.: 346 **Information Rank (1-5):** 3 **State:** DE

Date Planted: 05/01/92

Area Planted (1acre = 0.4047 ha): 0.202 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: US EPA Region III

Location of the marsh.

State: DE **County:** Kent **Latitude (N):** 39 07' 15" **Longitude (W):** 75 29' 30"

Bay and sub-bays: Delaware Bay; St. Jones River

Directions to the site: Take Rt. 13 north. Turn right onto Delaware Rt.10 northbound. Turn right onto Old Lebanon Road, after 500 ft. take the dirt road that bears left to the entrance of the landfill. The site is along the Wildcat Landfill on the St. Jones River.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- *Spartina alterniflora* was planted on 2-ft centers. Marsh shape: tidal pond, shoreline. Built by Environmental Concern, Inc. Interagency Agreement # DW96934592-0. This was an EPA hazardous waste site clean-up project.

- Marsh name: Wildcat Landfill.

Marsh Inventory No.: 313 **Information Rank (1-5):** 3 **State:** DE

Date Planted: 07/01/85

Area Planted (1acre = 0.4047 ha): 1.214 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: DE DOT, Office of Administration

Location of the marsh.

State: DE **County:** New Castle **Latitude (N):** 39 05' 50" **Longitude (W):** 75 27' 23"

Bay and sub-bays: Delaware Bay; St. Jones River

Directions to the site: Along each side of US Hwy 113 for about 1 km N of the bridge at Barkers Landing.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- *Spartina alterniflora* was planted on 2-ft centers. Marsh shape: rectangular, long. Built by Environmental Concern, Inc. FHWA Project # F-10005(11). DEDOT Contract # 82-109-04.

- This was a restoration after removal of a temporary access road along the Rt. 113 bridge at the St. Joans River. Marsh name: DE Route 113.

Marsh Inventory No.: 333 **Information Rank (1-5):** 3 **State:** DE

Date Planted: 05/01/83

Area Planted (1acre = 0.4047 ha): 0.243 hectares

COE Permit No.: NAPOP-R-83-0093-C **Date of COE Permit:** 1983

Originator or Applicant for permit: William K. McMahon, Rd. 2, Box 237, Lewes, DE 19958

Location of the marsh.

State: DE **County:** Sussex **Latitude (N):** 38 41' 35" **Longitude (W):** 75 04' 15"

Bay and sub-bays: Atlantic Ocean; Rehobeth Bay; Head of Bay Cove.

Directions to the site: Bay shore in front of wooden bulkhead on property due W of Bayard Ave. at New Orleans St., Dewey Beach, DE.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* : *Spartina patens*

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- *Spartina alterniflora* was planted on 2-ft centers. Marsh shape: irregular. Built by Environmental Concern, Inc.

- Marsh name: Dewey Beach.

Marsh Inventory No.: 314 **Information Rank (1-5):** 3 **State:** DE

Date Planted: 04/01/87

Area Planted (1acre = 0.4047 ha): 0.526 hectares

COE Permit No.: NAPOP-R-DV84-081 **Date of COE Permit:** 1984

Originator or Applicant for permit: Jerry Adkins real estate, Selbyville, DE 19975
(302) 436-5981.

Location of the marsh.

State: DE **County:** Sussex **Latitude (N):** 38 27' 38" **Longitude (W):** 75 05' 00"

Bay and sub-bays: Assawoman Bay; Fenwick Sound, Roy Creek

Directions to the site: Fenwick Island off East Stoney Run.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc.,
P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- *Spartina alterniflora* was planted on 2-ft centers. Marsh shape: rectangular.
- Built by Environmental Concern, Inc. This was a restoration required in a consent order.
- Marsh name: Keenwick Sound.

Marsh Inventory No.: 328 **Information Rank (1-5):** 3 **State:** DE

Date Planted: 06/01/89

Area Planted (1acre = 0.4047 ha): 1.052 hectares

COE Permit No.: NAPOR-85-1107-1 **Date of COE Permit:** 1985

Originator or Applicant for permit: Gray Farms Partnership

Location of the marsh.

State: DE **County:** Sussex **Latitude (N):** 38 28' 22" **Longitude (W):** 75 06' 30"

Bay and sub-bays: Assawoman Bay; Fenwick Sound, Roy Creek

Directions to the site: Adjacent to a tributary of Roy Creek on a tract of land formerly known as Gray Farm, along the north side of Rt. 54 immediately west of Treasure Beach Campground, near Bayville, Sussex County, DE.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- *Spartina alterniflora* was planted on 2-ft centers. Marsh shape: tidal pond, shoreline. Built by Environmental Concern, Inc. DE Wetlands Permit # 5P-0710/84 and WE-1609/85.

- Marsh name: Mallard Lakes.

Marsh Inventory No.: 338 **Information Rank (1-5):** 3 **State:** DE

Date Planted: 05/01/89

Area Planted (1acre = 0.4047 ha): 0.121 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Bay Colony Marina

Location of the marsh.

State: DE **County:** Sussex **Latitude (N):** 38 34' 58" **Longitude (W):** 75 09' 25"

Bay and sub-bays: Atlantic Ocean; Indian River Bay

Directions to the site: Along the sides of a small inlet and basin just 0.5 km SW of Grays Point on the S shore of Indian River Bay about 2 km N of Blackwater Village.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- *Spartina alterniflora* was planted on 2-ft centers. Marsh shape: shoreline. Built by Environmental Concern, Inc.
- Marsh name: Bay Colony.

Marsh Inventory No.: 298 **Information Rank (1-5):** 3 **State:** DE

Date Planted: 06/01/81

Area Planted (1acre = 0.4047 ha): 2.792 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: US Army COE, Philadelphia District

Location of the marsh.

State: DE **County:** Sussex **Latitude (N):** 38 37' 35" **Longitude (W):** 75 05' 30"

Bay and sub-bays: Atlantic Ocean; Indian River Bay.

Directions to the site: On bay-side of an island that is 1 km SE of Bluff Point, where Rehoboth Bay narrows to connect with Indian River Bay.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- *Spartina alterniflora* was planted on 2-ft centers. Marsh shape: irregular. Built by Environmental Concern, Inc. COE Contract # DACW61-81-C-0062.

- Marsh name: Indian River.

Marsh Inventory No.: 748 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 09/01/93

Area Planted (1acre = 0.4047 ha): 1.670 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit:

Location of the marsh.

State: FL **County:** Brevard **Latitude (N):** 28 11' **Longitude (W):** 80 37'

Bay and sub-bays: Banana River

Directions to the site: Satellite Beach; 4 miles N of Eau Gallie Causeway, entry road from South Patrick Drive. Perimeter marsh along E, W and S sides of old spoil island.

Source of planting information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

Source of subsequent assessment information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

References or reports:

Other species planted in addition to *Spartina alterniflora* : *Paspalum vaginatum* and *Distichlis spicata* were planted on 2-ft centers; mangroves were planted on 4-ft centers.

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? No

General comments about the marsh:

- Freshly planted in Sept-Dec 1993; appears to be doing well, no mortality as of Feb 1994. *Spartina alterniflora* clumps were planted on 2-ft centers.
- Marsh name: Lansing Island

Marsh Inventory No.: 749 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 05/01/93

Area Planted (1acre = 0.4047 ha): 5.260 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit:

Location of the marsh.

State: FL **County:** Brevard **Latitude (N):** 28 8' 30" **Longitude (W):** 80 36' 30"

Bay and sub-bays: Banana River

Directions to the site: Melbourne; access by boat only, N of Eau Gallie Causeway; created wetlands consisting of excavated tidal creeks and tidal flats. Entire W side of island returned to wetland elevations.

Source of planting information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

Source of subsequent assessment information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

References or reports:

Other species planted in addition to *Spartina alterniflora* : *Paspalum vaginatum*, *Sporobolus virginicus*, and *Batis maritima* were planted on 2-ft centers.

Percent vegetative cover attained by the planted marsh: 85% cover, 95% survival; no die back seen after 6 mo. Dec-93

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? No

General comments about the marsh:

- *Spartina alterniflora* clumps were planted on 2-ft centers beginning in May and continuing through Dec 1993. Marsh appears to be doing well, no mortality as of Feb 1994.

- Marsh name: Samson Island

Marsh Inventory No.: 751 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 09/01/86

Area Planted (1acre = 0.4047 ha): 1.670 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: David McWilliams (407)777-4111 is the contact person

Location of the marsh.

State: FL **County:** Brevard **Latitude (N):** 28 11' **Longitude (W):** 80 37'

Bay and sub-bays: Banana River

Directions to the site: Satellite Beach; 3.5 miles N of Eau Gallie Causeway, entry road off South Patrick Blvd. Perimeter marsh along E, W and S sides of old spoil island.

Source of planting information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

Source of subsequent assessment information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

References or reports:

Other species planted in addition to *Spartina alterniflora* : *Rhizophora mangle*, *Avicennia mangle*, and *Laguncularia racemosa* were planted on 5-ft centers; *Sporobolus virginicus* was planted on 2-ft centers.

Percent vegetative cover attained by the planted marsh: 100% cover by *Spartina* and mangroves.

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Not assessed.

General comments about the marsh:

- *Spartina alterniflora* clumps were planted on 2-ft centers. Some *Spartina* is being displaced by volunteer mangroves. After a slow start the grasses have filled in to almost complete coalescence.

- The mangroves have formed a 6-10' hedge along the entire shoreline. Marsh name: Windward Cove

Marsh Inventory No.: 783 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 08/01/87

Area Planted (1acre = 0.4047 ha): 0.490 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Jack Bennett (407) 783-3060 is the contact person.

Location of the marsh.

State: FL **County:** Brevard **Latitude (N):** 28 21' **Longitude (W):** 80 36' 30"

Bay and sub-bays: Banana River

Directions to the site: Cocoa Beach; A-1-A half mile S of State Rd. 520; narrow fringing site along W and S shoreline protected by coquina boulder revetment.

Source of planting information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

Source of subsequent assessment information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

References or reports:

Other species planted in addition to *Spartina alterniflora* : *Paspalum vaginatum* was planted on 2-ft centers; *Avicennia germinans* and *Laguncularia racemosa* were planted on 5-ft centers.

Percent vegetative cover attained by the planted marsh: 95-100% as of May-93

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Not assessed.

General comments about the marsh:

- Much of the *Spartina alterniflora* has been replaced by *Scirpus robustus* and mangroves. This is a replacement planting where mangroves had displaced the marsh, and then been killed by freezes.

- Marsh name: Rock Pointe. *S. alterniflora* clumps were planted 2' on center (=o.c.)

Marsh Inventory No.: 786 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 08/01/90

Area Planted (1acre = 0.4047 ha): 0.648 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Arnold Mallard (407) 730-2404 is the contact person.

Location of the marsh.

State: FL **County:** Brevard **Latitude (N):** 28 32' **Longitude (W):** 80 36' 40"

Bay and sub-bays: Banana River

Directions to the site: Cape Canaveral; adjacent to Titan missile complex (restricted access); rectangular created wetland connected to the Banana River.

Source of planting information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

Source of subsequent assessment information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

References or reports:

Other species planted in addition to *Spartina alterniflora* : *Bacopa carolinianum*, *Eleocharis cellulosa*, *Sesuvium portulacastrum*, *Batis maritima*, *Salicornia virginica*, *Sporobolus virginicus*, *Distichlis spicata*, and *Spartina bakeri* were planted on 3-ft centers; *Laguncularia racemosa* on 5-ft centers.

Percent vegetative cover attained by the planted marsh: 60-70% cover as of Oct-93

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Not assessed.

General comments about the marsh:

- *Spartina alterniflora* occupies only the waterward edge; the majority of the site is covered by *Distichlis* and other high marsh species. Mangroves have survived. Growth is good and coverage should reach 90-100% by 1995.

- Marsh name: Canaveral AFS-Solid Motor Assembly Building. *S. alterniflora* clumps were planted on 3-ft centers.

Marsh Inventory No.: 773 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 08/01/84

Area Planted (1acre = 0.4047 ha): 0.138 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Bill Moord (407) 783-0037 is the contact person.

Location of the marsh.

State: FL **County:** Brevard **Latitude (N):** 28 23' 25" **Longitude (W):** 80 36' 50"

Bay and sub-bays: Banana River

Directions to the site: Cape Canaveral; A-1-A; 1 mile S of State Rd. 528; narrow shelf along a canal bordered by wooden bulkhead.

Source of planting information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

Source of subsequent assessment information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

References or reports:

Other species planted in addition to *Spartina alterniflora* : *Rhizophora mangle*

Percent vegetative cover attained by the planted marsh: 80-90% as of March-93

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Not assessed.

General comments about the marsh:

- Planted between Aug-84 and June-85. Most *Spartina* has been displaced by a dense mangrove hedge. Some *Paspalum* and *Distichlis* invasion on the shelf, and isolated heavy patches of *Typha* combine to cover 90% of the shelf with 100% cover.

- Marsh name: Treasure Island Club *S. alterniflora* clumps were planted 2'on center (=o.c.)

Marsh Inventory No.: 186 **Information Rank (1-5):** 5 **State:** FL

Date Planted:

Area Planted (1acre = 0.4047 ha): 0.040 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Mariner's Square

Location of the marsh.

State: FL **County:** Brevard **Latitude (N):** 28 21' 50" ? **Longitude (W):** 80 40' 50" ?

Bay and sub-bays: Banana River

Directions to the site: Mariner's Square; Cocoa Beach

Source of planting information: Dr. Tom Roberts, Tennessee Tech University, Cookeville, TN 38505

Source of subsequent assessment information: (Roberts, 1991)

References or reports: (Roberts, 1991)

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh: 49%, 1987

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Yes. Fisheries species, birds, mammals, and epibenthic snails.

General comments about the marsh:

- No birds; a mouse; only a few killifish. Very small fringe marsh lacks habitat complexity needed to support fish; should not have been in the study because it was so small.

Marsh Inventory No.: 250 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 1983

Area Planted (1acre = 0.4047 ha): 0.400 hectares

COE Permit No.: None **Date of COE Permit:**

Originator or Applicant for permit: Costa del Sol Condominiums

Location of the marsh.

State: FL **County:** Brevard **Latitude (N):** 28 22' 15" **Longitude (W):** 80 36' 18"

Bay and sub-bays: Banana River; Banana River Lagoon

Directions to the site: Banana River Lagoon, Cocoa Beach

Source of planting information: Dr. Robin Lewis, III., Lewis Environmental Services, Inc., P.O. Box 20005, Tampa, FL 33622-0005, (813) 889-9684

Source of subsequent assessment information: (Crewz & Lewis, 1991; T. H. Roberts, 1991)

References or reports: (Crewz & Lewis, 1991; T. H. Roberts, 1991)

Other species planted in addition to *Spartina alterniflora* : *Rhizophora mangle*

Percent vegetative cover attained by the planted marsh: 30% (Crews & Lewis); 67% (Roberts); 1986

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Yes, by Roberts. Fisheries species, birds, mammals, and epibenthic snails.

General comments about the marsh:

- *Spartina alterniflora* limited by poor flushing; bad topography. Area is receiving much freshwater inflow and southern cattail, *Typha domingensis*, is invading. Mixed failure. Roberts found a few birds, a rat, signs of raccoons, and very many killifish.

Marsh Inventory No.: 251 **Information Rank (1-5):** 5 **State:** FL

Date Planted: 1983

Area Planted (1acre = 0.4047 ha): 0.120 hectares

COE Permit No.: None **Date of COE Permit:**

Originator or Applicant for permit: Fountain Cove Condominiums

Location of the marsh.

State: FL **County:** Brevard **Latitude (N):** 28 20' 47" **Longitude (W):** 80 36' 32"

Bay and sub-bays: Banana River; Banana River Lagoon

Directions to the site: Fountain Cove, Banana River Lagoon, Cocoa Beach

Source of planting information: Dr. Robin Lewis, III., Lewis Environmental Services, Inc., P.O. Box 20005, Tampa, FL 33622-0005, (813) 889-9684

Source of subsequent assessment information: (Crewz & Lewis, 1991; T. H. Roberts, 1991)

References or reports: (Crewz & Lewis, 1991; T. H. Roberts, 1991)

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh: 34% (Crewz & Lewis), 51% (Roberts); 1986

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Yes. Fisheries species, birds, mammals, and epibenthic snails.

General comments about the marsh:

- *Spartina alterniflora* limited by poor flushing and topography. Deemed a failure by Crewz and Lewis. Roberts found no birds or mammals, but very many killifish in the marsh.

Marsh Inventory No.: 785 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 06/01/79

Area Planted (1acre = 0.4047 ha): 0.554 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Bob Lynds (407) 723-2095 is the contact person.

Location of the marsh.

State: FL **County:** Brevard **Latitude (N):** 28 22' 20" **Longitude (W):** 80 42' 10"

Bay and sub-bays: Banana River; Newfound Harbor

Directions to the site: Merritt Island; State Rd. 3, 4 miles N of State Rd. 520; fringing intertidal shoreline along the Indian River and boat basin.

Source of planting information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

Source of subsequent assessment information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

References or reports:

Other species planted in addition to *Spartina alterniflora* : *Spartina patens*, *Paspalum vaginatum* and *Distichlis spicata* were planted on 2-ft centers.

Percent vegetative cover attained by the planted marsh: 90-100% cover by combined species as of March-93

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Not assessed.

General comments about the marsh:

- *Spartina alterniflora* clumps were planted on 2-ft centers. *S. alterniflora* is being replaced by high marsh grasses as the elevation has built up behind the coquina boulder revetment.

- Marsh name: Egret's Cove.

Marsh Inventory No.: 775 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 02/01/89

Area Planted (1acre = 0.4047 ha): 1.570 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Ron Taylor (407) 783-1320 is the contact person.

Location of the marsh.

State: FL **County:** Brevard **Latitude (N):** 28 21' 30" **Longitude (W):** 80 41' 05"

Bay and sub-bays: Banana River; Newfound Harbor

Directions to the site: Merritt Island; State Rd 520; behind and S of Merritt Square Mall; large cove connected by wide inlet to Newfound Harbor.

Source of planting information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

Source of subsequent assessment information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

References or reports:

Other species planted in addition to *Spartina alterniflora* : *Juncus roemerianus* was planted on 3-ft centers; *Rhizophora mangle* was also planted.

Percent vegetative cover attained by the planted marsh: 90-100% cover as of Aug-93

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Not assessed.

General comments about the marsh:

- The large tidal flats are completely grown in with *Spartina*, *Juncus* and other high marsh species. Volunteer mangroves (mostly white) are taking over in some areas. There is a sparse area in the NW corner or the planted shelf.

- Marsh name: Harbor Woods *S. alterniflora* clumps were planted on 3-ft centers.

Marsh Inventory No.: 761 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 03/01/85

Area Planted (1acre = 0.4047 ha): 1.137 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: William Lambert (407) 452-7946 is the contact person.

Location of the marsh.

State: FL **County:** Brevard **Latitude (N):** 28 24' 15" **Longitude (W):** 80 41' 55"

Bay and sub-bays: Banana River; Newfound Harbor; Sykes Creek

Directions to the site: Merritt Island; Hwy 528 at S.R. 3 on Merritt Island; enclosed pond connected to Newfound Harbor via culverts.

Source of planting information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

Source of subsequent assessment information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

References or reports:

Other species planted in addition to *Spartina alterniflora* : *Distichlis spicata* was planted on 3-ft centers.

Percent vegetative cover attained by the planted marsh: Much of the *Spartina* has been replaced by *Eleocharis* and *Scirpus robustus*, but the marsh as a whole has achieved 85-100% cover, as of May-92.

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Not assessed.

General comments about the marsh:

- Freshwater species have invaded. *Typha* has taken over a large area. There are periodic heavy freshwater inflows.
- Marsh name: Lambert Seafoods, Inc. *S. alterniflora* clumps were planted on 3-ft centers.

Marsh Inventory No.: 767 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 12/01/89

Area Planted (1acre = 0.4047 ha): 1.267 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Reese Kessler (407) 696-4222 is the contact person.

Location of the marsh.

State: FL **County:** Brevard **Latitude (N):** 28 26' 00" **Longitude (W):** 80 41' 50"

Bay and sub-bays: Banana River; Newfound Harbor; Sykes Creek

Directions to the site: Merritt Island; State Rd. 3, 3 miles N of Hwy 528; created wetlands, mostly high marsh adjacent to tidal ditch connected to Sykes Creek.

Source of planting information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

Source of subsequent assessment information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

References or reports:

Other species planted in addition to *Spartina alterniflora* : *Distichlis spicata*, *Salicornia virginica*, and *Batis maritima* were planted on 2-ft centers; *Avicennia germinans* and *Spartina bakeri*.

Percent vegetative cover attained by the planted marsh: 100% by high marsh species, and by fring *S. alterniflora* as of Oct-93

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Not assessed.

General comments about the marsh:

- *Spartina alterniflora* clumps were planted on 2-ft centers. Growing well.
- Marsh name: Savannahs

Marsh Inventory No.: 195 **Information Rank (1-5):** 5 **State:** FL

Date Planted: 1986

Area Planted (1acre = 0.4047 ha): 0.010 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit:

Location of the marsh.

State: FL **County:** Brevard **Latitude (N):** **Longitude (W):**

Bay and sub-bays: Canaveral Bay

Directions to the site: Canaveral Bay; Cocoa Beach

Source of planting information: Dr. Tom Roberts, Tennessee Tech University, Cookeville, TN 38505

Source of subsequent assessment information: (Roberts, 1991)

References or reports: (Roberts, 1991)

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh: 73%, 1987

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Yes. Fisheries species, birds, mammals, and epibenthic snails.

General comments about the marsh:

- No birds; no rats; raccoon tracks; many small killifish.

Marsh Inventory No.: 772 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 01/01/88

Area Planted (1acre = 0.4047 ha): 0.271 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Stan Rosenbaum (904) 446-0346 is the contact person.

Location of the marsh.

State: FL **County:** Brevard **Latitude (N):** **Longitude (W):**

Bay and sub-bays: ICW

Directions to the site: Flagler Beach; A-1-A 3 miles N of Hwy 100; created intertidal shelf along ICW protected by coquina rock revetment.

Source of planting information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

Source of subsequent assessment information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

References or reports:

Other species planted in addition to *Spartina alterniflora* : *Sporobolus virginicus*, *Distichlis spicata*, *Juncus roemerianus* and *Paspalum vaginatum* were planted on 2-ft centers.

Percent vegetative cover attained by the planted marsh: 100% cover by *Spartina* and highmarsh species.

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Not assessed.

General comments about the marsh:

- Erosion due to boat wakes is severe in this region, and is why the rock break water was built to protect the planting and shoreline. The planted intertidal community is healthy and has generally coalesced to complete ground coverage.

- The *Juncus* marsh planted adjacent to an existing natural needlerush marsh has also achieved 100% aerial coverage and is indistinguishable from the natural marsh. Marsh name: Ocean Marina.

Marsh Inventory No.: 757 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 04/01/90

Area Planted (1acre = 0.4047 ha): 1.012 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Bill Sargent (904) 329-4500 is the contact person.

Location of the marsh.

State: FL **County:** Brevard **Latitude (N):** **Longitude (W):**

Bay and sub-bays: Indian River

Directions to the site: Several dredged material deposition islands and peninsulas in the Indian River, 7 sites.

Source of planting information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

Source of subsequent assessment information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

References or reports:

Other species planted in addition to *Spartina alterniflora* : *Avicennia germinans* was planted on 4-ft centers.

Percent vegetative cover attained by the planted marsh: 100% cover in 20% of the planted area; 25% of the area was lost to erosion; remaining 55% are coalescing and have reached about 70% cover as of June-93.

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Not assessed.

General comments about the marsh:

- *Spartina alterniflora* clumps were planted on 2-ft centers. Some areas were subject to severe erosion and were washed out. No protective devices were used to protect the plantings.

- Project name: S.W.I.M. in Brevard and Indian River counties.

Marsh Inventory No.: 762 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 02/01/90

Area Planted (1acre = 0.4047 ha): 1.728 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Mark Hesse (407) 452-7631 is the contact person.

Location of the marsh.

State: FL **County:** Brevard **Latitude (N):** 28 26' 30" **Longitude (W):** 80 43' 25"

Bay and sub-bays: Indian River

Directions to the site: Merritt Island; State Road 3, 6 miles N of Hwy 528; protected cove and 3 sites on the Indian River.

Source of planting information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

Source of subsequent assessment information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

References or reports:

Other species planted in addition to *Spartina alterniflora* : *Distichlis spicata* and *Paspalum vaginatum* were planted on 3-ft centers; *Avicennia germinans* was planted from 1-gal containers on 10-ft centers.

Percent vegetative cover attained by the planted marsh: 80-100% cover attained in most *Spartina* areas as of May-92

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Not assessed.

General comments about the marsh:

- The grasses are healthy in all of the planted areas. The retention area and the Big Lake shoreline are experiencing some nuisance species invasion. The mangroves were all killed by freezes, but some mangroves are growing in the *Spartina*.

- Marsh name: Indian Bay *S. alterniflora* clumps were planted on 3-ft centers.

Marsh Inventory No.: 776 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 05/01/86

Area Planted (1acre = 0.4047 ha): 0.465 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Ed Poe (407) 267-1841 is the contact person.

Location of the marsh.

State: FL **County:** Brevard **Latitude (N):** 28 37' **Longitude (W):** 80 48' 10"

Bay and sub-bays: Indian River

Directions to the site: Titusville; State Rd. 406 at Indian River; shoreline marsh created behind a protective coquina rock revetment.

Source of planting information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

Source of subsequent assessment information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

References or reports:

Other species planted in addition to *Spartina alterniflora* : *Distichlis spicata*, *Sporobolus virginicus*, *Paspalum vaginatum* were planted on 2-ft centers; *Laguncularia racemosa*.

Percent vegetative cover attained by the planted marsh: 95% cover by the combined species as of Sept-93

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Not assessed.

General comments about the marsh:

- The *Spartina* has diminished in favor of the other high marsh grass species. Most of the mangroves were killed by freezes. Potential erosion by wave action was averted by a coquina boulder break water.

- Marsh name: Harbor Town. *S. alterniflora* clumps were planted on 2-ft centers.

Marsh Inventory No.: 780 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 09/01/82

Area Planted (1acre = 0.4047 ha): 0.255 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Steve Peffer (407) 453-9517 is the contact person.

Location of the marsh.

State: FL **County:** Brevard **Latitude (N):** 28 04' 40" **Longitude (W):** 80 35' 40"

Bay and sub-bays: Indian River

Directions to the site: Melbourne; U.S. Hwy 192 at U.S. 1; small pocket marsh created behind coquina breakwater.

Source of planting information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

Source of subsequent assessment information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

References or reports:

Other species planted in addition to *Spartina alterniflora* : *Distichlis spicata* and *Paspalum vaginatum* were planted on 2-ft centers; *Rhizophora mangle* and *Laguncularia racemosa*.

Percent vegetative cover attained by the planted marsh: 100% as of Sept-93

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Not assessed.

General comments about the marsh:

- *Spartina* is being replaced by volunteer mangroves. Most planted mangroves died in subsequent freezes. Elevations have favored a transition to high marsh dominated by *Distichlis* and *Paspalum*.

- Marsh name: Front Street Park. *S. alterniflora* clumps were planted on 2-ft centers.

Marsh Inventory No.: 221 **Information Rank (1-5):** 5 **State:** FL

Date Planted: 1985

Area Planted (1acre = 0.4047 ha): 0.120 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Harbor Isles

Location of the marsh.

State: FL **County:** Brevard **Latitude (N):** 28 21' 50" ? **Longitude (W):** 80 40' 50" ?

Bay and sub-bays: Newfound Harbor, Banana River;

Directions to the site: Harbor Isles; Cocoa Beach

Source of planting information: Dr. Tom Roberts, Tennessee Tech University, Cookeville, TN 38505

Source of subsequent assessment information: (Roberts, 1991)

References or reports: (Roberts, 1991)

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh: 79%, 1987

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Yes. Fisheries species, birds, mammals, and epibenthic snails.

General comments about the marsh:

- Marsh was successful. No birds; no rats; raccoon tracks; many small killifish.
-

Marsh Inventory No.: 750 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 07/01/90

Area Planted (1acre = 0.4047 ha): 3.950 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Paul Douglas (305) 254-2106 is the contact person

Location of the marsh.

State: FL **County:** Dade **Latitude (N):** 25 44' 00?" **Longitude (W):** 80 13' 00?"

Bay and sub-bays: Biscayne Bay; Deering Bay; flushing channel

Directions to the site: Miami; Old Cutler Road at Chapman Field; flushing channel connecting bay to Biscayne Bay.

Source of planting information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

Source of subsequent assessment information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

References or reports:

Other species planted in addition to *Spartina alterniflora* : *Sporobolus virginicus* was planted on 2-ft centers; *Eleocharis cellulosa*, *Sagittaria lancifolia*, *Scirpus validus*, *Rhizophora mangle*.

Percent vegetative cover attained by the planted marsh: After replanting: 70-100% aerial coverage.

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Not assessed.

General comments about the marsh:

- Water levels and salinities have fluctuated wildly at the site. 25,000 plants have died. Remedial action was to replant with *Spartina alterniflora* and *Sporobolus virginicus*. The *Spartina* is beginning to trap mangrove seeds.

- Marsh name: Deering Bay *S. alterniflora* clumps were planted on 2-ft centers.

Marsh Inventory No.: 774 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 06/01/87

Area Planted (1acre = 0.4047 ha): 0.567 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Jack Tallon (904) 829-6727 is the contact person.

Location of the marsh.

State: FL **County:** Duval **Latitude (N):** 30 17' 00?" **Longitude (W):** 81 25' 00?"

Bay and sub-bays: Pablo Creek

Directions to the site: Jacksonville; St. John's Bluff Rd.; 1 mile N of Butler Blvd.; created extension to natural tidal marsh.

Source of planting information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

Source of subsequent assessment information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

References or reports:

Other species planted in addition to *Spartina alterniflora* : *Juncus roemerianus*

Percent vegetative cover attained by the planted marsh: 100% cover as of Apr-93

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Not assessed.

General comments about the marsh:

- *Spartina alterniflora* clumps were planted on 2-ft centers. Blended into the adjacent natural marshes; are indistinguishable.
- Marsh name: Coastal Point

Marsh Inventory No.: 763 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 10/01/84

Area Planted (1acre = 0.4047 ha): 0.417 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Mike Antonopoulos (904) 353-8471 is the contact person.

Location of the marsh.

State: FL **County:** Duval **Latitude (N):** 30 33' 00?" **Longitude (W):** 81 36' 30"

Bay and sub-bays: St. John's River

Directions to the site: Jacksonville; N end of University Blvd.; semi-enclosed cove; connected to the St. John's River by shallow inlets.

Source of planting information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

Source of subsequent assessment information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

References or reports:

Other species planted in addition to *Spartina alterniflora* : *Sporobolus virginicus* was planted on 2-ft centers.

Percent vegetative cover attained by the planted marsh: 100% cover where elevation was high enough. May-92

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Not assessed.

General comments about the marsh:

- High water levels due to poorly constructed flushing connection were corrected; healthy saltmarsh community with areas of open water was established with open pools in the deeper areas.
- Marsh name: Sandy Shores *S. alterniflora* clumps were planted on 3-ft centers.

Marsh Inventory No.: 771 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 04/01/85

Area Planted (1acre = 0.4047 ha): 0.202 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Bruce Taylor (904) 731-7040 is the contact person.

Location of the marsh.

State: FL **County:** Duval **Latitude (N):** 30 33' 00?" **Longitude (W):** 81 31' 40?"

Bay and sub-bays: St. John's River

Directions to the site: Jacksonville; end of Ft. Caroline Rd.; wide planted area along the St. John's River protected by granite rock revetment, at St. John's Landing.

Source of planting information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

Source of subsequent assessment information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

References or reports:

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh: 100% cover as of April-93.

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Not assessed.

General comments about the marsh:

- Large tidal amplitudes and currents at this location. A granite break water surrounds the planted area, and acted as protection from wave erosion. The marsh reached coalescence within 3 months of planting, and has matured into a dynamic ecosystem.
- Marsh name: St. John's Landing *S. alterniflora* clumps were planted on 3-ft centers.

Marsh Inventory No.: 787 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 10/01/88

Area Planted (1acre = 0.4047 ha): 3.966 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Steve Duval (904) 642-8507 is the contact person.

Location of the marsh.

State: FL **County:** Duval **Latitude (N):** 30 31' 00?" **Longitude (W):** 81 34' 00?"

Bay and sub-bays: St. John's River

Directions to the site: Jacksonville; Hwy. 9A over St. John's River; wide marsh at S abutment and central island.

Source of planting information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

Source of subsequent assessment information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

References or reports:

Other species planted in addition to *Spartina alterniflora* : *Distichlis spicata* and *Sporobolus virginicus* were planted on 3-ft centers.

Percent vegetative cover attained by the planted marsh: 95-100% cover as of June-93.

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Not assessed.

General comments about the marsh:

- *Spartina alterniflora* clumps were planted on 3-ft centers. Both areas have matured into dense healthy saltmarsh communities.
- Marsh name: Dames Point Bridge - South Abutment and Nichols Creek.

Marsh Inventory No.: 241 **Information Rank (1-5):** 5 **State:** FL

Date Planted:

Area Planted (1acre = 0.4047 ha): 0.200 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: St. John's Landing

Location of the marsh.

State: FL **County:** Duval **Latitude (N):** **Longitude (W):**

Bay and sub-bays: St. John's River

Directions to the site: St. John's Landing; Jacksonville

Source of planting information: Dr. Tom Roberts, Tennessee Tech University, Cookeville, TN 38505

Source of subsequent assessment information: (Roberts, 1991)

References or reports: (Roberts, 1991)

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh: 63%, 1987

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Yes. Birds, mammals, and epibenthic snails.

General comments about the marsh:

- Few birds; a mouse;

Marsh Inventory No.: 782 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 05/01/87

Area Planted (1acre = 0.4047 ha): 0.352 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Pete Peterson (904) 264-2011 is the contact person.

Location of the marsh.

State: FL **County:** Duval **Latitude (N):** 30 24' 30" **Longitude (W):** 81 40' 00?"

Bay and sub-bays: St. John's River; Trout River

Directions to the site: Jacksonville; Hecksher Dr. W of I-95; created extension of natural marsh connected to the Trout River by an excavated tidal creek.

Source of planting information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

Source of subsequent assessment information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

References or reports:

Other species planted in addition to *Spartina alterniflora* : *Juncus roemerianus* was planted on 3-ft centers.

Percent vegetative cover attained by the planted marsh: 95-100% cover as of June-93

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Not assessed.

General comments about the marsh:

- *Spartina alterniflora* clumps were planted on 3-ft centers. Saltmarsh created next to a natural *Spartina-Juncus* marsh was very successful.
- Marsh name: River Crossing.

Marsh Inventory No.: 373 **Information Rank (1-5):** 3 **State:** FL

Date Planted: 1976

Area Planted (1acre = 0.4047 ha): hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit:

Location of the marsh.

State: FL **County:** Hillsborough **Latitude (N):** 27 48' 28" **Longitude (W):** 82 25' 00"

Bay and sub-bays: Tampa Bay

Directions to the site: North side of Fishhook Island

Source of planting information: Dr. Robin Lewis, III., Lewis Environmental Services, Inc., P.O. Box 20005, Tampa, FL 33622-0005, (813) 889-9684

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh: 1986

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Marsh name: Fishhook Island.

Marsh Inventory No.: 374 **Information Rank (1-5):** 3 **State:** FL

Date Planted: 1991

Area Planted (1acre = 0.4047 ha): hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit:

Location of the marsh.

State: FL **County:** Hillsborough **Latitude (N):** 27 44' 33" **Longitude (W):** 82 28' 15"

Bay and sub-bays: Tampa Bay

Directions to the site: Pelican Cove, Simmons Park

Source of planting information: Dr. Robin Lewis, III., Lewis Environmental Services, Inc., P.O. Box 20005, Tampa, FL 33622-0005, (813) 889-9684

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh: 1986

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Marsh name: Simmons Park.

Marsh Inventory No.: 376 **Information Rank (1-5):** 3 **State:** FL

Date Planted:

Area Planted (1acre = 0.4047 ha): 0.004 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Tampa Electric ?

Location of the marsh.

State: FL **County:** Hillsborough **Latitude (N):** 27 38' 40" **Longitude (W):** 82 33' 25"

Bay and sub-bays: Tampa Bay

Directions to the site: Ft. Lonely

Source of planting information: Dr. Robin Lewis, III., Lewis Environmental Services, Inc., P.O. Box 20005, Tampa, FL 33622-0005, (813) 889-9684

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Marsh name: Tampa Electric Shoreline.

Marsh Inventory No.: 263 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 1985

Area Planted (1acre = 0.4047 ha): 1.214 hectares

COE Permit No.: 83D-2158 **Date of COE Permit:** 1983

Originator or Applicant for permit: Harbour Island

Location of the marsh.

State: FL **County:** Hillsborough **Latitude (N):** 27 56' 08" **Longitude (W):** 82 26' 52"

Bay and sub-bays: Tampa Bay area; Sparkman Channel

Directions to the site: Seddon Island; along Sparkman Channel; E-side of Harbor Island, Site A.

Source of planting information: Dr. Robin Lewis, III., Lewis Environmental Services, Inc., P.O. Box 20005, Tampa, FL 33622-0005, (813) 889-9684

Source of subsequent assessment information: (Crewz & Lewis, 1991)

References or reports: (Crewz & Lewis, 1991)

Other species planted in addition to *Spartina alterniflora* : *Rhizophora mangle*

Percent vegetative cover attained by the planted marsh: 8%, 1986

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? no

General comments about the marsh:

- *Spartina alterniflora* was growing, but ship wakes and flotsam debris appears to have damaged patches of the growth. Mixed success.

Marsh Inventory No.: 264 **Information Rank (1-5):** 5 **State:** FL

Date Planted: 1985

Area Planted (1acre = 0.4047 ha): 0.400 hectares

COE Permit No.: 83D-2158 **Date of COE Permit:** 1983

Originator or Applicant for permit: Harbour Island

Location of the marsh.

State: FL **County:** Hillsborough **Latitude (N):** 27 56' 08" **Longitude (W):** 82 26' 52"

Bay and sub-bays: Tampa Bay area; Sparkman Channel

Directions to the site: Seddon Island; along Sparkman Channel; E-side of Harbor Island, Site B.

Source of planting information: Dr. Robin Lewis, III., Lewis Environmental Services, Inc., P.O. Box 20005, Tampa, FL 33622-0005, (813) 889-9684

Source of subsequent assessment information: (Crewz & Lewis, 1991); T. H. Roberts, 1991)

References or reports: (Crewz & Lewis, 1991); T. H. Roberts, 1991)

Other species planted in addition to *Spartina alterniflora* : *Rhizophora mangle*

Percent vegetative cover attained by the planted marsh: 27% (Crewz & Lewis), 60% (Roberts); 1986

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Yes. Fisheries species, birds, mammals, and epibenthic snails.

General comments about the marsh:

- *Spartina alterniflora* was growing, but uneven elevations created pooling that drowned some plants. This area was behind a berm from Site A, thus semi-protected from ship wakes. Deemed a mixed success by Crewz and Lewis.

- Roberts found a few birds, raccoon and rabbit tracks, and many killifish.

Marsh Inventory No.: 375 **Information Rank (1-5):** 4 **State:** FL

Date Planted:

Area Planted (1acre = 0.4047 ha): 0.004 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit:

Location of the marsh.

State: FL **County:** Hillsborough **Latitude (N):** 27 41'15" **Longitude (W):** 82 31' 15"

Bay and sub-bays: Tampa Bay; Cockroach Channel

Directions to the site: At end of Cockroach Bay Road

Source of planting information: Dr. Robin Lewis, III., Lewis Environmental Services, Inc., P.O. Box 20005, Tampa, FL 33622-0005, (813) 889-9684

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Marsh name: Cockroach Bay Boat Ramp.

Marsh Inventory No.: 368 **Information Rank (1-5):** 3 **State:** FL

Date Planted: 1989

Area Planted (1acre = 0.4047 ha): 1.740 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit:

Location of the marsh.

State: FL **County:** Hillsborough **Latitude (N):** 27 53' 10" **Longitude (W):** 82 24' 15"

Bay and sub-bays: Tampa Bay; Hillsborough Bay.

Directions to the site: Off Tamiami Trail, south of Port Sutton.

Source of planting information: Dr. Robin Lewis, III., Lewis Environmental Services, Inc., P.O. Box 20005, Tampa, FL 33622-0005, (813) 889-9684

Source of subsequent assessment information: Dr. Brandt Henningsen, SW Florida Water Management District, Surface Water Improvement and Management Program, 7601 Highway 301 N, Tampa, FL 33637, (813) 985-7481 ext. 2202

References or reports: (Crewz & Lewis, 1991)

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Yes. Birds and fish.

General comments about the marsh:

- Some sampling has been done by SWIM. Comments that the marsh is coalescing nicely and more animals are using the area.
- Marsh name: Delaney Creek.

Marsh Inventory No.: 367 **Information Rank (1-5):** 3 **State:** FL

Date Planted: 1991

Area Planted (1acre = 0.4047 ha): hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Tampa Port Authority

Location of the marsh.

State: FL **County:** Hillsborough **Latitude (N):** 27 53' 50" **Longitude (W):** 82 24' 45"

Bay and sub-bays: Tampa Bay; Hillsborough Bay.

Directions to the site: Off US Bus.45(= Rd. 41), south of Port Sutton.

Source of planting information: Dr. Robin Lewis, III., Lewis Environmental Services, Inc., P.O. Box 20005, Tampa, FL 33622-0005, (813) 889-9684

Source of subsequent assessment information:

References or reports: (Crewz & Lewis, 1991)

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh: 1986

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Marsh name: Tampa Port Authority

-

Marsh Inventory No.: 647 **Information Rank (1-5):** 3 **State:** FL

Date Planted: 1993

Area Planted (1acre = 0.4047 ha): 4.897 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: SWIM and City of Tampa

Location of the marsh.

State: FL **County:** Hillsborough **Latitude (N):** 27 50' 00" **Longitude (W):** 82 28' 15"

Bay and sub-bays: Tampa Bay; Hillsborough Bay.

Directions to the site: At the end of Interbay Peninsula, on the MacDill AFB, in the SE corner.

Source of planting information: Dr. Brandt Henningsen, SW Florida Water Management District, Surface Water Improvement and Management Program, 7601 Highway 301 N, Tampa, FL 33637, (813) 985-7481 ext. 2202

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Will be completed in Dec '93 or spring '94.
- Marsh name: MacDill AFB, No.2 - N of Gadsden Point.

Marsh Inventory No.: 677 **Information Rank (1-5):** 3 **State:** FL

Date Planted: 12/01/93

Area Planted (1acre = 0.4047 ha): 7.123 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: U.S. Air Force and SWIM

Location of the marsh.

State: FL **County:** Hillsborough **Latitude (N):** 27 49' 25" **Longitude (W):** 82 23' 30"

Bay and sub-bays: Tampa Bay; Hillsborough Bay.

Directions to the site: MacDill Air Force Base; Gadsden Point, Tampa.

Source of planting information: Dr. Brandt Henningsen, SW Florida Water Management District, Surface Water Improvement and Management Program, 7601 Highway 301 N, Tampa, FL 33637, (813) 985-7481 ext. 2202

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

-
-

Marsh Inventory No.: 371 **Information Rank (1-5):** 3 **State:** FL

Date Planted:

Area Planted (1acre = 0.4047 ha): hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit:

Location of the marsh.

State: FL **County:** Hillsborough **Latitude (N):** 27 50' 56" **Longitude (W):** 82 24' 30"

Bay and sub-bays: Tampa Bay; Hillsborough Bay.

Directions to the site: North side of Bird Island

Source of planting information: Dr. Robin Lewis, III., Lewis Environmental Services, Inc., P.O. Box 20005, Tampa, FL 33622-0005, (813) 889-9684

Source of subsequent assessment information:

References or reports: (Crewz & Lewis, 1991)

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh: 1986

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Marsh name: Bird Island.

Marsh Inventory No.: 617 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 1979

Area Planted (1acre = 0.4047 ha): 0.405 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit:

Location of the marsh.

State: FL **County:** Hillsborough **Latitude (N):** 27 52' 24" **Longitude (W):** 82 25' 45"

Bay and sub-bays: Tampa Bay; Hillsborough Bay.

Directions to the site: Dredged material Island 2D

Source of planting information: Dr. Robin Lewis, III., Lewis Environmental Services, Inc., P.O. Box 20005, Tampa, FL 33622-0005, (813) 889-9684

Source of subsequent assessment information: Dr. Brandt Henningsen, SWIM, (813) 985-7481 ext. 2202

References or reports:

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Furnished *Spartina alterniflora* transplant material for another project. This marsh's recovery was evaluated (Courtney 1991). The marsh filled in the 1-m wide harvested zones within a growing season.

- Henningsen said this marsh was coalescing nicely; successful.

Marsh Inventory No.: 679 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 1992

Area Planted (1acre = 0.4047 ha): 1.214 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit:

Location of the marsh.

State: FL **County:** Hillsborough **Latitude (N):** 27 51' 20" **Longitude (W):** 82 32' 50"

Bay and sub-bays: Tampa Bay; Hillsborough Bay.

Directions to the site: A little N of the end of Interbay Peninsula and west of MacDill AFB.

Source of planting information: Dr. Brandt Henningsen, SW Florida Water Management District, Surface Water Improvement and Management Program, 7601 Highway 301 N, Tampa, FL 33637, (813) 985-7481 ext. 2202

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Will be finished in 1993.
- Marsh name: Picnic Island.

Marsh Inventory No.: 370 **Information Rank (1-5):** 4 **State:** FL

Date Planted:

Area Planted (1acre = 0.4047 ha): hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit:

Location of the marsh.

State: FL **County:** Hillsborough **Latitude (N):** 27 52' 00" **Longitude (W):** 82 24' 13"

Bay and sub-bays: Tampa Bay; Hillsborough Bay.

Directions to the site: West shoreline of East Tampa.

Source of planting information: Dr. Robin Lewis, III., Lewis Environmental Services, Inc., P.O. Box 20005, Tampa, FL 33622-0005, (813) 889-9684

Source of subsequent assessment information:

References or reports: (Crewz & Lewis, 1991; Henningsen, 1993)

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh: 1986 and 1993

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Marsh is doing well (filling in).
- Marsh name: Gypsum Stack.

Marsh Inventory No.: 365 **Information Rank (1-5):** 5 **State:** FL

Date Planted: 1983?

Area Planted (1acre = 0.4047 ha): 2.800 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit:

Location of the marsh.

State: FL **County:** Hillsborough **Latitude (N):** 27 49' 35" **Longitude (W):** 82 29' 05"

Bay and sub-bays: Tampa Bay; Hillsborough Bay.

Directions to the site: At the end of Interbay Peninsula, on the MacDill AFB, a little W of the SE corner.

Source of planting information: Dr. Robin Lewis, III., Lewis Environmental Services, Inc., P.O. Box 20005, Tampa, FL 33622-0005, (813) 889-9684

Source of subsequent assessment information: (Roberts, 1991)

References or reports: (Roberts, 1991)

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh: 60%, 1987

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Yes. Fisheries species, birds, mammals, and epibenthic snails.

General comments about the marsh:

- Few birds; a rat; raccoon tracks; many killifish;
- Marsh name: MacDill AFB, No.1

Marsh Inventory No.: 266 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 1979

Area Planted (1acre = 0.4047 ha): 1.700 hectares

COE Permit No.: None **Date of COE Permit:**

Originator or Applicant for permit: Sunken Island

Location of the marsh.

State: FL **County:** Hillsborough **Latitude (N):** 27 48' 25" **Longitude (W):** 82 26' 01"

Bay and sub-bays: Tampa Bay; Hillsborough Bay; Alafia River Mouth.

Directions to the site: Northwest end of Sunken Island

Source of planting information: Dr. Robin Lewis, III., Lewis Environmental Services, Inc., P.O. Box 20005, Tampa, FL 33622-0005, (813) 889-9684

Source of subsequent assessment information: (Crewz & Lewis, 1991)

References or reports: (Crewz & Lewis, 1991)

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh: 26%, 1986

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? no

General comments about the marsh:

- *Spartina alterniflora* was growing well at low elevations; black and white mangrove had invaded as expected, and was doing very well amongst the *S. alterniflora* at higher elevations. Successful.

- *S. alterniflora* colonizes an area but then loses dominance as mangroves invade and displace the cordgrass.

Marsh Inventory No.: 670 Information Rank (1-5): 3 State: FL

Date Planted: 1978

Area Planted (1acre = 0.4047 ha): 0.405 hectares

COE Permit No.: Date of COE Permit:

Originator or Applicant for permit:

Location of the marsh.

State: FL County: Hillsborough Latitude (N): 27 51' 49" Longitude (W): 82 23' 40"

Bay and sub-bays: Tampa Bay; Hillsborough Bay; Archie Creek

Directions to the site: Archie Creek between 45 and 78 th St.

Source of planting information: Dr. Robin Lewis, III., Lewis Environmental Services, Inc., P.O. Box 20005, Tampa, FL 33622-0005, (813) 889-9684

Source of subsequent assessment information:

References or reports: (Crewz & Lewis, 1991; Henningsen, 1993)

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh: 1986; 1993

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Marsh name: One-acre Marsh

Marsh Inventory No.: 369 **Information Rank (1-5):** 3 **State:** FL

Date Planted: 1988

Area Planted (1acre = 0.4047 ha): 0.405 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit:

Location of the marsh.

State: FL **County:** Hillsborough **Latitude (N):** 27 51' 49" **Longitude (W):** 82 23' 40"

Bay and sub-bays: Tampa Bay; Hillsborough Bay; Archie Creek

Directions to the site: Archie Creek between 45 and 78 th St.

Source of planting information: Dr. Robin Lewis, III., Lewis Environmental Services, Inc., P.O. Box 20005, Tampa, FL 33622-0005, (813) 889-9684

Source of subsequent assessment information:

References or reports: (Crewz & Lewis, 1991; Henningsen, 1993)

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh: 1986; 1993

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Marsh name: One-acre Marsh

Marsh Inventory No.: 262 **Information Rank (1-5):** 5 **State:** FL

Date Planted: 1978

Area Planted (1acre = 0.4047 ha): 2.500 hectares

COE Permit No.: 76-074 **Date of COE Permit:** 1976

Originator or Applicant for permit: The Gardinier, Inc.

Location of the marsh.

State: FL **County:** Hillsborough **Latitude (N):** 27 51' 49" **Longitude (W):** 82 23' 40"

Bay and sub-bays: Tampa Bay; Hillsborough Bay; Archie Creek.

Directions to the site: East Tampa, at The Gardinier, Inc. property on Archie Creek off the Alafia River

Source of planting information: Crewz and Lewis, and Dr. Tom Roberts, Tennessee Tech University, Cookeville, TN 38505

Source of subsequent assessment information: (Roberts, 1991)

References or reports: (Crewz & Lewis, 1991; Roberts, 1991)

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh: 84%, 1986 (Crewz & Lewis); 72%, 1987 (Roberts)

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Yes, (Roberts, 1991). Fisheries species, birds, mammals, and epibenthic snails.

General comments about the marsh:

- Roberts found many birds, several rats and mice, and many killifish. Crewz and Lewis also surveyed this marsh and deemed it successful, however, fine silt drainage from this area appeared to be negatively impacting a natural marsh downstream.

Marsh Inventory No.: 366 **Information Rank (1-5):** 5 **State:** FL

Date Planted: 1992

Area Planted (1acre = 0.4047 ha): 1.214 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: City of Tampa

Location of the marsh.

State: FL **County:** Hillsborough **Latitude (N):** 27 57' 05" **Longitude (W):** 82 24' 50"

Bay and sub-bays: Tampa Bay; Hillsborough Bay; McKay Bay

Directions to the site: Northernmost branch of McKay Bay, south of Rd. 618.

Source of planting information: Dr. Robin Lewis, III., Lewis Environmental Services, Inc., P.O. Box 20005, Tampa, FL 33622-0005, (813) 889-9684

Source of subsequent assessment information: Dr. Brandt Henningsen, SWIM, (813) 985-7481 ext. 2202

References or reports: (Crewz & Lewis, 1991)

Other species planted in addition to *Spartina alterniflora* : *Spartina patens* and *Paspalum distichum*.

Percent vegetative cover attained by the planted marsh: 1986 and 1993.

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Yes. Fisheries studies with monthly sampling since Mar'92

General comments about the marsh:

- Marsh doing well, and being used by birds, reptiles and fish.
- Marsh name: McKay Bay

Marsh Inventory No.: 255 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 1984

Area Planted (1acre = 0.4047 ha): 0.142 hectares

COE Permit No.: 84W-0514 **Date of COE Permit:** 07/03/84

Originator or Applicant for permit: Wilson Company

Location of the marsh.

State: FL **County:** Hillsborough **Latitude (N):** 27 57' 52" **Longitude (W):** 82 33' 08"

Bay and sub-bays: Tampa Bay; Old Tampa Bay

Directions to the site: Fish Creek off 589-Bayport C

Source of planting information: Dr. Robin Lewis, III., Lewis Environmental Services, Inc., P.O. Box 20005, Tampa, FL 33622-0005, (813) 889-9684

Source of subsequent assessment information: (Crewz & Lewis, 1991)

References or reports: (Crewz & Lewis, 1991)

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh: 2%, 1986

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? no

General comments about the marsh:

- Planted *Spartina alterniflora* was dying; substrate was too low, and there was too much wave impact. Failure.

Marsh Inventory No.: 254 Information Rank (1-5): 4 State: FL

Date Planted: 1984

Area Planted (1acre = 0.4047 ha): 0.405 hectares

COE Permit No.: 84W-0514 Date of COE Permit: 07/03/84

Originator or Applicant for permit: Wilson Company

Location of the marsh.

State: FL County: Hillsborough Latitude (N): 27 57' 52" Longitude (W): 82 33' 08"

Bay and sub-bays: Tampa Bay; Old Tampa Bay

Directions to the site: Fish Creek off 589-Bayport B

Source of planting information: Dr. Robin Lewis, III., Lewis Environmental Services, Inc., P.O. Box 20005, Tampa, FL 33622-0005, (813) 889-9684

Source of subsequent assessment information: (Crewz & Lewis, 1991)

References or reports: (Crewz & Lewis, 1991)

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh: 45%, 1986

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? no

General comments about the marsh:

- *Spartina alterniflora* was growing well, good topography, but there was no room for expansion with sealevel rize. Successful.

Marsh Inventory No.: 256 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 1985

Area Planted (1acre = 0.4047 ha): 0.202 hectares

COE Permit No.: 84W-0514 **Date of COE Permit:** 07/03/84

Originator or Applicant for permit: Wilson Company

Location of the marsh.

State: FL **County:** Hillsborough **Latitude (N):** 27 58' 52" **Longitude (W):** 82 33' 08"

Bay and sub-bays: Tampa Bay; Old Tampa Bay

Directions to the site: Fish Creek off 589-Bayport D

Source of planting information: Dr. Robin Lewis, III., Lewis Environmental Services, Inc., P.O. Box 20005, Tampa, FL 33622-0005, (813) 889-9684

Source of subsequent assessment information: (Crewz & Lewis, 1991)

References or reports: (Crewz & Lewis, 1991)

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh: 1%, 1986

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? no

General comments about the marsh:

- *Spartina alterniflora* was growing poorly; site's substrate was too low and had poor tidal flushing. Failure.

Marsh Inventory No.: 53 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 1985

Area Planted (1acre = 0.4047 ha): 0.202 hectares

COE Permit No.: 84W-0514 **Date of COE Permit:** 07/03/84

Originator or Applicant for permit: Wilson Company

Location of the marsh.

State: FL **County:** Hillsborough **Latitude (N):** 27 57' 52" **Longitude (W):** 82 33' 08"

Bay and sub-bays: Tampa Bay; Old Tampa Bay

Directions to the site: Fish Creek off 589-Bayport A

Source of planting information: Dr. Robin Lewis, III., Lewis Environmental Services, Inc., P.O. Box 20005, Tampa, FL 33622-0005, (813) 889-9684

Source of subsequent assessment information: (Crewz & Lewis, 1991)

References or reports: (Crewz & Lewis, 1991)

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh: 16%, 1986

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? no

General comments about the marsh:

- *Spartina alterniflora* was growing, but humans were trampling it. Mixed success.

-

Marsh Inventory No.: 372 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 1983

Area Planted (1acre = 0.4047 ha): 0.100 hectares

COE Permit No.: 82(?) -0048 **Date of COE Permit:** 06/04/05

Originator or Applicant for permit: Gandy Bridge Boat Ramp

Location of the marsh.

State: FL **County:** Hillsborough **Latitude (N):** 27 53' 25" **Longitude (W):** 82 32' 05"

Bay and sub-bays: Tampa Bay; Old Tampa Bay.

Directions to the site: Under the east end of Gandy Bridge off the Interbay Peninsula, Tampa.

Source of planting information: Dr. Robin Lewis, III., Lewis Environmental Services, Inc., P.O. Box 20005, Tampa, FL 33622-0005, (813) 889-9684

Source of subsequent assessment information: (Crewz & Lewis, 1991; Roberts, 1991)

References or reports: (Crewz & Lewis, 1991; Roberts, 1991)

Other species planted in addition to *Spartina alterniflora* : *Rhizophora mangle*

Percent vegetative cover attained by the planted marsh: 25% 1986 (Crewz & Lewis); 50% 1987 (Roberts)

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Yes. Fisheries species, birds, mammals, and epibenthic snails.

General comments about the marsh:

- This area had too low of elevations in its interior for cordgrass establishment, and was deemed a failure by Crewz and Lewis (1991) in their survey made in the spring of 1986.

- No birds, a rat, signs of rabbits, and many killifish were found by Roberts (1991).

Marsh Inventory No.: 253 Information Rank (1-5): 4 State: FL

Date Planted: 1985

Area Planted (1acre = 0.4047 ha): 0.202 hectares

COE Permit No.: None Date of COE Permit:

Originator or Applicant for permit: Seagrove

Location of the marsh.

State: FL County: Indian River Latitude (N): 27 37' 10" Longitude (W): 80 21' 20"

Bay and sub-bays: Indian River

Directions to the site: Indian River Lagoon

Source of planting information: Dr. Robin Lewis, III., Lewis Environmental Services, Inc., P.O. Box 20005, Tampa, FL 33622-0005, (813) 889-9684

Source of subsequent assessment information: (Crewz & Lewis, 1991)

References or reports: (Crewz & Lewis, 1991)

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh: 37%, 1986

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? no

General comments about the marsh:

- *Spartina alterniflora* and mangrove well established; good flushing and topography. Successful.

Marsh Inventory No.: 752 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 07/01/85

Area Planted (1acre = 0.4047 ha): 0.640 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Andreas Gotzfried (305) 867-2000 is the contact person

Location of the marsh.

State: FL **County:** Indian River **Latitude (N):** 27 34' 00?" **Longitude (W):** 80 19' 50"

Bay and sub-bays: Indian River

Directions to the site: Vero Beach; Hwy. A-1-A 5 miles S of State Rd 60; a tidal creek-loop system connected through a fringing mangrove swamp to the Indian River.

Source of planting information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

Source of subsequent assessment information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

References or reports:

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh: 100% March-93

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Not assessed.

General comments about the marsh:

- *Spartina alterniflora* clumps were planted on 3-ft centers. Thick band of *Spartina* along edge. *Spartina* inside has largely (90%) been replaced by mangroves. The area also has *Batis*, *Salicornia*, *Sesuvium*, and *Sueda*.

- Marsh name: Sea Grove

Marsh Inventory No.: 753 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 04/01/86

Area Planted (1acre = 0.4047 ha): 0.250 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Henry Muller or Steve Moler (407) 231-4223 are the contact people.

Location of the marsh.

State: FL **County:** Indian River **Latitude (N):** 27 51' 05" **Longitude (W):** 80 29' 30"

Bay and sub-bays: Indian River

Directions to the site: Sebastian; US 1 at S end of city limits; created on a peninsula between harbor and Indian River.

Source of planting information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

Source of subsequent assessment information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

References or reports:

Other species planted in addition to *Spartina alterniflora* : *Paspalum vaginatum*

Percent vegetative cover attained by the planted marsh: 100% July-93

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Not assessed.

General comments about the marsh:

- *Spartina alterniflora* clumps were planted on 3-ft centers. Saltmarsh grasses were planted as an initial stabilizer community and as a nursery for mangrove establishment. *Spartina* has been mostly replaced by mangroves in the interior.
- Marsh name: River Run Condo

Marsh Inventory No.: 778 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 10/01/81

Area Planted (1acre = 0.4047 ha): 1.404 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Bob Lynds is the contact person.

Location of the marsh.

State: FL **County:** Indian River **Latitude (N):** 28 04' 40" **Longitude (W):** 80 36' 10"

Bay and sub-bays: Indian River

Directions to the site: Melbourne; Hwy 192 at the Indian River; fringing shoreline with coquina rock revetment.

Source of planting information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

Source of subsequent assessment information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

References or reports:

Other species planted in addition to *Spartina alterniflora* : *Spartina patens* and *Distichlis spicata* were planted on 2-ft centers.

Percent vegetative cover attained by the planted marsh: 95-100% cover as of Sept-93

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Not assessed.

General comments about the marsh:

- Coquina boulder break water was installed at the toe zone to prevent erosion. Mangroves are beginning to invade the marsh, especially in a 1 acre marsh on the N side of the point (about 50% mangrove as of Sept-93).

- Marsh name: Melbourne Harbor *S. alterniflora* clumps were planted on 2-ft centers.

Marsh Inventory No.: 779 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 03/01/90

Area Planted (1acre = 0.4047 ha): 0.210 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: George Fleming (407) 783-6153 is the contact person.

Location of the marsh.

State: FL **County:** Indian River **Latitude (N):** 27 36' 00?" **Longitude (W):** 80 20' 00?"

Bay and sub-bays: Indian River

Directions to the site: Vero Beach; A-1-A 5 miles S of State Rd. 60; fringing marsh along shoreline.

Source of planting information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

Source of subsequent assessment information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

References or reports:

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh: 80-90% cover as of Jun-93

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Not assessed.

General comments about the marsh:

- *Spartina alterniflora* clumps were planted on 2-ft centers. The *Spartina* marsh is being invaded by mangroves. The marsh was established for erosion control and intertidal habitat.

- Marsh name: Sand Pointe

Marsh Inventory No.: 765 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 07/01/88

Area Planted (1acre = 0.4047 ha): 28.410 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Ron Andrews (407) 778-0503 is the contact person.

Location of the marsh.

State: FL **County:** Indian River **Latitude (N):** 27 37' 30?" **Longitude (W):** 80 22' 45"

Bay and sub-bays: Indian River

Directions to the site: Vero Beach; U.S.1 N of Vero Beach and between 45th and 51st streets; rehabilitated tidal marsh in an old mosquito impoundment.

Source of planting information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

Source of subsequent assessment information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

References or reports:

Other species planted in addition to *Spartina alterniflora* : *Spartina bakeri*, *Iris virginica*, *Pontederia cordata*, *Sagittaria lancifolia*, *Distichlis spicata*, *Sporobolus virginicus*, *Spartina patens*, *Sesuvium portulacastrum*; *Juncus roemerianus*, *Batis maritima*, *Salicornia virginica* were planted on 1.5-ft centers.

Percent vegetative cover attained by the planted marsh: 95-100% cover by marsh plants, as of Nov-93

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Not assessed.

General comments about the marsh:

- The estuarine and freshwater marshes have reached nearly 100% ground cover throughout the project. Additional species have invaded from adjacent areas. Mangrove invasion in some areas.

- Marsh name: Grand Harbor. *S. alterniflora* clumps were planted on 1.5-ft centers.

Marsh Inventory No.: 766 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 03/01/89

Area Planted (1acre = 0.4047 ha): 2.975 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Reese Kessler (407) 699-4222 is the contact person.

Location of the marsh.

State: FL **County:** Indian River **Latitude (N):** 27 34' 00?" **Longitude (W):** 80 21' 40"

Bay and sub-bays: Indian River

Directions to the site: Vero Beach; U.S.1 S of Vero Beach, 1 mile S of Oslo Rd.; enclosed created tidal wetland, tidal flats.

Source of planting information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

Source of subsequent assessment information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

References or reports:

Other species planted in addition to *Spartina alterniflora* : *Avicennia germinans*, *Scirpus validus*, *Sagittaria lancifolia*, *Pontederia cordata*; *Batis maritima* and *Salicornia virginica* were planted on 2-ft centers.

Percent vegetative cover attained by the planted marsh: 100% by *S. alterniflora*; 100% cover by FW planting.

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Not assessed.

General comments about the marsh:

- *Spartina alterniflora* clumps were planted on 2-ft centers. Mangroves are beginning to dominate some areas of the *Spartina* marsh. Freshwater plantings have covered their areas and are spreading further.

- Marsh name: Grove Isle.

Marsh Inventory No.: 618 **Information Rank (1-5):** 5 **State:** FL

Date Planted:

Area Planted (1acre = 0.4047 ha): 0.450 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Sea Grove West

Location of the marsh.

State: FL **County:** Indian River **Latitude (N):** 27 38' 00" ? **Longitude (W):** 80 22' 30" ?

Bay and sub-bays: Indian River;

Directions to the site: Sea Grove West; Vero Beach

Source of planting information: Dr. Tom Roberts, Tennessee Tech University, Cookeville, TN 38505

Source of subsequent assessment information: (Roberts, 1991)

References or reports: (Roberts, 1991)

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh: 40%, 1987

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Yes. Fisheries species, birds, mammals, and epibenthic snails.

General comments about the marsh:

- Modest marsh establishment. No birds; a couple of rats; raccoon tracks; few killifish.

-

Marsh Inventory No.: 261 **Information Rank (1-5):** 5 **State:** FL

Date Planted: 05/01/76

Area Planted (1acre = 0.4047 ha): 0.202 hectares

COE Permit No.: Yes, but not known **Date of COE Permit:** 1976

Originator or Applicant for permit: Private land developer

Location of the marsh.

State: FL **County:** Indian River **Latitude (N):** 27 40' 57" **Longitude (W):** 80 22' 12"

Bay and sub-bays: Indian River; McCullers Cove

Directions to the site: NE corner of McCullers Cove, about 5 km N of Riomar which is E of Vero Beach. At the SW tip of the residential development bulkhead.

Source of planting information: A. Banner, 1977

Source of subsequent assessment information: (Banner, 1977)

References or reports: (Banner, 1977)

Other species planted in addition to *Spartina alterniflora* : *Rhizophora mangle*, *Laguncularia racemosa*, *Avicennia nitida*

Percent vegetative cover attained by the planted marsh: 90%, 1986

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? A general sampling of fish, shellfish, and benthos was made.

General comments about the marsh:

- *Spartina alterniflora* and mangrove well established; good flushing and topography. Successful.
- Many organisms typical of the estuary were found in the restored area, fewer benthic organisms were found in the thick *Spartina* marsh than away from it.

Marsh Inventory No.: 756 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 04/01/82

Area Planted (1acre = 0.4047 ha): 39.050 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Bob Snyder (407) 746-7290 is the contact person.

Location of the marsh.

State: FL **County:** Martin **Latitude (N):** 26 55' 00?" **Longitude (W):** 80 04' 20"

Bay and sub-bays: Loxahatchee River

Directions to the site: Jupiter; Alternate A-1-A, 1 mile S of S.R. 706; interconnecting system of canals and tidal marsh shelves, connected by 4 inlets.

Source of planting information: Mr. Steve Beeman, Ecoshores, Inc. Port Orange, FL (904) 767-6232. *S. alterniflora* clumps were planted 2'on center (=o.c.)

Source of subsequent assessment information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

References or reports:

Other species planted in addition to *Spartina alterniflora* : *Batis maritima*, *Sporobolus virginicus*, *Distichlis spicata*, *Juncus roemerianus*, *Avicennia germinans* and *Rhizophora mangle*. The grasses were palnted on 2-ft centers.

Percent vegetative cover attained by the planted marsh: 100% cover in 90 acres by mangroves or saltmarsh species.

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Not assessed.

General comments about the marsh:

- Most of the *Spartina* has been replaced (displaced) by volunteer mangroves. Cordgrass fring dominates the waterward edges.
- Marsh name: Admirals Cove

Marsh Inventory No.: 257 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 1981

Area Planted (1acre = 0.4047 ha): 3.500 hectares

COE Permit No.: 80M-0276 **Date of COE Permit:** 1980

Originator or Applicant for permit: Florida Keys Aqueduct Authority

Location of the marsh.

State: FL **County:** Monroe **Latitude (N):** 25 06' 39" **Longitude (W):** 80 24' 52"

Bay and sub-bays: Key Largo

Directions to the site: Parallel to Highway A1A for several miles on Key Largo.

Source of planting information: Dr. Robin Lewis, III., Lewis Environmental Services, Inc., P.O. Box 20005, Tampa, FL 33622-0005, (813) 889-9684

Source of subsequent assessment information: (Crewz & Lewis, 1991)

References or reports: (Crewz & Lewis, 1991); A. Mager

Other species planted in addition to *Spartina alterniflora* : *Avicennia germinans*

Percent vegetative cover attained by the planted marsh: 12% - cordgrass; 8% - mangroves, 1986

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? no

General comments about the marsh:

- *Spartina alterniflora* was growing poorly; site's substrate was too low and had poor tidal flushing. The mangroves' roots could not penetrate very far in the rocky substrate. Failure.

80

Marsh Inventory No.: 764 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 12/01/79

Area Planted (1acre = 0.4047 ha): 5.463 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Don Shakelford (407) 627-5110 is the contact person.

Location of the marsh.

State: FL **County:** Palm Beach **Latitude (N):** 26 53' 30" **Longitude (W):** 80 05' 00?"

Bay and sub-bays: ICW

Directions to the site: Palm Beach Gardens; Donald Ross Rd., W of ICW; loop system of created tidal canals with narrow tidal beaches.

Source of planting information: Mr. Steve Beeman, Ecoshores, Inc. Port Orange, FL (904) 767-6232. *S. alterniflora* clumps were planted 2'on center (=o.c.)

Source of subsequent assessment information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

References or reports:

Other species planted in addition to *Spartina alterniflora* : *Spartina patens*, *Sporobolus virginicus*, *Distichlis spicata*, and *Paspalum vaginatum* were planted on 2-ft centers; *Avicennia germinans*, *Rhizophora mangle*, and *Laguncularia racemosa*.

Percent vegetative cover attained by the planted marsh: 80-100% *Spartina* cover; thinner areas of mangrove cover, as of May-93

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Not assessed.

General comments about the marsh:

- Planted over a long time: Dec-79 thru May-90. Most of the intertidal shelves are successfully vegetated, although some areas are sparse; heavy invasion by mangrove volunteers.

- Marsh name: Frenchman's Creek.

Marsh Inventory No.: 755 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 11/01/82

Area Planted (1acre = 0.4047 ha): 0.500 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Don Shackelford (305) 627-5110 is the contact person.

Location of the marsh.

State: FL **County:** Palm Beach **Latitude (N):** 26 53' 00?" **Longitude (W):** 80 05' 05?"

Bay and sub-bays: ICW

Directions to the site: Palm Beach Gardens; tidal canal connencted to the Intracoastal via culverts; just N of Donald Ross Rd. on Cypress Creek RD; fringing marsh along steep bank.

Source of planting information: Mr. Steve Beeman, Ecoshores, Inc. Port Orange, FL (904) 767-6232. *S. alterniflora* clumps were planted 2'on center (=o.c.)

Source of subsequent assessment information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

References or reports:

Other species planted in addition to *Spartina alterniflora* : *Juncus roemerianus*, *Avicennia germinans*, *Laguncularia racemosa* and *Rhizophora mangle* were planted on 2-ft centers. *Rhizophora* was in 1-gal containers.

Percent vegetative cover attained by the planted marsh: 100% cover, but is all mangrove. As of Jan-92

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Not assessed.

General comments about the marsh:

- Most of the grasses and rushes have been displaced by the mangroves. The mangroves have formed a 10-15' tall dense hedge.
- Marsh name: Cypress Creek

Marsh Inventory No.: 754 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 06/01/83

Area Planted (1acre = 0.4047 ha): 0.100 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Robert Snyder (305) 746-7290 is the contact person.

Location of the marsh.

State: FL **County:** Palm Beach **Latitude (N):** 26 56' 00?" **Longitude (W):** 80 04' 20?"

Bay and sub-bays: ICW

Directions to the site: Jupiter; S side of Jupiter Inlet on a high energy shoreline.

Source of planting information: Mr. Steve Beeman, Ecoshores, Inc. Port Orange, FL (904) 767-6232. *S. alterniflora* clumps were planted 1.5'on center (=o.c.)

Source of subsequent assessment information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

References or reports:

Other species planted in addition to *Spartina alterniflora* : *Rhizophora mangle* was planted on 3-ft centers.

Percent vegetative cover attained by the planted marsh: 70% cover by *Spartina*; mangroves about 100% cover as of Aug-93

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Not assessed.

General comments about the marsh:

- The mangroves and *Spartina* have grown and spread, holding erosion in check and covering the shoreline. There was partial protection of the planting site by a rock revetment.

- Marsh name: Harpoon Louie's

Marsh Inventory No.: 357 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 1990

Area Planted (1acre = 0.4047 ha): 0.003 hectares

COE Permit No.: **Date of COE Permit:** 1990

Originator or Applicant for permit: Lakeside Palms Development

Location of the marsh.

State: FL **County:** Palm Beach **Latitude (N):** **Longitude (W):**

Bay and sub-bays: Lake Worth

Directions to the site: I 95 to 6 th Ave South. East to South Palm Way or to lakeside. Go South. Lakeside Palms is on the east 8-10 blocks away.

Source of planting information: Caroline Hanes, Florida Dept. of Environmental Regulation, SE District Office, 1900 S. Congress Ave., Suite A, West Palm Beach, FL 33406, (407) 433-2650.

Source of subsequent assessment information: Caroline Hanes, Florida Dept. of Environmental Regulation, SE District Office, 1900 S. Congress Ave., Suite A, West Palm Beach, FL 33406, (407) 433-2650.

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- *Spartina* well setablished on 5/92. Status reports availabel. Mitigation is considered successful as of 5/92. File # 501750906. 2 Sites, 150 sq. ft. and 225 sq. ft. planted.

Marsh Inventory No.: 356 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 1991

Area Planted (1acre = 0.4047 ha): 0.004 hectares

COE Permit No.: **Date of COE Permit:** 1989

Originator or Applicant for permit: Russell Wilson

Location of the marsh.

State: FL **County:** Palm Beach **Latitude (N):** **Longitude (W):**

Bay and sub-bays: Loxatchee River Cove

Directions to the site: South of Jupiter inlet, east of Loxatchee River. Take I 95 to Indian Town Road. Go East to A1A, north to Saturn St., east on Saturn to end of the road. Located Between some townhouses and a house at the north end of Saturn St.

Source of planting information: Caroline Hanes, Florida Dept. of Environmental Regulation, SE District Office, 1900 S. Congress Ave., Suite A, West Palm Beach, FL 33406, (407) 433-2650.

Source of subsequent assessment information: Caroline Hanes, Florida Dept. of Environmental Regulation, SE District Office, 1900 S. Congress Ave., Suite A, West Palm Beach, FL 33406, (407) 433-2650.

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh: 75%

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? no

General comments about the marsh:

- As of 10/91 *Spartina* had grown to 4'. No status reports have been submitted, but the site has been monitored. File # 501426206. 400 sq. ft. planted.

Marsh Inventory No.: 379 **Information Rank (1-5):** 3 **State:** FL

Date Planted: 1987

Area Planted (1acre = 0.4047 ha): 0.004? hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit:

Location of the marsh.

State: FL **County:** Pinellas **Latitude (N):** 27 42' 00" **Longitude (W):** 82 39' 00"

Bay and sub-bays: Tampa Bay; Main Channel

Directions to the site: South of Pinellas Point Drive, between Maxima Point and Point Pinellas.

Source of planting information: Dr. Robin Lewis, III., Lewis Environmental Services, Inc., P.O. Box 20005, Tampa, FL 33622-0005, (813) 889-9684

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Marsh name: Pinellas Point.

Marsh Inventory No.: 377 **Information Rank (1-5):** 3 **State:** FL

Date Planted: 1986

Area Planted (1acre = 0.4047 ha): hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit:

Location of the marsh.

State: FL **County:** Pinellas **Latitude (N):** 27 57' 30" **Longitude (W):** 82 42' 40"

Bay and sub-bays: Tampa Bay; Old Tampa Bay

Directions to the site: Bayview, south of the Courtney Campbell Parkway

Source of planting information:

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Marsh name: Bayview.

Marsh Inventory No.: 259 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 1984

Area Planted (1acre = 0.4047 ha): 2.500 hectares

COE Permit No.: 83T-0476 **Date of COE Permit:** 1983

Originator or Applicant for permit: Feather Sound

Location of the marsh.

State: FL **County:** Pinellas **Latitude (N):** 27 54' 15" **Longitude (W):** 82 39' 35"

Bay and sub-bays: Tampa Bay; Old Tampa Bay

Directions to the site: Feather Sound Site B; On the west side of Big Island; St. Petersburg

Source of planting information: Dr. Robin Lewis, III., Lewis Environmental Services, Inc., P.O. Box 20005, Tampa, FL 33622-0005, (813) 889-9684

Source of subsequent assessment information: (Crewz & Lewis, 1991)

References or reports: (Crewz & Lewis, 1991)

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh: 11%, 1986

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? no

General comments about the marsh:

- *Spartina alterniflora* was growing at the higher elevation; drainage creek blocked flushing, drowning the lower *S. alterniflora* was replanted after the Crewz & Lewis survey. Mixed failure.

Marsh Inventory No.: 249 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 1984

Area Planted (1acre = 0.4047 ha): 0.400 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit:

Location of the marsh.

State: FL **County:** Pinellas **Latitude (N):** 27 54' 20" **Longitude (W):** 82 39' 45"

Bay and sub-bays: Tampa Bay; Old Tampa Bay

Directions to the site: Feather Cove;

Source of planting information: Dr. Robin Lewis, III., Lewis Environmental Services, Inc., P.O. Box 20005, Tampa, FL 33622-0005, (813) 889-9684

Source of subsequent assessment information: Robin Lewis, (813) 889-9684

References or reports: (Crewz & Lewis, 1991)

Other species planted in addition to *Spartina alterniflora* : *Rhizophora mangle*

Percent vegetative cover attained by the planted marsh: 13% - cordgrass, 2% - mangroves; 1986

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? no

General comments about the marsh:

- Too much freshwater and nutrients in the area allowed souther cat-tail to become established and to crowd out the cordgrass and stunt the red mangroves. Failure.

Marsh Inventory No.: 258 **Information Rank (1-5):** 5 **State:** FL

Date Planted: 1983

Area Planted (1acre = 0.4047 ha): 0.6-Lewis; 0.53-Roberts hectares

COE Permit No.: 83T-0476 **Date of COE Permit:** 1983

Originator or Applicant for permit: Feather Sound

Location of the marsh.

State: FL **County:** Pinellas **Latitude (N):** 27 54' 15" **Longitude (W):** 82 39' 35"

Bay and sub-bays: Tampa Bay; Old Tampa Bay;

Directions to the site: Feather Sound Site A; On the west side of Big Island; St. Petersburg

Source of planting information: Dr. Robin Lewis, III., Lewis Environmental Services, Inc., P.O. Box 20005, Tampa, FL 33622-0005, (813) 889-9684

Source of subsequent assessment information: Robin Lewis, (813) 889-9684 ; (Roberts, 1991)

References or reports: Crewz and Lewis, 1991; (Roberts, 1991)

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh: 87% (Crews & Lewis); 55% (Roberts); 1986.

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Yes, by Roberts. Birds, mammals, and epibenthic snails.

General comments about the marsh:

- Few birds; many rats; raccoon tracks. Crewz and Lewis found the marsh and mangroves to be growing well enough to consider this planting a success. Marsh may convert to black mangrove forest later.

Marsh Inventory No.: 265 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 1983

Area Planted (1acre = 0.4047 ha): 0.040 hectares

COE Permit No.: 82C-1099 **Date of COE Permit:** 1982

Originator or Applicant for permit: Las Fontanas

Location of the marsh.

State: FL **County:** Pinellas **Latitude (N):** 27 54' 50" **Longitude (W):** 82 42' 00"

Bay and sub-bays: Tampa Bay; Old Tampa Bay; Cross Bayou Canal.

Directions to the site: East of 686 where the road turns south at High Point

Source of planting information: Dr. Robin Lewis, III., Lewis Environmental Services, Inc., P.O. Box 20005, Tampa, FL 33622-0005, (813) 889-9684

Source of subsequent assessment information: (Crewz & Lewis, 1991)

References or reports: (Crewz & Lewis, 1991)

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh: 91%, 1986

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? no

General comments about the marsh:

- *Spartina alterniflora* was growing where Brazilian pepper had been removed; some colonization by red mangrove had started. Successful.

Marsh Inventory No.: 260 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 1984

Area Planted (1 acre = 0.4047 ha): 0.202 hectares

COE Permit No.: 82D-0339 **Date of COE Permit:** 1984

Originator or Applicant for permit: FM 92 Radio Tower

Location of the marsh.

State: FL **County:** Pinellas **Latitude (N):** 28 00' 54" **Longitude (W):** 82 41' 04"

Bay and sub-bays: Tampa Bay; Old Tampa Bay; Safety Harbor

Directions to the site: Radio tower site east of 590, north of Phillippe Point.

Source of planting information: Dr. Robin Lewis, III., Lewis Environmental Services, Inc., P.O. Box 20005, Tampa, FL 33622-0005, (813) 889-9684

Source of subsequent assessment information: Robin Lewis, (813) 889-9684

References or reports: (Crewz & Lewis, 1991)

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh: 34%, 1986

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? no

General comments about the marsh:

- *Spartina alterniflora* was growing at lower and higher elevation areas; black mangrove was starting to colonize the higher area. Mixed success.

Marsh Inventory No.: 378 **Information Rank (1-5):** 5 **State:** FL

Date Planted: 1992

Area Planted (1acre = 0.4047 ha): 4.047 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: SWIM and City of St. Petersburg

Location of the marsh.

State: FL **County:** Pinellas **Latitude (N):** 27 49' 10" **Longitude (W):** 82 37' 07"

Bay and sub-bays: Tampa Bay; Placido Bayou

Directions to the site: W of Riveria Bay, N of the Mangrove Bay golf course, S of Gandy Blvd., and adjacent and S of the E end of 78th Ave NE

Source of planting information: Dr. Brandt Henningsen, SW Florida Water Management District, Surface Water Improvement and Management Program, 7601 Highway 301 N, Tampa, FL 33637, (813) 985-7481 ext. 2202

Source of subsequent assessment information: Dr. Brandt Henningsen, SWIM, (813) 985-7481 ext. 2202

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Yes. Fish, invertebrates and birds.

General comments about the marsh:

- This was a successful restoration and creation project. Many fisheries organisms have been caught in monitoring sampling. Bird sitings confirm bird utilization of the marsh.

- Marsh name: Mangrove Bay.

Marsh Inventory No.: 619 **Information Rank (1-5):** 5 **State:** FL

Date Planted: 1986

Area Planted (1acre = 0.4047 ha): 0.610 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Thunder Bay

Location of the marsh.

State: FL **County:** Pinellas ? **Latitude (N):** **Longitude (W):**

Bay and sub-bays: Tampa Bay; Thunder Bay

Directions to the site: Thunder Bay; St. Petersburg

Source of planting information: Dr. Tom Roberts, Tennessee Tech University, Cookeville, TN 38505

Source of subsequent assessment information: (Roberts, 1991)

References or reports: (Roberts, 1991)

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh: 52%, 1987

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Yes. Fisheries species, birds, mammals, and epibenthic snails.

General comments about the marsh:

- Marsh was successful. Few birds; a rat; many killifish.

Marsh Inventory No.: 760 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 01/01/83

Area Planted (1acre = 0.4047 ha): 0.631 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Randall Gentry (904) 824-0888 is the contact person.

Location of the marsh.

State: FL **County:** St. Johns **Latitude (N):** 29 54' 00?" **Longitude (W):** 81 18' 30?"

Bay and sub-bays: St. Augustine Inlet

Directions to the site: St. Augustine; N side of St. Augustine Inlet; cove and fringing marsh.

Source of planting information: Mr. Steve Beeman, Ecoshores, Inc. Port Orange, FL (904) 767-6232. *S. alterniflora* clumps were planted 2' on center (=o.c.)

Source of subsequent assessment information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

References or reports:

Other species planted in addition to *Spartina alterniflora* : *Distichlis spicata* and *Juncus roemerianus* were planted on 2-ft centers.

Percent vegetative cover attained by the planted marsh: 100% cover by *Spartina* and high marsh. as of may-93

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Not assessed.

General comments about the marsh:

- Used a coquina rock revetment which subsided after awhile, but not before the marsh became established. Revetment placed at MLW broke the waves that were causing severe shoreline erosion.

- Marsh name: Porpoise Point

Marsh Inventory No.: 758 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 02/01/88

Area Planted (1acre = 0.4047 ha): 0.560 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: David Lavine (407) 452-2580 is the contact person.

Location of the marsh.

State: FL **County:** Volusia **Latitude (N):** 29 11' 35" **Longitude (W):** 81 00' 00?"

Bay and sub-bays: Halifax River

Directions to the site: Daytona Beach; Palmetto Ave N of Hwy 400; border of Daytona Beach and South Daytona; perimeter marsh along canal.

Source of planting information: Mr. Steve Beeman, Ecoshores, Inc. Port Orange, FL (904) 767-6232. *S. alterniflora* clumps were planted 2' on center (=o.c.)

Source of subsequent assessment information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

References or reports:

Other species planted in addition to *Spartina alterniflora* : *Distichlis spicata*, *Sporobolus virginicus*, *Sesuvium portulacastrum*, *Avicennia germinans*.

Percent vegetative cover attained by the planted marsh: 100% cover by *Spartina* and high marsh species, as of Oct-93.

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Not assessed.

General comments about the marsh:

- Mangroves died during subsequent freezes, but black mangrove seedlings are now re-establishing in the *Spartina*.
- Marsh name: Post Cove

Marsh Inventory No.: 759 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 04/01/88

Area Planted (1acre = 0.4047 ha): 8.721 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Steve Tonjes (904) 736-5445 is the contact person.

Location of the marsh.

State: FL **County:** Volusia **Latitude (N):** 29 00' 00" **Longitude (W):** 80 54' 00?"

Bay and sub-bays: Halifax River

Directions to the site: New Smyrna Beach; several sites in SE Volusia county; 7 sites mostly spoil islands that had been denuded by frost damage to mangrove forests.

Source of planting information: Mr. Steve Beeman, Ecoshores, Inc. Port Orange, FL (904) 767-6232. *S. alterniflora* clumps were planted 3'on center (=o.c.)

Source of subsequent assessment information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

References or reports:

Other species planted in addition to *Spartina alterniflora* : *Sporobolus virginicus* was planted on 3-ft centers; *Avicennia germinans* seedlings were planted on 10-ft centers.

Percent vegetative cover attained by the planted marsh: 85-100% cover achieved by all planted areas, as of Feb-1992

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Not assessed.

General comments about the marsh:

- Planting extended from April-Dec in 1988-90. Most of the mangroves died in freezes, but natural colonization has more than replaced them. Some *Spartina* areas are being displaced by mangroves volunteer seedlings.

- Marsh name: Lytle Ave. Bridge

Marsh Inventory No.: 777 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 06/01/78

Area Planted (1acre = 0.4047 ha): 0.061 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Mrs. Coleman (904) 252-7246 is the contact person.

Location of the marsh.

State: FL **County:** Volusia **Latitude (N):** 29 08' 30" **Longitude (W):** 80 58' 00"

Bay and sub-bays: Halifax River

Directions to the site: Daytona Beach; Peninsula Drive 4 miles S of State Rd. 400; fringing shoreline marsh.

Source of planting information: Mr. Steve Beeman, Ecoshores, Inc. Port Orange, FL (904) 767-6232. *S. alterniflora* clumps were planted 2'on center (=o.c.)

Source of subsequent assessment information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

References or reports:

Other species planted in addition to *Spartina alterniflora* : *Spartina patens* and *Paspalum vaginatum* were planted on 2-ft centers.

Percent vegetative cover attained by the planted marsh: 100% cover as of July-93

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Not assessed.

General comments about the marsh:

- *Spartina* has filled in completely and has spread 25' to the west. The marsh has protected the land from erosion.
- Marsh name: Henry Coleman

Marsh Inventory No.: 784 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 03/01/91

Area Planted (1acre = 0.4047 ha): 0.287 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Jeanne Tolley (904) 258-3117 is the contact person.

Location of the marsh.

State: FL **County:** Volusia **Latitude (N):** 29 13' 00?" **Longitude (W):** 81 01' 20"

Bay and sub-bays: Halifax River

Directions to the site: Daytona Beach; South Beach Street half mile S of U.S. Hwy 92; fringing intertidal shoreline along outer perimeter of boat harbor.

Source of planting information: Mr. Steve Beeman, Ecoshores, Inc. Port Orange, FL (904) 767-6232. *S. alterniflora* clumps were planted 2'on centers.

Source of subsequent assessment information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

References or reports:

Other species planted in addition to *Spartina alterniflora* : *Sporobolus virginicus* was planted on 2-ft centers; *Spartina bakeri*.

Percent vegetative cover attained by the planted marsh: 50-75% as of Oct-93

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Not assessed.

General comments about the marsh:

- The high marsh and *Spartina alterniflora* are growing, but coverage is only 50-75%. There is no breakwater and the substrate is rocky which make complete coverage by the marsh difficult. Some replanting of *S. alterniflora* was needed.

- *S. alterniflora* clumps were planted 2'on centers. Marsh name: Halifax Harbor

Marsh Inventory No.: 621 **Information Rank (1-5):** 5 **State:** FL

Date Planted: 1985

Area Planted (1acre = 0.4047 ha): 0.280 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit:

Location of the marsh.

State: FL **County:** Volusia **Latitude (N):** 29 08' 45" ? **Longitude (W):** 80 58' 55" ?

Bay and sub-bays: Halifax River

Directions to the site: Bouchelle Island; Port Orange

Source of planting information: Dr. Tom Roberts, Tennessee Tech University, Cookeville, TN 38505

Source of subsequent assessment information: (Roberts, 1991)

References or reports: (Roberts, 1991)

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh: 74%, 1987

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Yes. Fisheries species, birds, mammals, and epibenthic snails.

General comments about the marsh:

- Marsh was successful. No birds; a few rats; several typically estuarine fish .

Marsh Inventory No.: 769 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 11/01/86

Area Planted (1acre = 0.4047 ha): 0.251 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Paul Kosmas (904) 427-6892 is the contact person.

Location of the marsh.

State: FL **County:** Volusia **Latitude (N):** 29 01' 00" **Longitude (W):** 80 54' 00"

Bay and sub-bays: ICW (Indian River North)

Directions to the site: New Smyrna Beach; State Rd. 44 (North Causeway); fringing marsh along ICW (Indian River North) and high marsh in protected cove.

Source of planting information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

Source of subsequent assessment information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

References or reports:

Other species planted in addition to *Spartina alterniflora* : *Sporobolus virginicus* and *Distichlis spicata* were planted on 2-ft centers.

Percent vegetative cover attained by the planted marsh: 95-100% cover by *Spartina* and high marsh, as of April-93

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Not assessed.

General comments about the marsh:

- *Spartina alterniflora* clumps were planted on 2-ft centers. Dense saltmarsh established to replace freeze killed mangroves.
- Marsh name: Marina Bay.

Marsh Inventory No.: 768 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 03/01/83

Area Planted (1acre = 0.4047 ha): 0.328 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Alice Williamson (904) 756-0171 is the contact person.

Location of the marsh.

State: FL **County:** Volusia **Latitude (N):** 29 12' 00?" **Longitude (W):** 81 00' 00"

Bay and sub-bays: Indian River North

Directions to the site: Daytona Beach Shores; S. Peninsula Dr., 4 miles N of Dunlawton Ave.; created intertidal shoreline behind coquina rock revetment.

Source of planting information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

Source of subsequent assessment information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

References or reports:

Other species planted in addition to *Spartina alterniflora* : *Sporobolus virginicus* was planted on 1-ft centers.

Percent vegetative cover attained by the planted marsh: 100% by each species

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Not assessed.

General comments about the marsh:

- *Spartina alterniflora* clumps were planted on 1-ft centers. The revetment has allowed the plants from the planted area to expand down to the MLW line.
- Marsh name: C.B. Williamson

Marsh Inventory No.: 770 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 07/01/89

Area Planted (1acre = 0.4047 ha): 0.109 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: David Hopple (904) 253-5204 is the contact person.

Location of the marsh.

State: FL **County:** Volusia **Latitude (N):** 29 01' 40" **Longitude (W):** 80 55' 00?"

Bay and sub-bays: Indian River North

Directions to the site: New Smyrna Beach; N end of Atlantic Ave (A-1-A), narrow planting area between two coquina rock revetments.

Source of planting information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

Source of subsequent assessment information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

References or reports:

Other species planted in addition to *Spartina alterniflora* : *Sporobolus virginicus* was planted on 1.5-ft centers.

Percent vegetative cover attained by the planted marsh: 90-100% as of Sept-93

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Not assessed.

General comments about the marsh:

- *Spartina alterniflora* clumps were planted on 1.5-ft centers. Shoreline stabilization project and native grass stormwater retention area.
- Marsh name: Spy Glass

Marsh Inventory No.: 781 **Information Rank (1-5):** 4 **State:** FL

Date Planted: 12/01/85

Area Planted (1acre = 0.4047 ha): 1.599 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Harriet hickox (904) 427-1239 is the contact person.

Location of the marsh.

State: FL **County:** Volusia **Latitude (N):** 29 04' 00?" **Longitude (W):** 80 58' 00?"

Bay and sub-bays: Indian River; Callalosa Creek

Directions to the site: New Smyrna Beach; State Rd. 44 between Indian River North and Callalosa Creek; fringing marsh completely surrounding N, E, and S sides of dredged material deposition island.

Source of planting information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

Source of subsequent assessment information: Mr. Steve Beeman, Ecoshores, Inc., Shoreline Associates, 3881 South Nova Road, Port Orange, FL 32127, (904) 767-6232.

References or reports:

Other species planted in addition to *Spartina alterniflora* : *Distichlis spicata*, *Spartina patens*, *Paspalum vaginatum*, *Sporobolus virginicus*, *Batis maritima* and *Salicornia virginica* were planted on 2-ft centers; *Avicennia germinans*.

Percent vegetative cover attained by the planted marsh: 85-100% cover as of Nov-93.

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Not assessed.

General comments about the marsh:

- *Spartina alterniflora* clumps were planted on 2-ft centers. The construction of the sloping wood-slat breakwater has allowed the creation of a vegetated intertidal shoreline where severe wave erosion had existed.

- Marsh name: Bouchelle Island.

104

Marsh Inventory No.: 299 **Information Rank (1-5):** 3 **State:** ME

Date Planted: 1979

Area Planted (1acre = 0.4047 ha): 1.376 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Portland Water District

Location of the marsh,

State: ME **County:** Cumberland **Latitude (N):** **Longitude (W):**

Bay and sub-bays: Atlantic Ocean; Fore River

Directions to the site: From stations 1+75 to station 22+45 north and south of Congress St., Portland, ME.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Planting density of *S. alterniflora* was 10 pure live seed per sq. ft. Marsh shape: irregular. Built by Environmental Concern, Inc.
- Marsh name: Fore River.

Marsh Inventory No.: 292 **Information Rank (1-5):** 3 **State:** ME

Date Planted: 05/01/78

Area Planted (1acre = 0.4047 ha): 1.619 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Portland Water District

Location of the marsh.

State: ME **County:** Cumberland **Latitude (N):** 43 40' 20?" **Longitude (W):** 70 15' 40?"

Bay and sub-bays: Casco Bay; Back Cove

Directions to the site: Between Vannah and George Sts., Portland, ME, on Back Cove.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- *S. alterniflora* was planted on 2-ft centers. Marsh shape: cove shoreline (linear). ME DEP Permit # 03-2153-05170. Bare spots noted in marsh at end of first growing season. Built by Environmental Concern, Inc.
- Marsh name: Back Cove

Marsh Inventory No.: 296 **Information Rank (1-5):** 3 **State:** ME

Date Planted: 1978

Area Planted (1acre = 0.4047 ha): 0.417 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Wells Sanitary District, P.O. Box 425, Wells ME 04090

Location of the marsh.

State: ME **County:** York **Latitude (N):** **Longitude (W):**

Bay and sub-bays: Atlantic Ocean; Wells Harbor

Directions to the site: Contact the Wells Sanitary District (207) 646-5906 for specific directions.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- *S. alterniflora* was planted on 2-ft centers. Marsh shape: irregular. Built by Environmental Concern, Inc.

- Marsh name: Wells Sanitary District.

Marsh Inventory No.: 308 **Information Rank (1-5):** 3 **State:** MD

Date Planted: 04/01/82

Area Planted (1 acre = 0.4047 ha): 0.809 hectares

COE Permit No.: NABOP-FN-80-0085-2 **Date of COE Permit:**

Originator or Applicant for permit: Locke Insulators Inc.

Location of the marsh.

State: MD **County:** Baltimore City **Latitude (N):** **Longitude (W):**

Bay and sub-bays: Chesapeake Bay; Patapsco River

Directions to the site: Between Potee and Hanover Sts., south of Reedbird Ave., Baltimore MD.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- *S. alterniflora* was planted on 2-ft centers. Marsh shape: rectangular. Built by Environmental Concern, Inc. .The site was destroyed by a suspected herbicide dumping.
- Marsh name: Locke Insulator

108

Marsh Inventory No.: 304 **Information Rank (1-5):** 3 **State:** MD

Date Planted: 06/01/82

Area Planted (1acre = 0.4047 ha): 0.372 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: MD State Highway Administration

Location of the marsh.

State: MD **County:** Baltimore City **Latitude (N):** **Longitude (W):**

Bay and sub-bays: Chesapeake Bay; Patapsco River

Directions to the site: Patapsco River at 6386 York Rd, Baltimore, MD.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* : 5,363 *Scirpus americanus* were also planted on 1.5-ft centers.

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- *S. alterniflora* was planted on 1.5-ft centers. Marsh shape: rectangular. Built by Environmental Concern, Inc. Application # NABOP-F80-013202. Permit # BC 246-42-815.

- Marsh name: City Garage Baltimore.

Marsh Inventory No.: 330 **Information Rank (1-5):** 3 **State:** MD

Date Planted: 05/01/87

Area Planted (1acre = 0.4047 ha): 3.035 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: MD State Highway Administration

Location of the marsh.

State: MD **County:** Baltimore City **Latitude (N):** 39 12' 50" **Longitude (W):** 76 32' 15"

Bay and sub-bays: Chesapeake Bay; Patapsco River

Directions to the site: Southwestern shoulder of the Francis Scott Key Bridge at Hawkins Point.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- *S. alterniflora* was planted on 2-ft centers. Marsh shape: linear. Built by Environmental Concern, Inc. Federal Assistance Program # 1-95-4 (45) 31. MDSHA# BC-246-131-815.

- Marsh name: Hawkins Point.

Marsh Inventory No.: 329 **Information Rank (1-5):** 3 **State:** MD

Date Planted: 09/01/88

Area Planted (1acre = 0.4047 ha): 0.971 hectares

COE Permit No.: NACOP-FW(Helfrich, G.W.)80-0244-2 **Date of COE Permit:**

Originator or Applicant for permit: George Helfrich, 1320 N. Monroe, Baltimore, MD 21217.

Location of the marsh.

State: MD **County:** Baltimore City **Latitude (N):** 39 15' 20" **Longitude (W):** 76 37' 40"

Bay and sub-bays: Chesapeake Bay; Patapsco River, Middle Branch

Directions to the site: Take the Baltimore-Washington Parkway east to the Waterview Ave. exit. Take Waterview Ave. east. The site is at the bottom slope of a landfill. Park at old landfill entrance and walk north, entrance is on the left.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- *S. alterniflora* was planted on 2-ft centers. Marsh shape: shoreline. Built by Environmental Concern, Inc. . Garbage has floated into the marsh, covering some of it (March, 1993).

- Marsh name: Tower Marina

Marsh Inventory No.: 316 **Information Rank (1-5):** 3 **State:** MD

Date Planted: 04/01/87

Area Planted (1acre = 0.4047 ha): 3.035 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Baltimore RESCO

Location of the marsh.

State: MD **County:** Baltimore City **Latitude (N):** **Longitude (W):**

Bay and sub-bays: Chesapeake Bay; Patapsco River; Middle Branch.

Directions to the site: Adjacent to I-95, E of Annapolis Rd, and W of Swann Park, Baltimore MD.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- *S. alterniflora* was planted on 2-ft centers. Marsh shape: 2 sites: oval. Built by Environmental Concern, Inc.

- Marsh name: Rust International.

112

Marsh Inventory No.: 283 **Information Rank (1-5):** 3 **State:** MD

Date Planted: 05/01/87

Area Planted (1acre = 0.4047 ha): 0.809 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Baltimore District Army Corps of Engineers

Location of the marsh.

State: MD **County:** Dorchester **Latitude (N):** 38 30' 10" **Longitude (W):** 76 17' 07"

Bay and sub-bays: Chesapeake Bay; Little Choptank River; Slaughter Creek.

Directions to the site: Cove facing Travers Cove on the S side of Hooper Point on Taylors Island and at the mouth of Slaughter Creek.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information: Environmental Effects of Dredging Vol. D-88-6, Dec. 1988 WES, Vicksburg, MS.

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Planting density of *S. alterniflora* was 10 pure live seed per sq. ft. Marsh shape: semi-circle. Built by Environmental Concern, Inc. Contract # DACW 31-86-M-0339. This is a beneficial use of dredge spoil project.
- Marsh name: Slaughter Creek.

Marsh Inventory No.: 281 **Information Rank (1-5):** 3 **State:** MD

Date Planted: 05/01/82

Area Planted (1acre = 0.4047 ha): 6.475 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Baltimore District Army Corps of Engineers.

Location of the marsh.

State: MD **County:** Dorchester **Latitude (N):** 38 20' 21" **Longitude (W):** 76 15' 30"

Bay and sub-bays: Chesapeake Bay; Tar Bay.

Directions to the site: In Tar Bay just 1 km off the NE tip of Barren Island.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information: (Earhart & Garbisch,1983)

References or reports: (Earhart & Garbisch,1983)

Other species planted in addition to *Spartina alterniflora* : *Spartina patens*

Percent vegetative cover attained by the planted marsh: Estimated to be >50% as of 1983

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? no

General comments about the marsh:

- Planting density of *S. alterniflora* was 10 pure live seed per sq. ft. Marsh shape: irregular. Built by Environmental Concern, Inc. This was a beneficial use of dredged material project.

- Marsh name: Barren Island #1.

Marsh Inventory No.: 315 **Information Rank (1-5):** 3 **State:** MD

Date Planted: 07/01/83

Area Planted (1acre = 0.4047 ha): 6.475 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Baltimore District Army Corps of Engineers.

Location of the marsh.

State: MD **County:** Dorchester **Latitude (N):** 38 20' 20" **Longitude (W):** 76 15' 31"

Bay and sub-bays: Chesapeake Bay; Tar Bay.

Directions to the site: Northeast side of Barren Island, facing Tar Bay.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Planting density of *S. alterniflora* was 10 pure live seed per sq. ft. Marsh shape: irregular. Built by Environmental Concern, Inc. Contract # DACW31-81-C-0073. This was a beneficial use of dredge spoil project.

- Marsh name: Barren Island #2.

Marsh Inventory No.: 317 **Information Rank (1-5):** 3 **State:** MD

Date Planted: 1985

Area Planted (1 acre = 0.4047 ha): 0.931 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: US Army COE, Baltimore District

Location of the marsh.

State: MD **County:** Northumberland **Latitude (N):** **Longitude (W):**

Bay and sub-bays: Chesapeake Bay; Little Wicomico River

Directions to the site: Rt. 652 at Smith Point at the mouth of the Little Wicomico River.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- *S. alterniflora* was planted on 2-ft centers. Marsh shape: irregular. Built by Environmental Concern, Inc. COE Contract # DACW31-85-C-0015. This is a dredge spoil planting project.

- Marsh name: Little Wicomico Planting.

Marsh Inventory No.: 324 **Information Rank (1-5):** 3 **State:** MD

Date Planted: 05/01/87

Area Planted (1acre = 0.4047 ha): 0.769 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Dutton and Garfield Inc., P.O. Box 746, Manchester, NH 03105.

Location of the marsh.

State: MD **County:** Queen Annes **Latitude (N):** 38 58' 25" **Longitude (W):** 76 15' 00"

Bay and sub-bays: Chesapeake Bay; Eastern Bay; Prospect Bay; Kent Narrows

Directions to the site: In the SE corner of Egg Harbor just W of Kent Narrows Rd. and N of US Hwy. 50.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* : 0.8 acres of *Typha angustifolia* and 0.8 acres of *Scirpus americanus*.

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Planting density of *S. alterniflora* was 10 pure live seed per sq. ft. Marsh shape: irregular. Built by Environmental Concern, Inc.
- Marsh name: Eastern Shore Outlet Center.

Marsh Inventory No.: 322 **Information Rank (1-5):** 3 **State:** MD

Date Planted: 06/01/86

Area Planted (1acre = 0.4047 ha): 0.607 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Ellwood Thompson

Location of the marsh.

State: MD **County:** Queen Annes **Latitude (N):** 38 57' 12" **Longitude (W):** 76 13' 45"

Bay and sub-bays: Chesapeake Bay; Eastern Bay; Prospect Bay; Marshy Creek

Directions to the site: Cove on S margin of Marshy Creek just 4 km WSW of Grasonville.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Planting density of *S. alterniflora* was 10 pure live seed per sq. ft. Marsh shape: half-moon. Built by Environmental Concern, Inc.
- Marsh name: Oyster Cove.

118

Marsh Inventory No.: 289 **Information Rank (1-5):** 3 **State:** MD

Date Planted: 04/01/76

Area Planted (1acre = 0.4047 ha): 0.018 hectares

COE Permit No.: NABOP-F/4 (Corinthian Yacht Club) 73-30 **Date of COE Permit:**

Originator or Applicant for permit: Corinthian Yacht Club, Ridge, MD. 20680

Location of the marsh.

State: MD **County:** St. Mary's **Latitude (N):** 38 07' 05" **Longitude (W):** 76 23' 48"

Bay and sub-bays: Chesapeake Bay; Smith Creek; Jutland Creek; Deep Cove

Directions to the site: At the eastern corner of the mouth of Deep Cove as it enters Jutland Creek, about 1 km NE of Wynne, St. Mary's County, MD. Planted behind a bulkhead.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- *S. alterniflora* was planted on 1-ft centers. Marsh shape: rectangular. Built by Environmental Concern, Inc.

- Marsh name: Corinthian Yacht Club.

Marsh Inventory No.: 327 **Information Rank (1-5):** 3 **State:** MD

Date Planted: 06/01/88

Area Planted (1acre = 0.4047 ha): 0.344 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: MD State Highway Administration.

Location of the marsh.

State: MD **County:** Talbot **Latitude (N):** 38 35' 45" **Longitude (W):** 76 01' 00?"

Bay and sub-bays: Chesapeake Bay; Choptank River, Bolingbroke Creek

Directions to the site: W shore of Bolingbroke Creek about 0.5 km N of where it opens into the Choptank River, at Highlys Beach.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- *S. alterniflora* was planted on 2-ft centers. Marsh shape: shoreline. Built by Environmental Concern, Inc.

- Marsh name: Choptank River Bridge.

120

Marsh Inventory No.: 341 **Information Rank (1-5):** 3 **State:** MD

Date Planted: 05/01/91

Area Planted (1acre = 0.4047 ha): 0.016 hectares

COE Permit No.: CENAB-OP-RS(BALDWIN,WH) 89-1221-4 **Date of COE Permit:**

Originator or Applicant for permit: William Baldwin

Location of the marsh.

State: MD **County:** Talbot **Latitude (N):** 38 51' 08" **Longitude (W):** 76 15' 55"

Bay and sub-bays: Chesapeake Bay; Harris Creek, Dogwood Harbor

Directions to the site: Between Dogwood Harbor Rd and South Absontown Rd, at Dogwood Harbor on Tilghman Island, MD.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- *S. alterniflora* was planted on 2-ft centers. Marsh shape: shoreline, rectangular. Built by Environmental Concern, Inc.
- Marsh name: Blue Haven.

Marsh Inventory No.: 320 **Information Rank (1-5):** 3 **State:** MD

Date Planted: 04/01/86

Area Planted (1acre = 0.4047 ha): 0.607 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Walter Johnston Seedling

Location of the marsh.

State: MD **County:** Talbot **Latitude (N):** 38 47' 02" **Longitude (W):** 76 10' 50"

Bay and sub-bays: Chesapeake Bay; Miles River

Directions to the site: On Marengo Farm Rd. at Hunting Creek.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- *S. alterniflora* was planted on 2-ft centers. Marsh shape: irregular. Built by Environmental Concern, Inc. This was a MD State Tidelands violation and settlement approved by the Baltimore District COE.

- Marsh name: Johnston. MD State Wetlands License # 81-WL-0111(R). Notice of violation #'s 85WV051 and 85WV052. Waterfowl damage happened after planting.

Marsh Inventory No.: 325 **Information Rank (1-5):** 3 **State:** MD

Date Planted: 05/01/73

Area Planted (1acre = 0.4047 ha): 0.809 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: US Army COE, Fort Belvoir, VA

Location of the marsh.

State: MD **County:** Talbot **Latitude (N):** 38 51' 20" **Longitude (W):** 76 15' 55"

Bay and sub-bays: Chesapeake Bay; San Domingo Creek

Directions to the site: 9 km NNW of St. Michaels, on the NW facing shore about 1.5 km SW of Tilghman Point,

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information: (Garbisch, Woller, & McCallum, 1975)

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- *S. alterniflora* was planted on 2-ft centers. Marsh shape: irregular. Built by Environmental Concern, Inc.
- Marsh name: Hambleton Island Breach. COE Contract # DACW72-73-C-0009.

Marsh Inventory No.: 282 **Information Rank (1-5):** 3 **State:** MD

Date Planted: 05/01/73

Area Planted (1acre = 0.4047 ha): 0.809 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: US Army COE, Fort Belvoir, VA

Location of the marsh.

State: MD **County:** Talbot **Latitude (N):** 38 45' 15" **Longitude (W):** 76 13' 55"

Bay and sub-bays: Chesapeake Bay; San Domingo Creek

Directions to the site: 3.5 km S of St. Michaels in the mouth of San Domingo Creek at the breach in Hambleton Island.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information: (Garbisch, Woller, & McCallum, 1975)

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- *S. alterniflora* was planted on 2-ft centers. Marsh shape: irregular. Built by Environmental Concern, Inc.
- Marsh name: Hambleton Island Breach. COE Contract # DACW72-73-C-0009.

Marsh Inventory No.: 626 **Information Rank (1-5):** 3 **State:** MD

Date Planted: 05/01/73

Area Planted (1acre = 0.4047 ha): 0.040 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit:

Location of the marsh.

State: MD **County:** Talbot **Latitude (N):** 38 41' 10" **Longitude (W):** 76 11' 40"

Bay and sub-bays: Chesapeake Bay; Tred Avon River

Directions to the site: Directly W across the Tred Avon River from Oxford, MD.; on Ferry Neck peninsula.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- *S. alterniflora* was planted on 2-ft centers. Marsh shape: shoreline. Built by Environmental Concern, Inc.

- Marsh name: Bachelor Point No.1.

Marsh Inventory No.: 343 **Information Rank (1-5):** 3 **State:** MD

Date Planted: 05/01/87

Area Planted (1acre = 0.4047 ha): 0.040 hectares

COE Permit No.: NABOP-RS-(Bachelor's Point) 86-0741-1 **Date of COE Permit:**

Originator or Applicant for permit: Bachelor Point Harbor Marina

Location of the marsh.

State: MD **County:** Talbot **Latitude (N):** 38 40' 32" **Longitude (W):** 76 10' 35"

Bay and sub-bays: Chesapeake Bay; Tred Avon River

Directions to the site: Directly S from Oxford, MD., on the Tred Avon River shore at the Bachelor Point Harbor.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- *S. alterniflora* was planted on 2-ft centers. Marsh shape: shoreline. Built by Environmental Concern, Inc.
- Marsh name: Bachelor Point No.2.

126

Marsh Inventory No.: 318 **Information Rank (1-5):** 3 **State:** MD

Date Planted: 1985

Area Planted (1acre = 0.4047 ha): 2.926 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Merritt Commercial Saving and Loan Association.

Location of the marsh.

State: MD **County:** Worcester **Latitude (N):** **Longitude (W):**

Bay and sub-bays: Assawoman Bay

Directions to the site: Between North and South Heron Drives, Ocean City, MD.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Planting density of *S. alterniflora* was 10 pure live seed per sq. ft. Marsh shape: oval. Built by Environmental Concern, Inc.

- Marsh name: Heron Harbor. MD Tidal Division Case # 82-WL-0255 (R). Some seeding failure occurred and peat potted plants were later installed.

Marsh Inventory No.: 303 **Information Rank (1-5):** 3 **State:** MA

Date Planted: 05/01/81

Area Planted (1acre = 0.4047 ha): 1.093 hectares

COE Permit No.: none **Date of COE Permit:**

Originator or Applicant for permit: Massachusetts Metropolitan District Commission

Location of the marsh.

State: MA **County:** Suffolk **Latitude (N):** 42 23' 35" **Longitude (W):** 70 59' 35"

Bay and sub-bays: Massachusetts Bay; Boston Harbor; President Roads; Winthrop Bay; Belle Isle Inlet.

Directions to the site: Belle Isle marsh reservation park, East Boston MA. Rt. 60 in Boston, just past Suffolk Downs Race track, left at traffic light. At next major intersection take left on Bennington St. Go about 1 mile. Site entrance on right across from the track.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- *S. alterniflora* was planted on 2-ft centers, Marsh shape: fringe on shoreline of a canal. Built by Environmental Concern, Inc. Initial planting problems occurred due to high salinity and rubble substrate.
- Marsh name: Belle Isle Marsh, MA.

128

Marsh Inventory No.: 305 **Information Rank (1-5):** 3 **State:** NJ

Date Planted: 1981

Area Planted (1acre = 0.4047 ha): 0.405 hectares

COE Permit No.: NAPOP-R-N80-17 **Date of COE Permit:**

Originator or Applicant for permit: Scarborough Corp., P.O. Box 387, Marlton, NJ 08053.

Location of the marsh.

State: NJ **County:** Atlantic **Latitude (N):** 39 25' 10" **Longitude (W):** 74 30' 00?"

Bay and sub-bays: Absecon Creek

Directions to the site: Just south (approx. 150') of Cortez Ave. and the intersection of Lisbon Ave., Absecon City, NJ.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* : *Spartina patens*

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- *S. patens* was planted on 2-ft centers. Marsh shape: irregular. Built by Environmental Concern, Inc.

- Marsh name: Scarborough.

Marsh Inventory No.: 332 **Information Rank (1-5):** 3 **State:** NJ

Date Planted: 05/01/81

Area Planted (1acre = 0.4047 ha): 0.081 hectares

COE Permit No.: NAPOP-R-78-12226-3 **Date of COE Permit:** 1978

Originator or Applicant for permit: Harrah's Resorts.

Location of the marsh.

State: NJ **County:** Atlantic **Latitude (N):** 39 22' **Longitude (W):** 74 25'

Bay and sub-bays: Absecon Inlet

Directions to the site: The marina at Harrah's Resort, Atlantic City, NJ.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- *S. alterniflora* was planted on 2-ft centers. Marsh shape: irregular. Built by Environmental Concern, Inc.

- Marsh name: Harrah's Marina.

Marsh Inventory No.: 342 **Information Rank (1-5):** 3 **State:** NJ

Date Planted: 05/01/83

Area Planted (1acre = 0.4047 ha): 0.073 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Scarboro-DiSabatino Construction, Suite 25, Barclay Pavilion, Rt. 70, Cherry Hill, NJ.

Location of the marsh.

State: NJ **County:** Atlantic **Latitude (N):** 39 19' 02" **Longitude (W):** 74 36' 50"

Bay and sub-bays: Gt. Egg Harbor Inlet Ship Channel; Bass Harbor.

Directions to the site: In the Somers Point Yacht Harbor.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- *S. alterniflora* was planted on 2-ft centers. Marsh shape: oval. Built by Environmental Concern, Inc.

- Marsh name: Somers Point Yacht Harbor.

Marsh Inventory No.: 340 **Information Rank (1-5):** 3 **State:** NJ

Date Planted: 1991

Area Planted (1acre = 0.4047 ha): 0.040 hectares

COE Permit No.: CENAB-OP-R-80-03968-12 **Date of COE Permit:**

Originator or Applicant for permit: Shapiro Ripps, Trump Tower, 725 5 th Ave., NY, NY 10022.

Location of the marsh.

State: NJ **County:** Atlantic **Latitude (N):** **Longitude (W):**

Bay and sub-bays:

Directions to the site: Lot 103, Block A-59, Holgate, NJ. At the end of Rosemma Ave., in Holgate, in back of the lot at the center of the Cul-de-sac.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- *S. alterniflora* was planted on 2-ft centers. Marsh shape: rectangular. Built by Environmental Concern, Inc. Marsh name: Shapiro Ripps. This was a restoration project required for illegal fill activity.

Marsh Inventory No.: 319 **Information Rank (1-5):** 3 **State:** NJ

Date Planted: 1986

Area Planted (1 acre = 0.4047 ha): 30.353 hectares

COE Permit No.: 12746 **Date of COE Permit:**

Originator or Applicant for permit: Hartz Mountain Development

Location of the marsh.

State: NJ **County:** Bergen **Latitude (N):** 40 48' 03" **Longitude (W):** 74 33' 05"

Bay and sub-bays: Newark Bay; Hackensack River; Cromakill Creek.

Directions to the site: Secaucus, east side of the eastern spur of the NJ Toll Road.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Planting density of *S. alterniflora* was 10 pure live seed per sq. ft. Built by Environmental Concern, Inc. Vegetation establishment has been problematic in some areas.

- Marsh name: Hartz Mountain Industries Eastern Brackish Marsh. Don Smith, Hackensack Meadowland Development Commission (201) 460-1700 can give a tour of the site.

Marsh Inventory No.: 311 **Information Rank (1-5):** 3 **State:** NJ

Date Planted: 06/01/83

Area Planted (1acre = 0.4047 ha): 1.174 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: NJ Turnpike Authority

Location of the marsh.

State: NJ **County:** Bergen **Latitude (N):** 40 50' 05" **Longitude (W):** 74 33' 00"

Bay and sub-bays: Newark Bay; Hackensack River

Directions to the site: Immediately south of the Vince Lombardi service area at the junction of the eastern and western spurs of the NJ Turnpike, south of exit 18E.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- *S. alterniflora* was planted on 2-ft centers. Built by Environmental Concern, Inc. NJ Turnpike Authority Contract # 780. Originally, the planting failed due to deconsolidated sediments which would not support plant overwintering.

- Goose and Muskrat predation was also a problem. The site is now a mixture of mudflat, Eleocharis flat, and Cordgrass. Marsh shape-basin. Marsh name: NJ Turnpike Authority.

Marsh Inventory No.: 331 **Information Rank (1-5):** 3 **State:** NJ

Date Planted: 1989

Area Planted (1acre = 0.4047 ha): 0.405 hectares

COE Permit No.: 14622 **Date of COE Permit:**

Originator or Applicant for permit: Hackensack Meadowlands Development Commission.

Location of the marsh.

State: NJ **County:** Bergen **Latitude (N):** **Longitude (W):**

Bay and sub-bays: Newark Bay; Mill Creek, Hackensack River

Directions to the site: At the end of Valley Brook Ave., Lyndhurst, NJ.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- *S. alterniflora* was planted on 2-ft centers. Marsh shape: shoreline. Built by Environmental Concern, Inc. Application # 87-0596-JI. Goose damage was a problem at this site.

- Don Smith, Hackensack Meadowland Development Commission (201) 460-1700 can give a tour of the site. Marsh name: Hackensack Meadowlands Development Commission 440 Shoreline.

Marsh Inventory No.: 684 **Information Rank (1-5):** 3 **State:** NJ

Date Planted: 1990

Area Planted (1acre = 0.4047 ha): 3.642 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Hackensack Meadowlands Development Commission.

Location of the marsh.

State: NJ **County:** Bergen **Latitude (N):** 40 48' 50" **Longitude (W):** 74 08' 03"

Bay and sub-bays: Newark Bay; Mill Creek, Hackensack River

Directions to the site: Lyndhurst, NJ, next to the NJ Turnpike and bordered by Berry's Creek on the N and a branch of Mary Ann's Creek to the S.

Source of planting information: Dr. Michael P. Bontje, B. Laing Associates, Environmental Science, 225 Main Street, Northport, NY 11768, (516) 261-7170

Source of subsequent assessment information: (Bontje, 1991)

References or reports: (Bontje, 1991)

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? no

General comments about the marsh:

- *S. alterniflora* was planted on 3-ft centers using 3-4 stem peat pot. Marsh borders two tidal creek channels. Built by Environmental Concern, Inc.
- Appeared to be growing, but was too early to tell. Bird population had already increased, so people were optimistic about the future success of the project.

136

Marsh Inventory No.: 306 **Information Rank (1-5):** 3 **State:** NJ

Date Planted: 04/01/81

Area Planted (1acre = 0.4047 ha): 1.619 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: US Army COE, New York District.

Location of the marsh.

State: NJ **County:** Essex **Latitude (N):** 40 38' 00?" **Longitude (W):** 74 15' 00?"

Bay and sub-bays: Newark Bay; Elizabeth River

Directions to the site: The Elizabeth River at the Joint Meeting sewage treatment plant, South First St., Elizabeth, NJ.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- *S. alterniflora* was planted on 2-ft centers. Marsh shape: rectangular. Built by Environmental Concern, Inc. COE Contract # DACW-78-C-0050
- Marsh name: Elizabeth Marsh.

Marsh Inventory No.: 323 **Information Rank (1-5):** 3 **State:** NJ

Date Planted: 1985

Area Planted (1acre = 0.4047 ha): 30.353 hectares

COE Permit No.: 12746 **Date of COE Permit:**

Originator or Applicant for permit: Hartz Mountain Development

Location of the marsh.

State: NJ **County:** Hudson **Latitude (N):** 40 48' 03" **Longitude (W):** 74 33' 08"

Bay and sub-bays: Newark Bay; Hackensack River; Mill Creek.

Directions to the site: Secaucus, west side of the eastern spur of the NJ Toll Road.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information: (Bontje, 1988)

References or reports: (Bontje, 1988)

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Yes. Fish, birds and mammals were assessed (Bontje, 1988).

General comments about the marsh:

- Planting density of *S. alterniflora* was 10 pure live seeds per sq. ft. Built by Environmental Concern, Inc. Fairly well established cordgrass population. Animal damage is prevalent in some areas.

- Marsh name: Hartz Mountain Western Brackish Marsh. Don Smith, Hackensack Meadowland Development Commission (201) 460-1700 can give a tour of the site.

Marsh Inventory No.: 344 **Information Rank (1-5):** 3 **State:** NJ

Date Planted: 05/01/89

Area Planted (1acre = 0.4047 ha): 0.243 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Dresdner, Robin and Associates

Location of the marsh.

State: NJ **County:** Hudson **Latitude (N):** 40 41' 50" **Longitude (W):** 74 03' 30"

Bay and sub-bays: Upper New York Bay; Hudson River.

Directions to the site: Along the south and east facing shoreline at the Water's Edge renovation area a few blocks N of the Morris Canal Basin, Jersey City.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- *S. alterniflora* was planted on 2-ft centers. Marsh shape: shoreline. Built by Environmental Concern, Inc.

- Marsh name: Newport.

Marsh Inventory No.: 301 **Information Rank (1-5):** 3 **State:** NJ

Date Planted: 04/01/78

Area Planted (1acre = 0.4047 ha): 0.890 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: William R. Cesare or James C. DeCesare

Location of the marsh.

State: NJ **County:** Ocean **Latitude (N):** 39 41' 16" **Longitude (W):** 74 09' 00"

Bay and sub-bays: Barnegat Bay (S of inlet)

Directions to the site: Point of land immediately east of Lagoon Drive West, Harvey Cedars NJ.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- *S. alterniflora* was planted on 2-ft centers. Marsh shape: irregular. Built by Environmental Concern, Inc. NJ Federal District Court Action # 77-1176.
- Marsh name: Long Beach NJ.

140

Marsh Inventory No.: 334 **Information Rank (1-5):** 3 **State:** NJ

Date Planted: 08/01/86

Area Planted (1acre = 0.4047 ha): 0.243 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: West Bay Point Inc.

Location of the marsh.

State: NJ **County:** Ocean **Latitude (N):** 39 51' 27" **Longitude (W):** 74 08' 05"

Bay and sub-bays: Barnegat Bay N of Barnegat Bay Inlet; Laurel Harbor

Directions to the site: Along the shoreline of Laurel Harbor Subdivision, Lacey Township, Ocean County, NJ.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- *S. alterniflora* was planted on 2-ft centers. Marsh shape: horseshoe. Built by Environmental Concern, Inc. NJ CAFRA Permit # 85-0716-5.

- Marsh name: Laurel Harbor.

Marsh Inventory No.: 337 **Information Rank (1-5):** 3 **State:** NJ

Date Planted: 05/01/89

Area Planted (1acre = 0.4047 ha): 0.130 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Michael Reed Construction Co.

Location of the marsh.

State: NJ **County:** Ocean **Latitude (N):** 39 44' 20" **Longitude (W):** 74 07' 05"

Bay and sub-bays: Barnegat Bay S of Barnegat Inlet

Directions to the site: Block T-166, Lots 6 and 7 on the bay shore just S of Holly Lagoons housing development, Long Beach Township, Ocean County, NJ.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- *S. alterniflora* was planted on 2-ft centers. Marsh shape: rectangular. Built by Environmental Concern, Inc. US District Court (NJ) Action # 87-552(GEB).
- Marsh name: Michael Reed.

142

Marsh Inventory No.: 295 **Information Rank (1-5):** 3 **State:** NJ

Date Planted: 05/01/78

Area Planted (1acre = 0.4047 ha): 0.405 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Ocean County Sewerage Authority

Location of the marsh.

State: NJ **County:** Ocean **Latitude (N):** 39 39' 16 **Longitude (W):** 74 11' 00"

Bay and sub-bays: Barnegat Bay; Manahawkin Bay

Directions to the site: Along sewer line on Bonnett Island and along access road at Ship Bottom shore, near where Rt. 72 crosses Manahawkin Bay to Ship Bottom, on Long Beach Island, NJ.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- *S. alterniflora* was planted on 2-ft centers. Marsh shape: linear. Built by Environmental Concern, Inc. EPA sewage treatment grant, project # C-34-371-01. Contract #'s SI-2,SI-3,SI-6.

- High salinity at sights required replanting in October of 1978. Marsh name: Ocean County Sewerage Authority.

Marsh Inventory No.: 294 **Information Rank (1-5):** 3 **State:** NJ

Date Planted: 05/01/78

Area Planted (1 acre = 0.4047 ha): 1.619 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Ocean County Sewerage Authority

Location of the marsh.

State: NJ **County:** Ocean **Latitude (N):** 39 39' 33" **Longitude (W):** 74 11' 45"

Bay and sub-bays: Barnegat Bay; Manahawkin Bay

Directions to the site: Along the footing of the NJ Rt. 72 causeway on Cedar Bonnett Island in Manahawkin Bay just W of Ship Bottom, NJ. Work was done on northern edge of Rt. 72.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- *S. alterniflora* was planted on 2-ft centers. Marsh shape: linear. Built by Environmental Concern, Inc. EPA sewage treatment grant, project # C-34-371-01. Contract #'s SI-2,SI-3,SI-6.

- High salinity at sights required replanting in October of 1978. Marsh name: Ocean County Sewerage Authority.

144

Marsh Inventory No.: 293 **Information Rank (1-5):** 3 **State:** NJ

Date Planted: 05/01/78

Area Planted (1acre = 0.4047 ha): 0.809 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Ocean County Sewerage Authority

Location of the marsh.

State: NJ **County:** Ocean **Latitude (N):** 39 40' 10" **Longitude (W):** 74 12' 58"

Bay and sub-bays: Barnegat Bay; Manahawkin Bay

Directions to the site: Along the base of the causeway where Rt. 72 begins to cross Manahawkin Bay over to Ship Bottom, NJ. Work was done on northern edge of Rt. 72.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- *S. alterniflora* was planted on 2-ft centers. Marsh shape: linear. Built by Environmental Concern, Inc. EPA sewage treatment grant, project # C-34-371-01. Contract #'s SI-2,SI-3,SI-6.

- High salinity at sights required replanting in October of 1978. Marsh name: Ocean County Sewerage Authority.

Marsh Inventory No.: 335 **Information Rank (1-5):** 3 **State:** NY

Date Planted: 1986

Area Planted (1acre = 0.4047 ha): 0.202 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: NY City DEP

Location of the marsh.

State: NY **County:** Richmond **Latitude (N):** **Longitude (W):**

Bay and sub-bays: Great Kills Harbor

Directions to the site: South of Tennyson Drive, east of Robinson Ave., west of Glovar Ave., Staten Island, NY.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- *S. alterniflora* was planted on 2-ft centers. Marsh shape: oval. Built by Environmental Concern, Inc. New York City Environmental Conservation Order on Consent R2 0155-85-11.

- Marsh name: Staten Island, NY.

146

Marsh Inventory No.: 339 **Information Rank (1-5):** 3 **State:** NY

Date Planted: 06/01/89

Area Planted (1acre = 0.4047 ha): 0.073 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Northport Marina Associates

Location of the marsh.

State: NY **County:** Suffolk **Latitude (N):** 40 53' 30" **Longitude (W):** 73 21' 26"

Bay and sub-bays: Long Island Sound; Huntington Bay; Northport Bay; Northport Harbor.

Directions to the site: Shore immediately N of the Admirals Yacht Club and Marina and about 25 m S of the Scudder Park boat launch ramp.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- *S. alterniflora* was planted on 2-ft centers. Marsh shape: oval. Built by Environmental Concern, Inc. NY DEC # 10-86-0780.

- Marsh name: Northport Admirals Yacht Club and Marina.

Marsh Inventory No.: 627 **Information Rank (1-5):** 4 **State:** NC

Date Planted: 1973

Area Planted (1acre = 0.4047 ha): hectares

COE Permit No.: none **Date of COE Permit:**

Originator or Applicant for permit: C.S. Benner, research

Location of the marsh.

State: NC **County:** **Latitude (N):** 36 10' 45" **Longitude (W):** 75 45' 30"

Bay and sub-bays: Albermarle Sound; Currituck Sound

Directions to the site: On the Currituck Sound shore of the barrier island, at the COE CERC Field Research Facility, Duck, NC.

Source of planting information: (C.S. Benner et al, 1982)

Source of subsequent assessment information: (Benner et al, 1982)

References or reports: (Benner et al, 1982)

Other species planted in addition to *Spartina alterniflora* : *Spartina patens*, *Spartina cynosuroides*, *Juncus roemerianus*, *Fimbristylis spadicea*, *Phragmites australis*, *A. breviligulata*, *Panicum amarulum*, and *Typha latifolia*.

Percent vegetative cover attained by the planted marsh: >70%

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? no

General comments about the marsh:

- Several profile surveys have been made of the planted area, one in 1978, 1979, 1980 and 1981. The vegetation continued to expand each year, but the unvegetated control area remained bare during this same period.

148

Marsh Inventory No.: 307 **Information Rank (1-5):** 3 **State:** SC

Date Planted: 1981

Area Planted (1acre = 0.4047 ha): 3.238 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: City of North Charleston

Location of the marsh.

State: SC **County:** Dorchester **Latitude (N):** **Longitude (W):**

Bay and sub-bays: Cooper River

Directions to the site: Basin #1, east of Spruill Ave. and north of Token St. Basin #2, east of Chicora Ave., and north of Quitman St. and Baxter St., west of Spruill Ave. Basin #3, west of Chicora Ave., south of Orvid St., North Charleston, SC.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Planting density of *S. alterniflora* was 10 pure live seed per sq. ft. Marsh shape: 3 oval basins. Built by Environmental Concern, Inc. This was HUD project Chicora Retention Basin, Project # B-80-MC-45-0005-28.

- Marsh name: City of North Charleston, SC.

Marsh Inventory No.: 664 **Information Rank (1-5):** 3 **State:** TX

Date Planted: 1984

Area Planted (1acre = 0.4047 ha): 0.121 hectares

COE Permit No.: 15105/1 **Date of COE Permit:** 1983

Originator or Applicant for permit: Dorchester Exploration, Inc.

Location of the marsh.

State: TX **County:** Aransas **Latitude (N):** 27 56' 00" **Longitude (W):** 97 02' 32"

Bay and sub-bays: Aransas Bay

Directions to the site: ST. 260, about 6 mi S. of Rockport; on the SW corner of Mud Island.

Source of planting information: Mr. Gary Galbraith, Espey, Huston & Ass, Inc., (512) 327-6840

Source of subsequent assessment information: Gary Galbraith

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh: 50%

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Planted by Espey Huston. Evaluated in 1987, lots of erosion, wave energy to high. Western 400' of 871' X 50' planted has failed.

150

Marsh Inventory No.: 179 Information Rank (1-5): 3 State: TX

Date Planted:

Area Planted (1 acre = 0.4047 ha): 0.040 hectares

COE Permit No.: 18172 Date of COE Permit: 11/03/87

Originator or Applicant for permit: City of Rockport

Location of the marsh.

State: TX County: Aransas Latitude (N): 28 01' 56" Longitude (W): 97 02' 55"

Bay and sub-bays: Aransas Bay

Directions to the site: Along the W shore of Little Bay at Rockport, TX

Source of planting information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

Marsh Inventory No.: 686 **Information Rank (1-5):** 4 **State:** TX

Date Planted: 1988

Area Planted (1acre = 0.4047 ha): 0.040 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Kontiki Condos

Location of the marsh.

State: TX **County:** Aransas **Latitude (N):** 28 06' 05" **Longitude (W):** 97 01' 47"

Bay and sub-bays: Aransas Bay

Directions to the site: In a bayou just SW of Live Oak Pt, on Aransas Bay 0.5 mi S of the base of the causeway.

Source of planting information: Mr. Charles E. Belaire, Belaire Consulting, Inc., P.O. Box 741, Rockport, TX 78382, (512) 729-2948

Source of subsequent assessment information: Mr. Charles E. Belaire, Belaire Consulting, Inc., P.O. Box 741, Rockport, TX 78382, (512) 729-2948

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh: >70%, 1992

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Very successful (>70% cover).
-

152

Marsh Inventory No.: 239 **Information Rank (1-5):** 3 **State:** TX

Date Planted: 1983

Area Planted (1acre = 0.4047 ha): 0.101 hectares

COE Permit No.: 13679 **Date of COE Permit:**

Originator or Applicant for permit: Bishop Ship-building Corp.

Location of the marsh.

State: TX **County:** Aransas **Latitude (N):** 27 54' 47" **Longitude (W):** 97 07' 50"

Bay and sub-bays: Aransas Bay; Redfish Bay

Directions to the site: N end of the harbor along the NE side; in Aransas Pass

Source of planting information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

-
-

Marsh Inventory No.: 177 **Information Rank (1-5):** 3 **State:** TX

Date Planted:

Area Planted (1acre = 0.4047 ha): 0.405 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: COE ?

Location of the marsh.

State: TX **County:** Aransas **Latitude (N):** 27 52' 10" **Longitude (W):** 97 05' 55"

Bay and sub-bays: Aransas Bay; Redfish Bay

Directions to the site: Stedman Island, SW side.

Source of planting information:

Source of subsequent assessment information: Dr. Thomas Minello, National Marine Fisheries Service, 4700 Ave. U, Galveston, TX 77551, (409) 766-3506

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh: >70%, 1992

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Yes. Aquatic organisms.

General comments about the marsh:

- The marsh is well established. It was sampled for fish, shellfish, and benthic invertebrates, as well as macro-organic matter

Marsh Inventory No.: 236 **Information Rank (1-5):** 3 **State:** TX

Date Planted:

Area Planted (1acre = 0.4047 ha): 0.061 hectares

COE Permit No.: 11405 **Date of COE Permit:**

Originator or Applicant for permit: Mitchell Energy Exp.

Location of the marsh.

State: TX **County:** Aransas **Latitude (N):** 28 10' 45" **Longitude (W):** 96 57' 45"

Bay and sub-bays: Aransas Bay; St. Charles Bay

Directions to the site: State tracts (ST.) 69 and 387, near Rockport

Source of planting information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

Marsh Inventory No.: 685 **Information Rank (1-5):** 3 **State:** TX

Date Planted: 1988

Area Planted (1acre = 0.4047 ha): 0.101 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Northpoint Marina

Location of the marsh.

State: TX **County:** Aransas **Latitude (N):** 28 03' 00" **Longitude (W):** 97 13' 45"

Bay and sub-bays: Copano Bay

Directions to the site: In Port Bay on N side on small peninsula between Italian Bend and Rattlesnake Point.

Source of planting information: Mr. Charles E. Belaire, Belaire Consulting, Inc., P.O. Box 741, Rockport, TX 78382, (512) 729-2948

Source of subsequent assessment information: Mr. Charles E. Belaire, Belaire Consulting, Inc., P.O. Box 741, Rockport, TX 78382, (512) 729-2948

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Partial Success (>10% cover).

-

156

Marsh Inventory No.: 690 **Information Rank (1-5):** 3 **State:** TX

Date Planted: 1992

Area Planted (1acre = 0.4047 ha): 0.004 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: SCS - Bayside, TX

Location of the marsh.

State: TX **County:** Aransas **Latitude (N):** 28 05' 45" **Longitude (W):** 97 12' 20"

Bay and sub-bays: Copano Bay

Directions to the site: Along the NW shore of Copano Bay at Bayside

Source of planting information: Mr. Charles E. Belaire, Belaire Consulting, Inc., P.O. Box 741, Rockport, TX 78382, (512) 729-2948

Source of subsequent assessment information: Mr. Charles E. Belaire, Belaire Consulting, Inc., P.O. Box 741, Rockport, TX 78382, (512) 729-2948

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Partial Success (>10% cover). 600 sq. ft.

Marsh Inventory No.: 689 **Information Rank (1-5):** 4 **State:** TX

Date Planted: 1992

Area Planted (1acre = 0.4047 ha): 0.121 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: SCS - Bayside, TX

Location of the marsh.

State: TX **County:** Aransas **Latitude (N):** 28 05' 30" **Longitude (W):** 97 12' 30"

Bay and sub-bays: Copano Bay

Directions to the site: Along the NW shore of Copano Bay at Bayside

Source of planting information: Mr. Charles E. Belaire, Belaire Consulting, Inc., P.O. Box 741, Rockport, TX 78382, (512) 729-2948

Source of subsequent assessment information: Mr. Charles E. Belaire, Belaire Consulting, Inc., P.O. Box 741, Rockport, TX 78382, (512) 729-2948

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh: >70%, 1992

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Very successful.

-

158

Marsh Inventory No.: 688 **Information Rank (1-5):** 4 **State:** TX

Date Planted: 1992

Area Planted (1acre = 0.4047 ha): 0.121 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: SCS - Bayside, TX

Location of the marsh.

State: TX **County:** Aransas **Latitude (N):** 28 05' 15" **Longitude (W):** 97 12' 45"

Bay and sub-bays: Copano Bay

Directions to the site: Along the NW shore of Copano Bay at Bayside

Source of planting information: Mr. Charles E. Belaire, Belaire Consulting, Inc., P.O. Box 741, Rockport, TX 78382, (512) 729-2948

Source of subsequent assessment information: Mr. Charles E. Belaire, Belaire Consulting, Inc., P.O. Box 741, Rockport, TX 78382, (512) 729-2948

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh: >70%, 1992

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Very successful.

-

Marsh Inventory No.: 687 **Information Rank (1-5):** 4 **State:** TX

Date Planted: 1992

Area Planted (1acre = 0.4047 ha): 0.008 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: SCS - Bayside, TX

Location of the marsh.

State: TX **County:** Aransas **Latitude (N):** 28 05' 05" **Longitude (W):** 97 12' 55"

Bay and sub-bays: Copano Bay

Directions to the site: Along the NW shore of Copano Bay at Bayside

Source of planting information: Mr. Charles E. Belaire, Belaire Consulting, Inc., P.O. Box 741, Rockport, TX 78382, (512) 729-2948

Source of subsequent assessment information: Mr. Charles E. Belaire, Belaire Consulting, Inc., P.O. Box 741, Rockport, TX 78382, (512) 729-2948

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh: >70%, 1992

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Very successful, 1,000 sq. ft.

160

Marsh Inventory No.: 248 **Information Rank (1-5):** 3 **State:** TX

Date Planted: 1986

Area Planted (1acre = 0.4047 ha): 0.304 hectares

COE Permit No.: 17075 **Date of COE Permit:**

Originator or Applicant for permit: Wilmar Development Corp.

Location of the marsh.

State: TX **County:** Aransas **Latitude (N):** 28 01' 45" **Longitude (W):** 97 07' 52"

Bay and sub-bays: Copano Bay; Port Bay

Directions to the site: Cape Valero Subdiv., about 5 mi W. of Rockport

Source of planting information: Mr. Charles E. Belaire, Belaire Consulting, Inc., P.O. Box 741, Rockport, TX 78382, (512) 729-2948

Source of subsequent assessment information: Mr. Charles E. Belaire, Belaire Consulting, Inc., P.O. Box 741, Rockport, TX 78382, (512) 729-2948

References or reports:

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Very successful (>70% cover).
-

Marsh Inventory No.: 692 **Information Rank (1-5):** 4 **State:** TX

Date Planted: 1992

Area Planted (1acre = 0.4047 ha): 0.364 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Mitchell Energy Exp.

Location of the marsh.

State: TX **County:** Aransas **Latitude (N):** 28 10' 10" **Longitude (W):** 96 52' 20"

Bay and sub-bays: Mesquite Bay

Directions to the site: Mesquite Bay side of spoil banks, S of ANWR.

Source of planting information: Mr. Charles E. Belaire, Belaire Consulting, Inc., P.O. Box 741, Rockport, TX 78382, (512) 729-2948

Source of subsequent assessment information: Mr. Charles E. Belaire, Belaire Consulting, Inc., P.O. Box 741, Rockport, TX 78382, (512) 729-2948

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Very successful (>70% cover).
-

162

Marsh Inventory No.: 691 **Information Rank (1-5):** 4 **State:** TX

Date Planted: 1992

Area Planted (1acre = 0.4047 ha): 3.035 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Mitchell Energy Exp.

Location of the marsh.

State: TX **County:** Aransas **Latitude (N):** 28 09' 56" **Longitude (W):** 96 52' 18"

Bay and sub-bays: Mesquite Bay

Directions to the site: S. side of the Aransas National Wildlife Refuge.

Source of planting information: Mr. Charles E. Belaire, Belaire Consulting, Inc., P.O. Box 741, Rockport, TX 78382, (512) 729-2948

Source of subsequent assessment information: Mr. Charles E. Belaire, Belaire Consulting, Inc., P.O. Box 741, Rockport, TX 78382, (512) 729-2948

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Very successful (>70% cover).
-

Marsh Inventory No.: 715 **Information Rank (1-5):** 3 **State:** TX

Date Planted: 1992

Area Planted (1acre = 0.4047 ha): 0.809 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: COE

Location of the marsh.

State: TX **County:** Brazoria **Latitude (N):** 29 10' 28" **Longitude (W):** 95 05' 33"

Bay and sub-bays: Galveston Bay; West Bay.

Directions to the site: Along the NW shore of West Bay in front of the SE corner of Halls Lake.

Source of planting information: Mr. Charles E. Belaire, Belaire Consulting, Inc., P.O. Box 741, Rockport, TX 78382, (512) 729-2948

Source of subsequent assessment information: Mr. Charles E. Belaire, Belaire Consulting, Inc., P.O. Box 741, Rockport, TX 78382, (512) 729-2948

References or reports:

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh: <10

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Too steep a slope and short a planting area; too deep to put in a breakwater so the 4-5 mi fetch in the prevailing wind direction was too much. The planting washed out within a few months. Failure.

164

Marsh Inventory No.: 705 **Information Rank (1-5):** 3 **State:** TX

Date Planted: 06/01/83

Area Planted (1acre = 0.4047 ha): 6.071 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: COE

Location of the marsh.

State: TX **County:** Brazoria **Latitude (N):** 29 10' 42" **Longitude (W):** 95 06' 55"

Bay and sub-bays: Galveston Bay; West Bay; Chocolate Bay.

Directions to the site: Along the SE shore of Chocolate Bay at Alligator Point.

Source of planting information: Dr. James W. Webb, Texas A&M Univ. at Galveston, P.O. Box 1675, Galveston, TX 77553, (409)740-4542

Source of subsequent assessment information: Dr. Thomas Minello, National Marine Fisheries Service, 4700 Ave. U, Galveston, TX 77551, (409) 766-3506

References or reports: (Espey, Huston & Assoc., Inc., 1988)

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Yes. Fisheries and benthic organisms.

General comments about the marsh:

- Fairly well protected, only subject to N wind waves.

Marsh Inventory No.: 698 **Information Rank (1-5):** 3 **State:** TX

Date Planted: 07/01/92

Area Planted (1acre = 0.4047 ha): 0.008 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Seadrift

Location of the marsh.

State: TX **County:** Calhoun **Latitude (N):** 28 24' 30" **Longitude (W):** 96 42' 30"

Bay and sub-bays: San Antonio Bay

Directions to the site: Along the N shore (facing S) just to the E of the boat docks and fish houses.

Source of planting information: Mr. Charles E. Belaire, Belaire Consulting, Inc., P.O. Box 741, Rockport, TX 78382, (512) 729-2948

Source of subsequent assessment information: Mr. Charles E. Belaire, Belaire Consulting, Inc., P.O. Box 741, Rockport, TX 78382, (512) 729-2948

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

-

-

166

Marsh Inventory No.: 680 Information Rank (1-5): 3 State: TX

Date Planted: 1989

Area Planted (1acre = 0.4047 ha): 0.040 hectares

COE Permit No.: Date of COE Permit:

Originator or Applicant for permit: Harlingen St. Bank

Location of the marsh.

State: TX County: Cameron Latitude (N): 26 06' 48" Longitude (W): 97 10' 25"

Bay and sub-bays: Laguna Madre

Directions to the site: 3 mi N of new causeway on the lagoon side of Padre Island.

Source of planting information: Mr. Charles E. Belaire, Belaire Consulting, Inc., P.O. Box 741, Rockport, TX 78382, (512) 729-2948

Source of subsequent assessment information: Mr. Charles E. Belaire, Belaire Consulting, Inc., P.O. Box 741, Rockport, TX 78382, (512) 729-2948

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh: 15

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Partial Success (>10% cover). 900 sq. ft.

Marsh Inventory No.: 681 **Information Rank (1-5):** 3 **State:** TX

Date Planted: 1990

Area Planted (1acre = 0.4047 ha): 0.040 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: NCNB, Texas

Location of the marsh.

State: TX **County:** Cameron **Latitude (N):** 26 04' 40" **Longitude (W):** 97 10' 03"

Bay and sub-bays: Laguna Madre

Directions to the site: South of the old Queen Isabella Causeway.

Source of planting information: Mr. Charles E. Belaire, Belaire Consulting, Inc., P.O. Box 741, Rockport, TX 78382, (512) 729-2948

Source of subsequent assessment information: Mr. Charles E. Belaire, Belaire Consulting, Inc., P.O. Box 741, Rockport, TX 78382, (512) 729-2948

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh: 15

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Partial Success (>10% cover). 2,000 sq. ft.

168

Marsh Inventory No.: 246 **Information Rank (1-5):** 3 **State:** TX

Date Planted:

Area Planted (1acre = 0.4047 ha): 0.004 hectares

COE Permit No.: 16564 **Date of COE Permit:**

Originator or Applicant for permit: J.M. Inhofe

Location of the marsh.

State: TX **County:** Cameron **Latitude (N):** (26 04' 30") **Longitude (W):** (97 10' 00")

Bay and sub-bays: Laguna Madre; Lower Laguna Madre

Directions to the site: W. end of Capricorn St., S. Padre Isl.

Source of planting information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

-
-

Marsh Inventory No.: 244 **Information Rank (1-5):** 3 **State:** TX

Date Planted:

Area Planted (1acre = 0.4047 ha): 0.004 hectares

COE Permit No.: 15883 **Date of COE Permit:**

Originator or Applicant for permit: Settlement II (Laguna del Sol)

Location of the marsh.

State: TX **County:** Cameron **Latitude (N):** (26 04' 30") **Longitude (W):** (97 10' 00")

Bay and sub-bays: Laguna Madre; Lower Laguna Madre

Directions to the site: Laguna del Sol Condo's, W end of Venus and Jupiter Sts., S. Padre Isl.

Source of planting information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

-
-

170

Marsh Inventory No.: 243 **Information Rank (1-5):** 3 **State:** TX

Date Planted:

Area Planted (1acre = 0.4047 ha): 0.004 hectares

COE Permit No.: 15832 **Date of COE Permit:**

Originator or Applicant for permit: Settlement II (Laguna del Sol)

Location of the marsh.

State: TX **County:** Cameron **Latitude (N):** (26 04' 30") **Longitude (W):** (97 10' 00")

Bay and sub-bays: Laguna Madre; Lower Laguna Madre

Directions to the site: N. end of Laguna Circle North, S. Padre Isl.

Source of planting information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

-
-

Marsh Inventory No.: 242 **Information Rank (1-5):** 3 **State:** TX

Date Planted:

Area Planted (1acre = 0.4047 ha): 0.012 hectares

COE Permit No.: 15341 **Date of COE Permit:**

Originator or Applicant for permit: Starrett Construction Co.

Location of the marsh.

State: TX **County:** Cameron **Latitude (N):** (26 04' 30") **Longitude (W):** (97 10' 00")

Bay and sub-bays: Laguna Madre; Lower Laguna Madre

Directions to the site: W. end of Constellation Dr., S. Padre Isl.

Source of planting information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

Marsh Inventory No.: 269 **Information Rank (1-5):** 4 **State:** TX

Date Planted: 1987

Area Planted (1acre = 0.4047 ha): 0.040 hectares

COE Permit No.: none **Date of COE Permit:**

Originator or Applicant for permit: Jackson Ranch

Location of the marsh.

State: TX **County:** Chambers **Latitude (N):** 29 34' 10" **Longitude (W):** 94 40' 00"

Bay and sub-bays: Galveston Bay

Directions to the site: Lake Steveson, southwest corner. Lone Oak Bayou worked into Lake Stevenson.

Source of planting information: Mr. Eddie Seidensticker, U.S. Soil Conservation Service, 17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

Source of subsequent assessment information: Mr. Eddie Seidensticker, U.S. Soil Conservation Service, 17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- 5% *S. alterniflora* shoot survival 5 years later; 60% shoots with seed. California bulrush (*Scirpus californica*) is out competing *S. alterniflora* Failure (0-15%).
- Seidensticker's Plot#3

Marsh Inventory No.: 717 **Information Rank (1-5):** 3 **State:** TX

Date Planted: 1970

Area Planted (1acre = 0.4047 ha): 0.040 hectares

COE Permit No.: none **Date of COE Permit:**

Originator or Applicant for permit: Private owner

Location of the marsh.

State: TX **County:** Chambers **Latitude (N):** 29 31' 20" **Longitude (W):** 94 45' 55"

Bay and sub-bays: Galveston Bay; East Bay

Directions to the site: S facing shore of Smith Point.

Source of planting information: Dr. James W. Webb, Texas A&M Univ. at Galveston, P.O. Box 1675, Galveston, TX 77553, (409)740-4542

Source of subsequent assessment information: Dr. James W. Webb, Texas A&M Univ. at Galveston, P.O. Box 1675, Galveston, TX 77553, (409)740-4542

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

-
-

Marsh Inventory No.: 714 **Information Rank (1-5):** 3 **State:** TX

Date Planted: 1973

Area Planted (1acre = 0.4047 ha): 0.405 hectares

COE Permit No.: none **Date of COE Permit:**

Originator or Applicant for permit: U.S. FWS Anahuac National Wildlife Refuge; J. Webb; COE

Location of the marsh.

State: TX **County:** Chambers **Latitude (N):** 29 33' 45" **Longitude (W):** 94 29' 15"

Bay and sub-bays: Galveston Bay; East Bay

Directions to the site: Anahuac Refuge, along the SE shore of the refuge just 0.25 mi W of mouth of Oyster Bayou

Source of planting information: Dr. James W. Webb, Texas A&M Univ. at Galveston, P.O. Box 1675, Galveston, TX 77553, (409)740-4542

Source of subsequent assessment information: Dr. James W. Webb, Texas A&M Univ. at Galveston, P.O. Box 1675, Galveston, TX 77553, (409)740-4542

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- This was Webb's dissertation study area. Many of the plantings were washed out.

Marsh Inventory No.: 196 **Information Rank (1-5):** 3 **State:** TX

Date Planted: 1978

Area Planted (1acre = 0.4047 ha): 0.005 hectares

COE Permit No.: none **Date of COE Permit:**

Originator or Applicant for permit: U.S. FWS Anahuac National Wildlife Refuge;
Boyscouts

Location of the marsh.

State: TX **County:** Chambers **Latitude (N):** 29 34' 05" **Longitude (W):** 94 32' 00"

Bay and sub-bays: Galveston Bay; East Bay

Directions to the site: Anahuac Refuge, West Refuge Road until it hits the bay;
planting is on the bay side of the road.

Source of planting information: Dr. James W. Webb, Texas A&M Univ. at
Galveston, P.O. Box 1675, Galveston, TX 77553, (409)740-4542

Source of subsequent assessment information: Dr. James W. Webb, Texas A&M
Univ. at Galveston, P.O. Box 1675, Galveston, TX 77553, (409)740-4542

References or reports:

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh:

**Was there an assessment made of animal utilization of the marsh? If so,
which animals were surveyed, and when?**

General comments about the marsh:

- Boyscouts did the transplanting.

176

Marsh Inventory No.: 268 **Information Rank (1-5):** 4 **State:** TX

Date Planted: 1958

Area Planted (1acre = 0.4047 ha): 0.040 hectares

COE Permit No.: none **Date of COE Permit:**

Originator or Applicant for permit: Brown Foundation

Location of the marsh.

State: TX **County:** Chambers **Latitude (N):** 29 33' 25" **Longitude (W):** 94 36' 50"

Bay and sub-bays: Galveston Bay; East Bay

Directions to the site: West windmill 1957-58; on N side of East Bay.

Source of planting information: Mr. Eddie Seidensticker, U.S. Soil Conservation Service, 17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

Source of subsequent assessment information: Mr. Eddie Seidensticker, U.S. Soil Conservation Service, 17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- 85% *S. alterniflora* shoot survival several years later; spread from 1,300 to 1,800 ft width. Hurricane Carla wiped out a good portion of it. Good development.

Marsh Inventory No.: 273 **Information Rank (1-5):** 4 **State:** TX

Date Planted: 1986

Area Planted (1acre = 0.4047 ha): hectares

COE Permit No.: none **Date of COE Permit:**

Originator or Applicant for permit: U.S. FWS Anahuac National Wildlife Refuge

Location of the marsh.

State: TX **County:** Chambers **Latitude (N):** 29 34' 05" **Longitude (W):** 94 32' 00"

Bay and sub-bays: Galveston Bay; East Bay

Directions to the site: Just southeast of plot 7.

Source of planting information: Mr. Eddie Seidensticker, U.S. Soil Conservation Service, 17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

Source of subsequent assessment information: Mr. Eddie Seidensticker, U.S. Soil Conservation Service, 17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- 80% shoot survival 13 months later. Good success.
- Seidensticker's Plot#6

Marsh Inventory No.: 272 **Information Rank (1-5):** 4 **State:** TX

Date Planted: 1986

Area Planted (1acre = 0.4047 ha): 0.010 hectares

COE Permit No.: none **Date of COE Permit:**

Originator or Applicant for permit: U.S. FWS Anahuac National Wildlife Refuge

Location of the marsh.

State: TX **County:** Chambers **Latitude (N):** 29 34' 05" **Longitude (W):** 94 32' 00"

Bay and sub-bays: Galveston Bay; East Bay

Directions to the site: Anahuac Refuge, West Refuge Road until it hits the bay; planting is on the bay side of the road.

Source of planting information: Mr. Eddie Seidensticker, U.S. Soil Conservation Service, 17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

Source of subsequent assessment information: Mr. Eddie Seidensticker, U.S. Soil Conservation Service, 17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- 90% shoot survival 12 months later. Small area planted behind a rock pyramid wavebreak. Excellent (85-100%).
- Seidensticker's Plot#7

Marsh Inventory No.: 267 **Information Rank (1-5):** 4 **State:** TX

Date Planted: 1987

Area Planted (1acre = 0.4047 ha): 0.330 hectares

COE Permit No.: none **Date of COE Permit:**

Originator or Applicant for permit: Brown Foundation

Location of the marsh.

State: TX **County:** Chambers **Latitude (N):** 29 32' 30" **Longitude (W):** 94 40' 30"

Bay and sub-bays: Galveston Bay; East Bay

Directions to the site: On the north shore of East Bay, even with the east end of Lake Surprise

Source of planting information: Mr. Eddie Seidensticker, U.S. Soil Conservation Service, 17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

Source of subsequent assessment information: Mr. Eddie Seidensticker, U.S. Soil Conservation Service, 17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- 70% *S. alterniflora* shoot survival 5 yrs later; slight blight attack; moderate cattle grazing. Good (65-85%). 1,200 ft of plastic snow fence used as wave dampening device.
- Seidensticker's Plot#1

180

Marsh Inventory No.: 270 **Information Rank (1-5):** 4 **State:** TX

Date Planted: 1988

Area Planted (1acre = 0.4047 ha): 0.324 hectares

COE Permit No.: none **Date of COE Permit:**

Originator or Applicant for permit: Brown Foundation

Location of the marsh.

State: TX **County:** Chambers **Latitude (N):** 29 33' 25" **Longitude (W):** 94 36' 50"

Bay and sub-bays: Galveston Bay; East Bay

Directions to the site: West windmill on N side of East Bay; seawall addition in 1988 just to west of original planting.

Source of planting information: Mr. Eddie Seidensticker, U.S. Soil Conservation Service, 17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

Source of subsequent assessment information: Mr. Eddie Seidensticker, U.S. Soil Conservation Service, 17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- 85% *S. alterniflora* shoot survival 4 years later; spread and joined with the marsh that was planted in 1958. Good development.
- Seidensticker's Plot#2

Marsh Inventory No.: 271 **Information Rank (1-5):** 4 **State:** TX

Date Planted: 1989, 1990

Area Planted (1acre = 0.4047 ha): 0.600 hectares

COE Permit No.: none **Date of COE Permit:**

Originator or Applicant for permit: Brown Foundation

Location of the marsh.

State: TX **County:** Chambers **Latitude (N):** 29 34' 40" **Longitude (W):** 94 36' 40"

Bay and sub-bays: Galveston Bay; East Bay

Directions to the site: Lake Robinson, southwest side along a levee.

Source of planting information: Mr. Eddie Seidensticker, U.S. Soil Conservation Service, 17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

Source of subsequent assessment information: Mr. Eddie Seidensticker, U.S. Soil Conservation Service, 17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- 80% shoot survival 2 years later, but cattle have severely damaged the planted area. Stand looks very healthy. Erosion is severe on levee bank where plants are not established. Established areas not spreading due to cattle grazing.

- Seidensticker's Plot#5

182

Marsh Inventory No.: 633 **Information Rank (1-5):** 3 **State:** TX

Date Planted: 08/01/92

Area Planted (1acre = 0.4047 ha): 0.029 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Chambers-Liberty Nav. Dist.

Location of the marsh.

State: TX **County:** Chambers **Latitude (N):** **Longitude (W):**

Bay and sub-bays: Galveston Bay; Trinity Bay

Directions to the site: On a dredged material deposition island along the west side of the Anahuac River Channel at marker 17.

Source of planting information: Mr. Eddie Seidensticker, U.S. Soil Conservation Service, 17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

Source of subsequent assessment information: Mr. Eddie Seidensticker, U.S. Soil Conservation Service, 17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

References or reports:

Other species planted in addition to *Spartina alterniflora* : Giant cutgrass (*Zizaniopsis miliacea*)

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- This plot is in the area of Trinity Delta destroyed by nutria last winter. A vinyl coated chain link fence was erected to keep nutria out; achieved 100% shoot survival one month later.

- Seidensticker's Plot#18

Marsh Inventory No.: 276 **Information Rank (1-5):** 4 **State:** TX

Date Planted: 07/01/91

Area Planted (1acre = 0.4047 ha): 0.010 hectares

COE Permit No.: none **Date of COE Permit:**

Originator or Applicant for permit: Chambers-Liberty Nav. District

Location of the marsh.

State: TX **County:** Chambers **Latitude (N):** **Longitude (W):**

Bay and sub-bays: Galveston Bay; Trinity Bay

Directions to the site: On a dredged material deposition island along the west side of the Anahuac River, at Fish Pass.

Source of planting information: Mr. Eddie Seidensticker, U.S. Soil Conservation Service, 17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

Source of subsequent assessment information: Mr. Eddie Seidensticker, U.S. Soil Conservation Service, 17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Nutria completely ate this area. Failure
- Seidensticker's Plot#9

184

Marsh Inventory No.: 275 **Information Rank (1-5):** 4 **State:** TX

Date Planted: 07/01/91

Area Planted (1acre = 0.4047 ha): 0.050 hectares

COE Permit No.: none **Date of COE Permit:**

Originator or Applicant for permit: Chambers-Liberty Nav. District

Location of the marsh.

State: TX **County:** Chambers **Latitude (N):** **Longitude (W):**

Bay and sub-bays: Galveston Bay; Trinity Bay

Directions to the site: On a dredged material deposition island along the west side of the Anahuac River Channel at marker 17.

Source of planting information: Mr. Eddie Seidensticker, U.S. Soil Conservation Service, 17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

Source of subsequent assessment information: Mr. Eddie Seidensticker, U.S. Soil Conservation Service, 17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Nutria and possibly grass carp ate this area completely by the winter. Failure.
- Seidensticker's Plot#10

Marsh Inventory No.: 274 **Information Rank (1-5):** 4 **State:** TX

Date Planted: 07/01/93

Area Planted (1 acre = 0.4047 ha): 0.050 hectares

COE Permit No.: none **Date of COE Permit:**

Originator or Applicant for permit: Various Private

Location of the marsh.

State: TX **County:** Chambers **Latitude (N):** **Longitude (W):**

Bay and sub-bays: Galveston Bay; Trinity Bay

Directions to the site: Along the east shoreline of the old Anahuac Barge channel halfway between Round Point and Ash Point.

Source of planting information: Mr. Eddie Seidensticker, U.S. Soil Conservation Service, 17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

Source of subsequent assessment information: Mr. Eddie Seidensticker, U.S. Soil Conservation Service, 17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- 70% shoot survival. 3 years later; free of damage and pests. Good.
- Seidensticker's Plot#8

Marsh Inventory No.: 280 **Information Rank (1-5):** 4 **State:** TX

Date Planted: 08/01/90

Area Planted (1acre = 0.4047 ha): 0.109 hectares

COE Permit No.: none **Date of COE Permit:**

Originator or Applicant for permit: Chambers-Liberty Nav. District

Location of the marsh.

State: TX **County:** Chambers **Latitude (N):** **Longitude (W):**

Bay and sub-bays: Galveston Bay; Trinity Bay

Directions to the site: Along the shore of a dredged material deposition island at marker 9 on the Anahuac River Channel, adjacent to plot 13.

Source of planting information: Mr. Eddie Seidensticker, U.S. Soil Conservation Service, 17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

Source of subsequent assessment information: Mr. Eddie Seidensticker, U.S. Soil Conservation Service, 17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- 100% shoot survival. 2 weeks later; plants look good. Plastic snow fence used for protection against waves and nutria. Excellent.
- Seidensticker's Plot#14

Marsh Inventory No.: 279 **Information Rank (1-5):** 4 **State:** TX

Date Planted: 08/01/90

Area Planted (1acre = 0.4047 ha): 0.010 hectares

COE Permit No.: none **Date of COE Permit:**

Originator or Applicant for permit: Chambers-Liberty Nav. District

Location of the marsh.

State: TX **County:** Chambers **Latitude (N):** **Longitude (W):**

Bay and sub-bays: Galveston Bay; Trinity Bay

Directions to the site: Along the shore of a dredged material deposition island at marker 9 on the Anahuac River Channel.

Source of planting information: Mr. Eddie Seidensticker, U.S. Soil Conservation Service, 17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

Source of subsequent assessment information: Mr. Eddie Seidensticker, U.S. Soil Conservation Service, 17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- 100% shoot survival. 2 weeks later; fishermen had cut the temporary barrier fence, but no noticeable damage and lots of aquatic life in the plot now. A plastic snow fence was used to dampen waves, and to keep nutria out. Excellent.

- Seidensticker's Plot#13

188

Marsh Inventory No.: 278 **Information Rank (1-5):** 4 **State:** TX

Date Planted: 08/01/90

Area Planted (1acre = 0.4047 ha): 0.017 hectares

COE Permit No.: none **Date of COE Permit:**

Originator or Applicant for permit: Chambers-Liberty Nav. District

Location of the marsh.

State: TX **County:** Chambers **Latitude (N):** **Longitude (W):**

Bay and sub-bays: Galveston Bay; Trinity Bay

Directions to the site: Next to the public boat ramp at Ft. Anahuac.

Source of planting information: Mr. Eddie Seidensticker, U.S. Soil Conservation Service, 17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

Source of subsequent assessment information: Mr. Eddie Seidensticker, U.S. Soil Conservation Service, 17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Bulkheaded off by accident and destroyed.
- Seidensticker's Plot#12

Marsh Inventory No.: 277 **Information Rank (1-5):** 4 **State:** TX

Date Planted: 08/01/90

Area Planted (1acre = 0.4047 ha): 0.109 hectares

COE Permit No.: none **Date of COE Permit:**

Originator or Applicant for permit: Chambers-Liberty Nav. District

Location of the marsh.

State: TX **County:** Chambers **Latitude (N):** **Longitude (W):**

Bay and sub-bays: Galveston Bay; Trinity Bay

Directions to the site: Along the north shore at the outlet of Long Island Bayou.

Source of planting information: Mr. Eddie Seidensticker, U.S. Soil Conservation Service, 17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

Source of subsequent assessment information: Mr. Eddie Seidensticker, U.S. Soil Conservation Service, 17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Completely eaten by nutria. Failure
- Seidensticker's Plot#11

190

Marsh Inventory No.: 709 **Information Rank (1-5):** 3 **State:** TX

Date Planted: 1983

Area Planted (1acre = 0.4047 ha): 2.833 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: COE

Location of the marsh.

State: TX **County:** Galveston **Latitude (N):** 29 20" 51.5" **Longitude (W):** 94 49' 26.8"

Bay and sub-bays: Galveston Bay

Directions to the site: Southern portion of little Pelican Island (the portion N of the GIWW)

Source of planting information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- This marsh was smothered by dredged material when a dike broke.
- Marsh name: L. Pelican Island.

Marsh Inventory No.: 711 **Information Rank (1-5):** 3 **State:** TX

Date Planted:

Area Planted (1acre = 0.4047 ha): 0.405 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: COE

Location of the marsh.

State: TX **County:** Galveston **Latitude (N):** 29 22' 00" **Longitude (W):** 94 53' 00"

Bay and sub-bays: Galveston Bay

Directions to the site: Along the SE facing shoreline of Snake Island; just a 0.25 mi W of Texas City Channel and harbor.

Source of planting information: Dr. James W. Webb, Texas A&M Univ. at Galveston, P.O. Box 1675, Galveston, TX 77553, (409)740-4542

Source of subsequent assessment information: Dr. James W. Webb, Texas A&M Univ. at Galveston, P.O. Box 1675, Galveston, TX 77553, (409)740-4542

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

-

-

192

Marsh Inventory No.: 697 **Information Rank (1-5):** 4 **State:** TX

Date Planted: 1992

Area Planted (1acre = 0.4047 ha): 2.024 hectares

COE Permit No.: none **Date of COE Permit:**

Originator or Applicant for permit: COE / NMFS

Location of the marsh.

State: TX **County:** Galveston **Latitude (N):** 29 21' 00" **Longitude (W):** 94 49' 35"

Bay and sub-bays: Galveston Bay

Directions to the site: Pelican Spit, along the S shore of the thin westward projection of the NW spit; first portion of two.

Source of planting information: Mr. Charles E. Belaire, Belaire Consulting, Inc., P.O. Box 741, Rockport, TX 78382, (512) 729-2948

Source of subsequent assessment information: Mr. Charles E. Belaire, Belaire Consulting, Inc., P.O. Box 741, Rockport, TX 78382, (512) 729-2948

References or reports:

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

-

-

Marsh Inventory No.: 696 **Information Rank (1-5):** 4 **State:** TX

Date Planted: 1992

Area Planted (1acre = 0.4047 ha): 1.214 hectares

COE Permit No.: none **Date of COE Permit:**

Originator or Applicant for permit: COE / NMFS

Location of the marsh.

State: TX **County:** Galveston **Latitude (N):** 29 21' 00" **Longitude (W):** 94 49' 36"

Bay and sub-bays: Galveston Bay

Directions to the site: Pelican Spit, along the S shore of the thin westward projection of the NW spit; second portion of two.

Source of planting information: Mr. Charles E. Beldaire, Beldaire Consulting, Inc., P.O. Box 741, Rockport, TX 78382, (512) 729-2948

Source of subsequent assessment information: Mr. Charles E. Beldaire, Beldaire Consulting, Inc., P.O. Box 741, Rockport, TX 78382, (512) 729-2948

References or reports:

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

-
-

194

Marsh Inventory No.: 708 **Information Rank (1-5):** 5 **State:** TX

Date Planted: 04/01/87

Area Planted (1acre = 0.4047 ha): 2.618 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: COE

Location of the marsh.

State: TX **County:** Galveston **Latitude (N):** 29 20' 47.3" **Longitude (W):** 94 49' 50.0"

Bay and sub-bays: Galveston Bay

Directions to the site: Pelican Spit, the extension N of the NW corner of Pelican Island

Source of planting information: Mr. Charles E. Belaire, Belaire Consulting, Inc., P.O. Box 741, Rockport, TX 78382, (512) 729-2948

Source of subsequent assessment information: Mr. G. Galbraith, Espey-Huston & Associates, Inc.

References or reports: Tom Minello, (409) 766-3606

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh: 75

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Yes. Fisheries and benthic organisms.

General comments about the marsh:

- Sprigs were planted on 1-m centers. Excellent survival and growth. Later studied for fisheries organisms. Also, trenched to study edge effects for fish and shellfish.
- Marsh name: Pelican Spit.

Marsh Inventory No.: 642 **Information Rank (1-5):** 3 **State:** TX

Date Planted: 1992

Area Planted (1acre = 0.4047 ha): 0.004 hectares

COE Permit No.: none **Date of COE Permit:**

Originator or Applicant for permit: GBF

Location of the marsh.

State: TX **County:** Galveston **Latitude (N):** 29 33' **Longitude (W):** 95 02'

Bay and sub-bays: Galveston Bay; Clear Lake

Directions to the site: Clear lake Shores-Jabot Bayou

Source of planting information: Ms. Linda Shead, Galveston Bay Foundation,
17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

Source of subsequent assessment information: Ms. Linda Shead, Galveston Bay
Foundation, 17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh: 0%

**Was there an assessment made of animal utilization of the marsh? If so,
which animals were surveyed, and when?**

General comments about the marsh:

- Eaten by nutria, complete failure.

Marsh Inventory No.: 713 **Information Rank (1-5):** 3 **State:** TX

Date Planted: 1977

Area Planted (1acre = 0.4047 ha): 1.214 hectares

COE Permit No.: none **Date of COE Permit:**

Originator or Applicant for permit: COE

Location of the marsh.

State: TX **County:** Galveston **Latitude (N):** 29 25' 10" **Longitude (W):** 94 43' 57"

Bay and sub-bays: Galveston Bay; East Bay

Directions to the site: On the N shore of Bolivar Peninsula, about 5 mi E of the Houston Ship Channel

Source of planting information: Dr. James W. Webb, Texas A&M Univ. at Galveston, P.O. Box 1675, Galveston, TX 77553, (409)740-4542

Source of subsequent assessment information: Dr. James W. Webb, Texas A&M Univ. at Galveston, P.O. Box 1675, Galveston, TX 77553, (409)740-4542

References or reports: Webb, 1977; Allen et al., 1978.

Other species planted in addition to *Spartina alterniflora* : *Spartina patens*

Percent vegetative cover attained by the planted marsh: 50

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Transplanted *S. alterniflora* was tried at different elevations and under different fertilizer treatments; half the plots were protected behind a sandbag breakwater. This was part of a 7.3 ha study of marsh and upland habitat development.

Marsh Inventory No.: 710 **Information Rank (1-5):** 3 **State:** TX

Date Planted: 1984

Area Planted (1acre = 0.4047 ha): 0.202 hectares

COE Permit No.: none **Date of COE Permit:**

Originator or Applicant for permit: COE

Location of the marsh.

State: TX **County:** Galveston **Latitude (N):** 29 24' 43" **Longitude (W):** 94 44' 37"

Bay and sub-bays: Galveston Bay; East Bay

Directions to the site: On the N shore of Bolivar Peninsula, about 4 mi E of the Houston Ship Channel

Source of planting information: Dr. James W. Webb, Texas A&M Univ. at Galveston, P.O. Box 1675, Galveston, TX 77553, (409)740-4542

Source of subsequent assessment information: Dr. James W. Webb, Texas A&M Univ. at Galveston, P.O. Box 1675, Galveston, TX 77553, (409)740-4542

References or reports: Allen and Webb, 1993

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh: 50

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Two breakwaters were used, a floating tire and a fixed tire. Both worked well to start and 50% cover was obtained. But when the fixed tire breakwater broke, its cordgrass was destroyed in about 5 yrs. The floating tire marsh expanded beyond 100%.

Marsh Inventory No.: 712 **Information Rank (1-5):** 3 **State:** TX

Date Planted: 1985

Area Planted (1acre = 0.4047 ha): 0.121 hectares

COE Permit No.: none **Date of COE Permit:**

Originator or Applicant for permit: COE

Location of the marsh.

State: TX **County:** Galveston **Latitude (N):** 29 24' 48" **Longitude (W):** 94 44' 24"

Bay and sub-bays: Galveston Bay; East Bay

Directions to the site: On the N shore of Bolivar Peninsula, about 4 mi E of the Houston Ship Channel

Source of planting information: Dr. James W. Webb, Texas A&M Univ. at Galveston, P.O. Box 1675, Galveston, TX 77553, (409)740-4542

Source of subsequent assessment information: Dr. James W. Webb, Texas A&M Univ. at Galveston, P.O. Box 1675, Galveston, TX 77553, (409)740-4542

References or reports: Allen et al., 1986

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh: 50

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Transplants were protected for awhile by a breakwater made of tires.

Marsh Inventory No.: 226 **Information Rank (1-5):** 3 **State:** TX

Date Planted:

Area Planted (1acre = 0.4047 ha): 1.214 hectares

COE Permit No.: 17072 **Date of COE Permit:** 03/13/85

Originator or Applicant for permit: R. G. Parker

Location of the marsh.

State: TX **County:** Galveston **Latitude (N):** 29 21' 00" **Longitude (W):** 94 53' 40"

Bay and sub-bays: Galveston Bay; Swan Lake.

Directions to the site: Along N half of E margin of Swan Lake;

Source of planting information: Dr. James W. Webb, Texas A&M Univ. at Galveston, P.O. Box 1675, Galveston, TX 77553, (409)740-4542

Source of subsequent assessment information: Dr. James W. Webb, Texas A&M Univ. at Galveston, P.O. Box 1675, Galveston, TX 77553, (409)740-4542

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

-
-

200

Marsh Inventory No.: 636 **Information Rank (1-5):** 4 **State:** TX

Date Planted: 1992

Area Planted (1acre = 0.4047 ha): 0.040 hectares

COE Permit No.: none **Date of COE Permit:**

Originator or Applicant for permit: GBF - SCS

Location of the marsh.

State: TX **County:** Galveston **Latitude (N):** 29 12' 30" **Longitude (W):** 94 57' 30"

Bay and sub-bays: Galveston Bay; West Bay

Directions to the site: Along the SW shore of Lake Como on Galveston Island near Pirates housing development on the bay side.

Source of planting information: Ms. Linda Shead, Galveston Bay Foundation, 17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

Source of subsequent assessment information: Ms. Linda Shead, Galveston Bay Foundation, 17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh: 80%

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Planted on 0.5 m centers.

Marsh Inventory No.: 706 **Information Rank (1-5):** 3 **State:** TX

Date Planted:

Area Planted (1acre = 0.4047 ha): 0.016 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: COE ?

Location of the marsh.

State: TX **County:** Galveston **Latitude (N):** 29 13' 50" **Longitude (W):** 95 01' 30"

Bay and sub-bays: Galveston Bay; West Bay.

Directions to the site: Along the N shore of West Bay just S of Carancahua Lake.

Source of planting information: Dr. James W. Webb, Texas A&M Univ. at Galveston, P.O. Box 1675, Galveston, TX 77553, (409)740-4542

Source of subsequent assessment information: Dr. James W. Webb, Texas A&M Univ. at Galveston, P.O. Box 1675, Galveston, TX 77553, (409)740-4542

References or reports:

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Fringe marsh subject to 2 mi fetch across the bay, but somewhat protected by the Carancahua Reef in the bay.

202

Marsh Inventory No.: 718 **Information Rank (1-5):** 3 **State:** TX

Date Planted: 04/01/93

Area Planted (1acre = 0.4047 ha): 12.546 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: City of Hitchcock ?

Location of the marsh.

State: TX **County:** Galveston **Latitude (N):** 29 20' 30" **Longitude (W):** 94 58' 20"

Bay and sub-bays: Galveston Bay; West Bay; Highland Bayou

Directions to the site: About 3 miles up the bayou, in the city of Hitchcock

Source of planting information: Dr. James W. Webb, Texas A&M Univ. at Galveston, P.O. Box 1675, Galveston, TX 77553, (409)740-4542

Source of subsequent assessment information: Dr. James W. Webb, Texas A&M Univ. at Galveston, P.O. Box 1675, Galveston, TX 77553, (409)740-4542

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

Marsh Inventory No.: 719 **Information Rank (1-5):** 3 **State:** TX

Date Planted:

Area Planted (1acre = 0.4047 ha): 0.004 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: COE

Location of the marsh.

State: TX **County:** Galveston **Latitude (N):** 29 17' 00" **Longitude (W):** 94 54' 18"

Bay and sub-bays: Galveston Bay; West Bay; North Deer Island

Directions to the site: Along the E side of North Deer island; transplants were planted through slits in erosion control coconut fibre mats.

Source of planting information: Dr. James W. Webb, Texas A&M Univ. at Galveston, P.O. Box 1675, Galveston, TX 77553, (409)740-4542

Source of subsequent assessment information: Dr. James W. Webb, Texas A&M Univ. at Galveston, P.O. Box 1675, Galveston, TX 77553, (409)740-4542

References or reports:

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Fringe marsh subject to barge traffic in the GIWW that was only 0.2 mi away.
-

204

Marsh Inventory No.: 695 **Information Rank (1-5):** 3 **State:** TX

Date Planted:

Area Planted (1 acre = 0.4047 ha): 0.299 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: COE ?

Location of the marsh.

State: TX **County:** Galveston **Latitude (N):** 29 16' 40" **Longitude (W):** 94 51' 00"

Bay and sub-bays: Galveston Bay; West Bay; Offats Bayou; Lake Madelaine channel.

Directions to the site: Along the W side of the channel, protected by a floating tire wave breaking device.

Source of planting information: Dr. James W. Webb, Texas A&M Univ. at Galveston, P.O. Box 1675, Galveston, TX 77553, (409)740-4542

Source of subsequent assessment information: Dr. Thomas Minello, National Marine Fisheries Service, 4700 Ave. U, Galveston, TX 77551, (409) 766-3506

References or reports:

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Yes. Fisheries and benthic organisms.

General comments about the marsh:

- Fringe marsh subject to pleasure boats' low wake waves.

Marsh Inventory No.: 707 **Information Rank (1-5):** 3 **State:** TX

Date Planted: 1984

Area Planted (1acre = 0.4047 ha): 0.324 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Moody Foundation

Location of the marsh.

State: TX **County:** Galveston **Latitude (N):** 29 16' 25.5" **Longitude (W):** 94 50' 57.3"

Bay and sub-bays: Galveston Bay; West Bay; Offatts Bayou; Palm Beach.

Directions to the site: Shoreline along Offatts Bayou in front of Palm Beach.

Source of planting information:

Source of subsequent assessment information: Dr. Geoffrey Matthews, National Marine Fisheries Service, 4700 Ave. U, Galveston, TX 77551, (409) 766-3532

References or reports:

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh: 50%

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Marsh is becoming established at Palm Beach.

206

Marsh Inventory No.: 722 **Information Rank (1-5):** 3 **State:** TX

Date Planted: 1990

Area Planted (1acre = 0.4047 ha): 0.405 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: TX DOT ?

Location of the marsh.

State: TX **County:** Harris **Latitude (N):** 29 33' 25" **Longitude (W):** 95 01' 20"

Bay and sub-bays: Galveston Bay

Directions to the site: Just NE of the Clear Lake channel causeway, in a tidal pond.

Source of planting information: Dr. James W. Webb, Texas A&M Univ. at Galveston, P.O. Box 1675, Galveston, TX 77553, (409)740-4542

Source of subsequent assessment information: Dr. James W. Webb, Texas A&M Univ. at Galveston, P.O. Box 1675, Galveston, TX 77553, (409)740-4542

References or reports:

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh: 70

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- This marsh has its good and bad years, depending on circulation of water to the pond.

Marsh Inventory No.: 634 **Information Rank (1-5):** 3 **State:** TX

Date Planted: 1992

Area Planted (1acre = 0.4047 ha): 0.040 hectares

COE Permit No.: none **Date of COE Permit:**

Originator or Applicant for permit: GBF - SCS

Location of the marsh.

State: TX **County:** Harris **Latitude (N):** 29 40' **Longitude (W):** 94 59'

Bay and sub-bays: Galveston Bay

Directions to the site: Morgan's Point in front of Mansion

Source of planting information: Ms. Linda Shead, Galveston Bay Foundation, 17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

Source of subsequent assessment information: Ms. Linda Shead, Galveston Bay Foundation, 17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh: 80-0%

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Stand was doing very well until this springs extended high tides that completely killed the *Spartina*. They replanted in July 1993 and are doing more in August 1993.

208

Marsh Inventory No.: 721 **Information Rank (1-5):** 3 **State:** TX

Date Planted: 1993

Area Planted (1 acre = 0.4047 ha): 0.040 hectares

COE Permit No.: none **Date of COE Permit:**

Originator or Applicant for permit: GBF

Location of the marsh.

State: TX **County:** Harris **Latitude (N):** 29 38' **Longitude (W):** 95 01'

Bay and sub-bays: Galveston Bay

Directions to the site: Little Cedar Bayou-South bank of bayou about 0.7 km up into the bayou.

Source of planting information: Ms. Linda Shead, Galveston Bay Foundation, 17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

Source of subsequent assessment information: Ms. Linda Shead, Galveston Bay Foundation, 17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Planted on 0.5-m centers.

Marsh Inventory No.: 635 **Information Rank (1-5):** 4 **State:** TX

Date Planted: 1991

Area Planted (1acre = 0.4047 ha): 0.040 hectares

COE Permit No.: none **Date of COE Permit:**

Originator or Applicant for permit: GBF

Location of the marsh.

State: TX **County:** Harris **Latitude (N):** 29 38' **Longitude (W):** 95 01'

Bay and sub-bays: Galveston Bay

Directions to the site: Little Cedar Bayou-South bank of bayou about 0.5 km up into the bayou.

Source of planting information: Ms. Linda Shead, Galveston Bay Foundation, 17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

Source of subsequent assessment information: Ms. Linda Shead, Galveston Bay Foundation, 17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh: 75-90%

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Planted on 0.5-m centers.
-

210

Marsh Inventory No.: 720 **Information Rank (1-5):** 4 **State:** TX

Date Planted: 1992

Area Planted (1acre = 0.4047 ha): 0.040 hectares

COE Permit No.: none **Date of COE Permit:**

Originator or Applicant for permit: GBF

Location of the marsh.

State: TX **County:** Harris **Latitude (N):** 29 38' **Longitude (W):** 95 01'

Bay and sub-bays: Galveston Bay

Directions to the site: Little Cedar Bayou-South bank of bayou about 0.5 km up into the bayou.

Source of planting information: Ms. Linda Shead, Galveston Bay Foundation, 17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

Source of subsequent assessment information: Ms. Linda Shead, Galveston Bay Foundation, 17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh: 75-90%

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Planted on 0.5-m centers.
-

Marsh Inventory No.: 637 **Information Rank (1-5):** 3 **State:** TX

Date Planted: 1990

Area Planted (1acre = 0.4047 ha): 0.040 hectares

COE Permit No.: none **Date of COE Permit:**

Originator or Applicant for permit: GBF - SCS

Location of the marsh.

State: TX **County:** Harris **Latitude (N):** 29 48' 55" **Longitude (W):** 95 05'

Bay and sub-bays: Galveston Bay; San Jacinto River, White Lake

Directions to the site: San Jacinto River Railroad Island

Source of planting information: Ms. Linda Shead, Galveston Bay Foundation, 17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

Source of subsequent assessment information: Ms. Linda Shead, Galveston Bay Foundation, 17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh: 80% cover in 40% of the marsh

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Planted on 0.5-m centers. Predations by grass carp and swamp rabbits.

212

Marsh Inventory No.: 724 **Information Rank (1-5):** 3 **State:** TX

Date Planted: 1991

Area Planted (1acre = 0.4047 ha): 0.040 hectares

COE Permit No.: none **Date of COE Permit:**

Originator or Applicant for permit: GBF - SCS

Location of the marsh.

State: TX **County:** Harris **Latitude (N):** 29 48' 55" **Longitude (W):** 95 05'

Bay and sub-bays: Galveston Bay; San Jacinto River, White Lake

Directions to the site: San Jacinto River Railroad Island, northeast margin of the island.

Source of planting information: Ms. Linda Shead, Galveston Bay Foundation, 17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

Source of subsequent assessment information: Ms. Linda Shead, Galveston Bay Foundation, 17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh: 80% cover in 40% of the marsh

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Planted on 0.5-m centers. Predations by grass carp and swamp rabbits.

Marsh Inventory No.: 638 **Information Rank (1-5):** 3 **State:** TX

Date Planted: 1991

Area Planted (1 acre = 0.4047 ha): 0.040 hectares

COE Permit No.: none **Date of COE Permit:**

Originator or Applicant for permit: GBF - SCS

Location of the marsh.

State: TX **County:** Harris **Latitude (N):** 29 48' 29" **Longitude (W):** 95 04'

Bay and sub-bays: Galveston Bay; San Jacinto River, White Lake

Directions to the site: San Jacinto River-Little Island, Phase 1

Source of planting information: Ms. Linda Shead, Galveston Bay Foundation, 17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

Source of subsequent assessment information: Ms. Linda Shead, Galveston Bay Foundation, 17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh: 95%

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Planted on 0.5-m centers.

214

Marsh Inventory No.: 639 **Information Rank (1-5):** 3 **State:** TX

Date Planted: 1992

Area Planted (1acre = 0.4047 ha): 0.040 hectares

COE Permit No.: none **Date of COE Permit:**

Originator or Applicant for permit: GBF - SCS

Location of the marsh.

State: TX **County:** Harris **Latitude (N):** 29 48' 29" **Longitude (W):** 95 04'

Bay and sub-bays: Galveston Bay; San Jacinto River, White Lake

Directions to the site: San Jacinto River-Little Island, Phase 2: added on to the west.

Source of planting information: Ms. Linda Shead, Galveston Bay Foundation, 17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

Source of subsequent assessment information: Ms. Linda Shead, Galveston Bay Foundation, 17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh: 65%

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

-
-

Marsh Inventory No.: 214 **Information Rank (1-5):** 3 **State:** TX

Date Planted: 1992

Area Planted (1acre = 0.4047 ha): 0.040 hectares

COE Permit No.: none **Date of COE Permit:**

Originator or Applicant for permit: GBF - SCS

Location of the marsh.

State: TX **County:** Harris **Latitude (N):** 29 48' 55" **Longitude (W):** 95 05'

Bay and sub-bays: Galveston Bay; San Jacinto River, White Lake

Directions to the site: San Jacinto River Railroad Island, expanded to south.

Source of planting information: Ms. Linda Shead, Galveston Bay Foundation, 17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

Source of subsequent assessment information: Ms. Linda Shead, Galveston Bay Foundation, 17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh: 80% cover in 40% of the marsh

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Planted on 0.5-m centers. Predations by grass carp and swamp rabbits.

216

Marsh Inventory No.: 723 **Information Rank (1-5):** 3 **State:** TX

Date Planted: 1993

Area Planted (1acre = 0.4047 ha): 0.040 hectares

COE Permit No.: none **Date of COE Permit:**

Originator or Applicant for permit: GBF - SCS

Location of the marsh.

State: TX **County:** Harris **Latitude (N):** 29 48' 55" **Longitude (W):** 95 05'

Bay and sub-bays: Galveston Bay; San Jacinto River, White Lake

Directions to the site: San Jacinto River Railroad Island, expanded to south and northwest.

Source of planting information: Ms. Linda Shead, Galveston Bay Foundation, 17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

Source of subsequent assessment information: Ms. Linda Shead, Galveston Bay Foundation, 17324-A Hwy. 3, Webster, TX 77598, (713) 332-3381

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh: 80% cover in 40% of the marsh

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Planted on 0.5-m centers. Predations by grass carp and swamp rabbits.

Marsh Inventory No.: 694 **Information Rank (1-5):** 3 **State:** TX

Date Planted: 1990

Area Planted (1acre = 0.4047 ha): 1.012 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Transco

Location of the marsh.

State: TX **County:** Matagorda **Latitude (N):** 28 24' 30" **Longitude (W):** 96 22' 00"

Bay and sub-bays: Matagorda Bay

Directions to the site: Along the N shore of Decros Point at the S tip of Matagorda Peninsula, beginning about 2 km NE of Pass Cavallo.

Source of planting information: Mr. Charles E. Belaire, Belaire Consulting, Inc., P.O. Box 741, Rockport, TX 78382, (512) 729-2948

Source of subsequent assessment information: Mr. Charles E. Belaire, Belaire Consulting, Inc., P.O. Box 741, Rockport, TX 78382, (512) 729-2948

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh: 20

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Partial Success (>10% cover). Wide open to the N, fetch greater than 10 km.

Marsh Inventory No.: 716 **Information Rank (1-5):** 5 **State:** TX

Date Planted: 07/01/84

Area Planted (1acre = 0.4047 ha): 8.000 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit:

Location of the marsh.

State: TX **County:** Matagorda **Latitude (N):** 28 39' 00" **Longitude (W):** 96 03' 00"

Bay and sub-bays: Matagorda Bay; East Matagorda Bay

Directions to the site: Along the S-facing shoreline of a dredged material island on the NE margin of the bay.

Source of planting information: Dr. Thomas Minello, National Marine Fisheries Service, 4700 Ave. U, Galveston, TX 77551, (409) 766-3506

Source of subsequent assessment information: (Minello & Zimmerman, 1992)

References or reports: (Minello & Zimmerman, 1992)

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Yes. Fisheries and benthic organisms.

General comments about the marsh:

-
-

Marsh Inventory No.: 245 **Information Rank (1-5):** 3 **State:** TX

Date Planted:

Area Planted (1acre = 0.4047 ha): 0.040 hectares

COE Permit No.: 16075 **Date of COE Permit:**

Originator or Applicant for permit: Latitude 27, Inc. (Island Moorings Subdiv.)

Location of the marsh.

State: TX **County:** Nueces **Latitude (N):** **Longitude (W):**

Bay and sub-bays: Corpus Christi Bay

Directions to the site: W. side of Mustang Isl., S. of Port Aransas

Source of planting information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

220

Marsh Inventory No.: 240 **Information Rank (1-5):** 3 **State:** TX

Date Planted:

Area Planted (1acre = 0.4047 ha): 0.004 hectares

COE Permit No.: 14454(02) **Date of COE Permit:**

Originator or Applicant for permit: Mustang Beach Devel. Corp. (Sunny Castor)

Location of the marsh.

State: TX **County:** Nueces **Latitude (N):** 27 49' 53" **Longitude (W):** 97 06' 40"

Bay and sub-bays: Corpus Christi Bay

Directions to the site: W. side of Mustang Isl, S. of Port Aransas; 400 ft SW from the canal to a mud flat.

Source of planting information: Mr. Lloyd Mullins, Texas General Land Office, 111 W. Wilson, 2nd floor Naylor Bldg., Aransas Pass, TX 78336, (512) 758-7228

Source of subsequent assessment information: Ms. Lynda Kahn, Texas General Land Office, 111 W. Wilson, 2nd floor Naylor Bldg., Aransas Pass, TX 78336, (512) 758-7228

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh: 100

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Impossible to distinguish from the natural marsh.

Marsh Inventory No.: 238 **Information Rank (1-5):** 3 **State:** TX

Date Planted:

Area Planted (1 acre = 0.4047 ha): 0.008 hectares

COE Permit No.: 13625 **Date of COE Permit:**

Originator or Applicant for permit: R.E. Jenkins

Location of the marsh.

State: TX **County:** Nueces **Latitude (N):** **Longitude (W):**

Bay and sub-bays: Corpus Christi Bay

Directions to the site: Port Aransas Municipal Boat Harbor

Source of planting information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

-

-

222

Marsh Inventory No.: 191 **Information Rank (1-5):** 3 **State:** TX

Date Planted:

Area Planted (1acre = 0.4047 ha): 0.154 hectares

COE Permit No.: 18175 **Date of COE Permit:** 10/26/87

Originator or Applicant for permit: TX State Aquarium Association

Location of the marsh.

State: TX **County:** Nueces **Latitude (N):** 27 51' 00" **Longitude (W):** 97 21' 20"

Bay and sub-bays: Corpus Christi Bay

Directions to the site:

Source of planting information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

Marsh Inventory No.: 665 **Information Rank (1-5):** 4 **State:** TX

Date Planted: 1981

Area Planted (1acre = 0.4047 ha): 0.040 hectares

COE Permit No.: 14454(06) **Date of COE Permit:**

Originator or Applicant for permit: Mustang Beach Ltd.

Location of the marsh.

State: TX **County:** Nueces **Latitude (N):** 27 48' 10" **Longitude (W):** 97 05' 45"

Bay and sub-bays: Corpus Christi Bay; Elizabeth Cove

Directions to the site: Mustang Island, Mustang Beach, Tx. 400 sq. ft. from end of canal sw to a mud flat.

Source of planting information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

-
-

224

Marsh Inventory No.: 683 **Information Rank (1-5):** 3 **State:** TX

Date Planted: 1990

Area Planted (1acre = 0.4047 ha): 12.141 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: COE

Location of the marsh.

State: TX **County:** Nueces **Latitude (N):** 27 52' 00" **Longitude (W):** 97 27' 10"

Bay and sub-bays: Corpus Christi Bay; Nueces Bay

Directions to the site: West end of Nueces Bay.

Source of planting information: Mr. Charles E. Belaire, Belaire Consulting, Inc., P.O. Box 741, Rockport, TX 78382, (512) 729-2948

Source of subsequent assessment information: Mr. Charles E. Belaire, Belaire Consulting, Inc., P.O. Box 741, Rockport, TX 78382, (512) 729-2948

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh: 2

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Failure (<5% cover). Scrape-down operation removed too much substrate so the area elevation was too low to support *Spartina*. Now *Ruppia* is growing, and a plover is using the area, probably no corrective measures will be done.

Marsh Inventory No.: 682 **Information Rank (1-5):** 3 **State:** TX

Date Planted: 1992

Area Planted (1acre = 0.4047 ha): 0.040 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Charlie Craveas

Location of the marsh.

State: TX **County:** Nueces **Latitude (N):** 27 38' 15" **Longitude (W):** 97 14' 46"

Bay and sub-bays: Laguna Madre

Directions to the site: S shoreline of the John F. Kennedy Causeway, 0.2 mi W of the GIWW bridge

Source of planting information: Mr. Charles E. Belaire, Belaire Consulting, Inc., P.O. Box 741, Rockport, TX 78382, (512) 729-2948

Source of subsequent assessment information: Mr. Charles E. Belaire, Belaire Consulting, Inc., P.O. Box 741, Rockport, TX 78382, (512) 729-2948

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh: 2

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Failure (<10% cover).
-

226

Marsh Inventory No.: 247 **Information Rank (1-5):** 3 **State:** TX

Date Planted:

Area Planted (1acre = 0.4047 ha): 0.004 hectares

COE Permit No.: 16889 **Date of COE Permit:**

Originator or Applicant for permit: G. Goodman

Location of the marsh.

State: TX **County:** Nueces **Latitude (N):** **Longitude (W):**

Bay and sub-bays: Laguna Madre; Upper Laguna Madre

Directions to the site: Under and S. of the JFK Causeway bridge over the GIWW

Source of planting information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

Marsh Inventory No.: 693 **Information Rank (1-5):** 4 **State:** TX

Date Planted: 1992

Area Planted (1acre = 0.4047 ha): 0.040 hectares

COE Permit No.: none **Date of COE Permit:**

Originator or Applicant for permit: Refugio County

Location of the marsh.

State: TX **County:** Refugio **Latitude (N):** 28 23' 30" **Longitude (W):** 96 50' 20"

Bay and sub-bays: San Antonio Bay

Directions to the site: Bayfront at Austwell, TX; NW of boat docks.

Source of planting information: Mr. Charles E. Belaire, Belaire Consulting, Inc., P.O. Box 741, Rockport, TX 78382, (512) 729-2948

Source of subsequent assessment information: Mr. Charles E. Belaire, Belaire Consulting, Inc., P.O. Box 741, Rockport, TX 78382, (512) 729-2948

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh: 75

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- Very successful (>70% cover).
-

228

Marsh Inventory No.: 237 **Information Rank (1-5):** 3 **State:** TX

Date Planted:

Area Planted (1acre = 0.4047 ha): 0.040 hectares

COE Permit No.: 12825 **Date of COE Permit:**

Originator or Applicant for permit: Ingelside Offshore Services

Location of the marsh.

State: TX **County:** San Patricio **Latitude (N):** **Longitude (W):**

Bay and sub-bays: Corpus Christi Bay

Directions to the site: Jewel Fulton Canal, at Ingleside

Source of planting information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

-

-

Marsh Inventory No.: 725 **Information Rank (1-5):** 5 **State:** VA

Date Planted: 10/01/84

Area Planted (1acre = 0.4047 ha): 2.833 hectares

COE Permit No.: ? **Date of COE Permit:** 1984

Originator or Applicant for permit: Navy

Location of the marsh.

State: VA **County:** **Latitude (N):** 36 57' 45" **Longitude (W):** 76 16' 12"

Bay and sub-bays: Chesapeake Bay; Hampton Roads; Willoughby Bay

Directions to the site: Just WNW of the City of Norfolk's Visitor Center.

Source of planting information: Mr. Carvel Blair, Norfolk Wetlands Board

Source of subsequent assessment information: Mr. Carvel Blair, Norfolk Wetlands Board

References or reports: (Blair, 1991)

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh: 64% as of 1989

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when? Yes. Feigenbaum & Swift (1989) studied fish and invertebrates; Blair (1989) studied the birds.

General comments about the marsh:

- *S. alterniflora* was planted on 2-ft centers. Marsh shape: rectangular; innudation channels form an H with the cross-bar leading to the culverts that connect to the bay.
- Non-quantitative monitoring showed presence of benthic invertebrates, fish and birds. Animal appeared to be as abundant in the created marsh as in the nearby natural marsh.

Marsh Inventory No.: 287 **Information Rank (1-5):** 3 **State:** VA

Date Planted: 04/01/75

Area Planted (1acre = 0.4047 ha): 2.833 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Norfolk District, Army COE

Location of the marsh.

State: VA **County:** Accomack **Latitude (N):** 37 28' 40" **Longitude (W):** 75 44' 30"

Bay and sub-bays: Atlantic Ocean; Quinby Inlet; North Inlet; Cunjer Channel; Sloop Channel.

Directions to the site: On the W shore of an island located at Bouy 171 of Sloop Channel, and about 9 km SE of Willis Wharf.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information: (Garbisch, Woller and McCallum, 1976)

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- *S. alterniflora* was planted on 2-ft centers. Marsh shape: irregular. Built by Environmental Concern, Inc. COE Contract # DACW65-74-C-0062. These were dredge disposal sites.

- Marsh name: Sloop Channel

Marsh Inventory No.: 286 **Information Rank (1-5):** 3 **State:** VA

Date Planted: 04/01/75

Area Planted (1acre = 0.4047 ha): 1.214 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Norfolk District, Army COE

Location of the marsh.

State: VA **County:** Accomack **Latitude (N):** 37 39' 20" **Longitude (W):** 75 38' 06"

Bay and sub-bays: Atlantic Ocean; Wachapreague Inlet; Burtons Bay.

Directions to the site: On the SW-facing shore at the N end of Burtons Bay, about 9 km NE of Wachapreague.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information: (Garbisch, Woller and McCallum, 1976)

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- *S. alterniflora* was planted on 2-ft centers. Marsh shape: irregular. Built by Environmental Concern, Inc. COE Contract # DACW65-74-C-0062. These were dredge disposal sites.

- Marsh name: Burtons Bay

Marsh Inventory No.: 336 **Information Rank (1-5):** 3 **State:** VA

Date Planted: 06/01/87

Area Planted (1acre = 0.4047 ha): 0.283 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: Virginia DOT

Location of the marsh.

State: VA **County:** Accomack **Latitude (N):** 37 56' 00?" **Longitude (W):** 75 37' 00?"

Bay and sub-bays: Chesapeake Bay; Pocomoke Sound; Powell Bay; Holdens Creek

Directions to the site: Scrape-down area where old Rte. 702 crossed creek near and E of intersection of 702 and 701, north of Jenkins Bridge.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* :

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- *S. alterniflora* was planted on 2-ft centers. Marsh shape: linear. Built by Environmental Concern, Inc.

- Marsh name: Lance Eller Inc.

Marsh Inventory No.: 326 **Information Rank (1-5):** 3 **State:** VA

Date Planted: 07/01/87

Area Planted (1acre = 0.4047 ha): 0.890 hectares

COE Permit No.: **Date of COE Permit:**

Originator or Applicant for permit: City of Hampton, VA

Location of the marsh.

State: VA **County:** City of Hampton **Latitude (N):** 37 03' 45" **Longitude (W):** 76 17' 05"

Bay and sub-bays: Chesapeake Bay; Horseshoe Shoal; Long Creek

Directions to the site: E-side of Long Creek, east of Salt Ponds Rd. and about 8 km NE of Hampton, VA.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- *S. alterniflora* was planted on 2-ft centers. Built by Environmental Concern, Inc. Permit # 2 SD OX2 1 002355.

- Marsh name: Hampton, VA.

234

Marsh Inventory No.: 345 **Information Rank (1-5):** 3 **State:** VA

Date Planted: 06/01/84

Area Planted (1 acre = 0.4047 ha): 0.648 hectares

COE Permit No.: 84-0166-12 **Date of COE Permit:** 1984

Originator or Applicant for permit: Vidco of Virginia

Location of the marsh.

State: VA **County:** Isle of Wight **Latitude (N):** 36 48' 45" **Longitude (W):** 76 17' 20"

Bay and sub-bays: Chesapeake Bay; Hampton Roads; Elizabeth River; Bell's Mill Creek

Directions to the site: Residential development mitigation area on Bell's Mill Creek and Cedar Road (Rte. 168?), Chesapeake, VA.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* : *Spartina cynosuroides*, *Spartina patens*

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- *S. alterniflora* was planted on 2-ft centers. Marsh shape: rectangular. Built by Environmental Concern, Inc.
- Marsh name: Vidco. Permit # 84-0166-12.

Marsh Inventory No.: 312 **Information Rank (1-5):** 3 **State:** VA

Date Planted: 07/01/84

Area Planted (1acre = 0.4047 ha): 1.295 hectares

COE Permit No.: NADOP-P79-0316-06 **Date of COE Permit:**

Originator or Applicant for permit: Town of Tangier, VA.

Location of the marsh.

State: VA **County:** James City **Latitude (N):** 37 12' 50" **Longitude (W):** 76 43' 30"

Bay and sub-bays: Chesapeake Bay; James River; The Thorofare.

Directions to the site: Jamestown National Historic Site shoreline, along the N 1.5 miles of shoreline on The Thorofare, between College Creek and Mill Creek.

Source of planting information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Source of subsequent assessment information:

References or reports:

Other species planted in addition to *Spartina alterniflora* : none

Percent vegetative cover attained by the planted marsh:

Was there an assessment made of animal utilization of the marsh? If so, which animals were surveyed, and when?

General comments about the marsh:

- *S. alterniflora* was planted on 2-ft centers. Marsh shape: shoreline. Built by Environ. Concern Inc. EPA Project # C-510527-03.
- Marsh name: Tangier Island Sewer.

Marshes with Minimal Information

Marsh Inventory No.: 1 **COE Permit No.:** AL84-00128-L **Size (ha):** 0.003
State: AL **County:** Baldwin **Bay :** Cotton Bayou
Source of information:
Comments: .

Marsh Inventory No.: 4 **COE Permit No.:** AL90-00065-B **Size (ha):** 0.004
State: AL **County:** Baldwin **Bay :** Perdido Bay
Source of information:
Comments: .

Marsh Inventory No.: 3 **COE Permit No.:** AL84-00109-A **Size (ha):** 0.020
State: AL **County:** Baldwin **Bay :** Perdido Bay; Arnica Bay
Source of information:
Comments: .

Marsh Inventory No.: 658 **COE Permit No.:** AL87-01424-U **Size (ha):** 18.070
State: AL **County:** Mobile **Bay :** Mississippi Sound, West Fowl River Bay
Source of information: Dr. Barry A. Vittor, Barry A. Vittor and Ass., Inc.,
Environmental Research & Consulting, 8060 Cottage Hill Road, Mobile, AL 36695, (205)
633-6100
Comments: Marsh shape: protected canal system. Planting density of *S. alterniflora* : 4
stems per sq. m. 5 site evaluations were performed from 1988-1989.. Marsh name: West
Fowl River Marsh.

Marsh Inventory No.: 2 **COE Permit No.:** AL84-00039-L-AF **Size (ha):** 0.008
State: AL **County:** Mobile **Bay :** Mississippi Sound; Bayou Coden
Source of information:
Comments: .

Marsh Inventory No.: 184 **COE Permit No.:** **Size (ha):** 0.350
State: AL **County:** **Bay :** Mississippi Sound; Portersville Bay
Source of information: (Allen, Shirley and Webb, 1982) (Allen, Shirley & Webb,
1982)

Comments: This was a test of new design breakwaters. The floating tire breakwater
successfully withstood the wave energy, but caused sediment accretion and smothered the
sprigs planted in June 1985. *S. alterniflora* was planted again in Aug '85. . Survival of
the 1985 planting was about 30-40% behind the breakwaters as of Nov '85.

Marsh Inventory No.: 673 **COE Permit No.:** **Size (ha):** 1.550

State: AL **County:** **Bay :** Mobile Bay

Source of information: (Allen and Webb, 1982) (Allen & Webb, 1982)

Comments: This was a test of new design breakwaters. The floating tire breakwater
successfully withstood the wave energy and protected the *S. alterniflora* planting..

Marsh Inventory No.: 655 **COE Permit No.:** **Size (ha):** 0.081

State: CT **County:** **Bay :** Mianus River

Source of information: Mr. Matthew J. Popp or Ms. Judith A. Slayback,
Environmental Design Associates, P.C., P.O. Box 247, 78 Danbury Road, Wilton, CT
06897, (203) 762-8020

Comments: Some problems with predation by Canadian geese. Initially the new plants showed vigorous growth..

Marsh Inventory No.: 656 **COE Permit No.:** **Size (ha):** 0.040

State: CT **County:** **Bay :**

Source of information: Mr. Matthew J. Popp or Ms. Judith A. Slayback,
Environmental Design Associates, P.C., P.O. Box 247, 78 Danbury Road, Wilton, CT
06897, (203) 762-8020

Comments: .

Marsh Inventory No.: 405 **COE Permit No.:** **Size (ha):** 0.081

State: DE **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc. . Marsh name: Beachgrass
planting.

Marsh Inventory No.: 310 **COE Permit No.:** **Size (ha):**

State: DE **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620

Comments: .

Marsh Inventory No.: 482 **COE Permit No.:** **Size (ha):** 0.162

State: DE **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc. . Marsh name: Shore planting.

Marsh Inventory No.: 510 **COE Permit No.:** **Size (ha):** 0.040

State: DE **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc. . Marsh name: Y. Usuki.

Marsh Inventory No.: 509 **COE Permit No.:** **Size (ha):** 0.040

State: DE **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc. . Marsh name: N.
Bunting/Vines Creek.

Marsh Inventory No.: 541 **COE Permit No.:** **Size (ha):** 0.567

State: DE **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc. . Marsh name: Hudson.

Marsh Inventory No.: 558 **COE Permit No.:** **Size (ha):** 0.121

State: DE **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc. . Marsh name: Phil Short.

Marsh Inventory No.: 595 **COE Permit No.:** **Size (ha):** 0.040

State: DE **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc. . Marsh name: RV Park.

Marsh Inventory No.: 5 **COE Permit No.:** 84W-5163 **Size (ha):** 0.057

State: FL **County:** **Bay :** AIWW

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 39 **COE Permit No.:** 85IPX-20324 **Size (ha):** 0.769

State: FL **County:** **Bay :** St. Andrew Bay

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 38 **COE Permit No.:** 82R-1605 **Size (ha):** 0.202

State: FL **County:** **Bay :** St. Andrew Bay

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 12 **COE Permit No.:** 87IPX-20545 **Size (ha):** 0.405

State: FL **County:** Brevard **Bay :** Banana River

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 11 **COE Permit No.:** 86-77-083 **Size (ha):** 0.890

State: FL **County:** Brevard **Bay :** Banana River

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 24 **COE Permit No.:** 87IPD-21185 **Size (ha):** 2.173

State: FL **County:** Brevard **Bay :** Indian River

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,

Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: .

Marsh Inventory No.: 22 **COE Permit No.:** 84F-4096 **Size (ha):** 0.049

State: FL **County:** Brevard **Bay :** Indian River

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: .

Marsh Inventory No.: 21 **COE Permit No.:** 84X-4629 **Size (ha):** 0.002

State: FL **County:** Brevard **Bay :** Indian River

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: .

Marsh Inventory No.: 20 **COE Permit No.:** 84M-2530 **Size (ha):** 0.012

State: FL **County:** Brevard **Bay :** Indian River

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: .

Marsh Inventory No.: 252 **COE Permit No.:** SAI-44 **Size (ha):** 0.202

State: FL **County:** Brevard **Bay :** Indian River, Indian River Lagoon

Source of information: Dr. Robin Lewis, III., Lewis Environmental Services, Inc.,
P.O. Box 20005, Tampa, FL 33622-0005, (813) 889-9684 (Crewz & Lewis, 1991)

Comments: *Spartina alterniflora* growing well, but fringe was covered by fill. Mixed
success. .

Marsh Inventory No.: 15 **COE Permit No.:** 84D-5195 **Size (ha):** 1.044

State: FL **County:** Camden **Bay :** Cumberland Sound; Amelia River

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: .

Marsh Inventory No.: 14 **COE Permit No.:** 84J-0062 **Size (ha):** 1.340

State: FL **County:** Camden **Bay :** Cumberland Sound; Amelia River

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: .

Marsh Inventory No.: 18 **COE Permit No.:** 85-47005 **Size (ha):** 0.004

State: FL **County:** Dixie **Bay :** Horseshoe Cove

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: .

Marsh Inventory No.: 6 **COE Permit No.:** 85IPX-21013 **Size (ha):** 0.028
State: FL **County:** Duval **Bay :** AIWW; Smith Creek
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.
Comments: .

Marsh Inventory No.: 31 **COE Permit No.:** 88IPK-20374 **Size (ha):** 0.012
State: FL **County:** Duval **Bay :** Nassau Sound; Nassau River
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.
Comments: .

Marsh Inventory No.: 362 **COE Permit No.:** **Size (ha):** 3.764
State: FL **County:** Duval **Bay :** St. John's River; Nichols Creek
Source of information: Ms. Lisa Adams, Florida Dept. of Environmental Regulation,
NE District Office, 7825 Baymeadows Way, Suite 200B, Jacksonville, FL 32256-7577,
(904) 448-4300.
Comments: Very successful, monitoring complete and reports in file. File #
161219709..

Marsh Inventory No.: 48 **COE Permit No.:** 89IPH-90034 **Size (ha):** 0.809
State: FL **County:** Duval **Bay :** St. Johns River
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.
Comments: .

Marsh Inventory No.: 47 **COE Permit No.:** 88-87-026 **Size (ha):** 0.085
State: FL **County:** Duval **Bay :** St. Johns River
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.
Comments: .

Marsh Inventory No.: 46 **COE Permit No.:** 86IPP-21097 **Size (ha):** 0.077
State: FL **County:** Duval **Bay :** St. Johns River
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.
Comments: .

Marsh Inventory No.: 45 **COE Permit No.:** 87IPP-20004 **Size (ha):** 0.001
State: FL **County:** Duval **Bay :** St. Johns River
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.
Comments: .

Marsh Inventory No.: 44 **COE Permit No.:** 86IPP-20625 **Size (ha):** 2.428
State: FL **County:** Duval **Bay :** St. Johns River

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 42 **COE Permit No.:** 84T-2871 **Size (ha):** 0.007

State: FL **County:** Duval **Bay :** St. Johns River

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 49 **COE Permit No.:** 90IPH-02068 **Size (ha):** 0.223

State: FL **County:** Duval **Bay :** St. Johns River; AIWW

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 43 **COE Permit No.:** 85IPX-20475 **Size (ha):** 0.004

State: FL **County:** Duval **Bay :** St. Johns River; Mill Cove

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 41 **COE Permit No.:** 83C-1603 **Size (ha):** 0.056

State: FL **County:** Duval **Bay :** St. Johns River; Pablo Creek

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 40 **COE Permit No.:** 83X-2396 **Size (ha):** 0.038

State: FL **County:** Duval **Bay :** St. Johns River; Pablo Creek

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 616 **COE Permit No.:** **Size (ha):** 0.280

State: FL **County:** Escambia **Bay :**

Source of information: Dr. Tom Roberts, Tennessee Tech University, Cookeville, TN 38505 (Roberts, 1991)

Comments: Many birds; several rats and mice; many killifish..

Marsh Inventory No.: 9 **COE Permit No.:** 87-67-002 **Size (ha):** 0.008

State: FL **County:** Franklin **Bay :** Apalachicola Bay

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 10 **COE Permit No.:** 89IPZ-20173 **Size (ha):** 0.040
State: FL **County:** Franklin **Bay :** Apalachicola Bay; East Bay
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: .

Marsh Inventory No.: 54 **COE Permit No.:** 86IPE-20122 **Size (ha):** 0.130
State: FL **County:** Hillsborough **Bay :** Tampa Bay
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: .

Marsh Inventory No.: 13 **COE Permit No.:** 85IPE-20730 **Size (ha):** 0.073
State: FL **County:** Lee **Bay :** Caloosahatchee River
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: .

Marsh Inventory No.: 55 **COE Permit No.:** 86IPE-20495 **Size (ha):** 0.057
State: FL **County:** Manatee **Bay :** Tampa Bay
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: .

Marsh Inventory No.: 51 **COE Permit No.:** 81C-1699 **Size (ha):** 0.017
State: FL **County:** Manatee **Bay :** Tampa Bay; Manatee River
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: .

Marsh Inventory No.: 7 **COE Permit No.:** 88IPJ-20334 **Size (ha):** 0.008
State: FL **County:** Martin **Bay :** AIWW
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: .

Marsh Inventory No.: 363 **COE Permit No.:** **Size (ha):** 2.428
State: FL **County:** Nassau **Bay :** ICW and Walker Creek
Source of information: Ms. Lisa Adams, Florida Dept. of Environmental Regulation,
NE District Office, 7825 Baymeadows Way, Suite 200B, Jacksonville, FL 32256-7577,
(904) 448-4300.

Comments: Very successful, no monitoring required in permit, no reports in file. File # 450804829..

Marsh Inventory No.: 30 **COE Permit No.:** 85IPF-20609 **Size (ha):** 1.044
State: FL **County:** Nassau **Bay :** Nassau Sound
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,

Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: .

Marsh Inventory No.: 32 **COE Permit No.:** 89IPV-90847 **Size (ha):** 2.226

State: FL **County:** Nassau **Bay :** Nassau Sound; Nassau River

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: .

Marsh Inventory No.: 57 **COE Permit No.:** 86IPE-20613 **Size (ha):** 0.020

State: FL **County:** Pinellas **Bay :** Tampa Bay

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: .

Marsh Inventory No.: 56 **COE Permit No.:** 86IPT-20286 **Size (ha):** 0.012

State: FL **County:** Pinellas **Bay :** Tampa Bay

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: .

Marsh Inventory No.: 37 **COE Permit No.:** 89-67-008 **Size (ha):** 0.093

State: FL **County:** Santa Rosa **Bay :** Santa Rosa Sound

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: .

Marsh Inventory No.: 36 **COE Permit No.:** 84D-3163 **Size (ha):** 0.081

State: FL **County:** Santa Rosa **Bay :** Santa Rosa Sound

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: .

Marsh Inventory No.: 620 **COE Permit No.:** **Size (ha):** 0.490

State: FL **County:** St. Johns **Bay :**

Source of information: Dr. Tom Roberts, Tennessee Tech University, Cookeville, TN
38505 (Roberts, 1991)

Comments: Marsh was fairly well established. Few birds; several rats; many killifish..

Marsh Inventory No.: 8 **COE Permit No.:** 90IPH-03073 **Size (ha):** 0.575

State: FL **County:** St. Johns **Bay :** AIWW

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: .

Marsh Inventory No.: 29 **COE Permit No.:** J199100788 **Size (ha):** 1.202

State: FL **County:** St. Johns **Bay :** Matanzas River

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,

Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: .

Marsh Inventory No.: 28 **COE Permit No.:** J199004572 **Size (ha):** 1.032

State: FL **County:** St. Johns **Bay :** Matanzas River

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: .

Marsh Inventory No.: 27 **COE Permit No.:** 88IPX-20182 **Size (ha):** 0.214

State: FL **County:** St. Johns **Bay :** Matanzas River

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: .

Marsh Inventory No.: 26 **COE Permit No.:** 85IPK-20997 **Size (ha):** 0.166

State: FL **County:** St. Johns **Bay :** Matanzas River

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: .

Marsh Inventory No.: 25 **COE Permit No.:** 84E-0916 **Size (ha):** 0.332

State: FL **County:** St. Johns **Bay :** Matanzas River

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: .

Marsh Inventory No.: 35 **COE Permit No.:** 84W-2418 **Size (ha):** 0.081

State: FL **County:** St. Johns **Bay :** San Sebastian River

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: .

Marsh Inventory No.: 34 **COE Permit No.:** 84W-5172 **Size (ha):** 0.060

State: FL **County:** St. Johns **Bay :** San Sebastian River

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: .

Marsh Inventory No.: 33 **COE Permit No.:** 84W-0524 **Size (ha):** 0.169

State: FL **County:** St. Johns **Bay :** San Sebastian River

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: .

Marsh Inventory No.: 50 **COE Permit No.:** J199100155 **Size (ha):** 0.040
State: FL **County:** St. Johns **Bay :** St. Johns River
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
 Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
 (813) 893-3503.
Comments: .

Marsh Inventory No.: 59 **COE Permit No.:** 88IPX-20575 **Size (ha):** 0.567
State: FL **County:** St. Johns **Bay :** Tolomato River
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
 Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
 (813) 893-3503.
Comments: .

Marsh Inventory No.: 58 **COE Permit No.:** 87IPL-21114 **Size (ha):** 0.008
State: FL **County:** St. Johns **Bay :** Tolomato River
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
 Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
 (813) 893-3503.
Comments: .

Marsh Inventory No.: 16 **COE Permit No.:** 86IPL-21086 **Size (ha):** 0.174
State: FL **County:** Volusia **Bay :** Halifax River
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
 Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
 (813) 893-3503.
Comments: .

Marsh Inventory No.: 622 **COE Permit No.:** **Size (ha):** 0.040
State: FL **County:** Volusia **Bay :** Halifax River
Source of information: Dr. Tom Roberts, Tennessee Tech University, Cookeville, TN
 38505 (Roberts, 1991)
Comments: No birds; no mammal signs or catches; few killifish. Small fringe marsh
 lacks habitat complexity needed to support fish..

Marsh Inventory No.: 23 **COE Permit No.:** 87IPK-20697 **Size (ha):** 7.285
State: FL **County:** Volusia **Bay :** Indian River
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
 Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
 (813) 893-3503.
Comments: .

Marsh Inventory No.: 17 **COE Permit No.:** 85IPW-90007 **Size (ha):** 0.057
State: FL **County:** Walton **Bay :** Hogtown Bayou
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
 Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
 (813) 893-3503.
Comments: .

Marsh Inventory No.: 625 **COE Permit No.:** **Size (ha):** 0.020
State: FL **County:** **Bay :**
Source of information: Mr.Dr. Tom Roberts, Tennessee Tech University, Cookeville,
 TN 38505 Roberts,1991

Comments: No birds; no mammal signs; few fish. Small fringe marsh lacks habitat complexity needed to support fish..

Marsh Inventory No.: 19 **COE Permit No.:** 82E-0666 **Size (ha):** 0.364

State: FL **County:** Bay : Indian River

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 623 **COE Permit No.:** **Size (ha):** 0.490

State: FL **County:** Bay : Tampa Bay

Source of information: Dr. Tom Roberts, Tennessee Tech University, Cookeville, TN 38505 (Roberts, 1991)

Comments: Many birds; no rats; signs of rabbits and raccoons; many killifish..

Marsh Inventory No.: 624 **COE Permit No.:** **Size (ha):** 0.280

State: FL **County:** Bay : Tampa Bay;

Source of information: Dr. Tom Roberts, Tennessee Tech University, Cookeville, TN 38505 (Roberts, 1991)

Comments: Few birds; no rats; signs of rabbits; very many killifish..

Marsh Inventory No.: 52 **COE Permit No.:** 82U-1047 **Size (ha):** 0.911

State: FL **County:** Bay : Tampa Bay; Old Tampa Bay

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 364 **COE Permit No.:** **Size (ha):** 0.405

State: FL **County:** Bay :

Source of information: Ms. Lisa Adams, Florida Dept. of Environmental Regulation, NE District Office, 7825 Baymeadows Way, Suite 200B, Jacksonville, FL 32256-7577, (904) 448-4300.

Comments: Successful. Monitoring complete, reports available but need time to locate. File # 551270462..

Marsh Inventory No.: 360 **COE Permit No.:** **Size (ha):**

State: FL **County:** Bay :

Source of information: Ms. Lucy Blair, Florida Dept. of Environmental Regulation, S District Office, 2295 Victoria Ave., Suite, Fort Myers, FL 33901, (813) 332-6975.

Comments: 2 areas were planted. Reports are required semiannually for the life of the permit. File # 361392379..

Marsh Inventory No.: 359 **COE Permit No.:** **Size (ha):** 0.728

State: FL **County:** Bay :

Source of information: Ms. Lucy Blair, Florida Dept. of Environmental Regulation, S District Office, 2295 Victoria Ave., Suite, Fort Myers, FL 33901, (813) 332-6975.

Comments: Monitoring reports to be submitted semiannually for 3 years. File # 89-0795..

Marsh Inventory No.: 361 **COE Permit No.:** **Size (ha):**

State: FL **County:** Bay :

Source of information: Ms. Lucy Blair, Florida Dept. of Environmental Regulation, S District Office, 2295 Victoria Ave., Suite, Fort Myers, FL 33901, (813) 332-6975.

Comments: Reports to be submitted annually, but none have been received. File # 361552385..

Marsh Inventory No.: 358 **COE Permit No.:** **Size (ha):**

State: FL **County:** Bay :

Source of information: Ms. Lucy Blair, Florida Dept. of Environmental Regulation, S District Office, 2295 Victoria Ave., Suite, Fort Myers, FL 33901, (813) 332-6975.

Comments: 2 of 4 mitigation areas planted with *S. alterniflora* in Aug 1991. 2 reports were submitted in 1992 and the latest report was submitted in March 1993. Semiannual reports are required for the life of the permit. File # 111793975..

Marsh Inventory No.: 69 **COE Permit No.:** 7189 **Size (ha):** 0.299

State: GA **County:** Chatham **Bay :** Wassaw Sound

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 68 **COE Permit No.:** 7075 **Size (ha):** 0.016

State: GA **County:** Chatham **Bay :** Wassaw Sound

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 67 **COE Permit No.:** 6852 **Size (ha):** 1.659

State: GA **County:** Chatham **Bay :** Wassaw Sound

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 66 **COE Permit No.:** 5922 **Size (ha):** 1.412

State: GA **County:** Chatham **Bay :** Wassaw Sound: Wilmington River

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 63 **COE Permit No.:** 6614 **Size (ha):** 18.899

State: GA **County:** Glynn **Bay :** St. Andrew Sound

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 65 **COE Permit No.:** 6879 **Size (ha):** 0.036

State: GA **County:** Glynn **Bay :** St. Simons Sound

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: . . .

Marsh Inventory No.: 60 **COE Permit No.:** 6790 **Size (ha):** 0.081
State: GA **County:** McIntosh **Bay :** Sapello Sound
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
 Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
 (813) 893-3503.
Comments: . . .

Marsh Inventory No.: 61 **COE Permit No.:** 074 OYN 004576 **Size (ha):** 0.044
State: GA **County:** Bay : St. Andrew Sound; Little Satilla River
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
 Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
 (813) 893-3503.
Comments: . . .

Marsh Inventory No.: 62 **COE Permit No.:** 4585 **Size (ha):** 1.562
State: GA **County:** Bay : St. Andrew Sound; Satilla River
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
 Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
 (813) 893-3503.
Comments: . . .

Marsh Inventory No.: 64 **COE Permit No.:** 074 OYN 004420 **Size (ha):** 0.971
State: GA **County:** Bay : St. Simons Sound; Brunswick River
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
 Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
 (813) 893-3503.
Comments: . . .

Marsh Inventory No.: 646 **COE Permit No.:** **Size (ha):**
State: LA **County:** Cameron **Bay :** Black Lake
Source of information: Mr. Ronald Marcantel, U.S. Soil Conservation Service, Lake
 Charles Field Office, 1400 Highway 14, Lake Charles, LA 71302, (318) 436-1483
Comments: Site was evaluated in 1988 and 1989..

Marsh Inventory No.: 96 **COE Permit No.:** **Size (ha):** 0.567
State: LA **County:** Cameron **Bay :** Calcasieu Lake
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
 Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
 (813) 893-3503.
Comments: Permit # OYSTER BAYOU-8.

Marsh Inventory No.: 97 **COE Permit No.:** **Size (ha):** 3.399
State: LA **County:** Cameron **Bay :** Calcasieu Lake; Calcasieu River
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
 Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
 (813) 893-3503.
Comments: Permit # LTCS-35.

Marsh Inventory No.: 124 **COE Permit No.:** **Size (ha):** 0.648
State: LA **County:** Cameron **Bay :**
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,

Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: Permit # CAMERON-815.

Marsh Inventory No.: 116 **COE Permit No.:** **Size (ha):** 284.909

State: LA **County:** Cameron **Bay :**

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: Permit # CREOLE CANAL-2.

Marsh Inventory No.: 700 **COE Permit No.:** **Size (ha):** 0.364

State: LA **County:** Iberia **Bay :** Vermillion Bay; Weeks Bay

Source of information: Mr. Brad Broussard, U.S. Soil Conservation Service, New
Iberia Field Office, 300 Iberia Street, New Iberia, LA 70560 (318) 369-6623

Comments: Marsh name: Weeks Bay..

Marsh Inventory No.: 703 **COE Permit No.:** **Size (ha):** 0.283

State: LA **County:** Iberia **Bay :** Vermillion Bay; Weeks Bay

Source of information: Mr. Brad Broussard, U.S. Soil Conservation Service, New
Iberia Field Office, 300 Iberia Street, New Iberia, LA 70560 (318) 369-6623

Comments: Marsh name: Bayou Carlin..

Marsh Inventory No.: 702 **COE Permit No.:** **Size (ha):** 0.049

State: LA **County:** Iberia **Bay :** Vermillion Bay; Weeks Bay

Source of information: Mr. Brad Broussard, U.S. Soil Conservation Service, New
Iberia Field Office, 300 Iberia Street, New Iberia, LA 70560 (318) 369-6623

Comments: Marsh name: Bayou Petite Anse-Avery Island. Single stems of *Spartina alterniflora* were planted 0.3 m apart, using an offset of 0.15 m between rows..

Marsh Inventory No.: 701 **COE Permit No.:** **Size (ha):** 0.049

State: LA **County:** Iberia **Bay :** Vermillion Bay; Weeks Bay

Source of information: Mr. Brad Broussard, U.S. Soil Conservation Service, New
Iberia Field Office, 300 Iberia Street, New Iberia, LA 70560 (318) 369-6623

Comments: Marsh name: Bayou Petite Anse-ICW..

Marsh Inventory No.: 83 **COE Permit No.:** **Size (ha):** 0.405

State: LA **County:** Jefferson **Bay :** Barataria Bay

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: Permit # CAMINADA BAY-50/1.

Marsh Inventory No.: 80 **COE Permit No.:** **Size (ha):** 1.214

State: LA **County:** Jefferson **Bay :** Barataria Bay

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: Permit # BLIND BAY-31.

Marsh Inventory No.: 78 **COE Permit No.:** **Size (ha):** 1.619

State: LA **County:** Jefferson **Bay :** Barataria Bay

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,

Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: Permit # BAYOU DUPONT-49.

Marsh Inventory No.: 76 **COE Permit No.:** **Size (ha):** 1.862

State: LA **County:** Jefferson **Bay :** Barataria Bay

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: Permit # BAYOU RIGAUD-57.

Marsh Inventory No.: 101 **COE Permit No.:** **Size (ha):** 0.486

State: LA **County:** Jefferson **Bay :** Gulf of Mexico; Caminada Bay

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: Permit # CAMINADA BAY-52/1.

Marsh Inventory No.: 130 **COE Permit No.:** **Size (ha):** 0.093

State: LA **County:** Jefferson **Bay :**

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: Permit # JEFFERSON-84.

Marsh Inventory No.: 87 **COE Permit No.:** **Size (ha):** 849.870

State: LA **County:** Lafourche **Bay :** Barataria Bay

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: Permit # LAFOURCHE-480/4.

Marsh Inventory No.: 82 **COE Permit No.:** **Size (ha):** 0.809

State: LA **County:** Lafourche **Bay :** Barataria Bay

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: Permit # LAFOURCHE-655.

Marsh Inventory No.: 85 **COE Permit No.:** **Size (ha):** 1.700

State: LA **County:** Lafourche **Bay :** Barataria Bay; Bayou Lafourche

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: Permit # BAYOU LAFOURCHE-1072.

Marsh Inventory No.: 77 **COE Permit No.:** **Size (ha):** 1424.544

State: LA **County:** Lafourche **Bay :** Barataria Bay; Caminada Bay

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: Permit # LAFOURCHE-517.

Marsh Inventory No.: 100 **COE Permit No.:** **Size (ha):** 10.118
State: LA **County:** Lafourche **Bay :** Gulf of Mexico
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.
Comments: Permit # FLOTATION CANAL-1.

Marsh Inventory No.: 110 **COE Permit No.:** **Size (ha):** 2.979
State: LA **County:** Lafourche **Bay :** Terrebonne Bay
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.
Comments: Permit # GRAND BAYOU BLUE-15.

Marsh Inventory No.: 111 **COE Permit No.:** **Size (ha):** 4.533
State: LA **County:** Lafourche **Bay :** Timbalier Bay; Flotation Canal
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.
Comments: Permit # FLOTATION CANAL-1/1.

Marsh Inventory No.: 128 **COE Permit No.:** **Size (ha):** 5.666
State: LA **County:** Lafourche **Bay :**
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.
Comments: Permit # LAFOURCHE-663/2.

Marsh Inventory No.: 126 **COE Permit No.:** **Size (ha):** 0.809
State: LA **County:** Lafourche **Bay :**
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.
Comments: Permit # LAFOURCHE-683.

Marsh Inventory No.: 125 **COE Permit No.:** **Size (ha):** 0.162
State: LA **County:** Lafourche **Bay :**
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.
Comments: Permit # LAFOURCHE-649.

Marsh Inventory No.: 121 **COE Permit No.:** **Size (ha):** 704.178
State: LA **County:** Lafourche **Bay :**
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.
Comments: Permit # LAFOURCHE-577.

Marsh Inventory No.: 119 **COE Permit No.:** **Size (ha):** 1.093
State: LA **County:** Lafourche **Bay :**
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;

(813) 893-3503.

Comments: Permit # LAFOURCHE-560.

Marsh Inventory No.: 117 **COE Permit No.:** **Size (ha):** 849.870

State: LA **County:** Lafourche **Bay :**

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: Permit # LAFOURCHE-480/1.

Marsh Inventory No.: 114 **COE Permit No.:** **Size (ha):** 5.625

State: LA **County:** Lafourche **Bay :**

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: Permit # LAFOURCHE-482.

Marsh Inventory No.: 104 **COE Permit No.:** **Size (ha):** 2.788

State: LA **County:** Orleans **Bay :** Lake Pontchartrain

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: Permit # Lake Pontchartrain-5.

Marsh Inventory No.: 84 **COE Permit No.:** **Size (ha):** 0.283

State: LA **County:** Plaquemines **Bay :** Barataria Bay

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: Permit # BARATARIA BAY-194.

Marsh Inventory No.: 81 **COE Permit No.:** **Size (ha):** 4.856

State: LA **County:** Plaquemines **Bay :** Barataria Bay

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: Permit # GRAND BAYOU-164.

Marsh Inventory No.: 79 **COE Permit No.:** **Size (ha):** 0.405

State: LA **County:** Plaquemines **Bay :** Barataria Bay

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: Permit # BAY BATISTE-18.

Marsh Inventory No.: 86 **COE Permit No.:** **Size (ha):** 3.764

State: LA **County:** Plaquemines **Bay :** Barataria Bay; Bayou Grand Cheni

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: Permit # BAYOU GRAND CHENIER-.

Marsh Inventory No.: 106 **COE Permit No.:** **Size (ha):** 0.607

State: LA **County:** Plaquemines **Bay :** Mississippi Delta
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.
Comments: Permit # CHICHARAS BAY-3.

Marsh Inventory No.: 105 **COE Permit No.:** **Size (ha):** 1.214
State: LA **County:** Plaquemines **Bay :** Mississippi Delta
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.
Comments: Permit # BAY EARANE-1.

Marsh Inventory No.: 352 **COE Permit No.:** **Size (ha):**
State: LA **County:** Plaquemines **Bay :**
Source of information: Mr. John E. Boatman, U.S. Soil Conservation Service,
Plaquemines Soil and Water Conservation District Office, 104 Hebert Boulevard, Belle
Chase, LA 70037, (504) 394-7741
Comments: Planting density of *Spartina alterniflora* 1 stem/ft. 600 linear ft. planted.
Marsh name: Lake Hermitage. To be planted in Sept. 1993. 5 board wave dampening
fences to be constructed prior to planting..

Marsh Inventory No.: 351 **COE Permit No.:** **Size (ha):**
State: LA **County:** Plaquemines **Bay :**
Source of information: Mr. John E. Boatman, U.S. Soil Conservation Service,
Plaquemines Soil and Water Conservation District Office, 104 Hebert Boulevard, Belle
Chase, LA 70037, (504) 394-7741
Comments: Planting density of *Spartina alterniflora* 1 stem/ft. 600 linear ft. planted.
Marsh name: Lake Laurier. To be planted in Sept. 1993. 5 board wave dampening fences
to be constructed prior to planting..

Marsh Inventory No.: 129 **COE Permit No.:** GP890950 **Size (ha):** 0.405
State: LA **County:** Plaquemines **Bay :**
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.
Comments: .

Marsh Inventory No.: 127 **COE Permit No.:** **Size (ha):** 0.567
State: LA **County:** Plaquemines **Bay :**
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.
Comments: Permit # PLAQUEMINES-505.

Marsh Inventory No.: 120 **COE Permit No.:** **Size (ha):** 0.243
State: LA **County:** Plaquemines **Bay :**
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.
Comments: Permit # PLAQUEMINES-427.

Marsh Inventory No.: 118 **COE Permit No.:** **Size (ha):** 8.094

State: LA **County:** Plaquemines **Bay :**

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: Permit # PLAQUEMINES-391.

Marsh Inventory No.: 115 **COE Permit No.:** **Size (ha):** 0.931

State: LA **County:** Plaquemines **Bay :**

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: Permit # PLAQUEMINES-330.

Marsh Inventory No.: 71 **COE Permit No.:** **Size (ha):** 0.931

State: LA **County:** Plaquemines **Bay :**

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: Permit # PLAQUEMINES-284.

Marsh Inventory No.: 347 **COE Permit No.:** **Size (ha):** 0.121

State: LA **County:** Plaquemines **Bay :**

Source of information: Mr. John E. Boatman, U.S. Soil Conservation Service, Plaquemines Soil and Water Conservation District Office, 104 Hebert Boulevard, Belle Chase, LA 70037, (504) 394-7741

Comments: Planting density of *Spartina alterniflora* 1 stem/ft. 13,050 linear ft. planted. Marsh name: Myrtle Grove. This project was completed in 1991. Survival rate only 2% due to nutria predation. Only 10% of plants were caged. .

Marsh Inventory No.: 350 **COE Permit No.:** none **Size (ha):** 0.017

State: LA **County:** Plaquemines **Bay :**

Source of information: Mr. John E. Boatman, U.S. Soil Conservation Service, Plaquemines Soil and Water Conservation District Office, 104 Hebert Boulevard, Belle Chase, LA 70037, (504) 394-7741

Comments: Planting density of *S. alterniflora* 1 stem/ft. 1,875 linear ft. planted. Marsh name: Yellow Cotton Bay. 5/93 Survival Rate 50%. SE (General Permit) NOD-100. Survival rate higher in this region probably because of greater mineral content of soil.. Plantings were done along a washed-out spoil bank along a canal. Plants showed good to excellent vigor but growth was confined to within nutria-exclusion devices (NED'S). Salinity levels in the area range from 3-10 ppt.

Marsh Inventory No.: 349 **COE Permit No.:** none **Size (ha):** 0.019

State: LA **County:** Plaquemines **Bay :**

Source of information: Mr. John E. Boatman, U.S. Soil Conservation Service, Plaquemines Soil and Water Conservation District Office, 104 Hebert Boulevard, Belle Chase, LA 70037, (504) 394-7741

Comments: Planting density of *Spartina alterniflora* 1 stem/ft. 2,022 linear ft. planted. Marsh name: Bayou Dupont 3/93 Survival rate 10%. Damage due to Hurricane Andrew and nutria. SE (General Permit) NOD-100..

Marsh Inventory No.: 348 **COE Permit No.:** none **Size (ha):** 0.012

State: LA **County:** Plaquemines **Bay :**

Source of information: Mr. John E. Boatman, U.S. Soil Conservation Service,

Plaquemines Soil and Water Conservation District Office, 104 Hebert Boulevard, Belle Chase, LA 70037, (504) 394-7741

Comments: Planting density of *Spartina alterniflora* 1 stem/ft. 1,320 linear ft. planted. Marsh name: Round Lake. Survival rate poor (3%). Causes include nutria predation, shoreline erosion, Hurricane Andrew, soils. . . Although 70% of plants were caged, once cages deteriorated or were opened by Hurricane Andrew, nutria ate the plants. Only plants away from the shoreline and still in cages survived. Also, plants not on firm soils failed to establish.

Marsh Inventory No.: 91 **COE Permit No.:** **Size (ha):** 2.428

State: LA **County:** St. Bernard **Bay :** Breton Sound

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: Permit # MRGO-70.

Marsh Inventory No.: 92 **COE Permit No.:** **Size (ha):** 2.428

State: LA **County:** St. Bernard **Bay :** Breton Sound; Lake Machias

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: Permit # LAKE MACHIAS-19.

Marsh Inventory No.: 93 **COE Permit No.:** **Size (ha):** 6.071

State: LA **County:** St. Bernard **Bay :** Breton Sound; Yscloskey Bayou

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: Permit # YSCLOSKEY BAYOU-14.

Marsh Inventory No.: 98 **COE Permit No.:** **Size (ha):** 1.943

State: LA **County:** St. Bernard **Bay :** Chandeleur Sound

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: Permit # ST. BERNARD-43.

Marsh Inventory No.: 113 **COE Permit No.:** **Size (ha):** 0.850

State: LA **County:** St. Bernard **Bay :**

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: Permit # ST. BERNARD-47.

Marsh Inventory No.: 112 **COE Permit No.:** **Size (ha):** 0.809

State: LA **County:** St. Bernard **Bay :**

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: Permit # ST. BERNARD-45.

Marsh Inventory No.: 75 **COE Permit No.:** **Size (ha):** 16.188

State: LA **County:** St. Martin **Bay :** Atachafalaya Floodway

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: Permit # LITTLE BAYOU CHEVREA.

Marsh Inventory No.: 643 **COE Permit No.:** **Size (ha):**

State: LA **County:** St. Tammeny **Bay :** Lake Pontchartrain

Source of information: Mr. Tony Beauboef, SCS, Franklinton, LA, (504) 839-5688

Comments: Site was evaluated in 1988 and 1989..

Marsh Inventory No.: 95 **COE Permit No.:** **Size (ha):** 0.975

State: LA **County:** Terrebonne **Bay :** Caillou Bay

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: Permit # TERREBONNE-763.

Marsh Inventory No.: 94 **COE Permit No.:** **Size (ha):** 1.214

State: LA **County:** Terrebonne **Bay :** Caillou Bay

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: Permit # CHARLEYS BAY-3.

Marsh Inventory No.: 644 **COE Permit No.:** **Size (ha):**

State: LA **County:** Terrebonne **Bay :** Lake DeCade

Source of information: Mr. Michael Tullos, U.S. Soil Conservation Service, Houma Field Office, Box 1266, Houma, LA 10361, (504) 872-5609

Comments: Site was evaluated in 1988 and 1989..

Marsh Inventory No.: 102 **COE Permit No.:** **Size (ha):** 80.940

State: LA **County:** Terrebonne **Bay :** Lake Pelto

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: Permit # GOM-3450.

Marsh Inventory No.: 109 **COE Permit No.:** **Size (ha):** 10.846

State: LA **County:** Terrebonne **Bay :** Terrebonne Bay

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: Permit # LITTLE BAYOU CHEVREA.

Marsh Inventory No.: 108 **COE Permit No.:** **Size (ha):** 0.040

State: LA **County:** Terrebonne **Bay :** Terrebonne Bay

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: Permit # TERREBONNE-536A.

Marsh Inventory No.: 107 **COE Permit No.:** **Size (ha):** 1.295

State: LA **County:** Terrebonne **Bay :** Terrebonne Bay

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: Permit # TERREBONNE-519/1.

Marsh Inventory No.: 123 **COE Permit No.:** **Size (ha):** 0.526

State: LA **County:** Terrebonne **Bay :**

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: Permit # TERREBONNE-798.

Marsh Inventory No.: 122 **COE Permit No.:** **Size (ha):** 1.862

State: LA **County:** Terrebonne **Bay :**

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: Permit # JACK STOUT BAY-1.

Marsh Inventory No.: 74 **COE Permit No.:** **Size (ha):** 1.821

State: LA **County:** Terrebonne **Bay :**

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: Permit # TERREBONNE-583.

Marsh Inventory No.: 73 **COE Permit No.:** **Size (ha):** 0.405

State: LA **County:** Terrebonne **Bay :**

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: Permit # TERREBONNE-559.

Marsh Inventory No.: 72 **COE Permit No.:** **Size (ha):** 0.809

State: LA **County:** Terrebonne **Bay :**

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: Permit # TERREBONNE-536.

Marsh Inventory No.: 70 **COE Permit No.:** **Size (ha):** 8.175

State: LA **County:** Terrebonne **Bay :**

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: Permit # TERREBONNE-528.

Marsh Inventory No.: 645 **COE Permit No.:** **Size (ha):**

State: LA **County:** Vermilion **Bay :** Rollover Bayou

Source of information: Mr. Donald Menard, U.S. Soil Conservation Service, Abbeville Field Office, P.O. Box 68, Abbeville, LA 70511-0068, (318) 893-5664

Comments: Site was evaluated in 1988 and 1989..

Marsh Inventory No.: 89 **COE Permit No.:** **Size (ha):** 0.931

State: LA **County:** **Bay :** Bayou De La Valle

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: Permit # BAYOU DE LA VALLE-7.

Marsh Inventory No.: 88 **COE Permit No.:** **Size (ha):** 0.809

State: LA **County:** **Bay :** Bayou De La Valle

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: Permit # BAYOU DE LA VALLE-4.

Marsh Inventory No.: 90 **COE Permit No.:** **Size (ha):** 16.188

State: LA **County:** **Bay :** Bayou Terrebonne

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: Permit # TERREBONNE-690.

Marsh Inventory No.: 99 **COE Permit No.:** **Size (ha):** 4.047

State: LA **County:** **Bay :** Grand Bayou

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: Permit # GRAND BAYOU-155.

Marsh Inventory No.: 103 **COE Permit No.:** **Size (ha):** 0.283

State: LA **County:** **Bay :** Lake Pontchartrain

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: Permit # Lake Pontchartrain-5.

Marsh Inventory No.: 667 **COE Permit No.:** **Size (ha):** 8.903

State: ME **County:** **Bay :** Webhannet River

Source of information: Mr. Bill Hubbard, COE, (617) 647-8552

Comments: Built by Environmental Concern, Inc.. Marsh name: Webhannet River Marsh.

Marsh Inventory No.: 393 **COE Permit No.:** **Size (ha):** 0.040

State: ME **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Kittery Restoration.

Marsh Inventory No.: 480 **COE Permit No.:** **Size (ha):** 0.040

State: ME **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Leather Loft.

Marsh Inventory No.: 585 **COE Permit No.:** **Size (ha):** 0.081
State: MD **County:** Kent **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: Little Neck Farm.

Marsh Inventory No.: 651 **COE Permit No.:** **Size (ha):**
State: MD **County:** Queen Anne **Bay :** Broad Bay
Source of information: Donald W. Hamer, U.S. Soil Conservation Service, 1536 Rt.
9 North, Cape May Court House, NJ 08210 (609) 465-5901
Comments: .

Marsh Inventory No.: 527 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** Talbot **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: Roderick Hickey,
Little Neck Creek.

Marsh Inventory No.: 564 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** Talbot **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: Bruce Hackett/Island
Creek.

Marsh Inventory No.: 589 **COE Permit No.:** **Size (ha):** 0.081
State: MD **County:** Talbot **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: John Swope/Irish
Creek.

Marsh Inventory No.: 650 **COE Permit No.:** **Size (ha):**
State: MD **County:** **Bay :** Chesapeake Bay
Source of information: Donald W. Hamer, U.S. Soil Conservation Service, 1536 Rt.
9 North, Cape May Court House, NJ 08210 (609) 465-5901
Comments: No *S. alterniflora* planted..

Marsh Inventory No.: 654 **COE Permit No.:** **Size (ha):**
State: MD **County:** **Bay :** Chester River
Source of information: Donald W. Hamer, U.S. Soil Conservation Service, 1536 Rt.
9 North, Cape May Court House, NJ 08210 (609) 465-5901
Comments: .

Marsh Inventory No.: 653 **COE Permit No.:** **Size (ha):**
State: MD **County:** **Bay :** Choptank River
Source of information: Donald W. Hamer, U.S. Soil Conservation Service, 1536 Rt.
9 North, Cape May Court House, NJ 08210 (609) 465-5901
Comments: .

Marsh Inventory No.: 380 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: San Domingo.

Marsh Inventory No.: 387 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Hambleton Island SD.

Marsh Inventory No.: 386 **COE Permit No.:** **Size (ha):** 0.081

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Sand Spit.

Marsh Inventory No.: 385 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Hambleton Island Salt Pan.

Marsh Inventory No.: 384 **COE Permit No.:** **Size (ha):** 0.121

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Long Point Island.

Marsh Inventory No.: 383 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Tred Avon River.

Marsh Inventory No.: 382 **COE Permit No.:** **Size (ha):** 0.081

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Rich Neck.

Marsh Inventory No.: 381 **COE Permit No.:** **Size (ha):** 0.081

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Hambleton Island.

Marsh Inventory No.: 284 **COE Permit No.:** **Size (ha):** 1.619

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Dredged material stabilization project.. Marsh name: Tar Bay

Marsh Inventory No.: 392 **COE Permit No.:** **Size (ha):** 0.081
State: MD **County:** Bay :
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: Hambleton Island BC.

Marsh Inventory No.: 391 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** Bay :
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: Harris Creek.

Marsh Inventory No.: 390 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** Bay :
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: Cober.

Marsh Inventory No.: 288 **COE Permit No.:** **Size (ha):** 0.809
State: MD **County:** Bay :
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Dredged material stabilization project.. Marsh name: Knapps Narrows -
COE

Marsh Inventory No.: 395 **COE Permit No.:** **Size (ha):** 0.081
State: MD **County:** Bay :
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: Ray's Point.

Marsh Inventory No.: 394 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** Bay :
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: Slaughter Creek-
Shore.

Marsh Inventory No.: 398 **COE Permit No.:** **Size (ha):** 0.081
State: MD **County:** Bay :
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: Bliss.

Marsh Inventory No.: 397 **COE Permit No.:** **Size (ha):** 0.081
State: MD **County:** Bay :
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: Hambleton Island
1977 Breach.

Marsh Inventory No.: 396 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** Bay :

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: R. Warbasse.

Marsh Inventory No.: 403 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Berlin.

Marsh Inventory No.: 402 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Francis Somers.

Marsh Inventory No.: 401 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Miller.

Marsh Inventory No.: 400 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Nevius.

Marsh Inventory No.: 399 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Turkey Neck Point.

Marsh Inventory No.: 408 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: St. Mary's College.

Marsh Inventory No.: 407 **COE Permit No.:** **Size (ha):** 0.121

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Nelson Point Farm.

Marsh Inventory No.: 406 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Bullard.

Marsh Inventory No.: 413 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** Bay :

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Dr. G. Callender.

Marsh Inventory No.: 412 **COE Permit No.:** **Size (ha):** 0.324

State: MD **County:** Bay :

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: E.G. Uhl.

Marsh Inventory No.: 411 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** Bay :

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Dunmoyle.

Marsh Inventory No.: 410 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** Bay :

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: William Hahn.

Marsh Inventory No.: 419 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** Bay :

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: G.A. Van Lennep, Jr.

Marsh Inventory No.: 418 **COE Permit No.:** **Size (ha):** 0.081

State: MD **County:** Bay :

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: S. Hamblin.

Marsh Inventory No.: 417 **COE Permit No.:** **Size (ha):** 0.081

State: MD **County:** Bay :

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: R. Dresser.

Marsh Inventory No.: 416 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** Bay :

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Miles River Bridge.

Marsh Inventory No.: 429 **COE Permit No.:** **Size (ha):** 0.243

State: MD **County:** Bay :

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: E.B.M. White.

Marsh Inventory No.: 428 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: K. White.

Marsh Inventory No.: 427 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: Struse-Arrowhead
Farm.

Marsh Inventory No.: 426 **COE Permit No.:** **Size (ha):** 0.081
State: MD **County:** **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: J.R. Griswold.

Marsh Inventory No.: 425 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: D.M. Palmer.

Marsh Inventory No.: 424 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: B.E. Blake.

Marsh Inventory No.: 423 **COE Permit No.:** **Size (ha):** 0.081
State: MD **County:** **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: H.T. Winner.

Marsh Inventory No.: 422 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: Bob Harris.

Marsh Inventory No.: 421 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: A. Van Dyke.

Marsh Inventory No.: 420 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: John Masone.

Marsh Inventory No.: 309 **COE Permit No.:** **Size (ha):** 8.701

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Dredged material stabilization project.. Marsh name: Barren Island

Marsh Inventory No.: 444 **COE Permit No.:** **Size (ha):** 0.081

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: L.G. Wineland.

Marsh Inventory No.: 443 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: T. Morison.

Marsh Inventory No.: 442 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Benjamin Florence.

Marsh Inventory No.: 441 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: D. Gregg III.

Marsh Inventory No.: 440 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: R. B. Scofield, Jr.

Marsh Inventory No.: 439 **COE Permit No.:** **Size (ha):** 0.324

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Kent Narrows.

Marsh Inventory No.: 438 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: W.II. Van Dusen-Spitzer.

Marsh Inventory No.: 437 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: E. Miller (Emerson Pt.).

Marsh Inventory No.: 436 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: R. King.

Marsh Inventory No.: 435 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: L.G. Conner.

Marsh Inventory No.: 434 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: R.C. Morris.

Marsh Inventory No.: 433 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: R. Kleffer, Jr.

Marsh Inventory No.: 432 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: S.M. Bothe.

Marsh Inventory No.: 431 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: C.C. Bond.

Marsh Inventory No.: 430 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Dennis Berg.

Marsh Inventory No.: 479 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: M. Millennann.

Marsh Inventory No.: 478 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** Bay :
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: R. Gladstone.

Marsh Inventory No.: 477 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** Bay :
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: G. Bunting.

Marsh Inventory No.: 476 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** Bay :
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: R. Barrett.

Marsh Inventory No.: 475 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** Bay :
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: M.D. Clark.

Marsh Inventory No.: 474 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** Bay :
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: W.T. Hershey.

Marsh Inventory No.: 473 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** Bay :
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: A.J. Clark.

Marsh Inventory No.: 472 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** Bay :
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: C.W. King.

Marsh Inventory No.: 471 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** Bay :
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: W.M. Cooper.

Marsh Inventory No.: 470 **COE Permit No.:** **Size (ha):** 0.121
State: MD **County:** Bay :
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: T.M. Rauch.

Marsh Inventory No.: 469 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: S. Gilliece.

Marsh Inventory No.: 468 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: R. Polevoy.

Marsh Inventory No.: 467 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: Dr. A. Kaplan.

Marsh Inventory No.: 466 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: T. McMahon.

Marsh Inventory No.: 465 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: Eco Smugglers Cove.

Marsh Inventory No.: 464 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: Dr. P.D. Kay.

Marsh Inventory No.: 463 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: Ms. P. Bryan.

Marsh Inventory No.: 462 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: J.B. Robertson.

Marsh Inventory No.: 461 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: J.J. Cottrell.

Marsh Inventory No.: 460 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: E.L. Strohbahn.

Marsh Inventory No.: 459 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: W. Brennen.

Marsh Inventory No.: 458 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: S.M. Margolis.

Marsh Inventory No.: 457 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Albert G. Gipe.

Marsh Inventory No.: 456 **COE Permit No.:** **Size (ha):** 0.081

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: G. G. Tucker.

Marsh Inventory No.: 455 **COE Permit No.:** **Size (ha):** 0.081

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: DNR Wye Island Demo-SCS.

Marsh Inventory No.: 454 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Dr. E. Throop.

Marsh Inventory No.: 453 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Hartages Yacht Club.

Marsh Inventory No.: 452 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: J.J. Young

Marsh Inventory No.: 451 **COE Permit No.:** **Size (ha):** 0.081
State: MD **County:** **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: J.K. Hyatt.

Marsh Inventory No.: 450 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: M. Sami.

Marsh Inventory No.: 449 **COE Permit No.:** **Size (ha):** 0.081
State: MD **County:** **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: Hooper Island Marsh.

Marsh Inventory No.: 448 **COE Permit No.:** **Size (ha):** 0.081
State: MD **County:** **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: Thomas Point Restoration.

Marsh Inventory No.: 446 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: C.H. Hardesty.

Marsh Inventory No.: 445 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: Mrs. R. H. Lee.

Marsh Inventory No.: 505 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: M. Franks.

Marsh Inventory No.: 504 **COE Permit No.:** **Size (ha):** 0.202
State: MD **County:** **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: Nassetta-Phillips.

Marsh Inventory No.: 503 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** Bay :
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: Jensens-Ed Fisher.

Marsh Inventory No.: 502 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** Bay :
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: P.J. Janson.

Marsh Inventory No.: 501 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** Bay :
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: General A.J. Goodpaster.

Marsh Inventory No.: 500 **COE Permit No.:** **Size (ha):** 0.081
State: MD **County:** Bay :
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: R. Timbie.

Marsh Inventory No.: 499 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** Bay :
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: P. Wilmerding.

Marsh Inventory No.: 498 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** Bay :
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: Mr. L. Root.

Marsh Inventory No.: 497 **COE Permit No.:** **Size (ha):** 0.081
State: MD **County:** Bay :
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: G.R. Kilbourn, Jr.

Marsh Inventory No.: 496 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** Bay :
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: S. Shaare.

Marsh Inventory No.: 495 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** Bay :
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: G. Taylor.

Marsh Inventory No.: 494 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: R.R. Price.

Marsh Inventory No.: 493 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: J.B. Wilson.

Marsh Inventory No.: 492 **COE Permit No.:** **Size (ha):** 0.081

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: J.K. Hyatt.

Marsh Inventory No.: 491 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: R.L. Counts.

Marsh Inventory No.: 490 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: E.A. Rommel, III.

Marsh Inventory No.: 489 **COE Permit No.:** **Size (ha):** 0.081

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: C.E. Peck.

Marsh Inventory No.: 488 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: L. Beaumont.

Marsh Inventory No.: 487 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: R.A. Simmons.

Marsh Inventory No.: 486 **COE Permit No.:** **Size (ha):** 0.081

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box

P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Horn Point.

Marsh Inventory No.: 485 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Mr. L. Tucker.

Marsh Inventory No.: 484 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: M.S. Kozel.

Marsh Inventory No.: 483 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: D. Proctor.

Marsh Inventory No.: 538 **COE Permit No.:** **Size (ha):** 0.081

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: D. Hamilton.

Marsh Inventory No.: 537 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: F.V. Lawrence III.

Marsh Inventory No.: 536 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Ali Mehrizi.

Marsh Inventory No.: 535 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Dr. E.H. Breen.

Marsh Inventory No.: 534 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Dr. W. Jewell.

Marsh Inventory No.: 533 **COE Permit No.:** **Size (ha):** 0.121

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: J.J. Roberts.

Marsh Inventory No.: 532 **COE Permit No.:** **Size (ha):** 0.081

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Doreen Hamilton, Irish Creek.

Marsh Inventory No.: 531 **COE Permit No.:** **Size (ha):** 0.081

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: A.E. MacKay.

Marsh Inventory No.: 530 **COE Permit No.:** **Size (ha):** 0.243

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Robert Gladstone.

Marsh Inventory No.: 529 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: W. Gettsinger.

Marsh Inventory No.: 528 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Karl Krieger.

Marsh Inventory No.: 526 **COE Permit No.:** **Size (ha):** 0.081

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: C.J. Zamoiske.

Marsh Inventory No.: 525 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Ships Point.

Marsh Inventory No.: 524 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: M. Harriman.

Marsh Inventory No.: 523 **COE Permit No.:** **Size (ha):** 0.081
State: MD **County:** Bay :
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: R. Garrett.

Marsh Inventory No.: 522 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** Bay :
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: Don Olson.

Marsh Inventory No.: 521 **COE Permit No.:** **Size (ha):** 0.121
State: MD **County:** Bay :
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: Anchorage Marina.

Marsh Inventory No.: 520 **COE Permit No.:** **Size (ha):** 0.081
State: MD **County:** Bay :
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: Rock Harbor Planting.

Marsh Inventory No.: 519 **COE Permit No.:** **Size (ha):** 0.081
State: MD **County:** Bay :
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: Church Creek Marsh.

Marsh Inventory No.: 518 **COE Permit No.:** **Size (ha):** 0.202
State: MD **County:** Bay :
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: Bodkin Island.

Marsh Inventory No.: 517 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** Bay :
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: G. Carlson.

Marsh Inventory No.: 516 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** Bay :
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: J. Cobbs.

Marsh Inventory No.: 515 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** Bay :
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: C. Petty.

Marsh Inventory No.: 514 **COE Permit No.:** **Size (ha):** 0.243
State: MD **County:** Bay :
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: Mario-Boiardi-Wye
Mills.

Marsh Inventory No.: 513 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** Bay :
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: B. DuPont.

Marsh Inventory No.: 512 **COE Permit No.:** **Size (ha):** 0.202
State: MD **County:** Bay :
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: Cheston on Wye.

Marsh Inventory No.: 511 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** Bay :
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: J. MacMartin.

Marsh Inventory No.: 321 **COE Permit No.:** **Size (ha):** 0.405
State: MD **County:** Bay :
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: . Marsh name: D. B. Cober

Marsh Inventory No.: 556 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** Bay :
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: Thomas
Boone/Maxmore Creek.

Marsh Inventory No.: 555 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** Bay :
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: Fred C. Meensden/San
Domingo Creek.

Marsh Inventory No.: 554 **COE Permit No.:** **Size (ha):** 0.081
State: MD **County:** Bay :
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: Laurence E. Diggs III.

Marsh Inventory No.: 553 **COE Permit No.:** **Size (ha):** 0.040

State: MD County: Bay :

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Stephanie Trivis/Island Creek.

Marsh Inventory No.: 552 COE Permit No.: Size (ha): 0.040

State: MD County: Bay :

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: R.D. Young/Broad Creek.

Marsh Inventory No.: 551 COE Permit No.: Size (ha): 0.081

State: MD County: Bay :

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Joseph L. Morris.

Marsh Inventory No.: 550 COE Permit No.: Size (ha): 0.040

State: MD County: Bay :

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Richard D. Richards.

Marsh Inventory No.: 549 COE Permit No.: Size (ha): 0.121

State: MD County: Bay :

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: John Nord/Cummings Creek.

Marsh Inventory No.: 548 COE Permit No.: Size (ha): 0.040

State: MD County: Bay :

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: H.W. Barrick/Wye River.

Marsh Inventory No.: 547 COE Permit No.: Size (ha): 0.040

State: MD County: Bay :

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Gilbert Hahn/Harris Creek.

Marsh Inventory No.: 546 COE Permit No.: Size (ha): 0.040

State: MD County: Bay :

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: John Schreiner/Harris Creek

Marsh Inventory No.: 545 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: Edgar Garbisch/San
Domingo Creek.

Marsh Inventory No.: 544 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: Paul
Makosky/Barrett's Cove.

Marsh Inventory No.: 543 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: John Reihl III/
Maxmore Creek.

Marsh Inventory No.: 542 **COE Permit No.:** **Size (ha):** 0.567
State: MD **County:** **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: Rouse Corporation.

Marsh Inventory No.: 582 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: J.F.
Moore/Trippe Creek.

Marsh Inventory No.: 581 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: Robert Price III/Wye
River.

Marsh Inventory No.: 580 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: C.J.
Lambertson/Ledenham Creek.

Marsh Inventory No.: 579 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: Edgar Garbisch

Marsh Inventory No.: 578 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** Bay :
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: William S. Gaines/San
Domingo Creek.

Marsh Inventory No.: 577 **COE Permit No.:** **Size (ha):** 0.081
State: MD **County:** Bay :
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: Bishop Sorge/Island
Creek.

Marsh Inventory No.: 576 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** Bay :
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: W.R. Simpson/Miles
River.

Marsh Inventory No.: 575 **COE Permit No.:** **Size (ha):** 0.081
State: MD **County:** Bay :
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: John Wilson/Harris
Creek.

Marsh Inventory No.: 574 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** Bay :
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: George
White/Wye River.

Marsh Inventory No.: 573 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** Bay :
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: Donald Zeman/Ball's
Creek.

Marsh Inventory No.: 572 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** Bay :
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: Richard M.
Siegal/Island Creek.

Marsh Inventory No.: 571 **COE Permit No.:** **Size (ha):** 0.081
State: MD **County:** Bay :
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box

P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Robert M. Valliant/Plaindealing Creek.

Marsh Inventory No.: 570 **COE Permit No.:** **Size (ha):** 0.202

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Howard W. Williams/Island Creek.

Marsh Inventory No.: 569 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: James Arnold/Glebe Creek.

Marsh Inventory No.: 568 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Colin Ferenback/Plaindealing Creek.

Marsh Inventory No.: 567 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Richard Crowley/LeGates Cove.

Marsh Inventory No.: 566 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: John L. Cobbs/San Domingo Creek.

Marsh Inventory No.: 565 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Clinton Taylor/Folly's Cove.

Marsh Inventory No.: 563 **COE Permit No.:** **Size (ha):** 0.202

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Alan Geller/Huntington Creek.

Marsh Inventory No.: 562 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: Charles Peck/Solitude
Creek.

Marsh Inventory No.: 561 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: L. David
Reynolds/San Domingo Creek.

Marsh Inventory No.: 560 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: Muriel
Blumenthal/Trippe Creek.

Marsh Inventory No.: 559 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: Janet Brown/Goose
Cove.

Marsh Inventory No.: 594 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: Craig Henry/Broad
Creek.

Marsh Inventory No.: 593 **COE Permit No.:** **Size (ha):** 0.081
State: MD **County:** **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: Sumner
Park/Shipshead Creek.

Marsh Inventory No.: 592 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: Edward Dwyer/Oak
Creek.

Marsh Inventory No.: 591 **COE Permit No.:** **Size (ha):** 0.081
State: MD **County:** **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Peter Black/Maxmore Creek.

Marsh Inventory No.: 590 **COE Permit No.:** **Size (ha):** 0.081

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Richard Granville/Tar Creek.

Marsh Inventory No.: 588 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Stephen Miller.

Marsh Inventory No.: 587 **COE Permit No.:** **Size (ha):** 0.162

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Clayton Hurt.

Marsh Inventory No.: 586 **COE Permit No.:** **Size (ha):** 0.081

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Michael Abramo/Wye River.

Marsh Inventory No.: 584 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Archie MacKay/Snug Harbor.

Marsh Inventory No.: 583 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Oscar Carlson.

Marsh Inventory No.: 599 **COE Permit No.:** **Size (ha):** 0.081

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Robert Simmons/Cummings Creek.

Marsh Inventory No.: 598 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Langly Shook/San Domingo Creek.

Marsh Inventory No.: 597 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: William Baldwin.

Marsh Inventory No.: 596 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Combined Sewer Overflow Project.

Marsh Inventory No.: 614 **COE Permit No.:** **Size (ha):** 0.081

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Mary Hagy.

Marsh Inventory No.: 613 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Carol McGowan.

Marsh Inventory No.: 612 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Sergio Proserpi.

Marsh Inventory No.: 611 **COE Permit No.:** **Size (ha):** 0.162

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Paul Fleagane.

Marsh Inventory No.: 610 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Jan Gervin.

Marsh Inventory No.: 609 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Elliott Bliss.

Marsh Inventory No.: 608 **COE Permit No.:** **Size (ha):** 0.081

State: MD **County:** Bay :

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Cove View Condos.

Marsh Inventory No.: 607 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** Bay :

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Maritime Museum/Chesapeake Bay.

Marsh Inventory No.: 606 **COE Permit No.:** **Size (ha):** 0.162

State: MD **County:** Bay :

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: John Reilly.

Marsh Inventory No.: 605 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** Bay :

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Joseph Schuble.

Marsh Inventory No.: 604 **COE Permit No.:** **Size (ha):** 0.081

State: MD **County:** Bay :

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Garth Harding.

Marsh Inventory No.: 603 **COE Permit No.:** **Size (ha):** 0.040

State: MD **County:** Bay :

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: John Thorman.

Marsh Inventory No.: 602 **COE Permit No.:** **Size (ha):** 0.121

State: MD **County:** Bay :

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Willard Parker.

Marsh Inventory No.: 601 **COE Permit No.:** **Size (ha):** 0.121

State: MD **County:** Bay :

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: John Akridge.

Marsh Inventory No.: 600 **COE Permit No.:** **Size (ha):** 0.081

State: MD **County:** Bay :

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc.. Marsh name: Robert Kennedy.

Marsh Inventory No.: 447 **COE Permit No.:** **Size (ha):** 0.040
State: MD **County:** Bay :
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: Queenslanding.

Marsh Inventory No.: 672 **COE Permit No.:** **Size (ha):** 0.708
State: MA **County:** Suffolk **Bay :** Boston Harbor
Source of information: Ms. Laurie Cullen, Massport, (617) 973-5389
Comments: . Marsh name: Massport/Logan Airport.

Marsh Inventory No.: 615 **COE Permit No.:** **Size (ha):**
State: MA **County:** Suffolk / Norfolk **Bay :**
Source of information: Dr. Lars H. Carlson, Sverdrup Corp., 2 Center Plaza, Boston, MA 02108-1906, (617) 742-8060; or Dr. Norton H. Nickerson, Dept. of Biology, Tufts Univ., Medford, MA 02155, (617) 627-3544 FEIS/Report, Old Colony Railroad Rehabilitation Project. (EOEA 5840).
Comments: . Marsh name: Old Colony Railroad rehabilitation project.

Marsh Inventory No.: 415 **COE Permit No.:** **Size (ha):** 0.202
State: MA **County:** Bay :
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc. , Inc.. Marsh name: Quincy MA.

Marsh Inventory No.: 131 **COE Permit No.:** MS87-00966-Y **Size (ha):** 0.040
State: MS **County:** Hancock **Bay :** Mississippi Sound
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.
Comments: .

Marsh Inventory No.: 133 **COE Permit No.:** MS91-01502-I **Size (ha):** 0.259
State: MS **County:** Harrison **Bay :** Mississippi Sound
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.
Comments: .

Marsh Inventory No.: 132 **COE Permit No.:** MS90-02404-H **Size (ha):** 0.004
State: MS **County:** Jackson **Bay :** Mississippi Sound; Bayou Cassotte
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.
Comments: .

Marsh Inventory No.: 134 **COE Permit No.:** MS91-01824-U **Size (ha):** 0.028
State: MS **County:** Jackson **Bay :** Mississippi Sound; Pascagoula River
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.
Comments: .

Marsh Inventory No.: 668 **COE Permit No.:** **Size (ha):** 0.405
State: NH **County:** Rockingham **Bay :** Little Harbor
Source of information: Mr. Frank Richardson, NH Wetlands Board, (603) 271-2147
Comments: . Marsh name: Wentworth Hotel.

Marsh Inventory No.: 669 **COE Permit No.:** **Size (ha):** 0.405
State: NH **County:** Rockingham **Bay :** North Mill Pond
Source of information: Mr. Dan McHugh, Great Meadow Farms, Sawyer Island,
Rowley, MA 01969, (508) 948-7870
Comments: . Marsh name: NH Port Authority.

Marsh Inventory No.: 671 **COE Permit No.:** **Size (ha):** 1.214
State: NH **County:** Rockingham **Bay :** Rye Harbor
Source of information: Mr. Pete Helm, New Hampshire Office of State Planning,
Coastal Program, 2-1/2 Beacon Street, Concord, NH 03301, (603) 271-2155
Comments: . Marsh name: Awcomin Marsh.

Marsh Inventory No.: 388 **COE Permit No.:** **Size (ha):** 0.081
State: NJ **County:** **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc. . Marsh name: Tidal Stream.

Marsh Inventory No.: 404 **COE Permit No.:** **Size (ha):** 0.040
State: NJ **County:** **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc. . Marsh name: Sloop Creek
(C1-3).

Marsh Inventory No.: 409 **COE Permit No.:** **Size (ha):** 0.081
State: NJ **County:** **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc. . Marsh name: Monmouth
County.

Marsh Inventory No.: 302 **COE Permit No.:** **Size (ha):** 0.405
State: NJ **County:** **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Tidal marsh mitigation project.. Marsh name: SI-3B OSCA

Marsh Inventory No.: 300 **COE Permit No.:** **Size (ha):** 2.266
State: NJ **County:** **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box
P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc.. Marsh name: R-1 Contract

Marsh Inventory No.: 414 **COE Permit No.:** **Size (ha):** 0.121
State: NJ **County:** **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box

P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc. . Marsh name: Shark River Island.

Marsh Inventory No.: 481 **COE Permit No.:** **Size (ha):** 2.064

State: NJ **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc. . Marsh name: Newton Creek.

Marsh Inventory No.: 506 **COE Permit No.:** **Size (ha):** 2.266

State: NJ **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc. . Marsh name: Pennsauken Wetland.

Marsh Inventory No.: 540 **COE Permit No.:** **Size (ha):** 0.162

State: NJ **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc. . Marsh name: Tozour.

Marsh Inventory No.: 539 **COE Permit No.:** **Size (ha):** 1.416

State: NJ **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Built by Environmental Concern, Inc. . Marsh name: NJ Route 130.

Marsh Inventory No.: 353 **COE Permit No.:** **Size (ha):**

State: NY **County:** Bronx **Bay :** Pelham Bay

Source of information: Mr. Marc Matsil, Director of the Natural Resources Group, City of New York Parks and Recreation, 1234 5th Avenue, Rm. 233, New York, NY 10029, (212) 360-1417

Comments: This is a restoration project. Marsh was not planted as of 3/93.. Marsh name: Twin Island Marsh.

Marsh Inventory No.: 354 **COE Permit No.:** **Size (ha):**

State: NY **County:** Queens **Bay :**

Source of information: Mr. Marc Matsil, Director of the Natural Resources Group, City of New York Parks and Recreation, 1234 5th Avenue, Rm. 233, New York, NY 10029, (212) 360-1417

Comments: . Marsh name: Alley Creek.

Marsh Inventory No.: 355 **COE Permit No.:** **Size (ha):** 0.405

State: NY **County:** Richmond **Bay :** Arthur Kill; Old Place Creek

Source of information: Mr. Marc Matsil, Director of the Natural Resources Group, City of New York Parks and Recreation, 1234 5th Avenue, Rm. 233, New York, NY 10029, (212) 360-1417

Comments: .

Marsh Inventory No.: 285 **COE Permit No.:** **Size (ha):** 0.809

State: NY **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Shoreline erosion control project.. Marsh name: Centerport Harbor

Marsh Inventory No.: 290 **COE Permit No.:** **Size (ha):** 5.666

State: NY **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Dredged material stabilization project.. Marsh name: Seaford.

Marsh Inventory No.: 297 **COE Permit No.:** **Size (ha):** 4.047

State: NY **County:** **Bay :**

Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620

Comments: Tidal marsh restoration project.. Marsh name: Port Washington - Solomon

Marsh Inventory No.: 139 **COE Permit No.:** 88-N-010-0329 **Size (ha):** 0.728

State: NC **County:** Brunswick **Bay :** AIWW

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 138 **COE Permit No.:** 86-N-010-0178 **Size (ha):** 0.355

State: NC **County:** Brunswick **Bay :** AIWW

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 140 **COE Permit No.:** 85-N-015-0276 **Size (ha):** 0.012

State: NC **County:** Camden **Bay :** Albemarle Sound; Pasquotank River

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 146 **COE Permit No.:** 87-N-016-0307 **Size (ha):** 0.607

State: NC **County:** Carteret **Bay :** Newport River

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 147 **COE Permit No.:** 88-N-016-0372 **Size (ha):** 0.425

State: NC **County:** Carteret **Bay :** Pamlico Sound

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 148 **COE Permit No.:** 90-N-028-0301 **Size (ha):** 0.012

State: NC **County:** Dare **Bay :** Pamlico Sound

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 141 **COE Permit No.:** 91-N-065-0091 **Size (ha):** 0.045

State: NC **County:** New Hanover **Bay :** Cape Fear River

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 143 **COE Permit No.:** 85-N-065-0146 **Size (ha):** 0.005

State: NC **County:** New Hanover **Bay :** Masonboro Sound

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 145 **COE Permit No.:** 89-N-067-0381 **Size (ha):** 0.035

State: NC **County:** Onslow **Bay :** New River

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 142 **COE Permit No.:** 86-N-010-0057 **Size (ha):** 0.024

State: NC **County:** Pasquotank **Bay :** Little River

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 137 **COE Permit No.:** 86-N-071-0047 **Size (ha):** 0.486

State: NC **County:** Pender **Bay :** AIWW

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 136 **COE Permit No.:** 84-N-065-0033 **Size (ha):** 0.051

State: NC **County:** Bay : AIWW

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 135 **COE Permit No.:** 83-N-010-0103 **Size (ha):** 0.202

State: NC **County:** Bay : AIWW

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .**Marsh Inventory No.:** 649 **COE Permit No.:** **Size (ha):****State:** NC **County:** Bay : Albermarle Sound**Source of information:** Donald W. Hamer, U.S. Soil Conservation Service, 1536 Rt. 9 North, Cape May Court House, NJ 08210 (609) 465-5901**Comments:** No *S. alterniflora* planted..**Marsh Inventory No.:** 628 **COE Permit No.:** **Size (ha):** 0.405**State:** NC **County:** Bay : Bogue Sound**Source of information:** Dr. Stephen S. Broome, North Carolina State Univ., Dept. of Soil Science, Box 7619, Raleigh, NC 27695-7619, (919) 515-2643 (Broome, 1986)**Comments:** Was planted on 0.45, 0.6, and 0.9-m centers. Stand appears to have filled in well, forming a border marsh in front of a bulkhead..**Marsh Inventory No.:** 631 **COE Permit No.:** **Size (ha):****State:** NC **County:** Bay : Cape Fear River; Snow's Cut.**Source of information:** Dr. Christopher B. Craft, Duke Univ. Wetland Center, School of the Environment, Duke University, Durham, NC 27706, (919) 684-2619 (Craft, 1988)**Comments:** Was reported in a study comparing the organics and nutrients of man-made and natural *S. alterniflora* marshes. This marsh was about 13 years old..**Marsh Inventory No.:** 144 **COE Permit No.:** 84-N-065-0009 **Size (ha):** 0.006**State:** NC **County:** Bay : Middle Sound; Motts Channel**Source of information:** Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.**Comments:** .**Marsh Inventory No.:** 666 **COE Permit No.:** **Size (ha):** 1.214**State:** NC **County:** Bay : Pamlico Sound**Source of information:** Dr. Stephen S. Broome, North Carolina State Univ., Dept. of Soil Science, Box 7619, Raleigh, NC 27695-7619, (919) 515-2643 (Broome, 1989)**Comments:** Thought to be planted on 0.7-m centers. Stand appears to have filled in well, forming a well developed marsh along a tidal channel..**Marsh Inventory No.:** 629 **COE Permit No.:** **Size (ha):** 0.040**State:** NC **County:** Bay : Pamlico Sound; Neuse River**Source of information:** Dr. Stephen S. Broome, North Carolina State Univ., Dept. of Soil Science, Box 7619, Raleigh, NC 27695-7619, (919) 515-2643 (Broome, 1983)**Comments:** Test of fertilizers on *S. alterniflora* growth..**Marsh Inventory No.:** 632 **COE Permit No.:** **Size (ha):****State:** NC **County:** Bay : Pamlico Sound; Oregon Inlet.**Source of information:** Dr. Christopher B. Craft, Duke Univ. Wetland Center, School of the Environment, Duke University, Durham, NC 27706, (919) 684-2619 (Craft, 1988)**Comments:** Was reported in a study comparing the organics and nutrients of man-made and natural *S. alterniflora* marshes. This marsh was about 15 years old..**Marsh Inventory No.:** 640 **COE Permit No.:** **Size (ha):****State:** NC **County:** Bay : Pamlico Sound; Pamlico River; Durham Creek.**Source of information:** Dr. Christopher B. Craft, Duke Univ. Wetland Center, School of the Environment, Duke University, Durham, NC 27706, (919) 684-2619 (Craft, 1988)

Comments: Was reported in a study comparing the organics and nutrients of man-made and natural *S. alterniflora* marshes. This marsh was about 4 years old..

Marsh Inventory No.: 641 **COE Permit No.:** **Size (ha):**

State: NC **County:** **Bay :** Pamlico Sound; Pamlico River; Durham Creek.

Source of information: Dr. Christopher B. Craft, Duke Univ. Wetland Center, School of the Environment, Duke University, Durham, NC 27706, (919) 684-2619 (Craft, 1988)

Comments: Was reported in a study comparing the organics and nutrients of man-made and natural *S. alterniflora* marshes. This marsh was a little over 1-yr old..

Marsh Inventory No.: 630 **COE Permit No.:** **Size (ha):** 0.405

State: NC **County:** **Bay :** Roanoke Sound

Source of information: Dr. Stephen S. Broome, North Carolina State Univ., Dept. of Soil Science, Box 7619, Raleigh, NC 27695-7619, (919) 515-2643 (Broome, 1987)

Comments: This marsh was still surviving and protecting the shore from erosion in 1987..

Marsh Inventory No.: 657 **COE Permit No.:** **Size (ha):** 0.202

State: RI **County:** Washington **Bay :** Winnapaug Pond

Source of information: Mr. Paul J. Shea, Independent Environmental Consultants, Inc., Exeter, RI, (401) 294-3909

Comments: This was an inkind mitigation/restoration.

Marsh Inventory No.: 152 **COE Permit No.:** 90-2X-018 **Size (ha):** 0.121

State: SC **County:** Beaufort **Bay :** AIWW

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 168 **COE Permit No.:** 88-2A-101 **Size (ha):** 0.923

State: SC **County:** Beaufort **Bay :** Port Royal Sound

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 167 **COE Permit No.:** 90-3A-383 **Size (ha):** 0.032

State: SC **County:** Beaufort **Bay :** Port Royal Sound

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 166 **COE Permit No.:** 91-2A-004 **Size (ha):** 0.028

State: SC **County:** Beaufort **Bay :** Port Royal Sound

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 165 **COE Permit No.:** 84-2Z-031 **Size (ha):** 0.291

State: SC **County:** Beaufort **Bay :** Port Royal Sound; Beaufort River

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 153 **COE Permit No.:** 82-4C-294 **Size (ha):** 0.364

State: SC **County:** Charleston **Bay :** Ashley River

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 158 **COE Permit No.:** 89-2T-262 **Size (ha):** 6.071

State: SC **County:** Charleston **Bay :** Charleston Harbor

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 157 **COE Permit No.:** 90-2A-189 **Size (ha):** 0.291

State: SC **County:** Charleston **Bay :** Charleston Harbor

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 156 **COE Permit No.:** 88-2B-234 **Size (ha):** 0.486

State: SC **County:** Charleston **Bay :** Charleston Harbor

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 155 **COE Permit No.:** 86-2B-331 **Size (ha):** 2.226

State: SC **County:** Charleston **Bay :** Charleston Harbor

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 154 **COE Permit No.:** 85-3T-316 **Size (ha):** 0.081

State: SC **County:** Charleston **Bay :** Charleston Harbor

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 161 **COE Permit No.:** 85-3S-211 **Size (ha):** 0.049

State: SC **County:** Charleston **Bay :** Cooper River; Shem Creek

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 160 **COE Permit No.:** 84-3Z-320 **Size (ha):** 0.006
State: SC **County:** Charleston **Bay :** Cooper River; Wando River
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.
Comments: .

Marsh Inventory No.: 163 **COE Permit No.:** 90-2A-025 **Size (ha):** 0.405
State: SC **County:** Charleston **Bay :** North Edisto River
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.
Comments: .

Marsh Inventory No.: 171 **COE Permit No.:** 86-3D-219 **Size (ha):** 0.008
State: SC **County:** Charleston **Bay :** Stono River
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.
Comments: .

Marsh Inventory No.: 170 **COE Permit No.:** 85-2D-114 **Size (ha):** 0.005
State: SC **County:** Charleston **Bay :** Stono River
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.
Comments: .

Marsh Inventory No.: 169 **COE Permit No.:** 84-2Z-058 **Size (ha):** 0.728
State: SC **County:** Charleston **Bay :** Stono River
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.
Comments: .

Marsh Inventory No.: 149 **COE Permit No.:** 89-3X-217 **Size (ha):** 0.012
State: SC **County:** Charleston **Bay :**
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.
Comments: .

Marsh Inventory No.: 174 **COE Permit No.:** 89-3T-289 **Size (ha):** 0.174
State: SC **County:** Georgetown **Bay :** Winyah Bay
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.
Comments: .

Marsh Inventory No.: 173 **COE Permit No.:** 87-3D-298 **Size (ha):** 0.295
State: SC **County:** Georgetown **Bay :** Winyah Bay
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,

Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: .

Marsh Inventory No.: 151 **COE Permit No.:** 83-4Z-278 **Size (ha):** 0.093

State: SC **County:** Bay : AIWW

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: .

Marsh Inventory No.: 150 **COE Permit No.:** 82-3C-172 **Size (ha):** 0.202

State: SC **County:** Bay : AIWW

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: .

Marsh Inventory No.: 159 **COE Permit No.:** 82-3C-221 **Size (ha):** 0.232

State: SC **County:** Bay : Cooper River

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: .

Marsh Inventory No.: 162 **COE Permit No.:** 83-3Z-218 **Size (ha):** 0.102

State: SC **County:** Bay : Jarvis Creek

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: .

Marsh Inventory No.: 164 **COE Permit No.:** 83-5D-110 **Size (ha):** 0.809

State: SC **County:** Bay : Port Royal Sound; Broad River

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: .

Marsh Inventory No.: 172 **COE Permit No.:** 83-3Z-203 **Size (ha):** 0.019

State: SC **County:** Bay : Wando River

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: .

Marsh Inventory No.: 180 **COE Permit No.:** 18189 **Size (ha):** 0.243

State: TX **County:** Aransas **Bay :** Aransas Bay

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: .

Marsh Inventory No.: 178 **COE Permit No.:** 16622(01) **Size (ha):** 0.069
State: TX **County:** Aransas **Bay :** Aransas Bay
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.
Comments: .

Marsh Inventory No.: 176 **COE Permit No.:** 17326 **Size (ha):** 0.020
State: TX **County:** Aransas **Bay :** Aransas Bay
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.
Comments: .

Marsh Inventory No.: 175 **COE Permit No.:** 16622 **Size (ha):** 0.008
State: TX **County:** Aransas **Bay :** Aransas Bay
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.
Comments: .

Marsh Inventory No.: 227 **COE Permit No.:** 16830 **Size (ha):** 0.894
State: TX **County:** Brazoria **Bay :** Galveston Bay; West Bay; Oyster Creek
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.
Comments: .

Marsh Inventory No.: 215 **COE Permit No.:** 17026 **Size (ha):** 0.304
State: TX **County:** Calhoun **Bay :** Matagorda Bay; Carancahua Bay
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.
Comments: .

Marsh Inventory No.: 213 **COE Permit No.:** 16519 **Size (ha):** 0.020
State: TX **County:** Calhoun **Bay :** Matagorda Bay; Carancahua Bay
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.
Comments: .

Marsh Inventory No.: 216 **COE Permit No.:** 14541(02) **Size (ha):** 0.809
State: TX **County:** Calhoun **Bay :** Matagorda Bay; Lavaca Bay
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.
Comments: .

Marsh Inventory No.: 224 **COE Permit No.:** 17955 **Size (ha):** 0.202
State: TX **County:** Calhoun **Bay :** San Antonio Bay
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;

(813) 893-3503.

Comments: .

Marsh Inventory No.: 704 **COE Permit No.:** 15341(01) **Size (ha):**

State: TX **County:** Cameron **Bay :** Laguna Madre

Source of information: Mr. Gary Galbraith, Espey, Huston & Ass, Inc., (512) 327-6840

Comments: *S. alterniflora* was planted on 3-ft centers..

Marsh Inventory No.: 211 **COE Permit No.:** 17533 **Size (ha):** 0.040

State: TX **County:** Cameron **Bay :** Laguna Madre

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 209 **COE Permit No.:** 17526 **Size (ha):** 0.263

State: TX **County:** Cameron **Bay :** Laguna Madre

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 208 **COE Permit No.:** 16147 **Size (ha):** 0.527

State: TX **County:** Cameron **Bay :** Laguna Madre

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 699 **COE Permit No.:** 15537(05) **Size (ha):** 1.222

State: TX **County:** Cameron **Bay :** Laguna Madre; Lower

Source of information: Phoenix Huston Construction Co., Inc., P.O. Box 73562, Houston, TX 77273, (713) 580-4776

Comments: *S. alterniflora* was planted as single stems on 3-ft centers.. Marsh name: Boardwalk.

Marsh Inventory No.: 201 **COE Permit No.:** 17837 **Size (ha):** 3.238

State: TX **County:** Chambers **Bay :** Galveston Bay

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 198 **COE Permit No.:** 17664 **Size (ha):** 0.040

State: TX **County:** Chambers **Bay :** Galveston Bay

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 193 **COE Permit No.:** 17318 **Size (ha):** 1.113

State: TX **County:** Chambers **Bay :** Galveston Bay; East Bay; GIWW

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 205 **COE Permit No.:** 18052 **Size (ha):** 0.004

State: TX **County:** Galveston **Bay :** Galveston Bay

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 203 **COE Permit No.:** 17848 **Size (ha):** 0.567

State: TX **County:** Galveston **Bay :** Galveston Bay

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 200 **COE Permit No.:** 17962 **Size (ha):** 0.036

State: TX **County:** Galveston **Bay :** Galveston Bay

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 183 **COE Permit No.:** 13636(02) **Size (ha):** 0.040

State: TX **County:** Galveston **Bay :** Galveston Bay; Clear Lake

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 182 **COE Permit No.:** 13926(01) **Size (ha):** 0.186

State: TX **County:** Galveston **Bay :** Galveston Bay; Clear Lake; Jarbo Bayou

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 197 **COE Permit No.:** 16379 **Size (ha):** 0.405

State: TX **County:** Galveston **Bay :** Galveston Bay; Dickinson Bay

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 194 **COE Permit No.:** 18034 **Size (ha):** 0.022

State: TX **County:** Galveston **Bay :** Galveston Bay; East Bay

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 206 **COE Permit No.:** 18891 **Size (ha):** 0.081
State: TX **County:** Galveston **Bay :** Galveston Bay; Jones Bay
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.
Comments: .

Marsh Inventory No.: 234 **COE Permit No.:** 17800 **Size (ha):** 0.809
State: TX **County:** Galveston **Bay :** Galveston Bay; West Bay
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.
Comments: .

Marsh Inventory No.: 233 **COE Permit No.:** 18958 **Size (ha):** 1.918
State: TX **County:** Galveston **Bay :** Galveston Bay; West Bay
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.
Comments: .

Marsh Inventory No.: 231 **COE Permit No.:** 17875 **Size (ha):** 0.239
State: TX **County:** Galveston **Bay :** Galveston Bay; West Bay
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.
Comments: .

Marsh Inventory No.: 230 **COE Permit No.:** 17937 **Size (ha):** 0.004
State: TX **County:** Galveston **Bay :** Galveston Bay; West Bay
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.
Comments: .

Marsh Inventory No.: 225 **COE Permit No.:** 16677 **Size (ha):** 0.405
State: TX **County:** Galveston **Bay :** Galveston Bay; West Bay
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.
Comments: .

Marsh Inventory No.: 235 **COE Permit No.:** 18982 **Size (ha):** 0.081
State: TX **County:** Galveston **Bay :** Galveston Bay; West Bay; Offatts Bayou
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.
Comments: .

Marsh Inventory No.: 232 **COE Permit No.:** 18330/1 **Size (ha):** 2.258
State: TX **County:** Galveston **Bay :** Galveston Bay; West Bay; Offatts Bayou
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,

Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: .

Marsh Inventory No.: 228 **COE Permit No.:** 17073 **Size (ha):** 0.405
State: TX **County:** Galveston **Bay :** Galveston Bay; West Bay; Offatts Bayou
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: .

Marsh Inventory No.: 229 **COE Permit No.:** 17240 **Size (ha):** 0.567
State: TX **County:** Galveston ? **Bay :** Galveston Bay; West Bay
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: .

Marsh Inventory No.: 204 **COE Permit No.:** 18195 **Size (ha):** 0.040
State: TX **County:** Harris **Bay :** Galveston Bay
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: .

Marsh Inventory No.: 202 **COE Permit No.:** 17759 **Size (ha):** 1.578
State: TX **County:** Harris **Bay :** Galveston Bay
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: .

Marsh Inventory No.: 199 **COE Permit No.:** 17866 **Size (ha):** 0.093
State: TX **County:** Harris **Bay :** Galveston Bay
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: .

Marsh Inventory No.: 181 **COE Permit No.:** 16601(01) **Size (ha):** 0.012
State: TX **County:** Harris ? **Bay :** Galveston Bay; Clear Lake; Clear Creek
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: .

Marsh Inventory No.: 223 **COE Permit No.:** 19377 **Size (ha):** 0.121
State: TX **County:** Jefferson **Bay :** Sabine Lake
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.

Comments: .

Marsh Inventory No.: 190 **COE Permit No.:** 16533 **Size (ha):** 0.389
State: TX **County:** Nueces **Bay :** Corpus Christi Bay
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.
Comments: .

Marsh Inventory No.: 189 **COE Permit No.:** 17371 **Size (ha):** 0.809
State: TX **County:** Nueces **Bay :** Corpus Christi Bay
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.
Comments: .

Marsh Inventory No.: 188 **COE Permit No.:** 13192(05) **Size (ha):** 1.214
State: TX **County:** Nueces **Bay :** Corpus Christi Bay
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.
Comments: .

Marsh Inventory No.: 185 **COE Permit No.:** 16816 **Size (ha):** 0.567
State: TX **County:** Nueces **Bay :** Corpus Christi Bay
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.
Comments: .

Marsh Inventory No.: 187 **COE Permit No.:** 17191 **Size (ha):** 0.036
State: TX **County:** Nueces **Bay :** Corpus Christi Bay; La Quinta Channel
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.
Comments: .

Marsh Inventory No.: 217 **COE Permit No.:** 18020 **Size (ha):** 3.035
State: TX **County:** Nueces **Bay :** Corpus Christi Bay; Nueces Bay
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.
Comments: .

Marsh Inventory No.: 218 **COE Permit No.:** 16213 **Size (ha):** 1.680
State: TX **County:** Nueces **Bay :** Corpus Christi Bay; Oso Bay
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;
(813) 893-3503.
Comments: .

Marsh Inventory No.: 210 **COE Permit No.:** 17181 **Size (ha):** 0.101
State: TX **County:** Nueces **Bay :** Laguna Madre; GIWW
Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters,
Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702;

(813) 893-3503.

Comments: .

Marsh Inventory No.: 222 **COE Permit No.:** 18286 **Size (ha):** 0.279

State: TX **County:** Orange **Bay :** Sabine Lake

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 220 **COE Permit No.:** Permit # G16475 **Size (ha):** 0.142

State: TX **County:** San Patricio **Bay :** Aransas Bay; Redfish Bay

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 219 **COE Permit No.:** 16412 **Size (ha):** 0.186

State: TX **County:** San Patricio **Bay :** Aransas Bay; Redfish Bay

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 212 **COE Permit No.:** 16560 **Size (ha):** 3.359

State: TX **County:** San Patricio **Bay :** Aransas Bay; Redfish Bay; GIWW

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 192 **COE Permit No.:** 17847/1 **Size (ha):** 22.259

State: TX **County:** San Patricio **Bay :** Corpus Christi Bay

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 207 **COE Permit No.:** 16656 **Size (ha):** 0.101

State: TX **County:** Bay : Gulf of Mexico

Source of information: Mr. Andreas Mager, Jr., NMFS SE Region Headquarters, Habitat Conservation Division, 9721 Executive Center Drive, St. Petersburg, FL 33702; (813) 893-3503.

Comments: .

Marsh Inventory No.: 648 **COE Permit No.:** **Size (ha):**

State: VA **County:** Bay : Broad Bay

Source of information: Donald W. Hamer, U.S. Soil Conservation Service, 1536 Rt. 9 North, Cape May Court House, NJ 08210 (609) 465-5901

Comments: No *S. alterniflora* planted..

Marsh Inventory No.: 726 **COE Permit No.:** **Size (ha):**

State: VA **County:** Bay : Chesapeake Bay, James River

Source of information: (Hardaway et al, 1985)

Comments: A fringe marsh of *S. alterniflora* and *S. patens* was planted on 1.5-ft centers to control shore erosion.. Marsh name: Camp Chanco

Marsh Inventory No.: 731 **COE Permit No.:** **Size (ha):**

State: VA **County:** **Bay :** Chesapeake Bay;

Source of information: (Hardaway et al, 1985)

Comments: A fringe marsh of *S. alterniflora* and *S. patens* was planted on 1.5-ft centers to control shore erosion.. Marsh name: Tankard

Marsh Inventory No.: 745 **COE Permit No.:** **Size (ha):**

State: VA **County:** **Bay :** Chesapeake Bay;

Source of information: (Hardaway et al, 1985)

Comments: A fringe marsh of *S. alterniflora* was planted on 1.5-ft centers to control shore erosion. The marsh was replanted with *S. alterniflora* in June 1983.. Marsh name: Vanderslice

Marsh Inventory No.: 675 **COE Permit No.:** **Size (ha):**

State: VA **County:** **Bay :** Chesapeake Bay; Cherrystone Inlet; Wescoat Cove

Source of information: (Knudsen et al, 1982)

Comments: This is likely the oldest man-made *S. alterniflora* marsh created to protect the shore from erosion. The marsh was used as a test site to gage wave energy dissipation by a marsh..

Marsh Inventory No.: 737 **COE Permit No.:** **Size (ha):**

State: VA **County:** **Bay :** Chesapeake Bay; Fleets Bay; Tabbs Creek

Source of information: (Hardaway et al, 1985)

Comments: A fringe marsh of *S. alterniflora* was planted on 1.5-ft centers to control shore erosion. The marsh was replanted with *S. alterniflora* in June 1983.. Marsh name: Poole

Marsh Inventory No.: 742 **COE Permit No.:** **Size (ha):**

State: VA **County:** **Bay :** Chesapeake Bay; James River

Source of information: (Hardaway et al, 1985)

Comments: A fringe marsh of *S. alterniflora* and *S. patens* was planted on 1.5-ft centers to control shore erosion. The marsh was replanted with both species in June 1983, and again in July 1983.. Marsh name: King

Marsh Inventory No.: 741 **COE Permit No.:** **Size (ha):**

State: VA **County:** **Bay :** Chesapeake Bay; James River

Source of information: (Hardaway et al, 1985)

Comments: A fringe marsh of *S. alterniflora* and *S. patens* was planted on 1.5-ft centers to control shore erosion. The marsh was replanted with *S. alterniflora* in June 1983.. Marsh name: Mariner's Museum

Marsh Inventory No.: 739 **COE Permit No.:** **Size (ha):**

State: VA **County:** **Bay :** Chesapeake Bay; James River

Source of information: (Hardaway et al, 1985)

Comments: A fringe marsh of *S. alterniflora* was planted on 1.5-ft centers to control shore erosion. The marsh was replanted with *S. alterniflora* in June 1983.. Marsh name: Eley

Marsh Inventory No.: 674 **COE Permit No.:** **Size (ha):**

State: VA **County:** Bay : Chesapeake Bay; James River

Source of information: (Hardaway et al, 1985)

Comments: A fringe marsh of *S. alterniflora* was planted on 1.5-ft centers to control shore erosion.. **Marsh name:** Hog Island

Marsh Inventory No.: 676 **COE Permit No.:** **Size (ha):**

State: VA **County:** Bay : Chesapeake Bay; Kings Creek

Source of information: (Knudsen et al, 1982)

Comments: The marsh was used as a test site to gage wave energy dissipation by a marsh..

Marsh Inventory No.: 740 **COE Permit No.:** **Size (ha):**

State: VA **County:** Bay : Chesapeake Bay; Lynnhaven Roads; Lynnhaven Bay; Broad Bay

Source of information: (Hardaway et al, 1985)

Comments: A fringe marsh of *S. alterniflora* and *S. patens* was planted on 1.5-ft centers to control shore erosion. The marsh was replanted with *S. alterniflora* in June 1983.. **Marsh name:** Broad Bay Manor

Marsh Inventory No.: 744 **COE Permit No.:** **Size (ha):**

State: VA **County:** Bay : Chesapeake Bay; Lynnhaven Roads; Lynnhaven Bay; Broad Bay; Linkhorn Bay

Source of information: (Hardaway et al, 1985)

Comments: A fringe marsh of *S. alterniflora* was planted on 1.5-ft centers to control shore erosion. The marsh was replanted with *S. alterniflora* in June 1983.. **Marsh name:** Johnsen

Marsh Inventory No.: 678 **COE Permit No.:** **Size (ha):**

State: VA **County:** Bay : Chesapeake Bay; Mojack Bay

Source of information: (Hardaway et al, 1985)

Comments: A fringe marsh of *S. alterniflora* was planted on 1.5-ft centers to control shore erosion.. **Marsh name:** Mountjoy

Marsh Inventory No.: 743 **COE Permit No.:** **Size (ha):**

State: VA **County:** Bay : Chesapeake Bay; Occohannock Creek

Source of information: (Hardaway et al, 1985)

Comments: A fringe marsh of *S. alterniflora* and *S. patens* was planted on 1.5-ft centers to control shore erosion.. **Marsh name:** Marshall

Marsh Inventory No.: 736 **COE Permit No.:** **Size (ha):**

State: VA **County:** Bay : Chesapeake Bay; Piankatank River

Source of information: (Hardaway et al, 1985)

Comments: A fringe marsh of *S. alterniflora* and *S. patens* was planted on 1.5-ft centers to control shore erosion. *S. alterniflora* was replanted in May 1982, and both species were replanted in June 1983.. **Marsh name:** Murphy

Marsh Inventory No.: 729 **COE Permit No.:** **Size (ha):**

State: VA **County:** Bay : Chesapeake Bay; Potomac River

Source of information: (Hardaway et al, 1985)

Comments: A fringe marsh of *S. alterniflora* and *S. patens* was planted on 1.5-ft centers to control shore erosion.. **Marsh name:** Gill

Marsh Inventory No.: 735 **COE Permit No.:** **Size (ha):**

State: VA **County:** Bay : Chesapeake Bay; Rappahannock River

Source of information: (Hardaway et al, 1985)

Comments: A fringe marsh of *S. alterniflora* and *S. patens* was planted on 1.5-ft centers to control shore erosion. Was replanted with both species in May 1982 and again in June 1983.. Marsh name: Garrett

Marsh Inventory No.: 734 **COE Permit No.:** **Size (ha):**

State: VA **County:** Bay : Chesapeake Bay; Rappahannock River

Source of information: (Hardaway et al, 1985)

Comments: A fringe marsh of *S. alterniflora* and *S. patens* was planted on 1.5-ft centers to control shore erosion. *S. alterniflora* was replanted with both species in May 1982.. Marsh name: Durham West

Marsh Inventory No.: 733 **COE Permit No.:** **Size (ha):**

State: VA **County:** Bay : Chesapeake Bay; Rappahannock River

Source of information: (Hardaway et al, 1985)

Comments: A fringe marsh of *S. alterniflora* and *S. patens* was planted on 1.5-ft centers to control shore erosion. The marsh was replanted with both species in May 1982.. Marsh name: Durham South

Marsh Inventory No.: 732 **COE Permit No.:** **Size (ha):**

State: VA **County:** Bay : Chesapeake Bay; Rappahannock River

Source of information: (Hardaway et al, 1985)

Comments: A fringe marsh of *S. alterniflora* was planted on 1.5-ft centers to control shore erosion. *S. alterniflora* was replanted in June 1983. Marsh name: Wellford

Marsh Inventory No.: 727 **COE Permit No.:** **Size (ha):**

State: VA **County:** Bay : Chesapeake Bay; Rappahannock River

Source of information: (Hardaway et al, 1985)

Comments: A fringe marsh of *S. alterniflora* and *S. patens* and *S. patens* was planted on 1.5-ft centers to control shore erosion. The marsh was replanted in May, 1982.. Marsh name: Windmill Point

Marsh Inventory No.: 728 **COE Permit No.:** **Size (ha):**

State: VA **County:** Bay : Chesapeake Bay; Rappahannock River; Corrotoman River

Source of information: (Hardaway et al, 1985)

Comments: A fringe marsh of *S. alterniflora* and *S. patens* was planted on 1.5-ft centers to control shore erosion. *S. alterniflora* and *S. patens* was replanted in June, 1983.. Marsh name: Lee

Marsh Inventory No.: 746 **COE Permit No.:** **Size (ha):**

State: VA **County:** Bay : Chesapeake Bay; York River

Source of information: (Hardaway et al, 1985)

Comments: A fringe marsh of *S. alterniflora* was planted on 1.5-ft centers to control shore erosion. The marsh was replanted with *S. alterniflora* in June 1983.. Marsh name: Collier

Marsh Inventory No.: 738 **COE Permit No.:** **Size (ha):**

State: VA **County:** Bay : Chesapeake Bay; York River

Source of information: (Hardaway et al, 1985)

Comments: A fringe marsh of *S. alterniflora* was planted on 1.5-ft centers to control shore erosion. The marsh was replanted with *S. alterniflora* in June 1983.. Marsh name: York River State Park

Marsh Inventory No.: 747 **COE Permit No.:** **Size (ha):**
State: VA **County:** **Bay :** Chesapeake Bay; York River; Pamunkey River
Source of information: (Hardaway et al, 1985)
Comments: A fringe marsh of *S. alterniflora* was planted on 1.5-ft centers to control shore erosion.. **Marsh name:** Davis

Marsh Inventory No.: 730 **COE Permit No.:** **Size (ha):**
State: VA **County:** **Bay :** Chesapeake Bay; mouth of Occohannock Creek
Source of information: (Hardaway et al, 1985)
Comments: A fringe marsh of *S. alterniflora* and *S. patens* was planted on 1.5-ft centers to control shore erosion.. **Marsh name:** Hickman

Marsh Inventory No.: 652 **COE Permit No.:** **Size (ha):**
State: VA **County:** **Bay :** Occohannock Creek
Source of information: Donald W. Hamer, U.S. Soil Conservation Service, 1536 Rt. 9 North, Cape May Court House, NJ 08210 (609) 465-5901
Comments: .

Marsh Inventory No.: 389 **COE Permit No.:** **Size (ha):** 0.040
State: VA **County:** **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc. . **Marsh name:** White Stone.

Marsh Inventory No.: 291 **COE Permit No.:** **Size (ha):** 0.405
State: VA **County:** **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620
Comments: Shoreline erosion control project.. **Marsh name:** Eastville.

Marsh Inventory No.: 508 **COE Permit No.:** **Size (ha):** 0.162
State: VA **County:** **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc. . **Marsh name:** Chincoteague Mitigation.

Marsh Inventory No.: 507 **COE Permit No.:** **Size (ha):** 0.040
State: VA **County:** **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc. . **Marsh name:** Willis Wharf.

Marsh Inventory No.: 557 **COE Permit No.:** **Size (ha):** 0.202
State: VA **County:** **Bay :**
Source of information: Dr. Edgar Garbisch, Environmental Concern, Inc., P.O.Box P, St. Michaels, MD 21663, (410) 745-9620
Comments: Built by Environmental Concern, Inc. . **Marsh name:** VIDCO.

SECTION III

DIRECTORY OF HUMAN RESOURCES

Table of Contents

Section III -- Directory of Human Resources

FOREWORD	iii
ACKNOWLEDGMENTS	v
DIRECTORY	1
Alabama	1
California	2
Connecticut	3
Delaware	5
District of Columbia	7
Florida	7
Georgia	12
Louisiana	12
Maine	18
Maryland	19
Massachusetts	21
Mississippi	23
New Hampshire	25
New Jersey	25
New York	27
North Carolina	27
Oregon	29
Pennsylvania	30
Rhode Island	30
South Carolina	31
Texas	32
Virginia	37
Washington	38
DIRECTORY FORM	43
INDEX of PARTICIPANTS.	45

Foreword

Each year, the coastal areas of the United States are burdened by increasing populations, industrial growth, and expanded recreational use. These growths lead to increased use of estuaries that often threatens the well being and even the existence of coastal wetland habitats. The salt marshes that abounded along most of our estuaries have often been dredged or filled and thus converted for man's use. Many of these changes to salt marshes occurred before these marshes had been shown to serve many of the beneficial functions they do, such as: protect shorelines from erosion, reduce the impacts of floods, remove toxic wastes, and provide valuable habitat that supports fish and wildlife. Our increased understanding of the way these marshes function has promoted a national desire to preserve and restore coastal wetlands.

The purpose of this directory is to provide our readers with a list of people involved in salt marsh ecology, with particular emphasis on those working on enhancement, restoration, and creation of *Spartina alterniflora* marshes. We hope and expect the directory will be of use to aid in information exchange among scientists working in this field, and between scientists and lay people concerned about their local estuary or marsh, coastal landowners interested in preserving their property, and companies needing assistance with habitat mitigation. This directory is not a "complete" listing of all persons and organizations that are involved and have pertinent knowledge on our subject. It is a list of those who kindly responded to our inquiries and who recognize the directory as another tool to help in the preserving and restoring of salt marshes and coastal wetlands. The directory is intended to be "living" in the sense that periodic updates are anticipated. We will be pleased to add people we missed if you will fill in a directory form (at the back of this directory) and return it to us.

The directory is organized alphabetically by state. People are listed alphabetically by last name under the state in their address. An alphabetical index of participants is included, and is ordered by last names and initials. A directory form is included at the back.

Acknowledgments

To those of you included in this directory, our sincere thanks for your participation. And if we have omitted an important player, it was purely accidental.

We wish to thank our four regional assistants who helped find many participants and recorded the early entries.

Northeast:	Mr. Chris Luis Sardella
Atlantic:	Mr. Andrew L. Bunch
Gulf:	Dr. Joseph L. Staton
West:	Mr. Russell E. DiFiori

We also thank the NMFS Southeast Fisheries Science Center and the Coastal Ocean Program for sponsoring this directory project as part of a grant to study the technology and success in restoration, enhancement, and creation of salt marshes in the United States.

Good Planting!

Directory

Alabama

Mr. Hugh M. Dowling
Dowling Environmental Services
P.O. Box 66003
Mobile, AL 36660
Work phone: (205) 476-2010
FAX: (205) 476-2011

President.

Projects:

Has planted 30-40 projects over the last 10 years. Marsh plantings are guaranteed.

Mr. Steve Heath
Alabama Marine Resources
P.O. Box 189
Dauphin Island, AL 36528
Work phone: (205) 861-2882
FAX: (205) 861-8741

Fisheries resource and habitat manager; coastal waters and wetlands, and their fisheries.

Projects:

1. Reviews proposed alterations of coastal habitat to preserve fisheries and wildlife habitat.
2. Monitors progress of mitigation efforts, some involving *Spartina alterniflora* salt marshes.
3. Assists in evaluating fisheries, and in developing fisheries management strategies.

Mr. Arthur L. Middleton
U.S. Army Corps of Engineers, Mobile District
Operations Division, Regulatory Branch
P.O. Box 2288
Mobile, AL 36628-0001
Work phone: (205) 694-3786
FAX: (205) 694-4264

Supervisory Ecologist.

Projects:

1. Presently (1992) installing in Mobile, Alabama, Navy home port site, a *Spartina alterniflora* marsh and a *Juncus* marsh of about 25 acres (unsure about percentages of each marsh type).
2. Delta Port development community along the west of Fowl River. Combination of *Spartina alterniflora* and *Juncus*, about 45 acres installed since 1987.

Mr. Edwin R. "Randy" Roach
U.S. Fish and Wildlife Service
Division of Ecological Services
P.O. Box 1197
Daphne, AL 36526
Work phone: (205) 690-2181
FAX: -

Senior Field Biologist.

Projects:

Naval Station, Mobile, AL. Marsh restoration with 75% *Juncus roemerianus* and 25% *Spartina alterniflora*.

Dr. Judy P. Stout
University of Southern Alabama
Marine Environmental Sciences Consortium, Dauphin Island Sea Lab
P.O. Box 369-370
Dauphin Island, AL 36528
Work phone: (205) 861-7529
FAX: (205) 861-4646

Associate Director; Estuarine Ecologist.

Projects:

1. Dauphin Island Sea Lab; 1-1.5 acre marsh restoration began in January 1993; plantings of 50% *S. alterniflora* and 50% *J. roemerianus*.
2. Evaluation of coastal wetlands from mitigation in coastal Alabama; final report available and includes created *S. alterniflora* wetlands.

Mr. Thomas D. Thornhill
U.S. Fish and Wildlife Service
P.O. Drawer 1190
Daphne, AL 36526
Work phone: (205) 690-2181
FAX: (205) 694-4222

Fish and Wildlife Biologist.

Projects:

1. Ongoing monitoring of mitigation project for Chevron of Pascagoula; planted earlier and destroyed by hurricane--replanted in 1988 (approx.) *Spartina* and *Juncus* marsh that is surviving well.
2. Planned site for Gulfport not yet complete by Vitor and Associates.

Dr. Barry A. Vittor

Barry A. Vittor & Associates, Inc.
Environmental Research & Consulting
8060 Cottage Hill Road
Mobile, AL 36695

Work phone: (205) 633-6100

FAX: (205) 633-6738

Director; wetlands creation/restoration.

Projects:

1. Restoration and creation of wetlands including *Spartina alterniflora* marsh along the West Fowl River near Mobile, AL.
2. Port Authority, Gulfport; tidal wetland creation at Bayou Portage.
3. Naval Station at Mobile; nursery work for *S. alterniflora* for a 25-acre tidal marsh on Mobile Bay.
4. LPM, Inc.; tidal influenced fresh marsh restoration at Chickasaw Creek (Mobile River).
5. AmWest Savings; creation of a tidal influenced marsh in a drainage swale on Perdido Bay, FL.

California**Mr. James R. Bybee**

National Marine Fisheries Service, SW Region
Habitat Conservation Division
777 Sonoma Ave., Room 325
Santa Rosa, CA 95404

Work phone: (707) 578-7513

FAX: (707) 578-3435

Environmental Coordinator, Northern California.

Projects:

1. Habitat protection for fisheries and fisheries related populations.

Mr. Curtis Daehler

University of California, Davis
Department of Zoology
P.O. Box 247 Bodega Marine Lab
Bodega Bay, CA 94923

Work phone: (707) 875-2022

FAX: (707) 875-2089

Graduate Student in Zoology; wetland entomology.

Projects:

1. Location, spread, genetics and ecology of alien *Spartina alterniflora* in Pacific marshes, especially San Francisco and San Pablo bays.
2. Effects of native herbivores on this alien weed: seed set and spread as affected by herbivory.

Ms. Francesca C. Demgen

Woodward-Clyde Consultants
Environmental
2014 Indiana
Vallejo, CA 94590

Work phone: (510) 874-1731

FAX: (510) 874-3268

Wetland design and monitoring specialist;
wastewater, freshwater, and estuarine.

Projects:

1. 1993. Research metal concentrations in a 145 acre man-made fresh and brackish water wetland which receives treated secondary effluent. Plant emergent vegetation to enhance nitrification.
2. 1990-93. Design, oversee construction, and monitor success of 19 acre freshwater marsh and creek restoration. Successful planting of riparian trees and shrubs, hydroseeding of emergents, and redistribution of site soils with seed stock.
3. 1990-93. Study uptake of copper, lead, and zinc in emergent vegetation and sediments in creek and marsh receiving urban run-off.

Mr. Russel E. DiFiori

5460 55th Street, #45
San Diego, CA 92115

Work phone: (619) 594-6522

421-6700 Ext.5286

FAX: -

Botanist. Wetlands ecology.

Projects:

1. Assisted in NMFS directory of people involved with restoration and creation of *Spartina spp.* marshes.

Mr. Brian Fink

Habitat Restoration
3234 "H" Ashford St.
San Diego, CA 92111

Work phone: (619) 279-8769

FAX: -

Owner/manager; horticulturist, wetlands.

Projects:

1. Wetland plants research: their growing requirements and pollination.
2. Contract growing of *Cordylanthus maritimus maritimus* and other sensitive plants.
3. Cultivation of *Spartina foliosa* studied for Caltrans.
4. Establishment of *Salicornia virginica* is being studied for the Pacific Estuarine Research Lab.

Mr. Robert S. Hoffman

National Marine Fisheries Service, SW Region
Habitat Conservation Div.
501 W. Ocean Blvd., Suite 4200
Long Beach, CA 90802-4213

Work phone: (310) 980-4000

FAX: (310) 980-4018

Fishery biologist; seagrass (*Zostera*) ecology.

Projects:

Evaluates habitat impacts of coastal waters, particularly as they relate to fisheries organisms. Involved with the long-term monitoring of a *Spartina* transplant project in San Diego Bay.

Dr. Michael Josselyn

San Francisco State University
Romberg Tiburon Center
P.O. Box 855
Tiburon, CA 94920

Work phone: (415) 435-7126

FAX: (415) 435-7120

Estuarine ecologist; wetland evaluations, design and restoration.

Projects:

1. Ecological consequences of introduced *Spartina alterniflora* on San Francisco Bay. 2. Control mechanisms to eradicate *S. alterniflora*; carbohydrate chemistry and means of control.

Mr. James R. Smith

Pacific Gas & Electric Co.
3400 Crow Canyon Rd.
San Ramon, CA 94583

Work phone: (510) 866-5828

FAX: (510) 866-5915

Biologist.

Projects:

1. Pittsburg (6/1/92); restoration of brackish tidal marshlands and native uplands in Suisun Bay, San Francisco, CA; about 1500 acres.
2. McArthur Swamp (1/2/92); restoration of permanent and seasonal freshwater wetlands in Fall River Valley, Shasta County, CA; about 1000 acres.

Mr. Douglas Spicher

Wetlands Research Associates, Inc.
2169-G East Francisco Blvd.
San Rafael, CA 94901

Work phone: (415) 454-8868

FAX: (415) 454-0129

Projects:

Studied the distribution and taxonomy of *Spartina* (Gramineae) in northern California.

Dr. Donald R. Strong

University of California, Davis
Bodega Marine Laboratory
P.O. Box 247, Bodega Marine Lab
Bodega Bay, CA 94923

Work phone: (707) 875-2022

FAX: (707) 875-2089

Professor; Zoology, wetland entomology.

Projects:

1. Location, spread, genetics and ecology of alien *Spartina alterniflora* in Pacific marshes, especially San Francisco and San Pablo bays.

2. Effects of native herbivores on this alien weed: seed set and spread as affected by herbivory.

Dr. Sue L. Williams

San Diego State University
Pacific Estuarine Research Lab
Biology Department
San Diego, CA 92182-0057

Work phone: (619) 594-6522

FAX: (619) 594-5676

Professor/Researcher; wetland ecosystems.

Projects:

1. Research needs for restoring sustainable coastal ecosystems on the Pacific Coast.

Dr. Joy B. Zedler

San Diego State University
Pacific Estuarine Research Lab
Biology Department
San Diego, CA 92182-0057

Work phone: (619) 594-5809

FAX: (619) 594-5676

Director PERL, Professor/Researcher; wetland ecosystems.

Projects:

1. Assessed functional equivalency of marshes in the San Diego area; the Caltrans Connector Marsh compared with the Sweetwater River Wetlands Complex.
2. Canopy architecture of natural and planted cordgrass marshes: selecting habitat evaluation criteria.
3. Research needs for restoring sustainable coastal ecosystems on the Pacific Coast.

Mr. John J. Zentner

Zentner & Zentner
925 Ygnacio Valley Rd, Suite 250
Walnut Creek, CA 94596

Work phone: (510) 947-6888

FAX: (510) 947-1798

Principal; Wetland restorations

Projects:

1. Warm Springs I, Fremont, CA. Created salt marsh (muted tidal).
2. Mariner's Cove, Vallejo, CA. Created tidal salt marsh.

Connecticut

Mr. Peter F. Alexander, ASLA, APA

Sound Shore Institute, Site Design
6 West Putnam Ave.
Greenwich, CT 06830

Work phone: (203) 869-6895

FAX: (203) 869-2005

Landscape architect; GIS based CAD designer/planner

Projects:

1. Katama Bay replanting, 1974.
2. Metzger Marsh at Stony Creek, 1979.
3. Byram Mianus River restoration, 1989.
4. Grass Island, 1992.

Mr. Michael A. Aurelia

Town of Greenwich
Inland Wetlands and Watercourses Agency
101 Field Point Road
Greenwich, CT 06830

Work phone: (203) 622-7736

FAX: (203) 622-3767

Director, regulatory planning and management; freshwater and marine wetlands scientist.

Projects:

1. Initiated the Greenwich Point project on Long Island Sound, where 2 acres of intertidal beach was planted with *Spartina alterniflora* by E. Garbisch and Environmental Concern back in 1975. Continues to monitor the well being of this marsh.
2. Involved with developing regulatory policy regarding freshwater and marine wetlands, their use and their preservation.

Mr. Mark Heller

S.E. Minor and Co., Inc.
81 Holly Hill Lane
Greenwich, CT 06830

Work phone: (203) 869-0136

FAX: (203) 869-7869

Landscape architect; design of coastal and freshwater wetlands.

Projects:

1. Coscob Harbor; Greenwich, CT. Designed a coastal *Spartina alterniflora* marsh for mitigation. Project involved dredging an area of marsh where *Phragmites australis* had taken over, to restore tidal salt water flows.
2. Designed several other freshwater and salt water marsh creation/restoration projects.

Mr. Richard E. Kent

Environmental Design Associates, P.C.
P.O. Box 247, 78 Danbury Road
Wilton, CT 06897

Work phone: (203) 762-8020

FAX: (203) 762-7881

Director; wetlands and uplands landscape architect and designer.

Projects:

1. Mianus River Bridge project to create a *Spartina alterniflora* salt marsh; 0.3 acres of intertidal area were planted with *S. alterniflora*, good growth was obtained before the Canadian geese ate the tops off.
2. Conair Corp.- restored a 0.2 acre *Spartina alterniflora* marsh and wading pool from a *Phragmites* colony.
3. Mansfield Landing- restored a tidal wetland by excavating fill material to appropriate grades; allowing revegetation through natural seeding from nearby marshes.

Dr. William A. Niering

Connecticut College
Botany Department
270 Mohegan Ave.
New London, CT 06320

Work phone: (203) 439-2143

FAX: (203) 439-2700

Professor; botany, wetlands ecology.

Projects:

1. Connecticut shoreline marshes, evaluation of plant succession.
2. Plant community change with the restoration of tidal flows to a 20 hectare coastal marsh in Stonington, Connecticut. GIS analysis of the changes showed *Typha angustifolia* habitat was greatly reduced while *Spartina alterniflora* habitat was greatly increased. *Phragmites* habitat increased only slightly.

Mr. Matthew J. Popp

Environmental Design Associates, P.C.
P.O. Box 247, 78 Danbury Road
Wilton, CT 06897

Work phone: (203) 762-8020

FAX: (203) 762-7881

Environmental analyst and landscape architect; site development and environmental planning.

Projects:

1. Mianus River Bridge project to create a *Spartina alterniflora* salt marsh.
2. Conair Corp.- restored a *Spartina alterniflora* marsh and wading pool from a *Phragmites* colony.
3. Mansfield Landing- restored a tidal wetland by excavating fill material to appropriate grades; allowing revegetation through natural seeding from nearby marshes.

Mr. Erwin N. Potter

Erwin Potter and Associates, Environmental System Planning, Ltd.
P.O. Box 839
Sandy Hook, CT 06482

Work phone: (203) 426-5936

FAX: (203) 426-8917

Landscape architect. Coastal landscape design.

Mr. Ron Rozsa

Connecticut Dept. of Environmental Protection,
Office of the Long Island Sound Program
79 Elm Street
Hartford, CT 06106
Work phone: (203) 566-7404

Ms. Judith A. Slayback

Environmental Design Associates, P.C.
P.O. Box 247, 78 Danbury Road
Wilton, CT 06897
Work phone: (203) 762-8020
FAX: (203) 762-7881

Senior Environmental Analyst; environmental planning and restoration, freshwater and marine.

Projects:

1. Mianus River Bridge project to create a *Spartina alterniflora* salt marsh; 0.3 acres of intertidal area were planted with *S. alterniflora*, good growth was obtained before the Canadian geese ate the tops off.
2. Conair Corp.- restored a 0.2 acre *Spartina alterniflora* marsh and wading pool from a *Phragmites* colony.
3. Mansfield Landing- restored a tidal wetland by excavating fill material to appropriate grades; allowing revegetation through natural seeding from nearby marshes.

Mr. Thomas J. Steinke

Conservation Commission, Town of Fairfield
Independence Hall, 725 Old Post Road
Fairfield, CT 06430

Work phone: (203) 256-3071
FAX: (203) 256-3080

Conservation Director for the Town of Fairfield.

Projects:

1. Turney Creek: 1973, restored tidal flows using self-regulating tide gates (SRT) and open marsh water management (OMWM) to a 16-acre *Phragmites* marsh diked for flood control: elimination of marsh fires, mosquitoes, reduction of *Phragmites*; self-regulating tidegate.
2. Ash Creek: 1974, restored tidal flows (SRT and OMWM) to a 14-acre *Phragmites* marsh diked for flood control and marina development. Elimination of *Phragmites* and fires, reduction of mosquitoes, restoration of site.
3. Pine Creek: 1979, restored tidal flows (SRT, dikes, & OMWM) to a 130-acre marsh diked for flood control, roads, garbage dump, subdivisions, parks. Elimination of fires, *Phragmites*, odors, flooding, and mosquitoes.
4. Sasco Creek: 1987, restored tidal flows to 30+ acres of salt marsh isolated by causeway, highway and railroad bridges. Rebuilt larger highway bridge and cleaned out 140 years of debris from RR bridge. Greatly improved flushing eliminated fires, reduced mosquitoes and *Phragmites*.
5. Great Creek (Milford, Connecticut) 1990. Assisted State of Connecticut in Silver

Sands State Park flood control and 150-acre marsh restoration effort with self-regulating tidegates. This solved the flooding, fire and mosquito problems, and restored shellfish, waterfowl, and fish habitats.

Dr. R. Scott Warren

Connecticut College
Botany Department
270 Mohegan Avenue
Box 5362, Conn. College
New London, CT 06320-4196

Work phone: (203) 439-2132
FAX: (203) 439-2519

Professor of Botany; tidal marsh ecology, physiological ecology; fish and invertebrate ecology in coastal wetlands.

Projects:

1. Restoration of an impounded tidal marsh at the Barn Island Wildlife Management Area - Stonington, CT. Effects of restored tidal flows to the area have been monitored for changes in flora and fauna. 1980 to present.
2. Marsh restoration through dredged material removal at Mumford Cove, Blaff Point Coastal Reserve - Groton, CT. Material was removed and surface was re-graded before dikes were breached to restore tidal exchange. Flora and fauna changes have been monitored since 1989.

Delaware**Mr. Bennett N. Anderson**

Delaware Dept. of Natural Resources & Environmental Control, Division of Water Resources, Watershed Assessment Branch
P.O. Box 1401, 89 Kings Highway
Dover, DE 19901

Work phone: (302) 739-4590
FAX: (302) 739-6140

Environmental Scientist. Estuarine Science; Oil Spill Response and Restoration

Projects:

Various EPA superfund and oil spill studies.

Mr. John H. Clark

Delaware Dept. of Natural Resources & Environmental Control, Division of Fish and Wildlife
P.O. Box 1401, 89 Kings Highway
Dover, DE 19903

Work phone: (302) 739-4782
FAX: (302) 739-6780

Environmental scientist.

Projects:

1. 1989-1994. Impoundment - estuarine interactions. Examining the effects of recently implemented water management plans enhancing tidal exchange for two high level tidal impoundments. Studying the in and out

movements and survival of fish associated with these impoundments.
 2. 1993-1994. Comparing fish communities in the impoundments with those in adjacent natural marshes.

Dr. John L. Gallagher

University of Delaware
 College of Marine Studies, Halophyte Biology Lab
 700 Pilottown Rd.
 Lewes, DE 19958

Work phone: (302) 645-4366

FAX: (302) 645-4028

Professor. Biochemistry of halophyte and marsh ecology.
 Projects:

1. Phenotypic growth of *S. alterniflora*.
2. Monitoring restored salt marsh growth and function.

Dr. George W. Luther, III

University of Delaware
 College of Marine Studies
 Lewes, DE 19958

Work phone: (302) 645-4208

FAX: (302) 645-4007

Professor; Geochemistry of salt marshes; iron and sulfur chemistry.

Projects:

1. Geochemical studies of iron and sulfur in various salt marshes in N.J., in the Great Sippewissett, MA, and in the Great Marsh, DE.
2. Additional biogeochemical studies were done in Delaware Bay, Chesapeake Bay, Newark Bay, and inland bays of Delaware.
3. Mid-Atlantic Bight Coastal work from NY to Cape Hatteras involved geochemical analyses of iron and sulfur in sediments and the water column.

Dr. Evelyn Maurmeyer

Coastal and Estuarine Research
 Marine Studies Complex
 P.O. Box 674
 Lewes, DE 19958

Work phone: (302) 645-9610

FAX: (302) 645-4006

Marine scientist; coastal geology, wetlands mitigation consulting.

Projects:

1. Delaware Bay shoreline erosion control near Lower Township, Cape May, NJ.
2. Wetlands delineations for coastal areas in Delaware.
3. Mitigation projects restoring coastal marshes using *Spartina alterniflora* and *S. patens*.

Mr. Richard McCorkle

U.S. Fish and Wildlife Service
 Delaware Bay Estuary Project
 RD #1 Box 146A
 Smyrna, DE 19977

Work phone: (302) 653-9152

FAX: (302) 653-9421

Fish and wildlife biologist

Projects:

1. Partners for Wildlife Program. "Gordon's Pond" project enhanced the salt marsh habitat in part of a 350-acre track by renovating water control structures to allow more tidal exchange. This reduced the *Phragmites* area, and increased the *Spartina alterniflora* and *S. patens* area.
2. At the Eugene Mills site, saltmarsh was restored on privately owned land by excavating the area to remove *Phragmites* and to improve freshwater inflow as well as tidal saltwater inflow.

Dr. Kurt R. Phillipp

Wetlands Research Associates, Inc.
 102 E Main Street, Suite 305
 Newark, DE 19711-7319

Work phone: (302) 738-7535

FAX: (302) 738-9173

Wetlands ecologist; consultant.

Projects:

1. Historic review of tidal wetlands of the Delaware Estuary.
2. Assist the state of Delaware in developing a tidal wetlands characterization and preservation plan.
3. Evaluate impacts to tidal wetlands from hazardous waste sites; Delaware, and New Jersey, and restore tidal wetlands.
4. Develop freshwater wetlands to mitigate impacts of the Dulles Toll Road in northern Virginia, and of road work by the Delaware DOT in northern Delaware.

Dr. Denise M. Seliskar

University of Delaware
 College of Marine Studies, Halophyte Biology Laboratory
 700 Pilottown Road
 Lewes, DE 19958

Work phone: (302) 645-4366

FAX: (302) 645-4028

Associate Research Scientist.

Projects:

1. Growth characteristics and tissue culture of *Spartina alterniflora* and *S. patens*.
2. Accelerating salt marsh functional development through plant genotype selection: intra-specific diversity from natural populations and tissue culture.
3. The effect of plant genotype on the food web in a created marsh.

Ms. Tracey Skrabal

Dept. of Natural Resources and Environmental Control,
 Division of Water Resources, Wetlands and Aquatic
 Protection Branch
 89 Kings Highway, Box 1401
 Dover, DE 19903

Work phone: (302) 739-4691

FAX: (302) 739-3491
Environmental Scientist.

Dr. William R. Witman

Delaware Dept. of Natural Resources & Environmental
Control, Division of Fish and Wildlife
P.O. Box 1401, 89 Kings Highway
Dover, DE 19901

Work phone: (302) 739-5297

FAX: (302) 739-6157

Biologist. Waterfowl management and habitat
conservation.

Projects:

1. Compiling, editing and publishing a waterfowl habitat management guide for the Atlantic Flyway.
2. Participant in the USFWS's Partners for Wildlife Program. Habitat advisor.
3. Technical representative to the North American Waterfowl Management Plan; including serving on the Policy Committee.

District of Columbia

Dr. Mary C. Barber

Science and Policy Associates
West Tower, Suite 400
1333 H Street NW
Washington, DC 20005

Work phone: (202) 789-1201

FAX: (202) 789-1206

Consultant-Senior Environmental Scientist.

Projects:

1. Competition between salt marsh plants, 1982 (thesis-not published)

Florida

Ms. Lisa Adams

Florida Dept. of Environmental Regulation,
NE District Office
7825 Baymeadows Way, Suite 200B
Jacksonville, FL 32256-7577

Work phone: (904) 448-4300

FAX: -

Environmental Specialist / Compliance of Permits

Projects:

1. Ocean Highway and Port Authority, 1993. Mitigation project involving scrape-down and planting of *Spartina alterniflora*.
2. Dames Point project. Establishment of *S. alterniflora* marshes. Has filled in well.

3. Amelia Island Yacht Club project. Restoration of marshes along a dredged material island and access road.
4. Anastasia Sanitary District pipeline project. Establish marsh over a sewer pipeline running through a marsh; is slowly filling in.

Mr. Carl Bates

Plants for Tomorrow, Inc.
16361 Norris Road
Loxahatchee, FL 33470-9430

Work phone: (800) 448-2525

FAX: (407) 790-1916

Grower/planter/designer; *Spartina* restorations.

Projects:

1. Grows and supplies *Spartina bakeri*, *S. alterniflora*, and *S. patens* for restoration of habitats. Has done many projects.
2. Currently installing a large *S. alterniflora* marsh at Morris Island near Charleston, SC.

Mr. Steve Beeman

Ecoshores, Inc.
Shoreline Associates
3881 South Nova Road
Port Orange, FL 32127

Work phone: (904) 767-6232

FAX: (904) 759-9895

Grower/supplier of *Spartina spp.*

Projects:

1. Grows and supplies *Spartina alterniflora*, *S. bakeri*, and *S. patens*.
2. Designs and builds coastal marshes and wetlands. Over 50 well established.
3. A varied wetlands habitat project at Admirals Cove near Jupiter, FL. Includes *S. alterniflora* marsh creation in the 60-acre site.

Mr. James William Beever, III

Florida Game & Freshwater Fish,
Office of Environmental Services
29200 Tuckers Grade
Punta Gorda, FL 33955

Work phone: (813) 639-3515

FAX: -

Biologist. Mangrove ecosystem and high marsh ecology.

Mr. W.Y. Bennett

Florida Aquatics of Avon Park, Inc.
3846 Menendez Drive
Pensacola, FL 32503

Work phone: (904) 438-2585

FAX: -

Grower/supplier of *Spartina alterniflora*.

Projects:

Suppliers of *Spartina alterniflora*.

Mr. David M. Biggar, Jr.

Liner Farm, Inc.
P.O. Box 701369
Saint Cloud, FL 34770-1369

Work phone: (407) 892-1484

FAX: (407) 892-3593

Grower/supplier of *Spartina spp.*

Projects:

Grow and supplies *Spartina alterniflora*, *S. patens*, and *S. bakeri* for restoration and other plantings.

Mr. William F. Bissett

The Natives
2929 J.B. Carter Road
Davenport, FL 33837

Work phone: (813) 422-6664

FAX: -

Landscape architect/builder.

Ms. Nancy Bissett

The Natives
2929 J.B. Carter Road
Davenport, FL 33837

Work phone: (813) 422-6664

FAX: -

Horticulturist.

Projects:

Grow *Spartina bakeri*, sand cordgrass.

Ms. Lucy Blair

Florida Dept. of Environmental Regulation,
S. District Office
2295 Victoria Ave., Suite 364
Fort Myers, FL 33901

Work phone: (813) 332-6975

FAX: -

Environmental Specialist

Projects:

Mr. Ronald E. Brockmeyer, Jr.

Harbor Branch Oceanographic Institution, Inc.
5600 U.S. 1, North
Ft. Pierce, FL 34946

Work phone: (407) 465-2400

Ext. 668

FAX: (407) 465-2446

Biologist. Fish biology and marsh ecology.

Mr. Otto M. Bundy

Nautilus Environmental Services, Inc.
P.O. Box 497
Parish, FL 34219

Work phone: (800) 771-4114, (800) 422-

6287 (in Florida)

FAX: (813) 766-2410

Grower/Sales Rep.

Projects:

1. Consults in the planting of *Spartina spp.*
2. Grows *Spartina alterniflora*, *S. patens*, *S. bakeri* and *S. spartinae* for restoration projects.

Mr. Allen G. Burdett, Jr.

Florida Dept. of Environmental Regulation
Water Management
4520 Oak Fair Blvd.
Tampa, FL -

Work phone: (813) 620-6121

FAX: (813) 622-6123

Environmental Restoration Coordinator. Habitat restoration, creation and resource management.

Projects:

Involved with planning and managing of several projects of coastal restoration.

Mr. Douglas B. Carlson

Indian River Mosquito Control District
5655 South Gifford Road
P.O. Box 670

Vero Beach, FL 32961-0670

Work phone: (407) 562-2393

FAX: (407) 562-9619

Assistant Director. Mosquito control and salt marsh/wetland management.

Projects:

1. Developing, reviewing and implementing management plans for salt marshes (high marsh - *Batis*, *Salicornia*, *Distichlis* and White and black mangrove) along the Indian River lagoon for multipurpose use.
2. Chairman of the Subcommittee on Managed Marshes, a legislatively established interagency committee responsible for helping develop and review marsh management plans.

Mr. Kim Cash

Florida Shrub Growers
730 Myakka Road
Sarasota, FL 34240

Work phone: (813) 322-1915

FAX: -

Salesperson.

Projects:

Grows and supplies *Spartina alterniflora*, smooth cord grass, and *S. bakeri*.

Mr. Jerry Cordy

CZR, Inc.
4494 Southside Blvd., Suite 200
Jacksonville, FL 32216

Work phone: (904) 642-8507

FAX: (904) 642-8911

Biologist, consulting, wetland delineation, *Spartina alterniflora* marsh creation.

Projects: (more)

1. Dames Point Bridge, May 1987; created a 9 acre *S. alterniflora* marsh from upland area to offset impacts from highway bridge construction. Jacksonville, Florida.
2. River Crossing, July 1987; create a 1 acre *S. alterniflora* marsh of offset impacts from project entrance road. Jacksonville, Florida.

Mr. David W. Crewz

Florida Marine Research Institute (FMRI)
100 8th Avenue SE
St. Petersburg, FL 33701
Work phone: (813) 896-8626
FAX: -

Associate Research Scientist.

Projects:

1. Evaluated several created *Spartina alterniflora* marshes in Florida. Florida Sea Grant Publication Tech No. 60 (w/ R. "Robin" Lewis, III)
2. Coastal wetland evaluation work for FMRI.
3. Caryopsis production in natural and artificially created *Spartina alterniflora* Loisel marshes in Tampa Bay, FL.
4. Caryopsis production in *Spartina spartinae*.

Dr. William P. Davis

Environmental Protection Agency
Environmental Research Lab
1 Fabine Island Drive
Gulf Breeze, FL 32561-9201
Work phone: (904) 0934-9312
FAX: (904) 934-9201

Aquatic Research Ecologist. Mangrove ecology of mangroves, coral reefs, fish and invertebrates.

Projects:

1. Bear Bluffs in S.C. Transplanted to dikes from ponds.

Mr. David M. Drylie

Green Images
8100 Curry Ford Road
Orlando, FL 32822
Work phone: (407) 568-1333
FAX: (407) 568-2061

Sales representative.

Growers/suppliers of *Spartina alterniflora*, smooth cordgrass, *S. patens*, marsh hay cordgrass, and *S. bakeri*, sand cordgrass.

Ms. Shelley J. DuPuy

National Marine Fisheries Service
SEFC Panama City Lab, Habitat Conservation Division
3500 Delwood Beach Road
Panama City, FL 32408
Work phone: (904) 234-5061
FAX: (904) 235-3559 (but call 904-235-6591 to

make sure it's operable)

Fishery Biologist. Habitat protection.

Projects:

Protection of estuarine and marine habitats for fisheries populations.

Mr. Mark O. Friedemann

Florida Dept. of Environmental Regulation
2600 Blair Stone Road
Tallahassee, FL 32399-2400
Work phone: (904) 488-0130
FAX: (904) 488-6579

Compliance Coordinator.

Projects:

1. In charge of vegetative monitoring of mitigation efforts.

Ms. Joyce Gann

Gann's Tropical Greenery and Natives
22140 SW 152nd Avenue
Goulds, FL 33170
Work phone: (305) 248-5529
FAX: -

Owner.

Growers/suppliers of *Spartina patens*, marsh hay cordgrass, and *S. bakeri*.

Mr. George D. Gann-Matzen

Ecohorizons
22601 SW 152nd Avenue
Goulds, FL 33170
Work phone: (305) 989-5100
FAX: -

President.

Projects:

1. Works with restoration of wetlands using *Spartina patens*, *S. bakeri*, *S. spartinae*.
2. Has a database of vascular plants of Florida.
3. Primarily works with rebuilding plant communities for State, private, and non-profit groups; works toward restoration of damaged communities rather than the creation of new habitats.

Mr. Joseph L. Gilio

Gone Native Nursery
Wetlands Management, Inc.
P.O. Box 1122
Jensen Beach, FL 34958
Work phone: (407) 334-1643
FAX: (407) 335-1644

Grower/supplier of *Spartina alterniflora*, *S. patens*, and *S. bakeri*; wetlands consultant and designer.

Projects:

1. Mobil Oil (Sailfish Point), planted *Spartina alterniflora* along 2 miles of shoreline.
2. Planted seven salt-tolerant species, including *Spartina alterniflora*, at the Breakers (1.4 ac).
3. Planted *Spartina alterniflora* at Jensen Beach Club and Island Shoppes.
4. Also planted *S. bakeri* at many littoral lake sites.

Dr. R. Grant Gilmore

Harbor Branch Oceanographic Institute
5600 Old Dixie Highway
Ft. Pierce, FL 34946

Work phone: (407) 465-2400
ext. 203

FAX: (407) 465-2446

Administration and Research Scientist. Fish ecology.

Ms. Caroline Hanes

Florida Dept. of Environmental Regulation
SE District Office
1900 S. Congress Ave., Suite A
West Palm Beach, FL 33406

Work phone: (407) 433-2650

FAX: -

Environmental Specialist

Dr. Brandt F. Henningsen

SW Florida Water Management District
Surface Water Improvement and Management Program
7601 Highway 301 N
Tampa, FL 33637

Work phone: (813) 985-7481 ext. 2202

FAX: (813) 985-7481 ext. 2069

Environmental Scientist. Estuarine wetland restoration and preservation.

Projects:

Monitoring 15 estuarine marshes, mostly *S. alterniflora*, and several were man-made.

Mr. Robert D. Karafel

Metropolitan Dade County
Environmental Resources Management
Metro-Dade Center
111 N.W. first Str., Suite 1310
Miami, FL 33128

Work phone: ???-375-3305

FAX: -

Code enforcement officer - environmental

Dr. Edwin J. Kepner

National Marine Fisheries Service
SEFC Panama City Lab, Habitat Conservation Division
3500 Delwood Beach Rd.
Panama City, FL 32407

Work phone: (904) 234-5061

FAX: (904) 235-3559

Habitat Conservation Division Station Chief

Projects:

Coordinates habitat conservation efforts of NMFS in the NE Gulf of Mexico.

Dr. Thomas C. Kosier

Plants for Tomorrow, Inc.
16361 Norris Road
Loxahatchee, FL 33470

Work phone: (407) 790-1422

FAX: (407) 790-1916

Head of Research and Development; vice President.

Projects:

Growers/suppliers of *Spartina alterniflora*, smooth cordgrass; *S. bakeri*, sand cordgrass; and *S. patens*, marsh hay cordgrass. Most of material grown for in-house contracting projects. Has done some consulting work with Robin Lewis. Approximately 5% of consulting is *Spartina alterniflora*.

Dr. William L. Kruczynski

U.S. Environmental Protection Agency
Gulf Breeze Environmental Research Laboratory
Sabine Island

Gulf Breeze, FL 32561-3999

Work phone: (904) 934-9279

FAX: (904) 934-9201

Wetlands Expert.

Projects:

1. Planting of a spoil island using *Spartina alterniflora* at Apalachicola, FL, 1976.
2. Planting of a spoil island using *Spartina alterniflora* at Dickerson Bay, FL, 1975.

Mr. Mark Latch

Florida Dept. of Environmental Regulation,
Water Management
2600 Blair Stone Rd.
Tallahassee, FL 32399-2400

Work phone: (904) 488-0130

FAX: (904) 488-6579

Division Director. Aquatic ecology and wetland resource management.

Dr. Roy Robin Lewis, III

Lewis Environmental Services, Inc.
P.O. Box 20005
Tampa, FL 33622-0005

Work phone: (813) 889-9684

FAX: (813) 881-1586

Wetlands consultant; Ecological consulting, *Spartina* and mangrove marsh enhancement, restoration and creation.

Projects:

1. Developed methods for creation and restoration of *Spartina alterniflora* and mangrove marshes in Florida.
2. Designed and created *Spartina alterniflora* marshes as mitigation for impacts to estuarine marshes in Tampa Bay area for several companies.

Mr. Brightman Logan

Association of Florida Native Nurseries
P.O. Box 1045

San Antonio, FL 33576-1045

Work phone: (904) 588-3687

FAX: (904) 588-4552

Secretary of Association (1992).

Projects:

Produce the "Plant and Service Directory" annually; about \$2.50; lists sources of *Spartina spp.* and companies that grow and install it.

Mr. Andreas Mager, Jr.

National Marine Fisheries Service, SE Region
Habitat Conservation Division
9721 Executive Center Drive, North
St. Petersburg, FL 33702

Work phone: (813) 893-3503

FAX: (813) 893-3111

Assistant Regional Director, SE Region. Resource management and Fisheries.

Projects:

1. Records and tracks water development projects (including proposed wetland mitigation) in the coastal area of the NMFS SE region using R-base (database system) on an IBM compatible PC.
2. Manages various NMFS habitat mitigation programs for the NMFS SE region (e.g. Coastal Wetland Planning, Protection and Restoration Act).

Mr. Frank Melton

Conservation Consultants, Inc. / Native Plants
P.O. Box 248
321 61st St. E. / Palm View Rd.
Palmetto, FL 34220

Work phone: (813) 729-5015

FAX: (813) 729-8384

Horticulturist; growing wetland plants, and restoring wetland areas.

Projects:

1. Grower and supplier of *Spartina alterniflora*, *S. bakeri*, and *S. patens*, as well as other wetland plants such as mangroves, *Rhizophora* and *Avicennia*.
2. Created marsh with *Spartina alterniflora* and red mangrove plants at Barefoot Beach near Bonito Springs.
3. Created 5 acres of marsh at Perico Harbor near Sarasota Bay. Site located at West Manatee Ave. and Bradenton.

Mr. Robert Patten

Coastal Dunes, Inc.
1605 Main St. #800
Sarasota, FL 34236

Work phone: (813) 366-6538

FAX: (813) 954-6036

President and Founder. Salt marsh and coastal habitat restoration.

Projects:

Various restoration projects.

Ms. Ann Redmond

Florida Dept. of Environmental Regulation
2600 Blair Stone Road
Tallahassee, FL 32399-2400

Work phone: (904) 488-0130

FAX: (904) 488-6579

Mitigation Coordinator.

Projects:**Ms. Pamela Reeder**

Coastal Revegetation, Inc.
Shoreline Associates
5 Northeast 2nd Street
Delray Beach, FL 33444

Work phone: (407) 495-0198

FAX: (407) 278-7513

Growers/suppliers of *Spartina alterniflora*, *S. patens*, and *S. bakeri*.

Projects:

1. Munyon Island restoration project for the Palm Beach County Dept. of Environmental Resources Management.
2. Ash Landfill mitigation for Broward County Office of Integrated Waste Management.
3. Many other *Spartina* planting projects since 1981.

Mr. Michael E. Reily

The Wetlands Company, Inc.
7650 S. Tamiami Trail
Sarasota, FL 34231

Work phone: (813) 921-6609

FAX: (813) 921-7265

Grower/supplier of *Spartina alterniflora*, *S. patens*, and *S. bakeri*.

Mr. Rhodes Robinson

Environmental Services, INC.
8711 Perimeter Park Blvd., Suite 11
Jacksonville, FL 32216

Work phone: (800) 443-3158

FAX: (904) 645-9954

President. Specialties are Mitigation and Wetlands. Various, about 75 projects completed.

Mr. Douglas M. Scheidt

Harbor Branch Oceanographic Institution
5600 U.S. 1 North
Ft. Pierce, FL 34946

Work phone: (407) 465-2400 Ext. 668

FAX: (407) 465-2446

Fish Biologist. Fish ecology and marsh ecology.

Mr. Michael D. Schroer

Breedlove, Dennis & Associates, Inc.
4301 Metric Drive
Winter Park, FL 32792

Work phone: (407) 677-1882

FAX: (407) 657-7008

Associate Scientist II; environmental consulting.

Mr. Carl Terwilliger

Meadow Beauty Nursery
7210 Saddle Road
Lake Worth, FL 33463

Work phone: (407) 969-9250

FAX: -

Grower/supplier of *Spartina alterniflora* and *S. bakeri*;
environmental consulting.

Projects:

1. Growing plants for restoration projects.
2. Bank stabilization of lake and water retention ponds at The Mounts Botanical Garden in West Palm Beach, FL, using *Spartina alterniflora*.

Mr. Thomas W. Tito

P.O. Box 14116
St. Petersburg, FL 33733

Work phone: (813) 895-0446

FAX: -

Supplier of *Spartina patens*, marsh hay cordgrass.

Projects:

1. Installation of *Spartina* and black mangroves at Big Bayou, Tampa Bay, 1979-1985.
2. North Clearwater Beach at St. Joseph's Sound, restoration/creation project to start soon, 1993.
3. Involved with designing and installing wetland and upland native plants for residential and commercial landscaping projects.

Dr. David A. Tomasko

Sarasota National Estuary Program
1550n Thompson Parkway
Sarasota, FL 34236

Work phone: (813) 361-6133

FAX: (813) 361-6165

Senior Scientist.

Projects:

1. City Island, Fall '91.
2. New College, Spring '92.
3. Leffis Key, Fall '91.

Mr. David J. Wesley

U.S. Fish and Wildlife Service, Jacksonville Field Office
3100 University Blvd., #120
Jacksonville, FL 32216

Work phone: (904) 232-2580

FAX: (904) 232-2404

Fish and wildlife biologist

Review of permit requests to alter coastal wetlands.

Dr. Quinton A. White

Jacksonville University
Department of Biology and Marine Science
2800 University North
Jacksonville, FL 32211

Work phone: (904) 744-3950 ext. 6391

FAX: (904) 744-0101

Professor of Biology and Marine Science, Chairman of the department.

Mr. Robert Whitman

9720 Princess Palm Ave., Suite 106
Tampa, FL 33619

Work phone: (813) 626-5400

FAX: (813) 623-1034

Co-president. Estuarine habitat restoration and fisheries.

Mr. Mike Zaffke

Indian Trails Native Nursery
10629 Anderson Lane
Lake Worth, FL 33467

Work phone: (407) 641-9488

FAX: -

Grower/supplier of wetland plants.

Projects:

Grows and supplies wetland plants for restoration or any landscaping projects.

Dr. Nancy C. Zyski

Mitigation Services, Inc.
8711 Perimeter Park Blvd., Suite 11
Jacksonville, FL 32216

Work phone: (800) 443-3158

FAX: (904) 645-9954

President, Specialties-Wetlands Creation and Monitoring Wetlands

Projects:

1. Anastasia Sanitary District, St. Augustine, FL. Restoration and monitoring for a wastewater pipe installation project across several thousand feet of salt marsh associated with the Matanzas River. The area substrate was compacted during pipe installation, but *Spartina* is growing out into the area.
2. The Greens or Marsh Landing. A salt marsh creation and monitoring project, as mitigation for the development of an apartment complex. *Spartina alterniflora* is growing rapidly and filling in where expected.
3. Amelia Island Yacht Club. A *Spartina alterniflora* creation and monitoring project to mitigate for road widening impacts associated with upgrading access to a dredged material deposition site. Monitoring was required for a three year period. Again, *Spartina* is growing and filling in very well.

Georgia**Mr. Oscar Fultz**

Chatham County Mosquito Control Commission
1321 Eisenhower Drive
Savannah, GA 31406

Work phone: (912) 355-6320

FAX: -

Director. Mosquito control.
 Projects:
 Some projects affect salt marsh ecology and alteration.

Mr. Jim Lanier
 Aquascape Environmental Services, Inc.
 2550 Sandy Plains Road, Suite 320-291
 Marietta, GA 30066
 Work phone: (404) 578-0333
 FAX: -

President; wetlands consulting/ wetland restoration.
 Projects:
 Primary focus is in freshwater wetland restorations.

Mr. James F. Renner, P.G.
 Kemron Environmental Services
 2987 Clairmont Road, Suite 150
 Atlanta, GA 30329
 Work phone: (404) 636-0928
 FAX: (404) 636-7162
 Senior Geologist; consulting.

Louisiana

Mr. John E. Boatman
 U.S. Soil Conservation Service
 Plaquemines Soil and Water Conservation District Office
 104 Hebert Boulevard
 Belle Chase, LA 70037
 Work phone: (504) 394-7741
 FAX: -
 District Conservationist.
 Projects:
 Assess the region for possible revegetation rehabilitation sites, and coordinate plantings of *Spartina alterniflora* for the CRD/DNR (Steyer's group).

Mr. Gerald W. Bodin
 U.S. Fish and Wildlife Service
 825 Kaliste Saloom Road
 Brandywine Bldg. II, Suite 102
 Lafayette, LA 70508
 Work phone: (318) 264-6630
 FAX: (318) 264-6663
 Fish and Wildlife Biologist.
 Projects:
 Permitting and recommendations; most of these are implemented by individual refuge managers.

Mr. Allen Bolotte
 U.S. Soil Conservation Service
 New Orleans Field Office
 P.O. Box 531
 Boutte, LA 70039

Work phone: (504) 758-2162
 FAX: -

District Conservationist.
 Projects:
 1. Resource conservation planning in St. John, St. Charles, St. Bernard, Jefferson, and Orleans Parishes in Louisiana.
 2. Managing several shoreline planting projects in the field office area.

Mr. Wayne Bordelon
 U.S. Soil Conservation Service
 Thibodaux Field Office
 204 East Bayou Road
 Thibodaux, LA 70302
 Work phone: (504) 447-3871
 FAX: -

District Conservationist.
 Projects:
 1. 9-14 July 1992 plant 1600 1-gallon containers of *Spartina alterniflora*. 800 at Catahoula Bay and 800 at Temple Bay, Louisiana, east side of Bayou LaFourche. Use "Nutria-excluders" to prevent grazing of plants by *Nutria*.
 2. 1991- planted 8000 single-stem plants at Bay Laurf, Louisiana, on the east side of Bayou LaFourche. Much of this was destroyed by *Nutria* grazing.

Mr. Edgar S. Bordes
 New Orleans Mosquito Control Board
 6601 Lake Shore Drive
 New Orleans, LA 70126
 Work phone: (504) 241-2370
 FAX: (504) 244-4662

Director.
 Projects:
 Manipulation of dredged and levied areas in fresh water marsh for the most part, but some survey work in natural marsh.

Mr. Brad L. Broussard
 U.S. Soil Conservation Service
 New Iberia Field Office
 300 Iberia Street
 Suite 320 Courthouse Building
 New Iberia, LA 70560
 Work phone: (318) 369-6623
 FAX: -

District Conservationist.
 Projects:
 Identify areas for possible revegetation and rehabilitation, and coordinate plantings of *Spartina alterniflora* for the Coastal Restoration Division of the Department of Natural Resources.

Mr. Daniel L. Broussard

Coastal Plants
2320 W. Alcide
Abbeville, LA 70510

Work phone: (318) 893-2202
FAX: (318) 898-3047

Owner/Manager.

Projects:

1. Experimenting with planting protocols to increase viability of plants propagated from the wild.
2. Sells single stem and gallon containers of *Spartina* (both wild varieties and cultivars from the PMC in Galliano, LA); supplier to state and federal contracts.

Dr. Donald R. Cahoon

U.S. Fish and Wildlife Service
National Wetlands Research Center
700 Cajun Dome Blvd.
Lafayette, LA 70506

Work phone: (318) 266-8634
FAX: (318) 266-8513

Research Scientist.

Projects:

1. Marsh resiliency with respect to sea level rise; measurement of accretion/subsidence with respect to elevation and sea level of natural marsh nationwide; is interested in expansion of project to incorporate transplanted marsh.

Mr. Richard H. Day

U.S. Fish and Wildlife Service
National Wetland Research Center
700 Cajun Dome Blvd.
Lafayette, LA 70506

Work phone: (318) 266-8557
FAX: (318) 266-8513

Geographer/Marine Ecologist.

Projects:

1. Global Climate Change- remote sensing (satellite) of marshes compared to near (aerial) spectral analysis of marsh - coordinated with ground structure of canopy, etc. Measure with spectral radiometer which integrates over 2 nm wavelengths to create a spectral signature. Satellite's 7 wide bands are then compared to 350+ bands of near sensing (between 400-1100 nm).

Dr. Kittie Sue Derstine

U.S. Soil Conservation Service
Golden Meadow Plant Materials Center
P.O. Box 2202
Galliano, LA 70354

Work phone: (504) 475-5280
FAX: (504) 475-6545

Research Coordinator/Botanist.

Projects:

1. Primary work focuses on collection of assemblies and cultivation of the four species of *Spartina* (*S. alterniflora*, *S. patens*, *S. spartinae*, and *S. cynosuroides*) in the range from Texas to Mississippi. These assemblies are evaluated in field trials for a range of parameters: above and below ground biomass, salt tolerance, growth rates, vigor, etc. From these, cultivars are selected to create germ plasm for release. Germ plasm is available free for propagation for vegetative increase. One cultivar is currently available for distribution: *Spartina alterniflora* Vermilion.
2. Dr. Derstine is currently creating a list of guidelines for nursery preparation to enhance eventual success of transplanted propagules.

Ms. Donna J. Devlin

4561 John Koonce Rd.
Lake Charles, LA 70611

Work phone: (318) 475-5917
FAX: -

Estuarine ecologist.

Projects:

1. 1989-present. Studies of a 10-year old mangrove creation site in SW Florida: plant survival, growth, reproduction, and succession. Some faunal data.
2. 1993-present. Studies on survival, growth, and succession in a Tampa Bay, FL created salt marsh. Secondary aspect evaluates the use of citizen volunteers to "monitor" such sites.
3. 1990-1993. Role of the litter-grazing snail, *Melampus coffeus* L. in the decomposition of mangrove leaf litter, and coupling export of detrital energy to the estuary.

Mr. William P. "Judge" Edwards

Vermilion Corporation
P.O. Box 27
Abbeville, LA 70511

Work phone: (318) 893-0268
FAX: -

Assistant Manager of Vermilion Corp. private hunting lands.

Projects:

1. Manages lands for hunting and fishing by Vermilion Corp.
2. Uses transplantation of *Spartina* along private canals which have accumulated silt to stabilize soil.

Mr. Gary Fine

U.S. Soil Conservation Service
Golden Meadow Plant Materials Center
P.O. Box 2202
Galliano, LA 70354

Work phone: (504) 475-5280
FAX: (504) 475-6545

Manager.

Projects: (more)

1. Manage the facility for the cultivation of the four species of *Spartina* (*S. alterniflora*, *S. patens*, *S. spartinae*, and *S. cynosuroides*) in the range from Texas to Mississippi. Germ plasm is available free for propagation for vegetative increase. One cultivar is currently available for distribution: *Spartina alterniflora* Vermilion.

Mr. Sherwood Gagliano

Coastal Environments, Inc.

1260 Main St.

Baton Rouge, LA 70802

Work phone: (504) 383-7455

FAX: (504) 383-7925

President.

Projects:

Primary focus is on monitoring water use (aquatic resource management and multi-use programs) effects on riverine and coastal areas through aerial and remote sensing

Mr. Thomas G. Hargis

University of Southwestern Louisiana

Department of Biology

P.O. Box 42451 USL

300 E St. Mary Street

Lafayette, LA 70504-2451

Work phone: (318) 231-5248

FAX: (318) 231-5834

Graduate Researcher.

Projects:

Master's research at USL: Site characterization of marsh ecology at Lafitte (Barataria Parish) and Houma (Terrebonne Parish), Louisiana, measuring sulfide, nutrients, bulk density, and salinity.

Mr. Richard D. Hartman

National Marine Fisheries Service

Habitat Conservation Division

c/o Louisiana State University

Baton Rouge, LA 70803-7535

Work phone: (504) 389-0508

FAX: (504) 389-0506

Fisheries biologist, estuarine ecologist.

Projects:

1. Wetland creation/enhancement in fresh to saline tidal marshes. Survey, design, evaluate, and monitor restoration activities in Louisiana.
2. Protect fisheries habitats.

Dr. Suzanne "Sue" R. Hawes

U.S. Army Corps of Engineers, New Orleans District

Planning Division

P.O. Box 60267

New Orleans, LA 70160-0267

Work phone: (504) 862-2518

FAX: (504) 862-2572

Project Manager for the Environment.

Projects:

Helps plan marsh creation with dredged material projects for the New Orleans District Corps of Engineers.

Dr. William H. Herke

Louisiana State University

School of Forestry, Wildlife and Fisheries Sciences

Cooperative Fish and Wildlife Research Unit

Baton Rouge, LA 70803-6202

Work phone: (504) 388-4180

FAX: (504) 388-4227

Assistant leader Louisiana Cooperative Fish and Wildlife Research Unit.

Projects:

1. Monitoring fisheries interactions with natural and managed marshes-- typically not those that are transplanted.
2. Studies the life history of marine transients while using the marsh nursery, and the effects of water-control structures and levees on their use of the nursery.

Dr. H. Dickson Hoes

University of Southwestern Louisiana

Department of Biology

P.O. Box 42451 USL

300 E St. Mary Street

Lafayette, LA 70504-2451

Work phone: (318) 231-6494

FAX: (318) 231-5834

Professor; ichthyology and wetlands biology.

Projects:

1. Primary work is on the ichthyology of natural marshes, weir structures and impoundments, 1990. Interim report: fisheries studies on the state wildlife refuge.
2. Comparison of effects of water control structures on fisheries in the Rockefeller Wildlife Refuge.
3. Cameron-Creole watershed management fisheries study.

Mr. John Hook

University of Southwestern Louisiana

Department of Biology

P.O. Box 42451 USL

Lafayette, LA 70504-2451

Work phone: (318) 383-1972

FAX: (318) 231-5834

Graduate researcher.

Mr. Craig A. Johnson

U.S. Fish and Wildlife Service

National Wetlands Research Center

700 Cajun Dome Blvd.

Lafayette, LA 70506

Work phone: (318) 266-8695

FAX: (318) 266-8513

Geographer

Mrs. Peggy M. Jones

National Marine Fisheries Service
SEFC Habitat Conservation Division
c/o Louisiana State University
Baton Rouge, LA 70803-7535
Work phone: (504) 389-0508
FAX: (504) 389-0506

Fisheries biologist, estuarine ecologist.

Projects:

1. Planning and evaluation subcommittee member representing DOC for Coastal Planning, Protection, and Restoration Act, 1990-1993.
2. Review and evaluate proposed alterations to coastal wetlands to protect habitat supportive of marine fishery resources, 1980-present.

Dr. Mark A. Konikoff

Univ. of Southwestern Louisiana
Department of Biology
P.O. Box 42451 USL
300 E St. Mary Street
Lafayette, LA 70504-2451
Work phone: (318) 231-6754
FAX: (318) 231-5834

Professor.

Projects:

1. Six year project on beneficial uses of dredge material deposition areas. Containment Area Aquaculture Program (CAAP) at WES.

Mr. Ronald J. Marcantel

U.S. Soil Conservation Service
Lake Charles Field Office
1400 Highway 14
Lake Charles, LA 70601
Work phone: (318) 436-1483
FAX: -

District Conservationist.

Projects:

- Coordinates regional plantings for CRD through SCS in Crowley for the Lake Charles Area.

Mr. Michael Materne

U.S. Soil Conservation Service
Plant Materials Center
P.O. Box 16030, University Station
Baton Rouge, LA 70893
Work phone: (504) 389-0335
FAX: -

Plant Materials Specialist.

Projects:

1. Plant materials database for nurseries of wetland plants in Louisiana and the U.S.

Dr. Linda Glenboski Mathies

U.S. Army Corps of Engineers, New Orleans District,
CELMN-OD-OR

P.O. Box 60267
New Orleans, LA 70115-0267
Work phone: (504) 862-2318
FAX: (504) 862-2317

Environmental Resources Specialist

Projects:

1. Mississippi River; Gulf Outlet, LA, marsh creation using dredged material.
2. Mississippi River; Baton Rouge to Gulf of Mexico, LA, marsh creation using dredged material (1974).
3. Houma Navigation Canal, LA, marsh restoration using dredged material (1990).
4. Calcasieu River and Pass, LA, marsh restoration using dredged material (1993).

Mr. Larry McNease

Rockefeller Wildlife Refuge
Louisiana Dept. of Wildlife and Fisheries
Rt. 1, Box 20-B
Grande Chenier, LA 70643
Work phone: (318) 538-2276
FAX: -

Biologist.

Projects:

- Study of the vegetation associated with structural coastal marsh management.

Mr. Donald E. Menard

U.S. Soil Conservation Service
Abbeville Field Office
P.O. Box 68
Abbeville, LA 70511-0068
Work phone: (318) 893-5664
FAX: -

District Conservationist.

Projects:

- Coordinates plantings for Louisiana's Department of Natural Resources, Coastal Restoration Division, in the Abbeville area.

Dr. Irving A. Mendelssohn

Louisiana State University
Wetland Biogeochemistry Institute, and Dept. of
Oceanography and Coastal Sciences
Baton Rouge, LA 70803
Work phone: (504) 388-6425
FAX: (504) 388-6423

University researcher and professor.

Projects:

1. Research involves the understanding of causes underlying wetland loss in Louisiana. Primarily subsidence loss with factors influencing heterogeneous vigor deterioration using natural gradients and monitoring and experimental manipulations of factors (elevation, salinities, etc.)
2. Past studies on transplanting.

Mr. Jeff Murphy

Blacklake Marsh, Inc.
1200 Paris Street
Lake Charles, LA 70605

Work phone: (318) 762-4242

FAX: Call ahead to FAX.

Owner/president; fresh, brackish and salt marsh restorations, grows and installs.

Projects:

1. Grows and supplies *Spartina alterniflora*, smooth cordgrass, *S. patens*, marsh hay cordgrass, *Scirpus californicus* and many more species. Can provide single stems or multi-stem trade-gallons in large quantities.
2. Planted terraces in Sabine National Wildlife Refuge with *Spartina alterniflora* (Vermillion strain). Trade-gallons were used on 5-ft centers on one side; later, single stems were planted on the opposite side for comparison. Great success, and the sides are indistinguishable.
3. Wave barrier wooden fencing (2300 ft) was used at the Brannan Ditch project to break waves and flood water currents impacting the south side of the ICWW. Single stems of *Spartina alterniflora* were planted shoreward of the fence, and rapidly became established. They acted as a sediment trap in combination with the fence.

Mr. Michael D. Nichols

U.S. Soil Conservation Service
Water Resources Planning Staff
3737 Government Street
Alexandria, LA 71302

Work phone: (318) 473-7688

FAX: (318) 473-7771

Wildlife biologist.

Projects:

1. Planning of coastal restoration and protection projects connected with the Coastal Wetlands Planning, Protection and Restoration Act.
2. Watershed and river basin planning in the coastal zone of Louisiana.
3. Assisting district offices of the USDA Soil Conservation Service with revegetation efforts throughout coastal Louisiana.

Mr. Will Nidecker

Sabine National Wildlife Refuge
Hwy. 27 South
300 Main Street
Hackberry, LA 70645

Work phone: (318) 762-3816

FAX: (318) 762-3780

Refuge Manager.

Projects:

Spartina alterniflora transplanting and establishment within the refuge. Project A: Mitigation wave break levees in Units 6 & 7. Project B: terracing near Old Hog Bayou.

Mr. Kenneth E. Owen

University of New Orleans
Earl K. Long Library
New Orleans, LA 70146

Work phone: (504) 286-7275

FAX: (504) 286-7277

Librarian.

Projects:

Head of the Louisiana Collection, an extensive set of references about Louisiana, including many about wetlands.

Mr. W. Guthrie Perry

Louisiana Department of Wildlife and Fisheries
Rockefeller Wildlife Refuge
Rt. 1, Box 20-B
Grande Chenier, LA 70643

Work phone: (318) 538-2276

FAX: -

Biologist.

Projects:

Study of the aquatic organisms associated with structural coastal marsh management.

Dr. S. R. Pezeshki

Louisiana State University
Center for Coastal Energy and Environmental Resources
Wetland Biogeochemistry Institute
Baton Rouge, LA 70803-7511

Work phone: (504) 388-6430

FAX: (504) 388-6423

Associate Professor of Wetland Resources; plant ecophysiological. more...

Projects:

1. Physiological responses of wetland plants to flood and salt stresses: soil - plant interactions. Work with R.D. De Laune and W.H. Patrick, 1993.
2. Wetland soil development. Work with R.D. De Laune and W.H. Patrick, 1988-1993.
3. Field studies of beach cones as coastal erosion reversal devices for areas with significant oil and gas activities. Work with R.D. De Laune, 1992-1994.

Dr. Edward Proffitt

Louisiana Environmental Research Center
McNeese State University
P.O. Box 90655
Lake Charles, LA 70609-0655

Work phone: (318) 475-5917

FAX: (318) 475-5122

Director of the Research Center; Associate Professor; mangrove and salt marsh ecology

Projects:

1. 1988-present. Studies of a 10-year old mangrove creation site in SW Florida: plant survival, growth, reproduction, and succession. Some faunal data.(more)

2. 1993-present. Studies on survival, growth, and succession in a Tampa Bay, FL created salt marsh. Secondary aspect evaluates the use of citizen volunteers to "monitor" such sites.
3. 1990-1993. Role of the litter-grazing snail, *Melampus coffeus* L. in the decomposition of mangrove leaf litter, and coupling export of detrital energy to the estuary.

Ms. Patricia S. Rafferty
University of Southwestern Louisiana
Department of Biology
P.O. Box 42451 USL
300 E St. Mary Street
Lafayette, LA 70504-2451

Work phone: (318) 231-5248
FAX: (318) 231-5834

Graduate Researcher.

Projects:

1. Doctoral research focuses on interaction between rhizosphere and sediment as it affects nitrogen cycling (denitrification rates) at Otter Bayou (with USFWS) and Fourleague Bay (Sea Grant project), 1993.

Dr. Elijah W. Ramsey
U.S. Fish and Wildlife Service
National Wetland Research Center
700 Cajun Dome Blvd.
Lafayette, LA 70506

Work phone: (318) 266-8575
FAX: (318) 266-8513

Oceanographer.

Mr. John E. Reddoch
U.S. Army Corps of Engineers, New Orleans District
Operations & Readiness Div., Regulatory Functions Br.
P.O. Box 60267
New Orleans, LA 70160-0267

Work phone: (504) 862-2277
FAX: (504) 862-2289

Environmental Resource Specialist.

Mr. Bryan J. Reed
The Wetlands Foundation
3841 Veterans Blvd., Suite 201
Metairie, LA 70002

Work phone: (504) 888-9944
FAX: (504) 888-9976

Executive Director.

Dr. Denise Janet Reed
Louisiana Universities Marine Consortium
Marine Research and Education Center
Chauvin, LA 70344

Work phone: (504) 851-2800
FAX: (504) 851-2874

Associate Professor; Salt marshes, tidal flows and sediment dynamics.

Projects:

1. Tidal flows in salt marshes, and their roles in sediment transport and accretion in the marshes.
2. Understanding interactions between biological communities and depositional processes in coastal marshes.

Mr. Rickey N. Reubsamen
National Marine Fisheries Service
Habitat Conservation Division
c/o Louisiana State University
Baton Rouge, LA 70803-7535

Work phone: (504) 389-0508
FAX: (504) 389-0506

Branch Chief; Fisheries biology and estuarine ecology.

Projects:

1. Wetland creation/enhancement in fresh to saline tidal marshes. Five projects designed and funded for construction in coastal Louisiana under the Coastal Wetland Planning, Protection, and Restoration Act. 1990-1993.
2. Seagrass meadow creation (*Halodule*) in Laguna Madre, Texas, 1985.
3. Wetland creation (*S. alterniflora*) at Lake Como, Galveston, Texas, 1984.
4. Estuarine brackish marsh enhancement for the Nueces River delta, Texas, 1983.

Mr. Mark Schleifstein
The Times-Picayune
3800 Howard Ave.

New Orleans, LA 70140-1097

Work phone: (504) 826-3327
FAX: (504) 826-3007

Reporter.

Reports on environmental issues and projects such as marsh loss and marsh restoration in Louisiana and the north-central coast of the Gulf of Mexico.

Mr. Chris Simon
Louisiana Department of Natural Resources
Coastal Restoration Division
200 Lafayette Street

Baton Rouge, LA 70804-9396

Work phone: (504) 342-7308
FAX: (504) 342-9417

Geoscience Specialist.

Projects:

1. Sabine terraces for *Spartina alterniflora* marsh creation and wave reduction.
2. Falgout canal/levee grass planting.

Ms. Ruth Spell
U.S. Fish and Wildlife Service
National Wetland Research Center
700 Cajun Dome Blvd.
Lafayette, LA 70506

Work phone: (318) 266-8571
FAX: (318) 266-8513

Geographer.

Mr. Gregory Dean Steyer

Louisiana Department of Natural Resources
Coastal Restoration Division
200 Lafayette St., P.O. Box 94396
Baton Rouge, LA 70804-9417

Work phone: (504) 342-9435
FAX: (504) 342-9417

Biological Program Manager.

Projects:

Directs the CRD, and initiates many coastal projects to protect and restore wetlands--both fresh and salt water. Many projects are listed in "Recommended three year coastal wetland vegetation program".

Ms. Faye A. Talbot

U.S. Soil Conservation Service
Crowley Area Office
P.O. Drawer 387
Crowley, LA 70527

Work phone: (318) 783-1257
FAX: (318) 783-5608

Resource Conservationist.

Projects:

1. Collects reports from state soil and water conservation districts, and compiles information for the Coastal Restoration Division of the Dept. Natural Resources as a yearly report.
2. Functions as a liaison between CRD and SCS on vegetation projects.
3. Serves as water resources staff leader at the Area III 3 level (coastal marshes) for the Ponchartrain, Breton Sound, Barataria, Mississippi River, Terrebonne and Tech/Vermilion Basin. Responsible for development of plans and projects for the SCS in these basins.

Mr. Michael D. Tullos

U.S. Soil Conservation Service
Houma Field Office
Box 1266
Houma, LA 10361

Work phone: (504) 872-5609
FAX: -

District Conservationist.

Projects:

1. Provide technical assistance for Terrebonne Parish in identifying potential rehabilitation projects.

Dr. Robert R. Twilley

University of Southwestern Louisiana
Department of Biology
P.O. Box 42451 USL
300 E St. Mary Street
Lafayette, LA 70504-2451

Work phone: (318) 231-6146
FAX: (318) 231-5834

Professor.

Projects:

1992. An ecological analysis of marsh management in coastal Louisiana: Processes related to the fertility of marsh ecosystems.

Maine

Mr. Stewart I. Fefer

U.S. Fish and Wildlife Service
Gulf of Maine Coastal & Estuary Project
312 Canco Road
Portland, ME 04103

Work phone: (207) 828-1080
FAX: (207) 828-4001

Restoration of tidal flow to salt marshes in northern New England.

Projects:

1. Awcomin Marsh at Rye, NH, 1993. 20 acres of *Spartina* marsh were restored by excavation of channels for inundation.
2. Marsh at Stratham, NH, 1993-94. 12 acres of marsh were restored through placement of large culverts to enhance tidal flow.
3. Lewis Lake at Winthrop, MA, 1993-94. 8 acres enhanced by restored tidal flows.
4. Hatches Harbor site at Provincetown, MA. 200 acres of salt marsh restored through placement of culverts to allow tidal inundations.

Maryland

Ms. Candy C. Bartoldus

Environmental Concern, Inc.
P.O. Box P, 210 West Chew Ave
St. Michaels, MD 21663

Work phone: (410) 745-9620
FAX: (410) 745-3517

Senior Associate. Wetland evaluation procedures.

Projects:

Senior author of Evaluation for Planned Wetlands.

Mr. Robert Blama

U.S. Army Corps of Engineers, Baltimore District
Operations Division, Navigation Branch
P.O. Box 1715
Baltimore, MD 21203 (more)

Work phone: (410) 962-3663
FAX: (410) 962-6039

Ecologist/Project Manager. Habitat creation and beneficial use of dredged material. (more)

Projects:

1. Barren Island, MD. 1987-88. About 10 acres of *Spartina alterniflora* and *S. patens* marsh were created.
2. Eastern Neck Wildlife Refuge. 1993. 5 acres of *Spartina alterniflora* and *S. patens* marsh were created.

Dr. Donald F. Boesch

University of Maryland, Horn Point Lab
Center for Estuarine and Environmental Study
P.O. Box 775
Cambridge, MD 21613

Work phone: (301) 228-9250

FAX: -

Director; biology-ecology, wetlands, community ecosystems, resource management

Projects:

Can direct you to the source.

Ms. Dawn Renee Briggs

Environmental Concern, Inc.
P.O. Box P, 210 West Chew Ave
St. Michaels, MD 21663

Work phone: (410) 745-9620

FAX: (410) 745-3517

Associate Wetland Designer. Landscape architecture and ecological design.

Projects:

Research and design of various wetland types, including shoreline restorations with *Spartina alterniflora* and *S. patens*.

Mr. Carlo R. Brunori

Maryland Department of Natural Resources
Forest, Park and Wildlife Service
Tawes State Office Building, E-1
580 Taylor Avenue
Annapolis, MD 21401

Work phone: (410) 974-3195

FAX: Chief, Technical Services; Water quality assessment, and wetland planning and design for preservation of wildlife habitat.

Projects:

1. Involved with water quality assessments and management in South, Gunpowder, Lower Susquehanna and Bush rivers in Maryland.
2. Develops water quality plans for river basins, including wetland creation or modification for wildlife enhancement.
3. Interagency coordination for water quality issues.

Dr. Leon M. Cammen

NOAA/National Sea Grant College Program
Marine Ecology
R/OR1, 133 East-West Highway
Silver Spring, MD 20910

Work phone: (301) 427-2435

FAX: (301) 427-2799

Associate Program Director for Environmental Studies.

Projects:

Program manager for Coastal Ocean Program's Estuarine Habitat Program.

Ms. Mary Ellen Dore

Maryland Department of Natural Resources
580 Taylor Ave. B-3
Annapolis, MD 21401

Work phone: (410) 974-2784

FAX: (410) 974-2833

Manager, Chesapeake Natural Estuarine Research Reserve. Resource management, environmental policy, and ecology.

Mr. H. Glenn Earhart

U.S. Army Corps of Engineers, Baltimore District
P.O. Box 1715
Baltimore, MD 21203

Work phone: (302) 962-3369

FAX: (301) 962-3966

Environmental Program Manager/Biologist. Habitat development and hazardous/toxic waste management.

Projects:

N/A

Dr. R. Michael Erwin

U.S. Fish and Wildlife Service
Patuxent Wildlife Research Center
Laurel, MD 20708

Work phone: 301-498-0380

FAX: 301-498-0438

Projects:

1. Chesapeake Bay, Delaware Bay, Eastern Shore of VA, Southern New Jersey, and Outer Banks, NC.

Mr. Thomas Filip

U.S. Army Corps of Engineers
Baltimore District - Regulatory Branch
P.O. Box 1715

Baltimore, MD 21203-1715

Work phone: (410) 962-3670

FAX: (410) 962-6024

Asst. Chief of Regulatory; wetlands chemist

Projects:

1. 119th Street marsh creation in Ocean City, MD.
2. Isle of Wright Bay *Spartina alterniflora* marsh establishment; very successful.
3. Other more recent plantings in the area.

Dr. Edgar W. Garbisch

Environmental Concern Inc.
P.O. Box P

St. Michaels, MD 21663 (more)

Work phone: (410) 745-9620

FAX: (410) 745-3517

President; environmental consulting; marsh restoration and creation. (more)

Projects:

1. *Spartina alterniflora* and *S. patens* have been used in 203 shore restoration and marsh construction for bank erosion control projects during the period of 1972-1992.
2. *Spartina alterniflora* and *S. patens* have been used in 13 beneficial uses of dredged material projects, 15 regulatory restoration projects, 2 voluntary restoration projects, 63 wetland mitigation projects, and 10 experimental wetland construction projects from 1972-1992.
3. Designed and created or restored 43 freshwater marshes of various types.

Mr. Tim Goodger

National Marine Fisheries Service
Habitat and Protected Resources Div., NE Region
Oxford Field Office
904 South Morris Street
Oxford, MD 21654

Work phone: (410) 226-5771

FAX: (410) 226-5417

Assistant coordinator; resource management and estuarine ecology.

Projects:

Habitat protection; coastal marshes for fisheries and fisheries related populations.

Dr. Michael S. Kearney

University of Maryland
Dept. of Geography
College Park, MD 20742

Work phone: (301) 405-4057

FAX: (301) 314-9299

Associate Professor.

Projects:

1. Chesapeake Bay, Delaware Bay, Mississippi Delta, VA Barrier Islands, Assateague Island, Fire Island, NY

Mr. Christopher P. Kilgore, ASLA

Environmental Concern, Inc.
P.O. Box P, 210 West Chew Ave
St. Michaels, MD 21663

Work phone: (410) 745-9620

FAX: (410) 745-3517

Senior Associate; Wetland evaluation, design and delineation.

Projects:

1. Assisted in the design and construction of 15 projects using *Spartina alterniflora* for shoreline protection primarily in the Chesapeake Bay area.
2. Assisted with the design and construction of a tidal estuarine marsh using *Spartina alterniflora*.
3. Assisted with the design and construction of four freshwater marsh projects.

Dr. Mark L. Kraus

Environmental Concern, Inc.
P.O. Box P, 210 West Chew Ave
St. Michaels, MD 21663

Work phone: (410) 745-9620

FAX: (410) 745-3517

Senior Associate, Director of Education. Wetland evaluation, design and delineation.

Projects:

1. Designed and constructed 11 tidal estuarine marshes where *Spartina alterniflora* was the major species planted. Creation and restoration projects were primarily in the Chesapeake Bay area.
2. Assisted with the design and construction of 11 freshwater marsh projects.

Mr. Donald Robert MacLean

University of Maryland, College Park Campus, Natural Resources Management and Water Resources Engineering.
9023 Queen Maria Court
Columbia, MD 21045

Work phone: (410) 997-3513

FAX: -

Consulting Ecologist and Teaching Assistant. Wetland mitigation, restoration, botany and delineation.

Projects:

1. C&F Realty Marsh Restoration Project, May '91.
2. Amerada Hess Wetland Mitigation Project, Jan '91.
3. Fort Meade Wetland Mitigation Project, Aug '90.
4. Charles County Wetland Mitigation Project, Aug '90.

Ms. Tamara L. McCandless

U.S. Fish and Wildlife Service
Chesapeake Bay Program,
Living Resources Branch
180 Admiral Cochrane Dr., Suite 535
Annapolis, MD 21401

Work phone: (410) 224-2732

FAX: (410) 224-2781

Habitat restoration and protection specialist.

Projects:

1. Develop wetland/riparian and stream restoration program.
2. 1993. Marsh protection and restoration project using brush-fencing technique at Blackwater National Wildlife Refuge.

Dr. Robert W. Paul

St. Mary's College of Maryland
Division of Natural Science and Math
St. Mary's City, MD 20686

Work phone: (301) 862-0362

FAX: (301) 862-0999

Professor; ecology of salt marshes.

Projects:

1. Ecological base line studies on Chesapeake Bay saltmarshes, gastropods. (more)
2. Consulting for the Navy on environmental contamination issues; hazardous waste.

Mr. Marc Seelinger

Environmental Concern, Inc.
P.O. Box P, 210 West Chew Ave
St. Michaels, MD 21663

Work phone: (410) 745-9620
FAX: (410) 745-3517

Senior Associate; Wetland design and construction.
Projects:

1. Assisted in the design and construction of a tidal estuarine marsh where *Spartina alterniflora* was the major species planted.
2. Assisted with the design and construction of 5 freshwater marsh projects.

Dr. J. Court Stevenson

University of Maryland, Horn Point Laboratory
Center for Estuarine and Environmental Study
Cambridge, MD 21613-0775

Work phone: (301) 228-8200
FAX: (301) 476-5490

Professor, researcher; wetlands.

Mr. Charles Utermohle

Whitman Requardt
2315 St. Paul Street
Baltimore, MD 21218

Work phone: (410) 235-3450 Ext.206
FAX: (410) 243-5716

Massachusetts**Mr. Richard Albano**

Sanford Ecological Services
30 Turnpike Road
Southborough, MA 01772

Work phone: (508) 460-9900
FAX: (508) 481-2009

Wetlands ecologist.

Projects:

Consulting for wetlands mitigation.

Mr. Thomas E. Bigford

National Marine Fisheries Service, NE Region,
Habitat and Protected Resources Div.

One Blackburn Drive
Gloucester, MA 01930-2298

Work phone: (508) 281-9300
FAX: (508) 281-9301

Division Chief.

Habitat protection for fisheries populations.

Dr. Burton Bryan

The BSC Group
425 Summer Street
Boston, MA 02210

Work phone: (617) 345-4025
FAX: (617) 345-8008

Senior Biologist; wetlands and water quality impacts
Projects:

Central Artery/Tunnel project in Boston harbor, planning for marsh restoration to compensate for project impacts.

Dr. Lars H. Carlson

Sverdrup Corporation
2 Center Plaza

Boston, MA 02108-1906

Work phone: (617) 742-8060
FAX: (617) 742-8830

Environmental Scientist; consulting on wetlands and environmental permitting.

Projects:

1. Restoration design of a *Spartina* marsh for mitigation of impacts resulting from construction of a new MBTA railroad bridge between Boston and Quincy, MA.

Dr. Jerome B. Carr

Carr Research Laboratory, Inc.
17 Waban Street
Wellesley, MA 02181

Work phone: (508) 651-7027
FAX: (508) 651-7028

Environmental research and consulting; hydrology and wetland ecology.

Projects:

Freshwater wetlands analyzed, and restored.

Ms. Cheryl C. Deshaies

Lelito Environmental Consultants
100 Corporate Place, Suite 103
Peabody, MA 01960

Work phone: (508) 535-7861
FAX: (508) 535-0073

Restoration biologist; wildlife management, freshwater and coastal wetland restorations.

Projects:

1. *Phragmites* eradication and *Spartina* restoration at Post Island Marsh in Quincy, Massachusetts.
2. Restoration of a filled *Spartina* marsh on Sea Pit Road, Falmouth, MA.
3. Sand dune restoration at Sippewissett Hills in Falmouth, MA: replacement of sand by grain size and color, and revegetation with dune grass.
4. Analysis of the direct and indirect effects of a rod and gun club on the plants and the wildlife in a salt marsh in Martha's Vineyard.

Mr. Daniel L. Falcone
OHM Corporation
88-C Elm Street
Hopkinton, MA 01748
Work phone: (508) 435-9561
FAX: -

Dredging.
Projects:

Mr. George Gavutis
U.S. Fish and Wildlife Service - Northeast, Region 5
Refuges and Wildlife Program
300 Westgate Center Drive
Hadley, MA 01034-9589
Work phone: (413) 253-8574
FAX: -

Ms. Wendi Goldsmith
Bestmann Green Systems, Inc.
7 Mall Street
Salem, MA 01970
Work phone: (508) 741-1166
FAX: (508) 741-3780

Bioengineer; erosion and water quality control through restoration of vegetation.

Projects:

1. Nursery facility supplies plants grown in biodegradable coconut fiber matrix for erosion protection. Various fresh, brackish, and salt species are available as single- and multi-stemmed plugs as well as sods which tolerate considerable wave impact and other stress.
2. Restoration of a *Spartina alterniflora* marsh that was impacted by barge wakes at Fresh Kills landfill, New York.
3. Restoration and enhancement of high and low marshes on tidal systems and bay frontage on private lots, public lands, and demonstration projects from New York to Maine.

Dr. William B. Kerfoot
KV Associates, Inc.
281 Main Street
Falmouth, MA 02540
Work phone: (508) 540-0561
FAX: (508) 457-9053

President; Biology, hydrology and geohydrology.

Projects:

1. Reverse layering, an alternative approach to dredging for Lake Ana coastal restoration, 1988-1991. Demonstration project.
2. Red Lily Pond diagnostic and feasibility study.
3. Subsidence dredging at Green Pond, 1991-present.

Mr. Paul R. Lelito
Lelito Environmental Consultants

P.O. Box 1525
North Falmouth, MA 02556
Work phone: (508) 564-5606
FAX: (508) 563-7669

President; biologist, consulting, wetlands restoration.

Projects:

1. *Phragmites* eradication and *Spartina* restoration at Post Island Marsh in Quincy, Massachusetts.
2. Restoration of a filled *Spartina* marsh on Sea Pit Road, Falmouth, MA.
3. Sand dune restoration at Sippewissett Hills in Falmouth, MA: replacement of sand by grain size and color, and revegetation with dune grass.
4. Analysis of the direct and indirect effects of a rod and gun club on the plants and the wildlife in a salt marsh in Martha's Vineyard.

Mr. Lee D. Lyman
Lycott Environmental Research, Inc.
600 Charlton Street
Southbridge, MA 01550
Work phone: (508) 765-0101, 1-800-462-8211
FAX: (508) 765-1352

President; environmental assessments and restorations.

Projects:

1. Musquashicut Lake; Scituate, MA
2. Environmental analysis of several coastal ponds of Rhode Island.

Mr. Christopher Mantzaris
National Marine Fisheries Service, NE Region
Habitat and Protected Resources Div.
One Blackburn Drive
Gloucester, MA 01930-2298
Work phone: (508) 281-9346
FAX: (508) 281-9301

Habitat Program coordinator for the NE; Wetlands ecology and fisheries biology.

Projects:

1. Tracks various coastal habitat restoration/creation projects in the NE.
2. Evaluates permits to protect critical marine and estuarine habitat for fisheries organisms.
3. Placement of shell in Long Island Sound to restore oyster beds.
4. Restoration of 13+ acres of combined habitats (mudflats with tidal creeks, *Spartina alterniflora* marsh, and salt pans) in Revere, MA.

Mr. Michael J. Marcus
New England Environmental, Inc.
800 Main Street
Amherst, MA 01002
Work phone: (413) 256-0202
FAX: (413) 256-1092

Senior Biologist; consulting.

Mr. David Martin

Wm. M. Warwick and Associates Inc.
P.O. Box 801, 213 Old Main Road
North Falmouth, MA 02556

Work phone: (508) 563-7777

FAX: -

Civil Engineer; coastal and wetland issues.

Projects:

1. Inspection of docks and other coastal structures for meeting engineering codes for permits.
2. Bank stabilization projects. Rip-rap was used to stabilize the bank, and the shore-side was filled to desired height and planted with *Spartina* for biological stabilization of the shore. This was done for three adjacent properties in West Falmouth, Massachusetts. The adjacent marsh, used for access, was also planted to restore it where it had been damaged.

Mr. Dan McHugh

Great Meadow Farms
Sawyer Island
Rowley, MA 01969

Work phone: -

FAX: -

Projects:

1. Creation of a 20-acre *Spartina alterniflora* salt marsh from a dredge material deposition area - the Awconim Marsh Restoration project in New Hampshire.

Dr. Norton H. Nickerson

Tufts University
Department of Biology
Medford, MA 02155

Work phone: (617) 627-3544

FAX: (617)627-3805

Professor; research and teaching about inland and coastal wetlands.

Projects:

1. Creation of salt marsh along the Neponset River near the new railroad bridge; between Boston and Quincy, MA.
2. Study of sulfide relationships to habitat in *Spartina alterniflora* in Cape Cod marshes.
3. Study of habitat differences in sulfide concentrations on habitat and successional patterns of the red mangrove (*Rhizophora mangle*) and the black mangrove (*Avicennia germinans*) in the Florida Everglades, the Bahamas, and Belize.

Dr. Douglas S. Sparrow

Louis Berger & Associates, Inc.
303 Bear Hill Road
Waltham, MA 02154

Work phone: (617) 890-8999

FAX: (617) 890-7959

Manager; wetlands specialist.

Projects:

1. Tidal marsh enhancement in Stonington, CT, 1992-present. Feasibility of expanding *Spartina* growth is a sub component in a tidal cove restoration project.

Ms. Susan Taylor

OHM Corporation
88-C Elm Street
Hopkinton, MA 01748

Work phone: (508) 435-9561

FAX: -

Dredging.

Dr. John M. Teal

Woods Hole Oceanographic Institute
Biology
Water Street
Woods Hole, MA 02543

Work phone: (508) 457-2090

FAX: (508) 457-2169

Researcher; biologist and ecologist.

Projects:

1. Long-term ecological studies of the Great Sippewissett Salt Marsh (GSSM), Buzzards Bay, Massachusetts.
2. Nitrogen cycling in GSSM.
3. Long-term pollution effects in GSSM.
4. Natural and artificial marsh restoration consulting in Los Angeles delta (CA), Delaware Bay (NJ), Copper River delta (AK), and Indian River (FL).

Mississippi**Mr. Hollis H. Allen**

U.S. Army Corps of Engineers
Waterways Experiment Station
3909 Halls Ferry Road
Vicksburg, MS 39180-6199

Work phone: (601) 634-3845

FAX: (601) 634-4016

Wetlands engineer; restores wetlands and uses bio-engineering to control erosion.

Projects:

Numerous freshwater, brackish and salt water marsh creation and bank stabilization projects.

Mr. Ellis J. Clairain, Jr.

U.S. Army Corps of Engineers
Waterways Experiment Station, Wetlands Ecology Group
3909 Halls Ferry Road
Vicksburg, MS 39180-6199

Work phone: (601) 634-3774

FAX: (601) 634-4016

Wetlands biologist.

Projects:

Development of wetland evaluation techniques.

Dr. Douglas G. Clarke

US Army Corps of Engineers
 Waterways Experiment Station, Environmental
 Laboratory
 3909 Halls Ferry Road
 Vicksburg, MS 39180-6199
 Work phone: (601) 634-3770
 FAX: (601) 634-4016

Oceanographer.

Projects:

1. Principal investigator of Corps WRP research area on coastal intertidal wetlands.

Site A - Winyah Bay, S. Carolina. *Spartina* marsh naturally established (not transplanted) on dredged material from maintenance dredging projects. Cooperative study of fauna and flora in various age zones in the established marsh. Participants include the Univ. of N. Carolina at Wilmington and the Univ. of S. Carolina Belle Baruch Marine Institute.

Site B - Pelican Spit, Galveston Bay, Texas. Transplanted marsh on dredged material derived from maintenance dredging of intra-coastal waterway. Cooperative project with National Marine Fisheries Service Galveston Laboratory (Tom Minello).

Mr. Gary J. Cuevas

Mississippi Department of Wildlife, Fisheries and Parks
 Bureau of Marine Resources
 2620 Beach Blvd.
 Biloxi, MS 39531

Work phone: (601) 385-5884

FAX: (601) 385-5864

Marine Program Manager.

Projects:

1. Marsh site at Pass Christian on Menge Avenue, a small mitigation project.

Dr. Lionel Eleuterius

Gulf Coast Research Laboratory
 P.O. Box 7000
 Ocean Springs, MS 39564

Work phone: (601) 872-4200

FAX: (601) 872-4204

Dr. Mary C. Landin

U.S. Army Corps of Engineers
 Waterways Experiment Station, Environmental
 Laboratory
 3909 Halls Ferry Road
 Vicksburg, MS 39180-6199

Work phone: (601) 634-2942, (800) 522-6937
 ext. 2942

FAX: (601) 634-4016

Research Biologist.

Projects:

1. Designs, conducts and coordinates many wetland restoration and creation projects involving the use of *Spartina alterniflora* on dredged material placement sites (for the past 20 years) and mitigation projects.
2. Manages COE nation-wide wetlands restoration, protection and establishment research program.
3. Is a member of the National Academy of Sciences committee on the role of technology in Coastal Habitat Restoration and Protection. Producing a final book in the summer, 1993.
4. Manages the monitoring program where *Spartina alterniflora* was used for coastal restoration/creation, sites include: 1) Buttermilk Sound, GA; 2) Apalachicola Bay, FL; 3) Gaillard Island, AL; 4) Bolivar Peninsula, TX; 5) several sites in Chesapeake Bay, MD & VA; 6) Winyah Bay, SC (natural colonization by *S.a.*); 7) several sites in LA & MS; 8) several sites in Tampa Bay, FL.
5. Two EPA-funded restoration studies along the Gulf of Mexico; mostly using *Spartina alterniflora*.

Dr. Mark W. LaSalle

Mississippi State University
 Coastal Research and Extension Center
 2710 Beach Blvd., Suite 1-E
 Biloxi, MS 39531

Work phone: (601) 388-4710

FAX: (601) 388-1375

Marine Resource Specialist, Mississippi Sea Grant Office.

Projects:

1. Evaluating transplanted marshes in Mississippi (some *Spartina* but mostly *Juncus*).
2. With COE - coastal wetland management protocol techniques for evaluating relative functional level of marsh - 1 year to completion.
3. Evaluation of different wetland type functions - preliminary work.
4. Evaluating thin layer disposal of dredge materials in marshes as a soil accretion mitigation tool.

Mr. John H. Lowe

Chevron USA Products
 Pascagoula Refinery
 P.O. Box 1300
 Pascagoula, MS 39568-1300

Work phone: (601) 938-4600

(601) 938-4363

FAX: (601) 938-4230

Environmental Specialist.

Projects:

1. Graded-down 35 acres of pine plantation into 25 acres of *Juncus/Spartina alterniflora* marsh; 1985.
2. Seeking permit for creation of a 1.4 acre *Spartina alterniflora* marsh; mitigation coupled to a water divergence plan; 1993.

Dr. L. Jean O'Neil

U.S. Army Corps of Engineers
 Waterways Experiment Station, Stewardship Branch
 CEWESEN-S
 3909 Halls Ferry Road
 Vicksburg, MS 39180
 Work phone: (601) 634-3641
 FAX: -

Mr. Michael R. Waring

U.S. Army Corps of Engineers
 Waterways Experiment Station
 P.O. Box 631
 Vicksburg, MS 39180
 Work phone: (601) 634-2290
 FAX: -

Biologist; Environmental laboratory.

Projects:

1. Whooping crane habitat characterization, Aransas National Wildlife Refuge, Texas.

New Hampshire**Ms. Sarah D. Allen**

Normandeau Associates, Inc.
 25 Nashua Road
 Bedford, NH 03110
 Work phone: (603) 472-5191
 FAX: (603) 472-7052

Wetland ecologist.

Projects:

1. A manual for monitoring mitigation and restoration projects of New Hampshire's salt marshes; 1992. Includes sampling methods, data collection forms and assessment criteria, plus extensive background and literature on local functions and technical considerations.

Dr. Alan Phelps Ammann

U.S. Soil Conservation Service
 New Hampshire State Office
 Federal Building
 Durham, NH 03824
 Work phone: (603) 868-7581
 FAX: (603) 868-5301

Biologist; wetlands, water quality and wildlife habitat responsibilities.

Projects:

- Rye Harbor. 1992. Restored a portion of salt marsh formerly used as harbor dredge material deposition site. This project breached the dike to allow saltwater access to the interior. *Spartina patens* is becoming established in the high marsh area, and the borders of the canals are supporting small stands of *Spartina alterniflora*, as are presently growing around the dike perimeter. *Phragmites* should be reduced as saltwater infiltrates.

Dr. David M. Burdick

University of New Hampshire
 Dept. of Natural Resources
 Jackson Estuarine Laboratory
 85 Adams Point Road, R.F.D. 2
 Durham, NH 03824-3406
 Work phone: (603) 862-2175
 FAX: (603) 862-1101

Research/ coastal plant ecology, landscape ecology.

Projects:

1. Halophyte Biology Lab saltmarsh, 3/90.
2. Awcomin Restoration, 4/92
3. Portsmouth Port Expansion, N. Mill Pond, proposed for 6/93.

Mr. Peter S. Helm

New Hampshire, Office of State Planning,
 Coastal Program
 2-1/2 Beacon Street
 Concord, NH 03301
 Work phone: (603) 271-2155
 FAX: (603) 271-1728

Director of Coastal Program; management, wetland restoration planning.

Projects:

1. Overseeing coastal restoration activities for the state of New Hampshire.
2. Managing the efforts to restore a 35-acre dredged material deposition site to a *Spartina alterniflora* salt marsh -- the Awcomin Marsh Restoration Project.
3. Project management of a 15-acre saltmarsh restoration of a former dredged material disposal site along the NM seacoast. Material removal and channel excavation is required; extensive monitoring is also part of the project.
4. Stuart Farm Marsh Restoration project, 1993. Installing of large box culverts will reintroduce tidal exchange to a 10-acre area; project will be monitored for marsh development.

New Jersey**Dr. James F. De Bouno, Ph.D.**

James De Bouno & Associates, Inc.
 Environmental Engineering
 P.O. Box 265, 28 Teabury Court
 Marlton, NJ 08053
 Work phone: (609) 985-0455
 FAX: (609) 228-7811

Environmental Scientist/Engineer; wetlands, wildlife, and plant studies.

Projects:

- more...
1. Inland Project, California Villa wetland restoration in North Hanover, Burlington County, NJ, 1992.
- Rechannelization of stream, restored grades and wetlands

in a small unnamed stream off the North Run River to help remedy the problem of unauthorized disturbance. (more)

2. Birch Tree Developers restoration; Bergen County, NJ, 1991. Channel bank restoration by fill removal, grading, and replanting wetland.

Mr. Oliver T. Edstrom

Associated Environmental Applications
901 Clarks Landing Road
Egg Harbor City, NJ 08215

Work phone: (609) 965-4034

FAX: (call work phone to access FAX)

Consultant; wetlands, enhancement, bank protection, restoration of coastal marshes.

Projects:

1. Several projects involving the planting of *Spartina alterniflora* in coastal areas.

-60 acres of coastal wetlands in New Jersey.

-3.5 acres: intercepting lot, dredge/disposal area.

-5.5 acres: commercial lot.

-1.5 acres: Ferry Terminal enlargement.

-7 acres: high school area.

2. Restoration after construction projects

-projects involved a variety of restoration plans including sites with intercepting pipelines and roadways, aqueduct installations, and force mains.

Mr. James M. Eisenhardt

Joseph L. Lomax and Associates, Inc.
P.O. Box 9

Cape May Court House, NJ 08210

Work phone: (609) 465-9857

FAX: (609) 465-2449

Vice-president; wetlands biologist, delineation, design, mitigation restoration.

Projects:

1. Mitigation project where 3-4 acres of coastal land was restored back to tidal marsh from higher elevations that had been taken over by *Phragmites*.

2. Several projects where 3-10 acre areas of uplands were converted to tidal marsh, with successful establishment of *Spartina*.

3. Many projects restoring freshwater wetlands.

Mr. Donald W. Hamer

U.S. Soil Conservation Service
Cape May Plant Materials Center
1536 Route 9N

Cape May Court House, NJ 08210

Work phone: (609) 465-5901

FAX: (609) 465-9284

Agronomist/manager; tidal bank restoration and stabilization.

Projects:

1. Developing hardy strains of *Spartina*.

2. Evaluation of *S. patens* for tidal bank stabilization: Choptank River, MD., Broad Bay, VA., Albemarle Sound, NC., and Chesapeake Bay, MD.

3. Evaluation of *S. alterniflora* for tidal bank stabilization: Broad Bay, MD., Occohannock Creek, VA., Choptank River, MD., and Chester River, MD.

Mr. Joseph L. Lomax

Joseph L. Lomax and Associates, Inc.
P.O. Box 9

Cape May Court House, NJ 08210

Work phone: (609) 465-9857

FAX: (609) 465-2449

Vice-president; wetlands biologist, delineation, design, mitigation restoration.

Projects:

1. Mitigation project where 3-4 acres of coastal land was restored back to tidal marsh from higher elevations that had been taken over by *Phragmites*.

2. Several projects where 3-10 acre areas of uplands were converted to tidal marsh, with successful establishment of *Spartina*.

3. Many projects restoring freshwater wetlands.

Mr. Christopher F. Miller

U.S. Soil Conservation Service
1370 Hamilton Street
Somerset, NJ 08873

Work phone: (908) 246-1205, Ext. 144

FAX: (908) 246-2358

Plant Materials Specialist, Certified. Stabilization of dunes and tidal shoreline.

Projects:

1. Develop and release improved varieties of *Spartina* for use in shoreline stabilization.

2. Evaluate and develop bioengineering techniques utilizing *Spartina*.

3. Provide technical assistance to commercial growers of *Spartina*.

Mr. Ronald W. Prann

International Technology Corporation
Wetland Division
165 Fieldcrest Avenue
Edison, NJ 08837

Work phone: (908) 225-2000

FAX: (908) 225-1691

Wetland delineations and land use issues.

Projects:

1. Formosa *Spartina* creation project, 1987. Arthur Kill, NJ/NY Waterway, Cateret, NJ.

2. Navy *Spartina* restoration, 1992, Norfolk, VA.

3. *Spartina* damage assessment following 567,000 gallon No.2 fuel oil spill, Arthur Kill, 1990-1992.

Dr. Joseph K. Shisler

Shisler Environmental Consultants, Inc.
23 Running Brook Drive
Hightown, NJ 08520 (more)

Work phone: (908) 446-3669
FAX: (908) 446-2381

Wetlands Ecologist; Mitigation and Restoration.

Projects:

1. Mores Beach, 1972, 600 acres of tidal restoration of a salt hay farm, Cumberland County, NJ.
2. Island Beach State Park, 5 acres of restoration of a wetland system, Ocean County, NJ.
3. Wheelers Hay-Farm, 300 acres of tidal restoration to a salt hay farm, Cape May County, NJ.

New York**Ms. Betsy Adamson**

New York City Dept. of Environmental Protection
Natural Systems Planning
59-17 Junction Blvd.
Elmhurst, NY 11373

Work phone: (718) 595-3509
FAX: (718) 595-3557

Director; wetland restoration, enhancement, inventory of natural resources, and impact analysis.

Projects:

1. Several projects were implemented where sewer outfall construction requires regrading, restoration or enhancement of degraded shoreline, 1993.
2. April-June, 1993, enhancement of shoreline near the Pelham Municipal Landfill for the treatment of subsurface seeps of landfill leachate.
3. Proposed project - Restoration of high marsh area in Broad Channel, an inhabited island in the middle of Jamaica Bay Wildlife Refuge, where intertidal and high marsh have been lost from filling.

Mr. Michael P. Bontje

B. Laing Associates,
Environmental Science
225 Main Street
Northport, NY 11768

Work phone: (516) 261-7170
FAX: (516) 261-7454

President; Environmental design and wetlands restorations.

Projects:

Mr. Sven Hoeger

Creative Habitat Corp.
253 Old Tarrytown Road
White Plains, NY 10603

Work phone: (914) 948-4389
FAX: (914) 681-5136

President; Hydro/soil bioengineer and wetlands ecologist.

Projects:

1. Tidal creek salt marsh restoration by Fresh Kills Landfill, 1990.

Mr. Leonard Houston

U.S. Army Corps of Engineers
Special Projects Section - Environmental Branch
26 Federal Plaza
New York, NY 10278

Work phone: (212) 264-1275
FAX: (212) 264-5472

Supervisory fisheries biologist; restoration and monitoring.

Projects:

1. Hemstead Harbor, NY. Replanting areas of existing marsh with *Spartina alterniflora*; about 2 acres. 1993.

Mr. Marc A. Matsil

Director Natural Resources Group
City of New York - Parks and Recreation
1234 5th Avenue, Rm. 233
New York, NY 10029

Work phone: (212) 360-1417
FAX: (212) 360-1426

Director of NRG; restoration ecology.

Projects:

1. NYC Parks Harbor Herons Wetlands Preserve. Restoration of salt marshes damaged by the 1990 Exxon oil spill; also involves a 6 year restoration, research and monitoring plan.
2. Alley Creek, Alley Pond Park, Queens, NY; *Spartina* salt marsh restoration; completion expected in 1994.
3. Plum Beach salt marsh restoration and dune restoration project; a collaborative effort between NYC Parks Natural Resources Group and the National Parks Service (Gateway NRA); completion expected in 1993.

North Carolina**Dr. Stephen W. Broome**

North Carolina State University
Department of Soil Science, Box 7619
Raleigh, NC 27695-7619

Work phone: (919) 515-2643
FAX: (919) 515-2167

Associate Professor of biology and ecology; ecosystems, wetlands, macrophytes, soil science, restoration, mitigation.

Projects:

1. North Carolina, South Carolina, Louisiana, and France (Brittany Coast, site of Amoco Cadiz oil spill): effects of oil spills and cleanup activities on coastal marshes (more)
2. Past projects encompassed developing methods for

Spartina alterniflora marsh creation and determining productivity and functional equivalence.

Dr. Christopher B. Craft

Duke University Wetland Center
School of the Environment
Duke University
Durham, NC 27706

Work phone: (919) 684-2619

FAX: -

Research Assistant Professor; Biogeochemical cycling in natural and created marshes.

Projects:

1. Dredge spoil stabilization with *Spartina alterniflora*, Spring '87. Hacker's Island-Swansboro-Sneads Ferry, North Carolina. Cooperative project with USCOE and NMFS.
2. Mitigation project, creation of *S. alterniflora* marsh, Spring '89. Cape Fear, North Carolina. Cooperative project with NC Ports Authority.

Ms. Carolyn Currin

National Marine Fisheries Service
SEFC Beaufort Laboratory
101 Pivers Island Rd.
Beaufort, NC 28516

Work phone: (919) 728-8749

FAX: (919) 728-8784

Microbiologist; Nitrogen cycling and wetlands restoration.

Projects:

1. Accelerated functional development in transplanted salt marshes. 6/90-93.
2. Evaluation of restored wetlands and enhancement methods for existing restorations. 1992.

Mr. Mark S. Fonseca

National Marine Fisheries Service
SEFC Beaufort Laboratory
101 Pivers Island Rd.
Beaufort, NC 28516

Work phone: (919) 728-8729

FAX: (919) 728-8784

Ecologist. Seagrass ecology and restoration.

Projects:

1. NMFS/CORPS MOA-North Carolina, 1985 to present; research involving marsh restoration at Harker Island, Swansboro, and Snead's Ferry, N.C.
2. NMFS-North Carolina, Maryland, Florida, St. Croix, Texas, California, Washington and SE Asia, 1977-present; research in seagrass ecology and restoration including hydrodynamics and landscape pattern.

Mr. Jeffrey C. Furness

Texasgulf Inc.
P.O. Box 48
Aurora, NC 27806

Work phone: (919) 322-4111

FAX: (919) 322-4444

Environmental Scientist; consulting, marsh creation and monitoring.

Projects:

1. Planted 10.1 acres of coastal marsh using *Spartina alterniflora*, *S. patens*, *S. cynosuroides*, and *Juncus roemerianus* adjacent to South Creek, a tributary of the Pamlico River, near Aurora, NC. Studies were made of vegetative growth and soil development over several years following planting.
2. Planted 1.6 acres of marsh using the three *Spartina* species in the same vicinity. Studies are being made of the fish and benthic populations in the marsh, comparing them with those in a nearby natural marsh.
3. Planted 8.8 acres of the *Spartina spp.* and *Juncus*. Studies of vegetative growth, fish populations, and invertebrate colonization have been done comparing the man-made to natural marsh habitat.

Mr. Douglas Huggett

N.C. Dept. of Transportation, Division of Highways & Planning, Environmental Branch
P.O. Box 25201
Raleigh, NC 27611

Work phone: (919) 733-9770

FAX: (919) 733-9794

Environmental Specialist.

Projects:

1. Onslow County, N.C. Created 1.3 acres of *Spartina alterniflora* and *Spartina patens* marsh to offset impacts of construction over 0.5 acres of natural marsh at New River inlet.
2. Dare County, N.C. Created 0.6 acres of *Juncus* marsh to offset impacts of two new bridge constructions at Nag's Head, and created 1.5 acres of *Juncus* marsh near Mantia to offset impacts of a bridge.
3. Working with U.S. Fish and Wildlife Service to put in *Scirpus* marsh at Pea Island Wildlife Refuge.
4. Carteret County, N.C. Creating 2.0 acres of *Juncus* marsh at Cedar Island to mitigate construction impacts on natural *Juncus* marsh.

Dr. Judson W. Kenworthy

National Marine Fisheries Service
SEFC Beaufort Laboratory
101 Pivers Island Rd.
Beaufort, NC 28516

Work phone: (919) 728-8750

FAX: (919) 728-8784

Research Fisheries Biologist. Ecology and seagrasses.

Projects:

North Carolina/ Corps MOA 1985 to Present.

Mr. David L. Meyer
National Marine Fisheries Service
SEFC Beaufort Laboratory
101 Pivers Island Rd.
Beaufort, NC 28516

Work phone: (919) 728-8714
FAX: (919) 728-8784

Fisheries Biologist. Invertebrate zoology and estuarine ecology.

Projects:

1. NMFS/CORPS MOA-1985 through present.
2. Evaluation of restored wetlands and enhancement methods for existing restored marshes. 6-92 Start.

Dr. Ernest D. Seneca
North Carolina State University
Botany Department
P.O. Box 7612
Raleigh, NC 27695-7612

Work phone: (919) 515-2727
FAX: (919) 515-3436

Professor and Head of Dept. Establishment, rehabilitation and restoration of coastal dunes and salt marshes.

Projects:

1. Numerous coastal marsh establishments and restorations among the Atlantic estuaries of the US, using *Spartina alterniflora*.
2. Many publications about materials and methods to create or restore salt marshes using *S. alterniflora*, primarily.
3. Restoration of salt marshes along the Brittany coast of France following the Amoco Cadiz oil spill.

Dr. Gordon W. Thayer
National Marine Fisheries Service
SEFC Beaufort Laboratory
101 Pivers Island Rd.
Beaufort, NC 28516

Work phone: (919) 228-8747
FAX: (919) 728-8784

SEFSC Habitat Coordinator, Fishery habitat Research Team Leader, and Fisheries ecologist. Also serves as the Research Coordinator for the NOAA Restoration Center.

Projects:

1. Studying seagrass and salt marsh habitat and restoration from N.C. to Florida from a fisheries perspective.
2. Studying mangrove habitats of South Florida.
3. Advises on salt marsh ecology, mitigation, and restoration, particularly along the mid-Atlantic coastal states.

Ms. Gwendolyn Ann Thunhorst
University of North Carolina
Department of Biology/Ecology
424 Hickory Drive
Chapel Hill, NC 27514

Work phone: FAX: Graduate Student.
Wetland ecology and botany research.

Projects:

Hartz Mountain Wetland Restoration Research-
Determination of Reasons for the failure of seeded
Spartina alterniflora (Environmental Concerns, INC.) May
1991

Oregon

Mr. Robert L. Emmett
National Marine Fisheries Service
Hammond Laboratory
P.O. Box 187
Hammond, OR 97121

Work phone: (503) 861-1818
FAX: -

Fisheries research; estuarine salmonid habitat.

Projects:

1. Interested in usage of *Spartina alterniflora* marshes by fish and invertebrates.
2. Cooperative study with the Corps of Engineers concerning stranding of juvenile salmonids by ship wakes in the Columbia River.
3. Potential use of dredged material to rebuild intertidal areas that have been scoured, and possible stabilization using marsh grasses.

Ms. Paula Levin
U.S. Fish and Wildlife Service
Habitat Conservation / Federal Activities
911 NE 11th Ave.
Portland, OR 97232

Work phone: (503) 231-2068
FAX: (503) 231-2080

Coastal Program Coordinator; wetlands and regulatory activities.

Mr. John Marshall
U.S. Fish and Wildlife Service
Oregon State Office
2600 SE 98th Ave., Suite 100
Portland, OR 97266

Work phone: (503) 231-6179
FAX: (503) 231-6195

Biologist; coastal permits and NEPA review.

Projects:

1. Proposed salt marsh restoration by dike breaching on the Salmon River Estuary. U.S. Forest Service.
2. Restoration of tidal exchange to a tidal salt marsh in Trestle Bay by breaching a jetty and a causeway. U.S. Army Corps of Engineers and Oregon State Parks Dept.

Mr. Richard P. Novitzki
ManTech Environmental Technology, Inc.
1600 SW Western Blvd.
Corvallis, OR 97333

Work phone: (503) 754-4666
 FAX: (503) 754-4335
 Senior Scientist; EMAP wetlands.
 Projects:

Pennsylvania

Dr. Keith L. Bildstein

Hawk Mountain Sanctuary
 R.R. 2, Box 191
 Kempton, PA 19526

Work phone: (215) 756-6961
 FAX: (215) 756-4468

Director of Research. Avian ecology and fiddler crab ecology.

Projects:

1. Long term study of white ibis, their breeding and feeding ecology in S. Carolina freshwater, brackish water and *Spartina alterniflora* marshes.
2. Long term effects of hurricane Hugo on the breeding ecology of wading birds in coastal salt marshes in S. Carolina.
3. Ecological studies of fiddler crabs, their burrowing behavior in *Spartina* and brackish marshes.

Mr. Edward R. Bonner

U.S. Army Corps of Engineers, Philadelphia District
 Regulatory Branch
 100 Penn Square East, Wanamaker Building
 Philadelphia, PA 19107-3390

Work phone: (215) 656-6728
 FAX: (215) 656-6724

Senior biologist; permitting and mitigation.

Projects:

1. Delaware DOT mitigation on the St. Jones River established about 4 acres of *Spartina alterniflora* marsh in 1986. Planted by Environmental Concern, Inc.
2. At Siliguini, about 1 acre of irregularly flooded salt marsh having *Spartina alterniflora*, *Salicornia* sp., and *Distichlis spicata*.

Mr. Kevin W. Dougherty

A.D. Marble & Company
 1062 Lancaster Ave.
 Rosemont, PA 19010

Work phone: (215) 527-9311
 FAX: (215) 527-5646

Senior environmental planner, and Certified Ecologist, ESA; freshwater and tidal wetlands.

Projects:

1. A 0.5 acre low marsh of *Spartina alterniflora* in Cape May County, NJ; design approved, construction scheduled for March 1993.

2. A 2.3 acre tidal freshwater marsh restoration in Prince George's County, MD; design approved August 1991.
3. A 0.2 acre tidal freshwater marsh restoration for oil damage in Monmouth County, NJ, in 1990.

Mr. Barry Dubinski

Roy F. Weston, Inc.
 1 Weston Way

West Chester, PA 19380

Work phone: (215) 430-3137
 FAX: (215) 430-7401

Dr. Michael A. Hardisky

University of Scranton
 Biology Department
 Scranton, PA 18510

Work phone: (717) 941-7514
 FAX: (717) 941-6369

Professor; researcher.

Dr. Willard C. McCartney

Michael Baker, Jr., Inc.
 Transportation Planning (Environmental)
 Airport Office Park, 420 Rouser Rd., Bldg.#3
 Coraopolis, PA 15108

Work phone: (412) 269-4601
 FAX: (412) 269-2048

Environmental Manager; Vertebrate and wetland ecology.

Projects:

1. Plum Beach Restoration, 9-92. Shoreline stabilization and marsh restoration.
2. Several freshwater wetland restoration projects.

Mr. Schucet

Baker Engineers
 Building 3, 420 Rouser Road
 Coraopolis, PA 15108

Work phone: (412) 269-2048
 FAX: (412) 269-4604

Projects in both New York and Virginia.

Rhode Island

Mr. Joseph Dowhan

U.S. Fish and Wildlife Service
 South New England Estuary Project
 Shoreline Plaza
 P.O. Box 307

Charlestown, RI 02813

Work phone: (401) 364-9124
 FAX: (401) 364-0170

Dr. Frank Golet
 University of Rhode Island
 Department of Natural Resources Science
 Woodward Hall
 Kingston, RI 02881

Work phone: (401) 792-2916
 FAX: (401) 792-4561

Professor and research scientist; avian and wetland ecology.

Projects:

1. Narragansett, RI. Baseline ecological study of the 125 acre Galilee Bird Sanctuary regarding tidal regime, ground water levels, vegetation types and habitats, salinity profiles, and use of the area by migratory and locally breeding populations of birds. The sanctuary, which was largely cut off from tidal flow in 1956--and is now brackish--will be restored to salt marsh in 1994. GIS technology is being used to compare historic conditions to current conditions to aid in designing the restoration plan.
2. 25 years of research studies involving the ecology of forested wetlands.
3. Recent studies of bird usage of coastal marshes, with varying degrees of mosquito control ditching.

Mr. Scott S. Hobson
 Environmental Scientific Corporation
 Division of Keyes Associates
 10 Lincoln Center Boulevard
 Lincoln, RI 02865

Work phone: (401) 333-0100 ext. 27
 FAX: (401) 333-4556

Environmental scientist; wetlands delineation and permitting.

Projects:

Assist individuals, companies and agencies with delineation of wetlands, mitigation activities including restoration design and obtaining permits from local, state, and federal agencies when necessary.

Mr. Paul J. Shea, MA
 Independent Environmental Consultants, Inc.
 32 Levitt Lane
 Exeter, RI 02822

Work phone: (401) 294-3909
 FAX: -

Biologist, hydrologist. Environmental assessments; coastal and freshwater wetlands.

Projects:

1. Coastal restoration of Winnapaug Pond, assessment of critical area, 1992.
2. Numerous freshwater wetland restorations.

Ms. Dana Zewinski
 Environmental Scientific Corporation
 Division of Keyes Associates
 10 Lincoln Center Boulevard
 Lincoln, RI 02865

Work phone: (401) 333-0100
 FAX: (401) 333-4556

Environmental scientist; wetlands delineation and permitting.

Projects:

Assist individuals, companies and agencies with delineation of wetlands, mitigation activities including restoration design and obtaining permits from local, state, and federal agencies when necessary.

South Carolina

Dr. Vincent J. Bellis, Jr.
 East Carolina University, Department of Biology
 P.O. Box 4353
 Greenville, SC 27858-4353

Work phone: (919) 757-6296
 FAX: (919) 757-4178

Professor of Biology; Wetlands and resource management.

Projects:

Primary production in saltmarshes (*Spartina patens* and *S. cynosuroides*) by algae and diatoms.

Dr. Elizabeth R. Blood
 University of South Carolina
 Department of Environmental Health Services
 Columbia, SC 29208

Work phone: (803) 777-6994
 FAX: (803) 777-4783

Associate Professor. Biogeochemistry and nitrogen cycling.

Projects:

Nitrogen cycling in *Spartina* marshes in the Charleston, S.C. area.

Mr. Andrew L. Bunch
 R. S. Industries
 P.O. Box 1594
 Georgetown, SC 29442-1594

Work phone: (803) 527-2204
 FAX: -

Executive, Foreign Relations Technical Div. Database research/compilation and military assessments and affairs.

Projects:

1. Wetlands database compilation, 8-91.
2. *Spartina alterniflora* database and directory construction, 6-92.

Dr. Robert R. Christian
 East Carolina University, Biology Department
 P.O. Box 4353
 Greenville, SC 27858

Work phone: (919) 757-6287
 FAX: (919) 757-4178

Professor of Biology; Estuarine microbial ecology and system ecology.

Projects:

1. High marsh dynamics and the influence of tidal inundation.
2. Below ground processes of production and decomposition.
3. Nitrogen cycling and exchanges between sediment and water.

Mr. John Hensel

S.C. Coastal Council, Permitting
4130 Faber Place, Suite 300
Charleston, SC 29405

Work phone: (803) 744-5838

FAX: (803) 744-5847

Chief of Permit Enforcement

Projects:

1. 1993-present. SCDHPT Highway and bridge construction project. Grade down old causeway to elevation of adjacent marsh to allow revegetation by marsh plants of 4.5 acres of intertidal zone.
2. 1993-present. North Charleston Sewer District mitigation for fill associated with the expansion of a waste treatment facility. Wooded upland (0.8 acre) was graded to intertidal elevations to allow revegetation by surrounding marshes in the Ashley River.
3. Mitigation by Walling Grove Development Co. for fill allowed to widen an existing causeway. Project involves establishing brackish marsh in an area of graded down upland; 0.1 acre.

Dr. Eric Thor Koepfler

University of South Carolina - Coastal Carolina College
P.O. Box 1954
Conway, SC 29526

Work phone: (803) 349-2222

FAX: (803) 349-2990

Assistant Professor.

Projects:

Testing a theory between the interplay between ecosystem maturation and physical development coastal systems.

Dr. Richard H. Moore

University of South Carolina
Coastal Carolina College
Box 1954
Conway, SC 29526

Work phone: (803) 349-2050

FAX: (803) 349-2900

Professor and Ass. Vice Chancellor

Bio-Ecology; community, wetlands, nekton, fish.

Projects:

1. A community ecology study of estuarine fishes in Singleton Swash, SC, a *Spartina* marsh inlet.

2. Estuarine study of the biology of the spotfin mojarra, *Eucinostomus argenteus*.

Mr. Steve Moore

South Carolina Coastal Council
4130 Faber Place Suite 300
Charleston, SC 29405

Work phone: (803) 744-5838

FAX: (803) 744-5847

Permit Administrator.

Projects:

Administers permits for wetlands

Dr. James T. Morris

University of South Carolina
Department of Biological Sciences
Columbia, SC 29208

Work phone: (803) 777-3940

FAX: (803) 777-4002

Professor. Wetlands ecology and plant physical ecology.

Projects:

1. Effect of nitrogen loading on wetlands ecosystem: particular reference to atmospheric deposition. 1992

Dr. Harold Ornes

University of South Carolina
Department of Biology and Geology
171 University Parkway
Aiken, SC 29801

Work phone: (803) 648-6851

FAX: (803) 641-3382

Chairman, Department of Biology & Ecology; Ecology of wetlands (algae, macrophytes, nutrients, sediments).

Projects:

1. Nutrient status of *Spartina* in North Inlet, S.C.
2. Live and dead stems and roots nutrient composition of *Spartina* at North Inlet, S.C.
3. Monitoring of the vegetation after cleanup and restoration of Lost Lake, a Carolina Bay at the Savannah River Site, Aiken, S.C.

Mr. Mark Purcell

U.S. Army Corps of Engineers, Regulatory Branch
P.O. Box 919
Charleston, SC 29402

Dr. Robert F. Van Dolah

S.C. Wildlife & Marine Resource Dept.
Marine Resource Research Institute
P.O. Box 12559
Charleston, SC 29412

Work phone: (803) 762-5048

FAX: (803) 762-5110

Assistant Director. Benthic ecology and environmental assessment.

Texas

Mr. John S. Adams

The Conrad Blucher Institute for Surveying & Science
Texas A & M University - Corpus Christi
6300 Ocean Drive

Corpus Christi, TX 78412

Work phone: (512) 994-2376

FAX: (512) 994-2715

Biologist; estuarine ecosystems.

Projects: (more)

1. Applications of telemetric collection of coastal environmental data.
2. Monitoring of 'attempted' creation of a *Spartina alterniflora* salt marsh in the Nueces River delta, Texas, 1990-1991.
3. Creation of a 'mini' *Spartina alterniflora* salt marsh in relation to water level fluctuations and soil elevation in the Nueces River delta.

Mr. Charles E. Belaire

Belaire Consulting, Inc.

P.O. Box 741

Rockport, TX 78382

Work phone: (512) 729-2948

more...

FAX: (512) 729-2970

President; wetlands planning, design, and construction.

Projects:

1. Creation of 7.5 acres of *Spartina alterniflora* marsh in Mesquite Bay, Texas, 1992. A mitigation project by Mitchell Energy Corp.
2. Creation of two *S. alterniflora* marshes totaling 5 acres on Pelican Spit in Galveston Bay, Texas, 1991. COE projects utilizing dredged material to fill low areas.
3. Kon Tiki Condos mitigation project in Aransas Bay created a small 0.1 acre wetland.
4. Planning, permitting, installation, and monitoring of 54 acres of seagrass habitat to restore and mitigate impacts due to construction of a 20" Transco pipeline in the Upper Laguna Madre, Texas, 1989-1990.
5. Planting of 45 acres of seagrass in the Lower Laguna Madre to restore impacts due to dredge material disposal for Energy Development Corp., 1990.

Mr. Paul Carangelo, R.E.M.

Island Botanics, Environmental Consultants

3734 Flour Bluff Dr.

Corpus Christi, TX 78418

Work phone: (512) 937-4873

FAX: -

Environmental and regulatory management; Ecological consulting, habitat restoration, wetland and seagrass bed evaluation, restoration, and construction.

Projects:

1. Texas State Aquarium. Designed and built a 1.25 acre marsh exhibit which included transitional habitats from uplands to *Spartina alterniflora* marsh to subtidal seagrass beds and oyster reefs.
2. San Patricio County Navigation District No.1. Designed and built a *Spartina alterniflora* wetlands for mitigation of marina facility expansion; Redfish Bay, Texas.
3. Confidential client. Major oil company. Designed and built a seagrass bed at a mitigation site in Laguna Madre, Texas.
4. Lake Padre Development Corp. Designed and built a 7.0 acre *Spartina alterniflora* / *Batis maritima*, brackish and fresh marsh and submergent habitat wetlands on North Padre Island, Texas.
5. Completed certification as a Registered Environmental Manager, Registration #3011.

Mrs. Patricia B. Clements

U.S. Fish and Wildlife Service

Texas A & M University - Corpus Christi

Campus Box 338

6300 Ocean Drive

Corpus Christi, TX 78412

Work phone: (512) 994-9005

FAX: (512) 994-8262

Fishery Biologist

Projects:

Evaluations of projects impacting fish and wildlife habitats.

Mr. Dolan Dunn

U.S. Army Corps of Engineers, Galveston District
Regulatory Branch

P.O. Box 1229

Galveston, TX 77553

Work phone: (409) 766-3935

FAX: (409) 766-3931

Chief, Special Actions Section.

Projects:

1. Jurisdiction from Cameron Parish, Louisiana, to Mexico for the Department of the Army Permit Program.

Mr. Rudy G. Esquivel

U.S. Soil Conservation Service

Plant Materials Center

Texas A&I University Campus Box 218

Kingsville, TX 78363

Work phone: (512) 595-3960

FAX: (512) 595-3713

Assistant Manager.

Projects:

1. Primary practices rearing of prairie grasses and forbes (switch grass, green sprinkle grass, Russian olive, kidney woods, *Pasculum*, alkali sacaton)
2. Starting *Spartina* germ plasm propagation project.

Mr. Philip O. Glass

U.S. Fish and Wildlife Service
17629 El Camino Real, Suite 211
Houston, TX 77058

Work phone: (713) 286-8282

FAX: (713) 488-5882

Fish and wildlife biologist

Projects:

1. Houston Ship Channel Widening Project. Creation of saltmarsh habitat on dredged material islands.
2. Colonial waterbird colony census and research in Galveston Bay.

Ms. Alisha R. Goldberg

National Marine Fisheries Service
SEFC Galveston Laboratory
4700 Ave U
Galveston, TX 77551-5997

Work phone: (409) 766-3550

FAX: (409) 766-3508

Fishery Biologist; benthic infauna, transplanted saltmarsh.

Projects:

1. Galveston Island State Park marsh evaluation for fisheries species.
2. A study of the development of benthic infauna in a transplanted *Spartina alterniflora* marsh, 1992-1993.

Mr. Richard Harrington

Texas Parks & Wildlife Department
Resource Protection Div., Habitat Assessment Branch
Texas A & M University - Corpus Christi
6300 Ocean Drive, Campus Box 317
Corpus Christi, TX 78412

Work phone: (512) 993-4491

FAX: (512) 993-4597

Regional biologist; protect wildlife and fishery resources and habitats.

Projects:

1. Provide recommendations regarding avoiding, minimizing, and compensating for wetland impacts.
2. Monitor required wetland, marsh, and seagrass bed mitigation projects for compliance and success.
3. Coordinate with the SCS concerning use of *Spartina alterniflora* for shoreline stabilization in middle and south Texas bays.
4. Recommend areas for beneficial use of dredged material, including areas for marsh creation, e.g. the 13 acres of marsh created by the Mitchell Energy Corporation in Mesquite Bay, Texas, which increases feeding habitat for the endangered whooping crane.

Ms. Lynnda M. Kahn

Texas General Land Office
111 West Wilson, Naylor Bldg.
Aransas Pass, TX 78336

Work phone: (512) 758-7228

FAX: (512) 758-1430

Environmental Quality Specialist; field inspections, including wetlands and coastal marshes.

Projects:

1. Clark Island mitigation in lower Laguna Madre involved excavation and planting of a 3 acre site with *S. alterniflora* and *Halodule wrightii*, 1989.
2. Pelone Island in Corpus Christi Bay had a 3 acre site excavated and 1.5 acres of it planted with *Halodule*, 1988.
3. Evaluation of a 20 acre excavated site in the upper Laguna Madre, which was done for the establishment of seagrass habitat. No planting was required, 1991.

Mr. Paul Lazarene

U.S. Army Corps of Engineers, Galveston District
Corpus Christi Area Office
P.O. Box 2948
Corpus Christi, TX 78403-2948

Work phone: (512) 880-8135

FAX: (512) 884-4959

Wetlands Biologist, regulatory.

Projects:

1. Wetland delineations, various.
2. Investigation of unauthorized activities in the coastal area.

Mr. John Lloyd-Reilley

U.S. Soil Conservation Service
Plant Materials Center
R.R. 1, Box 608-T
Kingsville, TX 78363

Work phone: (512) 595-1313

FAX: (512) 595-3713

Manager.

Projects:

1. Evaluates growth characteristics of grasses and forbes for soil and water conservation (sand dune stabilization, saline soil regeneration, water quality improvement for sewage treatment plants, wildlife habitat and cool-season forages).
2. *Spartina* plant propagation.
3. Out-planting for shoreline erosion and evaluating *Rhizoctenia* resistance and salinity tolerance.

Dr. Geoffrey A. Matthews

National Marine Fisheries Service
SEFC Galveston Laboratory
4700 Ave U
Galveston, TX 77551-5997

Work phone: (409) 766-3532

FAX: (409) 766-3508

Biological oceanographer; Habitat research; Forecast modeling.

Projects:

1. Marsh REC directory, annotated bibliography, and techniques evaluation.

2. Created/restored *Spartina alterniflora* marsh inventory and monitoring.

Mr. John E. Mauney

Number "1" Nursery
Landscape Division
131 Rabon Chapel Road
Montgomery, TX 77356

Work phone: (409) 588-2508,
1-800-288-2508

FAX: -

Owner/manager; Contract growing, reforestation, landscape consulting.

Projects:

1. Trout Creek Flood Control, Kirbyville, Texas. Nov. 1990 - reforestation using 800 1-gal trees 1-5 feet tall.
2. Harris County Flood Control District, Unit No.E500-04-00, mitigation planting and maintenance, Job No: 93/0094; planting of 5000 1-gal trees.
3. Growing *Spartina alterniflora* for restoration projects.

Mr. Rick Medina

U.S. Army Corps of Engineers, Galveston Dist.
Environmental Resources Branch
P.O. Box 1229
Galveston, TX 77550-1229

Work phone: (409) 766-3044
FAX: (409) 766-3064

Chief, Environmental Resources Branch

Projects:

1. Pelican Spit *Spartina alterniflora* marsh creation.
2. East Matagorda Bay *Spartina alterniflora* marsh creation.
3. Chocolate Bayou *Spartina alterniflora* marsh creation.

Dr. Thomas J. Minello

National Marine Fisheries Service
SEFC Galveston Laboratory
4700 Ave U
Galveston, TX 77551-5997

Work phone: (409) 766-3506
FAX: (409) 766-3508

Research Ecologist; habitat interactions with fishery species.

Projects:

1. Marsh REC directory, bibliography & evaluation of successful techniques.
2. Comparisons of fishery value in natural and created marshes.
3. Relationships between sediments and benthic infauna in created marshes.

Mr. Donald Moore

National Marine Fisheries Service
SEFC Habitat Protection Division
4700 Avenue U
Galveston, TX 77551-5997

Work phone: (409) 766-3699

FAX: (409) 766-3575

Director of NMFS Habitat Conservation office for Texas; fisheries and wetlands.

Projects:

1. Evaluates proposed changes to fisheries habitats, and makes recommendations which will minimize impacts to the habitats; else, seeks mitigation when habitat damages occur.
2. Mouth of the Colorado River, Texas, Diversion Project, June, 1989. Promoted the diversion of the river back into Matagorda Bay.
3. Corpus Christi, Texas, Ship Channel Project. Actions prevented deposition of dredged material which would have covered one fifth of Nueces Bay.
4. Houston-Galveston, Texas, Ship Channel Project. Delayed authorization for dredging until studies of the environmental impacts of using dredged material in the bay can be evaluated; authorization delayed until 1996.

Mr. Lloyd Mullins

Texas General Land Office, Field Operations
111 W. Wilson, 2nd floor Naylor Bldg.
Aransas Pass, TX 78336

Work phone: (512) 758-7228
FAX: (512) 758-1430

Office Manager and Field Biologist; wetland mitigation and permitting.

Projects:

1. Clark Island mitigation in lower Laguna Madre involved excavation and planting of a 3 acre site with *S. alterniflora* and *Halodule wrightii*, 1989.
2. Pelone Island in Corpus Christi Bay had a 3 acre site excavated and 1.5 acres of it planted with *Halodule*, 1988.
3. Evaluation of a 20 acre excavated site in the upper Laguna Madre, which was done for the establishment of seagrass habitat. No planting was required, 1991.

Mr. Robert William Nailon

ENSR Consulting and Engineering
Toxicology & Risk Management
3000 Richmond Ave.
Houston, TX 77098

Work phone: (713) 520-9900
FAX: (713) 520-6802

Senior Fisheries Biologist/Wetlands.

Projects:

1. East Galveston Bay, 1985, *Spartina alterniflora* marsh created on north shoreline; 9 mile fetch; difficult to establish but now functional marsh with fisheries organisms.
2. San Jacinto River, 1988, *Spartina alterniflora* marsh created on dredge material islands; difficult to establish without fencing plots, i.e.. predation by rabbits, grass carp, nutria, and grasshoppers.
3. Taylor Lake, 1991 (EPA grant); *Spartina alterniflora* marsh.

4. Swan Lake, Texas City area, 1992 (EPA grant); *Spartina alterniflora* marsh shows initial success.
5. Trinity River delta, 1991-1992 (EPA grant); *Spartina alterniflora* marsh showed moderate success but nutrient predation was a problem.
6. LaPorte, Little Cedar Bayou, 1991 (EPA grant); *Spartina alterniflora* marsh successful, but ship wake problem exists.
7. Morgans Point, 1991 (EPA grant); *Spartina alterniflora* marsh; initial success; ship wakes are again a problem.

Mr. Brien A. Nicolau

Texas A & M University - Corpus Christi
Center for Coastal Studies
6300 Ocean Drive
Corpus Christi, TX 78412
Work phone: (512) 994-2736 or 2465
FAX: -

Graduate Research Assistant. Benthos and epifauna of coastal wetlands.

Projects:

1. Monitoring of the benthos and epifauna in the area of a failed attempt at creating a *Spartina alterniflora* salt marsh in the Nueces River delta, Texas, 1991-1993.
2. Creation of a 'mini' *Spartina alterniflora* salt marsh in relation to water level fluctuations and soil elevation in the Nueces River delta.

Mrs. Katie Northrup

Northrup Associates, Inc.
9328 Westview Drive, Suite 100
Houston, TX 77055

Work phone: (713) 461-3489
FAX: (713) 461-2671

Environmental Planner; consulting.

Dr. Lawrence P. Rozas

National Marine Fisheries Service
SEFC Galveston Laboratory
4700 Avenue U
Galveston, TX 77551-5997

Work phone: (409) 766-3532
FAX: (409) 766-3508

Ecologist; Estuarine/wetlands, fisheries ecology.

Projects:

1. Influence of hydroperiod on use of Gulf coast marshes by fishes and macro crustaceans. 1990-93 studying relationship between salt marsh elevation and use by nekton; *Spartina alterniflora* and *Distichlis spicata* marshes.
2. Potential for enhancement of fisheries habitat by filling OCS pipeline canals. 1990-93 studying relationship between canal depth and habitat utilization in brackish and salt marshes.
3. Nekton use of the marsh surface: a comparison between channeled and natural marshes. 1989-1992 examined

Spartina alterniflora marsh-surface used by nekton on marshes adjacent to pipeline canals and on natural marshes.

4. Evaluation of marsh creation on Atkinson Island dredged material with respect to establishing a fisheries habitat.

Mr. Eddie Seidensticker

U.S. Soil Conservation Service
17324-A Hwy. 3
Webster, TX 77598

Work phone: (713) 332-3381

FAX: (713) 332-3253

Resource Conservationist/ vegetative shoreline erosion control specialist.

Projects:

1. Morgan's Point, shoreline stabilization using *Spartina alterniflora*; 1992.
2. Little Cedar Bayou, south shore stabilization with *S. alterniflora*; 1991, 1992.
3. Several plantings of *S. alterniflora* along the San Jacinto River; 1990-1992.
4. At least 12 other projects for shoreline stabilization using *Spartina alterniflora*, in the Galveston Bay area.

Ms. Linda Shead

Galveston Bay Foundation
17324-A Hwy. 3
Webster, TX 77598

Work phone: (713) 332-3381

FAX: (713) 332-3153

Executive Director.

Projects:

1. Morgan's Point, shoreline stabilization using *Spartina alterniflora*; 1992.
2. Little Cedar Bayou, south shore stabilization with *S. alterniflora*; 1991, 1992.
3. Several plantings of *S. alterniflora* along the San Jacinto River; 1990-1992.

Dr. R. Michael Smart

U.S. Army Corps of Engineers
Waterways Experiment Station
RR 3, Box 446
Lewisville, TX 75056-9720

Work phone: (214) 436-2215

FAX: (214) 436-1402

Research Team Leader; Aquatic plant ecologist.

Projects:

1. Studies with freshwater plants.

Mr. R. Darrell Smith

Jones/Smith Environmental Services, Inc.
P.O. Drawer M
Dickinson, TX 77539

Work phone: (713) 585-1766

FAX: (713) 331-4119 (more)

Aquatic Ecologist; wetlands and endangered species.
Mitigation plan for *Spartina* planting project. Located at the mouth of the San Jacinto River, 1993.

Mr. James A. Sutherlin

Texas Parks and Wildlife Department
J.D. Murphree Wildlife Management Area
10 Parks and Wildlife Drive
Port Arthur, TX 77640

Work phone: (409) 736-2551

FAX: (409) 736-0382

Wildlife Biologist.

Projects:

1. J.D. Murphree Wildlife Management Area, Jefferson County.
2. Lower Neches Wildlife Management Area, Orange County.

Mr. Russell E. Swafford

National Marine Fisheries Service
SEFC Galveston Laboratory, Habitat Conservation Division
4700 Ave U
Galveston, TX 77551-5997

Work phone: (409) 766-3699

FAX: (409) 766-3575

Fisheries Habitat Conservationist; wetlands, fisheries ecology.

Projects:

1. Wetlands mitigation/restoration; many projects such as a 5 acre plot along Swan Lake where a *Spartina alterniflora* marsh was restored by Amoco Oil in 1992, and the Pierce Marsh creation of a *Spartina alterniflora* - *S. patens* marsh by Sandifer Petroleum in 1992.
2. Seagrass mitigation/restoration; many projects including the Padre Isles Subdivision restoration of about 2 acres of seagrass beds in Upper Laguna Madre.
3. Beneficial uses of dredged material; monitoring of 250 acres at Atkinson Island and West Bay demonstration projects involving marsh creation and stabilization of shorelines using dredged material and *Spartina alterniflora*.

Mr. David L. Trimm

ENTRIX, Inc.
515 Post Oak Blvd., Suite 400
Houston, TX 77027

Work phone: (713) 960-9733

FAX: (713) 960-9461

Senior Staff Scientist/ Marine fisheries/ Ecology/
Wetlands ecology.

Dr. John W. Tunnell, Jr.

Texas A & M University - Corpus Christi
Center for Coastal Studies
6300 Ocean Drive
Corpus Christi, TX 78412

Work phone: (512) 994-2768

FAX: (512) 994-2770

Director of the Center for Coastal Studies; Research and teaching of coastal biology.

Projects:

1. Mitigation monitoring of a salt marsh creation project, Nueces River Delta, TX, 1989-present.
2. Environmental impact and recovery of the EXXON Pipeline oil spill and burn site (high marsh), Copano Bay, 1992-present.

Ms. Mary Ellen Vega

Texas Parks & Wildlife Department
Resource Protection Div., Habitat Assessment Branch
c/o Corpus Christi State Univ., Campus Box 317
6300 Ocean Drive
Corpus Christi, TX 78412

Work phone: (512) 993-4491

FAX: (512) 993-4597

Biologist; protect wildlife and fishery resources and habitats.

Projects:

1. Provide recommendations regarding avoiding, minimizing, and compensating for wetland impacts.
2. Monitor required wetland, marsh, and seagrass bed mitigation projects for compliance and success.
3. Coordinate with the SCS concerning use of *Spartina alterniflora* for shoreline stabilization in middle and south Texas bays.
4. Recommend areas for beneficial use of dredged material, including areas for marsh creation, e.g. the 13 acres of marsh created by the Mitchell Energy Corporation in Mesquite Bay, Texas, which increases feeding habitat for the endangered whooping crane.

Dr. James W. Webb, Jr.

Texas A&M University at Galveston
P.O. Box 1675
Galveston, TX 77553

Work phone: (409) 740-4542

FAX: (409) 740-5001

Professor; Research on *Spartina* marsh restoration and creation.

Projects:

1. An ecosystem comparison of transplanted and native salt marshes: the chronological development of habitat value for fishery species. NOAA Coastal Ocean Program, Estuarine Habitat Research Program. 1990-93.
2. Habitat availability and utilization by benthos and nekton in Hall's Lake and West Galveston Bay. NMFS/US Army COE program. 1991.
3. Salt marsh mitigation for Motco Superfund site.

Mr. Frederick T. Werner, Jr.

U.S. Fish and Wildlife Service
17629 El Camino Real, Suite 211
Houston, TX 77058

Work phone: (713) 286-8282

FAX: (713) 488-5882

Fish and wildlife biologist

Projects:

1. Seabrook Slough; federal project to mitigate impacted wetlands. *Spartina alterniflora* was planted to establish saltmarsh.
2. Saltmarsh restoration at a site along the north shore of Clear Lake; property owned by Harry Reed.

Mr. Richard M. Yuill

ENTRIX, Inc.

515 Post Oak Blvd., Suite 400

Houston, TX 77027

Work phone: (713) 960-9733

FAX: (713) 960-9461

Environmental Sciences Group Manager; Estuarine benthos, wetlands ecology

Projects:

Dr. Roger J. Zimmerman

National Marine Fisheries Service

SEFC Galveston Laboratory

4700 Ave. U

Galveston, TX 77551-5997

Work phone: (409) 766-3500

FAX: (409) 766-3508

Acting Lab Director, Galveston Lab; Research ecologist, habitats for fisheries organisms.

Projects:

1. Galveston Island State Park marsh evaluation for fisheries species.
2. Galveston Bay habitat evaluation for placement of dredged material from the Houston Ship Channel widening.
3. Feeding preferences of juvenile penaeid shrimp.

Virginia

Dr. Linda K. Blum

University of Virginia-Charlottesville

Department of Environmental Science

Clark Hall

Charlottesville, VA 22903

Work phone: (804) 924-0560

FAX: (804) 982-2137 more...

Research Assistant Professor. Microbial and estuarine ecology.

Projects:

1. A study of root dynamics and microbial processes involved with carbon and nitrogen changes in decay and production by short and tall forms of *Spartina alterniflora*.
2. A comparison of the contributions of *Spartina alterniflora* and phytoplankton to the dissolved organic carbon pool used by bacteria in a tidal creek.

Mr. Terry Getchell

U.S. Army Corps of Engineers

Civil Programs Branch

803 Front St.

Norfolk, VA 23510

Work phone: (804) 441-7617

Environmental scientist; wetlands and hardwood swamps.

Projects:

1. Various sites on the Elizabeth River had marsh creation/restoration actions; total area of about 10 acres.
2. Hampton Roads sites with marsh creation; total area of about 20 acres; from 1987-present.
3. Proposed project to convert a barrow pit into about 20 acres of *Spartina alterniflora* marsh and 10 acres of hardwood swamp; proposed for 1994-95.

Mrs. Robin V. Morgan

Williamsburg Environmental Group, Inc.

P.O. Box 3584

Williamsburg, VA 23187

Work phone: (804) 220-6869

FAX: (804) 229-4507

Ecologist; consulting.

Projects:

1. Restoration of a 0.5 acre marsh along the Elizabeth River in Tidewater, Virginia, by sprigging with *Spartina alterniflora*.
2. Several palustrine mitigation projects that involved design and creation; emergent and forested wetlands mainly in central Virginia.

Mr. A. Keith Taniguchi

U.S. Fish and Wildlife Service,

Habitat Conservation, Coastal & Wetland Resources

4401 N. Fairfax Drive, 400 ARLSQ

Arlington, VA 22203

Work phone: (703) 358-2201

FAX: (703) 358-2232

Information specialist, wetlands.

Washington

Dr. David A. Armstrong

University of Washington

School of Fisheries, WH-10

Seattle, WA 98195

Work phone: (206) 543-6132

FAX: (206) 685-3224

Professor; invertebrate ecology.

Projects:

1. Ecology of tidal flats in NW estuaries.
2. Use of intertidal oyster shell to enhance recruitment and survival of 0+ dungeness crab (*Cancer magister*) in Grays Harbor Estuary. This project has led to construction of over 8 hectares of shell habitat, and includes provisions to

limit movement of *Spartina* seed from Willapa Bay with shell transported from that estuary.

3. Shell, shrimp and eelgrass interactions in the Willapa estuary. The healthy shell/eelgrass tideflats lacks high densities of thalassinid shrimp, and may reduce proliferation of exotic *Spartina alterniflora* in estuaries of the Pacific Northwest.

Dr. Douglas A. Bulthuis

Padilla Bay National Estuarine Research Reserve
Washington State Department of Ecology
1043 Bay View - Edison Road
Mount Vernon, WA 98273

Work phone: (206) 428-1558

FAX: (206) 428-1491

Research Coordinator; seagrasses, estuarine biology.

Projects:

1. Assess the effectiveness of glyphosate on killing *Spartina alterniflora* in Padilla Bay, and the effects on adjacent and intermixed native salt marsh plants. June 1992.

Mr. Brett R. Dumbauld

Washington State Department of Fisheries
Marine Fish/Shellfish
P.O. Box 190
Ocean Park, WA 98640

Work phone: (206) 665-4166

FAX: (206) 665-6716

Fisheries Biologist; shellfish life history, estuarine ecology of Willapa Bay.

Projects:

1. Willapa Bay habitat inventory; aerial overflights and ground-truthing to assess the *Spartina* and other habitat acreage; create a GIS database. 1993-1994.
2. Effects of introduced *Spartina alterniflora* on native bivalve populations in Willapa Bay; 1992-1993.

Mr. Steve L. Foss

Washington State Department of Agriculture
Pesticide Management Division
P.O. Box 42589
Olympia, WA 98504-2589

Work phone: (206) 902-2049

FAX: (206) 902-2093

Pesticide Specialist; Agriculture, education.

Projects:

1. Education: *Spartina* - Threat to Washington's saltwater habitat.

Mr. James A. Hidy

U.S. Fish and Wildlife Service
Willapa National Wildlife Refuge & Satellites
HC 01 Box 910
Ilwaco, WA 98624-9707

Work phone: (206) 484-3482

FAX: (206) 484-3109

Refuge manager; *Spartina* control.

Projects:

1. Control of *Spartina alterniflora*; possible elimination of it from areas of Willapa Bay. Mechanical control, 1989.
2. Fish use of cordgrass, 1990.
3. Use of glyphosate on *Spartina*, efficacy, impacts to non-target organisms, fate.

Dr. Thomas F. Mumford

Washington Department of Natural Resources, Division
of Aquatic Lands
P.O. Box 47028
Olympia, WA 98504-7028

Work phone: (206) 902-1079

FAX: (206) 902-1786

Section Head, Environmental Information; algology and geographical information systems.

Projects:

1. Characterization of nearshore habitats in Puget Sound, WA. 1990. Mapping includes *Spartina* spp. marshes.
2. Programmatic EIS for nations emergent aquatic plants in Washington; 1991, for management of *Spartina* and loosestrife.
3. Inventory of *Spartina* spp. in Washington, 1991-92, using ground and aerial surveys.
4. Environmental impacts of different control methods for controlling the spread of *Spartina* spp. in Washington. 1991-92.

Ms. Sharon R. Riggs

Washington State Department of Ecology
Padilla Bay National Estuarine Research Reserve
1043 Bayview-Edison Road
Mount Vernon, WA 98273

Work phone: (206) 428-1558

FAX: (206) 428-1491

Program Coordinator; administration and estuarine ecological research.

Projects:

1. Distribution of *Spartina alterniflora* in Padilla Bay, Wash., 1991, Pad. Bay Ntl. Est. Res. Reserve Tech. Rpt. No. 3.
2. Net aerial primary productivity of *S. alterniflora* in Pad. Bay, Wash., 1992-1993. Used a non-destructive method to measure net aerial primary productivity over a year.

Ms. Kathleen A. Sayce

Shoalwater Botanical
P.O. Box 91
Nahcotta, WA 98637

Work phone: (206) 665-5292

FAX: (206) 665-5292

Botanist; salt marshes in estuaries.

Projects:

1. A study of *Spartina alterniflora* in Willapa Bay, WA. USFWS study 1987-88.
(more)

2. Survey of *S. alterniflora* in Willapa Bay includes annual seed set and vertical placement in the intertidal zone, 1988- .
3. Monitoring of diatoms in Willapa Bay, planktonic, benthic and epiphytic.

Mr. Charles A. Simenstad

University of Washington
Wetlands Ecosystem Team
Fisheries Research Institute, WH-10
Seattle, WA 98195

Work phone: (206) 543-7185

FAX: (206) 685-7471

Marine Biologist; Ecosystems, wetlands, fisheries biology.

Projects:

Studies of the invertebrates in Willapa Bay.

Mr. Curtis D. Tanner

U.S. Fish and Wildlife Service
3704 Griffin Ln. SE, Suite 102
Olympia, WA 98501-2192

Work phone: (206) 753-9440

FAX: (206) 753-9008

Wildlife biologist; wetlands.

Projects:

Coastal America intertidal habitat restoration projects in the Duwamish River estuary, Seattle, WA; (7/1/90). Projects were designed to 1) increase the amount and quality of resource base in the Duwamish River estuary; 2) investigate novel approaches to habitat restoration/enhancement in an urban environment; and 3) develop an inter-agency approach to non-compensatory habitat restoration.

Dr. Ronald M. Thom

Battelle Marine Sciences Laboratory
Marine & Wetland Resources Group
1529 W. Sequim Bay Road
Sequim, WA 98382

Work phone: (206) 681-3657

FAX: (206) 681-3681

Group leader; estuarine and marine plant ecology; wetlands, coastal systems; consultant.

Projects:

1. Sea level rise effects on coastal marshes. March 1991.
2. Effects of graveling beaches on nutrient flux rate and benthic primary productivity. March 1991.
3. Effects of oil on kelp. June 1992.
4. *Spartina alterniflora* ecology in Willapa Bay; project to start July, 1993.

Mr. Steve Wirth

Newstart Nursery
565 Juniper Beach Road
Camano Island, WA 98292

Work phone: (206) 629-3751

FAX: -

Manager/owner of a nursery; also is licensed to apply herbicides.

Projects:

1. Pilot study to use Rodeo to eradicate *Spartina townsendi* at Livingston Bay Park; 1992.

INDEX of PARTICIPANTS

Adams, L. 7
Adams, J. 34
Adamson 28
Albano 22
Alexander 4
Allen, H. 24
Allen, S. 26
Ammann 26
Anderson 5
Armstrong 39
Aurelia 4
Barber 7
Bartoldus 19
Bates 7
Beeman 7
Beever 7
Belaire 34
Bellis 32
Bennett 7
Bigford 22
Biggar 8
Bildstein 31
Bissett 8
Blair 8
Blama 19
Blood 32
Blum 39
Boatman 13
Bodin 13
Boesch 20
Bolotte 13
Bonner 31
Bontje 28
Bordelon 13
Bordes 13
Briggs 20
Brockmeyer 8
Broome 28
Broussard, B. 13
Broussard, D. 14
Brunori 20
Bryan 22
Bulthuis 40
Bunch 32
Bundy 8
Burdett 8
Burdick 26
Bybee 2
Cahoon 14
Cammen 20
Carangelo 34
Carlson, D. 8
Carlson, L. 22
Carr 22
Cash 8
Christian 32
Clairain 24
Clark 5
Clarke 25
Clements 34
Cordy 8
Craft 29
Crewz 9
Cuevas 25
Currin 29
Daehler 2
Davis 9
Day 14
De Bouno 26
Demgen 2
Derstine 14
Deshaies 22
Devlin 14
DiFiori 2
Dore 20
Dougherty 31
Dowhan 31
Dowling 1
Drylie 9
Dubinski 31
Dumbauld 40
Dunn 34
DuPuy 9
Earhart 20
Edstrom 27
Edwards 14
Eisenhardt 27
Eleuterius 25
Emmett 30
Erwin 20
Esquivel 34
Falcone 23
Fefer 19
Filip 20
Fine 14
Fink 2
Fonseca 29
Foss 40
Friedemann 9
Fultz 12
Furness 29
Gagliano 15
Gallagher 6
Gann 9
Gann-Matzen 9
Garbisch 20
Gavutis 23
Getchell 39
Gilio 9
Gilmore 10
Glass 35
Goldberg 35
Goldsmith 23
Golet 32
Goodger 21
Hamer 27
Hanes 10
Hardisky 31
Hargis 15
Harrington 35
Hartman 15
Hawes 15
Heath 1
Heller 4
Helm 26
Henningsen 10
Hensel 33
Herke 15
Hidy 40
Hobson 32
Hoeger 28
Hoese 15
Hoffman 2
Hook 15
Houston 28
Huggett 29
Johnson 15
Jones 16
Josselyn 3
Kahn 35
Karafel 10
Kearney 21
Kent 4
Kenworthy 29
Kepner 10
Kerfoot 23
Kilgore 20

Koepfler 33	Murphy 17	Slayback 5
Konikoff 16	Nailon 36	Smart 37
Kosier 10	Nichols 17	Smith, J. 3
Kraus 21	Nickerson 24	Smith, D. 37
Kruczynski 10	Nicolau 37	Sparrow 24
Landin 25	Nidecker 17	Spell 18
Lanier 13	Niering 4	Spicher 3
LaSalle 25	Northrup 37	Steinke 5
Latch 10	Novitzki 30	Stevenson 22
Lazarene 35	O'Neil 26	Steyer 19
Lelito 23	Ornes 33	Stout 1
Levin 30	Owen 17	Strong 3
Lewis 10	Patten 11	Sutherland 38
Lloyd-Reilly 35	Paul 21	Swafford 38
Logan 10	Perry 17	Talbot 19
Lomax 27	Pezeshki 17	Taniguchi 39
Lowe 25	Phillipp 6	Tanner 41
Luther 6	Popp 4	Taylor 24
Lyman 23	Potter 4	Teal 24
MacLean 21	Prann 27	Terwilliger 12
Mager 11	Proffitt 17	Thayer 30
Mantzaris 23	Purcell 33	Thom 41
Marcantel 16	Rafferty 18	Thornhill 1
Marcus 23	Ramsey 18	Thunhorst 30
Marshall 30	Reddoch 18	Tito 12
Martin 24	Redmond 11	Tomasko 12
Materne 16	Reed, B. 18	Trimm 38
Mathies 16	Reed, D. 18	Tullos 19
Matsil 28	Reeder 11	Tunnell 38
Matthews 35	Reily 11	Twilley 19
Mauney 36	Renner 13	Utermohle 22
Maurmeyer 6	Reubsamen 18	Van Dolah 33
McCandless 21	Riggs 40	Vega 38
McCartney 31	Roach 1	Vittor 2
McCorkle 6	Robinson 11	Waring 26
McHugh 24	Rozas 37	Warren 5
McNease 16	Rozsa 5	Webb 38
Medina 36	Sayce 40	Werner 38
Melton 11	Scheidt 11	Wesley 12
Menard 16	Schleifstein 18	White 12
Mendelssohn 16	Schroer 11	Whitman 12
Meyer 30	Schucet 31	Williams 3
Middleton 1	Seelinger 22	Wirth 41
Miller 27	Seidensticker 37	Witman 7
Minello 36	Seliskar 6	Yuill 39
Moore, R. 33	Seneca 30	Zaffke 12
Moore, S. 33	Shea 32	Zedler 3
Moore, D. 36	Shead 37	Zentner 3
Morgan 39	Shisler 28	Zewinski 32
Morris 33	Simenstad 41	Zimmerman 39
Mullins 36	Simon 18	Zyski 12
Mumford 40	Skrabal 6	

**Other Titles in the
Decision Analysis Series**

No. 1. Able, Kenneth W. and Susan C. Kaiser. 1994 Synthesis of Summer Flounder Habitat Parameters.