

Author Statement

Cathryn Wood: Formal analysis, Writing - Original Draft, Visualization

George H. Balazs: Conceptualization, Methodology, Resources, Data Curation, Writing - Review & Editing, Supervision

Marc Rice: Methodology, Resources, Supervision

Thierry M. Work: Methodology, Resources, Data Curation, Writing - Review & Editing, Supervision

T. Todd Jones: Resources, Data Curation, Writing - Review & Editing, Supervision

Eleanor Sterling: Conceptualization, Methodology, Resources, Data Curation, Writing - Review & Editing, Supervision, Funding acquisition



Tammy M. Summers: Methodology, Resources, Data Curation, Writing - Review & Editing, Supervision

John Brooker: Validation, Investigation, Data Curation, Writing - Review & Editing





Lauren Kurpita: Methodology, Resources, Data Curation, Supervision

Cheryl S. King: Methodology, Resources, Data Curation, Writing - Review & Editing

Jennifer M. Lynch: Conceptualization, Methodology, Validation, Formal analysis, Investigation, Resources, Data Curation, Writing - Original Draft, Writing - Review & Editing, Visualization, Supervision, Project administration



Green sea turtle
Hawksbill sea turtle



1 **Sea turtles across the North Pacific are exposed to perfluoroalkyl substances**

2
3 Cathryn Wood¹, George H. Balazs², Marc Rice³, Thierry M. Work⁴, T. Todd Jones⁵, Eleanor
4 Sterling⁶, Tammy M. Summers⁷, John Brooker⁸, Lauren Kurpita⁹, Cheryl S. King¹⁰, and Jennifer
5 M. Lynch^{1,11*}

6
7 1 Hawai‘i Pacific University, Center for Marine Debris Research, Waimānalo, HI, USA

8 2 Golden Honu Services of Oceania, Honolulu, HI, USA

9 3 Hawai‘i Preparatory Academy, Waimea, HI, USA

10 4 U.S. Geological Survey, National Wildlife Health Center, Honolulu Field Station, Honolulu,
11 HI, USA

12 5 NOAA Pacific Islands Fisheries Science Center, Honolulu, HI, USA

13 6 American Museum of Natural History, New York, NY

14 7 Rainbow Connection Research, Guam, USA

15 8 College of Charleston, Charleston, SC, USA

16 9 Hawai‘i Island Hawksbill Turtle Recovery Project, Hawai‘i National Park, HI, USA

17 10 Hawaiian Hawksbill Conservation, Kihei, HI, USA

18 11 National Institute of Standards and Technology, Chemical Sciences Division, Waimānalo, HI,

19 USA E-mail: Jennifer.lynch@nist.gov, *corresponding author, previously Jennifer M. Keller

20
21 **Abstract**

22 Perfluorinated alkyl substances (PFASs) are global, persistent, and toxic contaminants.
23 We assessed PFAS concentrations in green (*Chelonia mydas*) and hawksbill (*Eretmochelys*
24 *imbricata*) turtles from the North Pacific. Fifteen compounds were quantified via liquid

25 chromatography tandem mass spectrometry from 62 green turtle and 6 hawksbill plasma samples
26 from Hawai'i, Palmyra Atoll, and the Northern Marianas Islands. Plasma from 14 green turtles
27 severely afflicted with fibropapillomatosis, and eggs from 12 Hawaiian hawksbill nests from 7
28 females were analyzed. Perfluorooctane sulfonate (PFOS) predominated in green turtle plasma;
29 perfluorononanoic acid (PFNA) predominated in hawksbill tissues. Concentrations were greater
30 in hawksbill than green turtle plasma ($p < 0.05$), related to trophic differences. Green turtle plasma
31 PFOS concentrations related to human populations from highest to lowest: Hawai'i, Marianas,
32 Palmyra. Influence on fibropapillomatosis was not evident. PFASs were maternally transferred
33 to hawksbill eggs, with decreasing concentrations with distance from airports and with clutch
34 order from one female. A risk assessment of PFOS showed concern for immunosuppression in
35 Kailua green turtles and alarming concern for hawksbill developmental toxicity.
36 Perfluoroundecanoic (PFUnA) and perfluorotridecanoic (PFTriA) acid levels were correlated
37 with reduced emergence success ($p < 0.05$). Studies should further examine toxicity of PFASs on
38 sea turtle development.

39

40 **Capsule:**

41 Endangered or threatened sea turtle species at remote Pacific islands contain perfluoroalkyl
42 substances in blood and offload them to eggs, at concentrations known to be detrimental in birds.

43

44 **Keywords:** Perfluorinated contaminants, marine turtles, Pacific islands, reptile, maternal transfer

45

46 **Introduction**

47 Perfluoroalkyl substances (PFASs) are a chemical family used in many industrial and
48 commercial products (Lau et al. 2007). Their chemical makeup typically consists of a fluorinated
49 carbon backbone (4 to 14 in length) and a charged functional moiety, such as a carboxylate. The
50 carbon-fluorine bond is incredibly stable, which gives PFASs extreme persistence and both
51 hydrophobic and lipophobic properties (Buck et al. 2011). Since their invention in the 1930s and
52 commercialization in the 1950s, PFASs have been incorporated into every-day consumer
53 products including coatings, fabrics, grease-proof papers, soil repellents, and aqueous film-
54 forming foam fire suppressants (AFFFs) (Moody and Field 2000; Buck et al. 2011; Ahrens and
55 Bundschuh 2014; Interstate Technology Regulatory Council 2020). AFFFs used for firefighting
56 and training at military bases and airports are large point sources of PFAS contamination
57 (Schultz et al. 2004). Their nearly non-biodegradable properties, along with global (Yamashita et
58 al. 2008) render PFASs highly persistent pollutants in the environment worldwide (Lau et al.
59 2007) with detectable baseline concentrations in the most remote regions (Young et al. 2007).

60 PFASs are prominent contaminants in wildlife and human tissues, and can result in
61 toxicological effects, including immune, developmental, hepatic, and endocrine disruptions (Lau
62 et al. 2007; DeWitt 2015). In 2001, 3M Corporation, a major producer of perfluorooctane
63 sulfonate (PFOS), began phasing out production of this chemistry. In 2006, eight manufacturers
64 committed to reduce perfluorooctanoate (PFOA) production by 2015 through a US EPA
65 Stewardship program (US EPA 2018). Additional policies have focused on eliminating further
66 production of PFASs in Europe and Asia, but PFASs remain prominent in the environment
67 (McCarthy 2017).

68 Because sea turtles are rare and protected, understanding their exposure to toxic
69 contaminants is important for the recovery of these species (NMFS and US FWS 1998b, a).
70 Moreover, certain life stages of sea turtles can indicate the contaminant levels in their respective
71 foraging grounds, because they integrate those contaminants over several years (O'Connell et al.
72 2010; Keller 2013). The generalized long life history of green (*Chelonia mydas*) and hawksbill
73 (*Eretmochelys imbricata*) sea turtles includes an ontogenetic switch that separates two juvenile
74 phases (Bolten 2003). The youngest turtles forage in open ocean habitats, and transition to
75 foraging in nearshore benthic habitats as older juveniles where they grow into subadults and
76 adults. As adults, they migrate from foraging grounds to inter-nesting habitat where they do not
77 eat much. The green turtle is the lowest trophic level sea turtle species (Bjorndal 1997). They
78 transition from omnivorous pelagic juveniles (Parker et al. 2011) to herbivory in the neritic
79 stages (Arthur and Balazs 2008; Summers et al. 2017). Green turtles are highly migratory, across
80 entire ocean basins, but have strong site fidelity to foraging grounds within a life stage (Summers
81 et al. 2017; Shimada et al. 2020). Hawksbill sea turtles are less migratory (Gaos et al. 2020b) and
82 continue to feed at higher trophic levels after their ontogenetic switch to eating primarily sponges
83 in the neritic phase (Bjorndal 1997; Summers et al. 2017).

84 Six studies have reported PFAS concentrations in tissues of six sea turtle species (Keller
85 et al. 2005; O'Connell et al. 2010; Keller et al. 2012; Guerranti et al. 2013; Guerranti et al. 2014;
86 Pasanisi et al. 2016), but none from turtles across the North Pacific. Most analyzed plasma
87 (Keller et al. 2005; O'Connell et al. 2010; Keller et al. 2012), and none have analyzed eggs.
88 Differences in concentrations among five species provide evidence that PFASs biomagnify in the
89 turtles' distinct food webs and that the major route of exposure is diet (Keller et al. 2012). For

90 example, PFAS concentrations are greater in carnivorous loggerhead (*Caretta caretta*) sea turtles
91 than herbivorous green turtles from the same neritic habitat (Keller et al. 2012).

92 The toxicological effects of PFASs on sea turtles are still largely undetermined.
93 Immunosuppressive contaminants have long been a suspected contributing influence in a
94 debilitating, tumor-forming, viral disease, fibropapillomatosis (FP) (Herbst and Klein 1995;
95 Keller 2013), but to date, no convincing causal relationship between exposure to toxicants and
96 development of FP has been made. For instance, when levels of other types of persistent organic
97 pollutants (POPs) were measured in the plasma of many of the same turtles in the current study,
98 the levels did not relate to severity of FP in Hawaiian green turtles. PFASs were not included in
99 that analysis (Keller et al. 2014a). Furthermore, maternal offloading of POPs occurs during yolk
100 deposition (Keller 2013; Munoz and Vermeiren 2020), but transfer of PFASs to eggs and effects
101 on hatching success and embryonic development are unknown.

102 In this study, we measured concentrations of 15 PFASs in the plasma of neritic-phase
103 green and hawksbill sea turtles across three North Pacific regions: the Main Hawaiian Islands
104 (MHI), the Commonwealth of the Northern Marianas Islands (CNMI), and Palmyra Atoll. Eggs
105 from excavated hawksbill nests from MHI were also analyzed We hypothesized that (1) PFOS
106 will predominate like in most other studies, (2) higher trophic positioned, spongivorous
107 hawksbills will have greater plasma PFAS concentrations than herbivorous green turtles, (3)
108 PFAS concentrations in turtles will correlate to human population or proximity to military bases
109 or airports (and in the lack of local point sources, remote Palmyra Atoll will represent globally
110 diffuse contamination levels transported by air and ocean currents), (4) PFASs will be detected
111 in the sampled hawksbill eggs, revealing maternal offloading, (5) PFAS concentrations in eggs

112 will negatively correlate to nest success variables, and (6) eggs from successive clutches from
113 the same female within a nesting season will have stable PFAS concentrations.

114

115 **2. Materials and Methods**

116 *2.1 Study Sites*

117 Samples were collected on beaches and coastal waters from three Pacific Island regions:
118 the Main Hawaiian Islands (MHI), the Commonwealth of the Northern Marianas Islands
119 (CNMI), and Palmyra Atoll (Figure 1). The MHIs have roughly 1.4 million people, three
120 international airports, and 11 military bases (US Census Bureau 2019a). CNMI has about 57,000
121 people (United Nations Dept. of Economic and Social Affairs 2019), three international airports,
122 and no active military bases, but a historically large military presence from World War II.
123 Palmyra Atoll is a remote U.S. National Wildlife Refuge with less than 20 staff, no commercial
124 airport, and a decommissioned military base.

125 *2.2 Sample Collection*

126 Cryogenically archived sea turtle samples were selected from the NIST Biorepository's
127 Biological and Environmental Monitoring and Archival of Sea Turtle tissues project (BEMAST).
128 Sample collection and processing methods are detailed in Keller et al. (2014b). Sample metadata
129 are provided (Tables S1, S2). Plasma samples were selected to investigate spatial, species, and
130 health status differences and included 62 samples from 61 free-ranging green sea turtles from
131 across three study sites (Table 1, Figure 1): 39 from MHI (n=13 from three sites), 12 from CNMI
132 (n=6 from Saipan and Tinian), and 10 from Palmyra Atoll (n=2 from each segment of the refuge,
133 with one turtle sampled twice over two years). Plasma from 14 severely tumored, MHI-stranded
134 turtles were also analyzed. All MHI plasma samples came from the same turtles that were

135 analyzed previously for other POPs (Keller et al. 2014a), representing four groups of green
136 turtles with varying prevalence of FP: Kiholo Bay (0% FP), Kailua Bay (low to moderate FP 0%
137 to 5%), Kapoho Bay (higher FP 35%) and tumored-stranded (100% FP). All six hawksbill
138 plasma samples available in the BEMAST inventory from CNMI (n=4, two each from Saipan
139 and Tinian) and Palmyra (n=2) were included. All 12 hawksbill nests available in BEMAST
140 inventory at the time were included. They were from beaches on Maui, Hawai'i, and Kauai
141 (Figure 1) in 2012 and represent seven nesting females (Table S2). Unhatched eggs were
142 collected from each nest upon excavation after hatchling emergence. Egg contents from at most
143 three unhatched eggs per nest were pooled, homogenized, aliquoted, and cryogenically archived.
144 The number of eggs laid, days of incubation, hatching success, and emergence success were
145 calculated per nest according to Miller (1997) to inspect the fitness of each nest in relation to
146 PFAS concentrations. The percent of eggs laid that developed to at least stage 23 of
147 development, as defined by Miller et al. (2017) was also calculated (Table S2).

148 *2.3 PFAS Measurements*

149 Detailed methods are provided in Supplemental Information. Samples were measured
150 using methods modified from Keller et al. (2012). Concentrations of 15 PFASs were quantified
151 using the internal standard approach (Table S3). Field blanks were prepared with the same lot
152 number of blood collection supplies (Table S4). Samples were extracted via sonication with
153 acetonitrile and purified on 250 mg Envi-carb columns (Supelco, Bellefonte, Pennsylvania).
154 Methanol extracts (20 µL) were injected on an Agilent Zorbax Eclipse Plus C18 column using a
155 liquid chromatograph (Agilent 1100 HPLC, Palo Alto, CA) negative electrospray ionization
156 tandem mass spectrometer (API 4000, Applied Biosystems-MDS Sciex, Foster City, CA). The
157 mobile phase consisted of methanol and 20 mmol/L ammonium acetate in water. Two transitions

158 for PFOS were monitoring (499 → 99 and 499 → 80) to ensure bile acids did not interfere.
159 Concentrations are totals of linear and branched isomers. Reporting limits (RLs) were
160 determined according to O'Connell et al. (2010). The mass fractions of PFASs measured in NIST
161 SRM 1957 Organic Contaminants in Non-Fortified Human Serum and SRM 1947 Lake
162 Michigan Fish Tissue were within the uncertainty of the reference values (Table S5).

163 *2.4 Data Analyses*

164 R software package, NADA, was used whenever possible when samples were <RL as
165 suggested by Helsel (2005), otherwise JMP (SAS Institute Inc. Cary, NC) was used with values
166 <RL substituted with half the RL. Significance was determined as $p < 0.05$. Data was assessed for
167 assumptions of normality and equal variances. Non-parametric ANOVAs in NADA tested
168 differences among sites for green turtle plasma PFAS concentrations, followed by Steel Dwass
169 non-parametric multiple comparison tests using JMP. Wilcoxon t-tests in JMP compared
170 hawksbill plasma PFAS concentrations between Palmyra and CNMI, except PFUnA required
171 NADA's Wilcoxon t-test. Non-parametric ANOVAs in NADA tested differences among sites
172 and FP incidence (grouped as in Keller et al. (2014a)) for green turtle plasma PFOS
173 concentrations, followed by Steel Dwass non-parametric multiple comparison tests using JMP.
174 Species differences were assessed with CNMI data.

175 Amount of PFASs offloaded into each nest was estimated by multiplying the total
176 number of eggs laid by the average grams of each pooled egg contents from that nest, and by the
177 measured PFAS concentration (ng/g). The nanograms of PFASs offloaded were then compared
178 across clutches from one nesting season of a mother using linear regressions. PFAS
179 concentrations in the eggs were compared to nest success variables using Kendall's tau
180 correlations in NADA.

181 To assess toxicological risk, estimated margins of safety (EMOS) were calculated as
182 described in Keller et al. (2012), as the ratio of the average plasma, serum, or egg PFOS
183 concentration in laboratory-exposed animals in the lowest observed adverse effect level
184 (LOAEL) dose group to either the average or maximum sea turtle plasma or egg PFOS
185 concentration. EMOS below 100 were considered of concern (Faustman and Omenn 1996).

186

187 **3. Results and Discussion**

188 *3.1 Plasma PFAS Concentrations*

189 PFAS mass fractions (hereafter called concentrations) in ng/g wet mass for each turtle
190 plasma sample are provided (Table S6). Summary statistics for all possible groupings of Pacific
191 turtles are compared to turtles along the Eastern U.S. (Table S7).

192 *3.1.1 Green Turtle Plasma PFAS Concentrations*

193 PFOS comprised >96% of the Σ PFAS concentrations in green turtle plasma regardless of
194 capture location (Table 1; Figures 2 and S1), similar to previous reports for green, leatherback,
195 loggerhead, and Kemp's ridley turtles from the Eastern U.S. (O'Connell et al. 2010; Keller et al.
196 2012). PFOS concentrations in green turtle plasma were significantly different among the Pacific
197 Island regions and increased with human population (Figure 3a). This trend continues when
198 including previously published PFOS concentrations in green turtles from Core Sound, North
199 Carolina (mean=2.41 ng/g; (Keller et al. 2012) Figure 3a), a watershed with \approx 6,390,000 people
200 (O'Connell et al. 2010). The MHI with 1,400,000 people had the next highest mean PFOS
201 concentration at 1.14 ng/g and significantly greater than CNMI (57,000 people) at 0.524 ng/g
202 and Palmyra (20 people) at 0.155 ng/g. This relationship between PFAS plasma concentration
203 and human population corroborates previous relationships for sea turtle contaminant exposure

204 (O'Connell et al. 2010; Alava et al. 2011), is expected for these man-made chemicals, and
205 suggests that local or regional sources add to the diffuse, globally distributed PFAS levels in
206 nearshore marine organisms.

207 Green turtles from the MHIs were categorized into four groups based on their capture
208 location and incidence and severity of FP using the same grouping as Keller et al. (2014a). These
209 were No FP (Kiholo), Low FP (Kailua), High FP (Kapoho), and FP stranded. The FP stranded
210 turtles were alive, severely afflicted with FP tumors, and exceedingly emaciated because of the
211 disease. Significant differences in PFOS plasma concentrations were noted among the groups
212 ($p=5e-9$), but not in a dose-dependent manner if PFOS was contributing to FP (Figure S2). The
213 low FP Kailua group (mean=2.45 ng/g), from the most urbanized area sampled, had significantly
214 higher concentrations than the other three groups and was relatively similar to those in Core
215 Sound, NC (2.41 ng/g) (Table 1, Table S7) (Keller et al. 2012). This is likely due to O'ahu being
216 a heavily populated island (974,563 people), and hosting numerous military bases, airports, and
217 other point-sources (US Census Bureau 2019a). The Kailua sampling site is adjacent to Marine
218 Corps Base Hawai'i, with an active airstrip and firefighting training sites. Additionally, the
219 Kawainui watershed empties into Kailua Bay at this sampling site and contains 19.1% urban
220 development, while Kapoho and Kiholo watersheds had 7.8% and 0% urban development,
221 respectively (Parham et al. 2008a, b, c). Given that the MHIs are one of the most isolated
222 archipelagos in the world, approximately 4,000 km away from the nearest continent (National
223 Academy of Sciences 2004), air and ocean circulation transport a baseline level of globally
224 diffuse PFASs to Hawaiian waters. The differences among turtle groups here indicate that local
225 sources compound global sources of PFASs in Hawai'i.

226 The lack of a dose-dependent relationship between PFASs and FP is similar to results for
227 other POPs that were measured in these same turtles (Keller et al. 2014a) in which Kailua turtles
228 with low FP incidence had higher concentrations than the other two locations. What differs
229 between the two studies is the relative difference in the FP stranded turtles. For protein-
230 associating PFOS, the FP stranded turtles had levels that were close to average among the MHI
231 turtles (Table 1). For lipophilic POPs, the FP stranded turtles had elevated levels compared to the
232 other MHI turtles (Keller et al. 2014a). This can be explained by the different distribution of
233 these compounds during weight loss and lipid mobilization. The lipophilic POPs flooded into the
234 blood of the emaciated FP stranded group upon weight loss. With no lipid to associate within the
235 thin turtles, those POPs continued to circulate in the blood at high levels. Conversely, PFOS
236 preferentially associates with proteins in the blood and liver. Upon weight loss, protein
237 concentrations in these tissues do not change as much as lipid levels. Thus, PFOS remains
238 associated with proteins, even in emaciated animals, resulting in less mobilization of PFOS into
239 the blood upon weight loss.

240 Green and hawksbill turtles inhabit the coral reefs of CNMI (Summers et al. 2017), a
241 U.S. Commonwealth comprised of 14 islands. They face several threats, including illegal harvest
242 and marine debris entanglement (Summers et al. 2018). Sampling sites were on the archipelago's
243 two most populated islands, Saipan with 48,200 residents and Tinian with 3,056 residents
244 (Department of Commerce Central Statistics Division 2017). Green turtles from Saipan had
245 significantly greater plasma PFOS and Σ PFAS concentrations than from Tinian ($p=0.02$); (Table
246 1), corroborating other human population trends.

247 Palmyra Atoll is extremely remote in the Central Pacific at 5.9° N 162.1° W, with a
248 history of extensive human disturbance during World War II in the 1940s when construction of a

249 U.S. naval base necessitated the remodeling of the small islets (Collen et al. 2009). Construction
250 of >100 buildings and two airstrips occurred between 1940 and 1944. Dredging in the West
251 Lagoon, as well as causeway construction across multiple reef flats, roughly doubled the land
252 area and resulted in major hydrodynamic changes in the lagoon (Collen et al. 2009). By 1945,
253 most military activity had ceased and personnel were evacuated, which prompted many natural
254 changes in the atoll's topography (Collen et al. 2009). Since 2001, the atoll has been protected as
255 a U.S. Fish and Wildlife Refuge.

256 No uses or disposal of PFASs are known on Palmyra Atoll. AFFFs were invented 15
257 years after the U.S. Navy abandoned the atoll (Moody and Field 2000). A cursory spatial
258 comparison using green turtle plasma PFAS concentrations showed no evidence of a significant
259 point source on the atoll (see Supplemental Information). The comparison was confounded by
260 life stage with the greatest PFAS burden in a young green turtle that recently recruited from the
261 pelagic carnivorous stage into the nearshore herbivorous stage. New recruits favor the eastern
262 region of Palmyra (Sterling et al. 2013), and may carry a greater PFAS burden to Palmyra, which
263 dilutes with growth thereafter (Keller 2013). The current study suggests that remote Palmyra
264 Atoll is a control reference site with little to no local point sources of PFASs, reflecting baseline
265 globally diffuse concentrations. The first study to measure chemical contaminants in Palmyra sea
266 turtles found elevated aluminum and iron McFadden et al. (2014).

267

268 *3.1.2 Hawksbill Plasma PFAS Concentrations*

269 Perfluorononanoic acid (PFNA) predominated in hawksbill plasma from CNMI (n=4)
270 and Palmyra (n=2), with means of 1.08 ng/g and 3.40 ng/g, respectively (Figures 2 and 3b). The
271 same PFAS profile was seen in hawksbills from Juno Beach, FL (a typo in in Keller et al. (2012)
272 should read 17.3 ng/g for mean PFNA). It is interesting that hawksbill turtles from such distant

273 regions have similar PFAS profiles (PFNA predominating), while green turtles from these
274 disparate regions show PFOS dominating. The reasons for the species differences in PFAS
275 profiles are difficult to explain, especially since marine mammals stranded in Hawai'i show still
276 another profile (perfluoroundecanoic acid (PFUnA) dominated (Kurtz et al. 2019)). Differences
277 may be explained by elimination mechanisms, prey selection, or migratory pathways.

278 CNMI and Palmyra hawksbill plasma had lesser PFAS concentrations than Juno Beach
279 hawksbills, in line with the drastic differences in human population (Figure 3b). This finding is
280 congruent with the green turtle spatial differences described above and those in loggerhead
281 turtles from the U.S. East coast (O'Connell et al. 2010). Together these findings support the idea
282 that PFAS levels, biomagnifying in marine organisms inhabiting developed nearshore regions,
283 are influenced more from local or regional land-based sources than diffuse sources from global
284 air and ocean currents. Juno Beach, FL, is within the St. Lucie-Loxahatchee Watershed which
285 has 6,400,000 human residents (South Florida Water Management District 2009; US Census
286 Bureau 2019b, c). Surprisingly, Palmyra hawksbills (only 20 human residents) had a greater
287 mean concentration of PFNA, PFUnA, and \sum PFASs than those in CNMI, but low sample size
288 prevented statistical analyses. The reasons for the observed spatial differences could include prey
289 selection, migratory pathways, or sources of contamination.

290

291 *3.1.3 Species Comparisons of Plasma PFAS Concentrations*

292 The two hawksbill turtles from Palmyra were both captured in the eastern region of the
293 atoll, and had PFAS concentrations one order of magnitude greater than the Palmyra green turtle
294 mean (Tables 1 and S6). In CNMI, \sum PFAS concentrations in hawksbill plasma was significantly
295 greater than green turtles (Figure S3). These results corroborate previous results showing Eastern
296 U.S. hawksbills had greater plasma PFOS levels than three other species, including green turtles

297 (Keller et al. 2012). Similarly, in Australia and Japan, hawksbills had greater concentrations of
298 POPs (PFASs were not tested) than green turtles (Hermanussen et al. 2008; Malarvannan et al.
299 2011). The trophic position of the two species may explain greater biomagnification of PFASs in
300 hawksbills. Omnivorous hawksbills primarily prey on sponges, but also forage on other
301 invertebrates and marine vegetation, placing them higher on the food web than the herbivorous
302 stage of green turtles studied (Bjorndal 1997).

303

304 *3.2 Egg PFAS Concentrations*

305 Eleven PFAS compounds were detected in at least one hawksbill nest (Tables 1 and S2).
306 Other POPs are known to be deposited into eggs from mother sea turtles during egg production,
307 rather than crossing the eggshell in the nest environment (Keller 2013; Munoz and Vermeiren
308 2020). Therefore, POP concentrations in eggs represent the mother's exposure from her foraging
309 grounds (Alava et al. 2011). It is expected that PFASs deposit into eggs in a similar fashion as
310 maternal offloading has been documented in other egg-laying species (Wilson et al. 2020). The
311 presence of PFASs in hawksbill eggs reveals, for the first time, offloading from females to their
312 eggs and, to our knowledge, is the first report of PFAS offloading in any reptile species. These
313 data fill an important gap in understanding exposure of adult female hawksbills and their
314 developing embryos to PFASs.

315 \sum PFAS concentrations were greatest in the nest laid in Wailua, Kauai, and lowest in the
316 four nests from three mothers in Pohue, Hawai'i (Figure S4). Nest PFAS concentrations were
317 negatively, significantly ($p=0.030$) correlated with the distance over water from the nests to the
318 nearest international airport (Figure S4). This preliminary finding may be explained by airports
319 and military bases being some of the largest point sources of PFASs due to firefighting training

320 (Schultz et al. 2004; Houtz et al. 2016). The relationship was driven by the Wailua nest, and a
321 larger sample size should be analyzed with future satellite tracks of females.

322 PFAS profiles in eggs when averaged across the seven mothers were dominated by PFOS
323 (28.5% of \sum PFASs), PFUnA (24.7%), and PFNA (23.8%) (Figures 2 and S1). More
324 interestingly, the eggs from different mothers displayed drastic differences in PFAS profiles
325 (Table S2, Figure S5). The nest laid in Wailua, Kauai, had the highest predominance of PFOS
326 (88%) and PFTA (3.5%) and the lowest PFNA contribution (0.9%), a profile that reflects older,
327 phased-out formulations of AFFFs (Place and Field 2012). The four nests laid by three females
328 in Pohue, Hawai'i had an intermediate profile with PFUnA dominating, followed by either PFOS
329 (16% to 31%) or PFDA. The five nests laid by one female (ID 19591-04) on Maui were shifted
330 towards PFNA > PFOS (22.6%) > PFUnA > PFDA. Finally, the two nests from likely different
331 females on Apua and Kamehame beaches on Hawai'i had a profile most different from the Kauai
332 nest, dominated by PFNA > PFUnA > PFDA, with PFOS comprising only 6%. The extreme
333 spectrum of PFOS contributions, from 88.3% in Kauai to 6% in Hawai'i, suggests that adult
334 female Hawaiian hawksbill turtles forage in distinct areas that have different exposure profiles.
335 This suggestion is partially explained by understanding that POP concentrations in sea turtle
336 eggs originate from the mother's diet from her foraging grounds (Alava et al. 2011; Keller 2013).
337 Therefore, the interpretation of the extreme spectrum of PFOS contribution must be placed in
338 context of hawksbill migration and foraging selection, as described next.

339 Though hawksbill turtles are capable of long-distance migrations (>2,000 km) (Vaughan
340 and Spring 1980), both foraging juveniles and nesting adults exhibit natal philopatry with
341 relatively small foraging ranges (Gaos et al. 2017; Wood et al. 2017; Gaos et al. 2018; Gaos et al.
342 2020b). Hawaiian hawksbills have recently been recognized as a genetically distinct

343 management unit, with little connectivity to other populations (Gaos et al. 2020a). Satellite tracks
344 of nine Hawaiian hawksbills show that their nesting and foraging grounds are close, on the same
345 or neighboring island (Parker et al. 2009). One Maui nester (ID 19591-04) migrated to O‘ahu to
346 forage in 2004 (Parker et al. 2009). Four years later, she nested again in the same region on Maui
347 and her post-nesting track (King et al. unpublished data) showed the same path to the same
348 O‘ahu foraging site (Figure S6). Four more years later, she laid five nests on Maui that were
349 sampled in the current study. This strong site fidelity to foraging grounds, relatively near the
350 nesting beaches, with potentially different local point sources across the Main Hawaiian Islands
351 may explain the extreme differences in PFAS profiles observed between the nests laid farthest
352 from each other: Kauai vs. Kamehame. Future studies could test this theory by measuring PFAS
353 concentrations and profiles in marine environmental samples, such as sediment or prey in known
354 hawksbill foraging grounds.

355 Of the 12 nests sampled in 2012, five were laid during the same nesting season (22 days
356 to 25 days apart) by Turtle 19591-04 in Makena, Maui. Two other nests were laid by another
357 female (ID 112) 43 days apart at Pohue, Hawai‘i; it is likely this turtle laid at least one more nest
358 between these dates but it was not sampled. Two additional Hawai‘i Island nests came from
359 turtles with unique IDs, and three nests were laid by unknown females (Table S2). The amounts
360 of PFASs in nests of Turtle 19591-04 significantly declined with clutch order (Figure 4),
361 indicating maternally offloaded contaminant concentrations may be a function of time within a
362 nesting season. However, a decline in transferred PFAS amounts was not apparent in the two
363 nests from Turtle 112.

364 When sea turtles arrive at their nesting site, they are equipped with all lipid-rich follicles
365 ready to become the yolk of all eggs to be laid that nesting season (Miller 1997). The follicles for

366 a single nest transit the oviduct where they are fertilized and surrounded by protein-rich albumen
367 (Miller 1997). After laying this clutch, hawksbills prepare the next clutch which is laid 14 to 25
368 days later. Because PFASs associate with serum albumin and fatty acid binding proteins (Ahrens
369 and Bundschuh 2014) rather than lipids, they should deposit more in egg albumen than follicles.
370 Since the albumen is deposited just before each nest is laid, and because in general mother turtles
371 fast during nesting (Miller 1997; Hays et al. 2002; Guirlet et al. 2008; Guirlet et al. 2010), the
372 mother's body could have less PFASs to transfer through albumen into successive clutches.
373 Theoretically, females would offload a greater portion of her body burden of PFASs into the first
374 clutch. Turtle 19591-04 offloaded a total of 367 µg of PFASs into these five clutches in one year,
375 with approximately 25% of that into the first clutch and 12% in the fifth clutch (Table S8).
376 Previously, a decrease in ΣPCBs, ΣHCHs and ΣDDTs yolk concentrations from successive
377 leatherback clutches suggested that reproductive lipid investment into eggs decreases as the
378 maternal lipid stores decrease (Guirlet et al. 2010). The current findings suggest that PFASs are
379 offloaded through albumen, and that Turtle 19591-04 was fasting during this nesting season
380 while Turtle ID 112 may have been foraging. These interpretations are supported by changes in
381 chicken egg PFAS concentrations during and after exposure of hens to PFASs in drinking water
382 (Wilson et al. 2020).

383 Nest success variables were examined for relationships with PFAS concentrations (n=11
384 nests). Only two significant correlations were observed ($p < 0.05$). Emergence success was
385 negatively correlated with concentrations of two contaminants: PFUnA and perfluorotridecanoic
386 acid (PFTriA) (Figure 5). Few studies exist on developmental effects of PFASs, and no
387 toxicology studies are available for reptiles. In chickens, hatching success was significantly
388 reduced to 61.4 % by a 100 ng/g injection of PFOS into eggs compared to 85.7 % in controls

389 (Molina et al. 2006). In tree swallow (*Tachycineta bicolor*) nests, hatching success was
390 significantly, negatively correlated with PFOS concentrations, and complete nest failure was
391 observed in three nests with concentrations at or above 150 ng/g (Custer et al. 2012). While we
392 saw no negative correlation between egg burdens of PFOS and hatching success, PFASs may
393 have more insidious reproductive effects for hawksbill turtles. These novel results indicate a
394 potential consequence of PFAS maternal offloading to embryonic sea turtles, and are supported
395 by our risk assessment. Future studies should address the developmental effects of PFASs in
396 turtles. It is possible that the correlations in this study between emergence success and PFASs are
397 confounded by the decline in PFASs in successive clutches, but to the authors' knowledge, no
398 studies have examined whether emergence success increases in successive clutches of sea turtles.

399 *3.3 Risk Assessment*

400 Using surrogate species is the only available option for sea turtle toxicology risk
401 assessments, but because reptiles can be more sensitive than other taxa (Weir et al. 2010) a wider
402 margin of safety (<100) should be used. Keller et al. (2012) estimated margins of safety for
403 toxicological effects of PFOS based on plasma concentrations in five species of sea turtles along
404 the Eastern U.S. All five species had margins of safety <100, which indicate a risk of at least
405 immunosuppressive effects. Using the same method for PFOS plasma concentrations, average
406 green turtles from Kailua Bay and the maximum green and hawksbill turtles from CNMI were at
407 risk of immunosuppression (Table 2). Likewise for eggs, the average hawksbill nest had a
408 margin of safety of only six, indicating heightened risk of reduced hatching success (Table 2).
409 More concerningly, the maximum nest from Kauai was nearly equal to the PFOS concentrations
410 (no margin of safety) that cause reduced hatching success in chickens and tree swallows (Molina
411 et al. 2006; Custer et al. 2012). This represents the first PFAS risk assessment for embryonic

412 stage sea turtles and suggests that sea turtles inhabiting regions close to military bases and
413 airports, even on remote islands, could be at risk of PFAS toxicity.

414 **4. Conclusion**

415 Reptiles are significantly under-studied in toxicology, particularly for PFASs (Reiner and
416 Place 2015). This is the first report of plasma PFAS concentrations in Pacific sea turtles, plus the
417 first to document maternal offloading of PFASs into eggs of any sea turtle species. The results
418 reveal contamination patterns similar to those documented along the Eastern U.S., with PFOS
419 predominating in green turtles, hawksbills accumulating greater levels than green turtles, and
420 PFAS concentrations being related to human population and specific land uses. The PFAS
421 concentrations in Pacific turtles are generally less than those along the Eastern U.S., which can
422 be attributed to the remoteness and smaller human populations of the islands studied. Across the
423 study sites, islands with greater population densities and closer proximity to military bases and
424 airports rendered greater PFAS concentrations in turtles. Incidence and severity of FP did not
425 relate to PFAS concentrations, so the search continues for environmental stressors that may
426 contribute to this viral disease. No prior study has analyzed sea turtle tissues concurrently with
427 prey items for PFAS levels; future studies on this would improve our understanding of trophic
428 transfer. The PFAS egg concentrations are novel for reptilian species and show maternal
429 offloading is strongest in the first clutch of a season and egg concentrations were highest in nests
430 laid nearest international airports. Two contaminants (PFUnA and PFTriA) were related to
431 reduced emergence success of hatchlings, which aligns with the risk assessment showing
432 hawksbill egg PFOS concentrations are concerningly near concentrations causing developmental
433 toxicity in birds. Future studies should address the toxicological impacts that PFASs may have
434 on sea turtle development.

435

436 Acknowledgements

437 All activities performed to collect sea turtle samples were permitted by National Oceanic and
438 Atmospheric Administration (NOAA) National Marine Fisheries Service (NMFS) permits: 1581,
439 1581-01, 15685, 10027-05, 15661-01, and 17022; U.S. Fish and Wildlife Service permits: TE-
440 70288A, 12533-12008, and 12533-13001; and State of Hawai'i permit 2013-32. Many thanks to
441 dedicated employees, students, and volunteers that helped collect samples from turtles and nests;
442 to Denise Parker for tracking information; and to CNMI Department of Land and Natural
443 Resources for sharing data. The Hawaiian turtle sampling was approved by the NOAA Pacific
444 Islands Fisheries Science Center's Institutional Animal Care and Use Committee. The Palmyra
445 turtle sampling was approved by the American Museum of Natural History's Institutional
446 Animal Care and Use Committee and funded by NOAA NMFS grants NA07NMF4540185,
447 NA10NMF4540299. We thank the Palmyra Atoll Research Consortium, The Nature
448 Conservancy, and USFWS for facilitating this research. This is Palmyra Atoll Research
449 Consortium publication number PARC-158. The statements, findings, conclusions, and
450 recommendations are those of the authors and do not necessarily reflect the views of NOAA or
451 the US Department of Commerce. Any use of trade, firm, or product names is for descriptive
452 purposes only and does not imply endorsement by the U.S. Government.

453

454 **References**

- 455
- 456
- 457 Ahrens, L. and Bundschuh, M. (2014). "Fate and Effects of Poly- and Perfluoroalkyl Substances
458 in the Aquatic Environment: A Review." Environmental Toxicology and Chemistry **33**(9):
459 1921-1929.
- 460 Alava, J.J., Keller, J.M., Wyneken, J.M., Crowder, L.B., Scott, G. and Kucklick, J.R. (2011).
461 "Geographical Variation of Persistent Organic Pollutants in Eggs of Threatened
462 Loggerhead Sea Turtles (*Caretta caretta*) from Southeastern United States."
463 Environmental Toxicology and Chemistry **30**(7): 1677-1688.
- 464 Arthur, K.E. and Balazs, G.H. (2008). "A comparison of immature green turtle (*Chelonia mydas*)
465 diets among seven sites in the main Hawaiian Islands." Pacific Science **62**(4): 205-217.
- 466 Bjorndal, K.A. (1997). Foraging Ecology and Nutrition of Sea Turtles. The Biology of Sea
467 Turtles. Lutz, P.L. and Musick, J.A., CRC Press: 199-231.
- 468 Bolten, A.B. (2003). Variation in sea turtle life history patterns: Neritic vs. oceanic
469 developmental stages. The Biology of Sea Turtles Volume II. Lutz, P.L., Musick, J.A. and
470 Wyneken, J. Boca Raton, FL, USA, CRC Press: 243-257.
- 471 Buck, R.C., Franklin, J., Berger, U., Conder, J.M., Cousins, I.T., Voogt, P.D., Jensen, A.A.,
472 Kannan, K., Mabury, S.A. and van Leeuwen, S.P.J. (2011). "Perfluoroalkyl and
473 Polyfluoroalkyl Substances in the Environment: Terminology, Classification, and Origins."
474 Integrated Environmental Assessment and Management **7**(4): 513-541.
- 475 Collen, J.D., Garton, D.W. and Gardner, J.P.A. (2009). "Shoreline Changes and Sediment
476 Redistribution at Palmyra Atoll (Equatorial Pacific Ocean): 1874 – Present." Journal of
477 Coastal Research **25**(3): 711-722.
- 478 Custer, C.M., Custer, T., W., Schoenfuss, H., L., Poganski, B.H. and Solem, L. (2012).
479 "Exposure and effects of perfluoroalkyl compounds on tree swallows nesting at Lake
480 Johanna in east central Minnesota, USA." Reproductive Toxicology **33**(4): 556-562.
- 481 Department of Commerce Central Statistics Division (2017). 2016 Commonwealth of the
482 Northern Mariana Islands Household Income and Expenditures Survey (HIES) Report:
483 269. <https://commerce.gov.mp/sites/default/files/2016-hies-final-report.pdf>
- 484 DeWitt, J.C., Ed. (2015). Toxicological Effects of Perfluoroalkyl and Polyfluoroalkyl
485 Substances. Molecular and Integrative Toxicology. Springer International Publishing,
486 Springer, Cham.
- 487 Faustman, E.M. and Omenn, G.S. (1996). Risk Assessment. Casarett and Doull's Toxicology the
488 Basic Science of Poisons Klaassen, C.D. New York, NY, McGraw-Hill.
- 489 Gaos, A.R., LaCasella, E.L., Kurpita, L., Balazs, G., Hargrove, S., King, C., Bernard, H., Jones,
490 T.T. and Dutton, P.H. (2020a). "Hawaiian hawksbills: a distinct and isolated nesting
491 colony in the Central North Pacific Ocean revealed by mitochondrial DNA." Conservation
492 Genetics **21**: 771-783.
- 493 Gaos, A.R., Lewison, R.L., Jensen, M.P., Liles, M.J., Henriquez, A., Chavarria, S., Pacheco,
494 C.M., Valle, M., Melero, D., Gadea, V., Altamirano, E., Torres, P., Vallejo, F., Miranda,
495 C., LeMarie, C., Lucero, J., Ocegüera, K., Chacon, D., Fonseca, L., Abrego, M., Seminoff,
496 J.A., Flores, E.E., Llama, I., Donadie, R., Pena, B., Muñoz, J.P., Ruales, D.A., Chaves,
497 J.A., Otterstrom, S., Zavala, A., Hart, C.E., Brittain, R., Alfaro-Shigueto, J., Mangel, J.,
498 Yanez, I.L. and Dutton, P.H. (2017). "Natal foraging philopatry in eastern Pacific
499 hawksbill turtles." Royal Society Open Science **4**: 1-8.

500 Gaos, A.R., Lewison, R.L., Jensen, M.P., Liles, M.J., Henriquez, A., Chavarria, S., Pacheco,
501 C.M., Valle, M., Melero, D., Gadea, V., Altamirano, E., Torres, P., Vallejo, F., Miranda,
502 C., Lemarie, C., Lucero, J., Ocegüera, K., Chácon, D., Fonseca, L., Abrego, M., Seminoff,
503 J.A., Flores, E.E., Llamas, I., Donadi, R., Peña, B., Muñoz, J.P., Ruales, D.A., Chaves,
504 J.A., Otterstrom, S., Zavala, A., Hart, C.E., Brittain, R., Alfaro-shigueto, J., Mangel, J.,
505 Yañez, I.L. and Dutton, P.H. (2018). "Rookery contributions, movements and
506 conservation needs of hawksbill turtles at foraging grounds in the eastern Pacific Ocean."
507 Marine Ecology Progress Series **586**: 203-216.

508 Gaos, A.R., Martin, S.L. and Jones, T.T. (2020b). Sea Turtle Tagging in the Mariana Islands
509 Training and Testing (MITT) Study Area Programmatic Report. PIFSC Data Report, DR-
510 20-003. <https://doi.org/10.25923/qq2e-e198>: 1-48.

511 Guerranti, C., Ancora, S., Bianchi, N., Perra, G., Fanello, E.L., Corsolini, S., Fossi, M.C. and
512 Focardi, S.E. (2013). "Perfluorinated compounds in blood of *Caretta caretta* from the
513 Mediterranean Sea." Marine Pollution Bulletin **73**(1): 98-101.

514 Guerranti, C., Bains, M., Casini, S., Focardi, S.E., Giannetti, M., Mancusi, C., Marsili, L., Perra,
515 G. and Fossi, M.C. (2014). "Pilot study on levels of chemical contaminants and porphyrins
516 in *Caretta caretta* from the Mediterranean Sea." Marine Environmental Research **100**: 33-
517 37.

518 Guirlet, E., Das, K. and Girondot, M. (2008). "Maternal transfer of trace elements in leatherback
519 turtles (*Dermochelys coriacea*) of French Guiana." Aquatic Toxicology **88**: 267-276.

520 Guirlet, E., Das, K., Thomé, J.-p. and Girondot, M. (2010). "Chemosphere Maternal transfer of
521 chlorinated contaminants in the leatherback turtles, *Dermochelys coriacea*, nesting in
522 French Guiana." Chemosphere **79**(7): 720-726.

523 Hays, G.C., Broderick, A.C., Glen, F. and Godley, B.J. (2002). "Change in body mass associated
524 with long-term fasting in a marine reptile: the case of green turtles (*Chelonia mydas*) at
525 Ascension Island." Canadian Journal of Zoology **80**: 1299-1302.

526 Helsel, D.R. (2005). Nondetects and data analysis. Statistics for censored environmental data.
527 Hoboken, Wiley-Interscience.

528 Herbst, L.H. and Klein, P.A. (1995). "Green turtle fibropapillomatosis: Challenges to assessing
529 the role of environmental cofactors." Environmental Health Perspectives **103**(SUPPL. 4):
530 27-30.

531 Hermanussen, S., Matthews, V., Pöpke, O., Limpus, C.J. and Gaus, C. (2008). "Flame retardants
532 (PBDEs) in marine turtles, dugongs and seafood from Queensland, Australia." Marine
533 Pollution Bulletin **57**(6-12): 409-418.

534 Houtz, E.F., Sutton, R., Park, J.S. and Sedlak, M. (2016). "Poly- and perfluoroalkyl substances in
535 wastewater: Significance of unknown precursors, manufacturing shifts, and likely AFFF
536 impacts." Water Research **95**: 142-149.

537 Interstate Technology Regulatory Council. (2020). "PFAS Technical and Regulatory Guidance
538 Document and Fact Sheets PFAS-1." Retrieved October 22, 2020, from [https://pfas-](https://pfas-1.itrcweb.org/)
539 [1.itrcweb.org/](https://pfas-1.itrcweb.org/).

540 Keller, J.M. (2013). Exposure to and Effects of Persistent Organic Pollutants. The Biology of
541 Sea Turtles Volume III. Wyneken, J., Lohmann, K.J. and Musick, J.A. Boca Raton, FL,
542 CRC Press Taylor and Francis Group. **3**: 285-328.

543 Keller, J.M., Balazs, G.H., Nilsen, F., Rice, M., Work, T.M. and Jensen, B.A. (2014a).
544 "Investigating the Potential Role of Persistent Organic Pollutants in Hawaiian Green Sea
545 Turtle Fibropapillomatosis." Environmental Science & Technology **48**: 7807-7816.

546 Keller, J.M., Kurunthachalam, K., Taniyasu, S., Yamashita, N., Day, R.D., Arendt, M.D., Segars,
547 A.L. and Kucklick, J.R. (2005). "Perfluorinated Compounds in the Plasma of Loggerhead
548 and Kemp's Ridley Sea Turtles from the Southeastern Coast of the United." Environmental
549 Science & Technology **39**(23): 9101-9108.

550 Keller, J.M., Ngai, L., Braun McNeill, J., Wood, L.D., Stewart, K.R., O'Connell, S.G. and
551 Kucklick, J.R. (2012). "Perfluoroalkyl Contaminants in Plasma of Five Sea Turtle Species:
552 Comparisons in Concentration and Potential Health Risks." Environmental Toxicology and
553 Chemistry **31**(6): 1223-1230.

554 Keller, J.M., Pugh, R.S. and Becker, P.R. (2014b). Biological and Environmental Monitoring
555 and Archival of Sea Turtle Tissues (BEMAST): Rationale, Protocols, and Initial
556 Collections of Banked Sea Turtle Tissues. NIST IR **7996**.
557 <https://nvlpubs.nist.gov/nistpubs/ir/2014/NIST.IR.7996.pdf>

558 Kurtz, A.E., Reiner, J.L., West, K.L. and Jensen, B.A. (2019). "Perfluorinated Alkyl Acids in
559 Hawaiian Cetaceans and Potential Biomarkers of Effect: Peroxisome Proliferator-Activated
560 Receptor Alpha and Cytochrome P450 4A." Environmental Science and Technology **53**(5):
561 2830-2839.

562 Lau, C., Anitole, K., Hodes, C., Lai, D., Pfahles-Hutchens, A. and Seed, J. (2007).
563 "Perfluoroalkyl Acids: A Review of Monitoring and Toxicological Findings."
564 Toxicological Sciences **99**(2): 366-394.

565 Malarovannan, G., Takahashi, S., Isobe, T., Kunisue, T., Sudaryanto, A., Miyagi, T., Nakamura,
566 M., Yasumura, S. and Tanabe, S. (2011). "Levels and distribution of polybrominated
567 diphenyl ethers and organochlorine compounds in sea turtles from Japan." Marine
568 Pollution Bulletin **63**(5-12): 172-178.

569 McCarthy, C. (2017). "Ecological Considerations of Per- and Polyfluoroalkyl Substances
570 (PFAS)." Current Pollution Reports **3**(4): 289-301.

571 McFadden, K.W., Gomez, A., Sterling, E., J. and Naro-Maciel, E. (2014). "Potential impacts of
572 historical disturbance on green turtle health in the unique & protected marine ecosystem of
573 Palmyra Atoll (Central Pacific)." Marine Pollution Bulletin **89**(1-2): 160-167.

574 Miller, J.D. (1997). Reproduction in Sea Turtles. The Biology of Sea Turtles. Lutz, P.L. and
575 Musick, J.A., CRC Press: 51-81.

576 Miller, J.D., Mortimer, J.A. and Limpus, C.J. (2017). "A Field Key to the Developmental Stages
577 of Marine Turtles (Cheloniidae) with Notes on the Development of Dermochelys."
578 Chelonian Conservation and Biology **16**(2): 111-122.

579 Molina, E.D., Balander, R., Fitzgerald, S.D., Giesy, J.P., Kannan, K., Mitchell, R. and Bursian,
580 S.J. (2006). "Effects of air cell injection of perfluorooctane sulfonate before incubation on
581 development of the white leghorn chicken (*Gallus domesticus*) embryo." Environmental
582 Toxicology and Chemistry **25**(1): 227-232.

583 Moody, C.A. and Field, J.A. (2000). "Perfluorinated Surfactants and the Environmental
584 Implications of Their Use in Fire-Fighting Foams." Environmental Science and
585 Technology **34**(18): 3864-3870.

586 Munoz, C.C. and Vermeiren, P. (2020). "Maternal Transfer of Persistent Organic Pollutants to
587 Sea Turtle Eggs: A Meta-Analysis Addressing Knowledge and Data Gaps Towards an
588 Improved Synthesis of Research Outputs." Environmental Toxicology and Chemistry
589 **39**(1): 9-29.

590 National Academy of Sciences (2004). Evolution in Hawaii: A Supplement to 'Teaching About
591 Evolution and the Nature of Science'. Washington, DC, The National Academies Press.

592 [https://www.nap.edu/catalog/10865/evolution-in-hawaii-a-supplement-to-teaching-about-
594 evolution-and](https://www.nap.edu/catalog/10865/evolution-in-hawaii-a-supplement-to-teaching-about-
593 evolution-and)
594 NMFS and US FWS (National Marine Fisheries Service and U.S. Fish and Wildlife Service)
595 (1998a). Recovery Plan for U.S. Pacific Populations of the Green Turtle (*Chelonia mydas*).
596 National Marine Fisheries Service. Silver Spring, MD.
597 [https://www.fisheries.noaa.gov/resource/document/recovery-plan-us-pacific-populations-
599 green-turtle-chelonia-mydas](https://www.fisheries.noaa.gov/resource/document/recovery-plan-us-pacific-populations-
598 green-turtle-chelonia-mydas)
599 NMFS and US FWS (National Marine Fisheries Service and U.S. Fish and Wildlife Service).
600 (1998b). Recovery Plan for U.S. Pacific Populations of the Hawksbill Turtle (*Eretmochelys* 601
602 *imbricata*). National Marine Fisheries Service. Silver Spring, MD.
603 [https://www.fisheries.noaa.gov/resource/document/recovery-plan-us-pacific-populations-
605 hawksbill-turtle-eretmochelys-imbricata](https://www.fisheries.noaa.gov/resource/document/recovery-plan-us-pacific-populations-
604 hawksbill-turtle-eretmochelys-imbricata)
604 O'Connell, S.G., Arendt, M., Segars, A., Kimmel, T., Braun-Mcneill, J., Avens, L., Schroeder,
605 B., Ngai, L., Kucklick, J.R. and Keller, J.M. (2010). "Temporal and spatial trends of
606 perfluorinated compounds in juvenile loggerhead sea turtles (*Caretta caretta*) along the
607 east coast of the United States." Environmental Science and Technology **44**(13): 5202-
608 5209.
609 Parham, J.E., Higashi, G.R., Lapp, E.K., Kuamo'o, D.G.K., Nishimoto, R.T., Hau, S.,
610 Fitzsimons, J.M., Polhemus, D.A. and Devick, W.S. (2008a). Atlas of Hawaiian
611 watersheds and their aquatic resources: Kapoho, Hawai'i. Honolulu, HI, Bishop Museum
612 and State of Hawai'i Division of Aquatic Resources: 1121-1128.
613 <https://hawaiiwatershedatlas.com/watersheds/hawaii/83037.pdf>
614 Parham, J.E., Higashi, G.R., Lapp, E.K., Kuamo'o, D.G.K., Nishimoto, R.T., Hau, S.,
615 Fitzsimons, J.M., Polhemus, D.A. and Devick, W.S. (2008b). Atlas of Hawaiian
616 watersheds and their aquatic resources: Kawainui, O'ahu. Honolulu, HI, Bishop Museum
617 and State of Hawai'i Division of Aquatic Resources: 257-266.
618 <https://hawaiiwatershedatlas.com/watersheds/oahu/32013.pdf>
619 Parham, J.E., Higashi, G.R., Lapp, E.K., Kuamo'o, D.G.K., Nishimoto, R.T., Hau, S.,
620 Fitzsimons, J.M., Polhemus, D.A. and Devick, W.S. (2008c). Atlas of Hawaiian
621 watersheds and their aquatic resources: Pu'uwa'awa'a, Hawai'i. Honolulu, HI, Bishop
622 Museum and State of Hawai'i Division of Aquatic Resources: 1215-1222.
623 <https://hawaiiwatershedatlas.com/watersheds/hawaii/85042.pdf>
624 Parker, D.M., Balazs, G.H., King, C.S., Katahira, L. and Gilmartin, W. (2009). "Short-Range
625 Movements of Hawksbill Turtles (*Eretmochelys imbricata*) from Nesting to Foraging
626 Areas within the Hawaiian Islands 1." Pacific Science **63**(3): 371-382.
627 Parker, D.M., Dutton, P.H. and Balazs, G.H. (2011). "Oceanic Diet and Distribution of
628 Haplotypes for the Green Turtle, *Chelonia mydas*, in the Central North Pacific." Pacific
629 Science **65**(4): 419-431.
630 Pasanisi, E., Cortés-Gómez, A.A., Pérez-López, M., Soler, F., Hernández-Moreno, D., Guerranti,
631 C., Martellini, T., Fuentes-Mascorro, G., Romero, D. and Cincinelli, A. (2016). "Levels of
632 perfluorinated acids (PFCAs) in different tissues of *Lepidochelys olivacea* sea turtles from
633 the Escobilla beach (Oaxaca, Mexico)." The Science of the Total Environment **572**: 1059-
634 1065.
635 Place, B.J. and Field, J.A. (2012). "Identification of Novel Fluorochemicals in Aqueous Film-
636 Forming Foams Used by the US Military." Environmental Science & Technology **46**:
637 7120-7127.

638 Reiner, J.L. and Place, B.J. (2015). Perfluorinated Alkyl Acids in Wildlife. Toxicological Effects
639 of Perfluoroalkyl and Polyfluoroalkyl Substances. DeWitt, J.C., Humana Press, Cham.
640 Schultz, M.M., Barofsky, D.F. and Field, J.A. (2004). "Quantitative Determination of
641 Fluorotelomer Sulfonates in Groundwater by LC MS/MS." Environmental Science and
642 Technology **38**(6): 1828-1835.

643 Shimada, T., Limpus, C.J., Hamann, M., Bell, I., Esteban, N., Groom, R. and Hays, G.C. (2020).
644 "Fidelity to foraging sites after long migrations." Journal of Animal Ecology **89**(4): 1008-
645 1016.

646 South Florida Water Management District. (2009). "St. Lucie River Watershed Protection Plan."
647 Retrieved October 23, 2020, from
648 https://www.sfwmd.gov/sites/default/files/documents/ne_slrwpp_main_123108.pdf.

649 Sterling, E.J., McFadden, K.W., Holmes, K.E., Vintinner, E.C., Arengo, F. and Naro-Maciel, E.
650 (2013). "Ecology and Conservation of Marine Turtles in a Central Pacific Foraging
651 Ground." Chelonian Conservation and Biology **12**(1): 2-16.

652 Summers, T.M., Jones, T.T., Martin, S.L., Hapedei, J.R., Ruak, J.K. and Lepczyk, C.A. (2017).
653 "Demography of Marine Turtles in the Nearshore Environments of the Northern Mariana
654 Islands." Pacific Science **71**(3): 269-286.

655 Summers, T.M., Kelly, I.K., Work, T.M., Ruack, J.K. and Hapdei, J.R. (2018). "Human induced
656 trauma and directed take inhibits sea turtle recovery in the Commonwealth of the Northern
657 Mariana Islands." Micronesia **8**: 1-19.

658 United Nations Dept. of Economic and Social Affairs. (2019). "World Population Prospects 2019
659 Volume I: Comprehensive Tables." Retrieved October 22, 2020, from
660 [https://population.un.org/wpp/Publications/Files/WPP2019_Volume-I_Comprehensive-](https://population.un.org/wpp/Publications/Files/WPP2019_Volume-I_Comprehensive-Tables.pdf)
661 [Tables.pdf](https://population.un.org/wpp/Publications/Files/WPP2019_Volume-I_Comprehensive-Tables.pdf).

662 US Census Bureau. (2019a). "QuickFacts Honolulu County, Hawaii; Hawaii; United States."
663 Retrieved October 23, 2020, from
664 <https://www.census.gov/quickfacts/fact/table/honolulucountyhawaii,HI,US/PST045219>.

665 US Census Bureau. (2019b). "QuickFacts Okeechobee County, Florida; Palm Beach County,
666 Florida; Martin County, Florida; St. Lucie County, Florida." Retrieved October 22, 2020,
667 from
668 [https://www.census.gov/quickfacts/fact/table/okeechobeecountyflorida,martincountyflorida](https://www.census.gov/quickfacts/fact/table/okeechobeecountyflorida,martincountyflorida,sluciecountyflorida,palmbeachcountyflorida/PST045219)
669 [,sluciecountyflorida,palmbeachcountyflorida/PST045219](https://www.census.gov/quickfacts/fact/table/okeechobeecountyflorida,martincountyflorida,sluciecountyflorida,palmbeachcountyflorida/PST045219).

670 US Census Bureau. (2019c). "QuickFacts St. Lucie County, Florida; Port St. Lucie city, Florida."
671 Retrieved October 22, 2020, from
672 <https://www.census.gov/quickfacts/fact/table/stluciecountyflorida,portstluciecityflorida/PS>
673 [T045219](https://www.census.gov/quickfacts/fact/table/stluciecountyflorida,portstluciecityflorida/PS).

674 US EPA. (2018). "Fact Sheet: 2010/2015 PFOA Stewardship Program." Retrieved October 20,
675 2020, from [https://www.epa.gov/assessing-and-managing-chemicals-under-tsca/fact-sheet-](https://www.epa.gov/assessing-and-managing-chemicals-under-tsca/fact-sheet-20102015-pfoa-stewardship-program)
676 [20102015-pfoa-stewardship-program](https://www.epa.gov/assessing-and-managing-chemicals-under-tsca/fact-sheet-20102015-pfoa-stewardship-program).

677 Vaughan, P. and Spring, S. (1980). Long Distance Hawksbill Recovery. Marine Turtle
678 Newsletter. **16**: 6-7.

679 Weir, S.M., Suski, J.G. and Salice, C.J. (2010). "Ecological risk of anthropogenic pollutants to
680 reptiles: Evaluating assumptions of sensitivity and exposure." Environmental Pollution
681 **158**: 3596-3607.

682 Wilson, T.B., Stevenson, G., Crough, R., de Araujo, J., Fernando, N., Anwar, A., Scott, T.,
683 Quinteros, J., Scott, P.C. and Archer, M.J.G. (2020). "Evaluation of Residues in Hen Eggs

684 after Exposure of Laying Hens to Water Containing Per- and Polyfluoroalkyl Substances."
685 Environmental Toxicology and Chemistry <https://doi.org/10.1002/etc.4723>.
686 Wood, L.D., Brunnick, B. and Milton, S.L. (2017). "Home Range and Movement Patterns of
687 Subadult Hawksbill Sea Turtles in Southeast Florida." Journal of Herpetology **51**(1): 58-
688 67.
689 Yamashita, N., Taniyasu, S., Petrick, G., Wei, S., Gamo, T., Lam, P.K.S. and Kannan, K. (2008).
690 "Perfluorinated acids as novel chemical tracers of global circulation of ocean waters."
691 Chemosphere **70**: 1247-1255.
692 Young, C.J., Furdui, V.I., Franklin, J., Koerner, R.M., Muir, D.C.G. and Mabury, S.A. (2007).
693 "Perfluorinated Acids in Arctic Snow: New Evidence for Atmospheric Formation."
694 Environmental Science & Technology **41**: 3455-3461.

695 Table 1. Samples sizes and mass fractions (ng/g wet mass) of predominant perfluoroalkyl

696 substances (PFASs) in plasma of North Pacific green and hawksbill sea turtles.

697

Green turtle	Plasma	Kapoho (live captured)	2011-2012	39.2 - 83.4	13	0.407	0.796	0.734	0.196 - 2.19	100	0.407	0.796	0.734	0.196-2.19	100	
Green turtle	Plasma	Kailua (live captured)	2011-2012	42.9 - 80.3	13	2.11	2.45	1.32	0.865 - 4.52	100	2.11	2.45	1.32	0.865-4.52	100	
Green turtle	Plasma	MHI FP stranded	2011-2012	35.5 - 70.2	14	0.242	0.902	1.04	<0.054 - 2.77	85.7	0.242	0.816	0.912	<0.054-2.52	85.7	
Green turtle	Plasma	CNMI	2013		12	1.74	0.562	1.12	0.141 - 4.11	100	0.634	0.524	0.987	0.141-3.65	100	
Green turtle	Plasma	Saipan	2013	46.3 - 85.6	6	0.295	0.926	1.56	0.175 - 4.11	100	0.295	0.850	1.37	0.175-3.65	100	
Green turtle	Plasma	Tinian	2013	47.7 - 63.7	6	0.204	0.199	0.045	0.141 - 0.270	100	0.204	0.199	0.045	0.141-0.27	100	
Green turtle	Plasma	Palmyra Atoll	2012-2013	44.3 - 87.6	10	0.155	0.171	0.051	<0.085 - 0.265	80	0.155	0.171	0.051	<0.085-0.265	80	
Green turtle	Plasma	Core Sound, NC, USA (Keller et al., 2012)	2006	within 25-70	10	2.92	3.39	1.74	0.871 - 6.09	100	1.12	2.41	2.37	0.871 - 3.87	100	
Hawksbill turtle	Plasma	CNMI	2013	35.1 - 48.3	4	1.74	1.99	0.951	1.19 - 3.29	100	0.634	0.654	0.285	0.334-1.01	100	0.972
Hawksbill turtle	Plasma	Palmyra	2013	54.4 - 55.5	2		4.18	N/A	3.373 - 4.978	100		0.300	N/A	0.253-0.348	100	
Hawksbill turtle	Plasma	Juno Beach, FL, USA (Keller et al., 2012)	2006	within 25-70	5	33.5	44.1	23.5	24.8 - 79.0	100	11.9	11.9	6.27	5.45 - 21.2	100	17.0
Hawksbill turtle	Egg contents	MHI	2012	unknown	7*	31.0	35.0	34.1	5.46 - 106	100	2.92	16.3	34.2	0.862 - 93.8	100	1.28

698

699 Abbreviations: Commonwealth of the Northern Marianas Islands (CNMI), Main Hawaiian

700 Islands (MHI), North Carolina (NC), Florida (FL), % >R = percent of samples

701 above the reporting limit.

702 *Eleven nests from seven mothers were analyzed. Multiple nests from the same mom were

703 averaged before the summary statistics shown here were calculated for all seven moms.

704 NC and FL data taken from Keller et al. (2012).

705

706 Table 2. Estimated margins of safety (EMOS) for average/maximum sea turtle exposure to
 707 perfluorooctane sulfonate based on the lowest adverse effect levels in lab-exposed animals.

708

		LOEAL (ng/g or ng/mL)	85000	1500	2290	1000	5
		LOEAL in	serum	serum	serum	serum	se
		LOEAL in	rat/mouse	rat	rat	zebrafish	m
		Reference	Bjorket al 2008; Lau et al 2003	Currane et al 2008	Wanget al 2011	Zhang et al 2011	Peden- 2
Species	Tissue	Turtle Grouping	Estimated margin of safety for average / maximum turtle				
Green turtle	Plasma	MHI Live captured, no FP stranded	74561 / 18805	1316 / 332	2009 / 507	877 / 221	80*
Green turtle	Plasma	Kiholo (live captured)	464481 / 151786	8197 / 2679	12514 / 4089	5464 / 1786	50C
Green turtle	Plasma	Kapoho (live captured)	106784 / 38813	1884 / 685	2877 / 1046	1256 / 457	11E
Green turtle	Plasma	Kailua (live captured)	34694 / 18805	612 / 332	935 / 507	408 / 221	37*
Green turtle	Plasma	MHI FP stranded	104167 / 33730	1838 / 595	2806 / 909	1225 / 397	112
Green turtle	Plasma	CNMI	162214 / 23288	2863 / 411	4370 / 627	1908 / 274	17E
Green turtle	Plasma	Palmyra Atoll	497076 / 320755	8772 / 5660	13392 / 8642	5848 / 3774	53!
Hawksbill turtle	Plasma	CNMI	129969 / 84158	2294 / 1485	3502 / 2267	1529 / 990	14C
Hawksbill turtle	Plasma	Palmyra	283333 / 244253	5000 / 4310	7633 / 6580	3333 / 2874	30E
Hawksbill turtle	Egg contents	MHI					


709

710


711

712

* concern of risk when EMOS <100, ** heightened concern of risk when EMOS <10.




713 Figure 1. Green and hawksbill sea turtle sampling sites and sample sizes. Dots show capture sites
 714 of free-ranging green (green) and hawksbill (purple) turtles and nest excavation sites of
 715 hawksbill nests (yellow). Fourteen stranded green turtles from MHI severely afflicted with FP
 716 plasma are not mapped.




717

718 Figure 2. PFAS profiles detected in Pacific Island sea turtles. MHI = Main Hawaiian Islands,
 719 CNMI = Commonwealth of the Northern Marianas Islands. *All green turtles from MHI, CNMI
 720 and Palmyra were combined excluding the MHI FP-stranded green turtles. Perfluoroalkyl
 721 sulfonates (blue shades) are visually distinguished from perfluoroalkyl carboxylates (other
 722 colors).




723
 724 Figure 3. Mean (one standard deviation) PFAS concentrations (ng/g) in (a) green sea turtle and
 725 (b) hawksbill sea turtle plasma (bars) in relation to human population (dots). Commonwealth of
 726 the Northern Marianas Islands (CNMI), Main Hawaiian Islands (MHI), North Carolina (NC),
 727 Florida (FL). Sample sizes were 10, 12, 39, and 10 green turtles, and 2, 4, and 5 hawksbills,
 728 respectively. NC and FL PFOS data are from Keller et al. (2012). Population data are from

729 O'Connell et al. (2010) for NC, and summed for counties in the St. Lucie-Loxahatchee
730 Watershed (US Census Bureau 2019b, c) for FL. Stranded green turtles with severe
731 fibropapillomatosis were excluded from the MHI data. The asterisk indicates a difference in
732 PFOS concentration from other Pacific sites ($p < 0.05$).



733

734 Figure 4. PFAS quantities (ng) estimated in the entire clutch of eggs from five clutches laid by
 735 hawksbill turtle 19591-04 on Maui. Linear regression statistics for ng vs. clutch order are shown
 736 for each compound.



737

738 Figure 5. Significant correlations between perfluoroundecanoate (PFUnA) (A) and
 739 perfluorotridecanoate (PFTriA) (B) concentrations (ng/g) in eggs from 11 hawksbill sea turtle
 740 nests from the Main Hawaiian Islands and emergence success.

741

Table 1. Samples sizes and mass fractions (ng/g wet mass) of predominant perfluoroalkyl substances (PFASs) in plasma of North Pacific green and hawksbill sea turtles.

Species	Tissue	Grouping	Year	SCL range (cm)	n	TOTAL PFASs					PFOS					PFNA				
						Median	Mean	SD	Min-Max	%>RL	Median	Mean	SD	Min-Max	%>RL	Median	Mean	SD	Min-Max	%>RL
Green turtle	Plasma	MHI Live captured, no FP stranded	2011-2012	35.5 - 83.4	39	0.447	1.14	1.29	<0.063 - 4.52	92.3	0.447	1.14	1.29	<0.063- 4.52	92.3				<0.188 - <1.18	0
Green turtle	Plasma	Kiholo (live captured)	2011-2012	43.7 - 65.1	13	0.126	0.183	0.121	<0.063 - 0.560	76.9	0.126	0.183	0.121	<0.063- 0.560	76.9				<0.188 - <1.03	0
Green turtle	Plasma	Kapoho (live captured)	2011-2012	39.2 - 83.4	13	0.407	0.796	0.734	0.096 - 2.19	100	0.407	0.796	0.734	0.196- 2.19	100				<0.239 - <0.893	0
Green turtle	Plasma	Kailua (live captured)	2011-2012	42.9 - 80.3	13	2.11	2.45	1.32	0.865 - 4.52	100	2.11	2.45	1.32	0.865- 4.52	100				<0.233 - <1.18	0
Green turtle	Plasma	MHI FP stranded	2011-2012	35.5 - 70.2	14	0.242	0.902	1.04	<0.054 - 2.77	85.7	0.242	0.816	0.912	<0.054- 2.52	85.7				<0.162 - <1.21	0
Green turtle	Plasma	CNMI	2013		12	1.74	0.562	1.12	0.141 - 4.11	100	0.634	0.524	0.987	0.141- 3.65	100				<0.301 - <0.377	0
Green turtle	Plasma	Saipan	2013	46.3 - 85.6	6	0.295	0.926	1.56	0.175 - 4.11	100	0.295	0.850	1.37	0.175- 3.65	100				<0.326 - <0.377	0
Green turtle	Plasma	Tinian	2013	47.7 - 63.7	6	0.204	0.199	0.045	0.141 - 0.270	100	0.204	0.199	0.045	0.141- 0.27	100				<0.301 - <0.369	0
Green turtle	Plasma	Palmyra Atoll	2012-2013	44.3 - 87.6	10	0.155	0.171	0.051	<0.085 - 0.265	80.0	0.155	0.171	0.051	<0.085- 0.265	80.0				<0.273 - <8.70	0
Green turtle	Plasma	Core Sound,	2006	25 - 70	10	2.92	3.39	1.74	0.871 - 6.09	100	1.12	2.41	2.37	0.871 - 3.87	100				<0.070 -	10

		NC, USA (Keller et al., 2012)																	0.182	
Hawksbill turtle	Plasma	CNMI	2013	35.1 - 48.3	4	1.74	1.99	0.951	1.19 - 3.29	100	0.634	0.654	0.285	0.334-1.01	100	0.972	1.08	0.516	0.582 - 1.79	100
Hawksbill turtle	Plasma	Palmyra	2013	54.4 - 55.5	2		4.18	N/A	3.373 - 4.978	100		0.300	N/A	0.253-0.348	100		3.40	N/A	2.90 - 3.90	100
Hawksbill turtle	Plasma	Juno Beach, FL, USA (Keller et al., 2012)	2006	25 - 70	5	33.5	44.1	23.5	24.8 - 79.0	100	11.9	11.9	6.27	5.45 - 21.2	100	17.0	17.3	1.21	3.87 - 30.8	100
Hawksbill turtle	Egg contents	MHI	2012	unknown	7*	31.0	35.0	34.1	5.46 - 106	100	2.92	16.3	34.2	0.862 - 93.8	100	1.28	7.44	8.90	0.756 - 22.8	100

Abbreviations: Commonwealth of the Northern Marianas Islands (CNMI), Main Hawaiian Islands (MHI), North Carolina (NC), Florida (FL), % >R

= percent of samples

above the reporting limit.

*Eleven nests from seven mothers were analyzed. Multiple nests from the same mom were averaged before the summary statistics shown here were calculated for all seven moms.

NC and FL data taken from Keller et al. (2012).

Table 2. Estimated margins of safety (EMOS) for average/maximum sea turtle exposure to perfluorooctane sulfonate based on the lowest adverse effect levels in lab-exposed animals.

		Adverse effect	Neonate mortality, altered liver histology and gene expression	Increased liver weight	Altered thyroid hormones	Altered development of motor neurons	Decreased T-cell dependent IgM antibody response (immunosuppression)	Decreased hatching success
		LOEAL (ng/g or ng/mL)	85000	1500	2290	1000	91.5	100
		LOEAL in	serum	serum	serum	serum	serum	eggs
		LOEAL in	rat/mouse	rat	rat	zebrafish	mouse	chicken
		Reference	Bjork et al 2008; Lau et al 2003	Currane et al 2008	Wang et al 2011	Zhang et al 2011	Peden-Adams et al 2008	Molina et al 2006
Species	Tissue	Turtle Grouping	Estimated margin of safety for average / maximum turtle					
Green turtle	Plasma	MHI Live captured, no FP stranded	74561 / 18805	1316 / 332	2009 / 507	877 / 221	80* / 20*	
Green turtle	Plasma	Kiholo (live captured)	464481 / 151786	8197 / 2175	12514 / 4089	5464 / 1786	500 / 163	
Green turtle	Plasma	Kapoho (live captured)	106784 / 38813	1884 / 685	2877 / 1046	1256 / 457	115 / 42*	
Green turtle	Plasma	Kailua (live captured)	34694 / 18805	612 / 332	935 / 507	408 / 221	37* / 20*	
Green turtle	Plasma	MHI FP stranded	104167 / 33730	1838 / 595	2806 / 909	1225 / 397	112 / 36*	
Green turtle	Plasma	CNMI	162214 / 23288	2863 / 411	4370 / 627	1908 / 274	175 / 25*	
Green turtle	Plasma	Palmyra Atoll	497076 / 320755	8772 / 5660	13392 / 8642	5848 / 3774	535 / 345	
Hawksbill turtle	Plasma	CNMI	129969 / 84158	2294 / 1485	3502 / 2267	1529 / 990	140 / 91*	
Hawksbill turtle	Plasma	Palmyra	283333 / 244253	5000 / 4310	7633 / 6580	3333 / 2874	305 / 263	
Hawksbill turtle	Egg contents	MHI						6** / 1**

* concern of risk when EMOS <100, ** heightened concern of risk when EMOS <10.

Highlights

- Perfluorinated alkyl substances were detected in sea turtles from the North Pacific
- PFAS levels in sea turtles were correlated with human population and land use
- Maternal offloading of PFASs was detected in unhatched hawksbill eggs from Hawaii
- Levels in eggs were near those which have caused developmental toxicity in birds
- No correlation was found between fibropapillomatosis (FP) and PFAS concentrations

1 **Supplemental Information for**

2
3 **Sea turtles across the North Pacific are exposed to perfluoroalkyl substances**

4
5 Cathryn Wood¹, George H. Balazs², Marc Rice³, Thierry M. Work⁴, T. Todd Jones⁵, Eleanor
6 Sterling⁶, Tammy M. Summers⁷, John Brooker⁸, Lauren Kurpita⁹, Cheryl S. King¹⁰, and Jennifer
7 M. Lynch^{1,11*}

8
9 1 Hawai‘i Pacific University, Center for Marine Debris Research, Waimānalo, HI, USA

10 2 Golden Honu Services of Oceania, Honolulu, HI, USA

11 3 Hawai‘i Preparatory Academy, Waimea, HI, USA

12 4 U.S. Geological Survey, National Wildlife Health Center, Honolulu Field Station, Honolulu,
13 HI, USA

14 5 NOAA Pacific Islands Fisheries Science Center, Honolulu, HI, USA

15 6 American Museum of Natural History, New York, NY

16 7 Rainbow Connection Research, Guam, USA

17 8 College of Charleston, Charleston, SC, USA

18 9 Hawai‘i Island Hawksbill Turtle Recovery Project, Hawai‘i National Park, HI, USA

19 10 Hawaiian Hawksbill Conservation, Kihei, HI, USA

20 11 National Institute of Standards and Technology, Chemical Sciences Division, Waimānalo, HI,

21 USA E-mail: Jennifer.lynch@nist.gov, *corresponding author

22 Table of Contents

23 Detailed methods of laboratory analysis and data analysis 3

24 Palmyra Atoll Spatial Comparison 6

25 Figure S1. PFAS profiles in Pacific Islands sea turtle tissues 8

26 Figure S2. PFOS concentrations in plasma of 4 green turtle groups from MHI related to
27 fibropapillomatosis (FP) and percent of urban development of each watershed9

28 Figure S3. PFAS concentrations in green and hawksbill sea turtle plasma from CNMI..... 10

29 Figure S4. Σ PFAS concentrations in hawksbill eggs in relation to distance to nearest
30 international airport 11

31 Figure S5. PFAS profiles in individual hawksbill nests from Main Hawaiian Islands..... 12

32 Figure S6. A hawksbill sea turtle post-nesting migrations and dive behavior profile 13

33 References 14

34 Tables S1-S8..... Separate Supporting Information Excel Spreadsheet file

35 Detailed methods of laboratory analysis and data analysis

36 PFAS Measurements

37 Each plasma and egg sample was measured for concentrations of 15 PFASs according to
38 methods modified from (Keller et al. 2012). The calibration series consisted of NIST Reference
39 Materials 8446 Perfluorinated Carboxylic Acids and Perfluorooctane Sulfonamide in Methanol
40 and 8447 Perfluorinated Sulfonic Acids in Methanol. The internal standard (IS) solution
41 consisted of 11 mass-labeled PFASs purchased from Wellington Laboratories (Guelph Ontario,
42 Canada), RTI International (Research Triangle Park, NC, USA) and Cambridge Isotope Labs
43 (Tewksbury, MA, USA) gravimetrically combined in methanol. Compound names,
44 abbreviations, and IS choices are in Table S3. As part of the BEMAST protocols, three field
45 blanks were prepared by pulling Millipore water stored in a polyethylene bottle through the same
46 lot number of blood collection needles, tubes, pipets, and storage vials. Control materials were
47 five replicates of SRM 1957 Organic Contaminants in Non-Fortified Human Serum (≈ 1 g each)
48 and three replicates of SRM 1947 Lake Michigan Fish Tissue (≈ 1 g each). Plasma samples and
49 field blanks (≈ 1 g each) archived in plastic cryovials were thawed, mixed, transferred to glass
50 culture tubes gravimetrically, and 0.6 mL of IS solution was added gravimetrically to provide ≈ 3
51 ng of each mass-labeled compound to each sample. The calibration curve consisted of six tubes
52 with gravimetric dilutions from ≈ 0.05 ng to 275 ng of each compound. The IS solution and 1 g of
53 Millipore water were added to calibrants, and to three laboratory blanks. Tubes were vortexed,
54 and equilibrated at room temperature for 1.5 h or longer. Samples were extracted with 4 mL of
55 acetonitrile and 30 min of sonication. The supernatant, after centrifuging the tubes at 157 rad/s
56 (1500 rpm) for 5 min, was transferred to clean glass tubes. SRM 1957 replicates were processed
57 the same way.

58 Egg samples archived in plastic cryovials were thawed (≈ 1 g each), transferred to glass
59 tubes gravimetrically, centrifuged at 157 rad/s for 5 min, and IS solution (0.6 mL) was added
60 gravimetrically to provide ≈ 3 ng of each mass-labeled compound to each sample. Tubes were
61 vortexed for 20 s and allowed to equilibrate for 1.5 h or longer. Samples were extracted with 6
62 mL of acetonitrile and 20 s of vortex and 30 min of sonication. The supernatant, after
63 centrifuging the tubes at 157 rad/s for 5 min, was transferred to clean glass tubes. Samples were
64 extracted a second time with 6 mL acetonitrile, and extracts were combined with the first
65 extraction. After freezing, the extracts were no longer cloudy. The precipitate was discarded after
66 centrifugation at 157 rad/s for 5 min and transferring the supernatant to a clean culture tube.
67 SRM 1947 replicates were processed the same way.

68 Extracts were solvent exchanged to methanol using nitrogen evaporation being careful to
69 not evaporate to dryness. They were purified using 250 mg Envi-carb columns (Supelco,
70 Bellefonte, Pennsylvania) by loading the 3 mL sample, a 1 mL sample rinse of methanol, and 6
71 mL of methanol. The extracts were concentrated under nitrogen, and transferred to glass
72 autosampler vials for a final volume of 0.5 mL in methanol. Samples were analyzed using a
73 liquid chromatograph (Agilent 1100 HPLC, Palo Alto, CA) interfaced to a negative electrospray
74 ionization tandem mass spectrometer (API 4000, Applied Biosystems-MDS Sciex, Foster City,
75 CA). Samples (20 μ L) were injected onto an analytical column (Agilent Zorbax Eclipse Plus
76 C18, 150 mm x 2.1 mm x 5 μ m). The solvent gradient at a flow rate of 0.25 mL/min (all
77 mixtures expressed in volume fractions) started at 50 % methanol and 50 % 20 mmol/L
78 ammonium acetate in water and increased to 75% methanol by 20 min, held for 5 min, and then
79 increased to 95 % methanol by 35 min, held for 5 min, before reverting back to original
80 conditions at 40.5 min with a 9.5 min hold time. The MS/MS method was divided into six

81 windows and two of the most abundant transitions for each PFAS were monitored. The internal
82 standard approach was used to quantify each compound amount. Amounts of each analyte were
83 calculated using the slope and y-intercept of at least a three point calibration curve that bracketed
84 the peak area ratios observed in the samples. Concentrations were determined by dividing the
85 calculated analyte mass by the extracted sample mass. Reporting limits (RLs) were determined
86 using the highest calculated RL of two methods described in O'Connell et al. (2010).
87 Concentrations are totals of linear and branched isomers.

88 **Data Analyses**

89 JMP statistical software (SAS Institute Inc. Cary, NC) was used when all samples were
90 >RLs, and on occasion when required, with values <RL substituted with half the RL. When
91 some samples were <RLs, every attempt was made to use the R software package, NADA, as
92 suggested by Helsel (2005). Means, medians and standard deviations were calculated with either
93 Kaplan-Meier or regression on order statistical models. Significance was determined as $p < 0.05$.
94 Data normality and homoscedasticity were tested with Shapiro-Wilk W tests and O'Brien or
95 Bartlett tests, respectively. If assumptions were violated even with transformed data, non-
96 parametric tests were used. The cendiff function (non-parametric ANOVA) in R NADA was
97 used to test significant differences among sites for green sea turtle plasma PFAS concentrations,
98 followed by the Steel Dwass non-parametric multiple comparison test using JMP. The Wilcoxon
99 t-test in JMP was used to compare hawksbill plasma PFOS, PFNA, and \sum PFAS plasma
100 concentrations between Palmyra Atoll and CNMI, while the cendiff function was necessary for
101 PFUnA. Species differences were assessed only within CNMI data. PFOS and \sum PFAS plasma
102 concentrations between species could be compared using Wilcoxon tests in JMP, and PFNA and
103 PFUnA were tested with cendiff function. MHI green turtle plasma samples were grouped

104 according to location and FP incidence (same as in Keller et al. (2014)). Differences among
105 groups in PFOS concentrations was assessed with the cendiff function in R NADA, followed by
106 Steel-Dwass multiple comparison tests in JMP.


107 Amount of PFASs offloaded into each nest was estimated by multiplying the total
108 number eggs laid by the average grams of each pooled egg contents from that nest, and by the
109 measured PFAS concentration (in ng/g). The nanograms of PFASs offloaded were then
110 compared across clutches laid within one nesting season from a single mother using linear
111 regressions. PFAS concentrations in the eggs were compared to nest success variables using the
112 cenken function in R NADA, a version of Kendall's tau correlation.

113 To assess toxicological risk, estimated margins of safety (EMOS) were calculated as
114 described in Keller et al. (2012), as the ratio of the average plasma, serum, or egg PFOS
115 concentration in laboratory-exposed animals in the lowest observed adverse effect level
116 (LOAEL) dose group to either the average or maximum sea turtle plasma or egg PFOS
117 concentration. EMOS below 100 were considered of concern (Faustman and Omenn 1996).

118 **Palmyra Atoll Spatial Comparison**


119 If local sources released PFASs at Palmyra Atoll, which are unexpected, they could be
120 from runoff/sewage from the scientific field station or non-military use of AFFFs on the airstrips
121 from Cooper Island in the northwestern portion of the atoll. Surface currents nearshore and
122 within the lagoon flow westward (Gardner et al. 2011), so we hypothesized that turtles sampled
123 from the lagoon or west would have greater PFAS concentrations. Sample sizes prevented
124 statistical testing, but a cursory comparison was attempted of green turtle PFAS concentrations
125 on the four sides and within the lagoon to investigate the possibility of a point source on the
126 atoll. \sum PFASs in green turtles were greatest in a turtle (MT13W003) from the east (0.265 ng/g)

127 and least in the turtle captured twice (MT12W064 and MT13W014, adult female) in the lagoon
128 (<0.088 ng/g and 0.155 ng/g), refuting the hypothesis. No evidence of a significant PFAS point
129 source on Palmyra Atoll was observed. The small difference in PFAS concentrations between the
130 eastern and lagoon green turtles is likely related to life stage. The green turtle in the east was
131 only 44.3 cm straight carapace length, indicating recent recruitment to Palmyra. The lagoon
132 green turtle was an adult female and was captured twice over two years, suggesting residency for
133 at least a year. Green turtles undergo an ontogenetic shift from a carnivorous pelagic juvenile
134 phase to the herbivorous benthic juvenile phase when they recruit into nearshore habitats
135 (Bjorndal 1997). Because of their higher trophic status in the pelagic phase, green turtles may
136 recruit into Palmyra carrying a greater PFAS burden, which dilutes with growth through the next
137 benthic herbivorous phase (Keller 2013). The lesser concentrations in the adult female in the
138 lagoon may have resulted from her ability to offload PFASs to eggs. The eastern region of
139 Palmyra is known habitat for the smallest, recently recruited green turtles from the pelagic-phase
140 (Sterling et al. 2013) and green turtles in this region may enter with greater PFAS concentrations.
141 Spatial trends across the atoll are confounded by these life history traits. The data in the current
142 study suggest that remote Palmyra Atoll is a control reference site for PFAS with little to no
143 local point sources. McFadden et al. (2014) was the first study to measure chemical contaminants
144 (heavy metals) in Palmyra sea turtles and found elevated aluminum and iron.




145
 146 Figure S1. Mean and one standard deviation of PFAS profiles in sea turtle tissues from Pacific
 147 Islands. MHI = Main Hawaiian Islands, CNMI = Commonwealth of the Northern Marianas
 148 Islands. The y-axis values were derived by first calculating the percent of \sum PFAS in each
 149 individual plasma sample per compound. Those values were averaged across individual plasma
 150 samples. Samples that had no detectable PFASs were excluded. For hawksbill nests, the percent
 151 of \sum PFAS was calculated per nest for each compound. Percentages from multiple nests from the
 152 same mothers were averaged for the same compound, because these were similar. Then,
 153 percentages were averaged across the seven mothers for this graph. *All green turtles from MHI,

154 CNMI and Palmyra were combined, excluding the MHI FP-stranded green turtles.




155

156 Figure S2. Mean and one standard deviation of PFOS concentrations (ng/g) in plasma of four
157 green turtle groups from MHI related to fibropapillomatosis (FP) and urban development
158 percentage of each watershed. Mean PFOS concentrations increased with watershed
159 urbanization, with the highest being in Kailua, Hawai'i, rather than with FP incidence. FP
160 incidence is noted by more red colored bars. Black circles indicate % urbanization for each
161 Hawaiian watershed (Parham et al. 2008a; b; c). Different letters above bars indicate a significant
162 difference between groups ($p < 0.05$).




163

164 Figure S3. Mean and one standard deviation PFAS concentrations (ng/g) in green and hawksbill
 165 sea turtle plasma from CNMI. Patterned bars indicate that means fell below the detection
 166 limits. Sample sizes are 12 and 4. * indicates difference between species ($p < 0.05$).



167

168 Figure S4. Σ PFAS concentrations (ng/g wet mass) in hawksbill eggs in relation to the
 169 approximate distance (km) between the nests and the nearest international airport. When females
 170 laid multiple clutches, the average Σ PFAS concentrations of all clutches from that female were
 171 used. Pearson correlation statistics are shown.



173


174

Figure S5. PFAS profiles in egg contents of nests from individual hawksbill sea turtles laid in

175


different Hawaiian Island locations.

176 a.



177

178 b.



179

180 Figure S6. Hawksbill sea turtle 19591-04's a) post-nesting migrations, satellite tracked in 2004
181 (white dots, black line and arrows; base map modified from Figure 3C within Parker et al.
182 (2009)) and again in 2008 (yellow arrows) from Makena, Maui, to foraging grounds near
183 Kahuku, Oahu, and b) dive behavior profile for the 2008 track.

184 References

- 185 Bjorndal, K.A. (1997). Foraging Ecology and Nutrition of Sea Turtles. The Biology of Sea Turtles. Lutz,
186 P.L. and Musick, J.A., CRC Press: 199-231.
- 187 Faustman, E.M. and Omenn, G.S. (1996). Risk Assessment. Casarett and Doull's Toxicology the Basic
188 Science of Poisons. Klaassen, C.D. New York, NY, McGraw-Hill.
- 189 Gardner, J.P.A., Garton, D.W. and Collen, J.D. (2011). "Near-surface mixing and pronounced deep-water
190 stratification in a compartmentalised, human-disturbed atoll lagoon system." Coral Reefs **30**: 271-
191 282.
- 192 Helsel, D.R. (2005). Nondetects and data analysis. Statistics for censored environmental data. Hoboken,
193 Wiley-Interscience.
- 194 Keller, J.M. (2013). Exposure to and Effects of Persistent Organic Pollutants. The Biology of Sea Turtles
195 Volume III. Wyneken, J., Lohmann, K.J. and Musick, J.A. Boca Raton, FL, CRC Press Taylor
196 and Francis Group. **3**: 285-328.
- 197 Keller, J.M., Balazs, G.H., Nilsen, F., Rice, M., Work, T.M. and Jensen, B.A. (2014). "Investigating the
198 Potential Role of Persistent Organic Pollutants in Hawaiian Green Sea Turtle
199 Fibropapillomatosis." Environmental Science & Technology **48**: 7807-7816.
- 200 Keller, J.M., Ngai, L., Braun McNeill, J., Wood, L.D., Stewart, K.R., O'Connell, S.G. and Kucklick, J.R.
201 (2012). "Perfluoroalkyl Contaminants in Plasma of Five Sea Turtle Species: Comparisons in
202 Concentration and Potential Health Risks." Environmental Toxicology and Chemistry **31**(6):
203 1223-1230.
- 204 McFadden, K.W., Gomez, A., Sterling, E., J. and Naro-Maciel, E. (2014). "Potential impacts of historical
205 disturbance on green turtle health in the unique & protected marine ecosystem of Palmyra Atoll
206 (Central Pacific)." Marine Pollution Bulletin **89**(1-2): 160-167.
- 207 O'Connell, S.G., Arendt, M., Segars, A., Kimmel, T., Braun-Mcneill, J., Avens, L., Schroeder, B., Ngai,
208 L., Kucklick, J.R. and Keller, J.M. (2010). "Temporal and spatial trends of perfluorinated
209 compounds in juvenile loggerhead sea turtles (*Caretta caretta*) along the east coast of the United
210 States." Environmental Science and Technology **44**(13): 5202-5209.
- 211 Parham, J.E., Higashi, G.R., Lapp, E.K., Kuamo'o, D.G.K., Nishimoto, R.T., Hau, S., Fitzsimons, J.M.,
212 Polhemus, D.A. and Devick, W.S. (2008a). Atlas of Hawaiian watersheds and their aquatic
213 resources: Kapoho, Hawai'i. Honolulu, HI, Bishop Museum and State of Hawai'i Division of
214 Aquatic Resources: 1121-1128. <https://hawaiiwatershedatlas.com/watersheds/hawaii/83037.pdf>
- 215 Parham, J.E., Higashi, G.R., Lapp, E.K., Kuamo'o, D.G.K., Nishimoto, R.T., Hau, S., Fitzsimons, J.M.,
216 Polhemus, D.A. and Devick, W.S. (2008b). Atlas of Hawaiian watersheds and their aquatic
217 resources: Kawainui, O'ahu. Honolulu, HI, Bishop Museum and State of Hawai'i Division of
218 Aquatic Resources: 257-266. <https://hawaiiwatershedatlas.com/watersheds/oahu/32013.pdf>
- 219 Parham, J.E., Higashi, G.R., Lapp, E.K., Kuamo'o, D.G.K., Nishimoto, R.T., Hau, S., Fitzsimons, J.M.,
220 Polhemus, D.A. and Devick, W.S. (2008c). Atlas of Hawaiian watersheds and their aquatic
221 resources: Pu'uwa'awa'a, Hawai'i. Honolulu, HI, Bishop Museum and State of Hawai'i Division
222 of Aquatic Resources: 1215-1222.
223 <https://hawaiiwatershedatlas.com/watersheds/hawaii/85042.pdf>
- 224 Parker, D.M., Balazs, G.H., King, C.S., Katahira, L. and Gilmartin, W. (2009). "Short-Range Movements
225 of Hawksbill Turtles (*Eretmochelys imbricata*) from Nesting to Foraging Areas within the
226 Hawaiian Islands 1." Pacific Science **63**(3): 371-382.
- 227 Sterling, E.J., McFadden, K.W., Holmes, K.E., Vintinner, E.C., Arengo, F. and Naro-Maciel, E. (2013).
228 "Ecology and Conservation of Marine Turtles in a Central Pacific Foraging Ground." Chelonian
229 Conservation and Biology **12**(1): 2-16.

231

Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: