S2 Appendix Diagnostic plots from fitted VAST models




Fig AA. QQ residual plots for arrowtooth flounder, darkblotched rockfish, Dover sole, English sole, Pacific grenadier, and lingcod.




Fig BB. QQ residual plots for longspine thornyhead, Pacific sanddab, petrale sole, Pacific hake, sablefish, sharpnose rockfish, and shortspine thorryhead.




Fig CC. Spatial residuals for occurrence for arrowtooth flounder.






Fig DD. Spatial residuals for occurrence for darkblotched rockfish.



 $\label{eq:Fig-EE} \textbf{Fig-EE. Spatial residuals for occurrence for Dover sole.}$



 $Fig\ FF.\ Spatial\ residuals\ for\ occurrence\ for\ English\ sole.$




Fig GG. Spatial residuals for occurrence for Pacific greenling.




Fig HH. Spatial residuals for occurrence for Pacific hake.




Fig II. Spatial residuals for occurrence for Lingcod.




Fig JJ. Spatial residuals for occurrence for longspine thornyhead.




Fig KK. Spatial residuals for occurrence for Petrale sole.





Fig LL. Spatial residuals for occurrence for sablefish.



FigM M. Spatial residuals for occurrence for Pacific sanddab.




Fig NN. Spatial residuals for occurrence for sharpnose rockfish.




Fig O. Spatial residuals for occurrence for shortspine thornyhead.