

1 GLOBAL CONSERVATION STATUS OF THE WORLD'S MOST PROMINENT FORAGE
2 FISHES (TELEOSTEI: CLUPEIFORMES)

3
4 Birge, T.L.^{a,b}, Ralph, G.M.^{a,b}, Di Dario, F.^c, Munroe, T.A.^d, Bullock, R.W.^{b,e} Maxwell, S.M.^f, Santos,
5 M.D.^g, Hata, H.^h, Carpenter, K.E.*^{a,b}

- 6 a. IUCN Marine Biodiversity Unit, Biological Sciences, Old Dominion University, Norfolk,
7 VA, USA
8 b. Member of IUCN Species Survival Commission
9 1. The views expressed in this publication do not necessarily reflect those of IUCN.
10 2. The designation of geographical entities in this paper, and the presentation of the
11 material, do not imply the expression of any opinion whatsoever on the part of IUCN
12 concerning the legal status of any country, territory, or area, or of its authorities, or
13 concerning the delimitation of its frontiers or boundaries.
14 c. Universidade Federal do Rio de Janeiro (UFRJ), Instituto de Biodiversidade e
15 Sustentabilidade (NUPEM). CP 119331, CEP 27910-970, Macaé, RJ, Brazil
16 d. National Systematics Laboratory, NMFS/NOAA/OS&T, National Museum of Natural
17 History, Smithsonian Institution, Washington, D.C., USA
18 e. The Deep, Tower Street, Hull HU1 4DP. England
19 f. School of Interdisciplinary Arts and Science, University of Washington, Bothell, Bothell,
20 WA USA
21 g. National Fisheries Research and Development Institute, 101 Mother Ignacia St., Quezon
22 City 1103 Philippines
23 h. Center for Molecular Biodiversity Research, National Museum of Nature and Science,
24 Tsukuba, 305-0005 Japan

25 *Corresponding Author: Kent E. Carpenter, +1 (757) 683-4197, kcarpent@odu.edu, IUCN Marine
26 Biodiversity Unit, Biological Sciences, Old Dominion University, Norfolk, VA 23529-0266, USA.

27 **HIGHLIGHTS**

- 28 - Clupeiforms have the lowest percentage of elevated conservation concern species
29 (11%), but the highest percentage of species evaluated as Data Deficient (28%),
30 compared to other fish groups assessed.
31 - Global species richness of Clupeiforms is highest in the Indo-Malay- Philippine
32 Archipelago and the Caribbean.
33 - Major threats include overexploitation, pollution, and habitat modifications from dams.
34 - Increased and improved fisheries management measures and intensive habitat
35 restoration is urgent.

36

37 **ABSTRACT**

38 Understanding the extinction risk of taxonomic groups increases our ability to prioritize
39 efforts to address biodiversity loss. Over 400 species of herrings, shads, sardines, anchovies,
40 menhadens, and relatives belong to the Order Clupeiformes and include many of the most
41 important forage fishes. These small, schooling fishes are ecologically, economically, and
42 culturally significant. However, despite their global contribution to fisheries and our increasing
43 reliance on them for food and modern commodities, we lack critical information regarding basic
44 biology and population trends for most species. We applied the IUCN Red List methodology, a
45 comprehensive and systematic approach to assess extinction risk, to all clupeiform species. The
46 best estimate suggests nearly 11% of species are of elevated conservation concern, although
47 this could be as high as 36%. Two regions, the Caribbean and the Indo-Malay-Philippine
48 Archipelago have high concentrations of threatened and Data Deficient species and are areas of
49 conservation concern. Major threats include overexploitation, pollution, and habitat
50 modification. Immediate conservation priorities include: 1) increasing research and mitigative
51 action directed toward species assessed as threatened or Data Deficient; 2) improving fisheries
52 management regulations for the understudied but heavily exploited species, and 3) promoting
53 local, intensive habitat restoration to reduce pollution and remove dams. These extinction risk
54 assessments and subsequent analyses should be used as an informative tool for fisheries and
55 conservation managers and to monitor conservation progress.

56

57 Keywords: Extinction, baitfish, Red List, threatened

58

59 **1. INTRODUCTION**

60 Forage fishes are a crucial link between primary production and keystone predators in
61 aquatic environments (Pikitch et al., 2014). These typically abundant small- to medium-sized
62 pelagic species feed at the base of the food web and serve as a predominate prey source for
63 numerous larger predators, such as piscivorous fishes, mammals, squids, and seabirds, many of
64 which are commercially important (Cury et al., 2011; Smith et al., 2011; Pikitch et al., 2014;
65 Hilborn et al., 2017). Forage fishes include a diverse array of bony fishes and invertebrates such
66 as krill and squid (Pikitch et al., 2014; Rountos, 2016). Many species also support the global
67 economy by directly and indirectly sustaining several fisheries (Pikitch et al., 2014) and
68 contribute 20-30% to the annual global marine catch (Alder et al., 2008; Smith et al., 2011,
69 Pikitch et al., 2014).

70 Species of the Clupeiformes (Teleostei), commonly known as herrings, shads,
71 menhadens, sardines, anchovies, and their relatives, are a major component of forage fishes in
72 coastal ecosystems and dominate worldwide forage fish landings (Tacon and Metian, 2009a;
73 FAO, 2020). Additional to providing ecological and economic support, clupeiforms contribute to
74 global food security given their abundance, easy access, and exceptionally high nutrient content
75 (FAO, 2018). In some human communities, clupeiforms comprise the major or the sole protein
76 source (Alder et al., 2008; Mohanty et al., 2019). Historically, clupeiform presence has been
77 associated with persistent human settlement, growth, and survival (e.g., Bloch, 1809; Thornton
78 et al., 2010; Levin et al., 2016). To meet the needs of a rising human population (United
79 Nations, Department of Economic and Social Affairs, Population Division, 2017), demand for
80 fisheries resources is expected to continue growing (FAO, 2018). Given the overall ecological,
81 cultural, nutritional, and economic importance of clupeiforms worldwide, their conservation
82 status warrants greater attention.

83 The Clupeiformes includes 415 species that are globally distributed with tropical,
84 temperate, and sub-Arctic representatives (Whitehead, 1985; Whitehead et al., 1988; Lavoué et
85 al., 2013). Clupeiform fishes are ecologically diverse and span all aquatic habitats, including
86 coastal and open marine environments, oceanic islands, estuaries, and freshwater rivers and
87 lakes (Whitehead, 1985; de Pinna and Di Dario, 2003; Lavoué et al., 2013; Bloom and Egan,
88 2018). Species can be restricted to marine, estuarine, or fresh waters, or they can be
89 euryhaline, where a subset exhibit diadromy (Whitehead, 1985). Strictly marine clupeiforms
90 (33.7% of all species) are distributed in every ocean, except for the Southern Ocean
91 (Whitehead, 1985), while strictly freshwater species (17.8% of all species) are found on every
92 continent except for Antarctica (Bloom and Lovejoy, 2012, 2014; Bloom and Egan, 2018).

93 Despite the global importance of clupeiforms, basic biological information, fisheries
94 data, and management efforts are severely deficient compared to those of other commercially
95 important fishes, such as tunas and billfishes. This disparity may be due in part to perception of
96 extinction resistant traits or may result from the taxonomic complexity of clupeiforms
97 (Whitehead, 1985; Alder et al., 2008). Value per pound for clupeiforms is also far less than that
98 for other commercial fishes, which may further disincentivize the contribution of resources to
99 research and conservation for the clupeiforms. For example, the average commercial landed

100 value of all tunas in the U.S. for 2017 was about USD \$2.8/pound, while the average value for
101 clupeiforms was roughly USD \$0.09/pound (NOAA Fisheries, 2019). The paradox between
102 worldwide clupeiform importance and lack of available resources and reliable data reinforces
103 the need to invest effort into understanding the current conservation status of the members of
104 this group.

105 The International Union for Conservation of Nature (IUCN) Red List of Threatened
106 Species provides a key starting point for highlighting and addressing conservation needs for
107 species (Mace et al., 2008). The IUCN Red List, an open-access repository of species-specific
108 assessments, categorizes a species conservation status by interpreting its risk to extinction
109 (Rodrigues et al., 2006; Vié et al., 2009). Red List assessments are the most widely accepted
110 standard for species-level risk evaluations (Hoffman et al., 2008). By illuminating gaps in
111 conservation knowledge for species, assessments can be used to inform and influence decisions
112 regarding biodiversity conservation (Rodrigues et al., 2006; Mace et al., 2008; Vié et al., 2009).

113 Limited species-specific conservation information on clupeiforms hampers our ability to
114 proactively manage and conserve these essential components of aquatic food webs. To address
115 this gap, we applied the IUCN Red List methodology to assess the extinction risk of the 415 valid
116 clupeiform species. The assessments and accompanying data were used to evaluate: 1)
117 variability in the proportion of species at an elevated risk of extinction as a function of family,
118 and habitat; 2) major threats; and 3) spatial patterns in species richness. These analyses provide
119 a baseline from which to monitor changes in conservation status and are used to identify
120 conservation priorities and research needs.

121 **2. METHODS**

122 *2.1 Taxonomic Scope*

123 Phylogenetic relationships among the main groups of the Clupeiformes are contentious,
124 resulting in different proposals of taxonomic classifications (e.g., Di Dario, 2002, 2009;
125 Miyashita, 2010; Lavoué et al., 2014). Overall, the order is divided into the Denticipitoidei, with
126 a single living representative (*Denticeps clupeoides*) in the Denticipitidae, and the Clupeoidei
127 (Grande, 1985), which includes all remaining 414 species of the Clupeiformes assessed here.
128 The Clupeoidei has been traditionally divided into four families: the Chirocentridae,
129 Pristigasteridae, Engraulidae, and the Clupeidae (e.g., Whitehead, 1985). However,
130 morphological characters and molecular evidence indicates that the Clupeidae, which includes
131 about half of all currently valid species of the Clupeiformes (Nelson et al., 2016), is not a
132 monophyletic group (summarized in Lavoué et al., 2014). To partially acknowledge that, we
133 provisionally accept the classification of Lavoué et al. (2014), which includes *Dussumieria*,
134 *Etrumeus*, *Spratelloides*, and *Jenkinsia*, in a distinct family (Dussumieriidae). *Sundasilanx* is a
135 paedomorphic genus of freshwater clupeiforms of unknown relationships within the Clupeoidei
136 (Siebert, 1997; Lavoué et al., 2013, 2014). The genus is generally regarded as a member of the
137 Clupeidae (Siebert, 1997; Lavoué et al., 2014; Nelson et al., 2016), but given its controversial
138 position in the Clupeoidei, the Sundasalangidae is also provisionally recognized as a distinct
139 family (Van der Laan et al. 2014). In this arrangement, the Clupeidae includes 188 species.
140 Therefore, for the purposes of this study, seven families of the Clupeiformes are recognized:

141 Denticipitidae (1 species), Pristigasteridae (37 species), Engraulidae (161 species),
142 Chirocentridae (2 species), Clupeidae (188 species), Dussumieriidae (19 species), and
143 Sundasalangidae (7 species).

144 *2.2 Quantifying Extinction Risk*

145 We compiled a species list based on the online version of the Catalog of Fishes up to
146 March 2020 (Fricke et al. 2020) and in consultation with taxonomic experts. Individual
147 clupeiform species' assessments were based on available information from peer-reviewed and
148 grey literature regarding geographic distribution, population status, life history, utilization and
149 quality of habitat, potential threats, and known conservation measures. The assessment
150 process included involvement from 132 international experts from more than 20 countries with
151 regional or species expertise. We identified potential experts to be involved in the assessments
152 from the authors of peer-reviewed publications, FAO fisheries identification guides, and
153 through the IUCN Species Survival Commission network. All 415 species were assessed against
154 the IUCN Red List criteria (Mace et al., 2008; IUCN 2012) at workshops and through online
155 collaborations. Draft assessments go through multiple rounds of review by species experts and
156 the Red List process prior to publication. As of July 2020, all species assessments included in this
157 analysis are published on the Red List website (www.iucnredlist.org).

158 The IUCN Red List includes eight global levels of extinction risk: Extinct (EX), Extinct in
159 the Wild (EW), Critically Endangered (CR), Endangered (EN), Vulnerable (VU), Near Threatened
160 (NT), Least Concern (LC), and Data Deficient (DD: IUCN, 2012). A taxon is considered EX when
161 there is no reasonable doubt the last individual has died or is EW if it is only known to survive in
162 cultivation, captivation, or in naturalized populations outside of its historic range (IUCN 2012). A
163 species can reasonably be presumed EX or EW when exhaustive surveys fail to report it (IUCN
164 2012). To qualify for a threatened category (CR, EN, VU) a species must meet at least one of the
165 five quantitative thresholds under IUCN Criteria (A – E: Mace et al., 2008). The criteria evaluate
166 population decline (A), restricted geographic distribution (B), small population size and decline
167 (C), very small or restricted population size (D), and the high probability of potential extinction
168 (E: Akçakaya et al., 2000; Mace et al., 2008).

169 For each assessment, experts evaluated the species-specific data available against all
170 five Red List Criteria (IUCN, 2001; 2012). Almost all species were assessed under criteria A
171 (population decline) or B (restricted geographic range). Data required to assess a species under
172 the remaining criteria (C, D or E) were often unavailable. Species were assigned to the highest
173 threat category for which the available data met or exceeded the associated thresholds and
174 conditions (IUCN, 2001; 2012; IUCN Standards and Petitions Subcommittee, 2016). A category
175 of NT was applied if the quantified estimates of population decline or geographic range size
176 nearly meet the thresholds for assigning a threatened category under at least one of the
177 criteria. A species was listed as LC if it did not qualify for a threatened or NT listing based on the
178 available data. Finally, the DD category was applied if a species is known from few specimens,
179 lacks information to assess under any of the criteria, or there is uncertainty regarding its
180 taxonomic status (IUCN, 2001; 2012; IUCN Standards and Petitions Subcommittee, 2016). This

181 category was also applied if declines were likely due to a known but unquantified threat (e.g.,
182 fishing pressure), such that a more appropriate category could not be assigned.

183 Direct threats impacting each species were identified from the published literature
184 (peer-reviewed and grey), verified by species or regional experts, and categorized within each
185 species assessment using the standardized IUCN Threat Classification Scheme (version 3.2:
186 IUCN-CMP, 2016). These coded threats and full bibliographies are available as part of the
187 assessment for each species. Major threats were summarized across species as a function of
188 primary habitat system (marine, euryhaline, freshwater). The proportion of species listed as
189 threatened (CR, EN, VU) and NT, herein referred to as species of elevated concern, was also
190 explored as a function of family and major habitat system. The proportion of species of
191 elevated concern is expressed using both a midpoint and a range to address the uncertainty
192 surrounding the true status of DD species. The midpoint was calculated by removing the species
193 listed as DD, whereas the lower and upper bounds were calculated by excluding or including
194 the DD species with the threatened and NT, respectively. The lower boundary assumes that
195 none of the DD species are of an elevated concern, while the upper boundary assumes that all
196 DD species are of an elevated concern (IUCN, 2016).

197 A species was assigned a major habitat category using the information in the Red List
198 assessments. Given the known or suspected tolerance for salinity fluctuations exhibited by
199 many clupeiforms, we modified the IUCN Red List system classification scheme from two
200 aquatic categories (freshwater, including inland estuarine waters; and marine, including coastal
201 estuarine waters) to three categories. Therefore, the freshwater system includes those species
202 known to occupy only freshwater environments and the marine system includes species
203 restricted to marine waters. The third, euryhaline category includes estuarine species,
204 diadromous species, and species known or suspected to tolerate changes in salinity.

205 *2.3 Distribution Maps and Spatial Analyses*

206 Maps were created for each species using ArcMap 10.3 based on occurrence records,
207 habitat preferences, and depth limits and were reviewed by species experts. As marine
208 clupeiforms are primarily coastal, the distribution polygons for strictly marine species were
209 standardized using a base map that represents either the 200 m bathymetric line or 100 km
210 from the shore, whichever was further from the coast. Bathymetric layers were extracted from
211 two global level sources, the National Geophysical Data Center's ETPO1 (Amante and Eakins,
212 2009) and the General Bathymetric Chart of the Oceans (GEBCO: IOC et al., 2003). Maps for
213 freshwater species were created using hydrobasins, because these areas are considered as
214 minimum management units for freshwater conservation (Lévêque et al., 2008; Carrizo et al.,
215 2013). For species that utilize both marine and freshwater habitats (e.g., diadromous species),
216 maps separately followed the marine and freshwater protocols, and were combined to
217 encompass the entirety of the species' range.

218 Global maps of overall species richness, DD richness, and richness of elevated concern
219 species were also created using ArcMap 10.3. Species with a freshwater distribution were
220 summarized within the Global HydroBASINS (Lehner and Grill, 2013), using the largest river
221 basins of each continent. Species with a marine distribution were summarized within the

222 Marine Ecosystems of the World at the province level (Spalding et al., 2007). This shapefile was
223 modified to include a region for the Caspian Sea, as it is excluded from the Global HydroBASINS
224 and Marine Ecosystems of the World.

225 **3. RESULTS**

226 *3.1 Global IUCN Red List Status of Clupeiforms*

227 The best estimate of the proportion of clupeiforms of elevated concern is 11%. Given
228 the uncertainty of an appropriate Red List Category for all DD species, the true proportion of
229 elevated concern species could lie between 8 – 36%. Of all species (n = 415), three (0.7%) are
230 listed as CR, 11 (2.7%) as EN, 13 (3.1%) as VU, and five (1.2%) as NT. No species were listed as
231 EX or EW. Species are primarily listed as elevated concern either due to a restricted range size
232 with an ongoing threat (criterion B; n = 17) or due to population decline (criterion A; n = 10);
233 two species (*Sardinella tawilis* and *Alosa vistonica*) are listed under both criteria A and B. Three
234 species are listed as VU given a very restricted range and a serious plausible future threat
235 (criterion D). Of the remaining 383 species, 267 (64.3%) are categorized as LC, and 116 (28.0%)
236 are considered DD.

237 Among families of the Clupeiformes, the Denticipitidae consists of only one species,
238 *Denticeps clupeoides*, which is listed as VU. As such, this family has the highest proportion of
239 elevated concern species overall (Fig. 1). Excluding *D. clupeoides*, the Clupeidae has the highest
240 proportion of elevated concern species (25 of 188 species; midpoint = 16.7%), followed by the
241 Engraulidae (5 of 161 species; midpoint = 4.9%), and the Pristigasteridae (1 of 37 species;
242 midpoint = 3.8%). None of the Chirocentridae (n = 2), Dussumieriidae (n = 19) or
243 Sundasalangidae (n = 7) are listed as threatened. However, the high proportion of DD species,
244 especially within the Sundasalangidae, may be obscuring the actual conservation status of these
245 families.

246 Species classified as euryhaline (i.e., diadromous or estuarine) constituted nearly half of
247 all clupeiforms (n = 201; 48.4%), followed by marine (n = 140; 33.7%) and freshwater species (n
248 = 74; 17.8%) (Fig. 2). Euryhaline habitats harbor the largest proportion of LC species (n = 147;
249 73.1%) followed by marine habitats (n = 80; 57.1%), and then freshwater habitats (n = 40;
250 54.1%). Despite having the lowest number of representatives, freshwater clupeiforms have the
251 highest proportion of elevated concern species (16 of 74 species; midpoint = 28.6%), more than
252 three times the proportion in marine environments (7 of 140 species; midpoint = 8.0%), and
253 four times the proportion of elevated concern species found in euryhaline environments (9 of
254 201 species; midpoint = 5.7%). Additionally, all species assessed as CR (n = 3), the highest threat
255 level, are found in freshwater habitats.

256 *3.2 Major threats*

257 Of the 415 species, 144 have at least one identified threat. The remaining 271 species
258 have either no major threats causing significant impacts, or threats to these species are
259 unknown. The most prominent threat to clupeiforms in all habitats is overexploitation,
260 impacting 107 species overall (Fig. 3). Pollution and natural system changes (e.g., dams) impact
261 nearly the same number of species (47 and 42, respectively). However, despite having the

262 highest proportion of LC species (Fig. 2), the majority of species impacted by pollution or
263 natural system changes are euryhaline (Fig. 3). Of the species impacted by at least one threat,
264 roughly the same proportions of freshwater and euryhaline species are impacted by pollution
265 and natural system changes overall (84 and 76%, respectively). The proportion of marine
266 species impacted by climate change (36%) is more than two times the proportion of euryhaline
267 (11%) and freshwater (6%) species, while invasive species impact a higher proportion of
268 freshwater species (18%) relative to the proportion of marine and euryhaline species (11% and
269 7%, respectively).

270 *3.3 Spatial Analyses*

271 Global species richness of clupeiforms follows two distribution patterns; a longitudinal
272 gradient, where the highest tropical richness is within the Indo-West Pacific, and a latitudinal
273 gradient where richness decreases with increasing distance from the tropics. The highest
274 species richness of all clupeiforms is along coastal India and throughout the Indo-West Pacific
275 from the eastern Andaman Sea, east to the Philippines, Indonesia, and northeastern Papua New
276 Guinea (Fig. 4A). High richness also occurs in the central eastern Pacific from Mexico to
277 northern Peru, and the central western Atlantic from the greater Caribbean to northern Brazil.
278 Areas of lowest species richness are within the northern and southernmost limits of the global
279 range for clupeiforms (e.g., the Arctic and north of the Southern Ocean), in inland rivers, and off
280 Polynesian Islands.

281 In general, DD species richness closely follows that of the total species richness (Fig. 4B).
282 However, DD species richness is higher in northern Australian rivers relative to the total species
283 richness. In contrast, the high species richness in Europe, eastern United States, and South
284 American rivers is not mirrored by high DD species richness.

285 Conversely, the highest richness of species of elevated concern ($n = 32$) occurs within
286 the greater Caribbean (Fig. 4C). Other areas of high richness for species of elevated concern are
287 along the western Pacific continental coast (Russia south to Indonesia), and inland areas
288 including the Caspian Sea and the Congo River in Central Africa. A low richness of elevated
289 concern species is scattered along regions such as the northeastern United States, the eastern
290 and southern coasts of South America, western Africa, and parts of Europe and Asia.

291 **4. DISCUSSION**

292 Major threats to clupeiforms are similar to those found for other groups of fishes (e.g.,
293 Roberts and Hawkins, 1999; Reynolds et al., 2005; Dulvy et al., 2009; Harnik et al., 2012), with
294 overexploitation as the leading threat for all clupeiforms in all habitats. While overexploitation
295 may be the most prolific threat by impacting the highest number of clupeiforms, pollution may
296 be the most detrimental, as it affects greater numbers of CR species. When compared to other
297 economically and ecologically important fish groups globally assessed using the IUCN Red List
298 methodology, clupeiforms have the lowest estimated percentage of threatened and NT species
299 overall. Using the midpoint of species evaluated as elevated concern, roughly 11% are currently
300 at high risk compared to approximately 22% of tunas and billfishes (Collette et al., 2011), 19%

301 of sparids (Comeros-Raynal et al., 2016), and 19% of groupers (Sadovy de Mitcheson et al.,
302 2020).

303 The lower proportion of threatened species in clupeiforms may be a function of
304 uncertainty and is likely an underestimate of the true conservation status for many of these
305 species. The high percentage of DD clupeiforms (28%) surpasses that of the tunas and billfishes
306 (Collette et al., 2011), sparids (Comeros-Raynal et al., 2016), and groupers (Sadovy de
307 Mitcheson et al., 2020), each with less than 20% of those species evaluated as DD. A DD listing
308 is most often related to taxonomic uncertainty, low number of known specimens, unknown
309 geographical range, or inability to quantify a threat or decline in population (IUCN, 2012), all of
310 which are common within the Clupeiformes. Continued taxonomic research will likely identify
311 additional cryptic species (e.g., recent revisions of species of *Sardinella*, *Stolephorus* and
312 *Encrasicolina* – Thomas et al., 2014; Hata and Motomura 2019a,b,c), clarifying our current
313 understanding of the complex taxonomy and biodiversity of this group and influencing the
314 assessments of some species.

315 If the DD species were evenly distributed relative to total richness, we would expect that
316 all areas would have about 28% DD species. Instead, we found high variation in both the
317 numbers and proportion of DD species. For example, a few freshwater river basins (in eastern
318 and northwestern Africa; southern U.S. and northern Mexico; and Borneo) are, or are nearly
319 100% DD. However, these areas are characterized by low clupeiform richness, with only one or
320 two species occurring in each of these regions. The highest number of clupeiform species
321 evaluated as DD generally coincides with geographic areas of both high clupeiform biodiversity
322 and areas of low per-capita income. For example, the Coral Triangle is the epicenter of marine
323 biodiversity (Carpenter and Springer 2005; Sanciangco et al., 2013) and is a hotspot for
324 clupeiform species (up to 81), which are heavily relied on for subsistence in local fisheries.

325 In general, global biodiversity is unevenly distributed; the most biodiverse places are
326 often areas of high human populations of relatively low per capita income (Baille et al., 2004;
327 Brooks et al., 2006) and tend to have the highest number of threatened species (Hoffmann et
328 al., 2010; Baille et al., 2004). Countries with high human populations and high biodiversity are
329 less likely to have financial resources available for research and conservation purposes (Baille et
330 al., 2004), and may rely more heavily on local marine resources for livelihood (Creel 2003;
331 Ferrol-Schulte et al., 2015). In contrast, countries such as those in the advanced economies of
332 Europe invest substantially in conservation research and management and have few globally
333 threatened species (Baille et al., 2004), including those among the clupeiforms where both the
334 number and proportion of threatened and DD are very low.

335 In many parts of the world, particularly in highly biodiverse areas, clupeiform stock
336 assessments and fishery effort data are lacking or are unreported. Where data are available, it
337 is often in the form of raw fishery landings (FAO, 2016) or reconstructed catches (Pauly and
338 Zeller, 2016a). These landings frequently aggregate several species because those that co-occur
339 often school together and are difficult to identify (e.g., species of sardines and anchovies:
340 Bakun and Cury, 1999). Teasing apart landings from multi-species fisheries is a difficult task and
341 identifications that contain many errors can lead to false estimations of species-specific catch

342 data (Gaichas et al., 2012). Overexploitation is a major threat to over 25% of clupeiform
343 species, but this likely underestimates the impact given uncertainties in landings and the
344 population status of species evaluated as DD. Clupeiforms also contribute to many unreported
345 artisanal fisheries (Whitehead, 1985; Whitehead et al., 1988), represent a significant portion of
346 bycatch in other industrial trawl fisheries (e.g., Stobutzki et al., 2001), and are taken in illegal,
347 unreported and unregulated fisheries (IUU: Agnew et al., 2009). Accidental and IUU fishing,
348 along with aggregated landings, adversely affect our ability to quantify global fishing pressure
349 on these species. It can further impact conclusions drawn regarding population trends by
350 underestimating true catches (Pauly and Zeller, 2016b), which ultimately impacts the efficacy of
351 conservation or management decisions.

352 The highest concentration of threatened species is centered in the Caribbean region;
353 however, the highest species richness overall and of DD species is concentrated in the central
354 Indo-West Pacific region. Therefore, only about one-tenth of the Caribbean species are
355 assessed as DD compared to roughly one-third of Indo-West Pacific species, highlighting our
356 increased knowledge of Caribbean species. Currently, clupeiforms in the Caribbean would
357 benefit most from threat mitigation, while emergent research to fill in our knowledge gaps in
358 the Indo-West Pacific region should be prioritized. As more data become available to
359 adequately assess species currently listed as DD, it is likely that we may find a higher proportion
360 of elevated concern species within the Indo-West Pacific, relative to that reported from the
361 Caribbean.

362 In addition to the high proportion of DD species, traditional perceptions of intrinsic life
363 history traits have impeded the conservation of clupeiforms. Their typically high fecundity,
364 multiple spawning, and early age of maturation are regarded as resilience factors, even though
365 these traits often do not reflect lower vulnerability to extinction (Jennings et al., 1998;
366 Kindsvater et al., 2016; Sadovy, 2001; Juan-Jorda et al., 2012; Comeros-Raynal et al., 2016). For
367 example, the widely distributed Pacific herring (*Clupea pallasii*) is exploited to varying degrees
368 throughout a large portion of its range. In some regions where this species has experienced
369 drastic declines, subpopulations have not recovered even decades after fishing pressure has
370 ceased (see Hay et al., 2001 for description of Yellow Sea and Hokkaido – Sakhalin herring).
371 Overall, intrinsic life history characteristics of many clupeiforms and likely other important
372 forage groups may provide a buffer against extinction (compared to long-lived taxa such as
373 sharks, rays, tunas, billfishes, and groupers), but this buffer does not hold for all clupeiform
374 subpopulations.

375 Synergistic influences of threats can be detrimental to the survival of a population
376 (Brook et al., 2008). Often, freshwater and euryhaline clupeiforms are threatened by both
377 pollution and natural system modifications, indicating a potential for increased cumulative
378 effects. Many anadromous representatives in genera such as *Alosa* and *Tenualosa* appear to be
379 most negatively impacted by one or both threats (e.g., Freyhof and Kottelat, 2008a;
380 NatureServe, 2013; Di Dario, 2018; Mohd Arshaad et al., 2018). In line with previous studies of
381 other freshwater fishes (e.g., Collen et al., 2014), freshwater clupeiforms have roughly four
382 times the proportion of elevated concern species compared with that among marine and
383 euryhaline representatives within the group. Given that all species listed as CR are freshwater

384 clupeids, the responses to multiple stresses by all freshwater clupeids should be examined
385 more closely. Additionally, the freshwater denticle herring, *Denticeps clupeoides*, is the only
386 member of the Denticipitoidei, a very distinct and presumably old (ca. 126–121 Mya) lineage of
387 clupeiform fishes (Malabarba and Di Dario, 2017). This species is a relict that inhabits a few
388 isolated coastal streams of West Africa (Teugels, 2003), a region heavily impacted by
389 agricultural and urban developments (Lalèyè et al., 2010). Immediate implementation of
390 strategies aimed at the conservation of *D. clupeoides* and other threatened freshwater
391 clupeiforms is highly recommended.

392 **5. CONCLUSION**

393 Despite the relatively lower percentage of threatened species compared to that of other
394 fish groups of similar economic value, the overall ecological importance of clupeiform fishes
395 and their ubiquity as an essential fishery resource warrants conservation concern. At a local
396 level, species with limited ranges, such as *Alosa killarnensis*, *Denticeps clupeoides*, and
397 *Sardinella tawilis*, may require stringent protection and improvement of habitat quality
398 (Freyhof and Kottelat, 2008b; Lalèyè et al., 2010; Santos et al., 2018). Additionally, though some
399 species threatened with overexploitation have localized management and monitoring in place,
400 such as *Sardinella lemuru* in the southern Philippines (Rola et al., 2018), the efficacy of current
401 measures need to be evaluated. An increase in species-specific landings and catch statistics,
402 coupled with effort data, would also further improve future assessments of exploited species,
403 especially in developing countries. Large-scale industrial fisheries, such as those for the
404 Peruvian anchoveta (*Engraulis ringens*) and the Pacific herring (*Clupea pallasii*), may benefit
405 from increased multi-national cooperative regulations. Species of elevated conservation
406 concern are also potential targets for improved and more stringent monitoring. Given the
407 limited resources available, research and conservation prioritization can be difficult in areas of
408 high biodiversity; however, mitigation of anthropogenic stressors in these areas where elevated
409 concern species are distributed is critical. Fishery managers and funding agencies in regions
410 with large proportions of exploited DD species may also consider prioritizing research initiatives
411 to fill gaps in our understanding of these species.

412 **ACKNOWLEDGEMENTS**

413 This project was funded by the Toyota Motor Corporation with additional funding and
414 workshop logistical support provided by the Philippines' Bureau of Fisheries and Aquatic
415 Resources. This research was conducted as part of the master's thesis of T. Birge. The
416 statements made herein are solely the responsibility of the authors. We thank Christi Linardich
417 for contributions to data collection and workshop facilitation. This project could not have been
418 completed without the 132 species and regional experts who contributed to the completion
419 and review of the Red List assessments. We thank the IUCN Red List Unit, particularly Janet
420 Scott, for reviewing the Red List assessments. Additionally, we thank two anonymous reviewers
421 for their comments that improved this manuscript.

422

423 **REFERENCES**

- 424 Agnew, D.J., Pearce, J., Pramod, G., Peatman, T., Watson, R., Beddington, J.R., Pitcher, T.J.,
425 2009. Estimating the worldwide extent of illegal fishing. PLoS ONE 4. doi:
426 10.1371/journal.pone.0004570.
427
- 428 Akçakaya, H.R., Ferson, S., Murgman, M.A., Keith, D.A., Mace, G.M., Todd, C.R., 2000. Making
429 consistent IUCN classifications under uncertainty. Conservation Biology 14, 1001-1013. doi:
430 10.1046/j.1523-1739.2000.99125.x.
431
- 432 Alder, J., Campbell, B., Karpouzi, V., Kaschner, K., Pauly, D., 2008. Forage Fish: From Ecosystems
433 to Markets. Annual Review of Environment and Resources 33, 153-166. doi:
434 10.1146/annurev.enviro.33.020807.
435
- 436 Amante, C., Eakins, B.W., 2009. ETOPO1 1 Arc-Minute Global Relief Model: Procedures, Data
437 Sources and Analysis NOAA Technical Memorandum NESDIS NGDC-24.
438 doi:10.7289/V7285C8276M: National Geophysical Data Center, NOAA.
439
- 440 Baille, J. Hilton-Taylor, C., Stuart, S.N., 2004. 2004 IUCN Red List of Threatened Species: A Global
441 Species Assessment. IUCN, Gland, Switzerland and Cambridge, UK.
442
- 443 Bakun, A., Cury, P., 1999. The 'school trap': A mechanism promoting large-amplitude out-of-
444 phase population oscillations of small pelagic fish species. Ecology Letters 2, 349-351. doi:
445 10.1046/j.1461-0248.1999.00099.x.
446
- 447 Bloch, M., 1809. Natural history of the herring. The Belfast Monthly Magazine 2, 241-245.
448
- 449 Bloom, D.D., Egan, J.P., 2018. Systematics of Clupeiformes and testing for ecological limits on
450 species richness in a trans-marine/freshwater clade. Neotropical Ichthyology 16, e180095. doi:
451 10.1590/1982-0224-20180095.
452
- 453 Bloom, D.D., Lovejoy, N.R., 2012. Molecular phylogenetics reveals a pattern of biome
454 conservatism in New World anchovies (family Engraulidae). Journal of Evolutionary Biology 25,
455 701 - 775. doi: 10.1111/j.1420-9101.2012.02464.x.
456
- 457 Bloom, D.D., Lovejoy, N.R., 2014. The evolutionary origins of diadromy inferred from a time-
458 calibrated phylogeny for Clupeiformes (herring and allies). Proceedings of the Royal Society B
459 281, 20132081. doi: 10.1098/rspb.2013.2081.
460
- 460 Brook, B.W., Sodhi, N.S., Bradshaw, C.J.A., 2008. Synergies among extinction drivers under
461 global change. Trends in Ecology and Evolution 23, 453 – 460. doi: 10.1016/j.tree.2008.03.011.
462
- 462 Brooks, T.M., Mittermeier, R.A., da Fonseca, G.A.B., Gerlach, J., Hoffmann, M., Lamoreux, J.F.,
463 Mittermeier, C.G., Pilgrim, J.D., Rodrigues, A.S.L., 2006. Global biodiversity conservation
464 priorities. Science 313, 58-61. doi: 10.1126/science.1127609.

465
466 Carpenter, K.E., Springer, V.G., 2005. The center of the center of marine shore fish biodiversity:
467 the Philippine Islands. *Environmental Biology of Fishes* 72, 467 – 480.
468
469 Carpenter, K.E., Abrar, M., Aeby, G., Aronson, R.B., Banks, S., Bruckner, A., . . . Wood, E., 2008.
470 One-third of reef-building corals face elevated extinction risk from climate change and local
471 impacts. *Science* 321, 560-563. doi: 10.1126/science.1159196.
472
473 Carrizo, S.F., Smith, K.G., Darwall, W.R.T., 2013. Progress towards a global assessment of the
474 status of freshwater fishes (Pisces) for the IUCN Red List: application to conservation
475 programmes in zoos and aquariums. *International Zoo Yearbook* 47, 46 – 64. doi:
476 10.1111/izy.12019.
477
478 Collen, B., Whitton, F., Dyer, E.E., Baillie, J.E.M., Cumberlidge, N., Darwall, W.R.T., Pollock, C.,
479 Richman, N.I., Soulsby, A.-M., Böhm, M., 2014. Global patterns of freshwater species diversity,
480 threat and endemism. *Global Ecology and Biogeography* 23, 40 – 51. doi: 10.1111/geb.12096.
481
482 Collette, B.B., Carpenter, K.E., Polidoro, B.A., Juan-Jordá, M.J., Boustany, A., Die, D.J., ... Yáñez,
483 E., 2011. High value and long life-double jeopardy for tunas and billfishes. *Science* 333, 291-292.
484
485 Comeros-Raynal, M.T., Polidoro, B.A., Broatch, J., Mann, B.Q., Gorman, C., Buxton, C.D.,
486 Goodpaster, A.M., Iwatsuki, Y., MacDonald, T.C., Pollard, D., Russell, B., Carpenter, K.E., 2016.
487 Key predictors of extinction risk in sea breams and porgies (Family: Sparidae). *Biological*
488 *Conservation* 202, 88 – 98. doi: 10.1016/j.biocon.2016.08.027.
489
490 Creel, L., 2003. Ripple effects: population and coastal regions. Population Reference Bureau.
491 Washington, D.C. 1 – 7.
492
493 Cury, P.M., Boyd, I.L., Bonhommeau, S., Anker-Nilssen, T., Crawford, R.J.M., Furness, R.W.,
494 Mills, J.A., Murphy, E.J., Osterblom, H., Paleczny, M., Piatt, J.F., Roux, J.-P., Shannon, L.,
495 Sydeman, W.J., 2011. Global seabird response to forage fish depletion -- one-third for the birds.
496 *Science* 334, 1703-1706. doi: 10.1126/science.1212928.
497
498 de Pinna, M.C.C., Di Dario, F. 2003. *Family Pristigasteridae (Pristigasterids)*, in: R. E. Reis, S. O.
499 Kullander, C. Ferraris. (Org.). Check List of Freshwater Fishes of South and Central America.
500 Porto Alegre: EDIPUCRS, p. 43-45.
501
502 Di Dario, F., 2018. *Tenualosa toli* (errata version published in 2019). The IUCN Red List of
503 Threatened Species 2018: e.T187944A143832449. doi: 10.2305/IUCN.UK.2018-
504 2.RLTS.T187944A143832449.en.
505

506 Dulvy, N.K., Pinnegar, J.K., Reynolds, J.D., 2009. Holocene extinctions in the sea. Chapter 6, in:
507 Turvey, S.T. (Ed.) Holocene Extinctions. Oxford Biology. Oxford University Press, pp. 129 – 150.
508

509 FAO, 2016. Fishery and Aquaculture Statistics. Global capture production 1950-2014 (FishStatJ).
510 FAO Fisheries and Aquaculture Department [online or CD-ROM], Rome. Updated 2016.
511 <http://www.fao.org/fishery/>.
512

513 FAO, 2018. The State of World Fisheries and Aquaculture 2018 – Meeting the sustainable
514 development goals. Rome. License: CC BY-NC-SA 3.0 IGO.
515

516 FAO, 2020. The State of World Fisheries and Aquaculture 2020. Sustainability in action. Rome.
517 Licence: CC BY-NC-SA 3.0 IGO. <http://doi.org/10.4060/ca9229en>.
518

519 Ferrol-Schulte, D., Gorris, P., Baitoningsih, W., Adhuri, D.S., Ferse, S.C.A., 2015. Coastal
520 livelihood vulnerability to marine resource degradation: a review of the Indonesian national
521 coastal and marine policy framework. Marine Policy 52 (2015): 163 – 171.
522

523 Freyhof, J., Kottelat, M., 2008a. *Alosa immaculata*. The IUCN Red List of Threatened
524 Species 2008. e.T907A13093654. doi: [10.2305/IUCN.UK.2008.RLTS.T907A13093654.en](https://doi.org/10.2305/IUCN.UK.2008.RLTS.T907A13093654.en).
525

526 Freyhof, J., Kottelat, M., 2008b. *Alosa killarnensis*. The IUCN Red List of Threatened
527 Species 2008. e.T135582A4152432. doi: [10.2305/IUCN.UK.2008.RLTS.T135582A4152432.en](https://doi.org/10.2305/IUCN.UK.2008.RLTS.T135582A4152432.en).
528

529 Fricke, R., Eschmeyer, W.N., Van der Laan, R. (eds.), 2020. Catalog of Fishes: genera, species,
530 references. Accessed March
531 2020. <http://researcharchive.calacademy.org/research/ichthyology/catalog/fishcatmain.asp>.
532

533 Gaichas, S., Gamble, R., Fogarty, M., Benoit, H., Essington, T., Fu, C., Koen-Alonso, M., Link, J.,
534 2012. Assembly rules for aggregate-species production models: simulations in support of
535 management strategy evaluation. Marine Ecology Progress Series 459, 275 – 292. doi:
536 [10.3354/meps09650](https://doi.org/10.3354/meps09650).
537

538 Grande, L., 1985. Recent and fossil clupeomorph fishes with materials for revision of the
539 subgroups of clupeoids. Bulletin of the American Museum of Natural History 181 (2), 231 - 372
540

541 Hata, H., Motomura, H., 2019a. A new species of sardine, *Sardinella pacifica* from the
542 Philippines (Teleostei, Clupeiformes, Clupeidae). ZooKeys 829: 75 – 83.
543 <https://doi.org/10.3897/zookeys.829.30688>.
544

545 Hata, H., Motomura, H., 2019b. A new species of sardine, *Sardinella electra* (Teleostei:
546 Clupeiformes: Clupedidae), from the Ryukyu Islands, Japan. Zootaxa 4565(2): 274 – 280.
547 <https://doi.org/10.11646/zootaxa.4565.2.11>.

548
549 Hata, H., Motomura, H., 2019c. *Sardinella alcyone* n. sp., a new sardine (Teleostei:
550 Clupeiformes: Clupeidae) from the northwestern Pacific Ocean. Zootaxa 4702(1): 019 – 025.
551 <https://doi.org/10.11646/zootaxa.4702.1.6>.
552
553 Hay, D.E., Torenson, R., Stephenson, R., Thompson, M., Claytor, R., Funk, F., Ivshina, E.,
554 Jakobsson, J., Kobayashi, T., McQuinn, I., Melvin, G., Molloy, J., Naumenko, N., Oda, K.T.,
555 Parmanne, R., Power, M., Radchenko, V., Schweigert, J., Simmonds, J., Sjöstrand, B., Stevenson,
556 D.K., Tanasichuk, R., Tang, Q., Watters, D.L., Wheeler, J., 2001. Taking Stock: An Inventory and
557 Review of World Herring Stocks in 2001. Herring: Expectations for a New Millennium. Alaska
558 Sea Grant College Program. AK-SG-01-04.
559
560 Harnik, P.G., Lotze, H.K., Anderson, S.C., Finkel, Z.V., Finnegan, S., Lindberg, D.R., Liow, L.H.,
561 Lockwood, R., McClain, C.R., McGuire, J.L., O’Dea, A., Pandolfi, J.M., Simpson, C., Tittensor, D.
562 P., 2012. Extinctions in ancient and modern seas. Trends in Ecology & Evolution 27, 608-617.
563 doi: 10.1016/j.tree.2012.07.010
564
565 Hilborn, R., Amoroso, R.O., Bogazzi, E., Jensen, O.P., Parma, A.M., Szuwalski, C., Walters, C.J.,
566 2017. When does fishing forage species affect their predators? Fisheries Research 191, 211-221.
567 doi: 10.1016/j.fishres.2017.01.008.
568
569 Hoffmann, M., Brooks, T.M., da Fonseca, G.A., Gascon, C., Hawkins, A.F.A., James, R.E.,
570 Langhammer, P., Mittermeier, R.A., Pilgrim, J.D., Rodrigues, A.S.L., Silva, J.M.C., 2008.
571 Conservation planning and the IUCN Red List. Endangered Species Research 6, 113-125.
572
573 Hoffmann, M., Hilton-Taylor, C., Angulo, A., Böhm, M., Brooks, T.M., Butchart, S.H., ... Darwall,
574 W.R., 2010. The impact of conservation on the status of the world’s vertebrates. Science 330,
575 1503-1509.
576
577 IOC, IHO, BODC, 2003. Centenary Edition of the GEBCO Digital Atlas, published on CD-ROM on
578 behalf of the Intergovernmental Oceanographic Commission and the International
579 Hydrographic Organization as part of the General Bathymetric Chart of the Oceans. British
580 Oceanographic Data Centre, Liverpool.
581
582 IUCN, 2001. IUCN Red List Categories and Criteria: Version 3.1. First edition. IUCN, Gland,
583 Switzerland and Cambridge, UK.
584
585 IUCN, 2012. IUCN Red List Categories and Criteria: Version 3.1. Second edition. IUCN, Gland,
586 Switzerland and Cambridge, UK, pp. 32.
587
588 IUCN, 2016. Guideline for Appropriate Uses of IUCN Red List Data. Annex 1. Guidelines for
589 Reporting on Proportion Threatened. Version 1.1. Adopted by the IUCN Red List Committee on
590 October 2016.

591 https://nc.iucnredlist.org/redlist/content/attachment_files/Guidelines_for_Reporting_Proporti
592 [on_Threatened_ver1_1.pdf](https://nc.iucnredlist.org/redlist/content/attachment_files/Guidelines_for_Reporting_Proporti)
593
594 IUCN-CMP, 2016. Unified Classification of Direct Threats. Version 3.2.
595 <https://www.iucnredlist.org/resources/threat-classification-scheme>.
596
597 IUCN Standards and Petitions Subcommittee, 2016. Guidelines for using the IUCN Red List
598 Categories and Criteria. Version 12. Prepared by the Standards and Petitions Subcommittee.
599
600 Jennings, S., Reynolds, J.D., Mills, S.C., 1998. Life history correlates of responses to fisheries
601 exploitation. *Proceedings of the Royal Society B* 265, 333–339.
602
603 Juan-Jorda, M.J., Mosquiera, I., Freire, J., Dulvy, N.K., 2012. Life history correlates of marine
604 fisheries vulnerability: a review and a test with tunas and mackerel species, in: Briand, F. (Ed.),
605 Marine extinctions – patterns and processes. CIESM Workshop Monograph 45, 113 – 128.
606 CIESM Publisher, Monaco.
607
608 Kindsvater, H.K., Mangel, M., Reynolds, J.D., Dulvy, N.K., 2016. Ten principles from evolutionary
609 ecology essential for effective marine conservation. *Ecology and Evolution* 6, 2125–2138. doi:
610 10.1002/ece3.2012.
611
612 Lalèyè, P., Moelants, T., Olaosebikan, B.D. 2010. *Denticeps clupeioides*. *The IUCN Red List of*
613 *Threatened Species* 2010: e.T182459A7890688. [https://dx.doi.org/10.2305/IUCN.UK.2010-](https://dx.doi.org/10.2305/IUCN.UK.2010-3.RLTS.T182459A7890688.en)
614 [3.RLTS.T182459A7890688.en](https://dx.doi.org/10.2305/IUCN.UK.2010-3.RLTS.T182459A7890688.en). Downloaded on 08 May 2020.
615
616 Lavoué, S., Miya, M., Musikasinthorn, P., Chen, W.-J., Nishida, M., 2013. Mitogenomic evidence
617 for an Indo-West Pacific origin of the Clupeoidei (Teleostei: Clupeiformes). *PLoS ONE* 8, e56485.
618 doi: 10.1371/journal.pone.0056485.
619
620 Lavoué, S., Konstantinidis, P., Chen, W.-J. 2014. Progress in clupeiform systematics, in: Ganas,
621 K. (Ed.), *Biology and ecology of sardines and anchovies*. Boca Raton: CRC Press, 3– 42.
622
623 Lehner, B., Grill, G., 2013. Global river hydrography and network routing: baseline data and new
624 approaches to study the world’s large river systems. *Hydrological Processes* 27, 2171 – 2186.
625 doi: 10.1002/hyp.9740. Data is available at www.hydrosheds.org.
626
627 Lévêque, C., Oberdorff, T., Paugy, D., Stiassny, M.L.J., Tedesco, P.A., 2008. Global diversity of
628 fish (Pisces) in freshwater. *Hydrobiologia* 595, 545 – 567. doi: 10.1007/s10750-007-9034-0.
629
630 Levin, P.S., Francis, T.B., Taylor, N.G., 2016. Thirty-two essential questions for understanding
631 the social-ecological system of forage fish: the case of Pacific Herring. *Ecosystem Health and*
632 *Sustainability* 2, e01213. doi:10.1002/ehs2.1213.
633

634 Mace, G.M., Collar, N.J., Gaston, K.J., Hilton-Taylor, C., Akcakaya, H.R., Leader-Williams, N.,
635 Milner-Gulland, E.J., Stuart, S.N., 2008. Quantification of Extinction Risk: IUCN's system for
636 classifying threatened species. *Conservation Biology* 22, 1424-1442. doi: 10.1111/j.1523-
637 1739.2008.01044.x
638

639 Malabarba, M. C., Di Dario, F. 2017. A new predatory herring-like fish (Teleostei: Clupeiformes)
640 from the Early Cretaceous of Brazil, and implications for the relationships in the Clupeoidei.
641 *Zoological Journal of the Linnean Society* 180: 175-194.
642

643 Mohanty, B.P., Ganguly, S., Mahanty, A., Mitra, T., Patra, S., Karunakaran, D., Mathew, S.,
644 Chakraborty, K., Paul, B.N., Sarma, D., Dayal, S., Singh, S., Ayyappan, S., 2019. Fish in human
645 health and nutrition. *Advances in Fish Research* 7, 189 – 218.
646

647 Mohd Arshaad, W., Gaughan, D., Munroe, T.A., 2018. *Tenualosa macrura* (errata version
648 published in 2019). The IUCN Red List of Threatened Species 2018:
649 e.T98842673A143840186. doi: [10.2305/IUCN.UK.2018-2.RLTS.T98842673A143840186.en](https://doi.org/10.2305/IUCN.UK.2018-2.RLTS.T98842673A143840186.en).
650

651 NatureServe, 2013. *Alosa aestivalis*. The IUCN Red List of Threatened Species 2013:
652 e.T201946A2730890. doi: [10.2305/IUCN.UK.2013-1.RLTS.T201946A2730890.en](https://doi.org/10.2305/IUCN.UK.2013-1.RLTS.T201946A2730890.en).
653

654 NOAA Fisheries, 2019. National Oceanic and Atmospheric Administration: Commercial landings.
655 <https://foss.nmfs.noaa.gov/apexfoss/f?p=215:200:11056708372560>
656

657 Pauly, D., Zeller, D., 2016a. Sea Around Us Concepts, Design and Data. Available at:
658 searoundus.org.
659

660 Pauly, D., Zeller, D., 2016b. Catch reconstructions reveal that global marine fisheries catches are
661 higher than reported and declining. *Nature Communications* 7, 1 – 9. doi:
662 10.1038/ncomms10244.
663

664 Pikitch, E.K., Rountos, K.J., Essington, T.E., Santora, C., Pauly, D., Watson, R., Sumalia, U.R.,
665 Boersma, P.D., Boyd, I.L., Conover, D.O., Cury, P., Heppell, S.S., Houde, E.D., Mangel, M.,
666 Plaga'nyi, E., Sainsbury, K., Steneck, R.S., Geers, T.M., Gownaris, N., Munch, S.B., 2014. The
667 global contribution of forage fish to marine fisheries and ecosystems. *Fish and Fisheries* 15: 43-
668 64. doi: 10.1111/faf.12004.
669

670 Polidoro, B.A., Carpenter, K.E., Collins, L., Duke, N.C., Ellison, A.M., Ellison, J.C., Farnsworth, E.J.,
671 Fernando, E.S., Kathiresan, K., Koedam, N.E., Livingstone, S.R., Miyago, T., Moore, G.E., Ngoc
672 Nam, V., Eong Ong, J., Primavera, J.H., Salmo, S.G., Sanciangco, J.C., Sukardjo, S., Wang, Y., Hong
673 Yong, J.W., 2010. The loss of species: mangrove extinction risk and geographic areas of global
674 concern. *Public Library of Science One* 5, e10095. doi: 10.131/journal.pone.0010095.
675

676 Rola, A.C., Narvaez, T.A., Naguit, M.R.A., Elazegui, D.D., Brilloa, B.B.C., Paunlagui, M.M., Jalotjot,
677 H.C., Cervantes, C.P. 2018 Impact of the closed fishing season policy for sardines in Zamboanga
678 Peninsula, Philippines. *Marine Policy*. 40 – 50. doi: 10.1016/j.marpol.2017.09.029.
679

680 Reynolds, J.D., Dulvy, N.K., Goodwin, N.B., Hutchings, J.A., 2005. Biology of extinction risk in
681 marine fishes. *Proceedings of the Royal Society B: Biological Sciences* 272, 2337-2344.
682

683 Roberts, C.M., Hawkins, J.P., 1999. Extinction risk in the sea. *Trends in Ecology & Evolution* 14,
684 241-246. doi: 10.1016/S0169-5347(98)01584-5.
685

686 Rodrigues, A.S.L., Pilgrim, J.D., Lamoreux, J.F., Hoffmann, M., Brooks, T.M., 2006. The value of
687 the IUCN Red List for conservation. *Trends in Ecology & Evolution* 21, 71-76. doi:
688 10.1016/j.tree.2005.10.010.
689

690 Rountos, K., 2016. Defining forage species to prevent a management dilemma. *Fisheries* 41 (1),
691 16-17. doi.org/10.1080/03632415.2015.1110791.
692

693 Sadovy. Y., 2001. The threat of fishing to highly fecund fishes. *Journal of Fish Biology* 59, 90 –
694 108. doi: 10.1006/jfbi.2001.1760.
695

696 Sadovy de Mitcheson, Y., Linardich, C., Barreiros, J.P., Ralph, G.M., Aguilar-Perera, A., Afonso,
697 P., . . . Craig, M.T., 2020. Valuable but vulnerable: Over-fishing and under-management
698 continue to threaten groupers so what now? *Marine Policy* 116, 1 – 10. doi:
699 10.1016/j.marpol.2020.103909.
700

701 Sanciangco, J.C., Carpenter, K.E., Etnoyer, P.J., Moretzsohn, F., 2013. Habitat availability and
702 heterogeneity and the Indo-Pacific warm pool as predictors of marine species richness in the
703 tropical Indo-Pacific. *PLoS ONE* 8(2): e56245. doi:10.1371/journal.pone.0056245.
704

705 Santos, M., Munroe, T.A., Di Dario, F., Hata, H., Torres, F., Quilang, J.P., 2018. *Sardinella tawilis*
706 (errata version published in 2019). The IUCN Red List of Threatened Species 2018:
707 e.T98836352A143839946. doi: 10.2305/IUCN.UK.20182.RLTS.T98836352A143839946.en.
708

709 Schipper, J., Chanson, J.S., Chiozza, F., Cox, N.A., Hoffmann, M., Katariya, V., . . . Young, B.E.,
710 2008. The status of the world's land and marine mammals: Diversity, threat, and knowledge.
711 *Science* 322, 225-230. doi: 10.1126/science.1165115.
712


713 Short, F.T., Polidoro, B., Livingstone, S.R., Carpenter, K.E., Bandeira, S., Bujang, J.S.,
714 Calumpong, H.P., Carruthers, T.J.B., Coles, R.G., Dennison, W.C., Erftemeijer, P.L.A., Fortes,
715 M.D., Freeman, A.S., Jagtap, T.G., Kamal, A.H.M., Kendrick, G.A., Kensworthy, W.J., La Nafie,
716 Y.A., Nasution, I.M., Orth, R.J., Prathep, A., Sanciangco, J.C., van Tussenbroek, B., Vergara, S.G.,
717 Waycott, M., Zieman, J.C., 2011. Extinction risk assessment of the world's seagrass species.
718 *Biological Conservation* 144, 1961 – 1971.

719
720 Siebert, D. J., 1997. Notes on the anatomy and relationships of *Sundasalanx* Roberts (Teleostei,
721 Clupeidae), with descriptions of four new species from Borneo. Bulletin of the American
722 Museum of Natural History London (Zool.) 63(1), 13-26.
723
724 Smith, A.D.M., Brown, C.J., Bulman, C.M., Fulton, E.A., Johnson, P., Kaplan, I.C., Lozano-Montes,
725 H., Mackinson, S., Marzloff, M., Shannon, L.J., Sjin, Y.-J., Tam, J., 2011. impacts of fishing low-
726 trophic level species on marine ecosystems. Science 333, 1147 – 1150. doi:
727 10.1126/science.1209395.
728
729 Spalding, M.D., Fox, H.E., Allen, G.R., Davidson, N., Ferdaña, Z.A., Finlayson, M., . . . Robertson,
730 J., 2007. Marine ecoregions of the world: A bioregionalization of coastal and shelf areas.
731 BioScience 57, 573-583. doi: 10.1641/b570707.
732 Stobutzki, I., Miller, M., Brewer, D., 2001. Sustainability of fishery bycatch: a process for
733 assessing highly diverse and numerous bycatch. Environmental Conservation 28, 167 – 181. doi:
734 10.1017/S0376892901000170.
735
736 Tacon, A.G.J., Metian, M., 2009a. Fishing for feed of fishing for food: increasing global
737 competition for small pelagic forage fish. AMBIO: A Journal of the Human Environment 38, 294-
738 302.
739
740 Teugels, G. 2003. Denticipitidae, in: D. Paugy; C. Lévêque & Teugels,
741 G.G. (Eds.), Poissons d’eaux douces et saumâtres de l’Afrique de l’Ouest. Tome I. Collection
742 Faune et Flore tropicales 40. Institut de Recherche pour le Développement, and Muséum
743 National d’Histoire Naturelle, Paris, France; Musée Royal de l’Afrique Centrale, Tervuren,
744 Belgique. Paris.
745
746 Thomas Jr., R.C., Willette, D.A., Carpenter, K.E., Santos, M.D., 2014. Hidden diversity in sardines:
747 genetic and morphological evidence for cryptic species in the Goldstripe Sardinella, *Sardinella*
748 *gibbosa*, (Bleeker, 1849). PLoS ONE 9(1): e84719. doi:10.1371/journal.pone.0084719
749
750 Thornton, T.F., Moss, M.L., Butler, V.L., Hebert, J., Funk, F., 2010. Local and traditional
751 knowledge and the historical ecology of Pacific herring in Alaska. Journal of Ecological
752 Anthropology 14, 81 – 88.
753
754 United Nations, Department of Economic and Social Affairs, Population Division, 2017. World
755 population prospects: The 2017 Revision, World Population 2017 Wallchart. AT/ESA/SER.A/398.
756
757 Vié, J.-C., Hilton-Taylor, C., Stuart, S.N., 2009. Wildlife in a Changing World - An analysis of
758 the 2008 IUCN Red List of Threatened Species. IUCN, Gland, Switzerland.
759
760 Whitehead, P.J.P., 1985. FAO species catalogue. Vol. 7. Clupeoid fishes of the world (suborder
761 Clupeoidei). An annotated and illustrated catalogue of the herrings, sardines, pilchards, sprats,

762 shads, anchovies and wolf-herrings. Part 1 - Chirocentridae, Clupeidae and Pristigasteridae.
763 Food and Agricultural Organization (FAO) Fisheries Synopsis, Rome, Italy.
764
765 Whitehead, P.J.P., Nelson, G.J., Wongratana, T., 1988. FAO species catalogue. Vol. 7. Clupeoid
766 fishes of the world (Suborder Clupeioidi). An annotated and illustrated catalogue of the
767 herrings, sardines, pilchards, sprats, shads, anchovies and wolf-herrings. Part 2 - Engraulididae.
768 Food and Agricultural Organization (FAO) Fisheries Synopsis, Rome, Italy.
769
770


771 **FIGURES**

772 Figure 1 color and grayscale versions should be 1.5 columns (color version **only** for online
773 version).



774

775 **Fig. 1.** Proportion of species listed in Red List Categories partitioned by family. Abbreviations of
776 Red List Categories are as follows: EC = elevated concern (includes species evaluated as
777 Critically Endangered, Endangered, Vulnerable, or Near Threatened), LC = Least Concern and DD
778 = Data Deficient. The total number of species in each family is represented by the number at
779 the top of each bar. The midpoint is represented by the black bar and was calculated by the
780 following equation: $(CR + EN + VU + NT) / (Total - DD)$.




781

782 **Fig. 1.** Proportion of species listed in Red List Categories partitioned by family. Abbreviations of
 783 Red List Categories are as follows: EC = elevated concern (includes species evaluated as
 784 Critically Endangered, Endangered, Vulnerable, or Near Threatened), LC = Least Concern and DD
 785 = Data Deficient. The total number of species in each family is represented by the number at
 786 the top of each bar. The midpoint is represented by the black bar and was calculated by the
 787 following equation: $(CR + EN + VU + NT) / (Total - DD)$.

788

789


790 Figure 2 color and grayscale versions should be one column (color version **only** for online
791 version).



792

793 **Fig. 2.** Proportion of species listed in Red List Categories by major habitat system (freshwater,
794 euryhaline, or marine). Abbreviations of Red List Categories are as follows: EC = elevated
795 concern (includes species evaluated as Critically Endangered, Endangered, Vulnerable, or Near
796 Threatened), LC = Least Concern and DD = Data Deficient. The total number of species in each
797 family is represented by the number at the top of each bar. The midpoint is represented by the
798 black bar and was calculated by the following equation: $(CR + EN + VU + NT) / (Total - DD)$.

799




800


801 **Fig. 2.** Proportion of species listed in Red List Categories by major habitat system (freshwater,
 802 euryhaline, or marine). Abbreviations of Red List Categories are as follows: EC = elevated
 803 concern (includes species evaluated as Critically Endangered, Endangered, Vulnerable, or Near
 804 Threatened), LC = Least Concern and DD = Data Deficient. The total number of species in each
 805 family is represented by the number at the top of each bar. The midpoint is represented by the
 806 black bar and was calculated by the following equation: $(CR + EN + VU + NT) / (Total - DD)$.

807

808 Figure 3 color and grayscale versions should be 1.5 columns (color version **only** for online
 809 version).



810
 811 **Fig. 3.** Number of clupeiform species impacted by major threats. Each threat is represented by
 812 the number of species impacted separated by major habitat system (freshwater, euryhaline, or
 813 marine). Threats impacting less than ten species (Mining, Development, Human intrusion, and
 814 Transportation) are excluded.



815
 816 **Fig. 3.** Number of clupeiform species impacted by major threats. Each threat is represented by
 817 the number of species impacted separated by major habitat system (freshwater, euryhaline, or
 818 marine). Threats impacting less than ten species (Mining, Development, Human intrusion, and
 819 Transportation) are excluded.

820 Figure 4 color and grayscale should be two columns (color version **only** for online version).




Fig. 4. Number of clupeiform species in each Large Marine Ecoregion and freshwater hydrobasin for A) All species, B) all Data Deficient species, and C) all species of elevated concern (Critically Endangered, Endangered, Vulnerable, Near Threatened). Colors correspond to numbers of species listed at the bottom of each map. The Marine Ecosystems of the World at the province level (Spalding et al., 2007) was used for marine species, Global HydroBASINS at level three (Lehner and Grill, 2013) was used for freshwater species. The freshwater and marine extents were created separately and merged to represent the total global extent for euryhaline species.




Fig. 5. Number of clupeiform species in each Large Marine Ecoregion and freshwater hydrobasin for A) All species, B) all Data Deficient species, and C) all species of elevated concern (Critically Endangered, Endangered, Vulnerable, Near Threatened). Colors correspond to numbers of species listed at the bottom of each map. The Marine Ecosystems of the World at the province level (Spalding et al., 2007) was used for marine species, Global HydroBASINS at level three (Lehner and Grill, 2013) was used for freshwater species. The freshwater and marine extents were created separately and merged to represent the total global extent for euryhaline species.

822 **SUPPLEMENTARY INFORMATION**

823 **Table A1.** List of all 415 clupeiform species alphabetical by family and then by species name.
824 The global IUCN Red List categories and criteria are listed: CR = Critically Endangered, EN =
825 Endangered, VU = Vulnerable, NT = Near Threatened, LC = Least Concern, DD = Data Deficient,
826 NE = Not Evaluated. Criterion A = population decline in the past, present or future, B =
827 restricted range, C = small population size and decline, D = very small or restricted population, E
828 = quantitative analysis of extinction probability. For further information available on categories
829 and criteria, visit the IUCN Red List website (www.iucnredlist.org). The preferred habitat system
830 is also listed; F = Freshwater, M = Marine, E = Euryhaline which includes estuarine species and
831 diadromous species.

FAMILY	SPECIES NAME	GLOBAL CATEGORY & CRITERIA	SYSTEM
Chirocentridae	<i>Chirocentrus dorab</i>	LC	M
Chirocentridae	<i>Chirocentrus nudus</i>	LC	M
Clupeidae	<i>Alosa aestivalis</i>	VU A2b	E
Clupeidae	<i>Alosa agone</i>	LC	F
Clupeidae	<i>Alosa alabamae</i>	NT A2ac	E
Clupeidae	<i>Alosa algeriensis</i>	DD	E
Clupeidae	<i>Alosa alosa</i>	LC	E
Clupeidae	<i>Alosa braschnikowi</i>	DD	E
Clupeidae	<i>Alosa caspia</i>	LC	E
Clupeidae	<i>Alosa chrysochloris</i>	LC	E
Clupeidae	<i>Alosa curensis</i>	DD	E
Clupeidae	<i>Alosa fallax</i>	LC	E
Clupeidae	<i>Alosa immaculata</i>	VU B2ab(v)	E
Clupeidae	<i>Alosa kessleri</i>	LC	E
Clupeidae	<i>Alosa killarnensis</i>	CR B1ab(iii)	F
Clupeidae	<i>Alosa macedonica</i>	VU D2	F
Clupeidae	<i>Alosa maeotica</i>	LC	E
Clupeidae	<i>Alosa mediocris</i>	LC	E
Clupeidae	<i>Alosa pontica</i>	LC	E
Clupeidae	<i>Alosa pseudoharengus</i>	LC	E
Clupeidae	<i>Alosa sapidissima</i>	LC	E
Clupeidae	<i>Alosa saposchnikowii</i>	DD	E
Clupeidae	<i>Alosa sphaerocephala</i>	LC	E
Clupeidae	<i>Alosa suworowi</i>	DD	E
Clupeidae	<i>Alosa tanaica</i>	LC	E
Clupeidae	<i>Alosa vistonica</i>	CR A2ace; B1ab(iii,v)	F
Clupeidae	<i>Alosa volgensis</i>	EN B2ab(iii,v)	E
Clupeidae	<i>Amblygaster clupeoides</i>	LC	M
Clupeidae	<i>Amblygaster indiana</i>	DD	M

FAMILY	SPECIES NAME	GLOBAL CATEGORY & CRITERIA	SYSTEM
Clupeidae	<i>Amblygaster leiogaster</i>	LC	M
Clupeidae	<i>Amblygaster sirm</i>	LC	M
Clupeidae	<i>Anodontostoma chacunda</i>	LC	E
Clupeidae	<i>Anodontostoma selangkat</i>	LC	E
Clupeidae	<i>Anodontostoma thailandiae</i>	LC	E
Clupeidae	<i>Brevoortia aurea</i>	LC	E
Clupeidae	<i>Brevoortia gunteri</i>	LC	M
Clupeidae	<i>Brevoortia patronus</i>	LC	E
Clupeidae	<i>Brevoortia pectinata</i>	LC	E
Clupeidae	<i>Brevoortia smithi</i>	LC	E
Clupeidae	<i>Brevoortia tyrannus</i>	LC	E
Clupeidae	<i>Clupanodon thrissa</i>	LC	E
Clupeidae	<i>Clupea harengus</i>	LC	M
Clupeidae	<i>Clupea pallasii</i>	DD	M
Clupeidae	<i>Clupeichthys aesarnensis</i>	LC	F
Clupeidae	<i>Clupeichthys bleekeri</i>	VU B1ab(iii)	F
Clupeidae	<i>Clupeichthys goniognathus</i>	LC	E
Clupeidae	<i>Clupeichthys perakensis</i>	LC	E
Clupeidae	<i>Clupeoides borneensis</i>	LC	E
Clupeidae	<i>Clupeoides hypselosoma</i>	DD	F
Clupeidae	<i>Clupeoides papuensis</i>	DD	F
Clupeidae	<i>Clupeoides venulosus</i>	VU B2ab(iii,v)	F
Clupeidae	<i>Clupeonella abrau</i>	CR B1ab(ii,iii,v)+2ab(ii,iii,v)	F
Clupeidae	<i>Clupeonella caspia</i>	LC	E
Clupeidae	<i>Clupeonella cultriventris</i>	LC	E
Clupeidae	<i>Clupeonella engrauliformis</i>	EN A2bde	M
Clupeidae	<i>Clupeonella grimmi</i>	EN A2bde	M
Clupeidae	<i>Clupeonella muhlisi</i>	EN B1ab(iii)+2ab(iii)	F
Clupeidae	<i>Clupeonella tscharchalensis</i>	LC	E
Clupeidae	<i>Congothrissa gossei</i>	DD	F
Clupeidae	<i>Corica laciniata</i>	DD	F
Clupeidae	<i>Corica soborna</i>	LC	E
Clupeidae	<i>Dayella malabarica</i>	LC	E
Clupeidae	<i>Dorosoma anale</i>	LC	F
Clupeidae	<i>Dorosoma cepedianum</i>	LC	E
Clupeidae	<i>Dorosoma chavesi</i>	NT B1ab(iii)	F
Clupeidae	<i>Dorosoma petenense</i>	LC	E
Clupeidae	<i>Dorosoma smithi</i>	DD	F
Clupeidae	<i>Ehirava fluviatilis</i>	DD	E
Clupeidae	<i>Escualosa elongata</i>	DD	M

FAMILY	SPECIES NAME	GLOBAL CATEGORY & CRITERIA	SYSTEM
Clupeidae	<i>Escualosa thoracata</i>	LC	E
Clupeidae	<i>Ethmalosa fimbriata</i>	LC	E
Clupeidae	<i>Ethmidium maculatum</i>	DD	M
Clupeidae	<i>Gilchristella aestuaria</i>	LC	E
Clupeidae	<i>Gonialosa manmina</i>	LC	E
Clupeidae	<i>Gonialosa modesta</i>	DD	E
Clupeidae	<i>Gonialosa whiteheadi</i>	DD	E
Clupeidae	<i>Gudusia chapra</i>	LC	F
Clupeidae	<i>Gudusia variegata</i>	LC	F
Clupeidae	<i>Harengula clupeola</i>	LC	E
Clupeidae	<i>Harengula humeralis</i>	LC	E
Clupeidae	<i>Harengula jaguana</i>	LC	M
Clupeidae	<i>Harengula thrissina</i>	LC	E
Clupeidae	<i>Herklotsichthys blackburni</i>	DD	E
Clupeidae	<i>Herklotsichthys castelnaui</i>	LC	E
Clupeidae	<i>Herklotsichthys collettei</i>	LC	M
Clupeidae	<i>Herklotsichthys dispilonotus</i>	LC	M
Clupeidae	<i>Herklotsichthys gotoi</i>	LC	E
Clupeidae	<i>Herklotsichthys koningsbergeri</i>	LC	E
Clupeidae	<i>Herklotsichthys lippa</i>	LC	M
Clupeidae	<i>Herklotsichthys lossei</i>	LC	M
Clupeidae	<i>Herklotsichthys ovalis</i>	DD	M
Clupeidae	<i>Herklotsichthys punctatus</i>	LC	M
Clupeidae	<i>Herklotsichthys quadrimaculatus</i>	LC	M
Clupeidae	<i>Herklotsichthys spilurus</i>	LC	M
Clupeidae	<i>Hilsa kelee</i>	LC	E
Clupeidae	<i>Hyperlophus translucidus</i>	LC	E
Clupeidae	<i>Hyperlophus vittatus</i>	LC	E
Clupeidae	<i>Konosirus punctatus</i>	LC	E
Clupeidae	<i>Laeviscutella dekimpei</i>	LC	E
Clupeidae	<i>Lile gracilis</i>	LC	E
Clupeidae	<i>Lile nigrofasciata</i>	LC	E
Clupeidae	<i>Lile piquitinga</i>	LC	E
Clupeidae	<i>Lile stolidifera</i>	LC	E
Clupeidae	<i>Limnothrissa miodon</i>	LC	E
Clupeidae	<i>Limnothrissa stappersii</i>	DD	F
Clupeidae	<i>Microthrissa minuta</i>	VU D2	F
Clupeidae	<i>Microthrissa royauxi</i>	LC	F
Clupeidae	<i>Microthrissa whiteheadi</i>	LC	F
Clupeidae	<i>Minyclupeoides dentibranchialis</i>	LC	E

FAMILY	SPECIES NAME	GLOBAL CATEGORY & CRITERIA	SYSTEM
Clupeidae	<i>Nannothrissa parva</i>	LC	F
Clupeidae	<i>Nannothrissa stewarti</i>	EN B1ab(v)	F
Clupeidae	<i>Nematalosa arabica</i>	DD	M
Clupeidae	<i>Nematalosa come</i>	LC	M
Clupeidae	<i>Nematalosa erebi</i>	LC	F
Clupeidae	<i>Nematalosa flyensis</i>	DD	F
Clupeidae	<i>Nematalosa galathea</i>	LC	E
Clupeidae	<i>Nematalosa japonica</i>	DD	M
Clupeidae	<i>Nematalosa nasus</i>	LC	E
Clupeidae	<i>Nematalosa papuensis</i>	DD	F
Clupeidae	<i>Nematalosa persara</i>	DD	M
Clupeidae	<i>Nematalosa resticularia</i>	DD	M
Clupeidae	<i>Nematalosa vlaminghi</i>	LC	E
Clupeidae	<i>Odaxothrissa ansorgii</i>	LC	F
Clupeidae	<i>Odaxothrissa losera</i>	DD	F
Clupeidae	<i>Odaxothrissa mento</i>	LC	F
Clupeidae	<i>Odaxothrissa vittata</i>	LC	F
Clupeidae	<i>Opisthonema berlangai</i>	VU D2	M
Clupeidae	<i>Opisthonema bulleri</i>	LC	M
Clupeidae	<i>Opisthonema libertate</i>	LC	M
Clupeidae	<i>Opisthonema medirastre</i>	LC	M
Clupeidae	<i>Opisthonema oglinum</i>	LC	E
Clupeidae	<i>Pellonula leonensis</i>	LC	E
Clupeidae	<i>Pellonula vorax</i>	LC	E
Clupeidae	<i>Platanichthys platana</i>	LC	E
Clupeidae	<i>Poecilothrissa centralis</i>	LC	F
Clupeidae	<i>Poecilothrissa congica</i>	LC	F
Clupeidae	<i>Poecilothrissa moeruensis</i>	VU B1ab(v)	F
Clupeidae	<i>Potamalosa richmondia</i>	LC	E
Clupeidae	<i>Potamothrissa acutirostris</i>	LC	F
Clupeidae	<i>Potamothrissa obtusirostris</i>	LC	F
Clupeidae	<i>Potamothrissa whiteheadi</i>	DD	F
Clupeidae	<i>Ramnogaster arcuata</i>	LC	M
Clupeidae	<i>Ramnogaster melanostoma</i>	LC	F
Clupeidae	<i>Rhinosardinia amazonica</i>	LC	E
Clupeidae	<i>Rhinosardinia bahiensis</i>	LC	E
Clupeidae	<i>Sardina pilchardus</i>	LC	M
Clupeidae	<i>Sardinella albella</i>	LC	M
Clupeidae	<i>Sardinella alcyone</i>	DD	E
Clupeidae	<i>Sardinella atricauda</i>	LC	M

FAMILY	SPECIES NAME	GLOBAL CATEGORY & CRITERIA	SYSTEM
Clupeidae	<i>Sardinella aurita</i>	LC	M
Clupeidae	<i>Sardinella brachysoma</i>	LC	M
Clupeidae	<i>Sardinella brasiliensis</i>	DD	E
Clupeidae	<i>Sardinella dayi</i>	DD	M
Clupeidae	<i>Sardinella electra</i>	DD	M
Clupeidae	<i>Sardinella fijiense</i>	LC	M
Clupeidae	<i>Sardinella fimbriata</i>	LC	E
Clupeidae	<i>Sardinella gibbosa</i>	LC	M
Clupeidae	<i>Sardinella goni</i>	DD	M
Clupeidae	<i>Sardinella hualiensis</i>	LC	M
Clupeidae	<i>Sardinella jussieui</i>	DD	M
Clupeidae	<i>Sardinella lemuru</i>	NT A2bd	M
Clupeidae	<i>Sardinella longiceps</i>	LC	M
Clupeidae	<i>Sardinella maderensis</i>	VU A2d	M
Clupeidae	<i>Sardinella marquesensis</i>	LC	M
Clupeidae	<i>Sardinella melanura</i>	LC	E
Clupeidae	<i>Sardinella neglecta</i>	LC	M
Clupeidae	<i>Sardinella pacifica</i>	DD	M
Clupeidae	<i>Sardinella richardsoni</i>	DD	M
Clupeidae	<i>Sardinella rouxi</i>	DD	M
Clupeidae	<i>Sardinella sindensis</i>	LC	E
Clupeidae	<i>Sardinella tawilis</i>	EN A2bd; B1ab(iii,v)+2ab(iii,v)	F
Clupeidae	<i>Sardinella zunasi</i>	LC	M
Clupeidae	<i>Sardinops sagax</i>	LC	M
Clupeidae	<i>Sauvagella madagascariensis</i>	LC	E
Clupeidae	<i>Sauvagella robusta</i>	EN B2ab(iii)	F
Clupeidae	<i>Sierrathrissa leonensis</i>	LC	F
Clupeidae	<i>Spratellomorpha bianalis</i>	DD	E
Clupeidae	<i>Sprattus antipodum</i>	LC	M
Clupeidae	<i>Sprattus fuegensis</i>	LC	M
Clupeidae	<i>Sprattus muelleri</i>	LC	M
Clupeidae	<i>Sprattus novaehollandiae</i>	LC	E
Clupeidae	<i>Sprattus sprattus</i>	LC	E
Clupeidae	<i>Stolothrissa tanganicae</i>	LC	F
Clupeidae	<i>Strangomera bentincki</i>	LC	M
Clupeidae	<i>Tenualosa ilisha</i>	LC	E
Clupeidae	<i>Tenualosa macrura</i>	NT B2ab(iii)	E
Clupeidae	<i>Tenualosa reevesii</i>	DD	E
Clupeidae	<i>Tenualosa thibaudeaui</i>	VU A2bcd	F

FAMILY	SPECIES NAME	GLOBAL CATEGORY & CRITERIA	SYSTEM
Clupeidae	<i>Tenualosa toli</i>	VU B2ab(iii,v)	E
Clupeidae	<i>Thrattidion noctivagus</i>	DD	F
Denticipitidae	<i>Denticeps clupeoides</i>	VU B2ab(iii)	F
Dussumieriidae	<i>Dussumieria acuta</i>	LC	M
Dussumieriidae	<i>Dussumieria elopsoides</i>	LC	M
Dussumieriidae	<i>Etrumeus acuminatus</i>	LC	M
Dussumieriidae	<i>Etrumeus golanii</i>	DD	M
Dussumieriidae	<i>Etrumeus jacksoniensis</i>	LC	M
Dussumieriidae	<i>Etrumeus makiawa</i>	LC	M
Dussumieriidae	<i>Etrumeus micropus</i>	LC	M
Dussumieriidae	<i>Etrumeus sadina</i>	LC	M
Dussumieriidae	<i>Etrumeus whiteheadi</i>	LC	M
Dussumieriidae	<i>Etrumeus wongratanae</i>	DD	M
Dussumieriidae	<i>Jenkinsia lamprotaenia</i>	LC	M
Dussumieriidae	<i>Jenkinsia majua</i>	LC	M
Dussumieriidae	<i>Jenkinsia parvula</i>	DD	M
Dussumieriidae	<i>Jenkinsia stolifera</i>	LC	M
Dussumieriidae	<i>Spratelloides atrofasciatus</i>	LC	M
Dussumieriidae	<i>Spratelloides delicatulus</i>	LC	M
Dussumieriidae	<i>Spratelloides gracilis</i>	LC	M
Dussumieriidae	<i>Spratelloides lewisi</i>	LC	M
Dussumieriidae	<i>Spratelloides robustus</i>	LC	E
Engraulidae	<i>Amazonsprattus scintilla</i>	LC	F
Engraulidae	<i>Anchoa analis</i>	DD	E
Engraulidae	<i>Anchoa argentivittata</i>	LC	M
Engraulidae	<i>Anchoa belizensis</i>	LC	F
Engraulidae	<i>Anchoa cayorum</i>	LC	M
Engraulidae	<i>Anchoa chamensis</i>	DD	M
Engraulidae	<i>Anchoa choerostoma</i>	EN B1ab(v)+2ab(v)	M
Engraulidae	<i>Anchoa colonensis</i>	LC	M
Engraulidae	<i>Anchoa compressa</i>	LC	E
Engraulidae	<i>Anchoa cubana</i>	LC	E
Engraulidae	<i>Anchoa curta</i>	LC	E
Engraulidae	<i>Anchoa delicatissima</i>	LC	E
Engraulidae	<i>Anchoa eigenmannia</i>	LC	M
Engraulidae	<i>Anchoa exigua</i>	LC	M
Engraulidae	<i>Anchoa filifera</i>	LC	E
Engraulidae	<i>Anchoa helleri</i>	LC	M
Engraulidae	<i>Anchoa hepsetus</i>	LC	E
Engraulidae	<i>Anchoa ischana</i>	LC	M

FAMILY	SPECIES NAME	GLOBAL CATEGORY & CRITERIA	SYSTEM
Engraulidae	<i>Anchoa januaria</i>	LC	E
Engraulidae	<i>Anchoa lamprotaenia</i>	LC	M
Engraulidae	<i>Anchoa lucida</i>	LC	E
Engraulidae	<i>Anchoa lyolepis</i>	LC	M
Engraulidae	<i>Anchoa marinii</i>	LC	E
Engraulidae	<i>Anchoa mitchilli</i>	LC	E
Engraulidae	<i>Anchoa mundeola</i>	LC	E
Engraulidae	<i>Anchoa mundeoloides</i>	LC	E
Engraulidae	<i>Anchoa nasus</i>	LC	M
Engraulidae	<i>Anchoa panamensis</i>	LC	E
Engraulidae	<i>Anchoa parva</i>	LC	E
Engraulidae	<i>Anchoa pectoralis</i>	LC	E
Engraulidae	<i>Anchoa scofieldi</i>	LC	E
Engraulidae	<i>Anchoa spinifer</i>	LC	E
Engraulidae	<i>Anchoa starksi</i>	LC	E
Engraulidae	<i>Anchoa tricolor</i>	LC	E
Engraulidae	<i>Anchoa trinitatis</i>	DD	M
Engraulidae	<i>Anchoa walkeri</i>	LC	E
Engraulidae	<i>Anchovia clupeoides</i>	LC	E
Engraulidae	<i>Anchovia landivarensis</i>	DD	E
Engraulidae	<i>Anchovia macrolepidota</i>	LC	E
Engraulidae	<i>Anchovia surinamensis</i>	LC	E
Engraulidae	<i>Anchoviella alleni</i>	LC	F
Engraulidae	<i>Anchoviella balboae</i>	DD	M
Engraulidae	<i>Anchoviella blackburni</i>	DD	E
Engraulidae	<i>Anchoviella brevirostris</i>	LC	E
Engraulidae	<i>Anchoviella carrikeri</i>	LC	F
Engraulidae	<i>Anchoviella cayennensis</i>	LC	E
Engraulidae	<i>Anchoviella elongata</i>	LC	E
Engraulidae	<i>Anchoviella guianensis</i>	LC	F
Engraulidae	<i>Anchoviella hernanni</i>	LC	F
Engraulidae	<i>Anchoviella jamesi</i>	LC	F
Engraulidae	<i>Anchoviella juruasanga</i>	LC	F
Engraulidae	<i>Anchoviella lepidentostole</i>	LC	E
Engraulidae	<i>Anchoviella manamensis</i>	LC	F
Engraulidae	<i>Anchoviella miarcha</i>	DD	E
Engraulidae	<i>Anchoviella perezi</i>	DD	F
Engraulidae	<i>Anchoviella perfasciata</i>	LC	M
Engraulidae	<i>Anchoviella sanfranciscana</i>	DD	E
Engraulidae	<i>Anchoviella vaillanti</i>	LC	F

FAMILY	SPECIES NAME	GLOBAL CATEGORY & CRITERIA	SYSTEM
Engraulidae	<i>Cetengraulis edentulus</i>	LC	E
Engraulidae	<i>Cetengraulis mysticetus</i>	LC	M
Engraulidae	<i>Coilia borneensis</i>	DD	E
Engraulidae	<i>Coilia coomansi</i>	DD	E
Engraulidae	<i>Coilia dussumieri</i>	LC	E
Engraulidae	<i>Coilia grayii</i>	LC	E
Engraulidae	<i>Coilia lindmani</i>	LC	E
Engraulidae	<i>Coilia macrognathos</i>	DD	E
Engraulidae	<i>Coilia mystus</i>	EN A2bd	E
Engraulidae	<i>Coilia nasus</i>	EN A2bd	E
Engraulidae	<i>Coilia neglecta</i>	LC	E
Engraulidae	<i>Coilia ramcarati</i>	DD	E
Engraulidae	<i>Coilia rebentischii</i>	DD	E
Engraulidae	<i>Coilia reynaldi</i>	LC	E
Engraulidae	<i>Encrasicholina auster</i>	DD	M
Engraulidae	<i>Encrasicholina gloria</i>	DD	M
Engraulidae	<i>Encrasicholina heteroloba</i>	LC	M
Engraulidae	<i>Encrasicholina intermedia</i>	DD	M
Engraulidae	<i>Encrasicholina macrocephala</i>	DD	M
Engraulidae	<i>Encrasicholina oligobranchus</i>	DD	M
Engraulidae	<i>Encrasicholina pseudoheteroloba</i>	LC	M
Engraulidae	<i>Encrasicholina punctifer</i>	LC	M
Engraulidae	<i>Encrasicholina purpurea</i>	LC	E
Engraulidae	<i>Engraulis albidus</i>	DD	E
Engraulidae	<i>Engraulis anchoita</i>	NT A2bd	M
Engraulidae	<i>Engraulis australis</i>	LC	E
Engraulidae	<i>Engraulis capensis</i>	LC	M
Engraulidae	<i>Engraulis encrasicolus</i>	LC	E
Engraulidae	<i>Engraulis eurystole</i>	LC	M
Engraulidae	<i>Engraulis japonicus</i>	LC	M
Engraulidae	<i>Engraulis mordax</i>	LC	M
Engraulidae	<i>Engraulis ringens</i>	LC	M
Engraulidae	<i>Jurengraulis juruensis</i>	LC	F
Engraulidae	<i>Lycengraulis batesii</i>	LC	E
Engraulidae	<i>Lycengraulis figueiredoi</i>	LC	F
Engraulidae	<i>Lycengraulis grossidens</i>	LC	E
Engraulidae	<i>Lycengraulis limnichthys</i>	DD	E
Engraulidae	<i>Lycengraulis poeyi</i>	LC	E
Engraulidae	<i>Lycotrissa crocodilus</i>	LC	F
Engraulidae	<i>Papuengraulis micropinna</i>	DD	E

FAMILY	SPECIES NAME	GLOBAL CATEGORY & CRITERIA	SYSTEM
Engraulidae	<i>Pseudosetipinna haizhouensis</i>	DD	M
Engraulidae	<i>Pterengraulis atherinoides</i>	LC	E
Engraulidae	<i>Setipinna breviceps</i>	LC	E
Engraulidae	<i>Setipinna brevifilis</i>	DD	F
Engraulidae	<i>Setipinna melanochir</i>	DD	E
Engraulidae	<i>Setipinna paxtoni</i>	DD	M
Engraulidae	<i>Setipinna phasa</i>	LC	E
Engraulidae	<i>Setipinna taty</i>	LC	E
Engraulidae	<i>Setipinna tenuifilis</i>	DD	E
Engraulidae	<i>Setipinna wheeleri</i>	DD	F
Engraulidae	<i>Stolephorus advenus</i>	DD	M
Engraulidae	<i>Stolephorus andhraensis</i>	LC	E
Engraulidae	<i>Stolephorus apiensis</i>	LC	M
Engraulidae	<i>Stolephorus babarani</i>	DD	M
Engraulidae	<i>Stolephorus baganensis</i>	LC	M
Engraulidae	<i>Stolephorus bataviensis</i>	DD	M
Engraulidae	<i>Stolephorus baweanensis</i>	DD	M
Engraulidae	<i>Stolephorus bengalensis</i>	LC	M
Engraulidae	<i>Stolephorus brachycephalus</i>	LC	E
Engraulidae	<i>Stolephorus carpentariae</i>	LC	E
Engraulidae	<i>Stolephorus chinensis</i>	LC	E
Engraulidae	<i>Stolephorus commersonii</i>	LC	M
Engraulidae	<i>Stolephorus continentalis</i>	DD	M
Engraulidae	<i>Stolephorus dubiosus</i>	LC	E
Engraulidae	<i>Stolephorus holodon</i>	LC	E
Engraulidae	<i>Stolephorus indicus</i>	LC	E
Engraulidae	<i>Stolephorus insignis</i>	DD	M
Engraulidae	<i>Stolephorus multibranchus</i>	DD	M
Engraulidae	<i>Stolephorus nelsoni</i>	DD	E
Engraulidae	<i>Stolephorus oceanicus</i>	DD	M
Engraulidae	<i>Stolephorus pacificus</i>	DD	M
Engraulidae	<i>Stolephorus ronquilloi</i>	DD	E
Engraulidae	<i>Stolephorus shantungensis</i>	DD	E
Engraulidae	<i>Stolephorus tamilensis</i>	DD	M
Engraulidae	<i>Stolephorus teguhi</i>	DD	E
Engraulidae	<i>Stolephorus tri</i>	DD	M
Engraulidae	<i>Stolephorus waitei</i>	DD	M
Engraulidae	<i>Thryssa adelae</i>	DD	M
Engraulidae	<i>Thryssa aestuaria</i>	LC	E
Engraulidae	<i>Thryssa baelama</i>	LC	E

FAMILY	SPECIES NAME	GLOBAL CATEGORY & CRITERIA	SYSTEM
Engraulidae	<i>Thryssa brevicauda</i>	LC	E
Engraulidae	<i>Thryssa chefuensis</i>	DD	E
Engraulidae	<i>Thryssa cultella</i>	DD	M
Engraulidae	<i>Thryssa dayi</i>	DD	M
Engraulidae	<i>Thryssa dussumieri</i>	LC	E
Engraulidae	<i>Thryssa encrasicholoides</i>	DD	M
Engraulidae	<i>Thryssa gautamiensis</i>	DD	E
Engraulidae	<i>Thryssa hamiltonii</i>	LC	E
Engraulidae	<i>Thryssa kammalensis</i>	DD	E
Engraulidae	<i>Thryssa kammalensoides</i>	DD	E
Engraulidae	<i>Thryssa malabarica</i>	DD	E
Engraulidae	<i>Thryssa marasriae</i>	LC	E
Engraulidae	<i>Thryssa mystax</i>	LC	E
Engraulidae	<i>Thryssa polybranchialis</i>	DD	M
Engraulidae	<i>Thryssa purava</i>	DD	M
Engraulidae	<i>Thryssa rastrosa</i>	EN B1ab(i,ii,iii,v)+2ab(i,ii,iii,v)	F
Engraulidae	<i>Thryssa scratchleyi</i>	DD	E
Engraulidae	<i>Thryssa serena</i>	DD	M
Engraulidae	<i>Thryssa setirostris</i>	LC	E
Engraulidae	<i>Thryssa spinidens</i>	DD	M
Engraulidae	<i>Thryssa stenosoma</i>	DD	M
Engraulidae	<i>Thryssa vitrirostris</i>	LC	E
Engraulidae	<i>Thryssa whiteheadi</i>	LC	M
Pristigasteridae	<i>Chirocentrodon bleekermanus</i>	LC	E
Pristigasteridae	<i>Ilisha africana</i>	LC	E
Pristigasteridae	<i>Ilisha amazonica</i>	LC	F
Pristigasteridae	<i>Ilisha compressa</i>	LC	M
Pristigasteridae	<i>Ilisha elongata</i>	LC	E
Pristigasteridae	<i>Ilisha filigera</i>	DD	E
Pristigasteridae	<i>Ilisha fuerthii</i>	LC	E
Pristigasteridae	<i>Ilisha kampeni</i>	LC	E
Pristigasteridae	<i>Ilisha lunula</i>	DD	E
Pristigasteridae	<i>Ilisha macrogaster</i>	DD	E
Pristigasteridae	<i>Ilisha megaloptera</i>	LC	E
Pristigasteridae	<i>Ilisha melastoma</i>	LC	E
Pristigasteridae	<i>Ilisha novacula</i>	LC	F
Pristigasteridae	<i>Ilisha obfuscata</i>	DD	M
Pristigasteridae	<i>Ilisha pristigastroides</i>	DD	E
Pristigasteridae	<i>Ilisha sirishai</i>	DD	M

FAMILY	SPECIES NAME	GLOBAL CATEGORY & CRITERIA	SYSTEM
Pristigasteridae	<i>Ilisha striatula</i>	DD	E
Pristigasteridae	<i>Neopisthopterus cubanus</i>	VU B2ab(i,ii,iii)	E
Pristigasteridae	<i>Neopisthopterus tropicus</i>	LC	E
Pristigasteridae	<i>Odontognathus compressus</i>	LC	E
Pristigasteridae	<i>Odontognathus mucronatus</i>	LC	E
Pristigasteridae	<i>Odontognathus panamensis</i>	LC	E
Pristigasteridae	<i>Opisthopterus dovii</i>	LC	E
Pristigasteridae	<i>Opisthopterus effulgens</i>	DD	E
Pristigasteridae	<i>Opisthopterus equatorialis</i>	LC	M
Pristigasteridae	<i>Opisthopterus macrops</i>	LC	M
Pristigasteridae	<i>Opisthopterus tardoore</i>	LC	E
Pristigasteridae	<i>Opisthopterus valenciennesi</i>	DD	E
Pristigasteridae	<i>Pellona castelnaeana</i>	LC	E
Pristigasteridae	<i>Pellona dayi</i>	DD	M
Pristigasteridae	<i>Pellona ditchela</i>	LC	E
Pristigasteridae	<i>Pellona flavipinnis</i>	LC	F
Pristigasteridae	<i>Pellona harroweri</i>	LC	E
Pristigasteridae	<i>Pliosteostoma lutipinnis</i>	LC	E
Pristigasteridae	<i>Pristigaster cayana</i>	LC	F
Pristigasteridae	<i>Pristigaster whiteheadi</i>	LC	F
Pristigasteridae	<i>Raconda russeliana</i>	LC	E
Sundasalangidae	<i>Sundasalanx malletti</i>	DD	F
Sundasalangidae	<i>Sundasalanx megalops</i>	DD	F
Sundasalangidae	<i>Sundasalanx mekongensis</i>	LC	F
Sundasalangidae	<i>Sundasalanx mesops</i>	DD	F
Sundasalangidae	<i>Sundasalanx microps</i>	DD	F
Sundasalangidae	<i>Sundasalanx platyrhynchus</i>	DD	F
Sundasalangidae	<i>Sundasalanx praecox</i>	LC	F