

Importance of U.S. West Coast Shellfish

West Coast shellfish (wild-harvest and aquaculture) provide significant socioeconomic and recreational benefits to coastal communities in Washington, Oregon and California, in addition to providing nutritious, delicious seafood. Equally important, problems with shellfish portend potentially significant changes in marine ecosystems.

Wild harvest shellfish - Average annual (1998-2009) revenue to fishermen (dockside value) from shellfish fisheries is \$159 million for Washington, Oregon and California combined.

Shellfish contribute about 50% of total commercial fishery dockside revenue in Oregon and Washington, and about 60% in California.

- 🐚 In Washington, Oregon and Northern California, the most important species for commercial fisheries are Dungeness crab and Pacific Ocean (pink) shrimp.
- 🐚 In Southern California, market squid, sea urchin, spiny lobster and spot prawn are important commercial fisheries. Shellfish represent 62% of total commercial fishery revenue in this region with most of that value contributed by squid.
- 🐚 Recreationally important species include clams, oysters and geoducks in Washington and Oregon, abalone in Northern California, and spiny lobster in Southern California.

Aquaculture shellfish production - Annual producer revenue is \$117 million to coastal economies in Washington, Oregon and California (\$278 million wholesale).

- 🐚 Accounts for 3,000+ jobs in coastal communities.
- 🐚 Cultured species are oysters, clams, geoducks, mussels and scallops.
- 🐚 Pacific oyster production (a 2-4 year crop) declined 22% between 2005 and 2009.
- 🐚 In Willapa Bay, Washington, the largest oyster-producing region on the West Coast, wild oyster larvae failed to "set" in six consecutive years, resulting in steep declines in harvest rates.
- 🐚 In Washington and Oregon, two of the three major West Coast oyster seed hatcheries experienced production declines of up to 80% from 2006 to 2009.



Recommendations and Next Steps...

West Coast Ocean Acidification-Shellfish Workshop participants reviewed existing research, identified information gaps, and proposed modifications and potential solutions to improve the state of knowledge of ocean acidification on the West Coast and to address problems faced by shellfish hatcheries, growers and harvesters.

Recommendations included:

(1) A West Coast Ocean Acidification Research Coordination Working Group* should be established to provide recommendations on best practices and standardized research methods to achieve a coordinated network of ocean observations and experimental studies, in order to observe, understand and quantify the evolving threats posed by ocean acidification along the West Coast of the United States.

(2) A West Coast Ocean Acidification Data Exchange, integrated with IOOS, and consisting of a single website that serves as an entry portal to data and information, should be established to provide easy access to available data sets and information regarding ocean acidification processes affecting West Coast fish and shellfish.

(3) The proposed NOAA West Coast Ocean Acidification Observing Network should be enhanced and expanded to include sites on the West Coast where in-field biological response studies are conducted.

The workshop organizing committee thanks all the workshop participants and invites others to participate in future efforts.

*Nine participants volunteered to serve on a steering committee to develop the framework for the working group.

All workshop materials and related information are posted online at:
www.sccwrp.org/Meetings/Workshops/OceanAcidificationWorkshop.aspx

Workshop Organizing Committee:

Robin Downey, Pacific Coast Shellfish Growers Association
Richard Feely, NOAA Pacific Marine Environmental Laboratory
Russ Moll, California Sea Grant
Penny Dalton, Washington Sea Grant
Steve Brandt, Oregon Sea Grant
Phyllis Grifman, USC Sea Grant
Diane Pleschner-Steele, California Wetfish Producers Association
Bruce Steele, California sea urchin diver
Libe Washburn, University of California, Santa Barbara
Stephen Weisberg, Southern California Coastal Water Research Project Authority

Workshop sponsors:



Photography courtesy of: Richard Wilson (cover/background: Willapa Bay, WA), Washington Department of Fish and Wildlife (cover: clamming) Pauline Yu (pluteus-stage sea urchin larvae), Pacific Coast Shellfish Growers Association (shellfish), Dave Rudie (urchin diver), Washington Sea Grant (wave), Richard Feely, Dana Greeley, Dr. Steve Schroeter, Jesse Vance

Is Ocean Acidification Affecting Shellfish?

A NOAA Sea Grant West Coast Workshop Seeks Answers

July 7-8, 2010, Costa Mesa, CA

*No new Pacific oysters recruiting to oyster "beds."
Shellfish larvae dying at commercial hatcheries.
Corrosive, acidified seawater measured for the
first time off the Pacific Coast.*

These and other observations, beginning in 2005, prompted members of the West Coast shellfish industry to seek help from scientists to explore what is causing the shellfish losses, what role ocean acidification and other factors might be playing, and how to adapt to sustain West Coast shellfish resources.

An organizing committee planned a face-to-face meeting as a logical first step to frame the problems, assess what information is available and what is needed, and to suggest future actions.

The West Coast Ocean Acidification-Shellfish Workshop was held July 7-8, 2010, in Costa Mesa, CA, with 51 invited participants, including biologists, oceanographers, state and federal agency managers, industry members and representatives from the Sea Grant Programs in Washington, Oregon and California.

This brochure summarizes the information shared and recommended next steps.




Increasing levels of carbon dioxide (CO₂) emissions into the atmosphere, a consequence of fossil fuel use, are causing immediate, measurable impacts on ocean chemistry.

About one-third of the carbon dioxide produced every day around the world is absorbed by the oceans. As CO₂ reacts with seawater, it lowers seawater pH and reduces the concentration of carbonate ions, an essential component in the calcium carbonate that makes up the shells of shellfish and the skeletons of corals. Recently scientists have begun measuring changes in ocean chemistry and investigating the possible consequences to marine life, food webs and people. These studies find that ocean acidification is ongoing and may have significant biological impacts. The West Coast is vulnerable to the enhanced ocean acidification associated with seasonal upwelling, potentially causing serious impacts to ecosystems and some recreationally and commercially important shellfish.

Consider:

- Low pH seawater found in the deep ocean, low enough to be corrosive to the shells of some marine organisms, is expanding toward the ocean surface at the rate of 1-2 meters/year in the North Pacific.
- Some of the most vulnerable species may be tiny, shell-forming animals at the base of the food web, which provide food for many larger species.
- Also at risk are some shellfish commonly grown or harvested for human consumption, including oysters and sea urchins.
- Scientists are now conducting laboratory studies to “stress test” marine larvae (including oysters, abalone, sea urchins and other shellfish) at different CO₂ levels. Studies indicate significantly reduced shell growth for oysters and urchins in higher CO₂ environments.
- Options for an organism faced with environmental stress, such as ocean acidification, include migration, acclimatization (tolerance), adaptation (which may take generations if possible at all) and extinction.
- Demonstrated effects of ocean acidification on some species include reduced tolerance to temperature increase, impaired reproduction, inefficient cell function, impaired growth, larval shell dissolution, disease susceptibility and higher mortality rates at early life stages.
- Generally, there is a higher “cost of living” for many organisms subjected to a high CO₂ marine environment.



A 2007 NOAA research cruise measured evidence of acidified seawater in surface waters off the Pacific Coast. Observed levels of “corrosive” seawater had not been predicted to occur for another 100 years. diagram: courtesy of Richard A. Feely and Dana Greeley, NOAA/PMEL

Shellfish Industry Challenges

Shellfish recruitment problems challenging West Coast growers and harvesters may be associated with ocean acidification, *Vibrio tubiashii* infections, low-oxygen “dead zones,” or all three in combination. More research is clearly needed. Research and observations include:

- A 20-year study in California found significant differences in annual larval settlement of red and purple sea urchins, crabs and clams between northern and southern sites.
- In Northern California, where high upwelling is common in spring and summer, sea urchin larval settlement declined during upwelling periods.
- Oyster larval rearing problems were first observed at Hatfield Marine Science Center at Oregon State University in 2005.
- Whiskey Creek Hatchery in Oregon, and Taylor Shellfish Hatchery in Washington, subsequently experienced massive larval mortalities leading to steep declines in production over the past four years.

Shellfish Industry Response

Shellfish growers launched an initiative to research and develop adaptive solutions. Some hatcheries, including Whiskey Creek, now employ continuous seawater monitoring systems. Potential hatchery remedies to avoid the effects of acidified seawater:

- Install water monitoring and treatment systems for seawater intakes
- Pump seawater into hatcheries only during low CO₂ periods
- Culture larvae during periods with no upwelling
- Engineer treatment systems to modify seawater chemistry for rearing shellfish larvae
- Investigate evidence of local adaptation and vulnerability to changes in water chemistry and develop resistant oyster larval strains



Canary in the coal mine? Pacific oysters in Willapa Bay, Washington, have experienced six years of recruitment failure. Similar periods of low recruitment have been recorded in the past but the cause is unknown. photo: courtesy of Richard Wilson



Common scrub brushes have proved to be efficient and cost-effective collectors to study larval settlement in marine field studies. photo: courtesy of Dr. Steve Schroeter



Whiskey Creek and other hatcheries have installed continuous water monitoring and treatment systems. photo: courtesy of Jesse Vance

Mounting evidence suggests that ocean acidification may change the structure, function and biodiversity of marine ecosystems. In 2009, Congress passed the Federal Ocean Acidification Research and Monitoring (FOARAM) Act, authorizing federal ocean acidification research funding through NOAA, NASA and the National Science Foundation. The Act also established an interagency committee to develop a national ocean acidification strategic research and monitoring plan, expected to be completed in spring 2011. The national strategic plan process is now seeking recommendations, and this workshop is an important step toward developing a regional plan for the West Coast.

Future Collaboration

Addressing questions about ocean acidification requires integration of:

- ocean observing measurements
- laboratory exposure studies
- shellfish recruitment and production data
- field studies of organism performance in relation to ocean conditions

These data are collected by different sectors that to date have had limited interaction.

The West Coast Ocean Acidification-Shellfish Workshop stimulated discussion among a balanced group representing academia, government and industry and laid the groundwork for future collaboration.

Significant workshop findings:

Existing water quality and biological data sets are physically and spatially disassociated.

There is a need to:

- coordinate chemical data collection at the best biological monitoring sites, leveraging long-term data sets and employing standardized methods.
- supplement correlative data with an understanding of biological processes.
- develop predictive models at smaller scales.

Different communities must work collaboratively to resolve these issues.

The momentum generated at this workshop needs to be maintained.

