



JUN - 7 2011

To All Interested Government Agencies and Public Groups:

Under the National Environmental Policy Act, an environmental review has been performed on the following action.

TITLE: Amendment 2 to the Fishery Ecosystem Plan for Pelagic Fisheries of the Western Pacific Region, as Partially Approved; Prohibiting Longline Fishing within 30 nm of the CNMI (RIN 0648-AW67)

LOCATION: U.S. EEZ around the Commonwealth of the Northern Mariana Islands (CNMI)

SUMMARY: The Western Pacific Fishery Management Council (Council) prepared Amendment 2 to address pelagic fishing concerns in the Mariana Islands. Pelagic fisheries in federal waters around the Mariana Islands (Guam and the CNMI) consist of small trolling fleets, a few pelagic longline vessels, and purse seine vessels based there but not fishing near the islands. After considering a range of alternatives and deliberating the proposal in public meetings, the Council recommended prohibiting longline fishing within 30 nm of the CNMI and U.S. purse seine vessels from fishing within the EEZ around Guam and the CNMI.

NMFS proposes to partially approve Amendment 2. NMFS proposes to approve the portion of the amendment that establishes a longline fishing prohibited area in the CNMI (Alternative 1C). NMFS has determined that the proposed purse seine measures are not consistent with National Standard 2 of the Magnuson-Stevens Fishery Conservation and Management Act (MSA) and, therefore, are subject to disapproval. Because NMFS will not approve the portions of Amendment 2 relating to prohibitions on purse seine fishing, NMFS' disapproval effectively implements the relevant no-action alternatives (Alternatives 2A and 3A).

The proposed action will not result in a substantial change to subject fisheries. No increases in the number of vessels, the intensity of fishing, or large changes to where vessels fish are expected. As a result, no significant affects on the human environment are expected.

The combined FEP Amendment and EA, identified by RIN 0648-AW67, is available from www.regulations.gov ; or by mail at the following address:

RESPONSIBLE


OFFICIAL: Michael D. Tosatto
Regional Administrator
Pacific Islands Region
National Marine Fisheries Service, NOAA
1601 Kapiolani Blvd. 1110
Honolulu, HI 96814
Tel (808) 944-2200; Fax (808) 973-2941



The environmental review process led us to conclude that approving and implementing a prohibited area for longline fishing within 30-nm around the CNMI will not have a significant effect on the human environment. Therefore, an environmental impact statement will not be prepared. A copy of the finding of no significant impact (FONSI) including the supporting environmental assessment is enclosed for your information.

Although NOAA is not soliciting comments on this completed EA/FONSI, we will consider any comments submitted that would assist us in preparing future NEPA documents. Please submit any written comments to the Responsible Official named above.

Sincerely,

A handwritten signature in blue ink, appearing to read "P. Doremus", with a stylized flourish at the end.

Paul N. Doremus, Ph.D.
NOAA NEPA Coordinator

Enclosure



**Amendment 2 to the Fishery Ecosystem Plan for Pelagic Fisheries
of the Western Pacific Region**

**Establishment of Longline and Purse Seine Prohibited Areas
in the Mariana Archipelago**

**Including an Environmental Assessment, Regulatory Impact Review, and
Preliminary Regulatory Flexibility Act Analysis**

March 4, 2011

**Western Pacific Regional Fishery Management Council
1164 Bishop St. Suite 1400
Honolulu, HI 96813**

**Environmental Assessment Summary
For**

Establishment of Longline and Purse Seine Prohibited Areas in the Mariana Archipelago

**Amendment 2 to the Pelagics Fishery Ecosystem Plan for
Fisheries of the Western Pacific Region**

March 4, 2011

Responsible Agency:

Pacific Islands Region
National Marine Fisheries Service (NMFS)
National Oceanographic and Atmospheric Administration (NOAA)
Honolulu, Hawaii

Responsible Official: Michael D. Tosatto

Regional Administrator
Pacific Islands Region, NMFS, NOAA

For Further Information Contact:

Michael D. Tosatto
NMFS Regional Administrator
1601 Kapiolani Blvd. 1110
Honolulu, HI 96814
(808) 944-2200

Responsible Council and Contact:

Kitty M. Simonds
Executive Director
Western Pacific Fishery Management Council
1164 Bishop St. Suite 1400
Honolulu, HI 96813
(808) 522-8220

Project No.: RIN 0648-AW67

Abstract

Amendment 2 to the Pelagics Fishery Ecosystem Plan was recommended by the Western Pacific Fishery Management Council (Council) and would provide for the establishment of a longline fishing prohibited area in the Commonwealth of the Northern Mariana Islands (CNMI) and purse seine fishing prohibited areas in the U.S. EEZ around the CNMI and Guam. The Council recommendations would promote sustained participation in fishing by the Guam and CNMI fishing communities which are made up almost exclusively of small vessel trollers that have a strong cultural and economic dependence on inshore pelagic catches. In CNMI, the troll fishery is vulnerable to potential catch competition and gear conflicts with longliners in areas where both fleets fish. To reduce potential conflicts, the Council is recommending a 30 nm prohibited area for longline fishing around the CNMI.

Although no purse seine catches have been reported from within the EEZ waters around either Guam or CNMI in recent years, the Council is proposing to prevent purse seine fishing catches around the CNMI and Guam to minimize gear conflicts and catch competition among fishery sectors, while addressing potential unsustainable catches of undersized bigeye tuna. This would have an added benefit of potentially reducing catch competition for skipjack tuna, which the troll fishery targets and reducing the potential for impacts of purse seining on longliners.

Amendment 2 includes an environmental assessment (EA) that was prepared by the Council in close coordination with NMFS. The proposed action would result in a geographic separation of some fishing fleets. None of the proposed alternatives for the proposed fishing prohibited areas were found to result in a large change to troll, longline or purse seine fishing including the number of vessels used in the fishery, where the vessels fish, the time of year they fish, or gear used. There is currently no purse seine fishing occurring in the U.S. EEZ around the Mariana Islands and purse seiners would continue to fish in other areas. Because there would not be any large changes to fishing operations as a result of implementing prohibited fishing areas, none of the alternatives would result in overfishing of target, non-target, bycatch stocks or new impacts to protected species or ocean habitats.

NMFS is seeking public comments on Amendment 2 and on the analysis of potential environmental impacts of the proposed prohibited areas. The proposed rule and instructions on how to comment on the document, and how to obtain copies of the EA can be found by searching on RIN 0648-AW67 at www.regulations.gov; or by contacting the responsible official at the above address.

1.0 Executive Summary

The Western Pacific Fishery Management Council (Council) prepared this amendment to the Fishery Ecosystem Plan for Pelagic Fisheries of the Western Pacific Region (Pelagic FEP) to balance the needs of the small and locally-based pelagic fishing fleets of the Mariana Archipelago while 1) allowing for development of a longline fishery around the Commonwealth of the Northern Mariana Islands (CNMI), and 2) minimizing localized depletion¹ and catch competition between U.S. purse seine vessels and other pelagic fishing fleets operating in the U.S. EEZ around the CNMI and Guam.

The troll fleet is the main small-scale pelagic fishery in the Mariana Islands and was estimated to consist of over 430 vessels in 2008. It is of historic, cultural, and economic importance to the people of the Mariana Islands. The troll fleet mainly targets skipjack and yellowfin tuna within 30 nm of shore among the offshore banks and seamounts that tend to congregate pelagic fish. Pelagic longline, another important local pelagic fishing technique, has only recently begun in Guam and the CNMI targeting skipjack and yellowfin tuna. There is growing interest to increase longline fishing in EEZ waters around the Mariana Archipelago. Currently, seven longline vessels have western Pacific general longline permits to fish in the U.S. EEZ around the Mariana Archipelago.

The Council is concerned that future expansion of longline fishing around the Mariana Archipelago may result in adverse impacts to the troll fleet and the communities that rely on a continued supply of locally-caught fresh fish. The trolling fleet would likely be impacted by gear conflicts, localized stock depletion and competition for target species if longline vessels were to fish in the nearshore waters where the trollers fish, as occurred in Hawaii and American Samoa waters and which led to the establishment of large vessel prohibited areas around those areas. To address this concern, the Council is recommending the establishment of a longline fishing prohibited area in portions of the U.S. EEZ around the CNMI to address potential gear conflicts, localized stock depletion, and competition for fishery resources between the local CNMI pelagic trolling and charter fleets, and the developing longline fishery.

Since 2005 the U.S. purse seine fleet operating in the western and central Pacific Ocean (WCPO) increased from 15 to 36 vessels and U.S. landings of skipjack tuna has doubled. There is potential for U.S. purse seine vessels to begin fishing in U.S. EEZ waters around the Mariana Archipelago since catch competition for skipjack tuna with foreign purse seine fleets of the WCPO has increased in recent years. Due to the potentially large harvests that can be made by purse seiners, coupled with their inability to target specific and mature fish, an emergence of purse seine fishing in the U.S. EEZ around the Mariana Archipelago has the potential to negatively impact the viable continuation of the local troll and longline fisheries. Purse seining may result in localized, temporary stock depletion and catch competition with local longline and troll fisheries that target the same pelagic species. In addition, concerns surround the catch of bigeye tuna by the U.S. purse seine fleet. Overfishing the WCPO bigeye tuna stock has prompted the WCPO tuna regional fishery management organization (RFMO), the Western and Central

¹ Localized fish depletion occurs when a stock in a small area is reduced by the removal of large amounts of fish, thereby temporarily depleting the availability of the stock to fishing activity or other predators in that area.

Pacific Fisheries Commission (WCPFC) to aim for a 30 percent reduction in catch and has led to longline fishery bigeye catch limits whereas the U.S. purse seine catch more than doubled from 2007 to 2008. Furthermore, if purse seine fishing and longline fishing activities were to occur in close proximity there is potential for gear conflicts as longline and purse seine fishing gear are both extensive in size and can drift between deployment and retrieval.

Fishery management alternatives were developed that considered the needs and concerns of CNMI's and Guam's small and locally-based pelagic troll fishing fleets with the development of longline fisheries in waters around the Mariana Archipelago and purse seine fishing in the western Pacific, are described and their environmental impacts analyzed in this document. The Council recommended implementation of Alternative 1C, a 30-nm longline prohibited area around the CNMI. Alternative 1C was selected as the Council's preferred alternative because it would best protect the needs of the small boat fleet while allowing for development of a longline fishery around the CNMI.

To reduce the potential for localized depletion and catch competition between purse seine vessels and the pelagic troll and longline fishing fleets of CNMI and Guam, the Council recommended implementation of Alternatives 2D and 3D that would establish a purse seine fishing prohibited area to include all U.S. EEZ waters around the CNMI and Guam, respectively. The Council determined that Alternatives 2D and 3D would best-protect the fishery resources within the EEZ while providing for the viability of the local troll and longline fleets around CNMI and Guam. .

The environmental analysis showed that none of the alternatives would have an adverse impact on resources of management concern including target stocks, non-target stocks, bycatch or protected resources. Current levels of troll and longline fishing in CNMI and Guam are expected to continue under all alternatives, and the purse seiners will continue their fishing in other areas of the western Pacific. There is currently no purse seine fishing reported within the U.S. EEZs around the Mariana Islands. The analysis found that none of the alternatives would result in a change of fishing intensity by any of the three fleets. Furthermore, although the Council and NMFS are also considering an expanded purse seine fishing prohibited area around American Samoa, the domestic purse seine fleet will continue to be able to fish in the U.S. EEZ around the Pacific remote island areas, in foreign EEZ waters under existing fishing agreements, and in the high seas. The proposed closure of the U.S. EEZ waters of the Mariana Archipelago to purse seine fishing under the Council's preferred alternative is not expected to result in a large change to current fishing intensity or locations by purse seiners. The analysis also found no large or adverse cumulative effects of the proposed alternatives on the environment or on any of the three pelagic fisheries. The proposed action and alternatives were considered in light of a potential purse seine prohibited area in the American Samoa EEZ, recent closure of portions of recently established Pacific marine national monuments to commercial fishing, potential FAD management measures, FAD seasonal closures, and the military buildup on Guam. These past, present, and reasonably foreseeable actions by the Council and NMFS and others would not compound the very low expected effects of the proposed action or alternatives on the environment. The analysis did not show that there were climate change impacts or impacts of the proposed alternatives on climate change, nor were there and disproportionately high and adverse impacts on members of Environmental Justice populations, or marine protected areas.

2.0 Table of Contents

1.0 Executive Summary	4
2.0 Table of Contents	6
2.1 List of Figures	10
2.2 List of Tables	11
3.0 Introduction	12
4.0 Purpose and Need for Action	16
5.0 Initial Actions	17
6.0 Objective	18
7.0. Description of Alternatives	19
7.1 Topic 1: CNMI Longline Prohibited Area	19
7.1.1 Alternative 1A: No Action	19
7.1.2 Alternative 1B: 25 nm Longline Prohibited Area	19
7.1.3 Alternative 1C: 30 nm Longline Prohibited Area (Preferred)	19
7.1.4 Alternative 1D: 50 nm Longline Prohibited Area for Large Vessels	19
7.1.5 Alternative 1E: 75 nm Longline Prohibited Area	19
7.1.6 Alternative 1F: 100 nm Longline Prohibited Area	19
7.2 Topic 2: CNMI Purse Seine Prohibited Area	20
7.2.1 Alternative 2A: No Action	20
7.2.2 Alternative 2B: 50 nm Purse Seine Prohibited Area	20
7.2.3 Alternative 2C: 100 nm Purse Seine Prohibited Area	20
7.2.4 Alternative 2D: Entire EEZ Purse Seine Prohibited Area (Preferred)	20
7.3 Topic 3: Guam Purse Seine Prohibited Area	20
7.3.1 Alternative 3A: No Action	20
7.3.2 Alternative 3B: 50nm Purse Seine Prohibited Area	20
7.3.3 Alternative 3C: 100 nm Purse Seine Prohibited Area	20
7.3.4 Alternative 3D: Entire EEZ Purse Seine Prohibited Area (Preferred)	20
8.0 Description of the Affected Environment	20
8.1 U.S. EEZ around the Mariana Archipelago and the Western Pacific Warm Pool	20
8.2 Pelagic Fishing in CNMI	22
8.3 Pelagic Fishing in Guam	30
8.4 WCPO Purse Seine Tuna Fishery	42
8.4.1 U.S. Purse Seine Fishery-History	54
8.4.2 U.S. Fishing in the PRIA	55
8.4.3 U.S. Purse Seine Fishery-Present and Future	57
8.4.4 Fishery Interactions Between Various Gear Types	61
8.4.5 Foreign Fishing in U.S. EEZ Waters	63
8.5 Protected Species	64
8.5.1 Sea Turtles	64
8.5.2 Marine Mammals	69
8.5.3 Seabirds	71
8.6 Status of Target Stocks	73
8.7 Regional Tuna Fisheries Management	76
8.8 CNMI Longline Fishing	80

8.9 Climate Change Background	82
9.0 Impacts of the Alternatives	83
9.1 Topic 1: CNMI Longline Prohibited Area.....	83
9.1.1 Impacts on Target Stocks.....	83
9.1.2 Impacts on Non-target Stocks.....	84
9.1.3 Impacts on Protected Species.....	85
9.1.4 Impacts on Marine Habitat.....	86
9.1.5 Impacts on Public Health and Safety	86
9.1.6 Impacts on Fishery Participants and Fishing Communities.....	87
9.1.7 Impacts on Biodiversity and Ecosystem Functions	88
9.1.8 Impacts on Administration and Enforcement	88
9.1.9 Cumulative Effects of the Proposed CNMI Longline Prohibited Areas.....	89
9.1.9 Environmental Justice Effects of the Proposed CNMI Longline Prohibited Areas.....	90
9.1.10 Climate Change Impacts of the Proposed CNMI Longline Prohibited Areas.....	90
9.1.11.....	91
9.2 Topic 2: CNMI Purse Seine Prohibited Area	92
9.2.1 Impacts on Target Stocks.....	92
9.2.2 Impacts on Non-target Stocks.....	93
9.2.3 Impacts on Protected Species.....	94
9.2.4 Impacts on Marine Habitat.....	95
9.2.5 Impacts on Public Health and Safety	96
9.2.6 Impacts on Fishery Participants and Communities.....	96
9.2.7 Impacts on Biodiversity and Ecosystem Functions	98
9.2.8 Impacts on Administration and Enforcement	98
9.2.9 Cumulative Effects of the Proposed CNMI Purse Seine Prohibited Areas	99
9.2.10 Environmental Justice Effects of the Proposed CNMI Purse Seine Prohibited Areas	99
9.2.11 Climate Change Impacts of the Proposed CNMI Purse Seine Prohibited Areas.....	99
9.2.12 Reasons the Council Selected the CNMI Purse Seine Prohibited Area as its Preferred Alternative.....	99
9.3 Topic 3: Impacts of Proposed Alternatives for a Guam Purse Seine Prohibited Area	101
9.3.1 Impacts of the Proposed Guam Purse Seine Prohibited Areas on Target Stocks	101
9.3.2 Impacts on Non-target Stocks.....	103
9.3.3 Impacts on Protected Species.....	103
9.3.4 Impacts on Marine Habitat.....	104
9.3.5 Impacts on Public Health and Safety	104
9.3.6 Impacts on Fishery Participants and Communities.....	105
9.3.7 Impacts on Biodiversity and Ecosystem Functions	107
9.3.8 Impacts on Administration and Enforcement	107
9.3.9 Cumulative Effects of the Proposed Guam Purse Seine Prohibited Areas.....	107
9.3.10 Environmental Justice Effects of the Proposed Guam Purse Seine Prohibited Areas	107
9.3.11 Climate Change Impacts of the Proposed Guam Purse Seine Prohibited Areas	108
9.3.12 Reasons the Council Selected the Guam Purse Seine Prohibited Area as the Preferred Alternative.....	108
9.4 Essential Fish Habitat	109

9.7	Other environmental consequences	111
10.0	Consistency with the MSA and Other Laws	111
10.1	Consistency with National Standards	111
10.2	National Environmental Policy Act	115
10.3	Regulatory Impact Review	116
10.4	Administrative Procedure Act	117
10.5	Coastal Zone Management Act	117
10.6	Information Quality Act	117
10.7	Paperwork Reduction Act	117
10.8	Regulatory Flexibility Act	118
10.9	Endangered Species Act	118
10.10	Marine Mammal Protection Act	119
11.0	Proposed Regulations	120
12.0	References	126
	Appendix A	136
	Regulatory Impact Review and	136
	Preliminary Regulatory Flexibility Act Analysis	136
2.0	Purpose and Need for Action	137
3.0	Description of Alternatives Considered	138
	Three sets of alternatives were developed, each designed to provide long-term viability of CNMI and Guam’s small and locally-based fishing fleets, while balancing the needs and concerns of the fishing industry regarding the development of larger-scale domestic fisheries in U.S. EEZ waters around the Mariana Archipelago.	138
3.1	Topic 1: CNMI Longline Fishing Prohibited Area	138
	Alternative 1A: No Action	138
	Alternative 1B: 25 nm Longline Fishing Prohibited Area	138
	Alternative 1C: 30 nm Longline Fishing Prohibited Area (Preferred)	138
	Alternative 1D: 50 nm Longline Fishing Prohibited Area for Large Vessels	138
	Alternative 1E: 75 nm Longline Fishing Prohibited Area	138
	Alternative 1F: 100 nm Longline <i>Fishing</i> Prohibited Area	138
3.2	Topic 2: CNMI Purse Seine Fishing Prohibited Area	138
	Alternative 2A: No Action	138
	Alternative 2B: 50 nm Purse Seine Fishing Prohibited Area	139
	Alternative 2C: 100 nm Purse Seine Fishing Prohibited Area	139
	Alternative 2D: Entire U.S. EEZ Purse Seine Fishing Prohibited Area (Preferred)	139
3.3	Topic 3: Guam Purse Seine Fishing Prohibited Area	139
	Alternative 3A: No Action	139
	Alternative 3B: 50nm Purse Seine Fishing Prohibited Area	139
	Alternative 3C: 100 nm Purse Seine Fishing Prohibited Area	139
	Alternative 3D: Entire U.S. EEZ Purse Seine Fishing Prohibited Area (Preferred)	139
4.0	Description of Potentially Affected Fisheries	139
4.1	CNMI-based boats	139
	Table 1. Annual Participation, Catch and Revenue for CNMI-based Commercial Pelagic Vessels	141
4.2	Guam-based boats	142

Table 2. Annual Participation, Catch and Revenue for Guam-based Commercial Pelagic Vessels	143
4.3 WCPO Domestic Purse Seine Tuna Fishery.....	144
4.4 WCPO Domestic Purse Seine Fishery	145
5.0 Potential for Gear Conflicts between Purse Seine and Small Boat Fisheries	147
6.0 Impacts of the Alternatives	148
6.1 Topic 1: CNMI Longline Prohibited area.....	148
Table 4: Catch Rates for Longliners Operating in the FSM (lb/hook)	148
Table 5. Typical American Samoa Tuna Longline Vessel Operations.....	149
Table 6. Projected Annual Landings for One CNMI Longline Vessel.....	149
6.3 Topic 3: Guam Purse Seine Prohibited Area	152
7.0 Impacts of the Preferred Alternatives on National Costs and Benefits	153
Preliminary Regulatory Flexibility Act Analysis	154
1.0 Description of Small Businesses to Which the Rule Would Apply.....	154
2.0 Skills Necessary to Meet Compliance Requirements	155
3.0 Identification of Duplicating, Overlapping, and Conflicting Federal Rules.....	155

2.1 List of Figures

Figure 1: EEZ boundaries in the western Pacific.....	15
Figure 2: U.S. EEZ waters around the Mariana Archipelago.....	16
Figure 3: CNMI Annual Commercial Pelagic Trolling Landings 1983-2008.....	26
Figure 4: CNMI Annual Commercial Landings of Skipjack and Yellowfin Tuna by Trolling Vessels 1983-2008.....	27
Figure 5: CNMI Active Commercial Pelagic Trolling Vessels from 1983-2008.....	28
Figure 6: Number of Pelagic Fishing Trips (Trolling) in CNMI, 1983-2008.....	29
Figure 7: Trolling Catch Rates for Skipjack and Yellowfin Tunas in CNMI.....	29
Figure 8: Guam fishery landings by origin (2001-2006).....	34
Figure 9: Guam Annual Estimated Total Landings: All Pelagics, Tuna PMUS, and Non-Tuna PMUS 1980-2008.....	36
Figure 10: Guam Annual Estimated Total Tuna PMUS Landings 1982-2008.....	36
Figure 11: Guam Annual Estimated Total Skipjack Landings 1982-2008.....	37
Figure 12: Guam Estimated Number of Trolling Boats 1982-2008.....	38
Figure 13: Guam Estimated Number of Troll Trips.....	39
Figure 14: Guam Trolling CPUE (lb/hour) 1982-2008.....	39
Figure 15: Skipjack CPUE (lb/hour): All, Non-charter, and Charter Vessels 1982-2008.....	40
Figure 16 Yellowfin CPUE (lb/hour): All, Non-charter, and Charter Vessels 1982-2008.....	40
Figure 17: Wahoo CPUE (lb/hour): All, Non-charter, and Charter Vessels 1982-2008.....	41
Figure 18: Blue Marlin CPUE (lb/hour): All, Non-charter, and Charter Vessels 1982-2008.....	41
Figure 19: Mahimahi CPUE (lb/hour): All, Non-charter, and Charter Vessels 1982-2008.....	42
Figure 20: WCPO Purse Seine Vessels 1972-2008.....	43
Figure 21: Catches of tuna in the WCPFC Statistical Area, by species.....	44
Figure 22: Catches of tuna in the WCPFC Statistical Area, by gear type.....	44
Figure 23: Japanese Purse Seine Catch by Location, 2007.....	47
Figure 24: Japan Purse Seine Catch by Location, 2007 and 2008.....	47
Figure 25: Time series showing the percentage of total sets by school type for the major purse-seine fleets operating in the WCPO.....	49
Figure 26: The distribution of total purse seine effort in the WCPO, 2003-8, by days fishing....	50
Figure 27: The WCPFC Convention Area. The PNA Exclusive Economic Zones (EEZs) are ...	52
Figure 28: Overall Average Annual Catch from the EEZ Waters around Jarvis Island, Howland and Baker Islands, and Kingman Reef and Palmyra Atoll, and Average Annual Catch from within 50 nm, (i.e., Monument Waters), 1988-2007.....	56
Figure 29: Distribution of Effort by the U.S Purse Seine Fleet, 2007 and 2008.....	58
Figure 30: U.S. Purse Seine Catch Rates in the WCPO.....	58
Figure 31: U.S. Purse Seine SKJ Catch from Howland and Baker Islands, PRIA.....	59
Figure 32: Skipjack Catch by Gear (mt) in WCPO.....	76
Figure 33: Catch Composition of Longline Vessels Operating in the FSM.....	81

2.2 List of Tables

Table 1: Annual catches (mt) of bigeye (BET), blue marlin (BLZ), yellowfin (YFT) and skipjack (SKJ) by foreign fleets in the vicinity of CNMI (15°-25°N, 140°-150°E) 1962-1997.....	22
Table 2: CNMI 2008 Commercial Pelagic Trolling Landings, Revenues and Price.....	24
Table 3: CNMI Total Commercial Trolling Landings (lb) 1983-2008.....	25
Table 4: Annual catches (metric tons) of bigeye, blue marlin, yellowfin and skipjack by foreign fleets in the vicinity of Guam (10°-15°N, 140°-150°E), 1962-1997.	31
Table 5. Estimated Total Guam Pelagic Landings (lb) 1982-2008	34
Table 6: Guam 2008 Landings and Prices	35
Table 7: Estimated Number of Guam Trolling Vessels.....	37
Table 8: Numbers and Landings of Active DWFN Purse Seine Vessels in the WCPO, 2007.....	45
Table 9: Japan's Average Distant Water Purse Seine Catches, 2004-2006	46
Table 10: Reported Annual Landings (metric tons) for U.S. Purse Seine Vessels Operating in the WCPO, 1999-2008.....	57
Table 11: Purse Seine Discards by U.S. Vessels Fishing in the WCPO, 2005 and 2006.....	60
Table 12: 2005 U.S. Purse Seine Discards, (average/vessel)	61
Table 13: Percent of Catch Discarded by U.S. Purse Seine Fleet.....	61
Table 14: Non-ESA Listed Marine Mammals	71
Table 15: Catch Rates for Longliners Operating in the FSM (lbs/hook).....	81
Table 16: Typical American Samoa Tuna Longline Vessel Operations.....	81
Table 17: Projected Annual Landings for One CNMI Longline Vessel.....	81
Table 18. Essential Fish Habitat (EFH) and Habitat Areas of Particular Concern (HAPC) for species managed under the Fishery Ecosystem Plans.....	110

3.0 Introduction

The Magnuson-Stevens Fishery Conservation and Management Act (MSA) contains several provisions specifically aimed at fishery development and management in the Pacific Insular Areas which includes recognizing the cultural and economic importance of maintaining local fisheries. Section 305 (2)(E) states “Notwithstanding any other provision of this Act, the Western Pacific Council shall take into account traditional indigenous fishing practices in preparing any fishery management plan.” Section 2 (a)(10) states “Pacific Insular Areas contain unique historical, cultural, legal, political, and geographical circumstances which make fisheries resources important in sustaining their economic growth.”

Section 301 of the MSA describes the ten national standards that fishery management plans must be consistent with. National Standard 8 states “Conservation and management measures shall, consistent with the conservation requirements of this Act (including the prevention of overfishing and rebuilding of overfished stocks), take into account the importance of fishery resources to fishing communities by utilizing economic and social data that meet the requirements of paragraph (2), in order to (A) provide for the sustained participation of such communities, and (B) to the extent practicable, minimize adverse economic impacts on such communities.” The objective of the alternatives considered here is to minimize adverse impacts to the troll fleet and the communities which rely on a continued supply of locally-caught fresh fish by reducing chances of gear conflicts and catch competition among the various fishery sectors operating within the U.S. EEZ around the Mariana Archipelago.

Longline fishing has the potential to result in adverse impacts on troll fishing through localized stock depletion and catch competition, and in gear conflicts. The proposed action would provide some spatial separation between fishing by trollers and longliners around the CNMI to reduce the possibility of potential adverse interactions between the fleets.

Due to the potentially large harvests that can be made by purse seiners, coupled with their inability to target specific and mature fish, an increase in purse seine fishing in the EEZ around CNMI and Guam has the potential to negatively impact the viable continuation of the troll and longline fisheries. Purse seining may result in localized, temporary stock depletion and catch competition with local fisheries that target the same pelagic species. The proposed action would provide spatial separation between purse seiners and troll and longline fishing vessels to reduce the impacts of purse seiners on the troll and longline fishing fleets.

3.1 Background

Historically, few U.S. vessels have conducted pelagic longline fishing based out of the Mariana Archipelago. In 1992, a 50-nautical mile prohibited area for longline fishing around Guam was implemented. However no such prohibited area exists in waters around nearby CNMI and there is growing interest by both U.S. and foreign companies to begin longline fishing in the U.S. EEZ around the Mariana Archipelago, including domestic longline vessels currently based in Hawaii. In 2006, the Guam Fishermen’s Cooperative began operating a 57 ft longline vessel in the EEZ

around Guam to train local fishermen in longline fishing techniques with the objective of increasing locally-based longline operations. In 2009, NMFS issued five Western Pacific General Longline permits (two in CNMI, one in American Samoa, and two U.S. west coast longliners). These general longline permits may be used to fish in EEZ waters around CNMI where there is far less competition than in the Hawaii-based fishery. The EEZ around Guam is adjacent to that of the CNMI (Figure 1 and Figure 2) and, therefore, fishery or other activities in one area have a high potential to affect the other. As of 2010, a new longline company, based out of Saipan, with four permitted vessels, is attempting to get off the ground and expand into the frozen fish export and local fresh fish market (D. Lewis, longline fisherman, pers. comm. April 2010).

Trolling is the primary fishing method utilized in the locally-based pelagic fisheries in the Mariana Archipelago and the troll fishery is comprised of approximately 430 vessels. The pelagic trolling fishing fleet consists primarily of vessels less than 24 ft in length that generally take one-day trips within 30nm around the islands where they find abundant skipjack tuna. These vessels have a limited travel and fishery participants necessarily rely on catches from waters within their reach.

There is one Guam Fishermen's Cooperative longline vessel and four longline vessels in CNMI. Longline vessel operators want to keep fuel costs down and may, therefore, fish in waters within 30 nm of shore where the trollers fish. During the early years of longline fishing around Hawaii and American Samoa, vessels were fishing close to shore which prompted the Council and NMFS to establish prohibited areas around those archipelagos to protect sensitive resources and prevent gear conflicts.

Expansion of longline fishing around the Mariana Archipelago in nearshore waters would likely result in adverse impacts to the troll fleet and the communities which rely on a continued supply of locally-caught fresh fish, due to gear conflicts and possible localized stock depletion. The nearshore waters, within 25-30 nm from shore, include the reefs and banks where the troll fleet fishes.

The Council has also recognized the potential for purse seine vessels to expand their fishing grounds which could include EEZ waters around the Mariana Archipelago and elsewhere as the U.S. fleet has more than doubled in fleet size over the past few years although only U.S. hull vessels can fish in the U.S. EEZ. If purse seiners were to fish in U.S. EEZ waters there is potential for gear interactions and catch competition between small boats targeting skipjack and seiners. Fishery interactions between purse seiners and other sectors (including longline, troll, and artisanal) have been observed in several research studies, described in Section 8.4.4. For example, Sibert and Hampton (2003) concluded that in addition to the need for international cooperation in the conserving tuna stocks, "... Pacific Island countries can implement effective domestic management policies to promote conservation and sustainable utilization of tuna stocks within their EEZs," which highlights the Council's intention in recommending this proposed action.

Local troll vessels have a limited travel and fishing range and fishery participants necessarily rely on catches from waters within their range. Continuing to allow purse seine fishing around

the Mariana Archipelago is predicted to result in adverse impacts to the troll fleet through localized stock depletion of skipjack tuna, the target species of both sectors. Demand for skipjack tuna is high and therefore catches in the WCPO are increasing with 2007 showing the highest catch recorded in the WCPO (WCPFC 2009). The skipjack market is expected to continue to expand and with it interest in fishing for skipjack in new areas, such as waters around the Mariana Archipelago. Worldwide, the majority of skipjack tuna is harvested by purse seine vessels and therefore it is prudent to take measures to ensure the locally-based fleets in Guam and CNMI are not adversely impacted by future increased skipjack harvests.

In addition, regulating purse seine fishing within U.S. EEZ waters around the Mariana Islands is consistent with the need to limit impacts of purse seine fishing on juvenile recruitment of bigeye tuna while maintaining fishing opportunities for longline fishing on older fish, which may have had the opportunity to spawn. The recent renaissance of the U.S. purse seine fleet means that it is one of the largest in the Western Pacific and typically has fished extensively with untethered fish aggregating devices (FADs), in some years making up to 90% of sets on FADs, which captures juvenile bigeye along with the targeted skipjack.

Accordingly, the U.S. EEZ around the Mariana Archipelago, American Samoa and the Pacific Remote Island Areas (PRIA), outside of the newly designated marine national monument areas, may become increasingly attractive for the U.S. purse seine fleet to exploit. In the Mariana Archipelago, purse seine fishing has not been recorded, and under the proposed action, purse seine fishing would be prohibited within the U.S. EEZ, while providing the maximum opportunity for the fledgling longline fishery out of Saipan to develop. Moreover, skipjack is the dominant fish caught by several hundred small vessel trollers in Saipan and Guam and thus would be easily outcompeted by the larger catching power of purse seiners, which are also more mobile than the two local fleets and can still fish in large areas across the western and central Pacific Ocean including in the U.S. EEZ around the PRIA.




Figure 1: EEZ boundaries in the western Pacific.

The proposed action focuses on the waters of the U.S. EEZ around the Northern Mariana Islands and Guam.

CNMI and Guam and their associated EEZ waters (Figure 2) lie within the northern boundaries of the Western Pacific Warm Pool, which is the source of the largest proportion of the tuna catch in the Pacific Ocean. Therefore, a precautionary approach to managing tuna resources in the WCPO would include protecting this area. Prohibiting fishing activities capable of harvesting large amounts of tuna, such as purse seine fishing, could contribute to protecting the area from localized depletion while allowing small scale local fisheries to continue. With the near tripling of the size of the U.S. purse seine fleet in the past several years coupled with reductions in available areas due to establishment of marine national monuments, U.S. vessels may look to expand their fishing grounds and U.S. hulls could expand into the large area of U.S. EEZ waters around the Mariana Archipelago. In addition, U.S. purse vessels with valid fishery endorsements may fish in other areas of the WCPO including exclusive rights to fish in U.S. EEZ waters around the Pacific Remote Islands Areas (PRIA), except those parts recently designated as a marine national monument (see Section 8.4.2).




Figure 2: U.S. EEZ waters around the Mariana Archipelago

Source: NMFS, Pacific Islands Fisheries Science Center, Western Pacific Fisheries Information Network

4.0 Purpose and Need for Action

The purpose of this proposed action is to prevent or minimize adverse impacts to the troll fleet and the communities which rely on a continued supply of locally-caught fresh fish by reducing chances of gear conflicts and catch competition among fishery sectors while allowing for continued longline fisheries development. The proposed 30-nm longline closed area around the CNMI is intended to avoid gear conflicts and catch competition in the nearshore waters between longline fishing vessels and the CNMI-based small boat fishing fleet, primarily troll vessels.

The second purpose of this amendment is to prevent catch competition due to localized depletion of the skipjack tuna stock between purse seine vessels and local longline and troll fishing vessels by implementing a purse seine fishing prohibition for all U.S. Exclusive Economic Zone (EEZ) waters around the islands of the Mariana Archipelago. By prohibiting purse seine harvest in U.S. EEZ waters around the Mariana Archipelago, this action would serve to reduce impacts on bigeye tuna in the WCPO and preempt catch competition between seiners and smaller vessels in the Mariana Archipelago. The alternatives considered here are intended to comply with the MSA, while balancing the needs and concerns of Guam and CNMI's small troll and pelagic longline fishing fleet; and in effect, aid in the conservation of important stocks such as bigeye, yellowfin and skipjack tuna throughout their range in the Pacific Ocean.

5.0 Initial Actions

The Council has taken a series of management actions to avoid gear conflicts in waters close to island areas in the western Pacific, to protect species or habitats, and to facilitate the continuation and emergence of small-scale localized fishing effort in the various island areas of the Council's jurisdiction.

In 1991, Amendment 3 (56 FR 52214, October 18, 1991) to the Pelagics Fishery Management Plan (PFMP) created a protected species zone around the NWHI to protect endangered Hawaiian monk seals. This is a contiguous area extending 50 nm from named features in the NWHI and connected by 100-nm wide corridors between those areas where the 50-nm-radius areas do not intersect.

In 1992, Amendment 5 (57 FR 45989, March 4, 1992) created a domestic longline vessel prohibited area around the Main Hawaiian Islands ranging from 50 to 75 nm, and a similar 50 nm prohibited area around Guam and its offshore banks. The prohibited areas were designed primarily to prevent gear conflicts and vessel safety issues arising from interactions between longline vessels and smaller fishing boats which had arisen with the rapid growth of the Hawaii-based longline fleet early in the fishery. A seasonal reduction in the size of the closure was subsequently implemented in October 1992; between October and January, longline fishing is prohibited within 25 nm of the windward shores of all the main Hawaiian Islands except Oahu, where it is prohibited within 50 nm from the shore.

In 2002, the Council closed waters within 3-50 nm around American Samoa to pelagic fishing by any vessel greater than 50 ft in length (67 FR 4369, January 30, 2002). This action was taken to prevent potential gear conflicts and catch competition between large fishing vessels and locally-based small vessels.

Amendment 14 addressed overfishing of WCPO yellowfin and Pacific bigeye tuna and was partially approved by the Secretary of Commerce. It contained recommendations regarding both international and domestic management, including a mechanism by which the Council could participate in international negotiations regarding these stocks. Amendment 14 also contained measures to implement control dates (June 2, 2005) for Hawaii's non-longline commercial pelagic vessels (70 FR 47781) and purse seine and longline vessels (70 FR 47782), as well as requirements for federal permits and reporting for Hawaii-based non-longline commercial pelagic vessels. NMFS disapproved the Amendment's international measures as premature given ongoing international negotiations as well as the development of a memorandum of understanding by the Councils and the Secretary of Commerce, in consultation with the Secretary of State, regarding participation in U.S. delegations and other issues. The implementation of a control date is in recognition of the fact that unlimited expansion of purse seining and longline fishing is untenable with the conservation of bigeye and yellowfin tuna. NMFS also noted that Amendment 14 met the requirements of the Magnuson-Act regarding overfishing of fisheries that have been determined to be subject to overfishing due to excessive international fishing pressure.

At the 96th meeting of the Council's Scientific and Statistical Committee (SSC), held September 25-27, 2007, Council staff presented an options paper describing several alternatives to meet the objective described below. These included establishing longline prohibited areas of varying sizes (25, 50, or 100 nm) around the islands. The discussion focused on a 30 nm prohibited area that would fully encompass historic fishing grounds for the troll fleet by including popular banks (e.g., Esmeralda Bank), without creating a zone that would be onerously far offshore for development of local longline fishing. The SSC recommended consideration of a 30 nm zone.

Scoping meetings to discuss proposed exclusion areas and determine issues of concern were held in Saipan, Tinian, and Rota on September 4, 5, and 6, 2007, respectively to solicit comments on the proposed prohibited areas in Guam and CNMI. The meetings were attended primarily by fishermen. For the longline prohibited area around CNMI, the majority of participants preferred a 25 nm closure (Alternative 1B). Those in favor of this alternative claimed that it would still provide an incentive to those interested in longline fishing in the area, but the local pelagic trollers would have access to most of the banks and seamounts located within 25 nm. They felt that allowing longline fishing outside this 25 nm area would help the fishermen and the ailing economy while preventing gear conflict. Attendees stated that a larger prohibited area could be impracticable for developing a local longline fishery as the cost of fuel is now prohibitive. Others at the meeting opposed Alternative 1B and supported a bigger prohibited area such as 30 nm or 50 nm (i.e., Alternative 1C and 1D). These participants contended that a bigger zone would include all of the favorite trolling fishing areas, further reduce the potential for gear conflict, and reduce catch competition for targeted species. They also argued that unlike the small boats, longline vessels are designed to fish further out in the open waters. For the purse seine prohibited area, all in attendance at the meetings supported closing the entire EEZ to purse seine fishing around CNMI and Guam (Alternatives 2D and 3D). Meeting participants expressed that purse seine fishing catches large volumes of fish and they were concerned about the depletion of target species (i.e., skipjack tuna).

At the 139th Council meeting held October 9-12, 2007, the Council took initial action to recommend establishment of a 30-nm longline prohibited area around CNMI and a purse seine fishing prohibition for all EEZ waters around the Mariana Archipelago. At the 140th Council meeting held March 17-21, 2008, in Tumon, Guam, and Saipan, CNMI, the Council took action to recommend adopting Alternative 1C, establishment of a 30-nm longline prohibited area around CNMI; and Alternatives 2D and 3D, prohibitions on purse seine fishing in all EEZ waters around CNMI and Guam, respectively. Also at this meeting, the Council recommended prohibiting purse seine fishing within 75 nm of American Samoa to reduce potential catch competition and localized depletion in the U.S. EEZ waters around American Samoa.

6.0 Objective

As discussed in Section 4 (Purpose and Need for Action) above, the objective of the alternatives considered here are to:

1. Prevent or minimize adverse impacts to the troll fleet and the communities which rely on a continued supply of locally-caught fresh fish by reducing chances of gear conflicts and catch competition among fishery sectors while allowing for continued longline fisheries development.
2. Prevent catch competition between purse seine vessels and Guam and Saipan-based longline and troll fishing vessels that could occur due to localized depletion of skipjack and yellowfin tuna stocks by implementing a purse seine fishing prohibition for all U.S. Exclusive Economic Zone (EEZ) waters around the islands of the Mariana Archipelago.

7.0. Description of Alternatives

The following alternatives are intended to meet the purpose and need and objective described above. For ease of analysis and consideration of alternatives, the proposed actions were divided into three separate topics with a series of alternatives for each topic. Alternatives were developed by Council staff by examining the most recent fisheries data and relevant studies, consideration of public comment, and consideration of recommendations made by fishing communities and participants at public hearings including the 139th Council meeting held in the Mariana Archipelago in Saipan and Guam.

7.1 Topic 1: CNMI Longline Prohibited Area

7.1.1 Alternative 1A: No Action

Under Alternative 1A (No Action), no new regulations would be made and U.S. longline vessels would not be prohibited from fishing within U.S. EEZ waters around CNMI.

7.1.2 Alternative 1B: 25 nm Longline Prohibited Area

Under Alternative 1B, all vessels would be prohibited from pelagic longline fishing within 25 nm of CNMI.

7.1.3 Alternative 1C: 30 nm Longline Prohibited Area (Preferred)

Under Alternative 1C, all vessels would be prohibited from pelagic longline fishing within 30 nm of CNMI.

7.1.4 Alternative 1D: 50 nm Longline Prohibited Area for Large Vessels

Under Alternative 1D, vessels greater than 50 ft in length would be prohibited from pelagic longline fishing within 50 nm of CNMI.

7.1.5 Alternative 1E: 75 nm Longline Prohibited Area

Under Alternative 1E, all vessels would be prohibited from pelagic longline fishing within 75 nm of CNMI.

7.1.6 Alternative 1F: 100 nm Longline Prohibited Area

Under Alternative 1F, all vessels would be prohibited from pelagic longline fishing within 100 nm of CNMI.

7.2 Topic 2: CNMI Purse Seine Prohibited Area

7.2.1 Alternative 2A: No Action

Under Alternative 2A (No Action), no new regulations would be made and pelagic purse seine vessels would not be prohibited from fishing within U.S. EEZ waters around CNMI.

7.2.2 Alternative 2B: 50 nm Purse Seine Prohibited Area

Under Alternative 2B, all vessels would be prohibited from pelagic purse seine fishing within 50 nm of CNMI.

7.2.3 Alternative 2C: 100 nm Purse Seine Prohibited Area

Under Alternative 2C, all vessels would be prohibited from pelagic purse seine fishing within 100 nm of CNMI.

7.2.4 Alternative 2D: Entire EEZ Purse Seine Prohibited Area (Preferred)

Under Alternative 2D, all vessels would be prohibited from pelagic purse seine fishing within all U.S. EEZ waters around CNMI.

7.3 Topic 3: Guam Purse Seine Prohibited Area

7.3.1 Alternative 3A: No Action

Under Alternative 3A (No Action), no new regulations would be made and pelagic purse seine vessels would not be prohibited from fishing within U.S. EEZ waters around Guam

7.3.2 Alternative 3B: 50nm Purse Seine Prohibited Area

Under Alternative 3B, all vessels would be prohibited from pelagic purse seine fishing within 50 nm of Guam.

7.3.3 Alternative 3C: 100 nm Purse Seine Prohibited Area

Under Alternative 3C, all vessels would be prohibited from pelagic purse seine fishing within 100 nm of Guam.

7.3.4 Alternative 3D: Entire EEZ Purse Seine Prohibited Area (Preferred)

Under Alternative 3D, all vessels would be prohibited from pelagic purse seine fishing within all U.S. EEZ waters around Guam.

8.0 Description of the Affected Environment

8.1 U.S. EEZ around the Mariana Archipelago and the Western Pacific Warm Pool

The Mariana Archipelago is composed of 18 islands and extends north to south over a distance of 350 miles. Guam is the southernmost island of the Mariana Archipelago, located at 13° N latitude, 144° E longitude (Bureau of Statistics and Plans, 2006). U.S. EEZ waters around Guam are approximately 82,400 square miles whereas U.S. EEZ waters around CNMI are 3.5 times the size at 291,800 square miles.

CNMI, Guam, and their associated EEZ waters (Figure 1 and Figure 2) lie within the northern boundaries of the Western Pacific Warm Pool, which has the highest mean annual sea surface temperatures on earth ranging from 28°C to greater than 29.5°C. The largest proportion of the tuna catch in the Pacific Ocean originates from the Warm Pool, even though, paradoxically, this is a region of low primary productivity.

The Mariana Archipelago EEZ waters extend over varying degrees of latitude and oceanographic conditions which has the potential to cause variations in the catchability of target tuna species (OFP 1998). The EEZ waters around Guam have a shallower and steeper thermocline than those around CNMI indicating that yellowfin and bigeye catchability might be greater in Guam's EEZ than in the waters around the CNMI (OFP 1998). However, there are also localized effects of the many submarine features which lead to increased productivity in certain areas.

A major oceanographic and climatic phenomenon which drives variation in the Warm Pool is the El Niño–Southern Oscillation (ENSO). ENSO is linked to climatic shifts that are part of a normal and prominent weather pattern in the Pacific and Indian Oceans. ENSO, which can occur every 2–10 years, results in the reduction of normal trade winds which reduces the intensity of the westward flowing equatorial surface current (Sturman and McGowan 2003). In turn, the eastward flowing countercurrent tends to dominate circulation, bringing warm, low-salinity low-nutrient water to the eastern margins of the Pacific Ocean. As the easterly trade winds are reduced, the normal nutrient-rich upwelling system does not occur, leaving warm surface water pooled in the eastern Pacific Ocean. During strong ENSO events, the area where tunas typically school at the surface, such as skipjack and yellowfin tuna, may shift eastward by thousands of kilometers. Highly mobile fishing fleets of purse seiners and longliners are able to move to follow the fish schools, while Guam and CNMI's small-boat fleets must continue to fish in whatever conditions prevail within their limited range.

Larval and juvenile yellowfin stay in surface waters while adults are increasingly found at greater depths. Juvenile yellowfin tuna form a major component of surface landings in the Pacific and form an economically and socially important component of domestic, artisanal and subsistence fisheries in the Pacific, particularly in small island areas such as the Mariana Archipelago. In particular, although the primary target fish is skipjack tuna, small-scale troll and surface handline fisheries in the Mariana Islands generally catch juvenile yellowfin less than 100 cm in length.

Large-scale purse seine fishing operations using fish aggregation devices (FADs) in the WCPO have generated management concern due to their tendency to aggregate juvenile and commercially undersized tuna as well as bycatch species, e.g. billfish, oceanic sharks, and marine turtles (Hampton and Bailey 1993). Of particular management concern is the unintended catch of juvenile bigeye tuna by WCPO purse seine fleets as well as elevated landings of juvenile yellowfin tuna in drifting object sets². Juvenile yellowfin are also regularly taken as an incidental byproduct in skipjack pole and line fisheries when fishermen target effort around FADs. Large, mature-sized yellowfin tuna are caught by higher value sub-surface fisheries, primarily longline fleets landing sashimi grade product. Adult yellowfin tuna aggregate to

² http://www.soest.hawaii.edu/PFRP/biology/holland_itano_png.html

drifting flotsam and anchored buoys, though to a lesser degree than juvenile fish. Large yellowfin also aggregate over deep seamount and ridge features where they are targeted by some longline and handline fisheries. It appears that juvenile yellowfin tuna are particularly vulnerable to being caught by longline gear from around 55 cm and may be retained or discarded depending on the market characteristics of the fishery. Increasing harvests of juvenile yellowfin have been a concern for many years as evidenced by the IATTC's 1999 Resolution to implement catch limits which states that "fishing for juvenile yellowfin has increased considerably during 1999, and that these small fish must be protected"³. At the WCPFC's December 2006 3rd regular session, a conservation and management measure (CMM 2006-01) was adopted which says that "management plans for the use of FADs shall include strategies to limit the interaction with juvenile bigeye and yellowfin tuna". Recognizing the urgent need to reduce fishing mortality of juvenile bigeye and yellowfin tuna from fishing on FADs, in December 2008, the Commission adopted measures (CMM-2008-01) to reduce juvenile bigeye and yellowfin mortalities from fishing effort on FADs through a FAD seasonal closure (August 1 - September 30) in the convention area bounded by 20°N and 20°S; and provided guidelines for the development of FAD management plans"⁴.

8.2 Pelagic Fishing in CNMI

Foreign Vessels

Since 1980 foreign vessels have been prohibited from fishing in CNMI's EEZ waters. Prior to the MSA and the establishment of U.S EEZ boundaries in 1980⁵, foreign longline and pole-and-line fishing fleets operated in CNMI waters. A Japanese pole-and-line fishery and longliners from distant water fishing nations (DWFN) caught relatively large amounts of skipjack and yellowfin in CNMI waters. Japanese pole-and-line catch of skipjack (bonito) in the vicinity of CNMI ranged as high as 17,606 metric tons⁶ with an average closer to 5,000-6,000 mt (Table 1). The Japanese fishing activities (pole-and-line and longline fishing) within the 200 miles of the Northern Mariana Islands waters from 1973 to 1975 (between 5,700 and 7,700 mt) far exceeded local catches of tuna, marlins, etc. (Source: American Embassy - Japan). Longline catches of yellowfin, bigeye and blue marlin were relatively insignificant, with a peak level of 1,718 metric tons in 1962 (SPC 1999). In the 1990's there was some purse seine fishing effort outside of the U.S. EEZ around CNMI. To date, no purse seine fishing activity has been recorded in the U.S. EEZ waters or in the vicinity of CNMI.

Table 1: Annual catches (mt) of bigeye (BET), blue marlin (BLZ), yellowfin (YFT) and skipjack (SKJ) by foreign fleets in the vicinity of CNMI (15°-25°N, 140°-150°E) 1962-1997.
Source: OFP 1998

³ <http://www.iattc.org/PDFFiles/C-99-08%20YFT%20resolution%20Oct%2099.pdf>

⁴ <http://www.wcpfc.int/doc/cmm-2008-01/conservation-and-management-measure-bigeye-and-yellowfin-tuna-western-and-central-pa>

⁵ Based on a recent court decision EEZ waters around CNMI are defined to begin at the shoreline, and there are no territorial waters.

⁶ Note: 1 mt = 2,204.62 lb.

YEAR	LONGLINE				POLE AND LINE		PURSE SEINE		TOTAL				
	BET	BLZ	YFT	TOTAL	SKJ	YFT	SKJ	YFT	BET	BLZ	SKJ	YFT	TOTAL
1962	623	140	955	1,718					623	140		955	1,718
1963	316	43	270	629					316	43		270	629
1964	77	52	74	203					77	52		74	203
1965	129	47	80	256					129	47		80	256
1966	37	57	95	189					37	57		95	189
1967	112	38	46	196					112	38		46	196
1968	60	31	37	128					60	31		37	128
1969	69	23	52	144					69	23		52	144
1970	114	109	288	510					114	109		288	510
1971	131	78	230	439					131	78		230	439
1972	78	36	128	242	6,676	77			78	36	6,676	204	6,995
1973	51	15	25	91	7,495	347			51	15	7,495	372	7,932
1974	29	41	30	100	4,893	330			29	41	4,893	360	5,322
1975	45	28	65	137	6,039	284			45	28	6,039	348	6,460
1976	66	32	114	212	3,123	311			66	32	3,123	425	3,646
1977	121	24	173	318	3,159	455			121	24	3,159	627	3,931
1978	75	40	421	536	4,418	429			75	40	4,418	849	5,383
1979	40	38	78	156	6,330	497			40	38	6,330	575	6,982
1980	75	40	304	418	5,283	230			75	40	5,283	533	5,931
1981	31	22	53	105	7,026	374			31	22	7,026	427	7,505
1982	23	19	36	78	7,551	193			23	19	7,551	230	7,822
1983	49	42	124	215	6,941	328			49	42	6,941	451	7,483
1984	48	13	79	140	17,606	537			48	13	17,606	615	18,283
1985	24	4	46	75	4,979	345			24	4	4,979	392	5,398
1986	85	35	138	258	3,813	314			85	35	3,813	452	4,305
1987	115	40	107	262	6,184	85			115	40	6,184	192	6,531
1988	34	23	81	139	4,834	286			34	23	4,834	367	5,259
1989	11	6	25	42	6,194	183			11	6	6,194	207	6,419
1990	39	8	33	80	6,441	117			39	8	6,441	150	6,638
1991	62	19	130	211	3,569	202			62	19	3,569	332	3,982
1992	95	24	126	245	3,646	88	4	3	95	24	3,650	216	3,985
1993	118	29	125	272	5,277	124	2	2	118	29	5,279	250	5,675
1994	68	31	129	229	3,281	125	7	14	68	31	3,288	268	3,656
1995	24	20	203	247			15	5	24	20	15	208	267
1996	19	20	166	204			434	21	19	20	434	187	659
1997													

Domestic fishing in the CNMI

For a complete description of troll, purse seine and longline gear please refer to section 4.1.1 of the Fishery Ecosystem Plan for Pelagic Fisheries of the Western Pacific (WPFMC 2009a).

With the exception of the purse seine support base on Tinian discussed above, CNMI has never had a large infrastructure dedicated to commercial fishing. The majority of boats in the local fishing fleet are small, outboard engine-powered vessels. The harvest of pelagic species by CNMI-based vessels has always been small, around 100 metric tons annually, caught with trolling gear. Both supply and demand conditions direct the majority of domestic commercial fishing effort in CNMI toward reef fish and bottomfish. There is less seasonality in these fisheries, and they require shorter offshore trips; moreover, their market value is often much higher than that of the commonly caught pelagic fish.

CNMI's pelagic fishery occurs primarily from waters off the island of Farallon de Medinilla south to the Island of Rota. The pelagic fishing fleet consists primarily of trolling vessels less than 24 ft in length which generally take one-day trips within 30 nm around the islands where

they find abundant skipjack tuna. These vessels have a limited travel and fishing range and fishery participants necessarily rely on catches from waters within their reach.

Currently there is a new longline fishing company located on Saipan which, as of 2010, has four longliners fishing waters around the Mariana Archipelago, beyond 30 nm from shore but within EEZ waters. During the development of this document, there are no catch data available on this new fishery. Interest in longline fishing in CNMI has been variable with the issuance of eight, four, and five Western Pacific General Longline permits from 2007 through 2009, respectively.

The pelagic fishery is monitored using data in the Commercial Purchase Data Base which currently documents landings on Saipan where the majority of the CNMI's population and fishery participants live. Staff from the Department of Lands and Natural Resources, Division of Fish and Wildlife (DFW) routinely distribute and collect invoice books from 30 participating local fish purchasers on Saipan that record all fish purchases by species categories. The establishment of data collection systems for the islands of Tinian and Rota are in process. It is believed that the commercial purchase database landings include around 90 percent of all commercial landings on Saipan. There is also an unquantified subsistence fishery on Saipan where income is made by selling a small portion of catches door-to-door to cover fishing expenses.

The primary target and most marketable species for the pelagic fleet is skipjack tuna. In 2008, skipjack tuna continued to dominate the pelagic landings, comprising around 80 percent of commercial pelagic landings and revenues totaling about \$317,000 (Table 2). Schools of skipjack tuna have historically been common in nearshore waters, providing an opportunity for trollers to catch numerous fish with a minimum of travel time and fuel costs. Yellowfin tuna and mahimahi are also easily marketable species but are seasonal. Peak mahimahi catches are usually from February through April while the yellowfin season usually runs from April through September.

In 2008, skipjack tuna landings decreased when compared to landings in 2007, while yellowfin tuna and mahimahi ranked second and third in total landings. However mahimahi and yellowfin landings decreased 58 and 53 percent respectively from the previous year. The total weight of commercial pelagic landings decreased 37 percent in 2008 and fell below the 26 year mean of 242,276 lb (Table 3, Figure 3).

Table 2: CNMI 2008 Commercial Pelagic Trolling Landings, Revenues and Price⁷

Species	Landing (lb)	Value (\$)	Avg Price (\$/lb)
Skipjack Tuna	157,708	244,652	1.55
Yellowfin Tuna	16,344	32,149	1.97
Saba (kawakawa)	2,155	3,182	1.48
Tunas (misc.)	45	68	1.50

⁷ The year 2008 is the most recent for which we have comprehensive information about all fisheries considered in this amendment. Therefore, we use 2008 as the baseline year for analyzing the potential effects of this action. Although more recent troll, longline, and purse seine fishery information is available, inclusion of more recent information would not change the Council's analyses or management objectives

Species	Landing (lb)	Value (\$)	Avg Price (\$/lb)
Tuna PMUS	176,252	280,050	1.59
Mahimahi	11,169	20,428	1.83
Wahoo	1,388	2,881	2.08
Blue Marlin	1,098	1,331	1.21
Sickle Pomfret (w/woman)	481	1,060	2.21
Non-tuna PMUS	14,136	25,700	1.82
Dogtooth Tuna	3,457	5,399	1.56
Rainbow Runner	3,095	6,056	1.96
Barracuda	58	88	1.50
Troll Fish (misc.)	15	38	2.50
Non-PMUS Pelagics	6,625	11,580	1.75
Total Pelagics	197,013	317,330	1.61

Source: WPRFMC 2010

Table 3: CNMI Total Commercial Trolling Landings (lb) 1983-2008

Year	Catch lbs						Total
	Mahimahi	Wahoo	Blue Marlin	Skipjack	Yellowfin	Other species	
1983	13,939	8,760	3,787	183,411	21,281	14807	245,985
1984	7,614	14,087	1,544	290,843	19,580	7468	341,136
1985	12,955	18,251	1,860	177,344	12,466	11302	234,178
1986	17,796	9,062	2,654	254,362	16,917	6668	307,459
1987	9,502	13,404	2,460	161,504	10,454	7744	205,068
1988	30,799	11,697	1,309	266,497	15,375	8846	334,523
1989	7,320	1,571	5,704	257,703	10,109	4377	286,784
1990	10,439	3,462	2,034	147,962	10,468	6085	180,450
1991	33,756	1,521	1,568	115,802	13,042	22872	188,561
1992	26,257	17,172	6,603	82,280	25,687	41229	199,228
1993	37,545	2,779	3,687	97,268	14,898	25151	181,328
1994	15,063	3,863	2,635	92,212	13,445	20111	147,329
1995	23,321	5,722	6,619	131,377	20,918	12223	200,180
1996	35,655	10,783	8,593	165,037	38,043	23166	281,277
1997	31,277	7,580	7,068	133,446	21,352	18150	218,873
1998	25,375	6,299	4,201	167,114	14,570	22704	240,263
1999	12,882	8,063	3,541	106,297	24,419	21829	177,031
2000	7,324	4,097	3,608	140,389	17,673	14204	187,295

Year	Catch lbs						Total
	Mahimahi	Wahoo	Blue Marlin	Skipjack	Yellowfin	Other species	
2001	14,229	4,550	1,924	133,769	14,543	10166	179,181
2002	18,042	8,212	1,261	179,966	30,017	19484	256,982
2003	7,357	7,950	1,130	171,574	26,042	14363	228,416
2004	35,808	6,936	2,001	148,328	27,548	18386	239,007
2005	26,891	3,349	1,595	260,614	52,014	27912	372,375
2006	17,360	3,267	1,402	273,715	43,220	17742	356,706
2007	26,410	2,504	76	238,972	34,894	9698	312,554
2008	11,169	1,388	1,098	157,708	16,344	9306	197,013
Average	19,849	7,167	3,075	174,442	21,743	16,000	242,276

Source: WPRFMC 2010




Figure 3: CNMI Annual Commercial Pelagic Trolling Landings 1983-2008

Source WPRFMC (2010)

In 2005, skipjack landings showed a significant increase of 75 percent over the previous year, and in 2006 landings again increased slightly. The 2007 landings decreased slightly by 6 percent but this decline continued in 2008 decreasing by 34 percent compared with 2007, falling below the 26 year average (Figure 4).

Although more highly prized than skipjack, yellowfin tuna are not as common in CNMI's fisheries and their average size tends to be smaller when compared with yellowfin tuna from other geographic areas. Total landings for yellowfin tuna increased in 2002 by 51 percent over the previous year (Figure 4). This increase was partly due to landings from the Northern Islands bottom-fishing fleet and a longline fishing experiment by one fishing company. The longline operation did not continue longlining in 2003 and this led to a 13 percent decrease in yellowfin landings in 2003. Troll boat landings then increased in 2004 and 2005. Since then, 2008 yellowfin landings have declined by 57 percent.




Figure 4: CNMI Annual Commercial Landings of Skipjack and Yellowfin Tuna by Trolling Vessels 1983-2008

Source: WPRFMC 2010

The number of vessels making commercial pelagic landings in the CNMI was relatively constant from 1988-1991, but a record high number was recorded for 1992. Part of this increase was attributed to an influx of new fishing boats, but it was also discovered that some fishermen were using several different boats, thus artificially inflating the total number of boats concurrently making pelagic landings. Many of the new boats were believed to have left the fishery during 1993. It has been suggested that the increase in active fishing vessels from 1994 to 1997 might be due to the re-entry of boats from the 1992 fleet.

In 2006, the number of active commercial fishing vessels landing pelagic species decreased by 16 percent as compared to 2005 (Figure 5). This decrease was partly due to an increase in fuel prices, as well as a decline in the price received for skipjack tuna. It is also believed that the decline shown since 2001 is partly due to vendors entering the landings of multiple fishing boats on a single receipt and at times combining monthly total landings onto a single receipt. Fishing activity has also been affected by bad weather which plagued the Mariana Archipelago throughout 2003 and early 2004. The continued increase in fuel price also has affected many fishing boats in the CNMI.




Figure 5: CNMI Active Commercial Pelagic Trolling Vessels from 1983-2008.

Source: WPRFMC 2010

Although the number of vessels (Figure 5) and trips (Figure 6) has decreased since 2000, landings have fluctuated, especially for skipjack tuna (Figure 3 and Figure 4). The reason for this is not fully understood. Creel survey catch rates show catch rates oscillating between 50 and 100 lb/trip both before and after 1991 whereas, the Commercial Purchase data indicate sustained high catch rates before, and low catch rates after 1991. Skipjack catch rates increased between 2004 and 2007 to 114 lb/trip but declined to 96 lb/trip in 2008 (Figure 7). Previous discussions have suggested that non-tuna PMUS may be increasing in value and a slight shift in target troll fish may be occurring. In 2003, yellowfin catch rates remained relatively stable at 12 lb/trip despite bad weather (e.g. typhoons) that plagued the Mariana Islands nearly the entire 2003. In 2004, yellowfin catch rates decreased to eight lb/trip but in 2005 and 2006 they again increased. In

2007, yellowfin catch rates increased slightly by 6 percent to 17 lb/trip but declined in 2008 by 41 percent to 10 lb/trip (Figure 7). The troll catch rate of skipjack has continued a steady decline from 2006 through 2008, based on creel survey data (WPRFMC 2010).




Figure 6: Number of Pelagic Fishing Trips (Trolling) in CNMI, 1983-2008

Source: WPRFMC 2010




Figure 7: Trolling Catch Rates for Skipjack and Yellowfin Tunas in CNMI

Source: WPRFMC 2010

8.3 Pelagic Fishing in Guam

For a complete description of troll, purse seine and longline gear please refer to section 4.1.1 of the Fishery Ecosystem Plan for Pelagic Fisheries of the Western Pacific (WPFMC 2009a).

Aside from the pelagic troll fishery discussed below, there is currently one longline vessel and two purse seine vessels based out of Guam. Pelagic fishing vessels based on Guam are classified into two general groups:

1) distant-water purse seiners and longliners (foreign and domestic) that fish outside Guam's economic exclusive zone (EEZ) and transship through the island and;

2) small, primarily recreational, trolling boats that are either towed to boat launch sites or marina-berthed charter boats and fish only within local waters, either within Guam's EEZ or on some occasions in the adjacent EEZ of the Northern Mariana Islands. Most fishermen sell a portion of their catch at one time or another and it is difficult to make a distinction between recreational, subsistence, and commercial fishers. There are currently 15 civilian charter vessels on Guam and one charter operation run by the U.S. military from Sumay Cove (John Calvo, personal communication.) A summary of the catches by the Guam charter fleet is given in WPRFMC (2010). A feature of the Guam charter industry is that catches are often served as sashimi to the patrons, most of whom are Japanese.

Landings consist primarily of five major species: mahimahi (*Coryphaena hippurus*), wahoo (*Acanthocybium solandri*), bonita or skipjack tuna (*Katsuwonus pelamis*), yellowfin tuna (*Thunnus albacares*), and Pacific blue marlin (*Makaira mazara*). Other minor species caught include rainbow runner (*Elagatis bipinnulatus*), kawakawa (*Euthynnus affinis*), dogtooth tuna (*Gymnosarda unicolor*), double-lined mackerel (*Grammatorcynus bilineatus*), and oilfish (*Ruvettus pretiosus*).

High value is placed on sharing one's fish catch with relatives and friends. The social obligation to share one's fish catch extends to part-time and full-time commercial fishermen (Amesbury and Hunter-Anderson, 1989). In a study conducted by Rubinstein (2001), nearly all fishermen (96 percent) reported that they share fish regularly, giving fish to family (36 percent), friends (13 percent) or both (47 percent). A majority (53 percent) said they did not give fish to people other than family and close friends; of those who did occasionally, the main recipients were church fiestas (32 percent) and other church events or organizations (20 percent). A 2005 survey of Guam households found that out of the fish consumed by households, a little more than half (51 percent) was purchased at a store or restaurant and 9 percent was purchased at a flea market or from a roadside stand. Nearly one-quarter (24 percent) of the fish consumed was caught by the respondent or an immediate family member, and an additional 14 percent was caught by a friend or extended family member (Beukering et al., 2007 in Allen and Bartrum 2008).

Foreign Vessels

Since 1980, foreign vessels are prohibited from fishing in Guam's EEZ waters, however, prior to this the Japanese pole-and-line fishery and longliners from DWFN caught relatively large amounts of skipjack and yellowfin. Since 1980 there has been some purse seine fishing effort outside of the U.S. EEZ around Guam. Table 4 gives the annual catches made by foreign vessels in the U.S. EEZ around Guam before 1980 and outside the EEZ after 1980.

Table 4: Annual catches (metric tons) of bigeye, blue marlin, yellowfin and skipjack by foreign fleets in the vicinity of Guam (10°-15°N, 140°-150°E), 1962-1997.

Source: OFP 1998

YEAR	LONGLINE				POLE AND LINE		PURSE SEINE		TOTAL				
	BET	BLZ	YFT	TOTAL	SKJ	YFT	SKJ	YFT	BET	BLZ	SKJ	YFT	TOTAL
1962	204	166	544	914					204	166		544	914
1963	103	39	143	284					103	39		143	284
1964	165	122	303	589					165	122		303	589
1965	118	119	367	604					118	119		367	604
1966	74	88	295	457					74	88		295	457
1967	113	58	115	286					113	58		115	286
1968	113	65	232	410					113	65		232	410
1969	106	27	136	268					106	27		136	268
1970	77	53	198	329					77	53		198	329
1971	41	33	60	134					41	33		60	134
1972	42	19	51	111	649	4			42	19	649	55	764
1973	133	77	207	417	764	9			133	77	764	216	1,189
1974	431	206	760	1,396	199	1			431	206	199	761	1,597
1975	467	263	1,085	1,815	1,752	1			467	263	1,752	1,086	3,568
1976	474	152	1,136	1,762	199	3			474	152	199	1,139	1,964
1977	1,503	294	3,542	5,339	3,861	3			1,503	294	3,861	3,546	9,204
1978	585	244	4,196	5,026	1,753	41			586	244	1,753	4,237	6,820
1979	675	157	2,006	2,838	5,209	11			675	157	5,209	2,016	8,058
1980	168	63	527	757	1,289	24	4	0	168	63	1,293	551	2,075
1981	141	59	341	541	606	4	0	0	141	59	606	345	1,151
1982	105	53	136	293	914	4	0	0	105	53	914	140	1,211
1983	231	34	944	1,208	2,035	18	9	9	231	34	2,044	970	3,279
1984	346	118	517	981	202	15	54	54	346	118	256	586	1,307
1985	192	51	289	531	280	13	13	2	192	51	293	304	839
1986	215	56	408	678	3,740	39	164	103	215	56	3,904	550	4,725
1987	601	150	1,055	1,805	9,340	24	344	133	601	150	9,684	1,212	11,647
1988	368	94	684	1,146	840	18	506	183	368	94	1,346	885	2,693
1989	186	42	372	600	202	5	20	20	186	42	222	397	848
1990	286	21	185	492	3,257	14	0	0	286	21	3,257	200	3,764
1991	642	52	612	1,306	1,139	11	62	28	642	52	1,201	651	2,545
1992	500	83	745	1,328	282	0	10	5	500	83	292	750	1,625
1993	832	130	850	1,813	3,225	18	209	139	832	130	3,425	1,008	5,395
1994	332	29	381	743	2,295	15	127	13	332	29	2,422	410	3,193
1995	1,139	104	1,001	2,243	0	0	1,397	271	1,139	104	1,397	1,272	3,911
1996	505	75	555	1,134			764	81	505	75	764	636	1,980
1997	123	5	52	181	0	0	13	13	123	5	13	65	206

Domestic pelagic fisheries in Guam

There are three sources of locally-caught fish in Guam's commercial market: (1) full-time commercial fishermen; (2) part-time commercial fishermen; and (3) subsistence or recreational "expense" fishermen who frequently sell portions of their catch to help defray costs

Licenses are not required to sell fish in Guam, nor are there any reporting requirements for those selling fish.

Prior to establishment of the Guam Fishermen's Cooperative Association (GFCA) in 1979 there was no central place to sell fish, and fishermen had to develop their own markets and peddle fish after each trip. In 1982, the Western Pacific Fisheries Information Network (WPacFIN) began working with the GFCA to improve their invoicing system and to obtain data on all fish purchases. As time progressed, other fish markets began to operate, and DAWR and WPacFIN staff worked with GFCA to obtain data through the voluntary receipt book program. Although a proposed law has been introduced several times that would require reporting by dealers and possibly commercial fishermen, it has never made it through the legislative process, and the commercial landings data collection system remains voluntary.

The composition of Guam's fishery landings by fishing sector has troll fishing for pelagics forming almost 80 percent of the landings (Figure 8), with just over 40 percent of total troll landings coming from non-commercial troll fishing. The remainder of the landings includes charter fishing (7%) and bottomfish fishing (19%). Guam's fishing activity can be somewhat constrained by seasonality due to weather. In general, lower fishing activity is seen during the November to March period when sea conditions are at their worst.

Estimated annual total pelagic landings and tuna landings have varied widely, ranging between 301,504 and 935,809 lb in the 27-year time series for the total catch (Table 5, Figures 10-11). The 2008 total pelagic landings were approximately 551,504 lb, a decrease of 1.5 percent compared with 2007, and with an overall average price per pound of \$1.81, and value just under \$1.0 million (Table 6). Aggregate landings of tuna PMUS increased and non-tuna PMUS decreased. There are general wide year-to-year fluctuations in the estimated landings of the five major pelagic species. In 2008, 95 percent of tuna were caught by non-charter boats. Total tuna and non-charter tuna landings increased by 52 percent and 58 percent respectively while charter tuna landings decreased by 18 percent from 2007 totals. Skipjack tuna catch has fluctuated over the reporting period (since 1982), peaking in 2001. Total skipjack tuna landings and non-charter landings increased in 2008 by 89 percent and 99 percent respectively (Figure 11), while charter landings decreased by 11 percent.

The 2008 mahimahi catch decreased more than 57 percent from 2007, wahoo catch totals increased 123 percent, and Pacific blue marlin catch decreased 49 percent. The inflation-adjusted price of tuna and other non-tuna PMUS has shown a dramatic decline since data on the pelagic fishery were first collected in 1980. In 2008, the adjusted price for all PMUS increased 59 percent, 59 percent for tuna PMUS, and 49 percent for non-tuna PMUS. It should be recalled that these values are based on partial year data, and may not accurately reflect the market on Guam. All three prices are well below their 29 year averages. Locally caught pelagic fish continues to have to compete with cheaper imported fish.

Despite decreasing revenues with decreased commercial landings over the past decade, pelagic fishing continues, as a majority of trollers do not rely on the catch or selling of fish as their primary source of income. Several factors in recent years have negatively affected trolling activity and may affect fishing activity in the future. The price of fuel has increased significantly; making it more costly to fish and also more attractive to sell fish to recoup costs.

More than two-thirds of the FADS have been lost and not replaced, and difficulties with procurement have prevented timely redeployment of these aids to fishing. Trolling activity occurs regularly at FADs, and is reported to have occurred with regularity at offshore banks. However, fishermen also reported more interaction with sharks at offshore banks.

Another factor that will have a direct affect on localized fishery resources in Guam is the likely increase in fishing pressure (e.g., trolling) associated with the Guam military buildup, as well as an expected increase in market demand to supply both local and imported fish products to incoming workers and residents. While the levels of additional trolling and demand of addition fish product can't be quantified at this time, future analyses and data monitoring programs will be increasingly important in understanding the changing dynamics associated therein. No additional longline or purse seine vessels are expected as a direct result of the military buildup.

The number of boats involved in Guam's pelagic trolling or open ocean troll fishery gradually increased from 193 vessels in 1983 to a peak of 469 in 1998 (Table 7). This number decreased until 2001, but has generally been stable since that year. There were 385 boats involved in Guam's pelagic troll fishery in 2008, an increase of 4 percent from 2007 of 370 boats (Table 7 and Figure 13). A majority of the fishing boats are less than 10 meters (33 feet) in length and are usually owner-operated by fishermen who earn a living outside of fishing. Most fishermen sell a portion of their catch at one time or another. And a small, but significant, segment of the pelagic group is made up of marina-berthed charter boats that are operated primarily by full-time captains and crews.

Non-charter and charter troll trips generally increased for the first 15 years of the 24-year time series (since 1982) shown in Figure 14. The number of troll trips began to decline in 1999, due to a number of factors including a continuing economic recession on the island, a decline in Asian visitors for charter boats, and an increase in cost to maintain, repair, and fuel boats for the average fishermen compared with fish caught for sale to make up for expenses. Almost three times as many non-charter trips were taken (5,057) than charter trips (1,891) in 2008. In 2008, the total number of troll trips increased by 9 percent, while the number of charter trips increased by 0.8 percent and non-charter trips increased by 12 percent.

In 2008, trolling catch rates (pounds per hour fished) declined compared to 2007 (Figure 15). Catch rates for skipjack tuna by the Guam troll fleet have fluctuated widely over the past 25 years (Figure 15). The wide fluctuations are probably due to the high variability in the year-to-year abundance and availability of the stocks, although skipjack tuna is caught year round. However, it is not possible to allocate species-specific effort, since effort used to target other species can result in an artificially high or low catch rate for a given species. In 2008, total overall, non-charter, and charter trolling catch rate decreased 15 percent, 12 percent, and 39 percent, respectively. Although total skipjack CPUE has increased, the overall decreases are primarily a reflection of the low CPUEs for yellowfin tuna, mahi mahi and blue marlin (Figures 17 through 20). Charter catch rates have generally been lower than catch rates of non-charter boats, probably due to their shorter fishing time, and non-charter boats beginning earlier in the morning and ending as late as early evening.




Figure 8: Guam fishery landings by origin (2001-2006)
Source: WPacFin data

Table 5. Estimated Total Guam Pelagic Landings (lb) 1982-2008

Year	Catch lbs						Total
	Mahimahi	Wahoo	Blue Marlin	Skipjack	Yellowfin	Other species	
1982	112,202	55,909	21,787	126,652	112,654	17792	446,996
1983	156,340	86,530	30,402	97,802	65,996	13753	450,823
1984	26,080	53,847	49,711	218,556	68,048	8614	424,856
1985	72,699	130,304	54,319	107,815	93,018	18999	477,154
1986	102,921	69,583	57,105	77,735	55,611	18395	381,350
1987	79,601	86,203	49,508	61,767	41,475	19896	338,450
1988	332,874	96,996	60,863	210,185	84,102	27941	812,961
1989	95,975	127,096	86,163	128,015	40,355	28193	505,797
1990	140,293	85,280	94,796	149,312	72,314	17370	559,365
1991	415,007	59,132	96,962	122,378	44,068	15171	752,718
1992	85,017	79,894	82,965	123,903	134,653	16927	523,359
1993	234,979	62,550	57,992	109,582	50,350	32842	548,295
1994	138,014	50,457	76,633	188,784	71,221	20808	545,917
1995	327,394	77,369	76,703	178,635	93,424	27864	781,389
1996	327,604	146,926	64,527	238,409	107,023	51320	935,809
1997	265,157	65,034	90,777	219,177	90,167	29620	759,932
1998	264,695	158,230	43,872	201,666	137,392	35957	841,812
1999	161,935	76,338	80,537	124,452	128,026	61066	632,354
2000	85,585	70,484	86,565	267,541	76,606	27929	614,710

Year	Catch lbs						Total
	Mahimahi	Wahoo	Blue Marlin	Skipjack	Yellowfin	Other species	
2001	183,278	119,765	33,302	331,768	57,929	28986	755,028
2002	173,130	72,643	53,761	176,356	45,089	13899	534,878
2003	84,739	64,266	68,204	185,575	71,626	40410	514,820
2004	195,340	120,288	38,845	168,232	104,845	65667	693,217
2005	105,715	43,906	9,270	99,391	24,884	18338	301,504
2006	162,512	105,878	29,222	146,658	28,049	38289	510,608
2007	258,260	44,354	18,994	156,651	47,833	33799	559,891
2008	111,811	98,345	9,704	295,250	19,888	15083	550,081
Average	174,043	85,467	56,426	167,120	72,839	27,590	583,484

Source WPRFMC 2010

Table 6: Guam 2008 Landings and Prices

Fish	Landing (lb)	Value (\$)	Avg Price (\$/lb)
Skipjack Tuna	296,121	467,871	1.58
Yellowfin Tuna	19,899	41,588	2.09
Tunas (misc.)	4,459	89,18	2.00
Tuna PMUS	320,479	551,223	1.72
Mahimahi	111,746	208,965	1.87
Wahoo	98,865	197,730	2.00
Blue Marlin	9,707	13,783	1.42
Sailfish	283	464	1.64
Oceanic sharks	508		n/a
Pomphrets	2,672		n/a
Non-tuna PMUS	223,781	396,092	1.77
Dogtooth Tuna	2,622	4,011	1.53
Rainbow Runner	2,455	5,351	2.18
Barracuda	2,167	4,290	1.98
Non-PMUS Pelagics	7,244	14,053	1.94
Total Pelagics	551,504	998,222	1.81

Source: WPRFMC 2010




Figure 9: Guam Annual Estimated Total Landings: All Pelagics, Tuna PMUS, and Non-Tuna PMUS 1980-2008
 Source: WPRFMC 2009




Figure 10: Guam Annual Estimated Total Tuna PMUS Landings 1982-2008
 Source: WPRFMC 2010




Figure 11: Guam Annual Estimated Total Skipjack Landings 1982-2008

Source: WPRFMC 2010

Table 7: Estimated Number of Guam Trolling Vessels

Year	Estimated number of boats	Upper 95% CL	Lower 95% CL
1982	199	280	165
1983	193	242	168
1984	219	267	193
1985	276	323	249
1986	246	284	226
1987	219	244	201
1988	320	353	297
1989	329	374	303
1990	352	467	299
1991	349	422	309
1992	332	405	294
1993	346	401	316
1994	369	439	329
1995	427	476	393
1996	466	572	415
1997	449	572	393

Year	Estimated number of boats	Upper 95% CL	Lower 95% CL
1998	469	537	430
1999	449	510	415
2000	416	470	385
2001	375	429	345
2002	375	464	330
2003	371	492	312
2004	401	568	326
2005	358	498	293
2006	386	527	321
2007	370	485	315
2008	385	523	322

Source: WPRFMC 2009




Figure 12: Guam Estimated Number of Trolling Boats 1982-2008

Source: WPRFMC 2009




Figure 13: Guam Estimated Number of Troll Trips
 Source: WPRFMC 2009




Figure 14: Guam Trolling CPUE (lb/hour) 1982-2008
 Source: WPRFMC 2010




Figure 15: Skipjack CPUE (lb/hour): All, Non-charter, and Charter Vessels 1982-2008
 Source: WPRFM 2010




Figure 16 Yellowfin CPUE (lb/hour): All, Non-charter, and Charter Vessels 1982-2008
 Source: WPRFMC 2010




Figure 17: Wahoo CPUE (lb/hour): All, Non-charter, and Charter Vessels 1982-2008
 Source: WPRFMC 2010




Figure 18: Blue Marlin CPUE (lb/hour): All, Non-charter, and Charter Vessels 1982-2008
 Source: WPRFMC 2009




Figure 19: Mahimahi CPUE (lb/hour): All, Non-charter, and Charter Vessels 1982-2008
 Source: WPRFMC 2009

Guam domestic longline and purse seine fishing operations

Guam has one longliner and no domestic purse seine vessels in current operation. Catches from the single longline vessel cannot be reported due to confidentiality requirements.

8.4 WCPO Purse Seine Tuna Fishery

Purse seine fishing for tuna involves setting a net vertically in the water, with floats attached to the upper edge and chains for weight on the lower edge. A series of rings is attached to the lower edge of the net, and a pursing cable passes through the rings, enabling a winch on board the vessel to draw the net closed on the bottom. Purse seine nets can be up to 1,600 meters (m) or more in length and 220 m in depth. When the net is deployed from the purse seine vessel, a large skiff carrying the end of the net is then released from the stern of the fishing vessel. While the skiff holds one end of the net, the purse seine vessel circles and encloses the school of tuna, keeping it in visual contact if on the surface, or by using sonar if below the surface, and then retrieves most of the net onto the vessel. The fish are confined in the “sack” portion of the net, which consists of finer mesh webbing that prohibits their escape. The catch is removed from the sack onto the vessel with large “scoops” holding one mt or more, and then is placed in brine tanks for freezing and later storage (NMFS 2006). Purse seiners are one of the most complex classes of fishing vessels in terms of both technology and machinery.

Fishing by U.S. vessels in the WCPO purse seine tuna fishery currently occurs outside of the U.S. EEZ around Guam and CNMI. The purse seine tuna fishery is monitored by and operates under regulations promulgated by NMFS in accordance with the WCPFC.

The estimated delivered value of the purse seine tuna catch in the WCPFC area for 2008 was \$3,124 million (\$3.1 billion) which exceeded the previous year's record level of \$2,393 million (\$2.4 billion) (Williams and Terawasi 2008). In NOAA's 2009 annual report to the WCPFC, the domestic purse seine fishery was the largest U.S. fishery in 2008, accounting for 89 percent of the total U.S. catch of skipjack, yellowfin, bigeye, and albacore tuna in the WCPO. Total U.S. purse seine landings increased to 157,849 metric tons (mt) (or 78 percent over 2007 landings) in 2008. Yellowfin tuna catches in the fishery increased from 10,541 mt in 2007 to 23,801 mt in 2008, and skipjack tuna catches increased from 75,210 mt in 2007 to 127,307 mt in 2008.

The fishery operated mainly in areas between 5° N and 10° S latitude and 150° E and 180° longitude in 2008; below the U.S. EEZ around the Mariana Islands which resides between 11° N and 24° N (Figure 1). Before 1995, the fleet in the WCPO fished mainly on free-swimming schools of tunas. Though highly variable during the last 5 years, the fleet has been fishing equally on free-swimming schools and schools associated with floating objects, including logs and fish aggregating devices (FADs). In 2008 there were about 225+ purse seine vessels in the region. Thirty two of those vessels were U.S. permitted purse seine vessels.

The majority of the catch is taken by the distant water fishing nations (DWFN) fleets of Japan, Korea, Chinese Taipei, and the U.S. comprising of 130 vessels in 2008, down from 147 in 1995. In recent years, increasing numbers of Pacific Island domestic or joint venture vessels (i.e., Australia, New Zealand) have joined the fishery as well as some new distant water entrants (e.g., China). There has been a steady increase in the number of vessels from Pacific Island fleets, which totaled 59 vessels in 2008 (Williams and Terawasi 2008). Skipjack makes up 70-85 percent of the total WCPO purse seine fishery landings with yellowfin accounting for 15-30 percent (Williams and Reid 2007; Williams and Terawasi 2008: Figures 20-22




Figure 20: WCPO Purse Seine Vessels 1972-2008

Source: Williams and Terawasi. 2008




Figure 21: Catches of tuna in the WCPFC Statistical Area, by species
Source: WCPFC 2009




Figure 22: Catches of tuna in the WCPFC Statistical Area, by gear type
Source: WCPFC 2009

The numbers and landings of active distant water fishing nations' purse seine vessels in the WCPO during 2007 are listed in descending order of catch in Table 8.

Table 8: Numbers and Landings of Active DWFN Purse Seine Vessels in the WCPO, 2007

Country	vessels	BET Catch (mt)	YFT Catch (mt)	SKJ Catch (mt)	TOTAL
Korea	28	1,775	41,469	214,933	258,177
Japan distant water	35	4,883	24,390	215,310	252,692
Chinese Taipei	34	2,386	21,147	209,002	232,535
PNG	41	4,464	40,216	174,957	220,054
Philippines domestic purse seine	(164)*	3,418	39,308	128,178	170,904
USA	13	7,625	3,938	60,641	72,204
Vanuatu	10	391	7,030	59,589	67,010
Marshall Islands	5	2,118	3,370	53,916	59,485
China	10	768	5,428	48,745	54,941
Philippines distant-water	12	864	11,792	21,562	34,413
New Zealand	10	431	1,509	28,622	30,562
Philippines domestic ringnet	(164)*	713	6,652	16,629	23,994
Spain	3	3,040	4,019	12,688	19,747
Solomon Islands	5	817	6,326	10,164	17,307
FSM	4	196	1448	11,853	13,497
Kiribati	1	103	1,169	4,178	5,450
Japan coastal**	18	2	87	1,024	1,113
Totals		33,994	219,298	1,271,991	1,525,283

Source: Lawson (WCPFC) 2009

* This number is divided between domestic purse seine and ringnet gear vessels

**data from 2004

Purse seine fishing targeting skipjack is currently and has historically occurred near EEZ waters around the Mariana Archipelago, although no catches have been reported from within the EEZ waters. Japan is the primary nation purse seining in closest proximity to the Mariana Islands, both to the north and the south (Figure 24 and 25). This document utilizes Japan's purse seine catch data (CPUE and species composition) as a proxy for being most representative of what purse seine catches from EEZ waters around the Mariana Islands would be like, if purse seine fishing were to occur there (Table 9). Based on the WCPFC 2006 Tuna Fishery Yearbook, there were 35 active Japanese distant water purse seine vessels in 2002 through present data (WCPFC

2007). The three-year averages (2004 – 2006) are shown in Table 9 below. The average number of days fished by the Japanese DWFN vessels over the three years was 235 days per vessel.

The Federated States of Micronesia (FSM) has a relatively large-scale fishing operation with annual purse seine catches exceeding 200,000 mt in some years; however, most of this catch is taken by foreign vessels fishing under charter access arrangements. In 2004 there were six FSM flag purse seiners which caught a total of 27,000 mt. Since then, two vessels have left and one has sunk; the three remaining vessels are partly owned by State governments, with shareholding recently acquired by management of the two companies⁸.

Table 9: Japan's Average Distant Water Purse Seine Catches, 2004-2006

	Skipjack Catch (t)		Yellowfin Catch (t)		Bigeye Catch (t)		Total Catch (t)	
	Per Year	Per Day	Per Year	Per Day	Per Year	Per Day	Per Year	Per Day
Fleet	216,029	919	27,442	117	4,887	21	263,670	1,123
One Vessel	6,172	26	784	3	140	0.55	7,533	32

Source: WCPFC Yearbook 2007

⁸ <http://www.ffa.int/system/files/Trip+report++Palau,+FSM,+RMI.pdf>




Figure 23: Japanese Purse Seine Catch by Location, 2007

Source: WCPFC Yearbook 2007




Figure 24: Japan Purse Seine Catch by Location, 2007 and 2008

Source: Williams and Terawasi 2008

FAD Fishing

There have been increases in the amount of purse seine fishing effort around FADs especially by the U.S. fleet. In 2006, approximately 80 percent of U.S. purse seine sets were associated sets where they set on FADs or other floating objects (Reid 2007).

Deployment of FADs leads to an increased risk that the migratory behavior of tuna in the western Pacific might be affected as FADs may retain tuna in areas where they would otherwise quickly pass through, and not be enticed by concentrated forage to remain. This behavioral modification could affect their biological parameters (growth, maturity, survival) and population dynamics. These potential effects on the population dynamics of tunas and on their ecosystem are largely unknown and require research attention. Most large skipjack tuna (i.e. over 70 cm) are caught by unassociated sets whereas FAD sets and pole-and-line fishing account for most of the smaller sized skipjack catch in the WCPO tuna fishery (OFP 2007). In addition, purse seine catch composition is more varied when fishing around FADs and their use may have higher impacts on biodiversity as compared to purse seine fishing unassociated with FADs (Langley et al. 2006).

As a tuna fishery conservation and management measure, the WCPFC has been considering taking further action to reduce fishing on FADs by prohibiting FAD sets in EEZs and on the high seas, as was proposed at the fourth Commission meeting (WCPFC 2007). If this is finalized, such a measure as this could have a substantial impact on the U.S. fleet which is highly reliant on FAD sets (Figure 26)

Currently, the WCPFC implemented a 2-month FAD closure during August-September (WCPFC 2009). As of August, 2010, there has not yet been a WCPFC regulation prohibiting FAD sets in EEZs or in the high seas. It should be noted, however, that even if a prohibition on FAD sets in the EEZs and high seas were to be implemented by the WCPFC, this would not result in an increase in purse seine fishing in the U.S. EEZ of the Mariana Islands. Figure 27 below shows the distribution of total purse seine effort in the WCPO, 2003-08, by days fishing. The pink shading shows the extent of average sea surface temperature (SST) $>28.5^{\circ}\text{C}$




Figure 25: Time series showing the percentage of total sets by school type for the major purse-seine fleets operating in the WCPO

Source: Williams and Terawasi 2008




Figure 26: The distribution of total purse seine effort in the WCPO, 2003-8, by days fishing

WCPO Purse Seine Fishery Capacity

Through the Palau Arrangement and the FFA, the Pacific Island countries (PICS) maintains a register of fishing vessels allowed to operate in the WCPO region which is currently set at 205 purse seine vessels allowed to operate in the region at any one time. Of the 205 total, the U.S. is allotted up to 40 licenses (+ 5 reserved for vessels owned by joint ventures between U.S. and PIC interests) in accordance with the SPTT⁹. In the event that the U.S. does not fully utilize its full complement of 40 licenses in any fishing year, the unused portion of the licenses is added to “the special arrangements category”. These are licenses that may be issued under the FSM Arrangement¹⁰ or on a bilateral basis to get to the 205 total for the year. In addition to limiting the number of purse seine vessels operating in the WCPO region to 205, the PICS implemented a vessel day scheme (VDS). Under the VDS, each purse seine vessel operating in the region will be allocated a given number of days to fish within the EEZ of the relevant PICS issuing the days. According to the PICS, the number of fishing days allotted to a vessel will correlate to the size of the vessel and the conservation concerns of the PICS. U.S. vessels are not subject to the VDS.

Some Pacific Island nations are beginning to restrict access to fish in their EEZ waters, such as the Cook Islands who now prohibit purse seine fishing in their waters except by SPTT nations [of which the U.S. is a member]; the Kiribati Islands and other members of the Parties to the Nauru Agreement (PNA Group)¹¹. These measures are being implemented to protect the tuna stocks of the PICS and possibly reduce impacts on small-scale and artisanal fleets which target skipjack and yellowfin.

In addition, to strengthen conservation and management of tuna in the Pacific Islands, fishery management measures taken in the WCPO are also moving toward imposing restrictions to distant water fishing nations (DWFNs), including bans on high seas fishing in areas between their EEZs¹².

⁹ See http://www.fpir.noaa.gov/IFD/ifd_sptt.html

¹⁰ See <http://www.ffa.int/taxonomy/term/443>

¹¹ <http://www.ffa.int/node/1101>

¹² See <http://www.ffa.int/node/330>




Figure 27: The WCPFC Convention Area. The PNA Exclusive Economic Zones (EEZs) are shown in yellow (light), Archipelagic Waters and Territorial Seas for Pacific Island Countries and Territories, Australia and New Zealand are shown in white within the EEZs. Coordinates for Archipelagic Waters and Territorial Seas for other CCMs were not available and are not shown. The high-seas pockets wholly enclosed by EEZs between 20 deg N and 20 deg S are shown in black¹³.
 Source: (WCPFC 2008)

All fishing vessels, especially vessels fishing for highly migratory fish need to roam over the ocean looking for fish. The closure of the two high seas pockets denies the purse seine fleets a substantial area of ocean in which they have previously operated looking for skipjack schools. The WCPFC high seas pocket closure on January 2010 forces purse seiners to concentrate their fishing within the EEZs of the Pacific Island nations clustered around the Western Pacific warm pool, or fish in high seas in the Central Pacific.

As discussed in Section 8.4.1., the U.S. fishery currently fishes under a treaty with the Pacific Island Nations (PINs), which expires in 2013. This treaty allows access to the EEZ waters of the PINs, including those of the PNA Group, and the ability to fish on the high seas. Like other purse

¹³ The map in Figure 27 indicative maritime boundaries only. It is presented without prejudice to any past, current or future claims by any State. It is not intended for use to support any past, current or future claims by any State or territory in the western and central Pacific or east Asian region. Individual States are responsible for maintaining the coordinates for their maritime claims. It is the responsibility of flag States to ensure their vessels are informed of the coordinates of maritime limits within the Convention Area. Coastal States are invited to register the coordinates for their negotiated and agreed maritime areas with the Commission secretariat
 Source: http://www.spc.int/coastfish/countries/nauru/nfmra/laws/WCPFC_CMM_2008-01.pdf

seine fleets the U.S. fleet must abide by the Conservation and Management Measures of the WCPFC and is thus now unable to fish in the high seas pockets. The pocket that lies between the Federated States of Micronesia and Papua New Guinea has in the past been an important fishing ground for the U.S. purse seine fleet. Regardless, it is unlikely that the closure of the high seas pockets will greatly change the fishing behavior of the U.S. pure seine fleet, given its access to the PNA Group EEZs.

However, as will be discussed later under Section 9.2.6, the PNA Group intends to prohibit all high seas fishing as part of access agreements to their EEZs. Moreover, the renewal of the treaty beyond 2013 will likely be contingent on issues such as high seas fishing and the amount of vessel days allocated to the U.S. fleet. An unfavorable outcome may make the U.S. EEZ waters a more attractive proposition for fishing by the U.S. fleet, especially the Central Pacific equatorial EEZ segments (Howland and Baker, Jarvis, Palmyra and Kingman Reef), or around the Mariana Islands, the southern part of which borders the warm pool, and has proven skipjack resources (OFP 1998).

High Seas Fishery Compliance Act (HSFCA) permits are issued to vessels that carry an SPTT license. As of November 2009, 39 HSFCA permits with five-year durations have been issued. Two of these purse seine vessels were owned by a Guam limited liability company with majority U.S. ownership. These vessels transhipped their catches primarily where they fished, i.e. ports in the Federated States of Micronesia and Kiribati and have made few port calls into Guam or American Samoa. One vessel sank in June 2010, and the owners have indicated to NMFS that they are interested in replacing the lost vessel (Source: NMFS PIRO Division of International Fisheries). The remaining purse seine vessels are home ported in Pago Pago, American Samoa.

WCPO Purse Seine Vessel Capacity

Gillett and Lewis (2003) estimated the numbers and carrying capacities (amount of fish a vessel can store) of purse seine vessels participating in the tuna fishery of the WCPO during 1988, 1995 and 2003. They considered any vessel with a capacity greater than 400 cubic meters that fished during the year to be participating in the fishery in that year, and excluded vessels that fished only in the EEZs of Indonesia, the Philippines, Australia, New Zealand and other countries of the WCPO. For 1988, they estimated that there were 136 purse-seine vessels with a combined capacity of 140,000 cubic meters (average capacity equal to 1,073 cubic meters). For 1995, they estimated that there were 175 vessels, with a combined capacity of 200,000 cubic meters (average capacity equal to 1143 cubic meters). By 2003 the number of vessels had increased to 191, with a combined carrying capacity of 233,000 cubic meters. This represents a growth of 66 percent between 1988 and 2003 in the capacity of the purse-seine fleet in the WCPO.

In 2009, the U.S. domestic vessels had an average length of 212 ft (65 m) and an average gross registered tonnage (GRT) of 1,500. Fish carrying capacity was estimated to be 34,833 t for the U.S. fleet as a whole. This was an increase from 1995 when the average GRT was 1,181 with an average vessel length of 64.1 meters.

8.4.1 U.S. Purse Seine Fishery-History

Prior to beginning purse seine fishing operations in the Western Pacific, the U.S. fleet had been fishing out of California in areas of the eastern Pacific for decades. The main impetus for the transition from fishing in the eastern Pacific to the western Pacific was due to economic (overcapitalization) reasons, eroding relations with central America states over fishing access issues, increased management controls enacted by the IATTC, and difficulties over environmental concerns associated with fishing on tuna associated with dolphins. During the years when the fleet transitioned from fishing in the eastern Pacific to western Pacific operations U.S. vessels made several gear changes including deepening nets, installing larger power blocks and winches to accommodate larger seines, and using helicopters to spot schools of fish, among other changes (Gillett et al. 2002).

Under the South Pacific Tuna Treaty (SPTT), U.S. tuna fishing vessels are allowed access to fish in vast areas of the western and central Pacific Ocean (subject to certain conditions), including the exclusive economic zones of several Pacific Island States. In addition to the United States, the Parties to the Treaty include the 16 States of the South Pacific Forum Fisheries Agency (FFA): Australia, Cook Islands, Federated States of Micronesia, Fiji, Republic of Kiribati, Republic of the Marshall Islands, Republic of Nauru, New Zealand, Niue, Republic of Palau, Papua New Guinea, Solomon Islands, Kingdom of Tonga, Tuvalu, Republic of Vanuatu and Samoa. The Treaty entered into force on June 15, 1988. While the Treaty itself does not expire, the operational provisions were originally designed for a five-year period. In 1993, the operation of the Treaty was extended for 10 years, until June 14, 2003. The United States and the Pacific Island Parties extended operation of the Treaty for an additional 10-year period, through June 14, 2013.

Under the SPTT and corresponding South Pacific Tuna Act (SPTA), the U.S. provides, via an Economic Assistance Agreement associated with the SPTT, \$18 million annually to the South Pacific Parties to the Treaty for economic development purposes. This is approximately 10 percent of the total amount of U.S. assistance provided to Pacific Islands with almost \$190 million in assistance provided in FY 2006. Of this amount, about \$150 million was comprised of grants from the United States to the Federated States of Micronesia, the Republic of the Marshall Islands, and Palau under the Compacts of Free Association administered by the Department of the Interior. The SPTT governs the conduct of U.S. fishing vessel operations in the Treaty Area. The Treaty Area, which is defined at 50 CFR 300.31, encompasses approximately 10 million square miles (26 million square kilometers). Licenses are issued by the Pacific Islands Forum Fisheries Agency (FFA), based in Honiara, Solomon Islands, which acts as the Treaty administrator on behalf of the Pacific Island Parties to the Treaty (PIPs). Currently, the Treaty allows for a maximum of 45 licenses to U.S. purse seine fishing vessels to fish in the Licensing Area of the Treaty. Of the 45 licenses, 5 are reserved for "joint venture" arrangements with PIPs.

NMFS has announced that persons who enter the purse seine fishery in the WCPO, as managed under the SPTA, the Western and Central Pacific Convention Implementation Act (WCPFCIA) and other laws, after March 28, 2008, are not guaranteed future participation in the fishery if

NMFS decides to revise the criteria and procedures used to process license applications and/or to limit further the number of licenses available in the fishery. NMFS is considering the need to undertake such actions in order to provide greater clarity about the process used and thus help license holders and prospective license applicants in making business decisions, as well as to fulfill the obligations of the United States under international agreements to which it is party.

Requirements for U.S. purse seine vessels fishing under the SPTT including 2007 regulatory changes including: (1) vessel monitoring system (VMS) requirements, (2) vessel reporting requirements, and (3) area restrictions may be found in NOAA's "Small Entity Compliance Guide for the Regulations Implementing the Third Extension of the South Pacific Tuna Treaty for Purse Seine Vessels," June 14, 2007¹⁴. For comprehensive regulatory information please see 50 CFR 300 subpart D.

Foreign purse seine vessels have infrequently fished in the vicinity of Guam, as shown in Table 4, with effort less than 50 boat days since 1980 and an average catch of 500 mt and a high catch of 1,675 mt (OFP 1998). In the vicinity of CNMI, purse seine activity is rare as shown in Table 1 and was reported for only five years since 1980 with the largest catch reported by foreign fleets of 455 mt (434 SKJ and 21 YFT)(OFP 1998).

Before 1995 the purse seine fleet in the WCPO fished mainly on free-swimming schools of tunas; i.e., unassociated sets, perhaps because these sets result in increased value of catch due to large size of fish and the possibility of setting on pure yellowfin schools despite the lower success rates of these types of sets (Gillett et al. 2002). This trend peaked in 1991 when 90 percent of sets were made on unassociated schools and the fleet achieved its highest catches of 216,000 tons (Coan et al. 2002 in Gillett et al. 2002). In 2007, the U.S. fleet made 65 percent of all sets on drifting FADs which is a considerably higher percentage than other nations fishing in the WCPO (OFP 2007b). Skipjack is the dominant component of the catch ranging from approximately 70–74 percent of the total catch, with yellowfin and bigeye tuna comprising the majority of the remaining retained catch.

Activity by the U.S. purse seine fleet in the western and central Pacific Ocean (WCPO) increased from 32 U.S. vessels in 2007 to 39 in 2009. As of 2010, 36 vessels were operating in the WCPO and range in length from 50-115 m with the largest being able to hold up to 3,000 mt or more of frozen fish. Most tuna seiners are 70-80 m in length and can carry approximately 800-1,500 mt of frozen fish. Some vessels also carry helicopters that can improve their ability to find schools of fish and assist in keeping track of the school once the net is set (NMFS 2006).

8.4.2 U.S. Fishing in the PRIA

On January 6, 2009, President Bush established three new National Marine Monuments in the Western Pacific region by issuing Proclamations published in the Federal Register. These monuments were designated by the President using his authority under the Antiquities Act, a 1906 land-oriented statute which does not require public participation or compliance with the National Environmental Policy Act (NEPA). Proclamation 8336 (74 FR 1565) established the Pacific Remote Islands Marine National Monument located in waters around the Pacific Remote

¹⁴ http://www.fpir.noaa.gov/Library/IFD/Compl_guide_third_extension_PS.pdf

Islands Areas (PRIA) of Wake, Baker, Howland, and Jarvis Islands, Johnston Atoll, Kingman Reef, and Palmyra Atoll. The monument prohibits all commercial fishing from within 50 nm around the islands in the PRIA.

As commercial fishing is prohibited in monument waters, this monument designation resulted in displacement of longliners, purse seiners and some bottomfish fishing vessels. U.S. EEZ waters around the PRIA have been fished by the longline and the purse seine fleets based out of Hawaii and American Samoa, respectively. In some years, waters around Palmyra and Kingman Reef are important fishing grounds for the Hawaii-based longline fleet targeting bigeye and yellowfin tuna, however, effort shifts annually (Dowdell et al. 2009). For example, in 2002, more than 20% of the yellowfin and bigeye catch for the Hawaii longline fleet was harvested from within U.S. EEZ waters around Kingman Reef and Palmyra Island.

The U.S. purse seine fishery has historically and exclusively (foreign fishing is prohibited) fished in U.S. EEZ waters around the PRIA. On average, from 5 to 12 percent of the U.S. WCPO annual purse seine catch is harvested from within U.S. EEZ waters around Jarvis, and Howland and Baker Islands, respectively. In 1997, however, almost 20% of the total U.S. purse seine catch in the WCPO was caught from waters of the U.S. EEZ around Howland and Baker Islands. Figure 29 shows the overall average annual catch from the U.S. EEZ waters around Jarvis Island, Howland and Baker Islands, and Kingman Reef and Palmyra Atoll, compared against the average annual catch from within 50 nm (i.e., monument waters) for the period 1988-2007. This figure shows that most of the catch comes from waters outside the new monument designation.




Figure 28: Overall Average Annual Catch from the EEZ Waters around Jarvis Island, Howland and Baker Islands, and Kingman Reef and Palmyra Atoll, and Average Annual Catch from within 50 nm, (i.e., Monument Waters), 1988-2007

8.4.3 U.S. Purse Seine Fishery-Present and Future

Currently, most of the fishing activity by U.S. purse seine vessels occurs in areas between 5° N and 10° S latitude and 150° E and 170° W longitude in the EEZ waters surrounding Papua New Guinea, Federated States of Micronesia and other Pacific island nations. During El Nino events, however, purse seine fishing activity shifts east to the equatorial central Pacific. The percentage of yellowfin tuna in their catches often increases with this shift. U.S. purse seine catch locations for 2007 and 2008 and amount of effort, illustrated in (Figure 30) show the large increase in effort and for 2008 a shift to the west, closer to the Mariana Archipelago.

The purse seine catch is stored as a frozen whole product. Most of the catch has historically been off-loaded to the canneries in Pago Pago, American Samoa, however more and more vessels are transshipping catches in Pacific Islands ports for canning and loining destinations in Southeast Asia and Latin America. The final product that is canned in American Samoa is typically destined for the domestic U.S. canned tuna markets. Frozen non-tuna catches may be processed locally (e.g., wahoo) or transshipped to foreign destinations (e.g., billfish and shark)(NMFS 2009).

The number of U.S. vessels active in the WCPO purse seine fleet declined from 39 active vessels in 1998 to 15 in 2005 (WCPFC 2006b) with catches following suit. Increased investment and participation in the U.S. purse seine fishery is evident (NMFS 2009). However, since 2005, the U.S. purse seine fleet has more than doubled to 32 vessels in 2008. The most recent information regarding purse seine activity is that there are 36 purse seiners licensed in the U.S. for the 2009/2010 licensing year; the U.S. may license up to 40 vessels.

The U.S. purse seine catch of 157,849 mt in 2008 was composed primarily of skipjack tuna, with smaller catches of yellowfin and bigeye (Table 10). Total catches and effort increased significantly from 2007. Yellowfin tuna catches in the fishery increased from 10,541 mt in 2007 to 23,801 mt in 2008 and catches of skipjack have doubled since 2005 to 127,307 mt in 2008. Catch rates in terms of mt of fish caught per day by U.S. purse seiners are shown in Figure 31. The average catch rate for 2006 and 2007, based on data in WCPFC 2008 was 23.1 and 3.3 mt/day respectively for skipjack and yellowfin tuna.

Table 10: Reported Annual Landings (metric tons) for U.S. Purse Seine Vessels Operating in the WCPO, 1999-2008

Year	# Active Vessels	Skipjack (mt)	Yellowfin (mt)	Bigeye (mt)
1999	36	129,262	34,529	18,694
2000	33	81,368	29,961	13,886
2001	32	85,539	24,143	6,176
2002	29	88,535	27,191	4,889
2003	26	62,907	20,079	4,470
2004	21	47,896	14,492	5,031
2005	15	62,379	17,685	6,108
2006	13	55,633	8,448	4,364
2007	21	75,210	10,541	2,985
2008	32	127,307	23,801	6,741

Source: WCPFC 2008b




Figure 29: Distribution of Effort by the U.S Purse Seine Fleet, 2007 and 2008

Note: Mariana Archipelago is located in the mid upper left-hand quadrant.

Source: Williams and Terawasi 2008




Figure 30: U.S. Purse Seine Catch Rates in the WCPO

Source: WCPFC 2008

To date no U.S. purse seine catches have been recorded in U.S. EEZ waters around CNMI. Although catches by the Japanese pole-and-line fleet from the 1980's show that these waters have the capacity to produce tunas in large quantities (see Table 1). As noted earlier, two of these purse seine vessels were owned by a Guam limited liability company with majority U.S. ownership. These vessels transhipped their catches primarily where they fished, i.e. ports in the Federated States of Micronesia and Kiribati and have made few port calls into Guam or American Samoa. One vessel sank in June 2010, and the owners have indicated to NMFS that they are interested in replacing the lost vessel (Source: NMFS PIRO Division of International Fisheries).

The largest element of purse seine catches in U.S. EEZ waters of the Western Pacific is from purse seining around Howland and Baker Islands (Figure 31), where this segment of the U.S. EEZ straddles the equator in the Central Pacific. Catches from these waters have ranged between 500 to 23,500 mt with the annual average catches of about 8,500 mt (PIFSC data).




Figure 31: U.S. Purse Seine SKJ Catch from Howland and Baker Islands, PRIA

Bycatch

Purse seine floating object sets are known to result in greater overall amounts and species of bycatch than unassociated sets. Non-target species caught and discarded by the U.S. purse seine fleet in 2004 and 2005 included: marlins, silky shark, wahoo, mackerel scad, mahimahi, oceanic triggerfishes, rainbow runner, and others. Table 12 shows the amount of discard per vessel calculated from the data provided in Table 11. Based on observer data collected under the U.S. Multilateral Treaty the overall discard levels by the U.S. fleet for tunas varied from 2.4 percent to 20.2 percent (Tables 12 & 13 Source: NMFS 2007, SPC 2009). The main reasons for discard of tunas, including skipjack are: small size (i.e. too small for canning), gear damaged, and vessel fully loaded (OFP 2007). Associated-school sets accounted for higher rates of tuna discards than unassociated-school sets (i.e. more small fish are associated with drifting FADs than unassociated schools).

Nearly all the bycatch and discard species are important food fishes for the people of the Mariana Archipelago. Blue marlin is an important pelagic species for the troll and charter fleets in the Pacific Islands. The most recent stock assessment was in 2002 (Kleiber et al 2002) which concluded that current fishing effort in the early part of this decade is producing close to the maximum sustainable yield (MSY). However, because of uncertainty in projected yield curves, and because of sensitivity of the model to assumptions such as selectivity, the situation may well be more optimistic with current levels significantly less than those that would produce MSY. An

update of the stock assessment is anticipated to be conducted by the International Scientific Committee for Tuna and Tuna-like Species in the North Pacific Ocean (ISC 2010).

Table 11: Purse Seine Discards by U.S. Vessels Fishing in the WCPO, 2005 and 2006

Species or species group	Metric tons			
	2005	2006	2007	2008
Target species discards				
Total:	3,900	1,245	1,560	245
Skipjack tuna (<i>Katsuwonus pelamis</i>)	3,796	1,190	1,417	223
Yellowfin tuna (<i>Thunnus albacares</i>)	104	55	143	22
Other discards				
Billfish:	16.09	18.1	7.5	9.8
Black marlin (<i>Makaira indica</i>)	0.17	0.34	3.01	2.93
Sailfish (<i>Istiophorus platypterus</i>)	0.08	0.12	0.15	0.15
Marlin (Istiophoridae)	15.84	17.64	4.34	6.74
Sharks/Rays:	11.18	1.06	9.53	9.38
Oceanic whitetip shark (<i>Carcharhinus longimanus</i>)	0	0.03	0.21	1.62
Silky shark (<i>Carcharhinus falciformis</i>)	0.23	1.03	8.73	6.19
Shark	10.76	0	0.05	0.52
Rays, skates, mantas (Rajiformes)	0.19	0	0.54	1.05
Other tunas/Tuna-like species:	10.13	3.87	3.45	3.72
Albacore tuna (<i>Thunnus alalunga</i>)	4.50	0	0.04	0.5
Tuna unspecified (tribe Thunnini)	2	0	0	0.21
Mackerel (Scombridae)	3.23	3.51	0.35	
Wahoo (<i>Acanthocybium solandri</i>)	0.40	0.36	3.06	3.01
Other fish:	234.94	80.22	97.5	94.04
Mahimahi (<i>Coryphaena hippurus</i>)	0.69	0.62	2.96	12.79
Ocean sunfish (<i>Mola mola</i>)	0	0.02	0	0.07
Rainbow runner (<i>Elagatis bipinnulatus</i>)	2.65	5.02	78.63	36.65
Triggerfish (Balistidae)	0.69	0.58	15.93	1.90
Yellowtail (<i>Seriola lalandi</i>)	11.61	14.79	0.01	0.28
Other unspecified	219.99	59.19	25.01	42.35

Source: NMFS 2007, SPC 2009

Table 12: 2005 U.S. Purse Seine Discards, (average/vessel)

FISH	Amount Discarded (lb)
Skipjack	573,196
Yellowfin	15,285
Marlins	2,352
Albacore	6,614
Wahoo	59
Mahimahi	101
Misc. spp.	31,158

Table 13: Percent of Catch Discarded by U.S. Purse Seine Fleet

Year	Skipjack tuna	Yellowfin tuna	Bigeye tuna
2005	14.1	7.7	20.2
2006	2.6	4.5	2.4
2007	8.8	6.4	10.2
2009	1.2	0.9	4.4

Source: OFP (2007). Note: Data based on 20% observer coverage.

8.4.4 Fishery Interactions Between Various Gear Types

Determining the degree of fishery interactions between purse seining and longline or other small-scale pelagic fisheries has been studied extensively (see references cited below). However, the demonstration of a correlation of the negative impacts of one sector upon another is difficult and the results of such studies far from definitive. The studies cited below are all drawn from data on various commercial fisheries and not the results of controlled experiments where factors such as seasonal tuna availability, differences in skill levels between fishery participants and variable oceanographic conditions may confound the results. Nevertheless, the examples summarized in this section do indicate that there may be competition between different pelagic fisheries operating within proximity to one another.

A 1995 study investigated the impact of purse seine fishing on troll and domestic pole-and-line and artisanal fisheries in waters around Kiribati (Hampton et al. 1995). Over large areas, e.g., within radii of 300-600 nm of the islands, artisanal catch rates and purse seine catches were generally positively correlated, suggesting that, on this scale, variations in the abundance or catchability of yellowfin affect both purse seiners and artisanal catches in the same way. However, some negative correlations were found for smaller area (60 nm) and time scales, indicating that localized effects may occur.

Hampton et al. 1995 also conducted an analyses of tagging data to estimate the average impact of purse seine fishing on the Kiribati pole-and-line and artisanal catch rates. A spatially aggregated model for the Gilbert Islands area indicated only a modest overall impact of local purse seine

catches on pole-and-line and artisanal catch rates. A regional model with 1° square spatial structure indicated a slightly higher overall impact of the regional purse seine fishery on skipjack catches by the Kiribati pole-and-line fishery. The results of the correlation and tagging data analyses suggest that adverse impacts of purse seine fishing on artisanal and pole and- line catches in the Gilbert Islands were more likely to occur at a small scale (1° square or less) due to local concentrations of purse seine effort, rather than at a regional scale or on a scale of tens of degrees.

In addition, there is some indication from additional preliminary modeling of theoretical skipjack tuna movement that purse seine fishing could have an impact on the availability of skipjack tuna for small-scale trolling from equivalent levels of fishing effort occurring as far away as 600-800 nm (P. Kleiber, NOAA PIFSC, pers. comm.). Taken together the Hampton et al (1995) and Kleiber studies suggest that every mile of separation between purse seine fisheries and small-scale troll fisheries is beneficial for troll fisheries.

Another study was undertaken to examine the influence of purse seine fishing on longline catches of yellowfin tuna in the Indian Ocean (Nishida 1995). Using a spatial analysis this study's data analysis showed that the catch of the purse seine fishery affects the longline fishery when the purse seine catch exceeds 1 ton/month per 5° x 5° area. There was a modest decrease in longline CPUE when data for the heavily fished purse seine area were examined for impacts to the longline fishery. The author of this study does point out, however, that a decrease in CPUE of the longline caught yellowfin tuna does not necessarily imply impacts of the purse seine fishery; other factors such as environmental conditions and fish availability may be responsible for the changes in CPUE (Nishida 1995). Although this study occurred on a fishery in the Indian Ocean it is likely that similar outcomes may be expected on western Pacific tuna fisheries.

A similar study on interactions between Mexican longline and purse seine fisheries in the eastern Pacific concluded that the information was not sufficient to define an interaction; it was possible to suggest a "consecutive" interaction (Ortega-Garcia 1995). Another study looked at skipjack movement with regards to fisheries interactions in the western Pacific (Sibert et al. 1995). This study predicted declining yields of 20 percent in the pole-and-line fishery due to purse seine fishing activities, with greatest declines for pole-and-line fleets operating closest to the purse seine fleets. The authors point out that a change of this magnitude would be difficult to detect in actual recorded catches and to be separated from inter-annual variability in the fisheries. At the time of this study (late 80's – early 90's) pole-and-line and purse seine were the two principal gear types catching skipjack in the western Pacific. They also point out that potential fishery interaction estimates depend on the natural mortality rate as does the estimated movement pattern used in their analysis (Sibert et al. 1995).

More recently, the Council has examined the interactions of purse seine, longline and troll vessels in the U.S. EEZ around American Samoa as part of an amendment to implement a 75 nm purse seine area closure around the American Samoa archipelago¹⁵. A statistically significant correlation was found which accounted for about 40% of the variation between skipjack and

¹⁵ Establishment of Purse Seine Fishing Prohibited Areas Around American Samoa. Draft Amendment to the Fishery Ecosystem Plan for Pelagic Fisheries of the Western Pacific Region Including an Environmental Assessment November 11, 2010

yellowfin troll CPUE versus skipjack and yellowfin purse seine catch. Each 100 mt of yellowfin and skipjack caught by purses seining reduces the combined troll CPUE for these two species by about 30%. Regressions of skipjack troll CPUE versus skipjack purse seine catch and total troll CPUE versus skipjack and yellowfin purse seine catch were borderline significant, i.e. with probability values between 0.1 and 0.05 levels of significance. Interestingly, the relationships between skipjack, yellowfin and total troll CPUE and longline yellowfin and skipjack catches in isolation or combined with purse seine catches yielded regressions with flat or positive slopes.

These relationships may be statistical artifacts, but the possibility that purse seine catches in the U.S. EEZ around American Samoa, especially of skipjack, do have some weak but detectable influence on troll CPUEs. However, the mechanism by which this occurs relative to the much larger catches of skipjack and yellowfin made by the American Samoa longline fishery is unknown. One potential hypothesis is that large catches of yellowfin and skipjack may remove fish which are abundant at the surface and available to troll fishermen thus depressing the troll CPUE.

In summary the conclusions of the studies cited above and others contained in Shomura et al. (1994; 1996) indicate that detection of fishery interactions, where one tuna fishery can be shown to be affecting another, is generally difficult and the degree of correlation is often weak.

A study examined the migratory nature of some tunas, specifically skipjack and yellowfin, and the implications for fisheries management (Sibert and Hampton 2003). They found that the recoveries of most skipjack tags from two major tuna tagging programs in the Western & Central Pacific spanning three decades (1970s-1990s) were within 200 nautical miles of the release point. Moreover the median displacement¹⁶ of skipjack was only 28 nautical miles in the earliest study and 158 miles in a later study. The later study also produced a median displacement of yellowfin of 132 nautical miles. Sibert and Hampton (2003) also estimated the displacements of all tagged fish in their lifetimes. Skipjack were found to have median lifetime displacements ranging from from 420-470 nm and for yellowfin approximately 20% less (about the extent of an EEZ unconstrained by a neighboring EEZ). They also concluded that fishing reduces the median displacement. The key conclusion from this study was that a substantial proportion of skipjack and yellowfin population production takes place within a single EEZ, particularly for skipjack. Longer periods of residency provide the opportunity for implementation of effective unilateral domestic policies. It should be possible to base successful domestic fisheries on the expected yield from that portion of the population that persists within EEZs.

8.4.5 Foreign Fishing in U.S. EEZ Waters

Section 204 of the MSA authorizes the Secretary of State to enter into a Pacific Insular Area Fishery Agreement (PIAFA) with a foreign nation that would allow vessels of that nation to fish in EEZ waters around Guam, CNMI, American Samoa, or other U.S. possessions in the Pacific. Fishing under a PIAFA must comply with any applicable FMP or FEP, and harvest levels must fall within the total allowable level of foreign fishing (TALFF). A PIAFA can be effective for up

¹⁶ The median displacement is the displacement exceeded by 50% of tagged fish. It can be expressed empirically, i.e. the distance travelled between mark and recapture, or estimated from the tagging data for the lifetime of the fish from the point of release.

to three years, and may be concluded only with the concurrence of the appropriate Governor, if any. A PIAFA may be concluded only after a Marine Conservation Plan (MCP) for the insular area has been developed by the Governor (in the case of Guam, CNMI, and American Samoa) or by the Western Pacific Fishery Management Council (in the case of other western Pacific territories). An MCP identifies conservation and management objectives for the marine and coastal environment of the island area, and prioritizes those objectives. Neither the Guam nor the CNMI MCP, completed in 2009, contain provisions for purse seine fishing. Fees collected under a PIAFA for fishing around Guam, CNMI, or American Samoa are deposited in the treasury of the island government; the Governor may spend the funds to implement an approved MCP, to compensate the Council or the State Department for various administrative or travel expenses, or to carry out other purposes of section 204(e). Currently, there are no PIAFAs in effect between CNMI/Guam and foreign countries.

8.5 Protected Species

8.5.1 Sea Turtles

All Pacific sea turtles are designated under the Endangered Species Act (ESA) as either threatened or endangered (except for the flatback turtle). The breeding populations of Mexico's olive ridley sea turtles (*Lepidochelys olivacea*) are currently listed as endangered, while all other ridley populations are listed as threatened. Leatherback sea turtles (*Dermochelys coriacea*) and hawksbill turtles (*Eretmochelys imbricata*) are also classified as endangered. Loggerhead (*Caretta caretta*) and green sea turtles (*Chelonia mydas*) are listed as threatened (the green sea turtle is listed as threatened throughout its Pacific range, except for the endangered population nesting on the Pacific coast of Mexico). These five species of sea turtles are highly migratory, or have a highly migratory phase in their life history (NMFS 2001).

Leatherback Sea Turtles

Leatherback turtles (*Dermochelys coriacea*) are widely distributed throughout the oceans of the world, and are found in waters of the Atlantic, Pacific, and Indian Oceans; the Caribbean Sea; and the Gulf of Mexico (Dutton et al. 1999). Increases in the number of nesting females have been noted at some sites in the Atlantic (Dutton et al. 1999), but these are far outweighed by local extinctions, especially of island populations, and the demise of once-large populations throughout the Pacific, such as in Malaysia (Chan and Liew 1996) and Mexico (Sarti et al. 1996; Spotila et al. 1996). In other leatherback nesting areas, such as Papua New Guinea, Indonesia, and the Solomon Islands, there have been no systematic, consistent nesting surveys, so it is difficult to assess the status and trends of leatherback turtles at these beaches. In all areas where leatherback nesting has been documented, current nesting populations are reported by scientists, government officials, and local observers to be well below abundance levels of several decades ago. The collapse of these nesting populations was most likely precipitated by a tremendous overharvest of eggs coupled with incidental mortality from fishing (Sarti et al. 1996).

Leatherback turtles lead a completely pelagic existence, foraging widely in temperate waters, except during the nesting season when gravid females return to tropical beaches to lay eggs. Males are rarely observed near nesting areas, and it has been proposed that mating most likely takes place outside of tropical waters, before females move to their nesting beaches (Eckert and

Eckert 1988). Leatherbacks are highly migratory, exploiting convergence zones and upwelling areas in the open ocean, along continental margins, and in archipelagic waters (Eckert 1998). In a single year, a leatherback may swim more than 10,000 kilometers (Eckert 1998).

Satellite telemetry studies indicate that adult leatherback turtles follow bathymetric contours over their long pelagic migrations and typically feed on cnidarians (jellyfish and siphonophores) and tunicates (pyrosomas and salps), and their commensals, parasites, and prey (NMFS 1998). Females are believed to migrate long distances between foraging and breeding grounds, at intervals of typically two or four years (Spotila et al. 2000). In the western Pacific, nesting peaks on Jamursba-Medi Beach (Papua, Indonesia) from May to August, on War-Mon Beach (Papua) from November to January (Starbird and Suarez 1994), in peninsular Malaysia during June and July (Chan and Liew 1989), and in Queensland, Australia in December and January (Limpus and Reimer 1994).

Migratory routes of leatherback turtles originating from eastern and western Pacific nesting beaches are not entirely known. However, satellite tracking of post-nesting females and genetic analyses of leatherback turtles caught in U.S. Pacific fisheries or stranded on the west coast of the U.S. presents some strong insights into at least a portion of their routes and the importance of particular foraging areas.

Leatherback Sea Turtles in the Mariana Archipelago

There have been occasional sightings of leatherback turtles around Guam (Eldredge 2003); however, the extent to which (i.e. preferred location, abundance, seasonality) leatherback turtles are present around the Mariana Archipelago is unknown (WPRFMC 2007b).

Loggerhead Sea Turtles

Loggerheads can be divided into five regions: the Atlantic Ocean, Pacific Ocean, Indian Ocean, Caribbean Sea and Mediterranean Sea. These regions may be further divided into nesting aggregations. In the Pacific Ocean, loggerhead turtles are represented by a northwestern Pacific nesting aggregation (located in Japan) which may be comprised of separate nesting groups (Hatase et al. 2002) and a smaller southwestern nesting aggregation that occurs in Australia (Great Barrier Reef and Queensland) and New Caledonia (Limpus, 2006).

North Pacific loggerhead turtles nest in Japan, undertake trans-Pacific developmental migrations in the waters of the Central North Pacific, Mexico, and U.S. territorial waters throughout the Eastern Pacific, and reside as adults in waters of the Asian region. Loggerheads originating in Japan travel westward, move seasonally north and south primarily through the region 28–40°N, and occupy sea surface temperatures (SST) of 15–25°C. Their dive depth distribution indicated that they spend 40 percent of their time at the surface and 90 percent of their time at depths <40 m. Loggerheads are found in association with fronts, eddies, and geostrophic currents. Specifically, the North Pacific Transition Zone Chlorophyll Front and the southern edge of the Kuroshio Extension Bifurcation Region appear to be important forage and migration habitats for loggerheads (Polovina et al., 2004 and 2007).

The loggerhead sea turtle is listed as threatened under the ESA throughout its range, primarily due to direct take, incidental capture in various fisheries, and the alteration and destruction of its

habitat. In the South Pacific, Limpus (1982) reported an estimated 3,000 loggerheads nesting annually in Queensland, Australia during the late 1970s. However, long-term trend data from Queensland indicate a 50 percent decline in nesting by 1988–89 due to incidental mortality of turtles in the coastal trawl fishery. This decline is corroborated by studies of breeding females at adjacent feeding grounds (Limpus and Reimer 1994). Currently, approximately 300 females nest annually in Queensland, mainly on offshore islands (Capricorn-Bunker Islands, Sandy Cape, Swains Head; Dobbs 2001). In southern Great Barrier Reef waters, nesting loggerheads have declined approximately 8 percent per year since the mid-1980s (Heron Island), while the foraging ground population has declined 3 percent and comprised less than 40 adults by 1992. Researchers attribute the declines to recruitment failure due to fox predation of eggs in the 1960s and mortality of pelagic juveniles from incidental capture in longline fisheries since the 1970s (Chaloupka and Limpus 2001). Of the loggerheads taken in the Hawaii-based longline fishery, all have been determined to have originated from Japanese nesting beaches, based on genetic analyses (P. Dutton, NMFS, pers. comm., August 2005).

Loggerhead Sea Turtles in the Mariana Archipelago

There are no known reports of loggerhead turtles in waters around the Mariana Archipelago (WPRFMC 2007b).

Green Sea Turtles

Green turtles were listed as threatened under the ESA on July 28, 1978, except for breeding populations found in Florida and the Pacific coast of Mexico, which were listed as endangered. Using a precautionary estimate, the number of nesting female green turtles has declined by 48 to 67 percent over the last three generations (approx. 150 years; Troeng and Rankin 2005). Causes for this decline include harvest of eggs, subadults, and adults; incidental capture by fisheries; loss of habitat; and disease. The degree of population change is not consistent among all index nesting beaches or among all regions. Some nesting populations are stable or increasing (Balazs and Chaloupka 2004; Chaloupka and Limpus 2001; Troeng and Rankin 2005). However, other populations or nesting stocks have markedly declined. Because many of the threats that have led to these declines have not yet ceased, it is evident that green turtles face a measurable risk of extinction (Troeng and Rankin 2005).

Green sea turtles are a circumglobal and highly migratory species, nesting and feeding in tropical/subtropical regions. Their range can be defined by a general preference for water temperature above 20° C. Green sea turtles are known to live in pelagic habitats as post-hatchlings/juveniles, feeding at or near the ocean surface. The non-breeding range of this species can lead a pelagic existence many miles from shore while the breeding population lives primarily in bays and estuaries, and are rarely found in the open ocean. Most migration from rookeries to feeding grounds is via coastal waters, with females migrating to breed only once every two years or more (Bjorndal 1997).

Tag returns of eastern Pacific green turtles (often reported as black turtles) establish that these turtles travel long distances between foraging and nesting grounds. In fact, 75 percent of tag recoveries from 1982–1990 were from turtles that had traveled more than 1,000 kilometers from Michoacán, Mexico. Even though these turtles were found in coastal waters, the species is not confined to these areas, as indicated by sightings recorded in 1990 from a NOAA research ship.

Observers documented green turtles 1,000–2,000 statute miles from shore (Eckert 1993). The east Pacific green is also the second-most sighted turtle in the east Pacific during tuna cruises; they frequent a north–south band from 15° N to 5° S along 90° W and an area between the Galapagos Islands and the Central American Coast (NMFS 1998).

The non-breeding range of green turtles is generally tropical, and can extend approximately 500–800 miles from shore in certain regions (Eckert 1993). The underwater resting sites include coral recesses, undersides of ledges, and sand bottom areas that are relatively free of strong currents and disturbance from natural predators and humans. In the Pacific, the only major (> 2,000 nesting females) populations of green turtles occur in Australia and Malaysia. Smaller colonies occur in the insular Pacific islands of Polynesia, Micronesia, and Melanesia (Wetherall 1993) and on six small sand islands at French Frigate Shoals, a long atoll situated in the middle of the Hawaii Archipelago (Balazs et al. 1994).

Green Sea Turtles in the Mariana Archipelago

Based on nearshore surveys conducted jointly between the CNMI–DFW and NMFS around the Southern Mariana Islands (Rota and Tinian 2001; Saipan 1999), an estimated 1,000 to 2,000 green sea turtles forage in these areas (Kolinski 2001). The green sea turtle is a traditional food of the native population and although harvesting them is illegal, divers have been known to take them at sea and others have been taken as nesting females (NMFS and USFWS 1998a). Turtle eggs are also harvested in CNMI. Nesting beaches and seagrass beds on Tinian and Rota are in good condition but beaches and seagrass beds on Saipan have been impacted by hotels, golf courses and general tourist activities.

Nesting surveys for green sea turtles have been done on Guam since 1973 with the most consistent data collected since 1990. There have been up to 60 nesting females observed annually, with a generally increasing trend over the past 12 years aerial surveys done in 1999–2000 also found an increase in green sea turtle sightings around Guam (Cummings 2002).

Hawksbill Sea Turtles

The hawksbill turtle is listed as endangered throughout its range. In the Pacific, this species is rapidly approaching extinction primarily due to the harvesting of the species for its meat, eggs, and shell, as well as the destruction of nesting habitat by human occupation and disruption. Along the eastern Pacific Rim, hawksbill turtles were common to abundant in the 1930s (Cliffton et al. 1982). By the 1990s, the hawksbill turtle was rare to absent in most localities where it was once abundant (Cliffton et al. 1982).

Hawksbills are circumtropical in distribution, generally occurring from latitudes 30° N to 30° S within the Atlantic, Pacific, and Indian Oceans and associated bodies of water (NMFS 1998). Hawksbills have a unique diet of sponges (Meylan 1985, 1988). While data are somewhat limited on their diet in the Pacific, it is well documented that in the Caribbean hawksbill turtles are selective spongivores, preferring particular sponge species over others (Van Dam and Diez 1997). Foraging dive durations are often a function of turtle size, with larger turtles diving deeper and longer. As a hawksbill turtle grows from a juvenile to an adult, data suggest that the turtle switches foraging behaviors from pelagic surface feeding to benthic reef feeding (Limpus 1992). Within the Great Barrier Reef of Australia, hawksbills move from a pelagic existence to a

“neritic” life on the reef at a minimum CCL of 35 centimeters. The maturing turtle establishes foraging territory and will remain in this territory until it is displaced (Limpus 1992). As with other sea turtles, hawksbills will make long reproductive migrations between foraging and nesting areas (Meylan 1999), but otherwise they remain within coastal reef habitats. In Australia, juvenile turtles outnumber adults 100:1. These populations are also sex biased, with females outnumbering males 2.57:1 (Limpus 1992).

Along the far western and southeastern Pacific, hawksbill turtles nest on the islands and mainland of southeast Asia, from China to Japan, and throughout the Philippines, Malaysia, Indonesia, Papua New Guinea, the Solomon Islands (McKeown 1977), and Australia (Limpus 1982).

Hawksbill Sea Turtles in the Mariana Archipelago

Although hawksbill turtles have occasionally been sighted in the past around CNMI they were not observed in a detailed assessment conducted in 1999, nor were they observed in 10 aquatic surveys along the shores of Tinian in 1995. According to the 1998 Pacific Sea Turtle Recovery Team Recovery Plan for the hawksbill turtle (NMFS and USFWS 1998b), there are no reports of nesting in CNMI. This does not rule out the possibility of a few hawksbill nests, as nesting surveys on small pocket beaches in remote areas of CNMI have never been done. A single hawksbill sighting occurred in 1996 during the detonation of an unexploded ordinance off of Rota. The turtle was recovered near the explosion sight and subsequently died, apparently from internal injuries incurred from the blast (Trianni 1998). One hawksbill sea turtle nest was found in November 1991 on Guam (NMFS and USFWS 1998b); however this was highly unusual as nesting individuals are otherwise virtually unknown on Guam (Eldredge 2003).

Olive Ridley Sea Turtles

Olive ridley turtles are olive or grayish green above, with a greenish white plastron, and adults are moderately sexually dimorphic (NMFS and FWS 1998b). Olive ridleys lead a highly pelagic existence (Plotkin 1994). These sea turtles appear to forage throughout the eastern tropical Pacific Ocean, often in large groups, or flotillas. Olive ridleys generally have a tropical range, however, individuals do occasionally venture north, some as far as the Gulf of Alaska (Hodge and Wing 2000). The postnesting migration routes of olive ridleys, tracked via satellite from Costa Rica, traversed thousands of kilometers of deep oceanic waters ranging from Mexico to Peru and more than 3,000 kilometers out into the central Pacific (Plotkin 1994). Stranding records from 1990–1999 indicate that olive ridleys are rarely found off the coast of California, averaging 1.3 strandings annually (J. Cordaro, NMFS, personal communication, NMFS 2004). At least one olive ridley was reported in Micronesia (Yap) in 1973 (Falanruw et al. 1975).

The olive ridley turtle is omnivorous, and identified prey include a variety of benthic and pelagic prey items such as shrimp, jellyfish, crabs, snails, and fish, as well as algae and seagrass (Marquez 1990). It is also not unusual for olive ridley turtles in reasonably good health to be found entangled in scraps of net or other floating synthetic debris. Small crabs, barnacles, and other marine life often reside on debris and are likely to attract the turtles. Olive ridley turtles also forage at great depths, as a turtle was sighted foraging for crabs at a depth of 300 meters (Landis 1965, in Eckert et al. 1986). The average dive lengths for adult females and males are reported to be 54.3 and 28.5 minutes, respectively (Plotkin 1994, in Lutcavage and Lutz 1997).

Olive Ridley Sea Turtles in the Mariana Archipelago

There are no known reports of olive ridley turtles in waters around the Mariana Archipelago (WPRFMC 2007b).

8.5.2 Marine Mammals

Cetaceans listed as endangered under the ESA that have been observed in Mariana Archipelago include the humpback whale (*Megaptera novaeangliae*), sperm whale (*Physeter macrocephalus*), and sei whale (*Balaenoptera borealis*) (WPRFMC 2007b). Other ESA listed marine mammals that may occur in the EEZ around the Mariana Archipelago include the blue whale (*Balaenoptera musculus*) and the fin whale (*Balaenoptera physalus*).

Blue Whale

The blue whale is listed as endangered under the ESA and is the largest living animal. Blue whales can reach lengths of 30 meters and weights of 160 tons (320,000 lb), with females usually being larger than males of the same age. They occur in all oceans, usually along continental shelves, but can also be found in the shallow inshore waters and on the high seas. No sightings or strandings of blue whales have been reported in Hawaii, but acoustic recordings made off Oahu and Midway Islands have reported blue whales somewhere within the EEZ around Hawaii. The stock structure of blue whales in the North Pacific is uncertain. The status of this species in Hawaii waters relative to the optimum sustainable population is unknown, and there are insufficient data to evaluate trends in abundance.

Prior to whaling, the worldwide population of blue whales is believed to have been about 200,000 animals. Only 8,000-12,000 are estimated to be alive today. Blue whales have always been more abundant in the Antarctic than in the northern hemisphere. An estimated 4,900 to 6,000 blue whales are believed to have inhabited the North Pacific prior to whaling. The North Pacific population is now estimated at 1,200 to 1,700 animals.

Fin Whales

Fin whales are listed as endangered under the ESA and found throughout all oceans and seas of the world from tropical to polar latitudes. Although it is generally believed that fin whales make poleward feeding migrations in summer and move toward the equator in winter, few actual observations of fin whales in tropical and subtropical waters have been documented, particularly in the Pacific Ocean away from continental coasts. There have only been a few sightings of fin whales in Hawaii waters. There is insufficient information to accurately determine the population structure of fin whales in the North Pacific, but there is evidence of multiple stocks.

Humpback Whales

Humpback whales can attain lengths of 16 meters. Humpback whales winter in shallow nearshore waters of 100 fathoms or less. Mature females are believed to conceive on the breeding grounds one winter and give birth the following winter. At least six well-defined breeding stocks of humpback whales occur in the Southern Hemisphere Genetic and photo identification studies indicate that within the U.S. EEZ in the North Pacific, there are at least

three relatively separate populations of humpback whales that migrate between their respective summer/fall feeding areas to winter/spring calving and mating areas (Hill and DeMaster 1999). The Central North Pacific stock of humpback whales winters in the waters of the Main Hawaiian Islands (Hill et al. 1997). The humpbacks that winter in the Mariana Archipelago are believed to be part of the “Asian” stock, which migrate from the Bonin (Ogasawara) Islands (Eldredge 2003). Humpback whales have been sighted around Guam and CNMI (Eldredge 2003), however, the number of whales that winter in the Mariana Archipelago each year is unknown.

There is no precise estimate of the worldwide humpback whale population. The humpback whale population in the North Pacific Ocean basin is estimated to contain 6,000–8,000 individuals (Calambokidis et al. 1997). The Central North Pacific stock appears to have increased in abundance between the early 1980s and early 1990s; however, the status of this stock relative to its optimum sustainable population size is unknown (Hill and DeMaster 1999).

Sei Whales

Sei whales have a worldwide distribution but are found mainly in cold temperate to subpolar latitudes rather than in the tropics or near the poles (Horwood 1987). They are distributed far out to sea and do not appear to be associated with coastal features. Two sei whales were tagged in the vicinity of the Northern Mariana Islands (Reeves et al. 1999). The International Whaling Commission considers there to be one stock of sei whales in the North Pacific, but some evidence exists for multiple populations (Forney et al. 2000). In the southern Pacific most observations have been south of 30° (Reeves et al. 1999).

Sperm Whales

Sperm whales are found in tropical to polar waters throughout the world (Rice 1989). They are among the most abundant large cetaceans in the region. Sightings of sperm whales were made during May–July in the 1980s around Guam, and in recent years stranding of dwarf and pygmy sperm whales have been reported on Guam (Reeves et al. 1999).

Dugong

Dugongs are members of the Sirenia order, which include sea cows and manatees, and have a distribution from the east African coast to islands in the southwestern Pacific. In general, dugongs remain in coastal areas, such as protected bays and mangrove channels, feeding primarily on seagrass. Dugongs can be found further offshore in areas that have wide, shallow, and protected continental shelves (Marsh et al. 2002).

A single dugong (*Dugon dugong*) was observed in Cocos Lagoon, Guam in 1975 (Randall et al. 1975). Several sightings were reported in 1985 on the southeastern side of Guam (Eldredge 2003). Since that time, however no reports of dugong sightings have been made. No observations of dugongs have been reported for CNMI.

Other Marine Mammals

Table 14 lists known non-ESA listed marine mammals that have been observed in the Mariana Archipelago and all protected by the Marine Mammal Protection Act (MMPA).

Table 14: Non-ESA Listed Marine Mammals

Common Name	Scientific Name
Blainville's beaked whale	<i>Mesoplodon densirostris</i>
Bottlenose dolphin	<i>Tursiops truncatus</i>
Bryde's whale	<i>Balaenoptera edeni</i>
Common dolphin	<i>Delphinus delphis</i>
Cuvier's beaked whale	<i>Ziphius cavirostris</i>
Dwarf sperm whale	<i>Kogia simus</i>
False killer whale	<i>Pseudorca crassidens</i>
Fraser's dolphin	<i>Lagenodelphis hosei</i>
Killer whale	<i>Orcinus orca</i>
Longman's beaked whale	<i>Indopacetus pacificus</i>
Melon-headed whale	<i>Peponocephala electra</i>
Minke whale	<i>Balaenoptera acutorostrata</i>
Northern Elephant Seal	<i>Mirounga angustirostris</i>
Pantropical Spotted Dolphin	<i>Stenella attenuate</i>
Pygmy killer whale	<i>Feresa attenuata</i>
Pygmy sperm whale	<i>Kogia breviceps</i>
Risso's dolphin	<i>Grampus griseus</i>
Rough-toothed dolphin	<i>Steno bredanensis</i>
Short-finned pilot whale	<i>Globicephala macrorhynchus</i>
Spinner dolphin	<i>Stenella longirostris</i>
Spotted dolphin	<i>Stenella attenuata</i>
Striped dolphin	<i>Stenella coeruleoalba</i>

Source: Eldredge 2003

8.5.3 Seabirds¹⁷

The following seabirds are considered residents of CNMI: wedge-tailed shearwater (*Puffinus pacificus*), white-tailed tropicbird (*Phaethon lepturus*), red-tailed tropicbird (*Phaethon rubricauda*), masked booby (*Sula dactylatra*), brown booby (*Sula leucogaster*), red-footed booby (*Sula sula*), white tern (*Gygis alba*), sooty tern (*Sterna fuscata*), brown noddy (*Anous*

¹⁷ The USFWS is the primary federal agency with authority and responsibility to manage ESA listed seabirds.

stolidus), black noddy (*Anous minutus*), and the great frigatebird (*Fregata minor*). There are no known interactions between seabirds and any of the Mariana Archipelago pelagic fisheries (WPRFMC 2007b).

The following seabirds have been sighted and are considered visitors (some more common than others) to CNMI; short-tailed shearwater (*Puffinus tenuirostris*; common visitor), Newell's shearwater (*Puffinus auricularis*; rare visitor), Audubon's shearwater (*Puffinus iherminieri*), Leach's storm-petrel (*Oceanodroma leucorhoa*), and the Matsudaira's storm-petrel (*Oceanodroma matsudairae*). Of these, only the Newell's shearwater is listed as endangered. There have been no sightings of the endangered short-tailed albatross (*Phoebastria albatrus*) in CNMI although CNMI is within the range of the only breeding colony at Tora Shima, Japan (WPRFMC 2007b).

Newell's shearwater

The Newell's shearwater is listed as threatened under the ESA. The Newell's shearwater is listed, because of its small population of approximately 14,600 breeding pairs, its isolated breeding colonies, and the numerous land-based hazards affecting them at their breeding colonies (Ainley et al. 1997). The Newell's shearwater breeds only in colonies on the main Hawaiian Islands, especially Kauai, from April to October-November (Sincock and Swedberg 1969). They are present from about 18° to 25° N and from about 160° to 120° W (Ainley et al. 1997). They have been associated with the North Equatorial Counter Current (NECC) directly south of Hawaii, and from about 160° to 120° W, with occasional sighting throughout the tropical Pacific (USFWS 1983; Spear et al. 1995; Ainley et al. 1997; N. Holmes, KESRP, pers. comm. June 2009). No breeding colonies or resident population of Newell's shearwater is known to exist in the CNMI or Guam. This bird is considered a rare visitor to the CNMI and has not been observed in Guam.

Shearwaters are most active in the day and skim the ocean surface while foraging. During the breeding season, shearwaters tend to forage within 50–62 miles (80–100 km) of their nesting burrows (Harrison 1990). Shearwaters also tend to be gregarious at sea, and the Newell's shearwater is known to occasionally follow ships (Harrison 1990). Shearwaters feed by surface seizing and pursuit plunging. Often shearwaters will dip their heads under the water to sight their prey before submerging (Warham 1990).

Based on observer data from August 1994 to January 2009 from the SPC there have been no recorded interactions between U.S. purse seine vessels or foreign longline fisheries with Newell's shearwater (SPC 2010). No interactions with Newell's shearwater have been reported or observed in the Guam or CNMI longline fisheries. Since no purse seine activity is taking place in the U.S. EEZ around the Mariana Archipelago, and limited longline activity (less than 30 trips since 2007) is taking place in the EEZ, it is highly unlikely that these fisheries would interact with Newell's shearwaters.

Short-tailed albatross

Short-tailed albatross breed primarily on Torishima, Japan, and the Senkaku Islands. In the western Pacific, they generally range north of 15° N. No known breeding or resident populations of short-tailed albatross exist, and, to date, there have been no sightings of them in the CNMI or Guam.

No interactions with short-tailed albatross have been reported in Federal logbooks for Guam or the CNMI longline fisheries or in the SPC observer data from 1994 to 2009 for foreign longliners operating outside the U.S. EEZ (SPC 2010). Since purse seine activity is not taking place in the U.S. EEZ around the Mariana Archipelago and longline activity (less than 30 trips since 2007) is limited in the EEZ, it is highly unlikely that these fisheries would interact with short-tailed albatross.

According to Wiles et al. (2003), the only resident seabirds on Guam are the brown noddy and the white tern. Common visitors to Guam include the following seabirds: black noddy the short-tailed shearwater. Other less common or rare visitors include: brown and red-footed boobies, wedge-tailed shearwater, Matsudaira's storm-petrel, white-tailed and red-tailed tropicbirds, great frigatebird, gulls, and terns.

8.6 Status of Target Stocks

Bigeye Tuna

The Secretary of Commerce determined that overfishing of bigeye tuna (*Thunnus obesus*) was occurring Pacific-wide¹⁸ in 2004 (69 FR 78397). Pacific bigeye tuna are caught by a suite of domestic and foreign purse seiners and longliners, with small amounts also taken by handline and troll vessels. Until recently, the majority of the bigeye catch was taken by longliners, primarily for the Japanese sashimi market. However, during the past 10 years, catches of bigeye by purse seiners have increased considerably. This is not due to deliberate targeting of bigeye tuna by purse seiners, but as an incidental catch when purse seiners are targeting skipjack (*Katsuwonus pelamis*) and juvenile yellowfin tuna (*Thunnus albacares*) around FADs with larger and deeper purse seine nets. Stock assessments for bigeye tuna in the eastern and western Pacific, conducted in 2003 and 2004, showed that the level of fishing mortality had exceeded the fishing mortality associated with maximum sustainable yields (F_{msy}). This level of fishing mortality is one of the limit reference points of the Council's overfishing control rule for bigeye tuna and other pelagic fishes. The Pacific-wide stock itself is not yet overfished, but could become so if levels of fishing mortality are not reduced.

The status of bigeye tuna stocks in the Pacific Ocean has typically been assessed for two separate regions, the eastern Pacific Ocean (EPO) and the WCPO. That approach is based on the areas of authority and interest of the regional fishery management organizations, particularly the Inter-American Tropical Tuna Commission in the EPO and the Western and Central Pacific Fisheries Commission in the WCPO and assumed population structure of bigeye tuna in the Pacific (which is not well known). The 2010 WCPO stock assessment concluded that overfishing of bigeye tuna is occurring, the stock is approaching an overfished state, if it hasn't already reached a slightly overfished condition (Harley et al. 2010). The same authors also concluded that MSY levels

¹⁸ A stock is considered to be subject to overfishing whenever it is subjected to a rate of fishing mortality that jeopardizes its capacity to produce MSY on a continuing basis (50 CFR 600.310(d)(ii)). See Section 5.4 of this document for further information.

would increase if the mortality of small fish were reduced, which would allow greater overall yields to be sustainably harvested. NMFS makes a Pacific-wide bigeye tuna stock status determination, updated quarterly as applicable, based on assessments in both the EPO and WCPO.

The third quarter 2010 status determination lists Pacific bigeye tuna as subject to overfishing, but not overfished, nor approaching an overfished condition. (A fishery classified as approaching an overfished condition is estimated to become overfished within two years, based on fishing effort trends, fishery resource size, and other appropriate factors; Magnuson-Stevens Act 304(e)(1)) Fishing mortality for adult and juvenile bigeye tuna is estimated to have increased continuously since the beginning of industrial tuna fishing. Overall, depletion is estimated to have been rapid, particularly since the mid-1980s. While total biomass has remained relatively stable since 1970, it appears to have been sustained by above average recruitment, particularly since 1995. The assessment indicates that recruitment may have returned to the long-term average level (although recent recruitment estimates have high uncertainty) and, if recruitment remains at that level, biomass would decline rapidly at current exploitation rates. The current level of biomass is 20–26 percent of the unexploited level with higher depletion estimated from the model as longline catchability increases.

The attribution of depletion to various fisheries or groups of fisheries indicates that the Pacific-wide longline fisheries have the greatest impact throughout the model domain. The purse seine and Philippines/Indonesian domestic fisheries also have substantial impact in the western Pacific and to a lesser extent in the central Pacific; and the Japanese coastal pole-and-line and purse-seine fisheries are also having a significant impact near Japan (Langley et al. 2008).

Yellowfin Tuna

In August 2009, the Scientific Committee of the WCPFC reviewed a stock assessment for yellowfin tuna in the western and central Pacific (Langley et al. 2009). The main conclusions indicated that recent recruitment of yellowfin tuna is estimated to be considerably lower than the long-term average and fishing mortality is estimated well below the stocks fishing mortality maximum sustainable yield. The estimates of MSY for the four principal models are 552,000–637,000 mt and considerably higher than recent catches estimates for yellowfin (430,000 mt, source WCPFC Yearbook 2007). Therefore the WCPO yellowfin tuna stock is not experiencing overfishing or in an overfished state.

Yellowfin tuna are caught primarily by purse seine and ring nets in the Western and Central Pacific, and by purse seines in the Eastern Pacific Ocean. Substantial volumes of yellowfin tuna are also caught by pole-and-line fleets in Indonesia and by handliners in the Philippines. Yellowfin catches by longlines comprise a significant catch of yellowfin tuna, but longlining is a much smaller component of the fishing mortality on this species compared to bigeye tuna.

Skipjack Tuna

The major conclusions of the 2010 skipjack assessment (Hoyle et al 2010) are similar to those of the last four assessments (Hampton 2002; Langley et al. 2003b; Langley et al. 2005b; Langley and Hampton 2008).

Similar to other tropical tunas, estimates of natural mortality are strongly age-specific, with higher rates estimated for younger skipjack. Stocks of skipjack tuna in the WCPO appear to be robust at this time with increased landings and catch rates being experienced for the past few years in the WCPO. The 2008 catch of skipjack tuna in the WCPO was the second highest ever continuing a trend begun in 2002 (Figure 33)

The most recent assessment of skipjack tuna in the WCPO was conducted in 2010 (and included data from 1972 to 2009 (Hoyle et al. 2010). Current fishing mortality rates for skipjack tuna are estimated to be well below the F_{MSY} reference point, and therefore, overfishing is not occurring (i.e., current fishing mortality is less than F_{MSY}). The total biomass of skipjack tuna has fluctuated above the biomass based reference point B_{MSY} and recent biomass levels are estimated to be well above the B_{MSY} level. According to the authors, these conclusions appear relatively robust (i.e., scientifically valid), at least within the statistical uncertainty of the current assessment. Recruitment variability, influenced by environmental conditions, will continue to be the primary influence on stock size and fishery performance.

The model estimates significant seasonal movements between the western and eastern equatorial regions. The performance of the fishery in the eastern region has been shown to be strongly influenced by the prevailing environmental conditions with higher stock abundance and/or availability associated with El Niño conditions (Lehodey et al. 1997). This is likely to be at least partly attributable to an eastward displacement of the skipjack biomass due to the prevailing oceanographic conditions, although this dynamic is unlikely to be captured by the parameterisation of movement in the current model.

Recruitment showed an upward shift in the mid-1980s and is estimated to have remained at a higher level since that time. Recruitment in the eastern equatorial region is more variable with recent peaks in recruitment occurring in 1998 and 2004–2005 following strong El Niño events around that time. Conversely, the lower recruitment in 2001–2003 followed a period of sustained La Nina conditions. Recent recruitment is estimated to be at a high level, but is poorly determined due to limited observations from the fishery.

Biomass trends are driven largely by recruitment and fishing mortality. The highest biomass estimates for the model period occurred in 1998–2001 and in 2005–2007, immediately following periods of sustained high recruitment within the eastern equatorial region.

The biomass trajectory is influenced by the underlying assumptions regarding the treatment of the various fishery-specific catch and effort data sets within the model. The Japanese pole-and-line fisheries are all assumed to have constant catchability, with any temporal trend in efficiency assumed to have been accounted for by the standardization of the effort series. The CPUE trends are influential regarding the general trend in both recruitment and total biomass over the model period. In all regions there is a relatively good fit to the observed CPUE data. This indicates reasonable consistency between the CPUE series and the other sources of data, especially the size data, within the assessment model. The standardized CPUE indices appear to represent a substantially more consistent index of stock abundance than the indices used in previous years.

The model also incorporates a considerable amount of tagging data that provides information concerning absolute stock size during the main tag recovery period. For the equatorial regions,

the most recent data included in the model are from an intensive tagging programme that ceased in the early 1990s with most tag recoveries occurring over the following 18 months. Further analyses should be carried out to integrate the PTTP data into the stock assessment as soon as possible. Initial analyses of the data suggest results consistent with evidence from the CPUE time series. However, integrating the PTTP data into the model is likely to improve the accuracy and precision of estimates, particularly in the eastern equatorial region.

Within the equatorial region, fishing mortality increased throughout the model period and is estimated to be highest in the western region in the most recent years. The impact of fishing is predicted to have reduced recent biomass by about 50% in the western equatorial region and 25% in the eastern region. For the entire stock, the depletion is estimated to be approximately 40%.

The principal conclusions are that skipjack is currently exploited at a moderate level relative to its biological potential. Furthermore, the estimates of and indicate that overfishing of skipjack is not occurring in the WCPO, nor is the stock in an overfished state. These conclusions appear relatively robust, at least within the statistical uncertainty of the current assessment. Fishing pressure and recruitment variability, influenced by environmental conditions, will continue to be the primary influences on stock size and fishery performance.




Figure 32: Skipjack Catch by Gear (mt) in WCPO

Source: Williams and Terawasi 2008

8.7 Regional Tuna Fisheries Management

Management of pelagic fisheries is complicated by the nature of the targeted “highly migratory species” whose life histories span the arbitrary jurisdictional boundaries used in management, such as EEZs, of many nations. U.S. pelagic fisheries do much of their fishing on the high seas outside the EEZ of the U.S. and other nations. Management includes trans-boundary issues, both because fish stocks may straddle EEZs and because fishing may occur in international waters

where no nation has comprehensive jurisdiction. For this reason, fishery managers and their governments have sought to establish stable multilateral arrangements, but establishing such a regime is always difficult for “open access” resources. In Ward et al. (1998) a review of regional organizations established to manage or research living marine resources, 25 organizations were identified. The following cover the Pacific region and target species relevant to Council/NMFS managed pelagic fisheries:

The Western and Central Pacific Fisheries Commission (WCPFC). The WCPFC was established by the coastal States of the Western and Central Pacific and States fishing in that region through the Convention on the Conservation and Management of Highly Migratory Fish Stocks in the Western and Central Pacific Ocean. The Convention was opened for signature at Honolulu on September 5, 2000. The objective of the Convention is to ensure, through effective management, the long-term conservation and sustainable use of highly migratory fish stocks in the WCPO in accordance with the 1982 United Nations Convention on the Law of the Sea and the 1995 U.N. Fish Stocks Agreement. For this purpose, the Convention establishes a Commission for the Conservation and Management of Highly Migratory Fish Stocks in the Western and Central Pacific Ocean.

The Convention was open for signature for 12 months from September 5, 2000, by the States that participated in the Multilateral High-Level Conference on the Conservation and Management of Highly Migratory Fish Stocks in the Western and Central Pacific. These nations include: Australia, Canada, China, the Cook Islands, the Federated States of Micronesia, the Fiji Islands, France, Indonesia, Japan, the Republic of Kiribati, the Republic of the Marshall Islands, the Republic of Nauru, New Zealand, Niue, the Republic of Palau, the Independent State of Papua New Guinea, the Republic of the Philippines, the Republic of Korea, the Independent State of Samoa, the Solomon Islands, the Kingdom of Tonga, Tuvalu, the United Kingdom of Great Britain and Northern Ireland in respect of Pitcairn, Henderson, Ducie and Oeno Islands, the United States of America and the Republic of Vanuatu. The depositary for the Convention is the Government of New Zealand.

The WCPFC implemented its first conservation measure for WCPO bigeye tuna in December 2005, when the Commission promulgated its first resolution for tuna management (Conservation and Management Measures for Bigeye and Yellowfin Tuna in the Western and Central Pacific Ocean). Under this measure, members of the Commission and cooperating non-members, agreed to establish catch limits to reduce fishing mortality on bigeye for the years (2006-2008) to the average annual bigeye catch for the years 2001-2004 or the year 2004 (the year 2004 only applies to the U.S. and China). This measure did not apply to members or cooperating non-members that caught less than 2,000 mt in 2004; those that caught less than 2,000 mt of bigeye in 2004 was required to ensure that their catch did not exceed 2,000 mt in each of the next three years. This measure was maintained in 2006 at the third meeting of the WCPFC under Conservation and Management Measure 2006-01. In December 2008, the WCPFC adopted CMM 2008-01 which limited the Hawaii longline fishery to its 2004 bigeye tuna total catch (4,159 mt) and allocated a catch quota of 2,000 mt each to the U.S. territories of American Samoa, Guam and the Northern Mariana Islands.

The Inter-American Tropical Tuna Commission (IATTC). The IATTC, established by international convention in 1950, is responsible for the conservation and management of

fisheries for tunas and other species taken by tuna-fishing vessels in the Eastern Pacific Ocean (generally speaking, off the coast of Latin America west to 150° W. longitude). It collects data, assesses stock status, provides management advice to member nations, operates an observer program and conducts fishery research. In addition to target species issues, incidental dolphin mortality has been a major problem in the Eastern Tropical Pacific (ETP) and the IATTC has established a program to monitor incidental mortality and make conservation recommendations. The member countries of the IATTC are: Costa Rica, Ecuador, El Salvador, France, Guatemala, Japan, Mexico, Nicaragua, Panama, Peru, Republic of Korea, Spain, United States, Vanuatu, and Venezuela. Belize, Canada, China, Cook Islands, the European Union, Honduras and Chinese Taipei are Cooperating Non Parties or Cooperating Fishing Entities.

The South Pacific Tuna Treaty (SPTT). The SPTT entered into force in 1988 and after an initial 5-year agreement was extended in 1993 and again in March 2002. In 2002, the Parties agreed to amend and extend the Treaty and to extend the related Economic Assistance Agreement between the United States and the Forum Fisheries Agency (FFA) beyond the June 2003 expiration date, for a term of 10 years. The 2002 extension provides licenses for up to 40 U.S. purse seiners, with an option for 5 additional licenses reserved for joint venture arrangements, to fish for tuna in the EEZs of the Pacific Island Parties. It also contains a number of amendments to the Treaty and its annexes, such as updating the methods available for reporting; a revised procedure for amending the annexes; a revised observer program fee formula; provisions on the use of a vessel monitoring system (VMS); and general provisions on fishing capacity, revenue sharing, and linkages between the Treaty and the Western and Central Pacific Tuna Convention (WCPTC), among others. The SPTT agreement expires on June 14, 2013. The parties involved in the SPTT include: Australia, Cook Islands, Federated States of Micronesia, Fiji, Kiribati, Marshall Islands, Nauru, New Zealand, Niue, Palau, Papua New Guinea, Samoa, Solomon Islands, Tonga, Tuvalu, United States and Vanuatu.

The SPTT's implementing legislation is the South Pacific Tuna Act of 1988 as amended (54 FR 4033, January 27, 1989; 56 FR 19312, April 26, 1991). Of the total cost for access under the SPTT, the U.S. tuna industry, as coordinated by the American Tunaboat Association, provides up to \$3 million each year to the Forum Fisheries Agency (FFA) located in Honiara, Solomon Islands. The FFA Director and staff act as the SPTT Administrators for the Pacific Island Parties to the agreement. The FFA deducts a small amount (approx. \$500,000) for treaty administration, after which 15 percent of the revenue is divided equally among FFA members, with the remaining balance (85 percent) distributed on a *pro rata* basis depending on the weight of tuna landed in each respective EEZ.

Also associated with the SPTT is an Economic Assistance Agreement between the U.S. Government (U.S. Agency for International Development) and the FFA. The U.S. Government pays \$18 million annually, subject to the availability of appropriated funds for this purpose, into an economic development fund administered by the FFA. The FFA ensures that the fund is used to support economic development programs in the region. Payments to the Pacific Island Countries under the Economic Assistance Agreement are now the only significant source of U.S. economic support for the stability and security of the region outside the assistance provided to the Freely Associated States. Under the terms of the SPTT, both the U.S. tuna industry and the U.S. Government annual payments must be made before any fishing licenses will be issued. In

addition to paying access fees, the U.S. tuna industry also pays the FFA costs associated with observer coverage (including training), VMS deployment and associated recurring costs, and a regional registration fee.

U.S. operational, administrative, and enforcement commitments under the SPTT are carried out by NOAA, specifically its regional offices located in Honolulu, Hawaii. Issues that arise typically are addressed in formal annual consultations between U.S. Government and Pacific Island States representatives, or during informal discussions which also have taken place on an annual basis. The Department of State has specific authority to act for the United States.

The Oceanic Fisheries Program (OFP). The OFP is a component of the Secretariat of the Pacific Community (SPC), based in Noumea, New Caledonia. All Pacific island nations and several metropolitan states are members. American Samoa, CNMI and Guam have member status separate from U.S. membership. The OFP developed from SPC tuna assessment programs that began in the late 1970s. The program gathers fishery statistics and conducts research, including stock assessments for major Western Pacific tuna stocks. It advises member states by providing detailed scientific information on stock status.

The International Scientific Committee (ISC). The ISC was established in 1995 through an intergovernmental agreement between Japan and the United States of America (USA). Since its establishment and first meeting in 1996, the ISC has undergone a number of changes to its charter and name (from the Interim Scientific Committee to the International Scientific Committee) and has adopted guidelines for its operations. The two main goals of the ISC are to 1) to enhance scientific research and cooperation for conservation and rational utilization of the species of tuna and tuna-like fishes which inhabit the North Pacific Ocean during a part or all of their life cycle; and 2) to establish the scientific groundwork, if at some point in the future, it is decided to create a multilateral regime for the conservation and rational utilization of these species in this region. The ISC provides scientific advice on the stocks and fisheries of tuna and tuna-like species in the North Pacific to the Member governments and regional fisheries management organizations. (ISC Report, July 2008)

The Forum Fisheries Agency (FFA). Ten Pacific island countries founded the FFA in 1979 under the auspices of the South Pacific Islands Forum (now known as the Pacific Islands Forum). It has subsequently expanded to include 16 countries in the western and central Pacific. It is governed by the Forum Fisheries Committee with representation from each member country; they direct a Secretariat based in Honiara, Solomon Islands. The FFA assists member countries to conserve and manage their tuna stocks, for example in its early days by advising emerging Pacific islands on the establishment of their EEZs. With Secretariat assistance, member countries negotiated the United States Multilateral Fisheries Treaty, also known as the South Pacific Tuna Treaty discussed above, which was finalized in 1987. This treaty establishes a license regime for U.S. purse seine vessels that fish in member countries' EEZs. More recently the FFA has emphasized sustainable management of tuna resources, particularly through multi-lateral arrangements, and securing greater economic benefits from tuna resources for member nations.

8.8 CNMI Longline Fishing

There is currently a 50 nautical mile prohibited area for longline fishing around Guam implemented in 1993 following the expansion of the Hawaii longline fishery. However no comparable prohibited area exists in waters around nearby CNMI and there is growing interest in longline fishing in EEZ waters around the Mariana Archipelago. During the first nine months of 2007, NMFS issued eight Western Pacific General Longline permits including two in CNMI. These general longline permits may be used to fish in EEZ waters around CNMI where there is far less competition than in the Hawaii-based fishery.

As of 2010, there are four permitted longline vessels in CNMI, however, due to the limited number vessels in the fishery over the past decade, landings data are scarce or unable to be reported here due to data confidentiality issues¹⁹. Due to the lack of catch and effort data obtained from longline vessels around CNMI, the Council developed a proxy model using targeted catch, interactions with protected species and fish bycatch data from the nearby Federated States of Micronesia (FSM) fishery. Like FSM, longline vessels that operate around CNMI target yellowfin and bigeye, along with a variety of PMUS including albacore, yellowfin, blue marlin, mahimahi, skipjack, bigeye, and spearfish. The waters around FSM and the Mariana Archipelago are all within the equatorial band and have similar oceanic climates and oceanographic features (Bigelow et al. 1999). Vessels used in FSM are comparable in size to what would be expected in Mariana Archipelago longline operations and both would be expected to target bigeye and yellowfin. Catch data from longline vessels operating in FSM (Table 15) were therefore used to establish this proxy. These vessels are larger than 50 ft and are crewed by local Micronesians. They operate in oceanic habitats similar to the waters around CNMI and make deep sets to maximize catches of bigeye and yellowfin tuna, as would be anticipated of longline vessels around CNMI. Approximately 86 percent of FSM catch is bigeye and yellowfin, with bigeye tuna comprising almost two thirds of the catch

The catch rates for the FSM longliners were combined with typical American Samoa longline vessel operating patterns (Table 16) to generate projections of potential annual fish catches for a single longline vessel fishing within EEZ waters around CNMI (Table 17).

Longline fishing catches taken near the Mariana Islands (5-25°N, 140-150°E) recorded by OFP for fishing by U.S. vessels (21 sets; 45,038 hooks) showed the highest CPUE for bigeye followed by yellowfin tuna (Table 4 in OFP 1998). They had high catches of sharks, lancetfish, pomfrets, and other pelagic species; however, if we only look at the tuna and tuna-like species catch then 62 percent were bigeye, 27 percent yellowfin and 6 percent wahoo. This is similar to the FSM catch composition shown in Figure 34.

¹⁹ The Saipan-based longline fishing company, USA Islands Seafood, recently authorized the release by WPacFIN of the top ten species retained by the company's fishing vessels between 2007-2010, which were as follows: Yellowfin Tuna 30%, Albacore Tuna 22%, Mahimahi 17%, Bigeye Tuna 8%, Oilfish 6%, Skipjack Tuna 4%, Pomfret, 4%, Wahoo 2%, Blue Marlin 1%, Spearfish 1%. Clearly there is difference in the species composition of the CNMI longliners versus the vessels operating in the FSM used as a proxy in the amendment. However, in both cases tunas comprise the majority of the catch (64% in Saipan versus 86% in the FSM) but yellowfin and albacore forming just over 50% of the Saipan-based longliners versus 64% bigeye in the FSM. Without further information it is not possible to determine whether these differences are due to location of fishing, differences in gears and longline deployment or a combination of these factors.




Figure 33: Catch Composition of Longline Vessels Operating in the FSM.
Source: WCPFC 2005

Table 15: Catch Rates for Longliners Operating in the FSM (lbs/hook)

Year	Bigeye	Yellowfin	Black marlin	Blue marlin	Other marlins	Swordfish	Others	Total
2001	0.29	0.15	0.03	0.00	0.00	0.01	0.01	0.49
2002	0.28	0.06	0.02	0.00	0.00	0.01	0.00	0.36
2003	0.28	0.12	0.02	0.00	0.00	0.00	0.04	0.46
2004	0.28	0.10	0.00	0.03	0.00	0.00	0.03	0.44
Average	0.28	0.11	0.02	0.01	0.00	0.01	0.02	0.44

Source: WCPFC 2006

Table 16: Typical American Samoa Tuna Longline Vessel Operations

Effort measure	Numbers
Annual number of trips	10
Number of sets/trip	20
Hooks/set	2,200
Annual number of hooks	440,000

Source: WPRFMC 2007

Table 17: Projected Annual Landings for One CNMI Longline Vessel

Species	Projected Annual Landings (lb)
Bigeye	124,085
Yellowfin	46,658
Black marlin	7,568
Blue marlin	3,752
Other marlins	83
Swordfish	2,965
Others	8,046
Total	193,157

8.9 Climate Change Background

A memorandum dated February 18, 2010, issued by the Chair, Council on Environmental Quality (Sutley 2010) provides draft guidance to Federal departments and agencies on ways to improve consideration of effects of greenhouse gas (GHG) emissions and climate change in their evaluation of proposals under the National Environmental Policy Act (NEPA). Effects to be considered include the potential effects of agency action on global climate change and potential effects of climate change on a proposed project. This section provides a general discussion of climate change concepts as it may be tuna stocks, protected resources, and fisheries considered in this proposed action. Climate change considerations for the proposed action and alternatives are addressed in the impacts section, section 9, below.

The global mean temperature has increased by 0.76°C over the last 150 years, and the linear trend of temperature over the last 50 years is nearly twice that for the last 100 years (IPCC 2007a). Ample evidence now exists supporting the wide-ranging and variable impacts from climate change which can affect the phenology (timing of recurring natural phenomena such as life stages), range and distribution of species, and composition and dynamics of communities (Walther et al. 2002). Observed changes in marine systems associated with climate change include increasing water temperatures; sea level rise; changes in ice cover, salinity, oxygen levels, circulation; modification of shorelines; and increased ocean acidification. Changes to marine systems include shifts in ranges; changes in algal, plankton, and fish abundance (IPCC 2007b); damage to coral reefs (Scavia et al. 2002), prey abundance, and other impacts. A more complete summary of climate change and climate change impacts can be found online at http://www.ipcc.ch/publications_and_data/publications_and_data_reports.shtml.

In general, large scale climate cycles can impact winds, currents, ocean mixing, temperature regimes, nutrient recharge, and affect the productivity of all trophic levels in the North Pacific Ocean (Polovina et al. 1994). These impacts are expressed as variability in stock size, recruitment, growth rates, or other factors. Highly migratory pelagic fishes can be affected through changing distribution and abundance of prey species. Bleaching and the effects of ocean acidification on coral reef ecosystems are other climate change-related concerns that can affect food webs of tuna prey species.

Climate change may impact the availability of tunas and this may, in turn, impact all pelagic fisheries; however, how and to what extent is not known. Distinguishing the effects of climate change from natural variability is challenging. NMFS and other federal agencies will continue to conduct prioritized research in order to facilitate ocean resources stewardship. Osgood (2008) provides a review of marine ecosystem climate research priorities and concerns.

Climate change has the potential to impact protected resources that interact with fisheries including marine mammals, seabirds, and sea turtles. Impacts can include changes in prey abundance and distribution, and impacts of sea level rise and temperature changes on nesting seabirds, sea turtles and monk seal pupping sites. These changes will be monitored by NMFS, the Council, local governments and USFWS and information can be used to modify fishery operations, as needed.

Although direct information may not be available, the potential impacts of climate change on pelagic fisheries in the Mariana Archipelagic EEZ will continue to be noted indirectly through ongoing monitoring of fish stocks as well as fish harvests and monitoring of protected resources that interact with the pelagic fisheries. Climate change impacts on listed species and their habitats, are considered in section 7 consultations undertaken on the fisheries of the western Pacific region. The Council and NMFS will continue to implement responsive fishery management measures as needed to manage western Pacific fisheries sustainably.

9.0 Impacts of the Alternatives

9.1 Topic 1: CNMI Longline Prohibited Area

Alternative 1A: No Action

Under Alternative 1A (No Action), no new regulations would be made no vessels would be prohibited from longline fishing within the EEZ waters around CNMI.

Alternative 1B: 25nm Longline Prohibited Area

Under Alternative 1B all vessels would be prohibited from pelagic longline fishing within 25 nm of CNMI.

Alternative 1C: 30 nm Longline Prohibited Area (Preferred)

Under Alternative 1C all vessels would be prohibited from pelagic longline fishing within 30 nm of CNMI.

Alternative 1D: 50 nm Large Vessel Longline Prohibited Area

Under Alternative 1D vessels greater than 50 ft in length would be prohibited from pelagic longline fishing within 50 nm of CNMI.

Alternative 1E: 75 nm Longline Prohibited Area

Under Alternative 1E all vessels would be prohibited from pelagic longline fishing within 75 nm of CNMI.

Alternative 1F: 100 nm Longline Prohibited Area

Under Alternative 1F, all vessels would be prohibited from pelagic longline fishing within 100 nm of CNMI.

9.1.1 Impacts on Target Stocks

Under Alternative 1A (No Action) pelagic landings and fishing effort in CNMI would continue to fluctuate based on weather, catchability, fuel prices, and other factors. Catches of pelagic species by the troll and longline fleets would be expected to be small in comparison to the larger WCPO pelagic fishery, which in 2009 caught a total of 2,467,903 mt, the highest annual catch recorded. During 2009, the purse seine fishery accounted for an estimated 1,894,500 mt (77% of the total catch, and another record for this fishery) and the the longline fishery caught an estimated 223,792 mt (9%) (Williams & Terawasi 2009). The 2008 troll catch for the CNMI was

197,013 lb (Table 3) or about 89 mt. Based on Table 17, the four-vessel CNMI longline fleet is estimated to have an annual catch of 772,628 lb or about 350 mt, which would likely be sustainable given that there are no substantial pelagic fisheries within the U.S. EEZs around the Mariana Archipelago.

Regardless of which alternative is selected, development of the longline fishery could continue in the CNMI with a fleet of about four longline vessels. These vessels would continue to target bigeye tuna, and based on Table 17, the four vessel longline fleet would be expected to catch annually about 224 mt of bigeye. Under current regulations WCPO longline harvests of bigeye are currently managed by international quotas, limiting longline catches in CNMI to a maximum of 2,000 mt of bigeye per year. The projected level of catch of BET by the 4 longline vessels is just over 10% of the catch limit, with little to no bigeye landed by troll vessels. None of the alternatives is likely to affect the number of longline vessels, or increase longline fishing intensity; therefore, none of the alternatives is expected to have a large or adverse effect on BET stocks.

The second highest catch by longline fishing would most likely be yellowfin tuna which is currently a significant component of catches made by CNMI trolling vessels. Based on Table 17, the four vessel longline fleet would be expected to catch annually about 84 mt of yellowfin, while troll catches in 2008 amounted 16,344 lbs or 7.3 mt. Based on projections related to longline fishing near FSM, it is estimated that in one year the longline fleet may catch over ten times the volume of the commercial troll fleet. The decision by the Council to separate the two fleets through the adoption of a longline prohibited area is a way to minimize catch competition between the two fleets.

Under current regulations WCPO longline catch of yellowfin tuna is not to be increased in the longline fishery from the 2001-2004 levels (WCPFC 2008). However, this provision does not apply to members and participating territories that caught less than 2,000 mt in 2004.

All of the action alternatives would reduce the potential for localized stock depletion and catch competition except for alternative 1D which would allow longline vessels less than 50 ft in length to fish in the 50 nm prohibited area. If this were to occur, as is true with the no-action alternative, there could be some localized stock depletion of target stocks (bigeye and yellowfin) depending on the level of fishing and location of longline effort relative to troll fishing. The localized stock depletion would not result in target species overfishing as the anticipated catches for all alternatives is expected to be within (under) the quota limits.

The proposed shift in longline fishing areas under the remaining action alternatives is not expected to change either fishing intensity or catch rates. Therefore, there would be no change in impacts to sustainability of target species as a result of approving and implementing any of the proposed longline prohibited area alternatives in the CNMI.

9.1.2 Impacts on Non-target Stocks

Under Alternative 1A (No Action) fishing impacts on non-target stocks by the existing troll fleet would be expected to continue to be quite small based on historic catch data (Table 1).

However, additional longline vessels fishing in the EEZ under Alternative 1A, 1B, 1C, 1D, 1E, or 1F could each have the potential to catch and retain up to 22,000 pounds of non-target species annually (Table 17). For the four existing longline vessels, it is estimated that up to 88,000 pounds of non-target species are caught annually, and this level is sustainable. None of the non-target fish stocks that may be caught by the CNMI longliners is overfished or in an overfishing conditions, with the possible exception of North Pacific striped marlin (Brodziak and Piner 2010). However catches of marlins other than blue marlin would likely not exceed 83 lb/vessel or 332 lb for the fleet.

Little information is available on the bycatch (discards) associated with tropical deep set longline fishing in the WCPO. Molony (2005) analyzed longline observer data from across the entire western Pacific, for both shallow and deep set longline fishing, and identified 53 species and species groups in the limited observer data. Of these, about 30 percent were identified as likely to be discards and of these, about 80 percent were sharks, with blue sharks comprising more than half. Based on the information in Table 16 and in Molony (2005)²⁰ one longline vessel setting 440,000 hooks per year would be projected to catch 479 sharks, or just under 2000 sharks for a 4-vessel longline fleet. It is assumed that most sharks would be released alive, given the U.S. ban on finning and the low value for shark meat in CNMI. In this case impacts to non-target species (both retained and discarded) by one longline vessel fishing in EEZ waters around CNMI would be expected to be minimal.

The impacts of longline fishing on populations of non-target species in CNMI are largely unknown, however it could be expected that some localized stock depletion may be possible given any increase in fishing effort, particularly in high use areas.

Regardless of which action alternative is selected, there would be no change to the level of bycatch and fish discards that is currently occurring under the status quo. Therefore, the proposed establishment of longline closed areas under alternatives 1B, 1C, 1D, 1E, and 1F would not result in large or adverse effects to non-target fish stocks and the impact of troll and longline fishing on non-target species would continue to be sustainable. As with current management under the no-action alternative, the Council, NMFS, and the CNMI government would continue to monitor the status of non-target stocks and impacts of domestic fisheries on these resources and would take appropriate management actions if additional conservation measures were found to be needed.

9.1.3 Impacts on Protected Species

Under Alternative 1A (No Action), pelagic fishing around CNMI would continue to develop with concomitant impacts on protected species expected to continue to be nonexistent to minimal. The troll fishery around CNMI was included in NMFS' 2002 and 2009 biological opinions (BiOp) on pelagic fisheries of the Western Pacific Region.

²⁰ Mean observed tropical deep set longline shark catch rates in Molony (2006) \approx 1.09 sharks per 1000 hooks

The 2009 BiOp considered potential stressors and impacts related to vessel collisions, hooking and entanglement with fishing gear, human disturbance, exposure to vessel waste, direct and indirect competition with foraging grounds and exposure to marine mammal deterrents. Based on the 2009 BiOp, NMFS determined that troll and handline fisheries in the CNMI are not likely to adversely affect ESA listed species. A small potential exists that trolling gear may incidentally hook or entangle a sea turtle or marine mammal, however, NMFS and the Council consider this type of interaction extremely rare and therefore unlikely to adversely affect listed species.

Regardless of which alternative for a longline prohibited area in the CNMI is selected, existing longline fishing regulations would continue to apply for all longline vessels. All longline vessels would be required to adhere to all seabird, sea turtle, and other protected species mitigation and avoidance measures, including carrying an observer if requested to do so by NMFS, currently in effect for deep set longline fishing activities by general longline permit holders.

None of the proposed action alternatives that would establish a longline prohibited area in the CNMI are expected to change the conduct or intensity of fishing by longline vessels as compared to the current levels. Moving longline fishing out further from the coast under the various alternatives is not expected to increase the likelihood of interactions with marine mammals or sea turtles. Under existing regulations, interactions with protected species are required to be reported on Federal logbooks and to date there have been no reported interactions with marine mammals, seabirds or turtles in CNMI or Guam.

Seabird interactions with longline vessels are minimal in tropical latitudes (Molony 2005). Interactions with seabirds are not expected to change as a result of establishing longline fishing prohibited areas. Therefore, large adverse impacts to seabirds would not be expected under any of the alternatives considered here.

9.1.4 Impacts on Marine Habitat

The area affected by this action includes areas that have been identified as EFH for pelagic management unit species managed in the western Pacific. There are currently no known adverse effects of the troll fishery or the longline fishery on EFH or habitat areas of particular concern. None of the proposed alternatives is expected to change the intensity of fishing and therefore, none of the alternatives is likely to lead to substantial physical, chemical, or biological alterations to the habitat. None of the alternatives would be expected to significantly alter fishing operations, and therefore, there is no increased likelihood of gear loss in either the troll or longline fisheries for any of the alternatives.

9.1.5 Impacts on Public Health and Safety

There are no known impacts on public health or safety that are associated with either the troll or longline fisheries in the CNMI. If the longline fishery continues developing in CNMI establishing a longline fishing prohibited area under Alternatives 1B, 1C, 1D, 1E, or 1F would potentially have positive impacts on the health and safety of the small boat fishery participants, as compared to the no-action alternative, by minimizing gear or catch conflicts with longline

vessels when fishing inside the prohibited area. However, if longline fishing were to occur within small boat pelagic fishing areas under the no-action alternative, this could decrease catches (perceived or actual) and potentially cause the small boats to fish farther from land and in unfamiliar waters resulting in a decrease in safety for those participants.

Under all other alternatives, the larger the prohibited area, the greater the positive impacts would be to the troll fishing sector. Under Alternatives 1B, 1C, 1D, 1E, or 1F, longline vessels would be required to fish in waters farther from shore, however, longline fishing in general is usually conducted farther from shore than small boat fisheries and vessels are better equipped to safely fish offshore waters. Therefore, none of the action alternatives is likely to result in large adverse effects to public health or safety at sea.

9.1.6 Impacts on Fishery Participants and Fishing Communities

Under Alternative 1A (No Action), CNMI pelagic fishery participants would continue to be affected by factors such as weather, catchability, fuel prices, market prices and other fishery dependent and independent factors. Under the no-action alternative, a longline fishery could continue develop and expand in the CNMI. If the longline fishery is allowed to fish in nearshore waters, the troll fishing community could be directly impacted by localized stock depletion, catch competition and potential gear interactions between the two fisheries. Although the majority of the longline catch is projected to be bigeye tuna, which is not targeted by CNMI's trolling vessels, the second highest longline catch is projected to be yellowfin tuna which is the trolling fleet's second most important species. Based on the projections made here, one longline vessel could have an annual catch of yellowfin tuna (approximately 47,000 lb.) that is about double the entire average commercial troll fleet (approximately 22,000 lb, Table 3). In addition, a greater number of longline vessels could potentially cause the troll fleet to absorb an increase in costs associated with catching target species, and could expect lower prices due to an increase of market supply over what has been historically been afforded with species such as yellowfin tuna.

All of the proposed alternatives would result in a separation of the two fleets in the waters around the CNMI. None of the action alternatives would prevent a longline fishery from developing in the CNMI. Under Alternatives 1C, 1D, 1E or 1F, the troll fishing community would likely be less impacted by catch competition and potential gear interactions between the two fisheries because the majority of small boats fish within 30 nm of shore. Potential impacts from longline vessels on the troll fleet would vary depending on the size of the longline prohibited area (i.e., the larger the zone the more that impacts (i.e., catch competition and localized depletion would be reduced). Under Alternative 1B, the 25 nm prohibited area would not include some of the reefs and banks where the fleet typically fishes and, therefore, the troll fleet participants would continue to be adversely impacted by longline fishing in these areas.

Under Alternative 1D, a 50 nm prohibited area would be established prohibiting longline fishing by vessels greater than 50 ft in length. Therefore, longline fishing vessels less than 50 ft length would be able to fish within nearshore waters and without direct competition from large longline vessels, which would benefit those participants. This scenario would, however, be likely to

impact the small vessel troll fleet through catch competition and gear conflict in nearshore waters (i.e. within 30 nm of shore) where the troll fleet fishes.

Under all alternatives, the impacts associated with the development of a longline fleet in and around CNMI largely depend on the extent to which each vessel would harbor, fuel, land, transship and/or market their catches in CNMI. Benefits to the community may be accrued through development of a longline fleet in CNMI including the creation of jobs, boat and land based, as well as the establishment of a variety of markets for longline caught species that are not currently realized or available. The effect on prices from a large supply of longline caught fish to local markets would need to be considered, particularly in light of the current military buildup and the projected increase in workers and operations in the region. For example, an increase in supply without an increase in demand would result in lower prices at the market; allowing local people the opportunity to better afford PMUS, yet negatively affecting the ex-vessel price fishermen would expect when selling their catch locally.

9.1.7 Impacts on Biodiversity and Ecosystem Functions

Under the no action alternative (1A), extraction of fishery resources would continue to occur as the CNMI troll and longline fleets fish around the CNMI. These fisheries are managed with the best available scientific information and there have been no known issues related to large and adverse impacts to the biodiversity or ecosystem functions.

None of the proposed action alternatives that would establish longline prohibited areas around the CNMI are expected to change the intensity of fishing and therefore, there are no expected large or adverse impacts to biodiversity and ecosystem functions that would result from implementing any of the action alternatives.

9.1.8 Impacts on Administration and Enforcement

In general, administrative and enforcement activities that are required in managing the current CNMI troll and longline fisheries include creel censuses of troll and longline fishing, tracking of sales receipts, law enforcement by the U.S. Coast Guard and the NMFS Office of Law Enforcement (OLE) and local CNMI officials, issuance of Federal permits for longline fishing, conduct of fishery workshops, monitoring of Federal logbooks, and informing all fishery participants through a variety of public announcements, forums and workshops.

Alternatives 1B, 1C, 1D, 1E and 1F would all place an additional burden on law enforcement, U.S. Coast Guard and NOAA Office of Law Enforcement (OLE), in enforcing the respective prohibited areas. Increasing the use of vessel monitoring systems (VMS), patrol boats, aerial monitoring support and local enforcement would all add to the impacts of enforcement in CNMI. Currently, vessels operating under a general longline permit for the Western Pacific are not required to carry a VMS; however vessels that fish on the high seas must have a WCPFC fishery endorsement and are required to carry VMS. Elsewhere in the region, i.e. in Hawaii and American Samoa, longline spatial management zone compliance is monitored through VMS. The USCG conducts patrols of the U.S. EEZ around Guam and CNMI with sea-going and fixed wing airplane assets based on Guam. As with longline fleets elsewhere in the Western Pacific,

the USCG would make boardings of longline vessels at sea and at dockside to ensure compliance with various regulations and statutes. All of these activities will incur additional enforcement costs. The amount of expenditure required, would depend on the need for additional enforcement effort.

9.1.9 Cumulative Effects of the Proposed CNMI Longline Prohibited Areas

The proposed CNMI longline prohibited area would separate longliners from areas that are preferentially fished by small trollers to reduce gear conflicts and address concerns with localized depletion and catch competition between the two fleets.

None of the proposed alternatives is likely to change the conduct of the fishery or the volume of fishing by either trollers or longliners. The low level of environmental impacts of the CNMI longline prohibited area alternatives on target species, non-target species, bycatch, protected resources, and other features of the affected environment would not be increased when the proposed action is considered together with other fishery management activities that are being considered that may affect the same fisheries or area.

Establishing purse seine closed areas in the CNMI and Guam (Topics 2 and 3) would not in themselves change fishing activity by trollers or longliners in the CNMI or Guam. However, it is important to frame this conclusion in light of other measures that may influence these fisheries in future.

The 1996 reauthorization of the Magnuson Stevens Act requires the Regional Fishery Management Councils to manage fisheries under federal jurisdiction through annual catch limits (ACLs). Certain exceptions to ACLs may be applied to species with a one year life span or if they are being managed under an international fishery convention.

The target species of both the troll and longline fisheries are pelagic highly migratory fish, which fall under the purview of the Western and Central Pacific Fishery Commission. The Council at its 149th meeting in October 2010 recommended applying the international management exception to all pelagic management unit species (PMUS), except squid, for which the one year life span would be applied. If the Council's ACL amendment is approved by the Secretary of Commerce, the Council will not be required to establish ACLs for pelagic fish caught in federal waters around the CNMI. Any limitation on catches would stem from Conservation and Management Measures implemented by the Western and Central Pacific Fisheries Commission, such as CMM 2008-01, which limited catches of bigeye and yellowfin tuna for the period 2009-2011.

Currently, the CNMI may have a catch limit of 2000 mt of bigeye and a 2000 mt limit of yellowfin tuna or no limits if pursuing responsible fishery development²¹. As noted in Section

²¹ See WCPFC CMM 2008-01. The language of the Conservation and Management Measure is somewhat ambiguous about whether the limit for yellowfin is the average of the 2001-2004 levels or 2,000mt but in case the longline catch limits for bigeye and yellowfin contained in CMM 2008-01 "shall not apply to small island developing State members and participating territories in the Convention Area undertaking responsible development of their domestic fisheries."

9.1.1 catches of these species are expected to be significantly lower than these catch limits, i.e. 224 mt of longline caught bigeye and a combined troll and longline catch of about 91 mt of yellowfin tuna. The WCPFC limits apply until 2011 and new conservation and management measures in the future may apply further limitations to longline fisheries, depending on stock status of bigeye and yellowfin tuna.

It should be noted, however, that further longline catch limit implementation may also be an opportunity for the CNMI fishery, since limits on bigeye, yellowfin and potentially other species like striped marlin may have greater impacts to the U.S. Hawaii-based longline fishery, which operates under WCPFC catch limits for bigeye and yellowfin tuna in the WCPO and an IATTC EPO bigeye limit (for vessel > 24 m). Hawaii's domestic fisheries supply only about only 20% of the total demand for seafood in the state, especially fresh tuna (WPRFMC in prep)²². Exports of fresh and frozen tuna have increased sharply over the past decade (coming from countries as far away as the Marshall Islands, Federated States of Micronesia, Philippines and Indonesia (WPRFMC in prep). Given the excellent airline connections between Guam and Hawaii, there may be opportunities for the CNMI fleet to supply the Hawaii seafood market.

As is discussed in more detail in Section 9.2.6, the Parties to the Nauru Agreement or PNA Group is taking a more aggressive approach to fishery management in their respective EEZ waters, which includes implementation of a vessel day scheme (VDS) for longline fisheries²³. As with purse seine fisheries a longline VDS is a scheme where vessel owners would purchase and trade days fishing at sea in places subject to the Parties to the Nauru Agreement (PNA). Whether or not the WCPFC will adopts a VDS for all longline vessels operating in the WCPO remains to be seen, but if this were to occur, it may affect the operations of the longline fleet operating in the U.S. EEZ around CNMI.

Other pelagic fishery management amendment that are being developed by the Council (e.g., establishing purse seine closed areas in American Samoa and managing FAD use by purse seiners in the U.S. EEZ) would likely not affect the activity of fishing by trollers or longliners in the CNMI or Guam.

9.1.9 Environmental Justice Effects of the Proposed CNMI Longline Prohibited Areas

Environmental Justice policies intend to promote environmental protection by focusing attention on potential environmental harms and risks that may disproportionately and adversely affect minority, low-income and Tribal populations. None of the proposed alternatives to establish longline prohibited areas would have a high adverse health or environmental effect on members of environmental justice populations. The proposed action is intended to reduce gear conflicts and reduce catch competition by separating the troll and longline fleets, and, as shown in the preceding analyses, would not adversely affect target, non-target, or protected species resources.

9.1.10 Climate Change Impacts of the Proposed CNMI Longline Prohibited Areas

²² Amendment 20 to the Pelagics FMP

²³ PNA press release November 1, 2010, posted on www.atuna.com.

The Council considered several alternatives for the establishment of prohibited areas for longline and for purse seine fishing that vary in terms of the distance from shore. The larger the prohibited area the more fuel could be required by longliners or purse seiners to reach available fishing areas. However, under the proposed alternative, as described in section 9.1.11, CNMI longline vessels generally fish beyond 35 miles from shore, and so, under the preferred alternative, there would be no large change in the amount of fuel consumed or greenhouse gases emitted by longline vessels. For all of the longline prohibited area alternatives, longline vessels currently fish throughout the EEZ and our analysis shows that none of the alternatives would result in a large change to fuel consumption or greenhouse gas emissions by longline fishing vessels. The 30nm longline fishing prohibited area is the least burdensome in terms of transit requirements of all the alternatives.

Potential impacts of climate change (e.g., ocean acidification, sea level changes, warming temperatures, changes in circulation patterns, etc.) are not expected to change the effectiveness of any of the alternatives with respect to achieving the fishery objectives and meeting the purpose and need for action because the proposed longline prohibited area measure is intended to reduce the potential for gear interactions, reduce catch competition between trollers and longliners, and reduce the potential for longlining to affect troll catches due to localized depletion of fish in and around preferred troll fishing areas. Climate change effects would act upon the proposed action under any of the alternatives. If climate change were to affect fish stocks, this would continue to be detected through catch reporting and monitoring, and management measures could subsequently be recommended by the Council to address these changes in fish stocks.

Climate change impacts would not interact with the proposed prohibited longline fishing areas in a way that would change the low level of environmental impact expected to occur in implementing any of the alternatives.

9.1.11 Reasons for the Council's Selection of the CNMI Longline Prohibited Area as the Preferred Alternative

Alternative 1C, whereby all vessels would be prohibited from pelagic longline fishing within 30 nm of the CNMI, was selected by the Council as the preferred alternative because it best balances the needs of the small boat fleet for the reasons described below, while allowing for development of a longline fishery in waters around CNMI. The non-longline small boat fleet's needs include the ability to continue their small-scale fishing operations in waters around CNMI without localized stock depletion, catch competition, or gear conflicts with longline vessels.

The preferred alternative's 30 nm prohibited area encompasses the majority of the submarine habitat containing banks and reefs where the small boat fleet typically concentrates their fishing efforts. Making these areas off limits to longline activity would prevent gear and fishing conflicts between the two sectors and preclude longline gear entanglement with the banks and reefs. The preferred alternative may also prevent potential safety impacts to participants in the troll fishery which could arise if, without the prohibited area, the small boat operators felt compelled to fish farther from land and in unfamiliar waters to avoid gear conflicts or catch competition with longline vessels.

Also, the few longline vessels that have fished in CNMI in the past few years, both reported to be fishing outside of 35 miles from shore (J. Ogumoro, CNMI Island Coordinator, WPRFMC, pers. comm.). This supports selection of the 30 nm closure area with regards to avoiding gear conflicts and catch competition while not discouraging increased participation and sustained community participation in local, developing pelagic fisheries operations. Current longline operators in CNMI have also expressed their interest in purchasing fresh fish from the small boat fleet which would also be an economic benefit to CNMI.

In addition, as described in Section 5.0 of this document, a prohibited area of 25-30 nm was generally preferred by the fishing community in CNMI. It was thought that a 25 nm longline prohibited area would best provide an incentive to those interested in longline fishing in the area, while the local pelagic trollers would maintain exclusive access to most of the banks and seamounts which are located within the 25nm. They felt that allowing for longline fishing to occur outside the 25 nm area would help fishery development and the ailing economy while preventing gear conflicts. Others supported a larger prohibited area, such as 30nm or 50nm, as this would encompass all of the favorite trolling fishing areas, further reduce the potential for gear conflicts, and reduce catch competition for targeted species. They also argued that unlike the small boats, longline vessels are designed to fish further out in the open waters.

9.2 Topic 2: CNMI Purse Seine Prohibited Area

Alternative 2A: No Action

Under Alternative 2A (No Action) no new regulations would be made and pelagic purse seine vessels would not be prohibited from fishing within EEZ waters around CNMI.

Alternative 2B: 50 nm Purse Seine Prohibited Area

Under Alternative 2B, purse seining would be prohibited within 50 nm of CNMI.

Alternative 2C: 100 nm Purse Seine Prohibited Area

Under Alternative 2C, purse seining would be prohibited within 100 nm of CNMI.

Alternative 2D: Entire EEZ Purse Seine Prohibited Area (Preferred)

Under Alternative 2D, purse seining would be prohibited within all EEZ waters around CNMI.

9.2.1 Impacts on Target Stocks

Under Alternative 2A (No Action) pelagic landings and fishing effort by CNMI's existing pelagic troll fleet would continue and the current level of fishing is considered sustainable with respect to all target stocks. However, under any of the alternatives in which purse seine could take place in the EEZ, troll fishing could be impacted by gear interactions and decreased catches should purse seining expand into the EEZ around CNMI. Catches of pelagic species by the troll fleet under all alternatives would be expected to continue to be small, with a long term average of 224,276 lb (102 mt) which would have no significant adverse impact on target species.

If a purse seine fishery were to develop in EEZ waters around CNMI under Alternatives 2A, 2B, or 2C, the majority of the purse seine catch is projected to be skipjack tuna, which is also the most important target species for CNMI's troll fleet. The second highest purse seine catch would likely be yellowfin tuna which is the trolling fleet's second most important species. Based on 2004-06 Japanese purse seine operations (which are considered most representative of catches in this area, see Section 8.4), a single purse seine vessel can catch up to 29 mt of tuna per day, representing approximately 33 percent of the long term mean annual skipjack and yellowfin catch by CNMI's entire commercial troll fleet (89 mt, Table 3).

If a single purse seine vessel were to fish for an entire year in EEZ waters around CNMI under Alternatives 2A, 2B, or 2C it could be expected to catch up to 7,533 mt of tunas. Of this total, approximately 84 percent could be expected to be skipjack, 9 percent yellowfin tuna and the remainder bigeye and other mixed species. In comparison, this collectively adds up to about 85 times the entire CNMI troll fleet tuna harvests in one year. The assumption that a purse seine vessel would spend 100 percent of the time fishing within the EEZ around CNMI is unrealistic, as purse seining is a highly mobile fishery which harvests migratory fish species across vast portions of the ocean. Because of this, catches within the EEZ of CNMI are expected to be lower than projected in this analysis. However, in relation to historical catches of CNMI's troll fleet, any purse seining within the EEZ of CNMI would be expected to yield significantly greater catches than that caught by the troll fleet, and therefore may impact species composition and stock abundance within waters around CNMI.

There is currently no purse seine fishing in the U.S. EEZ around CNMI. If purse seining were to occur, the actual impacts of purse seine fishing in EEZ waters around CNMI on the stocks of WCPO skipjack, bigeye and yellowfin are unknown. There is likely to be some localized stock depletion under alternatives 2A, 2B, and 2C, but the degree of which would depend on the level and location of fishing effort. If purse seiners were to set on FADs or other floating objects they would be likely to harvest juvenile bigeye tuna which has been shown to contribute significantly to bigeye overfishing occurring in the WCPO due to concomitant impacts on spawning potential and recruitment of bigeye (Hampton et al. 2005). Overfishing issues related to unregulated FAD use in the WCPO are currently being addressed by the Council in a separate action. Nonetheless, the Pacific-wide impact would depend on whether already existing purse seine vessel(s) moved to CNMI from other areas of the WCPO, or new vessels were built for CNMI fishery. The former case would represent a transfer of existing impacts to CNMI waters, while the latter would result in new impacts.

Regardless of which alternative is selected, purse seine fishing will continue to be regulated by the WCPFC. None of the alternatives is expected to change the intensity or location of purse seine fishing, and none is expected to result in overfishing of target stocks.

9.2.2 Impacts on Non-target Stocks

Troll and longline impacts on non-target stocks are considered sustainable and none of the alternatives is expected to change the conduct of these fisheries, so there would be no change on impacts to non-target stocks from troll or longline fishing.

There is currently no purse seine fishing in the U.S. EEZ around CNMI. Under Alternative 2D, no resultant purse seine catches or associated non-target stock impacts would occur in this area as a result of purse seining. However, purse seining is expected to continue to occur at the same levels of intensity and in the same areas that it is currently occurring.

Under any of the alternatives that would allow purse seine vessels to fish in EEZ waters around CNMI (Alternatives 2A, 2B, or 2C) they would be expected to catch primarily target species, however, they would also catch some non-target species. Between 2005 and 2008 purse seine fleet non-target catches included: black marlin, blue marlin, silky shark, kawakawa, wahoo, mackerel scad, mahimahi, oceanic triggerfishes and rainbow runner (Table 11). Wahoo, mahimahi and blue marlin are retained by troll fishermen and collectively account for about 7% of the total pelagic fish catch on CNMI in 2008 (Table 3) and on average 12% of the annual pelagic troll catch between 1982 and 2008 (WPRMC 2010).

The majority of these species were discarded by the U.S. purse seine fleet and similar purse seine catches and discards could be expected under all alternatives. The impacts of purse seine fishing under Alternative 2A, 2B, or 2C in EEZ waters around CNMI on WCPO populations of non-target species are unknown; however, because of the large capacity and inability to target specific fishes, localized stock depletion may occur for non-target species, including those which have economic value. Purse seine discards could still occur under alternative 2D. There is no change to current purse seine fishing that would affect non-target species. Management of purse seine impacts on non-target species will continue to be the purview of the WCPFC, regardless of which alternative is selected.

9.2.3 Impacts on Protected Species

Under all of the alternatives, pelagic fishing around CNMI would continue to be conducted by trolling in small boats with concomitant impacts on protected species expected to continue to be nonexistent to minimal. The troll fishery around CNMI was included in NMFS' 2002 and 2009 biological opinions on pelagic fisheries of the Western Pacific Region which concluded that continued operation was not likely to jeopardize the continued existence of any listed sea turtles or marine mammals. A small potential exists that the trolling gear may incidentally hook or entangle a sea turtle or marine mammal, however, NMFS and the Council consider this type of interaction extremely rare and therefore unlikely to adversely affect listed species. None of the alternatives is expected to result in a change to impacts of trolling on protected species.

Under all of the alternatives, longline fishing would continue around the CNMI at a projected level of four longline vessels. None of the proposed alternatives is expected to change the level or intensity of longline vessels, and therefore, there would be no change in impacts to protected resources, which were described in the no-action alternative (section 9.1.3 above).

Under all of the alternatives that allow purse seine fishing to continue to be conducted in the U.S. EEZ, there is a potential for impacts on protected resources. In November 2006, NMFS issued a biological opinion on the effects of the U.S. purse seine fishery in the WCPO on listed sea turtles and whales. Potential adverse effects on listed species were analyzed on aspects of the fishery

identified to have potential adverse impacts including vessel traffic, gear deployment and retrieval, entanglement in FADs, and removal of fish biomass from the pelagic ecosystem (NMFS 2006). However, the relative impact of these interactions would depend on whether already existing purse seine vessel(s) moved to CNMI from other areas of the WCPO, or new vessels were built for CNMI fishery. The former case would represent a transfer of existing impacts to CNMI waters, while the latter would result in new impacts to protected species. Marine mammal and sea turtle interactions are reported to be disproportionately higher in sets associated with floating objects (associated sets) than in free swimming (unassociated sets) schools (NMFS 2006). Interaction information used in the biological opinion's analysis of effects was derived from observer data.

Molony (2005) found that the majority of interactions between turtles and marine mammals occur in the western tropical WCPO west of 170° W around the EEZs of Papua New Guinea, FSM, the Solomon Islands, and Nauru. The NMFS biological opinion concluded that the fishery is not likely to jeopardize the continued existence of any of the threatened or endangered species found in the area. The biological opinion's Incidental Take Statement expects 14 green, 14 hawksbill, 11 leatherback, 11 loggerhead, and 11 olive ridley sea turtles per year to be incidentally taken in the U.S. WCPO purse seine fishery. The biological opinion indicates zero mortality to result from interactions. Alternatives 2A, 2B, and 2C would all allow purse seine fishing within all or some EEZ waters around CNMI, with impacts to protected species likely to increase within the EEZ of CNMI proportional to the extent at which purse seine fishing were to occur, and proximity to shore where turtles such a greens and hawksbills are likely to be more abundant.

None of the alternatives would result in a change of the number of purse seiners, fishing intensity or activity of purse seining, or areas purse seiners are fishing in, so none of the alternatives would result in a change in impacts to protected species. Under Alternative 2D, no purse seine fishing would be allowed in EEZ waters around CNMI although purse seining would continue to occur in other areas of the western Pacific. There would be no impacts on protected resources in the waters of the CNMI or in other areas where purse seine fishing is occurring as a result of implementing the preferred alternative.

9.2.4 Impacts on Marine Habitat

Purse seine vessels are believed to normally have little impact on marine habitat as they deploy nets in the water column that do not contact the seabed, however, the EEZ waters around the Mariana Archipelago contain many submarine banks and seamount features which could be impacted by deployment and retrieval of a large purse seine net should it come into contact with these shallow features²⁴. Alternative 2A would allow purse seine fishing within zones in CNMI where seamounts are known to exist, Alternatives 2B, and 2C would all allow purse seine fishing within 50nm or 100nm, respectively, from shore within EEZ waters around CNMI. Since the majority of seamounts for fishing are within 30nm of landmasses in CNMI, alternatives 2B and 2C would have little or no impact on marine habitat. Under Alternative 2D, no purse seine fishing would be allowed in EEZ waters around CNMI and there would be no such impacts.

²⁴ Purse seine nets employed in the WCPO extend between 220 to 309 meters and are fished at depths ranging from 65 to 226 meters (Delgado de Molina et al. 2010)

The area affected by this action has been identified as EFH for pelagic management unit species managed in the western Pacific. The action is not likely to lead to substantial physical, chemical, or biological alterations to the habitat. The action in the context of the fishery as a whole will not change the intensity of fishing or fishing operations, therefore, the proposed alternatives would not result in a large or adverse impact on EFH; therefore, an EFH consultation is not required.

9.2.5 Impacts on Public Health and Safety

Allowing purse seine activities to occur within small boat pelagic fishing areas, under Alternative 2A, could decrease catches and potentially cause the small boats to fish farther from land and in unfamiliar waters resulting in a decrease in safety for those participants. Under Alternatives 2B and 2C, the larger the prohibited area, the greater the expected positive impacts to the troll fishing sector. Prohibiting purse seine fishing from U.S. EEZ waters around CNMI is not expected to result in any safety impacts to purse seine fishing participants as it is currently conducted in all areas of the Pacific including the high seas and this would not change.

9.2.6 Impacts on Fishery Participants and Communities

If a purse seine fishery were to fish in the EEZ around CNMI under Alternatives 2A, 2B, or 2C, the troll fishing community could be impacted by localized stock depletion leading to catch competition between the two fisheries (Hampton et al. 1996). The majority of the purse seine catch is projected to be skipjack tuna, which is also the most important target species for CNMI's troll fleet. The second highest purse seine catch would be yellowfin tuna which is also the trolling fleet's second most important species. Greater amounts of purse seine effort closer to areas fished by the small boat fleets of CNMI could force the troll fleet to travel farther to maintain their current catches, and/or to lose revenue normally gained through skipjack and yellowfin catches. Under the preferred Alternative 2D, no purse seine fishing would be allowed in EEZ waters around CNMI and there would, therefore, be no associated impacts. As described in Section 9.2.1, if purse seine fishing occurred around CNMI under the no-action Alternative 2A, projected purse seine catches could be many times that of the troll fleets typical annual harvest which could negatively impact the troll fleet's catches (as was shown in scientific studies described in Section 8.4.4) and cause economic losses to the troll participants and community.

Under the preferred alternative, no purse seine fishing would be allowed to take place in the U.S. EEZ waters around CNMI and this would represent a loss of opportunity for U.S. purse seiners if they had intended to fish these waters. Impacts, however, are expected to be minimal because the highly mobile purse seine fleet typically fishes in large areas of the Pacific and has exclusive rights to fish in U.S. EEZ waters in the PRIA, except those waters now off limits due to the establishment of the PRIA Marine National Monument.

While the U.S. purse seine fleet does not currently fish in waters of or deliver to CNMI, future purse seine fishing near CNMI, or the ability to accept fish transshipped by nearby purse seine vessels could influence the development the infrastructure needed to process or export such species. Aside from the negative impacts discussed earlier associated with gear competition and

stock depletion, the support of a purse seine industry could provide positive benefits through increasing potential employment while opening up markets currently unexplored in CNMI. Although for this to occur, CNMI's marine-based infrastructure (i.e., dock space, fueling platforms, skilled labor, etc) would need to be significantly enhanced beyond current conditions and is not likely to happen in the near future.

Under Alternative 2D, no purse seine fishing would be allowed in EEZ waters around CNMI and there would be no resultant purse seine catches or associated stock impacts in this area. It may be argued that such a stringent purse seine measure is not required and that one of the intermediary options, e.g. Alternatives 2B and 2C might be preferable. Arguments supporting this view include the lack of any historical catches made in the U.S. EEZ around the entire Mariana Archipelago, and the fact that only about a third of the U.S. purse seine fleet has authorization to fish within U.S. EEZ waters.

However, as noted previously, the catching power of a single purse seiner is several orders of magnitude greater than the troll fisheries in either Guam or NMI. Moreover, the fledgling longline fishery based out of Saipan targets yellowfin and bigeye tuna which are caught by purses seiners, especially when fishing on FADs. Thus purse seine fishing poses the potential to competitively interact with and adversely affect the two main pelagic fisheries in the CNMI. Further, purse seine fishing is likely to become more restricted elsewhere, especially if measures being proposed by the PNA Group are adopted by the WCPFC. These include the doubling or tripling of the costs of access agreements to fish in PNA Group EEZs²⁵, and, as a condition of access, the requirement of purse seine access agreements to fish on the high seas between 10 deg N to 20 deg S latitude and 170 deg E to 150 deg W longitude²⁶. There has also been a proposal advanced at the 2010 WCPFC Technical and Compliance Committee (TCC) meeting (WCPFC 2010) by the PNA Group to extend an existing annual three-month ban on purse seining on FADs in PNA EEZs for an additional three months for purse seine fleets that catch in excess of 4,000MT of bigeye annually.

The U.S. fleet operates under the South Pacific Tuna Treaty (SPTT) which provides access to PNA Group waters. Other Pacific Island Nations (PINs) are also members of the Treaty, but it's the PNA Group countries, clustered around the Pacific Warm Pool that are the principal fishing grounds for U.S. and other nation's purse seine fleets. At the time of writing, the SPTT expires in 2013 and a new agreement has not yet been concluded. However, as noted above, the PNA Group is signaling its intentions to place greater restrictions on purse seine fleet operations for access to their waters. Further, the FFA member countries and the PNA group have elected to allocate access to their EEZs for purse seiners based on a Vessel Day Scheme (VDS). The current SPTT has provisions for U.S. 40 vessels to fish, with an additional 5 vessels allowed under joint venture arrangements. However, the PNA Group may want the U.S. vessels to fish under the VDS like other fleets. The number of Vessel Days allocated to the U.S. fleet may fall short of the number needed by a fleet currently with 36 active vessels.

²⁵ Islands Business November 2010 article: 'Pacific islands look at tuna fisheries'

²⁶ FFA Press release April 23, 2010, PNA announces date for closure of 4 million square kilometer high seas areas to purse seine fishing.

Taken together, the more aggressive attitude of the PNA Group to confine purse seine fishing to their EEZ waters, to increase access costs, to penalize purse seine fleets with domestic longline bigeye fisheries and the potential for the VDS allocation to be less than optimum for the U.S. fleet provides an incentive for those U.S. hulled vessels to explore options such as fishing in U.S. EEZ waters. There are proven skipjack resources in the U.S. EEZ around the Mariana Archipelago as evinced from the Japanese pole and line fishery that operated from Saipan between 1922 and 1942 (Higuchi 2007) and which at its peak landed 3,700 mt of skipjack. As noted above, a single purse seine vessel is capable of catching twice this amount annually.

Closing the entire U.S. EEZ around the Mariana Archipelago (Alternatives 2D and 3D) to purse seine fishing provides the maximum degree of protection for the small boat fleets of the two island groups, which collectively amount to over 430 small fishing vessels in Guam and CNMI (Figures 5 and 12), and protects the long terms continuity of small scale pelagic fisheries which have social, economic and cultural importance for the two U.S. territories. U.S. vessels with authorization to fish within U.S. EEZ waters would still have access to part of the U.S. EEZ around American Samoa and the Pacific Remote Islands Area, of which Howland and Baker, Jarvis and Palmyra and Kingman Reef. All of these areas have been fished by the U.S. fleet in the past and would continue to be accessible to vessels with U.S. hulls. American Samoa longline and troll fishermen had considered a complete closure of their EEZ waters to purse seine fishing for the same reasons, but recognized that the U.S. purse seine fleet has a history of fishing in this segment of the U.S. EEZ, and so elected for a compromise that closes waters out to 75 nm from their islands. As such the preferred alternatives 2D and 3D balances the needs of the CNMI and Guam fishermen to fish and provide a source of fresh tuna to the local populace, with those U.S. purse seine vessels who have authorization to fish within UZ EEZ waters still able to fish in other segments of the U.S. EEZ in the Western Pacific.

9.2.7 Impacts on Biodiversity and Ecosystem Functions

This action will have no known adverse impacts on biodiversity and ecosystem functions since the action of creating fishing prohibited zones would protect, rather than diminish, marine resources. If purse seiners were to begin fishing in U.S. EEZ waters around the CNMI then removal of top predators, including tuna, could potentially alter the trophic food chain and the energy flow through the ecosystem, however, this is not readily discernible and the potential impacts are unknown. Under the preferred Alternative 3D, no purse seine fishing would be allowed in EEZ waters around Guam and there would therefore be no associated impacts.

9.2.8 Impacts on Administration and Enforcement

There would be no impacts on administration and enforcement under Alternative 2A (No Action). Alternatives 2B, 2C, and 2D would all place similar administrative burden of publishing regulations associated with a prohibited area and informing all fishery participants. Alternatives 2B, 2C, and 2D would all place an additional burden on law enforcement, U.S. Coast Guard and NOAA Office of Law Enforcement (OLE), in enforcing the closed areas through VMS, which all U.S. purse seiners are obliged to carry.

9.2.9 Cumulative Effects of the Proposed CNMI Purse Seine Prohibited Areas

Establishment of a purse seine prohibited area in the CNMI would address concerns about the large harvests and harvests of juvenile tunas. Regardless of which alternative is selected for a purse seine prohibited area in the CNMI, none of the alternatives would likely change the intensity or location of fishing by trollers, longliners, or purse seiners. Other fishery management activities that are being proposed that affect purse seiners (e.g., establishing purse seine closed areas in American Samoa and managing FAD use by purse seiners) would not result in any of the alternatives having a large or negative environmental effect.

9.2.10 Environmental Justice Effects of the Proposed CNMI Purse Seine Prohibited Areas

Environmental justice policies intend to promote environmental protection by focusing attention on potential environmental harms and risks that may disproportionately and adversely affect minority, low-income and Tribal populations. None of the proposed alternatives to establish a purse seine prohibited area would have a high adverse health or environmental effect on environmental justice populations. The proposed action is intended to reduce gear conflicts, reduce catch competition, and prevent localized depletion by purse seining. None of the alternatives would change the current intensity or location of fishing by purse seine vessels, or troll or longline vessels, so there would not be an adverse environmental or health impact from implementing any of the alternatives.

9.2.11 Climate Change Impacts of the Proposed CNMI Purse Seine Prohibited Areas

The alternatives consider various distances from shore at which purse seiners could fish. There is no purse seine fishing occurring in the CNMI, and none of the alternatives would change the locations or intensity of fishing by purse seiners, or local troll and longline fishing vessels. None of the proposed alternatives would result in a federal contribution to greenhouse gas emissions greater than 25,000 mt carbon dioxide equivalents. Climate change impacts are not expected to affect the effectiveness of any of the alternatives with respect to achieving the fishery objectives and meeting the purpose and need for action. Climate change is also not expected to affect the environmental impact of implementing any of the alternatives which would establish a geographic separation between fishing fleets.

9.2.12 Reasons the Council Selected the CNMI Purse Seine Prohibited Area as its Preferred Alternative

The preferred Alternative 2D, recommended by the Council, would prohibit all vessels from pelagic purse seine fishing within U.S. EEZ waters around CNMI. This is intended to preserve the viability of smaller scale local pelagic fisheries in CNMI while potentially providing positive benefits to Pacific stocks of target tunas by preventing the harvest of undersized tunas within U.S. EEZ waters.

Purse seine vessels operating in the WCPO caught over 1.52 million mt (1.675 t) in 2005, a record high catch dominated by skipjack tuna at approximately 1.25 million mt (1.38 t) (OFF 2007). The provisional total WCPO tuna catch for 2009 was estimated at 2,467,903 mt, the

highest annual catch recorded and 70,000 mt higher the previous record in 2008 (2,398,664 mt) (Williams & Terasi 2010). Further, The 2009 WCPO catch of skipjack (1,789,979 mt – 73% of the total catch) was the highest recorded, and nearly 120,000 mt more than the previous record catch of 2007 (1,672,996 mt).

As discussed earlier, in a few days, a single purse seiner has the potential to catch tuna species well in excess of similar catches by the entire CNMI troll fleet for one year (see Table 3) Moreover, the conclusions from the Hampton et al (1995) and Kleiber (pers. comm.) studies, and others listed in Section 8.4.4 (Nishida 1995; Ortega-Garcia1995; Sibert et al. 1995; Shomura et al. 1996) suggest that every mile of separation between purse seine fisheries and other pelagic fisheries is beneficial in terms of the other fisheries catch rates.

Alternative 2D was selected because it best balances the needs of the existing small boat fleet around CNMI, allows for expansion of longline fishing, and creates a zone around CNMI where these sectors may operate without potential impacts of stock depletion and subsequent catch competition caused by purse seine fishing within the EEZ. There are several other compelling justifications that support a prohibition on purse seine fishing in EEZ waters around CNMI, including:

- (1) As noted above in this Section and as described in Section 8.4.3, several studies have shown direct impacts of purse seine fishing on tuna catches in longline and pole-and-line fleets;
- (2) Although no purse seine catches have yet been recorded from EEZ waters around CNMI the potential impacts to the small boat and fledgling longline fleet should this occur must be considered especially in light of the fact that the largest skipjack catch in world history was recorded in 2009, much of it from waters adjacent to the Mariana Archipelago;
- (3) The U.S. purse seine fleet has rebuilt and thus more vessels may lead to expansion of fishing grounds because of competition with other purse seine fleets especially since the Pacific-wide catch of skipjack is at an all-time high;
- (4) The U.S. purse seine fleet may be looking for new areas to fish to compensate for lost fishing grounds from Monument closures in the Pacific (PRIA, Rose Atoll and the Mariana Trench) and potential RFMO instituted high seas pockets closures;
- (5) CNMI small boats rely on catches of skipjack and other pelagics for sustenance and as an important part of their economy.

The cumulative effects of the expansion of purse seine fishing in both the WCPO and the EPO have contributed to overfishing of bigeye tuna across the Pacific, but especially in the EPO, where the stock may be approaching an overfished condition. FAD-associated purse seine fishing is contributing to excess catches of juvenile yellowfin tuna which is of concern Pacific-wide. In 2007 the U.S. fleet made 80 percent of all sets on drifting FADs which is a considerably higher percentage than other nations fishing in the WCPO (OFP 2007). It should be noted that

regulatory measures being considered and in place now (annual seasonal FAD closures) in the WCPO may further restrict FAD associated fishing of this type in the WCPO.

In addition, there are other suitable fishing areas the purse seine fleet can access due to their great mobility unlike the small vessel fleets which are travel restricted. Hence, the Council believes Alternative 2D is warranted to balance the needs of the existing small boat fleet around CNMI and allow for expansion of CNMI-based longline fishing while minimizing the potential for localized stock depletion associated catch competition with purse seine vessels.

9.3 Topic 3: Impacts of Proposed Alternatives for a Guam Purse Seine Prohibited Area

Alternative 3A: No Action

Under Alternative 3A (No Action) no new regulations would be made and pelagic purse seine vessels would not be prohibited from fishing within EEZ waters around Guam.

Alternative 3B: 50 nm Purse Seine Prohibited Area

Under Alternative 3B all vessels would be prohibited from pelagic purse seine fishing within 50 nm of Guam.

Alternative 3C: 100 nm Purse Seine Prohibited Area

Under Alternative 3C, all vessels would be prohibited from pelagic purse seine fishing within 100 nm of Guam.

Alternative 3D: Entire EEZ Purse Seine Prohibited Area

Under Alternative 3D, all vessels would be prohibited from pelagic purse seine fishing within all EEZ waters around Guam.

9.3.1 Impacts of the Proposed Guam Purse Seine Prohibited Areas on Target Stocks

Under Alternative 3A (No Action) pelagic landings and fishing effort by Guam's existing pelagic troll fleet would continue to fluctuate based on weather, catchability, fuel prices, and other factors. Catches of pelagic species by this troll fleet would be expected to continue to be sustainable because they are small and this fishery does not have a large and adverse impact on target species.

There is currently no purse seine fishing in the U.S. EEZ around Guam. If a purse seine fishery were to develop in EEZ waters around Guam under Alternatives 3A, 3B, or 3C, the majority of the purse seine catch is projected to be skipjack tuna, which is also the most important target species for Guam's troll fleet. Based on 2002 Japanese purse seine operations (which are considered most representative of catches in this area, see Section 8.4), if a single purse seine vessel were to fish for a single day in EEZ waters around Guam, it could be expected to catch up to 32 mt of tuna, including 26 mt of skipjack, 3 mt of yellowfin tuna, and 0.55 mt of bigeye and mixed species which represents approximately 36 percent of the long term mean annual skipjack and yellowfin catch (107 mt) by Guam's entire troll fleet (Table 5).

If a single purse seine vessel were to fish for an entire year in U.S. EEZ waters around Guam under Alternatives 3A, 3B, or 3C they could be expected to catch up to 7,533mt of tunas, assuming the relatively small EEZ around Guam could support such catch rates. Of this total, 84 percent could be expected to be skipjack, 9 percent yellowfin tuna and the remainder bigeye and other mixed species. Impacts arising from purse seine harvest, such as localized stock depletion, would be less pronounced with increased size of a prohibited area (i.e. Alternative 3A would be expected to have greater impact on target stocks than Alternative 3C). The assumption that a purse seine vessel would spend 100 percent of the time fishing within the EEZ around Guam is unrealistic, as purse seining is a highly mobile fishery which harvests migratory fish species across large areas of the ocean. Because of this, catches within the EEZ of Guam are expected to be lower than projected in this analysis. However, in relation to historical catches of Guam's troll fleet, any purse seining within the EEZ of Guam would be expected to yield significantly greater catches than that caught by the troll fleet, and therefore may impact species composition and stock abundance within waters around Guam.

The relative impact of purse seine fishing on Pacific stocks would depend on whether already existing purse seine vessel(s) moved to Guam from other areas of the WCPO, or new vessels were built for the Guam fishery. The former case would represent a transfer of existing impacts to Guam waters, while the latter would result in new impacts. Note also that the U.S. EEZ around Guam is very small, and is one of the smaller zones in the WCPO, although the entire U.S. EEZ around the Mariana Archipelago is large, amounting to about 182,000 square nautical miles. Moreover, the U.S. EEZ around Guam is immediately adjacent to the epicenter of the Warm Pool where purse seining is known to occur, thus any transfer of fishing effort into the southern part of the U.S. EEZ around the Mariana Archipelago would be expected to have an influence on skipjack and yellowfin abundance.

The impacts of purse seine fishing in EEZ waters around Guam on the WCPO skipjack, bigeye and yellowfin populations are unknown, however there is likely to be localized stock depletion, the degree of which would depend on the level and location of fishing effort. If purse seiners were to set on FADs or other floating objects they would be likely to harvest juvenile bigeye tuna which has been shown to contribute significantly to bigeye overfishing occurring in the WCPO due to concomitant impacts on spawning potential and recruitment of bigeye (Hampton et al. 2005). With larger prohibited areas, it is anticipated that the potential for adverse effects to stocks within the EEZ and localized depletion would be smaller. Under Alternative 3D, no purse seine fishing would be allowed in EEZ waters around Guam and there would be no resultant purse seine catches or associated stock impacts in this area.

Regardless of which alternative is selected, purse seine fishing outside of the U.S. EEZ around the Marianas Archipelago will continue to be regulated by the WCPFC. None of the alternatives is expected to change the intensity or location of purse seine fishing, longline fishing, or troll fishing, and none is expected to result in overfishing of target stocks.

9.3.2 Impacts on Non-target Stocks

There is currently no purse seine fishing in the U.S. EEZ around CNMI. If purse seine vessels were to begin fishing in EEZ waters around Guam under Alternatives 3A, 3B, or 3C they would be expected to catch primarily target species, however, they would also catch some non-target species. Between 2005 and 2008 purse seine fleet non-target catches included: black marlin, blue marlin, silky shark, kawakawa, wahoo, mackerel scad, mahimahi, oceanic triggerfishes and rainbow runner (Table 11). The majority of these were discarded and similar purse seine catches and discards could be expected under all Alternatives except 3D which would not allow any purse seine fishing in EEZ waters around Guam. Purse seine discards could occur outside of the EEZ under Alternative 3D. None of the alternatives is expected to increase purse seine impacts on non-target stocks including the amount of discards that are made.

Wahoo, mahimahi and blue marlin are an important component of the pelagic troll catch in Guam, forming about 40% of landings in 2008 (Table 5) and over 50% of landings on average between 1982 and 2008 (WPRFMC 2010). Thus the impact of purse seine fishing on non-target stocks may have an additional impact on the troll fishery given the importance of the non-tuna stocks in the troll catch. Further, Guam has a charter vessel fishery of about 16 boats for which blue marlin are an important catch component. The charter fleet being limited catches only about 9% of the total pelagic fish catch but 30% of the blue marlin catch (WPRFMC 2010)

The majority of these species were discarded by the U.S. purse seine fleet and similar purse seine catches and discards could be expected under all alternatives except the preferred alternative, Alternative 3D, which would prohibit purse seine fishing in the entire EEZ around Guam. The impacts of purse seine fishing under Alternative 3A, 3B, or 3C in EEZ waters around Guam on WCPO populations of non-target species are unknown, but fishery managers from the WCPFC track discards through logbook reports and the WCPFC is responsible for ensuring sustainability of the purse seiner fishery. If purse seine fishing were to develop in the EEZ however, localized stock depletion could occur for non-target species, including those which have economic value.

9.3.3 Impacts on Protected Species

Under Alternative 1A (No Action), pelagic fishing around Guam would continue to be conducted by trolling in small boats with concomitant impacts on protected species expected to continue to be minimal. The troll fishery around Guam was included in NMFS' 2002 biological opinion on pelagic fisheries of the Western Pacific Region which concluded that continued operation was not likely to jeopardize the continued existence of any listed sea turtles or marine mammals. A small potential exists that the trolling gear may incidentally hook or entangle a sea turtle or marine mammal, however, NMFS and the Council consider this type of interaction extremely rare and therefore unlikely to adversely affect listed species. None of the alternatives is expected to result in a change to impacts of trolling on protected species.

There is little available information on protected species interactions with WCPO purse seine vessels, however these vessels are under an international management regime which includes consultations under section 7 of the Endangered Species Act. In November 2006, NMFS issued its biological opinion on the effects of the U.S. purse seine fishery in the WCPO on listed sea turtles and whales. Potential adverse effects on listed species were analyzed on aspects of the fishery identified to have potential adverse impacts including vessel traffic, gear deployment and retrieval, entanglement in FADs, and removal of fish biomass from the pelagic ecosystem (NMFS 2006).

Marine mammal and sea turtle interactions are reported to be disproportionately higher in sets associated with floating objects (associated sets) than in free swimming (unassociated sets) schools (NMFS 2006). Interaction information used in the biological opinion's analysis of effects was derived from observer data. Molony (2005) found that the majority of interactions between turtles and marine mammals occur in the western tropical WCPO west of 170° W around the EEZs of Papua New Guinea, FSM, the Solomon Islands, and Nauru. The NMFS biological opinion concluded that the fishery is not likely to jeopardize the continued existence of any of the threatened or endangered species found in the area. The biological opinion's Incidental Take Statement expects 14 green, 14 hawksbill, 11 leatherback, 11 loggerhead, and 11 olive ridley sea turtles per year to be incidentally taken in the U.S. WCPO purse seine fishery, however, their data indicate zero mortality to result. For more detailed information refer to the 2006 biological opinion (NMFS 2006). Alternatives 3A, 3B, and 3C would all allow purse seine fishing within EEZ waters around Guam with their associated impacts to protected species. Under Alternative 3D, no purse seine fishing would be allowed in EEZ waters around Guam and there would, therefore, be no associated impacts to protected species from purse seining within the EEZ. Purse seiners are expected to continue fishing, even under Alternative 3D, so there is no change expected to impacts of the alternatives on protected resources for any of the alternatives.

9.3.4 Impacts on Marine Habitat

Purse seine vessels are believed to normally have little impact on marine habitat as they deploy nets in the water column that do not contact the seabed, however, the EEZ waters around the Mariana Archipelago contain many submarine banks and sea mount features which could be impacted by deployment and retrieval of a large purse seine net should it come into contact with these shallow features (see footnote 22). Alternatives 3A, 3B, and 3C would all allow purse seine fishing within EEZ waters around CNMI with their associated potential impacts to marine habitat. Under Alternative 3D, no purse seine fishing would be allowed in EEZ waters around Guam and there would be no associated impacts. None of the alternatives is expected to result in a change of fishing operations or fishing intensity, and therefore, none of the alternatives is expected to result in large or adverse effects on marine habitats.

9.3.5 Impacts on Public Health and Safety

If purse seine vessels were to begin fishing in Guam's EEZ waters under Alternatives 3B or 3C there would likely be positive impacts on the health and safety of the small boat fishery participants, as compared to the no-action alternative, as they would not experience catch conflicts with purse seine vessels when fishing inside the prohibited area. Without a purse seine

prohibited area such conflicts, perceived or actual, could cause the small boats to fish farther from land and in unfamiliar waters which could potentially result in a decrease in safety for those participants. The larger the prohibited area, the greater the positive impacts would be. Under Alternative 3D, no purse seine fishing would be allowed in EEZ waters around Guam and there would be no associated impacts.

9.3.6 Impacts on Fishery Participants and Communities

If a purse seine fishery were to fish in EEZ waters around Guam under Alternatives 3A, 3B, or 3C, the troll fishing community could be impacted by localized stock depletion leading to catch competition between the two fisheries (Hampton et al. 1996). As described in Section 9.3.1, if purse seine fishing occurred around Guam under Alternative 3A, purse seine catches could be expected to reduce the troll fleet's catches (as was shown in scientific studies described in Section 8.4.4). The majority of the purse seine catch is projected to be skipjack tuna, which is the most important species for Guam's trolling vessels. The second highest purse seine catch would be yellowfin which is also important to the trolling fleet. As discussed in Section 9.4.1 the catch of skipjack and yellowfin by one purse seine vessel fishing for one day (29 t) would be approximately 33% percent of Guam's entire troll fleet's (2006) annual catch of these species (approximately 89 tons, Table 5) and greater amounts of purse seine effort could force the troll fleet to travel farther to maintain their current catches, and to lose revenue normally gained through skipjack and yellowfin catches. Impacts arising from purse seine harvest, such as localized stock depletion leading to decreased catches for the small vessel Guam-based fleet, would be less pronounced with increased size of an prohibited area (i.e. Alternative 3A would be expected to have greater impact on target stocks than Alternative 3C). Under Alternative 3D, no purse seine fishing would be allowed in U.S.EEZ waters around Guam, therefore minimizing such impacts.

Under the preferred alternative, no purse seine fishing would be allowed to take place in the U.S. EEZ waters around Guam and it is possible there would be some concomitant impacts to the U.S. purse seine fleet if they had chosen to fish these waters. However, the majority of these impacts are expected to be minimal due to the highly mobile nature of the purse seine fleet, although the exclusive right of U.S. hull purse seiners to fish in waters around the PRIA has been diminished by establishment of the PRIA Marine National Monument which closes a portion of these waters to any fishing.

Under Alternative 3D, no purse seine fishing would be allowed in EEZ waters around Guam and there would be no resultant purse seine catches or associated stock impacts in this area. It may be argued that such a stringent purse seine measure is not required and that one of the intermediary options, e.g. Alternatives 2B and 2C might be preferable. Arguments in favor of this include the lack of any historical catches made in the U.S. EEZ around the entire Mariana Archipelago, and about only a third of the U.S. purse seine fleet has authorization to fish within U.S. EEZ waters.

However, as noted previously, the catching power of a single purse seiner is several orders of magnitude greater than the troll fisheries in either Guam or NMI. Moreover, the fledgling longline fishery based out of Saipan targets yellowfin and bigeye tuna which are caught by purses seiners, especially when fishing on FADs. Thus purse seine fishing poses the potential to

competitively interact with the two main pelagic fisheries in the CNMI. Further, purse seine fishing is likely to become more restricted, especially if measures being proposed by the PNA Group are adopted by the WCPFC. These include the doubling or tripling of the cost of access agreements to fish in PNA Group EEZs²⁷, and, as a condition of access, the requirement of purse seine access agreements to fish on the high seas between 10 deg N to 20 deg S latitude and 170 deg E to 150 deg W longitude²⁸. There has also been a proposal advanced at the 2010 WCPFC Technical and Compliance Committee (TCC) meeting (WCPFC 2010) by the PNA Group that a bans on FAD fishing in PNA EEZs be additional three months for purse seine fleets from flag States that operate longline vessels that catch in excess of 4,000MT of bigeye annually.

The U.S. fleet operates under the South Pacific Tuna Treaty (SPTT) which provides access to PNA Group waters. Other Pacific Island Nations (PINs) are also members of the Treaty, but the it's the PNA Group countries, clustered around the Pacific Warm Pool that are the principal fishing grounds for U.S. and other nation's purse seine fleets. At the time of writing, the SPTT expires in 2013 and a new agreement has not yet been concluded. However, as noted above, the PNA Group is signaling its intentions to place greater restrictions on purse seine fleet operations for access to their waters. Further, the FFA member countries and the PNA group have elected to allocate access to their EEZs for purse seiners based on a Vessel Day Scheme (VDS). The current SPTT has provisions for U.S. 40 vessels to fish, with an additional 5 vessels allowed under joint venture arrangements. However, the PNA Group may want the U.S. vessels to fish under the VDS like other fleets. The number of Vessel Days allocated to the U.S. fleet may fall short of the number needed by a fleet currently with 36 active vessels.

Taken together, the more aggressive attitude of the PNA Group to confine purse seine fishing to their EEZ waters, to increase access costs, to penalize purse seine fleets with domestic longline bigeye fisheries and the potential for the VDS allocation to be less than optimum for the U.S. fleet, provides and incentive for those U.S. hulled vessels to explore options such as fishing in U.S. EEZ waters. There are proven skipjack resources in the U.S. EEZ around the Mariana Archipelago as evinced from the Japanese pole and line fishery that operated from Saipan between 1922 and 1942, and which at its peak landed 3,700 mt of skipjack. As noted above, a single purse seine vessel is capable of catching twice this amount annually.

Closing the entire U.S. EEZ around the entire Mariana Archipelago (Alternatives 2D and 3D) to purse seine fishing provides the maximum degree of protection for the small boat fleets of the CNMI and Guam, which collectively amount to over 430 small fishing vessels, and protects the long term continuity of small scale pelagic fisheries which have social, economic and cultural importance for the two U.S. territories. U.S. vessels with authorization to fish within U.S. EEZ waters would still have access to part of the U.S. EEZ around American Samoa and the Pacific Remote Island Areas, which includes Howland and Baker, Jarvis and Palmyra and Kingman Reef. All of these areas have been fished by the U.S. fleet in the past and would continue to be accessible to vessels with U.S. hulls. American Samoa longline and troll fishermen had considered a complete closure of their EEZ waters to purse seine fishing for the same reasons, but recognized that the U.S. purse seine fleet has a history of fishing in this segment of the U.S.

²⁷ Islands Business November 2010 article: 'Pacific islands look at tuna fisheries'

²⁸ FFA Press release April 23, 2010, PNA announces date for closure of 4 million square kilometer high seas areas to purse seine fishing.

EEZ, and so elected for a compromise that closes waters out to 75 nm from their islands. As such the preferred alternatives 2D and 3D balance the needs of the CNMI and Guam fishermen to fish and provide a source of fresh tuna to the local populace. U.S. purse seine vessels who have authorization to fish within U.S. EEZ waters would still be able to fish in other segments of the U.S. EEZ in the Western Pacific.

9.3.7 Impacts on Biodiversity and Ecosystem Functions

This action will have no known adverse impacts on biodiversity and ecosystem functions since the action of creating fishing prohibited zones would protect, rather than diminish, marine resources. If purse seiners were to begin fishing in U.S. EEZ waters around Guam the removal of top predators, including tuna, could potentially alter the trophic food chain and the energy flow through the ecosystem, however, this is not readily discernible and the potential impacts are unknown. Under the preferred Alternative 3D, no purse seine fishing would be allowed in EEZ waters around Guam and there would therefore be no associated impacts.

9.3.8 Impacts on Administration and Enforcement

There would be no impacts on administration and enforcement under Alternative 3A (No Action). Alternatives 3B, 3C, and 3D would all place similar administrative burden of publishing the prohibited area and informing all fishery participants. Alternatives 3B, 3C, and 3D would all place similar burden on law enforcement, U.S. Coast Guard and NOAA Office of Law Enforcement (OLE), by enforcing the closed area.

9.3.9 Cumulative Effects of the Proposed Guam Purse Seine Prohibited Areas

Establishment of a purse seine prohibited area in Guam would address concerns about the large harvests and harvests of juvenile tunas. Regardless of which alternative is selected for a purse seine closed area in Guam, none of the alternatives would likely change the intensity or location of fishing by trollers, longliners, or purse seiners. Other fishery management activities that are being proposed that affect purse seiners (e.g., establishing purse seine closed areas in American Samoa and managing FAD use by purse seiners) would not result in any of the alternatives having a large or negative environmental effect.

9.3.10 Environmental Justice Effects of the Proposed Guam Purse Seine Prohibited Areas

Environmental justice policies intend to promote environmental protection by focusing attention on potential environmental harms and risks that may disproportionately and adversely affect minority, low-income and Tribal populations. None of the proposed alternatives to establish a purse seine prohibited area would have a high adverse health or environmental effect on environmental justice populations. The proposed action is intended to reduce gear conflicts, reduce catch competition, and prevent localized depletion by purse seining. None of the alternatives would change the current intensity or location of fishing by purse seine vessels, or

troll or longline vessels, so there would not be an adverse environmental or health impact from implementing any of the alternatives.

9.3.11 Climate Change Impacts of the Proposed Guam Purse Seine Prohibited Areas

The alternatives consider various distances from shore at which purse seiners could fish. There is no purse seine fishing occurring in the Guam, and none of the alternatives would change the locations or intensity of fishing by purse seiners, or local troll and longline fishing vessels. Because there is no expected change to current fishing operations from this proposed action, none of the proposed alternatives would result in a federal contribution to greenhouse gas emissions greater than 25,000 mt carbon dioxide equivalents. Climate change impacts are not expected to affect the effectiveness of any of the alternatives with respect to achieving the fishery objectives and meeting the purpose and need for action because the action is dependent on establishment of a separation between fishing fleets.

9.3.12 Reasons the Council Selected the Guam Purse Seine Prohibited Area as the Preferred Alternative

Preferred Alternative 3D would prohibit all vessels from pelagic purse seine fishing within all EEZ waters around Guam.

As shown in Table 9 a single purse seiner can catch on average 26 t of skipjack and 3 t of yellowfin tuna a day which far exceeds catches by the Guam-based troll fleet with on average 142 mt of skipjack and yellowfin tuna caught in 2008 and on average about 89 mt per year between 1982 and 2008. The provisional total WCPO tuna catch for 2009 was estimated at 2,467,903 mt, the highest annual catch recorded and 70,000 mt higher the previous record in 2008 (2,398,664 mt) (Williams & Terasi 2010). Further, the 2009 WCP-CA catch of skipjack (1,789,979 mt – 73% of the total catch) was the highest recorded, and nearly 120,000 mt more than the previous record catch of 2007 (1,672,996 mt).

As discussed earlier, in a few days, a single purse seiner has the potential to catch tuna species well in excess of similar catches by the entire Guam troll fleet for one year. Moreover, the conclusions from the Hampton et al (1995) and Kleiber (pers. comm.) studies, and others listed in Section 8.4.4 (Nishida 1995; Ortega-Garcia 1995; Sibert et al. 1995; Shomura et al. 1996) suggest that every mile of separation between purse seine fisheries and other pelagic fisheries is beneficial in terms of the other fisheries catch rates.

Alternatives 2D and 3D would implement consistent purse seine prohibited areas around Guam and CNMI which would facilitate compliance and enforcement by providing consistent fishery regulations.

In summary, there are a list of compelling justifications to a prohibition on purse seine fishing in EEZ waters around Guam including:

- 1) that several studies have shown direct impacts of purse seine fishing on tuna catches in longline and pole-and-line fleets, as described in Section 8.4.3, and

- 2) although no domestic purse seine catches have yet been recorded from U.S. EEZ waters around Guam the potential impacts to the small boat and fledgling longline fleet should this occur must be considered especially in light of the fact that the largest skipjack catch in world history was recorded in the most recent (2007) fishing year; and
- 3) FSM also has one of the largest resources of surface tunas in its extensive zone, whereas purse seine catches in Palau and Marshall Islands waters are much smaller and tend to fluctuate considerably from year to year²⁹ and Guam's EEZ waters are contiguous with FSM EEZ waters and could therefore be expected to have similar capacity; and
- 4) Guam has an extremely small segment of the U.S. EEZ as it is truncated by FSM's EEZ waters to the south and CNMI's portion of the U.S. EEZ to the north; and
- 5) the WCPO purse seine fleet may be at overcapacity³⁰ while U.S. vessels with an endorsement to fish in the U.S. EEZ may choose to fish in alternate areas of the U.S. EEZ without foreign vessel competition; and
- 6) Guam's small boats and charter industry relies on catches of skipjack and other PMUS for sustenance and as an important part of their economy, currently in a severe decline; and
- 7) Guam's small scale fisheries may already be experiencing some localized depletion due to the purse seine activities in FSM waters (annual catches >200,000 mt/yr)³¹; and
- 8) the Magnuson-Stevens Fishery Conservation and Management Act mandates consideration of the importance of fishery resources to fishing communities in order to provide for the sustained participation of such communities, and to the extent practicable, minimize adverse economic impacts on such communities.

9.4 Essential Fish Habitat

The preferred alternatives are not expected to have adverse impacts on essential fish habitat (EFH) or habitat areas of particular concern (HAPC) for species managed under the Pacific Pelagics, American Samoa Archipelago, Mariana Archipelago, Hawaii Archipelago, or Pacific Remote Island Areas Fishery Ecosystem Plans. EFH and HAPC for these species groups has been defined as presented in Table 18 and also described in detail in the Pelagics FEP (WPFMC 2009).

²⁹ <http://www.ffa.int/system/files/Trip+report+-+Palau,+FSM,+RMI.pdf>

³⁰ The increase in purse seine capacity in the WCPO in the past decade has prompted a proposal from Japan for the WCPFC to adopt a Conservation and Management Measure to ensure that the level of purse seine fishing capacity in the number of purse seine vessels from member countries and cooperating non-members fishing on the high seas does not increase from the current level, and to ensure that the level of purse seine fishing effort in days fished on the high seas does not increase from the levels of 2004 or the average of 2001-2004.

³¹ <http://www.ffa.int/system/files/Trip+report+-+Palau,+FSM,+RMI.pdf>

None of the alternatives would adversely affect EFH or HAPC for any managed species as none of the alternatives would change the intensity of fishing by the troll, longline, or purse seine fleets, and none of the alternatives would result in physical, chemical, or biological alterations to EFH or HAPC areas. The proposed changes in areas that are open to longliners and purse seiners and the separation of trollers and longliners would not result in damage to ocean or coastal habitats and none of the alternatives would result in an increase in gear loss that could affect EFH or HAPC.

Table 18. Essential Fish Habitat (EFH) and Habitat Areas of Particular Concern (HAPC) for species managed under the Fishery Ecosystem Plans.

All areas are bounded by the shoreline, and the outward boundary of the EEZ, unless otherwise indicated.

SPECIES GROUP	EFH (juveniles and adults)	EFH (eggs and larvae)	HAPC
Pelagics	water column down to 1,000 m	water column down to 200 m	water column down to 1,000 m that lies above seamounts and banks.
Bottomfish	water column and bottom habitat down to 400 m	water column down to 400 m	all escarpments and slopes between 40-280 m, and three known areas of juvenile opakapaka habitat
Seamount Groundfish	(adults only): water column and bottom from 80 to 600 m, bounded by 29°-35°N and 171°E -179°W	(including juveniles): epipelagic zone (0-200 m) bounded by 29°-35°N and 171°E - 179°W	not identified
Precious Corals	Keahole, Makapuu, Kaena, Wespac, Brooks, and 180 Fathom gold/red coral beds, and Milolii, S. Kauai and Auau Channel black coral beds	not applicable	Makapuu, Wespac, and Brooks Bank beds, and the Auau Channel
Crustaceans	bottom habitat from shoreline to a depth of 100 m	water column down to 150 m	all banks within the Northwestern Hawaiian Islands with summits less than 30 m
Coral Reef Ecosystems	water column and benthic substrate to a depth of 100 m	water column and benthic substrate to a depth of 100 m	all Marine Protected Areas identified in FEPs, all PRIAs, many specific areas of coral reef habitat (see FMP)

9.7 Other environmental consequences

None of the alternatives for any of the three proposed action topics would result in unavoidable adverse environmental consequences. None of the proposed alternatives is expected to change the intensity of fishing by the CNMI or Guam troll or longline fleets or the U.S. purse seine fleet which is not currently fishing in the U.S. EEZ surrounding the Mariana Islands. None of the proposed alternatives being considered here would result in irreversible or irretrievable commitment of resources; current and anticipated fishing would continue to be managed sustainably. There are no known controversies regarding the environmental impacts of the proposed actions.

10.0 Consistency with the MSA and Other Laws

10.1 Consistency with National Standards

Section 301 of the Magnuson-Stevens Act requires that regulations implementing any FMP, FEP or amendment be consistent with the ten national standards listed below.

National Standard 1 states that conservation and management measures shall prevent overfishing while achieving, on a continuing basis, the optimum yield from each fishery for the United States fishing industry.

The preferred alternatives considered in this amendment would establish prohibited areas for longline vessels around CNMI and purse seine vessels throughout the Mariana Archipelago and is not expected to change the intensity of fishing; therefore, it has been determined that the proposed prohibited areas would be unlikely to contribute to overfishing. The pelagic catch by the existing Mariana-based fisheries typically has annual catches averaging approximately 950,000 lb (430 mt) and this action is not expected to increase catches by the existing vessels. It is also intended to inhibit gear conflict between trollers and longliners in CNMI and to reduce the potential for stock depletion and corresponding catch competition which could arise if purse seine vessels were to fish within the EEZ waters. Minimizing pelagic fishing competition between all fisheries impacted by this action would allow local small scale pelagic fisheries to better achieve MSY, while allowing the continued establishment of a longline fishery that can operate towards an optimal yield. In addition, given that WCPO purse seine fisheries are not currently dependent on operating in the U.S. EEZ of CNMI or Guam, a prohibited area out to 200nm would have little or no effect on current yields achieved by WCPO purse seine fisheries.

National Standard 2 states that conservation and management measures shall be based upon the best scientific information available.

The preferred alternatives considered in this amendment are based on the best currently available information on pelagic fishing in EEZ waters around the Mariana Islands contained in logbook accounts from the small boat, longline, and purse seine fleets. Information from research studies

examining impacts of purse seining on other fisheries were considered when assessing potential impacts of the preferred alternatives (see Section 8.4.3) including information from a research study examining impacts of purse seining on fisheries in the Kiribati Islands (Hampton et al. 1996). This study documented the impacts of purse seine fishing on the catch rates of small scale artisanal fisheries around the islands of Kiribati. Similar small scale fisheries operate around the Mariana Archipelago, and as such it is reasonable to extrapolate the results of a study from one small Pacific Island to another. Supporting information to describe the fisheries in the U.S. EEZ around Guam and CNMI by longline vessels and troll fishery were obtained from logbooks, observer records and fisheries surveys. Other sources such as the Science Committee of the WCPFC and peer-reviewed scientific journals and reports were used in analyzing the impacts of the alternatives.

National Standard 3 states that, to the extent practicable, an individual stock of fish shall be managed as a unit throughout its range, and interrelated stocks of fish shall be managed as a unit or in close coordination.

The preferred alternatives considered in this amendment are not expected to have a significant effect on the management of fish stocks as a unit. The preferred alternatives are intended to avoid localized stock depletion of several tuna target species which are generally managed through international efforts throughout their western and central Pacific-wide range. The tuna PMUS stocks considered in this amendment range throughout western and central Pacific Ocean and the measures in waters around CNMI and Guam were considered together to account for this.

National Standard 4 states that conservation and management measures shall not discriminate between residents of different States. If it becomes necessary to allocate or assign fishing privileges among various United States fishermen, such allocation shall be (A) fair and equitable to all such fishermen; (B) reasonably calculated to promote conservation; and (C) carried out in such manner that no particular individual, corporation, or other entity acquires an excessive share of such privileges.

The preferred alternatives considered in this amendment do not discriminate between residents of different States. They establish prohibited areas in EEZ waters which are intended to prevent gear conflict and localized stock depletion, while at the same time allow for the development of longline fisheries and minimizing potential impacts.

The preferred alternative would assign ocean areas to different gear users in a fair and equitable manner. In this case, except for purse seine vessels, fishing privileges inside the U.S. EEZ would be provided to local pelagic fishing vessels to promote conservation of local stocks, maximize overall benefits, and promote conservation by optimizing the yield in terms of economic or social benefit of the available, local stock of skipjack tuna. Highly mobile purse seine vessels retain the opportunity to fish in the U.S. EEZ of the PRIA and other areas of international jurisdictions including high seas areas open to purse seining. The CNMI longline prohibited area of 0 to 30 nm would further assign privileges to small pelagic vessels fishing in nearshore waters to prevent gear conflicts with longline vessels. The initial estimate of the relative benefits and hardships imposed by the assignments are described in section 9.0, and compares the

consequences of the proposed action with alternative allocation schemes, including the status quo. No particular individual, corporation, or other entity would acquire an excessive share of such privileges under the preferred alternatives.

U.S. purse seine operators are eligible to fish under the South Pacific Tuna Treaty and may fish in other areas of the WCPO including exclusive rights to fish in U.S. EEZ waters around the Pacific Remote Islands Areas (PRIA), except those parts recently designated as a Marine National Monument (see Section 8.4.2). It might be argued that the preferred alternatives may be perceived as discriminatory towards the purse seine fishery, since the action is precautionary to avoid potential impacts as opposed to demonstrating that the troll and longline fisheries have been negatively impacted by the U.S. purse seine fishery. Indeed, any potential impacts of purse seining on local troll and longline fisheries are inferred from studies other locations where purse seine fishing has been more intensive and the impacts to troll and longline fisheries measurable (Hampton et al. 1996, Nishida 1996). The Council considered the relative size of harvests among each fishery as well as the inability of purse seine fishing to target adult fishes, in making its recommendation for the proposed purse seine exclusion zone.

The trends documented in section 8.4 which describes the current purse seine fishery in the WCPO, indicates that future international management of purse seine fishing in the WCPO is likely to become more restrictive, given the continuing concern for the overfishing of bigeye tuna. CMM 2008-01 closed the two major high seas pockets to all purse seine fishing from 2010 onwards. In addition, the Parties to the Nauru Agreement (PNA Group) have made it a condition of access to their waters that purse seine vessels would not be able to fish on the high seas between 10 degrees north latitude to 20 degrees south latitude and 170 degrees east to 150 degrees west in longitude, an area of 4,555,000 sq km. The WCPFC established closed seasons and is considering bans on the use of FADs. Increasing restrictions on purse seining are likely to make the U.S. EEZ waters more attractive to the U.S. fleet.

Given the current operational climate of the fisheries, the Council found that the preferred alternative would provide the maximum protection from competitive interactions among all three fisheries, without undue discrimination, and recognizes the greater reliance on the part of longliners to fish within EEZ waters and the total reliance on the part of the troll fishery to fish within Guam territorial waters and waters of the the U.S. EEZ waters as compared to the purse seine fleet.

National Standard 5 states that conservation and management measures shall, where practicable, consider efficiency in the utilization of fishery resources; except that no such measure shall have economic allocation as its sole purpose.

The preferred alternatives considered in this amendment include consideration of efficiency in the utilization of pelagic resources to the extent practicable by establishing prohibited areas in EEZ waters which are intended to prevent gear conflicts and localized stock depletion which could reduce fishery profitability between fishing sectors. Economic allocation is not the sole purpose of the preferred alternatives; the proposal to prohibit purse seine fishing in the U.S. EEZ around Guam and the CNMI has a conservation benefit in that it would enhance the potential for juvenile bigeye tuna in these areas to mature and reproduce.

National Standard 6 states that conservation and management action shall take into account and allow for variations among, and contingencies in, fisheries, fishery resources and catches.

The preferred alternatives considered in this amendment took into account variation among and contingencies in fisheries based out of Guam and CNMI. The preferred alternatives would provide a suitable buffer in favor of conservation by creating prohibited areas to prevent gear conflict and protect fishery stocks from localized depletion. Each fishery has its own characteristics based on gear type and island areas and these differences were considered in development of the preferred alternatives. In addition, there is little or no purse seine fishing taking place in or around the U.S. EEZ of Guam and CNMI, and establishing a clear boundary for those fisheries to fish beyond is ideal for the conservation of preserving local fishery resources, should purse seine effort migrate toward the Mariana Archipelago in the future.

National Standard 7 states that conservation and management measures shall, where practicable, minimize costs and avoid unnecessary duplication.

The preferred alternatives under consideration would not duplicate other fishery regulations. The preferred alternatives may add minor costs to existing or future longline fishing operations in EEZ waters around CNMI because they will have to travel farther than 30 nm from shore to fish. Establishment and implementation of purse seine prohibited areas would not cause fishery participants to incur any additional costs because there has been no domestic purse seining in the EEZ waters around the Mariana Archipelago. The purse seine fleet has many options for areas to fish outside of the EEZ around the Mariana Archipelago including exclusive access to U.S. EEZ waters within the PRIA.

National Standard 8 states that conservation and management measures shall, consistent with the conservation requirements of this Act (including the prevention of overfishing and rebuilding of overfished stocks), take into account the importance of fishery resources to fishing communities in order to (A) provide for the sustained participation of such communities, and (B) to the extent practicable, minimize adverse economic impacts on such communities.

National Standard 8 requires the consideration of impacts on fishery dependent communities where a fishing community is “a community which is substantially dependent on or substantially engaged in the harvesting or processing of fishery resources to meet social and economic needs, and includes fishing vessel owners, operators, and crew of U.S. fish processors that are based in such community.” Guam and CNMI are each defined as fishing communities under the MSA. The preferred alternatives considered in this amendment would be expected to beneficially impact these fishing communities by protecting Mariana Archipelago-based pelagic fishery participants which are almost exclusively small vessel trollers from potential stock depletions and reduced catch rates providing for sustained participation by these communities which are culturally and economically dependent on pelagic catches by the local fleet. In addition, longline operations would be allowed to continue in the offshore areas where they occur now, thus providing ongoing benefits to the fishing communities from that sector.

National Standard 9 states that conservation and management measures shall, to the extent practicable, (A) minimize bycatch and (B) to the extent bycatch cannot be avoided minimize the mortality of such bycatch.

The preferred alternatives considered in this amendment are not expected to increase bycatch of any species primarily because the prohibited areas are not expected to change the intensity of fishing by any of the three fleets. The prohibited areas would prevent fishing by certain sectors in portions of the EEZ including the longline sector in CNMI and the purse seine sector in the entire EEZ around the Mariana Archipelago. The purse seine sector catches a substantial amount of bycatch including juvenile bigeye tuna as described in Section 8.4 and shown in Table 11, and the preferred alternatives would remove the potential for this bycatch to occur in EEZ waters around the Mariana Archipelago.

National Standard 10 states that conservation and management measures shall, to the extent practicable, promote the safety of human life at sea.

The preferred alternatives considered are expected to increase the safety of human life at sea. Establishment of CNMI longline exclusion area would increase safety to small boats by excluding longline boats from fishing in the same waters. The small boats currently fish within 30 nm of CNMI and the longline exclusion zone would preserve this area for these vessels. Without this action small boats may be forced to seek target species farther out to sea which could cause a safety risk to the small boat participants. The proposed exclusion area affecting longline vessels in the CNMI is not expected to adversely affect the safety of the crew of these vessels because they are equipped to fish on the open ocean. Establishment of the purse seine prohibited areas is expected to increase safety to longline and small boat participants by eliminating the need for vessel operators to seek out other fishing grounds farther from shore if local stock depletions were to occur. Finally, purse seiners are not being displaced from their current operational areas, so no change to this fishery is expected to occur that would increase the risk to human safety at sea.

10.2 National Environmental Policy Act

This proposed amendment to the Council's Pelagics FEP has been written and organized to meet the requirements of the National Environmental Policy Act and thus is a consolidated document including an Environmental Assessment, as described in NOAA Administrative Order 216-6, Section 603.a.2. This document is part of the administrative record for rulemaking associated with U.S. Dept. of Commerce Regulatory Identifier No. (RIN) 0648-AW67.

10.2.1 Purpose and Need

The purpose and need for this action is described in Section 4.0.

10.2.2 Alternatives Considered

The alternatives considered for three proposed actions are described in Section 7.0. The document examines alternatives for three main topics. Topic 1) a range of longline prohibited

areas around the Northern Mariana Islands; Topic 2) a range of purse seine prohibited areas around the Northern Mariana Islands; and Topic 3) a range of purse seine prohibited areas around Guam.

10.2.3 Affected Environment

The affected environment for this action is described in Section 8.0. The main focus of the proposed action is the troll, longline, and purse seine fisheries in the waters of the U.S. EEZ surrounding Guam and the Northern Mariana Islands. Other areas of the Pacific Ocean were considered because the purse seine fleet would continue to operate in these areas under each of the alternatives.

10.2.4 Impacts of the Alternatives

The expected impacts of the alternatives considered in this action are described in Section 9.0. The analysis included a description of the baseline (no action) alternative and potential impacts of action alternatives on the fisheries and their target fish stocks, non-target fishes, bycatch, protected resources, Essential Fish Habitat (EFH) and Habitat areas of Potential Concern, and special resources or management areas. Direct, indirect, short-term, long-term, and cumulative impacts of each alternative were considered in the analysis in section 9. The impacts with respect to Environmental Justice and climate change were also addressed in section 9.

10.3 Regulatory Impact Review

Please see Appendix A for the Regulatory Impact Review of this action. To meet the requirements of Executive Order 12866 (E.O. 12866), the National Marine Fisheries Service (NMFS) requires that a Regulatory Impact Review (RIR) be prepared for all regulatory actions that are of public interest. This review provides an overview of the problem, policy objectives, and anticipated impacts of regulatory actions, and ensures that management alternatives are systematically and comprehensively evaluated such that the public welfare can be enhanced in the most efficient and cost effective way.

Based on these findings, this rule is determined to not be significant under E.O. 12866. In accordance with E.O. 12866, the following is set forth: (1) This rule is not expected to have an annual effect on the economy of more than \$100 million or to adversely affect in a material way the economy, a sector of the economy, productivity, jobs, the environment, public health or safety; or state, local or tribal governments or communities; (2) This rule is not likely to create any serious inconsistencies or otherwise interfere with any actions taken or planned by another agency; (3) This rule is not likely to materially alter the budgetary impact of entitlements, grants, user fees, or loan programs or the rights or obligations of recipients thereof; (4) This rule is not likely to raise novel or policy issues arising out of legal mandates, or the principles set forth in the Executive Order.

10.4 Administrative Procedure Act

All federal rulemaking is governed under the provisions of the Administrative Procedure Act (APA) (5 U.S.C. Subchapter II) which establishes a “notice and comment” procedure to enable public participation in the rulemaking process. Under the APA, NMFS is required to publish notification of proposed rules in the Federal Register and to solicit, consider and respond to public comment on those rules before they are finalized. The APA also establishes a 30-day wait period from the time a final rule is published until it becomes effective, with rare exceptions.

10.5 Coastal Zone Management Act

The Coastal Zone Management Act requires a determination that a recommended management measure has no effect on the land, water uses, or natural resources of the coastal zone or is consistent to the maximum extent practicable with an affected state’s enforceable coastal zone management program. A copy of this document will be submitted to the appropriate state government agencies in the Mariana Archipelago for review and concurrence with a determination that the preferred alternatives are consistent, to the maximum extent practicable, with their coastal zone management programs.

10.6 Information Quality Act

To the extent practicable, the information in this amendment complies with the Information Quality Act and NOAA standards (NOAA Information Quality Guidelines, September 30, 2002) which recognize information quality is comprised of three elements: utility, integrity, and objectivity. The information product was prepared by Council and NMFS staff based on information provided by NMFS Pacific Islands Fisheries Science Center (PIFSC) and NMFS Pacific Islands Regional Office (PIRO). The information product was reviewed by PIRO and PIFSC staff, and NMFS Headquarters (including the Office of Sustainable Fisheries). Legal review was performed by NOAA General Counsel Pacific Islands and General Counsel for Enforcement and Litigation for consistency with applicable laws, including but not limited to the Magnuson-Stevens Act, National Environmental Policy Act, Administrative Procedure Act, Paperwork Reduction Act, Coastal Zone Management Act, Endangered Species Act, Marine Mammal Protection Act, and Executive Orders 13132 and 12866.

10.7 Paperwork Reduction Act

The purpose of the Paperwork Reduction Act is to minimize the paperwork burden on the public resulting from the collection of information by or for the Federal government. It is intended to ensure the information collected under the proposed action is needed and is collected in an efficient manner (44 U.S.C. 3501(1)).

The preferred alternatives do not establish any new collection of information requirements for the purpose of the Paperwork Reduction Act.

10.8 Regulatory Flexibility Act

The Regulatory Flexibility Act (RFA) (5 U.S.C. 601 *et seq.*) requires government agencies to assess and present the impact of their regulatory actions on small entities including small businesses, small organizations, and small governmental jurisdictions. The assessment is done by preparing a Regulatory Flexibility Analysis when impacts are expected, however, the proposed alternatives, described in section 7.0, would have minimal impacts on small entities. Based on the preliminary evaluation of the economic impacts associated with the proposed alternatives (Appendix A), an initial regulatory flexibility analysis is not required and none has been prepared.

10.9 Endangered Species Act

The Endangered Species Act of 1973, as amended, (Public Law 93-205; 87 Stat. 884) prohibits the taking of any endangered species except under limited circumstances. Pursuant to Section 7 of the Endangered Species Act, in November 2006, NMFS prepared a biological opinion (BiOp) on the effects of the U.S. purse seine fishery in the WCPO on sea turtles and whales. The BiOp concluded that the fishery is not likely to jeopardize the continued existence of any of the threatened or endangered species found in the area or adversely modify their critical habitat.

A 2001 NMFS BiOp established incidental take statements for each longline fishery in Guam and CNMI. Although no longline fishery existed at the time, NMFS utilized fishery information from other sources to estimate incidental take and mortality of ESA listed species. The 2001 BiOp estimated annual take of 3 hardshell turtles and 1 leatherbackback turtle for each of the longline fisheries in Guam and CNMI. In addition, the estimated mortality level is 1 hardshell turtle in each of the longline fisheries in Guam and CNMI. To date there have been no records of interactions with ESA listed species.

In addition, the troll fisheries around CNMI and Guam were included in NMFS' 2009 biological opinion on pelagic fisheries of the western Pacific which concluded that continued operations were not likely to jeopardize the continued existence of any listed sea turtles or marine mammals. The 2009 BiOp considered potential stressors and impacts related to vessel collisions, hooking and entanglement with fishing gear, human disturbance, exposure to vessel waste, direct and indirect competition with foraging grounds and exposure to marine mammal deterrents. The 2009 BiOp did consider that a small potential exists that the trolling gear may incidentally hook or entangle a sea turtle or marine mammal; however, NMFS and the Council consider this type of interaction extremely rare and, therefore, unlikely to adversely affect listed species.

Section 8.5 describes the threatened and endangered species known to occur in the WCPO and Section 9.1.3, 9.2.3, and 9.3.3 describe the potential impacts the preferred alternatives may have on these listed species. There are no known interactions between seabirds and any of the Mariana Archipelago pelagic fisheries (WPRFMC 2007b). Based on the gear types used and the low likelihood of fishery interactions occurring under the preferred alternatives, the Council believes that the preferred alternatives will not jeopardize or adversely affect any populations or habitats of species listed as endangered or threatened under the ESA.

10.10 Marine Mammal Protection Act

The CNMI and Guam troll fisheries are classified as Category III under Section 118 of the Marine Mammal Protection Act (MMPA) (62 FR 28657, 27 May 1997), meaning that they have been determined by NMFS to have a remote likelihood of, or no known incidental mortality and serious injury of marine mammals (50 CFR 229.2). Vessel owners and crew that are engaged only in Category III fisheries may incidentally take marine mammals without registering or receiving an Authorization Certificate under the MMPA, but they are required to: 1) report all incidental mortality and injury of marine mammals to NMFS, 2) immediately return to the sea with minimum of further injury any incidentally taken marine mammal, 3) allow vessel observers if requested by NMFS, and 4) comply with guidelines and prohibitions under the MMPA when deterring marine mammals from gear, catch, and private property (50 CFR 229.5, 229.6, 229.7).

The List of Fisheries for 2010, published pursuant to 50 CFR 229, classifies the CNMI and Guam longline and U.S. purse seine fleets as Category II for all fisheries under Section 118 of the MMPA (74 FR 58859; November 16, 2009). The list includes mammal species and stocks incidentally killed or injured in each fishery based on observer data, logbook data, stranding reports, disentanglement network data, and fisher reports. The List of Fisheries in 2010 for the South Pacific Tuna Fisheries lists the marine mammal species killed or injured as “undetermined” until additional information on marine mammal populations and fishery interactions becomes available. The “undetermined” status has not changed from previous years List of Fisheries.

Please see Section 8.5.2 of this document for descriptions of marine mammals found around the Mariana Archipelago. Section 9.0 provides an analysis of the anticipated impacts on these species under each of the alternatives considered by the Council. Based on the gear types used and the low likelihood of fishery interactions occurring under the preferred alternatives, the Council believes that the preferred alternatives will not adversely affect any marine mammal populations or habitats.

11.0 Proposed Regulations

For the reasons set out in the preamble, 50 CFR chapter VI is proposed to be amended as follows:

PART 665--FISHERIES IN THE WESTERN PACIFIC

1. The authority citation for part 665 continues to read as follows:

Authority: 16 U.S.C. 1801 et seq.

2. In § 665.800, add a definition of “Purse seine” in alphabetical order to read as follows:

§ 665.800 Definitions.

* * * * *

Purse seine means a floated and weighted encircling net that is closed by means of a drawstring threaded through rings attached to the bottom of the net.

* * * * *

3. In § 665.802, revise paragraphs (v), (w), and (xx) to read as follows:

§ 665.802 Prohibitions.

* * * * *

(v) Use longline gear to fish within a longline fishing prohibited area in violation of § 665.806, except as allowed pursuant to an exemption issued under §§ 665.17 or 665.807.

(w) Use a purse seine to fish within a purse seine fishing prohibited area in violation of § 665.806, except as allowed pursuant to an exemption issued under § 665.17.

* * * * *

(xx) Use a large vessel to fish for western Pacific Pelagic MUS within an American Samoa large vessel prohibited area in violation of § 665.806, except as allowed pursuant to an exemption issued under §§ 665.17 or 665.818.

* * * * *

4. Revise § 665.806 to read as follows:

§ 665.806 Prohibited area management.

(a) Longline fishing prohibited areas. Longline fishing is prohibited in the longline fishing prohibited areas as defined in paragraphs (a)(1) through (a)(4) below.

(1) NWHI protected species zone. The NWHI protected species zone is the portion of the EEZ within 50 nm of the center geographical positions of certain islands and reefs in the NWHI, as follows:

Name	N. lat.	W. long.
Nihoa Island	23° 05'	161° 55'
Necker Island	23° 35'	164° 40'
French Frigate Shoals	23° 45'	166° 15'
Gardner Pinnacles	25° 00'	168° 00'
Maro Reef	25° 25'	170° 35'
Laysan Island	25° 45'	171° 45'
Lisianski Island	26° 00'	173° 55'
Pearl and Hermes Reef	27° 50'	175° 50'
Midway Island	28° 14'	177° 22'
Kure Island	28° 25'	178° 20'

Where the areas are not contiguous, parallel lines drawn tangent to and connecting those semicircles of the 50-nm areas that lie between Nihoa Island and Necker Island, French Frigate Shoals and Gardner Pinnacles, Gardner Pinnacles and Maro Reef, and Lisianski Island and Pearl and Hermes Reef, delimit the remainder of the NWHI longline protected species zone.

(2) Main Hawaiian Islands (MHI). (i) From February 1 through September 30 each year, the MHI longline fishing prohibited area is the portion of the EEZ around Hawaii bounded by straight lines connecting the following coordinates in the order listed:

Point	N. lat.	W. long.
A	18° 05'	155° 40'
B	18° 20'	156° 25'
C	20° 00'	157° 30'
D	20° 40'	161° 40'

E	21° 40'	161° 55'
F	23° 00'	161° 30'
G	23° 05'	159° 30'
H	22° 55'	157° 30'
I	21° 30'	155° 30'
J	19° 50'	153° 50'
K	19° 00'	154° 05'
A	18° 05'	155° 40'

(ii) From October 1 through the following January 31 each year, the MHI longline fishing prohibited area is the portion of the EEZ around Hawaii bounded by straight lines connecting the following coordinates in the order listed:

Point	N. lat.	W. long.
A	18° 05'	155° 40'
L	18° 25'	155° 40'
M	19° 00'	154° 45'
N	19° 15'	154° 25'
O	19° 40'	154° 20'
P	20° 20'	154° 55'
Q	20° 35'	155° 30'
R	21° 00'	155° 35'
S	22° 30'	157° 35'
T	22° 40'	159° 35'
U	22° 25'	160° 20'
V	21° 55'	160° 55'
W	21° 40'	161° 00'
E	21° 40'	161° 55'
D	20° 40'	161° 40'
C	20° 00'	157° 30'
B	18° 20'	156° 25'
A	18° 05'	155° 40'

(3) Guam. The Guam longline fishing prohibited area is the portion of the EEZ around Guam bounded by straight lines connecting the following coordinates in the order listed:

Point	N. lat.	E. long.
A	14° 25'	144° 00'
B	14° 00'	143° 38'
C	13° 41'	143° 33' 33"
D	13° 00'	143° 25' 30"
E	12° 20'	143° 37'
F	11° 40'	144° 09'
G	12° 00'	145° 00'
H	13° 00'	145° 42'
I	13° 27'	145° 51'

(4) CNMI. The CNMI longline fishing prohibited area is the portion of the EEZ around the CNMI bounded by straight lines connecting the following coordinates in the order listed:

Point	N. lat.	E. long.
A	14° 00'	144° 34'
B	15° 49'	145° 29'
C	16° 21'	145° 06'
D	17° 03'	145° 22'
E	19° 07'	145° 09'
F	20° 39'	144° 19'
G	21° 04'	145° 06'
H	19° 19'	146° 04'
I	16° 00'	146° 32'
J	13° 32'	145° 32'
A	14° 00'	144° 34'

(b) American Samoa large vessel prohibited areas. A large vessel of the United States may not be used to fish for western Pacific pelagic MUS in the American Samoa large vessel prohibited areas as defined in paragraphs (b)(1) and (b)(2) of this section, except as allowed pursuant to an exemption issued under § 665.818.

(1) Tutuila Island, Manua Islands, and Rose Atoll (AS-1). The Tutuila Island, Manua Islands, and Rose Atoll large vessel prohibited area is the portion of the EEZ around American Samoa enclosed by straight lines connecting the following coordinates:

Point	S. lat.	W. long.
AS-1-A	13°30'	167°25'
AS-1-B	15°13'	167°25'
and from Point AS-1-A westward along the latitude 13°30' S. until intersecting the U.S. EEZ boundary with Samoa, and from Point AS-1-B westward along the latitude 15°13' S. until intersecting the U.S. EEZ boundary with Samoa.		

(2) Swains Island (AS-2). The Swains Island large vessel prohibited area is the portion of the EEZ around American Samoa enclosed by straight lines connecting the following coordinates:

Point	S. lat.	W. long.
AS-2-A	11°48'	171°50'
AS-2-B	11°48'	170°20'
and from Point AS-2-A northward along the longitude 171°50' W. until intersecting the U.S. EEZ boundary with Tokelau, and from Point AS-2-B northward along the longitude 170°20' W. until intersecting the U.S. EEZ boundary with Tokelau.		

(c) Purse seine fishing prohibited areas. Purse seine fishing is prohibited in the EEZ around Guam and the CNMI.


5. Revise the section heading in § 665.807 to read as follows:

§ 665.807 Exemptions for Hawaii longline fishing prohibited areas; procedures.

* * * * *

6. Remove and reserve § 665.817 as follows:

§ 665.817 [Reserved]



12.0 References

- Ainley, D.G., Telfer, T.C., and Reynolds, M.H. 1997. Townsend's and Newell's shearwater: *Puffinus auricularis*. Birds of North America. 297 p.
- Balazs, G. H., P. Craig, B.R. Winton, and R.k. Miya. 1994. Satellite telemetry of green turtles nesting at French Frigate Shoals, Hawaii, and Rose Atoll, American Samoa. In: Bjorndal, K. A., A.B. Bolten, D.A. Johnson, and P.J. Eliazar. (Eds.), Proc. 14th Ann. Symp. on Sea Turtle Biology and Conservation. NOAA Tech. Memo. NMFS SEFSC-351., pp. 184–187.
- Balazs, G. H. and M. Chaloupka. 2004. Thirty-year recovery trend in the once depleted Hawaiian green sea turtle stock. Biological Conservation. 117:491–498.
- Bigelow, K. 1999. Fisheries and stock assessment of yellowfin tuna in the Western and Central Pacific. Second meeting of the Interim Scientific Committee for Tuna and Tuna-like species in the North Pacific Ocean, Honolulu. 15-23 January 1999, ISC2/99/PLEN/17.
- Bjorndal, K. A. 1997. Foraging ecology and nutrition of sea turtles. In P. L. Lutz and J. A. Musick (Eds.), The biology of sea turtles. Boca Raton, FL: CRC Press.
- Brodziak, J. and K. Piner. 2010. Model averaging and probable status of North Pacific striped marlin, *Tetrapturus audax*. Can. J. Fish. Aquat. Sci. 67:793-805.
- Calambokidis, J., G. Steiger, J. Straley, T. Quinn II, L. Herman, S. Cerchio, D. Salden, M. Yamaguchi, F. Sato, J. Urban, R. Jacobsen, O. von Ziegesar, K. Balcomb, C. Gabriele, M. Dahlheim, N. Higashi, S. Uchida, J. Ford, Y. Miyamura, P. de Guevara, S. Mizroch, L. Schlender, and K. Rasmussen. 1997. Abundance and population structure of Humpback whales in the North Pacific Basin (Final Report). Cascadia Research Collective. Contract #50ABNF500113 eport.
- Chaloupka, M. and C. Limpus. 2001. Trends in the abundance of sea turtles resident in southern Great Barrier Reef waters. Biological Conservation 102:235–249.
- Chan, E. and H. Liew. 1989. Charting the movements of a sea giant. In Research News, Universiti Pertanian Malaysia 3 (4): 7-8.
- Chan, E. and H. Liew. 1996. Decline of the leatherback population in Terengganu, Malaysia, 1956–1995. Chelonian Conservation Biology 2(2): 196–203.
- Cliffton, K., D.O. Cornejo, and R. S. Felger. 1982. Sea turtles of the Pacific coast of Mexico. In Bjorndal, K.A. (Ed.), Biology and conservation of sea turtles (pp. 199–209). Washington, DC: Smithsonian Institution Press.
- Cummings, V. 2002. Sea turtle conservation in Guam. In: I. Kinan (Ed.) Proc. Western

Pacific Sea Turtle Cooperative Research and Management Workshop, Western Pacific Regional Fishery Management Council, Honolulu, Hawaii, pp. 37-38.

Delgado de Molina, A., J. Ariz, C. Santana, and B. Sotillo. 2010. Analysis of the Catch Rate of Juvenile Bigeye Depending On the Depth of the Purse-seine Net Used by the Tropical Fleet. Western & Central Pacific Fishery Commission, Scientific Committee Sixth Regular Session Nuku'alofa, Tonga, 10-19 August 2010, WCPFC-SC6-2010/FT- WP-03.

Dutton, P., B. Bowen, D. Owens, A. Barragán, and S. Davis. 1999. Global phylogeography of the leatherback turtle (*Dermochelys coriacea*). *Journal of Zoology*. 248:397–409.

Eckert, K. L. 1993. The biology and population status of marine turtles in the North Pacific Ocean (NOAA Tech. Memo, NOAA-TM-NMFS-SWFSC-186, 156 pp.). La Jolla, CA: National Marine Fisheries Service, Southwest Region.

Eckert, S. A. 1998. Perspectives on the use of satellite telemetry and other electronic technologies for the study of marine turtles, with reference to the first year-long tracking of leatherback sea turtles, p. 294. In: Proceedings of the Seventeenth 21 Annual Sea Turtle Symposium. S. P. Epperly and J. Braun (Eds.). NOAA Technical Memorandum NMFS-SEFSC-415, Miami.

Eckert, K.L. and S.A. Eckert. 1988. Pre-reproductive movements of leatherback turtles (*Dermochelys coriacea*) nesting in the Caribbean. *Copeia* 1988(2):400-406.

Eldredge, L.G. 2003. The marine reptiles and mammals of Guam. *Micronesica* 35-36: 653-60.

Falanruw, M.V.C., M. McCoy, and Namlug. 1975. Occurrence of ridley sea turtles in the Western Caroline Islands. *Micronesica* (11)a:151-152.

FAO. 2005. Second meeting of the Technical Advisory Committee of the FAO project Management of tuna fishing capacity: conservation and socio-economics, March 15-18, 2004, Madrid Spain. Edited by W.H. Bayliff, J.I. de Leiva Moreno and J. Majkowski. Food and Agriculture Organization of the United Nations, Rome, 2005.

Forney, K., J. Barlow, M. Muto, M. Lowry, J. Baker, G. Cameron, J. Mobley, C. Stinchcomb, and J. Carreta. 2000. Draft U.S. Pacific Marine Mammal Stock Assessments: 2000. NMFS Southwest Fisheries Science Center: La Jolla.

Gillett, R. and A. Lewis. 2003. A Survey of Purse Seine Fishing Capacity in the Western and Central Pacific Ocean, 1988-2003. 17th Meeting of the Standing Committee on Tuna and Billfish Working Paper INF-FTWG-1c.

Hamnett, M. and W. Pintz. 1996. The contribution of tuna fishing and transshipment to the economies of American Samoa, the Commonwealth of the Northern Mariana Islands, and Guam. Pelagic Fisheries Research Program. SOEST 96-05. JIMAR Contribution 96-303.

Hampton J., T. Lawson, P. Williams, and J. Sibert. 1996. Interactions between small-scale fisheries in Kiribati and the industrial purse seine fishery in the western and central Pacific Ocean. 183-223. In: R. Shomura, J. Majkowski, and R. Harman (Eds.). Status of Interactions of Pacific Tuna fisheries in 1995. FAO Fisheries Tech. Paper 365. Rome.

Hampton, J., A. Langley, S. Harley, P. Kleiber, Y. Takeuchi, and M. Ichinokawa. 2005. Estimates of sustainable catch and effort levels for target species and the impacts on stocks of potential management measures. Oceanic Fisheries Programme. Secretariat of the Pacific Community. Noumea, New Caledonia. July 2005. 34 pp.

Hampton, J., A. Langley, and P. Kleiber. 2006. Stock Assessment of Yellowfin Tuna in the Western and Central Pacific Ocean, Including an Analysis of Management Options. WCPFC-SC2-2006/SA WP-1. Scientific Committee Second Regular Session, 7-18 August 2006, Manila, Philippines. July 2006. 99 pp.

Hampton, J., A. Langley, and P. Kleiber. 2006. Stock Assessment of Bigeye Tuna in the Western and Central Pacific Ocean, Including an Analysis of Management Options. WCPFC-SC2-2006/SA WP-2. Scientific Committee Second Regular Session, 7-18 August 2006, Manila, Philippines. July 2006. 91 pp.

Hampton, J. 2002. Stock assessment of skipjack tuna in the western and central Pacific Ocean. 15th Meeting of the Standing Committee on Tuna and Billfish working paper SKJ-1.

Harley, S., J. Hampton, S. Hoyle, A. Langley, and P. Kleiber. 2009. Stock assessment of bigeye tuna in the western and central Pacific Ocean. The Western and Central Pacific Fisheries Commission, Scientific Committee Fifth Regular Session. WCPFC-SC5-2009/SA-WP-4. August 2009. 98 pp.

Harrison, C.S. 1990. Seabirds of Hawaii: natural history and conservation. Cornell Univ. Press, Ithaca, New York. 249 p.

Higuchi, W. 2007. Pre-war Japanese fisheries in Micronesia: Focusing on bonito and tuna fishing in the Northern Mariana Islands. Immigration Studies No.3, pp. 49-68. Center for Migration Studies, University of the Ryukyus, Nishihara, Okinawa, Japan.

Hill P. and D. DeMaster. 1999. Alaska marine mammal stock assessments 1999. National Marine Mammal Laboratory, NMFS Alaska Fisheries Science Center. Seattle, WA.

Hill, P., D. DeMaster, and R. Small. 1997. Alaska Marine Mammal Stock Assessments, 1996. U.S. Pacific Marine Mammal Stock Assessments: 1996. U.S. Dept. of Commerce, NOAA, Tech. Memo., NMFS, NOAA-OTM-NMFS-AFSC-78. 149p.

Hodge, R. and B. Wing. 2000. Occurrence of marine turtles in Alaska waters: 1960-1998. Herpetological Review. 31:148-151.

Holmes, N. 2009. Kauai Endangered Seabird Recovery Project, pers. comm., June 2009

Horwood, J. 1987. *The Sei Whale: Population Biology, Ecology and Management*. Croom Helm. London.

Hoyle, S.D. and M. N. Maunder. 2005. Status of yellowfin tuna in the Eastern Pacific Ocean in 2004 and outlook for 2005. IATTC Stock Assessment Report 6. Inter-American Tropical Tuna Commission. La Jolla, California. 102 pp.

Hoyle, S., P. Kleiber, N. Davies, S. Harley, and J. Hampton. 2010. Stock Assessment of Skipjack Tuna in the Western And Central Pacific Ocean. Western and Central Pacific Fishery Commission, Sixth Scientific Committee, Tonga, WCPFC-SC6-2010/SA-WP-10 rev.1.

Intergovernmental Panel on Climate Change (IPCC). 2007a. *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. [Solomon, S., D. Qin, M. Manning, M. Marquis, K.B. Averyt, M. Tignor, and H.L. Miller (Eds.)]. Cambridge University Press, NY.

IPCC 2007b *Climate Change 2007: Working Group II: Impacts, Adaptation, and Vulnerability. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. [M.L. Parry, O.F. Canziani, J.P. Palutikof, P.J. van der Linden, and C.E. Handon (Eds.)]. Cambridge University Press, NY.

ISC. 2010. 10th Plenary Report of the Billfish Working Group Workshop. International Scientific Committee for Tuna and Tuna-like Species in the North Pacific Ocean, 15-22 April 2010, Hakodate, Hokkaido, Japan

Kleiber, P.J., J. Hampton, M.G. Hinton and Y. Uozumi. 2002. Update on blue marlin stock assessment. 15th Meeting of the Standing Committee on Tuna and Billfish, July 22, 2002. Honolulu, Hawaii.

Kolinski, S. 2001. Sea turtles and their marine habitats at Tinian and Aguijan, with projections on resident turtle demographics in the southern arc of the Commonwealth of the Northern Mariana Islands. NMFS Southwest Fisheries Science Center Administrative Report H-01—6C. 93 pp.

Landis, C. 1965. Research: new high pressure research animal? *Undersea Tech* 6: 21.

Langley, A., M. Ogura, and J. Hampton. 2003. Stock assessment of skipjack tuna in the western and central Pacific Ocean. 16th Meeting of the Standing Committee on Tuna and Billfish. SCTB16 Working Paper, June 2003. 43 pp.

Langley, A., P. Williams, and J. Hampton. 2006. *The Western and Central Pacific Tuna Fishery. 2005. Overview and Status of Stocks*. Ocean Fisheries Programme, Secretariat of the Pacific Community. Noumea, New Caledonia.

Langley, A., J. Hampton, P. Kleiber and S. Hoyle. 2007. Stock assessment of yellowfin tuna in the western and central Pacific Ocean, including an analysis of management options. The

Western and Central Pacific Fisheries Commission, Scientific Committee Third Regular Session. WCPFC-SC3-SA WP-1. July 2007. 36 pp.

Langley, A. and J. Hampton. 2008. Stock assessment of skipjack tuna in the western and central Pacific Ocean. The Western and Central Pacific Fisheries Commission, Scientific Committee Fifth Regular Session. WCPFC-SC4-2008/SA-WP-4. August 2008. 75 pp.

Langley, A., S. Harley, S. Hoyle, N. Davies, J. Hampton, J., and P. Kleiber. 2009. Stock assessment of yellowfin tuna in the western and central Pacific Ocean. The Western and Central Pacific Fisheries Commission, Scientific Committee Fifth Regular Session. WCPFC-SC5-2009/SA-WP-03. August 2009. 125 pp.

Lawson, T.A. 1997. Estimation of bycatch in central and western Pacific tuna fisheries: preliminary results. Oceanic Fisheries Programme Internal Report No. 33. South Pacific Commission, Noumea, New Caledonia. 27 pp.

Limpus, C. J. 1982. The status of Australian sea turtle populations. In K. A. Bjorndal (Ed.), *Biology and conservation of sea turtles*. Washington, DC: Smithsonian Institution Press

Limpus, C. 1992. The hawksbill turtle, *Eretmochelys imbricata*, in Queensland: Population structure within a southern Great Barrier Reef feeding ground. *Wildlife Research* (19): 489–506.

Limpus, C. J., and D. Reimer. 1994. The loggerhead turtle, *Caretta caretta*, in Queensland: A population in decline. In R. James (Compiler). *Proceedings of the Australian Marine Turtle Conservation Workshop: November 14–17, 1990* Canberra, Australia: Australian Nature Conservation Agency.

Marquez, M. 1990. Sea turtles of the world. An annotated and illustrated catalogue of sea turtle species known to date. *FAO species Catalog. FAO Fisheries Synopsis* 11(125). 81pp.

Maunder, M.N., and S.D. Hoyle. 2005. Status of Bigeye Tuna in the Eastern Pacific Ocean in 2004 and Outlook for 2005. IATTC Working Group on Stock Assessment Document SAR-06-07B.

McKeown, A. 1977. *Marine turtles of the Solomon Islands*. Honiara: Solomon Islands: Ministry of Natural Resources, Fisheries Division.

Meylan, A. 1985. The role of sponge collagens in the diet of the Hawksbill turtle, *Eretmochelys imbricata*. In A. Bairati and R. Garrone, (Eds.), *Biology of invertebrate and lower vertebrate collagens*. New York: Plenum Press.

Meylan, A. 1988. Spongivory in hawksbill turtles: A diet of glass. *Science* 239:393–395.

Meylan, A. 1999. International movements of immature and adult hawksbill turtles (*Eretmochelys imbricata*) in the Caribbean Region. *Chelonian Conservation and Biology* 3:189-194.

Moloney, B. 2004. Summarized fishery information for evaluating Alternatives for bigeye and yellowfin tuna. 1. Review of fleet capacity, catch, and effort of the longline fishery in the western and central Pacific Ocean, 17th Meeting of the Standing Committee on Tuna and Billfish, 9-18 August 2004, Majuro, Marshall Islands. Working Paper GEN-2. 31 pp.

Molony, B. 2005. Estimates of the mortality of non-target species with an initial focus on seabirds, turtles and sharks. A report of the Oceanic Fisheries Programme, Secretariat of the Pacific Community, Noumea, New Caledonia.

NMFS (National Marine Fisheries Service). 1998. Biological opinion on the fishery management plan for the pelagic fisheries of the Western Pacific Region: Hawaii Central North Pacific longline fishery. La Jolla, CA: National Marine Fisheries Service, Southwest Region.

NMFS and USFWS. 1998a. (National Marine Fisheries Service and U.S. Fish and Wildlife Service) Recovery Plan for U.S. Pacific Populations of the green turtle (*Chelonia mydas*). National Marine Fisheries Service. Silver Spring, MD. 84 pp.

NMFS and USFWS. 1998b. Recovery Plan for U.S. Populations of the hawksbill turtle (*Eretmochelys imbricata*). National Marine Fisheries Service. Silver Spring, MD. 82 pp.

NMFS. 2001. Final Environmental Impact Statement for the Fishery Management Plan for Pelagic Fisheries of the Western Pacific Region.

NMFS. 2006. Final Biological opinion under Section 7 of the Endangered Species Act on the effects of the U.S. tuna purse seine fishery in the western and central Pacific Ocean on listed sea turtles and marine mammals. National Marine Fisheries Service Pacific Islands Regional Office. Issued November 1, 2006. 185 pp.

NMFS. 2007. 2007 Annual Report to the Western and Central Pacific Fisheries Commission, Part 1. Information on Fisheries, Research, and Statistics (for 2006). PIFSC Data Report DR-07-005 Issued 13 July 2007, Revised August 1, 2007. 40 pp.

NMFS. 2008. 2008 Annual Report to the Western and Central Pacific Fisheries Commission, Part 1. Information on Fisheries, Research, and Statistics (for 2007). PIFSC Data Report DR-08-008 Issued 8 August 2008, Revised 25 September 2008. 40 pp.

NMFS. 2009 Annual Report to the Western and Central Pacific Fisheries Commission, Part 1. Information on Fisheries, Research, and Statistics (for 2008), WCPFC-SC5-AR/CCM-26. August 2009. 35 pp.

OFP (Oceanic Fisheries Programme).1998. Mariana Islands 1998 Pelagic Fisheries Assessment. Oceanic Fisheries Programme Country Report No. 12. Secretariat of the Pacific Community, Noumea, New Caledonia. 87 pp.

OFP. 2007. Preliminary review of the Western and Central Pacific Ocean purse seine fishery 2006. Secretariat of the Pacific Community. February 2007. 23 pp.

OFP. 2007b. Preliminary review of the Western and Central Pacific Ocean purse seine fishery 2007. Secretariat of the Pacific Community. February 2007. 23 pp.

Osgood, K.E. (Ed.). 2008. Climate Impacts on U.S. Living Marine Resources: National Marine Fisheries Service Concerns, Activities and Needs. U.S. Department of Commerce, National Oceanic and Atmospheric Administration, National Marine Fisheries Service. NOAA Technical Memorandum NMFS-F/SPO-89. August 2008.

URL: <http://spo.nmfs.noaa.gov/tm/TM%20SPO%2089.pdf>

Plotkin, P.T. 1994. The migratory and reproductive behavior of the olive ridley, *Lepidochelys olivacea* (Eschscholtz, 1829), in the eastern Pacific Ocean. Ph.D. Thesis, Texas A&M Univ., College Station, TX.

Polovina, J.J., G.T. Mitchum, N.E. Graham, M.P. Craig, E.E. DeMartini, and E.N. Flint. 1994. Physical and biological consequences of a climate event in the central North Pacific. *Fisheries Oceanography*, 3(1):15–21.

Randall, R.H., R.T. Tsuda, R.S. Jones, M.J. Gawel, J.A. Chase, and R. Rechebei. 1975. Marine biological survey of the Cocos barrier reefs and enclosed lagoon. University of Guam Marine Laboratory Technical Report 17. 160 pp.

Reeves, R., S. Leatherwood, G. Stone, and L. Eldridge. 1999. Marine mammals in the area served by the South Pacific Regional Environment Programme (SPREP). South Pacific Regional Environment Programme: Apia, Samoa. 48 pp.

Rice, D. 1989. Sperm whale *Physeter macrocephalus*. Academic Press. 442 pp.

Sarti, L., S. Eckert, N. Garcia, and A. Barragan. 1996. Decline of the world's largest nesting assemblage of leatherback turtles. *Marine Turtle Newsletter* 74:2–5.

Scavia, D., J. C. Field, D. F. Boesch, R. W. Buddemeier, V. Burkett, D. R. Cayan, M. Fogarty, M. A. Harwell, R. W. Howarth, C. Mason, D. J. Reed, T. C. Royer, A. H. Sallenger, and J. G. Titus. 2002. Climate change impacts on U.S. coastal and marine ecosystems. *Estuaries*, 25:149-164.

Sibert, J. and J. Hampton. 2003. Mobility of tropical tunas and the implications for fisheries management. *Marine Policy* 27: 87-95.

Sincock, J.L., and Swedberg, G.E. 1969. Rediscovery of the nesting grounds of Newell's Manx Shearwater (*Puffinus puffinus newelli*), with initial observations. The Elepaio 29(12). Hawai'i Audubon Society. Honolulu, Hawai'i. Reprinted from the January 1969 (Vol.71, No.1, pp.69-71) issue of The Condor. The Cooper Ornithological Society.

SPC (South Pacific Commission). 1999. Western and Central Pacific Fisheries Commission Tuna Fishery Yearbook 1998. Oceanic Fisheries Programme. Noumea, New Caledonia.

SPC. 2004. Western and Central Pacific Fisheries Commission Tuna Fishery Yearbook 2003. Oceanic Fisheries Programme. Noumea, New Caledonia.

SPC. 2005. Western and Central Pacific Fisheries Commission Tuna Fishery Yearbook 2004. Oceanic Fisheries Programme. Noumea, New Caledonia.

SPC. 2006. Western and Central Pacific Fisheries Commission Tuna Fishery Yearbook 2005. Oceanic Fisheries Programme. Noumea, New Caledonia.

SPC. 2010. Non-Target Species Interactions with the Tuna Fisheries of the Western and Central Pacific Ocean. Noumea, New Caledonia. WCPFC- SC6-2010/EB-IP-8. Scientific Committee Sixth Regular Session. Nuku'alofa, Tonga 10- 19 August 2010.

Spear, L.B., Ainley, D.G., Nur, N., and Howell, S.N.G. 1995. Population Size and Factors Affecting At-Sea Distributions of Four Endangered Procellariids in the Tropical Pacific. The Condor, 97(3): 613-638.

Spotila J., A. Dunham, A. Leslie, A. Steyermark, P. Plotkin, and F. Paladino. 1996. Worldwide population decline of *Dermodochelys coriacea*: Are leatherback turtles going extinct? Chelonian Conservation Biology 2(2):209–222.

Spotila, J.R., R.D. Reina, A.C. Steyermark, P.T. Plotkin, and F.V. Paladino. 2000. Pacific leatherback turtles face extinction. Nature 405:529-530.

Starbird, C. H. and M. M. Suarez. 1994. Leatherback sea turtle nesting on the north Vogelkop coast of Irian Jaya and the discovery of a leatherback sea turtle fishery on Kei Kecil Island. Fourteenth Annual Symposium on Sea Turtle Biology and Conservation (p. 143). March 1–5, 1994, Hilton Head, South Carolina.

Sturman, A. P. and H. McGowan. 2003. Climate. In: M. Rapaport (Ed.), The Pacific Islands: Environment and Society. Honolulu, HI: The Bess Press.

Trianni, M.S. 1998. Summary and further analysis of the nearshore reef fishery of the Northern Mariana Islands. CNMI Division of Fish and Wildlife. Technical Report 98-02. 64 pp.

Troeng, S. and E. Rankin. 2005. Long-term conservation efforts contribute to positive green turtle (*Chelonia mydas*) nesting trend at Tortuguero, Costa Rica. Biological Conservation 121:111–116.

- U.S. Fish and Wildlife Service (USFWS). 1983. Hawaiian Dark-rumped Petrel and Newell's Manx Shearwater: Recovery Plan. Portland, OR. 57 p.
- Van Dam, R.P. and C.E. Diez. 1997. Diving behavior of immature hawksbill turtles (*Eretmochelys imbricata*) in a caribbean reef habitat. *Coral Reefs*.16:133–138.
- Walther, G., E. Post, P. Convey, A. Menzel, C. Parmesan, T.J.C. Beebee, J. Fromentin, O. Hoegh-Guldberg, and F. Bairlein. 2002. Ecological responses to recent climate change. Review Article. *Nature*, 416: 389-395.
- Ward P., N. Tsirbasm, and B. Kearney. 1998. Getting Science into Regional Fishery Management A Global View. Bureau of Rural Sciences: Kingston, Australia.
- Warham, J. 1990. The shearwater *Fenus puffinus*. In *The petrels: Their ecology and breeding system* (pp. 157–170). San Diego, CA: Academic Press.
- WCPFC. 2005. Report of the First Regular Session of the Scientific Committee, Noumea, New Caledonia, 8-19 August, 2005, Western and Central Pacific Fisheries Commission.
- WCPFC. 2006. National Tuna Fishery Report Federated States of Micronesia. Second Regular Session of the Scientific Committee of the Western and Central Pacific Fisheries Commission.
- WCPFC. 2006b. Western and Central Pacific Fisheries Commission Tuna Fishery Yearbook 2005. Edited by T.A. Lawson. Western and Central Pacific Fisheries Commission.
- WCPFC. 2007. Western and Central Pacific Tuna Bulletin. July 2007. Western and Central Pacific Fisheries Commission. 40 pp.
- WCPFC. 2008. Western and Central Pacific Tuna Bulletin. July 2008. Western and Central Pacific Fisheries Commission. 44 pp.
- WCPFC. 2008b. Western and Central Pacific Fisheries Commission Tuna Fishery Yearbook 2007. Edited by T.A. Lawson. Western and Central Pacific Fisheries Commission. 2008 (Revised February 2009)
- WCPFC. 2009. Western and Central Pacific Fisheries Commission Scientific Committee Fifth Regular Session – Estimates of Annual Catches in the WCPFC Statistical Area. WCPFC-SC5-2009/ST IP-1, 28 pp.
- Wetherall, J. A. 1993. Pelagic distribution and size composition of turtles in the Hawaii longline fishing area. In G. H. Balazs and S. G. Pooley (Eds.), *Research plan to assess marine turtle hooking mortality: Results of an expert workshop held in Honolulu, Hawaii, November 16–18, 1993*. SWFSC Administrative Report H-93-18.
- Wiles, G., J. Bart, R. Beck, Jr. and C. Aguon. 2003. Impacts of the brown tree snake: patterns of decline and species persistence in Guam's avifauna. *Conservation Biology* 17(5): 1350-1360.

Williams, P. and C. Reid. 2005. Overview of tuna fisheries in the western and central Pacific Ocean, including economic conditions – 2004. 1st Meeting of the Scientific Committee of the Western and Central Pacific Fisheries Commission. WCPFC-SC1, Noumea, New Caledonia, 8 – 19 August 2005. GN WP-1. 39 pp.

Williams, P. and C. Reid. 2007. Overview of tuna fisheries in the western and central Pacific Ocean, including economic conditions—2006. Western and Central Pacific Fisheries Commission Scientific Committee Third Regular Session, August 2006. WCPFC-SC3-2007/GN WP-1. 45 pp.

Williams, P. and P. Terawasi. 2009. Overview of tuna fisheries in the western and central Pacific Ocean, including economic conditions. Western and Central Pacific Fisheries Commission Scientific Committee Fifth Regular Session, August 2009. WCPFC-SC5-2009/GN WP-1, 46 pp.

WPRFMC. 2007a. Pelagic Fisheries of the Western Pacific Region 2006 Annual Report. Western Pacific Regional Fishery Management Council.

WPRFMC. 2007b. Draft Programmatic Environmental Impact Statement: Towards an Ecosystem Approach for the Western Pacific Region: From Species-based Fishery Management Plans to Place-based Fishery Ecosystem Plans. Prepared by the Western Pacific Regional Fishery Management Council National Marine Fisheries Service, Pacific Islands Region. March 30, 2007. 547 pp.

WPRFMC. 2009. Fishery Ecosystem Plan for Pacific Pelagic Fisheries of the Western Pacific Region. Western Pacific Regional Fishery Management Council. Honolulu, Hawaii. September, 2009. 249 pp.

WPRFMC. 2010. Pelagic Fisheries of the Western Pacific Region 2008 Draft Annual Report. Western Pacific Regional Fishery Management Council.

Appendix A

**Regulatory Impact Review and
Preliminary Regulatory Flexibility Act Analysis**

**Amendment 2 to the Fishery Ecosystem Plan
for Pelagic Fisheries of the Western Pacific Region**

Regulatory Impact Review (RIR)

1.0 Introduction

To comply with Executive Order 12866, the National Marine Fisheries Service (NMFS) requires that a Regulatory Impact Review be prepared for all regulatory actions that are of public interest. This review provides an overview of the problem, policy objectives, and anticipated impacts of regulatory actions, and ensures that management alternatives are systematically and comprehensively evaluated such that the public welfare can be enhanced in the most efficient and cost effective way.

This document contains discussion of the economic impacts to fisheries, fishing communities, and the region due to the implementation of Amendment 2 to the Fisheries Ecosystem Plan for Pelagic Fisheries of the Western Pacific Region. This amendment contains three regulatory actions proposed for domestic pelagic fisheries operating or based in the U.S. exclusive economic zone (EEZ) waters of the Mariana Islands Archipelago which includes the Commonwealth of the Northern Mariana Islands (CNMI) and Guam. It also contains a discussion of the net national benefits of the proposed actions.

2.0 Purpose and Need for Action

The purpose of this proposed action is to prevent or minimize adverse impacts to the troll fleet and the communities which rely on a continued supply of locally-caught fresh fish by reducing chances of gear conflicts and catch competition among fishery sectors while allowing for continued longline fisheries development. The proposed 30-nm longline closed area around CNMI is intended to avoid gear conflicts and catch competition in the nearshore waters between longline fishing vessels and CNMI-based small boat fishing fleet, primarily troll vessels.

The second purpose of this amendment is to prevent catch competition due to localized depletion of the skipjack tuna stock between purse seine vessels and local longline and troll fishing vessels by implementing a purse seine fishing prohibition for all U.S. Exclusive Economic Zone (EEZ) waters around the islands of the Mariana Archipelago. By prohibiting purse seine harvest in U.S. EEZ waters around the Mariana Islands Archipelago, this action would serve to reduce impacts on bigeye tuna in the WCPO and preempt catch competition between seiners and smaller vessels in the Mariana Islands. The alternatives considered here are intended to comply with the MSA, while balancing the needs and concerns of Guam and CNMI's small troll and pelagic longline fishing fleet; and to reduce impacts on stocks of bigeye tuna in the WCPO.

3.0 Description of Alternatives Considered

Three sets of alternatives were developed, each designed to provide long-term viability of CNMI and Guam’s small and locally-based fishing fleets, while balancing the needs and concerns of the fishing industry regarding the development of larger-scale domestic fisheries in U.S. EEZ waters around the Mariana Archipelago.

3.1 Topic 1: CNMI Longline Fishing Prohibited Area

Alternative 1A: No Action

Under Alternative 1A (No Action), no new regulations would be made and U.S. longline vessels would not be prohibited from fishing within U.S. EEZ waters around CNMI.

Alternative 1B: 25 nm Longline Fishing Prohibited Area

Under Alternative 1B, all vessels would be prohibited from pelagic longline fishing within 25 nm of CNMI.

Alternative 1C: 30 nm Longline Fishing Prohibited Area (Preferred)

Under Alternative 1C, all vessels would be prohibited from pelagic longline fishing within 30 nm of CNMI.

Alternative 1D: 50 nm Longline Fishing Prohibited Area for Large Vessels

Under Alternative 1D, vessels greater than 50 ft in length would be prohibited from pelagic longline fishing within 50 nm of CNMI.

Alternative 1E: 75 nm Longline Fishing Prohibited Area

Under Alternative 1E, all vessels would be prohibited from pelagic longline fishing within 75 nm of CNMI.

Alternative 1F: 100 nm Longline *Fishing* Prohibited Area

Under Alternative 1F, all vessels would be prohibited from pelagic longline fishing within 100 nm of CNMI.

3.2 Topic 2: CNMI Purse Seine Fishing Prohibited Area

Alternative 2A: No Action

Under Alternative 2A (No Action), no new regulations would be made and pelagic purse seine vessels would not be prohibited from fishing within U.S. EEZ waters around CNMI.

Alternative 2B: 50 nm Purse Seine Fishing Prohibited Area

Under Alternative 2B, all vessels would be prohibited from pelagic purse seine fishing within 50 nm of CNMI.

Alternative 2C: 100 nm Purse Seine Fishing Prohibited Area

Under Alternative 2C, all vessels would be prohibited from pelagic purse seine fishing within 100 nm of CNMI.

Alternative 2D: Entire U.S. EEZ Purse Seine Fishing Prohibited Area (Preferred)

Under Alternative 2D, all vessels would be prohibited from pelagic purse seine fishing within all U.S. EEZ waters around CNMI.

3.3 Topic 3: Guam Purse Seine Fishing Prohibited Area

Alternative 3A: No Action

Under Alternative 3A (No Action), no new regulations would be made and pelagic purse seine vessels would not be prohibited from fishing within U.S. EEZ waters around Guam.

Alternative 3B: 50nm Purse Seine Fishing Prohibited Area

Under Alternative 3B, all vessels would be prohibited from pelagic purse seine fishing within 50 nm of Guam.

Alternative 3C: 100 nm Purse Seine Fishing Prohibited Area

Under Alternative 3C, all vessels would be prohibited from pelagic purse seine fishing within 100 nm of Guam.

Alternative 3D: Entire U.S. EEZ Purse Seine Fishing Prohibited Area (Preferred)

Under Alternative 3D, all vessels would be prohibited from pelagic purse seine fishing within all U.S. EEZ waters around Guam.

4.0 Description of Potentially Affected Fisheries

4.1 CNMI-based boats

In the 1990's there was some purse seine fishing effort outside of the U.S. EEZ around CNMI. To date, no purse seine fishing activity has been recorded in the U.S. EEZ waters or in the vicinity of CNMI. The CNMI has served as an important transshipment hub for many fishery products because it is exempt from the Jones Act, which requires the use of U.S. built and flagged vessels to carry cargo between U.S. ports. This exemption allows U.S. purse seine vessels fishing on the high seas to offload their catch at Tinian onto foreign vessels for shipment to tuna canneries in American Samoa. In the early 1980s, U.S. purse seine vessels established a transshipment operation at Tinian Harbor. In 1991, a second type of tuna transshipment operation was established on Saipan (Hamnett and Pintz 1996). This operation transshipped fresh longline-caught tuna in the Federated States of Micronesia from air freighters to wide-body jets bound for Japan. The volume of fish flown into and out of Saipan was substantial (Hamnett and Pintz 1996), but the contribution of this operation to the local economy was minimal and it is now defunct for economic reasons.

With the exception of the purse seine support base on Tinian, CNMI has never had a large infrastructure dedicated to commercial fishing. The majority of boats in the local fishing fleet are small, outboard engine-powered vessels. The harvest of pelagic species by CNMI-based vessels has always been small, around 100 metric tons annually, caught with trolling gear. Both supply and demand conditions direct the majority of domestic commercial fishing effort in CNMI toward reef fish and bottomfish. There is less seasonality in these fisheries, and they require shorter offshore trips; moreover, their market value is often much higher than that of the commonly caught pelagic fish.

CNMI's pelagic fishery occurs primarily from the island of Farallon de Medinilla south to the island of Rota. The pelagic fishing fleet consists primarily of trolling vessels less than 24 ft in length which generally take one-day trips within 20 miles around the islands where they find abundant skipjack tuna. These vessels have a limited travel and fishing range and fishery participants necessarily rely on catches from waters within their reach.

Currently there is a new longline fishing company located on Saipan which as of 2010 has four longliners fishing waters around the Mariana Archipelago, beyond 30 miles from shore but within EEZ waters. As of April 2010, there are not yet catch data available on this new fishery. Interest in longline fishing in CNMI has been variable with the issuance of eight, four, and five Western Pacific General Longline permits from 2007 through 2009, respectively.

The pelagic fishery is characterized using data in the Commercial Purchase Data Base which currently documents landings on Saipan where the majority of CNMI's population and fishery participants live. Staff from the Department of Lands and Natural Resources, Division of Fish and Wildlife (DFW) routinely distributes and collects invoice books from 30 participating local fish purchasers on Saipan that record all fish purchases by species categories. The establishment of data collection systems for the islands of Tinian and Rota are in process. It is believed that the commercial purchase database landings include around 90 percent of all commercial landings on Saipan. There is also an unquantified subsistence fishery on Saipan where income is made by selling a small portion of catches door-to-door to cover fishing expenses.

Table 1 summarizes annual participation, landings, and revenue for commercial pelagic vessels. The primary target and most marketable species for the pelagic fleet is skipjack tuna. Yellowfin tuna and mahimahi are also targeted by the pelagic fleet. In 2008 skipjack tuna continued to dominate the pelagic landings, comprising around 80 percent of commercial pelagic landings.

The number of vessels making commercial pelagic landings was relatively constant from 1988-1991, with a sudden jump in 1992. Part of this increase was attributed to an influx of new fishing boats, but it was also discovered that some fishermen were using several different boats, thus artificially inflating the total number of boats concurrently making pelagic landings. Many of the new boats were believed to have left the fishery during 1993. It has been suggested that the increase in active fishing vessels from 1994 to 1997 might be due to the re-entry of boats from the 1992 fleet.

Although the number of vessels and trips has generally trended downward since 2000, landings have fluctuated, especially for skipjack tuna. The reason for this is not fully understood. Creel survey catch rates show catch rates oscillating between 50 and 100 lb/trip both before and after 1991 whereas, the Commercial Purchase data indicate sustained high catch rates before, and low catch rates after 1991. Skipjack catch rates increased between 2004 and 2007 to 114 lb/trip but declined to 96 lb/trip in 2008. The troll catch rate of skipjack has continued a steady decline from 2006 through 2009, based on creel survey data (PIFSC 2010). In 2003, yellowfin catch rates remained relatively stable at 12 lb/trip despite bad weather (e.g. typhoons) that plagued the Mariana Islands nearly the entire year. In 2004, yellowfin catch rates decreased to eight lb/trip but in 2005 and 2006 they again increased. In 2007, yellowfin catch rates increased slightly by 6 percent to 17 lb/trip but declined the following year to 10 lb/trip.

Table 1. Annual Participation, Catch and Revenue for CNMI-based Commercial Pelagic Vessels

Year	Number of Vessels	Number of Trips	Pounds Sold	Adjusted Price per Pound (\$)	Adjusted Revenue (\$)	Adjusted Revenue Per Trip (\$)	Adjusted Revenue Per Vessel (\$)
1983	92	1,408	245,985	2.29	563,838	400	6,129
1984	99	1,621	341,136	2.02	690,231	426	6,972
1985	82	1,240	234,178	2.1	490,784	396	5,985
1986	96	1,356	307,459	2.13	654,181	482	6,814
1987	62	992	205,068	2.18	446,852	450	7,207
1988	78	1,298	334,523	2.18	728,154	561	9,335
1989	77	1,242	286,784	2.19	628,197	506	8,158
1990	79	888	180,450	2.62	473,329	533	5,992
1991	76	999	188,561	2.68	504,578	505	6,639
1992	104	1,419	199,228	2.6	517,468	365	4,976
1993	55	1,372	181,328	2.27	410,971	300	7,472
1994	65	1,218	147,329	2.25	332,122	273	5,110
1995	89	1,721	200,180	2.28	455,504	265	5,118
1996	114	2,249	281,277	2.34	658,346	293	5,775
1997	111	2,042	218,873	2.62	574,156	281	5,173

1998	92	2,223	240,263	2.5	600,949	270	6,532
1999	106	1,759	177,031	2.38	421,274	239	3,974
2000	113	2,095	187,295	2.19	410,048	196	3,629
2001	113	2,178	179,181	2.36	423,214	194	3,745
2002	90	1,835	256,982	2.32	597,436	326	6,638
2003	73	1,715	228,416	2.33	532,700	311	7,297
2004	71	1,801	239,007	2.35	562,321	312	7,920
2005	77	1,990	372,375	2.15	800,952	402	10,402
2006	69	1,463	356,706	1.79	637,137	436	9,234
2007	52	1,366	312,554	1.48	461,951	338	8,884
2008	47	989	197,013	1.61	317,330	321	6,752
Average	84	1,557	242276	2.24	534,386	361	6,610

(WPRFMC 2008)

Note: Inflation-adjusted prices and revenues were derived using the 2008 CNMI Annual Consumer Price Index (CPI).

4.2 Guam-based boats

Pelagic fishing vessels based on Guam are classified into two general groups: 1) distant-water purse seiners and longliners (foreign and domestic) that fish outside Guam’s economic exclusive zone (EEZ) and transship through the island and; 2) small, primarily recreational, trolling boats that are either towed to boat launch sites or marina-berthed charter boats and fish only within local waters, either within Guam’s EEZ or on some occasions in the adjacent EEZ of the Northern Mariana Islands. Most fishermen sell a portion of their catch at one time or another and it is difficult to make a distinction between recreational, subsistence, and commercial fishers. Landings consist primarily of five major species: mahimahi, wahoo, bonita or skipjack tuna, yellowfin tuna, and Pacific blue marlin.

There are three sources of locally-caught fish in Guam’s commercial market: (1) full-time commercial fishermen; (2) part-time commercial fishermen; and (3) subsistence or recreational “expense” fishermen who frequently sell portions of their catch to help defray costs. Licenses are not required to sell fish in Guam, nor are there any reporting requirements for those selling fish.

Prior to establishment of the Guam Fishermans Cooperative Association (GFCA) in 1979 there was no central place to sell fish, and fishermen had to develop their own markets and peddle fish after each trip. In 1982, the Western Pacific Fisheries Information Network (WPacFIN) began working with the GFCA to improve their invoicing system and to obtain data on all fish purchases. As time progressed, other fish markets began to operate, and Guam Department of Agriculture Division of Aquatic and Wildlife Resources (DAWR) and WPacFIN staff worked with them to obtain data through the voluntary receipt book program. Although a proposed law has been introduced several times that would require reporting by dealers and possibly commercial fishermen, it has never made it through the legislative process, and the commercial landings data collection system remains voluntary. Commercial landing data for 2008 show a dramatic decrease from 2007 levels. A primary reason for this is the difficulty in collecting commercial landing data. The largest commercial vendor on Guam, in terms of locally caught fish, ceased providing DAWR with commercial receipts in early 2008. Thus, commercial receipt totals for 2008 reflect only data collected for the first three months of the calendar year. Attempts

to convince the vendor to resume providing DAWR with commercial data have thus far failed. As providing this data is voluntary, DAWR does not have legal recourse to force the data to be submitted. Other vendors have been contacted and enrolled in the commercial receipts program, but the amount of data provided thus far has been minimal. All commercial data provided from 2008 should be regarded as partial at best.

The composition of Guam's fishery landings by fishing sector has troll fishing for pelagics forming almost 80 percent of the landings, with just over 40 percent of total landings coming from non-commercial troll fishing. Guam's fishing activity can be somewhat constrained by seasonality due to weather with generally lower fishing activity during the November to March period when sea conditions are at their worst.

Estimated annual pelagic landings have varied widely in the 27-year time series. Aggregate landings of tuna PMUS increased and non-tuna PMUS decreased. There are general wide year-to-year fluctuations in the estimated landings of the five major pelagic species. In 2008, 95 percent of tuna were caught by non-charter boats.

Table 2. Annual Participation, Catch and Revenue for Guam-based Commercial Pelagic Vessels

Year	Number of Boats	Number of Trips	Pounds Sold	Price Per Pound (2008 \$)	Revenue (2008 \$)	Revenue Per Trip (\$)	Revenue Per Vessel (\$)
1982	199	5,292	153,577	3.95	606,446	115	3,047
1983	193	5,339	285,118	3.83	1,093,163	205	3,047
1984	219	5,913	218,028	3.47	757,480	128	5,664
1985	276	7,454	237,695	4.00	951,683	128	3,459
1986	246	6,999	226,138	3.67	830,019	119	3,448
1987	219	6,776	242,444	3.43	832,514	123	3,374
1988	320	11,981	284,408	3.08	876,748	73	3,801
1989	329	10,669	242,554	2.83	686,650	64	2,740
1990	352	10,523	279,121	3.01	839,281	80	2,087
1991	349	9,870	285,696	2.61	746,438	76	2,384
1992	332	10,167	296,809	2.44	722,802	71	2,139
1993	346	10,295	351,201	2.11	741,769	72	2,177
1994	369	11,125	351,187	1.96	687,665	62	2,144
1995	427	15,562	389,849	1.52	591,140	38	1,864
1996	466	16,066	252,075	1.63	417,099	26	1,384
1997	449	14,313	307,754	1.78	548,780	38	895
1998	469	14,944	405,666	1.87	758,026	51	1,222
1999	449	14,848	260,669	1.91	496,949	33	1,616
2000	416	13,203	376,192	1.66	623,669	47	1,107
2001	375	11,977	399,471	1.69	673,656	56	1,499
2002	375	8,917	325,299	1.54	502,279	56	1,796
2003	371	6,991	272,633	1.43	390,389	56	1,339
2004	401	7,296	285,545	1.39	398,116	55	1,052
2005	358	6,344	228,936	1.32	301,927	48	993

2006	386	6,414	203,139	1.22	248,709	39	843
2007	370	6,383	266,964	1.18	313,946	49	644
2008	385	6,947	136,665	1.81	247,188	36	849
Average	350	9,726	280,179	2.31	625,353	72	642

(WPRFMC 2008)

Note: Inflation-adjusted prices and revenues were derived using the 2008 Guam Annual Consumer Price Index (CPI).

The inflation-adjusted price of tuna and other non-tuna PMUS has exhibited a downward trend since data on the pelagic fishery were first collected in 1980. Sales of locally caught pelagic fish has increasingly suffered due to the prevalence of cheaper pelagic fish caught by longliners.

Despite decreasing revenues with decreased commercial landings over the past decade, pelagic fishing continues, as a majority of trollers do not rely on the catch or selling of fish as their primary source of income. Several factors in recent years have negatively affected trolling activity and may affect fishing activity in the future. The price of fuel has increased significantly; making it more costly to fish and also more attractive to sell fish to recoup costs. More than two-thirds of the FADS are offline, and difficulties with procurement have prevented timely redeployment of these systems. Trolling activity occurs regularly at FADs, and reported to have occurred significantly at offshore banks. However, fishermen also reported more interaction with sharks at offshore banks.

The number of boats involved in Guam’s pelagic or open ocean fishery gradually increased from 193 in 1983 to a peak of 469 in 1998. This number decreased until 2001, but has generally been stable since that year. There were 385 boats involved in Guam’s pelagic fishery in 2008, an increase of 4 percent from 2007 of 370 boats. A majority of the fishing boats are less than 10 meters (33 feet) in length and are usually owner-operated by fishermen who earn a living outside of fishing. Most fishermen sell a portion of their catch at one time or another. And a small, but significant, segment of the pelagic group is made up of marina-berthed charter boats that are operated primarily by full-time captains and crews.

4.3 WCPO Domestic Purse Seine Tuna Fishery

The domestic purse seine fishery has become an increasing portion of all purse seine vessels fishing in WCPO (Figure 1). The estimated delivered value of the purse seine tuna catch in the WCPFC area for 2008 is \$3,124 million that exceeds the previous year’s record level of \$2,393 million (Williams and Terawasi 2008). In NOAA’s 2009 annual report to the WCPFC the domestic purse seine fishery was the largest U.S. fishery in 2008, accounting for 89 percent of the total U.S. catch of skipjack, yellowfin, bigeye, and albacore tuna in the WCPO. Total U.S. purse seine landings increased to 157,849 metric tons (mt) (or 78 percent over 2007 landings) in 2008. Yellowfin tuna catches in the fishery increased from 10,541 mt in 2007 to 23,801 mt in 2008, and skipjack tuna catches increased from 75,210 mt in 2007 to 127,307 mt in 2008.

The fishery operated mainly in areas between 5° N and 10° S latitude and 150° E and 180° longitude in 2008; below the U.S. EEZ around the Mariana Islands which resides between 11° N and 24° N. Before 1995, the fleet in the WCPO fished mainly on free-swimming schools of tunas. Though

highly variable during the last 5 years, the fleet has been fishing equally on free-swimming schools and schools associated with floating objects, including logs and fish aggregating devices (FADs). Since the 1990's there have been more than 200 purse seine vessels operating in the region (Figure 1).




Figure 1: WCPO Purse Seine Vessels 1972-2006

Source: Williams and Terawasi 2008.

4.4 WCPO Domestic Purse Seine Fishery

Prior to beginning purse seine fishing operations in the Western Pacific, the U.S. fleet had been fishing out of California in areas of the eastern Pacific for decades. Various economic, political and regulatory factors contributed to the transition from fishing in the eastern Pacific to the western Pacific, such as fishing overcapacity and increased management controls. Under the South Pacific Tuna Treaty (SPTT), U.S. tuna fishing vessels are able to gain access to fishing grounds in vast areas of the western and central Pacific Ocean (subject to certain conditions), including the exclusive economic zones of several Pacific Island States.

Currently, most of the fishing activity by U.S. purse seine vessels occurs in areas between 5° N and 10° S latitude and 150° E and 170° W longitude in the EEZ waters of Papua New Guinea, Federated States of Micronesia and other Pacific island nations. During El Nino events, however, fishing activity shifts east to the equatorial central Pacific. The percentage of yellowfin tuna in their catches often increases with this shift. U.S. purse seine catch locations for 2007 and 2008 and amount of effort show the large increase in effort and for 2008 a shift to the west, closer to the Mariana Islands Archipelago .

The purse seine catch is stored as a frozen whole product. Most of the catch has historically been off-loaded to the canneries in Pago Pago, American Samoa, however more and more vessels are transshipping catches in Pacific Islands ports for canning and loining destinations in Southeast Asia and Latin America. The final product that is canned in American Samoa is typically destined for the domestic U.S. canned tuna markets. Frozen non-tuna catches may be processed locally (e.g., wahoo) or transshipped to foreign destinations (e.g., billfish and shark)(NMFS 2009).

The number of U.S. vessels active in the WCPO purse seine fleet declined from 39 active vessels in 1998 to 15 in 2005 (WCPFC 2006b) with catches following suit. Increased investment and participation in the U.S. purse seine fishery is evident (NMFS 2009). However, since 2005, the U.S. purse seine fleet more than doubled to 32 vessels in 2008 (Table 3). The most recent information regarding purse seine activity is that there are 36 purse seiners licensed in the U.S. for the 2009/2010 licensing year; the U.S. may license up to 40 vessels.

The U.S. purse seine catch of 157,849 mt in 2008 was composed primarily of skipjack tuna, with smaller catches of yellowfin and bigeye tuna (Table 3). Total catches and effort increased significantly from 2007. Yellowfin tuna catches in the fishery increased from 10,541 mt in 2007 to 23,801 mt in 2008 and catches of skipjack have doubled since 2005 to 127,307 mt in 2008. Figure 2 shows the distribution of effort by the U.S. purse seine fleet in 2007 and 2008.

Table 3: Reported Annual Landings (metric tons) for U.S. Purse Seine Vessels Operating in the WCPO, 1999-2008

	# Active Vessels	Skipjack (mt)	Yellowfin (mt)	Bigeye (mt)
1999	36	129,262	34,529	18,694
2000	33	81,368	29,961	13,886
2001	32	85,539	24,143	6,176
2002	29	88,535	27,191	4,889
2003	26	62,907	20,079	4,470
2004	21	47,896	14,492	5,031
2005	15	62,379	17,685	6,108
2006	13	55,633	8,448	4,364
2007	21	75,210	10,541	2,985
2008	32	127,307	23,801	6,741

Source: WCPFC 2008b




Figure 2: Distribution of Effort by the U.S. Purse Seine Fleet, 2007 & 2008

Note: Mariana Archipelago is located in the mid upper left-hand quadrant.

Source: Williams & Terawasi 2009

The South Pacific Tuna Treaty governs the conduct of U.S. fishing vessel operations in the Treaty Area, which encompasses approximately 10 million square miles. Licenses are issued by the Pacific Islands Forum Fisheries Agency (FFA), based in Honiara, Solomon Islands, which acts as the Treaty administrator on behalf of the Pacific Island Parties to the Treaty (PIPs). Currently, the Treaty allows for a maximum of 45 licenses to U.S. purse seine fishing vessels to

fish in the Licensing Area of the Treaty. Of the 45 licenses, 5 are reserved for "joint venture" arrangements with PIPs.

5.0 Potential for Gear Conflicts between Purse Seine and Small Boat Fisheries

A 1996 study investigated the impact of purse seine fishing on domestic pole-and-line and artisanal fisheries in waters around Kiribati (Hampton et al 1996). Over large areas, e.g., within radii of 300-600 nm of the islands, artisanal catch rates and purse seine catches are generally positively correlated, suggesting that, on this scale, variations in the abundance or catchability of yellowfin affect both purse seiners and artisanal catches in the same way. However, some negative correlations were found for smaller area (50-60 nm) and time scales, indicating that localised effects may occur (Hampton et al. 1996). In addition, there is some indication from additional preliminary modeling of theoretical skipjack tuna movement that purse seine fishing could have an impact on the availability of skipjack tuna for small-scale trolling from equivalent levels of fishing effort occurring as far away as 600 -800 nm (P. Kleiber pers. comm.).

In addition, a 1998 OFP Report titled *Marianas Islands 1998 Pelagic Fisheries Assessment* in assessing the potential impact of allowing foreign longliners to fish in the EEZ around the Mariana Islands states that "...allowing foreign access to the Marianas Islands may cause negative interactions with the troll or artisanal fisheries. While negative interactions have been difficult to substantiate in a variety of fishery studies, the probability of an interaction increases as longline vessels operate in closer proximity to other fishery sectors." These same conclusions may be applied towards assessing potential impacts of domestic purse seining in EEZ waters around the Mariana Islands.

Another study was undertaken to examine the influence of purse seine fishing on longline catches of yellowfin tuna in the Indian Ocean (Nishida 1995). The spatial analysis showed that the catch of the purse seine fishery affects the longline fishery when the purse seine catch exceeds 1 ton/month/5° X 5° area. When data for the heavily fished purse seine area were examined for impacts to the longline fishery there was a modest decrease in longline CPUE evident. The author of this study does point out, however, that a decrease in CPUE of the longline caught yellowfin does not necessarily imply impacts of the purse seine fishery; that other factors such as environmental conditions and fish availability may be responsible for the changes in CPUE (Nishida 1995). Although this study occurred on a fishery in the Indian Ocean it is likely that similar outcomes may be expected on western Pacific tuna fisheries. A similar study on interactions between Mexican longline and purse seine fisheries in the eastern Pacific concluded that the information was not sufficient to define an interaction; it was possible to suggest a "consecutive" interaction (Ortega-Garcia 1995). Another study looked at skipjack movement with regards to fisheries interactions in the western Pacific (Sibert et al. 1995). This study predicted declining yields of 20 percent in the pole-and-line fishery due to purse seine fishing activities, with greatest declines for pole-and-line fleets operating closest to the purse seine fleets. The authors point out that a change of this magnitude would be difficult to detect in actual recorded catches and to be separated from interannual variability in the fisheries. At the time of this study (late 80's – early 90's) pole-and-line were the two principal gear types catching skipjack in the western Pacific. They also point out that potential fishery interaction estimates depend on the natural mortality rate as does the estimated movement pattern used in their analysis (Sibert et al. 1995).

Finally, a study examined the migratory nature of some tunas, specifically skipjack and yellowfin, and the implications for fisheries management (Sibert and Hampton 2003). They found that the median lifetime movement (i.e. displacement) of skipjack (as a measure of mobility) to be from 420-470 nm and for yellowfin approximately 20% less. They also concluded that fishing reduces this and as such only 50% of the fish move beyond the median displacement while the rest remain closer. In other words, a significant proportion of the population will move beyond most EEZs during their lifetime and therefore, both international and domestic (including sub-regional) conservation policies are necessary for conservation and management. The authors conclude in addition to the need for international cooperation "... results also suggest that Pacific Island countries can implement effective domestic management policies to promote conservation and sustainable utilization of tuna stocks within their EEZs".

6.0 Impacts of the Alternatives

6.1 Topic 1: CNMI Longline Prohibited area

As of 2010, there are four permitted longline vessels in CNMI, however, due to the limited number vessels in the fishery over the past decade, landings data are scarce or unable to be reported here due to confidentiality issues. Section 402 of the Magnuson Stevens Act (MSA) generally prohibits the release of confidential fishery information that is submitted to the Secretary in accordance with the MSA and is identifiable to an individual submitter. Because of the limited size of the longline fleet in the Marianas, NMFS is prohibited from releasing annual fishery information. Due to the lack of catch and effort data obtained from longline vessels around CNMI, the Council developed a proxy model using targeted catch, interactions with protected species and fish bycatch data from the nearby Federated States of Micronesia (FSM) fishery. Like FSM, longline vessels that operate around CNMI target yellowfin and bigeye, along with a variety of PMUS including albacore, yellowfin, blue marlin, mahimahi, skipjack, bigeye, and spearfish. The waters around FSM and the Mariana Archipelago are all within the equatorial band and have similar oceanic climates and oceanographic features (Bigelow 1999). Vessels used in FSM are comparable in size to what would be expected in Mariana Archipelago longline operations and both would be expected to target bigeye and yellowfin. Catch data from longline vessels operating in FSM (Table 4) were therefore used to establish this proxy. These vessels are larger than 50 ft and are crewed by local Micronesians. They operate in oceanic habitats similar to the waters around CNMI and make deep sets to maximize catches of bigeye and yellowfin tuna, as would be anticipated of longline vessels around CNMI. Approximately 86 percent of FSM catch is bigeye and yellowfin, with bigeye tuna comprising almost two thirds of the catch.

Table 4: Catch Rates for Longliners Operating in the FSM (lb/hook)

Year	Bigeye	Yellowfin	Black marlin	Blue marlin	Other marlins	Swordfish	Others	Total
2001	0.29	0.15	0.03	0.00	0.00	0.01	0.01	0.49
2002	0.28	0.06	0.02	0.00	0.00	0.01	0.00	0.36
2003	0.28	0.12	0.02	0.00	0.00	0.00	0.04	0.46
2004	0.28	0.10	0.00	0.03	0.00	0.00	0.03	0.44
Average	0.28	0.11	0.02	0.01	0.00	0.01	0.02	0.44

Source: WCPFC 2006

The catch rates for the FSM longliners (Table 4) were combined with typical American Samoa longline vessel operating patterns (Table 5) to generate projections of potential annual fish catches for a single longline vessel fishing within EEZ waters around CNMI (Table 6). American Samoa operations were used as they are believed to be most similar to those used in CNMI.

Table 5. Typical American Samoa Tuna Longline Vessel Operations

Effort measure	Numbers
Annual number of trips	10
Number of sets/trip	20
Hooks/set	2,200
Annual number of hooks	440,000

Source: WPRFMC 2007

Table 6. Projected Annual Landings for One CNMI Longline Vessel

Species	Projected Annual Landings (lb)
Bigeye	124,085
Yellowfin	46,658
Black marlin	7,568
Blue marlin	3,752
Other marlins	83
Swordfish	2,965
Others	8,046
Total	193,157

Under **Alternative 1A** (No Action) CNMI pelagic fishery participants would continue to be affected by factors such as weather, catchability, fuel prices, market prices and other fishery dependent and independent factors. However, if a longline fishery were to develop in CNMI under Alternative 1A, the troll fishing community could be adversely affected by localized stock depletion leading to catch competition and potential gear conflicts between the two fisheries. Although the majority of the longline catch is projected to be bigeye tuna, which is not targeted by CNMI's trolling vessels, the second highest longline catch is projected to be yellowfin tuna which is the trolling fleet's second most important species. In addition, a greater number of longline vessels could potentially cause the troll fleet to face higher fuel costs and other costs associated with having to spend a greater amount of time in seeking target species. Furthermore, the troll fleet could expect to receive lower landings prices due to an increase in the supply of pelagic fish landed by longline vessels. It is not unreasonable to expect that some fishermen would choose to no longer troll depending on the extent of the catch competition and gear conflicts with longline fishing vessels.

It is important to note, however, that allowing the development and expansion of a regulated domestic longline fishery in CNMI could yield positive economic and social benefits through the introduction of new product markets (including both forward and backward linkages throughout the local economy), the creation of jobs supporting the fishery, as well as the potential to strengthen and develop the regions basic fishery infrastructure (e.g., dock space, fuel availability, import/export facilities, processing plants, etc.) and to local consumers through lower pelagic fish prices.

Alternative 1B would prohibit longline fishing within 25 nm around CNMI which would provide an exclusive area for trolling vessels, but this alternative allows longliners to fish in some of the offshore reefs and banks located between 25-30 nm from shore where the troll fleet often fish. Thus trolling operations would experience some adverse impacts from the expansion of longline fishing in the area, but less so than under Alternative 1A.

Alternative 1C (Preferred) would prohibit longline fishing within 30 nm around CNMI which would fully protect the offshore reefs and banks currently used by trolling operations thus significantly reducing the likelihood of adverse impacts on the trolling fleet. The majority of small boats fish within 30 nm of shore, so the potential for gear conflicts and catch competition between longliners and trollers still exists under Alternative 1C, but to a much lesser extent than Alternatives 1A and 1B.

The preferred alternative's 30 nm prohibited area encompasses the majority of the submarine habitat containing banks and reefs where the small boat fleet typically concentrates their fishing efforts. Making these areas off limits to longline activity would prevent gear and fishing conflicts between the two sectors and preclude longline gear entanglement with the banks and reefs. The preferred alternative may also prevent potential safety impacts to participants in the troll fishery which could arise if, without the prohibited area, the small boats felt compelled to fish farther from land and in unfamiliar waters to avoid gear conflicts or catch competition with longline vessels.

The few longline vessels that have fished in CNMI in the past few years, both reported to be fishing outside of 35 miles from shore (J.Ogumoro, CNMI Island Coordinator, WPRFMC, pers.comm.). This supports selection of the 30 nm closure area with regards to avoiding gear conflicts and catch competition while not discouraging increased participation and sustained community participation in local, developing pelagic fisheries operations. Current longline operators in CNMI have also expressed their interest in purchasing fresh fish from the small boat fleet which would also be an economic benefit to CNMI.

Alternative 1D prohibits longline fishing by vessels greater than 50 feet in length within 50 nm of shore. Therefore, longline fishing vessels less than 50 ft length would be able to fish within nearshore waters and without direct competition from large longline vessels, which would benefit those participants. This scenario would, however, be likely to impact the small vessel troll fleet through catch competition and gear conflict with small longline vessels in nearshore waters (i.e. within 30 nm of shore) where the troll fleet fishes. Alternative D is the only one among the six alternatives that would place restrictions based on vessel size.

Under Alternative 1D, longline fishing vessels less than 50 ft in length would be able to fish within nearshore waters and without direct competition from large longline vessels, which may potentially have positive economic impacts, not only to the smaller longliners, but also through the expansion of fisheries related industry to support the potential larger number of landings.

Alternative 1E would prohibit longline fishing within 75 nm around CNMI which would fully protect the areas currently used by trolling operations thus further reducing the likelihood of

adverse impacts on the trolling fleet as compared to Alternative 1C. However, this alternative would have a greater negative economic impact to the longline fleet relative to Alternatives 1A-1C by the incurrence of higher travel costs.

Alternative 1F would prohibit longline fishing within 100 nm around CNMI which would fully protect the areas currently used by trolling operations thus further reducing the likelihood of adverse impacts on the trolling fleet as compared to Alternative 1A, 1B, 1C, and 1E, however the longline fleet would incur higher travel costs.

6.2 Topic 2: CNMI Purse Seine Prohibited area

If a purse seine fishery were to fish in the EEZ around CNMI under **Alternative 2A**, the troll fishing community could be adversely affected by localized stock depletion leading to catch competition between the two fisheries (Hampton et al. 1996). The majority of the purse seine catch is projected to be skipjack tuna, which is also the most important target species for CNMI's troll fleet. The second highest purse seine catch would be yellowfin tuna which is the trolling fleet's second most important species as well. Greater amounts of purse seine effort closer to areas fished by the small boat fleets of CNMI could force the troll fleet to travel farther to maintain their current catches, and/or to lose revenue normally gained through skipjack and yellowfin catches. The catch of skipjack and yellowfin by just one purse seine vessel fishing for just one day would be about 26 tons, which would be approximately 18 percent of the annual skipjack and yellowfin sold by CNMI's entire commercial troll fleet. Reductions in catch rates, landings, revenues and profits accruing to the commercial trolling fleet cannot be quantified due to a lack of detailed information on fishery operations and how these might change under any of the action alternatives. However these are small operations with average annual ex-vessel revenues of \$8,500 per vessel which likely have small profit margins, and even minor reductions in fishery revenues could be highly detrimental to this fishery. Greater numbers of purse seine vessels could also cause the troll fleet to travel farther to find new concentrations of fish, thus incurring increased fuel costs and further reducing their profits.

Alternative 2B would prohibit purse seine fishing within 50 nm around CNMI which would provide an exclusive area for trolling vessels. However given the high fishing capacity of purse seine vessels, depending on the amount of purse seine fishing effort localized stock depletion could still occur in the fishing grounds currently used by commercial trolling vessels. Thus trolling operations could experience some adverse impacts, but less than those under Alternative 2A.

Alternative 2C would prohibit purse seine fishing within 100 nm around CNMI which would further protect the areas currently used by trolling operations thus significantly reducing the likelihood of adverse impacts on the trolling fleet. However given the high fishing capacity of purse seine vessels, depending on the amount of purse seine fishing effort localized stock depletion could still occur in the fishing grounds currently used by commercial trolling vessels. Thus trolling operations could experience some adverse impacts, but less than those under Alternatives 2A or 2B.

Alternative 2D (Preferred) would prohibit purse seine fishing within U.S. EEZ waters around CNMI which would be expected to provide a sufficient buffer area to drastically reduce the likelihood of gear conflicts and catch competition between trollers and purse seiners. Purse seine fishing vessels currently do not fish within U.S. EEZ waters around CNMI, so Alternative 2D will not alter the decision of where to fish, at least in the short term. This may create an adverse impact to purse seine vessels in the future when they would choose to fish within the U.S. EEZ around CNMI, were it not for the implementation of this prohibited area.

Under the preferred alternative, no purse seine fishing would be allowed to take place in the U.S. EEZ waters around CNMI and it is possible there would be some concomitant impacts to the U.S. purse seine fleet if they had chosen to fish these waters, however, impacts are expected to be minimal because the highly mobile purse seine fleet typically fishes in large areas of the Pacific and has exclusive rights to fish in U.S. EEZ waters in the PRIA, except those waters now off limits due to the establishment of the PRIA Marine National Monument. It is also not known what impact upcoming FAD fishing regulations will have on the behavior of the WCPO purse seine fleets.

While the U.S. purse seine fleet does not currently fish in waters of or deliver to CNMI, future purse seine fishing near CNMI, or the ability to accept fish transshipped by nearby purse seine vessels could influence the development the infrastructure needed to process or export such species. Aside from the negative benefits discussed earlier associated with gear competition and stock depletion, the support of a purse seine industry could provide positive benefits through increasing potential employment while opening up markets currently unexplored in CNMI. Although for this to occur, CNMI's marine-based infrastructure (i.e., dock space, fueling platforms, skilled labor, etc) would need to be significantly enhanced beyond current conditions and is not likely to happen in the near future.

6.3 Topic 3: Guam Purse Seine Prohibited Area

If domestic purse seine vessels were to fish in EEZ waters around Guam under **Alternative 3A** (No Action) localized stock depletion could occur and lead to reduced catch rates and catch competition, ex-vessel revenues and profits for commercial trollers (Hampton et al. 1996). The majority of the purse seine catch is projected to be skipjack tuna, which is also the most important target species for Guam's troll fleet. The second highest purse seine catch would be yellowfin tuna which is also important to the trolling fleet. The catch of skipjack and yellowfin by just one purse seine vessel fishing for just one day (26 tons) would be approximately 27 percent of the annual skipjack and yellowfin caught by Guam's entire commercial troll fleet (approximately 96 tons, WPRFMC 2007). Reductions in catch rates, landings, revenues and profits accruing to the commercial trolling fleet cannot be quantified due to a lack of detailed information on fishery operations and how these might change under Alternative 3A (or any of the other alternatives). However these are very small operations with average annual ex-vessel revenues of \$2,780 per vessel which likely have small profit margins and even minor reductions in fishery revenues could be highly detrimental to this fishery. Greater numbers of purse seine vessels could also cause the troll fleet to travel farther to find new concentrations of fish, thus incurring increased fuel costs and further reducing their profits.

Alternative 3B would prohibit purse seine fishing within 50 nm around Guam which would provide an exclusive area for trolling vessels. However given the high fishing capacity of purse seine vessels, depending on the amount of purse seine fishing effort localized stock depletion could still occur in the fishing grounds currently used by commercial trolling vessels. Thus trolling operations could experience some adverse impacts, but less than those under Alternative 3A.

Alternative 3C would prohibit purse seine fishing within 100 nm around Guam which would further protect the areas currently used by trolling operations thus significantly reducing the likelihood of adverse impacts on the trolling fleet. However given the high fishing capacity of purse seine vessels, depending on the amount of purse seine fishing effort localized stock depletion could still occur in the fishing grounds currently used by commercial trolling vessels. Thus trolling operations could experience some adverse impacts, but less than those under Alternatives 3A or 3B.

Alternative 3D (preferred) would prohibit purse seine fishing within all U.S. EEZ waters around Guam which would be expected to provide a sufficient buffer area to eliminate the likelihood of adverse impacts on the trolling fleet. This Alternative addresses the needs of the existing small boat fleet around Guam while imposing little, if any, financial hardships on domestic purse seine operations. This Alternative is consistent with Magnuson-Stevens Act requirements that fishery managers take into account the importance of fishery resources to fishing communities such as Guam in order to provide for the sustained participation of such communities, and to the extent practicable, minimize adverse economic impacts on such communities.

Restricting or prohibiting purse seine fishing in EEZ waters around Guam is not expected to adversely impact the U.S. purse seine fleet given the lack of historical fishing effort associated with this area. Other suitable fishing areas in the proximity would remain available, as would EEZ waters around the Pacific Remote Island Areas (in which domestic purse seine vessels are known to fish).

7.0 Impacts of the Preferred Alternatives on National Costs and Benefits

In accordance with Executive Order 12866, the following is set forth: (1) This rule is not likely to have an annual effect on the economy of more than \$100 million or to adversely affect in a material way the economy, a sector of the economy, productivity, jobs, the environment, public health or safety, or state, local, or tribal governments or communities; (2) This rule is not likely to create any serious inconsistencies or otherwise interfere with any action taken or planned by another agency; (3) This rule is not likely to materially alter the budgetary impact of entitlements, grants, user fees or loan programs or the rights or obligations of recipients thereof; and (4) This rule is not likely to raise novel or policy issues arising out of legal mandates, or the principles set forth in the Executive Order.

As compared to the no action baseline, the implementation of the preferred alternatives would appropriately balance the needs and concerns of CNMI and Guam's small and locally-based fishing fleets with the development of larger-scale domestic longline and purse seine fisheries in waters around the Mariana Archipelago. This action is consistent with the Magnuson-Stevens Fishery Conservation and Management Act.

Preliminary Regulatory Flexibility Act Analysis

The Regulatory Flexibility Act (RFA) requires federal agencies to consider the impact of regulations on small entities in developing the proposed and final regulations. If a proposed rule is expected to have a significant economic impact on a substantial number of small entities, an initial regulatory flexibility analysis (IRFA) must be prepared. Under Section 603(b) of the RFA, each IRFA is required to address:

- (1) reasons why the agency is considering the action,
- (2) the objectives and legal basis for the proposed rule,
- (3) the kind and number of small entities to which the proposed rule will apply,
- (4) the projected reporting, record keeping and other compliance requirements of the proposed rule, and
- (5) any federal rules that may duplicate, overlap or conflict with the proposed rule.

A business is considered to be ‘small’ business by the Small Business Administration (SBA) if it has annual receipts less than \$4.0 million for fish-harvesting, average annual receipts less than \$6.5 million for charter/party boats, 100 or fewer employees for wholesale dealers, or 500 or fewer employees for seafood processors. The IRFA is conducted to comply with the Regulatory Flexibility Act (5 USC 601 et. seq.) and provides a description of the economic impacts of the various alternatives on small entities.

If a proposed rule is not expected to have a significant economic impact on a substantial number of small entities, either adversely or beneficially, then the agency is not required to perform an IRFA. In these instances, an agency must undertake a threshold (preliminary) analysis to determine the economic impact of a proposed rule on small entities. Once this preliminary analysis is undertaken, an agency then can determine whether to certify or undertake a complete IRFA.

1.0 Description of Small Businesses to Which the Rule Would Apply

Based on available information, NMFS has determined that there are no disproportionate economic impacts to small entities. CNMI longline and troll vessels are both considered small entities under the Small Business Administration definition of a small entity, i.e., they are engaged in the business of fish harvesting, are independently owned or operated, are not dominant in their field of operation, and have annual gross receipts not in excess of \$4 million³². Troll vessels are likely to benefit, and CNMI longline vessels could, but are not likely to, be slightly adversely affected. The longliners could benefit because any negative effects of purse seine fishing on the local availability of fish stocks that are targeted by both longliners and purse seiners would be diminished for the longline fleet. Purse seiners are not likely to be negatively

³² The annual total revenues for the commercial pelagic fleets in CNMI and Guam have always remained below \$1million during 1982-2008 (Table 1 and 2 in this Appendix).

affected because they do not currently fish in the areas from which they would be prohibited. Given the amount of landings from U.S. purse seiners, they are not likely to be considered small entities.

The alternatives in this Amendment 2 are primarily are intended to protect the interests of small vessels/fishers in the nearshore fisheries. It will affect all the troll vessels in the same manner and the displacement of few longliners to outside of 30nm from the U.S. EEZ in the CNMI may have insignificant economic impact to this fleet. All these vessels are considered small business entities by the U.S. Small Business Administration, as their annual sales receipts do not exceed \$4.0 million. Therefore, there would be no economic impacts resulting from disproportionality between small and large business entities under the proposed amendment. For these reasons, the actions and alternatives were not evaluated through an IRFA.

2.0 Skills Necessary to Meet Compliance Requirements

No special skills would be required to comply with any of the alternatives considered here.

3.0 Identification of Duplicating, Overlapping, and Conflicting Federal Rules

To the extent practicable, it has been determined that there are no federal rules that may duplicate, overlap, or conflict with this action.



U.S. DEPARTMENT OF COMMERCE
National Oceanic and Atmospheric Administration
NATIONAL MARINE FISHERIES SERVICE
Pacific Islands Regional Office
1601 Kapiolani Blvd., Suite 1110
Honolulu, Hawaii 96814-4700
(808) 944-2200 • Fax (808) 973-2941

FINDING OF NO SIGNIFICANT IMPACT

Amendment 2 to the Fishery Ecosystem Plan for Pelagic Fisheries of the Western Pacific Region, as Partially Approved; Prohibiting Longline Fishing within 30 nm of the CNMI (RIN 0648-AW67)

June 6, 2011

Introduction

This Finding of No Significant Impact (FONSI) was prepared according to the guidelines established in National Marine Fisheries Service (NMFS) Instruction 30-124-1 (July 22, 2005) and the requirements of National Oceanic and Atmospheric Administration (NOAA) Administrative Order 216-6 (NAO 216-6, May 20, 1999), concerning compliance with the National Environmental Policy Act (NEPA, 42 U.S.C. § 4321, *et seq.*). This FONSI is supported by the environmental impact analysis prepared in accordance with requirements of NEPA and documented in Amendment 2 to the Fishery Ecosystem Plan for Pelagic Fisheries of the Western Pacific Region: Establishment of Longline and Purse Seine Prohibited Areas in the Mariana Archipelago, including an Environmental Assessment, Regulatory Impact Review, and Preliminary Regulatory Flexibility Act Analysis," dated March 4, 2011 (attached).

After considering a range of alternatives developed in coordination with its plan team members, the Scientific and Statistical Committee and members of the public, the Western Pacific Fishery Management Council (Council) recommended in Amendment 2 that NMFS take the following actions:

- 1) Prohibit longline fishing within approximately 30 nm of the CNMI (Alternative 1C).
- 2) Prohibit purse seine fishing within the entire U.S. EEZ around the CNMI (Alternative 2D).
- 3) Prohibit purse seine fishing within the entire U.S. EEZ around Guam (Alternative 3D).

Alternatives considered for the three proposed fishery management topics are described in section 7 of the EA and include analyses of the no action alternatives.

Proposed Action

NMFS proposes to partially approve Amendment 2. NMFS proposes to approve the portion of the amendment that establishes a longline fishing prohibited area in the CNMI (Alternative 1C). NMFS has determined that the proposed purse seine measures are not consistent with National Standard 2 of the Magnuson-Stevens Fishery Conservation and Management Act (MSA) and, therefore, are subject to disapproval. Because NMFS will not approve the portions of Amendment 2 relating to prohibitions on purse seine fishing, NMFS' disapproval effectively implements the relevant no action alternatives (Alternatives 2A and 3A). This FONSI analyzes



the significance of the environmental impacts of the proposed action, specifically, implementing the longline fishing prohibited area around the CNMI, and taking no action regarding (i.e., not implementing) the proposed prohibition on purse seine fishing in the EEZ around the CNMI and Guam.

Public involvement

The Amendment document, including the EA, describes public involvement in section 5.0. NMFS published a notice of availability of the draft FEP Amendment and EA in the *Federal Register* (76 FR 13330; March 11, 2011) requesting public review and comments on the Amendment by May 10, 2011. In its proposed rule (76 FR 17811; March 31, 2011), NMFS requested comments by May 16, 2011. NMFS received comments from private individuals generally in support of the Council's recommended actions, and one comment in opposition of the proposed action from the American Tuna Association (ATA), an organization that represents owners of purse seine vessels that fish in the central Pacific Ocean. None of the comments identified any problems with the environmental analysis contained in the EA, so no changes were made as a result of the comments.

Significance Analysis

NAO 216-6 (May 20, 1999) contains criteria for determining the significance of the environmental impacts of a proposed action. In addition, the Council on Environmental Quality's (CEQ) regulations at 40 CFR 1508.27 state that the significance of an action should be analyzed both in terms of "context" and "intensity." Each criterion listed below is relevant in making a finding of no significant impact and has been considered individually, as well as in combination with the others. The significance of this action is analyzed based on the NAO 216-6 criteria and CEQ's context and intensity criteria. These include:

- 1) Can the proposed action reasonably be expected to jeopardize the sustainability of any target species that may be affected by the action?

No. None of the alternatives would result in large changes to current fishing operations. The proposed action will not result in an increase in the number of vessels, the intensity of fishing, or result in large changes to where vessels fish. The measures are intended to allow for long-term stability for troll and longline fleets as they fish within the U.S. EEZ around the CNMI and continuation of purse seine fishing in areas where the purse seiners are currently fishing.

The status of target stocks is summarized in section 8.6, and descriptions of the two affected pelagic fisheries in Guam and the CNMI and purse seining in the WCPO are provided in section 8.2.

With no expected increase in the number of vessels or in effort as a result of the measures, CNMI troll fisheries are expected to continue to catch pelagic species at current levels in nearshore areas and generally within 30 nm of shore. Skipjack tuna will continue to be the main target for troll fisheries (157,708 lb landed in 2008). Based on 2010 stock assessment of skipjack tuna, stocks are robust. The troll fishery catches little bigeye tuna, and some yellowfin tuna (16,

344 lb). The amount of anticipated catch of tuna by troll and longliners for all alternatives is not expected to change from current levels of catch, and will continue to be within the WCPFC quota limits. Catches of pelagic species by the CNMI troll fleet will be expected to continue to be small with a long term average of 224,276 lb which would have no significant adverse impact on target species.

Under the selected alternative, the longline fishery will continue operations outside of 30 nm in the CNMI with a fleet of about four longline vessels. These vessels will continue to target bigeye tuna, and the four-vessel longline fleet is expected to catch annually about 350 mt of pelagic fishes. Under current regulations, WCPO longline harvests of bigeye are currently managed by international quotas, limiting longline catches in CNMI to a maximum of 2,000 mt of bigeye per year. The projected level of catch of bigeye tuna by the four longline vessels (estimated to be 224 mt of bigeye) is just over 10% of the CNMI catch limit, with little to no bigeye landed by troll vessels. The fishery is expected to continue to catch approximately 84 mt of yellowfin tuna annually. The WCPO yellowfin tuna stock is not experiencing overfishing and is not in a state of being overfished.

Troll fishing and longline fishing in Guam are also expected to continue as in the current fishery. The approximately 385 troll vessels from Guam will continue to account for 80% of pelagic landings, and will continue to target skipjack tuna, mahimahi, with some landings of blue marlin. Based on 2010 stock assessment of skipjack tuna, stocks are robust. This fishery is relatively small and the tuna catches by the Guam troll fleet are sustainable and will continue to be sustainable after the management actions are implemented.

Longlining by the single fishing vessel will continue around Guam outside of the existing 50 nm prohibited area and catches of bigeye tuna will continue to be well below the international quota for Guam. The WCPO yellowfin tuna stock is not experiencing overfishing and is not in a state of being overfished. The single longline vessel fishery will continue to represent a very small amount of the fishing mortality and will continue to be managed by the Council to ensure it is a sustainable fishery.

The proposed action is not expected to change the current fishing activity of purse seiners. The U.S. EEZ is currently open to purse seine vessels and unaffected by this proposed action. In the western and central Pacific Ocean, U.S. purse seiners normally operate on the high seas and in the national waters of the parties to the South Pacific Tuna Treaty (SPTT). There is also some effort in the EEZ around the PRIA, and limited effort in the EEZ around American Samoa. No U.S. purse seine catches have been recorded from the EEZ around the CNMI, and no U.S. purse seine catches have been recorded from the EEZ around Guam in over 30 years. Purse seine fishing in the Western and Central Pacific Ocean will continue to be regulated by the Western and Central Pacific Fishery Commission. For these reasons, the purse seine no-action alternatives (2A and 3A) will not result in significant adverse effects on target species by purse seiners.

2) Can the proposed action reasonably be expected to jeopardize the sustainability of any non-target species?

No. None of the alternatives is expected to jeopardize the sustainability of any non-target species. The proposed action is not expected to jeopardize the sustainability of any non-target species because there will be no large increase in the number of vessels, intensity of fishing, or large changes to where vessels fish as a result of the establishment of a longline fishing prohibited area in the CNMI. The current level of bycatch by the CNMI troll and longline fleets is expected to continue and is quite small based on historic catch data.

Alternatives 2A and 3A (no action) will not change the areas in which purse seining is occurring or levels of intensity with which purse seine fishing is conducted because no purse seine vessels currently fish in the U.S. EEZ around CNMI and Guam. Purse seine discards will continue to be managed by the WCPFC and will be unchanged by the implementation of the proposed action. Therefore, there will be no change to impacts to non-target species related to purse seining as a result of implementing the proposed action.

3) Can the proposed action reasonably be expected to cause substantial damage to the ocean and coastal habitats and/or essential fish habitat (EFH) as defined under the Magnuson-Stevens Act and identified in Fishery Management Plans?

No. None of the alternatives would result in large changes to fishery operations that could have adverse impacts on essential fish habitat (EFH) or habitat areas of particular concern (HAPC) for species managed under the Pacific Pelagics or the Marianas Archipelago Fishery Ecosystem Plans. The establishment of a longline fishing prohibited area will not change the intensity of fishing by the troll, longline, or purse seine fleets or result in physical, chemical, or biological alterations to EFH or HAPC. The proposed action will provide spatial separation among fishing activities and reduce the potential for gear conflicts that could result in physical damage to the marine environment. The measure will not result in increases in gear loss that could affect EFH or HAPC or in ocean or coastal habitats. For these reasons, implementing the proposed action will not result in a significant adverse effect on EFH, HAPC, or oceanic or coastal habitats.

4) Can the proposed action reasonably be expected to have a substantial adverse impact on public health or safety?

No. There are no known impacts on public health or safety that are associated with any of the subject fisheries and there will be no large change to any of the fisheries as a result of any of the alternatives. The CNMI longline fishing prohibited area is expected to have positive impacts on the health and safety of the small boat fishery participants by reducing the potential for gear conflicts with longline vessels and allowing for the continuation of nearshore troll fishing at submerged seamounts and reefs that are preferred fishing areas. Longline and purse seine fishing is usually conducted farther from shore than small boat fisheries and these vessels are better equipped to safely fish in offshore waters. With no large changes to current fisheries, and with better separation between longliners and trollers around the CNMI, the proposed action will not result in significant effects on public health or safety at sea.

5) Can the proposed action reasonably be expected to adversely affect endangered or threatened species, marine mammals, or critical habitat of these species?

No. None of the alternatives is expected to adversely affect endangered or threatened species, marine mammals, or critical habitat. The 2009 Biological Opinion (BiOp) on pelagic fisheries of the Western Pacific Region included pelagic fisheries of Guam and the CNMI. The BiOp considered potential stressors and impacts to listed species related to vessel collisions, hooking and entanglement with fishing gear, human disturbance, exposure to vessel waste, direct and indirect competition with foraging grounds and exposure to marine mammal deterrents. Based on the 2009 BiOp, NMFS determined that troll and handline fisheries in the CNMI are not likely to adversely affect ESA listed species. A small potential exists that trolling gear may incidentally hook or entangle a sea turtle or marine mammal, however, NMFS and the Council consider this type of interaction extremely rare and therefore unlikely to adversely affect listed species. The management measures will not result in a change to the conduct of the CNMI or Guam troll fisheries, so there would be no change to the potential impacts of the measure on troll fishery interactions with listed species.

Existing longline fishing regulations would continue to apply for all longline vessels. Moving longline fishing out further from the coast in the CNMI is not expected to increase the likelihood of interactions with marine mammals or sea turtles. Under existing regulations, longline interactions with protected species are required to be reported on Federal logbooks and to date there have been no reported interactions with marine mammals, seabirds or turtles in CNMI or Guam. All longline vessels will continue to be required to adhere to all seabird, sea turtle, and other protected species mitigation and avoidance measures, including carrying an observer if requested to do so by NMFS, currently in effect for deep set longline fishing activities by general longline permit holders.

In November 2006, NMFS issued a biological opinion on the effects of the U.S. purse seine fishery in the WCPO on listed sea turtles and whales. The purse seine fishery was found to be not likely to adversely affect ESA listed species and is expected to continue to operate in areas of the western Pacific without large changes.

There are no known interactions between seabirds and any of the Mariana Archipelago pelagic fisheries. The management of longline fishing will not change impacts to seabirds in any of the three fisheries.

Based on the continuation of current fishing for three types of fishing gear, and the low likelihood of fishery interactions with protected species occurring in the fisheries, and according to the findings of the biological opinions conducted on these fisheries, implementing the proposed action will not jeopardize or adversely affect any populations or habitats of species listed as endangered or threatened under the ESA in a manner not previously considered.

6) Can the proposed action be expected to have a substantial impact on biodiversity and/or ecosystem function within the affected area (e.g., benthic productivity, predator-prey relationships, etc.)?

No. None of the alternatives is expected to have a substantial impact on biodiversity and/or ecosystem function. Prohibiting longline fishing within 30 nm of CNMI will not change the intensity of fishing by longline or troll vessels, which currently do not have a significant impact

on biodiversity and/or ecosystem function, and there are no expected significant adverse impacts from implementing this measure.

There is currently no purse seine fishing within the U.S. EEZ around CNMI and Guam. The fishery is expected to continue to operate outside the U.S. EEZ under the proposed action and no substantial impacts on biodiversity and/or ecosystem function are expected within the affected area as a result of maintaining the status quo for the purse seine fishery.

7) Are significant social or economic impacts interrelated with natural or physical environmental effects?

No. None of the alternatives would result in a large change in fishing operations by trollers, longliners, or purse seiners. The longline prohibited in the CNMI is intended to reduce the potential for adverse effects of longline fishing on trollers and it is intended to allow for continued fishing by the longline fleet while providing for long-term viability of the troll fleet by reducing the potential for catch competition, localized depletion and gear interactions.

Impacts associated with purse seine fishery will continue to be minimal because the highly-mobile purse seine fleet does not typically fish in the U.S. EEZ around Guam or the CNMI; the fleet fishes in large areas of the Pacific and has exclusive rights to fish in U.S. EEZ waters in the Pacific Remote Islands Area (PRIA), except those waters now off-limits due to the establishment of the PRIA Marine National Monument.

NMFS and the Council considered the proposed action in consideration with the potential closures of waters to FAD-fishing within the PRIA, as well as the future potential establishment of purse seine prohibited areas around American Samoa. None of the potential fishery management actions would lead to increased environmental impacts that would, in turn, result in significant social or economic impacts. The environmental impacts of the proposed action and other reasonably foreseeable (though not approved) management actions would likely be beneficial for fish stocks by improving recruitment and survival of sub-adult fishes. These positive impacts have benefits for fishermen as well as fishing communities.

8) Are the effects on the quality of the human environment likely to be highly controversial?

No. The proposed action is intended to provide for the long-term viability of the Guam and CNMI troll fishery while allowing development of the longline fishery in the CNMI. Through public meetings and opportunities for public and agency comment on the proposed rule, this project was coordinated with interested members of the public and other agencies and no controversy regarding effects on the quality of the human environment was revealed.

9) Can the proposed action reasonably be expected to result in substantial impacts to unique areas, such as historic or cultural resources, park lands, prime farmlands, wetlands, wild and scenic rivers, or ecologically critical areas?

No. None of the alternatives would result in large changes to fishing operations. The proposed prohibited area for longline fishing around the CNMI will not conflict with any existing federal

or state marine managed area and there will be no change to fisheries that would affect unique or protected areas.

10) Are the effects on the human environment likely to be highly uncertain or involve unique or unknown risks?

No. None of the alternatives would result in uncertain or unknown effects on the human environment. A longline prohibited area in the CNMI is expected to reduce the potential for catch competition, reduce localized depletion and prevent gear conflicts between trollers and longliners. As no large changes are expected to occur in the troll or longline fisheries as a result of establishing the proposed prohibited area, the effects of the proposed action are predictable and not likely to involve unique or unknown risks.

As no change is expected to occur in the purse seine fishery or troll and longline fisheries as a result of establishing the proposed prohibited areas, the environmental impacts of the proposed action in the U.S. EEZ around Guam and the CNMI are predictable and not likely to involve unique or unknown risks.

11) Is the proposed action related to other actions with individually insignificant, but cumulatively significant impacts?

No. The cumulative effects of the proposed action are discussed in question 16 below.

12) Is the proposed action likely to adversely affect districts, sites, highways, structures, or objects listed in or eligible for listing in the National Register of Historic Places or may cause loss or destruction of significant scientific, cultural, or historical resources?

No. There are no known historic, scientific, or cultural resources that are listed in the National Register of Historic Places, and no known significant resources of this type that would be affected by the proposed prohibited area. No large changes are expected to result in the conduct of the fisheries, and therefore, none of the alternatives would result in the loss or destruction of any significant historic, scientific, or cultural resources.

13) Can the proposed action reasonably be expected to result in the introduction or spread of a non-indigenous species?

No. No introductions or spread of non-indigenous species are expected as there would be no large changes to uses or activities in the affected areas under all alternatives. Current fishery operations are not known to introduce or spread alien species.

14) Is the proposed action likely to establish a precedent for future actions with significant effects or does the proposed action represent a decision in principle about a future consideration?

No. The Council previously recommended and NMFS approved fishery management actions that establish fishing prohibited areas within U.S. EEZ waters. For example, there are longline fishing and large vessel prohibited areas established around Hawaii, Guam, and American

Samoa. Establishment of a longline prohibited area will not set a precedent and does not represent a decision in principle about a future consideration.

15) Can the proposed action reasonably be expected to threaten a violation of federal, state, or local law or requirements imposed for the protection of the environment?

No. The fishery management measures will be consistent with all applicable federal laws and other requirements for environmental protections and was evaluated for compliance with the Magnuson-Stevens Act, the Endangered Species Act, the Marine Mammal Protection Act, and the Coastal Zone Management Act.

16) Can the proposed action reasonably be expected to result in cumulative adverse effects that could have a substantial effect on the target or non-target species?

No. None of the alternatives would result in cumulatively significant adverse effects on target or non-target species. The CNMI longline prohibited area would separate longliners from areas that are preferentially fished by small trollers to reduce gear conflicts and address concerns with localized depletion and catch competition between the two fleets. The longline prohibited area is not likely to change the conduct of the fishery or the volume of fishing by either trollers or longliners so impacts to target and non-target stocks will remain as they currently are, which are at sustainable levels. These impacts were considered in light of other fisheries across the Pacific that fish for the same species. The low level of environmental impacts resulting from establishing a CNMI longline fishing prohibited area in the CNMI would not be magnified when combined with the effects of other fisheries, other proposed or recent fishery management actions or other activities affecting the same fish stocks or marine resources.

The proposed action would not change the operations of the purse seine fishery. Other fishery management activities that are being proposed that affect purse seiners (e.g., establishing purse seine closed areas in American Samoa and managing FAD use by purse seiners) were considered but would not change the impact analysis for the current proposed action.

Other Findings

Climate Change Impacts

There are no specific studies about the impacts of ocean circulation pattern changes or climate change on tuna stocks of the western Pacific region. In general, it has been shown that large scale climate cycles can impact winds, currents, ocean mixing, temperature regimes, nutrient recharge, ocean acidity, sea level, and affect the productivity of all trophic levels in the Pacific Ocean. These impacts can result in variability in fish stock size, recruitment, growth rates, or other factors affecting fish stocks. Pelagic fish stocks, as well as protected species that interact with the fisheries, are currently affected by these large-scale climate fluctuations and will continue to be affected in the same way whether or not the longline and purse seine fishery prohibited areas are established in the Mariana Archipelago. Catch reports and fish stock assessments will continue to be used by the Council, NMFS, local fishery management agencies and others, to help manage western Pacific fisheries sustainably, even in the face of climate change impacts. Climate change

will not affect the environmental impact or effectiveness of implementing the measures which will establish geographic separations between fishing fleets but will otherwise not change fishing intensity, location of troll or purse seiners, or the number of participants. Because no large changes are expected to occur in any of the three fisheries, the proposed action will not result in a change to annual greenhouse gas emissions.

Environmental Justice

The EA considered whether the alternatives would result in environmental harm and risks that might disproportionately and adversely affect members of minority and low-income populations. The alternatives considered would not result in significant adverse environmental effects, so disproportionate high and adverse effects on members of environmental justice populations will not occur.

Determination

In view of the information presented in this document and the analysis contained in the supporting Environmental Assessment prepared for Amendment 2, I have determined that implementing the proposed action, establishing a 30-nm prohibited area for longline fishing around the CNMI (Alternative 1C) and not implementing the proposed purse seine prohibitions (Alternatives 2A and 3A), will not have a significant impact on the quality of the human environment as described above and in the supporting EA. In addition, all beneficial and adverse impacts of the proposed action have been addressed to reach the conclusion of no significant impacts. Accordingly, preparation of an Environmental Impact Statement for this action is not necessary.



Michael D. Tosatto
Regional Administrator
Pacific Islands Region, NMFS, NOAA

6 JUNE 2011

Date