

NOAA Technical Report NMFS SSRF-782



**A Five-Year Study
of Seasonal Distribution
and Abundance of Fishes
and Decapod Crustaceans
in the Cooper River and
Charleston Harbor, S.C.,
Prior to Diversion**

E. L. Wenner, W. P. Coon III,
M. H. Shealy, Jr., and P. A. Sandifer

July 1984

NATIONAL SEA GRANT DEPOSITORY
PEL LIBRARY BUILDING
URI, NARRAGANSETT BAY CAMPUS
NARRAGANSETT, RI 02882

U.S. DEPARTMENT OF COMMERCE
National Oceanic and Atmospheric Administration
National Marine Fisheries Service

NOAA TECHNICAL REPORTS

National Marine Fisheries Service, Special Scientific Report—Fisheries

The major responsibilities of the National Marine Fisheries Service (NMFS) are to monitor and assess the abundance and geographic distribution of fishery resources, to understand and predict fluctuations in the quantity and distribution of these resources, and to establish levels for optimum use of the resources. NMFS is also charged with the development and implementation of policies for managing national fishing grounds, development, and enforcement of domestic fisheries regulations, surveillance of foreign fishing off United States coastal waters, and the development and enforcement of international fishery agreements and policies. NMFS also assists the fishing industry through marketing service and economic analysis programs, and mortgage insurance and vessel construction subsidies. It collects, analyzes, and publishes statistics on various phases of the industry.

The Special Scientific Report—Fisheries series was established in 1949. The series carries reports on scientific investigations that document long-term continuing programs of NMFS, or intensive scientific reports on studies of restricted scope. The reports may deal with applied fishery problems. The series is also used as a medium for the publication of bibliographies of a specialized scientific nature.

NOAA Technical Reports NMFS SSRF are available free in limited numbers to governmental agencies, both Federal and State. They are also available in exchange for other scientific and technical publications in the marine sciences. Individual copies may be obtained from DR22, User Services Branch, Environmental Science Information Center, NOAA, Rockville, MD 20852. Recent SSRF's are:

722. Gulf menhaden, *Brevoortia patronus*, purse seine fishery: Catch, fishing activity, and age and size composition, 1964-73. By William R. Nicholson. March 1978, iii + 8 p., 1 fig., 12 tables.

723. Ichthyoplankton composition and plankton volumes from inland coastal waters of southeastern Alaska, April-November 1972. By Chester R. Maitson and Bruce L. Wing. April 1978, iii + 11 p., 1 fig., 4 tables.

724. Estimated average daily instantaneous numbers of recreational and commercial fishermen and boaters in the St. Andrew Bay system, Florida, and adjacent coastal waters, 1973. By Doyle F. Sutherland. May 1978, iv + 23 p., 31 figs., 11 tables.

725. Seasonal bottom-water temperature trends in the Gulf of Maine and on Georges Bank, 1963-75. By Clarence W. Davis. May 1978, iv + 17 p., 22 figs., 5 tables.

726. The Gulf of Maine temperature structure between Bar Harbor, Maine, and Yarmouth, Nova Scotia, June 1973-November 1976. By Robert J. Pawlowski. December 1978, iii + 10 p., 14 figs., 1 table.

727. Expendable bathythermograph observations from the NMFS/MARAD Ship of Opportunity Program for 1975. By Steven K. Cook, Barclay P. Collins, and Christine S. Carty. January 1979, iv + 93 p., 2 figs., 13 tables, 54 app. figs.

728. Vertical sections of semimonthly mean temperature on the San Francisco-Honolulu route: From expendable bathythermograph observations, June 1966-December 1974. By J. F. T. Saur, L. E. Eber, D. R. McLain, and C. E. Dorman. January 1979, iii + 35 p., 4 figs., 1 table.

729. References for the identification of marine invertebrates on the southern Atlantic coast of the United States. By Richard E. Dowds. April 1979, iv + 37 p.

730. Surface circulation in the northwest Gulf of Mexico as deduced from drift bottles. By Robert F. Temple and John A. Martin. May 1979, iii + 13 p., 8 figs., 4 tables.

731. Annotated bibliography and subject index on the shortnose sturgeon, *Acipenser brevirostrum*. By James G. Hoff. April 1979, iii + 16 p.

732. Assessment of the Northwest Atlantic mackerel, *Scomber scombrus*, stock. By Emory D. Anderson. April 1979, iv + 13 p., 9 figs., 15 tables.

733. Possible management procedures for increasing production of sockeye salmon smolts in the Naknek River system, Bristol Bay, Alaska. By Robert J. Ellis and William J. McNeil. April 1979, iii + 9 p., 4 figs., 11 tables.

734. Escape of king crab, *Paralithodes camtschatica*, from derelict pots. By William L. High and Donald D. Worlund. May 1979, iii + 11 p., 5 figs., 6 tables.

735. History of the fishery and summary statistics of the sockeye salmon, *Oncorhynchus nerka*, runs to the Chignik Lakes, Alaska, 1888-1956. By Michael L. Dahlberg. August 1979, iv + 16 p., 15 figs., 11 tables.

736. A historical and descriptive account of Pacific coast anadromous salmonid rearing facilities and a summary of their releases by region, 1960-76. By Roy J. Wahle and Robert Z. Smith. September 1979, iv + 30 p., 15 figs., 25 tables.

737. Movements of pelagic dolphins (*Stenella* spp.) in the eastern tropical Pacific as indicated by results of tagging, with summary of tagging operations, 1969-76. By W. F. Perrin, W. E. Evans, and D. B. Holly. September 1979, iii + 14 p., 9 figs., 8 tables.

738. Environmental baselines in Long Island Sound, 1972-73. By R. N. Reid, A. B. Frame, and A. F. Draxler. December 1979, iii + 31 p., 40 figs., 6 tables.

739. Bottom-water temperature trends in the Middle Atlantic Bight during spring and autumn, 1964-76. By Clarence W. Davis. December 1979, iii + 13 p., 10 figs., 9 tables.

740. Food of fifteen northwest Atlantic gadiform fishes. By Richard W. Langton and Ray E. Bowman. February 1980, iv + 23 p., 3 figs., 11 tables.

741. Distribution of gammaridean Amphipoda (Crustacea) in the Middle Atlantic Bight region. By John J. Dickinson, Roland L. Wigley, Richard D. Brodeur, and Susan Brown-Lieger. October 1980, iv + 46 p., 26 figs., 52 tables.

742. Water structure at Ocean Weather Station V, northwestern Pacific Ocean, 1966-71. By D. M. Husby and G. R. Seckel. October 1980, 18 figs., 4 tables.

743. Average density index for walleye pollock, *Theragra chalcogramma*, in the Bering Sea. By Loh-Lee Low and Ikuo Ikeda. November 1980, iii + 11 p., 3 figs., 9 tables.

NOAA Technical Report NMFS SSRF-782



**A Five-Year Study
of Seasonal Distribution
and Abundance of Fishes
and Decapod Crustaceans
in the Cooper River and
Charleston Harbor, S.C.,
Prior to Diversion**

**E. L. Wenner, W. P. Coon III,
M. H. Shealy, Jr., and P. A. Sandifer**

July 1984

U.S. DEPARTMENT OF COMMERCE
Malcolm Baldrige, Secretary
National Oceanic and Atmospheric Administration
John V. Byrne, Administrator
National Marine Fisheries Service
William G. Gordon, Assistant Administrator of Fisheries

The National Marine Fisheries Service (NMFS) does not approve, recommend or endorse any proprietary product or proprietary material mentioned in this publication. No reference shall be made to NMFS, or to this publication furnished by NMFS, in any advertising or sales promotion which would indicate or imply that NMFS approves, recommends or endorses any proprietary product or proprietary material mentioned herein, or which has as its purpose an intent to cause directly or indirectly the advertised product to be used or purchased because of this NMFS publication.

CONTENTS

Introduction	1
Study area	1
Methods and materials	3
Data collection	3
Data analysis	3
Results	4
Physicochemical parameters	4
Community composition and richness	4
Temporal and spatial distributions	10
<i>Stellifer lanceolatus</i>	10
<i>Anchoa mitchilli</i>	11
<i>Micropanchax undulatus</i>	11
<i>Brevoortia tyrannus</i>	11
<i>Leiostomus xanthurus</i>	11
<i>Syphurus plagiura</i>	11
<i>Bairdiella chrysoura</i>	11
<i>Cynoscion regalis</i>	12
<i>Urophycis regia</i>	12
<i>Trinectes maculatus</i>	12
<i>Penaeus setiferus</i>	12
<i>Penaeus aztecus</i>	13
<i>Xiphopenaeus kroyeri</i>	13
<i>Callinectes sapidus</i>	13
Biomass estimates	13
Discussion	14
Acknowledgments	15
Literature cited	15

Figures

1. Locations of six sampling stations in the Charleston Harbor-Cooper River estuarine system, S.C., during the 5-yr study from February 1973 to December 1977	2
2. Two-way coincidence tables of constancy and fidelity, comparing species associations among sampling stations in the Cooper River-Charleston Harbor estuarine system for fall, winter, spring, and summer collections, 1973-77	8-9
3. Abundance of 10 major fish species collected monthly in the channel of the Cooper River-Charleston Harbor estuarine system, 1973-77	10
4. Annual variation in the transformed mean number of individuals of 10 major fish species collected in the Cooper River-Charleston Harbor estuarine system, 1973-77	11
5. Abundance of four major species of decapod crustaceans collected in the channel of the Cooper River-Charleston Harbor estuarine system, 1973-77	12
6. Annual variation in the transformed mean number of individuals of four major decapod crustacean species collected in the Cooper River-Charleston Harbor estuarine system, 1973-77	13

Tables

1. Average water temperatures and salinities in the Charleston Harbor-Cooper River estuarine system, S.C., 1973-77	4
2. Total numbers and total weights of fish species collected 1973-77 in the Cooper River-Charleston Harbor estuarine system, S.C.	5
3. Total numbers and total weights of decapod crustacean species collected 1973-77 in the Cooper River-Charleston Harbor estuarine system, S.C.	6
4. Average numbers of species of fishes and decapod crustaceans collected 1973-77 at stations in the Cooper River-Charleston Harbor estuarine system, S.C.	6
5. Average numbers of individual fish and decapod crustaceans collected 1973-77 at stations in the Cooper River-Charleston Harbor estuarine system, S.C.	6
6. Species groups formed from seasonal cluster analyses of fishes and crustaceans collected in the Cooper River-Charleston Harbor estuarine system, 1973-77	7
7. Average seasonal biomass and density of fishes and decapod crustaceans collected at stations in the Cooper River-Charleston Harbor estuarine system, S.C., 1973-77	13

A Five-Year Study of Seasonal Distribution and Abundance of Fishes and Decapod Crustaceans in the Cooper River and Charleston Harbor, S.C., Prior to Diversion

E. I. WENNER, W. P. COON III, M. H. SHEALY, Jr., and P. A. SANDIFER¹

ABSTRACT

Fluctuations in the distribution and abundance of fishes and decapod crustaceans collected by a 6 m otter trawl net from the Cooper River-Charleston Harbor estuarine system (South Carolina, USA) were examined over a 5-year sampling period. A total of 101 fish species and 41 decapod crustacean species were collected. Species richness was greatest at stations nearest the harbor mouth. Annual fluctuations in species abundance were apparently related to low bottom-water temperatures which affected year-class strength. Ten species accounted for ~ 90% of the total number and ~ 71% of the total biomass of fin fishes collected in the estuary: *Stellifer lanceolatus*, *Anchoa mitchilli*, *Micropanchax undulatus*, *Brevoortia tyrannus*, *Leiostomus xanthurus*, *Syphodus plagiusa*, *Bairdiella chrysoura*, *Cynoscion regalis*, *Urophycis regia*, and *Trinectes maculatus*. The decapod crustaceans *Fenaeus setiferus*, *P. aztecus*, *Xiphopenaeus kroyeri*, and *Callinectes sapidus* dominated the fin fishes in abundance but not biomass. They composed ~ 96% by number and ~ 97% by weight of the total decapod fauna. The biomass of fishes from this study is lower than values reported for other estuaries along the Atlantic coast of the United States.

The Cooper River-Charleston Harbor estuarine system, an important nursery area for fishes and decapod crustaceans, is characterized by gradual changes in faunal assemblages and considerable overlap in spatial distributional patterns of resident and transient species. Numerically dominant species of fish and decapod crustaceans form assemblages which are spatially and temporally ubiquitous. Resident estuarine species and stenohaline marine species are more restricted in their distribution.

INTRODUCTION

Charleston Harbor and its tributary, the Cooper River, have been subjected to greatly increased man-made alterations since 1942. Prior to that time, the Cooper River was a relatively small coastal plains stream with a watershed of 1.86 million km². After construction of Pinopolis Dam across the upper watershed of the Cooper River and creation of Lake Moultrie, input of freshwater to the Cooper River increased, resulting in inundation of marshes and abandoned rice fields. Increased freshwater flow into Charleston Harbor decreased salinity (Zetler 1953) and formed density currents with a predominant upstream bottom flow throughout most of the lower 18 km of the harbor. As a consequence, sediments were trapped within the harbor and shoaling increased considerably (U.S. Army Corps of Engineers²). In turn, the shoaling has caused an increase in dredging costs and depletion of available disposal sites within the harbor. Because of this situation, the Army Corps of Engineers will redivert water flow in 1983 from Lake Moultrie into the Santee River system to effect a reduction of flow into the Cooper River.

The proposed rediversion probably will produce significant changes in estuarine habitat as well as in populations of estuarine organisms, such as fishes and decapod crustaceans. To assess possible effects of rediversion on population structure, spawning success, and distribution of these organisms, it is necessary to determine species composition, abundance, and distribution prior to the perturbation. This paper describes fluctuations in these parameters over a 5-yr period for fishes and decapod crustaceans in the Cooper River-Charleston Harbor estuarine system.

STUDY AREA

The Cooper River is classified as a mixohaline system, in which the salt wedge extends along the bottom to Big Island (Station C002) and bottom salinities decrease from about 27‰ at Cummings Point (Station J003) to freshwater at the Tee (Station C001) (Mathews and Shealy 1978) (Fig. 1). Charleston Harbor is a stratified or salt-wedge estuary with saltwater intrusion primarily a function of the tidal range and the amount of freshwater released by the Santee-Cooper Dam. A salinity differential between top and bottom strata of the harbor causes the bottom flow currents to predominate over the bottom ebb currents, with the result that upstream movement of the bottom currents within the saline region of the harbor forms a sediment trap (South Carolina Wildlife and

¹Marine Resources Research Institute, South Carolina Wildlife & Marine Resources Department, P.O. Box 12359, Charleston, SC 29412.

²U.S. Army Corps of Engineers. 1975. Final Environmental Statement: Cooper River Rediversion Project, Charleston Harbor, S.C. [U.S. Army Corps of Engineers, Charleston District Office, Charleston, SC 29403. 201 p.]




Figure 1. - Locations of six sampling stations in the Charleston Harbor - Cooper River estuary, S.C., during the 5-yr study from February 1973 to December 1977.

Marine Resources Department²). Extended periods of high river flow in the Cooper River frequently dilute water in Charleston Harbor and even in the vicinity of the harbor mouth (U.S. Army Corps of Engineers, footnote 2).

Coastal marshes cover approximately 20,230 ha in the entire Charleston Harbor system. Of the total marsh area, salt marshes compose about 48%, while freshwater marshes cover approximately 36%, brackish marshes make up 6%, and impoundment areas cover 10% (Tiner 1977). The marshes of the Cooper River reflect strong freshwater inflow, dominated by bulrushes (*Scirpus* sp.), cattail (*Typha* sp.), and giant cordgrass, *Spartina cynosuroides*. Smooth cordgrass, *Spartina alterniflora*, dominates the low salt-marsh habitats and is mixed with black needlerush, *Juncus roemerianus*, in upstream locations where salinity transitions occur (South Carolina Wildlife and Marine Resources Department, footnote 3).

METHODS AND MATERIALS

Data Collection

We sampled six fixed stations in the channel of the Cooper River-Charleston Harbor system (Fig. 1): C001 (The Tee), C002 (Big Island), C003 (North Charleston), C004 (Mouth of Cooper River), J001 (Charleston Harbor), and J003 (Cummings Point). Stations extended in a transect from the harbor mouth inland to above permanent freshwater. Each station was sampled once a month during the 5 yr from February 1973 through December 1977, with the following exceptions: C001 was sampled only during 1973 and January 1974 and was discontinued because of untrawlable bottom; a new station, J001, was established in January 1975. In addition, J003 was not sampled until May 1973.

All collections were made with a 6 m (20-ft) semiballoon otter trawl of 2.5 cm (1-in) stretch mesh. This net is particularly selective toward capture of juvenile fishes and is less effective in collection of older, larger fish and highly mobile decapod crustaceans. Twenty-minute tows were made against flood tide during daylight hours at a speed of 1.3 m/s (2.5 kn), resulting in a coverage of 1.5 ± 0.4 km/tow.

Bottom-water samples were collected 0.5 m above the bottom with Van Dorn bottles at each station prior to trawling. Water temperature was read from stem thermometers mounted within the Van Dorn bottles. Salinity was measured in the laboratory with a Beckman RS7B induction salinometer. Dissolved oxygen was determined by the Winkler-Carpenter method (Strickland and Parsons 1968). Turbidity was determined with a Hach Model 2100A turbidimeter. Winter sampling encompassed January-March; spring sampling April-June; summer July-September; and fall October-December.

Specimens collected were either processed in the field or preserved in 10% Formalin and returned to the laboratory for identification, counting, weighing (nearest 0.1 g), and measuring (total length for fishes; carapace width for crabs, measured as distance between tips of lateral spines; and total length for shrimp from tip of rostrum to tip of telson). We

recorded size measurements for all species numbering ≤ 50 specimens per tow. At stations where the trawl captured larger numbers of organisms, we subsampled the catch as follows: If ≥ 50 to ≤ 250 individuals were collected, a minimum of 50 randomly selected specimens was measured; if > 250 to ≤ 500 individuals were caught, a minimum of 20% was measured; when > 500 were caught, a minimum of 10% was measured.

Data Analysis

The degree of similarity among collections and among species was determined using normal and inverse cluster analyses, employing the Bray-Curtis similarity coefficient and "flexible sorting strategy" with the cluster intensity coefficient, β , set at the now standard value of -0.25 (Lance and Williams 1967; Williams 1971; Stephenson et al. 1972; Clifford and Stephenson 1975). Species which occurred in only one or two collections during a sampling period and collections which contained only one species were eliminated from the analyses. Abundances were logarithmically transformed ($\log_{10}(x + 1)$ where x is the number of individuals for a given species) in order to lessen the tendency of extremely abundant species to dominate the similarity matrix (Clifford and Stephenson 1975).

Two dendograms were generated for each season: 1) A dendrogram which indicated association of all sites by season during the 5-yr sampling period based on faunal similarity, and 2) a dendrogram which indicated association of all species collected each season during the 5-yr sampling period based on the abundance of species at sites where they were collected. Nodal analysis (Williams and Lambert 1961; Lambert and Williams 1962) was subsequently used to examine species group and station coincidences based on patterns of constancy and fidelity (Boesch 1977).

An index of abundance (Musick and McEachran 1972; Elliott 1977) was used to compare numbers and weights of selected dominant species and is expressed as:

$$\text{Index of abundance} = \frac{1}{N} \sum_{i=1}^N \log_{10}(x_i + 1)$$

where x = no. or weight of individuals of a given species in a chosen frame and N = no. of collections in that time frame

We determined biomass and density estimates for fishes and decapod crustaceans from computations of area swept by our trawl gear. Estimates of area swept (a) were determined by the following equation given by Klima⁴:

$$a = \frac{KM(0.6H)}{10,000 \text{ m}^2/\text{ha}}$$

where K is speed in meters per hour, M is time in hours fished, and H is headrope length in meters. The constant 0.6 designates an effective swath of about 60% of the headrope length as used by Roe (1969) and established by Wathne (1959). The area swept by our 6 m otter trawl was estimated by this method to be 0.54 ha/tow.

²South Carolina Wildlife and Marine Resources Department. 1972. A study of the Charleston Harbor estuary with special reference to deposition of dredged sediments. Unpubl. manuscr., unpaginated. Office of Marine Conservation, Management and Services, P.O. Box 12559, Charleston, SC 29412.

⁴Klima, E. F. 1976. A review of the fishery resources in the western central Atlantic. WECAF Studies No. 3, FAO No. 32975-76, 77 p. Available from UNIPUB, 1180 Avenue of the Americas, New York, NY 10036.

RESULTS

Physicochemical Parameters

Bottom-water temperatures were very similar among all stations with mean temperatures lowest but most variable in February and warmest in July, August, and September. Yearly average temperatures were lowest in 1976 and 1977 (Table 1).

Salinities measured monthly were highly variable at all stations; nonetheless, average salinities were sufficiently different between stations (Table 1) to justify classification of sites according to the Venice system (Anonymous 1958). Station C001 was classified as limnetic because salinity did not exceed 0.5‰ throughout the year it was sampled. Salinities at stations C002 and C003 ranged from 0.4 to 18.0‰, and these stations were characterized as limnetic-mesohaline. We classified station C004 as limnetic-polyhaline based on the salinity range of 0.67-26.2‰. Stations J001 (7.5-27.7‰) and J003 (19.4-33.3‰) had the highest salinities and were classified as mesopolyhaline and polyehaline, respectively. Average salinity varied also with season, being lowest in spring and highest in fall (Table 1).

Average dissolved oxygen concentrations were greatest at all stations in January and February and lowest in summer. No relation was apparent between dissolved oxygen concentrations and station or depth. The lowest average concentration measured in the Cooper River was 4.9 mg/l.

Although we did not specifically determine sediment characteristics for our fixed stations in the Cooper River-Charleston Harbor system, Mathews and Shealy (1978) reported the general bottom type to be as follows: C001 (hard-mud), C002 (sand), C003 (shell and sand), C004 (mud-shell-sand), J001 (mud and sand), and J003 (shell and mud).

Table 1.—Average water temperatures and salinities in the Charleston Harbor-Cooper River estuarine system, S.C., 1973-77.

Parameters	Environmental factors	
	Avg. temp (°C)	Avg. salinity (‰)
Year (all stations)		
1973	22.1	12.0
1974	20.2	14.1
1975	21.1	14.5
1976	19.9	15.7
1977	19.8	15.7
Station (all years, 1973-77)		
J003	20.9	27.6
J001	20.7	19.2
C004	19.8	12.6
C003	20.9	5.4
C002	20.5	1.8
C001	18.6	0.07
Season (all stations, 1973-77)		
Fall	19.1	15.6
Winter	11.7	14.5
Spring	21.7	13.7
Summer	28.4	14.9

Community Composition and Richness

A total of 101 species of fishes was collected from the Cooper River-Charleston Harbor system during the 1973-1977 sampling period (Table 2). Length, bottom salinity, and tem-

perature ranges, along with relative abundance of all species collected, are available upon request from the authors. Ten species accounted for 90% of the total number and 71% of the total biomass of fishes collected in this estuarine system: Star drum, *Stellifer lanceolatus*; bay anchovy, *Anchoa mitchilli*; Atlantic croaker, *Micropogonias undulatus*; Atlantic menhaden, *Brevoortia tyrannus*; spot, *Leiostomus xanthurus*; blackcheek tonguefish, *Syphodus plagiatus*; silver perch, *Bairdiella chrysoura*; weakfish, *Cynoscion regalis*; spotted hake, *Urophycis regia*; and hogchoker, *Trinectes maculatus*. *Stellifer lanceolatus* was the most abundant fish collected each year of the study, except in 1977 when *Brevoortia tyrannus* was most abundant.

During the 5-yr sampling period, we collected 44 decapod crustacean species (Table 3). Decapods dominated the fishes numerically but not in biomass. The numerical dominance was due to large numbers of white shrimp, *Penaeus setiferus*, collected in the Cooper River-Charleston Harbor system. This species constituted 83% of the total number and 69% of the total biomass of decapod crustaceans. *Penaeus setiferus* was numerically dominant during each of the 5 yr of our study, except in 1977 when *P. aztecus* was most abundant. These two penaeid shrimps, together with seabob, *Xiphopenaeus kroyeri*, and blue crab, *Callinectes sapidus*, composed about 96% by number and 97% by weight of total decapod fauna collected from the Cooper River-Charleston Harbor estuarine system.

Average numbers of species collected were greatest in 1976 at the higher salinity stations C004, J003, and J001 (Table 4), whereas species richness decreased along the transect of stations upriver. Mean numbers of individuals were greatest at the higher salinity stations (C004, J003, J001). In addition, more individuals were collected in 1975-76 than in 1973-74 and 1977 (Table 5). The fewest individuals were collected in 1977, probably because of prolonged periods of extremely low water temperatures during January and February 1977.

Normal classification analysis showed that collections were not distinctly grouped according to their location along the salinity gradient. During all seasons, collections from stations classified as limnetic and/or mesohaline were faunistically least similar to those from high-salinity sites, but overlap occurred in classification of collections in the mesopolyhaline and polyehaline range. Because groups broadly overlapped by stations and were not clearly separated by cluster analysis according to salinity regimes within the estuary, we compared collections from our fixed stations, rather than site groups as determined from cluster analysis, with the species groups listed in Table 6. In this way, seasonal comparisons among stations were facilitated by direct cross-referencing against the species assemblages at each station.

During all seasons, collections from higher salinity stations J003 and J001 were characterized by stenohaline marine species. These included black sea bass, *Centropristes striata*; searobins (*Prionotus* spp.); striped cusk-eel, *Ophidion marginatum*; lady crab, *Ovalipes ocellatus*; seabob, *Xiphopenaeus kroyeri*; swimming crabs (*Portunus* spp.); and hermit crabs, *Pugurus longicarpus* and *Clibanarius vittatus*. In fall, stenohaline marine species (Group B) displayed only moderate to low constancy and fidelity for collections at stations J003 and J001 (Fig. 2), while in winter, many of the same species and other marine transient species were still infrequently encountered but highly faithful to collections from station J003 (Fig. 2). Stenohaline marine species, which

Table 2.--Total numbers and total weights of fish species collected 1973-77 in the Cooper River-Charleston Harbor estuarine system, S.C. Species are listed in order of abundance, and data are pooled over the 5-yr sampling period.

Species	Number		Weight	
	Total	%	Total (kg)	%
<i>Stellifer lanceolatus</i>	22,932	33.33	162.882	18.90
<i>Anchoa mitchilli</i>	9,203	13.38	15.957	1.80
<i>Micropogonias undulatus</i>	7,862	11.43	108.374	12.20
<i>Bruvoutia tyrranum</i>	4,848	7.05	62.699	9.31
<i>Larimus minthuna</i>	4,228	6.15	56.200	6.33
<i>Syphurus plagiusa</i>	3,053	4.44	41.633	4.69
<i>Gairdiella chrysoura</i>	3,006	4.37	83.849	9.44
<i>Syngnathus regalis</i>	2,578	3.75	33.005	3.72
<i>Umbrinacirrata</i>	2,250	3.27	30.091	3.39
<i>Theranthes maculatus</i>	1,996	2.90	15.108	1.70
<i>Ictalurus catfish</i>	1,931	2.81	40.623	4.57
<i>Dorosoma petenense</i>	1,060	1.54	5.374	0.60
<i>Alosa aestivalis</i>	513	0.75	1.809	0.20
<i>Menticirrhus americanus</i>	310	0.45	5.057	0.57
<i>Opisturus tau</i>	282	0.41	38.124	4.29
<i>Paralichthys dentatus</i>	250	0.36	9.805	1.10
<i>Periplus alepidotus</i>	239	0.35	2.730	0.31
<i>Paralichthys lethostigmus</i>	215	0.31	39.568	4.45
<i>Periplus trispinatus</i>	184	0.27	1.730	0.19
<i>Chlorosomus chrysourus</i>	132	0.19	0.493	0.06
<i>Opisthomene ogilbini</i>	128	0.19	0.763	0.09
<i>Etrypus croesus</i>	126	0.18	0.842	0.09
<i>Ictalurus punctatus</i>	124	0.18	1.763	0.20
<i>Prionotus carolinus</i>	108	0.16	0.268	0.03
<i>Anchoa hepsetus</i>	103	0.15	0.818	0.09
<i>Pomatomus saltatrix</i>	91	0.13	2.306	0.26
<i>Anguilla rostrata</i>	74	0.11	9.959	1.12
<i>Centropristes philadelphica</i>	70	0.10	2.945	0.33
<i>Citharichthys spilopterus</i>	63	0.09	0.345	0.04
<i>Chaetodipterus faber</i>	56	0.08	0.617	0.07
<i>Scophthalmus aquosus</i>	54	0.08	0.484	0.05
<i>Ictalurus nebulosus</i>	53	0.08	2.557	0.29
<i>Alosa sagittaeina</i>	50	0.07	1.149	0.13
<i>Ancylorhynchus quadrirostrata</i>	43	0.06	0.709	0.08
<i>Trachinurus lepturus</i>	41	0.06	1.690	0.19
<i>Silene vomer</i>	40	0.06	0.377	0.04
<i>Prionotus suillus</i>	32	0.05	0.216	0.02
<i>Arius felis</i>	32	0.05	2.788	0.31
<i>Urophycis floridana</i>	29	0.04	0.978	0.11
<i>Hypsoblennius hentzii</i>	29	0.04	0.236	0.03
<i>Morone saxatilis</i>	23	0.03	0.154	0.02
<i>Opistionichthys marginatus</i>	22	0.03	0.433	0.05
<i>Ictalurus fuscus</i>	22	0.03	0.330	0.06
<i>Silene setapinnis</i>	21	0.03	0.096	0.01
<i>Cynoscion nebulosus</i>	21	0.03	1.037	0.12
<i>Lagodon rhomboides</i>	19	0.03	0.692	0.08
<i>Caranx hippos</i>	18	0.03	0.321	0.04
<i>Centropristes striata</i>	15	0.02	0.437	0.05
<i>Drophycis carilli</i>	12	0.02	0.203	0.02
<i>Dasyatis sabina</i>	12	0.02	11.731	1.32
<i>Pogonias cromis</i>	12	0.02	5.000	0.56
<i>Eucinostomus sp.</i>	11	0.02	0.107	0.01
<i>Leptoscarus cassus</i>	10	0.01	17.662	1.99
<i>Alotes mediterraneus</i>	9	0.01	0.011	0.01
<i>Centroscyllium striatum</i>	9	0.01	0.046	0.01
<i>Synodus microlepis</i>	9	0.01	0.214	0.02
<i>Lutjanus griseus</i>	9	0.01	0.107	0.01
<i>Fucinotomus argenteus</i>	7	0.01	0.062	0.01
<i>Prionotus sp.</i>	7	0.01	0.022	0.01
<i>Prionotus carolinus</i>	7	0.01	0.019	0.01
<i>Menticirrhus undulatus</i>	6	0.01	0.025	0.01
<i>Prionotus aculeatus</i>	5	0.01	0.038	0.01
<i>Morwong hispidus</i>	5	0.01	0.009	0.01
<i>Archosargus probatocephalus</i>	5	0.01	0.156	0.02
<i>Cynoscion notatus</i>	5	0.01	0.035	0.01
<i>Larimus fuscus</i>	4	0.01	0.027	0.01
<i>Astrocopus y-gracilis</i>	4	0.01	0.030	0.01
<i>Mugil curema</i>	4	0.01	0.078	0.01
<i>Cynoscion nebulosus</i>	4	0.01	18.883	2.13
<i>Synodus foetens</i>	4	0.01	0.187	0.02
<i>Brama brama</i>	3	0.01	1.297	0.15
<i>Apogonichthys oxyrhynchus</i>	3	0.01	13.835	1.56
<i>Bagre marinus</i>	3	0.01	0.041	0.01
<i>Chilomycterus orillarum</i>	3	0.01	0.006	0.01
<i>Gobionellus shufeldti</i>	3	0.01	0.006	0.01
<i>Syphodus civitatus</i>	3	0.01	0.026	0.01
<i>Sphoeroides maculatus</i>	3	0.01	0.133	0.01
<i>Perissodus microlepis</i>	3	0.01	0.024	0.01
<i>Lagodon rhomboides</i>	3	0.01	0.106	0.01
<i>Mugil cephalus</i>	3	0.01	0.028	0.01
<i>Gobionellus boliviensis</i>	3	0.01	0.006	0.01
<i>Hymenobranchus ionthas</i>	2	0.01	0.009	0.01
<i>Ictalurus platycephalus</i>	2	0.01	0.335	0.04
<i>Rhinoptera bonariensis</i>	2	0.01	-----	-----
<i>Dorosoma cepedianum</i>	2	0.01	0.035	0.01
<i>Leptoscarus punctatus</i>	1	0.01	0.030	0.01
<i>Elops saurus</i>	1	0.01	0.126	0.01
<i>Morone americana</i>	1	0.01	0.010	0.01
<i>Myctophopeltis microlepis</i>	1	0.01	0.073	0.01
<i>Syngnathus marmoratus</i>	1	0.01	0.177	0.02
<i>Raja eglanteria</i>	1	0.01	0.671	0.08
<i>Caranx ignobilis</i>	1	0.01	1.046	0.12
<i>Leptoscarus curitus</i>	1	0.01	0.008	0.01
<i>Ictalurus natalensis</i>	1	0.01	1.204	0.01
<i>Ictalurus spilus</i>	1	0.01	0.058	0.01
<i>Eleotris pisonis</i>	1	0.01	0.018	0.01
<i>Gobionellus heterotus</i>	1	0.01	0.003	0.01
<i>Sphyraena guachancho</i>	1	0.01	0.001	0.01
<i>Solenops caeruleus</i>	1	0.01	0.004	0.01
<i>Menticirrhus littoralis</i>	1	0.01	0.028	0.01
<i>Ammodytes glomeratus</i>	1	0.01	0.002	0.01
TOTAL	68,796		886.409	

Table 3.—Total numbers and total weights of decapod crustacean species collected 1973–77 in the Cooper River–Charleston Harbor estuarine system, S.C. Species are listed in order of abundance, and data are pooled over the 5-yr sampling period.

Species	Number		Weight	
	Total	%	Total (kg)	%
<i>Penaeus setiferus</i>	80,121	82.62	492.927	69.20
<i>Penaeus aztecus</i>	8,053	8.30	56.757	7.97
<i>Xiphopenaeus kroyeri</i>	2,657	2.74	4.928	0.69
<i>Callinectes sapidus</i>	1,914	1.97	137.936	19.36
<i>Callinectes similis</i>	1,438	1.48	11.029	1.55
<i>Trachypenaeus constrictus</i>	721	0.74	0.912	0.13
<i>Palamonaetes vulgaris</i>	388	0.40	0.199	0.03
<i>Portunus spinimanus</i>	243	0.25	2.308	0.32
<i>Pagurus longicarpus</i>	215	0.22	0.182	0.03
<i>Procturus gibbesii</i>	193	0.20	0.420	0.06
<i>Ovalipes stephensoni</i>	191	0.20	0.646	0.09
<i>Penaeus duorarum</i>	177	0.18	1.256	0.18
<i>Panopeus herbstii</i>	133	0.14	0.358	0.05
<i>Rhithropopenaeus harrisi</i>	82	0.08	0.045	0.01
<i>Palamonaetes pugio</i>	82	0.08	0.047	0.01
<i>Clibanarius vittatus</i>	77	0.08	0.054	0.01
<i>Ovalipes ocellatus</i>	69	0.07	0.426	0.06
<i>Macrobrachium chione</i>	47	0.05	0.165	0.02
<i>Pagurus politcaris</i>	25	0.03	0.086	0.01
<i>Palamonaetes</i> sp.*	24	0.02	0.014	<0.01
<i>Neopanope sayi</i>	20	0.02	0.019	<0.01
<i>Cancer irroratus</i>	19	0.02	0.248	0.03
<i>Alpheus normanni</i>	11	0.01	0.003	<0.01
<i>Menippe mercenaria</i>	10	0.01	1.066	0.15
<i>Libinia emarginata</i>	10	0.01	0.077	0.01
<i>Libinia dubia</i>	7	0.01	0.019	<0.01
<i>Hemipanopeus angustifrons</i>	7	0.01	0.020	<0.01
<i>Alpheus heterochaelis</i>	6	0.01	0.009	<0.01
<i>Palamonaetes intermedius</i>	5	0.01	0.003	<0.01
<i>Sicyonia lasiognatha</i>	5	0.01	0.005	<0.01
<i>Panopeus occidentalis</i>	5	0.01	0.005	<0.01
<i>Portunus</i> sp.*	5	0.01	—	—
<i>Eurypanopeus depressus</i>	4	<0.01	0.004	<0.01
<i>Callinectes ornatus</i>	2	<0.01	0.082	0.01
<i>Echippolytmata opipharoides</i>	3	<0.01	0.004	<0.01
<i>Lysmata wurdemanni</i>	1	<0.01	0.002	<0.01
<i>Sicyonia brevirostris</i>	1	<0.01	0.004	<0.01
<i>Hepatus epitheliticus</i>	1	<0.01	0.060	0.01
<i>Micropanope sp.</i>	1	<0.01	0.001	<0.01
<i>Libinia</i> sp.*	1	<0.01	0.001	<0.01
<i>Lanthididae</i> *	1	<0.01	—	—
<i>Penaeus</i> sp.*	1	<0.01	0.001	<0.01
<i>Procambarus clarkii</i>	1	<0.01	0.006	<0.01
<i>Callinectes</i> sp.*	1	<0.01	0.001	<0.01
TOTAL	96,978		712,330	

*Field identification or damaged specimen(s).

Table 4.—Average numbers of species of fishes and decapod crustaceans collected 1973–77 at stations in the Cooper River–Charleston Harbor estuarine system, S.C. Numbers in parentheses = standard error of the mean; n = number of samples per year.

Year	Average numbers of species/station						Grand Mean
	C001	C002	C003	C004	J003	J001	
1973	4 (0.81)	6 (1.05)	10 (0.89)	13 (1.56)	12 (2.35)	—	10
	n = 11	n = 11	n = 12	n = 11	n = 8	—	
1974	2 (0.81)	6 (1.23)	9 (2.04)	15 (2.14)	15 —	11	
	n = 1	n = 12	n = 12	n = 12	n = 12	—	
1975	— (0.94)	6 (1.53)	11 (1.41)	12 (2.12)	15 (1.79)	16	12
	n = 12	n = 12	n = 12	n = 12	n = 12	—	
1976	— (0.78)	8 (1.40)	12 (1.46)	18 (1.44)	18 (1.66)	16	14
	n = 12	n = 12	n = 12	n = 12	n = 12	—	
1977	— (0.75)	9 (1.38)	12 (1.79)	12 (1.77)	13 (1.27)	19	13
	n = 12	n = 12	n = 12	n = 12	n = 12	—	

Table 5.—Average numbers of individual fish and decapod crustaceans collected 1973–77 at stations in the Cooper River–Charleston Harbor estuarine system, S.C. Numbers in parentheses = standard error of the mean; n = number of samples per year.

Year	Average numbers of individuals/station						Grand Mean
	C001	C002	C003	C004	J003	J001	
1973	21 (6.70)	355 (218.97)	867 (429.07)	1,195 (473.38)	456 (146.88)	—	718
	n = 11	n = 11	n = 11	n = 11	n = 8	—	
1974	2 (43.92)	158 (309.96)	541 (94.50)	533 (180.18)	619 —	—	463
	n = 1	n = 12	n = 12	n = 12	n = 12	—	
1975	— (47.58)	80 (369.91)	554 (123.91)	519 (355.84)	1,233 (346.78)	1,467	771
	n = 12	n = 12	n = 12	n = 12	n = 12	n = 12	
1976	— (136.55)	319 (397.20)	715 (196.60)	1,018 (149.82)	688 (225.00)	922	732
	n = 12	n = 12	n = 12	n = 12	n = 12	n = 12	
1977	— (65.86)	206 (50.67)	263 (48.07)	171 (119.31)	299 (191.41)	503	288
	n = 12	n = 12	n = 12	n = 12	n = 12	n = 12	
Grand mean	229.4	586	687.2	639	964		

Table 6.--Species groups formed from seasonal cluster analyses of fishes and decapod crustaceans collected in the Cooper River-Charleston Harbor estuarine system, S.C., 1973-77.

FALL	WINTER	SPRING	SUMMER
Group A	Group A	Group A	Group A
<i>Callinectes sapidus</i> <i>Syphurus plagiatus</i> <i>Cynoscion regalis</i> <i>Saindiella chrysoura</i> <i>Anchoa mitchilli</i> <i>Stellifer lanceolatus</i> <i>Penaeus setiferus</i>	<i>Sicyonia laevigata</i> <i>Libinia dubia</i> <i>Mugil curema</i> <i>Neopanope sayi</i> <i>Alpheus normanni</i> <i>Libinia emarginata</i> <i>Prionotus acutulus</i> <i>Prionotus evolans</i>	<i>Stellifer lanceolatus</i> <i>Penaeus setiferus</i> <i>Urophycis regia</i> <i>Trachypenaeus constrictus</i> <i>Callinectes similis</i> <i>Brevoortia tyrannus</i> <i>Peprilus triacanthus</i> <i>Micropanogonias undulatus</i> <i>Leiostomus xanthurus</i> <i>Anchoa mitchilli</i> <i>Callinectes sapidus</i> <i>Syphurus plagiatus</i> <i>Trinectes maculatus</i>	<i>Leiostomus xanthurus</i> <i>Micropanogonias undulatus</i> <i>Cynoscion regalis</i> <i>Penaeus setiferus</i> <i>Anchoa mitchilli</i> <i>Penaeus setiferus</i> <i>Callinectes similis</i> <i>Trachypenaeus constrictus</i> <i>Stellifer lanceolatus</i>
Group B	Group B	Group B	Group B
<i>Hypsoblennius hentzii</i> <i>Ophidion marginatum</i> <i>Prionotus tribulus</i> <i>Penaeus duorarum</i> <i>Ovalipes ocellatus</i> <i>Portunus spinimanus</i> <i>Pagurus longicarpus</i> <i>Clibanarius vittatus</i> <i>Callinectes similis</i> <i>Menticirrhus americanus</i> <i>Trachypenaeus constrictus</i> <i>Portunus gibbesii</i> <i>Xiphopenaeus kroyeri</i>	<i>Centropristes striata</i> <i>Urophycis floridana</i> <i>Centropristes philadelphica</i> <i>Callinectes similis</i> <i>Xiphopenaeus kroyeri</i> <i>Portunus gibbesii</i> <i>Portunus spinimanus</i> <i>Trachypenaeus constrictus</i> <i>Gobiesox strumosus</i> <i>Ovalipes ocellatus</i> <i>Menticirrhus americanus</i> <i>Portunus gibbesii</i> <i>Xiphopenaeus kroyeri</i>	<i>Opaneus tau</i> <i>Paralichthys dentatus</i> <i>Penaeus aztecus</i> <i>Cynoscion regalis</i> <i>Bairdiella chrysoura</i> <i>Palamonetes vulgaris</i>	<i>Pagurus longicarpus</i> <i>Portunus gibbesii</i> <i>Portunus spinimanus</i> <i>Ophidion marginatum</i> <i>Pagurus pollicaris</i> <i>Ovalipes ocellatus</i>
Group C	Group C	Group C	Group C
<i>Anchoa hepsetus</i> <i>Chloroscombrus chrysurus</i> <i>Selene setapinnis</i> <i>Peprilus alepidotus</i> <i>Opisthonema oglinum</i> <i>Dasyatis sabina</i> <i>Arius felis</i> <i>Rhithropanopeus harrietti</i> <i>Chaetodipterus faber</i> <i>Penaeus aztecus</i>	<i>Anchoa mitchilli</i> <i>Brevoortia tyrannus</i> <i>Micropanogonias undulatus</i> <i>Bairdiella chrysoura</i> <i>Dorosoma petenense</i> <i>Penaeus setiferus</i> <i>Stellifer lanceolatus</i> <i>Leiostomus xanthurus</i> <i>Syphurus plagiatus</i> <i>Callinectes sapidus</i> <i>Paralichthys lethostigma</i> <i>Urophycis regia</i>	<i>Ancylomenes quadrocellata</i> <i>Clibanarius vittatus</i> <i>Pagurus longicarpus</i> <i>Prionotus tribulus</i> <i>Penaeus duorarum</i> <i>Prionotus carolinus</i>	<i>Palamonetes pugio</i> <i>Palamonetes vulgaris</i> <i>Paralichthys lethostigma</i> <i>Paralichthys dentatus</i> <i>Opaneus tau</i> <i>Callinectes sapidus</i> <i>Syphurus plagiatus</i> <i>Trinectes maculatus</i> <i>Bairdiella chrysoura</i> <i>Brevoortia tyrannus</i>
Group D	Group D	Group D	Group D
<i>Gobiesox strumosus</i> <i>Cynoscion nebulosus</i> <i>Selene vomer</i> <i>Pomatomus saltatrix</i>	<i>Dasyatis sabina</i> <i>Centropristes striata</i> <i>Arius felis</i> <i>Citharichthys spilopterus</i>	<i>Anchoa hepsetus</i> <i>Ovalipes ocellatus</i> <i>Trichurus lepturus</i>	<i>Peprius alepidotus</i> <i>Chloroscombrus chrysurus</i> <i>Opisthonema oglinum</i> <i>Anchoa hepsetus</i> <i>Selene setapinnis</i>
Group E	Group E	Group E	Group E
<i>Palamonetes vulgaris</i> <i>Eucinostomus argenteus</i> <i>Palamonetes pugio</i> <i>Caranx hippos</i> <i>Alosa aestivalis</i> <i>Dorosoma petenense</i> <i>Lutjanus griseus</i>	<i>Etropus crossotus</i> <i>Cynoscion regalis</i> <i>Hypsoblennius hentzii</i> <i>Scophthalmus aquosus</i> <i>Palamonetes pugio</i> <i>Lagodon rhomboides</i> <i>Prionotus tribulus</i> <i>Alosa aestivalis</i> <i>Nemidius menidia</i> <i>Clibanarius vittatus</i> <i>Alosa sapidissima</i> <i>Ancylomenes quadrocellata</i>	<i>Dorosoma petenense</i> <i>Prionotus volvus</i> <i>Rhithropanopeus harrietti</i> <i>Palamonetes pugio</i> <i>Pomatomus saltatrix</i> <i>Alosa aestivalis</i> <i>Lepisosteus osseus</i> <i>Alosa sapidissima</i> <i>Urophycis floridana</i> <i>Anguilla rostrata</i> <i>Pomacanthus occidentalis</i> <i>Scophthalmus aquosus</i> <i>Prionotus sp.</i>	<i>Selene vomer</i> <i>Trichurus lepturus</i> <i>Pomatomus saltatrix</i> <i>Pomacanthus herbeti</i> <i>Prionotus tribulus</i> <i>Penaeus duorarum</i> <i>Menticirrhus americanus</i> <i>Chaetodipterus faber</i> <i>Neopanope sayi</i>
Group F	Group F	Group F	Group F
<i>Paralichthys lethostigma</i> <i>Anguilla rostrata</i> <i>Ictalurus fuscus</i> <i>Eucinostomus sp.</i> <i>Leiostomus xanthurus</i> <i>Trinectes maculatus</i> <i>Ictalurus catfish</i> <i>Etropus crossotus</i> <i>Centropristes philadelphica</i> <i>Paralichthys dentatus</i> <i>Opaneus tau</i> <i>Nemidius menidia</i> <i>Micropanogonias undulatus</i> <i>Brevoortia tyrannus</i>	<i>Penaeus aztecus</i> <i>Palamonetes intermedius</i> <i>Rhithropanopeus harrietti</i> <i>Pogonias cromis</i> <i>Archosargus probatocephalus</i> <i>Pomacanthus herbeti</i> <i>Cynoscion nebulosus</i> <i>Palamonetes vulgaris</i> <i>Opaneus tau</i>	<i>Ictalurus punctatus</i> <i>Ictalurus nebulosus</i> <i>Anguilla rostrata</i> <i>Macrobrachium ohione</i> <i>Percis flavescens</i> <i>Morone saxatilis</i> <i>Lepisosteus osseus</i> <i>Alpheus heterochelus</i> <i>Ictalurus catfish</i> <i>Trinectes maculatus</i> <i>Acipenser oxyrinchus</i> <i>Ictalurus platycephalus</i>	<i>Dorosoma petenense</i> <i>Alosa aestivalis</i> <i>Morone saxatilis</i> <i>Peprilus triacanthus</i> <i>Scomberomorus maculatus</i> <i>Rhithropanopeus harrietti</i> <i>Macrobrachium ohione</i> <i>Ictalurus catfish</i>
Group G	Group G	Group G	Group G
		<i>Ophidion marginatum</i> <i>Pagurus pollicaris</i> <i>Ovalipes stephani</i> <i>Portunus spinimanus</i> <i>Portunus gibbesii</i> <i>Menticirrhus americanus</i>	







Figure 2.—Two-way coincidence tables of constancy and fidelity which compare species associations among sampling stations in the Cooper River-Charleston Harbor for fall, winter, spring, and summer collections, 1973-77. See Table 6 for species group lists.



were restricted in spring to collections at higher salinity stations but displayed only moderate to low constancy there, composed assemblages C, D, E, and H (Fig. 2). Species in these groups included fourspot flounder, *Ancylopsetta quadrocincta*; searobins (*Prionotus* spp.); black sea bass, *Centropristes striata*; lady crabs, *Ovalipes ocellatus* and *O. stephensi*; Atlantic cutlassfish, *Trichiurus lepturus*; and striped cusk-eel, *Ophidion marginatum*. Many of the same species composed groups B and C from our cluster analysis of summer data and were restricted but infrequently encountered at collections from stations J003 and J001 (Fig. 2).

The only species restricted to samples from the lowest salinity stations (C001, C002) formed group G in spring (Fig. 2). However, the resident estuarine species, *Ictalurus punctatus*, *I. catus*, and *Macrobrachium ohione*, which composed this group displayed only moderate constancy for collections from these low-salinity stations.

Ubiquitous species were present during all seasons in the Cooper River-Charleston Harbor estuarine system. These species included the numerically dominant fishes and decapod crustaceans. Although their penetration extended as far upriver as stations C002-C001, species in these assemblages were generally most constant in collections from stations J003, J001, and C004. In the fall, members of groups A and F were encountered at all stations; but only members of group A were consistently collected, as denoted by their very high constancy, at collections from stations J003, J001, C004, and C003 (Fig. 2). In our analysis of winter collections, only members of group C were eurytopic in station location. Species in this group were consistently represented in collections at stations J003, J001, and C004, but they were not restricted to these stations (Fig. 2). Members of group A in spring were generally found at all sites but were most consistently encountered at stations J003, J001, and C004 (Fig. 2). Our analysis of summer data showed that euryhaline species of Group A were con-

sistently present in collections from stations J003, J001, C004, and C003.

Other assemblages defined by our analysis included species tolerant of a wide salinity range and not restricted to any station location, but generally of low density. Included in these groups were anadromous species, *Alosa aestivalis* and *A. sapidissima*; the American eel, *Anguilla rostrata*; and the caridean shrimps, *Palaemonetes pugio* and *P. vulgaris*.

Temporal and Spatial Distributions

Patterns of distribution for the most abundant species of fishes at each station for each month of collection are shown in Figure 3, and fluctuations in their abundance over the 5-yr sampling period are shown in Figure 4. Length-frequency plots which were generated for selected species are not shown but are available from the authors upon request.

Stellifer lanceolatus, Star drum.—Star drum were most abundant at higher salinity stations C004, J001, and J003, whereas catches were negligible at stations farther upriver (C001, C002) (Fig. 3A). This species displayed seasonality in its abundance, with most individuals collected October through May. Catches of *S. lanceolatus* also underwent considerable annual variation and were greatest during the years 1974-76 (Fig. 4A). Length-frequency polygons for star drum over the 5-yr sampling period suggested a consistent influx of small fish (< 80 mm TL) into the population during summer with recruitment continuing into fall. Frequencies of these small fish were lower during winter and spring. Our results are consistent with those of Welsh and Breder (1923), Dahlberg and Odum (1970), and Shealy et al. (1974), who also noted that recruitment of young fish first occurred during summer after late-spring and early-summer spawning.




Figure 3.—Abundance, expressed as the antilog of the transformed $\log_{10}(x + 1)$ mean number of individuals, of 10 major fish species collected monthly in the channel of the Cooper River-Charleston Harbor estuarine system, 1973-77. Legend indicates four arbitrary levels of abundance, from rare or absent (0-1) to maximum abundance (51-315).

Anchoa mitchilli, Bay anchovy.—*Anchoa mitchilli* were collected at all stations in the Cooper River-Charleston Harbor system, except for the most limnetic station (C001) (Fig. 3B). Bay anchovy were most abundant at station J001. Monthly fluctuations in abundance indicated that little, if any, seasonality was associated with catches of bay anchovy, but catches did undergo annual fluctuations, being highest during 1974-76 (Fig. 4B).

Micropogonias undulatus, Atlantic croaker.—Atlantic croaker were collected at all stations in the Cooper River-Charleston Harbor system, but abundances were greatest from April through July at higher salinity stations, particularly those located near the mouth (J001, J003) (Fig. 3C). Annual variation in catches of croaker was small, with little fluctuation about the grand mean (Fig. 4C). Length-frequency distributions indicated that most estuarine croaker available to our trawls were <120 mm TL throughout the year. Although these smaller fish predominated in spring and summer catches, they were also present during other seasons, but in fewer numbers. Newly recruited fish (<30 mm TL) generally appeared first in fall and continued to appear in the population during winter and spring. The continued presence of small croaker during spring in South Carolina may reflect the slow growth of fish spawned in late winter or early spring (Chao and Musick 1977). By summer, few croaker <45 mm were collected from the Cooper River-Charleston Harbor system, and modal groups were in the 75-90 mm range. Although 1-yr-old fish (108-285 mm, from Chao and Musick 1977) were infrequently caught, probably due to gear avoidance (Wenner et al. 1982), their numbers were fewer in summer. Migration of yearling croaker from the estuarine environment during summer has been reported in the York River, Virginia (Chao and Musick 1977), and late summer and early fall in South Carolina (Bearden 1964) and Florida (Hansen 1969).

Brevoortia tyrannus, Atlantic menhaden.—Atlantic menhaden were collected at every station except for C001, the station farthest upriver (Fig. 3D). Menhaden displayed little change in abundance by month but were most consistently present in November, December, January, and June. Annual catches fluctuated moderately about the grand mean and were greatest in 1977 (Fig. 4D).

Leiostomus xanthurus, Spot.—Spot exhibited a distributional pattern similar to that of Atlantic croaker, being most abundant at stations J001 and J003 near the mouth. Spot also were most abundant during May-July (Fig. 3E). Annual catches of spot steadily increased over the 5-yr sampling period (Fig. 4E). The average size of spot was greatest in fall and winter. Length-frequency distributions indicated that spring and summer catches of spot were dominated by fishes in the 60-80 mm size range. Our data support results of other studies in South Carolina (Dawson 1958; Shealy et al. 1974), North Carolina (Hildebrand and Cable 1930), and the lower Chesapeake Bay (Hildebrand and Schroeder 1928; Chao and Musick 1977), which found that young-of-the-year spot first entered the estuary in April.

Syphurus plagiura, Blackcheek tonguefish.—*Syphurus plagiura* were most abundant in higher salinity areas of the Cooper River-Charleston Harbor system, although the species did penetrate into limnetic-mesohaline areas of the estuaries at station C003 (Fig. 3F). Over the 5-yr sampling period, abundance of *S. plagiura* was greatest in October and November; however, annual catches showed a consistent increase from 1973 to 1976 with a slight decrease in 1977 (Fig. 4F).

Bairdiella chrysoura, Silver perch.—Silver perch were collected at all stations except C001; and their abundance increased at stations downriver, especially at C004 and J001




Figure 4.—Annual variation in the transformed $[\log_{10}(x + 1)]$ mean number of individuals of 10 major fish species collected in the Cooper River-Charleston Harbor estuarine system, 1973-77.

(Fig. 3G). Although silver perch were present in the Cooper River-Charleston Harbor system throughout the year, abundance of the species was greatest from August through January. Annual catches decreased in 1977 (Fig. 4G). The presence of silver perch throughout the year in estuaries of South Carolina (Shealy et al. 1974; Wenner et al. 1982) differs from the seasonal pattern observed in Chesapeake Bay. Chao and Musick (1977) noted that most silver perch leave the York River estuary of Virginia by November. They collected no silver perch from January to March and suggested that the year-round presence of the species in estuaries south of Chesapeake Bay may be due to the higher salinity or temperature of those waters.

Cynoscion regalis, Weakfish.—Weakfish were collected at all stations except C001 and were most abundant during summer (Fig. 3H). Annual catches were fairly constant with little variation about the grand mean (Fig. 4H). The average length of weakfish did not differ markedly by season, but length-frequency distributions showed that the modal length for spring catches was usually smaller than for other seasons. This reduction in size is probably caused by increasing numbers of young-of-the-year weakfish, newly recruited from the May-August spawning period (Lunz and Schwartz 1970).

Urophycis regia, Spotted hake.—Spotted hake displayed the most seasonality in its distribution and abundance, being collected from February to May only (Fig. 3I). Their absence from South Carolina estuaries during the rest of the year is attributed to offshore migration to deeper water during warmer months (Hildebrand and Cable 1938). Spotted hake

were collected at the most seaward stations, with maximum abundance occurring at the mouth of the estuary. Little variation in annual catches of spotted hake was present in our samples (Fig. 4I).

Trinectes maculatus, Hogchoker.—*Trinectes maculatus* were collected sometime during the year at every station in the Cooper River-Charleston Harbor system. Hogchoker were most consistently abundant at stations upriver (C002 and C003) and displayed no apparent seasonality in abundance (Fig. 3J). Annual catches of hogchoker increased over the 5-yr sampling period and were greatest in 1976 and 1977 (Fig. 4J).

Penaeus setiferus, White shrimp.—Catches of white shrimp were seasonal with most individuals occurring late summer through fall. White shrimp were also most numerous at the downriver stations (Fig. 5A). Length-frequency distributions indicated that young-of-the-year white shrimp were present in the Cooper River-Charleston Harbor system during summer and fall. Most shrimp collected during these seasons were <80 mm TL, except in 1977 when a few shrimp >120 mm TL were collected in summer and fall. The absence of young-of-the-year shrimp in 1977 is also reflected in the annual catch data for that year when a marked decrease in abundance of white shrimp occurred (Fig. 6A). Modal lengths of white shrimp generally increased during spring to >100 mm TL. The larger size of shrimp during spring is attributable to shoreward migration of large shrimp from offshore waters (Williams 1955) or to the growth of overwintering shrimp to subadult size (Bishop and Shealy 1977).




Figure 5.—Abundance, expressed as the antilog of the transformed $\log_{10}(x + 1)$ mean number of individuals at each station each month, of four major species of decapod crustaceans collected in the channel of the Cooper River-Charleston Harbor estuarine system, 1973-77. Legend indicates four arbitrary levels of abundance, from rare or absent (0-1) to maximum abundance (101-1,340).




Figure 6.—Annual variation in the transformed $\log_{10}(x + 1)$ mean number of individuals of four major decapod crustacean species collected in the Cooper River-Charleston Harbor estuarine system, 1973-77.

Penaeus aztecus, Brown shrimp.—Brown shrimp were highly seasonal in their occurrence within the Cooper River-Charleston Harbor estuarine system. They were collected May to September and were most abundant during June and July at higher salinity stations (Fig. 5B). Annual catch rates were variable and highest in 1974 and 1976 (Fig. 6B). Juvenile brown shrimp entered the estuary in spring, remained through the summer, and were almost totally absent from fall and winter collections. This seasonal abundance pattern has also been noted in other South Carolina estuaries (Bishop and Shealy 1977), although the absence of brown shrimp from

these estuaries during winter is probably due to gear bias for larger-sized shrimp (Wenner et al. 1982).

Xiphopenaeus kroyeri, Seabob.—Seabob were limited to higher salinity areas of the estuary during fall and winter (Fig. 5C). Annual catches were low, and no seabob were collected in 1973 and 1977 (Fig. 6C). Although *X. kroyeri* occurs in the lower portion of estuaries, it is most commonly encountered in the near offshore coastal zone (Gunter 1950).

Callinectes sapidus, Blue crab.—Blue crab were collected throughout the Cooper River-Charleston Harbor system, but occurred year-round only at stations C004 and J001 (Fig. 5D). Crab were also least abundant at stations upriver. Annual catches increased over the 5-yr sampling period except for a slight decline in 1977 (Fig. 6D). Size-frequency distributions of blue crab covered a wide range of sizes from 10 to 100 mm carapace width with small crab (< 60 mm CW) present during all seasons. Average sizes of blue crab were generally larger in spring and summer.

Biomass Estimates

Biomass and density for fishes were greatest at higher salinity stations J001 and C004 during winter and spring (Table 7). Increased values during these time periods were coincident with the increased dominance of catches by *Stellifer lanceolatus*, *Brevoortia tyrannus*, and *Micropogonias undulatus*. Decapod biomass and density were greatest at station J001 in Charleston Harbor and during fall and summer for all stations combined. These seasonal peaks coincided with periods when young-of-the-year shrimp became vulnerable to our trawl gear.

Our mean total biomass and density estimates for all seasons and stations sampled in the Cooper River-Charleston Harbor system over the 5-yr study period were:

	Biomass (kg/ha)	Density (no./ha)
Fishes	6.04	471
Decapods	4.98	678

These estimates are comparable to those obtained by Wenner et al. (1982) for estuarine portions of the Santee River system of South Carolina and by Shealy et al. (1974) for estuarine portions of the Cooper River-Charleston Harbor and Edisto River systems.

Table 7.—Average seasonal biomass (kg/ha) and density (no./ha) of fishes and decapod crustaceans collected at stations in the Cooper River-Charleston Harbor estuarine system, S.C., 1973-77.

Season	Average biomass and density/station										Grand mean	
	C002		C003		C004		J001		J003			
	kg/ha	no./ha	kg/ha	no./ha	kg/ha	no./kg	kg/ha	no./ha	kg/ha	no./ha	kg/ha	no./ha
Fishes												
Fall	3.62	166	2.63	202	7.55	592	11.03	542	4.30	888	5.37	472
Winter	3.53	175	3.78	258	16.48	621	14.52	976	6.52	558	8.60	480
Spring	1.75	428	5.25	242	13.94	763	11.99	1,071	6.06	636	7.45	569
Summer	0.86	97	2.44	289	2.49	212	5.30	602	4.59	593	2.95	346
Decapods												
Fall	1.57	521	6.71	1,122	9.35	1,380	15.60	1,993	6.68	915	7.31	1,116
Winter	0.05	8	0.22	17	0.77	143	2.03	314	6.40	931	1.74	260
Spring	0.06	10	0.53	58	5.10	376	17.80	1,270	2.27	287	4.26	339
Summer	0.90	221	12.20	2,078	5.26	928	12.10	1,288	3.18	338	6.28	943

DISCUSSION

The Cooper River-Charleston Harbor estuarine system is characterized as mixohaline with gradual changes in faunal assemblages. The most striking differences in species composition occurred between those stations located at or near the mouth of the estuary and those located far upriver. The fish and decapod crustacean species assemblages associated with these two areas were primarily composed of stenohaline marine species and low-salinity resident estuarine species, respectively. Nevertheless, euryhaline species, which extended from the mouth of the estuary into brackish waters, were the dominant faunal component throughout the estuary as a whole. Except for a few freshwater species, resident estuarine species (e.g., *Trinectes maculatus*, *Anchoa mitchilli*, *Palaemonetes pugio*, *Ictalurus catus*) were found throughout the system, their distributions often overlapping with those of species derived from the marine environment. This distributional pattern is similar to that described by Weinstein et al. (1980) who noted considerable overlap in distributional patterns of resident fishes and stenohaline marine transients in the Cape Fear River, N.C.

The observed overlapping spatial distributional patterns of resident and transient fishes and decapod crustaceans can be related to salinity regimes within the estuary and to the physiological tolerances of component estuarine species to these regimes. In comparison with estuaries of the Middle Atlantic states, such as Chesapeake Bay, South Carolina estuaries are narrower, deeper, and shorter in length (Mathews and Shealy 1978). The physiography of estuaries (Pritchard 1954), in addition to other factors such as runoff, tidal action, and current velocity (Mathews and Shealy 1978), affect vertical mixing and, consequently, determine salinity regimes as well. The combined effect of these factors in South Carolina estuaries is a compression of the isohalines, with resultant overlap in the distributional patterns of many estuarine species. Ultimately, however, it is the physiological tolerances of component estuarine species which really determine their distribution. The spatial limits of freshwater species are maintained through physiological constraints, while other resident estuarine species are able to tolerate a wider range of salinity and apparently are not limited by competition and predation to the lower reaches of the estuary (Weinstein et al. 1980). Physiological tolerances are also important in determining the upestuary limits of species which are numerically dominant in the Cooper River-Charleston Harbor system. For the most part, these species were unable to penetrate into areas where the isohalines were $\leq 0.5^{\circ}/\text{oo}$ and were generally most abundant at stations in the mesopolyhaline zone.

The overlapping spatial distributions of many resident estuarine, stenohaline marine, and numerically dominant euryhaline species in the Cooper River-Charleston Harbor system are reflected in the greater species richness and abundance of individuals at stations in the mesopolyhaline zone. Assemblages at these stations were comparatively diverse, consisting of some resident estuarine and euryhaline species and many stenohaline marine species. Seasonal peaks in species diversity are largely attributable to those stenohaline marine transients which occur sporadically in low densities throughout the lower reaches of the Cooper River-Charleston Harbor system. Biological interactions such as predation and competition for space and food can also contribute to species

diversity and richness. Weinstein et al. (1980) noted that increased predation pressure probably enhanced species diversity in downstream marsh areas of the Cape Fear River, N.C., by preventing dominant competitors from monopolizing the major food and space resource. An alternative explanation is that enough food may be present in the lower reaches of the river to support a high diversity of species. Euryhaline species such as the sciaenids were numerically dominant in the Cooper River-Charleston Harbor system and were most abundant at downriver stations. Juvenile sciaenids feed opportunistically on a variety of infaunal and demersal species (Chao and Musick 1977). Their successful coexistence in higher salinity areas with stenohaline marine and other estuarine species may be attributed to utilization of food resources from different levels of the water column and to the abundant food resources of the estuarine system. In this case, food would not be a limiting resource and intrafamilial or interspecific competition would not be as important a factor (Chao and Musick 1977).

Temporal distributional patterns were another important aspect of the fish and decapod community of the Cooper River-Charleston Harbor system. Temporal changes in species associations and abundance were related primarily to fluctuations in abiotic variables. Bottom-water temperature in the channel of the estuarine system exerted a substantial influence on the abundance of species collected. The most noticeable decreases in abundance of fishes and decapods coincided with annual minimum temperatures, especially those experienced during the extremely harsh winter of 1977. These seasonal trends in abundance were especially evident for the sciaenid fishes and penaeid shrimps. For those species which may overwinter in the estuary, such as *Micropogonias undulatus* and *Penaeus* spp., extremely low winter temperatures can destroy an entire year-class (Massmann 1971; Farmer et al. 1977). Thus, temperature-related mortalities, as well as emigration of juveniles, probably contributed to the low abundance and biomass observed at that time. Similar explanations were suggested by Weinstein (1979) for decreased abundance of *Penaeus* spp. in the Cape Fear River, N.C.

Seasonal differences in species assemblages reflected changes in abundance as well as exclusion of some species from the estuary during part of the year. However, most species remained in the estuary throughout the year, while their abundances changed seasonally. Nevertheless, while faunal affinities varied throughout the year, as indicated by cluster analysis, the species composition of the estuarine system as a whole was not altered appreciably. Temporal fluctuations in abundance appear to be a means through which more species are able to utilize the estuary simultaneously by a reduction in densities and competition for food and space.

The importance of abiotic factors in determining the distributional patterns of estuarine biota has elicited concern about the effects of redirection on the integrity of species assemblages and, more importantly, on interspecific balance (Shealy and Bishop 1979). A restriction of freshwater inflow will probably cause salinities to be higher and, consequently, modify the existing salinity gradient. Additional consequences of a decrease in flow rate might include a decrease in nutrient and detritus influx, lowering of the water table, reduction in water turbidity, alteration of estuarine circulation, and reduction in the ability of organisms to withstand stresses of normal drought periods (Heald 1970; Keiser and Aldrich 1976). These alterations, should they occur in the Cooper River-Charleston

Harbor estuary, will undoubtedly affect the suitability of the estuary as a nurseryground.

A reduction of freshwater inflow will eventually increase the homeohalinity of this estuarine system. A displacement of the current mesopolyhaline zone further upstream will affect the distribution and abundance of larval and juvenile fishes and shrimps. Upstream marshes are critical areas for early developmental stages of fishes and shellfish (Weinstein 1979). The inflow of freshwater, which currently inundates marshes and abandoned rice fields in the Cooper River-Charleston Harbor estuarine system is more important in maintaining upstream marsh habitat suitable for fishes and decapod crustaceans. After rediversion, much of this habitat, currently subject to overflow, will no longer be available as a nursery due to lowered water levels and higher salinities (U.S. Army Corps of Engineers footnote 2). A substantial reduction in nursery habitat could affect the entire estuarine foodweb. Estuarine salt marshes are highly productive, being dominated by cordgrass (*Spartina*) which ultimately provides a source of food to organisms in the estuarine system (Massmann 1971). Thus, the nursery functions of estuaries are closely related to the viability of plant communities. Alteration of areas which supply much plant detritus may lower the numbers of detritus-algae consumers which, in turn, will eventually limit subsequent trophic levels. This may be particularly troubling from an economic point of view because the abundance of commercially valuable penaeid shrimp is directly related to the absolute area and type of estuarine-intertidal vegetation (Turner 1977).

The habitat of stenohaline marine species may not be affected deleteriously by rediversion. In fact, these species will probably penetrate even farther upriver than they currently do. Because numbers of species and individuals of fishes and decapod crustaceans are now higher in more saline reaches of the river, species diversity of areas in the Cooper River-Charleston Harbor which are currently lower in salinity could be increased by rediversion. Increases in diversity probably will be attributed to higher numbers of stenohaline marine species rather than euryhaline or resident estuarine species; however, many estuarine species, whether resident or transient, are living near the limit of their physiological tolerance of temperature or salinity, so further alteration of the environment may exclude some species permanently (Odum 1970).

In addition to changes in diversity, the species assemblages as we have defined them by station location will probably be altered following rediversion. Whether this alteration will entail a mere shifting of assemblages upriver or the introduction of completely different groupings of species will depend on the effects of rediversion on competition and predation. Food resources can be limiting in estuaries (Lasker 1975; Houda 1978; Laurence 1977). If habitat is lessened and the opportunity for spatial segregation becomes minimal, then seasonality and other forms of temporal segregation may be the only means of reducing competition among species with similar food requirements (Weinstein 1979). Seasonality, which includes differences in spawning periods as well as density-independent factors such as temperature, dissolved oxygen, and nutrient inputs, may mitigate any changes in competition or predation (Enright 1976) precipitated by man-made perturbations. In turn, the adverse effects of rediversion on the estuarine biota may be neither drastic nor irreversible, although this remains to be seen.

ACKNOWLEDGMENTS

We are grateful to all persons who assisted in collection of specimens and data in the field, especially J. Miglarese, R. Richter, and J. Bishop. We thank T. D. Mathews and K. H. Austin for analysis of water samples; K. Swanson for preparation of figures; N. Kopacka for assistance with computer programming; and B. J. Ashby and L. Hodges for typing and retyping the manuscript. The field study and manuscript benefited from comments by our colleagues: E. B. Joseph, V. G. Burrell, Jr., G. R. Sedberry, and C. Bearden.

The data were collected and automated by financial support from the Coastal Plains Regional Commission (Contract 10340031) as part of the Estuarine Survey Program of the South Carolina Marine Resources Center and were analyzed and published through funding from NOAA Office of Sea Grant (Grant 00140-93027).

LITERATURE CITED

ANONYMOUS.
1958. Symposium on the classification of brackish waters. The Venice system for the classification of marine waters according to salinity. *Oikos* 9:311-312.

BEARDEN, C. M.
1964. Distribution and abundance of Atlantic croaker, *Micropogonias undulatus*, in South Carolina. *Contrib. Beauf Bluff Lab.* 40, 23 p.

BISHOP, J. M., and M. H. SHEALY, Jr.
1977. Biological observations on commercial penaeid shrimps caught by bottom trawl in South Carolina estuaries—February 1973-January 1975. *S.C. Mar. Resour. Cent. Tech. Rep.* 25, 97 p.

BOESCH, D. F.
1977. Application of numerical classification in ecological investigations of water pollution. *Va. Inst. Mar. Sci. Spec. Sci. Rep.* 77, 125 p.

CHAO, L. N., and J. A. MUSICK.
1977. Life history, feeding habits, and functional morphology of juvenile sciaenid fishes in the York River estuary, Virginia. *Fish. Bull., U.S.* 75:657-702.

CLIFFORD, H. T., and W. STEPHENSON.
1975. An introduction to numerical classification. *Acad. Press, N.Y.* 229 p.

DAHLBERG, M. D., and E. P. ODUM.
1970. Annual cycles of species occurrence, abundance, and diversity in Georgia estuarine fish populations. *Am. Mid. Nat.* 83:382-392.

DAWSON, C. E.
1958. Study of the biology and life history of the spot, *Leiostomus xanthurus* Lacépède, with special reference to South Carolina. *Contrib. Beauf Bluff Lab.* 28, 48 p.

ELLIOTT, J. M.
1977. Some methods for the statistical analysis of samples of benthic invertebrates, 2nd ed. *Freshwater Biol. Assoc., Sci. Publ.* 25, 160 p.

ENRIGHT, J. T.
1976. Climate and population regulation: The biogeographer's dilemma. *Oecologia (Berl.)* 24:295-310.

FARMER, C. H., III, C. W. BOARDMAN, and J. D. WHITAKER.
1977. The South Carolina shrimp fishery. *S.C. Wildl. Mar. Resour. Div., Educ. Rep.* 8, 49 p.

GUNTER, G.
1950. Seasonal population changes and distributions as related to salinity, of certain invertebrates of the Texas coast, including commercial shrimp. *Publ. Inst. Mar. Sci., Univ. Tex.* 1(2):7-31.

HANSEN, D. J.
1969. Food, growth, migration, reproduction, and abundance of pinfish, *Lagodon rhomboides*, and Atlantic croaker, *Micropogonias undulatus*, near Pensacola, Florida, 1963-65. *Fish. Bull., U.S.* 68:135-146.

HEALD, E. J.
1970. The everglades estuary: An example of seriously reduced inflow of freshwater. *Trans. Am. Fish. Soc.* 99:847-848.

HILDEBRAND, S. F., and L. E. CABLE.
1950. Development and life history of fourteen teleostean fishes at Beaufort, N.C. *Bull. U.S. Bur. Fish.* 46:383-408.

1938. Further notes on the development and life history of some teleosts at Beaufort, N.C. Bull. U.S. Bur. Fish. 48:505-642.

HILDEBRAND, S. F., and W. C. SCHROEDER. 1928. Fishes of Chesapeake Bay. Bull. U.S. Bur. Fish. 43(1), 366 p.

HOODE, E. D. 1978. Critical food concentrations for larvae of three species of subtropical marine fishes. Bull. Mar. Sci. 28:395-411.

KEISER, R. K., Jr., and D. V. ALDRICH. 1976. Salinity preference of postlarval brown and white shrimp (*Penaeus aztecus* and *P. setiferus*) in gradient tanks. Texas A & M Sea Grant Publ. TAMU-75-208, 260 p.

LAMBERT, J. M., and W. T. WILLIAMS. 1962. Multivariate methods in plant ecology. IV. Nodal Analysis. J. Ecol. 50:775-802.

LANCE, G. N., and W. T. WILLIAMS. 1967. A general theory of classificatory sorting strategies. I. Hierarchical systems. Comput. J. (Lond.) 9:373-380.

LASKER, R. 1975. Field criteria for survival of anchovy larvae: The relation between inshore chlorophyll maximum layers and successful first feeding. Fish. Bull., U.S. 73:453-462.

LAURENCE, G. C. 1977. A bioenergetic model for the analysis of feeding and survival potential of winter flounder, *Pseudopleuronectes americanus*, larvae during the period from hatching to metamorphosis. Fish. Bull., U.S. 75:529-546.

LUNZ, G. R., and F. J. SCHWARTZ. 1970. Analysis of eighteen year trawl captures of seatrout (*Cynoscion* sp.: *Sciaenidae*) from South Carolina. Contrib. Bears Bluff Lab. 53, 29 p.

MASSMANN, W. H. 1971. The significance of an estuary on the biology of aquatic organisms of the Middle Atlantic Region. In P. A. Douglas and R. H. Stroud (editors), A symposium on the biological significance of estuaries, p. 96-109. Sport Fish. Inst. Publ., Wash. D.C.

MATHEWS, T. D., and M. H. SHEALY, Jr. 1978. Hydrography of South Carolina estuaries, with emphasis on the North and South Edisto and Cooper Rivers. S.C. Mar. Resour. Cent. Tech. Rep. 30, 148 p.

MUSICK, J. A., and J. D. MCEACHRAN. 1972. Autumn and winter occurrence of decapod crustaceans in Chesapeake Bay, U.S.A. Crustaceana 22:190-200.

ODUM, W. E. 1970. Insidious alteration of the estuarine environment. Trans. Am. Fish. Soc. 99:836-847.

PRITCHARD, D. W. 1954. A study of the salt balance in a coastal plain estuary. J. Mar. Res. 13:133-144.

ROE, R. B. 1969. Distribution of royal-red shrimp, *Hymenopenaeus robustus*, on three potential commercial grounds off the southeastern United States. Fish. Ind. Res. 5:161-174.

SHEALY, M. H., Jr., and J. M. BISHOP. 1979. Hydrographic and biological studies in relation to Cooper River low-flow conditions prior to river redirection. In Cooper River Controlled Low-Flow Study, p. 139-178. S.C. Water Resour. Comm. Rep. 131.

SHEALY, M. H., Jr., J. V. MIGLARESE, and E. B. JOSEPH. 1974. Bottom fishes of South Carolina estuaries—relative abundance, seasonal distribution, and length-frequency relationships. S.C. Mar. Resour. Cent., Tech. Rep. 6, 189 p.

STEPHENSON, W., W. T. WILLIAMS, and S. D. COOK. 1972. Computer analyses of Petersen's original data on bottom communities. Ecol. Monogr. 42:387-415.

STRICKLAND, J. D. H., and T. R. PARSONS. 1968. A practical handbook of seawater analysis. Fish. Res. Board Can., Bull. 167, 311 p.

TINER, R. W., Jr. 1977. An inventory of South Carolina's coastal marshes. S.C. Mar. Resour. Cent., Tech. Rep. 23, 31 p.

TURNER, R. E. 1977. Intertidal vegetation and commercial yields of penaeid shrimp. Trans. Am. Fish. Soc. 106:411-416.

WATHNE, F. 1959. Observations on trawl-door spread and a discussion of influencing factors. Commer. Fish. Rev. 21(10):7-15.

WEINSTEIN, M. P. 1979. Shallow marsh habitats as primary nurseries for fishes and shellfish, Cape Fear River, North Carolina. Fish. Bull., U.S. 77:339-357.

WEINSTEIN, M. P., S. L. WEISS, and M. F. WALTERS. 1980. Multiple determinants of community structure in shallow marsh habitats, Cape Fear River Estuary, North Carolina, U.S.A. Mar. Biol. (Berl.) 58:227-243.

WELSH, W. W., and C. M. BREDER, Jr. 1923. Contributions to life histories of *Sciaenidae* of the eastern United States Coast. Bull. U.S. Bur. Fish. 39:141-201.

WENNER, E. L., M. H. SHEALY, Jr., and P. A. SANDIFER. 1982. A profile of the fish and decapod crustacean community in a South Carolina estuarine system prior to flow alteration. U.S. Dep. Commer., NOAA Tech. Rep. NMFS SSRE-757, 17 p.

WILLIAMS, A. B. 1955. A contribution to the life histories of commercial shrimps (Penaeidae) in North Carolina. Bull. Mar. Sci. Gulf Caribb. 5:116-146.

WILLIAMS, W. T. 1971. Principles of clustering. Annu. Rev. Ecol. Syst. 2:303-326.

WILLIAMS, W. T., and J. M. LAMBERT. 1961. Multivariate methods in plant ecology. III. Inverse association analysis. J. Ecol. 49:717-729.

ZETLER, B. D. 1953. Some effects of the diversion of the Santee River on the waters of Charleston Harbor. Trans. Am. Geophys. Union 34:729-732.

NATIONAL SEA GRANT DEPOSITORY
PELL LIBRARY BUILDING
URI, NARRAGANSETT BAY CAMPUS
NARRAGANSETT, RI 02882

Guidelines for Contributors

CONTENTS OF MANUSCRIPT

First page. Give the title (as concise as possible) of the paper and the author's name, and footnote the author's affiliation, mailing address, and ZIP code.

Contents. Contains the text headings and abbreviated figure legends and table headings. Dots should follow each entry and page numbers should be omitted.

Abstract. Not to exceed one double-spaced page. Footnotes and literature citations do not belong in the abstract.

Text. See also Form of the Manuscript below. Follow the *U.S. Government Printing Office Style Manual*, 1973 edition. Fish names, follow the American Fisheries Society Special Publication No. 12, *A List of Common and Scientific Names of Fishes from the United States and Canada*, fourth edition, 1980. Use short, brief, informative headings in place of "Materials and Methods."

Text footnotes. Type on a separate sheet from the text. For unpublished or some processed material, give author, year, title of manuscript, number of pages, and where it is filed—agency and its location.

Personal communications. Cite name in text and footnote. Cite in footnote: John J. Jones, Fishery Biologist, Scripps Institution of Oceanography, La Jolla, CA 92037, pers. commun. 21 May 1977.

Figures. Should be self-explanatory, not requiring reference to the text. All figures should be cited consecutively in the text and their placement, where first mentioned, indicated in the left-hand margin of the manuscript page. Photographs and line drawings should be of "professional" quality—clear and balanced, and can be reduced to 42 pices for page width or to 20 picas for a single-column width, but no more than 57 picas high. Photographs and line drawings should be printed on glossy paper—sharply focused, good contrast. Label each figure. DO NOT SEND original figures to the Scientific Editor; NMFS Scientific Publications Office will request these if they are needed.

Tables. Each table should start on a separate page and should be self-explanatory, not requiring reference to the text. Headings should be short but amply descriptive. Use only horizontal rules. Number table footnotes consecutively across the page from left to right in Arabic numerals; and to avoid confusion with powers, place them to the left of the numerals. If the original tables are typed in our format and are clean and legible, these tables will be reproduced as they are. In the text all tables should be cited consecutively and their placement, where first mentioned, indicated in the left-hand margin of the manuscript page.

Acknowledgments. Place at the end of text. Give credit only to those who gave exceptional contributions and not to those whose contributions are part of their normal duties.

Literature cited. In text as: Smith and Jones (1977) or (Smith and Jones 1977); if more than one author, list according to years (e.g., Smith 1936; Jones et al 1975; Doe 1977). All papers referred to in the text should be listed alphabetically by the senior author's surname under the heading "Literature Cited"; only the author's surname and initials are required in the author line. The author is responsible for the accuracy of the literature citations. Abbreviations of names of periodicals and serials should conform to *Biological Abstracts List of Serials with Title Abbreviations*. Format, see a recent Technical Report.

Abbreviations and symbols. Common ones, such as mm, m, g, ml, mg, °C (for Celsius), %, %_{oo}, etc., should be used. Abbreviate units of measures only when used with numerals; periods are rarely used in these abbreviations. But periods are used in et al., vs., e.g., i.e., Wash. (WA is used only with ZIP code), etc. Abbreviations are acceptable in tables and figures where there is lack of space.

Measurements. Should be given in metric units. Other equivalent units may be given in parentheses.

FORM OF THE MANUSCRIPT

Original of the manuscript should be typed double-spaced on white bond paper. Triple space above headings. Send good duplicated copies of manuscript rather than carbon copies. The sequence of the material should be:

FIRST PAGE
CONTENTS
ABSTRACT
TEXT
LITERATURE CITED
TEXT FOOTNOTES
APPENDIX
TABLES (provide headings, including "Table" and Arabic numeral, e.g., Table 1.—, Table 2.—, etc.)
LIST OF FIGURE LEGENDS (entire legend, including "Figure" and Arabic numeral, e.g., Figure 1.—, Figure 2.—, etc.)
FIGURES

ADDITIONAL INFORMATION

Send ribbon copy and two duplicated copies of the manuscript to:

Dr. William J. Richards, Scientific Editor
Southeast Fisheries Center Miami Laboratory
National Marine Fisheries Service, NOAA
75 Virginia Beach Drive
Miami, FL 33149

Copies. Fifty copies will be supplied to the senior author and 100 to his organization free of charge.

NOAA SCIENTIFIC AND TECHNICAL PUBLICATIONS

The National Oceanic and Atmospheric Administration was established as part of the Department of Commerce on October 3, 1970. The mission responsibilities of NOAA are to assess the socioeconomic impact of natural and technological changes in the environment and to monitor and predict the state of the solid Earth, the oceans and their living resources, the atmosphere, and the space environment of the Earth.

The major components of NOAA regularly produce various types of scientific and technical information in the following kinds of publications:

PROFESSIONAL PAPERS—Important definitive research results, major techniques, and special investigations.

CONTRACT AND GRANT REPORTS—Reports prepared by contractors or grantees under NOAA sponsorship.

ATLAS—Presentation of analyzed data generally in the form of maps showing distribution of rainfall, chemical and physical conditions of oceans and atmosphere, distribution of fishes and marine mammals, ionospheric conditions, etc.

TECHNICAL SERVICE PUBLICATIONS—Reports containing data, observations, instructions, etc. A partial listing includes data serials; prediction and outlook periodicals; technical manuals, training papers, planning reports, and information serials; and miscellaneous technical publications.

TECHNICAL REPORTS—Journal quality with extensive details, mathematical developments, or data listings.

TECHNICAL MEMORANDUMS—Reports of preliminary, partial, or negative research or technology results, interim instructions, and the like.



Information on availability of NOAA publications can be obtained from:

PUBLICATIONS SERVICES BRANCH (E/AI 13)

NATIONAL ENVIRONMENTAL SATELLITE, DATA, AND INFORMATION SERVICE
NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION
U.S. DEPARTMENT OF COMMERCE

NATIONAL SEA GRANT DEPOSITORY

PEL LIBRARY BUILDING

PROVIDENCE, RHODE ISLAND 02882

3300 Whitehaven St.
Washington, DC 20586

RECEIVED

NATIONAL SEA GRANT DEPOSITORY
DATE: MAY 10 1986

APR 30 1986