


Technical Report Series  
Number 81-2

CIRCULATING COPY  
Sea Grant Depository

ADENYLATE ANALYSIS  
IN NATURAL WATERS

Walton B. Campbell

Georgia Marine Science Center  
University System of Georgia  
Skidaway Island, Georgia



ADENYLATE ANALYSIS IN NATURAL WATERS

Technical Report 81-2

by

Walton B. Campbell

Institute of Ecology  
University of Georgia  
Athens, Georgia 30602

The Technical Report Series of the Georgia Marine Science Center is issued by the Georgia Sea Grant Program and the Marine Extension Service of the University of Georgia on Skidaway Island (P. O. Box 13687, Savannah, Georgia 31406). It was established to provide dissemination of technical information and progress reports resulting from marine studies and investigations mainly by staff and faculty of the University System of Georgia. In addition, it is intended for the presentation of techniques and methods, reduced data, and general information of interest to industry, local, regional, and state governments and the public. Information contained in these reports is in the public domain. If this prepublication copy is cited, it should be cited as an unpublished manuscript. Reproduction within the guidelines of Title 17, U. S. Code is permitted; other forms of reproduction require written permission of the author.

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Department of Energy, nor any of their employees, nor any of their contractors, sub-contractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe upon privately-owned rights.

## CONTENTS

|   |    |
|---|----|
| Abstract                                  | 1  |
| Introduction                              | 1  |
| Applications of the reaction              | 2  |
| Assumptions                               | 3  |
| Firefly enzyme                            | 4  |
| Luciferin                                 | 5  |
| Luciferase                                | 6  |
| Energy charge conversion enzymes          | 8  |
| Pyruvate kinase                           | 8  |
| Adenylate kinase                          | 9  |
| Types of adenylate extraction             | 10 |
| Extraction efficiency                     | 11 |
| Concentration of aquatic samples          | 12 |
| Procedure                                 | 15 |
| Cleaning glassware                        | 15 |
| Experimental protocol                     | 16 |
| AMP correction                            | 17 |
| Evaluation of experimental protocol       | 17 |
| Stability of adenine nucleotides          | 19 |
| Optimization of light-producing reactants | 20 |
| FLE enzyme optimization                   | 21 |
| Effect of freezing FLE extracts           | 23 |
| Handling of energy charge reagents        | 27 |
| Preparation of standards                  | 27 |
| Calculation of samples                    | 33 |
| Bibliography                              | 35 |

## FIGURES

|   |    |
|---|----|
| 1. Schematic of the procedure for adenylate analysis        | 18 |
| 2. Stability of dilute solutions of ATP at room temperature | 22 |
| 3. Stability of ATP at 4° C                                 | 22 |
| 4. Effect of amount of glycylglycine on ATP assay           | 24 |
| 5. FLE extract dilution without added luciferin             | 25 |
| 6. FLE extract dilution with added luciferin                | 25 |
| 7. Freeze-thaw experiment with the FLE extract              | 26 |
| 8. Dilution series for ATP standards                        | 30 |
| 9. Dilution series for ATP standard curve                   | 31 |

## TABLES

|   |    |
|---|----|
| 1. Replication with and without the filter included | 15 |
| 2. Procedure for adenylate assay                    | 20 |

### ABSTRACT

The firefly luciferase assay is widely used by marine ecologists to measure both ATP and total adenylates. There are substantial variations in the procedures used in these analyses, some of which may lead to substantial errors. The preparation of reagents, analytical procedures, and standardization are discussed and sources of error evaluated. An extensive bibliography on the methods and their application is included.

### INTRODUCTION

During the past decade marine ecologists have been using the firefly luciferase assay for adenosine 5' triphosphate in determinations of microbial biomass. According to Watson (1978),

The ATP technique is a difficult technique and subject to many sources of error.

Judging from the number of conflicting reports in the literature, many investigators have this feeling. The introduction of total adenylate and adenylate pool energy charge ratio methods (Atkinson 1968) have both enhanced the information content of microbial adenylate pool measurements and increased the complexity of the assays with a concomitant increase in the sources of error.

This report is not intended to replace careful examination of the literature nor laboratory experience with these methods. The theoretical metabolic implications of adenylate pool measurements have been covered comprehensively by Knowles (1977). Methodological considerations are reviewed in this report toward the end that better measurements may be made by more investigators on diverse systems and measurements can be compared between labs and ecosystems. Our primary emphasis has been on problems associated with environmental sampling. Hopefully not all of the potential difficulties covered in this report will be manifest in a single set of samples. Each type of interference is discussed with respect to its site or mode of action. A variety of time and money saving experiments have been performed and are presented as an aid in adenylate pool analyses. Our experiments have been performed on both the JRB

model 1000 and SAI model 2000 ATP photometers. Most of the methodologies discussed could be adapted easily to another instrument.



I would like to express my gratitude to the Georgia Sea Grant Program for funding the publication of this report. Special thanks are due to co-workers who shared many long hours of assays and produced most of the discussions that provoked critical experiments covered in this document: T. R. Jacobsen, A. M. Wood, J. V. Nabholtz, R. A. Rabek. This research was funded by NIH grant 5-T01-GM01968 and DOE contract EY-76-S-09-639.

#### APPLICATIONS OF THE REACTION

Firefly luminescence has been studied for over a century (Harvey 1940). The dependence on ATP of the light output from crude firefly lantern preparations was elucidated by McElroy (1947), McElroy and Strehler (1949) and McElroy et al. (1969). Strehler and Totter (1952) first suggested an assay for ATP based upon this reaction. The absolute specificity of the enzyme, luciferase, for ATP was demonstrated by Green and McElroy (1956). Although assays of this type had been used in biochemistry, Holm-Hansen and Booth (1966) first suggested a biomass assay based upon the ATP content of microbes. Chapman et al. (1971) demonstrated a correspondence between the ratio of ATP, ADP and AMP pools and microbial metabolic condition in pure culture. Weibe and Bancroft (1975) proposed use of this pool ratio for ecological investigations of microbial community metabolism.

The firefly luciferin-luciferase assay is a very sensitive method of measuring ATP, other nucleotides, or other compounds which are enzymatically converted to ATP (e.g. cyclic AMP: Johnson et al. 1970, Shultz and Daly 1973; FMN and NAD: Stanley 1971a, b). Minimal detection ranges have been reported from  $10^{-8}$   $\mu\text{g}$  ATP (Chappelle and Levin 1968) to  $10^{-10}$  mg ATP (Wise 1974) to  $10^{-6}$   $\mu\text{g}$  (Cheer et al. 1974) to  $10^5$  bacteria/ml (Conn, et al. 1975).

The current hypothesis on the mechanism of the firefly enzyme (luciferase)-substrate (luciferin)-ATP-light reaction is that two steps are involved:



Assuming all reactants except ATP are present in excess, the light output should be proportional to ATP concentration (Karl and Holm-Hansen 1976, Strehler and Totter 1952; Seliger and McElroy 1960). A quantum yield of  $0.88 \pm 0.25$  photon per ATP molecule consumed has been calculated by Seliger and McElroy (1960) and this is the basis for the assay of ATP by this reaction (Strehler and McElroy 1957). The rate limiting step has been hypothesized to be the formation of the luciferyl-adenylate complex (E-LH<sub>2</sub>-AMP) (McElroy and DeLuca 1973).

#### ASSUMPTIONS

The principal assumptions of both biomass and metabolic measurements of microbial adenylate pools are: 1) adenylates are made only by living organisms, 2) energy-containing adenylates (ATP and ADP) are rapidly degraded upon cessation of life, 3) adenylates can be quantitatively extracted and accurately assayed, 4) the amount of adenine nucleotides per unit biomass is predictable, and 5) non-cellular adenylates are not a significant source of interference. While these assumptions may be closely approached for some laboratory cultures, there are several pitfalls awaiting the unwary sampler of natural systems.

The assumption that ATP arises only from biological sources is not totally justified on the basis of work with simulations of the reducing conditions thought to exist in the primitive earth atmosphere. The nucleotide adenine has been produced in reasonable yields (100 mg/l in 4 days at 27° C) without biological intervention (Oro 1960, Oro and Kimball 1962, Oro 1963, Lowe et al. 1963). Ultraviolet irradiation of dilute adenine and ribose solutions has been shown to produce the nucleoside adenosine by Ponnampertuma et al. (1963). The nucleoside phosphates ATP, ADP and AMP have been produced



abiologically in measurable quantities from ADP, AMP and adenosine respectively in primitive earth simulations (Ponnamperuma *et al.* 1963). The ATP produced was found to be active in producing light from purified firefly luciferin-luciferase enzyme system with time course curves identical to those from ATP extracted from a biological source. There is no reason to believe that interference from non-biogenic adenylate is a significant problem. However, the hypothesis that ATP is only associated with living organisms and is quickly hydrolyzed upon cessation of life processes (Hamilton and Holm-Hansen 1967) has been called into serious question. Hodson and Azam (1977) and Azam and Hodson (1977a) found waters from the aphotic zone off Southern California to contain ten times as much dissolved ATP as was found in organisms. According to the calculations of Hulett (1970), ATP dissolved in 21° C seawater at 21°C should have a half-life of 7-8 years in the absence of organisms. Azam and Hodson (1977a) have found concentrations of dissolved ATP in coastal seawater up to 500 ng/l. Their stability estimates for ATP in sterile seawater were for a half-life in excess of 100 days.

The process of nucleotide extraction seeks to inactivate ATPases and ADPases by denaturation of enzyme proteins. All ATP-ase activity is not associated with proteins (Lowenstein 1958,1960). Nucleotide tri- and di-phosphates have been found to bind to apatite crystals (Taqui, *et al.* 1962; Cohn and Hughes 1962). This not only causes an adsorptive loss of nucleotides from solution but also promotes, in the presence of inorganic phosphate and bivalent metal ions, trans-phosphorylations of the terminal nucleotide phosphate at physiological temperature and pH in the absence of proteins (Krane and Glimcher 1962). ATP hydrolysis is catalyzed by members of the 3d group, especially copper (Schneider and Britzinger, 1964; Tetas and Lowenstein, 1963 and Buisson and Sigel, 1972), but use of 2, 2' bipyridyl is reported to complex these ions effectively out of solution by means of mixed-ligand metal ion complexes.

#### FIREFLY ENZYME


The enzyme and substrate for the ATP to light reaction may be obtained in a variety of forms. The substrate, luciferin, can be synthesized or purchased

either as a synthetic product or in a partially purified mixture with its enzyme. The enzyme, luciferase, can be purified chromatographically (Green and McElroy 1956, Nielsen and Rasmussen 1968, Bekhor *et al.* 1977), by isoelectric focusing (Lundin 1977), purchased as a purified product or as a crude mixture. The commonly used and relatively inexpensive Sigma preparations, FLE 50 and FLE 250 have been found to be undersaturated in luciferin. Thus additional linearity and sensitivity can be obtained by supplementing the crude FLE with purified or synthetic D-luciferin (Jones and Simon 1977, Karl and Holm-Hansen 1976, Bekhor *et al.* 1977, Lundin *et al.* 1976, Neufeld *et al.* 1975, Chappelle and Levin 1968, Schram *et al.* 1970). These authors arrive at the rather different minimal values of luciferin necessary for maximum sensitivity of 100 µg/ml, 10 µg/ml, 5 µg/ml, 2.5 µg/ml, 1.0 µg/ml and 0.112 µg/ml respectively. All do agree that additional luciferin increases assay sensitivity up to some concentration, beyond which it becomes inhibitory. Such an increase in sensitivity of 100 fold (Karl and Holm-Hansen 1976) allows dilution of the preparation to such an extent that contaminating endogenous ATP within the preparation becomes so dilute as to substantially lower its contribution to the background light emission.

Sensitivity can also be increased by using the much more highly purified DuPont reagent (#760-145-902), either alone or as a mixture, with less refined preparations available from Sigma, Calbiochem, Worthington, Antonik or Lumac. This blending of enzymes has the advantage of enhancing the sensitivity of the crude preparations while increasing the stability of the highly purified enzyme substrate (usually less than 4 hours at room temperature unless made up in 17% glycerol as recommended by Chapman *et al.* 1971). Addition of 0.1% bovine serum albumin has also been reported to increase both enzyme stability and sensitivity (Jones and Simon 1977). Purified luciferase is available for custom blending with luciferin or any of the mixture preparations for the desired combination of sensitivity, background stability, and economy.

#### LUCIFERIN

The substrate for the enzyme luciferase, D-luciferin, was first synthesized and characterized by Seliger *et al.* (1961) as  $C_{13}H_{10}N_2S_2O_3$ :



It may be synthesized by the methods of Bitler and McElroy 1957, Seto et al. (1963) or White et al. (1963). It has a molecular weight of 280.33, a fluorescence peak at 535 nm (Seliger et al. 1961) and is quite stable to storage if frozen in the crystalline form.

The oxyluciferin (dehydroluciferin) produced during the assay reaction or during prolonged storage in aqueous solution is a potent competitive inhibitor of luciferin (80% inhibition at 3.2  $\mu\text{M}$ ) (DeLuca et al. 1964; Airth et al. 1958). This oxyluciferin catalyzes the breakdown of ATP in the presence of pyrophosphate without the emission of light (DeLuca and McElroy 1965). Natural firefly lanterns have a very high pyrophosphatase activity which counteracts the inhibition in vivo (Hastings 1968). Completely synthetic enzyme preparations lack this activity. The metabolic cofactor, Co-A ( $5 \times 10^{-5}\text{M}$ ), has been reported to remove oxyluciferin from its binding site on the luciferase enzyme surface by forming oxyluciferyl-Co-A. In the presence of cysteine this complex reportedly formed a stable N-oxyluciferyl cysteine which did not interfere with the light producing reaction (Airth et al. 1958).

#### LUCIFERASE

The enzyme, luciferase, approximately 50,000 mw., has been purified from natural sources and characterized by Green and McElroy (1956), Nielsen and Rasmussen (1968) and Bekhor et al. (1977). The temperature optimum for this enzyme has been found to be 23° C (Seliger and McElroy 1964; Green and McElroy

1957; Chappelle and Levin 1968). The emission peak of yellow-green light at 562nm is shifted toward the red and attenuated by a pH lower than 7.6 (Seliger and McElroy 1960, McElroy and Seliger 1962, McElroy and DeLuca 1973) and is adversely affected the presence of metallic ions ( $Zn^{+2}$ ,  $Cd^{+2}$ ,  $Hg^{+2}$ ) or halogens (Seliger and McElroy 1964, Aledort et al. 1966).

The quantum efficiency of the reaction has been reported to be decreased by a wide variety of natural and artificial chemicals. The herbicides isopropyl-3-chlorocarbanilate, isopropyl-5-chloro-2-hydroxycarbanilate and isopropyl-3-chloro-4-hydroxycarbanilate have been reported as competitive inhibitors with respect to D-luciferin (Rusness and Still 1974). Both 0.1  $\mu$ M phosphoenol pyruvate and 2  $\mu$ g/ml pyruvate kinase (which are used in energy charge assays) have been reported to be inhibitory to the luciferase light yield (Robertson and Wolfe 1970). In addition, the chlorine ion has been found to be inhibitory (65%) to the production of light at 97 mM concentration, and 20% ethanol has been reported inhibitory (50%) (Holmsen *et al.* 1966). Heavy metals, such as mercury, also inhibit the light yield from the luciferase assay by direct action on the enzyme-substrate complex (Patterson *et al.* 1970). Samples collected from near shore, swamp, or forest regions subject to leaching of water-soluble phenolic compounds may inhibit significantly the luciferin-luciferase reaction. These phenolic compounds have been reported to be sorbed from solution after extraction but prior to assay, by treatment with an insoluble polyvinylpyrrolidone (Guinn and Eidenbock 1972) but there has been some question about how complete this adsorption is in some systems (Cunningham and Wetzel 1978).

The presence of inorganic salts changes the ionic strength of the reaction mixture and can be inhibitory to the quantum light yield. Sodium chloride (10mM) has been found to be highly inhibitory (50%) (Strehler and Totter 1952, Holmsen *et al.* 1966). Magnesium ion concentration has been reported to have an optimal concentration of approximately 0.01 M and was found to be inhibitory both above and below this concentration (Green and McElroy 1956, Chappelle and Levin 1968). Measured with respect to 97 mM sodium chloride the cations  $Ca^{+2}$ ,  $K^{+2}$ ,  $Rb^{+2}$ , and  $Li^{+2}$  have been shown to be inhibitory (100% to 73%) to the light yield at concentrations of 97 mM (Aledort *et al.* 1966). The salts, NaBr,  $KClO_3$ , and  $KH_2PO_4$ , have been found to be inhibitory 53% to

93%; iodide ion was (100%) inhibitory and  $\text{NaHCO}_3$ , NaF, and KF salts produced stimulations of 127%, 222% at 97 mM with respect to the reaction yield in 97 mM NaCl (Aledort et al. 1966).

#### ENERGY CHARGE CONVERSION ENZYMES

The estimation of total adenylates and energy charge involve measurement of ATP as described and also ADP and AMP. The most common method of doing this is by performing enzymatic conversion on subsamples which are then processed as ATP assays. After enzymatic (pyruvate kinase) conversion of the ADP to ATP, the ADP concentration is obtained by subtraction. A second subsample is processed in the same manner as the ADP sample except that it also contains the enzyme, adenylate kinase (myokinase) to convert AMP to ATP and ADP. AMP concentration is then determined by another subtraction. A brief description of the interconversion enzymes should facilitate understanding both the mechanism of sample processing and precautions associated with their use on diverse types of samples.

#### PYRUVATE KINASE

Pyruvate kinase (ATP:pyruvate kinase EC.2.7.1.40) reversibly catalyzes the transfer of the phosphoryl group from phosphoenolpyruvate to ADP in the presence of magnesium and potassium to yield pyruvic acid and ATP (Lohmann and Meyerhof 1934). The mammalian enzyme has been reported to exist as several distinct isozymes (Osterman and Fritz 1972, Jimenez et al. 1971). Pyruvate kinase is probably a constituent of all living cells (Boyer 1962). It is easily obtained in good yields from skeletal muscle and is stable for over one year in high purity preparations (Kayne 1973). Preparation and purification are described by Boyer (1962) and Kayne (1973). This enzyme has preferential specificity of purine di- and tri- phosphates (the activity with inosine diphosphate is 75% and with guanosine di-phosphate being 60% of that with ADP (Davidson 1959). Maximum reaction velocities at pH 7.5 have been found to be  $\text{ADP} \cong \text{GDP} > \text{IDP} > \text{ADP} > \text{UDP} > \text{CDP} > \text{dCDP}$  (CDP being only 30% of that of ADP)

(Plowmann and Krall, 1965; Weber, 1969), while the  $K_m$  for ADP was one fourth that for GDP.

Pyruvate kinase is inhibited competitively by ATP (Meyerhof and Oesper 1949; Reynard *et al.* 1961). As with other kinases, pyruvate kinase has a requirement for the divalent cation, magnesium (Lohman and Meyerhof 1934), and additionally requires potassium or a related monovalent cation for activity (Boyer *et al.* 1942) such as  $\text{NH}_4^+$  or  $\text{Rb}^{+2}$  (Kachmar and Boyer 1953). It is strongly inhibited by calcium (Kachmar and Boyer 1953), copper, zinc, mercury (Boyer 1962) and p-mercuribenzoate (Solvonuk and Collier 1955). Diethyl-stilbesterol (Kimberg and Yielding 1962) and phenylethylbiguanide (Mildvan and Cohn 1965) have also been found to be competitively inhibitory. Fluoride, ferricyanide, iodoacetate, phenothazine, alloxan, urethan arsenate and 2,4-dinitrophenol were reported as having little or no inhibitory activity (Boyer 1962).

#### ADENYLATE KINASE

Adenylate kinase (myokinase, ATP:AMP phosphotransferase, EC.2.7.4.3) reversibly catalyzes the transfer of one phosphoryl group from ATP onto AMP to produce two molecules of ATP in the presence of magnesium as described by Kalckar (1943). Adenylate kinase also catalyzes the conversion of IDP to ITP by the same mechanism (Kalckar 1943). The forward reaction of adenylate kinase to produce ATP and AMP is inhibited by AMP, but no inhibition has been noted from IMP or adenosine (Colowick and Kalckar 1943). Karl and Holm-Hansen (1978a) have reported an inverse relationship between total adenylate concentration and the apparent  $K_m$  of adenylate kinase when used in a coupled reaction with pyruvate kinase and low total adenylate concentrations (<500ng/mg). For rabbit muscle adenylate kinase, a bivalent cation is required and the order of reactivity is  $\text{Mg}^{+2} > \text{Ca}^{+2} > \text{Mn}^{+2} > \text{Ba}^{+2}$  (O'Sullivan and Noda 1968; Mildvan 1970). Adenylate kinase is ubiquitous in living cells (Noda 1973). It's preferences for nucleotide triphosphates are  $\text{ATP} > 2'\text{dATP} > \text{CTP} > \text{GTP} > \text{UTP} > \text{ITP}$  (O'Sullivan and Noda 1968). The pH optimum is 7.5 (Kalckar 1943). Adenylate kinase is found in high concentrations in rabbit and frog muscle and is only reduced to 79% of it original activity after being subjected to 100° C

for 10 minutes in 0.1 N HCl (Colowick and Kalckar 1943). Electrophoretic resolution of adenylate kinase has revealed the existence of several isozymes from each source (Giblett 1969; Khoo and Russell 1972). Its preparations and purification are described by Colowick and Kalckar (1943) and Kalckar (1943).

#### TYPES OF ADENYLATE EXTRACTION

The principal function of adenylate extraction methods is to cause leakage of cellular nucleotide pools, usually through cell lysis, accompanied by a rapid inactivation of all enzyme systems which could produce, degrade or interchange (by means of transphosphorylations) of nucleotide phosphate bonds. A second aspect of adenylate extractions is to stabilize the extracted nucleotide phosphates against losses from solution by non-enzymatic hydrolysis, adsorption or precipitation while maintaining an environment which is compatible with subsequent assay techniques. Surveys of adenylate extraction techniques for different conditions of the medium have yielded different optimum methods for each matrix (Lundin and Thorne 1975a,b; Knust et al. 1975; Cunningham and Wetzel 1978). Three approaches to extraction are commonly employed. First are the organic solvent extractions which are usually carried out at or above room temperature; these include ionizing solvents such as dimethyl sulfoxide (Chappelle and Levin 1968; Searle 1975; Levin et al. 1967); and formamide (Knust et al. 1968; Knust et al. 1975) and nonionizing organic solvents such as chloroform (Paul and Johnson 1977; Dhople et al. 1971; Dhople and Hanks 1973a,b), boiling ethanol (St. John 1970; 1971) acetone (Chappelle and Levin 1968), acetone with ether (Clark et al. 1978), methylene chloride (Knust et al. 1975), butanol-octanol (Conklin and MacGregor 1972; D'Eustachio and Johnson 1968), n-butanol (Chappelle and Levin 1968) and n-bromo-succinimide (MacLeod et al. 1969). The hydrophobic organic solvents disrupt cell walls causing leakage and inactivation of enzymes through dehydration denaturation.

A second type of extraction uses acids and is generally performed at or near freezing temperatures. The types of acids used include perchloric (Picciolo et al. 1971; Albaum and Lipshitz 1950; Albaum et al. 1951; Abood and Goldman 1956; Cheng 1961; Chapman et al. 1971; Cole et al. 1967; Bagnara and

Finch 1972; Bächli and Ettliger 1973; Cocucci et al. 1973; Ball and Atkinson 1975; Parkinson and Medley 1972); trichloroacetic (Brooks and Lutwak-Maunn 1971; Brooks 1970; Bauchop and Elsdén 1960; Bomsel and Pradet 1968; Chalauatnatol and Haesungcharern 1977); sulfuric (Ausmus 1971, 1973; Hodson et al. 1976; Forrest and Walker 1965a, b; Belaich 1966; Lee et al. 1971a,b; Kari and LaRock 1975); formic (Klofat et al. 1969, Bomsel and Pradet 1968), and nitric (Picciolo et al. 1975b). Low temperatures are generally required (Azam and Hodson (1977a) to minimize acidic hydrolysis of the nucleotides while allowing oxidative denaturation of the enzymes and other cellular components.

The third type of extraction involves boiling aqueous buffered solutions such as those using TRIS [tris-hydroxymethyl-amino-methane] (Greengard et al. 1954; Beutler and Baluda 1964; Holm-Hansen and Booth 1966; Chappelle and Levin 1968; Bamstedt and Skjoldal 1976) boiling glycine buffer (Bedell and Govindjee 1973), boiling glycyglycine (Ching and Kronstad 1972), boiling water (Burn 1962; Aledort et al. 1966; Ching and Ching 1972; Cockburn 1974), boiling sodium bicarbonate (Bancroft et al. 1976), boiling phosphate (Bulleid 1978), boiling phosphate-citrate buffer (Bulleid 1978), or just boiling buffered growth medium (Billen et al. 1953).

Not all the above extractions are optimal for all systems and some of them may be considerably less than optimal for most systems (Lundin and Thore 1975b). According to Bulleid (1978) the two major requirements of extraction are: 1) the maximum amount of ATP is extracted from the sample and 2) substances interfering with the later stages of the assay are removed, minimized, or not extracted. As noted above, perchloric acid extractions seem to have remained quite popular even after reports that ATP is coprecipitated in a non-predictable manner when the perchloric acid is neutralized with base (Wiener et al. 1974).

#### EXTRACTION EFFICIENCY

The most common method of evaluating the efficiency of adenylate extraction methods is to measure the yield from each method as a percentage of the one giving the highest total yield. This is valid only so long as the tech-



nique giving the highest yield is doing so without changing the adenylate concentration from the other cellular components during extraction. The quantity of adenylates extracted from *E. coli* was found to be very sensitive to pH in that at a pH greater than 7.0 metabolic processes which generate ATP, ADP, and UTP may proceed during the extraction processes (Kaplan et al. 1951). Thus alkaline extractions such as boiling bicarbonate (Bancroft et al. 1976), boiling phosphate-citrate (Bulleid 1978), and boiling glycylglycine (Ching and Kronst 1972) may produce artificially elevated ATP and ADP yields since these can both be produced from AMP or UDP by NADH under alkaline extraction conditions (Lundin and Thore 1975).

Soils and sediments may contain phosphate binding sites which make adenosine phosphate extractions less efficient the greater the concentration of sites (Williams et al. 1970). This adsorption can be minimized by using 0.6 N sulfuric acid extraction (Lee et al. 1971b). Unfortunately, this solubilizes many metal ions which then must be removed by cation exchange resins since metal ions are inhibitory to the luciferin-luciferase assay reaction (Seliger and McElroy 1960). Metal ions such as  $\text{Cu}^{+2}$ ,  $\text{Zn}^{+2}$ ,  $\text{Ca}^{+2}$ , and  $\text{Ce}^{+3}$  have been shown to hydrolyze ATP and ADP in aqueous solution above pH 7 (Tetas and Lowenstein 1963). According to Karl and LaRock (1975) some of the ionic interferences may be reduced by the addition of EDTA to the extractant, but Lee et al. (1971b) dispute this. Addition of potassium phosphate buffer to a boiling TRIS buffer extraction has been reported to reduce ATP binding and give an increase in ATP extraction yield from the sediments (Moriarty 1975). Fulvic acids (>1000, <10,000 mw) have been reported to form irreversible complexes with ATP, preventing its interaction with the firely enzyme luciferase (Cunningham and Wetzel 1978). These complexes were found to be dependent only upon the concentration of fulvic acids.

#### CONCENTRATION OF AQUATIC SAMPLES

Aquatic samples often require concentration before extraction. In open ocean samples Sutcliff et al. (1976) found total ATP recovery from filtered sea water to be directly proportional to the volume of water filtered, but for inshore waters they noted an apparent loss of up to 50% to some undefined

absorption phenomenon associated with the increased particle content of in-shore water. Chapman et al. (1971) found that filtration of pure cultures caused a drop in energy charge of 0.5 (the entire energy charge scale spans from 0 to 1) and loss of 50% of total adenylates, while Jones and Simon (1977) reported 44-97% loss of ATP as a result of filtration concentration. Direct extraction in acid without filtering can be done to avoid loss of ATP or change in energy charge, but it must then be concentrated before assay. You can extract directly in acid without filtering to avoid losses of ATP or lowering of the adenylate energy charge - then concentrate. The speed with which adenine nucleotide pools can change has been found to be as little as one minute for an 80% decrease in ATP (Klofat et al. 1969), or energy charge may drop 50% in less than one minute (Cocucci et al. 1973). Karl and Holm-Hansen (1978a,b) found an asymptotic relationship between the decrease in energy charge and the volume of water filter-concentrated. They also noted the higher the starting energy charge, the greater the effect of filtration. The total adenylate pool remained constant while the individual nucleotides rearranged their respective proportions. The difference between leaving a small film of water on the filter as opposed to pulling it dry has been reported to cause a loss of up to 98.6% of the ATP pool (Cheer et al. 1974). Lundin and Thore (1975b) found that centrifugation harvesting resulted in a significant decrease of the total pool of adenine nucleotides and of the energy charge.

The potentially rapid response of cellular adenine nucleotide pools to even minor stress makes the measurement of ATP alone a rather questionable procedure. The theory of adenylate-pool energy charge reflecting cellular metabolic condition dictates that the proportions of the pool constituents be free to change in response to physiological needs (Bamstedt and Skjoldal 1976, Skjoldal and Bamstedt 1976). This has been repeatedly demonstrated (Strange et al. 1963; Forrest 1965; Klofat et al. 1969; Chapman et al. 1971a,b; Chapman et al. 1971; Wiebe and Bancroft 1975; Miovic and Gibson 1973). Cole et al. (1967) measured very large fluctuations in ATP levels in E. coli subjected to stress by cooling or abrupt changes in oxygen partial pressure. Shifting the conditions of the medium in E. coli cultures were followed by such large decreases in ATP levels that Cole et al. (1967) concluded "these results throw doubt on methods for assaying ATP in which a large lag exists between harvest-

ing and extracting for assay." This is not at all uncommon in field ecological sampling regimes. If filtered samples were frozen and extracted several days later, up to 70-80% of the ATP was lost, but if frozen in liquid nitrogen after extraction, little if any ATP was degraded (Cheer *et al.* 1974). The possibility of extracting unconcentrated samples and then concentrating the extract on activated charcoal (Rasmussen and Nielsen 1968; Hodson *et al.* 1976) has been proposed as a method of collecting adenylate data from minimally perturbed aquatic samples. Karl and Holm-Hansen (1978a) have reported nonquantitative recoveries of AMP, ADP, and ATP from such columns. Azam and Hodson (1977c) report 50-60% recovery on columns. They also report 13X higher recovery of ATP from water samples using acid extraction instead of TRIS. Other means of increasing sensitivity rely on making the luciferin-luciferase system more sensitive. This is discussed in the section covering luciferase.

The above observations have lead to the conclusions that: 1) adenylate samples should be extracted as soon as possible to limit sampling effects on the nucleotide pool proportions (Holm-Hansen *et al.* 1975; Hodson and Azam 1977; Knowles 1978), and 2) total adenylate pools are more likely to give a realistic estimate of biomass than any single component of the pools, much less the most labile pool constituent, ATP.

If filter-concentrated samples are harvested by extracting the filter, it can be excluded from the extract or allowed to be frozen with the concentrate. The main reason for including the filter in with the extract is that all of the extract should be present to be later assayed. This includes both the liquid absorbed into the filter itself (as much as 0.8 ml for a 47 mm cellulose acetate filter) and the extract captured by capillary-action between the walls of the extraction tube and the filter membrane (as much as 0.3 ml for the apparatus used in this laboratory. The logic for excluding the filter is the possibility of some variable adsorption or dephosphorylations of adenylates by the filter or the materials (eg. metals) concentrated thereon.

Samples frozen with the filter show slightly higher ATP and total adenylate concentration (Table 1). The energy charges from the samples frozen with the filter removed are approximately 10% higher than those with the filter

Table 1. Replicate samples frozen with and without concentrating filter included. Coefficients of variation for replicate ATP and total adenylate extraction were less than 3% and for energy charge less than 8%.

| Depth | Frozen with Filter |          |      | Filter Removed |          |      |
|-------|--------------------|----------|------|----------------|----------|------|
|       | nM ATP             | nM Total | E.C. | nM ATP         | nM Total | E.C. |
| 8 m   | 0.48               | 1.54     | 0.55 | 0.50           | 1.41     | 0.62 |
| 18 m  | 0.87               | 1.83     | 0.66 | 0.54           | 1.34     | 0.65 |
| 21 m  | 1.09               | 2.42     | 0.65 | 1.11           | 2.27     | 0.73 |
| 26 m  | 1.58               | 4.32     | 0.64 | 1.13           | 2.59     | 0.67 |

included. The reason for the lower energy charges in the samples was a consistently higher AMP concentration. If the included filter and contents were adsorbing adenylates, there would seem to be a slight preference for ATP over AMP, or else the removed filter retain more AMP than ATP. Possibly the filter and contents were leaching an inhibitor of adenylate kinase which interfered with the conversion reaction at the time of assay. Although these data are not extensive, workers using filtration of samples should consider the possibility of this discrepancy with respect to their own filtered samples.

## PROCEDURE

### Cleaning Glassware

Great care must be taken with the preparation of standards and chemicals for use in the luciferin-luciferase assay. Metallic ion residues left behind from chromic acid cleaning cling tenaciously to glass surfaces and are slowly released into solution (Laug 1934; Henry and Smith 1946; Richards 1936; Ames and Nesbett 1960). Metallic ions interfere with the enzyme-substrate reactions, as discussed under the description of luciferase, and can also cause hydrolysis of ATP in solution, as discussed under the section on extraction methods. Cleaning of glassware and utensils with dichromate solutions should be avoided. Soaking or rinsing in 0.1 N, HCl, followed by thorough rinsing and oven drying, has produced satisfactory results.

## Experimental Protocol

The experimental protocol used for energy charge determinations is summarized in Table 2 and pictorially outlined in Figure 1. It is essentially a modification of that described by Chapman *et al.* (1971) for use with an SAI model 2000 ATP photometer, or any other photometer which will accommodate 1.5 ml sample volumes and use peak detection mode. A total of 600 enzyme units of pyruvate kinase and 1800 enzyme units of myokinase are dialyzed separately for a minimum of four hours against 800 ml of 0.1 M potassium phosphate buffer pH 7.4 at 4° C.

The ATP determinations are performed on aliquots of a mixture of 100  $\mu$ l of a mixture A which contains 16 mM  $\text{MgSO}_4$  and 84 mM pH 7.4 potassium phosphate buffer incubated 15 min. at 35°C. This is kept on ice until time of assay (less than 60 min.) whereupon two 200  $\mu$ l subsamples are placed in 1.3 ml 7.5 mM pH 7.74 glycylglycine buffer contained in 20 ml scintillation vial. This is mixed by rapid swirling, lowered into the assay chamber of the photometer and mixed with an injected volume of 200  $\mu$ l firefly lantern extract. The assay count is begun at the instant of injection of FLE by an automatic switch and only the first major peak within the first 6 sec. is measured.

The ADP determinations are identical to those of the ATP except that 100  $\mu$ l of the sample is incubated with mixture A and a mixture B containing 16 mM  $\text{MgSO}_4$ , 76 mM pH 7.4 phosphate buffer, 0.8 mM phosphoenol pyruvate and 600 enzyme units of pyruvate kinase. The ATP concentration determined above is then subtracted from the ADP + ATP concentration determined in this step to yield the ADP concentration.

The measurement of AMP is identical to that of ADP except that 1800 enzyme units of myokinase are added to the pyruvate kinase, phosphoenol pyruvate, phosphate buffer and  $\text{MgSO}_4$  reaction mixture. The ADP + ATP concentration determined above is then subtracted from this ATP + ADP + AMP value to obtain the AMP concentration.

## AMP Correction

Occasionally a negative value for AMP will be calculated when dilute adenylate samples are assayed (Bamstedt and Skjoldal 1976). This is caused by the dependence on total adenylate concentration for the coupled adenylate kinase-pyruvate kinase reaction. The apparent adenylate kinase  $K_m$  for AMP is raised by lowering the concentration of ATP (Karl and Holm-Hansen 1978). This is not a problem in samples containing several micromolar total adenylates and can be corrected in samples in the nanomolar range by performing conversion efficiency assays on each sample. This is done by adding some unknown quantity (0.4-0.04 ng/ $\mu$ l or approximately 1-10 nM) of ATP to a duplicate AMP determination, incubation sample, and standard. A second standard curve for the ATP + ADP + AMP measurements will be generated by which more accurate AMP determinations may be made. Since AMP is usually less than 10% of the total adenylate pool of living cells, this additional processing may not be desirable for large-quantity, routine assays.

For adenylate samples near the lower limits of detection (less than 500 ng/ml), the impure nature of the inexpensive firefly lantern extracts necessitates a heat inactivation step of incubated ATP, ADP and AMP subsamples. A 2 min treatment before assay at 100° C has been suggested (Karl and Holm-Hansen 1978) as sufficient to denature the added pyruvate kinase and prevent instantaneous production of ATP from ADP introduced in the crude firefly reagent. This procedure has been found to increase the linearity and lower detection limits of the ATP vs light curve.

## EVALUATION OF EXPERIMENTAL PROTOCOL

A series of experimental evaluations of several practical considerations to routine adenylate pool measurements are discussed below. They involve those aspects of adenylate measurement as they pertain to the above protocol.




Figure 1. Schematic of the procedure for adenylate analysis.

## Stability of Adenine Nucleotides

The stability of dilute solutions of ATP at room temperature was measured to ascertain the necessity of maintaining samples at low temperatures during the assay. The concentration of a  $1.0 \times 10^{-7}$  M ATP (Fig. 2) standard decreased 10% within the first 3 hrs and fell 20% within 6 hrs at 25°C. The energy charge of the standard was down 28% by 6 hr and after 8 hr at room temperature the energy charge had dropped 0.35 units of its 1.0 span. A drop of 93% in ATP concentration occurred after 24 hr at 25° C, with an accompanying decrease of 0.8 energy charge units. Thus a dilute solution of nucleotide deteriorates over intervals as short as 3 hr at room temperature. Each data point was the mean of 10 replicates. Standard deviations were smaller than the point depicting the mean.

Changes over time of dilute aqueous solutions at 4° C were determined in order to evaluate the maximum time that samples can be held after thawing. Shown in Figure 3 is the deterioration curve for a  $1.0 \times 10^{-7}$  M ATP standard held in a 4° C ice bath for up to 65 hours after being thawed. In the first 24 hr the ATP had decreased 12% and energy charge had decreased 22%. At 65 hr ATP concentration was still within 69% of its original value while energy charge had dropped to 61%. Each data point represents the mean of 10 replicate samples, and one standard deviation for the ATP measurements is shown (brackets), while those for the energy charge were smaller than the symbols depicting the means.

It should be borne in mind that the two experiments shown above were performed with ATP standards in deionized pH 7.0 water and thus would indicate only an upper limit of holding characteristics. Real samples, containing cellular debris and possible inorganic materials would potentially contain adsorption and reaction sites which could seriously shorten the permissible holding time. If samples must be held for prolonged periods after thawing, the speed and degree of their deterioration at 4° C should be determined individually. Substantial decreases in the ATP content of ATP standards held in an ice bath for as little as 6 hours have been reported (Patterson et al. 1970).



Table 2. Procedure for adenylate assay.

Reagents

1000 ml potassium phosphate buffer 0.1 M pH 7.4; 1800 enzyme units myokinase (Sigma grade III); 1200 enzyme units pyruvate kinase (Sigma type II); 1.0 ml 0.01 M phosphoenolpyruvate; 100 ml 0.1 M  $MgSO_4$ ; 2000 ml 0.075 M glycylglycine (10 g/l + 60 ml 0.1 M  $MgSO_4$ ; adjust to pH 7.74 with NaOH)

|               | $MgSO_4$ | $PO_4$ Buffer | PEP           |               |      |
|---------------|----------|---------------|---------------|---------------|------|
| 20.0 ml Mix A | 3.2      | 16.8          | -             |               |      |
| 12.5 ml Mix B | 2.0      | 9.5           | 1.0           |               |      |
|               | Mix A    | Mix B         | Pyruv. Kinase | Myokinase     | HOH  |
| 10 ml ADP mix | 3.13     | 6.25          | 0.15 (approx) | -             | 0.52 |
| 10 ml AMP mix | 3.13     | 6.25          | 0.15 (approx) | 0.52 (approx) | -    |

120 ml firefly enzyme: 2 FLE-250 rehydrated to 30 ml each + 1.2 mg synthetic luciferin + 60 ml arsenate buffer (12 ml 0.1 M  $MgSO_4$  + 0.54 g  $KH_2AsO_4$ , adjust to pH 7.74 with NaOH, bring up to 60 ml)

Incubation

0.4 ml sample extract with 0.1 ml AMP, ADP, or Mix A (for AMP, ADP, or ATP). Agitate every 5 min and maintain at 30-35 C for 15 min, then hold on ice until assay.

Assay

0.2 ml of incubated mixture + 1.3 ml glycylglycine; inject 0.2 ml firefly enzyme and begin peak count.

## Optimization of Light-Producing Reactants

Batch preparations of ATP standard solutions may be divided into several aliquots and frozen for later use. Storage times of up to 6 months (Aledent

et al. 1960), and one year (Cheer et al. 1974) have been reported with no measureable change in ATP standards and we have stored frozen standards for over three years with no loss in ATP content. We do not recommend refreezing a thawed standard because some deterioration in ATP content may occur during the interval the standard is in use (Patterson et al. 1970). One must also ensure that frozen standards are well sealed to preclude freeze-dry concentration of the solution during storage. When ATP standards are thawed for calibration purposes they must not be brought to room temperature by heating above room temperature as this will accelerate the rate of ATP deterioration in a dilute solution. Frozen standards must be completely thawed (e.g., no remaining ice) before being sampled due to the likelihood of differential freeze exclusion of ATP from the ice. Depending upon the rate the standards was initially frozen, the rate that the ATP molecules come out of the ice may be different than the rate water molecules leave the ice crystals.

The optimal quantities of reagents for the firefly ATP assay depend upon the interactions between the optical geometry of the instrument and sample, the mixing characteristics of the reactants, the effect of these on reaction kinetics, and the total pH buffering ability of the reaction system. An experiment was conducted to determine the optimal glycylglycine buffer volume for 200  $\mu$ l of sample to which 200  $\mu$ l of firefly lantern extract was injected. The results (Fig. 4) imply that the buffer volume is rather critical to obtain good reproducibility. The extremely wide error bars around all quantities of glycylglycine less than 1.3 ml make these unacceptable. The range of 1.3 to 1.4 ml had much smaller standard errors and the highest sensitivity. The range of 1.5 to 2.0 fell on a broad plateau of acceptable standard errors but was less sensitive by 5.2%. Proportions other than those tested would require reevaluation.

#### FLE Enzyme Optimization

To optimize the economics, sensitivity, and low background counts of the firefly lantern enzyme assay of ATP, the crude preparation can be diluted (Chappelle and Levin 1969; Schram 1970; Karl and Holm-Hansen 1976). Different concentrations of FLE extract without additional luciferin were compared




Figure 2. Stability of dilute solutions of ATP at room temperature.




Figure 3. Stability of ATP at 4°C.

(Figure 5). When counts from  $1.0 \times 10^{-7}$  M ATP standard were assayed in 66% Sigma FLE (1:3 in 0.05 M arsenate buffer), the background was within 33% of the counts of the standard, while full strength FLE preparation had a background within 69% of the standard.

Figure 6 demonstrates the effects of added luciferin and dilution on crude FLE extract. For 100% FLE synthetic D-luciferin the  $1 \times 10^{-7}$  M ATP sample was 68% over background, and at 66%, the same standard was eight times the background. The total counts for this 1/3 dilution were still over 300% of those from the 100% unsupplemented FLE extract (Figure 4) while the background was 1500 counts lower with the luciferin-fortified, diluted FLE extract.

#### Effect of Freezing FLE Extracts

The utility of being able to prepare, in advance, a large quantity of firefly lantern extract, using only as much as is needed for an assay series, and refreezing the remainder, lead to an investigation of the stability of FLE extract to repeated cycles of refreezing. Shown in Figure 7 is the typical response of a oneliter volume of FLE extract used for routine assays and refrozen several times. The initial sensitivity of the enzyme is almost halved after the first freeze-thaw cycle, which is accompanied by a slight drop in background. The overall trend is for a generally stable response with a slight decline in background counts for each additional thawing. At some time, in this example the sixteenth cycle, the enzyme activity drops drastically and does not recover with further cycles of freezing. The exact time of this decline is not always sixteen but is usually more than ten cycles. Attempts to reactivate the enzyme preparation by addition of more luciferin produced no measureable increase in activity. Attempts to reactivate the enzyme activity through additions of cofactor Co-A either alone or with cysteine produced no measureable change (Airth et al. 1958). Addition of a small quantity (1 ml) of this inactive enzyme to 100 ml of freshly prepared FLE resulted in total inactivation of all of the new FLE extract.




Figure 4. Effect of amount of glyglycine on ATP assay.

There is apparently a very potent inhibitor produced during freezing, thawing and storage. The FLE enzyme solution for the data presented in Figure 7 was prepared as discussed earlier. The standard errors for samples are shown as bars while the standard error intervals for the background count were smaller than the symbol depicting the point. This FLE was used in routine assays, frozen to  $-30^{\circ}\text{C}$  for from 12 hours to several days, thawed for 2-3 hours in a  $25^{\circ}\text{C}$  water bath and this process repeated until loss of activity occurred. While the FLE was thawed and being used, an aliquot of approximately 100 ml was allowed to come to room temperature for the actual assays, since the temperature optimum for the reaction is  $23^{\circ}\text{C}$  (Green and McElroy 1956; Chappelle and Levin 1968). The remainder of the thawed enzyme was held in an ice bath until more was needed during the day. Any FLE remaining at the




Figure 5. FLE extract dilution without added luciferin.




Figure 6. FLE extract dilution with added luciferin.

end of a day was returned to the main volume and refrozen. The inhibitor could have been produced during any or all of the above steps and no evaluation was made to ascertain the relative importance of each of these. Freezing FLE in small batches, to eliminate the need for repeated thawing, is thus recommended.




Figure 7. Freeze-thaw experiment with the FLE extract.

#### Handling of Energy Charge Reagents

Additional observations on the handling of energy charge reagents have been made toward the end of easier routine sampling processing. Both pyruvate kinase and myokinase should be dialyzed a minimum of four hours (for high

concentrations of adenine nucleotide the contamination introduced by omitting dialysis is apparently negligible) but can be dialyzed for 6 days and retain  $97.8\% \pm 0.06\%$  of original activity. When maintaining the dialysis solution at  $0-4^{\circ}\text{C}$  it is possible to have ice formation on the surface of the solution. This has not been found harmful to the kinases as long as the portion of dialysis tubing containing the enzymes is not allowed to become frozen. After dialysis, the kinases can lose up to  $96.1\% \pm 0.21\%$  of their activity if allowed to become frozen.

Aqueous (pH 7) solutions of phosphoenol pyruvate can be stored at  $0-4^{\circ}\text{C}$  for up to 32 days and retain  $98.2\% \pm 0.51\%$  of original activity, but if allowed to freeze were found to drop to 0.1% of previous activity levels. The enzyme mixtures for ADP and AMP conversions dropped  $73.2\% \pm 0.33\%$  and  $70.1\% \pm 0.23\%$  respectively when stored at  $0-4^{\circ}\text{C}$  for 24 hours. Taking into account the above findings, maintenance of all adenylate standards and samples at or below  $4^{\circ}\text{C}$  at all times is advisable to minimize sample deterioration and freezing of reaction enzymes should be avoided.

#### Preparation of Standards

Very highly purified adenosine phosphate standards may be prepared from commercial preparations of "pure" ATP, ADP, and AMP by chromatographic re-purification on DOWEX-1, 10% crosslinked, 200-400 mesh anion exchange resin (Cohn and Carter 1950). Since the resin is supplied in the chloride form, it must be regenerated initially to remove the luciferase-inhibitory ion (Aledort et al. 1966) by exchange with 10% NaOH through the  $1\text{ cm}^2$ , 1 cm long bed. A solution of commercial adenosine nucleotide should be made up approximately  $10^{-2}\text{ M}$  and adjusted to pH 11 with NaOH. With a flow rate of  $3\text{ ml min}^{-1}$  the pH then should be dropped in steps by use of acetic acid. Adenine should elute at approximately pH 7, 0.01 M  $\text{NH}_4$  acetate; AMP at pH 3, 0.003 M acetic acid; ADP at pH 2.5 with 0.2 M Na acetate; and ATP at pH 2.5 with 0.2 M Na acetate. The pH of the resolved peaks should then be adjusted to pH 7.0 with NaOH and the concentrations precisely determined spectrophotometrically. Experience with purified adenine nucleotides from Sigma, Calbiochem and Boehringer would indicate that this step is superfluous. This procedure is not recommended for



use except in cases where there is some doubt about the purity of the standard ATP, ADP, or AMP.

Prepare a  $10^{-2}$  M solution disodium adenosine 5'-triphosphate (m.w.= 551.2) in deionized, autoclaved water at pH  $7 \pm 0.1$ . This should be serially diluted to give a  $1 \times 10^{-4}$  M solution. One further dilution will give a  $1 \times 10^{-5}$  M solution (this is roughly the equivalent of  $10^{10}$  cells of Enterobacter aerogenes in logarithmic growth phase). The concentration of this  $1 \times 10^{-5}$  M ATP solution should next be measured spectrophotometrically. Be sure to use 1 cm quartz spectrophotometric cells. The molar extinction coefficient of adenine nucleotides at pH 7.0 for the wavelength 259 nm is  $1.54 \times 10^4$  Kalckar (1947). Using the equation  $O.D._{259}/E = C$  where  $O.D._{259}$  is the optical density of the solution at 259 nm (a dimensionless quantity), C is the molar concentration of the solution and E is the molar extinction coefficient ( $\text{moles}^{-1} \text{ cm}^{-1}$ ) the actual concentration can be determined and noted before further serial dilutions are made. If the above manipulations have been carried out precisely, the  $O.D._{259}$  of a  $1 \times 10^{-5}$  M solution of ATP at pH 7.0 will be 0.154.

An extensive standard series of ATP solutions may be prepared by either of two different dilution methods from spectrophotometrically validated standards. Solutions of ATP should be diluted (Figure 8) to obtain the series of 1 through  $9 \times 10^{-5}$  M in increments of  $1.0 \times 10^{-5}$  M. These solutions are used as the stock for further dilutions. A 1 : 10 serial dilution is performed on each of these nine stock solutions and each again is checked for accuracy by optical density measurement at 259 nm. A different set of these (Figure 9) 1:10, 2:20, 3:10 etc. dilutions is carried out on the  $1 \times 10^{-5}$  M standard created above and then measured spectrophotometrically in a 10.0 cm cell. This second set of standards is serially diluted 1:10 for each unit step in molarity to generate a continuous series from  $1.0 \times 10^{-4}$  M ATP to  $1.0 \times 10^{-8}$  M.

Either of the two different dilution series may be used to bracket unknown sample values and intersperse several values between the extreme upper and lower boundaries. The original stock solutions of the dilution series should be assayed for ADP by reaction with pyruvate kinase and for AMP by

reaction with pyruvate kinase and myokinase and phosphoenol pyruvate (see Experimental Protocol section). This is performed as a check that the spectrophotometric measurements were on ATP and not ADP or AMP which have the same extinction coefficients as ATP. ADP and AMP standard curves are prepared following the same procedure as ATP preparations. Standards from  $5 \times 10^{-7}$  to  $1 \times 10^{-8}$  M range have been used by us to bracket the concentration range found in the continental shelf waters of the Southeastern U. S. and Gulf of Mexico when filtered concentrated 100-fold.

Once the standard concentrations of ATP are prepared, an instrumentenzyme standardization procedure must be undertaken. Standards close to the concentration of the unknown solutions should be run under the same assay conditions (temperature, pH and ionic strength) as the samples to which they are to be compared. The derivation of the standard curve cannot be performed after all the unknowns have been processed, because both the firefly enzyme activity and the residual light emitted by the enzyme preparation change with time (Booth 1975). Standards must be used periodically throughout the course of sample assays. The corrections for this enzyme drift will be smaller, the shorter the time necessary to process the unknown samples. It is recommended that all samples to be run at one time be ready for immediate assay to keep the run time as short as possible. Since the enzyme will continue to drift with time, if delays of hours are encountered during an assay run, standards should be checked periodically throughout the delay period to ensure a knowledge of the rate of change when the assay is resumed.

Since the firefly lantern extract changes its light yield characteristics with age, storage conditions, and physical handling, the assay for ATP must be calibrated by means of standard ATP solutions. The standard curve is prepared by plotting log ATP concentration against log counts from the photometer over the range of intended sample assays. The curve is not strictly linear and does not necessarily regress through an origin, so the approximate range of the unknown solutions should be bracketed by standard measurements periodically to ascertain the changing slope and intercept for accurate back calculation of unknown concentrations.




Figure 8. Dilution series for ATP standards.

The most common method of standard curve preparation is by means of log-log transformation. It should be emphasized that the logarithm of the ATP concentration is not really a linear function of the logarithm of the measured light output. Even cursory examination of the references cited below will reveal that this non-linearity is readily apparent for less than single-decade concentration ranges. The mechanical disturbance of injection will evoke light production for firefly lantern extracts in distilled, deionized, autoclaved water. Although the assumption that zero ATP concentration should give zero light output (as measured above background light from the enzyme preparation itself) is true, the functional converse of this statement is not necessarily true. That is, photometric devices have finite lower limits of sensi-

tivity for a particular wavelength. Zero-measured light output can occur at non-zero concentrations of ATP. Thus computations cannot assume that the line through the standard points also passes through the origin. Strictly speaking, a log-log plot has an undefined origin. As discussed in the section




Figure 9. Dilution series for ATP standard curve.

on enzyme preparation and purification, a change in the proportions of reagents or a different photodetector system may detect a smaller non-zero concentration.

Many authors utilize a linear fit either by a least-squares regression or best fit by eye. This is very simple and is the most rapid method of deter-

mining the approximate relative magnitudes of differences between samples. The most convincing standard curves are prepared sufficiently small that nonlinearities are not apparent (Karl and Holm-Hansen 1976; Lundin and Thore 1975; Law and Button 1977). Less compelling but realistic straight lines can be drawn through standard error bars or between several data points to produce the best-fit line (Chappelle and Levin 1968; Patterson *et al.* 1970; Neufield *et al.* 1975; McLeod *et al.* 1969; Parkinson and Medley 1972; Lazdunski and Belaich 1972; Addanki *et al.* 1966; Schram 1970). The most accurate graphic portrayals of this non-linear correspondence is the method of linear interpolation between actual standard points (Strehler and Totter 1952; Strehler 1968); Picciolo *et al.* 1975, Cheer *et al.* 1974).

Attempts have been made to eliminate drawing the actual standard curve and rely on regression analysis commonly available on automatic computing devices. This may produce the appearance of increased precision without actually doing so. The use of a simple proportional relationship between the measured light output of an unknown and a standard (St. John 1975) assumes a linear relationship between the light and ATP without the benefit of a log-log transformation or any knowledge of the shape of the curve between determined points. This could be useful only over a very small concentration ranges for qualitative comparative purposes to determine which samples had more ATP without regard to actual quantity. Other workers have applied polynomial regressions for their standard curves (Booth 1975). Fitting a curve to the data will produce a smooth line among the data points, but would not be as accurate as individual short straight lines drawn between accurately prepared standard points.

To summarize the above points; too few data points in the standard curve can produce an apparent linearity which may appreciably affect the absolute accuracy of the results. This accuracy is a function of the span of the data points. A log-log transformation of the data will produce the smoothest, most continuous distribution of the standard points. The standard curve cannot be assumed to pass through the origin. A linear interpolation between actual data points will produce more accurate data than will an attempt to smooth a curve to compensate for presumed poor quality control of the standard solutions.

Each time an ATP standard curve is rerun during the course of a total adenylate or energy charge determination, and AMP standard curve of the same concentrations should be rerun so that any changes in the conversion efficiency for AMP, either with time or with concentration will be monitored and taken into account during calculations. The AMP+ADP+ATP samples can then be calculated using this curve, if it has a different slope at different concentrations.

#### Calculation of Samples

The simplest method of data reduction uses a log transformation of both machine counts and ATP concentrations for the standard curve. Several, usually three or more, short two-point straight-line sections should be constructed between standard concentrations and mean values of each set of machine counts for each of four or more half powers of ten increments bracketing the expected range of the sample concentrations. Slope and intercepts are calculated as follows:

$$\text{Slope}_{1,2} = \frac{\log \text{ATP}_1 - \log \text{ATP}_2}{\log(\text{counts}_1 - \text{background}) - (\log(\text{counts}_2 - \text{background}))}$$

$$\text{Intercept}_{1,2} = \log \text{ATP}_1 - [\text{Slope}_1 \times \log(\text{counts}_1 - \text{background})]$$

For each set of counts between any standard values (1) and (2) the sample ATP concentration is then calculated:

$$\text{ATP} = \text{ANTILOG}(\text{Slope}_{1,2}) \times (\text{counts} - \text{background}) + \text{Intercept}_{1,2}$$

where counts is the machine counts produced from an unknown sample. If the sample was concentrated and/or diluted, ATP must then be multiplied by the inverse of these concentration factors to yield the concentration in desired units. The units of the standards (nM, mg/l etc.) will be the units of the results unless subsequently transformed. Representative values of slope ( $1.01 \pm 0.2$ ) and intercept ( $-10.89 \pm 1.22$ ) were determined from 87 separate standard curves. Standard curves with background determinations should be performed

often (every 60-100 assays) to ensure accuracy and comparability of data. This is necessary, because all firefly enzyme mixtures have different ATP vs light curves and this will change with time (2-4 hours) enough to be measurable and thus affect the absolute accuracy of the determination.

## BIBLIOGRAPHY

- Abood, L. G. and E. Goldman. 1956. Inhibition of Phosphorylation During Electrical Excitation of Frog Nerves. *Amer. J. Physiol.* 184: 329-332.
- Addanki, S., J. F. Sotos and P. D. Rearick. 1966. Rapid determination of picomole quantities of ATP with a liquid scintillation counter. *Analyt. Biochem.* 14: 261-264.
- Airth, R. L., G. E. Foerster and R. Hinde. 1970. Bioluminescence. *In: Photobiology of Microorganisms.* Edited by P. Halldal. John Wiley & Sons, Inc., N.Y. pp. 417-446.
- Airth, R. L., W. C. Rhodes and W. D. McElroy. 1958. The function of coenzyme A in luminescence. *Biochim. Biophys. Acta.* 27:519-532.
- Aksenov, S. I. and L. N. Sosunova. 1975. Methods of Isolation of ATP from Microorganisms. *Mikrobiologiya* 44: 1002-1006.
- Albaum, H. G. and R. L. Lipshitz. 1950. Determination of adenosine triphosphate based on deamination rates. *Arch. Biochem.* 27:102-108.
- Albaum, H., T. Cayle and A. Shapiro. 1951. Blood levels of adenine nucleotides. *J. Clin. Invest.* 30: 525-530.
- Aledort, L. M., R. I. Weed and S. B. Troup. 1966. Ionic effects of firefly bioluminescence assay of red blood cell ATP. *Anal. Biochem.* 17: 268-277.
- Allen, P. D. 1973. Development of the Luminescence Biometer for Microbial Detection. *Dev. Ind. Microbial.* 14: 67-73.
- Ames, A. and F. B. Nesbet. 1960. Analysis for potassium, sodium, chloride, and water in a 2  $\mu$ l sample of extracellular fluid. *Anal. Biochem.* 1:1-7.
- Andersen, K. B. and K. von Meyenburg. 1977. Changes of nicotinamide adenine nucleotides and adenylate energy charge as regulatory parameters of the metabolism in *Escherichia coli*. *J. Biol. Chem.* 252: 4151-4156.
- Anderson, J. R. and P. I. Davies. 1973. Investigations on the extraction of adenosine triphosphate from soils. *Bull. Ecol. Res. Comm. (Stockholm)* 17: 271-273.
- Anon. 1970. Bacterial cell counts. *Process Biochem.* 5: 35.
- Anon. 1970. DuPont 760 Luminescence Biometer Instruction Manual. E. I. DuPont de Nemours & Co. (Inc.), Wilmington, Delaware 19898.
- Anon. Simple Chamber facilitates chemiluminescent detection of bacteria. Technical support package for Tech. Brief 70-10525. Technology Utilization Office. NASA Langley Research Center, Hampton, VA. 23365.



- Anon. 1974. Lab-Line. ATP Photometer Instruction Manual No. 9140. Lab-Line Instruments Inc., Melrose Park, Illinois 60160 (Also JRB ATP Photometer Brochure for Model 2000. SAI Technology--Previously JRB Incorporated, San Diego, California 92123).
- Anon. Biometer Flashes. E.I. DuPont de Nemours & Co., Inc., Wilmington, Delaware 19898. (i) Number 760BF1. 1970. Water ecology/waste water treatments. (ii) Number 760BF2. 1971. Monitoring cooling tower and water and biocide control. (iii) Number 760BF3. 1972a. Bacteria in cutting oils and hydraulic fluids. (iv) Number 760BF4. 1972b. Rapid determination of BOD problem. (v) Number 760BF5. 1972c. Bacterial screening of urine.
- Anon. 1972. Biomass Determination--A New Technique for Activated Sludge Control, Water Pollution Control Research Series 17050EoY 01/72. Environmental Protection Agency PB 211-127. Supt. of Documents, U. S. Government Printing Office, Washington, D. C. 20402.
- Anon. Biometer Flash Number BF6 (P. D. Allen). Fermentation monitoring. E. I. DuPont Nemours & Co., Inc. Wilmington, Delaware 19898.
- Anon. 1977. Proposed Test Method for Adenosine Triphosphate (ATP) Content of Microorganisms in Water, ASTM Annual Book of Standards, Part 31, Water p. 1043-1046, ASTM, Philadelphia, Pa.
- Ansell, A. D. 1977. The adenosine triphosphate content of some marine bivalve molluscs. *J. Exp. Mar. Biol. Ecol.* 28: 269-283.
- Atkinson, D. E. 1965. Biological feedback control at the molecular level. *Science* 150: 851-857.
- Atkinson, D. E. 1968a. Citrate and the citrate cycle in the regulation of energy metabolism. *In*: *The Metabolic Roles of Citrate*. Edited by T. W. Goodwin. Academic Press, N.Y., pp. 23-40.
- Atkinson, D. E. 1968b. The energy charge of the adenylate pool as a regulatory parameter. Interaction with feedback modifiers. *Biochemistry* 7: 4030-4034.
- Atkinson, D. E. 1968c. Regulation of energy metabolism. *Symp. Fundam. Cancer Res.* 22: 397-413.
- Atkinson, D. E. 1969. Regulation of enzyme function. *Ann. Rev. Microbiol.* 23: 47-68.
- Atkinson, D. E. 1971. Adenine nucleotides as stoichiometric coupling agents in metabolism and as a regulatory modifier. The adenylate energy charge. *In*: *Metabolic Pathways*. Vol. 5. D. M. Greenberg, Ed. Academic Press, N.Y., pp. 1-23.
- Atkinson, D. E. 1971. Adenine nucleotide as universal stoichiometric coupling agents. *In*: *Advances in Enzyme Regulation*. Vol. 9. G. Weber, Ed. Pergamon Press, N.Y., pp. 207-219.

- Atkinson, D. E. and N. J. Ball, Jr. 1975. Adenylate energy charge in Saccharomyces cerevisiae during starvation. *J. Bacteriol.* 121: 975-982.
- Atkinson, D. E. and L. Fall. 1967. Adenosine triphosphate in biosynthetic regulation. Escherichia coli phosphoribosylpyrophosphate synthase. *J. Biol. Chem.* 242: 3239-3241.
- Ausmus, B. S. 1971. Measurement of microbial activity using the luciferin-luciferase assay for adenosine triphosphate. U. S. IBP Deciduous Forest Biome. Memo Report No. 71-72. Oak Ridge Natl. Lab.
- Ausmus, B. S. 1973. The use of the ATP assay in terrestrial decomposition studies. *Bull. Ecol. Res. Comm. (Stockholm)* 17: 223-234.
- Axelrod, D. M., K. A. Moore and M. E. Bender. 1976. Nitrogen, phosphorus and carbon flux in Chesapeake Bay marshes. Bull 79, Virginia Water Resources Center, Blacksburg, Va.
- Azam, F. and R. E. Hodson. 1977a. Dissolved ATP in the sea and its utilisation by marine bacteria. *Nature* 267: 696-698.
- Azam, F. and R. E. Hodson. 1977b. Size distribution and activity of marine microheterotrophs. *Limnol. Oceanogr.* 22: 492-501.
- Azam, F. and R. E. Hodson. 1977c. Use of ATP as an indicator of bacterial biomass in sea water. In: G. A. Borun (Ed.), Second Bi-annual ATP Methodology Symposium, SAI Technology, San Diego.
- Bächi, B. and L. Ettliger. 1973. Influence of glucose on adenine nucleotide levels and energy charge in Acetobacter aceti. *Arch. Mikrobiol.* 93: 155-164.
- Bagnara, A. S. and L. R. Finch. 1972. Quantitative extraction and estimation of intracellular nucleoside triphosphates of Escherichia coli. *Anal. Biochem.* 45: 24-34.
- Balch, N. 1972. ATP content of Calanus finmarchicus. *Limnol. Oceanogr.* 17: 906-909.
- Balharry, G. J. E. and D. J. D. Nicholas. 1970. ATP-sulphurylase in spinach leaves. *Biochim. Biophys. Acta* 220: 513-524.
- Balharry, G. J. E. and D. J. D. Nicholas. 1971. New assay for ATP-sulphurylase using the luciferin-luciferase method. *Anal. Biochem.* 40: 1-17.
- Ball, W. J. and D. E. Atkinson. 1975. Adenylate energy charge in Saccharomyces cerevisiae during starvation. *J. Bacteriol.* 121: 975-982.
- Bamstedt, U. and H. R. Skjoldal. 1976. Studies on the deep-water pelagic community of Kors Fjorden, Western Norway. *Sarsia* 60: 63-80.

- Bancroft, K. 1974. Extraction and measurement of adenosine triphosphate from marine sediments with boiling bicarbonate solutions. M.S. Thesis, University of Georgia.
- Bancroft, K., E. A. Paul and W. J. Wiebe. 1976. The extraction and measurement of ATP from marine sediments. *Limnol. Oceanogr.* 21: 473-480.
- Banse, K. 1980. Microzooplankton interference with ATP estimates of plankton biomass. *J. Plankton. Res.* 2: 235-238.
- Barrenscheen, H. K. and W. Filz. 1932. Untersuchungen zur Frage der Co-Fermentwirkung. *Biochem. Z.* 250: 281-304.
- Barrett, J. 1973. Nucleoside triphosphate metabolism in the muscle tissue of Ascaris lumbricoides (Nematoda). *Int. J. Parasitol.* 3: 393-400.
- Bass, R. 1914. Über die Purinkörper des menschlichen Blutes und den wirkungsmodus der 2-Phenyl-4Chinolin-carbosaure (Atophan). *Arch. Exp. Path. u. Pharm.* 76: 40-64.
- Bauchop, T. and S. R. Elsen. 1960. The growth of microorganisms in relation to their energy supply. *J. Gen. Microbiol.* 23: 457-469.
- Bedell, G. W. II and Govindjee. 1973. Photophosphorylation in intact algae: Effects of inhibitors, intensity of lights, electron acceptor and donors. *Plant cell physiol.* 14: 1081-1097.
- Bekhor, I., J. K. Pollard, W. Drell. 1977. Purification and properties of luciferase from firefly lanterns, G. A. Borun, Ed. 2nd Bi-Annual ATP methodology symposium, San Diego, SAI Technology.
- Belaich, J. P. 1966. Sur une mutation thermosensible d'Escherichia Coli affectant une energetique. *C. R. Acad. Sci. Paris, T 263, serie D*, 886-888.
- Berland, B. R., D. J. Bonin, R. A. Daumas, P. L. Laborde et S.Y. Maestrini. 1970. Variations du comportement physiologique de l'algue Monallantus salina (xanthophyceae) en culture. *Mar. Biol.* 7: 82-92.
- Berland, B. R., D. J. Bonin, P. L. Laborde et S.Y. Maestrini. 1972. Variations de quelques facteurs estimatifs de la biomass, et en particulier de l'ATP, chez plusieurs algues marines planctoniques. *Mar. Biol.* 13: 338-345.
- Bernardi, A. 1974. A fast method of analysis of the 5 terminal nucleotides of deoxyribonucleotides. *Anal. Biochem.* 59: 501-507.
- Beutler, E. and M. C. Baluda. 1964. Simplified determination of blood adenosine triphosphate using the firefly system. *Blood* 23: 688-696.
- Beutler, E. and O. Duron. 1965a. Red cell ATP determination. *Blood* 25: 625.

- Beutler, E. and O. Duron. 1965b. Effect of pH on preservation of red cell ATP. *Transfusion* 5: 1724.
- Beutler, E. and K. Mathai. 1967. A comparison of normal red cell ATP levels as measured by the firefly system and the hexokinase system. *Blood* 20: 311-320.
- Billen, D., B. L. Strehler, G. E. Stapleton and E. Brigham. 1953. Post-irradiation release of ATP from Escherichia coli. *Arch. Biochem. Biophys.* 43: 1-10.
- Bitler, B. and W. D. McElroy. 1957. The preparation and properties of crystalline firefly luciferin. *Arch. Biochem. Biophys.* 72: 358-368.
- Bock, R. M., N. Ling, S. A. Morell and S. H. Lipton. 1956. Ultraviolet absorption spectra of ATP and related 5'-ribonucleosides. *Arch. Biochem. Biophys.* 62: 253-264.
- Bomsel, J. L. and A. Pradet. 1967. Étude des adenosine 5' mono-, di- et tri-phosphates dans les tissus vegetaux. II. Evolution in vivo de l'ATP, l'ADP et l'AMP dans les feuilles de biete en fonction de differentes conditions du milieu (AMP, ADP, and ATP in plant tissues. *Physiol. Veg.* 5: 223-236.
- Bomsel, J. L. and A. Pradet. 1968. Study of adenosine 5' mono-, di-, and triphosphate in plant tissues. IV. Regulation of the level of nucleotides, in vivo, by adenylate kinase: theoretical and experimental study. *Biochim. Biophys. Acta* 162: 230-242.
- Booth, C. R. 1975. Instrument development and data processing for the ATP photometer, Borun, G.A. [ed], ATP methodology Seminar, SAI, Technology, San Diego.
- Borun, G. A. [ed.] 1975. ATP methodology Seminar, San Diego, SAI Technology Co., San Diego.
- Borun, G. A. 1977. 2nd Bi-annual ATP methodology symposium, SAI Technology Co., San Diego.
- Bott, T. L. 1973. Bacteria and the assessment of water quality. *Biological Methods for the Assessment of Water Quality*, ASTM STP. 528, American Society for Testing and Materials, pp. 6175.
- Bowie, L. J., R. Irwin, M. Loken, M. DeLuca, L. Brand. 1973. Excited-state proton transfer and the mechanism of action of firefly luciferase. *Biochem.* 12: 1852-1857.
- Boyer, P. D. 1962. Pyruvate kinase. p. 95-113. In: P. D. Boyer, H. Lardy and K. Myrback (eds.), *The Enzymes* 2nd Edition, Vol. 6. Academic Press, N.Y.
- Boyer, P. D., H. A. Lardy and P. H. Phillips. 1942. The role of potassium in muscle phosphorylations. *J. Biol. Chem.* 146: 673-682.

- Brewer, G. J. and J. W. Eaton. 1971. Erythrocyte metabolism: Interaction with oxygen transport. *Science* 171: 1205-1211.
- Brewer, G. J. and C. A. Knutsen. 1966. A technique for the processing of blood samples for subsequent assay of ATP, and an investigation of the method of standardization of the firefly luciferase ATP assay. *Clin. Chim. Acta* 14: 836-839.
- Brewer, G. J. and R. D. Powell. 1964. The adenosine triphosphate content of G-6-PD deficient and normal erythrocytes including studies of a G-6-PD deficient man with elevated erythrocytic ATP. *J. Lab. Clin. Med.* 67: 726.
- Brewer, G. J. and R. D. Powell. 1965. A study of the relationship between the content of adenosine triphosphate in human red cells and the course of Falciparum Malaria. A new system may confer protection against malaria. *Proc. Nat. Acad. Sci. (U.S.)* 54: 741-745.
- Brezonik, P. L., F. X. Browne and J. L. Fox. 1975. Application of ATP to plankton biomass and bioassay studies. *Water Res.* 9: 155-162.
- Brezonik, P. L., and J. W. Patterson. 1971. Activated sludge ATP: Effects of environmental stress. *J. Sanit. Eng. Div., Am. Soc. Civ. Eng.* 97: 813-824.
- Brintzinger, H. 1963. The structures of adenosine triphosphatmetal (SIC) ion complexes in aqueous solution. *Biochim. Biophys. Acta* 77: 343-345.
- Browkaw, C. J. 1967. Adenosine triphosphate usage by flagella. *Science* 156: 76-78.
- Brooks, D. E. 1970. Content of ATP and ADP in bull spermatozoa using the firefly luciferase system. *J. Reprod. Fertil.* 23: 525-528.
- Brooks, D. E. and C. Lutwak-Maunn. 1971. Content of ATP and ADP in rabbit blastocysts. *Nature* 229: 202-203.
- Brown, E. G. and K. C. Short. 1969. The changing nucleotide pattern of sycamore cells during culture in suspension. *Phytochem.* 8:1365-1372.
- Brown, P. R. 1970. The rapid separation of nucleotides in cell extracts using high-pressure liquid chromatography. *J. Chromatog.* 52: 257-272.
- Browne, F. X. 1971. ATP measurements in laboratory and field populations of lake plankton. Ph.D. Thesis, University of Florida, Gainesville, Fla.
- Brundage, R. S., R. L. Karpel, K. Kustin and J. Weisel. 1972. Relaxation of spectra of copper II - ATP complexes at low pH. *Biochim. Biophys. Acta* 267: 258-267.
- Brunker, R. 1975. Autoinjector for the determination of picomolar quantities of ATP with a liquid scintillation counter. *Anal. Biochem.* 63: 418-422.

- Buck, J. B. 1948. The anatomy and physiology of the light organ in fireflies. *Ann. N.Y. Acad. Sci.* 49: 397.
- Buisson, D. H. and H. Sigel. 1972. Protection of adenosine 5'-triphosphate toward hydrolysis by the formation of a mixed ligand metal ion complex. *Angew. Chem. Internat. Edit.* 11: 1025.
- Bulleid, N. C. 1978. An improved method for the extraction of adenosine triphosphate from marine sediment and seawater. *Limnol. Oceanogr.* 23: 174-178.
- Burn, G. P. 1962. Adenosine triphosphate content and glucose uptake of human erythrocytes and the influence of insulin. *Biochim. Biophys. Acta* 59: 347-354.
- Burnison, B. K. 1974. Microbial ATP Studies. *Proc. 19th Congress International Association of Theoretical and Applied Limnology.* p. 23.
- Bush, V. M. 1973. The examination of urine: samples for pathogenic microbes by the luciferase assay for ATP. I. The effect of the presence of fungi fungal-like bacteria and kidney cells in urine samples. NASA Grant Report No. N73 18096. 1973.
- Bush, V. N., G. L. Picciolo and E. W. Chappelle. 1975. The effect of growth phase and medium on the use of the firefly adenosine triphosphate (ATP) assay for the quantitation of bacteria. NASA SP-388, p. 35-41.
- Butler, E. and C. K. Mattai. 1967. A Comparison of normal red cell ATP levels as measured by the firefly system and the hexokinase system. *Blood* 30: 311-319.
- Campbell, W. B., R. W. Harris, R. E. Benoit. 1972. Response of Alaskan tundra microflora to crude oil spill. *Proceedings of the symposium on the impact of oil resource development of northern plant communities.* Univ. of Alaska, Fairbanks. 96pp.
- Campbell, W. B., T. R. Jacobsen and A. M. Wood. 1979. Localization of marine microbial populations within vertical density gradients. *Bacteriol. Proc.*, p. 237.
- Campbell, W. B. and W. J. Wiebe. 1978. ATP standard curves: a precautionary note. *Bacteriol Proc.* 78: 205.
- Campbell, W. B., T. R. Jacobsen and L. R. Pomeroy. 1979. Heterotrophic-Photoautotrophic index: a qualitative parameter of microbial interactions applied to a Gulf Stream intrusion. *Mar. Sci. Commun.* 5:383-398.
- Carlson, A. D. and J. Copeland. 1978. Behavioral Plasticity in the flash communication systems of fireflies. *Amer. Scientist.* 66: 340-346.
- Carlucci, A. F., O. Holm-Hansen and P. M. Williams. 1971. Distribution biochemical activity, and chemical composition of plankton in Antarctic waters. *Antarct., J.* 6: 155-156.

- Cavari, B. 1976. ATP in Lake Kinneret: Indicator of microbial biomass or phosphorus deficiency? *Limnol. Oceanogr.* 21: 231-236.
- Chalauatnatol, M. and A. Haesungcharern. 1977. Stabilization of adenylate energy charge and its relation to human sperm motility. *J. Biol. Chem.* 252:8088-8091.
- Chambers, R. W., P. Shapiro, V. Kurkov. 1960. Synthesis of cytidine 5'-diphosphate and guanosine 5'-diphosphate. *J. Amer. Chem. Soc.* 82: 970-975.
- Chan, P. S., C. T. Black and B. J. Williams. 1973. Separation of cyclic 3', 5'-AMP from ATP, ADP, and 5'-AMP by precipitation with inorganic compounds. *Anal. Biochem.* 55: 16-25.
- Chapman, C. and W. Bartley. 1969. Adenosine phosphate and the control of glycolysis and gluconeogenesis in yeasts. *Biochem. J.* 111: 609-613.
- Chapman, A. G., L. Fall, D. E. Atkinson. 1971a. Adenylate energy charge in *Escherichia coli* during growth starvation. *J. Bacteriol.* 108: 1072-1086.
- Chapman, J. D., R. G. Webb and J. Borsa. 1971. ATP pool levels in synchronously growing Chinese hamster cells. *J. Cell. Biol.* 49: 229-233.
- Chappelle, J. B. 1977. ATP. Carolina Biological Supply, Burlington, North Carolina. 32 pp.
- Chappelle, E. W. 1977. Use of hexokinase for the reduction of inherent light levels in luciferase preparations. U. S. Patent 3437550, p. 3.
- Chappelle, E. W. 1971. Virus detection. U. S. Patent 3575812, p. 5.
- Chappelle, E. W. and D. G. Hoffman. 1970. Apparatus for detecting and measuring light emitted as a result of the ATP-luciferin-luciferase biochemical reaction. U. S. Patent 3520660, p. 6.
- Chappelle, E. W. and G. V. Levin. 1968. The use of the firefly bio-luminescent reaction for rapid detection and counting of bacteria. *Biochem. Med.* 2: 41-52.
- Chappelle, E. W. and G. V. Levin. 1971. Cancer detection. U. S. Patent 3575811, p. 4.
- Chappelle, E. W. and G. L. Picciolo. 1970. Bacterial adenosine triphosphate as a measure of urinary tract infection. (NASA, U. S. Patent -Appl. SN-60950). p. 31.
- Chappelle, E. W. and G. L. Picciolo. 1971a. Assay of flavine mononucleotide (FMN) and flavine adenine dinucleotide (FAD) using the bacterial luciferase reaction. In: *Methods of Enzymology*. S. Colowick and N. O. Colowick, Eds. 18 Part B, Academic Press, N.Y., p. 381-385.

- Chappelle, E. W. and G. L. Picciolo. 1971b. Bacterial phosphate as a measure of urinary tract infection. NASA Tech. Brief, 71: 10051, GSFC, Greenbelt, Maryland.
- Chappelle, E. W. and G. L. Picciolo, (Eds) 1975. Analytical applications of bioluminescence and Chemiluminescence, NASA #SP-388 Washington, D. C. 115 pp.
- Chappelle, E. W., and G. L. Picciolo and R. H. Altland. 1967. A. Sensitive assay for flavin mononucleotide using the bacterial bioluminescence reaction. Biochem. Med. 1: 252-260.
- Chappelle, E. W., G. L. Picciolo, C. A. Curtis, G. A. Knust, D. A. Nibley, R. B. Vance. 1975. Laboratory procedures manual for the firefly luciferase assay for adenosine triphosphate (ATP). NASA-GSFC TM X-70926. Greenbelt, MD.
- Chase, A. M. 1960. The measurement of luciferin and luciferase. In: Methods of Biochemical Analysis. Vol. 8. Edited by D. Glick. Interscience Publishers. John Wiley & Sons, Inc. N.Y., 61-118.
- Cheer, S., J. H. Gentile and C. S. Hegre. 1974. Improved methods for ATP analysis. Anal. Biochem. 60: 102-114.
- Cheng, S. 1961. A sensitive assay method for ADP and the determination of ATP, ADP, CrP in single nerve trunks. J. Neurochem. 7: 271-277.
- Chiaudani, J. B. and R. Pognotta. 1978. Ratio ATP/chlorophyll as an indicator of river's water quality. Verh. Internat. Verein. Limnol. 20:1897-1901.
- Ching, T. M. 1973. Adenosine triphosphate content and seed vigor. Plant Physiol. 51: 400-402.
- Ching, T. M. 1975. Bioluminescence for determining energy state of plants. In: Chappelle, E. W. and G. L. Picciolo (eds). Analytical applications of bioluminescence and chemiluminescence, NASA SP-338, Washington, D. C. p. 49-56.
- Ching, T. M. and K. K. Ching. 1972. Content of adenosine phosphate and adenylate energy charge in germinating ponderosa pine seeds. Plant Physiol. 50: 536-540.
- Ching, T. M. and R. Danielson. 1972. Seedling vigor and adenosine triphosphate level of lettuce seeds. Proc. Assoc. Off. Seed. Anal. 62: 116-124.
- Ching, T. M. and W. E. Kronstad. 1972. Varietal differences in growth potential, adenylate energy level and energy charge of wheat. Crop Sci. 12: 785-789.
- Chiu, S. Y. 1972. Modeling of kinetics and significance of ATP measurements in an activated sludge system. Ph.D. Thesis, Kansas State University, p. 158.



- Chiu, S. Y., I. C. Kao, L. E. Erickson and L. T. Fan. 1973. ATP pools in activated sludge. *J. Water Pollut. Control Fed.* 45: 1746-1758.
- Christian, R. R., K. Bancroft and W. J. Wiebe. 1974. Distribution of microbial adenosine triphosphate in salt marsh sediments. *Bacteriol. Proc.* 74: 54.
- Christian, R. R., K. Bancroft and W. J. Wiebe. 1975. Distribution of microbial adenosine triphosphate in salt marsh sediments at Sapelo Island, Georgia. *Soil Sci.* 119: 89-97.
- Chrzanowski, T. H., L. H. Stevenson and B. Kjerfve. 1979. Adenosine 5'-triphosphate flux through the North Inlet marsh system. *Appl. Environ. Microbiol.* 37: 841-848.
- Chulavantnatol, M. and A. Haesangcharern. 1977. Stabilization of adenylate energy charge and its relation to human sperm motility. *J. Biol. Chem.* 252: 8088-8091.
- Clark, J. R., D. I. Messenger, K. L. Dickson, J. Cairns. 1978. Extraction of ATP from Aufwuchs communities. *Limnol. Oceanogr.* 23: 1055-1059.
- Classman, T. A., C. Cooper, L. W. Harrison and T. J. Swift. 1971. A proton magnetic resonance study of metal ion-adenic ring interactions in metal ion-complexes with ATP. *Bot. Gaz.* 135: 368-372.
- Clegg, T. J. and J. L. Koenevig. 1974. The effect of four chlorinated hydrocarbon pesticide and one organophosphate pesticide on ATP levels in three species of photosynthesizing fresh water algae. *Bot. Gaz.* 135: 368-372.
- Cockburn, W. 1974. Simultaneous measurement of oxygen evolution and endogenous adenosine triphosphate levels in isolated "intact" chloroplast suspension. *J. Exp. Bot.* 25: 111-120.
- Cocucci, M., M. C. Cocucci and E. Marre. 1973. Growth inhibition without ATP decrease in *Rhodotorula gracilis* cultured under reduced oxygen pressure. *Plant Sci. Lett.* 1: 425-431.
- Cohn, M. and T. R. Hughes, Jr. 1962. Nuclear magnetic resonance spectra of adenosine Di-and- tri-phosphate II. Effect of complexing with divalent metal ions. *J. Biol. Chem.* 237: 176-180.
- Cohn, W. E. and C. E. Carter. 1950. The separation of adenosine poly-phosphates by ion exchange chromatography. *J. Amer. Chem. Soc.* 72: 4273-4275.
- Cohn, M. and T. R. Hughes, Jr. 1962. Nuclear magnetic resonance spectra of adenosine Di-and tri-phosphate II. Effect of complexing with divalent metal ions. *J. Biol. Chem.* 237: 176-180.
- Cole, H. A., J. M. Winpenny and D. E. Hughes. 1967. The ATP pool in *Escherichia coli*. I. Measurement of the pool using a modified luciferase assay. *Biochim. Biophys. Acta* 143: 445-453.

- Coleman, R. L. 1971. An investigation of the utility of parallel photometric analysis as a new approach for environmental determinations. Ph.D. Thesis, University of Tennessee. 115 pp.
- Colowick, S. P. and H. M. Kalckar. 1943. The role of myokinase in transphosphorylations. I. The enzymatic phosphorylation of hexoses by adenylyl pyrophosphate. *J. Biol. Chem.* 148: 117-126.
- Conklin, A. R. and A. N. MacGregor. 1972. Soil adenosine triphosphate: extraction recovery and half-life. *Bull. Environ. Contamination and Tox.* 7: 296-298.
- Conn, R. B., P. Charachs and E. W. Chappelle. 1975. Limits of applicability of the firefly and luminescence ATP assay for the detection of bacteria in clinical specimens. *Am. J. Clin. path.* 63:493-501.
- Coombs, J., P. J. Halicki, O. Holm-Hansen and B. E. Volcani. 1967a. Studies on the biochemistry and fine structure of silica shell formation in diatoms. I. Changes in concentration of nucleoside triphosphates during synchronized division of Cylindrotheca fusiformis Reimann and Lewin. *Exp. Cell. Res.* 47: 302-314.
- Coombs, J., P. J. Halicki, O. Holm-Hansen and B. E. Volcani. 1967b. Studies on the biochemistry and fine structure of silica shell formation in diatoms. II. Changes in concentration of nucleoside triphosphate in silicon-starvation synchrony of Navicula pelliculose (Breb.) Hilse. *Exp. Cell. Res.* 47: 315-328.
- Cooney, D. A., H. N. Jayram, E. R. Homan and C. F. Motley. 1974. A facile radiometric technique for measuring ATP. *Anal. Biochem.* 62: 157-165.
- Cormier, M. H., J. Lee and D. M. Hercules (Eds.). 1973. *Chemiluminescence and bioluminescence*. Plenum Press, New York.
- Cormier, M. J. and J. R. Totter. 1964. Bioluminescence. *Annu. Rev. Biochem.* 33: 431-458.
- Cunningham, H. W. and R. G. Wetzel. 1978. Fulvic acid interferences on ATP determinations in sediments. *Limnol. Oceanogr.* 23: 166-173.
- Daly, K. F. 1974. The luminescence biometer in the assessment of water quality and wastewater analysis. *American Laboratory* 6: 38-44.
- Damoglou, A. P. and E. A. Dawes. 1967. Studies on the lipid content and phosphate requirement of glucose- and acetate-grown Escherichia coli. *Proc. Biochem. Soc. 467th meeting*, 5-6 Jan. (*Biochem. J.* 102: 32P).
- Daulas, R. and M. Fiala. 1969. Evaluation de la matiere organique vivante dans les eaux marines par la mesure de l'adenosine triphosphate. *Mar. Biol.* 3: 243-246.
- Daumas, R. A. 1973. Evaluation de la teneur en adenosine triphosphate des organismes marins: possibilitees actuelles et limites de la methode. *Tethys* 5: 71-80.

- Davidson, E. A. 1959. Specificity of pyruvate kinase. *Biochim. Biophys. Acta.* 33: 238-240.
- Dawe, L. L. and W. R. Penrose. 1978. Bactericidal property of seawater: Death or debilitation? *Appl. Environ. Microbiol.* 35: 829-833.
- Dawes, E. A. and P. J. Large. 1970. Effect of starvation on the viability and cellular constituents of *Zymomonas anaerobia* and *Zymomonas mobilis*. *J. Gen. Microbiol.* 60: 31-42.
- D'Eustachio, A. J. and D. R. Johnson. 1968. An instrument approach to rapid microbiology. DuPont Internal Study. pp. 10.
- DeLeo, P. and G. G. Sermanni. 1968. Interferences of myokinase in the determination of ATP by luciferin assay. *Biochim. Biophys. Acta* 170: 208-210.
- DeLuca, M. 1969. Hydrophobic nature of the active site of firefly luciferase. *Biochemistry* 8: 160-166.
- DeLuca, M., L. Brand, T. A. Cebula, H. H. Seliger and A. F. Makula. 1971. Nanosecond time-resolved proton transfer studies with dehydroluciferin and its complex with luciferase. *J. Biol. Chem.* 246: 6702-6704.
- DeLuca, M. and M. E. Dempsey. 1972. Mechanism of bioluminescence and chemiluminescence elucidated by use of oxygen. *Chemiluminescence and Bioluminescence* 18: 345-360.
- DeLuca, M. and W. D. McElroy. 1965. The hydrolase properties of firefly luciferase. *Biochem. Biophys. Res. Commun.* 18: 836-842.
- DeLuca, M., G. W. Wirtz and W. D. McElroy. 1964. Role of sulfhydryl groups in firefly luciferase. *Biochemistry* 3: 935-939.
- Deming, J. W., M. W. McGarry, E. W. Chappelle and G. L. Picciolo. 1977. Procedures for the quantitation of bacteria in fluids: firefly luciferase ATP assay used with concentration procedures, in 2nd Bi-annual ATP Methodology Symposium (G. A. Borun, Ed.) SAI Technology, San Diego.
- Denburg, J. and M. DeLuca. 1968. Pseudo-allosteric behavior of firefly luciferase. *Biochem. Biophys. Res. Commun.* 31: 453-460.
- Denburg, J. L., R. T. Lee and W. D. McElroy. 1969. Substrate binding properties of firefly luciferase. I. Luciferin-binding site. *Arch. Biochim. Biophys.* 134: 381-394.
- Denburg, J. L. and W. D. McElroy. 1970. Catalytic subunit of firefly luciferase. *Biochemistry* 9: 4619-4624.
- Derr, R. F. and L. Zieve. 1972. Adenylate energy charge: Relation to guanylate energy charge and the adenylate kinase equilibrium constant. *Biochem. Biophys. Res. Commun.* 49: 1385-1390.

- Devol, A. H., T. T. Packard and O. Holm-Hansen. 1976. Respiratory electron transport activity and adenosine triphosphate in the oxygen minimum of the eastern tropical North Pacific. *Deep-Sea Res.* 23: 963-973.
- Devulder, V. K. 1969. Jahreszeitliche, tiefenabhängige und örtliche Vera im wechselseitigen Verhältnis von Nucleinsäuren, Eiweiß und Chlorophyll in Netzplankton. *Ber. Dt. wiss. Komm. Meeresforsch.* 20: 216-255.
- DeWeer, P. and A. C. Lowe. 1973. Myokinase equilibrium: An enzymatic method for the determination of stability constant of  $Mg^{++}$  complexes with ATP, ADP, and AMP in media of high ionic strength. *J. Biol. Chem.* 298: 2829-2835.
- Dhople, A. M., J. H. Hanks and E. W. Chappelle. 1971. Ultra sensitive method for detection of microbial and myobacterial ATP (abstr). *Bacteriol. Proc.* Vol. 77:11.
- Dhople, A. M. and J. H. Hanks. 1973a. Extraction of bacterial ATP from host-grown organisms. *Bacteriol. Proc.* 73, 15.
- Dhople, A. M. and J. H. Hanks. 1973b. Quantitative extraction of adenosine triphosphate from cultivable and host-grown microbes: calculation of adenosine triphosphate pools. *Appl. Microbiol.* 26: 399-403.
- Dhople, A. M. and J. H. Hanks. 1973. Microbes: Calculation of Adenosine triphosphate pools. *Appl. Microbiol.* 26: 399-403.
- Dhople, A. M. and J. H. Hanks. 1975. Energetics of host-dependent Mycobacterium lepraemarium during transition to a capacity for extracellular growth. *NASA SP-388*, p. 67-73.
- Douglas, W. W. and A. M. Poisner. 1966a. Evidence that the secreting adrenal chromaffin cell releases catecholamines directly from ATP-rich granules. *J. Physiol.* 183: 236-248.
- Douglas, W. W. and A. M. Poisner. 1966b. Relation between ATP splitting and secretion in the adrenal of chromaffin cell: extrusion of ATP (unhydrolyzed) during release of catecholamines. *J. Physiol.* 183: 249-256.
- Doxtader, K. G. 1969. Estimation of microbial biomass in soil on the basis of ATP measurements. *Bacteriol. Proc.*, 69, 14.
- Doxtader, K. G., J. W. Nyhan and E. M. Bautista. 1972. Microbial respiration and ATP levels in grassland soil. *Bacteriol. Proc.* 72, 13.
- Dufresne, L. and H. J. Gitelman. 1970. A semiautomated procedure for determination of ATP. *Anal. Biochem.* 37: 402-408.
- Ebadi, M. S., B. Weiss and E. Costa. 1971. Microassay of adenosine-3', 5'-monophosphate (cyclic AMP) in brain and other tissues by the luciferin-luciferase system. *J. Neurochem.* 18: 183-192.

- Eppley, R. W., A. F. Carlucci, O. Holm-Hansen, D. Kiefer, J. J. McCarthy and P. M. Williams. Evidence for eutrophication in the sea near Southern California coastal sewage outfalls. *Calif. Mar. Res. Commun.* 16: 74-83.
- Eppley, R. W., A. F. Carlucci, O. Holm-Hansen, D. Kiefer, J. J. McCarthy, E. Venrick and P. M. Williams. 1971. Phytoplankton growth and composition in shipboard cultures supplied with nitrate, ammonium, or urea as the nitrogen source. *Limnol. Oceanogr.* 16: 741-751.
- Eppley, R. W., W. G. Harrison, S. W. Chisholm and E. Stewart. 1977. Particulate organic matter in surface waters off Southern California and its relationship to phytoplankton. *J. Mar. Res.* 35: 671-696.
- Erkenbrecher, C. W., S. J. Crabtree and L. H. Stevenson. 1976. Computer-assisted analysis of adenosine triphosphate data. *Appl. Environ. Microbiol.* 32: 451-454.
- Erkenbrecher, C. W. and L. H. Stevenson. 1975. The influence of tidal flux on microbial biomass in salt marsh creeks. *Limnol. Oceanogr.* 20: 618-625.
- Erkenbrecher, C. W. and L. H. Stevenson. 1978. The transport of microbial biomass and suspended material in a high marsh creek. *Can. J. Microbiol.* 24: 839-846.
- Ernst, W. 1970. ATP als indikator für biomasse mariner sedimente. *Oecologia* 5: 56-60.
- Estabrook, R. W. and P. K. Maitra. 1962. A fluorimetric method for the quantitative microanalysis of adenine and pyridine nucleotides. *Anal. Biochem.* 3: 369-382.
- Euler, U. S., F. Lishajko and L. Stjarne. 1963. Catecholamines and adenosine triphosphate in isolated adrenergic nerve granules. *Acta Physiol. Scand.* 59: 495-496.
- Fanica-Gaignier, M., J. Clement-Metral and M. D. Kamen. Adenine nucleotide levels and photopigment synthesis in a growing photosynthetic bacterium. *Biochim. Biophys. Acta* 226: 135-143.
- Farber, E. 1973. ATP and cell integrity. *Fed. Proc.* 32: 1534-1539.
- Faupel, R. P., H. J. Seitz, W. Tarnowski, V. Thiemann and C. H. Weiss. 1972. The problem of tissue sampling from experimental animals with respect to freezing technique, anoxia, stress and narcosis. *Arch. Biochem. Biophys.* 148: 509-522.
- Ferguson, R. L. and M. B. Murdoch. 1972. Microbial biomass in the Newport River Estuary as measured by luciferin-luciferase-ATP reaction: Instrumentation, design, calibration and preliminary data. In: Center for Estuarine and Menhaden Research Annual Report to the Atomic Energy Commission. pp. 271-277.

- Ferguson, R. L. and M. B. Murdoch. 1973. Microbiology of the Newport River Estuary. N.O.A.A. Nat. Mar. Fish. Ser. Annual Report to Atomic Energy Commission.
- Ferguson, R. L. and M. B. Murdoch. 1975. Microbial ATP and organic carbon in sediments of the Newport River estuary, N. C. p. 229-250. In: L. E. Cronin (ed.), Estuarine research, VI. Academic Press.
- Ferguson, R. L. and P. L. Rublee. 1976. Contribution of bacteria to standing crop of coastal plankton. *Limnol. Oceanogr.* 21: 141-144.
- Forrest, W. W. 1965. Adenosine triphosphate pool during the growth cycle in Streptococcus faecalis. *J. Bacteriol.* 90: 1013-1016.
- Forrest, W. W. and D. J. Walker. 1965a. Synthesis of reserve materials for endogenous metabolism in Streptococcus faecalis. *J. Bacteriol.* 89: 1448-1452.
- Forrest, W. W. and D. J. Walker. 1965b. Control of glycolysis in washed suspensions of Streptococcus faecalis. *Nature* 207: 46-48.
- Forrest, W. W. 1969. Energetic aspects of microbial growth, *Symp. Soc. Gen. Microbiol.* 19: 65-86.
- Forsberg, C. W. and K. Lam. 1977. Use of adenosine 5-triphosphate as an indicator of the microbiota biomass in rumen contents. *Appl. Environ. Microbiol.* 33: 528-537.
- Fox, J. L. and M. S. Moyer. 1973. Some effects of a power plant on marine microbiota. *Chesapeake Sci.* 14: 1-10.
- Franzen, J. S. and S. B. Binkley. 1961. Comparison of the acid soluble nucleotide in Escherichia coli at different growth rates. *J. Biol. Chem.* 236: 515-519.
- Frei, J. 1962. Metabolisme energetique du leucocyte. I. Determination enzymatique par fluorimetrie des di- et triphosphates d'adenosine des leucocytes peritoneaux du cobaye. *Enzymol. Biol. Clin.* 2: 175-187.
- Fric, F. and A. Haspelova-Horvatovicova. 1967. Ein Beitrag zur schnellen quantitativen bestimmung von ATP, ADP and AMP in grünen pflanzenorganen. *Phytochemistry* 6: 633-640.
- Gabridge, M. G. and R. B. Polisky. 1977. Intracellular levels of adenosine triphosphate in hamster trachea organ cultures exposed to Mycoplasma pneumoniae cells or membranes. *In Vitro* 13: 510-516.
- Gadkari, D. and H. Stolp. 1975. Energy metabolism of Bdellovibrio bacteriovorus. I. Energy Production, ATP pool, energy charge. *Arch. Microbiol.* 102: 179-185.
- Gajdos, A. and M. Gajdox-Torok. 1969. The quantitative regulation of the biosynthesis of porphyrins by intracellular ATP concentration. *Biochem. Med.* 2: 372-388.

- Gemerden, H. van. 1968. On the ATP generation by Chromatium in darkness. Arch. Mikrobiol. 64: 170-176.
- Giblett, E. R. 1969. Glucose - 6 - phosphate dehydrogenase. p. 512. In: Genetic Markers in Human Blood. Blackwell, Oxford. 629 p.
- Gibson, J. and S. Morita. 1967. Changes in adenine nucleotides of intact Chromatium D produced by illumination. J. Bacteriol. 93: 1544-1550.
- Gilboe, D. P. and F. Q. Nuttall. 1972. The role of ATP and glucose 6-phosphate in the regulation of glycogen synthetase D phosphatase. Biochem. Biophys. Res. Commu. 48: 898-906.
- Glassman, T. A., C. Cooper, L. W. Harrison and T. J. Swift. 1971. A proton magnetic resonance study of metal ion-adenine ring interactions in metal ion complexes with adenosine triphosphate. Biochemistry 10: 843-851.
- Goerke, H. and W. Ernst. 1975. ATP content of estuarine nematodes: Contribution to the determination of meiofauna biomass by ATP measurements. In: Proc. 9th Europ. Mar. Biol. Symp. H. Barnes, (Ed.), Aberdeen Univ. Press, Aberdeen, pp. 13-15.
- Goto, Y. and I. Ito. 1957. Effect of Adrenaline in blood ATP level in diabetic patients. Tohuka J. Exp. Med. 67: 65.
- Goto, Y., I. Ito and Y. Ujiie. 1958. Effect of oral administration of glucose on blood ATP level in diabetic patients. Tohuka J. Exp. Med. 68: 51.
- Greaves, M. P., R. E. Wheatley, H. Shepherd and A. H. Knight. 1973. Relationship between microbial populations and adenosine triphosphate in a basin plat. Soil Biol. Biochem. 5: 685-687.
- Green, A. A. and W. D. McElroy. 1956. Crystalline firefly luciferase. Biochim. Biophys. Acta, 20: 170-176.
- Greengard, P., F. Brink and S. P. Colowick. 1954. Some relationships between action potential, oxygen consumption and coenzyme content in degenerating peripheral axons. J. Cell. Comp. Physiol. 44: 395-420.
- Grylls, F. S. M. 1961. The chemical composition of yeasts. In: Biochemists Handbook. C. Long (Ed.). Van Nostrand, N.Y., pp. 1050-1053.
- Grys, S. and A. Malinowska. 1972. The effect of tetracycline on the level of nicotinamide and adenine nucleotides and of flavins in the tissues of piglets. Pol. Arch. Weter. 15: 787-788.
- Guinn, G. and M. P. Eidenbock. 1972. Extraction, purification, and estimation of ATP from leaves, floral buds, and immature fruits of cotton. Anal. Biochem. 50: 89-97.
- Gurd, J. W. and P. G. Scholefield. 1971. The energy metabolism of Novikoff ascites hepatoma cells: I. The effects of glucose on the intracellular level of adenine nucleotides. Can. J. Biochem. 49: 686-694.

- Hamilton, R. D. and O. Holm-Hansen. 1967. Adenosine triphosphate content of marine bacteria. *Limnol. Oceanogr.* 12: 319-324.
- Hamilton, R. D., O. Holm-Hansen and J. D. H. Strickland. 1968. Notes on the occurrence of living microscopic organisms in deep water. *Deep-Sea Res.* 15: 651-656.
- Hammerstedt, R. H. 1973. An automated method for ATP analysis utilizing the luciferin-luciferase reaction. *Anal. Biochem.* 52: 449-455.
- Hanks, J. H. and A. M. Dhople. 1975. Ultrasensitive bioluminescent determinations of adenosine triphosphate (ATP) for investigating the energetics of host-grown microbes. In: Chappelle, E. W. and G. L. Picciolo (Eds.) Analytical applications of bioluminescence and chemiluminescence, NASA SP-388, Washington, D. C. p. 5765.
- Harrison, D. E. F. and P. Maitra. 1968. The role of ATP in the control of energy metabolism in growing bacteria. *J. Gen. Microbiol.* 53: vii-viii.
- Harrison, D. E. and P. K. Maitra. 1969. Control of respiration and metabolism in growing *Klebsiella aerogenes*. The role of adenine nucleotides. *Biochem. J.* 112: 597-630.
- Harvey, E. N. 1940. Living light. Princeton Univ. Press, 328 pp.
- Hastings, J. W. 1968. Bioluminescence. *Annu. Rev. Biochem.* 37:579-630.
- Heinle, D. R. and D. A. Flemer. 1976. Flows of materials between poorly flooded tidal marshes and an estuary. *Mar. Biol.* 35: 359-373.
- Hellman, B., L. A. Idahl and A. Danielson. 1966. Adenosine triphosphate levels of mammal  $\beta$ -cells after stimulation with glucose and sulfonylurea compounds. *Diabetes* 18: 509.
- Hempling, H. G., C. C. Stewart and G. Gasic. 1969. The effect of exogenous ATP on the electrolyte content of TA<sub>3</sub> ascites tumor cells. *J. Cell. Physiol.* 73: 133-140.
- Henry, R. J. and A. E. Smith. 1946. Use of sulfuric acid - dichromate mixture in cleaning glassware. *Science* 104: 426-427.
- Herbland, A. et J. Pages. 1975. L'adenosine triphosphate (ATP) dans le dome de Guinee distribution verticale et signification ecologique. *Cah. O.R.S.T.O.M., Ser. Oceanogr.* 13: 163-169.
- Hernandez, E. and M. J. Johnson. 1967. Energy supply and cell yield in aerobically grown microorganisms. *J. Bacteriol.* 94: 996-1001.
- Hiltibran, R. C. 1971. Effects of cadmium, zinc, manganese, and calcium on oxygen and phosphate metabolism in bluegill liver mitochondria. *J. Water Pollution Control Fed.* 43: 818-823.



- Hiltz, D. F., L. J. Bishop and W. J. Dyer. 1974. Accelerated nucleotide degradation and glycolysis during warming to and subsequent storage at  $-5^{\circ}\text{C}$  of prerigor, quick-frozen adductor muscle of the sea scallop (Placopecten magellanicus). *J. Fish. Res. Bd. Can.* 31: 1181-1187.
- Hinkle, P. C. and R. E. McCarty. 1978. How cells make ATP. *Scientific American* 238: 104-123.
- Ho, K. P. and W. J. Payne. 1979. Assimilation efficiency and energy contents of prototrophic bacteria. *Biotechnol. Bioengin.* 21: 787-802.
- Hobbie, J. E., O. Holm-Hansen, T. T. Packard, L. R. Pomeroy, R. W. Sheldon, J. P. Thomas and W. J. Wiebe. 1972. Distribution and activity of microorganisms on ocean water. *Limnol. Oceanogr.* 17: 544-555.
- Hobson, P. N. and R. Summers. 1972. ATP pool and growth yield in Selenomonas ruminantium. *J. Gen. Microbiol.* 70: 351-360.
- Hodson, R. E., O. Holm-Hansen and F. Azam. 1976. Improved methodology for ATP determination in marine environments. *Mar. Biol.* 34: 143-149.
- Hodson, R. E. and F. Azam. 1977. Determination and biological significance of dissolved ATP in seawater. p. 127-140. In: G. A. Borun (ed.), 2nd Bi-annual ATP Methodology Symposium, SAI Technology Co., San Diego.
- Holm-Hansen, O. 1968. Ecology, physiology, and biochemistry of blue-green algae. *Annu. Rev. Microbiol.* 22: 47-70.
- Holm-Hansen, O. 1969. Determination of microbial biomass in ocean profiles. *Limnol. Oceanogr.* 14: 740-747.
- Holm-Hansen, O. 1970a. ATP levels in algal cells as influenced by environmental conditions. *Plant Cell Physiol.* 11: 689-700.
- Holm-Hansen, O. 1970b. Determination of microbial biomass in deep ocean water. In: *Organic Matter in Natural Waters*. D. W. Hood (Ed.). Institute of Marine Science. Univ. of Alaska Press, College, Alaska. pp. 287-300.
- Holm-Hansen, O. 1970c. Microbial distribution in ocean water relative to nutrients and food sources. In: *Biological Sound Scattering in the Ocean*. G. B. Farquhar, Ed. Marine Center for Ocean Science, Washington, D. C., pp. 149-157.
- Holm-Hansen, O. 1971. Determination of microbial biomass in deep ocean profiles. In: *Fertility of the Sea*. J. D. Costlow, Jr. (Ed.). Gordon and Breach Ltd., London, pp. 197-207.
- Holm-Hansen, O. 1973a. The use of ATP determinations in ecological studies. *Bull. Ecol. Res. Comm. (Stockholm)* 17: 215-222.

- Holm-Hansen, O. 1973b. Determination of total microbial biomass by measurement of adenosine triphosphate. In: Estuarine Microbial Ecology, H. L. Stevenson and R. R. Colwell (Eds.). University of South Carolina Press, Columbia, S. C. pp. 73-89.
- Holm-Hansen, O. and C. R. Booth. 1966. The measurement of adenosine triphosphate in the ocean and its ecological significance. *Limnol. Oceanogr.* 11: 510-519.
- Holm-Hansen, O., R. Hodson, F. Azam. 1975. Applications of adenine nucleotide measurements in oceanography. In: Chappelle, E. W. and G. L. Picciolo (Eds.) Analytical applications of bioluminescence and chemiluminescence, NASA SP-388, Washington, D. C. p. 75-87.
- Holm-Hansen, O., T. T. Packard and L. R. Pomeroy. 1970. Efficiency of the reverse flow filter technique for concentration of particulate matter. *Limnol. Oceanogr.* 15: 832-835.
- Holm-Hansen, O. and H. W. Paerl. 1972. The applicability of ATP determination for estimation of microbial biomass and metabolic activity. In: Detritus and its ecological role in aquatic ecosystems. Mem. Ist. Ital. Idrobiol. 29 (Suppl.): 149-168.
- Holm-Hansen, O., M. B. Shattuck and D. M. Karl. 1977. Effects of luciferin concentration on light emission from luciferase-ATP preparations In: 2nd BiAnnual ATP Methodology Symposium, G. A. Borun (Ed.). SAI Technology, San Diego.
- Holm-Hansen, O., J. H. D. Strickland and P. M. Williams. 1966. A detailed analysis of biologically important substances in a profile off Southern California. *Limnol. Oceanogr.* 11: 548-561.
- Holms, W. H., I. D. Hamilton and A. G. Robertson. 1972. The rate of turnover of the adenosine triphosphate pool of *Escherichia coli* growing aerobically in simple defined media. *Arch. Microbiol.* 83: 95-109.
- Holmsen, H. 1972. Compartmentation of adenine nucleotide in platelets. In: Erythrocytes, Thrombocytes, Leucocytes. E. Gelach, K. Moser, E. Deutsch and W. Wilmanns (Eds.) 1 II. Int. Symp. Vienna, p. 321-327.
- Holmsen, J., I. Holmsen and A. Bernhardsen. 1966. Microdetermination of adenosine diphosphate and adenosine triphosphate in plasma with the firefly luciferase system. *Anal. Biochem.* 17: 456-473.
- Holmsen, H., E. Storm and H. J. Day. 1972. Determination of ATP and ADP in blood platelets: a modification of the firefly luciferase assay for plasma. *Anal. Biochem.* 46: 489-501.
- Holmsen, H., C. A. Setkowky and H. J. Day. 1974. Effects on antimycin and 2-deoxyglucose on adenine nucleotides in human platelets: Role of metabolic adenosine triphosphate in primary aggregation, secondary aggregation and shape change of platelets. *Biochem. J.* 144: 385-396.

- Hopkins, T. A., H. H. Seliger, E. H. White and M. W. Cass. 1967. The chemiluminescence of firefly luciferin. A model for the bioluminescent reaction and identification of the product excited state. *J. Am. Chem. Soc.* 89: 7148-7150.
- Horska, S. and D. Sofrova. 1971. Relative variations of ATP, ADP and AMP levels in the blood of mother and fetus exhibiting signs of intrauterine hypoxia during childbirth. *Biol. Neonate* 19: 194-199.
- Horska, S. D., D. Sofrova and Z. K. Stembera. 1969. Variation of the ATP content of foetal blood after hypoxia in childbirth. *Biol. Neonate*. 14: 279-285.
- Hoskins, D. D., D. T. Stephens and E. R. Casillas. 1971. Enzymic control of fructolysis in primate spermatozoa. *Biochim. Biophys. Acta* 237: 227-238.
- Howell, S. L. and R. B. L. Ewart. 1973. Synthesis and secretion of growth hormone in the rat anterior pituitary. *J. Cell. Sci.* 12: 23-35.
- Huennekens, F. M. and H. R. Whitely. 1970. Phosphoric acid anhydrides and other energy-rich compounds. In: *Comparative Biochemistry*. Vol. 1, M. Florkin and H. S. Mason (Eds.). Academic Press, Inc., New York pp. 107-180.
- Hulett, H. R. 1970. Non-enzymatic hydrolysis of adenosine phosphates. *Nature* 225: 1248-1249.
- Hysert, D. W., F. Kovacs and N. M. Morrison. 1976. A firefly bio-luminescence ATP assay method for rapid detection and enumeration of brewery microorganisms. *Amer. Soc. Brew. Chemists* 34: 145-150.
- Imai, S., A. L. Riley and R. M. Berne. 1964. Effect of ischemia on adenine nucleotides in cardiac and skeletal muscle. *Circulation Res.* 15: 443-450.
- Imshenetskii, A. A., M. D. Evdokimova and G. C. Sotnikov. 1971. Methods of rupturing microbial cells for determining their ATP content. *Mikrobiologiya* 40: 730-734.
- Imshenetskii, A. A., T. F. Koudrat'eva and G. G. Sotnikov. 1970. Effect of high vacuum on ATP stability in microorganisms. *Mikrobiologiya* 39: 986-989.
- Imshenetskii, A. A., S. V. Lysenko, G. G. Sotnikov and S. S. Abyzov. 1973. Influence of ultra low temperatures on the ATP of microorganisms. *Mikrobiologiya* 42: 579-581.
- Ireland, D. M. and D. C. B. Mills. 1966. Detection and determination of adenosine diphosphate and related substances in plasma. *Biochem. J.* 99: 283-296.
- Ivanovici, A. 1976. The adenylate energy charge as an estimator of stress in an estuarine mollusc. *Aust. Mar. Sci. Bull.*, Annual Conference of AMSA.

- Iwamura, T., T. Kanazawa and K. Kanazawa. 1963. Nucleotide metabolism in Chlorella. II. Quantitative changes in component nucleotides of the acid soluble fraction of alga cells during their life-cycle. In: Japanese Soc. Plant. Physiol. Studies on Microalgae and photosynthetic bacteria. Univ. Tokyo Press, Tokyo, pp. 587-596.
- Izzo, R. S. Firefly assay method for adenosine triphosphate. Aminco Reprint 454. Am. Inst. Co., Silver Spring, Maryland.
- Jaffe, M. J. 1973. The role of ATP in mechanically stimulated rapid closure of the Venus's Flytrap. Plant Physiol. 51: 17-18.
- Jallet, P.M.M., P. Laget, H. Guerin, et J. Piere. 1972. Determination de la constante d'equilibre du complexe MgATP a l'aide du system luciferin-luciferase. C. R. Acad. Sci. Paris, p. 274.D: 603-606.
- Janglova, Z., J. Suchy and Z. Vanek. 1969. Intracellular ATP concentration in Streptomyces aureofaciens. Folia Microbial. 14: 208-210.
- Jannasch, H. W. and G. E. Jones. 1959. Bacterial populations in sea water as determined by different methods of enumeration. Limnol. Oceanogr. 4: 128-139.
- Jimenez, L., E. Rozengurt, J. J. Devalle and H. Carminatti. 1971. Some kinetic differences between the M isoenzymes of pyruvate kinase from liver and muscle. Biochim. Biophys. Acta. 235:326-334.
- Johnson, F. H., H. C. Edward and Y. Haneda. 1961. The luciferin luciferase reactions. In: A Symposium on Light and Life. W. D. McElroy and B. Glass, Eds. Johns Hopkins Univ. Press, Baltimore, Maryland. pp. 206-218.
- Johnson, R., J. H. Gentile and S. Cheer. 1974. Automatic sample injector-its application in the analysis of adenosine triphosphate. Anal. Biochem. 60: 115-121.
- Johnson, R. A., J. G. Hardman, A. E. Broadus and E. W. Sutherland. 1970. Analysis of adenosine 3', 5'-monophosphate with luciferase luminescence. Anal. Biochem. 35: 91-97.
- Johansson, M. and E. Hagerby. 1974. Influence of growth conditions, metabolic inhibitors, and phenolic compounds on the ATP pool in Fomes annosus. Physiol. Plant. 32: 23-32.
- Jones, J. G. and B. M. Simon. 1977. Increased sensitivity in the measurement of ATP in freshwater samples with a comment on the adverse effect of membrane filtration. Freshwater Biol. 1: 253-260.
- Jones, P. C. T. 1969. The effect of temperature on ATP levels in cells of an ascitic hepatoma. J. Cell Physiol. 73: 37-42.
- Jones, P. C. T. 1970a. The effect of light, temperature, and anesthetics on ATP levels in the leaves of Chenopodium rubrum and Phaseolus vulgaris. J. Exp. Bot. 21: 58-63.

- Jones, P. C. T. 1970b. The interaction of light and temperature in determining ATP levels in the myxamoeba of the cellular slime mould Dictyostelium discoideum Acr. 12. Cytobios 2: 89-94.
- Jones, P. C. T. 1970c. Studies on the nature of circadian rhythms: I. Nucleoside phosphate levels in the tissue of wakeful and sleeping golden hamsters. Cytobios 2: 175-179.
- Kachmar, J. F. and P. D. Boyer. 1953. Kinetic analysis of enzyme reactions II. The potassium activation and calcium inhibition of pyruvic phosphorase. J. Biol. Chem. 200, 669-682.
- Kalbhen, D. A., R. Domenjoz and K. Ehlers. 1967. The effect of sodium salicylate on the ATP level in normal and inflamed tissue. Life Sci. (Oxford) 6: 1883-1886.
- Kalbhen, D. A., K. Ehlers and R. Domenjoz. 1967. Effect of sodium salicylate on dextran edema of rat paws and on the ATP content in inflamed tissues. Med. Pharmacol. Exp. 16: 552-560.
- Kalbhen, D. A. and H. J. Koch. 1967. Methodische untersuchungen zur quantitativen mikrobestimmung von ATP in biologischen material mit dem firefly-enzyme system. Zeit. Klin. Chem. Klin. Biochem. 5: 299-311.
- Kalbhen, D. A. and H. J. Koch. 1967. The quantitative microdetermination of ATP in biological material with the firefly enzyme system. Zeit. Kin. Chem. Klin. Biochem. 5: 299-304.
- Kalckar, H. M. 1943. The role of myokinase in transphosphorylations, II. The enzymatic action of myokinase on adenine nucleotides. J. Biol. Chem. 148, 127-137.
- Kalckar, H. M. 1947. Differential spectrophotometry of purine compounds by means of specific enzymes. II. Determination of adenine compounds. J. Biol. Chem. 167: 445-459.
- Kaplan, N. D., S. P. Colowick and C. C. Barnes. 1951. Effect of alkali on diphosphopyridine nucleotide. J. Biol. Chem. 191: 461-472.
- Karl, D. M. 1978. Occurrence and ecological significance of ATP in the ocean and microbial cells. Appl. Environ. Microbiol. 36: 349-355.
- Karl, D. M., J. A. Haugsness, L. Campbell and O. Holm-Hansen. 1978. Adenine nucleotide extraction from unicellular organisms and beach sand: ATP recovery, energy charge ratios and determination of carbon/ATP ratios. J. Exp. Mar. Biol. Ecol. 34: 163-181.
- Karl, D. M. and O. Holm-Hansen. 1976. Effects of luciferin concentration on the quantitative assay of ATP using crude luciferase preparations. Anal. Biochem. 75: 100-112.
- Karl, D. M. and O. Holm-Hansen. 1977. Adenylate energy charge measurements in natural sea water and sediment samples. In: 2nd Bi-Annual ATP methodology Symposium, G. A. Borun (Ed.). SAI Technology, San Diego.

- Karl, D. M. and O. Holm-Hansen. 1978a. ATP, ADP and AMP determinations in water samples and algal cultures. In: Handbook of Physiological Methods, Physiological and Biochemical Methods. J. A. Hellebust and J. S. Craigie (Eds.).
- Karl, D. M. and O. Holm-Hansen. 1978b. Methodology and measurement of adenylate energy charge ratios in environmental samples. *Mar. Biol.* 48: 185-199.
- Karl, D. M. and P. A. LaRock. 1974. Extraction and measurements of adenosine triphosphate in sediments and sands. *Bacteriol. Proc.* 74, 55.
- Karl, D. M. and P. A. LaRock. 1975. Adenosine triphosphate measurements in soil and marine sediments. *J. Fish. Res. Board Can.* 32: 599-607.
- Karl, D. M., P. A. LaRock, J. W. Morse and W. Sturges. 1976. Adenosine triphosphate in the North Atlantic Ocean and its relationship to the oxygen minimum. *Deep-Sea Res.* 23: 81-88.
- Karl, D. M., J. W. Morse, R. Scott and L. Butler. 1974. ATP and DOC measurements in the Mid-Atlantic. *Trans. Am. Geophys. Union* 55: 294.
- Karl, D. M., J. W. Morse, and W. Sturges. 1976. Adenosine triphosphate in the North Atlantic Ocean and its relationship to the oxygen minimum. *Deep-Sea Res.* 23: 81-88.
- Karl, D. M. 1978. Distribution, abundance, and metabolic states of microorganisms in the water column and sediments of the Black Sea. *Limnol. Oceanogr.* 23, 936-949.
- Karl, D. M. and D. B. Craven. 1980. Effects of Alkaline phosphatase activity on nucleotide measurements in aquatic microbial communities. *Appl. Environ. Microbiol.* 40:549-561.
- Kao, I. C., S. Y. Chiu, L. T. Fan and L. E. Erickson. 1973. ATP pools in pure and mixed cultures. *J. Water Poll. Control Fed.* 45: 926-931.
- Kayne, F. J. 1973. Pyruvate Kinase. p. 353-382. In: P. D. Boyer (ed.), The Enzymes, third edition, Vol. 8, Academic Press, N.Y.
- Kees, U., A. Lewenstein and R. Bachofen. 1975. ATP pools in activated sludge. *Eur. J. Appl. Microbiol.* 2: 59-64.
- Kelbaugh, Burton N., Grace L. Picciolo, Emmett Chappelle and Maurice E. Colbur, inventors (to NASA) 51 p. Filed 4 Aug. 1970 (NASA-Case-GSC-1116901:U.S.-Patent-Appl.-SN-60882). Avail: NTIS CSCL06M. Automatic Instrument for Chemical Processing to Detect Microorganisms in Biological Samples by Measuring Light Reactions.
- Kelbaugh, Burton N., inventor (to NASA), Grace L. Picciolo, inventor (to NASA), Emmette Chappelle, inventor (to NASA) Filed 30 Apr. 1971. (NASA-Case-GSC-11169-2: U.S.-Patent-Appl.-SN-139094) Avail: NTIS CSCL 06B #N71-34079 NASA GSFC. Automatic Instrument for Chemical Processing to Detect Microorganisms in Biological Samples by Measuring Light Reactions.

- Kelbaugh, Burton N., Grace L. Picciolo, Emmette W. Chappelle, and Maurice E. Colburn, inventors (to NASA) issued 4 Sept. 1973, 16 p., Filed 30 Apr. 1971. Supersedes N71-34079 (09 - 21 p. 3385) Continuation-in-part of abandoned U.S. Patent Appl. Sn-60882, filed 4 Aug. 1970. (NASA-Case-GSC 11169-2; U.S.-Patent-3, 756, 920; U.S. Pat. Appl. SN 139094; U.S. Pat. Class 195 127; U.S. Patent Appl. SN 60882) Avail: U.S. Patent Office CSCS 068. Automatic Instrument for Chemical Processing to Detect Microorganisms in Biological Samples by Measuring Light Reactions.
- Kelly, D. P. and P. J. Surett. 1966. Energy coupling during sulfur compound oxidation in Thiobacillus. *J. Gen. Microbiol.* 43: 109-118.
- Kennard, O., N. W. Issacs, J. C. Coppola, A. J. Kirby, S. Warren, W.D.S. Motherwell, D. G. Watson, D. L. Wampler, D. H. Chenery, A. C. Larson, K. A. Kerr and L. R. Di Sanseverion. 1970. Three dimensional structure of adenosine triphosphate. *Nature* 225: 333-336.
- Kerr, S. E. 1941a. On the preparation of adenosine triphosphate. *J. Biol. Chem.* 139: 121-130.
- Kerr, S. E. 1941b. Notes on the preparation of muscle adenylic acid. *J. Biol. Chem.* 139: 131-134.
- Keys, A. J. 1968. The intracellular distribution of free nucleotides in the tobacco leaf. *Biochem. J.* 108: 1-8.
- Khoo, J. C. and P. J. Russell. 1972. Isoenzymes of adenylate kinase in human tissue. *Biochem. Biophys. Acta* 268, 98-101.
- Khorana, H. G. 1954. Carbodiimides. Part V. A Novel Synthesis of Adenosine di- and triphosphate and  $P^1$ ,  $P^2$ - diadenosine-5'-pyrophosphate. *J. Amer. Chem. Soc.* 76: 3517-3522.
- Kilbert, L. H., Jr., M. S. Schiff and P. O. Foa. 1972. The measurement of ATP in pancreatic islets by means of the luciferin-luciferase reaction. *Horm. Metab. Res.* 4: 242-244.
- Kimberg, D. V. and K. L. Yielding. 1962. Pyruvate kinase, structural and functional changes induced by diethylstilbestrol and certain steroid hormones. *J. Biol. Chem.* 237, 3233-3239.
- Kirkpatrick, F. H., D. G. Willman and P. L. LaCelle, 1975. A23187 and red cells: changes in deformability  $K^+$ ,  $Mg^{2+}$ ,  $Ca^{2+}$ , and ATP. *Experientia* 131, 653-654.
- Klofat, W., G. Picciolo, E. W. Chappelle and E. Freese. 1969. Production of Adenosine triphosphate in normal cells and sporulation mutants of Bacillus subtilis. *J. Biol. Chem.* 244: 3270-3276.
- Klungsoyr, L., J. H. Hageman, L. Fall and D. E. Atkinson. 1968. Interaction between energy charge and product. Feedback in the regulation of biosynthetic enzymes. Aspartokinase, phosphoribosyl adenosine triphosphate synthetase, and phosphoribosyl pyrophosphate synthetase. *Biochemistry* 7, 4035-4040.

- Knauer, G. A. and A. V. Ayers. 1977. Changes in carbon, nitrogen, adenosine triphosphate, and chlorophyll a in decomposing Thalassia testudinum leaves. *Limnol. Oceanogr.* 22, 408-414.
- Knowles, C. J. 1977. Microbial metabolic regulation by adenine nucleotide pools. In: *Microbial Energetics* (B.A. Haddock and W. A. Hamilton, Eds. Cambridge Univ. Press, Cambridge. Vol. 27: 241-283.
- Knowles, C. J. and L. Smith. 1970. Measurements of ATP levels of intact Azotobacter vinelandii under different conditions. *Biochim. Biophys. Acta* 197: 152-160.
- Knust, E. A., E. W. Chappelle and G. L. Picciolo. 1975. A comparison of certain extracting agents for extraction of adenosine triphosphate (ATP) from microorganisms for use in the firefly luciferase ATP assay. p. 27-34. In: E. W. Chappelle and G. L. Picciolo (eds.), *Analytical Applications of Bioluminescence and Chemiluminescence*. NASA, SP-338. Washington, D. C.
- Kolburn, D. G. 1969. The energetics of mammalian cell growth. *J. Cell. Sci.* 4: 645-654.
- Krane, S. M. and M. J. Glimcher. 1962. Transphosphorylation from nucleoside di- and triphosphates by apatite crystals. *J. Biol. Chem.* 237: 2291-2998.
- Kullmann, L., G. E. Staal and H. W. Wouters. 1972. Increased ATP concentration in erythrocytes in neurofibromatosis. *Clin. Sci.* 42: 775-779.
- Kylin, A. and J. E. Tillberg. 1967. The relation between total photo-phosphorylation, level of ATP, and oxygen evolution in Scenedesmus as studied with DCMU and antimycin A. *Zeit. Pflanzphysiol.* 58: 165-174.
- Laborde, P. L. 1972. L'adenosine triphosphate des micro-organismes marins planctoniques. Rapports avec la biomasse et al productivité primaire. These de Doctorat, Universite d' Aix-Marseille. pp. 108.
- Laborde, P. L. and H. J. Minas. 1974. Influence des conditions hivernales sur les productions phyto-et zooplanctoniques en Mediterranée Nord-Occidentale. IV. Distribution verticale des taux d'adenosine triphosphate et de production primaire. *Marc. Bio.* 25: 203-212.
- Lannergren, C. and H. R. Skjoldal. 1976. The spring phytoplankton bloom in Lindispollene, a land-locked Norwegian Fjord. Autotrophic and heterotrophic activities in relation to nutrients. IN: *Proceedings of the Tenth European Marine Biology Symposium*, Vol. 12, p. 363-391. G. Persoone and E. Jaspers, Eds. Wetteren: Universa Press.
- Laug, E. P. 1934. Retention of dichromate by glassware. *J. Ind. Eng. Chem.* 6, 111-112.
- Law, A. T. and D. K. Button. 1977. "Low substrate concentration determinations for nutrient limited growth kinetic studies: I. Glucose Analysis with hexokinase." *Mar. Sci. Commu.* 3, 21-34.



- Lazdunski, A. and J. P. Belaich. 1972. Uncoupling in bacterial growth: ATP pool variation in Zymomonas mobilis cells in relation to different uncoupling conditions of growth. J. Gen. Microbiol. 70: 187-197.
- Lee, C. C., R. F. Harris, J. K. Syers and D. E. Armstrong. 1971a. Adenosine triphosphate content of Selenastrum capricornutum. Appl. Microbiol. 21: 957-958.
- Lee, C. C., R. F. Harris, J. D. H. Williams, D. E. Armstrong and J. K. Syers. 1971b. Adenosine triphosphate in lake sediments: I. Determination. Soil Sci. Soc. Am. Proc. 35: 82-86.
- Lee, C. C., R. F. Harris, J. D. H. Williams, J. K. Syers and D. E. Armstrong. 1971c. ATP in lake sediments. II. Origin and significance. Soil. Sci. Soc. Am. Proc. 35: 86-91.
- Lee, J., C. L. Murphy, G. J. Faini and T. L. Baucom. 1974. Bacterial bioluminescence and its application to analytical procedures, in Liquid scintillation counting, recent developments. Academic Press, New York p. 403-420.
- Lee, J. and H. H. Seliger. 1965. Absolute spectral sensitivity of phototubes and the application to the measurement of the absolute quantum yields of chemiluminescence and bioluminescence. Photochem. and Photobiol. 4: 1015-1948.
- Lee, R. T., J. L. Denburg and W. D. McElroy. 1970. Substrate-binding properties of firefly luciferase. II. ATP-binding site. Arch. Biochem. Biophys. 141: 38-52.
- Lee, R. T. and W. D. McElroy. 1969. Role and reactivity of sulfhydryl groups in firefly luciferase. Biochemistry 8: 130-136.
- Lee, R. T. and W. D. McElroy. 1971a. Effects of 5'-adenylic acid on firefly luciferase. Arch. Biochem. Biophys. 145: 78-84.
- Lee, R. T. and W. D. McElroy. 1971b. Isolation and partial characterization of a peptide derived from the luciferin binding site of firefly luciferase. Arch. Biochem. Biophys. 146: 551-556.
- Leenders, H. J., A. Kemp, A. F. Koninky and J. Rosing. 1974. Changes in cellular ATP, ADP, AMP levels following treatments affecting cellular respiration and the activity of certain nuclear genes in Drosophila salivary glands. Exp. Cell. Res. 86: 25-30.
- Leese, H. J. and J. R. Bronk. 1972. Automatic fluorometric analysis of micromolar quantities of ATP, glucose, and lactic acid. Anal. Biochem. 45: 211-221.
- Lehninger, A. L. 1965. Bioenergetics. Benjamin, Inc., New York. pp. 258.
- Lehninger, A. L. 1971. Bioenergetics. 2nd ed. W. A. Benjamin, Inc., Menlo Park, Calif. 245 pp.

- Leonard, F. and A. F. Hegyeli. 1973. A mechanism for topical cutaneous calcinosis. *Calcif. Tissue Research*, 13: 331-334.
- Leonard, N. J. and R. A. Laursen. 1965. Synthesis and properties of analogs of adenosine diphosphate, adenosine triphosphate, and nicotinamide adenine dinucleotide derived from 3- $\beta$ -D-ribofuranosyladenine (3-isoadenosine). *Biochemistry* 4: 365-376.
- Levin, G. V., C. S. Chen and G. Davis. 1967. Development of the firefly bioluminescence assay for the rapid, quantitative detection of microbial contamination in water. *Aerosp. Med. Res. Lab.*, TR-67-71.
- Levin, G. V., J. R. Clendenning, E. W. Chappelle, A. H. Heim, and E. Rocek. 1964. A rapid method for detection of microorganisms by ATP assay: its possible application in virus and cancer studies. *BioScience* 14: 37-38.
- Levin, G. V. and A. H. Heim. 1965. Gulliver and Diogenes - Exobiological antitheses. *In: Life and Space. Research Vol. III.* M. Florin, Ed. Academic Press, New York, pp. 105-119.
- Levin, G. C., J. R. Schrot and W. C. Hess. 1970. Methodology for adenosine triphosphate determination in waste water treatment. *Env. Sci. Technol.* 10, 961-965.
- Levin, G. A., E. Usdin and A. R. Slonim. 1968. Rapid detection of microorganisms in aerospace water systems. *Aerosp. Med.* 39: 14-16.
- Lewenstein, A. and R. Bachofen. 1972. Transient induced oscillations in the level of ATP in *Chlorella fusca*, *Biochim. Biophys. Acta* 267, 80-85.
- Lewenstein, A. and R. Bachofen. 1978. CO<sub>2</sub> - Fixation and ATP synthesis in continuous cultures of *Chlorella fusca*. *Arch. Microbiol.* 116, 169-173.
- Lieberman, M. A., M. Simon and J. Hong. 1977. Characterization of *Escherichia coli* mutant incapable of maintaining a transmembrane potential. *J. Biol. Chem.* 252, 4056-5067.
- Lin, S. and H. P. Cohen. 1968. Measurement of adenosine triphosphate content of crayfish stretch receptor cell preparations. *Anal. Biochem.* 24: 531-540.
- Lipmann, F. F. 1941. Metabolic generation and utilization of phosphate bond energy. *In: Adv. in Enzymology and Related Subjects.* F. F. Nord and C. H. Werkman, Eds., Vol. I, Interscience Publishers, Inc. p. 99-162.
- Litchfield, C. D. and P. L. Seyfried. 1979. Methodology for biomass determination and microbial activities in sediments. ASTM (STP673) Philadelphia. 199 p.
- Lohmann, K. and O. Meyerhof. 1934. Über die enzymatische Umwandlung von Phosphoglycerinsäure in Brenztraubensäure und Phosphorsäure. *Biochem. Zeit.* 273: 60-72.

- Loos, J. A., R. C. H. VanDoorn and D. Roos. 1973. A sensitive mechanized determination of ATP and ADP. *Anal. Biolchem.* 53: 309-312.
- Loppert, H. G. and E. Broda. 1973. ATP-Gehalt and ATP-Umsatz von *Chlorella* in Abhängigkeit von Belichtung und Belüftung. *Zeit. Allg. Mikrobiol.* 13: 499-506.
- Lowe, C. U., M. W. Rees and R. Markham. 1963. Synthesis of complex organic compounds from simple precursors: Formation of amino-acids, amino-acid polymers, fatty acids, and purines from ammonium cyanide. *Nature* 199: 219-222.
- Lowenstein, J. M. 1958. Transphosphorylations catalysed by bivalent metal ions. *Biochem. J.* 70: 222-230.
- Lowry, O. H., J. Carter, J. B. Ward and L. Glaser. 1971. The effect of carbon and nitrogen sources on the level of metabolic intermediates in *Escherichia coli*. *J. Biol. Chem.* 246: 6511-6521.
- Lowry, O. H., J. V. Passonneau, F. X. Hasselberger and D. W. Schulz. 1964. Effect of Ischemia on known substrates and cofactors of the glycolytic pathways in brain. *J. Biol. Chem.* 239: 18-30.
- Lundin, A., A. Rickardsson and A. Thore. 1976. Continuous monitoring of ATP-converting reactions by purified firefly luciferase. *Anal. Biochem.* 75: 611-620.
- Lundin, A., A. Rickardsson, and A. Thore. 1977. Substrate and enzyme determination by continuously monitoring the ATP level by purified luciferase reagent, in 2nd Biannual ATP methodology Symposium (G. A. Borun Ed.) SAI Technology, San Diego.
- Lundin, A. and A. Thore. 1975a. "Analytical information obtainable by evaluation of the time course of firefly bioluminescence in the assay of ATP." *Anal. Biochem.* 66: 47-63.
- Lundin, A. and A. Thore. 1975b. Comparison of methods for extraction of bacterial adenine nucleotides determined by firefly assay. *Appl. Microbiol.* 30: 713-721.
- Lutkenhaus, J., J. Ryan and M. Konrad. 1973. Kinetics of phosphate incorporation into adenosine triphosphate and guanosine triphosphate in bacteria. *J. Bacteriol.* 116: 1113-1123.
- Lyman, G. E. and T. P. DeVincenzo. 1967. Determination of picogram amounts of ATP using the luciferin-luciferase enzyme system. *Anal. Biochem.* 21: 435-443.
- MacLeod, N. H., E. W. Chappelle and A. M. Crawford. 1969. ATP assay of terrestrial soils: A test of an exobiological experiment. *Nature* 223: 267-268.
- Mague, T. H., N. M. Weare and O. Holm-Hansen. 1974. Nitrogen fixation in the North Pacific Ocean. *Mar. Biol.* 24: 109-119.

- Manandhar, M. S. P. and K. Van Dyke. 1974a. Adenosine tetraphosphate analysis: Polyethyleneimine (PEI) thin layer purification and firefly extract liquid scintillation quantitation. *Anal. Biochem.* 58: 368-375.
- Manandhar, M. S. P. and K. Van Dyke. 1974b. Guanosine triphosphate analysis: PEI-thin layer purification and luciferin-luciferase scintillation quantitation. *Anal. Biochem.* 60: 112-129.
- Manandhar, M. S. P. and K. Van Dyke. 1974c. Delayed luminescence analysis (DLA) of purine and pyrimidine ribose and deoxyribose nucleotide triphosphate in picomole quantities. *Microchem. J.* 19: 42-51.
- Mandelstam, J. and McQuillen. 1968. *Biochemistry of bacterial growth*. John Wiley and Son, New York. 540 pp.
- Manuels, M. W. and H. Postma. 1974. Measurements of ATP and organic carbon in suspended matter of the Dutch Wadden Sea. *Neth. J. Sea. Res.* 8: 292-311.
- Masterson, T. S., A. D. Mason and W. L. Brown. 1964. Instrument for the quantitation of ATP utilizing firefly bioluminescence. *Am. J. Med. Electron.* 3: 283-285.
- McCapra, F. 1973. Chemistry of bioluminescence. *Endeavour* 32: 139-145.
- McClare, C. W. F. 1972. In defense of the high energy phosphate bond. *J. Theor. Biol.* 35: 233-246.
- McElroy, W. D. 1947. The energy source for bioluminescence in an isolated system. *Proc. Nat. Acad. Sci. U. S.* 33: 342-345.
- McElroy, W. D. 1956. *Cypridina* and firefly luciferase. *In: Methods in Enzymology*. Vol. II. S. P. Colowick and N. O. Kaplan, Eds. Academic Press, Inc. New York. pp. 445-448.
- McElroy, W. D. 1963a. Crystalline firefly luciferase. *In: Methods in Enzymology*. Vol. VI. S. P. Colowick and N. O. Kaplan, Eds. Academic Press, Inc. New York. pp. 445-448.
- McElroy, W. D. 1963b. Preparation and properties of firefly luciferyl and oxyluciferyl adenylic acid. *In: Methods in Enzymology*. Vol. VI. S. P. Colowick and N. O. Kaplan, Eds. Academic Press, Inc., New York pp. 775-782.
- McElroy, W. D. and M. DeLuca. 1973. Chemical and enzymatic mechanisms of firefly luminescence. *In: Chemiluminescence and Bioluminescence*. M. J. Cormier, D. M. Hercules and J. Lee, Eds. Plenum Press, New York. pp. 285-312.
- McElroy, W. D. and H. B. Glass (Editors). 1961. *A Symposium on Light and Life*. John Hopkins Univ. Press, Baltimore, Maryland.
- McElroy, W. D. and C. S. Rainwater. 1948. Spectral energy distribution of the light emitted by firefly extracts. *J. Cell. Comp. Physiol.* 32: 421-425.

- McElroy, W. D. and H. H. Seliger. 1966. Firefly bioluminescence. In: Bioluminescence in Progress. F. H. Johnson and Y. Haneda, Eds. Princeton University Press, Princeton, New Jersey. pp. 427-458.
- McElroy, W. D., H. H. Seliger and E. H. White. 1969. Mechanism of bioluminescence, chemiluminescence and enzyme function in the oxidation of firefly luciferin. Photochem. Photobiol. 10: 153-170.
- McElroy, W. D. and B. L. Strehler. 1949. Factors influencing the response of the bioluminescent reaction of ATP. Arch. Biochem. Biophys. 22: 420-433.
- Meyerhof, O. and P. Oesper. 1949. The enzymatic equilibria of phospho(enol) pyruvate. J. Biol. Chem. 179: 1371-1385.
- Mildvan, A. S. 1970. Metals in enzyme catalysis. p. 445-536. In: P. D. Boyer (ed.). The Enzymes, third edition, Vol. 2. Academic Press, N.Y.
- Mildvan, A. S. and M. Cohn. 1965. Kinetic and magnetic resonance studies of the pyruvate kinase reaction. J. Biol. Chem. 240: 238-246.
- Milekhina, E. I., E. E. Rakhleeva, G. C. Sotnikov and I. O. Ivanov. 1970. Use of the luciferin-luciferase method for ATP determination in the cells of Azobacter vinelandii. Mikrobiologiya 39: 722-726.
- Mills, D. C. B. 1973. Changes in the adenylate energy charge in human blood platelets induced by adenosine diphosphate. Nature 243: 220-222.
- Miovic, M. L. and J. Gibson. 1971. Nucleotide pools in growing Chromatium strain D. J. Bacteriol. 108: 954-956.
- Miovic, M. L. and J. Gibson. 1973. Nucleotide pools and adenylate energy charge in balanced and unbalanced growth of Chromatium. J. Bacteriol. 114: 86-95.
- Mitchell, G. W. and J. W. Hastings. 1971. A stable, inexpensive, solid-state photomultiplier photometer. Anal. Biochem. 39: 243-250.
- Montague, M. D. and E. A. Dawes. 1974. The survival of Peptococcus prevotii in relation to the adenylate energy charge. J. Gen. Microbiol. 80: 291-299.
- Moriarty, D. J. W. 1975. A method for estimating the biomass of bacteria in aquatic sediments and its application to trophic studies. Oecologia. 20: 219-229.
- Morton, R. A., T. A. Hopkins and H. H. Seliger. 1968. The spectroscopic properties of firefly luciferin and related compounds. An approach to product emission. Biochemistry 8: 1598-1607.
- Munch-Petersen, A. and J. Neuhard. 1964. Studies on the acid-soluble nucleotide pool in thymine-requiring mutants of Escherichia coli during thymine starvation. I. Accumulation of deoxyadenosine triphosphate in Escherichia coli. Biochim. Biophys. Acta 80: 542-551.

- Nakao, M., T. Nakao, M. Tatibana and H. Yoshikawa. 1960. Phosphorus metabolism in human erythrocyte. III. Regeneration of adenosine triphosphate in long-stored erythrocyte by incubation with inosine and adenine. *J. Biochem.* 47: 661-671.
- Nakao, T., T. Wada, T. Kamiyama, M. Nakao and K. Nagano. 1962. A direct relationship between ATP levels and in vivo viability of erythrocytes. *Nature* 194: 877-878.
- Navon, S. and A. Agrest. 1963. ATP content in the central nervous system of rats exposed to chronic hypercapnea. *Am. J. Physiol.* 205: 957-958.
- Nazar, R. N., H. G. Lawford and J. T. Wong. 1970. An improved procedure for extraction and analysis of cellular nucleotides. *Anal. Biochem.* 35: 305-313.
- Neilson, A. H., O. Holm-Hansen and R. A. Lewin. 1972. An obligately autotrophic mutant of *Chlamydomonas dysosmos*: a biochemical elucidation. *J. Gen. Microbiol.* 71: 141-148.
- Neufeld, H. A., R. D. Towner and J. Pace. 1975. A rapid method for determining ATP by the firefly luciferin-luciferase system. *Experientia* 31: 391-392.
- Netwon, A. A. and Lord Rothschild. 1961. Energy-rich phosphate compounds in bull semen: comparison of their metabolism with anaerobic heat production and impedance change frequency. *Proc. Roy. Soc. Lond. Ser. B* 155: 183-194.
- Nibley, D. A., G. L. Picciolo, E. W. Chappelle and M. A. McGarry. 1977. Laboratory preparation and evaluation of purified, standardized firefly luciferase for low level ATP measurements, in 2nd Bi-Annual ATP Methodology Symposium (G.A. Borun ed.) SAI Technology, San Diego.
- Neilsen, R. and H. Rasmussen. 1968. Fractionation of extracts of firefly tails by gel filtration. *Acta Chem. Scand.* 22: 1757-1762.
- Nigam, V. N. 1969. ATP level and control of glycolysis in Novikoff ascites-hepatoma cells. *Enzymologia* 36: 257-268.
- Nilsson, L. 1970. The effect of anaesthetics on the tissue lactate, pyruvate, phosphocreatine, ATP, and AMP concentrations and on intracellular pH in the rat brain. *Acta Physiol. Scand.* 80: 142-144.
- Nobel, P. S., D. T. Chang, C. Wong, S. S. Smith and D. E. Barcus. 1968. Initial ATP formation, NADP reduction, CO<sub>2</sub> fixation, and chloroplast flattening upon illuminating pea leaves. <sup>2</sup>Dept. of Botanical Sciences and Molecular Biology Inst., Univ. of Calif.
- Noda, L. 1970. Adenylate kinase. *In*: P. D. Boyer (Ed.), *The Enzymes*, third edition, 8: 279-305. Academic Press, New York.
- Noda, L. 1973. Adenylate kinase. p. 279-305. *In*: P. D. Boyer (Ed.), *The Enzymes*, third edition, Vol. 8. Academic Press, N.Y.

- Nungester, W. J., L. J. Paradise and J. A. Adair. 1969. Loss of cellular ATP as a measure of cytolytic activity of antiserum. *Proc. Soc. Exp. Biol. Med.* 132: 582-586.
- Obendorf, R. L., S. Spiegel, and A. Marcus. 1974. Rapid increase in adenosine 5' triphosphate during early wheat embryo germination. *Plant Physiol.* 53 (Suppl.): 38.
- O'Brien, R. W. and J. G. Morris. 1971. Oxygen and the growth and metabolism of *Clostridium acetobutylicum*. *J. Gen. Microbiol.* 68: 307-318.
- Oeize, J. and M. C. Kamen. 1971. Adenosine triphosphate cellular levels in *Rhodospirillum rubrum* during transition from aerobic to anaerobic metabolism. *Biochim. Biophys. Acta* 234: 137-143.
- Olah, J. 1972. Leaching, colonization and stabilization during detritus formation. *Mem. Ist. Ital. Idrobiol.* 29 Suppl: 105-127.
- Osterman, J. and P. J. Fritz. 1972. Pyruvate kinase isozymes: species differences in mammals. *Fed. Proc.*, 31: 983 Abs.
- O'Sullivan, W. J. and L. Noda. 1968. Magnetic resonance and kinetic studies related to the manganese activation of the adenylate kinase reaction. *J. Biol. Chem.* 243: 1424-1433.
- Oleniacz, W. S., M. A. Pisano, M. H. Rosenfeld and R. L. Elgart. 1968. Chemiluminescent method for detecting microorganisms in water. *Environ. Sci. Tech.* 2: 1030-1033.
- Olson, O. E. and S. A. Kuby. 1964. Studies on ATP transphosphorylases. *J. Biol. Chem.* 239: 460-467.
- Oro, J. 1960. Synthesis of adenine from ammonium cyanide. *Biochem. Biophys. Res. Commun.* 2: 407-412.
- Oro, J. 1963. Studies in experimental organic cosmochemistry. *Ann. New York Acad. Sci.* 108: 464-481.
- Oro, J. and A. P. Kimball. 1962. Synthesis of purines under possible primitive earth conditions. II. Purine intermediates from hydrogen cyanide. *Arch. Biochem. Biophys.* 96: 293-313.
- Osterhelt, D., R. Gottschlich, R. Hartmann, H. Michel, G. Wagner. 1977. Light energy conversion in halobacteria in microbial energetics. (B. A. Haddock and W. A. Hamilton, eds.). Cambridge Univ. Press, p. 333-349.
- Owen, E. M. and M. L. Vasers. 1973. The adaptation of the biometer for screening antimicrobial agents for cosmetic use, in *Developments in Industrial Microbiology*. Vol. 14. 74-79, Amer. Inst. of Biol. Sci.
- Owen, E. M., P. C. Scarry and M. L. Vasers. 1972. Use of the biometer for evaluation of process. *Water Dev. Ind. Microbiol.* 13: 257-263.

- Pacold, I. and L. E. Anderson. 1973. Energy charge control of the calvin cycle enzyme 3-phosphoglyceric acid kinase. *Biochem. Biophys. Res. Commun.* 51: 139-143.
- Paerl, H. W. 1975. Microbial attachment to particles in marine and freshwater ecosystems. *Microbial Ecology* 2: 73-83.
- Paerl, H. W., M. M. Tilzer and C. R. Goldman. 1976. Chlorophyll a versus adenosine triphosphate as algal biomass indicators in lakes. *J. Phycol.* 12: 242-246.
- Palmiano, E. P. and B. O. Juliano. 1972. Biochemical changes in the rice grain during germination. *Plant Physiol.* 49: 751-756.
- Pamatmat, M. M. and H. R. Skjoldal. 1974. Dehydrogenase activity and adenosine triphosphate concentration of marine sediments in Lind and Spollene, Norway. *Sarsia* 56: 1-11.
- Parkinson, P. I. and E. Medley. 1972. The estimation of ATP, ADP, and AMP in human plasma using luciferin/luciferase and a scintillation counter. *Liquid Scintillation Counting*, 2: 109-113. (M. A. Crook, P. Johnson, B. Scales, eds.). Haydon & Son. 327 pp.
- Patterson, J. W., P. L. Breznoik and H. D. Putnam. 1969. Sludge activity parameters and their application to toxicity measurements in activated sludge. *Proc. 24th Purdue Industrial Waste Conference, Purdue University, Lafayette, Indiana.* pp. 127-154.
- Patterson, J. W., P. L. Brezonik and H. D. Putnam. 1970. Measurement and significance of adenosine triphosphate in activated sludge. *Environ. Sci. Technol.* 4: 569-575.
- Paul, E. A. and R. L. Johnson. 1977. Microscopic counting and adenosine 5'-triphosphate measurement in determining microbial growth in soils. *Appl. Environ. Microbiol.* 34: 263-269.
- Payne, W. J. 1970. Energy yields and growth of heterotrophs. *Ann. Rev. Microbiol.* 24: 17-52.
- Pederson, T. A., M. Kirk and J. A. Bassham. 1966. Light-dark transients in levels of intermediate compounds during photosynthesis in air-adapted Chlorella. *Physiol. Plant.* 19: 219-231.
- Peterson, R. N. and M. Freund. 1970. ATP synthesis and oxidative metabolism in human spermatozoa. *Biol. Reprod.* 3: 47-54.
- Phillips, R. C., P. George and R. J. Ruman. 1969. Thermodynamic data for the hydrolysis of ATP as a function of pH,  $Mg^{2+}$  ion concentration and ionic strength. *J. Biol. Chem.* 244: 3330-3342.
- Picciolo, G. L., E. W. Chappelle and A. J. Fleig. 1971a. An automated luciferase assay for bacteria in urine. *NASA Tech. Manual TM X-65521.*



- Picciolo, G. L., B. N. Kelbaugh, E. W. Chappelle and M. C. Colburn. 1971b. Automatic bio-sample bacteria detection system. NASA Tech. Brief 71-10055. Washington, D. C.
- Picciolo, G. L., B. N. Kelbaugh, E. W. Chappelle and A. J. Fleig. 1971c. An automatic luciferase assay of bacteria in urine. NASA, GSFC X-641-71-163. Greenbelt, Maryland.
- Picciolo, G. L., B. N. Kelbaugh, E. W. Chappelle and A. J. Fleig. 1971d. Automatic bio-sample bacteria detection system. NASA Tech. Brief 71-10055, GSFC, Greenbelt, Maryland.
- Picciolo, G. L., D. A. Nibley and E. W. Chappelle. 1975a. Characteristics of three (3) commercial photometers for use with liquid luminescence systems. ATP Methodology Conference, San Diego, California, G. A. Borun (ed.) SAI Technology, Inc., San Diego, CA.
- Picciolo, G. L., E. W. Chappelle, E. A. Knust, S. A. Tuttle, C. A. Curtis. 1975b. Problem areas in the use of the firefly luciferase assay for bacterial detection. In: Analytical Application of Bioluminescence and Chemiluminescence, Chappelle, E. W. and G. L. Picciolo (eds.). NASA SP-338, p. 1-26, Washington, D. C.
- Picciolo, G. L., E. W. Chappelle, R. R. Thomas and M. McGarry. 1977. Performance characteristics of a new photometer with a moving filter tape for luminescence assay. Appl. Environ. Microbiol. 34: 720-724.
- Plagemann, P. G. W. 1971. Nucleotide pools of Novikoff rat hepatoma cells growing in suspension culture, II. Independent nucleotide pools for nucleic acid synthesis. J. Cell. Physiol. 77: 241-285.
- Plant, P. J., E. H. White and W. D. McElroy. 1968. The decarboxylation of luciferin in firefly bioluminescence. Biochem. Biophys. Res. Commun. 31: 98-103.
- Plowman, K. M. and A. R. Krall. 1965. A kinetic study of nucleotide interactions with pyruvate kinase. Biochemistry. 4: 2809-2814.
- Polakis, E. S. and W. Bartley. 1966. Changes in the intracellular concentrations of adenosine phosphates and nicotinamide nucleotides during the aerobic growth cycle of yeast on different carbon sources. Biochem. J. 99: 521-533.
- Ponnamperuma, C., R. M. Lemmon, R. Mariner and M. Calvin. 1963. Formation of adenine by electron irradiation of methane, ammonia, and water. Proc. Nat. Acad. Sci. USA 49: 737-740.
- Ponnamperuma, C., R. Mariner and C. Sagan. 1963. Formation of adenosine by ultraviolet irradiation of a solution of adenine and ribose. Nature 198: 1199-1200.
- Ponnamperuma, C., S. Sagan and R. Mariner. 1963. Synthesis of ATP under possible primitive earth conditions. Nature 199: 220-226.

- Porumb, S. 1972. Influenta cimpurilor electromagnetice asupra continutului in ATP si ADP in hematite de iepure. Stud. Cercet. Bio. Ser. Zool. 24: 165-168.
- Pradet, A. 1967a. Études des adenosines-5'-mono, di- et tri-phosphates dans les tissus vegetaux. I. Dosage enzymatique. *Physiol. Veg.* 5: 209-221.
- Pradet, A. 1967b. Études des adenosines-5'-mono, di- et tri-phosphates dans les tissus vegetaux. V. Effect in vivo, sur le niveau de la charge energetique, d'un desequilibre induit entre fourniture et utilisation de l'energie dans les semences de laitue. *Physiol. Veg.* 7: 261-275.
- Pradet, A. and J. L. Bomsel. 1968. In vivo control of adenylic nucleotides by adenylate kinase. *C. R. Acad. Sci., Paris, Ser. D* 266: 2416-2418.
- Pradet, A., A. Narayanan and J. Vermeersch. 1968. Plant tissue AMP, ADP, and ATP. III. Energy metabolism during the first stages of germination of lettuce seeds. *Bull. Soc. Fr. Physiol. Veg.* 14: 107-114.
- Prager, J. C., C. S. Hegre, S. Cheer, P. Rogerson, J. Gentile. 1973. Power, Plankton, and Firefly Tails. *Underwater Naturalist* 8: 18-22.
- Prydz, H. and L. O. Froholm. 1964. The use of a recording Beckman DB spectrophotometer for the assay of ATP with a luciferin-luciferase reagent. *Acta Chem. Scand.* 18: 554-556.
- Quammen, M. L., P. A. LaRock and J. A. Calder. 1973. Environmental effects of pulp mill wastes. In: *Estuarine Microbiol. Ecology*. H. Stevenson and R. R. Colwell, Eds. Univ. of South Carolina Press, Columbia, South Carolina, pp. 329-343.
- Qureshi, A. A. and J. Patel. 1976. Adenosine triphosphate (ATP) levels in microbial cultures and a review of the ATP biomass estimation technique. *Scientific Series No. 63*, Environment Canada.
- Racker, E. 1972. Bioenergetics and the problem of tumor growth. *Am. Sci.* 60: 50-63.
- Ramaiah, A., J. A. Hataway and D. E. Atkinson. 1964. Adenylate as a metabolic regulator. *J. Biol. Chem.* 239: 3619-3622.
- Ramaiah, A. and J. D. Nicholas. 1964. The synthesis of ATP and the incorporation of  $^{32}\text{P}$  by cell free preparations from *Nitrosomonas europaea*. *Biochim. Biophys. Acta* 86: 459-465.
- Ramirez, J. and L. Smith. 1968. Synthesis of adenosine triphosphate in intact cells of *Rhodospirillum rubrum* and *Rhodopseudomonas spheroides* on oxygenation or illumination. *Biochim. Biophys. Acta* 153: 466-475.
- Rasmussen, H. and R. Neilsen. 1968. An improved analysis of adenosine triphosphate luciferase method. *Acta Chem. Scand.* 22: 1745-1762.
- Raymond, P., A. Narayanan and A. Pradet. 1973. Evidence for the presence of cyclic AMP in plant tissues. *Biochem. Biophys. Res. Commun.* 53: 1115-1121.

- Reasons, R. H. and R. S. Hikida. 1973. Biochemistry of adenosine triphosphate activity of avian fast and slow muscle. *Exp. Neurol.* 38: 27-32.
- Reynard, A. M., L. F. Hass, D. D. Jacobsen and P. D. Boyer. 1961. The correlation of reaction kinetics and substrate binding with the mechanism of pyruvate kinase. *J. Biol. Chem.* 236: 2277-2283.
- Reporter, M. 1972. An ATP pool with rapid turnover within the cell membrane. *Biochem. Biophys. Res. Commu.* 48: 598-604.
- Richards, O. W. 1936. Killing organisms with chromium as from incompletely washed bichromate-sulfuric-acid cleaned glassware. *Physiol. Zool.* 9: 246-253.
- Riemann, B. 1979. The occurrence and ecological importance of dissolved ATP in fresh water. *Freshwater Biol.* 9: 481-490.
- Riemers, H. J., M. A. Packham and J. F. Mustard. 1975. Labeling of the releasable adenine nucleotide part of human platelets. *Fed. Proc.* 34: 855.
- Rhee, S. G., M. I. Greifner and P. B. Chock. 1975. Determination of adenosine 5-triphosphate by the luciferin-luciferase system with a stopped-flow spectrometer. *Analyt. Biochem.* 66: 259-264.
- Rhodes, W. C. and W. D. McElroy. 1958. The synthesis and function of luciferyl-adenylate and oxylucifery-adenylate. *J. Biol. Chem.* 233: 1528-1536.
- Robertson, A. M. and R. S. Wolfe. 1970. Adenosine triphosphate pools in *Methanobacterium*. *J. Bacteriol.* 102: 43-51.
- Roman, M. R. 1978. Tidal resuspension in Buzzards Bay, Massachusetts. II. Seasonal changes in the size distribution of chlorophyll, particle, concentration, carbon and nitrogen in resuspended particulate matter. *Estuarine and Coastal Mar. Sci.* 6: 47-53.
- Rooney, D. W. and J. J. Eiler. 1969. Effects of division-synchronizing hypoxia and hyperthermic shocks upon *Tetrahymena* respiration and intracellular ATP pools. *J. Cell. Biol.* 41: 145-153.
- Rottino, A., G. T. Hoffman and H. Albaum. 1952. Adenine nucleotide content of red blood cells of normal individuals and of patients suffering from Hodgkin's disease and other diseases. *Blood* 7: 836-841.
- Rudd, J. W. M. and R. D. Hamilton. 1973. Measurement of adenosine triphosphate (ATP) in two Precambrian Shield lakes of Northwestern Ontario. *J. Fish. Res. Board Can.* 30: 1537-1546.
- Rusness, D. G. and G. G. Still. 1974. Firefly luciferase inhibition by isopropyl-3-chlorocarbonilate and isopropyl-3-chloro-hydroxycarbonilate analogues. *Pest. Biochem. Physiol.* 4: 109-119.

- Ryther, J. H. and W. M. Dunstan. 1971. Nitrogen, phosphorus, and eutrophication in the coastal marine environment. *Science* 171: 1008-1013.
- Sakshang, E. and O. Holm-Hansen. 1977. Chemical composition of *Skeletonema costatum* (Grev.) Cleve and *Povlova (Monochrysis) lutheri* (Droop) Green as a function of nitrate-, phosphate, and iron limited growth. *J. Exp. Mar. Biol. Ecol.* 29: 1-34.
- Salisbury, J. L. and G. L. Floyd. 1978. Calcium-induced contraction of the rhizoplast of a quadriflagellate green alga. *Science* 202: 975-977.
- Salzmann, E. W., D. A. Chambers and L. L. Neri. 1966. Possible mechanism of aggregation of blood platelets by adenosine diphosphate. *Nature* 210: 167.
- Sandmeier, M. and J. Ivart. 1972. Modification du taux des nucleotides adenyliques (ATP, ADP, AMP), par un eclairement de lumiere rouge clair (660 nm). *Photochem. Photobiol.* 16: 51-59.
- Santarius, J. A. and U. Heber. 1965. Changes in the intracellular levels of ATP, ADP, AMP and Pi and regulatory function of the adenylate system in leaf cells during photosynthesis. *Biochim. Biophys. Acta* 102: 39-54.
- Schlegel, J. U., G. E. Pipkin and G. N. Shultz. 1969. The etiology of bladder tumors. *Brit. J. Urol.* 41: 718-723.
- Schmidt, G. L. and M. D. Kamen. 1970. Variable cellular composition of *Chromatium* in growing cultures. *Arch. Mikrobiol.* 73: 1-18.
- Schmidt, G. L. and M. D. Kamen. 1971. Control of chlorophyll synthesis in *Chromatium vinosum*. *Arch. Mikrobiol.* 76: 51-64.
- Schmitz, H., R. B. Hurlbert, V. R. Potter. 1954. Nucleotide metabolism. III. Mono-, di-, and triphosphates of cytidine, guanosine, and uridine. *J. Biol. Chem.* 209: 41-54.
- Schneider, P. W. and H. Brintzinger. 1964. Zum Mechanismus der Metallionen-Katalysierten Hydrolyse von Adenosintriphosphat (ATP). *Helv. Chim. Acta* 47: 1717-1733.
- Schon, G. 1969. Der einfluss der kulturbedingungen auf den ATP, ADP, und AMP spiegel bei *Rhodospirillum rubrum*. *Arch. Microbiol.* 66: 348-364.
- Schram, E. 1970. Use of scintillation counters for bioluminescence assay of adenosine triphosphate (ATP). In: *The Current Status of Liquid Scintillation Counting*. E. D. Bransome, Jr., Grune and Stratton, Eds. New York. pp. 129-133.
- Schram, E., R. Cortenbosch, E. Gerlo and H. Roosens. 1970. Application of scintillation counters to the assay of bioluminescence. p. 125-135. In: *D. L. Horrocks and C. Peng (eds.), Organic scintillators and liquid scintillation counting*. Academic Press, New York.

- Schram, E. and H. Roosens. 1972. Semi-automatic microtransferator and cell for the bioluminescence assay of ATP and reduced NAD with scintillation counters. *In: Liquid Scintillation Counting. Vol. 2, M. A. Crook, P. Johnson, and B. Scales, Eds. Heydon and Son Ltd., London pp. 115-120.*
- Schram, E. and P. Stanley (eds.). 1978. International Symposium on analytical applications of bioluminescence and chemiluminescence. Vrije Universiteit Brussels. 20 pp.
- Schultz, J. and J. W. Daly. 1973. Cyclic adenosine 3'-5'-monophosphate in guinea pig cerebral cortical slices. *J. Biol. Chem.* 248: 843-852.
- Searle, N. D. 1975. Applications of chemiluminescence to bacterial analysis. *In: Chappelle, E. W. and G. L. Piccolo (eds.) Analytical applications of bioluminescence and chemiluminescence. NASA SP-338, 95-103.*
- Seitz, W. R. 1976. Chemiluminescence analysis in water pollution studies in Modern fluorescence spectroscopy (E. L. Wehry ed.) Vol. I. Plenum, New York. 238 pp.
- Seitz, W. R. and M. P. Neary. 1974. Chemiluminescence and bioluminescence in chemical analysis. *Anal. Chem.* 46: 188A-202A.
- Seliger, H. H. 1972. Applications of bioluminescence and chemiluminescence. *In: Chemiluminescence and Bioluminescence. M. J. Cormier, D. M. Hercules, and J. Lee, Eds. Plenum Press, New York, pp. 461-478.*
- Seliger, H. H., J. B. Beuk, W. G. Fastie and W. D. McElroy. 1964. Spectral distribution of firefly light. *J. Gen. Physiol.* 48: 94.
- Seliger, H. H. and W. D. McElroy. 1959. Quantum yield in the oxidation of firefly luciferin. *Biochem. Biophys. Res. Commun.* 1: 21-24.
- Seliger, H. H. and W. D. McElroy. 1960. Spectral emission and quantum yield of firefly bioluminescence. *Arch. Biochem. Biophys.* 88: 136-141.
- Seliger, H. H. and W. D. McElroy. 1964. The colors of firefly bioluminescence. Enzyme configuration and species specificity. *Proc. Nat. Acad. Sci.* 52: 75-81.
- Seliger, H. H. and W. D. McElroy. 1965. *Light: Physical and Biological Action.* Academic Press, New York, pp. 417.
- Seliger, H. H. and W. D. McElroy, E. H. White and G. D. Field. 1961. Stereospecificity and firefly bioluminescence, a comparison of natural and synthetic luciferins. *Proc. Nat. Acad. Sci.* 47: 1129-1134.
- Senez, J. C. 1962. Some considerations of the energetics of bacterial growth. *Bacteriol. Rev.* 26: 95-107.
- Seto, S., K. Ogura and Y. Nishiyama. 1963. A convenient synthetic method of 2-carbamoyl-6-methoxybenzothiazole, one of the intermediates for the synthesis of firefly luciferin. *Bull. Chem. Soc. Japan* 36: 331-333.

- Sharpe, A. N. 1973. Automation and instrumentation developments for the bacteriology laboratory. In: Sampling - Microbiological Monitoring of Environments. Edited by R. C. Board and D. W. Lovelock. Soc. Appl. Bacteriol. Tech. Ser. 7: 197-232.
- Sharpe, A. N., M. N. Woodrow and A. K. Jackson. 1970. Adenosine triphosphate (ATP) levels in foods contaminated by bacteria. J. Appl. Bacteriol. 33: 758-767.
- Sheldon, R. W. and W. H. Sutcliffe. 1978. Generation times of 3h for Sargasso Sea microplankton determined by ATP analysis. Limnol. Oceanogr. 23: 1051-1055.
- Shen, L., L. Fall, G. M. Walton and D. E. Atkinson. 1968. Interaction between energy charge and metabolite modulation in the regulation of enzymes of amphibolic sequences. Phosphofructokinase and pyruvate dehydrogenase. Biochemistry 7: 4041-4045.
- Shoaf, W. T. and B. W. Lium. 1976. The measurement of adenosine triphosphate in pure algal cultures and natural aquatic samples. J. Res. U. S. Geol. Survey, 4: 241-245.
- Shropshire, W. and K. Bergman. 1968. Light induced concentration changes of ATP from *Phycomyces sporangiophores*: A re-examination. Plant Physiol. 43: 1317-1318.
- Shropshire, W. and R. H. Gettens. 1966. Light induced concentration changes of adenosine triphosphate in *Phycomyces sporangiophores*. Plant Physiol. 41: 203-207.
- Sieburth, J. McN., K. M. Johnson, C. M. Burney and D. M. Lavoie. 1977. Estimation of in situ rates of heterotrophy using diurnal changes in dissolved organic matter and growth rates of picoplankton in diffusion culture. Helgolander wiss. Meeresunters. 30: 565-574.
- Sieburth, J. McN., P.-J. Willis, K. M. Johnson, C. M. Burney, D. M. Lavoie, K. R. Kinga, D. A. Caron, F. W. French III, P. W. Johnson and P. G. Davis. 1976. Dissolved organic matter and heterotrophic microneuston in the surface microlayers of the North Atlantic. Science 194: 1415-2428.
- Sikora, J. P., W. B. Sikora, C. W. Erkenbrecher and B. C. Coull. 1977. Significance of ATP, carbon, and caloric content of meiobenthic nematodes in partitioning benthic biomass. Mar. Biol. 44: 7-14.
- Skjoldal, H. R. and U. Bamstedt. 1976. Studies on the deep-water pelagic community of Korsfjorden, Western Norway. Adenosine phosphates and Nucleic acids: *Meganyctiphanes norvegica* (euphausiacea) in relation to the life cycle. Sarsia 61: 1-14.
- Skjoldal, H. R. and U. Bamstedt. 1977. Ecobiochemical studies on the deep-water pelagic community of Korsfjorden, Western Norway. Adenine Nucleotides in Zooplankton. Mar. Bio. 42: 197-211.

- Slayman, C. L. 1973. Adenine nucleotide levels in Neurospora, as influenced by conditions of growth and by metabolic inhibitors. J. Bacteriol. 114: 752-766.
- Smith, M. and H. G. Khorana. 1958. Nucleoside polyphosphates. VI. An improved and general method for the synthesis of ribo- and deoxyribo-nucleoside 5'triphosphates. J. Amer. Chem. Soc. 80: 1141-1145.
- Smith, R. C. and O. Maaloe. 1964. Effect of growth rate on the acid soluble nucleotide composition in Salmonella typhimurium. Biochim. Biophys. Acta 86: 229-234.
- Sofrova, D. and S. Leblova. 1970. Determination of adenosine triphosphate (ATP) in plant tissue. Photosynthetica 4: 162-184.
- Sojka, G. A., G. A. Din and H. Gest. 1967. Integration of energy conversion and biosynthetic processes in bacterial photosynthesis. Nature 216: 1021-1022.
- Sojka, G. A. and H. Gest. Integration of energy conservation and biosynthesis in the photosynthetic bacterium Rhodospseudomonas capsulata. Proc. Nat. Acad. Sci. 61: 1486-1493.
- Solvonuk, P. F. and H. B. Collier. 1955. The pyruvic phosphoferase of erythrocytes I. Properties of the enzyme and its activity in erythrocytes of various species. Can. J. Biochem. Physiol. 33: 38-45.
- Somlo, M. 1970. Respiration-linked ATP formation by an "Oxidative phosphorylation mutant" of yeast. Arch. Biochem. Biophys. 136: 122-133.
- Sorokin, Y. I. and H. Kadota (Editors). 1972. Total microbial biomass estimation by the ATP method. In: Techniques for the assessment of microbial production and decomposition in fresh waters. IBP Handbook No. 23, Blackwell Scientific Publications, Oxford, Eng. pp. 71-76.
- Sparrow, E. B. and K. G. Doxtader. 1973. Adenosine triphosphate (ATP) in grassland soil. Its relationship to microbial biomass and activity. Tech. Rep. No. 224 Grassland biome. U. S. International Biological Program. 161 pp.
- Spiro, T. G., W. A. Kjellstrom, M. Zeydel and R. A. Butow. 1968. On the kinetics of metal catalyzed ATP dephosphorylation. Biochemistry. 7: 859-865.
- Stadelman, P. and M. Munawar. 1974. Biomass parameters and primary production at a nearshore midlake station of Lake Ontario during IFYGL. Proc. 17th Conf. Great Lakes Res. 109-119.
- Stanford, J. A., J. T. Boswell, D. P. Wilcox and W. B. Perry. 1977. Limnological application of ATP assay in a mesotrophic, Texas reservoir. 2nd Bi-annual ATP methodology Symposium, G. A. Borun (Ed.) SAI Technology, San Diego.

- Stanley, P. E. 1971a. Determination of subpicomole levels of NADH and FMN using bacterial luciferase and the liquid scintillation spectro-photometer. *Anal. Biochem.* 39: 441-453.
- Stanley, P. E. 1971b. The use of the liquid scintillation spectrometer for measuring NADH and FMN by the photobacterium luciferase and ATP by the firefly luciferase, in *Organic Scintillators and Liquid Scintillation Counting* (D. L. Horrocks and C. Peng (eds.)). Academic Press, New York p. 60-616.
- Stanley, P. E. 1974. Analytical bioluminescence assays using the liquid scintillation spectrometer: a review, In: *Liquid Scintillation Counting*. Vol. 3. M. A. Crook and P. Johnson, Eds. Heyden and Son Ltd., London. pp. 253-272.
- Stanley, P. E. and S. G. Williams. 1969. Use of the liquid scintillation spectrometer for determining adenosine triphosphate by luciferase enzyme. *Anal. Biochem.* 29: 381-392.
- Statham, M. and D. Langton. 1975. The use of adenosine triphosphate measurements in the control of the activated sludge process by the solids retention time method. *Process. Biochem.* 10: 25-28.
- Stenlid, G. 1970. Flavenoids as inhibitors of the formation of adenosine triphosphate in plant mitochondria. *Phytochemistry* 9: 251-256.
- Stenlid, G. 1971. The effect of some inhibiting sugars upon the content of ATP in wheat roots. *Physiol. Plant.* 25: 397-401.
- Stewart, J. M. and G. Guinn. 1969. Chilling injury and changes in adenosine triphosphate of cotton seedlings. *Plant Physiol.* 44: 605-608.
- St. John, J. B. 1970. Determination of ATP in *Chlorella* with the luciferin-luciferase enzyme system. *Anal. Biochem.* 37: 409-416.
- St. John, J. B. 1971. Comparative effects of diuron and chloroprotham on ATP levels in *Chlorella*. *Weed Sci.* 19: 274-276.
- St. John, J. B. 1975. Application of the luciferin-luciferase enzyme system for determination of adenosine triphosphate (ATP) to studies on the mechanisms of herbicide action. In: Chappelle, E. W. and G. L. Picciolo (eds) *Analytical applications of bioluminescence and chemiluminescence*, NASA SP-338, Washington, D. C. p. 45-48.
- Strange, R. E., H. E. Wade and F. A. Dark. 1963. Effect of starvation on adenosine triphosphate concentration in *Aerobacter aerogenes*. *Nature* 199: 55-57.
- Strehler, B. L. 1953. Firefly luminescence in the study of energy transfer mechanisms. II. ATP and photosynthesis. *Arch. Biochem. Biophys.* 43: 67-69.
- Strehler, B. L. 1968. Bioluminescence assay: principles and practice. In: *Methods of Biochemical Analysis*. Vol. 16. D. Glick, Ed. Interscience Publishers, John Wiley and Sons, Inc., New York. pp. 99-181.



- Strehler, B. L. and W. D. McElroy. 1957. Assay of adenosine triphosphate. In: Methods in Enzymology. Vol. 3. S. P. Colowick and N. O. Kaplan, Eds. Academic Press, Inc., New York. pp. 871-873.
- Strehler, B. L. and R. Totter. 1952. Firefly luminescence in the study of energy transfer mechanisms: Firefly luminescence and other methods. I. Substrate and enzyme determination. Arch. Biochem. Biophys. 40: 28-41.
- Strehler, B. L. and J. R. Totter. 1954. Determination of ATP and related compounds. In: Methods of Biochemical Analysis. Vol. 1. D. Glick, Ed. Interscience Publishers, John Wiley & Sons, Inc., New York. pp. 341-356.
- Strickland, J. D. H. 1965. Production of organic matter in the primary stages of the marine food chain. In: Chemical Oceanography. Vol. 1. J. P. Riley and G. Skirrow, Eds. Academic Press, Inc. New York.
- Strickland, J. D. H. 1971. Microbial activity in aquatic environments. In: Microbes and Biological Productivity. 21st Symposium of the Society of General Microbiology. Cambridge University Press, Cambridge, Eng. pp. 231-253.
- Strickland, J. D. H., O. Holm-Hansen, R. W. Eppley and R. J. Linn. 1969. The use of a deep tank in plankton ecology. I. Studies of the growth and composition of phytoplankton crops at low nutrient levels. Limnol. Oceanogr. 14: 23-34.
- Strickland, J. D. H. and T. R. Parsons. 1972. Determination of adenosine triphosphate (ATP). In: A Practical Handbook of Seawater Analysis. Fish. Res. Board. Can., Bull. 167, pp. 245-249.
- Sunderland, D. W. and M. J. Merrett. 1967. Respiration rate, adenosine diphosphate and triphosphate concentrations of leaves showing necrotic local lesions following infection by tobacco mosaic virus. Physiol. Plant. 20: 368-372.
- Sutcliffe, W. H. 1972. Some relations of land drainage, nutrients, particulate material, and fish catch in two Eastern Canadian Bays. J. Fish. Res. Bd. Can. 29: 357-362.
- Sutcliffe, W. H., E. A. Orr and O. Holm-Hansen. 1976. Difficulties with ATP measurements in inshore waters. Limnol. Oceanogr. 21: 145-148.
- Suzuki, N. and T. Goto. 1972a. Studies of firefly bioluminescence. II. Tetrahedron, 28: 4075-4082.
- Suzuki, N. and T. Goto. 1972b. Synthesis of 4-thiazolone derivatives related to firefly oxyluciferin. Agr. Biol. Chem. 12: 2213-2221.
- Suzuki, N., M. Sato and T. Goto. 1972. Studies on firefly bioluminescence. I. Synthesis and spectral properties of firefly oxyluciferin, a possible emitting species in firefly bioluminescence. Tetrahedron 28: 4065-4075.
- Swedes, J. S., R. J. Sedo and D. E. Atkinson. 1975. Regulation of growth and protein synthesis to the adenylate energy charge in an adenine requiring mutant of Escherichia coli. J. Biol. Chem. 250: 6930-6938.

- Syrett, P. J. 1958. Respiration rate and internal adenosine triphosphate concentration in Chlorella. Arch. Biochem. Biophys. 75: 117-124.
- Szczecinska, O. and W. Leyko. 1966. Adenine nucleotide content of erythrocytes of diabetes mellitus patients. Clin. Chem. Acta 13: 131-132.
- Tal, E., S. Dickstein and F. G. Sulman. 1964. ATP determination with the Tricrab scintillation counter. Experientia 20: 652.
- Talwalkar, R. T. and R. L. Lester. 1973. The response of diphosphoinositide and triphosphoinositide to perturbations of the adenylate energy charge in cells of Saccharomyces cerevisiae Biochim. Biophys. Acta 306: 412-421.
- Taqi Kahn, M. M. and A. E. Martell. 1962. Metal chelates of adenosine triphosphate. J. Phys. Chem. 66: 10-15.
- Tarkkanen, P. R. Driesch and H. Greiling. 1978. A rapid enzymatic micro-method for the determination of intracellular and extracellular ATP and its clinical-chemical applications. Zeit. Anal. Chem. 290: 180.
- Tetas, M. and J. M. Lowenstein. 1963. The effect of bivalent metal ions on the hydrolysis of ADP and ATP. Biochemistry 2: 350-357.
- Thompson, F. M. and D. E. Atkinson. 1971. Response of nucleoside diphosphate kinase to the adenylate energy charge. Biochem. Biophys. Res. Commun. 45: 1581-1585.
- Thore, A. 1977. An overview of the work of the Swedish bioluminescence group, 2nd Bi-Annual ATP methodology symposium, G. A. Borun (ed.) SAI Technology, San Diego.
- Traganza, E. D. and K. J. Graham. 1977a. Carbon/adenosine triphosphate ratios in marine zooplankton. Deep-Sea Res. 24: 1187-1193.
- Traganza, E. D. 1977b. Carbon/adenosine triphosphate ratios in marine zooplankton. 2nd Bi-annual ATP methodology symposium, G. A. Borun (ed.) SAI Technology, San Diego.
- Travis, J. and W. D. McElroy. 1966. Isolation and sequence of an essential sulfhydryl peptide at the active site of firefly luciferase. Biochemistry 5: 2170-2176.
- Turnock, G. S., S. U. Erickson, B. A. C. Ackrell and B. Birch. 1972. A mutant of Escherichia coli with a defect in energy metabolism. J. Gen. Microbiol. 70: 507-515.
- Ustach, J. F. 1974. The role of organic debris and associated microorganisms in pelagic estuarine food chains. Appendix A. Problems with adenosine triphosphate measurements in the Patuxent River Marshes. Report to Office of Water Resources Research. NRI Ref. No. 74-159. Technical Report No. 29. 16 pp.

- Van Dyke, K., R. Stitzel, R. McClellan and C. Szustkiewicz. 1969a. An automated procedure for the sensitive and specific determination of ATP. *Clin. Chem.* 15: 3-14.
- Van Dyke, K., R. Stitzel, R. McClellan and C. Szustkiewicz. 1969b. Automated analysis of ATP; its application to on-line continuous-flow incubations and measurement of ATP-coupled enzyme system. *Advan. Automat. Anal. Technicon Inst. Congr.* 1: 47-53.
- Vanstaen, H. 1978. ATP as an activity parameter for waste water treatment. *Environ. Safety Control.* Jan-Feb.
- Van Tamelen, E. E., K. B. Sharples, R. Hanzlik, R. B. Clayton, A. L. Burlingame and P. C. Wszolek. 1967. The chemiluminescence of firefly luciferin. A model for the bioluminescent reaction and identification of the product excited state. *J. Amer. Chem. Soc.* 89: 7148-7151.
- Vechet, B., H. Brana and H. Kupkova. 1971. The adenosine triphosphate pool and the adenyl cyclase activity in *Escherichia coli* strains with different sensitivity. *Folia Microbiol.* 16: 31-34.
- Vellend, H., S. A. Tuttle, M. Burza, L. Weinstein, G. L. Picciolo and E. W. Chappelle. 1975. A rapid method for the determination of microbial susceptibility using the firefly luciferase assay for adenosine triphosphate (ATP). In: Chappelle, E. W. and G. L. Picciolo (eds.) *Analytical applications of bioluminescence and chemiluminescence*, NASA SP-388, Washington, D. C. p. 43-44.
- Vinkler, C., G. Rosen and P. D. Boyer. 1978. Light-driven ATP formation from  $^{32}\text{P}_i$  by chloroplast thylakoids without detectable labeling of ADP, as measured by rapid mixing and acid quench techniques. *J. Biol. Chem.* 253: 2507-2510.
- Vlodavsky, I., M. Inbar and L. Sachs. 1973. Membrane changes and adenosine triphosphate content in normal and malignant transformed cells. *Proc. Nat. Acad. Sci. U. S.* 70(6): 1780-1784.
- Wahl, R. and L. J. L. Kozloff. 1962. The nucleoside triphosphate content of various bacteriophages. *J. Biol. Chem.* 237: 1953-1960.
- Wampler, J. E. 1975. Simple photometer circuits using modular electronic components. In: Chappelle, E. W. and G. L. Picciolo (eds.) *Analytical applications of bioluminescence and chemiluminescence*, NASA SP-338, Washington, D. C. p. 105-110.
- Watanabe, S., L. Evenson and I. Gulz. 1963. Effect of  $\text{Mg}^{++}$  and  $\text{Ca}^{++}$  ions on the fluorescence of the protonated ring of adenine, adenosine, AMP, ADP, and ATP. *J. Biol. Chem.* 238: 324-327.
- Watson, S. W. 1978. Role of bacteria in an upwelling ecosystem. p. 139-154. In: R. Boje and M. Tomczak (eds.), *Upwelling Ecosystems*. Springer, Berlin.

- Weber, C. I. 1970. New plankton biomass parameters. Newsletter 4. Analytical Quality Control Laboratory, Federal Water Pollution Control Administration, Environ. Prot. Agency, Cincinnati, Ohio.
- Weber, C. I. 1973a. Recent developments in the measurement of the response of plankton and periphyton to changes in their environment. Environ. Prot. Agency Bioassay Techniques and Environmental Chemistry, pp. 119-138.
- Weber, C. I. 1973b. Biological monitoring of the aquatic environment. In: Biological Methods for the Assessment of Water Quality, ASTM STP, American Society for Testing and Materials, pp. 46-60.
- Weber, G. 1969. Inhibition of human brain pyruvate kinase and hexokinase by phenylalanine and phenylpyruvate: possible relevance to phenylketonuric brain damage. Proc. Nat. Acad. Sci. U. S. 63: 1365-1369.
- Weddle, C. L. and D. Jenkins. 1971. The viability and activity of activated sludge. Water Res. 5: 621.
- Weiler, C. S. and D. M. Karl. 1979. Diel changes in phase-dividing cultures of *Ceratium Furca* (Dinophyceae): Nucleotide triphosphates, adenylate energy charge, cell carbon, and patterns of vertical migration. J. Phycol. 15: 384-391.
- Weiss, B., R. Lehne and S. Strada. 1972. Rapid micro assay of adenosine 3', 5' monophosphate phosphodiesterase activity. Anal. Biochem. 45: 22235.
- Welsch, F. and L. Smith. 1969. Kinetics of synthesis and utilization of ATP by intact cells of *Rhodospirillum rubrum*. Biochemistry 8: 3403-3411.
- Wettermark, G., E. Borglund and S. E. Brolin. 1968. A regenerating system for studies of phosphoryl transfer from ATP. Anal. Biochem. 22: 211-218.
- Wettermark, G., H. Stymne, S. E. Brolin and B. Peterson. 1975. Substrate analyses in single cells. I. Determination of ATP. Anal. Biochem. 63: 293-307.
- Wettermark, G., L. Tegner, S. E. Brolin and E. Borglund. 1970. Photokinetic measurement of ATP and ADP levels in isolated islets of Langerhans, in the structure and metabolism of the Panchreactic Islets (Falkner, S., B. Hellman and I. B. Taljedal eds.) Wenner-Gren Symposium No. 16, p. 275-282.
- White, E. H., F. McCapra and W. D. McElroy. 1961. The structure and synthesis of firefly luciferin. Am. Chem. Soc. J. 83: 2402-2403.
- White, E. H., F. McCapra and G. F. Field. 1963. The structure and synthesis of firefly luciferin. Am. Chem. Soc. J. 85: 337-343.
- White, E. H., E. Rapport, T. A. Hopkins and H. H. Seliger. 1969. Chemi-and bioluminescence of firefly luciferin. J. Am. Chem. Soc. 91: 2178-2180.

- White, E. H., E. Rapport, H. H. Seliger and T. A. Hopkins. 1971. The chemi- and bioluminescence of firefly luciferin: An efficient chemical production of electronically excited states. *Bioorg. Chem.* 1: 92-122.
- Whittam, R. 1958. Potassium movements and ATP in human red cells. *J. Physiol.* 140: 479-497.
- Wiebe, W. J. and K. Bancroft. 1975. Use of the adenylate energy charge ratio to measure growth state to natural microbial communities. *Proc. Nat. Acad. Sci. US*, 72: 2112-2115.
- Wiener, S., R. Wiener, M. Urivetzky and E. Meilman. 1974. Coprecipitation of ATP with potassium perchlorate: the effect on the firefly enzyme assay of ATP in tissue and blood. *Anal. Biochem.* 59: 489-500.
- Wijsman, T. C. M. 1976. ATP content and mortality in *Mytilus edulis* from different habitats in relation to anaerobiosis. *Neth. J. Sea Res.* 10: 140-148.
- Wildish, D. J. 1976. Determination of Adenosine 5'- triphosphate in estuarine water and sediments by firefly bioluminescence assay. *Tech. Rep. Fish. Mar. Serv. Res. Dev.* 649: 49 pp.
- Williams, J. D. H., J. K. Syers, R. F. Harris and D. E. Armstrong. 1970. Adsorption and desorption of inorganic phosphorus by lake sediments in a 0.1 N NaCl system. *Environ. Sci. Technol.* 4: 517-519.
- Williams, M. L. B. 1971. The limitations of the DuPont luminescence biometer in the microbiological analysis of food. *Can. Inst. Food Technol. J.* 4: 187-189.
- Williams, S. G. 1970. Role of phytic acid in the wheat grain. *Plant Physiol.* 45: 376-381.
- Windisch, R. M., P. R. Pax and M. M. Bracken. 1970. Variation in blood ATP after oral administration of glucose in individuals diagnosed as normal equivocal or diabetic according to the glucose tolerance sum principle. *Clin. Chem.* 16: 941-944.
- Wise, R. H. 1974. "Off-the-shelf" analyzers for measuring adenosine triphosphate (ATP) in activated sludge. EPA-670/2 74-034, National Environmental Research Centre, U. S. Environmental Protection Agency, Cincinnati, Ohio, 45268.
- Witkamp, M. 1973. Compatibility of microbial measurements. *Bull. Ecol. Res. Comm. (Stockholm)* 17: 179-188.
- Witteveen, S. A. D. J., B. E. Sobel and M. DeLuca. 1974. Kinetic properties of the isoenzyme of human creatine phosphokinase. *Proc. Natl. Acad. Sci. U.S.* 71: 1384-1387.
- Wolf, P. L., T. Kearns, J. Neuhoff and J. Lauridson. 1974. Identification of CPK isoenzyme MB in myocardial infarction. *Lab. Med.* 5.

- Wood, L. and E. Beutler. 1974. The effect of ascorbate and dihydroxyacetone on the 2, 3-diphosphoglycerate and ATP levels of stored human red cells. *Transfusion* 14: 272-277.
- Yamada, S. 1971. The effect of adenosine triphosphate (ATP) contents of human erythrocytes on immune hemolysis. *J. Nara Med. Assoc.* 22: 28-35.
- Yamaguchi, Y. and H. Seki. 1977. Microbial biomass in the eutrophic Bay of Shimoda as compared by estimates of several biomass parameters. *J. Ocean. Soc. Japan* 33: 38-44.
- Yousten, A. A. and R. S. Hanson. Sporulation of tricarboxylic acid cycle mutants of *Bacillus subtilis*. *J. Bacteriol.* 109: 886-894.
- Zwetnow, N. N. 1970. The influence of an increased intracranial pressure on the lactate, pyruvate, bicarbonate, phosphocreatine, ATP, ADP and AMP concentrations of the cerebral cortex of dogs. *Acta Physiol. Scand.* 79: 158-166.
- Zilinsky, J. W., G. A. Sojka and H. Gest. 1971. Energy charge regulation in photosynthetic bacteria. *Biochem. Biophys. Res. Commun.* 42: 955-961.