



Original Article

Temperature-dependent effects on fecundity in a serial broadcast spawning fish after whole-life high CO₂ exposure

Callie A. Concannon¹, Emma L. Cross², Lucas F. Jones¹, Christopher S. Murray³, Catherine M. Matassa¹, Richard S. McBride⁴, and Hannes Baumann^{1,*}

¹Department of Marine Sciences, University of Connecticut, 1080 Shennecossett Road, Groton, CT, 06340, USA

²Department of the Environment, Geography, and Marine Sciences, Southern Connecticut State University, 501 Crescent Street, New Haven, CT, 06515, USA

³Washington Ocean Acidification Center, School of Marine and Environmental Affairs, University of Washington, G 3707 Brooklyn Ave NE, Seattle, WA, 98105, USA

⁴National Marine Fisheries Service, Northeast Fisheries Science Center, 166 Water Street, Woods Hole, MA, 02543, USA

*Corresponding author: tel: 860-405-9297; e-mail: hannes.baumann@uconn.edu

Concannon, C. A., Cross, E. L., Jones, L. F., Murray, C. S., Matassa, C., McBride, R. S., and Baumann, H. Temperature-dependent effects on fecundity in a serial broadcast spawning fish after whole-life high CO₂ exposure. – ICES Journal of Marine Science, 78: 3724–3734.

Received 9 September 2021; revised 18 October 2021; accepted 19 October 2021; advance access publication 2 November 2021.

Experiments examining fish sensitivities to future oceanic CO₂ levels have greatly expanded over past decades and identified many potentially affected traits. Curiously, data on reproductive trait responses to high CO₂ are still scarce, despite their strong link to Darwinian fitness and thus to population vulnerability to ocean acidification. We conducted two rearing experiments on the first broadcast-spawning marine fish model (Atlantic silverside, *Menidia menidia*) to examine how long-term and novel whole life-cycle exposures to predicted future CO₂ levels (~2,000 μatm) affect laboratory spawning, temperature-specific reproductive investment, fecundity, and size distributions of maturing oocytes. At low temperatures (17°C), female body size and therefore potential fecundity (F_{Pot}, oocytes/female) slightly increased with CO₂, while relative fecundity (F_{Rel}, oocytes/g female) remained unaffected. At high temperatures (24°C), high CO₂ substantially reduced both F_{Pot} (–19%) and F_{Rel} (–28%) relative to control treatments. Irrespective of CO₂, females at 24°C grew larger and heavier than those at 17°C, and although larger females produced larger oocytes at some developmental stages, they also had lower gonadosomatic indices and lower F_{Rel}. Our findings contrast with most previous studies and thus highlight the need to investigate reproductive impacts of increasing CO₂ on multiple fish species with contrasting life history strategies.

Keywords: asynchronous batch spawner, Atlantic silverside (*Menidia menidia*), fitness, ocean acidification, temperature.

Introduction

The threat of unfolding marine climate change has motivated rapidly expanding research on the CO₂ sensitivity of marine organisms in recent decades (Boyd *et al.*, 2018; Doney *et al.*, 2020). Most studies have used experimental approaches to reveal important, first-order insights into species and traits affected by predicted future CO₂ levels and into mechanisms causing such effects (Baumann, 2019). The rich amount of empirical data now contains a complex spectrum of negative, neutral, and positive responses to

acidification within as well as across traits and species (Kroeker *et al.*, 2013; Przeslawski *et al.*, 2015), which cumulatively suggest a shifting marine fitness landscape but still limit more specific predictions. Experiments on marine fishes have shown that short-term high CO₂ exposures can reduce survival in some species, but also that such effects are generally confined to the earliest, least developed life stages (Chambers *et al.*, 2014; Baumann *et al.*, 2018; Cattano *et al.*, 2018; Murray *et al.*, 2019). Most other documented responses to high CO₂ in fishes comprise non-lethal changes to e.g. metabolism (Munday *et al.*, 2009), growth (Murray and Baumann,

2020), behaviour (Ashur *et al.*, 2017), or otolith formation (Bignami *et al.*, 2013), which arise as by-products of efficient acid-base regulation and swift CO₂ acclimation (Esbaugh, 2018). While potentially important, the challenge for many of these CO₂ responses often lies in drawing unequivocal links to Darwinian fitness, which is what matters most, because it measures the success of an individual in contributing to the next generation.

Curiously, very few studies to date have examined fecundity or reproductive output in fishes under future CO₂ environments, even though reproductive traits are closely linked to Darwinian fitness and thus to overall species vulnerabilities to ocean acidification (Nagelkerken *et al.*, 2021). Such experiments are clearly more complex, require longer durations, or face challenges in identifying appropriate fish models. However, their scarcity has left a knowledge gap preventing better assessments of ocean acidification as an emerging threat for marine fishes. Pioneering work has exclusively tested fish species with parental care, where experimenters could select adults or sub-adults from the wild to then quantify frequency and sizes of egg clutches laid by breeding pairs under contrasting CO₂ conditions (Miller *et al.*, 2013; Welch and Munday, 2016; Faria *et al.*, 2018). No comparable data exist yet for broadcast spawners, a group that includes most commercial fishes. Moreover, no marine fish model has yet been reared from fertilization to maturity to examine how realistic whole-life CO₂ exposures may affect reproductive output.

To address this knowledge gap, we conducted two long-term rearing experiments on the Atlantic silverside (*Menidia menidia*, hereafter: silverside), a broadcast-spawning forage fish along the North-American Atlantic coast (Middaugh *et al.*, 1987) and well-studied model in evolutionary, ecological, and climate change research (e.g. Conover and Kynard, 1981; Billerbeck *et al.*, 2000; Conover *et al.*, 2009; Cross *et al.*, 2019). Silversides are particularly amenable to whole-life studies because of their relatively short, annual life span and the existence of established laboratory protocols for whole-life rearing (Conover and Munch, 2002). Furthermore, silversides affix fertilized eggs to submersed vegetation in the wild (or to artificial substrates in the laboratory), which facilitates off-spring collection and quantification. However, known challenges to whole-life rearing include their obligate schooling behaviour and batch spawning, which demand large rearing tanks (>500 L) and associated space-replication trade-offs. We therefore used two complementary approaches over two experimental seasons to quantify reproductive output under contrasting CO₂ environments in silversides. The first approach (hereafter: *spawning trial*) reared wild-caught early juveniles to maturation and then quantified embryo production in laboratory populations over a 2-month period. The second approach (hereafter: *fecundity trial*) reared silversides from fertilization to just prior to spawning, in order to quantify individual female reproductive investment, fecundity, and oocyte development via macroscopic and histological analyses. We hypothesized that long-term or whole-life exposures to future CO₂ conditions would have small but accruing metabolic costs, which would result in lower reproductive investment, lower fecundity, and/or smaller oocytes in silversides reared at high compared to control CO₂ conditions.

Material and methods

Experimental CO₂ and temperature treatments

The spawning trial reared wild-caught juveniles to maturity at 20°C under duplicate control (~600 µatm CO₂) and high CO₂ condi-

tions (~2200 µatm CO₂). The fecundity trial reared newly fertilized embryos to maturity under control vs. high CO₂ conditions and two temperatures (17°C: ~500, ~2000 µatm CO₂; 24°C: ~500, ~2600 µatm CO₂; Table 1). High CO₂ conditions were chosen to approximate the predicted surface ocean maximum in coming centuries (Caldeira and Wickett, 2003) that is a common benchmark in ocean acidification studies (Riebesell *et al.*, 2011). However, comparably high CO₂ levels already occur seasonally (late summer–fall) in nearshore silverside habitats (Baumann *et al.*, 2015) and may become more frequent and extreme in future decades (Shaw *et al.*, 2013; McNeil and Sasse, 2016). Temperature treatments were consistent with past experimental work, representing average thermal conditions at the beginning (17°C), height (20°C), and end (24°C) of the species' spawning season in Long Island Sound (Baumann *et al.*, 2015; Murray and Baumann, 2018), with 24°C being the thermal optimum for larval and juvenile growth (Middaugh *et al.*, 1987).

Experimental treatment tanks were continuously bubbled with mixes of air and 100% bone-dry CO₂ controlled by gas proportioners (ColeParmer®). In control CO₂ treatments, metabolic acidification by the fish was offset by bubbling CO₂-stripped air (~50 µatm CO₂) via diffuser tubes into the rearing tanks (Murray and Baumann, 2020). All treatments maintained near 100% dissolved oxygen saturations (~8.1, 7.6, and 7.1 mg L⁻¹ at 17°C, 20°C, and 24°C, respectively). Salinity was monitored daily via refractometer and adjusted as necessary to maintain a target of 31 psu. Temperature conditions were maintained by thermostats (Aqualogic®) connected to submersible heaters or in-line chillers (DeltaStar®), and target pH conditions were monitored daily using a handheld pH meter (Hach® Intellical PHC281 pH electrode with HQ11D handheld pH/ORP meter, calibrated bi-weekly using two-point NIST buffers). Achieved pCO₂ levels were calculated via CO₂SYS (v2.1) from total alkalinity (A_T) measurements in samples of tank and lab source water (N_{Spawn} = 12, N_{Fecund} = 17) measured via endpoint titration (G20 Potentiometric Titrator, Mettler Toledo®) with an accuracy of ± 1% (Dr Andrew Dickson's certified A_T reference, batch number 162). Using A_T, pH, temperature, salinity, and K₂ constants from Mehrbach *et al.* (1973) refitted by Dickson and Millero (1987) and Dickson (1990), we derived CO₂ partial pressure (pCO₂, µatm), dissolved inorganic carbon (C_T, µmol kg⁻¹), and carbonate ion concentrations (CO₃²⁻, µmol kg⁻¹) given in Table 1.

Offspring production and laboratory rearing

For the spawning trial, juveniles were caught on 25 September 2015 via beach seine (30 × 2 m) in Mumford Cove, a small seagrass-dominated embayment connected to eastern Long Island Sound (41.32°N, 72.02°W). Individuals were transported to the Rankin Seawater Facility at the University of Connecticut (Groton, CT, USA). A total of 100 individuals (mean total length [TL] = 4.5 cm) were randomly assigned to each of two 700-L replicate rearing tanks per CO₂ treatment and reared to maturity over the next five months at 20°C (N_{total} = 400). Maturation was achieved via photoperiod manipulation, i.e. by reducing daylight from 15 h to 12 h during months 3–4 followed by a reversal back to 15 h for the final (fifth) month (Conover and Munch, 2002; Salinas *et al.*, 2012). Fish were fed equal amounts of pelleted food (Otohime Marine Fish Diet, size B1, B2, C1, Reed Mariculture®), delivered continuously to each tank during light hours via belt feeders.

For the fecundity trial, spawning-capable silversides were captured in Mumford Cove on 29 May 2018, transported to the laboratory, separated by sex, and held overnight without food in

Table 1. Overview of experimental water chemistry during the spawning and fecundity trials, using measured temperature ($^{\circ}\text{C}$), pH (NIST), and total alkalinity (A_t , $\mu\text{mol kg}^{-1}$) to calculate achieved mean \pm SD of pCO_2 (μatm), dissolved inorganic carbon (C_t , $\mu\text{mol kg}^{-1}$), and carbonate ion concentrations (CO_3^{2-} , $\mu\text{mol kg}^{-1}$).

	CO_2	Temp	pH	A_t	pCO_2	C_t	CO_3^{2-}
Spawning trial	Control CO_2	20 ± 0.5	7.99 ± 0.08	$2\,035 \pm 74$	614 ± 147	$1\,893 \pm 90$	108 ± 12.2
			7.98 ± 0.04	$2\,044 \pm 49$	620 ± 87	$1\,905 \pm 59$	107 ± 5.9
	High CO_2		7.46 ± 0.02	$2\,030 \pm 57$	$2\,252 \pm 161$	$2\,050 \pm 60$	35 ± 1.0
			7.47 ± 0.04	$2\,054 \pm 70$	$2\,217 \pm 295$	$2\,071 \pm 80$	37 ± 1.2
Fecundity trial	Control CO_2	16.9 ± 0.6	8.07 ± 0.11	$2\,023 \pm 9$	489 ± 136	$1\,869 \pm 44$	115 ± 26
		23.7 ± 1.2	8.06 ± 0.13		538 ± 178	$1\,839 \pm 58$	135 ± 35
	High CO_2	16.9 ± 0.9	7.50 ± 0.09		$2\,024 \pm 445$	$2\,045 \pm 25$	34 ± 6
		23.6 ± 1.2	7.42 ± 0.12		$2\,628 \pm 882$	$2\,048 \pm 39$	36 ± 9

80-L aerated containers at ambient CO_2 and temperature conditions (17°C). The following day, 20 females and 27 males were strip-spawned together using established protocols (Baumann *et al.*, 2018; Murray and Baumann, 2020). Within 2 hours post fertilization, screens with attached embryos were randomly placed into 32 rearing containers (20-L, with flow-through screening), which were then distributed and placed within the four 700-L rearing tanks ($n = \text{eight containers/tank}$). Newly hatched larvae were fed *ad libitum* amounts of brine shrimp nauplii (*Artemia salina*, San Francisco strain, brineshrimpdirect.com) augmented initially by a powdered weaning diet (Otohime Marine Fish Diet, size A1).

After reaching approximately 30 mm TL (45 days post hatch, dph, for 24°C , 73 dph for 17°C), fish were counted, measured ($n = 10$), and divided randomly among three 80-L rearing containers (with flow-through screening) placed back into their original 700-L tank and corresponding CO_2 and temperature treatment. At this point, each of the two 24°C tanks contained 148 juveniles, while each of the two 17°C tanks contained 96 individuals. At 104 dph (17°C) and 111 dph (24°C), juveniles were counted and measured again then placed directly into their original 700-L tank and corresponding treatment. Fish were fed daily rations of pelleted food via belt feeders (Otohime Marine Fish Diet, size B1, B2, C1) corresponding to 20% of their estimated dry weight biomass per tank (dW_{all}). We readjusted dW_{all} daily using published temperature-specific TL growth data for silversides (Billbeck *et al.*, 2000) and the empirical relationship $\text{dW} = 0.0012 * \text{TL}^{2.997}$ (Murray *et al.*, 2017), multiplied by the number of fish in each tank. To avoid overcrowding as fish grew, at 245 dph half of the fish in each of the two 24°C tanks were randomly selected and moved into two new tanks receiving the corresponding temperature (24°C) and CO_2 treatments. Biological filtration (four-stage canister biofilters with 9-watt ultraviolet [UV] sterilizers, Polar Aurora[®]), regular solid waste removal, and weekly water changes with filtered (1 μm) and UV-sterilized seawater from Long Island Sound ensured undetectable ammonia concentrations throughout the experiment (<0.25 ppm, tested $3 \times$ weekly; Saltwater Master Test Kit, API[®]). A photoperiod of 15 h light, 9 h dark was maintained throughout the embryo, larval, and juvenile stages. To induce maturation, the photoperiod was reduced to 12 h light for 2 months beginning 245 dph, followed by reversal back to 15 h light until sampling. The fecundity trial was terminated before spawning occurred at 317–325 dph, when all fish were euthanized with MS-222 (Tricaine-S).

Trait measurements

For the spawning trial, spawning in each tank began when silversides were provided bundled threads of green yarn suspended mid-tank. Yarns were checked every 48 h for attached embryos (i.e. water hardened eggs), which were counted and subsampled ($n = 15$) to measure egg diameters via calibrated pictures ($4 \times$ magnification, ImagePro Premier, v9.0, Media Cybernetics[®]). The spawning trial was terminated after 60 days, which is approximately the extent of silverside spawning season in the wild (Pringle and Baumann, 2019). Surviving fish were euthanized, measured for TL (0.1 mm) and wet weight (wW, 0.01g), and their sex determined by gonad inspection under a stereo-microscope (Nikon SMZ-1000). The total number of embryos spawned was scaled to the cumulative wet weight of females in each tank to obtain a mass-specific estimate of embryo production ($n_{\text{emb}}/\text{g female}$) per replicate tank. This necessarily assumed that all females within a tank participated in spawning.

For the fecundity trial, euthanized fish were measured for TL and wW, followed by dissection for sex identification and gonad removal. Re-weighing yielded the gonad-free body weight ($\text{wW}_{\text{no gonad}}$), which was subtracted from wW to yield gonad weight (wW_{gonad}) and the gonado-somatic index ($\text{GSI} = 100 * \text{wW}_{\text{gonad}} / \text{wW}_{\text{no gonad}}$). Male gonads were discarded. Female gonads were preserved in 5% buffered formaldehyde/freshwater solution. One ovary lobe per female was used to quantify all secondary growth (i.e. vitellogenic) oocytes; the other was saved for histological analyses. To characterize secondary growth oocytes, the ovary lobe was divided into anterior, middle, and posterior sections that were each gently teased apart so that oocytes were evenly distributed among four rectangular 2 mL wells. Each well was photographed ($3 \times$ magnification) and analyzed via ImageJ (v1.52a, National Institute of Health) with the ObjectJ plugin (v1.04t, University of Amsterdam) and a customized macro to count and measure the diameter of all oocytes. The total number of oocytes counted per lobe was doubled to obtain each female's potential fecundity (F_{pot} , oocytes/female) and relative fecundity ($F_{\text{rel}} = F_{\text{pot}}/\text{wW}$, oocytes/g female). In addition, we averaged the size of the 25 largest oocytes per female ($\sim 99\%$ percentile) as a measure of maximum oocyte size.

The other ovary lobe from a subset of 17 females at control CO_2 treatments ($n_{17^{\circ}\text{C}} = 6$, $n_{24^{\circ}\text{C}} = 11$) and 20 females at high CO_2 treatments ($n_{17^{\circ}\text{C}} = 8$, $n_{24^{\circ}\text{C}} = 12$) were embedded in paraffin, sectioned, placed on glass slides, and stained with Hematoxylin & Eosin (H&E) (Horus Scientific, Worcester, MA). Slides were then

Table 2. *Menidia menidia*: mean ± SD of response traits during the spawning trial (2015–2016) and the fecundity trial (2018–2019).

CO ₂ treatment Temperature (°C)	Spawning trial		Fecundity trial			
	Control 20	High	Control 17	High	Control 24	High
N _{females}	53	65	40	17	11	17
TL (mm)	119.3 ± 9.9	115.5 ± 9.4	88.6 ± 8.2	94.2 ± 9.1	111.4 ± 6.7	115.0 ± 5.8
wW (g)	9.3 ± 2.1	8.7 ± 1.8	4.2 ± 1.1	5.4 ± 1.3	8.1 ± 1.4	9.1 ± 1.4
N _{embryos/fem}	550, 541*	174, 571*				
N _{embryos/g fem}	0.8, 2.5*	2.3, 2.3*	Not assessed			
Embryo diameter (µm)	1 131 ± 54	1 126 ± 50				
GSI (%)			18.9 ± 8.1	16.1 ± 5.4	8.1 ± 3.5	8.2 ± 1.8
Potential fecundity (oocytes/fem)			3 262 ± 1 348	4 355 ± 1 338	3 260 ± 747	2 631 ± 675
Relative fecundity (oocytes/g fem)			762 ± 221	807 ± 164	404 ± 53	291 ± 66
Maximum oocyte diameter (µm)	Not assessed		1 152 ± 139	1 113 ± 156	1 047 ± 219	917 ± 136
Median oocyte diameter (PG, CA, V1, V2, NM, H in µm)**			132, 289, 397, 577, 648, 804		135, 326, 435, 577, 601, 932	

*values given for each duplicate;

**values pooled across CO₂ treatments

photographed (4 × magnification) and six oocyte developmental stages identified: primary growth (PG), cortical alveolar (CA), early vitellogenesis (V1), late vitellogenesis (V2), nuclear migration (NM), and hydrating (H) oocytes (following Ganas *et al.*, 2004; Hyle *et al.*, 2014; Press *et al.*, 2014). We also checked for but did not find postovulatory follicles that would have indicated the beginning of spawning. Furthermore, we examined all histological images for the occurrence of alpha or beta atresia, following McElroy *et al.* (2016). Given that the incidence of atresia was not significantly different from zero (%) and also not significantly different between treatments, no adjustment to whole oocyte counts was necessary (see electronic Supplementary Material). Up to 56 oocytes per developmental stage per female were measured (mean = 13.5). Histological preparation deformed most oocytes to ellipses; therefore, we first calculated the area of each ellipse from its major and minor axis and then inferred the oocyte diameter (= size) from a circle with the same area. For each female, we determined the stage-specific median and maximum oocyte size, which were then compared across treatments.

Data analysis

Rearing large numbers of schooling fish to maturity is space demanding and, in a facility like ours, demanded experimental design trade-offs with respect to replication. With the replication unit being each 700-L tank (instead of each female), the spawning trial thus had duplicates for each CO₂ level (single temperature), while the fecundity trial had single replicates for each CO₂ × temperature combination. Although the 24°C treatments ended with two equal density groups per CO₂ level, these were not true replicates, because fish were only separated after 245 dph. Given that these duplicates were not significantly different from one another with respect to mean TL and wW (t-tests, both $p > 0.21$), they were pooled for all subsequent analyses. Lacking the treatment-level replication and normal distributions required by parametric statistical approaches (General Linear Model (GLM) and analysis of variance), we instead compared trait distributions between CO₂ and temperature treatments using non-parametric Mann–Whitney U tests (reporting Z-scores) with adjusted p -values for multiple comparisons during each trial (Benjamini and Hochberg, 1995). We focused on female traits because

reproductive investment is more costly and more tightly linked to body size and fitness for females than for males (Wootton, 1985). We examined female TL and wW (spawning + fecundity trials), number and size of spawned embryos (spawning trial), GSI, F_{Pot} , F_{Rel} , maximum oocyte size, as well as stage-specific oocyte sizes (fecundity trial). We also examined the relationships between F_{Pot} and wW using linear regression and compared CO₂ effects on these relationships with analysis of covariance (ANCOVA). Three females were removed from analyses of reproductive trait data because they had either no discernible or only immature oocytes in their gonads. Male trait distributions are presented as supplementary material (Supplementary Figure S1). Statistical analyses were performed in SPSS (V20, IBM).

Results

Spawning trial

Female TL and wW did not differ significantly between duplicates within each CO₂ treatment (control: $Z_{\text{TL}} = 0.15$, $Z_{\text{wW}} = -0.28$, $p > 0.87$; high CO₂: $Z_{\text{TL}} = 1.33$, $Z_{\text{wW}} = 1.30$, $p > 0.35$) or between CO₂ treatments when duplicate tanks were pooled ($Z_{\text{TL}} = 1.81$, $p = 0.35$; $Z_{\text{wW}} = 1.42$, $p = 0.35$, Table 2, Figure 1a and b). Each of the four laboratory populations continually spawned throughout the 60-day period, producing 0.1–3.6 embryos g-female⁻¹ during most days, interspersed by several large spawning peaks of 4.8–10.6 embryos g-female⁻¹ in three of the four populations (Figure 2a). One high CO₂ replicate did not produce any spawning peaks and thus produced fewer total embryos (174 per female) than the other high CO₂ replicate and both control CO₂ replicates (541–571 per female, Figure 2b, Table 2). The peaks in embryo production showed no apparent periodicity within or synchronicity across tanks. Overall, egg diameter did not differ between duplicates ($Z_{\text{con}} = -1.67$, $Z_{\text{high}} = 0.93$, $p > 0.35$), nor when pooled, between CO₂ treatments ($Z = -0.47$, $p = 0.82$). Mean (±SD) egg diameter across treatments was 1129 ± 52 µm (n = 416).

Fecundity trial

In both CO₂ treatments, female TL and wW increased with temperature ($Z_{\text{TL, con}} = 4.96$, $Z_{\text{TL, High}} = 4.71$, $Z_{\text{wW, con}} = 4.82$,




Figure 1. *M. menidia*. Adult female total length (a, c) and wet weight (b, d) distributions after the conclusion of the spawning and fecundity trials at control and high CO₂ conditions at 20°C (a, b—green), 17°C (c, d—blue) and 24°C (c, d—red). Distributions in (a) and (b) were pooled across statistically similar duplicates. Circles and error bars depict treatment-specific means \pm 1 SD. Asterisks denote statistically significant differences between adjacent treatment groups (Mann–Whitney U tests, $p < 0.05$).

$Z_{wW, high} = 4.75$, all $p < 0.001$; Figure 1c–d, Table 2). At 17°C, TL and wW increased with CO₂ ($Z_{TL, 17^\circ C} = 2.45$, $p = 0.024$; $Z_{wW, 17^\circ C} = 3.33$, $p = 0.002$), while CO₂ had no effect on body size at 24°C ($Z_{TL, 24^\circ C} = 1.44$, $p = 0.23$; $Z_{wW, 24^\circ C} = 1.55$, $p = 0.18$). GSI decreased with temperature at each CO₂ level ($Z_{con} = -3.97$; Z_{high}

$= -4.48$, both $p < 0.001$, Table 2, Figure 3a), but CO₂ had no effect on GSI at either temperature ($Z_{17^\circ C} = -1.34$, $p = 0.24$; $Z_{24^\circ C} = 0.31$, $p = 0.82$).

Distributions of F_{Pot} (oocytes/female) and F_{Rel} (oocytes/g female) suggested interacting temperature and CO₂ effects on fe-




Figure 2. *M. menidia*. Spawning trial: Time-series of embryo production in duplicate laboratory populations (R1, R2) spawning over a 2-month period at control (orange lines, circles and squares) and high CO₂ conditions (green lines, triangles and diamonds), with (a) embryo production per g female and (b) cumulative embryo production per female. Numbers denote females in each duplicate.

male fecundity (Figure 3b and c). At 17°C, high CO₂ increased F_{Pot} ($Z_{17^\circ C} = 2.69$, $p = 0.013$) but had no effect on F_{Rel} ($Z_{17^\circ C} = 0.51$, $p = 0.67$). At 24°C, however, both F_{Pot} and F_{Rel} distributions shifted lower in response to high CO₂ conditions (F_{Pot} : $Z_{24^\circ C} = -2.07$, $p = 0.059$; F_{Rel} : $Z_{24^\circ C} = -3.65$, $p < 0.001$), with average reductions of -19% and -28%, respectively. Warmer temperature negatively affected F_{Pot} at high CO₂ only ($Z_{high} = -4.01$, $p < 0.001$; $Z_{con} = -0.02$, $p = 0.98$; Figure 3b) but reduced F_{Rel} regardless of CO₂ level ($Z_{con} = -4.49$; $Z_{high} = -4.98$, both $p < 0.001$, Figure 3c).

At 17°C F_{Pot} increased linearly with female wW (Linear regression: $F_{Pot} = 871 * wW - 386$, $r^2 = 0.61$, $p < 0.001$), and this relationship did not differ between CO₂ treatments (ANCOVA: CO₂ × wW, $F_{1,56} = 0.015$, $p = 0.9$, Figure 4). At 24°C, however, F_{Pot} increased linearly with wW at only control CO₂ ($F_{Pot} = 413 * wW - 75$, $r^2 = 0.64$, $p = 0.003$) but not at high CO₂ levels (regression, $p = 0.094$).

Maximum oocyte size distributions were CO₂ independent at each temperature ($Z_{17^\circ C} = -0.67$; $Z_{24^\circ C} = -1.11$, $p > 0.34$), but at high CO₂ treatments they shifted lower at 24°C relative to 17°C ($Z_{high} = -3.13$, $p = 0.002$, Figure 3d). Distributions of temperature- and stage-specific median and maximum oocyte sizes did not differ between CO₂ treatments ($Z_{PG,CA,V1,V2,NM,H} = -1.48$ – 1.73 , $p > 0.26$). When pooled, median and maximum oocyte size distributions increased with temperature for some early developmental

stages (PG_{max}: $Z = 3.13$, $p = 0.002$; CA_{med}: $Z = 4.04$, $p < 0.001$; V1_{med}: $Z = 3.13$; $p = 0.002$) and the final, hydrating stage (H_{med,max}: $Z = 2.34$, 2.49 ; $p = 0.013$, $p = 0.019$). However, the latter is more uncertain, given the low sample size and the observation that many hydrating oocytes in the 17°C treatments collapsed during histological preparation and were therefore not measurable. The same developmental stages (i.e. CA, V1, and H, but not PG, V2, NM) increased in size with wW (all $p < 0.015$); therefore, larger females had larger, more advanced oocyte stages across temperature and CO₂ levels at the end of the experiment (Figure 5).

Discussion

Two complementary rearing experiments examined how long-term and whole life-cycle exposures to future oceanic CO₂ conditions affect female reproductive traits in a temperate marine fish model, the Atlantic silverside. We hypothesized that high CO₂ environments would demand additional metabolic costs due to increased acid-base regulation (Munday *et al.*, 2009; Heuer and Grosell, 2016), thereby diminishing resources for energetically demanding oocyte production, maturation, and spawning (McBride *et al.*, 2015). Our findings partially supported this hypothesis, because potential and relative fecundities shifted lower in response to whole-life, high CO₂ exposure at warm temperatures. At cold temperatures, rela-




Figure 3. *M. menidia*. Fecundity trial: temperature- and CO₂-specific distributions of (a) gonado-somatic index, (b) potential fecundity, (c) relative fecundity, and (d) maximum oocyte size in spawning-capable females. Blue colors and circles depict 17°C, red colors and diamonds depict 24°C treatments. Symbols and error bars depict treatment-specific means \pm 1 SD. Asterisks denote statistically significant differences between adjacent treatment groups (Mann-Whitney U tests, $p < 0.05$).

tive fecundity was unaffected, while body size and thus potential fecundity slightly increased with CO₂. Other assessed reproductive traits were sensitive to temperature but not CO₂.

Temperature-dependent CO₂ effects are common in ocean acidification experiments (Munday *et al.*, 2009; Harvey *et al.*, 2013; Przeslawski *et al.*, 2015; Murray and Baumann, 2018), presumably

because ectotherms tolerate additional stressors better when near their thermal optimum than when at the extremes of their thermal performance curves (Pörtner, 2008). Yet for silversides, the thermal optimum is often assumed to be around 24°C (Middaugh *et al.*, 1987), whereas 17°C is near their lower thermal limit, which at first seems inconsistent with our results. However, these ther-




Figure 4. *M. menidia*. Fecundity trial: temperature- and CO₂-specific relationships between female wet weight and potential fecundity. Relationships at 17°C were CO₂-independent; the pooled relationship is plotted ($p < 0.001$). At 24°C, the relationship was significant for the control CO₂ treatment ($p = 0.003$), but not for the high CO₂ treatment ($p = 0.094$). Shading depicts 95% confidence intervals from linear regressions.

mal optima pertain to somatic growth, not reproductive traits, and they have largely been derived from rearing larvae and juveniles (Conover and Present, 1990), which are adapted to developing in warm nearshore waters during summer and fall (Conover and Ross, 1982). Indeed, our growth data confirmed that silversides grew 24% larger and 79% heavier at 24°C than at 17°C, consistent with expectations based on thermal growth reaction norms. But sub-adults in the wild mature at much colder temperatures while overwintering offshore (Conover and Murawski, 1982), after which they return to nearshore waters and commence spawning as temperatures reach approximately 15°C again (at ~41°N, Pringle and Baumann, 2019). This means that our cold temperature treatment of 17°C was actually closer to the natural thermal conditions for maturing silversides than the warm 24°C treatment, which may explain why the latter not only led to lower reproductive investment (lower GSI) but also to significantly lower potential and relative fecundities in the presence of the additional stressor, high CO₂.

Conversely, high CO₂ increased potential fecundity at 17°C, but only because body size increased with CO₂, and larger females simply produced more oocytes. Oocyte production per unit weight remained unchanged at 17°C, as indicated by the linear relationship between weight and potential fecundity, and confirmed by CO₂-independent relative fecundities. At 24°C, however, the relationship between weight and potential fecundity disappeared at high CO₂ conditions, body size was independent of CO₂ treatment, and both potential and relative fecundities declined at high compared to control CO₂ conditions—suggesting a size-independent CO₂ effect. We note that the spawning trial at an intermediate temperature




Figure 5. *M. menidia*. Fecundity trial: relationships between wet weight and stage-specific median oocyte size in spawning capable females. Each symbol represents an individual female, with circles and squares depicting females reared at control and high CO₂ levels, respectively.

(20°C) found neutral CO₂ effects on embryo production, which is consistent but circumstantial evidence, because the two trials are not directly comparable. We infer that high temperature combined with high CO₂ conditions led to increased metabolic demands that compromised available resources for oocyte maturation in laboratory silverside populations.

Our findings contrast with the still limited empirical evidence that has so far suggested mostly stimulating effects of increased CO₂ on fish reproductive output. Pioneering work quantified breeding in wild-caught cinnamon clownfish (*Amphiprion melanopus*) over a 9-month period, reporting twice as many pairs breeding, producing twice as many clutches per pair and 67% more eggs per clutch in elevated compared to control CO₂ treatments (~400 vs. ~1000 μatm, Miller *et al.*, 2013). A follow-up study confirmed a CO₂-mediated increase in egg clutches and eggs per clutch for the congener *Amphiprion percula*, although it also found contrasting, negative CO₂ effects on produced egg clutches in a different reef fish, *Acanthochromis polyacanthus* (Welch and Munday, 2016). Schade *et al.* (2014) acclimated groups of adult three-spined sticklebacks (*Gasterosteus aculeatus*) caught in the German Bight (North Sea) to control and high CO₂ conditions (~1000 μatm) for three months, after which strip-spawning revealed a 20% increase in eggs per female from control to high CO₂ levels, despite similar female body size. In two-spotted gobies (*Gobiusculus flavescens*), wild-caught breeding pairs held for 4 months under high CO₂ levels (~2300 μatm) produced 50% more clutches and 44% more eggs per clutch than pairs under control conditions, although there was a trade-

off with smaller hatch size (Faria *et al.*, 2018). In all these experiments, breeding pairs were provided high or *ad libitum* food rations, which might have allowed females to increase consumption to sustain their CO₂-stimulated reproductive output. This point was recently underscored by Nagelkerken *et al.* (2021), who measured prey availability and reproductive investment in several small benthic fishes (triple-fin gobies, blenny) inhabiting contrasting CO₂ environments at natural CO₂ vents off the coast of New Zealand. Over several years, the authors found increased benthic prey abundance at vent sites compared to control sites, which coincided with higher foraging activities and higher reproductive investment in both male and female fishes (Nagelkerken *et al.*, 2021).

Interestingly, whole-life silverside rearing resulted in neutral (24°C) to weakly positive CO₂ effects (17°C) on adult female body size. This seems inconsistent with the neutral to negative growth responses to high CO₂ documented by two previous studies, which reared larger numbers of silversides to late juvenile stages (to ~130 dph) at comparable CO₂ and temperature levels (Murray *et al.*, 2017; Murray and Baumann, 2020). Contrasting intraspecific and across-experiment outcomes are not uncommon in ocean acidification studies (Parker *et al.*, 2011), and serial experimentation is often required for robust quantification of trait responses (Baumann *et al.*, 2018). In silverside experiments, additional variability is likely introduced by using wild-caught parental founders that may vary in genetic composition within and between spawning seasons (Murray *et al.*, 2014; Malvezzi *et al.*, 2015). It is also possible that positive CO₂ effects only arise after the period of fast larval and juvenile somatic growth ends, and fish start diverting resources towards reproduction and thus grow slower. For silversides in the wild, this coincides with seasonal declines in temperature and food availability (Schultz *et al.*, 1998), whereas laboratory populations experienced static thermal conditions and unchanged ~20% food rations. These food rations might be limiting larval and juvenile growth and therefore produce CO₂-induced growth reductions, but in maturing sub-adults they might have provided sufficient excess resources to benefit preferentially the fish maturing at high CO₂ treatments (Supplementary Figure S2).

Over the past decade, a large number of studies systematically examined many Atlantic silverside traits for their climate sensitivity, using short- and long-term as well as single- and multi-stressor exposures to static and fluctuating regimes of CO₂, temperature, and oxygen (e.g. Murray *et al.*, 2014; Depasquale *et al.*, 2015; Malvezzi *et al.*, 2015; Miller *et al.*, 2016; Baumann *et al.*, 2018; Cross *et al.*, 2019; Murray and Baumann, 2020; Schwemmer *et al.*, 2020). Silversides are appealing models, given their ecological importance as forage fish, their almost year-round, nearshore availability to experimenters, their short life span, and the existence of established laboratory spawning and rearing protocols (Middaugh *et al.*, 1987). They thus were also obvious candidates here for studying reproductive CO₂ sensitivities. However, the species also exhibits life history characteristics that are difficult to overcome experimentally and which limit conclusions particularly with regards to the effects of environmental conditions on reproduction.

As an obligatory schooling species, silversides must be held in large groups (demanding large enclosures, replication trade-offs) for quantifiable, semi-natural spawning activity to occur. This can be insightful (spawning trial) but it precludes resolving individual female contributions. We first tried to overcome this by experimenting with small silverside groups (female + three males), but this failed to elicit quantifiable spawning. The alternative was to quantify maturing oocytes in individual female gonads prior to spawn-

ing (fecundity trial), which provided a useful measure of potential fecundity, but not necessarily of realized fecundity that is ultimately of most interest. For example, realized embryo production during the spawning trial (~600 embryos female⁻¹) was five to seven times lower than potential fecundities estimated during the fecundity trial, which may be due to method differences between trials, but could also indicate that a large number of vitellogenic oocytes are resorbed again after spawning. In total spawners, i.e. fish species that release all eggs in one spawning event, potential and realized fecundity are tightly correlated (McBride *et al.*, 2015), but silversides are batch spawners that mature and release egg batches in semi-lunar intervals over the course of their spawning season in the wild (Conover, 1985). Moreover, overall oocyte size-distributions (Supplementary Figure S3) and stage-specific oocyte measurements showed overlapping size ranges of successive developmental stages, therefore suggesting that silversides are indeterminate batch spawners with asynchronous oocyte development (Ganias *et al.*, 2015; Ganias and Lowerre-Barbieri, 2018), where the number of maturing batches cannot be readily deduced from distinguishable oocyte cohorts. Asynchronicity might partially be an artifact of laboratory populations lacking periodic environmental cues (temperature, tides, moon light), which is supported by the aperiodic and asynchronous spawning patterns observed during the spawning trial (Conover and Kynard, 1984). A final difficulty in conducting whole-life experiments with silverside populations from mid-latitudes is their high degree of temperature-dependent sex determination (Conover and Kynard, 1981; Murray and Baumann, 2020), which produces roughly balanced sex ratios at low temperatures (i.e. 17°C), but results in impractically small female proportions at warmer temperatures (at 24°C, only 1–2 out of 10 fish become females).

In conclusion, our two complementary approaches suggested neutral CO₂ effects on silverside reproductive traits at low temperatures, but CO₂-induced reproductive depression at warm temperatures, which currently occur only during summer and not during overwinter maturation. Many reproductive traits depended on body size, thereby emphasizing the fitness relevance of growth and the general importance of examining growth-determining variables in climate sensitivity studies of fish. The short, annual life cycle of this species may also allow evolutionary responses to future high CO₂ environments (Malvezzi *et al.*, 2015), thereby potentially offsetting any temperature × CO₂-related reductions in fecundity. Future work should aim for further in-depth reproductive analyses, enhanced replication and statistical power, as well as seasonally varying thermal regimes. However, conclusive insights will likely depend on developing an alternative, non-schooling fish model that is a total broadcast spawner equally amenable to experimental manipulation.

Ethics

The Institutional Animal Care and Use Committee (IACUC) of the University of Connecticut approved fish husbandry and experimental protocols (#A14-032, #A17-043).

Supplementary Data

Supplementary material is available at the ICESJMS online version of the manuscript.

Data availability

Citable source data are available from the BCO-DMO database: Spawning trial DOI: 10.26008/1912/bco-dmo.845633.1, 10.26008/1912/bco-dmo.845804.1; Fecundity trial DOI: 10.26008/1912/bco-dmo.845906.1, 10.26008/1912/bco-dmo.845921.1

Funding

This study was funded by the National Science Foundation grant (OCE#1536165) to H.B.

Acknowledgements

We are grateful to C. Woods, C. Dyke, I. Mayo, and R. Moy-Huwylar for their assistance with experimental setups, egg production counts, and measurements. E. Tholke, Y. Press, and D. McElroy facilitated image analyses.

References

- Ashur, M. M., Johnston, N. K., and Dixon, D. L. 2017. Impacts of ocean acidification on sensory function in marine organisms. *Integrative and Comparative Biology*, 57: 63–80.
- Baumann, H. 2019. Experimental assessments of marine species sensitivities to ocean acidification and co-stressors: how far have we come? *Canadian Journal of Zoology*, 97: 399–408.
- Baumann, H., Cross, E. L., and Murray, C. S. 2018. Robust quantification of fish early life CO₂ sensitivities via serial experimentation. *Biology Letters*, 14: 20180408.
- Baumann, H., Wallace, R., Tagliaferri, T., and Gobler, C. J. 2015. Large natural pH, CO₂ and O₂ fluctuations in a temperate tidal salt marsh on diel, seasonal and interannual time scales. *Estuaries and Coasts*, 38: 220–231.
- Benjamini, Y., and Hochberg, Y. 1995. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society: Series B (Methodological)*, 57: 289–300.
- Bignami, S., Enochs, I. C., Manzello, D. P., Sponaugle, S., and Cowen, R. K. 2013. Ocean acidification alters the otoliths of a pantropical fish species with implications for sensory function. *Proceedings of the National Academy of Sciences*, 110: 7366–7370.
- Billerbeck, J. M., Schultz, E. T., and Conover, D. O. 2000. Adaptive variation in energy acquisition and allocation among latitudinal populations of the Atlantic silverside. *Oecologia*, 122: 210–219.
- Boyd, P. W., Collins, S., Dupont, S., Fabricius, K., Gattuso, J. P., Havenhand, J., Hutchins, D. A. *et al.* 2018. Experimental strategies to assess the biological ramifications of multiple drivers of global ocean change—a review. *Global Change Biology*, 24: 2239–2261.
- Caldeira, K., and Wickett, M. E. 2003. Anthropogenic carbon and ocean pH. *Nature*, 425: 365–365.
- Cattano, C., Claudet, J., Domenici, P., and Milazzo, M. 2018. Living in a high CO₂ world: a global meta-analysis shows multiple trait-mediated fish responses to ocean acidification. *Ecological Monographs*, 88: 320–335.
- Chambers, R. C., Candelmo, A. C., Habeck, E. A., Poach, M. E., Wicczorek, D., Cooper, K. R., Greenfield, C. E. *et al.* 2014. Effects of elevated CO₂ in the early life stages of summer flounder, *Paralichthys dentatus*, and potential consequences of ocean acidification. *Biogeosciences*, 11: 1613–1626.
- Conover, D., and Murawski, S. 1982. Offshore winter migration of the Atlantic silverside, *Menidia menidia*. *Fishery Bulletin*, 80: 145–150.
- Conover, D. O. 1985. Field and laboratory assessment of patterns in fecundity of a multiple spawning fish: the Atlantic silverside, *Menidia menidia*. *Fishery Bulletin*, 83: 331–341.
- Conover, D. O., Duffy, T. A., and Hice, L. A. 2009. The covariance between genetic and environmental influences across ecological gradients – reassessing the evolutionary significance of countergradient and cogradient variation. *Annals of the New York Academy of Sciences*, 1168: 100–129.
- Conover, D. O., and Kynard, B. E. 1981. Environmental sex determination: interaction of temperature and genotype in a fish. *Science*, 213: 577–579.
- Conover, D. O., and Kynard, B. E. 1984. Field and laboratory observations of spawning periodicity and behavior of a northern population of the Atlantic silverside, *Menidia menidia* (Pisces: atherinidae). *Environmental Biology of Fishes*, 11: 161–171.
- Conover, D. O., and Munch, S. B. 2002. Sustaining fisheries yields over evolutionary time scales. *Science*, 297: 94–96.
- Conover, D. O., and Present, T. M. C. 1990. Countergradient variation in growth rate: compensation for length of the growing season among Atlantic silversides from different latitudes. *Oecologia*, 83: 316–324.
- Conover, D. O., and Ross, M. R. 1982. Patterns in seasonal abundance, growth and biomass of the Atlantic silverside, *Menidia menidia*, in a New England estuary. *Estuaries*, 5: 275–286.
- Cross, E. L., Murray, C. S., and Baumann, H. 2019. Diel and tidal pCO₂ × O₂ fluctuations provide physiological refuge to early life stages of a coastal forage fish. *Scientific Reports*, 9: 18146.
- Depasquale, E., Baumann, H., and Gobler, C. J. 2015. Variation in early life stage vulnerability among Northwest Atlantic estuarine forage fish to ocean acidification and low oxygen. *Marine Ecology Progress Series*, 523: 145–156.
- Dickson, A., and Millero, F. 1987. A comparison of the equilibrium constants for the dissociation of carbonic acid in seawater media. *Deep Sea Research Part A. Oceanographic Research Papers*, 34: 1733–1743.
- Dickson, A. G. 1990. Standard potential of the reaction: $agCl(s) + 12H_2(g) = Ag(s) + HCl(aq)$, and the standard acidity constant of the ion HSO₄[–] in synthetic sea water from 273.15 to 318.15 K. *The Journal of Chemical Thermodynamics*, 22: 113–127.
- Doney, S. C., Busch, D. S., Cooley, S. R., and Kroeker, K. J. 2020. The impacts of ocean acidification on marine ecosystems and reliant human communities. *Annual Review of Environment and Resources*, 45: 83–112.
- Esbaugh, A. J. 2018. Physiological implications of ocean acidification for marine fish: emerging patterns and new insights. *Journal of Comparative Physiology B*, 188: 1–13.
- Faria, A. M., Lopes, A. F., Silva, C. S. E. d., Novais, S. C., Lemos, M. F., and Gonçalves, E. J. 2018. Reproductive trade-offs in a temperate reef fish under high pCO₂ levels. *Marine Environmental Research*, 137: 8–15.
- Ganias, K., Lowerre-Barbieri, S. K., and Cooper, W. 2015. Understanding the determinate–indeterminate fecundity dichotomy in fish populations using a temperature dependent oocyte growth model. *Journal of Sea Research*, 96: 1–10.
- Ganias, K., and Lowerre-Barbieri, S. 2018. Oocyte recruitment and fecundity type in fishes: refining terms to reflect underlying processes and drivers. *Fish and Fisheries*, 19: 562–572.
- Ganias, K., Somarakis, S., Machias, A., and Theodorou, A. 2004. Pattern of oocyte development and batch fecundity in the Mediterranean sardine. *Fisheries Research*, 67: 13–23.
- Harvey, B. P., Gwynn-Jones, D., and Moore, P. J. 2013. Meta-analysis reveals complex marine biological responses to the interactive effects of ocean acidification and warming. *Ecology and Evolution*, 3: 1016–1030.
- Heuer, R. M., and Grosell, M. 2016. Elevated CO₂ increases energetic cost and ion movement in the marine fish intestine. *Scientific Reports*, 6: 34480.
- Hyle, A. R., McBride, R. S., and Olney, J. E. 2014. Determinate versus indeterminate fecundity in American shad, an anadromous clupeid. *Transactions of the American Fisheries Society*, 143: 618–633.
- Kroeker, K. J., Kordas, R. L., Crim, R., Hendriks, I. E., Ramajo, L., Singh, G. S., Duarte, C. M. *et al.* 2013. Impacts of ocean acidification on marine organisms: quantifying sensitivities and interaction with warming. *Global Change Biology*, 19: 1884–1896.

- Malvezzi, A., Murray, C. S., Feldheim, K. A., Dibattista, J. D., Garant, D., Gobler, C. J., Chapman, D. D. *et al.* 2015. A quantitative genetic approach to assess the evolutionary potential of a coastal marine fish to ocean acidification. *Evolutionary Applications*, 8: 352–362.
- McBride, R. S., Somarakis, S., Fitzhugh, G. R., Albert, A., Yaragina, N. A., Wuenschel, M. J., Alonso-Fernández, A. *et al.* 2015. Energy acquisition and allocation to egg production in relation to fish reproductive strategies. *Fish and Fisheries*, 16: 23–57.
- McElroy, W. D., Wuenschel, M. J., Towle, E. K., and McBride, R. S. 2016. Spatial and annual variation in fecundity and oocyte atresia of yellowtail flounder, *Limanda ferruginea*, in US waters. *Journal of Sea Research*, 107: 76–89.
- McNeil, B. I., and Sasse, T. P. 2016. Future ocean hypercapnia driven by anthropogenic amplification of the natural CO₂ cycle. *Nature*, 529: 383–386.
- Mehrbach, C., Culberson, C., Hawley, J., and Pytkowicz, R. 1973. Measurement of the apparent dissociation constants of carbonic acid in seawater at atmospheric pressure. *Limnology and Oceanography*, 18: 897–907.
- Middaugh, D. P., Hemmer, M. J., and Goodman, L. R. 1987. Methods for Spawning, Culturing and Conducting Toxicity-Tests with Early Life Stages of Four Atherinid Fishes: the Inland Silverside, *Menidia beryllina*, Atlantic silverside, *M. menidia*, Tidewater Silverside, *M. peninsulae* and California grunion, *Leuresthes tenuis*. U.S. Environmental Protection Agency, Washington, DC.
- Miller, G. M., Watson, S.-A., McCormick, M. I., and Munday, P. L. 2013. Increased CO₂ stimulates reproduction in a coral reef fish. *Global Change Biology*, 19: 3037–3045.
- Miller, S. H., Breitburg, D. L., Burrell, R. B., and Keppel, A. G. 2016. Acidification increases sensitivity to hypoxia in important forage fishes. *Marine Ecology Progress Series*, 549: 1–8.
- Munday, P. L., Crawley, N. E., and Nilsson, G. E. 2009. Interacting effects of elevated temperature and ocean acidification on the aerobic performance of coral reef fishes. *Marine Ecology Progress Series*, 388: 235–242.
- Murray, C. S., and Baumann, H. 2018. You better repeat it: complex temperature × CO₂ effects in Atlantic silverside offspring revealed by serial experimentation. *Diversity*, 10: 1–19.
- Murray, C. S., and Baumann, H. 2020. Are long-term growth responses to elevated pCO₂ sex-specific in fish? *Plos One*, 15: e0235817.
- Murray, C. S., Fuiman, L., and Baumann, H. 2017. Consequences of elevated CO₂ exposure across multiple life stages in a coastal forage fish. *ICES Journal of Marine Science*, 74: 1051–1061.
- Murray, C. S., Malvezzi, A. J., Gobler, C. J., and Baumann, H. 2014. Offspring sensitivity to ocean acidification changes seasonally in a coastal marine fish. *Marine Ecology Progress Series*, 504: 1–11.
- Murray, C. S., Wiley, D., and Baumann, H. 2019. High sensitivity of a keystone forage fish to elevated CO₂ and temperature. *Conservation Physiology*, 7: 1–12.
- Nagelkerken, I., Alemany, T., Anquetin, J. M., Ferreira, C. M., Ludwig, K. E., Sasaki, M., and Connell, S. D. 2021. Ocean acidification boosts reproduction in fish via indirect effects. *PLoS Biology*, 19: e3001033.
- Parker, L., Ross, P., and O'Connor, W. 2011. Populations of the Sydney rock oyster, *Saccostrea glomerata*, vary in response to ocean acidification. *Marine Biology*, 158: 689–697.
- Pörtner, H.-O. 2008. Ecosystem effects of ocean acidification in times of ocean warming: a physiologist's view. *Marine Ecology Progress Series*, 373: 203–217.
- Press, Y., McBride, R., and Wuenschel, M. 2014. Time course of oocyte development in winter flounder *Pseudopleuronectes americanus* and spawning seasonality for the Gulf of Maine, Georges Bank and southern New England stocks. *Journal of Fish Biology*, 85: 421–445.
- Pringle, J. W., and Baumann, H. 2019. Otolith-based growth reconstructions in young-of-year Atlantic silversides *Menidia menidia* and their implications for sex-selective survival. *Marine Ecology Progress Series*, 632: 193–204.
- Przeslawski, R., Byrne, M., and Mellin, C. 2015. A review and meta-analysis of the effects of multiple abiotic stressors on marine embryos and larvae. *Global Change Biology*, 21: 2122–2140.
- Riebesell, U., Fabry, V. J., Hansson, L., and Gattuso, J. P. 2011. Guide to Best Practices for Ocean Acidification Research and Data Reporting. Publications Office of the European Union, Luxembourg.
- Salinas, S., Perez, K. O., Duffy, T. A., Sabatino, S. J., Hice, L. A., Munch, S. B., and Conover, D. O. 2012. The response of correlated traits following cessation of fishery-induced selection. *Evolutionary Applications*, 5: 657–663.
- Schade, F. M., Clemmesen, C., and Wegner, K. M. 2014. Within- and transgenerational effects of ocean acidification on life history of marine three-spined stickleback (*Gasterosteus aculeatus*). *Marine Biology*, 161: 1667–1676.
- Schultz, E. T., Conover, D. O., and Ehtisham, A. 1998. The dead of winter: size dependent variation and genetic differences in seasonal mortality among Atlantic silverside (Atherinidae: *Menidia menidia*) from different latitudes. *Canadian Journal of Fisheries and Aquatic Sciences*, 55: 1149–1157.
- Schwemmer, T., Baumann, H., Murray, C., Molina, A., and Nye, J. 2020. Acidification and hypoxia interactively affect metabolism in embryos, but not larvae, of the coastal forage fish *Menidia menidia*. *Journal of Experimental Biology*, 223: jeb228015.
- Shaw, E. C., Munday, P. L., and McNeil, B. I. 2013. The role of CO₂ variability and exposure time for biological impacts of ocean acidification. *Geophysical Research Letters*, 40: 4685–4688.
- Welch, M. J., and Munday, P. L. 2016. Contrasting effects of ocean acidification on reproduction in reef fishes. *Coral Reefs*, 35: 485–493.
- Wootton, R. 1985. Energetics of reproduction. *In Fish Energetics*, pp. 231–254. Ed. by Tytler, P. and Calow, P. Springer, Berlin, Germany.

Handling Editor: Howard Browman