

1 Observing post-release mortality for dusky sharks" *Carcharhinus obscurus*, captured in
2 the US pelagic longline fishery

3
4 James A. Sulikowski¹, Walt Golet², Eric R. Hoffmayer³, William B. Driggers III³, Lisa J.
5 Natanson⁴, Amy Carlson¹, and Brett B. Swezey¹,

6
7 ¹University of New England, Marine Science Center, Biddeford, ME 04005 USA

8 ²University of Maine, School of Marine Sciences, 5706 Aubert Hall, RM 360, Orono, ME
9 04469-5706

10 ³National Marine Fisheries Service, Southeast Fisheries Science Center, Mississippi
11 Laboratories, 3209 Frederic Street, Pascagoula, MS 39567, USA

12 ⁴National Marine Fisheries Service, Northeast Fisheries Science Center, Narragansett
13 Laboratory, 28 Tarzwell Drive, Narragansett, RI 02882, USA

14
15 Corresponding Author: Professor, School of Mathematical and Natural Sciences, New College of
16 Interdisciplinary Arts and Sciences, Arizona State University's West Campus

17 USA. tel: 602-543-6050: e-mail: james.sulikowski@asu.edu

18

19

20 **Abstract**

21 The latest stock assessment for the dusky shark, *Carcharhinus obscurus*, in the western North
22 Atlantic Ocean indicates the population is overfished and experiencing overfishing. As part of a
23 rebuilding plan, the commercial and recreational retention of dusky sharks has been prohibited
24 since 2000. Despite this prohibition, dusky sharks are bycatch in multiple fisheries, including
25 the pelagic longline fishery; however, post-release mortality rates (PRM) have not been
26 empirically determined for this gear. Herein we estimated PRM of dusky sharks captured by the
27 US pelagic long line fleet in the western North Atlantic Ocean utilizing pop-up satellite archival
28 transmitting (PSAT) tags. One hundred and twenty three dusky sharks were captured on
29 commercial pelagic long line gear and time on the hook, based on hook timer data, ranged from
30 0.8 to 8.1 hours (4.3 ± 0.28 S.D.). No at-vessel mortality (AVM) was observed for any dusky
31 sharks in this study. Prior to release, 50 PSAT LIFE tags (Lotek Inc.) were attached to dusky
32 sharks (females $n=12$, 209 ± 8 cm FL; males $n=4$, 198 ± 7 cm FL; unknown sex $n=34$, 214 ± 7
33 cm FL) to assess PRM rates in the pelagic long line fishery during a 30 day attachment period.
34 Forty-three of the 50 deployed tags reported data with deployment times ranging from one to 28
35 days (11.2 ± 9.8 days). Four dusky sharks were in poor condition at release, two individuals
36 suffered PRM, which occurred within two hours after release. Total mortality rate (AVM and
37 PRM) in the current study was 5.1%, far below estimates reported for bottom longline gear
38 (~97%), and reinforces the notion that PRM should be evaluated by species, season, and gear
39 type.
40

41

42 Key words: Dusky shark; bycatch; survival; satellite tag, pelagic long line

43

44 **Introduction**

45 Sharks represent a group of marine organisms that are exploited globally by recreational and
46 commercial fisheries as both the target species (e.g. Worm et al., 2013) and or as bycatch (e.g.
47 Molina and Cooke, 2012). Although bycatch is often discarded, the capture and handling can
48 often lead to death (e.g. Gallagher et al., 2014). The coupling of life history traits such as late
49 maturity and slow growth, limits the resiliency of this group of fish to recover from these
50 anthropogenic pressures (e.g Stevens, 2000; Musyl and Gilman, 2019) on a global scale
51 (Lewison et al., 2004; Ellis et al., 2017). These circumstances have mandated bycatch mitigation
52 measures and management plans in various regions of the world (e.g. Ellis 2017; Musyl and
53 Gilman, 2019).

54 The dusky shark, *Carcharhinus obscurus*, inhabits coastal and pelagic ecosystems
55 circumglobally in temperate, subtropical and tropical marine waters (Compagno 1984). In the
56 territorial waters of the United States (U.S.) within the western North Atlantic Ocean, dusky
57 sharks range from the New England states to Florida and throughout the northern Gulf of Mexico
58 (GOM) (Castro 2011). Based on genetic analyses (Benavides et al. 2011) and their highly
59 migratory nature (Kohler et al. 1998), dusky sharks are managed as a single stock in this region
60 (NMFS, 1999). Stock assessments indicate this population is overfished and has been
61 experiencing overfishing since 1990 (SEDAR, 2106). Even though dusky shark retention has
62 been prohibited for several decades (NMFS, 1999), stock assessment model analysis suggests
63 stock recovery will not occur until 2108 (SEDAR, 2016). A potential limiting factor in the
64 recovery process is the impact of incidental capture in multiple commercial (e.g. bottom and
65 pelagic longline; bottom long line and pelagic long line, respectively) and recreational fisheries
66 (Morgan et al., 2009; NMFS 2016). The U.S. pelagic long line fishery operates within the U.S.
67 Exclusive Economic Zone (EEZ) and on the high seas, employs thousands of people, and is a
68 high value fishery responsible for approximately 66% of the total highly migratory species
69 (HMS) landed value in U.S. waters (NOAA Safe Report 2017). The pelagic long line fishery is
70 highly regulated, including, but not limited to, mandatory maintenance of logbooks, 100%
71 electronic monitoring, exclusive use of circle hooks (since 2004, minimum hook size of 16/0),
72 utilization of several types of dehooking and leader cutting devices and a requirement to leave no
73 more than three feet of gangion material attached if a hook is left in place (NMFS, 2016). Based
74 on observer coverage, self-reporting and modeled extrapolation, the pelagic long line fleet
75 interacted with thousands of dusky sharks between 2008 and 2014; however, the fate of these
76 released sharks is unknown (NMFS, 2016).

77 At-vessel (AVM) and post-release (PRM) mortality are important metrics for estimating total
78 mortality, calculating population size and setting catch limits, especially for vulnerable species
79 such as dusky sharks. (e.g. Davis, 2002; Sulikowski et al., 2017). These inputs are essential for
80 calculating total fishing mortality and stock biomass and aid in the development of biologically
81 acceptable catch limits for a given fishery (Alverson, 1999; Davis, 2002). Fishery observers can
82 provide estimates of AVM, but obtaining PRM rates is more difficult given the fate of a released
83 shark is unknown. Consequently, marine fisheries management utilizes conservative (“worst-
84 case scenario”) PRM estimates when data are not available or surrogate rates are applied from
85 different species and/or gear types. For example, the Dusky Shark Working Group (SEDAR
86 2011) used the difference (6%) between AVM (13%) and PRM (19%) for blue sharks, *Prionace*
87 *glauca*, and added that difference to the AVM Pelagic Longline Observer Program data. The

88 use of this surrogate data, resulted in a 44.2% total discard mortality rate for dusky sharks
89 captured in the pelagic long line fishery. Using such proxies can have major consequences,
90 especially if the relationships between AVM and PRM rates vary substantially among species
91 and gear types. To prevent such a scenario, the National Marine Fisheries Service (NMFS) has
92 prioritized obtaining accurate PRM estimates for all HMS species and gear types. Recent
93 estimates of total discard mortality for dusky sharks captured in the bottom long line fishery are
94 between 88% and 97% (Morgan and Burgess, 2007, Marshall et al., 2012; Marshall 2015).
95 However, to date, no studies have directly quantified the mortality of dusky sharks associated
96 with pelagic long line fisheries. Given the status of the dusky shark population in the Western
97 North Atlantic Ocean and the lack of direct estimates of PRM rates in the pelagic long line
98 fishery, the objective of the current study was to estimate AVM and PRM rates for dusky sharks
99 captured on pelagic long line gear using survivorship pop-up archival satellite transmitting
100 (PSAT) tags.

101 **Methods**

102 Commercial pelagic long line gear was deployed from two vessels off the Wanchese, North
103 Carolina using experienced pelagic long line captains and crew. Each vessel conducted one day
104 and one night set approximately 48 km offshore over a 36-hour period in May 2016 (Figure 1).
105 With the exception of mainline length, gear configuration was kept identical to methods used in
106 that region to remove bias. Mainline length was 8-13km whereas commercial sets are typically
107 44-88km in length. Shorter mainlines were used so that each gangion on every set was equipped
108 with a hook timer (HT 600, Lindgren-Pitman, Inc.). Between 150-175 gangions, each
109 constructed of a 20 m 1.8 mm diameter monofilament leader (Lindgren-Pitman, Inc), weighted
110 swivel (Lindgren-Pitman, Inc.) and 16/0 circle hook (Mustad, # 39960-DC), were attached to the
111 3.5 mm diameter monofilament mainline (Lindgren-Pitman, Inc), spaced approximately 25 m
112 apart, and baited with whole squid (*Loligo* sp.). According to NMFS pelagic long line observer
113 data, average soak time within this fishery is approximately 8.3 hours, with an interquartile range
114 of 7.1 to 9.3 hours (Cushner S., NMFS, pers. comm.) and mainline length varies from 44 to 88
115 km in length (NMFS, 2017). Thus, for the purposes of the study herein, each longline set
116 soaked for nine hours (timing started when first hook was deployed). Hook timers were activated
117 when a fish applied tension to the leader and provided a detailed record for time on the line
118 (TOL) for each captured shark. Following the nine hour soak, the mainline was retrieved and the
119 status of each hook recorded (i.e. bare hook, bite off, fish captured). At haulback, the location of
120 capture, TOL, sea surface temperature (SST), estimated fork length (so sharks did not need to be
121 removed from the water) and injury condition of each dusky shark were recorded prior to
122 tagging. A shark was given an injury condition of (1) if there were no visible signs of trauma to
123 the body (e.g. no blood or skin abrasions) and the shark was hooked in the jaw; (2) if minor skin
124 abrasions or small lacerations were present on the body, multiple hooks observed in the jaw,
125 and/or trailing monofilament from a previous capture was observed; (3) if there were obvious
126 signs of trauma, such as lacerations on the body, broken jaw, or gut hooking; or (4) if the shark
127 was moribund (modified from Marshall et al., 2015).

128 Prior to release and regardless of assigned capture condition, the first 50 sharks that were
129 captured were tagged with a satellite tag to measure PRM. Here, a Lotek PSAT LIFE tag was
130 attached to each shark using a stainless-steel dart anchor (Hallprint®, SSD, 57 mm x 15 mm,
131 Victoria Harbor, Australia). The 13cm tether consisted of 136kg monofilament line (300-lb test

132 extra-hard Hi-Catch, Momoi Fishing Net Mfg. Co. Ltd., Ako City, Hyogo Prefecture, Japan)
133 with heat shrink tubing to minimize abrasions to the animal. Tags were inserted into the dorsal
134 musculature just below the first dorsal fin and in line with the insertion of the fin with a 2m
135 tagging pole following the protocols of Hoffmayer et al. (2014). All dusky sharks were tagged *in*
136 *situ* and at no point were removed from the water. Following tag attachment, the line was cut
137 less than 1m above the hook and a release condition adapted from Manire et al. (2001) (1- swim
138 burst, 2- strong swimming, 3- sluggish swimming, 4- sank with no visible swimming effort) was
139 assigned.

140 Two components of mortality were estimated, AVM and PRM. At vessel mortality was defined
141 as a shark that was dead upon capture, while PRM was defined as a shark spending three
142 consecutive days at a constant depth and temperature, as determined by tag data, after release
143 (Heberer et al., 2010; Marshall et al. 2015; Campana et al., 2016). If a mortality event occurred
144 prior to each tag's preprogrammed 28 day deployment duration, the PSAT LIFE tags were
145 equipped with a constant depth fail-safe release that jettisoned the tag from the shark if a
146 constant depth ($\pm 3\text{m}$) was maintained for 72 hours (indicative of a mortality event). Tags
147 collected daily minimum and maximum depth (up to 2000m) and ambient temperature (range -5
148 to 35°C). Following pop-off and data transmission, tag reports were downloaded from the
149 ARGOS website and post-processed using the Lotek TagTalk software (ver. 1.10.8.14). In
150 addition to determining the PRM rate, the binomial 95% confidence intervals were calculated to
151 determine the uncertainty around this estimate; however, due to the low sample size, more data
152 are required to support a more meaningful analysis. All analyses and figures were completed in
153 SigmaPlot v12.5, MATLAB v9.1 and ArcGIS v10.4. All means are reported with corresponding
154 standard errors, and statistical tests were considered significant at $\alpha = 0.05$.

155 **Results**

156 Six hundred and seventy three hooks were deployed during the four pelagic long line sets which
157 resulted in 175 hook bite-offs and the capture of 123 dusky sharks (202 ± 28 cm FL; range 167
158 cm-243 cm). Dusky shark TOL ranged from 0.8 to 8.1 hours with a mean of 4.3 ± 0.3 hours.
159 Mean injury and release condition codes for all captured dusky sharks were 1.8 ± 0.1 and 1.3 ± 0.1 ,
160 respectively (Table 1). No AVM was observed for dusky sharks caught in this study. Average
161 SST over the course of the study was $24 \pm 0.2^\circ\text{C}$.

162 Fifty PSAT LIFE tags were deployed on dusky sharks (females $n=12$, 209 ± 8 cm FL; males
163 $n=4$, 198 ± 7 cm FL; unknown sex $n=34$, 214 ± 7 cm FL). Forty-three of the 50 tags
164 subsequently transmitted data (86%) with deployment times ranging from 1 to 28 days (mean =
165 11.2 ± 9.8 days) (Table 2). Although the depths were not constant, four tags remained on sharks
166 less than three days, the minimum number of days to identify a PRM event, and were
167 subsequently removed from analyses. Thirty-one tags released prematurely (i.e. < 28 days) due
168 to an assumed tether failure. However, despite these premature detachments, the PSAT LIFE
169 tags captured daily activity patterns and vertical profiles from 37 tags, which indicated post-
170 release survival (Figure 2). Two sharks recorded constant depth readings immediately following
171 release that we considered PRM events. One shark that was at first considered an AVM (injury
172 code and release condition of 4), recovered and was at liberty for 8 days prior to premature
173 release of the satellite tag.

174 Despite four dusky sharks being assigned a release condition of 4, these sharks survived the
175 capture and tagging process. However, two sharks (1476 and 1513) with a release condition of 2
176 suffered PRM (5.1%). The 95% confidence intervals around the PRM rate ranged from 0.6 to
177 17.3%. Mean injury code, release condition code and TOL of those two dusky sharks were
178 2.0 ± 0.0 , 2.0 ± 1.4 , and 4.8 ± 1.7 hours, respectively. Soak times of 1–3 h resulted in 0% PRM,
179 while soak-times of 3–5 hours and ≥ 5 hours each accounted for one PRM event (the two sharks
180 listed previously; Figure 3ab). Regardless of TOL, the majority of the sharks exhibited minimal
181 signs of injury from pelagic long line capture (95%), and most (78%) were released in good
182 condition (Figure 3ab). Since only two PRM events were observed in the study, a larger sample
183 size is needed to better identify causal variables.

184 **Discussion**

185 One of the biggest challenges facing fisheries managers is reducing bycatch (Musick 1999;
186 Afonso et al. 2011). Understanding the fate (dead or live) of discarded fishes is essential for
187 properly characterizing total fishing mortality and its resulting implications for estimates of stock
188 status (Sulikowski et al., 2017). In addition, PRM estimates have a practical application that aids
189 managers in the development of biologically realistic catch limits for a given fishery (e.g.,
190 Alverson, 1999; Benoit et al., 2015; Sulikowski et al., 2017). Also vital to management is
191 understanding the relationship between mortality estimates and key variables linked to those
192 events. For example, gear configuration, bait type, environmental parameters, and biology of the
193 species of interest, can lead to best practices that will reduce overall mortality of that species
194 (e.g. Carruthers et al., 2009; Capizzano et al., 2016; Ellis et al., 2017) and extend fishing
195 opportunities. However, obtaining this information can be difficult due to challenges associated
196 with cost and the ability to collect representative samples. As such, AVM and PRM studies have
197 only been conducted for a limited number of elasmobranchs (Hoolihan et al. 2011), and even
198 fewer studies have been able to investigate the relationships between mortality and the
199 aforementioned variables in sharks (e.g. Dapp et al. 2016; Campana et al., 2016; Ellis et al.,
200 2017). While fishing practices vary (e.g., vessel size, mainline length, hook/bait configurations,
201 soak time) across all the different regions the pelagic long line fleet fishes (Musyl et al., 2011)
202 here gear was fished in line with how vessels typically fish off North Carolina. Thus, the AVM
203 and PRM presented herein, are representative of the fishery in the geographic region this study
204 was conducted. The current study represents the first empirically obtained PRM estimate for
205 dusky sharks within the western North Atlantic pelagic long line fishery and adds to the limited
206 body of knowledge for this species of cartilaginous fish.

207 Previous studies reporting fishing mortality rates of specific shark species in both pelagic long
208 line and bottom long line fisheries have documented a wide range of estimated rates (0%-90%)
209 among species and gear types (Beerkircher et al., 2002; Diaz and Serafy, 2005; Morgan and
210 Burgess, 2007; Campana et al. 2009; Morgan and Carlson, 2010; Hutchinson et al., 2015; Ellis et
211 al., 2017). The current findings represent the lowest published PRM estimates (5%) for dusky
212 sharks, a stark contrast to that of Marshall et al. (2015) who reported a 29% PRM rate for dusky
213 sharks caught on bottom long line gear. Moreover, no AVM (0%) was observed for any of the
214 123 dusky sharks caught in this study, which again is in contrast to the estimated AVM of nearly
215 40% (dead or moribund individuals) reported by Marshall et al. (2015) for dusky sharks captured
216 by bottom long line gear. Interestingly, although the values for AVM and PRM in the current
217 study were lower than for dusky sharks caught on bottom long line (Marshall et al. 2015), the

218 relationship between the two mortality types was similar ($PRM = AVM + 6\%$). Although the
219 $AVM + 6\%$ rule was used to estimate a PRM rate of 44.2% for dusky sharks in the pelagic long
220 line fishery (SEDAR 2011), that estimate is well above values obtained in the current study. In
221 addition, while limited, observer data obtained within our general sampling area and time frame
222 suggests that interaction with and AVM of dusky sharks is variable. For example, in March,
223 April, and May 0/1, 0/0 and 7/15 dusky sharks were observed/suffered AVM (Cushner S.,
224 NMFS, pers. comm.). Collectively, this information suggests that future work should focus on
225 the temporal and spatial variability in AVM/PRM within this fishery. Finally, the two
226 mortalities that were observed in the current study occurred within 24 hours of release, a time
227 frame widely observed for several sharks species captured in fixed gear including dusky sharks
228 (Heberer et al., 2010; Marshall et al. 2015; Campana et al., 2016; Whitney et al., (2016).

229 The knowledge of how a species' physiology is affected by the interaction with a specific gear
230 type is important as inferences can be made relative to the causes producing the observed PRM
231 rates (Marshall et al., 2015). For example, dusky sharks are obligate ram ventilators and need to
232 force oxygenated water over their gills in order to respire (Liem and Summers 1999). In general,
233 bottom long line gear is set on the substrate and has relatively short leader lengths (~2m;
234 Marshall et al., 2015) limiting a sharks movement and their ability to ram ventilate. Thus, sharks
235 that rely on ram ventilation must compensate for decreases in oxygen availability by increasing
236 swimming speed and/or mouth gape (Carlson and Parsons 2001). In addition, during the warmer
237 months of the year, hypoxic and anoxic conditions can occur at or near the bottom due to
238 eutrophication and water column stratification (Rabalais et al. 2002), which can further
239 exacerbate the respiratory stress. The combination of short gangion lengths and hypoxic and
240 anoxic conditions would result in rapid asphyxiation following hooking and would contribute to
241 mortality when sharks are caught during these conditions (Morgan and Burgess, 2007). In
242 comparison, pelagic long line gear drifts at or near the water's surface with longer leader lengths
243 (>20m), presumably allowing the sharks to remain swimming while on the line in more
244 oxygenated waters. While not directly comparable, these gear and environmental differences
245 between bottom long line and pelagic long line may have resulted in the much lower mortality
246 rates observed in the current study. However, since the current study was conducted during a
247 specific temporal period, further research as to the effects of temperature on the PRM of dusky
248 shark captured in the pelagic long line should be explored.

249 Previous research suggests soak time and water temperature are the main factors affecting
250 mortality for sharks captured with bottom long line gear (Morgan and Burgess, 2007; Morgan
251 and Carlson, 2010; Marshall et al., 2012; Gallagher et al., 2014). While, the depth the hooks
252 were fishing or the temperatures that they were experiencing are not known, given the length of
253 mainline, spacing of floats, gangion length, geographic location and the time of year did not
254 vary/were consistent during the study, all sharks tagged were assumed representative of
255 environmental conditions while on the line. Given this, and since total mortality was low,
256 temperature did not appear to be associated with the mortality in dusky sharks in the current
257 study. However, future studies should investigate how this abiotic parameter influences survival
258 over varying temporal regimes. Mean TOL for dusky sharks captured was 4.3 hours with no
259 AVM observed. Over this same period, 39% of the dusky sharks caught on bottom long line
260 were either moribund or dead (marshall et al., 2015). research has suggested that cooler water
261 temperatures positively correlate with increased survival of sharks captured on fixed gear. For
262 example, Gallagher et al., (2014) found a significant increase in survival of several shark species

263 (including dusky sharks) that were captured at deeper hook depths. Since dissolved oxygen
264 content increases with decreasing temperature the ability to overcome any oxygen deficits
265 produced by the capture event would be greater in cooler waters (e.g Skomal and Bernal, 2010)
266 and enhanced by the ability to actively ram ventilate via a longer gangion (Gallagher et al.,
267 2014). Collectively, the aforementioned data reinforce the possibility that the longer gangions
268 and mobility of the gear provided dusky sharks on pelagic long line the ability to reduce their
269 metabolic rate (swim down to cooler water; Skomal and Bernal, 2010) and ram ventilate (Liem
270 and Summers, 1999) allowing for higher survival and faster recovery than those observed from
271 previous bottom long line studies.

272 While direct comparisons among species and between gear types (bottom long line vs pelagic
273 long line) cannot be made (i.e different gear configurations and soak times), the limited
274 information that exists suggests that mitigation measures can be effective in reducing PRM in
275 sharks. For example, Moyes et al. (2006) found that moribund blue sharks had blood chemistry
276 values that were indicative of exhaustive exercise, a possible result of long soak times, and
277 suggested shorter soak times may lead to better survival rates. Using a generalized linear model,
278 Campana et al. (2009) found fishing gear and techniques appear to be the main factors
279 influencing hooking mortality in blue sharks, and health status of the shark at release contributed
280 to PRM. Afonso et al. (2011) reported that the use of circle hooks (as opposed to J hooks)
281 suspended in the middle of the water column reduced the bycatch of several demersal
282 elasmobranch species. Finally, results of Tolotti et al., (2013) indicated that setting longline
283 hooks at depths greater than 100m could reduce the bycatch of oceanic whitetip sharks. Based on
284 the aforementioned data and the observations from the current study, requirements already in
285 place seem sufficient to keep AVM and PRM low in the pelagic long line fishery and could be
286 sufficient mitigation measures to reduce mortality for dusky sharks.

287 The results herein illustrate the importance of providing empirical data when estimating AVM
288 and PRM on species and gear specific bases. “Borrowing” values from other species and/or gear
289 types, while necessary for some data poor situations, in this case, overestimates the discard
290 mortality for this species and gear type (e.g Gallagher et al., 2014). Relatively low AVM and
291 PRM rates reported herein indicate dusky sharks are resilient to the capture and handling stress
292 experienced in the pelagic long line fishery, especially when compared to estimates derived for
293 the same species using bottom long line gear (97%; Marshall et al., 2015). Estimating AVM and
294 PRM associated with the pelagic long line fishery across a larger sample size, a wider geographic
295 range, set of gear configurations and environmental conditions would provide a more
296 comprehensive evaluation and should be considered in future studies. Based on the total fishing
297 mortality estimates collected in this study (AVM and PRM) for dusky sharks captured on pelagic
298 long line fishing gear, the PRM rate occurring within the pelagic long line fishery will need to be
299 updated for future stock assessments.

300 **Acknowledgments**

301 This study would not have been possible without the expertise, hard work, and dedication of
302 Captains Dewey Hemilright and Jeff Oden. This project was supported by the National Oceanic
303 and Atmospheric Administration section Bycatch Reduction Grant (NA15NMF4720375) and
304 was approved by University of New England’ s Animal Care and Use Committee (Institutional
305 Animal Care and Use Committee protocol number UNE-20160510SULJ). This manuscript
306 represents Marine Science Center contribution number 124.

307 **Literature Cited**

- 308 Afonso, A.S., Hazin, F.H.V., Carvalho, F., Pachero, J.C., Hazin, H., Kerstetter, D.W., Murie, D.,
309 and Burgess, G.H. 2011. Fishing gear modifications to reduce elasmobranch mortality in pelagic
310 and bottom longline fisheries off Northeast Brazil. *Fish. Res.*, 108: 336-343.
- 311 Alverson, D.L. 1999. Some observations on the science of bycatch. *Mar. Technol. Soc.*, 33: 12–
312 16.
- 313
- 314 Benoît, H.P., Capizzano, C.W., Knotek, R.J., Rudders, D.B., Sulikowski, J.A., Dean, M.J.,
315 Hoffman, W.S., Zemeckis, D.R., Mandelman, J.W. 2015. A generalized model for longitudinal
316 short- and long-term mortality data for commercial fishery discards and recreational fishery
317 catch-and-releases. *ICES Jour. of Mar. Sci.*, doi:10.1093/icesjms/fsv039.
- 318
- 319 Benavides, M.T., Horn, R.L., Feldheim, K.A., Shivji, M.S., Clarke, S.C., Wintner, S., Natanson,
320 L., Braccini, M., Boomer, J.J., Gulak, S.J.B., Chapman, D.D. 2011. Global phylogeography of
321 the dusky shark *Carcharhinus obscurus*: implications for fisheries management and monitoring
322 the shark fin trade. *End. Spec. Res.*, 14:13-22
- 323 Beerkircher, L., Cortes, E., Shivji, M., 2002. Characteristics of shark bycatch on pelagic
324 longlines off the southeastern United States, 1992-2000. *Mar. Fish. Rev.*, 64: 40–49.
- 325 Campana, S.T., Joyce, W., Manning, M.J. 2009. Bycatch and discard mortality in commercially
326 caught blue sharks *Prionace glauca* assessed using archival satellite pop-up tags. *Mar. Ecol.*
327 *Prog. Ser.*, 387: 241-253
- 328 Campana, S.E., Joyce, W., Fowler, M., Showell, M. 2016. Discards, hooking and post-release
329 mortality of porbeagle (*Lamna nasus*), shortfin mako (*Isurus oxyrinchus*) and blue shark
330 (*Prionace glauca*) in the Canadian pelagic longline fishery. *ICES Jour. of Mar. Sci.*, 73: 520–528.
- 331 Capizzano, C.W., Mandelman, J.W., Hoffman, W.S., Dean, M.J., Zemeckis, D.R., Benoît H.P.,
332 Stettner, M.J., Kneebone, J., Buchan, N.C., Langan, J.A., and Sulikowski, J.A. 2015. Estimating
333 and mitigating post-release mortality of Atlantic cod in the Gulf of Maine’s recreational rod-and-
334 reel fishery. *ICES Jour. of Mar. Sci.*, doi:10.1093/icesjms/fsw058
- 335 Carruthers, E.H., Schneider, D.C., Neilson, J.D. 2009. Estimating the odds of survival and
336 identifying mitigation opportunities for common by-catch in pelagic longline fisheries. *Biol.*
337 *Con.*, 142: 2620–2630.
- 338 Carlson, J.K. and G.R. Parsons. 2001. The effects of hypoxia on 3 sympatric shark species:
339 physiological and behavioral responses. *Env. Bio. of Fish.*, 61:427–433.
- 340 Castro, J.I. 2011. *The sharks of North America*. Oxford University Press, New York, p 640.
- 341 Compagno, L.J.V. 1984. *FAO Species Catalogue*. Vol. 4. *Sharks of the world. An annotated and*
342 *illustrated catalogue of shark species known to date. Part 2 - Carcharhiniformes*. *FAO Fish.*
343 *Synop.*, 125(4/2):251-655. Rome: FAO.

344 Dapp, D.R., Huveneers, C., Walker, T.I., Drew, M., and Reina, R.D. 2016. Moving from
345 measuring to predicting bycatch mortality: predicting the capture condition of a longline-caught
346 pelagic shark. *Fron. Mar. Sci.*, 2:126. doi: 10.3389/fmars.2015.00126.

347 Davis, M.W. 2002. Key principles for understanding fish bycatch discard mortality. *Can. J. Fish.*
348 *Aq. Sci.*, 59:1834-1843.

349 Diaz, G.A. and Serafy, J.E. 2005. Longline-caught blue shark (*Prionace glauca*): factors affecting
350 the numbers available for live release. *Fish. Bull.*, 103: 720–724.

351 Ellis, J.R., McCully Phillips, S.R., Poisson, F. 2017. A review of capture and post-release
352 mortality of elasmobranchs. *J. Fish Biol.*, 90:653-722.

353 Gallagher, A.J., Orbesen, E.S., Hammerschlag, N., Serafy, J.E. 2014. Vulnerability of oceanic
354 sharks as pelagic longline bycatch. *Glob. Ecol. Conserv.*, 1: pp. 50-59

355 Heberer, C., Aalbers, S.A., Bernal, D., Kohin, S., DiFiore, B., Sepulveda, C.A. 2010. Insights
356 into catch-and-release survivorship and stress-induced blood biochemistry of common thresher
357 sharks (*Alopias vulpinus*) captured in the southern California recreational fishery. *Fish. Res.*,
358 106: 495–500.

359 Hoffmayer, E.R., Franks, J.S., Driggers III, W.B., McKinney, J.A. 2014. Habitat, movements
360 and environmental preferences of dusky sharks, *Carcharhinus obscurus*, in the northern Gulf of
361 Mexico. *Mar. Bio.*, 161 (4): 911-924.

362 Hoolihan, J.P., Luo, J., Abascal, F.J., Campana, S.E., Metrio, G.D., Dewar, H., Domeier, M.L.,
363 Howey, L.A., Lutcavage, M.E., Musyl, M.K., Neilson, J.D., Orbesen, E.S., Prince, E.D., Rooker,
364 J.R. 2011. Evaluating post-release behavior modification in large pelagic fish deployed with pop-
365 up satellite archival tags. *ICES Jour. of Mar. Sci.*, 68(5):880-889.

366 Hutchinson, M.R., Itano, D.G., Muir, J.A., Holland, K.N. 2015. Post-release survival of juvenile
367 silky sharks captured in a tropical tuna purse-seine fishery. *Mar. Ecol. Prog. Ser.*, 521: 143–154.

368 Kohler, N.E., Casey, J.G., Turner, P.A. 1998. NMFS Cooperative Shark Tagging Program,
369 1962–93: an atlas of shark tag and recapture data. *Mar. Fish. Rev.*, 60(2):1–87.

370 Liem, K.F. and Summers, A.P. 1999. Muscular system: gross anatomy and functional
371 morphology of muscles. *In: Sharks, skates, and rays: the biology of elasmobranch fishes* (W.C.
372 Hamlett, ed.), Johns Hopkins University Press., 93-114.

373 Manire, C., Hueter, R., Hull, E. and Spieler, R. 2001. Serological changes associated with
374 gill-net capture and restraint in three species of sharks. *Trans. Am. Fish. Soc.* 130: 1038–
375 1048.

376 Marshall, H., Field, L., Afadada, A., Sepulveda, C., Skomal, G., Bernal, D. 2012. Hematological
377 indicators of stress in longline-captured sharks. *Comp. Biochem. Physio.*, A 162: 121–129.

378 Marshall, H., Skomal, G., Ross, P.G., Bernal, D. 2015. At-vessel and post-release mortality of
379 the dusky (*Carcharhinus obscurus*) and sandbar (*C. plumbeus*) sharks after longline capture.
380 *Fish. Res.*, 172: 373–384.

381 Molina, J. M., & Cooke, S. J. (2012). Trends in shark bycatch research: Current status and
382 research needs. *Reviews in Fish Biology and Fisheries*, 22, 719–737.
383 <https://doi.org/10.1007/s11160-012-9269-3>

384 Morgan, A., Burgess, G.H. 2007. At-vessel fishing mortality for six species of sharks caught in
385 the northwest Atlantic and Gulf of Mexico. *Proceedings of the 59th. Annual Conference of the*
386 *Gulf and Caribbean Fisheries Institute.*, 19, 123–130.

387 Morgan, A., Cooper, P., Curtis, T., Burgess, G.H. 2009. An overview of the United States East
388 Coast Bottom Longline Shark-Fishery, 1994-2003. *Mar. Fish. Rev.*, 71:23-38.

389 Morgan, A. and Carlson, J.K. 2010. Capture time, size and hooking mortality of bottom longline-
390 caught sharks. *Fish. Res.*, 101:32-37.

391 Moyes, C.D., Fragoso, N., Musyl, M., Brill, R.W. 2006. Predicting postrelease survival in large
392 pelagic fish. *Transactions of the American Fisheries Society.*, 135:1389-1397.

393 Musick, J.A. 1999. Ecology and conservation of long-lived marine animals. *American Fisheries*
394 *Society Symposium.*, 23:1-10.

395 Musyl, M.K., Brill, R.W., Curran, D.S., Fragoso, N.M., McNaughton, L.M., Nielsen, A.,
396 Kikkawa, B.S., Moyes, C.D. 2011. Postrelease survival, vertical and horizontal movements and
397 thermal habitats of five species of pelagic sharks in the central Pacific Ocean. *Fish. Bull.*, 109,
398 341–368.

399 Musyl MK, Gilman EL. Meta-analysis of post-release fishing mortality in apex predatory pelagic
400 sharks and white marlin. *Fish Fish.* 2019;00:1–35. <https://doi.org/10.1111/faf.12358>

401 National Marine Fisheries Service (NMFS). 1999. Final fishery management plan for Atlantic
402 tunas, swordfish and sharks. US Dept. of Commerce, NMFS, Office of Sustainable Fisheries,
403 Silver Spring, MD, 854 p.

404 National Marine Fisheries Service (NMFS). 2016. Amendment 5b to the 2006 Consolidated
405 Atlantic Highly Migratory Species Fishery Management Plan. Highly Migratory Species
406 Management Division.

407 NOAA Fisheries. 2017. 2016 Stock Assessment and Fishery Evaluation (SAFE) Report for
408 Atlantic Highly Migratory Species. Atlantic Highly Migratory Species Management Division

409 Rabalais, N.N., Turner, R.E., Wiseman Jr., W.J. 2002. Gulf of Mexico Hypoxia, A.K.A. “The
410 Dead Zone”. *An. Rev. Ecol. System.*, 33:235-263.

411 Skomal, G.B., Bernal, D. 2010. Physiological responses to stress in sharks. In J. Carrier, J.A.
412 Musick, M. Heithaus (Eds.), *Sharks and Their Relatives II: Biodiversity, Adaptive Physiology,*
413 *and Conservation*, CRC Press, Boca Raton (2010), pp. 459-490.

414 Southeast Data, Assessment, and Review (SEDAR). 2011. HMS dusky shark stock assessment
415 report. SEDAR, North Charleston, SC.

416 Southeast Data, Assessment, and Review (SEDAR). 2016. HMS dusky shark. Update assessment
417 to SEDAR 21 stock assessment report. SEDAR, North Charleston, SC.

418 Sulikowski, J.A., Benoit, H.P., Capizzano, C.W., Knotek, R.J., Mandelman, J.W., Platz, T.,
419 Rudders, D.B. 2017. Evaluating the condition and discard mortality of winter skate, *Leucoraja*
420 *ocellata*, following capture and handling in the Atlantic monkfish sink gillnet fishery. *J. Fish.*
421 *Res.*, <http://dx.doi.org/10.1016/j.fishres.2017.10.001>

422 Stevens, J. (2000). The effects of fishing on sharks, rays, and chimaeras (chondrichthyans), and
423 the implications for marine ecosystems. *ICES Journal Marine Science*, 57, 476–494.
424 <https://doi.org/10.1006/jmsc.2000.0724>

425 Tolotti, M.T., Travassos, P., Frédou, F.L., Wor, C., Andrade, H.A., Hazin, F. 2013. Size
426 distribution and catch rates of the oceanic whitetip shark caught by the Brazilian tuna longline
427 fleet. *Fish. Res.*, 143 (2013), pp. 136-142.

428 Whitney N.M., White C.F., Gleiss A.C., Schwieterman G.D., Anderson P, Hueter R.E., and
429 Skomal, G.B. 2016. A novel method for determining post-release mortality, behavior, and
430 recovery period using acceleration data loggers. *Fisheries Research*. 183: 210-221.

431 Worm, B., Davis, B., Kettner, L., Ward-Paige, C. A., Chapman, D., Heithaus, M. R., & Gruber,
432 S. H. (2013). Global catches, exploitation rates, and rebuilding options for sharks. *Marine Policy*,
433 40, 194–204. <https://doi.org/10.1016/j.marpol.2012.12.034>

434

Table 1. Injury code and release condition of 50 dusky sharks, *Carcharhinus obscurus*, captured with pelagic longline gear and affixed with a LOTEK PSAT LIFE satellite tag. Injury codes and release conditions represent at-vessel observations. Sample size (**n**) and time on the line are provided for each code/condition.

Injury Code	n (%)	Time-on-the-Line (hour)	
		Range	Mean (\pm SD)
1 (No outward injury, hook in jaw)	40 (80.0)	1:53 – 6:13	4:13 \pm 0.05
2 (Lacerations on body, multiple hooks in jaw)	8 (16.0)	2:35 – 6:38	5:27 \pm 0.06
3 (Lacerations on body, broken jaw, gut hooked)	1 (2.0)	6:12	-
4 (Moribund)	1 (2.0)	6:05	-

Release Condition	n (%)	Time-on-the-Line (hour)	
		Range	Mean (\pm SD)
1 (Swim burst)	21 (52.0)	1:53 – 6:38	4:33 \pm 0.06
2 (Strong swimming)	13 (26.0)	2:30 – 6:13	4:55 \pm 0.05
3 (Sluggish swimming)	7 (14.0)	2:35 – 6:00	3:57 \pm 0.06
4 (Sank with no visible swimming effort)	4 (8.0)	3:00 – 6:05	4:05 \pm 0.06

Table 2. Biological, tag-deployment, and post-release outcomes for dusky sharks, *Carcharhinus obscurus*, captured with pelagic longline gear. Note: Fork length was estimated; * denotes sharks removed from mortality analysis.

Shark	ID	FL (cm)	Sex	TOL (min)	Date tagged	Tag latitude N	Tag longitude W	Time at liberty (days)	Pop-up latitude ° N	Pop-up longitude ° W	Release Code	Injury Code	Release Condition	Outcome
1	1452	NA	NA	NA	5/25/16	NA	NA	12	37.5708	72.0347	1	1	NA	Survived
2	1453	NA	NA	6:05	5/25/16	35.7167	74.865	8	36.6667	68.7589	1	1	A	Survived
3	1455	213	NA	6:07	5/25/16	35.7233	74.8633	3	36.1175	74.4236	1	2	A	Survived
4	1456	177	NA	3:09	5/25/16	35.6190	74.7869	4	36.5942	73.8517	3	1	A	Survived
5	1457*	NA	NA	3:11	5/24/16	35.6683	74.8267	<3	36.3553	74.6686	1	1	A	Survived
6	1458	175	F	1:53	5/25/16	35.7517	74.8733	28	36.9217	74.5728	1	1	A	Survived
7	1459	NA	F	4:14	5/24/16	35.695	74.7833	28	37.3353	75.0675	1	1	A	Survived
8	1460	NA	NA	6:05	5/25/16	35.7267	74.8633	8	37.0907	73.005	4	4	A	Survived
9	1461	244	NA	No timer	5/25/16	35.73	74.85	4	38.1867	69.9911	3	1	A	Survived
10	1463	NA	NA	NA	5/25/16	NA	NA	NR	NR	NR	1	1	A	NR
11	1464	NA	NA	5:46	5/25/16	35.73	74.85	NR	NR	NR	1	1	A	NR
12	1465	177	NA	6:00	5/25/16	35.6883	74.8009	NR	NR	NR	1	1	A	NR
13	1466	163	NA	3:31	5/25/16	35.6409	74.7974	28	36.2542	74.5681	1	1	A	Survived
14	1467	137	NA	5:13	5/24/16	35.6967	74.7983	18	37.0169	75.2356	1	1	A	Survived
15	1469	NA	NA	6:12	5/25/16	35.7167	74.865	26	36.9175	74.5867	1	1	A	Survived
16	1470	203	F	5:54	5/25/16	35.6807	74.7820	4	36.2389	74.4828	2	2	A	Survived
17	1471	125	NA	4:18	5/25/16	35.7517	74.8783	NR	NR	NR	1	1	A	NR
18	1472	190	M	5:54	5/25/16	35.6553	74.7361	28	58.4333	74.5861	2	1	A	Survived
19	1473	185	F	2:30	5/25/16	35.7517	74.8733	8	36.3000	74.4258	2	1	A	Survived
20	1474	177	F	6:13	5/25/16	35.6856	74.7972	12	37.6367	72.2594	2	1	A	Survived
21	1475	150	NA	4:44	5/25/16	35.7533	74.8783	10	36.6753	73.5319	1	1	A	Survived
22	1476	203	NA	6:12	5/24/16	35.6542	74.7389	0	35.6536	74.6617	2	3	A	Mortality
23	1477*	203	NA	5:48	5/25/16	35.6542	74.7389	<3	36.6383	74.6192	1	2	A	Survived
24	1478	305	F	6:38	5/25/16	35.7250	74.8633	7	38.3306	70.6875	1	2	A	Survived
25	1479	NA	NA	NA	5/25/16	NA	NA	28	38.7186	73.9261	1	1	NA	Survived
26	1480	126	NA	5:06	5/25/16	35.7653	74.7964	25	37.7186	75.6947	2	1	A	Survived
27	1481	183	NA	6:05	5/25/16	35.7217	74.8633	28	37.3575	75.5856	1	1	A	Survived
28	1485	NA	NA	4:10	5/25/16	35.7400	74.8717	5	36.2808	74.8894	4	1	A	Survived
29	1486	203	NA	3:01	5/25/16	35.6122	74.7799	28	38.0392	71.4203	1	1	A	Survived
30	1487	NA	NA	4:23	5/25/16	35.7367	74.8733	NR	NR	NR	1	2	A	NR
31	1488	164	NA	3:41	5/25/16	35.6372	74.7956	7	36.3936	74.4014	1	1	A	Survived
32	1489*	NA	NA	2:49	5/24/16	35.7050	74.8283	1	36.2117	74.4014	1	1	A	Survived
33	1490	190	NA	4:02	5/24/16	35.7183	74.8517	10	37.5358	66.6225	1	1	A	Survived
34	1491	NA	NA	4:53	5/25/16	35.7437	74.8700	5	38.1408	70.9686	1	1	A	Survived
35	1492	152	NA	2:10	5/24/16	35.7017	74.8200	12	36.2967	74.1144	1	1	A	Survived
36	1493*	177	F	3:26	5/25/16	35.6242	74.7894	3	36.0461	74.5911	1	1	A	Survived
37	1495*	151	F	6:05	5/25/16	35.6655	74.7801	<3	36.8217	74.1453	2	2	A	Survived
38	1496	NA	NA	4:09	5/25/16	35.8767	75.2767	<3	36.0800	74.6189	1	1	A	Survived
39	1498	137	NA	5:10	5/25/16	35.6680	74.7830	NR	NR	NR	1	1	A	NR
40	1499	164	M	3:00	5/25/16	36.6220	74.7879	8	36.1186	74.6000	4	1	A	Survived
41	1500	177	F	5:28	5/24/16	35.6521	74.7450	10	36.5502	73.8914	2	1	A	Survived
42	1502	164	NA	3:00	5/25/16	35.6116	74.7771	28	38.1989	73.6792	2	1	A	Survived
43	1503	177	NA	2:35	5/25/16	35.6334	74.7943	28	37.6339	75.3331	3	2	A	Survived
44	1506	177	NA	3:07	5/25/16	35.6178	74.7863	28	36.9422	74.5483	4	1	A	Survived
45	1507	177	NA	5:07	5/24/16	35.7653	74.7964	NR	NR	NR	1	1	A	NR
46	1510	125	NA	6:12	5/25/16	35.7350	74.8733	8	38.0706	70.2778	1	2	A	Survived
47	1511	NA	NA	5:19	5/25/16	35.6450	74.9133	16	34.3911	76.0492	1	1	A	Survived
48	1512	177	F	2:40	5/25/16	35.6334	74.7953	6	36.3286	74.3356	3	1	A	Survived
49	1513	138	F	3:42	5/25/16	35.6372	74.7956	0	36.5656	74.3328	2	1	A	Mortality
50	1514	190	F	3:37	5/25/16	35.6356	74.7944	28	37.1264	75.6044	2	1	A	Survived

Figure 1. Deployment (n=50, circles) and pop off (n=43, triangles) locations for dusky sharks captured and tagged with Lotek PSAT LIFE satellite tags. Seven tags failed to report data.

Figure 2. Daily mean max depth plots representing 41 dusky sharks that remained alive after release and whose satellite tags transmitted a daily minimum and maximum depth. Numbers above each day post-release indicate sample size/remaining attached tags on that given date. Change in sample size at any given day represents early tag detachment. Vertical bars above each data point on a given date represent the corresponding mean and standard error.

Figures 3a and 3b. At-vessel (a) release condition and (b) and injury code of the 43 dusky shark tags that transmitted data as a function of binned soak times. Release conditions we defined as (1) swim burst, (2) strong swimming, (3) sluggish swimming, (4) sank with no visible swimming effort while injury conditions of (1) represented no visible signs of trauma to the body, (2) if minor damage was present, (3) if there were obvious signs of trauma, and (4) if the shark was moribund (see methods for full description). Columns represent cumulative percentages.

Figure 1.




Figure 2





Figure 3



Figure






Figure
[Click here to download high resolution image](#)



Figure

[Click here to download high resolution image](#)



Figure

[Click here to download high resolution image](#)

