

Using Photo-Identification and Genetic Data to Examine Fine-Scale Population Structure of Common Bottlenose Dolphins (*Tursiops truncatus*) in the Estuarine Waters Surrounding Savannah, Georgia

Emily K. Griffin,¹ Patricia E. Rosel,² Brian C. Balmer,³ Robin M. Perrtree,⁴ and Tara M. Cox⁴

¹*Center for Environmental and Human Toxicology, Department of Physiological Sciences, College of Veterinary Medicine, University of Florida, Gainesville, FL, 32611, USA*

E-mail: emilygriffin@ufl.edu

²*NOAA National Marine Fisheries Service, 646 Cajundome Boulevard, Lafayette, LA 70506, USA*

³*National Marine Mammal Foundation, 3419 Maybank Highway Suite B, Johns Island, SC 29455, USA*

⁴*Savannah State University, Department of Marine and Environmental Sciences, 3219 College Street, Savannah, Georgia 31404, USA*

Abstract

Many marine mammal species exhibit complex patterns of population structure. Specifically, common bottlenose dolphin (*Tursiops truncatus*) populations in the southeastern United States display varying degrees of spatial overlap and residency, including some that have year-round site fidelity to localized bays, sounds, and estuaries. Evidence of resident estuarine animals along the U.S. Atlantic and Gulf of Mexico coastlines is supported through photo-identification and genetic studies, although currently few studies have integrated both methods. The purpose of this project was to couple long-term photo-identification data with spatial and genetic analyses to examine population structure of common bottlenose dolphins in northern Georgia. Genomic DNA was extracted from skin samples ($n = 69$) collected in the Northern Georgia/Southern South Carolina Estuarine System Stock, and a portion of the mitochondrial DNA control region was sequenced. To determine potential fine-scale geographic delineations within this stock, the study area was split into three regions: north, buffer, and south. No significant genetic differentiation was found when regions were compared by only sample collection location. The sighting locations of sampled dolphins that had been seen ≥ 10 times between 2009–2017 ($n = 45$) were mapped in *ArcMap 10.2*. For analysis of dolphins with only ≥ 10 sightings, a significant difference in F_{ST} was found between the north versus buffer area ($p = 0.0147$). When dolphins with ≥ 10 sightings ($n = 45$) were assigned to the region where $\geq 50\%$ of their sightings were located, a significant difference in F_{ST} was found between the north versus buffer and north versus south areas ($p = 0.0018$ and $p = 0.0164$, respectively). The addition of spatial data revealed genetic division among the three regions. Thus, the combination of photo-identification and genetic analyses used in this study may be useful in determining population structure in the future.

Key Words: common bottlenose dolphin, *Tursiops truncatus*, mitochondrial DNA, photo-identification, population structure

Introduction

Marine mammals can exhibit broad distributions and ranging patterns over extended distances resulting in complex patterns of population structure (Hoelzel et al., 1998). Despite their extensive ranging capabilities, many marine mammal species display population structure on a much finer spatial scale than their full distribution might suggest (Norris, 2000). Understanding aspects of a population such as geographic range, abundance, and demographic trends is necessary for proper conservation and management (Wade & Angliss, 1997; Hoelzel et al., 1998), which relies on correctly identifying population units (Baird et al., 2009). The U.S. Marine Mammal Protection Act (MMPA) defines a stock as “a group of marine mammals of the same species or smaller taxa in a common spatial arrangement that interbreed when mature” (MMPA, 1972 [16 U.S.C 1362]). Stocks should be, under most circumstances, demographically independent populations, and the primary goal of the MMPA is to conserve stocks as a significant functioning element of the ecosystem, which includes preventing any marine mammal stock from dropping below its optimum sustainable population level (Wade & Angliss, 1997). Therefore, accurately identifying demographically independent populations is a critical task for marine mammal management (Wade & Angliss, 1997).

An integrative approach that combines genetics with spatial data collected through photo-identification can provide additional insight into population structure. Evidence of such an approach has been exhibited in several marine mammal species including the short-finned pilot whale (*Globicephala macrorhynchus*; Alves et al., 2013) and the southern right whale (*Eubalaena australis*; Carroll et al., 2011). Torres et al. (2003) examined the distribution and overlap of two common bottlenose dolphin (*Tursiops truncatus*) ecotypes and found that combining spatial data and genetics can provide insight into the distribution of overlapping populations. Areas utilized only by the coastal ecotype or only by the offshore ecotype of common bottlenose dolphins were identified using spatial and molecular analyses (Torres et al., 2003). Furthermore, evidence of habitat partitioning by common bottlenose dolphins between estuarine and bordering coastal waters has been found through photo-identification (Fazioli et al., 2006) and genetic studies (Sellas et al., 2005) in the northern Gulf of Mexico. Consequently, the combination of spatial analyses with genetic data can provide evidence of multiple, demographically independent populations of common bottlenose dolphins (Rosel et al., 2017a).

Fine Scale Population Structure of Common Bottlenose Dolphins

Common bottlenose dolphins in the southeastern United States exhibit varying degrees of residency, including small-localized bay, sound, and estuary (BSE) populations (Shane, 1980; Wells et al., 1987; Scott et al., 1990). Evidence of residency in estuarine populations along the U.S. Atlantic and Gulf of Mexico coastlines is supported through photo-identification (Wells et al., 1980; Zolman, 2002; Mazzoil et al., 2005; Balmer et al., 2008; Balmer et al., 2014), genetic (Sellas et al., 2005; Rosel et al., 2009), and telemetry studies (Wells et al., 2017; Balmer et al., 2018,2018b). Multiple populations of BSE common bottlenose dolphins exist within the complex system of salt marshes and bays along the eastern coast of the U.S. (Hayes et al., 2017). The National Marine Fisheries Service (NMFS) has defined the Northern Georgia/Southern South Carolina Estuarine System (NGSSCES) Stock of common bottlenose dolphins as residing in the estuarine waters between the North Edisto River in South Carolina and northern Ossabaw Sound in Georgia. NMFS considers the NGSSCES Stock of common bottlenose dolphins to be a strategic stock because the estimated abundance of the stock is small, and few mortalities would surpass the potential biological removal level (Hayes et al., 2017). Based on recent mark-recapture data, it has been postulated that the current southern boundary of the NGSSCES Stock needs to be modified, and animals residing from the Savannah River to the Wilmington River may be distinct from the animals residing between the Wilmington River and northern Ossabaw Sound (Thompson et al., 2017). The incomplete understanding of the region inhabited by the NGSSCES Stock and the movement patterns of these dolphins make it difficult to implement long-term sustainability plans for this protected species and strategic stock.

Many BSE populations in the southeastern U.S. have minimal to no genetic and photo-identification data, which are the primary and supplemental type of data, respectively, used to identify population structure (Sellas et al., 2005; Rosel et al., 2009; Litz et al., 2012; Rosel 2017a). Using the geographic location where the sample was collected is the most conservative method of estimating where an individual dolphin may reside and/or identifying with which population the animal is associated. However, incorporation of long-term sighting histories into sampling design for genetic studies is a unique method and incorporates multiple types of data into population structure analyses. Combining multiple methods to investigate population structure has been found to be fruitful (Torres et al., 2003; Balmer et al., 2011). A study investigating residency patterns of common bottlenose dolphins in the Stono River estuary in South Carolina using photo-identification has been used to suggest that incorporating genetics could clarify population discreteness in future studies (Zolman, 2002). However, there have been few published studies combining long-term sighting histories and genetic data to examine population structure (Carroll

et al., 2011; Alves et al., 2013). The ability to integrate long-term sighting data with genetic information is a novel approach that may allow for a more comprehensive assessment of common bottlenose dolphin population structure in the southeastern U.S., and have potential applications for other marine mammal populations globally. The purpose of this study was to integrate multiple tools—long-term photo-identification and genetic data—to examine population structure of common bottlenose dolphins in the estuarine waters surrounding Savannah, GA.

Methods

The study area consisted of the estuarine waters south of Savannah, GA from the southern Savannah River channel to northern Ossabaw Sound. This area comprised the southern range of the NGSSCES Stock extending into the northernmost region of the Central Georgia Estuarine System (CGES) Stock (Hayes et al., 2017), and covered approximately 400 km² of small creeks, salt marshes, larger rivers, and sounds (Figure 1; defined in Perrtree et al., 2014). Photo-identification surveys were conducted from April 2009 to June 2015. Surveys were conducted at an on-effort speed of 33-40 km/hr along previously determined transects (place Figure 2 here) (Figure 2; methods reviewed in Perrtree et al., 2014). The dorsal fin of each dolphin within a group was photographed using digital single lens reflex (DSLR) cameras (Cannon EOS 40D and Nikon D90) with 70-300 mm or 70-400 mm zoom lenses. Dolphins were considered a group if they were within 100 m of each other, moving in the same direction, and engaging in similar behaviors (Shane, 1990). Images were matched to images collected from previous survey efforts, and a catalog was created for the Savannah area. All dolphins entered into the Savannah area catalog were given a unique numerical identification code. A total of 508 cataloged individuals are in this database.

To identify animals to target for remote biopsy, sighting history data for each individual dolphin were examined from the long-term photo-identification database for this study area. Cataloged individuals with 10 or more sightings were identified as high priority biopsy targets based upon the minimum number of sightings required for home range estimation by Urian et al. (2009). Individuals with less than 10 sightings were considered moderate priority targets. Neonates, young calves, and their mothers were not considered for biopsy. Individuals were identified as neonates if fetal folds or fetal lines were present, if the individual was less than half the size of an adult, or if the individual was observed swimming in the echelon position (Shane, 1990; Mann and Smuts, 1999; Thayer et al., 2003; McFee et al., 2014). Dolphins with non-distinctive fins were considered low priority targets because there likely were less sighting history data available for these individuals. Sighting locations for each dolphin with 10 or more sightings were mapped in *ArcGIS 10.2* (ESRI, Redlands, CA USA). The study area was

categorized into three regions: a) a northern region that consisted of the estuarine waters from the Savannah River to the Wilmington River; b) a buffer region that included the Wilmington River and Wassaw Sound; and c) a southern region that included the area between the Wilmington River and northern Ossabaw Sound (Figure 2). The regions were categorized based on previous photo-identification research (Thompson et al., 2017) which indicated that these regions could contain two separate populations with the buffer region acting as an area of overlap. The number of sightings in the north region and south region, as well as the number of sightings within small creeks, were counted and recorded for every individual with ≥ 10 sightings. Small creeks included all creeks within the study area excluding Wassaw and Ossabaw Sounds, the Savannah River, the Wilmington River, Vernon River, and portions of the Bull River and Oddingsell River (Figure 2).

To properly examine BSE populations only, it was necessary to exclude coastal animals—animals that live primarily in coastal waters and not the BSE waters—from the data set. Thus, criteria were developed to ensure individuals biopsied were likely from the resident NGSSCES Stock and not the adjacent coastal stock, whose members are known to make short-term (transient) forays into estuarine waters, particularly at the larger openings of the bays to coastal waters (Balmer et al., 2018b). Of the dolphins with 10 or more sightings, animals with 50% or more of their sightings within small creeks were identified. Identifying animals with a majority of their sightings in small creeks provided a greater chance that the target animals were truly estuarine dolphins. In addition, common bottlenose dolphins residing outside of estuarine environments are thought to enter estuaries primarily during warm summer months (Speakman et al., 2010; Balmer et al., 2013); thus, samples were collected in non-summer months: February, March, and September. Furthermore, dolphins with 50% or more of their sightings occurring in the north or south region and 15% or less occurring in the opposite region were identified as target animals for biopsy. These metrics were formulated with the goal of maximizing the percentage of sightings in the target area while minimizing the percentage of sightings in the non-target area, while also isolating small creek animals from coastal animals.

Remote biopsy sampling is a common tool for collecting tissue samples for genetic analyses (Sinclair et al., 2015). Remote biopsy sampling was conducted in September 2015 and February and March 2017. A total of 50 skin samples were collected using a crossbow and dart fitted with a 10 x 25 mm stainless steel sampling tip. The sampling tip contained a sharpened cutting edge angled inward to hold the sample in place. The size and type of sampling tip was chosen based on methods described in Sinclair et al. (2015). The crossbow was equipped with a

retainer to hold the dart in place and a battery powered sight to increase accuracy. The same digital DSLR cameras utilized during photo-identification surveys were used during remote biopsy sampling. When a group of dolphins was sighted, the group was approached to determine if an animal suitable for biopsy was present. Known target individuals identified using an onboard photo-identification catalog were prioritized for sampling. However, if no known dolphins were within the group, dolphins with unidentified yet distinctive fins were considered for biopsy. Dolphins with unidentified fins were only sampled if found in a small creek away from the open ocean to improve the chances that the sample was from an estuarine dolphin.

Sampling was attempted when the dolphin was perpendicular to the sampler and 4-10 m from the boat (Krützen et al., 2002). The sample was collected from below the dorsal fin on either side of the animal. After the dart came into contact with the animal and a sample was obtained, the dart floated near the surface and was retrieved. The sample was then removed from the sampling tip and cut into five separate pieces. One section—half of the skin—was placed directly into a 5 mL 20% DMSO/saturated NaCl vial for later genetic analysis and was stored at room temperature. Additionally, skin samples from stranded animals ($n = 19$) were provided by the Georgia Department of Natural Resources. These samples were collected using a scalpel blade to remove a small portion of skin from the body of the animal below the dorsal fin and placed into 5 mL 20% DMSO/saturated NaCl vials and stored at room temperature. Stranding samples were collected from April 2008–May 2017.

Genomic DNA was extracted from the skin samples using a Qiagen DNeasy Blood and Tissue extraction kit. Upon completion of extraction, the DNA was stored at 4°C. The quality of the extracted DNA was examined by gel electrophoresis. The quantity of DNA was measured via fluorometry on a Hoefer DyNA Quant 200 fluorometer. Extracted DNA was diluted to 25 ng/μl, but samples with less than 30 ng/μl DNA were not diluted. A 500 bp fragment of the 5' end of the mitochondrial DNA (mtDNA) control region and adjacent tRNAs were amplified using the polymerase chain reaction (PCR) with the primers L15824 (5'-CCTCACTCCTCCCTAAGACT-3') (Rosel et al., 1999) and H16498 (5'-CCTGAAGTAAGAACCA GATG-3') (Rosel et al., 1994). Extracted genomic DNA (25 ng) was added to a 24 μl PCR reaction mix that included 20 mM Tris HCl, 50 mM KCl, 1.5 mM MgCl₂, 0.3 μM of each primer, 150 μM dNTPs, and 5 U Taq DNA polymerase (Thermo Fisher Scientific, Waltham, MA USA). For samples with a DNA concentration less than 25 ng/μl, 0.24 mg ml⁻¹ of Bovine Serum Albumin (BSA) was added. The cycling profile was an initial cycle at 95°C for 30 seconds, followed by 30 cycles of 95°C for 30 seconds, 55°C for 30 seconds, and 72°C for 30 seconds. After 30

cycles the PCR continued at 72°C for 7 minutes and then was held at 4°C. PCR product was run on a 0.8% agarose gel to test the success of the reaction and, if successful, the remaining product was purified using a 0.8% low melting point agarose gel. Each band was excised from the gel under UV light with a razor blade and placed in a labeled 1.5 ml tube and digested using 2.5 U of agarase (Thermo Fisher Scientific) and held at 40°C overnight. A cycle sequencing PCR reaction was then prepared using the original primers, 5X buffer, and BigDye v1.1 dye terminator mix (Applied Biosystems, Foster City, CA USA). The cycling profile was an initial 96°C for one minute, followed by 25 cycles of 96°C for 10 seconds, 50°C for 5 seconds, 60°C for 4 minutes, and then held at 4°C. The cycle sequencing products were cleaned with Sephadex®, completely dried and stored at -20°C. The final sequencing products were loaded on an ABI 3130 Genetic Analyzer. All samples were sequenced in both forward and reverse directions. Sequences were edited and aligned creating consensus sequences using *Geneious 10.2.3*. Consensus sequences were aligned with a master alignment to determine haplotypes.

Nucleotide diversity (π) and haplotype diversity (h ; Nei, 1987), which measure the degree of polymorphism within a population, were estimated using *ARLEQUIN 3.5.2.2* (Excoffier & Lischer, 2010). Population differentiation, F_{ST} and Φ_{ST} were estimated in *ARLEQUIN* using an analysis of molecular variance (AMOVA) framework. Significance values were determined using 10,000 permutations. The program jModelTest (Guignon & Gascuel, 2003; Darriba et al., 2012) was used to determine the best evolutionary model for estimating Φ_{ST} . The Tamura & Nei (1993) model with a gamma correction ($\alpha = 0.5$) was used for the distance matrix in the analyses. F_{ST} and Φ_{ST} were calculated for regions north versus buffer, north versus south, and buffer versus south. The Bonferroni false discovery test was used to correct probability values for multiple comparisons.

Data were first analyzed based on the location where each sample was collected; F_{ST} and Φ_{ST} values were calculated comparing all samples collected in the north, buffer, or south region. Then, data were analyzed and F_{ST} and Φ_{ST} values were calculated based on sample collection location for only animals with ≥ 10 sightings. Lastly, based on sighting history data, samples from dolphins with ≥ 10 sightings were assigned to the region where 50% or more of the animal's sightings were located and F_{ST} and Φ_{ST} values were calculated. In summary, analyses were first conducted on all the animals based on sample collection location and then with the inclusion of sighting history data to observe how the addition of photo-identification data may have helped characterize population structure.

Results

A total of 69 skin samples for genetic analysis were collected in the estuarine waters surrounding Savannah, Georgia. Fifty samples were collected through biopsy efforts, and 19 samples were obtained from stranded animals. Twenty-one samples were collected from the north region, 14 samples were collected from the buffer region, and 28 samples were collected from the south region. Six samples were collected outside of the study area on Tybee Beach and upstream from the Savannah River and Ogeechee River. Samples collected outside of the study area were obtained from stranded individuals and were included into the closest study region from where the sample was collected for analyses. Fifty-eight samples were matched to known dolphins in the Savannah study area catalog, and 45 of those had 10 or more sightings.

Five haplotypes were identified from all 69 samples. All haplotypes were associated with the western North Atlantic coastal ecotype (Rosel et al. 2009). Four animals were identified as Ttr1, 4 were Ttr4, 2 were Ttr5, 43 were Ttr6, and 16 were identified as Ttr9 (haplotype names from Rosel et al. 2009) (place Table 1 here; place Figure 3 here) (Table 1; Figure 3). The most common haplotype, Ttr6, was found in all three study regions. The next most frequent haplotype, Ttr9, was also found in all three regions. Haplotype Ttr1 was found only in the north and south regions. Ttr4 was found only in the north region, and Ttr5 was found only in the buffer and south regions. The distribution of each haplotype is shown in Table 1. Haplotype and nucleotide diversity for all 69 samples based on sample collection location is shown in Table 2a, for animals with ≥ 10 sightings based on sample collection location is shown in Table 2b, and for animals with ≥ 10 sightings assigned to the location where $\geq 50\%$ of their sightings were located is found in Table 2c (place Table 2 here).

Genetic differentiation (F_{ST} and Φ_{ST}) was estimated for north versus south, north versus buffer, and buffer versus south utilizing three different data analyses. First, F_{ST} and Φ_{ST} were estimated for the sample collection location of all 69 samples, and no significant differences were observed for any comparison after Bonferroni correction (Table 3a) (place Table 3 here). Second, F_{ST} and Φ_{ST} were estimated based on the sampling collection location of only animals with ≥ 10 sightings ($n = 45$); only the estimate of F_{ST} between the north and buffer regions differed significantly from zero (Table 3b). A significant difference was found for F_{ST} between the north versus buffer and north versus south when the samples were assigned to location by sighting history rather than sample collection location (Table 3c). In this case, the percentage of sightings in each of the three regions was calculated for each of the 45 individuals, and they were assigned to the region where 50% or more of their sightings were found.

Discussion

In contrast to using sampling collection location alone as a proxy for regional affiliation where no significant differences were seen among regions, a significant difference in F_{ST} was found between the north region and the buffer region when using animals with ≥ 10 sightings in the analyses, and a significant difference was found between the north and the south regions with the inclusion of sighting history information. Thus, the addition of photo-identification data revealed a significant difference between the north and buffer region and the north and south region that was not identified using sample collection location alone. However, no difference was detected between the buffer and south region with any of the three data partitions. It is possible that instead of the Wilmington River being a buffer region, the area is alternatively a part of the southern region and both are separate from the north, thus creating two study regions instead of three. If true, the border of the NGSSCES and CGES Stocks might be more appropriate at the Wilmington River. Also, the sample size from the buffer region was the lowest, and this may have had an effect on the power to detect significant differences.

Another possible interpretation of the results from this study is that the larger tributaries, such as the Wilmington River (buffer region), are an area of BSE and coastal population overlap. An overlap of populations and sampling in these areas may explain why no genetic differentiation was found without the inclusion of photo-identification data. Balmer et al. (2013) suggested that based on photo-identification data from southern Georgia, dolphins from the western North Atlantic South Carolina/Georgia Coastal Stock were sighted primarily in larger tributaries, and these were potentially areas of shared habitat between the BSE and adjacent coastal population(s). This hypothesis was further strengthened by satellite telemetry data which demonstrated higher overlap between BSE and coastal dolphins in the larger tributaries and sounds of southern Georgia (Balmer et al., 2018b). Additionally, some common bottlenose dolphin populations are known to exhibit seasonal changes in distribution (Speakman et al., 2010). In some locations, coastal animals travel into bays, sounds, and estuaries in the warmer months (Speakman et al., 2010; Balmer et al., 2013). Remote biopsy sampling was conducted in February, March, and September, which are considered part of the winter and fall seasons, and samples collected from stranded animals were collected year-round. The animals biopsied in the buffer region may actually be coastal animals that had traveled inshore or be a mixture of coastal animals and estuarine residents. However, this would indicate that coastal animals travel into the bays in the cooler seasons as well, in contrast to the findings of Speakman et al. (2010) and Balmer et al. (2013). The inclusion of coastal animals provides a possible explanation for the lack of a significant difference found using only sample collection location. Without the inclusion of photo-identification

data, a mixture of coastal and estuarine animals may have been used in the analysis. Genetic analyses that include animals from the coastal stock may help sort out these different hypotheses. In addition, to fully compare the genetic differences among locations examined in this study, nuclear DNA analysis, such as, microsatellites, should be included. Unlike mtDNA, microsatellites are inherited from both parents and may provide a more thorough examination of relatedness and gene flow (Rosel et al., 2017b). Future studies should include microsatellite data or other nuclear data in addition to mtDNA to acquire a more comprehensive assessment of population structure and determine whether there are multiple demographically independent populations within the NGSSCES Stock. Nuclear microsatellite data would also allow examination of relatedness among the samples and ensure estimates of differentiation among regions are not influenced by groups of highly related individuals.

The dolphins in the Savannah area are known to interact with recreational and commercial fishing vessels (Perrtree et al., 2014). These human-interaction behaviors in the Savannah area were found at a much higher rate than in other areas with known human-interaction behaviors (Perrtree et al., 2014). Interactions between dolphins and humans can lead to serious injury (Powell et al., 2018; Balmer et al., 2019), mortality (McFee et al., 2006; Donaldson et al., 2010), and behavioral changes (Mann & Smuts, 1999; Orams, 2002; Hazelkorn et al., 2016). There is currently no abundance estimate nor estimate of Potential Biological Removal for the NGSSCES Stock; yet, a recent Atlantic Bottlenose Dolphin Take Reduction Team noted that the common bottlenose dolphins in the Savannah area may be close to exceeding PBR due to crab pot entanglements. With little known about the dolphins in the Savannah area coupled with the threat of cumulative anthropogenic impacts, it is imperative to use a multi-faceted approach, such as the combination of genetics and photo-identification, to better understand the population structure and thus the stock delineations of the NGSSCES. The inclusion of spatial data may give a more accurate representation of population structure and genetic make-up of a population. Therefore, methods in this study could be utilized by future studies in which long-term photo-identification data are available to improve our understanding of the complex population structure of common bottlenose dolphins and other marine mammals.

Acknowledgements

This research was supported by the National Oceanic and Atmospheric Administration through the Living Marine Resources Cooperative Science Center (LMRCSC) NA16SEC4810007 project #17- 04. We thank the Georgia Department of Natural Resources, in particular Clay George, Trip Kolkmeier, and Kate Sparks Sabbe, for providing

Fine Scale Population Structure of Common Bottlenose Dolphins

stranding samples. Zoe Wong, funded by NSF REU OCE-9402526, helped design the study; Zach Swaim collected biopsy samples in 2017. We also thank Lynsey Wilcox, Charlotte Sprehn, and Ana Costa for their time and assistance with all genetic analyses. Savannah State University Dolphin Science Laboratory data collected by past undergraduates and graduates funded by NOAA LMRCSC (NA05OAR4811017, NA16SEC4810007, and Environmental Entrepreneurship Program), EDGE (NSF grant #0914680), Title VII (Dept. of Ed. P382G090003), NSF GK12 (award # 0841372) and REU (NSF OCE-REU 1156525, 1460457), and Oak Foundation with boat time provided by the Savannah State University Marine Sciences Program. Data were collected in accordance with the Marine Mammal Protection Act under NMFS Letters of Confirmation #14219 and #18605 issued to Dr. Tara Cox. Remote biopsy sampling was conducted under permit #14450-01 issued to NMFS Southeast Fisheries Science Center. The scientific results and conclusions, as well as any views or opinions expressed herein, are those of the author(s) and do not necessarily reflect those of NOAA or the Department of Commerce.

Literature Cited

- Alves, F., Qué, S., Dinis, A., Nicolau, C., Ribeiro, C., Freitas, L., Kaufmann, M., & Fortuna, C. (2013). Population structure of short-finned pilot whales in the oceanic archipelago of Madeira based on photo-identification and genetic analyses: implications for conservation. *Aquatic Conservation*, 23(5), 758-776. <https://doi.org/10.1002/aqc.2332>
- Baird, R. W., Gorgone, A. M., McSweeney, D. J., Ligon, A. D., Deakos, M. H., Webster, D. L., ... Mahaffy, S. D. (2009). Population structure of island-associated dolphins: evidence from photo-identification of common bottlenose dolphins (*Tursiops truncatus*) in the main Hawaiian Islands. *Marine Mammal Science*, 25, 251-274. <https://doi.org/10.1111/j.1748-7692.2008.00257.x>
- Balmer, B. C., Wells, R. S., Nowacek, S. M., Nowacek, D. P., Schwacke, L. H., McLellan, W. A., ... Pabst, D. A. (2008). Seasonal abundance and distribution patterns of common bottlenose dolphins (*Tursiops truncatus*) near St. Joseph Bay, Florida, USA. *Journal of Cetacean Research and Management*, 10(2), 157-167.
- Balmer, B. C., Schwacke, L. H., Wells, R. S., George, R. C., Hoguet, J., Kucklick, J. R., ... Pabst, D. A. (2011). Relationship between persistent organic pollutants (POPs) and ranging patterns in common bottlenose dolphins (*Tursiops truncatus*) from coastal Georgia, USA. *Science of the Total Environment*, 409(11), 2094-2101. <https://doi.org/10.1016/j.scitotenv.2011.01.052>
- Balmer, B. C., Schwacke, L. H., Wells, R. S., Adams, J. D., George, R. C., Lane, S. M., ... Pabst, D. A. (2013). Comparison of abundance and habitat usage for common bottlenose dolphins between sites exposed to differential anthropogenic stressors within the estuaries of Southern Georgia, USA. *Marine Mammal Science*, 29(2), 114-135. <https://doi.org/10.1111/j.1748-7692.2012.00589.x>
- Balmer, B. C., Wells, R. S., Schwacke, L. H., Schwacke, J. H., Danielson, B., George, R. C., ... Zolman, E. S. (2014). Integrating multiple techniques to identify stock boundaries of common bottlenose dolphins (*Tursiops truncatus*). *Aquatic Conservation: Marine Freshwater Ecosystems*, 24(4), 511-521. <https://doi.org/10.1002/aqc.2357>
- Balmer, B. C., McDonald T., Hornsby, F., Adams, J., Allen, J., Barleycorn, A., ... Schwacke, L. (2018). Long-term trends in a northern Gulf of Mexico common bottlenose dolphin (*Tursiops truncatus*) population in the wake of the Deepwater Horizon oil spill. *Journal of Cetacean Research and Management*, 18, 1-9.
- Balmer, B. C., Zolman, E., Rowles, T., Smith, C., Townsend, F., Fauquier, D., ... Schwacke, L. (2018b). Ranging patterns, spatial overlap, and association with dolphin morbillivirus exposure in common bottlenose dolphins (*Tursiops truncatus*) along the Georgia, USA coast. *Ecology and Evolution*, 8, 12890-12904. <https://doi.org/10.1002/ece3.4727>
- Balmer, B. C., Zolman, E., Bolton, J., Fauquier, D., Fougères, E., George, R. C., ... Ylitalo, G. (2019). Ranging patterns and exposure to cumulative stressors of a *Tursiops truncatus* (Common Bottlenose Dolphin) in Georgia. *Southeastern Naturalist*, 18(1), page numbers?. <https://doi.org/10.1656/058.018.0112>
- Carroll, E. L., Patenaude, N. J., Childerhouse, S. J., Kraus, S. D., Fewster, S. D., & Baker, C. S. (2011). Abundance of the New Zealand subantarctic southern right whale population estimated from photo-identification and genotype mark-recapture. *Marine Biology*, 158(11), 2565-2575. <https://doi.org/10.1007/s00227-011-1757-9>
- Darriba, D., Tboada, G. L., Doallo, R., & Posada, D. (2012). JModelTest 2: more models, new heuristics and parallel computing. *Nature Methods*, 9(8), 772. <https://doi.org/10.1038/nmeth.2109>
- Donaldson, R. H., Finn, H., & Calver, M. (2010). Illegal feeding increases risk of boat-strike and entanglement in bottlenose dolphins in Perth, Western Australia. *Pacific Conservation Biology*, 16(3), 157-161. <https://doi.org/10.1071/pc100157>

Fine Scale Population Structure of Common Bottlenose Dolphins

- Excoffier, L., & Lischer, H. E. L. (2010). Arlequin suite version 3.5: A new series of programs to perform population genetics analyses under Linux and Windows. *Molecular Ecology Resources*, 10(3), 564-567. <https://doi.org/10.1111/j.1755-0998.2010.02847.x>
- Fazioli, K. L., Hoffman, S., & Wells, R. S. (2006). Use of Gulf of Mexico coastal waters by distinct assemblages of bottlenose dolphins (*Tursiops truncatus*). *Aquatic Mammals*, 32(2), 212-222. <https://doi.org/10.1578/am.32.2.2006.212>
- Guidon, S., & Gascuel, O. (2003). A simple, fast and accurate method to estimate large phylogenies by maximum-likelihood. *Systematic Biology*, 52(5), 696-704. <https://doi.org/10.1080/10635150390235520>
- Hayes, S. A., Josephson, E., Maze-Foley, K., & Rosel, P. E. (2017). US Atlantic and Gulf of Mexico marine mammal stock assessments – 2016. NOAA Technical Memorandum NMFS-NE-241. pp. 274.
- Hazelkorn, R. C., Schulte, B. A., & Cox, T. (2016). Persistent effects of begging on common bottlenose dolphin (*Tursiops truncatus*) behavior in an estuarine population. *Aquatic Mammals*, 42(4), 531-541. <https://doi.org/10.1578/am.42.4.2016.531>
- Hoelzel, A. R., Potter, C. W., & Best, P. B. (1998). Genetic differentiation between parapatric “nearshore” and “offshore” populations of the bottlenose dolphin. *Proceedings of the Royal Society of London B*, 265(1402), 1177-1183. <https://doi.org/10.1098/rspb.1998.0416>
- Krützen, M., Barre, L. M., Möller, L. M., Heithaus, M. R., Simms, C., & Sherwin, W. B. (2002). A biopsy system for small cetaceans: darting success and wound healing in *Tursiops* spp. *Marine Mammal Science*, 18(4), 385-394. <https://doi.org/10.1111/j.1748-7692.2002.tb01078.x>
- Litz, J. A., Hughes, C. R., Garrison, L. P., Fieber, L. A., Rosel, P. E. (2012). Genetic structure of common bottlenose dolphins (*Tursiops truncatus*) inhabiting adjacent South Florida estuaries – Biscayne Bay and Florida Bay. *Journal of Cetacean Research and Management*, 12(1), 107-117.
- Mann, J., & Smuts, B. B. (1999). Behavioral development in wild bottlenose dolphin newborns (*Tursiops* sp.). *Behavior*, 136(5), 529-566. <https://doi.org/10.1163/156853999501469>
- Marine Mammal Protection Act (MMPA). (1972). U.S. Marine Mammal Protection Act, amended 2007. 16 U.S.C. 1361 et seq., 1401-1407, 1538, 4107.
- Mazzoil, M., McCulloch, S. D., & Degran, R. H. (2005). Observations on the site fidelity of bottlenose dolphins (*Tursiops truncatus*) in the Indian River Lagoon, Florida. *Florida Scientist*, 68(4), 217-226.
- McFee, W. E., Hopkins-Murphy, S. R., & Schwacke, L. H. (2006). Trends in bottlenose dolphin (*Tursiops truncatus*) strandings in South Carolina, USA, 1997-2003: implications for the Southern North Carolina and South Carolina Management Units. *Journal of Cetacean Research and Management*, 8(2), 195-201.
- McFee, W. E., Speakman, T. R., Balthis, L., Adams, J. D., & Zolman, E. S. (2014). Reproductive seasonality of a recently designated bottlenose dolphin stock near Charleston, South Carolina, U.S.A. *Marine Mammal Science*, 30(2), 528-543. <https://doi.org/10.1111/mms.12055>
- Nei, M. (1987). *Molecular Evolutionary Genetics* (1st ed.) New York, New York: Columbia University Press.
- Norris, R.D. (2000). Pelagic species diversity, biogeography, and evolution. *Paleobiology*, 26(s4), 236-258. <https://doi.org/10.1017/s0094837300026956>
- Orams, M. B. (2002). Feeding wildlife as tourism attraction: a review of issues and impacts. *Tourism Management*, 23(3), 281-293. [https://doi.org/10.1016/s0261-5177\(01\)00080-2](https://doi.org/10.1016/s0261-5177(01)00080-2)

Fine Scale Population Structure of Common Bottlenose Dolphins

- Perrtree, R. M., Kovacs, C. J., & Cox, T. M. (2014). Standardization and application of metrics to quantify human-interaction behaviors by the bottlenose dolphin (*Tursiops spp.*) *Marine Mammal Science*, 30(4), 1320-1334. <https://doi.org/10.1111/mms.12114>
- Powell, J. R., Machernis, A. F., Engleby, L. K., Farmer, N. A., & Spradlin, T. R. (2018). Sixteen years later: an updated evaluation of the impacts of chronic human interactions with bottlenose dolphins (*Tursiops truncatus*) at Panama City, Florida, USA. *Journal of Cetacean Research and Management*, 19, 79-93.
- Rosel, P. E., Dizon, A. E., & Heyning, J. E. (1994). Genetic analysis of sympatric morphotypes of common dolphins (genus *Delphinus*). *Marine Biology*, 119(2), 159-167. <https://doi.org/10.1007/bf00349552>
- Rosel, P. E., France, S. C., Wang, J., & Kocher, T. D. (1999). Genetic structure of harbor porpoise *Phocoena phocoena* populations in the Northwest Atlantic based on mitochondrial and nuclear markers. *Molecular Ecology*, 8(1), 541-554. <https://doi.org/10.1046/j.1365-294x.1999.00758.x>
- Rosel, P. E., Hansen, L., & Hohn, A. A. (2009). Restricted dispersal in a continuously distributed marine species: common bottlenose dolphins, *Tursiops truncatus*, in coastal waters of the western North Atlantic. *Molecular Ecology*, 18(24), 5030-5045. <https://doi.org/10.1111/j.1365-294x.2009.04413.x>
- Rosel, P. E., Wilcox, L., Sinclair, C., Speakman, T., Tumlin, M., Litz, J., & Zolman, E. (2017a). Genetic assignment to stock of stranded common bottlenose dolphins in Southeastern Louisiana after the Deepwater Horizon oil spill. *Endangered Species Research*, 33(1), 221-234. <https://doi.org/10.3354/esr00780>
- Rosel, P. E., Taylor, B. L., Hancock-Hanser, B. L., Morin, P. A., Archer, F. I., Lang, A. R., . . . Martien, K. K. (2017b). A review of molecular genetic markers and analytical approaches that have been used for delimiting marine mammal subspecies and species. *Marine Mammal Science*, 33(1), 56-75. <https://doi.org/10.1111/mms.12412>
- Scott, G. P., Irvine, A., & Wells, R. S. (1990). A long-term study of bottlenose dolphins on the west coast of Florida. In S. Leatherwood & R. R. Reeves (Eds.), *The Bottlenose Dolphin* (1st ed., pp 35-244). San Diego, California: Academic Press.
- Sellas, A., Wells, R. S., & Rosel, P. E. (2005). Mitochondrial and nuclear DNA reveal fine scale geographic structure in bottlenose dolphins (*Tursiops truncatus*) in the Gulf of Mexico. *Conservation Genetics*, 6(5), 715-728. <https://doi.org/10.1007/s10592-005-9031-7>
- Shane, S. H. (1980). Occurrence, movements and distribution of bottlenose dolphin, *Tursiops truncatus*, in Southern Texas. *Fisheries Bulletin*, 78(3), 593-601.
- Shane, S. H. (1990). Behavior and ecology of the bottlenose dolphin at Sanibel Island, Florida. In S. Leatherwood & R. R. Reeves (Eds.), *The Bottlenose Dolphin* (1st ed., pp245-265). San Diego, California: Academic Press.
- Sinclair, C., Sinclair, J., Zolman, E., Martinez, A., Balmer, B. C., & Barry, K. (2015). Remote biopsy sampling field procedures for cetaceans used during the natural resource damage assessment of the MSC252 *Deepwater Horizon* oil spill. NOAA Technical Memorandum NMFS-SEFSC-670. 50.
- Speakman, T. R., Lane, S. M., Schwacke, L. H., Fair, P. A., & Zolman, S. (2010). Mark-recapture estimates of seasonal abundance and survivorship for bottlenose dolphins (*Tursiops truncatus*) near Charleston, South Carolina, USA. *Journal of Cetacean Research and Management*, 11(2), 153-162.
- Tamura, K., & Nei, M. (1993). Estimation of the number of nucleotide substitutions in the control region of mitochondrial DNA in Humans and Chimpanzees. *Molecular Biology and Evolution*, 10(3), 512-526. <https://doi.org/10.1093/oxfordjournals.molbev.a040023>

Fine Scale Population Structure of Common Bottlenose Dolphins

- Thayer, V. G., Read, A. J., Friedlaender, A. S., Colby, D. R., Hohn, A. A., McLellan, W. A., . . . Rittmaster, K. A. (2003). Reproductive seasonality of western Atlantic bottlenose dolphins off North Carolina, U.S.A. *Marine Mammal Science*, 19(4), 617-629. <https://doi.org/10.1111/j.1748-7692.2003.tb01120.x>
- Thompson, J., Cox, T. M., & Conroy, M. (2017, October). Estimation of site fidelity for common bottlenose dolphins *Tursiops truncatus* in a complex estuarine system using the Robust Design and multistate analysis. *The Society for Marine Mammalogy's 22nd Biennial Conference on the Biology of Marine Mammals*, Halifax, Nova Scotia, Canada.
- Torres, L. G., Rosel, P. E., D'Agrosa, C., & Read, A. J. (2003). Improving management of overlapping bottlenose dolphin ecotypes through spatial analyses and genetics. *Marine Mammal Science*, 19(3), 502-514. <https://doi.org/10.1111/j.1748-7692.2003.tb01317.x>
- Urian, K. W., Hofmann, S., Wells, R. S., & Read, A. J. (2009). Fine-scale population structure of bottlenose dolphins (*Tursiops truncatus*) in Tampa Bay, Florida. *Marine Mammal Science*, 25(3), 619-638. <https://doi.org/10.1111/j.1748-7692.2009.00284.x>
- Wade, P. R., & Angliss, R. P. (1997). Guidelines for assessing marine mammal stocks: report of the GAMMS workshop April 3-5, 1996, Seattle, Washington. NOAA Technical Memorandum NMFS-OPR-4. 121.
- Wells, R. S., Irvine, A. B., & Scott, M. D. (1980). The social ecology of inshore odontocetes. In L. Herman (Ed.), *Cetacean Behavior: Mechanisms and Functions* (1st ed., pp 263-317). New York, New York: John Wiley and Sons.
- Wells, R. S., Scott, M. D., & Irvine, A. B. (1987). The social structure of free-ranging bottlenose dolphins. In H. H. Genoways (Ed.) *Current Mammalogy* (1st ed., pp 247-305). New York, New York: Plenum Press.
- Wells, R. S., Schwacke, L. H., Rowles, T. K., Balmer, B. C., Zolman, E., Speakman, T., . . . (2017). Ranging patterns of common bottlenose dolphins *Tursiops truncatus* in Barataria Bay, Louisiana, following the Deepwater Horizon oil spill. *Endangered Species Research*, 33, 159-180. <https://doi.org/10.3354/esr00732>
- Zolman, E. S. (2002). Residence patterns of bottlenose dolphins (*Tursiops truncatus*) in the Stono river estuary, Charleston County, South Carolina, U.S.A. *Marine Mammal Science*, 18(4), 879-892. <https://doi.org/10.1111/j.1748-7692.2002.tb01079.x>

Table 1. Mitochondrial DNA control region haplotypes found among 69 common bottlenose dolphins (*Tursiops truncatus*) sampled in the estuarine waters of Savannah, Georgia. Total number of individuals with each haplotype, the percentage of each haplotype in the sample population, and the number of individuals with each haplotype exhibited in the north, buffer, and south regions.

Haplotype	Number of Individuals	Percentage of total sample population	Number of individuals in the north region	Number of individuals in the buffer region	Number of individuals in the south region
Ttr1	4	0.6%	2	0	2
Ttr4	4	0.6%	4	0	0
Ttr5	2	0.3%	0	1	1
Ttr6	43	62%	13	12	18
Ttr9	16	23%	7	1	8
Total	69		26	14	29

Table 2. Genetic diversity estimates in mtDNA control region sequences including number of haplotypes (H), mean haplotype diversity (h) and nucleotide diversity (π) based on a) sample collection location of all 69 samples, b) sample collection location of dolphins with ≥ 10 sightings, and c) sighting history data -- samples from dolphins with ≥ 10 sightings were assigned to the region where 50% or more of the animal's sightings were located, n sample size, SE standard error.

	n	H	h (SE)	π (SE)
a)				
All	69	5	0.6905 (0.5306)	0.0019 (0.0016)
North	26	4	0.8276 (0.6108)	0.0023 (0.0019)
Buffer	14	3	0.4285 (0.4107)	0.0012 (0.0013)
South	29	4	0.6748 (0.5313)	0.0019 (0.0016)
b)				
All	45	5	0.6767 (0.5270)	0.0019 (0.0016)
North	16	4	1.0166 (0.7182)	0.0028 (0.0022)
Buffer	11	2	0.1818 (0.2534)	0.0005 (0.0008)
South	18	3	0.7134 (0.5594)	0.0020 (0.0017)
c)				
All	45	5	0.6767 (0.5270)	0.0019 (0.0016)
North	18	4	1.0065 (0.7091)	0.0028 (0.0022)
Buffer	13	2	0.1538 (0.2286)	0.0043 (0.0007)
South	14	3	0.5494 (0.4796)	0.0015 (0.0015)

Table 3. Pairwise estimates of genetic differentiation between the north, buffer, and south regions in the Savannah study area based on a) sample collection location of all 69 samples; b) sample collection location of dolphins with ≥ 10 sightings; and c) sighting history data -- samples from dolphins with ≥ 10 sightings were assigned to the region where 50% or more of the animal's sightings were located. F_{ST} and Φ_{ST} were estimated among regions in ARLEQUIN 3.5.2.2. F_{ST} values are reported below the diagonal, and Φ_{ST} values are reported above. *Significant difference after Bonferroni correction for multiple comparisons.

	n	North	Buffer	South
a)				
North	26	-	0.0404 ($p = 0.1433$)	-0.0020 ($p = 0.4069$)
Buffer	14	0.1169 ($p = 0.0303$)	-	0.0307 ($p = 0.2401$)
South	29	-0.0040 ($p = 0.4219$)	0.0556 ($p = 0.1341$)	-
b)				
North	16	-	0.0634 ($p = 0.0978$)	-0.0101 ($p = 0.5072$)
Buffer	11	0.2237 ($p = 0.0147$)*	-	0.0168 ($p = 0.2844$)
South	18	0.0507 ($p = 0.1279$)	0.0500 ($p = 0.2036$)	-

c)

North	18	-	0.1241 (p = 0.0407)	0.0528 (p = 0.1371)
Buffer	13	0.2969 (p = 0.0018)*	-	-0.0430 (p = 0.9999)
South	14	0.1678 (p = 0.0164)*	-0.0229 (p = 0.5944)	-




Figure 1. The Northern Georgia/Southern South Carolina Estuarine System (NGSSCES) Stock of common bottlenose dolphins (*Tursiops truncatus*) on the eastern coast of the United States and the Savannah study area. The NGSSCES Stock is continuous in the estuarine waters from the North Edisto River, South Carolina to northern Ossabaw Sound, Georgia (Hayes et al., 2017). The Savannah study area includes the inshore waters of Savannah, Georgia from the Savannah River to northern Ossabaw Sound.




Figure 2. The Savannah study area was categorized into three segments: a) a north region that consisted of the estuarine waters between the Savannah River and the Wilmington River; b) a buffer region that included the Wilmington River; and c) a south region that included the area between the Wilmington River and northern Ossabaw Sound. Photo-identification survey transects are in red.




Figure 3. Biopsy (circles) and stranding samples (triangles) collected from common bottlenose dolphins in Savannah, Georgia from the Savannah River to northern Ossabaw Sound. A total of 69 samples were collected. Biopsy samples (n = 50) were collected in September 2015 and February and March 2017. Stranding samples (n = 19) were collected from July 2008 through April 2017. Sampling locations are color coded to match identified mitochondrial control region haplotype.