

Online Resource 3 – Bowhead whale (*Balaena mysticetus*) carcass drift directions, 2009-2019.

Bowhead whale (Balaena mysticetus) carcasses documented during the 2019 aerial surveys in the eastern Chukchi and western Beaufort seas: A follow up to evidence of bowhead whale and killer whale (Orcinus orca) co-occurrence during 2009–2018

Polar Biology

Willoughby, AL^{*1,2}, Ferguson MC^{2,3}, Stimmelmayer R^{4,5}, Brower AA^{1,2}

¹ University of Washington, Cooperative Institute for Climate, Ocean and Ecosystem Studies, Seattle, WA, USA

² NOAA, NMFS, Alaska Fisheries Science Center, Marine Mammal Laboratory, Seattle, WA, USA

³ University of Washington, School of Fishery and Aquatic Sciences, Seattle, WA, USA

⁴ North Slope Borough, Department of Wildlife Management, Utqiagvik, AK, USA

⁵ University of Alaska Fairbanks, Institute of Arctic Biology, Fairbanks, AK, USA

*Corresponding author: email: a.willowb@gmail.com




Fig. 3.1 Bowhead whale (*Balaena mysticetus*) carcass drift directions, 2009–2019. Sightings of individual carcasses sighted more than once are color-coded. The star symbol represents the initial sighting, the octagon symbols represent the second sighting, the cross symbols represent the third sighting, and the asterisk symbol represents the four sightings. Lines with triangles depict the drift direction of the carcass with the matching color. The boundary for the eastern Chukchi Sea (EC) study area is indicated in grey and the border for the western Beaufort (WB) Sea study area is shown in black (see Fig. 1 for a full view of the EC and WB study areas)