

Supplemental Materials for Data Rich but Model Resistant: An Evaluation of Data-Limited Methods to Manage Fisheries with Failed Age-based Stock Assessments

Christopher M. Legault¹, John Wiedenmann² Jonathan J. Deroba¹, Gavin Fay³, Timothy J. Miller¹, Elizabeth N. Brooks¹, Richard J. Bell⁴, Joseph A. Langan⁵, Jamie M. Cournane⁶, Andrew W. Jones¹, Brandon Muffley⁷

¹National Marine Fisheries Service, Northeast Fisheries Science Center, Woods Hole, MA, USA

²Department of Ecology, Evolution and Natural Resources, Rutgers, The State University of New Jersey, New Brunswick, New Jersey, USA

³University of Massachusetts Dartmouth, School for Marine Science and Technology, Massachusetts, USA

⁴The Nature Conservancy, Rhode Island, USA

⁵University of Rhode Island, Graduate School of Oceanography, Narragansett, RI, USA

⁶New England Fishery Management Council, Newburyport, MA, USA

⁷Mid-Atlantic Fishery Management Council, Dover, DE, USA

We provide additional details of the simulation study (Figures S1-S4) and results (Figures S5-S37) for readers seeking more granular results of the 16 scenarios. The distribution of Mohn's rho values is shown in Figure S1, illustrating that we achieved a median rho of 0.5 for SSB. The difference in true F_{MSY} relative to the fishing history scenarios is shown in Figure S2; this provides some insight into the choice for which reference point was used for calculating overfishing. A similar plot for spawning stock biomass is shown in Figure S3 demonstrating the starting conditions for the feedback portion of the simulations. Retrospective patterns continued in the feedback portion of the closed-loop simulations, as seen in two realizations for both sources of the retrospective pattern in Figure S4. An alternative way of viewing the median results from Figure 1 in a more summarized approach that does not include uncertainty from the 1000 realizations is seen in the plot of the first two principal components of Figure S5. Plots similar to Figure 5 are seen in Figures S6-S21, showing the catch and spawning stock biomass relative to their MSY reference points for each of the 1000 realizations with each figure representing a different scenario. Figures S22-S37 are similar to Figures S6-S21 except that the y-axis is now fishing mortality relative to F_{MSY} , making them similar to Kobe plots.




Figure S1. Distribution of 50 Mohn's rho values for no misspecification (top row), catch misspecification (middle row), and natural mortality misspecification (bottom row) for spawning stock biomass (SSB), fishing mortality rate (Fbar), and recruitment (R). Panels are defined by retrospective type (none, catch, natural mortality), fishing intensity (1=always overfishing, 2=overfishing reducing to F_{MSY}), and number of selectivity blocks (1 or 2).




Figure S2. Fishing mortality rate (F) during the base period (black line) for the 8 combinations of F history ($F = F_{MSY}$ in second half of base period, O = overfishing throughout), number of fishery selectivity blocks (1 or 2), and retrospective source (catch or natural mortality, M). The horizontal blue line denotes the F at maximum sustainable yield (F_{MSY}) computed from conditions in the first year of the base period (1970) while the red dashed horizontal line denotes the F_{MSY} computed from conditions in the terminal year in the base period (2019).




Figure S3. Spawning stock biomass (SSB) during the base period (black line) for the 8 combinations of F history ($F = F_{MSY}$ in second half of base period, O = overfishing throughout), number of fishery selectivity blocks (1 or 2), and retrospective source (catch or natural mortality, M). The horizontal blue line denotes the F at maximum sustainable yield (F_{MSY}) computed from conditions in the first year of the base period (1970) while the red dashed horizontal line denotes the F_{MSY} computed from conditions in the terminal year in the base period (2019).




Figure S4. Mohn's rho for spawning stock biomass calculated from the statistical catch at age model during the feedback period of the simulations for two realizations (columns) and the two sources of retrospective (catch and natural mortality).




Figure S5. Principal components analysis of the median values of the metrics shown in Figure 1.




Figure S6. Relationship between long-term average catch and spawning stock biomass relative to their reference points by method. Each point represents the average for years 21-40 in the feedback period for a single iteration of a scenario. The scenario shown is where catch was the source of the retrospective pattern with F reduced to F_{MSY} in the second half of the base period, there was a single selectivity block, and where no buffer was applied to the catch advice (catch multiplier = 1).

CF1R




Figure S7. Relationship between long-term average catch and spawning stock biomass relative to their reference points by method. Each point represents the average for years 21-40 in the feedback period for a single iteration of a scenario. The scenario shown is where catch was the source of the retrospective pattern with F reduced to F_{MSY} in the second half of the base period, there was a single selectivity block, and where a buffer was applied to the catch advice (catch multiplier = 0.75).

CF2A




Figure S8. Relationship between long-term average catch and spawning stock biomass relative to their reference points by method. Each point represents the average for years 21-40 in the feedback period for a single iteration of a scenario. The scenario shown is where catch was the source of the retrospective pattern with F reduced to F_{MSY} in the second half of the base period, there were two selectivity blocks, and where no buffer was applied to the catch advice (catch multiplier = 1).

CF2R




Figure S9. Relationship between long-term average catch and spawning stock biomass relative to their reference points by method. Each point represents the average for years 21-40 in the feedback period for a single iteration of a scenario. The scenario shown is where catch was the source of the retrospective pattern with F reduced to F_{MSY} in the second half of the base period, there were two selectivity blocks, and where a buffer was applied to the catch advice (catch multiplier = 0.75).




Figure S10. Relationship between long-term average catch and spawning stock biomass relative to their reference points by method. Each point represents the average for years 21-40 in the feedback period for a single iteration of a scenario. The scenario shown is where catch was the source of the retrospective pattern with F remaining at 2.5 times F_{MSY} in the second half of the base period, there was a single selectivity block, and where no buffer was applied to the catch advice (catch multiplier = 1).

CO1R




Figure S11. Relationship between long-term average catch and spawning stock biomass relative to their reference points by method. Each point represents the average for years 21-40 in the feedback period for a single iteration of a scenario. The scenario shown is where catch was the source of the retrospective pattern with F remaining at 2.5 times F_{MSY} in the second half of the base period, there was a single selectivity block, and where a buffer was applied to the catch advice (catch multiplier = 0.75).




Figure S12. Relationship between long-term average catch and spawning stock biomass relative to their reference points by method. Each point represents the average for years 21-40 in the feedback period for a single iteration of a scenario. The scenario shown is where catch was the source of the retrospective pattern with F remaining at 2.5 times F_{MSY} in the second half of the base period, there were two selectivity blocks, and where no buffer was applied to the catch advice (catch multiplier = 1).




Figure S13. Relationship between long-term average catch and spawning stock biomass relative to their reference points by method. Each point represents the average for years 21-40 in the feedback period for a single iteration of a scenario. The scenario shown is where catch was the source of the retrospective pattern with F remaining at 2.5 times F_{MSY} in the second half of the base period, there were two selectivity blocks, and where a buffer was applied to the catch advice (catch multiplier = 0.75).




Figure S14. Relationship between long-term average catch and spawning stock biomass relative to their reference points by method. Each point represents the average for years 21-40 in the feedback period for a single iteration of a scenario. The scenario shown is where natural mortality was the source of the retrospective pattern with F reduced to F_{MSY} in the second half of the base period, there was a single selectivity block, and where no buffer was applied to the catch advice (catch multiplier = 1).

MF1R




Figure S15. Relationship between long-term average catch and spawning stock biomass relative to their reference points by method. Each point represents the average for years 21-40 in the feedback period for a single iteration of a scenario. The scenario shown is where natural mortality was the source of the retrospective pattern with F reduced to F_{MSY} in the second half of the base period, there was a single selectivity block, and where a buffer was applied to the catch advice (catch multiplier = 0.75).

MF2A




Figure S16. Relationship between long-term average catch and spawning stock biomass relative to their reference points by method. Each point represents the average for years 21-40 in the feedback period for a single iteration of a scenario. The scenario shown is where natural mortality was the source of the retrospective pattern with F reduced to F_{MSY} in the second half of the base period, there were two selectivity blocks, and where no buffer was applied to the catch advice (catch multiplier = 1).




Figure S17. Relationship between long-term average catch and spawning stock biomass relative to their reference points by method. Each point represents the average for years 21-40 in the feedback period for a single iteration of a scenario. The scenario shown is where natural mortality was the source of the retrospective pattern with F reduced to F_{MSY} in the second half of the base period, there were two selectivity blocks, and where a buffer was applied to the catch advice (catch multiplier = 0.75).

MO1A




Figure S18. Relationship between long-term average catch and spawning stock biomass relative to their reference points by method. Each point represents the average for years 21-40 in the feedback period for a single iteration of a scenario. The scenario shown is where natural mortality was the source of the retrospective pattern with F remaining at 2.5 times F_{MSY} in the second half of the base period, there was a single selectivity block, and where no buffer was applied to the catch advice (catch multiplier = 1).

MO1R




Figure S19. Relationship between long-term average catch and spawning stock biomass relative to their reference points by method. Each point represents the average for years 21-40 in the feedback period for a single iteration of a scenario. The scenario shown is where natural mortality was the source of the retrospective pattern with F remaining at 2.5 times F_{MSY} in the second half of the base period, there was a single selectivity block, and where a buffer was applied to the catch advice (catch multiplier = 0.75).

MO2A




Figure S20. Relationship between long-term average catch and spawning stock biomass relative to their reference points by method. Each point represents the average for years 21-40 in the feedback period for a single iteration of a scenario. The scenario shown is where natural mortality was the source of the retrospective pattern with F remaining at 2.5 times F_{MSY} in the second half of the base period, there were two selectivity blocks, and where no buffer was applied to the catch advice (catch multiplier = 1).

MO2R




Figure S21. Relationship between long-term average catch and spawning stock biomass relative to their reference points by method. Each point represents the average for years 21-40 in the feedback period for a single iteration of a scenario. The scenario shown is where natural mortality was the source of the retrospective pattern with F remaining at 2.5 times F_{MSY} in the second half of the base period, there were two selectivity blocks, and where a buffer was applied to the catch advice (catch multiplier = 0.75).




Figure S22. Relationship between long-term average fishing mortality rate and spawning stock biomass relative to their reference points by method. Each point represents the average for years 21-40 in the feedback period for a single iteration of a scenario. The scenario shown is where catch was the source of the retrospective pattern with F reduced to F_{MSY} in the second half of the base period, there was a single selectivity block, and where no buffer was applied to the catch advice (catch multiplier = 1).




Figure S23. Relationship between long-term average fishing mortality rate and spawning stock biomass relative to their reference points by method. Each point represents the average for years 21-40 in the feedback period for a single iteration of a scenario. The scenario shown is where catch was the source of the retrospective pattern with F reduced to F_{MSY} in the second half of the base period, there was a single selectivity block, and where a buffer was applied to the catch advice (catch multiplier = 0.75).

CF2A




Figure S24. Relationship between long-term average fishing mortality rate and spawning stock biomass relative to their reference points by method. Each point represents the average for years 21-40 in the feedback period for a single iteration of a scenario. The scenario shown is where catch was the source of the retrospective pattern with F reduced to F_{MSY} in the second half of the base period, there were two selectivity blocks, and where no buffer was applied to the catch advice (catch multiplier = 1).

CF2R




Figure S25. Relationship between long-term average fishing mortality rate and spawning stock biomass relative to their reference points by method. Each point represents the average for years 21-40 in the feedback period for a single iteration of a scenario. The scenario shown is where catch was the source of the retrospective pattern with F reduced to F_{MSY} in the second half of the base period, there were two selectivity blocks, and where a buffer was applied to the catch advice (catch multiplier = 0.75).




Figure S26. Relationship between long-term average fishing mortality rate and spawning stock biomass relative to their reference points by method. Each point represents the average for years 21-40 in the feedback period for a single iteration of a scenario. The scenario shown is where catch was the source of the retrospective pattern with F remaining at 2.5 times F_{MSY} in the second half of the base period, there was a single selectivity block, and where no buffer was applied to the catch advice (catch multiplier = 1).




Figure S27. Relationship between long-term average fishing mortality rate and spawning stock biomass relative to their reference points by method. Each point represents the average for years 21-40 in the feedback period for a single iteration of a scenario. The scenario shown is where catch was the source of the retrospective pattern with F remaining at 2.5 times F_{MSY} in the second half of the base period, there was a single selectivity block, and where a buffer was applied to the catch advice (catch multiplier = 0.75).

CO2A




Figure S28. Relationship between long-term average fishing mortality rate and spawning stock biomass relative to their reference points by method. Each point represents the average for years 21-40 in the feedback period for a single iteration of a scenario. The scenario shown is where catch was the source of the retrospective pattern with F remaining at 2.5 times F_{MSY} in the second half of the base period, there were two selectivity blocks, and where no buffer was applied to the catch advice (catch multiplier = 1).

CO2R




Figure S29. Relationship between long-term average fishing mortality rate and spawning stock biomass relative to their reference points by method. Each point represents the average for years 21-40 in the feedback period for a single iteration of a scenario. The scenario shown is where catch was the source of the retrospective pattern with F remaining at 2.5 times F_{MSY} in the second half of the base period, there were two selectivity blocks, and where a buffer was applied to the catch advice (catch multiplier = 0.75).

MF1A




Figure S30. Relationship between long-term average fishing mortality rate and spawning stock biomass relative to their reference points by method. Each point represents the average for years 21-40 in the feedback period for a single iteration of a scenario. The scenario shown is where natural mortality was the source of the retrospective pattern with F reduced to F_{MSY} in the second half of the base period, there was a single selectivity block, and where no buffer was applied to the catch advice (catch multiplier = 1).

MF1R




Figure S31. Relationship between long-term average fishing mortality rate and spawning stock biomass relative to their reference points by method. Each point represents the average for years 21-40 in the feedback period for a single iteration of a scenario. The scenario shown is where natural mortality was the source of the retrospective pattern with F reduced to F_{MSY} in the second half of the base period, there was a single selectivity block, and where a buffer was applied to the catch advice (catch multiplier = 0.75).

MF2A




Figure S32. Relationship between long-term average fishing mortality rate and spawning stock biomass relative to their reference points by method. Each point represents the average for years 21-40 in the feedback period for a single iteration of a scenario. The scenario shown is where natural mortality was the source of the retrospective pattern with F reduced to F_{MSY} in the second half of the base period, there were two selectivity blocks, and where no buffer was applied to the catch advice (catch multiplier = 1).

MF2R




Figure S33. Relationship between long-term average fishing mortality rate and spawning stock biomass relative to their reference points by method. Each point represents the average for years 21-40 in the feedback period for a single iteration of a scenario. The scenario shown is where natural mortality was the source of the retrospective pattern with F reduced to F_{MSY} in the second half of the base period, there were two selectivity blocks, and where a buffer was applied to the catch advice (catch multiplier = 0.75).

MO1A




Figure S34. Relationship between long-term average fishing mortality rate and spawning stock biomass relative to their reference points by method. Each point represents the average for years 21-40 in the feedback period for a single iteration of a scenario. The scenario shown is where natural mortality was the source of the retrospective pattern with F remaining at 2.5 times F_{MSY} in the second half of the base period, there was a single selectivity block, and where no buffer was applied to the catch advice (catch multiplier = 1).

MO1R




Figure S35. Relationship between long-term average fishing mortality rate and spawning stock biomass relative to their reference points by method. Each point represents the average for years 21-40 in the feedback period for a single iteration of a scenario. The scenario shown is where natural mortality was the source of the retrospective pattern with F remaining at 2.5 times F_{MSY} in the second half of the base period, there was a single selectivity block, and where a buffer was applied to the catch advice (catch multiplier = 0.75).

MO2A




Figure S36. Relationship between long-term average fishing mortality rate and spawning stock biomass relative to their reference points by method. Each point represents the average for years 21-40 in the feedback period for a single iteration of a scenario. The scenario shown is where natural mortality was the source of the retrospective pattern with F remaining at 2.5 times F_{MSY} in the second half of the base period, there were two selectivity blocks, and where no buffer was applied to the catch advice (catch multiplier = 1).

MO2R




Figure S37. Relationship between long-term average fishing mortality rate and spawning stock biomass relative to their reference points by method. Each point represents the average for years 21-40 in the feedback period for a single iteration of a scenario. The scenario shown is where natural mortality was the source of the retrospective pattern with F remaining at 2.5 times F_{MSY} in the second half of the base period, there were two selectivity blocks, and where a buffer was applied to the catch advice (catch multiplier = 0.75).