



COMMENTARY

10.1029/2021GL097617

Special Section:

The Exceptional Arctic Polar Vortex in 2019/2020: Causes and Consequences

Key Points:

- The stratospheric polar vortex is a well-defined feature dominating the cool-season circulation in each hemisphere from ~15–50 km altitude
- The tropospheric circulation does not constitute a single coherent structure and is most aptly described by regional jet stream variations
- Accuracy in defining and describing the “polar vortex” and its effects is key to improving understanding by non-specialist audiences

Correspondence to:

G. L. Manney,
manney@nwra.com

Citation:

Manney, G. L., Butler, A. H., Lawrence, Z. D., Wargan, K., & Santee, M. L. (2022). What's in a name? On the use and significance of the term “polar vortex”. *Geophysical Research Letters*, 49, e2021GL097617. <https://doi.org/10.1029/2021GL097617>

Received 7 JAN 2022
 Accepted 12 MAY 2022

Author Contributions:

Conceptualization: Gloria L. Manney, Amy H. Butler, Zachary D. Lawrence, Krzysztof Wargan, Michelle L. Santee
Methodology: Gloria L. Manney, Amy H. Butler, Zachary D. Lawrence, Krzysztof Wargan, Michelle L. Santee
Software: Gloria L. Manney
Visualization: Gloria L. Manney
Writing – original draft: Gloria L. Manney

© 2022. The Authors.

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs License](https://creativecommons.org/licenses/by/4.0/), which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

What's in a Name? On the Use and Significance of the Term “Polar Vortex”

Gloria L. Manney^{1,2} , Amy H. Butler³ , Zachary D. Lawrence^{4,5,6} , Krzysztof Wargan^{7,8} , and Michelle L. Santee⁹ 

¹NorthWest Research Associates, Socorro, NM, USA, ²New Mexico Institute of Mining and Technology, Socorro, NM, USA, ³NOAA Chemical Sciences Laboratory, Boulder, CO, USA, ⁴Cooperative Institute for Research in Environmental Sciences, University of Colorado, Boulder, CO, USA, ⁵NOAA Physical Sciences Laboratory, Boulder, CO, USA, ⁶NorthWest Research Associates, Boulder, CO, USA, ⁷NASA Goddard Space Flight Center, Greenbelt, MD, USA, ⁸Science Systems and Applications, Inc., Lanham, MD, USA, ⁹Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA, USA

Abstract Mainstream and popular science media are rife with misunderstandings about what a “polar vortex” is. The term most aptly describes the stratospheric polar vortex, a single feature dominating the cool-season circulation from ~15–50 km. Regional jet stream variations dominate the tropospheric circulation, which is not well-described as a polar vortex; indeed, there is no single consistent definition of a tropospheric polar vortex in the literature. Stratospheric polar vortex disturbances profoundly influence extreme weather events, including cold air outbreaks (CAOs). How the stratospheric polar vortex affects tropospheric jets, whose local excursions drive CAOs, is not fully understood. Public-facing parts of publications describing research on this topic are not always clear about how the “polar vortex” is defined; greater clarity could improve communications both within the community and with non-specialist audiences.

Plain Language Summary What is a “polar vortex”? The atmospheric science community most commonly uses this term to describe the stratospheric polar vortex, a band of winds extending from about 15–50 km altitude that flows around the pole of each hemisphere during their respective fall through spring seasons. However, the term “polar vortex” has been used in mainstream media and popular science platforms to instead describe local variations in the upper tropospheric jet streams (winds that blow most strongly between about 8 and 13 km altitude) and even individual extreme cold weather events. We argue that the term should be used only in reference to the stratospheric polar vortex, which is a single feature that predominantly controls dynamical and chemical variability in the winter polar stratosphere. The stratospheric polar vortex is related to but distinct from more regional jet stream excursions and associated weather extremes; further study is needed to fully understand these relationships.

1. The Stratospheric Polar Vortex, Tropospheric Jet Streams, and Cold Air Outbreaks

This commentary appears in the Special Collection focusing on the Arctic stratospheric “polar vortex” in 2019/2020. But how clear are we about what constitutes a “polar vortex”? Confusion persists in the popular press about what a polar vortex is and how polar vortices relate to extreme weather events. This confusion stems in part from imprecise descriptions by the scientific community.

In January 2014, a cold air outbreak (CAO) set record-low minimum temperatures throughout the south central and eastern US (e.g., Screen et al., 2015). Headlines hailed it as “the polar vortex,” and this language became commonplace in news and popular science media. At the time, the term “polar vortex” in scientific literature typically described the stratospheric polar vortex (see, e.g., Lillo et al., 2021; Waugh et al., 2017, for discussion of this), but some studies used the term to describe the “tropospheric polar vortex” (e.g., Wallace et al., 2014; Yu & Zhang, 2015), in both cases often without further qualification. Waugh et al. (2017) sought to dispel confusion, describing the stratospheric and tropospheric “circumpolar” vortices as these terms had been commonly used in scientific literature, highlighting their differences and relationships to extreme weather events, and providing recommendations for describing them. While this work is widely cited, the two concepts are still often confused, including on educational websites and in climate change communication studies (e.g., Lyons et al., 2018; Shepherd, 2016; UCAR, 2021; UC Davis, 2019). Even recent papers within the atmospheric science community are

Writing – review & editing: Amy H. Butler, Zachary D. Lawrence, Krzysztof Wargan, Michelle L. Santee

not always clear about which circulation feature(s) they are discussing, and some use the term “polar vortex” to describe synoptic-scale disturbances associated with CAOs, echoing the inaccurate usage in popular media (e.g., Bushra & Rohli, 2019, 2021; Dai et al., 2021; Jiang, 2021; Juzbašić et al., 2021; Kömüščü & Oğuz, 2021; Nielsen-Gammon et al., 2021; Overland, 2021; Overland & Wang, 2019; Xiong et al., 2021; Zhang et al., 2021). Sometimes the most public-facing parts of research papers (abstracts, plain language summaries, key points) do not clearly define how the term “polar vortex” is used.

Figure 1 shows characteristics of the stratospheric and tropospheric circulation on two occasions the popular press described CAOs as a polar vortex “outbreak” or “attack,” but which were associated with very different stratospheric polar vortex conditions. This figure shows the stratospheric polar vortex, upper tropospheric jet streams, and the circulation that is sometimes described as a “tropospheric polar vortex” (i.e., the 2 and 3 PVU potential vorticity (PV) contours on the 330-K isentropic surface, one possible definition discussed by Waugh et al., 2017).

The stratospheric polar vortex is bounded by the polar night jet, a band of strong eastward winds throughout the stratosphere that forms in fall in each hemisphere and vanishes in spring. Different diagnostics of the stratospheric polar vortex edge (e.g., Lawrence & Manney, 2018) select similar physically meaningful boundaries (Figure 1a, left, defined using a PV contour coincident with the strongest PV gradients, as in Lawrence et al. (2018)). *The stratospheric polar vortex constitutes a single feature that dominates the circulation and transport throughout the polar stratosphere from fall through spring.*

The so-called “tropospheric polar vortex,” as most often defined, exists year-round, but no single definition uniquely identifies it or the altitude(s) at which it exists (the characteristics described herein do not depend substantially on which of numerous definitions is used). We show one common definition (Waugh et al., 2017, and references therein) whereby its edge follows the axis of an upper tropospheric jet on an isentropic surface in the middle to upper troposphere. The maximum winds of these jets are very localized in altitude compared to the stratospheric polar night jet, and they vary strongly with longitude (e.g., Manney, Hegglin, et al., 2011; Manney et al., 2014, Figure 1a). Because regional variability of discontinuous jet streams governs the extratropical tropospheric circulation, “tropospheric polar vortex” definitions do not describe a single dominant circumpolar circulation. Further confusion arises from the distinction between tropospheric “polar” (primarily eddy driven) and “subtropical” (largely radiatively driven) jets. While some recent papers and popular science pieces identify the “tropospheric polar vortex” with the tropospheric polar jet (e.g., Bushra & Rohli, 2021; UCAR, 2021; Waugh et al., 2017), numerous studies show that tropospheric jets are not well-represented by this simplified conceptual division but rather form a seasonally and regionally varying complex with hybrid radiatively and eddy-driven features that is rarely continuous around the globe (S. Lee & Kim, 2003; Manney et al., 2014; Spensberger & Spengler, 2020, and references therein).

These differences are reflected in windspeeds (Figures 1a and 1b), which peak sharply along the stratospheric polar vortex edge; in contrast, a “tropospheric polar vortex” defined as noted above meanders through regions of weak and strong winds, leading to a broad, flat distribution of “vortex-edge” windspeeds. Potential vorticity gradients (indicating polar vortex strength) are consistently strong along the circumference of the stratospheric polar vortex but have many localized maxima in small portions of the “tropospheric vortex” edge and elsewhere in the extratropics (Figure 1c). This results in relatively stronger mean PV gradients along the stratospheric vortex edge, versus weaker mean PV gradients and most frequent values near zero in the troposphere (Figure 1d). Further, tropospheric windspeeds (Figure 1a) often show a single jet (or no strong jet) because separate tropospheric polar and subtropical jets do not always exist. A “tropospheric polar vortex” might therefore follow the polar jet in one region but the subtropical jet in another, thus traversing regimes controlled by different dynamical processes.

The stratospheric polar vortex is critical for transport, chemical processing, confinement of processed air, and ozone loss. Processes promoting ozone depletion are commonly analyzed from a vortex-centered perspective because the stratospheric vortex represents a strong transport barrier, isolating air primed for ozone destruction (e.g., Manney et al., 2020; Manney, Santee, et al., 2011; Schoeberl et al., 1992); the amount of polar ozone loss in a given spring depends critically on the strength and coldness of the winter/spring stratospheric polar vortex. In contrast, upper tropospheric ozone variability is dominated by regional variations in stratosphere-troposphere exchange and the amount of lower stratospheric ozone available for transport into the troposphere (e.g., Albers




Figure 1.

et al., 2018; Breeden et al., 2021; Olsen et al., 2019). Figures 1e and 1f illustrate these differences: Ozone gradients change abruptly across the stratospheric polar vortex edge but are quite uniform within it. In contrast, ozone gradients are strong in many localized regions within the “tropospheric polar vortex,” with highly variable gradients often appearing well poleward of the “vortex” edge. These characteristics are reflected in sharply peaked ozone distributions along the stratospheric polar vortex edge and large variability in ozone along the “tropospheric vortex” edge (Figure 1f). Note that the broad change from uniform gradients to highly variable gradients across the “tropospheric vortex edge” as defined here is a reflection of vertical ozone gradients and the tilt of the 330-K isentropic surface in the subtropics.

Stratosphere-troposphere coupling (e.g., Baldwin & Dunkerton, 2001; Kidston et al., 2015) dynamically links variability of the polar vortex to extremes at the surface (e.g., Domeisen & Butler, 2020). For example, extreme stratospheric polar vortex disruptions (sudden stratospheric warmings, SSWs) are associated with increased risk of mid-latitude CAOs (e.g., Baldwin et al., 2021; Butler et al., 2017; Huang et al., 2021; King et al., 2019), and unusually strong stratospheric polar vortices are associated with anomalously high extratropical surface temperatures (including heat waves and destructive wildfires) (Lawrence et al., 2020; Limpasuvan et al., 2005; Overland & Wang, 2021). Because radiative timescales are longer in the lower stratosphere, disruptions to the circulation can persist there for weeks to months, potentially providing subseasonal-to-seasonal forecast skill for extremes like CAOs (e.g., Domeisen et al., 2019). Using information about the stratospheric polar vortex to predict CAOs is, however, complicated because the timing and location of individual CAOs varies significantly following polar vortex disruptions, perhaps related to details of the stratospheric polar vortex characteristics and evolution. Recent work suggests that Eurasian CAOs are more closely linked to SSWs, while North American CAOs are more strongly associated with stratospheric polar vortex elongation that might or might not accompany an SSW (e.g., Cohen, Agel, Barlow, Garfinkel, & White, 2021; Kretschmer et al., 2018; S. H. Lee et al., 2019). It is worth emphasizing that CAOs can occur during both strong and weak stratospheric polar vortex conditions (e.g., Cohen, Agel, Barlow, Furtado, et al., 2021; S. H. Lee et al., 2019): Figure 1 shows a CAO (January 2014) linked to a strong (but distorted) stratospheric vortex and one (February 2021) following an SSW.

CAOs are often termed “polar vortex events” in the news, popular science media, and less specialized peer-reviewed papers (e.g., Lyons et al., 2018, on communication of climate change risks), but the dynamical processes involved argue that they are best described as equatorward excursions of the tropospheric jets and southward advection of cold Arctic air. These features are not generally correlated with the strength of any globally defined “tropospheric polar vortex” (e.g., Bushra & Rohli, 2021; Cellitti et al., 2006; Waugh et al., 2017), so the utility of the latter concept in relation to CAOs is questionable. CAOs in some regions are indeed more likely, and more likely to be severe, following SSWs (e.g., Huang et al., 2021; King et al., 2019; S. H. Lee et al., 2019), explaining why the media often hails reports of an SSW with “the polar vortex is coming” even though an SSW actually represents a rapid deceleration, or disappearance, of the stratospheric polar vortex winds. While the relationship to stratospheric polar vortex disturbances can improve lead times for probabilistic forecasts of CAO occurrence, more extensive mechanistic understanding of how stratospheric polar vortex anomalies affect regional excursions of tropospheric jet streams is needed to further improve prediction of when and where CAOs will occur.

The term “polar vortex” is used in another way that is not directly related to any planetary-scale circumpolar vortex, but is related to many CAOs (e.g., Lillo et al., 2021). A “tropopause polar vortex” (TPV) is a sub-synoptic-scale feature characterized by a deep depression of the tropopause (sometimes to near the surface) bounded by an “Arctic jet stream” poleward of and below the tropospheric polar jet (Shapiro et al., 1987). Lillo et al. (2021) showed that the North American CAO in late January 2019 resulted directly from a TPV moving southward from its high-latitude origins; TPVs play a role in many (but by no means all) CAOs (e.g., Biernat et al., 2021; Papritz et al., 2019). While the existence of yet another feature termed a “polar vortex” may engender confusion, the direct link of these localized vortices to CAOs emphasizes the importance of local/regional circulation anomalies (and associated jet stream excursions) to extreme weather events.

Figure 1. Characteristics of (left to right) 6 January 2014 and 16 February 2021 stratospheric and upper tropospheric circulations: (a) Windspeeds (colorfill) and two potential vorticity (PV) contours representing the stratospheric polar vortex edge (green) and boundary of tropospheric “global” circulation (orange). (b) Windspeed histograms along the “vortex edge” (most equatorward PV contour shown in panel (a)); vertical lines show mean around that PV contour. (c) Normalized PV gradient magnitudes. (d) Normalized PV gradient magnitude along the “vortex edge” (hemispheric mean is 1 by definition; vertical lines as in panel (b)). (e) Normalized ozone gradient magnitudes. (f) Normalized ozone mixing ratios along the “vortex edge” (vertical lines as in panel (b)). Cyan contours in panels (c and e) show “vortex edge” PV. 600-K (330-K) fields are shown for stratosphere (troposphere), except windspeeds are at 345 K (near level of maximum tropospheric jet stream winds). Data are from MERRA-2 (Gelaro et al., 2017).

Points such as those above regarding the stratospheric polar vortex have been highlighted in studies using theoretical fluid-dynamical or dynamical systems approaches (e.g., Mester & Esler, 2020; Scott & Dritschel, 2006; Serra et al., 2017). It is not clear that similar approaches could usefully describe what some have termed a “tropospheric polar vortex.”

2. Best Practices for Describing the Polar Vortex

It is clearly appropriate and useful to describe the stratospheric polar vortex as dominating stratospheric cool-season variability and exerting influence on the surface on sub-seasonal to seasonal timescales, including probabilistic links to extreme weather events. Jet stream excursions and related troughs and ridges are suitable for describing the genesis and evolution of CAOs, whereas the concept of a “tropospheric polar vortex” is typically not helpful in describing extreme weather events or elucidating their causes. We conclude:

1. The term “polar vortex” is most appropriate for describing the stratospheric polar vortex, but given its broad use and misuse, “stratospheric” should be specified explicitly.
2. The stratospheric polar vortex is a climatological feature that exists throughout the cool seasons (though sometimes temporarily disrupted) and thus should not be described as an “event” with a sub-seasonal time scale.
3. The tropospheric circulation, especially in relation to extreme weather events, can most clearly be described in relation to the tropospheric jet streams, without invoking the term “tropospheric polar vortex.” More accurate and appropriate terminology for referring to such events would be “Arctic CAO” (or more simply a CAO) or a “polar front.”
4. While the term “tropopause polar vortex” has been used to describe sub-synoptic scale vortices that are sometimes linked to CAOs, local features might be more clearly described in relation to their provenance, for example, a “Canadian tropopause vortex.”
5. Scientists should be careful in the public-facing parts of our communications (e.g., titles, abstracts, plain language summaries, web sites) to be clear and precise about what we mean by the term “polar vortex.”
6. In communications with the media, atmospheric scientists should emphasize that stratospheric polar vortex variability is indeed helpful in predicting CAOs and other extreme weather events, but stratospheric influence is exerted via regional jet stream variations that cannot in themselves be called a “polar vortex.”

Further study is needed to elucidate the relationship of stratospheric polar vortex variations to underlying regional tropospheric jet stream variations and ultimately to extreme weather events. The stratospheric polar vortex and tropospheric jet streams play important, but distinct, roles in understanding and forecasting extreme weather events. Accurate description of these features is thus critical to improving communication, both within the scientific community and with the public, regarding events that can have profound human impacts.

Acknowledgments

We thank the Microwave Limb Sounder team at JPL, especially Brian Knosp, Luis Millán, and Ryan Fuller, for data management, processing, and analysis support; NASA’s GMAO for the MERRA-2 data set; Ken Minschwaner, Jessica George, and Kody Gray for helpful discussions; and two anonymous reviewers for their insightful comments. We are grateful to B. J. Hoskins, M. E. McIntyre, and A. W. Robertson (Hoskins et al., 1985) for inspiring the title of this commentary. G. L. Manney was partially supported by the JPL Microwave Limb Sounder team under a JPL subcontract to NWSA. Z. D. Lawrence was partially supported under a NWS OSTI Weeks 3–4 Project (NA20NWS4680051). G. L. Manney and Z. D. Lawrence were partially supported by NSF Climate and Large-scale Dynamics Grant #2015906. K. Wargan was supported by NASA’s GMAO core funding. Work at the Jet Propulsion Laboratory, California Institute of Technology, was carried out under a contract with the National Aeronautics and Space Administration (80NM0018D0004).

Data Availability Statement

The MERRA-2 data set used here is publicly available: <https://disc.gsfc.nasa.gov/datasets?project=MERRA-2> (Global Modeling and Assimilation Office (GMAO), 2015).

References

- Albers, J. R., Perlwitz, J., Butler, A. H., Birner, T., Kiladis, G. N., Lawrence, Z. D., et al. (2018). Mechanisms governing interannual variability of stratosphere-to-troposphere ozone transport. *Journal of Geophysical Research: Atmospheres*, 123(1), 234–260. <https://doi.org/10.1002/2017JD026890>
- Baldwin, M. P., Ayarzagüena, B., Birner, T., Butchart, N., Butler, A. H., Charlton-Perez, A. J., et al. (2021). Sudden stratospheric warmings. *Reviews of Geophysics*, 59(1), e2020RG000708. <https://doi.org/10.1029/2020RG000708>
- Baldwin, M. P., & Dunkerton, T. J. (2001). Stratospheric harbingers of anomalous weather regimes. *Science*, 294(5542), 581–584. <https://doi.org/10.1126/science.1063315>
- Biernat, K. A., Bosart, L. F., & Keyser, D. (2021). A climatological analysis of the linkages between tropopause polar vortices, cold pools, and cold air outbreaks over the central and eastern United States. *Monthly Weather Review*, 149(1), 189–206. <https://doi.org/10.1175/MWR-D-20-0191.1>
- Breeden, M. L., Butler, A. H., Albers, J. R., Sprenger, M., & Langford, A. O. (2021). The spring transition of the North Pacific jet and its relation to deep stratosphere-to-troposphere mass transport over western North America. *Atmospheric Chemistry and Physics*, 21(4), 2781–2794. <https://doi.org/10.5194/acp-21-2781-2021>

- Bushra, N., & Rohli, R. V. (2019). An objective procedure for delineating the circumpolar vortex. *Earth and Space Science*, 6(5), 774–783. <https://doi.org/10.1029/2019EA000590>
- Bushra, N., & Rohli, R. V. (2021). Relationship between atmospheric teleconnections and the Northern Hemisphere's circumpolar vortex. *Earth and Space Science*, 8(9), e2021EA001802. <https://doi.org/10.1029/2021EA001802>
- Butler, A. H., Sjöberg, J. P., Seidel, D. J., & Rosenlof, K. H. (2017). A sudden stratospheric warming compendium. *Earth System Science Data*, 9(1), 63–76. <https://doi.org/10.5194/essd-9-63-2017>
- Cellitti, M. P., Walsh, J. E., Rauber, R. M., & Portis, D. H. (2006). Extreme cold air outbreaks over the United States, the polar vortex, and the large-scale circulation. *Journal of Geophysical Research*, 111(D2), D02114. <https://doi.org/10.1029/2005JD006273>
- Cohen, J. L., Agel, L., Barlow, M., Garfinkel, C. I., & White, I. (2021). Linking Arctic variability and change with extreme winter weather in the United States. *Science*, 373(6559), 1116–1121. <https://doi.org/10.1126/science.abi9167>
- Cohen, J. L., Agel, L. A., Barlow, M. A., Furtado, J. C., Kretschmer, M., & Matthias, V. (2021). The “polar vortex” winter of 2013/14. AGU Fall Meeting.
- Dai, G., Li, C., Han, Z., Luo, D., & Yao, Y. (2021). The nature and predictability of the East Asian extreme cold events of 2020/21. *Advances in Atmospheric Sciences*. <https://doi.org/10.1007/s00376-021-1057-3>
- Domeisen, D. I., & Butler, A. H. (2020). Stratospheric drivers of extreme events at the Earth's surface. *Communications Earth & Environment*, 1(1), 59. <https://doi.org/10.1038/s43247-020-00060-z>
- Domeisen, D. I., Butler, A. H., Charlton-Perez, A. J., Ayarzagüena, B., Baldwin, M. P., Dunn-Sigouin, E., et al. (2019). The role of the stratosphere in subseasonal to seasonal prediction Part II: Predictability arising from stratosphere - troposphere coupling. *Journal of Geophysical Research: Atmospheres*, 125(2), e2019JD030923. <https://doi.org/10.1029/2019JD030923>
- Gelaro, R., McCarty, W., Suárez, M. J., Todling, R., Molod, A., Takacs, L., et al. (2017). The Modern-Era Retrospective analysis for Research and Applications, version-2 (MERRA-2). *Journal of Climate*, 30(14), 5419–5454. <https://doi.org/10.1175/JCLI-D-16-0758.1>
- Global Modeling and Assimilation Office (GMAO). (2015). *MERRA-2 inst3_3d_asm_nv: 3d, 3-hourly, instantaneous, model-level, assimilation, assimilated meteorological fields v5.12.4, Greenbelt, MD, USA*. Goddard Earth Sciences Data and Information Services Center (GES DISC). <https://doi.org/10.5067/WWQX8Q8IVFW8>
- Hoskins, B. J., McIntyre, M. E., & Robertson, A. W. (1985). On the use and significance of isentropic potential vorticity maps. *Quarterly Journal of the Royal Meteorological Society*, 111(470), 877–946. <https://doi.org/10.1002/qj.49711147002>
- Huang, J., Hitchcock, P., Maycock, A. C., McKenna, C. M., & Tian, W. (2021). Northern hemisphere cold air outbreaks are more likely to be severe during weak polar vortex conditions. *Communications Earth & Environment*, 2(1), 147. <https://doi.org/10.1038/s43247-021-00215-6>
- Jiang, J. H. (2021). *Polar vortex linked to atmospheric circulation at daily scale*. Retrieved from https://eos.org/editor-highlights/polar-vortex-linked-to-atmospheric-circulation-at-daily-scale?utm_campaign=eaalert
- Juzbašić, A., Kryjov, V. N., & Ahn, J. B. (2021). On the anomalous development of the extremely intense positive Arctic Oscillation of the 2019–2020 winter. *Environmental Research Letters*, 16(5), 055008. <https://doi.org/10.1088/1748-9326/abe434>
- Kidston, J., Scaife, A. A., Hardiman, S. C., Mitchell, D. M., Butchart, N., Baldwin, M. P., & Gray, L. J. (2015). Stratospheric influence on tropospheric jet streams, storm tracks and surface weather. *Nature Geoscience*, 8(6), 433–440. <https://doi.org/10.1038/ngeo2424>
- King, A. D., Butler, A. H., Jucker, M., Earl, N. O., & Rudeva, I. (2019). Observed relationships between sudden stratospheric warmings and European climate extremes. *Journal of Geophysical Research: Atmospheres*, 124(24), 13943–13961. <https://doi.org/10.1029/2019JD030480>
- Körmüscü, A. Ü., & Oğuz, K. (2021). Analysis of cold anomalies observed over Turkey during the 2018/2019 winter in relation to polar vortex and other atmospheric patterns. *Meteorology and Atmospheric Physics*, 133(4), 1327–1354. <https://doi.org/10.1007/s00703-021-00806-0>
- Kretschmer, M., Cohen, J., Matthias, V., Runge, J., & Coumou, D. (2018). The different stratospheric influence on cold-extremes in Eurasia and North America. *npj Climate and Atmospheric Science*, 1, 44. <https://doi.org/10.1038/s41612-018-0054-4>
- Lawrence, Z. D., & Manney, G. L. (2018). Characterizing stratospheric polar vortex variability with computer vision techniques. *Journal of Geophysical Research: Atmospheres*, 123(3), 1510–1535. <https://doi.org/10.1002/2017JD027556>
- Lawrence, Z. D., Manney, G. L., & Wargan, K. (2018). Reanalysis intercomparisons of stratospheric polar processing diagnostics. *Atmospheric Chemistry and Physics*, 18, 13547–13579. <https://doi.org/10.5194/acp-18-13547-2018>
- Lawrence, Z. D., Perlwitz, J., Butler, A. H., Manney, G. L., Newman, P. A., Lee, S. H., & Nash, E. R. (2020). The remarkably strong Arctic stratospheric polar vortex of winter 2020: Links to record-breaking Arctic oscillation and ozone loss. *Journal of Geophysical Research: Atmospheres*, 125(22), e2020JD033271. <https://doi.org/10.1029/2020JD033271>
- Lee, S., & Kim, H.-K. (2003). The dynamical relationship between subtropical and eddy-driven jets. *Journal of the Atmospheric Sciences*, 60(12), 1490–1503. [https://doi.org/10.1175/1520-0469\(2003\)060<1490:tdrbsa>2.0.co;2](https://doi.org/10.1175/1520-0469(2003)060<1490:tdrbsa>2.0.co;2)
- Lee, S. H., Furtado, J. C., & Charlton-Perez, A. J. (2019). Wintertime North American weather regimes and the Arctic stratospheric polar vortex. *Geophysical Research Letters*, 46(24), 14892–14900. <https://doi.org/10.1029/2019GL085592>
- Lillo, S. P., Cavallo, S. M., Parsons, D. B., & Riedel, C. (2021). The role of a tropopause polar vortex in the generation of the January 2019 extreme Arctic outbreak. *Journal of the Atmospheric Sciences*, 78(9), 2801–2821. <https://doi.org/10.1175/JAS-D-20-0285.1>
- Limpasuvan, V., Hartmann, D. L., Thompson, D. W. J., Jeev, K., & Yung, Y. L. (2005). Stratosphere-troposphere evolution during polar vortex intensification. *Journal of Geophysical Research: Atmospheres*, 110(D24), D24101. <https://doi.org/10.1029/2005JD006302>
- Lyons, B. A., Hasell, A., & Stroud, N. J. (2018). Enduring extremes? Polar vortex, drought, and climate change beliefs. *Environmental Communication*, 12(7), 876–894. <https://doi.org/10.1080/17524032.2018.1520735>
- Manney, G. L., Hegglin, M. I., Daffer, W. H., Santee, M. L., Ray, E. A., Pawson, S., et al. (2011). Jet characterization in the upper troposphere/lower stratosphere (UTLS): Applications to climatology and transport studies. *Atmospheric Chemistry and Physics*, 11(12), 6115–6137. <https://doi.org/10.5194/acp-11-6115-2011>
- Manney, G. L., Hegglin, M. I., Daffer, W. H., Schwartz, M. J., Santee, M. L., & Pawson, S. (2014). Climatology of upper tropospheric/lower stratospheric (UTLS) jets and tropopauses in MERRA. *Journal of Climate*, 27(9), 3248–3271. <https://doi.org/10.1175/jcli-d-13-00243.1>
- Manney, G. L., Livesey, N. J., Santee, M. L., Froidevaux, L., Lambert, A., Lawrence, Z. D., et al. (2020). Record-low Arctic stratospheric ozone in 2020: MLS observations of chemical processes and comparisons with previous extreme winters. *Geophysical Research Letters*, 47(16), e2020GL089063. <https://doi.org/10.1029/2020GL089063>
- Manney, G. L., Santee, M. L., Rex, M., Livesey, N. J., Pitts, M. C., Veefkind, P., et al. (2011). Unprecedented Arctic ozone loss in 2011. *Nature*, 478(7370), 469–475. <https://doi.org/10.1038/nature10556>
- Mester, M., & Esler, J. G. (2020). Dynamical elliptical diagnostics of the Antarctic polar vortex. *Journal of the Atmospheric Sciences*, 77(3), 1167–1180. <https://doi.org/10.1175/JAS-D-19-0232.1>
- Nielsen-Gammon, J., Bolinger, R., Attard, H., Bentley, A., Brown, V., Fuhrmann, C., et al. (2021). *Evaluation of the February 2021 south-central big freeze*. AGU Fall Meeting.

- Olsen, M. A., Manney, G. L., & Liu, J. (2019). The ENSO and QBO impact on ozone variability and stratosphere-troposphere exchange relative to the subtropical jets. *Journal of Geophysical Research: Atmospheres*, *124*(13), 7379–7392. <https://doi.org/10.1029/2019JD030435>
- Overland, J. E. (2021). Rare events in the Arctic. *Climatic Change*, *168*(3), 27. <https://doi.org/10.1007/s10584-021-03238-2>
- Overland, J. E., & Wang, M. (2019). Impact of the winter polar vortex on greater North America. *International Journal of Climatology*, *39*(15), 5815–5821. <https://doi.org/10.1002/joc.6174>
- Overland, J. E., & Wang, M. (2021). The 2020 Siberian heat wave. *International Journal of Climatology*, *41*(S1), E2341–E2346. <https://doi.org/10.1002/joc.6850>
- Papritz, L., Rouges, E., Aemisegger, F., & Wernli, H. (2019). On the thermodynamic preconditioning of Arctic air masses and the role of tropopause polar vortices for cold air outbreaks from Fram Strait. *Journal of Geophysical Research: Atmospheres*, *124*(21), 11033–11050. <https://doi.org/10.1029/2019JD030570>
- Schoeberl, M. R., Lait, L. R., Newman, P. A., & Rosenfield, J. E. (1992). The structure of the polar vortex. *Journal of Geophysical Research*, *97*(D8), 7859–7882. <https://doi.org/10.1029/91jd02168>
- Scott, R. K., & Dritschel, D. G. (2006). Vortex–vortex interactions in the winter stratosphere. *Journal of the Atmospheric Sciences*, *63*(2), 726–740. <https://doi.org/10.1175/JAS3632.1>
- Screen, J. A., Deser, C., & Sun, L. (2015). Reduced risk of North American cold extremes due to continued Arctic sea ice loss. *Bulletin of the American Meteorological Society*, *96*(9), 1489–1503. <https://doi.org/10.1175/BAMS-D-14-00185.1>
- Serra, M., Sathe, P., Beron-Vera, F., & Haller, G. (2017). Uncovering the edge of the polar vortex. *Journal of the Atmospheric Sciences*, *74*(11), 3871–3885. <https://doi.org/10.1175/JAS-D-17-0052.1>
- Shapiro, M. A., Hampel, T., & Krueger, A. J. (1987). The Arctic tropopause fold. *Monthly Weather Review*, *115*(2), 444–454. [https://doi.org/10.1175/1520-0493\(1987\)115<0444:TAFF>2.0.CO;2](https://doi.org/10.1175/1520-0493(1987)115<0444:TAFF>2.0.CO;2)
- Shepherd, M. (2016). *12 weather and climate concepts that confuse the public*. Retrieved from <https://www.forbes.com/sites/marshallshepherd/2016/12/13/12-weather-and-climate-concepts-that-confuse-the-public/?sh=2943e192350b>
- Spensberger, C., & Spengler, T. (2020). Feature-based jet variability in the upper troposphere. *Journal of Climate*, *33*(16), 6849–6871. <https://doi.org/10.1175/JCLI-D-19-0715.1>
- UCAR. (2021). *Why the polar vortex keeps breaking out of the Arctic*. Retrieved from <https://scied.ucar.edu/learning-zone/climate-change-impacts/why-polar-vortex-keeps-breaking-out-arctic>
- UCDavis. (2019). *What is the polar vortex?* Retrieved from <https://climatechange.ucdavis.edu/climate-change-definitions/what-is-the-polar-vortex/>
- Wallace, J. M., Held, I. M., Thompson, D. W., Trenberth, K. E., & Walsh, J. E. (2014). Global warming and winter weather. *Science*, *343*(6172), 729–730. <https://doi.org/10.1126/science.343.6172.729>
- Waugh, D. W., Sobel, A. H., & Polvani, L. M. (2017). What is the polar vortex and how does it influence weather? *Bulletin of the American Meteorological Society*, *98*(1), 37–441. <https://doi.org/10.1175/BAMS-D-15-00212.1>
- Xiong, X., Liu, X., Wu, W., Yang, Q., & Zhou, D. K. (2021). *Polar vortex outbreak air transport: Observation using satellite IR sounder derived ozone product and comparison with model*. AGU Meeting Fall.
- Yu, B., & Zhang, X. (2015). A physical analysis of the severe 2013/2014 cold winter in North America. *Journal of Geophysical Research: Atmospheres*, *120*(19), 10149–10165. <https://doi.org/10.1002/2015JD023116>
- Zhang, X., Fu, Y., Han, Z., Overland, J. E., Rinke, A., Tang, H., et al. (2021). Extreme cold events from East Asia to North America in winter 2020/21: Comparisons, causes, and future implications. *Advances in Atmospheric Sciences*. <https://doi.org/10.1007/s00376-021-1229-1>