




Shine a light: Under-ice light and its ecological implications in a changing Arctic Ocean

Giulia Castellani , Gaëlle Veyssi re, Michael Karcher, Julienne Stroeve, S. Neil Banas, A. Heather Bouman, S. Andrew Brierley, Stacey Connan, Finlo Cottier, Fabian Gro e, Laura Hobbs, Christian Katlein, Bonnie Light, David McKee, Andrew Orkney, Roland Proud, Vibe Schourup-Kristensen

Received: 30 April 2021 / Revised: 11 October 2021 / Accepted: 26 October 2021 / Published online: 25 November 2021

Abstract The Arctic marine ecosystem is shaped by the seasonality of the solar cycle, spanning from 24-h light at the sea surface in summer to 24-h darkness in winter. The amount of light available for under-ice ecosystems is the result of different physical and biological processes that affect its path through atmosphere, snow, sea ice and water. In this article, we review the present state of knowledge of the abiotic (clouds, sea ice, snow, suspended matter) and biotic (sea ice algae and phytoplankton) controls on the underwater light field. We focus on how the available light affects the seasonal cycle of primary production (sympagic and pelagic) and discuss the sensitivity of ecosystems to changes in the light field based on model simulations. Lastly, we discuss predicted future changes in under-ice light as a consequence of climate change and their potential ecological implications, with the aim of providing a guide for future research.

Keywords Arctic ecosystem · Arctic Ocean · Light transmission · Phytoplankton · Primary production · Sea ice algae

INTRODUCTION

The reduction of Arctic sea ice is one of the strongest manifestations of global climate change. Besides shrinking in extent, sea ice properties are also changing, as the Arctic Ocean shifts towards a thinner (e.g. Renner et al. 2014) and younger (Maslanik et al. 2011) ice pack. This shift has profound implications for the structure of the remaining sea ice, for melt pond development, and for the amount of snow that accumulates. The open water season has become longer because sea ice is forming later and melting earlier (e.g. Stroeve and Notz 2018). These changes in sea ice

strongly modulate the underwater light field (i.e. its intensity and spectral composition) leading to an increase in light penetrating through the ice cover into the water (Nicolaus et al. 2012).

Light and nutrients are key drivers of Arctic ecosystem dynamics. Primary producers within the sea ice (sea ice algae) and in the underlying ocean (phytoplankton) require light for growth (e.g. Michel et al. 1988). Therefore, changes in light availability can have a significant impact on Arctic primary production. Ice algae and phytoplankton form the basis of the Arctic marine food web, thus changes in primary production will have cascading effects on higher trophic level species such as fish, birds, and mammals (e.g. Steiner et al. 2019). However, food web responses may not be linear since the timing, as well as the magnitude of production are important. Mismatches in the timing of the blooms of sea ice algae and phytoplankton—principally regulated by light—and the timing of the zooplankton bloom (termed secondary production)—principally regulated by temperature (Richardson 2008)—may decouple primary and secondary production, with consequences for fish and higher trophic levels (S reide et al. 2010). Furthermore, many higher trophic level predator–prey interactions are themselves regulated by light (e.g. Hobbs et al. 2021), as the ability of visual predators (e.g. fish, birds) to detect prey is a function of available light (as well as visual acuity and prey size).

Monitoring under-ice light levels and ecosystem responses is crucial for better understanding the effects of ongoing changes on sympagic (ice associated) and pelagic (water column) Arctic ecosystems. Obtaining observations in extreme conditions, which are characteristic of the polar environment, remains a challenge. We are, therefore, largely reliant upon numerical models and satellite products to quantify large-scale changes in the light field and to

predict associated ecological implications. Recent modelling studies have shown a marked increase in light conditions favourable for under-ice blooms over the last two decades (Horvat et al. 2017) and pointed to the controlling role that shortwave radiation has on the magnitude of phytoplankton bloom (Popova et al. 2010). Moreover, CMIP5 simulations (Tedesco et al. 2019) indicate sympagic primary production, which is triggered by light availability, will begin earlier and increase at most latitudes under modelled climate change scenarios. Thus, the parameterization of the under-ice light field in numerical models is crucial to properly represent future trends in high-latitude ocean primary production. In this study, we review the physical and biological processes that alter and attenuate light in its journey through the atmosphere, the snow and ice cover, and the upper ocean (“Incoming light” to “Light propagation through seawater—the role and sources of CDOM and SPM” sections, Fig. 1). Focus is on the parameterizations of light transmission through sea ice and snow which are often used in large-scale sea ice-ocean models (e.g. CMIP6, Eyring et al. 2016) and recently applied to satellite data (Stroeve et al. 2021). We elucidate the role of primary producers on light absorption, as well as their dependency on light (“Sea ice algae and phytoplankton” section). By means of numerical simulations (“Sensitivity of sea ice algae and phytoplankton to light transmission parameterization” section), we show how the choice of light transmission parameterizations and their parameters affect the simulated under-ice light field and, as

consequence, the sympagic and pelagic ecosystems. Finally, we consider possible societal implications of future changes in the Arctic (“Expected future changes and socio-economic impact” section).

INCOMING LIGHT

Light availability at the ocean surface in the Arctic is primarily governed by seasonal changes in solar zenith angle and cloud cover (Fig. 1). In the range relevant for polar applications, incoming light decreases almost linearly with solar zenith angle above 50 degrees as illustrated in Fig. 2, upper panel. Clouds can be a dominant feature in the Arctic. Their impact remains limited for cloud cover up to ~ 30%, but it rapidly reduces surface irradiance for higher levels of cloud cover (Fig. 2, lower panel). 100% cloud cover reduces surface irradiance to ~ 20% of cloud free levels.

In coupled physical ecosystem models (e.g. SINMOD, Wassmann et al. 2006; BESTMAS + CLADACH, Banas et al. 2016) incoming light is often provided as shortwave radiation (300–3000 nm) by satellite available products and reanalysis data that usually include atmospheric conditions such as cloudiness. The part of the spectrum relevant for biological processes is termed PAR (Photosynthetically Active Radiation: 400 nm–700 nm) which is generally taken in the range 43–50% of the provided shortwave




Fig. 1 Schematic of light transmission through atmosphere, sea ice and snow, and ocean (credit: Andrew Orkney)




Fig. 2 Effect of varying solar zenith angle (upper panel), and cloud cover (lower panel) on surface PAR irradiance. Data are presented as the ratio of light intensity relative to the equator at noon on midsummer's day. The red box highlights solar zenith angles relevant for Arctic regions. All simulations done with Hydrolight Radiative Transfer Software that includes the RADTRAN sky model (Gregg and Carder 1990) and the cloud model from Kasten and Czeplak (1980)

radiation (e.g. Banas et al. 2016; Castellani et al. 2017; Schourup-Kristensen et al. 2018).

LIGHT ATTENUATION BY SEA ICE

Despite the notably high albedo (i.e. fraction of solar radiation reflected by the surface) of a sea ice cover, a fraction of the light is propagated through the ice, into the ocean. Sea ice attenuates light about ten times more strongly than clear ocean waters and in turn, snow attenuates ten times more strongly than the ice (Perovich 1996). Consequently, the under-ice light field averaged over large scale is only up to 20% of the incoming light (Katlein et al. 2019). Besides the thickness of the snow and ice cover, light transmission is strongly influenced by the internal structure of the ice and snow, and by the different surface elements that characterise the ice cover in different seasons (e.g. pressure ridges, melt ponds). While the large ice thickness in ridges reduces light levels underneath, their complex internal structure can lead to favourable light conditions within (Katlein et al. 2021). Melt ponds on top

of the ice can dramatically increase the transmittance of sea ice since they have a much lower albedo than the surrounding ice (Nicolaus et al. 2012; Light et al. 2015). Melt ponds can also introduce strong horizontal inhomogeneity to the light field in the water column (Frey et al. 2011). Further constituents deposited on top of the ice surface or within the ice matrix like sediment, Colored Dissolved Organic Matter (CDOM), soot, or algae can locally or temporally influence the intensity and spectral shape of the transmitted light.

Modelling light transmission through sea ice

In numerical ice-ocean models, such as those contributing to CMIP6 (Eyring et al. 2016), light attenuation in sea ice is represented at different levels of complexity. In the presence of sea ice, the amount of underwater light in each grid cell is the weighted average of light reaching the ice-free ocean surface and light transmitted through the ice cover. The simplest approach assumes zero transmittance for ice (bare, ponded, or snow-covered) and 0.93 (the difference between 1—maximum albedo—and an albedo of 0.07, the

value which is typically used for open water) transmittance for areas of open water (Perovich et al. 2007). More complex models use a variety of approximations and numerical techniques for computing light transmission in the ice-ocean domain. One such approach widely used in CMIP6 models, relies on a simple exponential description (Grenfell and Maykut 1977) with empirically derived extinction coefficients that represent the attenuation of light through snow and ice. The values of extinction coefficients vary according to season and surface characteristics. The value for bare ice is usually taken in the range 1.1 to 1.6 m⁻¹ (Perovich 1996; Grenfell and Maykut 1977), whereas the values for snow show a larger variability with values ranging from 4.3 to 40 m⁻¹ (Perovich 1996). In the case of bare summer ice, the absorption of solar radiation causes the above-freeboard ice to weather, become crumbly, and have a significantly larger air-ice interface, thus scattering light much more effectively than the ice beneath. The uppermost portion of this layer is termed the “surface scattering layer” (SSL), which is typically up to 10 cm thick (e.g. Light et al. 2008). To estimate light transmittance through the sea ice cover, the simple exponential models rely on an approximation that extinguishes a significant portion (up to 70%) of the incident light in the SSL. Thin ice with small freeboard, and in turn a rather thin SSL, forms a relevant fraction of the ice cover, especially in the marginal ice zone and in the seasons of ice formation/melt. Thus, the treatment of the SSL and of the light transmission through thin ice in models will impact the under-ice PAR, and consequently, the ecosystems (“[Sensitivity of sea ice algae and phytoplankton to light transmission parameterization](#)” section).

More sophisticated approaches include explicit treatment of multiple scattering and use inherent optical properties to compute the full radiation budget for all surface types present (Briegleb and Light 2007). Indeed, snow-covered ice, ponded ice, melting ice, and even bare ice all exhibit large vertical gradients in scattering. Such treatments are incorporated into only a few sea ice models (e.g. CICE, Holland et al. 2012).

LIGHT PROPAGATION THROUGH SEAWATER—THE ROLE AND SOURCES OF CDOM AND SPM

Once light reaches the ocean surface, it is further attenuated through absorption and scattering by sea water itself and by particles such as CDOM, Suspended Particulate Matter (SPM), and phytoplankton (“[Sea ice algae and phytoplankton](#)” section). Within the visible spectrum, absorption by pure water is two orders of magnitude stronger in the red than in the blue (e.g. Pope and Fry 1997), whereas scattering is an order of magnitude greater

in the blue than in the red. The scattering coefficient also depends on the salinity of the water, with values increasing by around 30% from freshwater to sea water (Morel 1966). CDOM primarily affects light propagation through absorption (Dall’Olmo et al. 2009), with CDOM absorption decreasing approximately exponentially from the blue to the red (e.g. Carder et al. 1989). Similarly, SPM absorbs highly in the blue and decays almost exponentially towards the red (Babin et al. 2003). Scattering by SPM exhibits similar behaviour in both organic and mineral form, with limited spectral variability in scattering coefficients (Lo Prejato et al. 2020).

CDOM in the Arctic is mainly provided by discharge from rivers. In open water, sources of SPM are mainly organic detritus, formed during phytoplankton blooms (Macquaker et al. 2010). In coastal regions, the sources of SPM are mainly mineral in origin, usually originating from coastal erosion, or run-off from rivers and land (Klein et al. 2019).

On a global scale, CDOM and SPM absorption in the blue is broadly equivalent to that of phytoplankton (Siegel et al. 2002). However, light attenuation by these constituents in marine ecosystem models is generally represented by very simplistic parameterizations. Commonly, attenuation in water in Arctic ecosystem models is a function of phytoplankton biomass with a constant background PAR attenuation, while the effect of CDOM/SPM in some instances is added as an average over a broad, heterogeneous ocean region (e.g. Banas et al. 2016). Other models represent attenuation by SPM and other dissolved organics as a function of salinity (e.g. Mei et al. 2010). However, in coastal regions, such as the Russian shelves highly affected by river discharge, the lack of an explicit treatment of CDOM might lead to biases in model simulations (e.g. Schourup-Kristensen et al. 2018).

SEA ICE ALGAE AND PHYTOPLANKTON

Light transmitted within or under sea ice may be absorbed or scattered by phytoplankton and ice algae (Kirk 1994). The absorption of visible radiation by algal cells is spectrally dependent and relies on the presence of a range of chlorophyll and carotenoid molecules and biliproteins, each with their own characteristic absorption spectrum. There are also strong differences in the efficiency of light absorption by pelagic and sympagic algae based on their intracellular packaging of pigment molecules and the size and morphology of the cells (e.g. Chase et al. 2013). It is the harvesting of sunlight by algal pigments that ultimately powers the metabolism of primary producers. In contrast to absorption, resulting mainly from chemicals stored in the cell interior, the amount, wavelength dependence, and

direction of scattering by phytoplankton cells depends on their size and exterior morphology.

Whilst there exists variation in the bio-optical properties of algal cells, the most important component of the light absorption spectra is the photosynthetic pigment chlorophyll *a* (chl *a*). Thus, the light absorptive properties of ice algae and phytoplankton, and the effect they have on the transmission of light through the water column, can largely be described as a function of chl *a* concentration. However, this may not be true for coastal waters, where CDOM derived from the land may enter the sea (“[Light propagation through seawater—the role and sources of CDOM and SPM](#)” section).

Photophysiology of Arctic algal assemblages—theoretical and modelling approach

The ways in which different algal assemblages interact with incident light fields vary. Hence, spatiotemporal variability in both the intensity and the spectral quality of light may favour some communities over others. Many Arctic algal communities also adopt different photophysiological strategies to cope with the highly variable light field in the Arctic. Despite the relief of darkness in spring, ice algae and many phytoplankton communities find themselves shaded beneath ice and snow. Shade-adapted algae include the diatoms *Nitzschia frigida*, the main species dominating sea ice algae communities, and *Melosira arctica* (Fig. 3). Such sympagic diatoms form mat-like colonies or filaments affixed to the sea ice subsurface. Despite very low light intensity, high pigmentation and occupying a stable band a few centimetres thick at the ice-water interface permit growth as early as February (Syvertsen 1991). During the summer months, many Arctic ice algae and phytoplankton assemblages have unique adaptations that let them survive relatively high light in summer without sacrificing their ability to grow in lower light in spring. An example are the photophysiological

adaptation strategies of diatom species: their intense pigmentation and concentrated communities result in self-shading, moderating their light environment (Barros et al. 2003). The ability of diatoms to adapt to a range of light environments, ranging from light-limited conditions beneath seasonal ice to intensely illuminated melt layers explains their widespread occurrence in both under-ice and ice-edge blooms (Degerlund and Eilertsen 2010).

Once a photon is absorbed by an algal cell, how efficiently the photosynthetic machinery of the cell can convert this into chemical energy (photophysiology) depends on several factors. Variation in algal photophysiology has been explicitly considered in Arctic ecosystem models for at least three decades (Slagstad and StØle-Hansen 1991). Many recent high-latitude models have represented the phytoplankton by two competing size classes (usually taken to represent diatoms and small flagellates), either with (Wassmann et al. 2006) or without (Vernet et al. 2017) including differences in photophysiology between the two groups. Banas et al. (2016) found that including seasonal photoacclimation in a one-phytoplankton model was sufficient, and also necessary, to capture the timing and magnitude of a high-latitude spring bloom in detail. The above cited model studies illustrate that the inclusion of flexible photophysiology within populations as well as photoresponse-based competition between functional types is required in high-latitude plankton models to properly represent timing and magnitude of the bloom.

SENSITIVITY OF SEA ICE ALGAE AND PHYTOPLANKTON TO LIGHT TRANSMISSION PARAMETERIZATION

Modelled PAR and growth of sea ice algae and phytoplankton are sensitive to light transmission parameterizations. In the case of the exponential model, largely used in CMIP6 models, the same holds for the chosen extinction




Fig. 3 Example of sea ice algae forming a mat-like colony under the ice (left panel, credit: Carsten Wancke), and of *Melosira arctica* forming filaments affixed to the sea ice subsurface (right panel, credit: Oliver Müller)

coefficients which determine the exponential decay in sea ice and snow, both of which are derived from observations, but are subject to large uncertainty (Katlein et al. 2019; Castellani et al. 2020). To illustrate the consequence of different treatments of light transmission through thin ice and through the SSL, and of the extinction coefficients, we present results of numerical experiments performed with the Finite Element Sea ice Ocean Model (FESOM) version 1.4 coupled to the ocean biogeochemical model REcoM2 (Schourup-Kristensen et al. 2018) and to the Sea Ice Model for Bottom Algae SIMBA (Castellani et al. 2017).

Experiment ‘standard’ assumes the standard thickness of the SSL of 10 cm (Light et al. 2008) for sea ice. Experiment ‘drainage’ investigates the impact of making the existence and thickness of the SSL dependent on the freeboard, assuming a linear increase of the SSL from 0 (for ice thickness below 50 cm) up to 10 cm (for sea ice thickness of 80 cm). The third experiment ‘drainage_2k_s’ investigates the effect of a doubled extinction coefficient for dry snow k_s from 10 to 20 m⁻¹. To highlight the sensitivity of the biological system to changes in light transmission parameterization, we focus on the seasonal evolution of in-ice chl *a* (as proxy for sea ice algae), and diatom net primary production (NPP) in the ocean (phytoplankton).

The introduction of a thickness-dependent SSL (‘drainage’) leads to a large increase of PAR in the summer months (Figs. 4 and 5). Particularly in the marginal areas, where ice is often thinner than 80 cm, under-ice PAR

increases up to twice that in the ‘standard’ simulation (Fig. 4). Differences in PAR start to appear in May–June (Fig. 5) and lead to a small increase in diatom NPP, limited to lower latitudes (70–85°N), but with no effect on the onset of the bloom. Differences remain negligible for sea ice algae. In contrast, a doubling of the extinction coefficient for snow (‘drainage_2xK_s’) already affects PAR in spring and early summer (Fig. 5), leading to a reduction of under-ice PAR in those months when snow did not yet melt completely. Using a larger k_s causes a delay of the sea ice algal bloom onset by up to 2 weeks in higher latitudes and a shift of the peak by more than a month (Fig. 5). The effects on the phytoplankton remain, however, negligible, since phytoplankton start to grow later in the season compared to sea ice algae, when the differences in PAR become negligible.

Our results show that the choice of parameterizations of light transmission and their coefficients affect underwater light, but differently according to season. The response of the ecosystems also differs, with sea ice algae being more sensitive than phytoplankton. Moreover, sea ice algae and phytoplankton have different photophysiology (“Sea ice algae and phytoplankton” section), which results in different timing of the bloom and thus different response to changes in the light field. These results show that how reliably we can project future changes in the magnitude and particularly in the timing of sea ice and ocean primary production is affected also by how we describe light transmission in models. With the aim to reliably project




Fig. 4 Under-ice PAR at the end of August 2012 for the case with a standard parameterization of the SSL (‘standard’ simulation, left hand side) and for ‘drainage’, when the SSL only exists if the sea ice is thicker than 50 cm (right hand side)




Fig. 5 Mean PAR (top row), mean sea ice algae chl *a* (middle row) and NPP of diatoms (bottom row) in latitude bands. The columns show on the left hand side: the standard parameterization of the SSL ('standard' simulation); in the middle: 'drainage'—existence of an SSL only if the sea ice thickness > 50 cm; and on the right hand side: 'drainage_2K_s' with a doubled extinction coefficients for snow K_s. For the simulations 'drainage' and 'drainage_2k_s', NPP and PAR are shown as differences 'drainage'—'standard' and 'drainage_k_s'—'drainage', respectively

future changes in sympagic and pelagic productivity, there is the need to theoretically constrain parameterizations (such as the representation of the SSL) and their parameters. Moreover, an intercomparison between model results adopting different parameterizations and parameters is needed in order to quantify the differences between formulations and their impact on ecosystems response.

EXPECTED FUTURE CHANGES AND SOCIO-ECONOMIC IMPACT

The current global warming trend is likely to result in future increase of light reaching the upper ocean (Lannuzel et al. 2020) with consequences for the temporal and spatial patterns of sea ice algae and phytoplankton growth. Phytoplankton growth is expected to shift northwards and occur earlier following a reduction in sea ice extent and thickness (Ardyna and Arrigo 2020). Climate models suggest that Arctic precipitation will transition from being snow to rain dominated leading to a reduced snow cover (Bintanja and Olivier 2017) and the increased likelihood that the sea ice algal bloom will happen earlier in the season (Post et al. 2013; Tedesco et al. 2019; Lannuzel et al. 2020). These shifts have the potential to significantly alter the composition and abundance of primary producers by favouring different photoadaptation strategies. Furthermore, such potential shifts in timing are likely to cause a mismatch between primary production and associated zooplankton grazing, thus compromising the life cycle of zooplankton (Sørreide et al. 2010).

Changes in the light field may also modify the vertical positioning of zooplankton as they negotiate the trade-offs between predation risk and feeding (Hobbs et al. 2021). Such migratory behaviour of zooplankton can actively draw carbon out of the surface waters making it an important contribution to the vertical carbon flux (Hansen and Visser 2016). Thus, a consequence of a change in light-mediated vertical migration is disruption of the biological pump.

The most substantial impacts on fish and seabirds are likely to be through visual predation. Increasing illumination will improve feeding conditions for epipelagic (i.e. that oceanic zone where enough light penetrates for photosynthesis) fish, and increased predation by fish may lead to changes in the size distribution of the zooplankton community (Varpe et al. 2015). Planktivorous fish themselves will be subject to increased predation risk, so increasing illumination may lead to wholesale changes in trophic transfer efficiency through food webs (Langbehn and Varpe 2017). However, observations and models both suggest that in some high-latitude regions the increases in primary production associated with low-ice conditions are actually inversely correlated with the success of fish, birds, and mammals, for reasons of plankton composition and timing (Banas et al. 2016). This gives the impression of a fragile ecosystem, but there are internal balances that may come into play to provide resilience in the form of variable life history strategies (Hobbs et al. 2020) and changes in species composition (Renaud et al. 2018) such that Arctic marine food webs may be more resilient to climate-related shifts than previously assumed.

Changes in both the physical environment (e.g. sea ice loss in coastal regions, loss of permafrost) and the ecosystem will affect ecosystem services, with strong societal and economic effects (O'Garra 2017). This is particularly relevant for communities based on fisheries, subsistence hunting and coastal infrastructure, but also on commercial activities making use of those ecosystem services. Further warming and ice loss may lead to further biogeographic shifts in fish distributions and perhaps to behavioural changes such as reduced school sizes (Brierley and Cox 2010). Following fish, larger Arctic species such as seals and whales are moving northwards into the Arctic basin. This will affect native populations, who rely on whales as a food source, as well as for their cultural heritage.

Increasing light intensity might lead via altered primary production to altered fish production, including a predicted increase in pelagic and planktivorous fish (Heath et al. this volume). There is growing concern that the next big global development in commercial fishing will target the meso-pelagic, where the estimated c. 10 GT fish biomass (Iri-goien et al. 2014) may be the planet's last remaining untapped source of protein. Fishing in the central Arctic Ocean is, however, prohibited until at least 2034 by binding legal agreement between multiple Arctic nations (Hoag 2017).

The complexity and interconnectivity of the social-ecological system in the Arctic, at a time where it experiences rapid changes, is thus calling for holistic studies to assess the impacts on the ecosystems and on human communities, as well as ways to respond.

Acknowledgements This study was supported by the Changing Arctic Ocean projects *EcoLight* (03V01465), *Arctic PRIZE* (NE/P00573X/1; NE/P005721/1), *Diatom-ARCTIC* (NE/R012849/1; 03F0810A), *MiMeMo* (NE/R012679/1), *CHASE*, *Peanuts* (NE/R012636/1) and *Diapod* (NE/P005985/1) jointly funded by the UKRI Natural Environment Research Council (NERC) and the German Federal Ministry of Education and Research (BMBF). VSK was funded by the FRontiers in Arctic marine Monitoring program (FRAM). This work benefited from the collaboration with the international working group BEPSII: Biogeochemical Exchange Processes at Sea Ice Interfaces.

Funding Open Access funding enabled and organized by Projekt DEAL.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright

holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

REFERENCES

- Ardyna, M., and K.R. Arrigo. 2020. Phytoplankton dynamics in a changing Arctic Ocean. *Nature Clinical Practice Endocrinology & Metabolism* 10: 892–903.
- Babin, M., A. Morel, V. Fournier-Sicre, F. Fell, and D. Stramski. 2003. Light scattering properties of marine particles in coastal and open ocean waters as related to the particle mass concentration. *Limnology and Oceanography* 48: 843–859.
- Banas, N., J. Zhang, R. Campbell, R. Sambrotto, M. Lomas, E. Sherr, B. Sherr, C. Ashjian, et al. 2016. Spring plankton dynamics in the Eastern Bering Sea, 1971–2050: Mechanisms of interannual variability diagnosed with a numerical model. *Journal of Geophysical Research: Oceans* 121: 1476–1501.
- Barros, M., M. Pedersén, P. Colepicolo, and P. Snoeijs. 2003. Self-shading protects phytoplankton communities against H₂O₂-induced oxidative damage. *Aquatic Microbial Ecology* 30: 275–282.
- Bintanja, R., and A. Olivier. 2017. Towards a rain-dominated Arctic. *Nature Climate Change* 7: 263–267.
- Briegleb, B. P. and Light, B. 2007. A delta-Eddington multiple scattering parameterization for solar radiation in the sea ice component of the community climate system model, NCAR/TN-472+STR.
- Brierley, A.S., and M.J. Cox. 2010. Shapes of krill swarms and fish schools emerge as aggregation members avoid predators and access oxygen. *Current Biology* 20: 1758–1762.
- Carder, K., R. Steward, G. Harvey, and P. Ortner. 1989. Marine humic and fulvic acids: Their effects on remote sensing of ocean chlorophyll. *Limnology and Oceanography* 34: 68–81.
- Castellani, G., M. Losch, B.A. Lange, and H. Flores. 2017. Modeling Arctic sea-ice algae: Physical drivers of spatial distribution and algae phenology. *Journal of Geophysical Research: Oceans*. <https://doi.org/10.1002/2017JC012828>.
- Castellani, G., F.L. Schaafsma, S. Arndt, B.A. Lange, I. Peeken, J. Ehrlich, C. David, R. Ricker, et al. 2020. Large-scale variability of physical and biological sea-ice properties in polar oceans. *Frontiers in Marine Science* 7: 536. <https://doi.org/10.3389/fmars.2020.00536>.
- Chase, A., E. Boss, R. Zaneveld, A. Bricaud, H. Claustre, J. Ras, G. Dall'Olmo, and T.K. Westberry. 2013. Decomposition of in situ particulate absorption spectra. *Methods in Oceanography* 7: 110–124.
- Dall'Olmo, G., T. Westberry, M. Behrenfeld, E. Boss, and W. Slade. 2009. Significant contribution of large particles to optical backscattering in the open ocean. *Biogeosciences* 6: 947–967.
- Degerlund, M., and H.C. Eilertsen. 2010. Main species characteristics of phytoplankton spring blooms in NE Atlantic and Arctic waters (68–80 N). *Estuaries and Coasts* 33: 242–269.
- Eyring, V., S. Bony, G.A. Meehl, C.A. Senior, B. Stevens, R.J. Stouffer, and K.E. Taylor. 2016. Overview of the coupled model intercomparison project phase 6 (CMIP6) experimental design and organization. *Geoscience Model Development* 9: 1937–1958. <https://doi.org/10.5194/gmd-9-1937-2016>.
- Frey, K.E., D.K. Perovich, and B. Light. 2011. The spatial distribution of solar radiation under a melting Arctic sea ice cover. *Geophysical Research Letters*. <https://doi.org/10.1029/2011GL049421>.

- Gregg, W., and K. Carder. 1990. A simple spectral solar irradiance model for cloudless maritime atmospheres. *Limnology and Oceanography* 35: 1657–1675.
- Grenfell, T.C., and G.A. Maykut. 1977. The optical properties of ice and snow in the arctic basin. *Journal of Glaciology* 18: 445–463.
- Hansen, A.N., and A.W. Visser. 2016. Carbon export by vertically migrating zooplankton. *Limnology and Oceanography* 61: 701–710.
- Hoag, H. 2017. Nations agree to ban fishing in Arctic Ocean for at least 16 years. *Science*. <https://doi.org/10.1126/science.aar6437>.
- Hobbs, L., N.S. Banas, F.R. Cottier, J. Berge, and M. Daase. 2020. Eat or sleep: Availability of winter prey explains mid-winter and spring activity in an Arctic Calanus population. *Frontiers in Marine Science* 7: 541564. <https://doi.org/10.3389/fmars.2020.541564>.
- Hobbs, L., N.S. Banas, J.H. Cohen, F.R. Cottier, J. Berge, and Ø. Varpe. 2021. A marine zooplankton community vertically structured by light across diel to interannual timescales. *Biology Letters* 17: 20200810.
- Holland, M.M., D.A. Bailey, B.P. Briegleb, B. Light, and E. Hunke. 2012. Improved sea ice shortwave radiation physics in CCSM4: The impact of melt ponds and aerosols on arctic sea ice. *Journal of Climate* 25: 1413–1430.
- Horvat, C., D.R. Jones, and S., Iams, Schroeder, D., Flocco, D., Feltham, D. 2017. The frequency and extent of sub-ice phytoplankton blooms in the Arctic Ocean. *Science Advances* 3: 3.
- Irigoin, X., T. Klevjer, A. Røstad, U. Martinez, G. Boyra, J.L. Acuña, A. Bode, F. Echevarria, et al. 2014. Large mesopelagic fishes biomass and trophic efficiency in the open ocean. *Nature Communications* 5: 3271.
- Katlein, C., S. Arndt, H.J. Belter, G. Castellani, and M. Nicolaus. 2019. Seasonal evolution of light transmission distributions through Arctic Sea Ice. *Journal of Geophysical Research: Oceans* 124: 5418–5435. <https://doi.org/10.1029/2018jc014833>.
- Katlein, C., J.-P. Langelier, A. Ouellet, F. Lévesque-Desrosiers, Q. Hisette, B.A. Lange, S. Lambert-Girard, M. Babin, and S. Thibault. 2021. Under Revision. The three-dimensional light field within sea ice ridges. *Earth and Space Science Open Archive*. <https://doi.org/10.1002/essoar.10506296.2>.
- Kasten, F., and G. Czeplak. 1980. Solar and terrestrial radiation dependent on the amount and type of cloud. *Solar Energy* 24: 177–189.
- Kirk, J.T. 1994. *Light and photosynthesis in aquatic ecosystems*. Cambridge University Press.
- Klein, K., H. Lantuit, B. Heim, F. Fell, D. Doxaran, and A. Irrgang. 2019. Long-term high-resolution sediment and sea surface temperature spatial patterns in Arctic Nearshore waters retrieved using 30-year landsat archive imagery. *Remote Sensing* 11: 2791.
- Langbehn, T.J., and Ø. Varpe. 2017. Sea-ice loss boosts visual search: Fish foraging and changing pelagic interactions in polar oceans. *Global Change Biology* 23: 5318–5330.
- Lannuzel, D., L. Tedesco, M. van Leeuwe, K. Campbell, H. Flores, B. Delille, L. Miller, J. Stefels, et al. 2020. The future of Arctic sea-ice biogeochemistry and ice-associated ecosystems. *Nature Clinical Practice Endocrinology & Metabolism* 10: 983–992.
- Light, B., T.C. Grenfell, and D.K. Perovich. 2008. Transmission and absorption of solar radiation by Arctic sea ice during the melt season. *Journal of Geophysical Research-Oceans*. <https://doi.org/10.1029/2006jc003977>.
- Light, B., D.K. Perovich, M.A. Webster, C. Polashenski, and R. Dacic. 2015. Optical properties of melting first-year Arctic sea ice. *Journal of Geophysical Research: Oceans* 120: 7657–7675. <https://doi.org/10.1002/2015JC011163>.
- Lo Prejato, M., D. McKee, and C. Mitchell. 2020. Inherent optical properties: Reflectance relationships revisited. *Journal of Geophysical Research: Oceans* 125: 10.
- Macquaker, J., M. Keller, and S. Davies. 2010. Algal Blooms and “marine snow”: Mechanisms that enhance preservation of organic carbon in ancient fine-grained sediments. *Journal of Sedimentary Research* 80: 934–942.
- Maslanik, J., J. Stroeve, C. Fowler, and W. Emery. 2011. Distribution and trends in Arctic sea ice age through spring 2011. *Geophysical Research Letters*. <https://doi.org/10.1029/2011GL047735>.
- Mei, Z., F. Saucier, V. Le Fouest, B. Zakardjian, S. Sennville, H. Xie, and M. Starr. 2010. Modeling the timing of spring phytoplankton bloom and biological production of the Gulf of St. Lawrence (Canada): Effects of colored dissolved organic matter and temperature. *Continental Shelf Research* 30: 2027–2042.
- Morel, A. 1966. Étude expérimentale de la diffusion de la lumière par l’eau, les solutions de chlorure de sodium et l’eau de mer optiquement pures. *Journal De Chimie Physique* 63: 1359–1367.
- Michel, C., L. Legendre, S. Demers, and J.-C. Therriault. 1988. Photoadaptation of sea-ice microalgae in springtime: Photosynthesis and carboxylating enzymes. *Marine Ecology Progress Series* 50: 177–185.
- Nicolaus, M., C. Katlein, J. Maslanik, and S. Hendricks. 2012. Changes in Arctic sea ice result in increasing light transmittance and absorption. *Geophysical Research Letters* 39: L24501. <https://doi.org/10.1029/2012gl053738>.
- O’Garra, T. 2017. Economic value of ecosystem services, minerals and oil in a melting Arctic: A preliminary assessment. *Ecosystem Services* 24: 180–186.
- Perovich, D. K. 1996. The optical properties of sea ice, *Monograph* 96-1.
- Perovich, D.K., B. Light, H. Eicken, K.F. Jones, K. Runciman, and S.V. Nghiem. 2007. Increasing solar heating of the Arctic Ocean and adjacent seas, 1979–2005: Attribution and role in the ice-albedo feedback. *Geophysical Research Letters* 34: L19505. <https://doi.org/10.1029/2007GL031480>.
- Pope, R., and E. Fry. 1997. Absorption spectrum (380–700 nm) of pure water II Integrating cavity measurements. *Applied Optics* 36: 8710.
- Popova, E.E., A. Yool, A.C. Coward, Y.K. Aksenov, S.G. Alderson, B.A. de Cuevas, and T.R. Anderson. 2010. Control of primary production in the Arctic by nutrients and light: Insights from a high resolution ocean general circulation model. *Biogeosciences* 7: 3569–3591.
- Post, E., U.S. Bhatt, C.M. Bitz, J.F. Brodie, T.L. Fulton, M. Hebblewhite, A.J. Kerby, S.J. Kutz, et al. 2013. Ecological consequences of sea-ice decline. *Science* 341: 519–524.
- Renaud, P.E., M. Daase, N.S. Banas, T.M. Gabrielsen, J.E. Søreide, Ø. Varpe, F. Cottier, Falk-Petersen, et al. 2018. Pelagic food-webs in a changing Arctic: A trait-based perspective suggests a mode of resilience. *ICES Journal of Marine Science*. <https://doi.org/10.1093/icesjms/fsy063>.
- Renner, A.H.H., S. Gerland, C. Haas, G. Spreen, J.F. Beckers, E. Hansen, M. Nicolaus, and H. Goodwin. 2014. Evidence of Arctic sea ice thinning from direct observations. *Geophysical Research Letters* 41: 5029–5036. <https://doi.org/10.1002/2014GL060369>.
- Richardson, A. 2008. In hot water: Zooplankton and climate change. *ICES Journal of Marine Science* 65: 279–295.
- Schourup-Kristensen, V., C. Wekerle, D.A. Wolf-Gladrow, and C. Völker. 2018. Arctic Ocean biogeochemistry in the high resolution FESOM 1.4-REcoM2 model. *Progress in Oceanography* 168: 65–81. <https://doi.org/10.1016/j.pocean.2018.09.006>.
- Siegel, D., S. Maritorena, N. Nelson, D. Hansell, and M. Lorenzi-Kayser, M. 2002. Global distribution and dynamics of colored dissolved and detrital organic materials. *Journal of Geophysical Research: Oceans*, 107(C12): 21-1-21-14.

- Slagstad, D., and K. Støle-Hansen. 1991. Dynamics of plankton growth in the Barents Sea: Model studies. *Polar Research* 10: 173–186.
- Søreide, J.E., E.V. Leu, J. Berge, M. Graeve, and S.T.I.G. Falk-Petersen. 2010. Timing of blooms, algal food quality and *Calanus glacialis* reproduction and growth in a changing Arctic. *Global Change Biology* 16: 3154–3163.
- Steiner, N.S., W.W.L. Cheung, A.M. Cisneros-Montemayor, H. Drost, H. Hayashida, C. Hoover, J. Lam, T. Sou, et al. 2019. Impacts of the changing ocean-sea ice system on the key forage fish Arctic cod (*Boreogadus saida*) and subsistence fisheries in the western Canadian Arctic—Evaluating linked climate, ecosystem and economic (CEE) models. *Frontiers in Marine Science* 6: 179.
- Stroeve, J., and D. Notz. 2018. Changing state of Arctic sea ice across all seasons. *Environmental Research Letters* 13: 103001.
- Stroeve, J., M. Vancoppenolle, G. Veyssiere, M. Lebrun, G. Castellani, M. Babin, M. Karcher, J. Landy, et al. 2021. A multi-sensor and modeling approach for mapping light under sea ice during the ice-growth season. *Frontiers Marine Science* 7: 592337.
- Syvertsen, E.E. 1991. Ice algae in the Barents Sea: Types of assemblages, origin, fate and role in the ice-edge phytoplankton bloom. *Polar Research* 10: 277–288.
- Tedesco, L., M. Vichi, and E. Scoccimarro. 2019. Sea-ice algal phenology in a warmer Arctic. *Science Advance* 5: 4830.
- Varpe, Ø., M. Daase, and T. Kristiansen. 2015. A fish-eye view on the new Arctic lightscape. *ICES Journal of Marine Science* 72: 2532–2538.
- Vernet, M., T.L. Richardson, K. Mefies, E.-M. Nothig, and I. Peeken. 2017. Models of plankton community changes during a warm water anomaly in Arctic waters show altered trophic pathways with minimal changes in carbon export. *Frontiers in Marine Science* 4: 160.
- Wassmann, P., D. Slagstad, C.W. Riser, and M. Reigstad. 2006. Modelling the ecosystem dynamics of the Barents Sea including the marginal ice zone: II. Carbon flux and interannual variability. *Journal of Marine Systems* 59: 1–24.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

AUTHOR BIOGRAPHIES

Giulia Castellani (✉) is a researcher at the Alfred-Wegener-Institute Helmholtz-Zentrum für Polar und Meeresforschung. Her research interests include sea ice physics, sea ice ecology and implication of ice loss for marine ecosystems in polar regions.
Address: Alfred-Wegener-Institute Helmholtz-Zentrum für Polar und Meeresforschung, Am Handelshafen 12, 27570 Bremerhaven, Germany.
 e-mail: giulia.castellani@awi.de

Gaëlle Veyssière is a Postdoctoral Researcher at the British Antarctic Survey. Her research interests include snow and sea ice physics as well as remote sensing of the cryosphere.
Address: British Antarctic Survey, High Cross Madingley Road, Cambridge CB3 0ET, UK.
 e-mail: gaevey@bas.ac.uk

Michael Karcher is a senior scientist at the Alfred-Wegener-Institut Helmholtz-Zentrum für Polar und Meeresforschung and co- leads the science company Ocean Atmosphere Systems GmbH. His research interests include the physics of the sea ice and ocean system.

Address: Alfred-Wegener-Institute Helmholtz-Zentrum für Polar und Meeresforschung, Am Handelshafen 12, 27570 Bremerhaven, Germany.

Address: Ocean Atmosphere Systems GmbH, Teweesteg 4, 20249 Hamburg, Germany.
 e-mail: Michael.Karcher@awi.de; michael@oasys-research.de

Julienne Stroeve is a Canada 150 Chair at the University of Manitoba, Professor at University College London and Senior Scientist at the National Snow and Ice Data Center and the Cooperative Institute for Research in Environmental Science at the University of Colorado. Her interests are focussed on remote sensing of polar regions and implications of Arctic ice loss.

Address: University College London, Gower St, London WC1E 6BT, UK.

Address: University of Manitoba, 66 Chancellors Cir, Winnipeg, MB R3T 2N2, Canada.

Address: National Snow and Ice Data Center CIRES, 449 UCB University of Colorado, Boulder, CO 80309-0449, USA.
 e-mail: j.stroeve@ucl.ac.uk

S. Neil Banas is a Reader at the University of Strathclyde. His research interests include biophysical ocean modelling, plankton dynamics and climate impacts on marine ecosystems.

Address: University of Strathclyde, Livingstone Tower, 26 Richmond Street, Glasgow G1 1XH, UK.

e-mail: neil.banas@strath.ac.uk

A. Heather Bouman is an Associate Professor of Marine Biogeochemistry at the University of Oxford. Her research interests include the measurement and modelling of marine primary production, phytoplankton ecophysiology, marine optics and remote sensing of ocean colour.

Address: University of Oxford, South Parks Road, Oxford OX1 3AN, UK.

e-mail: heather.bouman@earth.ox.ac.uk

S. Andrew Brierley is a Professor of Biology at the University of St Andrews. His research interests include spatial and temporal variability in communities of open-ocean zooplankton and fish, and the consequences of this variability for higher predators (and fishers) and vertical carbon flux.

Address: Pelagic Ecology Research Group, Scottish Oceans Institute, Gatty Marine Laboratory, School of Biology, University of St Andrews, Fife KY16 8LB, Scotland, UK.

e-mail: asb4@standrews.ac.uk

Stacey Connan is a doctoral candidate at the University of Strathclyde. Her research interests include optimised modelling approaches to the transmission and propagation of spectrally resolved light fields in the Arctic Ocean and the influence a changing light field will have on primary productivity and animal behaviour.

Address: University of Strathclyde, Livingstone Tower, 26 Richmond Street, Glasgow G1 1XH, UK.

e-mail: stacey.connan@strath.ac.uk

Finlo Cottier is a Professor of Oceanography at the Scottish Association for Marine Science and holds an adjunct Professorship in Arctic Oceanography at the Department of Arctic Marine Biology, The Arctic University of Norway, Tromsø. His research interests span the coupled physical and ecological processes occurring in polar seas, particularly Arctic shelves.

Address: Scottish Association for Marine Science, Oban, Argyll and Bute PA37 1QA, Scotland, UK.

e-mail: finlo.cottier@sams.ac.uk

Fabian Große is a Research Scientist at the Department for Microbiology at the German Federal Institute of Hydrology, and a Visiting Researcher at the Department of Mathematics and Statistics of the University of Strathclyde. His research includes coupled physical-biogeochemical modelling of Arctic and coastal ecosystems.

Address: University of Strathclyde, Livingstone Tower, 26 Richmond Street, Glasgow G1 1XH, UK.

Address: German Federal Institute of Hydrology, Department for Microbiology, Am Mainzer Tor 1, 56068 Koblenz, Germany.

e-mail: grosse@bafg.de

Laura Hobbs is a Postdoctoral Researcher at the Scottish Association for Marine Science and University of Strathclyde. Her interests include acoustic observations of zooplankton and fish, chronobiology, and modelling.

Address: University of Strathclyde, Livingstone Tower, 26 Richmond Street, Glasgow G1 1XH, UK.

e-mail: laura.hobbs@strath.ac.uk

Christian Katlein is a Sea Ice Physicist at the Alfred-Wegener-Institut Helmholtz-Zentrum für Polar und Meeresforschung. His research interests include sea ice physics and polar observation technology.

Address: Alfred-Wegener-Institute Helmholtz-Zentrum für Polar und Meeresforschung, Am Handelshafen 12, 27570 Bremerhaven, Germany.

e-mail: ckatlein@awi.de

Bonnie Light is a Senior Principal Scientist at the Polar Science Center at the Applied Physics Laboratory, University of Washington. Her research focuses on the partitioning of solar radiation by sea ice covers.

Address: University of Washington, Seattle, WA 98195, USA.

e-mail: bonlight@uw.edu

David McKee is a Reader in the Physics Department at the University of Strathclyde and holds an adjunct Associate Professorship at the Department of Arctic Marine Biology, The Arctic

University of Norway, Tromsø. His research interests span the measurement of light in the ocean and use of light to measure biogeochemical properties of the ocean.

Address: University of Strathclyde, Livingstone Tower, 26 Richmond Street, Glasgow G1 1XH, UK.

e-mail: david.mckee@strath.ac.uk

Andrew Orkney is a doctoral candidate at the department of Earth Sciences of the University of Oxford. His interests include Arctic ecosystem dynamics, remote sensing of Phytoplankton Functional Types, and vertebrate evolution.

Address: University of Oxford, South Parks Road, Oxford OX1 3AN, UK.

e-mail: andrew.orkney@earth.ox.ac.uk

Roland Proud is a postdoctoral research fellow at the University of St Andrews. His research interests include the ecology and biology of open-ocean zooplankton and micronekton. His research includes sampling zooplankton/fish using scientific echosounders and nets, and mining acoustic 'big data'.

Address: Pelagic Ecology Research Group, Scottish Oceans Institute, Gatty Marine Laboratory, School of Biology, University of St Andrews, Fife KY16 8LB, Scotland, UK.

e-mail: rp43@st-andrews.ac.uk

Vibe Schourup-Kristensen is a researcher at Aarhus University in the department of Applied Marine Ecology and Modeling. In her work she develops large-scale and coastal ocean biogeochemical models and applies them to increase our understanding of the impact of the changing physical environment in the Arctic Ocean on the marine biogeochemical cycles.

Address: Department of Applied Marine Ecology and Modeling, Aarhus University, Nordre Ringgade 1, 8000 Aarhus C, Denmark.

e-mail: vibe.schourup@ecos.au.dk