



RESEARCH ARTICLE

10.1029/2021JA030045

Improving the Thermosphere Ionosphere in a Whole Atmosphere Model by Assimilating GOLD Disk Temperatures

F. I. Laskar¹ , N. M. Pedatella² , M. V. Codrescu³ , R. W. Eastes¹ , and W. E. McClintock¹ 

¹Laboratory for Atmospheric and Space Physics, University of Colorado Boulder, Boulder, CO, USA, ²High Altitude Observatory, National Center for Atmospheric Research, Boulder, CO, USA, ³Space Weather Prediction Center, NOAA, Boulder, CO, USA

Key Points:

- A new approach has been developed to assimilate Global-Scale Observations of Limb and Disk (GOLD) T_{disk} in whole atmosphere community climate model with thermosphere-ionosphere eXtensions which is validated using independent measurements
- Analysis states of both the thermosphere and ionosphere show improved agreement with independent measurements
- Results demonstrate a great potential of the GOLD T_{disk} data to improve thermosphere-ionosphere data assimilation

Correspondence to:

F. I. Laskar,
Fazlul.Laskar@Lasp.colorado.edu

Citation:

Laskar, F. I., Pedatella, N. M., Codrescu, M. V., Eastes, R. W., & McClintock, W. E. (2022). Improving the thermosphere ionosphere in a whole atmosphere model by assimilating GOLD disk temperatures. *Journal of Geophysical Research: Space Physics*, 127, e2021JA030045. <https://doi.org/10.1029/2021JA030045>

Received 15 OCT 2021

Accepted 25 FEB 2022

Abstract Global-Scale Observations of Limb and Disk (GOLD) disk measurements of far ultraviolet molecular nitrogen band emissions are used to retrieve column integrated disk temperatures (T_{disk}), which are representative of the lower-and-middle thermosphere. The present work develops a new approach to assimilate the T_{disk} in the whole atmosphere community climate model with thermosphere-ionosphere eXtension using the data assimilation research testbed ensemble adjustment Kalman filter. Nine days of data, 1–9 November 2018, are assimilated. Analysis state variables such as thermospheric effective temperature (T_{eff} , airglow layer integrated temperature), ratio of atomic oxygen to molecular nitrogen column densities (O/N_2), and column electron content are compared with a control simulation that is only constrained up to ~ 50 km. It is observed that assimilation of the GOLD T_{disk} improves the analysis states when compared with the control simulation. The analysis and model states, particularly, T_{eff} , O/N_2 , and electron column density (ECD) are compared with their measurement counterparts for a validation of the assimilation. T_{eff} and O/N_2 are compared with GOLD T_{disk} and O/N_2 . While, the ECD is compared with ground based total electron content measurements from global navigational satellite system receivers. Root mean square error (RMSE) improvements in T_{eff} and O/N_2 are about 10.8% and 22.6%, respectively. The RMSE improvement in analyses ECD is about 10% compared to the control simulation.

Plain Language Summary Understanding the temperature and density variability of the thermosphere-ionosphere system is very important for satellite drag calculations and satellite communication. The thermosphere-ionosphere system is influenced by waves from the lower atmosphere and solar and geomagnetic forcing from above. For the characterization of this coupled system, realistic whole atmosphere ionosphere parameters are of great interest. The global-scale observations of limb and disk (GOLD) satellite mission provides daytime thermospheric temperature observations with unprecedented local time and spatial coverage. Including them with the lower and middle atmospheric observations in a whole atmosphere data assimilation system, we find that they improve the state of the thermosphere-ionosphere. This shows the promise of the GOLD disk temperatures in improving thermosphere-ionosphere states and their potential use to improve space weather forecast capabilities.

1. Introduction

Improvements in the satellite drag forecasts and satellite communication depend on a better understanding of the thermosphere-ionosphere (TI) system variability. Earth's TI system is coupled to the lower atmosphere by wave-dynamical forcing and to the solar and geomagnetic forcing from above. The lower atmospheric forcing also varies with location and time. Thus, for a better understanding of this coupled system, a global four dimensional data set with good temporal and spatial resolution is needed. Satellite measurements from low-Earth orbit can provide good spatial coverage, but they lack local time coverage, unless a constellation of satellites is used. Ground based observations on the other hand have good local time coverage, but they are not available globally due to the significant fraction of the Earth that is covered by ocean. Moreover, the currently available whole atmosphere ionosphere thermosphere observations have data gaps at different altitudes and geographic locations. However, the currently available observations and state-of-the-art whole atmosphere model simulations can be combined in a data assimilation framework. Data assimilation combines observations with model forecasts to produce analysis states that can better estimate the current state of the TI system.

©2022. The Authors.

This is an open access article under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

With time the whole atmosphere ionosphere thermosphere models are improving, and number of observations from the TI system and lower atmosphere are increasing. Therefore, we are in a great stage to do a whole atmosphere data assimilation by combing the models and the observations. There is a long-history of lower atmosphere data assimilation (Gelaro et al., 2017; Hersbach et al., 2020; Rienecker et al., 2011), but the whole atmosphere system data assimilation is relatively new. There have been significant developments in the assimilation of thermosphere-ionosphere observations such as, neutral density (Matsuo et al., 2013; Mehta et al., 2018; M. V. Codrescu et al., 2004; Ren & Lei, 2020; S. M. Codrescu et al., 2018; Sutton, 2018), thermospheric temperature (Laskar, Pedatella, et al., 2021), thermospheric airglow radiance (Cantrall et al., 2019), and electron content (Aa et al., 2016; Bust et al., 2004; Bust & Immel, 2020; Chen et al., 2016; Datta-Barua et al., 2013; Forsythe et al., 2021; He et al., 2020; Kodikara et al., 2021; Lee et al., 2012; Lin et al., 2015; Matsuo et al., 2013; Pedatella et al., 2020; Song et al., 2021). While these results were promising and showed that the assimilation of TI observations improves the model states, most were limited to using upper atmosphere only models or used limited thermospheric datasets from low-earth-orbit satellites or ionospheric only measurements or observing system simulation experiments. Furthermore, a majority of them have not combined lower, middle, and upper atmosphere data in the assimilation. Also, the spatial and temporal coverage of thermospheric data available earlier were limited.

Temperature is one of the basic parameters in whole atmosphere models. Neutral temperature retrieved from Global-scale Observations of Limb and Disk (GOLD) disk measurements have increased the number of thermospheric observation in the recent years, which enables scope for a better whole atmosphere data assimilation that can potentially improve the specification of the TI system. Laskar, Pedatella, et al. (2021) performed a set of Observing System Simulation Experiments (OSSEs) to evaluate the impact of assimilating GOLD disk temperature (T_{disk}) observations on thermospheric temperature and dynamics. They found that the OSSE that includes the GOLD T_{disk} improved the model temperature root mean square error (RMSE) and bias by 5% and 71% when compared with the forecast state, and the improvements are 20% and 94% when compared with lower atmosphere only assimilation. Laskar, Pedatella, et al. (2021) also found that the migrating diurnal tide (DW1) and local diurnal tide over Americas improve by about 8% and 17%, respectively, upon assimilation of GOLD disk temperature (T_{disk}) observations. In the current study we assimilate actual GOLD T_{disk} in a whole atmosphere data assimilation system and assess their impact on the thermosphere-ionosphere parameters by validating analysis states with their measurement counterparts.

2. Data, Models, and Methodology

The primary data set used is the GOLD T_{disk} , which has been assimilated in the whole atmosphere community climate model with thermosphere-ionosphere eXtension (WACCMX). In addition to T_{disk} , lower and middle atmosphere data have also been assimilated. For validation of the analysis states from the assimilation system, independent measurements of GOLD O/N₂ and Global Navigation Satellite System Total Electron Content (GNSS-TEC) are also used. Further details of these data and models are given below.

2.1. GOLD T_{disk}

GOLD observed the Earth's thermosphere in the far ultraviolet wavelengths for over 18.5 hr each day, from 0610 to 0040 Universal Time (UT) of the next day (Eastes et al., 2019, 2020; McClintock et al., 2020; Laskar et al., 2020). The primary GOLD observations are emission intensities in the far ultraviolet (FUV) range of 134.5–166.5 nm. Data for one full disk scan are available at every 30 min from 6 to 23 UT (Eastes et al., 2019, 2020; Laskar, Eastes, et al., 2021). The current investigation uses level 2 T_{disk} data (version 3), that are retrieved from 2×2 binned level-1C data, which are available in the GOLD web-page, <https://gold.cs.ucf.edu/> as 'Level 2—TDISK'. The retrieval algorithm is an improvement of the previously used methods for limb measurements (Aksnes et al., 2006; Krywonos et al., 2012).

The 2×2 binned data have a spatial resolution of 250×250 km near nadir and it gets slightly coarse at view angles higher than 45° from nadir. The GOLD daytime disk scans in N₂ Lyman-Birge-Hopfield (LBH) bands are used to retrieve T_{disk} data. Effective altitude and contribution function (CF) of the T_{disk} varies with solar zenith angle (SZA) and emission angle (EA). The SZA variation of the CF is well quantified (Laskar, Pedatella, et al., 2021) and thus is included in the present assimilation. However, the EA effects are not yet included in the assimilation. But, it has been observed that the EA does not have significant impact on the CF for EAs below




Figure 1. (a) Geo-locations, (b) altitude or pressure and number of observations that are assimilated successfully during a representative hour on a particular day are shown. The red points show the Global-scale Observations of Limb and Disk observations and blue points are the rest of the observations, which we term as lower atmosphere observations including SABER and MLS. MLS, Microwave Limb Sounder; SABER, Sounding of the Atmosphere using Broadband Emission Radiometry.

50° , so the T_{disk} data having $EA > 50^\circ$ are not included in this assimilation and analysis. This limit also restricts the latitude and longitude coverage, as shown in Figure 1, to about $\pm 50^\circ$ in latitude and about -10°W to -90°W in longitude. Also, for high SZA observations the signal-to-noise ratio (SNR) is low, which for the current V3 T_{disk} introduces a bias. Thus, the low SNR observations having $SZA > 65^\circ$ are not considered in the analysis and assimilation.

2.2. GOLD O/N_2

GOLD disk measurements of OI-135.6 nm emission and N_2 -LBH bands in the $\sim 134\text{--}162$ nm wavelength range are used to retrieve the ratio of atomic oxygen to molecular nitrogen column densities ($\Sigma O/\Sigma N_2$) (Correia et al., 2021). For simplicity we use the notation O/N_2 instead of $\Sigma O/\Sigma N_2$. The disk O/N_2 has the same spatial and temporal coverage as T_{disk} . O/N_2 data are used here only for the comparison and validation of the analyses O/N_2 . We use the 2×2 binned version 3 O/N_2 data, named as ‘Level 2— ON_2 ’ in the GOLD data repository. Also, as the GOLD O/N_2 is not optimized for auroral latitudes (Correia et al., 2021), the latitudes above $\pm 60^\circ$ are not used in the current analysis. Typical random, systematic, and model uncertainties in the GOLD O/N_2 are about 5%, 5%, and 30%–40%, respectively. Note that the model uncertainty is a bias with an unknown sign (Correia et al., 2021).

2.3. GNSS-TEC

The GNSS-TEC data used in this study are obtained from the madrigal database (<https://cedar.openmadrigal.org>). Madrigal TEC maps are derived from worldwide Global Navigational Satellite System (GNSS) ground-based receivers. The vertical TEC data are available at 5 min temporal and 1° by 1° spatial bins. Details on the TEC retrieval algorithm can be found in Rideout and Coster (2006) and Vierinen et al. (2016). In the current study the TEC maps are averaged over 20 min centered at every UT hour to compare them with the analysis ECD from assimilation. The 20 min averaging is chosen to get enough satellite passes over a particular spatial grid.

2.4. WACCMX

The WACCMX version 2.1 is a whole atmosphere general circulation model extending from the surface to the upper thermosphere (~500–700 km depending on solar activity) (Liu et al., 2018). Whole Atmosphere Community Climate Model with thermosphere-ionosphere eXtension includes the chemical, dynamical, and physical processes that are necessary to model the lower, middle, and upper atmospheres. The thermosphere and ionosphere processes in WACCMX are similar to those in the NCAR Thermosphere-Ionosphere-Electrodynamics General Circulation Model (TIE-GCM), including the transport of O⁺ and self-consistent electrodynamics as well as realistic solar and geomagnetic forcing. The model horizontal resolution is 1.9° × 2.5° in latitude and longitude, and the vertical resolution is 0.25 scale height above ~50 km.

2.5. SD-WACCMX

In this simulation the WACCMX horizontal winds and temperature are relaxed toward Modern-Era Retrospective analysis for Research and Applications, Version 2 (MERRA2) (Gelaro et al., 2017; Rienecker et al., 2011), so we name it as Specified Dynamics WACCMX (SD-WACCMX). The relaxation or nudging to MERRA2 is up to 50 km altitude, and the model is free-running above this altitude (Marsh, 2011). The SD-WACCMX is used in this study as a control simulation. In addition to MERRA2, SD-WACCMX simulations (often referred here as SD) also use operational solar F10.7 cm flux and geomagnetic K_p index for forcing and thus they can be used as a control simulation for the assessment of the data assimilation states.

2.6. WACCMX + DART

The data assimilation capability in WACCMX was initially implemented by Pedatella et al. (2018) using DART (J. Anderson et al., 2009), which uses the ensemble adjustment Kalman filter (J. L. Anderson, 2001). In the present work we assimilate lower and middle atmosphere as well as thermosphere observations in the WACCMX + DART. The lower atmosphere measurements include conventional meteorological observations (i.e., temperatures and winds from aircraft, radiosonde measurements, etc.), as well as GNSS radio occultation refractivity. Assimilation of these observations improves specifications of the troposphere-stratosphere globally, which is important for the studies of the vertical coupling of waves from lower-atmosphere to the thermosphere (McCormack et al., 2017; Pedatella et al., 2014, 2018; Wang et al., 2011).

In addition to lower altitude observations, middle atmosphere temperatures from Sounding of the Atmosphere using Broadband Emission Radiometry (SABER) instrument on the Thermosphere Ionosphere Mesosphere Energetics and Dynamics (TIMED) satellite and Aura Microwave Limb Sounder (Aura-MLS) are also used. Altitude coverage of temperature profiles extends from stratosphere to mesosphere-lower-thermosphere (MLT) altitudes (~15–105 km for TIMED-SABER and ~15–90 km for Aura-MLS). The latitude coverage of TIMED-SABER retrieved temperature alternates between 83°S and 52°N (south viewing mode) and 83°N–52°S (north-viewing mode) (Remsberg et al., 2008). We performed 9 days (1–9 November 2018) of data assimilation, during which TIMED-SABER was in the north-viewing mode on 1 November only. From 2 to 9 November it was in the south viewing mode. While for the Aura-MLS it varies from 82°S to 82°N (Schwartz et al., 2008). Though Aura-MLS and TIMED-SABER temperatures are middle atmospheric observations, for simplicity we refer to them here as part of lower atmosphere observations. Assimilation of these data has previously been demonstrated to improve specification of the MLT state and dynamics (Laskar et al., 2019; McCormack et al., 2017; Pedatella et al., 2014).

In addition to lower atmosphere observations, GOLD T_{disk} are used in the whole atmosphere assimilation. As the thermospheric dynamics can quickly change in response to changes in forcing conditions, we use a 1 hr assimilation frequency. Additionally, Pedatella et al. (2020) have shown that using a 1 hr data assimilation cycle and removal of second-order divergence damping in WACCMX + DART significantly improves tidal amplitudes, which were previously found to be too small (Pedatella et al., 2018). As full disk images are available at 30 min intervals during sunlit hours, a 1 hr interval will have sufficient data in the thermosphere. Also, the lower atmosphere analysis states in WACCMX + DART agree well with other lower atmospheric assimilations, for example, MERRA2 (McCormack et al., 2021).

Figure 1 shows the locations (in a) and altitude or pressure versus number of observations (in b) that are assimilated successfully during a representative hour on a particular day. The red points show the GOLD observations

Table 1

Whole Atmosphere Community Climate Model With Thermosphere-Lonosphere Extension Simulation and Data Assimilation Experiments Used in This Study are Listed

Experiment	Observations assimilated	Nudging used	Model states updated
SD (SD-WACCMX, Control experiment)	N/A	MERRA2 U, V, T up to 50 km	N/A
DA1 (WACCMX + DART experiment 1)	Meteorological, Aura-MLS-T, SABER-T, GOLD-T _{disk}	N/A	T
DA2 (WACCMX + DART experiment 2)	Same as DA1	N/A	T, O, O ₂ , O ⁺

Note. U, V, T, N/A, SD, and DA stands for zonal wind, meridional wind, temperature, not applicable, specified dynamics, and data assimilation, respectively. Also, O, O₂, and O⁺ refers to the mass mixing ratio of atomic oxygen, molecular oxygen, oxygen ion, respectively.

and blue points are the rest of the observations, which we term as lower atmosphere observations, including TIMED-SABER and Aura-MLS. Note that the peak altitude of the N₂-LBH emission is shown here as a representative altitude of about 150 km, but in the assimilation the impact of T_{disk} is distributed over altitudes based on the SZA dependent CF (Laskar, Pedatella, et al., 2021). One can see that about 70,000 observations per hour are assimilated. On average about 1.5 million observations per day are assimilated. The simulations used in this study are listed in Table 1. The SD-WACCMX is used in this study as the control simulation.

We have performed two WACCMX + DART assimilations. One that assimilates lower atmosphere and GOLD T_{disk} observations, but the direct impact of T_{disk} has been restricted only to the model temperature, referred to as DA1 in Table 1. The second experiment assimilates the same observations as the first experiment, but the T_{disk} observations directly impact the model T, O, O₂, and O⁺, referred to as DA2 in Table 1. We used 40 ensemble members in the assimilation. In order to achieve sufficient spread in the ensemble members, we used Gaussian distributions of solar and geomagnetic forcing parameters with mean as the actual value and standard deviations of 15 sfu for F10.7 cm flux and 1 for Kp index (i.e., $d_{F10.7} \sim N(F10.7, 15^2)$ and $d_{Kp} \sim N(Kp, 1^2)$). We reset any F10.7 value less than 60 sfu to 60 sfu and any negative Kp to 0. The forcing perturbation for each ensemble member remains the same for all the days. To avoid artifacts arising from initial ensemble members, the spinup duration for the two assimilation runs are about 2 weeks that is, each assimilation run starts from 15 October 2018.

3. Results

In order to assess and validate the performance of the assimilation we compare the ensemble averaged analysis states to their measurement counterparts. For example, effective temperature (T_{eff}) from model simulation is compared with GOLD T_{disk}; O/N₂ is compared with GOLD O/N₂; and Electron Column Density (ECD) is compared with the GNSS-TEC. Note that T_{eff} here refers to the vertically integrated GOLD equivalent temperature that is calculated by integrating the model temperature profile weighted by the SZA dependent CFs. Also, the ECD is similar to TEC, but the column integration is only to the topmost layer of WACCMX, which is about 480 km for the current cases. Figure 2 shows a comparison of the local time and latitude variation of the GOLD T_{disk} with T_{eff} from ensemble averaged states of the DA1 (DA1 T_{eff}) and SD-WACCMX (SD T_{eff}) for 2 different days. The latitudes and local times are restricted to only those locations and times where GOLD T_{disk} is being assimilated. Beyond those local time and latitudes GOLD data are available, but we are not using them in the assimilation as explained in Sections 2.1 and 2.2. Note that in this figure only a representative longitude of 48°W is shown, which is close to the sub-satellite point of GOLD.

It can be noted from Figure 2 that the broad variations between T_{disk} and DA1 T_{eff} are similar on both the days. On 5 November 2018 there was a moderate geomagnetic storm for which the average temperature is more than 100 K higher than 3 November 2018. Moreover, the morning temperatures are relatively warmer, particularly between 40° and 50°S. These variations of the GOLD T_{disk} during geomagnetic events have been reported and discussed in Laskar, Eastes, et al. (2021). These results suggest that the data assimilation is driving the model temperature in the right direction that is, closer to those observed. A quantitative estimate of the differences between them are given later. Note that since both the assimilation experiments updated temperature directly at every assimilation step, the T_{eff} are almost the same for both DA cases. So, the T_{eff} for only the DA1 is shown here.




Figure 2. Local time and latitude variation of the Global-scale Observations of Limb and Disk T_{disk} compared with T_{eff} from DA1 (DA1 T_{eff}) and specified dynamics-Whole Atmosphere Community Climate Model with thermosphere-ionosphere extension (SD T_{eff}).

A change in temperature also impacts other states by altering the model dynamics. Therefore, assimilation of T_{disk} can also impact the O/N_2 ratio, which is another primary data set from the GOLD mission. Figure 3 shows a comparison of GOLD O/N_2 with the O/N_2 from data assimilation and control simulation experiments, for the same 2 days shown in Figure 2. Note that the model O/N_2 values are calculated by integrating the O and N_2 profiles down to the altitude corresponding to $1.5 \times 10^{21} \text{ m}^{-2}$ of N_2 , instead of 10^{21} m^{-2} as suggested by Strickland et al. (1995). The resulting O/N_2 values closely correspond to those from GOLD. Unlike Figure 2, here the latitude range is extended to 60°N/S , as the GOLD O/N_2 are valid for those latitudes. We compare O/N_2 from the DA1 (c and d), DA2 (e and f), and SD (g and h) with the GOLD O/N_2 (a and b). Note that the GOLD O/N_2 observations have not been assimilated in any of the experiments. In the DA2 the GOLD T_{disk} observations also




Figure 3. Same as Figure 2 but for the column integrated O/N_2 ratio. In addition to the DA1 the DA2 O/N_2 is also shown in (e and f).




Figure 4. The root mean square error (RMSEs) in DA1 T_{eff} and DA2 T_{eff} with respect to Global-scale Observations of Limb and Disk T_{disk} are shown in (a) and similar RMSEs in O/N_2 are shown in (b). Note that the temperature RMSEs in the two DA runs, are clearly smaller than the SD. Also, the average O/N_2 RMSEs are better for the two assimilation runs compared to the SD, and DA2 has the best RMSE.

directly update the O, O_2 , and O^+ mass mixing ratios in addition to temperature. The direct updating of these quantities impacts the neutral composition and ionosphere at every assimilation step and thus they are expected to compare better than the indirectly updated states. It can be observed from Figure 3 that the broad variations in O/N_2 agree well between GOLD O/N_2 and the two assimilation experiments. Though interhemispheric features in SD, the assimilation experiments, and the observations match well, there are clear differences in magnitudes and large-scale structures between them. For the quiet-day of 3rd November the two assimilation experiments show better agreement with GOLD O/N_2 compared to the SD O/N_2 . The highest discrepancy in O/N_2 can be seen on the storm day (right panel) where the Northern higher-latitude depletion in the GOLD O/N_2 occurs relatively at higher latitudes in DA1 and DA2 and is weaker in the SD.

For a quantitative estimation of the above observed differences between actual measurements and their data assimilation equivalents we calculate the Root Mean Square Error (RMSE). The RMSE in SD T_{eff} , DA1 T_{eff} , and DA2 T_{eff} with respect to GOLD T_{disk} are shown in Figure 4a for all 9 days. The RMSE for each day is calculated over the whole disk and local time range as shown in Figure 2 for temperature and Figure 3 for O/N_2 . Note that the temperature RMSEs in the two data assimilation runs are clearly smaller than the SD. Also, the temperature RMSE for the two assimilation runs are almost the same, which is expected as both the assimilations updated model temperature directly. The RMSEs in O/N_2 are shown in Figure 4b. The average O/N_2 RMSEs are better for the two assimilation runs compared with the SD, and DA2 has the best RMSE. The pre-storm RMSEs are smaller compared with storm onset and recovery phase. Average RMSE improvements in effective temperature and O/N_2 compared to the SD are about 10.8% and 22.6%, respectively. The improvements of pre-storm RMSE in T_{disk} and O/N_2 are about 6.4% and 27.9% while during the storm they were about 15.5% and 17.4%, respectively. These results suggest that even though the storm times RMSEs are larger, the percentage improvements are larger too.

For a more robust diagnosis of the relationship between SD T_{eff} , DA1 T_{eff} , and DA2 T_{eff} with respect to GOLD T_{disk} for all the available latitudes and longitudes in the disk scans between 10 and 20 UT during 1–9 November 2018 we make scatter diagrams as shown in Figure 5, where the red color represents high density points. Red (solid) and blue (dashed) lines represent least square fitted straight line and one-to-one (45° slope or gradient equal to one line) relationship. Correlation coefficients and fitted linear equations are also given. From these scatter plots it can be seen that the majority of the T_{disk} versus DA2 T_{eff} points (in a) fall on the one-to-one line. But, for the T_{disk} versus SD T_{eff} (in e) comparison, the highest density observations (red points) deviate away from the one-to-one linear relationship. Also, the correlation coefficient and gradient of the fitted lines are better for the assimilation runs. Note that here also, only those observations are shown that fall within the 50° EA and 65° SZA limits. As the GOLD T_{disk} has higher spread compared to DA2 T_{eff} , DA1 T_{eff} , and SD T_{eff} the shape of the scatter plot is elongated toward the T_{disk} axis (in a, c, and e). Similar to temperature, the O/N_2 scatter diagrams are shown in Figures 5b, 5d, and 5f but the EA and SZA restrictions are not applied here. The correlation coefficients for O/N_2 are small, though they are statistically significant as p-values (probability that the correlation arises from noise) are zero, suggesting a weak linear relationship. As the high density (red) points are mostly located around a circle for the two assimilation cases, the linear correlation would not be a great measure of the relationship between them. Therefore, we calculated the RMSE for the two assimilations and SD with respect to GOLD O/N_2 . The RMSEs for the DA1, DA2, and SD with respect to GOLD are 0.20, 0.17, and 0.23, respectively, suggesting




Figure 5. Scatter diagram of the Global-scale Observations of Limb and Disk (GOLD) T_{disk} and O/N_2 compared to their DA2, DA1, and SD equivalents are shown. For this analysis all the disk scans between 10 and 20 UT during 1–9 November 2018 are used. The red regions in the scatter diagram represents highest density points. For the GOLD versus DA2 the highest density points distribute around the one-to-one line (dashed), particularly for the temperature. The comparisons with respect to SD for both temperature and O/N_2 , on the other hand, are not as good.

that the two DA runs perform better compared to SD. The distribution of points in the GOLD T_{disk} versus DA1 T_{eff} and GOLD T_{disk} versus DA2 T_{eff} is nearly identical because the temperature was updated directly in both the assimilations. However, the distributions in O/N_2 in Figures 5b and 5d are significantly different.

The 23% improvement in DA2 O/N_2 , as seen in Figure 4, motivated us to analyze the electron content derived from the assimilations and compare them with independent TEC measurements. Figure 6 shows a latitude versus day-to-day variation of ECD in SD (a), DA1 (b), DA2 (c), and GNSS-TEC (d) centered at around 60°W ($\pm 5^\circ$) longitude. This spatial bin has been chosen due to the greater availability of GNSS data in this region. As mentioned in Section 2.3, the GNSS-TEC data are averaged over 20 min duration centered at every hour. Note that even with the 20 min averaging, there are missing data, specifically between 20° and 40°N . This figure shows that the magnitudes of electron densities and some of the shape and temporal variabilities of Equatorial Ionization Anomaly (EIA) in DA2 has better agreement with GNSS-TEC compared to the DA1 and SD. Particularly, the northern mid-latitude enhanced DA2 ECDs are in better agreement with GNSS-TEC. A quantification of the improvements is given at the end of this section. Though there are improvements in DA2, the EIA latitude extent and hemispheric asymmetries are not yet well reproduced in the assimilations. This could be due to the fact that the temperature variability cannot fully reflect the changes in the ionosphere as the ionosphere is also influenced by E-region winds in addition to neutral and ionospheric composition changes. We expect to have better agreement in the future when the GOLD O/N_2 and other ionospheric data set are assimilated in addition to the T_{disk} .

For a qualitative assessment of the improvements seen in the ionospheric electron content, a comparison between SD-ECD (green), DA1 ECD (cyan), DA2 ECD (red), and GNSS-TEC (blue) for a limited spatial region is shown in Figure 7a. The RMSE (in Figure 7b) and bias (in Figure 7c) with respect to GNSS-TEC are also shown. Except for November 1st and second and the night hours of each day (shaded regions, when GOLD data are not




Figure 6. Latitude and day-to-day variability of SD electron column density (ECD) (a), DA1 ECD (b), DA2 ECD (c), and global navigation satellite system total electron content (d) averaged over 55°W to 65°W longitude.

assimilated), the other days' DA2 ECD has better agreement with GNSS-TEC as can be inferred from the smaller values of the RMSE and bias. Some of the local time variabilities also have better agreement with DA2. For example, the two-peak structures in daytime GNSS-TEC on days 3 and 5 are better reproduced in the DA2 ECD, while that on eighth has not been reproduced. The two peak structure is particularly strong on 3rd November as indicated by downward arrows. Note the dates are in local time at 60°W. Also, the broader shape of the local time variability in GNSS-TEC match better with DA2 ECD as can be seen on most days in Figure 7a. Except for 1st and 2nd November, the night sector (shaded regions) has higher RMSEs, in general during the last 4 hr of each day and particularly at the end of 6th November. This is expected because the GOLD temperature are assimilated only during daylight sector and therefore they are not able to constrain the night-time dynamics. Including ionospheric and O/N_2 observations in the assimilation would improve the results. The purpose of this comparison is to demonstrate that the ionosphere is also improved upon assimilation of GOLD T_{disk} , though there are still large RMSEs and biases. Quantitative estimates of the differences, that vary with latitude and time, are given in Figure 8 and its discussion as given below.

In Figure 3 we show that the GOLD O/N_2 has latitudinal differences from the DA O/N_2 . Also, we have seen in Figure 6 that the agreement between DA2 ECD and GNSS-TEC varies with latitude. To investigate these latitudinal differences in TEC we have calculated the RMSE and mean bias (GNSS-TEC minus DA-ECD) at every 10° latitude bin during the 9 days. The RMSE and mean bias in electron contents from SD, DA1, and DA2 with respect to GNSS TEC are shown in Figure 8. One can note that the lowest values of the RMSE and bias are observed for the DA2, the red lines marked with stars. The RMSE and bias at every latitude bin is calculated from all the $24 \times 9 = 216$ hr of data. The percentage improvements in RMSE for DA1 ECD and DA2 ECD with respect to GNSS-TEC are about 3% and 10%, respectively. The 9 days average mean biases with respect to GNSS-TEC




Figure 7. (a) Comparison of the SD electron column density (ECD) (green), DA1 ECD (cyan), DA2 ECD (red), and Global Navigation Satellite System Total Electron Content (GNSS-TEC) (blue) which are averaged over 10°S to 0°N and 55°W to 65°W. (b) RMSEs in GNSS-TEC versus SD ECD (blue), GNSS-TEC versus DA1 ECD (green), and GNSS-TEC versus DA2 ECD (red). (c) Mean bias in GNSS-TEC versus SD ECD (blue), GNSS-TEC versus DA1 ECD (green), and GNSS-TEC versus DA2 ECD (red) are shown. Two dashed arrows in (a) indicate example two-peak structure. The shaded regions represent nighttime, when Global-Scale Observations of Limb and Disk data are not assimilated.

for the SD, DA1, and DA2 are about 1.9, 0.5, and 0.2 TECu, respectively. Also, the latitudinal average of absolute-biases are 1.92, 1.37, and 1.44 for SD, DA1, and DA2, respectively. Though the latitudinal average of the mean biases is slightly smaller for the DA2 compared to DA1, it is clear, from the absolute values, that the biases are smaller for both the assimilations compared to SD. Also, the mean bias is positive at higher latitudes (>30°) as seen in Figure 8b. Since O/N_2 and TEC vary in proportion, to a large extent, the smaller O/N_2 (from GOLD as shown in Figure 3b at the higher latitudes compared to SD and DA) may produce the positive mean biases in TEC. Negative bias and high RMSE between 0 and 20°S for the DA2 also imply that the equatorial electrodynamic, which is controlled by ionospheric E-region winds and composition, are not well constrained in the assimilations. Also, the night-time (when GOLD data are not assimilated) electrodynamic, particularly pre-reversal enhancement that is highly variable, contributes to poorer low-latitude results. But, overall these results further emphasize that the DA2—Where in addition to temperature the O, O^+ , and O_2 mixing ratios are updated directly—has the most improved thermosphere and ionosphere. Overall, it can be observed that the RMSEs are lower in the Northern hemisphere compared to the Southern hemisphere, which suggests that the Northern hemispheric variabilities are better reproduced in the assimilation.

4. Conclusions

An investigation of the impact of GOLD T_{disk} assimilation on thermosphere-ionosphere states is carried out using WACCMX + DART analysis states, GOLD measurements, and GNSS-TEC. The salient results of this investigation are:




Figure 8. (a) Latitudinal variability of root mean square error (RMSE) and (b) mean bias for SD (blue), DA1 (green), and DA2 (red) with respect to Global Navigation Satellite System Total Electron Content during 1–9 November 2018 are shown. Clearly, for the DA2 the RMSE is smaller compared to other two cases and bias is closer to zero.

1. GOLD T_{disk} assimilation analysis states of the thermosphere-ionosphere show better agreement with independent measurements than the control simulation.
2. The GOLD T_{disk} and O/N_2 compare better with the WACCMX + DART analysis effective temperature and O/N_2 when compared with equivalent parameters from SD-WACCMX.
3. The RMSE (with respect to GOLD) improvements in the analyses effective temperature and O/N_2 , when compared to their SD-WACCMX equivalents, are about 10.8% and 22.6%, respectively.
4. The RMSE between GNSS-TEC and analysis electron column density (ECD) is improved compared to that between GNSS-TEC and SD-WACCMX ECD. The improvement is about 10% for the assimilation that updates the O, O^+ , and O_2 densities in addition to temperature.

These results indicate that the GOLD observations of the thermospheric temperature have a great potential to improve the operational and short term forecast of the thermosphere-ionosphere system.

Data Availability Statement

Computing and data storage resources, including the Cheyenne supercomputer (<https://doi.org/10.5065/D6RX99HX>), were provided by the Computational and Information Systems Laboratory (CISL) at NCAR. Whole Atmosphere Community Climate Model with thermosphere-ionosphere eXtension is part of the Community Earth System Model (CESM) and the source code is available at <http://www.cesm.ucar.edu>. DART is available at <https://www.image.ucar.edu/DAReS/DART/>. The Level 2 data used in this study are available at the GOLD Science Data Center (<https://gold.cs.ucf.edu/search/>) and at NASA's Space Physics Data Facility (<https://spdf.gsfc.nasa.gov/pub/data/gold/level2/disk/>). The assimilation and simulation data used in this work are available at <https://doi.org/10.5281/zenodo.5816381>.

Acknowledgments

This research was supported by NASA Contract 80GSFC18C0061 to the University of Colorado, Boulder. This material is also based upon work supported by the National Center for Atmospheric Research (NCAR), which is a major facility sponsored by the National Science Foundation under Cooperative Agreement No. 1852977.

References

- Aa, E., Liu, S., Huang, W., Shi, L., Gong, J., Chen, Y., et al. (2016). Regional 3-D ionospheric electron density specification on the basis of data assimilation of ground-based GNSS and radio occultation data. *Space Weather*, 14(6), 433–448. <https://doi.org/10.1002/2016sw001363>
- Aksnes, A., Eastes, R., Budzien, S., & Dymond, K. (2006). Neutral temperatures in the lower thermosphere from N_2 Lyman-Birge-Hopfield (LBH) band profiles. *Geophysical Research Letters*, 33(15), L15103. <https://doi.org/10.1029/2006gl026255>
- Anderson, J., Hoar, T., Raeder, K., Liu, H., Collins, N., Torn, R., & Avellano, A. (2009). The data assimilation research testbed: A community facility. *Bulletin of the American Meteorological Society*, 90(9), 1283–1296. <https://doi.org/10.1175/2009bams2618.1>
- Anderson, J. L. (2001). An ensemble adjustment kalman filter for data assimilation. *Monthly Weather Review*, 129(12), 2884–2903.

- Bust, G. S., Garner, T. W., & Gaussiran, T. L., II. (2004). Ionospheric data assimilation three-dimensional (IDA3D): A global, multisensor, electron density specification algorithm. *Journal of Geophysical Research*, *109*(A11), A11312. <https://doi.org/10.1029/2003ja010234>
- Bust, G. S., & Immel, T. J. (2020). IDA4D: Ionospheric data assimilation for the ICON mission. *Space Science Reviews*, *216*(3), 33. <https://doi.org/10.1007/s11214-020-00648-z>
- Cantrall, C. E., Matsuo, T., & Solomon, S. C. (2019). Upper atmosphere radiance data assimilation: A feasibility study for GOLD far ultraviolet observations. *Journal of Geophysical Research: Space Physics*, *124*(10), 8154–8164. <https://doi.org/10.1029/2019ja026910>
- Chen, C. H., Lin, C. H., Matsuo, T., Chen, W. H., Lee, I. T., Liu, J. Y., et al. (2016). Ionospheric data assimilation with thermosphere-ionosphere-electrodynamics general circulation model and GPS-TEC during geomagnetic storm conditions. *Journal of Geophysical Research: Space Physics*, *121*(6), 5708–5722. <https://doi.org/10.1002/2015ja021787>
- Codrescu, M. V., Fuller-Rowell, T. J., & Minter, C. F. (2004). An ensemble-type kalman filter for neutral thermospheric composition during geomagnetic storms. *Space Weather*, *2*(11). <https://doi.org/10.1029/2004sw000088>
- Codrescu, S. M., Codrescu, M. V., & Fedrizzi, M. (2018). An ensemble kalman filter for the thermosphere-ionosphere. *Space Weather*, *16*(1), 57–68. <https://doi.org/10.1002/2017sw001752>
- Correia, J., Evans, J. S., Lumpe, J. D., Krywonos, A., Daniell, R., Veibell, V., & Eastes, R. W. (2021). Thermospheric composition and solar EUV flux from the global-scale observations of the limb and disk (GOLD) mission. *Journal of Geophysical Research: Space Physics*, *126*(12). <https://doi.org/10.1029/2021ja029517>
- Datta-Barua, S., Bust, G. S., & Crowley, G. (2013). First storm-time plasma velocity estimates from high-resolution ionospheric data assimilation. *Journal of Geophysical Research: Space Physics*, *118*(11), 7458–7471. <https://doi.org/10.1002/2013ja019153>
- Eastes, R. W., McClintock, W. E., Burns, A. G., Anderson, D. N., Andersson, L., Aryal, S., et al. (2020). Initial observations by the GOLD mission. *Journal of Geophysical Research: Space Physics*, *125*(7). <https://doi.org/10.1029/2020ja027823>
- Eastes, R. W., Solomon, S. C., Daniell, R. E., Anderson, D. N., Burns, A. G., England, S. L., & McClintock, W. E. (2019). Global-scale observations of the equatorial ionization anomaly. *Geophysical Research Letters*, *46*(16), 9318–9326. <https://doi.org/10.1029/2019gl084199>
- Forsythe, V. V., Azeem, I., Blay, R., Crowley, G., Makarevich, R. A., & Wu, W. (2021). Data assimilation retrieval of electron density profiles from ionosonde virtual height data. *Radio Science*, *56*(5). <https://doi.org/10.1029/2021rs007264>
- Gelaro, R., McCarty, W., Suárez, M. J., Todling, R., Molod, A., Takacs, L., et al. (2017). The modern-era retrospective analysis for research and applications, version 2 (MERRA-2). *Journal of Climate*, *30*(14), 5419–5454. <https://doi.org/10.1175/jcli-d-16-0758.1>
- He, J., Yue, X., Le, H., Ren, Z., & Wan, W. (2020). Evaluation on the quasi-realistic ionospheric prediction using an ensemble kalman filter data assimilation algorithm. *Space Weather*, *18*(3). <https://doi.org/10.1029/2019sw002410>
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., et al. (2020). The ERA5 global reanalysis. *Quarterly Journal of the Royal Meteorological Society*, *146*(730), 1999–2049. <https://doi.org/10.1002/qj.3803>
- Kodikara, T., Zhang, K., Pedatella, N. M., & Borries, C. (2021). The impact of solar activity on forecasting the upper atmosphere via assimilation of electron density data. *Space Weather*, *19*(5). <https://doi.org/10.1029/2020sw002660>
- Krywonos, A., Murray, D. J., Eastes, R. W., Aksnes, A., Budzien, S. A., & Daniell, R. E. (2012). Remote sensing of neutral temperatures in the Earth's thermosphere using the Lyman-Birge-Hopfield bands of N₂: Comparisons with satellite drag data. *Journal of Geophysical Research*, *117*(A9). <https://doi.org/10.1029/2011ja017226>
- Laskar, F. I., Eastes, R. W., Codrescu, M. V., Evans, J. S., Burns, A. G., Wang, W., & Cai, X. (2021). Response of GOLD retrieved thermospheric temperatures to geomagnetic activities of varying magnitudes. *Geophysical Research Letters*, *48*(15). <https://doi.org/10.1029/2021gl093905>
- Laskar, F. I., Eastes, R. W., Martinis, C. R., Daniell, R. E., Pedatella, N. M., Burns, A. G., & Codrescu, M. V. (2020). Early morning equatorial ionization anomaly from GOLD observations. *Journal of Geophysical Research: Space Physics*, *125*(7). <https://doi.org/10.1029/2019ja027487>
- Laskar, F. I., McCormack, J. P., Chau, J. L., Pallamraju, D., Hoffmann, P., & Singh, R. P. (2019). Interhemispheric meridional circulation during sudden stratospheric warming. *Journal of Geophysical Research: Space Physics*, *124*(8), 7112–7122. <https://doi.org/10.1029/2018ja026424>
- Laskar, F. I., Pedatella, N. M., Codrescu, M. V., Eastes, R. W., Evans, J. S., Burns, A. G., & McClintock, W. (2021). Impact of GOLD retrieved thermospheric temperatures on a whole atmosphere data assimilation model. *Journal of Geophysical Research: Space Physics*, *126*(1). <https://doi.org/10.1029/2020ja028646>
- Lee, I. T., Matsuo, T., Richmond, A. D., Liu, J. Y., Wang, W., Lin, C. H., et al. (2012). Assimilation of formosat-3/cosmic electron density profiles into a coupled thermosphere/ionosphere model using ensemble kalman filtering. *Journal of Geophysical Research*, *117*(A10). <https://doi.org/10.1029/2012ja017700>
- Lin, C. Y., Matsuo, T., Liu, J. Y., Lin, C. H., Tsai, H. F., & Araujo-Pradere, E. A. (2015). Ionospheric assimilation of radio occultation and ground-based GPS data using non-stationary background model error covariance. *Atmospheric Measurement Techniques*, *8*(1), 171–182. <https://doi.org/10.5194/amt-8-171-2015>
- Liu, H.-L., Bardeen, C. G., Foster, B. T., Lauritzen, P., Liu, J., Lu, G., et al. (2018). Development and validation of the whole atmosphere community climate model with thermosphere and ionosphere extension (WACCM-x 2.0). *Journal of Advances in Modeling Earth Systems*, *10*(2), 381–402. <https://doi.org/10.1002/2017ms001232>
- Marsh, D. R. (2011). Chemical–dynamical coupling in the mesosphere and lower thermosphere. In *Aeronomy of the Earth's atmosphere and ionosphere* (pp. 3–17). Springer Netherlands. https://doi.org/10.1007/978-94-007-0326-1_1
- Matsuo, T., Lee, I.-T., & Anderson, J. L. (2013). Thermospheric mass density specification using an ensemble kalman filter. *Journal of Geophysical Research: Space Physics*, *118*(3), 1339–1350. <https://doi.org/10.1002/jgra.50162>
- McClintock, W. E., Eastes, R. W., Beland, S., Bryant, K. B., Burns, A. G., Correia, J., & Veibel, V. (2020). Global-scale observations of the limb and disk mission implementation: 2. Observations, data pipeline, and level 1 data products. *Journal of Geophysical Research: Space Physics*, *125*(5). <https://doi.org/10.1029/2020ja027809>
- McCormack, J. P., Harvey, V. L., Pedatella, N., Koshin, D., Sato, K., Coy, L., et al. (2021). Intercomparison of middle atmospheric meteorological analyses for the northern hemisphere winter 2009–2010. *Atmospheric Chemistry and Physics Discussions*, 1–48. <https://doi.org/10.5194/acp-2021-224>
- McCormack, J. P., Hoppel, K., Kuhl, D., de Wit, R., Stober, G., Espy, P., et al. (2017). Comparison of mesospheric winds from a high-altitude meteorological analysis system and meteor radar observations during the boreal winters of 2009–2010 and 2012–2013. *Journal of Atmospheric and Solar-Terrestrial Physics*, *154*, 132–166. <https://doi.org/10.1016/j.jastp.2016.12.007>
- Mehta, P. M., Linares, R., & Sutton, E. K. (2018). A quasi-physical dynamic reduced order model for thermospheric mass density via hermitian space-dynamic mode decomposition. *Space Weather*, *16*(5), 569–588. <https://doi.org/10.1029/2018sw001840>
- Pedatella, N. M., Anderson, J. L., Chen, C. H., Raeder, K., Liu, J., Liu, H.-L., & Lin, C. H. (2020). Assimilation of ionosphere observations in the whole atmosphere community climate model with thermosphere-ionosphere EXTension (WACCMX). *Journal of Geophysical Research: Space Physics*, *125*(9). <https://doi.org/10.1029/2020ja028251>

- Pedatella, N. M., Liu, H.-L., Marsh, D. R., Raeder, K., Anderson, J. L., Chau, J. L., et al. (2018). Analysis and hindcast experiments of the 2009 sudden stratospheric warming in WACCMX + DART. *Journal of Geophysical Research: Space Physics*, *123*(4), 3131–3153. <https://doi.org/10.1002/2017ja025107>
- Pedatella, N. M., Raeder, K., Anderson, J. L., & Liu, H.-L. (2014). Ensemble data assimilation in the whole atmosphere community climate model. *Journal of Geophysical Research: Atmospheres*, *119*(16), 9793–9809. <https://doi.org/10.1002/2014jd021776>
- Remsberg, E. E., Marshall, B. T., Garcia-Comas, M., Krueger, D., Lingenfeller, G. S., Martin-Torres, J., & Thompson, R. E. (2008). Assessment of the quality of the version 1.07 temperature-versus-pressure profiles of the middle atmosphere from TIMED/SABER. *Journal of Geophysical Research*, *113*(D17). <https://doi.org/10.1029/2008jd010013>
- Ren, D., & Lei, J. (2020). Evaluation of physics-based data assimilation system driven by neutral density data from a single satellite. *Space Weather*, *18*(8). <https://doi.org/10.1029/2020sw002504>
- Rideout, W., & Coster, A. (2006). Automated GPS processing for global total electron content data. *GPS Solutions*, *10*(3), 219–228. <https://doi.org/10.1007/s10291-006-0029-5>
- Rienecker, M. M., Suarez, M. J., Gelaro, R., Todling, R., Bacmeister, J., Liu, E., et al. (2011). MERRA: NASA's modern-era retrospective analysis for research and applications. *Journal of Climate*, *24*(14), 3624–3648. <https://doi.org/10.1175/jcli-d-11-00015.1>
- Schwartz, M. J., Lambert, A., Manney, G. L., Read, W. G., Livesey, N. J., Froidevaux, L., et al. (2008). Validation of the aura microwave limb sounder temperature and geopotential height measurements. *Journal of Geophysical Research*, *113*(D15), D15S11. <https://doi.org/10.1029/2007jd008783>
- Song, R., Hattori, K., Zhang, X., & Yoshino, C. (2021). The three-dimensional ionospheric electron density imaging in Japan using the approximate kalman filter algorithm. *Journal of Atmospheric and Solar-Terrestrial Physics*, *219*, 105628. <https://doi.org/10.1016/j.jastp.2021.105628>
- Strickland, D. J., Evans, J. S., & Paxton, L. J. (1995). Satellite remote sensing of thermospheric o/n2 and solar EUV: 1. Theory. *Journal of Geophysical Research*, *100*(A7), 12217. <https://doi.org/10.1029/95ja00574>
- Sutton, E. K. (2018). A new method of physics-based data assimilation for the quiet and disturbed thermosphere. *Space Weather*, *16*(6), 736–753. <https://doi.org/10.1002/2017sw001785>
- Vierinen, J., Coster, A. J., Rideout, W. C., Erickson, P. J., & Norberg, J. (2016). Statistical framework for estimating GNSS bias. *Atmospheric Measurement Techniques*, *9*(3), 1303–1312. <https://doi.org/10.5194/amt-9-1303-2016>
- Wang, H., Fuller-Rowell, T. J., Akmaev, R. A., Hu, M., Kleist, D. T., & Iredell, M. D. (2011). First simulations with a whole atmosphere data assimilation and forecast system: The January 2009 major sudden stratospheric warming. *Journal of Geophysical Research*, *116*(A12). <https://doi.org/10.1029/2011ja017081>