

## **Insights into Identifying Habitat Hotspots and Migratory Corridors of Green Turtles in the South China Region**

Authors & Affiliations with address:

Connie Ka Yan Ng<sup>1\*</sup>, He Xiang Gu<sup>2</sup>, Tsung Hsien Li<sup>3</sup>, Ming Bin Ye<sup>2</sup>, Zhong Rong Xia<sup>2</sup>, Fei Yan Zhang<sup>2</sup>, Jin Xia Duan<sup>2</sup>, Chung Kang Hsu<sup>4</sup>, George H. Balazs<sup>5</sup>, Margaret B. Murphy<sup>1\*</sup>

<sup>1</sup> Department of Biology and Chemistry and State Key Laboratory in Marine Pollution, City University of Hong Kong, Kowloon Tong, Hong Kong Special Administrative Region, People's Republic of China

<sup>2</sup> Guangdong Huidong Sea Turtle National Nature Reserve Bureau. China Sea Turtle Base. Sea Turtle Bay, Gangkou Town, Huidong County, Guangdong Province, 516359, People's Republic of China

<sup>3</sup> National Museum of Marine Biology and Aquarium, Checheng, Pingtung, 94450, Taiwan.

<sup>4</sup> Penghu Marine Biology Research Center, Fisheries Research Institute, 266, Shihli, Magong, Penghu, Taiwan

<sup>5</sup> NOAA Pacific Islands Fisheries Science Center, National Oceanic and Atmospheric Administration, 1845 Wasp Blvd. Honolulu, Hawaii, 96818 USA

\*Emails of corresponding authors

Connie Ka Yan Ng: [kayan.ng.connie@gmail.com](mailto:kayan.ng.connie@gmail.com)

Margaret B. Murphy: [mbmurphy.cityu@gmail.com](mailto:mbmurphy.cityu@gmail.com)

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as doi: [10.1002/aqc.2923](https://doi.org/10.1002/aqc.2923)

## Abstract

1. Sea turtles are globally endangered and face daily anthropogenic threats, such as direct take, bycatch, habitat degradation. Current research efforts on sea turtles in the South China Region mainly focus on captivity and husbandry, haematology and blood chemistry, and nesting ecology. Published information on the habitat use of wild populations is limited.
2. This situation therefore creates a pressing need for scientific research on free-ranging sea turtles as a foundation for habitat management and species protection in South China. In this study, habitat use and oceanic movement of nesting, and by-catch or stranded green turtles were determined by satellite tracking combined with home range analysis.
3. Coupled with previous findings, foraging grounds of several sea turtle species (green turtle *Chelonia mydas*, hawksbill turtle *Eretmochelys imbricata* and loggerhead *Caretta caretta*) were mainly distributed along the coasts of Hainan Island Province and Guangdong Province, mainland China, as well as Taiwan and the Philippines, and the outlying islands in the South China Sea and East China Sea.
4. Habitat hotspots and migratory corridors of green turtles, in particular nesting turtles in South China, were identified. Coastal waters near Wanning City of Hainan Island, eastern Leizhou Peninsula, Iriomote-jima and Ishigaki-shima of the Ryukyu Islands of Japan and Dao Bach Long Vi of Vietnam serve as foraging grounds for nesting green turtles from different origins in South China. Moreover, the Paracel (Xisha) and Pratas (Dongsha) Islands in the South China Sea, Huidong Gangkou and its vicinity in mainland China, Liouciou Island and Penghu Island of Taiwan contain both nesting sites and foraging grounds for green turtles.
5. These sites that are associated with the migratory corridors, in particular Hainan Island, eastern Leizhou Peninsula and Liouciou Island, which currently lack conservation plans for sea turtles, should be given higher priority for habitat and species protection.

### Keywords (Mandatory)

Ocean, Coastal, Habitat mapping, Endangered species, Reptiles, Fishing

### Keywords (Set by authors)

Green turtle, Habitat use, Home range, Migratory corridor, Satellite telemetry, South China Region

## Introduction

Of the seven sea turtles species in the world, five species are found in the South China Sea. They are green turtle (*Chelonia mydas*), leatherback (*Dermochelys coriacea*), olive ridley (*Lepidochelys olivacea*), loggerhead (*Caretta caretta*), hawksbill (*Eretmochelys imbricata*) (Chan et al., 2007; Wang, 1993). These sea turtle species are migratory. Adults travel hundreds to thousands of kilometres between nesting beaches and foraging grounds. Among the five sea turtle species recorded in South China, the green turtle is the most common and is the only species that nests in the area (Chan et al., 2007; Ng, Dutton et al., 2014; Wang, 1993; Wang & Li, 2008). The green turtle is a globally endangered species (IUCN, 2015) and faces various anthropogenic threats, such as direct take and by-catch (Cheng & Chen, 1997; Wang & Li, 2008), trade pressure (Lam, Xu, Takahashi, & Burges, 2011; Pilcher, Chan, & Hiew, 2009) and habitat degradation (Wang & Li, 2008). Although green turtle nesting beaches on Wan-an Island of Taiwan, the Gangkou National Sea Turtle Nature Reserve of mainland China and Sham Wan of Hong Kong are protected by local ordinances (IUCN & UNEP-WCMC, 2018), nesting populations of green turtles in South China have been dwindling for decades (Chan et al., 2007; Wang & Li, 2008).

Habitat degradation as a result of rapid coastal development and human activities, e.g. fishing pressure, remains one of the key drivers in the dwindling populations of sea turtles worldwide (Bohm et al., 2013; Casale et al., 2010; IUCN Marine Turtle Specialist Group, 2013; Mazaris, Matsinos, & Pantis, 2009; McClellan & Read, 2009). As an effective and adaptive measure to conserve these migratory species, it is essential to identify and establish activity hotspots, such as habitat use in both the neritic and pelagic ocean and migratory pathways (Arendt et al., 2012; Craig, Parker, Brainard, Rice, & Balazs, 2004; Scales et al., 2011) to develop protection measures, such as delineation of management units, marine protected areas (MPAs) and threat mitigation (Casale, Affronte et al., 2012; Howell, Kobayashi, Parker, Balazs, & Polovina, 2008; Howell et al., 2015; Polovina et al., 2006; Rees, Al-Kiyumi, Broderick, Papathanasopoulou, & Godley, 2012; Scott et al., 2012).

Using satellite telemetry to track oceanic movements of sea turtles helps identify their habitat hotspots at different life stages, notably migratory pathways (i.e. movement between nesting sites and foraging grounds), inter-nesting grounds (i.e. areas where nesting turtles rest underwater between consecutive nesting activities) and foraging sites. Home range analysis (Casale, Broderick et al., 2012; Gaos et al., 2012; Hart & Fujisaki, 2010; Seminoff, Resendiz, & Nichols, 2002) of sea turtle activity hotspots, such as inter-nesting grounds and foraging sites, describes the area traversed by a sea turtle during the inter-nesting period or while foraging, excluding migrations or erratic movements. The location, extent and characteristics of home range and habitat use of sea turtles provide a crucial foundation for wildlife conservation and management, e.g. delineation of MPAs to protect habitats and food sources of species of concern (Craig et al., 2004; Gaos et al. 2012; Hart, Lamont, Fujisaki, Tucker, & Cathy 2012; Seminoff et al., 2002), and mitigation of threats such as unfavourable fisheries interactions (Hays

2008; Kobayashi et al., 2011; Polovina et al., 2006).

In China, tracking studies have primarily focused on post-nesting movements of green turtles (Figure 1). Chan, Chan, Lo, and Balazs (2003) and Ng, Dutton et al. (2014) determined post-nesting migratory pathways of nesting green turtles from Hong Kong to their foraging grounds in eastern Hainan Island, China and Dao Bach Long Vi in Vietnam. Nesting green turtles from Gangkou National Sea Turtle Reserve in China travelled to their foraging pastures on the east coast of Leizhou Peninsula and Okinawa Island in southern Japan (Song et al., 2002). After nesting, green turtles from Wan-an Island, Taiwan migrated to their foraging aggregations in coastal waters off northern Taiwan, the Guangdong coastline, Hainan Island, the northern Philippines and the Ryukyu Archipelago of Japan (Cheng, 2000); other nesting green turtles were found to migrate within Taiwanese waters (Chan et al., 2007). Cheng (2007) also determined post-nesting movement of green turtles from Taipin Island of the Nansha Archipelago in the South China Sea to coastal waters of the Philippines and Malaysia.

Although these studies have provided important information about the movement and distribution of green turtles in South China, there are severe knowledge gaps in habitat use of green turtles at different life stages (e.g. juvenile, sub-adult) and home range extent of key at-sea habitats in this region. Nesting green turtles are vital to sustain populations, while juvenile and non-breeding green turtles are also important considering that they comprise the majority of turtle populations. Baseline information on their habitat use and movement is hence essential for conservation, such as in terms of spatial management of habitat protection and threat mitigation. This study therefore aims to determine the movement patterns, migratory corridors and habitat hotspots of by-catch or stranded and nesting green turtles and to estimate home range extent of foraging/ inter-nesting habitats used by green turtles in the South China Region that covers Hong Kong, Guangdong and Taiwan. Research priorities for conserving and managing migratory green turtles will also be identified.

## Methods

Habitat use and hotspots of green turtles were identified by satellite telemetry aided by home range analysis. In this study, green turtles were sourced from: (i) by-catch (e.g. with direct reports from the fishermen involved) (n=6) or stranding incidents (n=17) after rehabilitation in the South China Region, and (ii) nesting in Hong Kong (n=1) and Taiwan (n=2) from 2006 to 2014. Home range extent of another nesting green turtle from Hong Kong was determined from tracking data published in Ng, Dutton et al. (2014). Tagging and satellite telemetry of these turtles generally followed methods described in Balazs, Miya, and Beavers (1996) and Balazs (1999). After a health assessment to ensure that each turtle was clinically healthy and physically fit for release, it was tagged with Inconel tags on its fore- and/ or hind-limbs for identification prior to release (Balazs, 1999). Its straight carapace length (SCL) /curved carapace length (CCL) was measured. Life stage was defined using carapace length intervals described for the best-available and geographically closest green turtle population, that in Hawaii

described by Balazs (1980). The following age classes were used: a juvenile is a post-hatchling to an individual of 65 cm straight carapace length (SCL); a sub-adult is an individual of SCL from 65 cm to 81 cm; and an adult is an individual of SCL > 81 cm and reproductively mature. According to the conversion between SCL and CCL where  $CCL = -0.414 + 1.039 \text{ SCL}$  (Bjorndal & Bolten, 1989), the life stage of green turtles based on CCL was defined as: juvenile, a post-hatchling to an individual of 67 cm CCL; sub-adult, CCL from 67 cm to 84 cm; and adult, CCL > 84 cm. A satellite Argos-linked transmitter was also attached to the carapace of each turtle with fiberglass resin following the protocol as described by Balazs et al. (1996). The weight of the transmitter package was less than 5% of the body weight of the turtles to minimize potential impact to their health (Watson & Granger, 1998). Of the 26 transmitters deployed, nine had GPS function.

Tracks of the oceanic movement of each turtle were plotted using Maptool (SEATURTLE.ORG, Inc.; <http://www.seaturtle.org/maptool/>) primarily with positional Argos data derived from the more accurate GPS and Location Class (LC) 1 to 3 signals; large spatial gaps were filled using data points of LC 0, A and B where appropriate following visual filtering for obviously inaccurate points (Chan et al., 2003; Parker, Balazs, King, Katabira, & Gilmartin, 2009; Witt et al., 2010; Yasuda & Arai, 2005), on the basis of excluding biologically unreasonable results of location points, including travel speed (>5 km/hr), points located on land, and a turn of greater than 90 degrees in less than a 24-hour period (Gaos et al., 2012; Parker et al., 2009). A maximum of one fix per day was selected with the highest accuracy LC, and if more than one fix had this LC, the one closest to midday was selected (Casale, Affronte et al., 2012; Casale, Broderick et al., 2012; Parker et al., 2009). Positional fixes during directional movement were connected with lines for ease of illustration. A bathymetry layer was included for movements in the open ocean. The end of a track was determined either by the last Argos position or when positional locations aggregated at a specific area 'inshore' if the turtle settled within estuaries, or 'near shore' if the turtle settled in areas along the open coast generally for approximately one month, implying that the tracked turtle had arrived and settled down at its inter-nesting ground (Gaos et al., 2012) or foraging ground (Parker et al., 2009). Speed of the tracked green turtle during inter-nesting, in transit or foraging was obtained by dividing the distance travelled during each specific activity by the corresponding duration of tracking. Two-sample t-tests were used to determine any significant differences ( $p < 0.05$ ) in speed during transit and foraging. Correlations between speed during transit and foraging, and carapace length as a proxy for age (Balazs, 1980; Hirth, 1997) were examined by parametric Pearson correlation analysis. Satellite tracks in the present study were compiled with relevant findings of previous studies to generate an overview of oceanic movements of sea turtles in the region. Threats to habitat hotspots and migratory corridors identified in this study were also reviewed and identified as far as possible.

Home range analysis in terms of Minimum Convex Polygon (MCP) and 50%, 90% and 95% utilization distribution (UD) of Kernel Density Estimate (KDE) to pinpoint any core-use area and/or overall home range was performed using the Home Range Tools

(HRT) extension of ArcView (ArcView 9, ESRI, Redlands, California, USA). No home ranges were calculated for turtles with less than 20 location points during the inter-nesting or foraging phases (Casale, Broderick et al., 2012; Gaos et al., 2012; Hart & Fujisaki, 2010; Rodgers, Carr, Smith, & Kie, 2005; Seminoff et al., 2002). MCP typically overestimates home ranges when the area is biased by extreme outliers of locality data (Hooge, Eichenlaub, & Solomon, 1999) while KDEs can be overestimated in cases of few or dispersed locations (Casale, Broderick et al., 2012). In view of the possible over-estimation of home range size by both MCP and KDE, the estimated home range extents in this study are provided for comparison with other studies to discuss conservation implications rather than for exact home range estimation.

## Results

The physical features of each released green turtle with tracking results and home range extent (if available) are shown in Table 1 of the Appendix. Home range extent at each respective foraging/ inter-nesting ground in this study and other studies is presented in Table 2 of the Appendix. Movements for all by-catch or stranded turtles and nesting green turtles by satellite tracking are exhibited in Figures A and B of the Appendix, respectively.

### *By-catch or Stranded Green Turtles in Hong Kong, Guangdong and Taiwan*

A total of 23 by-catch (n=6) or stranded (n=17) green turtles were released after rehabilitation and tracked by satellite transmitters for their movements in this study. Among these turtles, 13 turtles were released from Hong Kong, three from Gangkou Reserve of Guangdong and one turtle from Yangjiang of Guangdong, five from National Museum of Marine Biology and Aquarium (NMMBA) of Taiwan, and one from Penghu Islands of Taiwan.

The results of the satellite tracking (Table 1A and Figure A of Appendix) revealed that green turtles arrived at their foraging grounds in the coastal waters of various places in the South China Sea: eastern Hong Kong waters (where Yan Chau Tong Marine Park and Tung Ping Chau Marine Park are located), the coastal areas of Fujian Province, Hainan Island, Pratas (Dongsha) Island, Taiwan coastal waters and as far as Luzon and Palawan in the Philippines (Table 2 of Appendix). The total integrated distance travelled by these turtles ranged from 13 km to 3232 km, and the duration of tracking varied from 18 days to 382 days. The speed of travel in transit (0.06 to 2 km/hr), i.e. during movement/ migration before reaching the foraging ground, was significantly higher than that after the turtles reached their foraging ground (0.01 to 1.37 km/hr) (n=18; two-sample *t*-test,  $p=0.024$ ). No significant correlations between carapace length of green turtles and speed of travel in transit (n=22,  $r=0.177$ ,  $p=0.399$ ) or during foraging (n=18,  $r=-0.101$ ,  $p=0.689$ ) were observed in this study.

There were no correlations between turtle size or tracking duration and the area of foraging home ranges (Blanco, Morreale, Bailey et al., 2012; Gaos et al., 2012; Hawkes et al., 2012). Results of home range extents of all green turtles were compiled in this

study (Table 2 of Appendix). Home range extent at the neritic foraging grounds ranged from 1 to 1017 km<sup>2</sup> (median: 203 km<sup>2</sup>) in terms of MCP, from 0.2 to 974 km<sup>2</sup> (median: 64 km<sup>2</sup>) for KDE 50%, from 1 to 3978 km<sup>2</sup> (median: 285 km<sup>2</sup>) for KDE 90%, and from 2 to 5148 km<sup>2</sup> (median: 355 km<sup>2</sup>) for KDE 95%.

***Nesting Green Turtles in Hong Kong and Liouciou Island, Taiwan***

Nesting green turtles in Hong Kong started their migration from the nesting site to the foraging grounds (Wanning City of Hainan Island, China and Dao Bach Long Vi in Vietnam) from September to October. (Ng, Dutton et al., 2014). The extent of inter-nesting area of a nesting green turtle in Hong Kong during its three nesting activities in 2003, 2008 and 2012 ranged from 27 to 376 km<sup>2</sup> (median: 52 km<sup>2</sup>) in terms of MCP, from 5 to 118 km<sup>2</sup> (median: 11 km<sup>2</sup>) for KDE 50%, from 45 to 560 km<sup>2</sup> (median: 68 km<sup>2</sup>) for KDE 90%, and from 59 to 719 km<sup>2</sup> (median: 86 km<sup>2</sup>) for KDE 95%. The inter-nesting areas were generally located in coastal waters around Lamma Island (where the turtle nested), with the core area in southern Lamma (Table 1B and Figures B of Appendix).

According to the satellite tracking results (Table 1B and Figures B of Appendix), two nesting green turtles migrated from the nesting site in Liouciou Island, Taiwan from September to October in 2013 and arrived at their respective foraging grounds at Iriomote-jima in the Ryukyu Islands, Japan and Dao Bach Long Vi Island, Vietnam (respective home range also reported in Table 2). The total integrated distance travelled by these nesting turtles was 862 km and 1506 km, respectively. Duration of tracking varied from 64 days to 177 days.

**Discussion****Habitat Hotspots and Migratory Corridors of Sea Turtles in the South China Sea**

Broderick, Coyne, Fuller, Glen, and Godley (2007) demonstrated that green turtles exhibited high levels of fidelity to migratory routes and foraging areas both between and within years and after successive breeding migrations. Foraging grounds of sea turtles identified by satellite telemetry, news reports mainly in Chinese and interviews with local people on sea turtle strandings or sightings in the South China Sea from the present study and previous studies are summarized in Table 3 of the Appendix and graphically presented in Figures 1 and 2.






Figure 1 Nesting sites, foraging grounds and migratory pathways of green turtles and loggerhead turtles determined by satellite telemetry in South China Sea. The red star and line denote a green turtle nesting site in Hong Kong and post-nesting movement; Yellow=Gangkou; Light green=Wan-an; Blue=Lanyu; White=Liouciou Island; Orange=Taipin, Nansha Island. Purple = loggerhead in Minabe, Japan. The numbers in closed circles represent foraging grounds: 1 Wanning of Hainan, 2 Dao Bach Long Vi in Vietnam, 3 East coast of Leizhou Peninsula, 4 Okinawa Island in southern Japan, 5 Coastal waters off northern Taiwan, 6 Qinpeng-Dao of Nanao Island at Shantou, 7 Huidong, 8 Pratas (Dongsha) Island, 9 Dangan Liedao of Wanshan Archipelago, 10 northern Philippines, 11 Iriomote-jima and Ishigaki-shima of Ryukyu Archipelago, 12 Koshiki, Okinawa Island in southern Japan, 13 Coral reefs in the southern Penghu Archipelago, north of Chimei Island, 14 Coastal waters off Palawan Island, the

Philippines; 15 the north coast of eastern Malaysia; 16 east coast of Luzon Island, the Philippines, 17 East China Sea. (References refer to Table 3, Appendix)




Figure 2 Distribution of major nesting and foraging grounds of sea turtles in South China Sea. Green closed circles denote green turtle foraging grounds; open circles denote nesting grounds. Orange closed triangles denote hawksbill turtle foraging grounds; open triangles denote nesting grounds. Blue closed pentagons denote olive ridley foraging grounds. Purple closed squares denote loggerhead turtle foraging grounds; open squares denote nesting grounds. Black stars denote leatherback turtle foraging grounds. Numbers in closed circle represents foraging grounds as described for Figure 1 (References refer to Table 3, Appendix).

Satellite telemetry in the present study supports the findings of previous studies on important foraging grounds of green turtles, in particular post-nesting individuals, e.g. coastal waters along Guangdong Province and Fujian Province (Chan et al., 2007; Frazier et al., 1988); the northern, eastern and southern coasts of Hainan Island (Chan et al., 2003; Cheng, 2000; Song et al., 2002); the Pratas (Dongsha) Island (Cheng, 2000); the Paracel (Xisha) Islands (Wang & Li, 2008); Taiwan coastal waters (Cheng, 2000; Cheng & Chen, 1997); and north and south of the Philippines (Cheng, 2000, 2007; Xia & Gu, 2012). Ng, Chen, & Balazs (2014) also identified the waters off Keelung, northern Taiwan associated with drifting mats of *Sargassum* as habitat used by pelagic-phase green turtles. Moreover, the genetic analysis by Ng et al. (2017) indicated coastal Guangdong, the Taiwan Strait and the East China Sea as habitat used by pelagic-phase green turtles hatched from nesting beaches in Taiwan and mainland China.

Moreover, the tracking results, together with site verification at locations where the tracking ended, allow identification of foraging grounds of green turtles on a finer scale which have not been documented previously. The tracking outcomes in this study confirm that coastal waters along Guangdong Province, specifically Gangkou/ Huidong (ID 104688, Table 1, Appendix), Shanwei (ID 60991, 60992), Wanshan Archipelago (ID 104684, 52099), Shantou (Nanao Island), Xuwen, Nao Zhou, Hainan Island, as well as southern Penghu (ID 129723, 88057), Taitung (East Taiwan) (ID 53748, 71914) and Liouciou Island in Taiwan, contain suitable foraging habitats for green turtles. The movement range of green turtles with satellite tag ID 76441 and 60995, together with reports of green turtle sightings by fishermen in this study, indicates possible year-round occupancy of green turtles in Mirs Bay and Daya Bay and the proximate waters.

Post-nesting green turtles travelled from August to October, and possibly before nesting from April to June, via various major pathways and straits to reach their respective foraging grounds (Table 3, Appendix and Figure 1): coastal waters off Guangdong and Fujian Province, Qiongzhou Strait between Leizhou Peninsula and Hainan Island, Taiwan Strait between mainland China and Taiwan, Taiwan coastal waters, Luzon Strait between Taiwan and the Philippines, coastal waters off Palawan and the northern Philippines, Lumbucan Channel connecting the South China Sea with the Sulu Sea, and the East China Sea. Gu (2014) and Ye, Chen, Gu, and Li (2015) also suggested the Taiwan Strait and nearshore sea between Hainan and Leizhou Peninsula as important migratory routes of green turtles released from Gangkou National Nature Reserve.

Based on the results of the present study and previous studies, a number of sites harbour foraging grounds for a mixed stock of nesting green turtles from different places (Figure 2). Nesting green turtles from Hong Kong and Gangkou Reserve of mainland China shared the same foraging areas near Wanning City on Hainan Island. The east coast of Leizhou Peninsula was also foraging grounds for nesting green turtles from Gangkou Reserve and Wan-an Island of Taiwan. The Ryukyu Islands of Japan served as foraging grounds for nesting green turtles from Wan-an Island and Liouciou Island, Taiwan. Dao Bach Long Vi Island in Vietnam also harboured foraging areas for nesting green turtles

from Hong Kong and Liouciou Island. In addition, several locations contain nesting sites and foraging grounds for juvenile and adult green turtles (Figures 1 and 2), namely Gangkou and its vicinity, Liouciou Island, Penghu Island, the Paracel (Xisha) Islands (also for hawksbill and loggerhead turtles) and Pratas (Dongsha) Island (also for hawksbill turtles). These sites associated with the migratory corridors should be given higher priority for habitat and species protection in the face of limited resources and when imminent threats to sea turtle population(s) are identified.

### **Home Range and its Conservation Implications**

Home range extent of green turtles at each foraging ground identified in this study generally fell within the range determined by other studies in the Pacific, Atlantic and the Mediterranean Sea (Table 2 of Appendix). Overall, differences in the usage of foraging areas could be due to different foraging strategies (Godley et al., 2003), food availability at different sites or within the same area (Berube, Dunbar, Rutzler, & Hayes, 2012; Cuevas, Abreu-Grobois, Guzman-Hernandez, Liceaga-Correa, & VanDam, 2008; Makowski, Seminoff, & Salmon, 2006; Seminoff et al., 2002) and seasonality, particularly in temperate zones (Hawkes et al., 2011; Shaver, Hart, Fujisaki, Rubio, & Sartain, 2013). Site fidelity to specific segments of foraging grounds among juvenile green turtles has been documented in several foraging areas such as Hawaii, South Texas, Florida and Mexico (e.g. Brill et al., 1995; Makowski et al., 2006; Renaud, Carpenter, & Williams, 1995; Seminoff et al., 2002; Senko, Lopez-Castro, Koch, & Nichols, 2010) and among nesting green turtles (Broderick et al., 2007; Rees, Hafez, Lloyd, Papathansopoulou, & Godley, 2013) and nesting hawksbill turtles (Berube et al., 2012; Parker et al., 2009). The home ranges of these juvenile and nesting green turtles are particularly small. Such specific distribution of foraging habitats could be attributed to clustered and localized food sources and quality of food (Berube et al., 2012; Brill et al., 1995; Cuevas et al., 2008; Hazel, Hamann, & Lawler, 2013; Makowski et al., 2006; Mendonca 1983; Renaud et al., 1995; Seminoff et al., 2002; Whiting & Miller, 1998). In general, the home range extent of foraging green turtles near islands (e.g. the Pratas (Dongsha) Island, Penghu Island, Wanshan Archipelago, Dao Bach Long Vi, Ryukyu Islands) was relatively smaller than that observed in coastal waters in this study (Table 2 of Appendix). This difference in home range extent among habitat types (i.e. a hundred to thousand km<sup>2</sup> of KDE 95% at islands versus up to several thousand km<sup>2</sup> of KDE 95% in coastal waters) may be due to variation in availability and spatial distribution of food and shelter.

Identification of specific sites of key habitat areas (e.g. foraging grounds) is conducive to delineating areas for strategic protection in MPAs (Hart & Fujisaki, 2010). Overlapping home ranges may be an indication of high-quality habitat in an area (Berube et al., 2012; Casale, Broderick et al., 2012; Hart & Fujisaki, 2002; Seminoff et al., 2002). The results in this study implied that Luzon in the northern Philippines, eastern Taiwan waters, the Pratas (Dongsha) Island, Penghu Island, Wanshan Archipelago, eastern Hong Kong waters and Dao Bach Long Vi, where overlapping use of habitats by green turtles was observed (Table 2 of Appendix), likely constitute high-quality habitat, and hence these areas should be accorded high priority for further

study on habitat characterization to better understand the ecological niches of green turtles for appropriate habitat management.

### **Plasticity in Movement and Feeding Behavior of Green Turtles**

Blanco, Morreale, Bailey et al. (2012) reported that the largest foraging ground of a nesting green turtle included a combination of both nearshore and offshore waters providing different food items (seagrass in nearshore areas v.s. plankton-like jellyfish offshore) in Central America. Similarly, a relatively large home range of foraging loggerhead turtles (varying from 8,702 to 60,797 km<sup>2</sup>) spanning neritic and oceanic zones was observed in the Mediterranean by Casale, Broderick et al. (2012). This observation is comparable to the home range extent of individuals that adopt plasticity in their feeding habits by alternating between neritic and pelagic environments, e.g. the green turtle that foraged in the Taiwan Strait (ID 104685 in Table 1 of Appendix) in this study. Moreover, a few green turtles (ID 60991, 104684, 104685) in the present study demonstrated cyclic movement with ocean currents; these currents create hotspots of high productivity (e.g. chlorophyll a) in the open ocean (Kobayashi et al., 2008, 2011). These green turtles likely adopt plasticity in their feeding habits alternately between the neritic and pelagic environments. The same pelagic foraging behaviour was also observed in a nesting green turtle from Gangkou Reserve of Guangdong China (Song et al., 2002) and an adult green turtle that resided in the middle of the Taiwan Strait for nearly four months (Wang & Li, 2008). The same strategy was reported in adult female green turtles in the South China Sea by Hatase, Sato, Tamaguchi, Takahashi, & Tsukamoto (2006) and loggerheads following mesoscale eddies in the East China Sea off the coast of Taiwan by Kobayashi et al. (2008, 2011). Further studies on the association of sea turtle movement with oceanographic features in the China region and the South China Sea should be pursued to characterize their pelagic habitats and ultimately assess interactions with human activities in these habitats.

### **Overview of Threats to Sea Turtle Hotspots and Migratory Corridors**

Successful tracking of the majority of sea turtles in this study provided evidence for the post-release survivorship of the rehabilitated sea turtles in the wild. However, according to the satellite tracking results, site visits of this and previous studies, and news reports, some tagged green turtles have been incidentally captured in China in the previous and the current studies. Ye et al. (2015) also highlighted the by-catch of green turtles released from Gangkou Reserve during a tracking study. The female green turtle released from Yangjiang (ID 52100 in Table 1 of Appendix) in this study was incidentally caught by fishermen in Xinliao, Xuwen County, China and then released immediately by fishermen and local authority. Another female green turtle released from Hong Kong (ID 104687) was also incidentally caught in the East Harbor of Fujian and subsequently released. Ng, Dutton et al. (2014) also reported that a female green turtle found to nest in Hong Kong was reported to be entangled in a fishing net on its migratory pathway back to its foraging ground in Dao Bach Long Vi of Vietnam, and was found dead when fishermen recovered the net in early October 2012.

Incidental capture and direct take of sea turtles for trading purposes have been

commonly reported in Hainan Island, the Dongsha, Nansha and Xisha Archipelagoes in the South China Sea; South-east Asia, in particular Vietnam, the Philippines, Malaysia and Indonesia (Chan et al., 2007; Cheng, 1996; Globalpost, 2014; Hamann, Cuong, Hong, Thuoc, & Thuhien, 2006; Lam et al., 2011; Wang & Li, 2008) and the coastal waters along Guangdong and Fujian Province such as Xuwen and Shantou as observed in the present study. These localities overlap with the habitat hotspots and migratory corridors of nesting green turtles identified in the present study. Incidental take of sea turtles in the South China Sea was further illustrated by satellite tracking of a green turtle rescued from a market in Hainan, China reported by Yeh, Balazs, Parker, Ng, and Shi (2014). According to the tracking data, the turtle was apparently caught in the waters near Palawan in the Philippines, one of the hotspots of sea turtle poaching to satisfy growing demand in the sea turtle trade in mainland China (Lam et al., 2011). Release of animals including sea turtles is a long-standing tradition in Chinese culture (Balazs et al., 2013; Kuo et al., 2017). In this study, two religious-release green turtles found in Hong Kong were tracked and were found to finally settle at Shanwei (ID 60991) and in the East China Sea offshore of Zhejiang (ID 134341). These findings imply that the turtles possibly originated from and were caught in these areas. Cross-boundary take of sea turtles is not uncommon. Lam et al. (2011) reported 128 seizures involving East Asian countries between 2000 and 2008, with a trade volume of over 9,180 marine turtle products including whole specimens (2062 turtles), crafted projects (6161 pieces) and raw shell (789 scutes of 919 kg). Moreover, the largest seizure reported by Lam et al. (2011) involved 387 dead turtles aboard a Chinese fishing vessel in the Derawan Archipelago in East Kalimantan of Indonesia; the poachers were presumed to have targeted locations widely distributed across the Sulu and Celebes Sea corresponding to a major migratory corridor used by nesting green turtles from Taipin Island in the Nansha Archipelago. Curbing the opportunistic and direct take of sea turtles for the booming trade demand rests on effective multi-national management of these migratory species.

### **Attributes of Satellite Tracking**

Speed of travel varies with the behaviour of individual green turtles, such as during inter-nesting and migration and foraging (Hays, Luschi, Papi, del Seppia, & Marsh, 1999; Hochscheid, 2014; Rice & Balazs, 2008). Blanco, Morreale, Bailey et al. (2012) reported that there was no correlation between turtle size and speed of travel. The same result was also obtained in the present study. The speed of travel of green turtles in transit was significantly higher than that after the turtles reached their foraging grounds in the present study, although this is based on a relatively small sample size ( $n=18$ ). The same pattern was also observed by Casale, Freggi, Cina, and Rocco (2013), Gaos et al. (2012) and Papi, Luschi, Crosio, and Hughes (1997) where submergences were shorter and more frequent during migration than during time spent at foraging grounds. Broderick et al. (2007) reported that short shallow dives in summer were indicative of active foraging. These differences in speed of travel, together with apparent residence at a specific area, help to characterize the behaviour of green turtles during a tracking study.

## Recommendations

The habitat hotspots and migratory corridors of green turtles, in particular nesting turtles, in South China Region that were identified by satellite tracking and site verification in this study and previous studies are under varying levels of protection (Figures 1 and 2). Of these key sites, the coastal waters of Dao Bach Long Vi in Vietnam, Iriomote-jima and Ishigaki-shima of the Ryukyu Islands in Japan, Gangkou and its bay area, the south-western part of Leizhou Peninsula in mainland China, Dazhoudao Island in south-eastern and Sanya in southern waters of Hainan Island, Wan-an of Penghu Island in Taiwan, Paracel (Xisha) Island and Pratas (Dongsha) Island have been designated as protected areas by the corresponding local governments (IUCN & UNEP-WCMC, 2018). Operation of gill nets is not allowed in coastal waters of Liouciou Island to avoid entanglement and drowning of foraging green turtles there, but the island is not formally a marine protected area. While these existing protected areas should be maintained or enhanced, more resources should be directed to protect other key sites which are currently lacking conservation management, in particular a large part of Hainan Island, the eastern Leizhou Peninsula and Liouciou Island, as they are part of a network of critical habitats used by migratory green turtles.

To advance our knowledge of the distribution of nesting and foraging grounds and to enrich our understanding of migratory corridors/connectivity of sea turtle populations in the South China Region, including their extent and seasonality, more tracking and/or genetic studies should be conducted on nesting sea turtles. In particular, major nesting grounds in the South China Sea, which are under-studied, such as for green turtles and hawksbills in the Pratas (Dongsha) (Cheng, 1995) and Paracel (Xisha) Islands (Anonymous, 1975; Wang & Li, 2008), as well as other potential nesting sites, should be studied. Tracking a large number of adults from one foraging area, such as Liouciou Island in Taiwan, may help to identify multiple breeding sites (Luschi & Casale, 2014). Stable isotope analysis can also be used to infer connectivity of foraging grounds with nesting grounds (Ceriani et al., 2014). Resources and effort should be devoted to the monitoring of nesting sea turtles at existing and potential nesting grounds, such as by fishermen and citizen scientists in coastal communities.

Further quantitative studies on the availability and spatial distribution of marine resources (Hart, Lamont, Sartain, Fujisaki, & Stephens, 2013; Hart, Zawada, Fujisaki, & Barbara, 2013) and dietary studies (Berube et al., 2012; Ng, Ang, Russell, Balazs, & Murphy, 2016) should be performed to characterize habitat requirements of sea turtles (e.g. food item availability, seasonality, landscape, etc.) at the major foraging grounds in South China Region identified in this study. Habitat characterization is complementary to the application of home range when designating MPAs in a strategic and cost-effective fashion. More information on home ranges of foraging sea turtles should also be acquired to identify a common and repeatable pattern of habitat use and residency at each specific foraging ground.



In addition to habitat protection and further research studies, threat of direct take or bycatch of sea turtles should be quantitatively assessed and mitigated at the activity hotspots and migratory corridors identified in South China Sea. The only published study was on the spatial and temporal distribution of incidental capture of sea turtles by coastal setnets in Taiwan conducted more than 20 years ago (Cheng & Chen, 1997). Observer programmes should be established in close liaison with fishermen to identify areas of high by-catch risk and quantify interactions between fisheries and sea turtles at a preliminary stage. Further quantitative studies on interactions of by-catch species with oceanographic features and fisheries could be pursued in these by-catch hotspots, if identified. An adaptive management tool similar to TurtleWatch to reduce by-catch of loggerhead (Howell et al., 2008) and leatherback turtles (Howell et al., 2015) by the Hawaiian fishery in the Pacific could then be developed and implemented for China. Moreover, to combat the poaching and trading pressure on sea turtles, legal enforcement and public awareness campaigns targeting the local public, tourists, vendors and fishermen involved in the illegal sale and/or capture of sea turtles should be implemented. Sea Turtles 911 (<http://www.seaturtles911.org/>), a sea turtle conservation organization based in Hainan, has been delivering educational talks to tourists and the local public to discourage illegal purchase of sea turtle products in collaboration with hotels in Hainan, one of the hotspots of the sea turtle trade. More importantly, trans-regional and multi-national commitments to deter the sea turtle trade should be strengthened across international boundaries, such as in the form of forums, workshops and memoranda. One recent workshop co-hosted by government representatives from Indonesia, Malaysia, the Philippines and Vietnam focused on improving intergovernmental efforts to curb the illegal trade of sea turtles being harvested in the Coral Triangle (Traffic, 2014). China's participation in multi-national taskforces would be an important step in combating the sea turtle trade in the South China Sea and beyond.

### **Acknowledgements**

Professional and continuous support from Prof. Paul K. S. Lam of the City University of Hong Kong (CityU), Dr. Thierry Work of the USGS National Wildlife Health Center, Yaru Cai, H. H. Chan, Raymond W.M. Chan, friends and colleagues of the State Key Laboratory in Marine Pollution of the CityU, Gangkou National Sea Turtle Nature Reserve of mainland China and NMMBA of Taiwan were indispensable to this study. We would also like to acknowledge the approval of the Agriculture, Fisheries and Conservation Department of the Government of the Hong Kong Special Administrative Region of the People's Republic of China for using the data and specimens of green turtles collected in Hong Kong for this study. We thank Prof. I-Jiunn Cheng and his marine turtle research team of Institute of Marine Biology, National Taiwan Ocean University, who attached transmitters on nesting green turtles on Liouciou Island, Taiwan and also Liouciou Township Office, local conservationists Tien-yun Tsai, Yao-wen Tsai, Tien-cheng Wang, Sophia Ching-ju Hung and Jian-wei Zeng for their great efforts to facilitate the tracking on Liouciou Island, Taiwan. Connie Ka-yan NG thanks Denise M. Parker for her professional scientific support of the tracking research.

## Supporting Information

Additional Supporting Information may be found online in the supporting information tab for this article.

## References

Anonymous. (1975). A preliminary observation on the reproductive behaviour of sea turtles in Xisha Archipelago – Report from the Biological Resources Group, South China Sea Institute of Oceanography. *Chinese Academy of Science. Zoological Bulletin*, 10, 34-35.

Arendt, M.D., Segars, A.L., Byrd, J.L., Boynton, J., Whitaker, J.D., Parker, L., ...Roberts, M.A. (2012). Distributional patterns of adult male loggerhead sea turtles (*Carretta caretta*) in the vicinity of Cape Canaveral, Florida, USA during and after a major annual breeding aggregation. *Marine Biology*, 159, 101-112.

Balazs, G.H. (1980). Synopsis of Biological Data on the Green Turtle in the Hawaiian Islands. NOAA Technical Memorandum NMFS. Hawaii: Southwest Fisheries Center, National Oceanic and Atmospheric Administration. U.S. Department of Commerce.

Balazs, G. H., Miya, R. K., & Beavers, S. C. (1996). Procedures to attach a satellite transmitter to the carapace of an adult green turtle, *Chelonia mydas*. Keinath, J. A., Barnard, D. E., Musick, J. A., & Bell, B. A. (Eds.), *Proceedings of the Fifteenth Annual Symposium on Sea Turtle Biology and Conservation, February 20-25, 1995* (pp. 21-26), South Carolina., U.S.

Balazs, G. H. (1999). Factors to consider in the tagging of sea turtles. Eckert, K. L., Bjorndal, K. A., Abreu-Grobois, F. A., & Donnelly, M. (Eds.), *Research and management techniques for the conservation of sea turtles* (pp. 101-109), IUCN/ SSC Marine Turtle Specialist Group Publication No. 4, Washington, US.

Balazs, G.H., Ng, K.Y., Gu, H.X., & Zhang, F.Y. (2013). The Xunlian Guangdong Province Experience: Releasing Sea Turtles For Restocking and Conservation Awareness in China. 33rd International Sea Turtle Symposium, Baltimore, U.S.

Benson, S.R., Eguchi, T., Foley, D.G., Forney, K.A., Bailey, H., Hitipeuw, C., ...Dutton, P.H. (2012). Large-scale movements and high-use areas of western Pacific leatherback turtles, *Dermochelys coriacea*. *Ecosphere*, 2, Article 84.

Berube, M.A., Dunbar, S.G., Rutzler, K., & Hayes, W.K. (2012). Home range and foraging ecology of juvenile hawksbill sea turtles (*Eretmochelys imbricata*) on in shore reefs of Honduras. *Chelonian Conservation and Biology*, 11, 33-43.

- Bjorndal, K.A., & Bolten, A. B. (1989). Comparison of straight-line and over-the-curve measurements for growth rates of green turtles, *Chelonia mydas*. *Bulletin of Marine Science*, 43, 189-192.
- Blanco, G.S., Morreale, S.J., Bailey, H., Seminoff, J.A., Paladino, F.V., & Spotila, J.R. (2012a). Post-nesting movements and feeding grounds of a resident East Pacific green turtle *Chelonia mydas* population from Costa Rica. *Endangered Species Research*, 18, 233-245.
- Blanco, G.S., Morreale, S.J., Seminoff, J.A., Paladino, F.V., Piedra, R., & Spotila, J.R. (2012b). Movements and diving behavior of interesting green turtles along Pacific Costa Rica. *Integrative Zoology*, 8, 293-306.
- Bohm, M., Collen, B., Baillie, J.E.M., Bowles, P., Chanson, J., Coz, N., ...Zug, G. (2013). The conservation status of the world's reptiles. *Biological Conservation*, 157, 372-385.
- Brill, R.W., Balazs, G.H., Holland, K.N., Chang, R.K.C., Sullivan, S., & George, J.C. (1995). Daily movements, habitat use, and submergence intervals of normal and tumor-bearing juvenile green turtles (*Chelonia mydas* L.) within a foraging area in the Hawaiian Islands. *Journal of Experimental Marine Biology and Ecology*, 185, 203-218.
- Broderick, A.C., Coyne, M.S., Fuller, W.J., Glen, F., & Godley, B.J. (2007). Fidelity and over-wintering of sea turtles. *Proceedings of the Royal Society B*, 274, 1533-1538.
- Casale, P., Affronte, M., Insacco, G., Freggi, D., Vallini, C., D'Astore, P.P., ...Argano, R. (2010). Sea turtle strandings reveal high anthropogenic mortality in Italian waters. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 20, 611-620.
- Casale, P., Affronte, M., Scaravelli, D., Lazar, B., Vallini, C., & Luschi, P. (2012a). Foraging grounds, movement patterns and habitat connectivity of juvenile loggerhead turtles (*Caretta caretta*) tracked from the Adriatic Sea. *Marine Biology*, 159, 1527-1535.
- Casale, P., Broderick, A.C., Freggi, D., Mencacci, R., Fuller, W.J., Godley, B.J., & Luschi, P. (2012b). Long-term residence of juvenile loggerhead turtles to foraging grounds: a potential conservation hotspot in the Mediterranean. *Aquatic Conservation: Marine Freshwater Ecosystem*, 22, 144-154.
- Casale, P., Freggi, D., Cina, A., & Rocco, M. (2013). Spatio-temporal distribution and migration of adult male loggerhead sea turtles (*Caretta caretta*) in the Mediterranean Sea: further evidence of the importance of neritic habitats off North Africa. *Marine Biology*, 160, 703-718.
- Ceriani, S.A., Roth, J.D., Sasso, C.R., McClellan, C.M., James, M.C., Haas, H.L.,... Weishampel, J.F. (2014). Modelling and mapping isotopic patterns in the Northwest

Atlantic derived from loggerhead sea turtles. *Ecosphere*, 5, 1-24.

Chan, S. K. F., Chan, J. K., Lo, L. T., & Balazs, G. H. (2003). Satellite tracking of the post-nesting migration of a green turtle (*Chelonia mydas*) from Hong Kong. *Marine Turtle Newsletter*, 102, 2-4.

Chan, S. K. F., Cheng, I. J., Zhou, T., Wang, H. J., Gu, H. X., & Song, X. J. (2007). A comprehensive overview of the population and conservation status of sea turtles in China. *Chelonian Conservation and Biology*, 6, 185-198.

Chen, W.L. (1995). Release of Green Turtle and Hawksbill at Tai Shan of South China. *Fisheries Science and Technology*, 4, 46.

Cheng, I.J. (1995). Sea Turtles at Dungsha Tao, South China Sea. *Marine Turtle Newsletter*, 70, 13-14.

Cheng, I.J. (1996). Sea Turtles at Taipin Tao, South China Sea. *Marine Turtle Newsletter*, 75, 6-8.

Cheng, I.J. (2000). Post-nesting migrations of green turtles (*Chelonia mydas*) at Wan-An Island, Penghu Archipelago, Taiwan. *Marine Biology*, 137, 747-754.

Cheng, I.J. (2007). Nesting Ecology and Postnesting Migration of Sea Turtles on Taipin Tao, Nansha Archipelago, South China Sea. *Chelonian Conservation and Biology*, 6(2), 277-282.

Cheng, I.J. (2013). Recovery plan of green turtles on Liouciou Island, Pingtung County. Final Report to the Liuqiu Township Office: Pingtung County, Taiwan.

Cheng, I. J., & Chen, T. H. (1997). The incidental capture of five species of sea turtle by coastal setnet fisheries in the eastern waters of Taiwan. *Biological Conservation*, 82, 255-259.

Craig, P., Parker, D., Brainard, R., Rice, M., & Balazs, G. (2004). Migrations of green turtles in the central South Pacific. *Biological Conservation*, 116, 433-438.

Cong, S., & Wang, Z.M. (1997). Study on Marine Turtle and its Raising along the Coasts of Shandong. *Transactions of Oceanology and Limnology*, 3, 76-80.

Cuevas, E., Abreu-Grobois, F.A, Guzman-Hernandez, V., Liceaga-Correa, M.A., & VanDam, R.P. (2008). Post-nesting migratory movements of hawksbill turtles *Eretmochelys imbricata* in waters adjacent to the Yucatan Peninsula, Mexico. *Endangered Species Research*, 10, 123-133.

Frazier, S.S., Frazier, J.G., Ding, H.B., Huang, Z.J., Zheng J., & Lu, L. (1988). Sea

turtles in Fujian and Guangdong Provinces. *Acta Herpetologica Sinica*, 7, 16-46.

Gaos, A.R., Lewison, R.L., Wallace, B.P., Yañez, Liles, M.Y., Nichols, W.J., Baquera, A.,...Seminoff, J.A. (2012). Spatial ecology of critically endangered hawksbill turtles *Eretmochelys imbricata*: implications for management and conservation. *Marine Ecology Progress Series*, 450, 181-194.

Globalpost. 2014. Philippine police find 120 captured turtles in swamp. <http://www.globalpost.com/dispatch/news/afp/140131/philippine-police-find-120-captured-sea-turtles-swamp> [3 February 2014].

Godley, B.J., Lima, E.H.S.M., Akesson, S., Broderick, A.C., Glen, F., Godfrey, M.H.,...Hays, G.C. (2003). Movement patterns of green turtles in Brazilian coastal waters described by satellite tracking and flipper tagging. *Marine Ecology Progress Series*, 253, 279-288.

Gu, H.X. (2014). Leizhou Peninsular as a migratory corridor of sea turtles. *Man and the Biosphere*, 2, 29-33.

Hamann, M., Cuong, C.T., Hong, N.D., Thuoc, P., & Thuhien, B.T. (2006). Distribution and abundance of marine turtles in the Socialist Republic of Viet Nam. *Biodiversity and Conservation*, 15, 3703-3720.

Hangzhou. 2012. Release of rehabilitated loggerhead and green turtle at Dongyushan, Hangzhou of Zhejiang. [http://ori.hangzhou.com.cn/ornews/content/2012-11/13/content\\_4473574.htm](http://ori.hangzhou.com.cn/ornews/content/2012-11/13/content_4473574.htm) [3 February 2014]

Hart, K.M., & Fujisaki, I. (2010). Satellite tracking reveals habitat use by juvenile green sea turtles *Chelonia mydas* in the Everglades, Florida, USA. *Endangered Species Research*, 11, 221-232.

Hart, K.M., Lamont, M.M., Fujisaki, I., Tucker, A.D., & Cathy, R.R. (2012). Common coastal foraging areas for loggerheads in the Gulf of Mexico: opportunities for marine conservation. *Biological Conservation*, 145, 185-194.

Hart, K.M., Lamont, M.M., Sartain, A.R., Fujisaki, I., & Stephens, B.S. (2013a). Movements and habitat-use of loggerhead sea turtles in the Northern Gulf of Mexico during the reproductive period. *PLoS ONE*, 8, e66921.

Hart, K.M., Zawada, D.G., Fujisaki, I., & Barbara, H.L. (2013b). Habitat use of breeding green turtles *Chelonia mydas* tagged in Dry Tortugas National Park: Making use of local and regional MPAs. *Biological Conservation*, 161, 142-154.

Hatase, H., Takai, N., Matsuzawa, Y., Sakamoto, W., Omuta, K., Goro, K., ...Fujiwara,

T. (2002). Size-related differences in feeding habitat use of adult female loggerhead turtles *Caretta caretta* around Japan determined by stable isotope analyses and satellite telemetry. *Marine Ecology Progress Series*, 233, 273-281.

Hatase, H., Sato, K., Tamaguchi, M., Takahashi, K., & Tsukamoto, K. (2006). Individual variation in feeding habitat use by adult female green sea turtles (*Chelonia mydas*): are they obligately neritic herbivores? *Oecologia*, 149, 52-64.

Hawkes, L.A., Witt, M.J., Broderick, A.C., Coker, J.W., Coyne, M.S., Dodd, M.,...Godley, B.J. (2011). Home on the range: spatial ecology of loggerhead turtles in Atlantic waters of the USA. *Diversity and Distributions*, 17, 624-640.

Hawkes, L.A., Tomas, J., Revuelta, O., Leon, Y.M., Blumenthal, J.M., Broderick, A.C., ...Godley, B.J. (2012). Migratory patterns in hawksbill turtles described by satellite tracking. *Marine Ecology Progress Series*, 461, 223-232.

Hays, G.C. (2008). Sea Turtles: a review of some key recent discoveries and remaining questions. *Journal of Experimental Marine Biology and Ecology*, 356, 1-7.

Hays, G.C., Luschi, P., Papi, F., del Seppia, C., & Marsh, R. (1999). Changes in behavior during the inter-nesting period and post-nesting migration for Ascension Island green turtles. *Marine Ecology Progress Series*, 189, 263-273.

Hazel, J., Hamann, M., & Lawler, I.R. (2013). Home range of immature green turtles tracked at an offshore tropical reef using automated passive acoustic technology. *Marine Biology*, 160, 617-627.

Hirth, H. F. (1997). Synopsis of the biological data on the green turtle *Chelonia mydas* (Linnaeus 1758). Biological Report. Washington, DC: Fish and Wildlife Service. U.S. Department of the Interior.

Hochscheid, S. (2014). Why we mind sea turtles' underwater business: a review on the study of diving behavior. *Journal of Experimental Marine Biology and Ecology*, 450, 118-136.

Hooge, P.N., Eichenlaub, W., & Solomon, E. (1999). The Animal Movement Program. United States Geologic Survey, Alaska Biological Science Center, US.

Howell, E.A., Hoover, A., Benson, S.R., Bailey, H., Polovina, J.J., Seminoff, J.A., & Dutton, P.H. (2015). Enhancing the TurtleWatch product for leatherback sea turtles, a dynamic habitat model for ecosystem-based management. *Fisheries Oceanography*, 24, 57-68.

Howell, E.A., Kobayashi, D.R., Parker, D.M., Balazs, G.H., & Polovina, J.J. (2008). TurtleWatch: a tool to aid in the bycatch reduction of loggerhead turtles *Caretta caretta*

in the Hawaii-based pelagic longline fishery. *Endangered species research*, 1033, 1-12.

Hua, H.L., & Yin, J.W. (1993). Protected animals in China. Shanghai: Shanghai Science and Technology Education Publishing Co.

Huang, C.C. (1979). Distribution of Sea Turtles in China Sea. K.A., Bjorndal. (Eds.), *Biology and Conservation of Sea Turtles* (pp. 321-322), Smithsonian Institution, U.S.

IUCN Marine Turtle Specialist Group. 2013. Hazards to Marine Turtles. <http://iucn-mtsg.org/about-turtles/hazards/> [20 January 2013]

IUCN Red List Of Threatened Species. 2015. All Sea Turtles Species and Python bivittatus. <http://www.iucnredlist.org/> [23 October 2015]

IUCN & UNEP-WCMC. 2018. The world database on Protected Areas (WDPA). Cambridge, UK. <http://www.protectedplanet.net> [1 February 2018]

Jinghua. 2011. Release of stranded green turtle from Beihai, Guangxi. <http://news.jinghua.cn/351/c/201106/21/n3380402.shtml> [23 October 2015]

Kobayashi, D.R., Polovina, J.J., Parker, D.M., Kamezaki, N., Cheng, I.J., Uchida, I.,...Balazs, G.H. (2008). Pelagic habitat characterization of loggerhead sea turtles, *Caretta caretta*, in the North Pacific Ocean (1997-2006): Insights from satellite tag tracking and remotely sensed data. *Journal of Experimental Marine Biology and Ecology*, 356, 96-114.

Kobayashi, D.R., Cheng, I.J., Parker, D.M., Polovina, J.J., Kamezaki, N., & Balazs, G.H. (2011). Loggerhead turtle (*Caretta caretta*) movement off the coast of Taiwan: characterization of a hotspot in the East China Sea and investigation of mesoscale eddies. *ICES Journal of Marine Science*, doi:10.1093/icesjms/fsq185.

Kuo, F.W., Fan, T.Y., Ng, C.K.Y., Cai, Y, Balazs, G.H., & Li, T.H. (2017) Tale of the unlucky tags: the story of a rescued, rehabilitated, and released green sea turtle (*Chelonia mydas*) in southern Taiwan. *Bulletin of Marine Science*, 93, <https://doi.org/10.5343/bms.2016.1108>.

Lam, T., Xu, L., Takahashi, S., & Burges, E.A. (2011). Market forces: an examination of marine turtle trade in China and Japan. TRAFFIC East Asia, Hong Kong.

Luschi, P., & Casale, P. (2014). Movement patterns of marine turtles in the Mediterranean Sea: a review. *Italian Journal of Zoology*, DOI:10.1080/11250003.2014.963714 : 1-18.

Makowski, C., Seminoff, J.A., & Salmon, M. (2006). Home range and habitat use of juvenile Atlantic green turtles (*Chelonia mydas* L.) on shallow reef habitats in Palm

Beach, Florida, USA. *Marine Biology*, 148, 1167-1179.

Mazaris, A. D., Matsinos, M. G, & Pantis, J.D. (2009). Evaluating the impacts of coastal squeeze on sea turtle nesting. *Ocean and Coastal Management*, 52, 139-145.

McClellan, C.M., & Read, A.J. (2009). Confronting the gauntlet: understanding incidental capture of green turtles through fine-scale movement studies. *Endangered Species Research*, 10, 165-179.

Mendonca, M.T. (1983). Movements and feeding ecology of immature green turtles (*Chelonia mydas*) in a Florida lagoon. *Copeia*, 1983, 1013-1023.

Ng, C.K.Y., Dutton, P.H., Chan, S.K.F., Cheung, K.S., Qiu, J.W., & Sun, Y.A. (2014a). Characterization and conservation concern of green turtles (*Chelonia mydas*) nesting in Hong Kong, China. *Pacific Science*, 68, 231-243.

Ng, K.Y., Chen, T.H., & Balazs, G.H. (2014b). Flying fish egg harvest off Keelung, Taiwan uncovers occurrence of pelagic-phase green turtles. *Marine Turtle Newsletter*, 143, 14-15.

Ng, C.K.Y., Ang, P.O., Russell, D.J., Balazs, G.H. & Murphy, B.M. (2016). Marine Macrophytes and Plastics Consumed by Green Turtles (*Chelonia mydas*) in Hong Kong, South China Sea Region. *Chelonian Conservation and Biology*, 15, 289-292.

Ng, C.K.Y., Dutton, P.H., Gu, H.X., Li, T.H., Ye, M.B., Xia, Z.R.,...& Murphy, B.M. (2017). Regional Conservation Implications of Green Turtle (*Chelonia mydas*) Genetic Stock Composition in China. *Chelonian Conservation and Biology*, 16, in press.

Papi, F., Luschi, P., Crosio, E., & Hughes, G.R. (1997). Satellite tracking experiments on the navigational ability and migratory behavior of the loggerhead turtle *Caretta caretta*. *Marine Biology*, 129, 215-220.

Parker, D.M., Balazs, G.H., King, C.S., Katabira, L., & Gilmartin, W. (2009). Short-range movements of Hawksbill turtles (*Eretmochelys imbricata*) from nesting to foraging areas within the Hawaiian Islands. *Pacific Science*, 63, 371-382.

Pilcher, N., Chan, E.H., & Hiew, K. (2009). Battling the direct poaching of sea turtles in south-east Asia. Workshop on Regional Cooperation to Address Poaching of Sea Turtles, Kuala Terengganu, Malaysia.

Polovina, J., Uchida, I., Balazs, G.H., Howell, E.A., Parker, D., & Dutton, P. (2006). The Kuroshio Extension Bifurcation Region: a pelagic hotspot for juvenile loggerhead sea turtles. *Deep-sea Research II*, 53, 326-339.

Rees, A.F., Al-Kiyumi, A., Broderick, A.C., Papathanasopoulou, N., & Godley, B.J.



(2012). Conservation related insights into the behavior of the olive ridley sea turtle *Lepidochelys olivacea* nesting in Oman. *Marine Ecology Progress Series*, ^, 195-205.

Rees, A.F., Hafez, A.A., Lloyd, J.R., Papathansopoulou N., & Godley, B.J. (2013). Green turtles, *Chelonia mydas*, in Kuwait: nesting and movements. *Chelonia Conservation and Biology*, 12, 157-163.

Renaud, M.L., Carpenter, J.A., & Williams, J. A. (1995). Activities of juvenile green turtle, *Chelonia mydas*, at a jettied pass in South Texas. *Fishery Bulletin*, 93, 586–593

Rice, M.R., & Balazs, G. H. (2008). Diving behavior of the Hawaiian green turtle (*Chelonia mydas*) during oceanic migrations. *Journal of Experimental Marine Biology and Ecology*, 356, 121-127.

Rodgers, A. R., Carr, A. P., Smith, L., & Kie, J. G. (2005). HRT: Home Range Tools for ArcGIS. Ontario Ministry of Natural Resources, Centre for Northern Forest Ecosystem Research, Thunder Bay, Ontario, Canada.

Scales, K.L., Lewis, J.A., Lewis, J.P., Castellanos, D., Godley, B.J. & Graham, R.T. (2011). Insights into habitat utilization of the hawksbill turtle, *Eretmochelys imbricata* (Linnaeus, 1766), using acoustic telemetry. *Journal of Experimental Marine Biology and Ecology*, 407, 122-129.

Scott, R., Hodgson, D.J., Witt, M.J., Coyne, M.S., Adnyana, W., Blumenthal, J.M.,...Godley, B.J. (2012). Global analysis of satellite tracking data shows that adult green turtles are significantly aggregated in Marine Protected Areas. *Global Ecology and Biogeography*, 21, 1053-1061.

Seminoff, J.A., Resendiz, A., & Nichols, W.J. (2002). Home range of green turtles *Chelonia mydas* at a coastal foraging area in the Gulf of California, Mexico. *Marine Ecology Progress Series*, 242, 253-265.

Senko, J., Lopez-Castro, M.C., Koch, V., & Nichols, W.J. (2010). Immature East Pacific green turtles (*Chelonia mydas*) use multiple foraging areas off the Pacific coast of Baja California Sur, Mexico: first evidence from mark-recapture data. *Pacific Science*, 64, 125-130.

Shaver, D.J., Hart, K.M., Fujisaki, I., Rubio, C., & Sartain, A.R. (2013). Movement mysteries unveiled: spatial ecology of juvenile green sea turtles. Lutterschemidt., W.I., (Eds.), *Reptiles in Research* (pp. 463-484), Nova Science Publishers, Inc.

Song, X.J., Wang, H.J., Wang, W.Z., Gu, H.X., Chan, S., & Jiang, H.S. (2002). Satellite Tracking of Post-Nesting Movements of Green Turtles *Chelonia mydas* from the Gangkou Sea Turtle National Nature Reserve, China, 2001. *Marine Turtle Newsletter*, 97, 8-9.

Traffic. 2014. Indonesia, Malaysia, Philippines and Viet Nam join forces to crack down on turtle trade. <http://www.traffic.org/home/2014/6/6/indonesia-malaysia-philippines-and-viet-nam-join> [20 October 2015]

Wang, Y. M. (1993). Current status and prospect of sea turtle resources and research in South China Sea. *Chinese Journal of Ecology*, 12, 60-61.

Wang, Y. M., & Li, W. (2008). Sea turtle survey, monitoring and awareness promotion programme in Mainland China. Final Report. Fish and Wildlife Service, US Department of Interior.

Watson, K.P., & Granger, R.A. (1998). Hydrodynamic effect of a satellite transmitter on a juvenile green turtle (*Chelonia mydas*). *Journal of Experimental Biology*, 201, 2496-2505.

Whiting, S.D., & Miller, J.D. (1998). Short term foraging ranges of adult green turtles (*Chelonia mydas*). *Journal of Herpetology*, 32, 330–337

Witt, M.J., Akesson, S., Broderick A.C., Coyne, M.S., Ellick, J., Formia, A., ...Godley, B.J. (2010). Assessing accuracy and utility of satellite-tracking data using Argos-linked Fastloc-GPS. *Animal Behaviour*, 80, 571-581.

Xia, Z.R., & Gu., H.X. (2012). Reports of satellite tracking green sea turtles in China. *Sichuan Journal of Zoology*, 31, 435-438.

Xu, G., & Zheng, M.Q. (2003). Achievement from sea turtle conservation in Nanao. *Fisheries Science and Technology*, 4, 14.

Yasuda, T., & Arai, N. (2005). Fine-scale tracking of marine turtles using GPS-Argos PTTs. *Zoological Science*, 22, 547-553.

Ye, M.B., Chen, H.L., Gu, H.X., & Li, P.P. (2015). Satellite tracking of migration for three captive-reared juvenile green turtles from Huidong National Sea Turtle Reserve. *Sichuan Journal of Zoology*, 34, 15-20.

Yeh, F.C., Balazs, G.H., Parker, D.M., Ng, C.K.Y., & Shi, H.T. (2014). Novel use of satellite tracking as a forensic tool to determine foraging ground of a rescued green turtle (*Chelonia mydas*). *Marine Turtle Newsletter*, 142, 1-3.




Figure 1 Nesting sites, foraging grounds and migratory pathways of green turtles and loggerhead turtles determined by satellite telemetry in South China Sea. The red star and line denote a green turtle nesting site in Hong Kong and post-nesting movement; Yellow=Gangkou; Light green=Wan-an; Blue=Lanyu; White=Liouciou Island; Orange=Taipin, Nansha Island. Purple = loggerhead in Minabe, Japan. The numbers in closed circles represent foraging grounds: 1 Wanning of Hainan, 2 Dao Bach Long Vi in Vietnam, 3 East coast of Leizhou Peninsula, 4 Okinawa Island in southern Japan, 5 Coastal waters off northern Taiwan, 6 Qinpeng-Dao of Nanao Island at Shantou, 7 Huidong, 8 Pratas (Dongsha) Island, 9 Dangan Liedao of Wanshan Archipelago, 10 northern Philippines, 11 Iriomote-jima and Ishigaki-shima of Ryukyu Archipelago, 12 Koshiki, Okinawa Island in southern Japan, 13 Coral reefs in the southern Penghu Archipelago, north of Chimei Island, 14 Coastal waters off Palawan Island, the Philippines; 15 the north coast of eastern Malaysia; 16 east coast of Luzon Island, the Philippines, 17 East China Sea. (References refer to Table 3, Appendix)




Figure 2 Distribution of major nesting and foraging grounds of sea turtles in South China Sea. Green closed circles denote green turtle foraging grounds; open circles denote nesting grounds. Orange closed triangles denote hawksbill turtle foraging grounds; open triangles denote nesting grounds. Blue closed pentagons denote olive ridley foraging grounds. Purple closed squares denote loggerhead turtle foraging grounds; open squares denote nesting grounds. Black stars denote leatherback turtle foraging grounds. Numbers in closed circle represents foraging grounds as described for Figure 1 (References refer to Table 3, Appendix).