




# NOAA Technical Memorandum NMFS-SEFC-116

## Size Composition of Monthly Catches of Brown Shrimp from the Texas Coast, Mississippi River to Texas, and Pensacola to the Mississippi River, 1960- 1981.



MAY 1983

U.S. DEPARTMENT OF COMMERCE  
National Oceanic and Atmospheric Administration  
National Marine Fisheries Service  
Southeast Fisheries Center  
Galveston Laboratory  
Galveston, Texas 77550



# **NOAA Technical Memorandum NMFS-SEFC-116**

## **Size Composition of Monthly Catches of Brown Shrimp from the Texas Coast, Mississippi River to Texas, and Pensacola to the Mississippi River, 1960- 1981.**

By

Charles Wax Caillouet, Jr.

and

Dennis Brian Koi

---

U. S. DEPARTMENT OF COMMERCE

Malcolm Baldrige, Secretary

National Oceanic and Atmospheric Administration

John V. Byrne, Administrator

National Marine Fisheries Service

William G. Gordon, Assistant Administrator for Fisheries

---

**MAY 1983**

---

Technical Memorandums are used for documentation and timely communication of preliminary results, interim reports, or special-purpose information, and have not received complete formal review, editorial control, or detailed editing.

NOTICE

This report should be cited as follows:

Caillouet, Charles Wax, Jr. and Dennis Brian Koi. 1983. Size composition of monthly catches of brown shrimp from the Texas Coast, Mississippi River to Texas, and Pensacola to the Mississippi River, 1960-1981. NOAA Technical Memorandum NMFS-SEFC-116, 73 p.

It can be obtained by writing to the National Technical Information Service, 5258 Port Royal Road, Springfield, Virginia 22161.

## INTRODUCTION

Christmas and Etzold (1977) and the Gulf of Mexico Fishery Management Council (GMFMC, 1980) have summarized much of the available information concerning the biology and population dynamics of brown shrimp in the context of management of the fishery for this species in the Gulf of Mexico. The size composition of the reported monthly catches of brown shrimp, Penaeus aztecus, reflects the combined effects of recruitment, growth and mortality, including losses due to natural causes and those caused by fishing. Annually recurring recruitment has an obvious effect of reducing the size of brown shrimp in the monthly catches, but the time-phasing of open seasons and the intensity of fishing also can alter the size composition patterns (Caillouet and Koi, 1981 and 1983; Caillouet, Patella and Jackson, 1979 and 1980).

The seasonal patterns in the monthly catches of brown shrimp vary from year to year and area to area, depending upon climatic conditions, recruitment, survival, and growth of the shrimp, the timing and duration of open seasons, and the intensity of fishing during open seasons set by State and Federal shrimp management agencies (Caillouet and Koi, 1982). Caillouet and Koi (1982) graphically portrayed the cumulative monthly catches, the cumulative ex-vessel value of the catches, and the average monthly ex-vessel price per pound for this species to elucidate these seasonal patterns, as well as year to year trends.

Ex-vessel value of the brown shrimp catch is strongly linked to the size composition of the catch (Caillouet and Koi, 1981, 1983; Caillouet and Patella, 1978; Caillouet, Patella and Jackson, 1979 and 1980) as well as total weight of the catch. In addition, the seasonal variations in local supply of shrimp influence the price, and thus the ex-vessel value of the catch (Poffenberger, 1982a, 1982b).

The purposes of this paper are to summarize available brown shrimp catch statistics by size categories, and to portray graphically the

monthly size composition of the catches from the Texas Coast, Mississippi River to Texas, and Pensacola to the Mississippi River for calendar years 1960 through 1981. For this purpose, we used combined inshore (landward of barrier islands) and offshore (seaward of barrier islands) catches. These graphically portrayed summaries should quickly convey information on the monthly patterns and trends in size composition of the brown shrimp catches. They also can be compared, month by month, with the information in our previously published graphs (Caillouet and Koi, 1982).

## METHODS

### Description of Data

Monthly summations of reported catches (inshore and offshore combined) of brown shrimp by size category were compiled from data files available from the NMFS, Southeast Fisheries Center (SEFC), Fisheries Information Management Division (FIMD), Miami, Florida. The weight of each monthly catch was expressed in pounds (heads off) distributed among eight size categories (<15, 15-20, 21-25, 26-30, 31-40, 41-50, 51-67, and ≥68 shrimp tails per pound), usually referred to as "count". As an example, the catch of brown shrimp from the Texas Coast in the month of August 1981 is shown by size categories in Table 1. Monthly summations were obtained for each of three coastal areas (Fig. 1) distinguished as follows:

- (1) Texas Coast (Statistical areas 18-21 combined),
- (2) Mississippi River to Texas (Statistical areas 13-17 combined), representing that part of the Louisiana coast west of the Mississippi River, and
- (3) Pensacola to the Mississippi River (Statistical areas 10-12 combined), representing that part of the Louisiana coast east

Table 1. Distribution of the brown shrimp catch (inshore and offshore combined) by size category for the Texas Coast in August 1981.

	Size Category (number of shrimp per pound, heads off)								
	≥68	51-67	41-50	31-40	26-30	21-25	15-20	≤15	Totals
Weight (thousands of pounds, heads off)	406	1,839	1,601	6,668	2,321	1,317	341	39	14,532
Percentage by weight	2.8	12.6	11.0	45.9	16.0	9.1	2.3	0.3	100.0
Cumulative percentage by weight	2.8	15.4	26.4	72.3	88.3	97.4	99.7	100.0	-




Figure 1. Boundaries of statistical areas 10-21, and three coastal regions (Texas Coast, Mississippi River to Texas, and Pensacola to the Mississippi River).

of the Mississippi River, the Mississippi coast, the Alabama coast, and a small part of the upper west coast of Florida (catches from Pensacola Bay are not included in this area; they are allocated to the adjacent Apalachicola area by the FIMD).

The percentages of the monthly catch (by weight) represented by each of the eight size categories were calculated, then the cumulative percentages by size category were derived for each month as exemplified in Table 1 for August 1981.

For each coastal area and year, the cumulative percentage (by weight) of the monthly catch of brown shrimp for each size category was plotted by month (see graphs). Thus, on any given graph, the bottommost line connects the monthly percentages for shrimp in the  $\geq 68$  count category. The line immediately above it connects the combined monthly percentages for both the  $\geq 68$  count and 51-67 count categories, and so on. The top line (100%) on each graph connects the combined monthly percentages for the all size categories. Thus, the distance between lines reflects the actual monthly percentage (by weight) taken in a given count category.

For each year, in consecutive order from 1960 through 1981, the graphs contained in this report are grouped in triplicates (one graph for each coastal area): the first for the Texas Coast, the second for the Mississippi River to Texas, and the third for Pensacola to the Mississippi River.

#### GENERAL OBSERVATIONS

The most prominent feature on all of the graphs was the dramatic upsurge in percentage of the catch represented by the smallest shrimp ( $\geq 68$  per pound, or  $\geq 68$  count), which usually (but not always) appeared in May or June. Minor upsurges in percentage of the smallest shrimp sometimes occurred in fall and winter, especially in the earlier years of the series. The major upsurges in percentage of



smallest shrimp obviously represented recruitment. In some years, a series of successive peaks, representing growth of the shrimp into successively lower count classes (i.e., consecutively larger sizes), occurred over several months.

There also were striking differences among the three coastal areas with regard to the magnitude of the upsurge in percentage of smallest shrimp ( $\geq 68$  count). The most pronounced upsurge in this percentage occurred in the Mississippi River to Texas coastal area. It was usually of an intermediate level for Pensacola to the Mississippi River, and was lowest for the Texas Coast (note: for the Texas Coast in 1960, this percentage was barely detectable). These differences undoubtedly reflected the well-known differences in fishing regulations and strategies among the three coastal areas (Caillouet and Koi, 1981, 1983; Caillouet and Patella, 1978; Caillouet et al, 1979, 1980).

The graphs in this report may be used in conjunction with those of our previous publication (Caillouet and Koi, 1982) which showed the changes in weight, value and price per pound of the monthly catch. The two papers together depict the fluctuations in catch and its size composition from month to month over the same period of years. Fisheries scientists and managers should find them useful in interpreting seasonal patterns in the contexts of the life cycle of brown shrimp and changes in State and Federal management practices over the years.

#### LITERATURE CITED

Caillouet, C. W., Jr. and D. B. Koi. 1981. Trends in ex-vessel value and size composition of reported May-August catches of brown shrimp and white shrimp from the Texas, Louisiana, Mississippi, and Alabama Coasts, 1960-1978. Gulf Research Reports 7(1):59-70.

\_\_\_\_\_ and \_\_\_\_\_. 1982. Cumulative monthly weight and ex-vessel value, and monthly price per pound, for brown shrimp catches from the northern Gulf of Mexico, 1960-1981. NOAA Technical Memorandum NMFS-SEFC-96, 6 p. plus 67 figures.

\_\_\_\_\_ and \_\_\_\_\_. 1983. Ex-vessel value and size composition of reported May-August catches of brown shrimp and white shrimp from 1960 to 1981 as related to the Texas closure. Gulf Research Reports 7(3):(in press).

\_\_\_\_\_ and F. J. Patella. 1978. Relationship between size composition and ex-vessel value of reported shrimp catches from two Gulf coast states with different harvesting strategies. Marine Fisheries Review 40(2):14-18.

\_\_\_\_\_, \_\_\_\_\_, and W. B. Jackson. 1979. Relationship between marketing category (count) composition and ex-vessel value of reported annual catches of shrimp in the eastern Gulf of Mexico. Marine Fisheries Review 41(5-6):1-7.

\_\_\_\_\_, \_\_\_\_\_, and \_\_\_\_\_. 1980. Trends toward decreasing size of brown shrimp, Penaeus aztecus, and white shrimp, Penaeus setiferus, in reported annual catches from Texas and Louisiana. U.S. Fishery Bulletin 77(4):985-989.


Christmas, J. Y. and D. J. Etzold (editors). 1977. The shrimp fishery of the Gulf of Mexico United States: a regional management plan. Technical Report Series No. 2, Gulf Coast Research Laboratory, Ocean Springs, MS. 128 pp.

Gulf of Mexico Fishery Management Council (GMFMC). 1980. Fishery management plan for the shrimp fishery of the Gulf of Mexico. Federal Register 45(218):74190-74308.

Poffenberger, J. R. 1982a. Estimated impacts on ex-vessel brown shrimp prices and value as a result of the Texas Closure regulation. Marine Fisheries Review 44(9-10):38-43.


\_\_\_\_\_. 1982b. Estimated impacts of Texas closure regulation on ex-vessel prices and value, 1981 and 1982. NOAA Technical Memorandum NMFS-SEFC-111, 34 p.

BROWN SHRIMP  
TEXAS COAST  
1960




BROWN SHRIMP  
MISSISSIPPI RIVER TO TEXAS  
1960


CUMULATIVE




BROWN SHRIMP  
PENSACOLA TO THE MISSISSIPPI RIVER  
1960




BROWN SHRIMP  
TEXAS COAST  
1961



BROWN SHRIMP  
MISSISSIPPI RIVER TO TEXAS  
1961




BROWN SHRIMP  
PENSACOLA TO THE MISSISSIPPI RIVER  
1961






BROWN SHRIMP  
TEXAS COAST  
1962




# BROWN SHRIMP MISSISSIPPI RIVER TO TEXAS 1962






BROWN SHRIMP  
TEXAS COAST  
1963

CUMULATIVE  
PERCENT




# BROWN SHRIMP MISSISSIPPI RIVER TO TEXAS 1963

CUMULATIVE  
PERCENTAGE




BROWN SHRIMP  
PENSACOLA TO THE MISSISSIPPI RIVER  
1963




BROWN SHRIMP  
TEXAS COAST  
1964

CUMULATIVE  
PERCENT  
20




BROWN SHRIMP  
MISSISSIPPI RIVER TO TEXAS  
1964







BROWN SHRIMP  
PENSACOLA TO THE MISSISSIPPI RIVER  
1964



BROWN SHRIMP  
TEXAS COAST  
1965




BROWN SHRIMP  
MISSISSIPPI RIVER TO TEXAS  
1965




BROWN SHRIMP  
PENSACOLA TO THE MISSISSIPPI RIVER  
1965

25


CUMULATIVE  
PERCENT




BROWN SHRIMP  
TEXAS COAST  
1966




BROWN SHRIMP  
MISSISSIPPI RIVER TO TEXAS  
1966



# BROWN SHRIMP PENSACOLA TO THE MISSISSIPPI RIVER 1966




BROWN SHRIMP  
TEXAS COAST  
1967






BROWN SHRIMP  
MISSISSIPPI RIVER TO TEXAS  
1967




# BROWN SHRIMP PENSACOLA TO THE MISSISSIPPI RIVER 1967


31




BROWN SHRIMP  
TEXAS COAST  
1968




# BROWN SHRIMP MISSISSIPPI RIVER TO TEXAS 1968




BROWN SHRIMP  
PENSACOLA TO THE MISSISSIPPI RIVER  
1968




BROWN SHRIMP  
TEXAS COAST  
1969



BROWN SHRIMP  
MISSISSIPPI RIVER TO TEXAS  
1969




# BROWN SHRIMP PENSACOLA TO THE MISSISSIPPI RIVER 1969






BROWN SHRIMP  
TEXAS COAST  
1970




BROWN SHRIMP  
MISSISSIPPI RIVER TO TEXAS  
1970

CUMULATIVE  
PERCENT




BROWN SHRIMP  
PENSACOLA TO THE MISSISSIPPI RIVER  
1970




BROWN SHRIMP  
TEXAS COAST  
1971


CUMULATIVE  
PERCENT  
41




# BROWN SHRIMP MISSISSIPPI RIVER TO TEXAS 1971



BROWN SHRIMP  
PENSACOLA TO THE MISSISSIPPI RIVER  
1971




BROWN SHRIMP  
TEXAS COAST  
1972



BROWN SHRIMP  
MISSISSIPPI RIVER TO TEXAS  
1972


45






BROWN SHRIMP  
PENSACOLA TO THE MISSISSIPPI RIVER  
1972


46  
CUMULATIVE PERCENT




BROWN SHRIMP  
TEXAS COAST  
1973



# BROWN SHRIMP MISSISSIPPI RIVER TO TEXAS 1973




BROWN SHRIMP  
PENSACOLA TO THE MISSISSIPPI RIVER  
1973




BROWN SHRIMP  
TEXAS COAST  
1974

CUMULATIVE  
PERCENT




BROWN SHRIMP  
MISSISSIPPI RIVER TO TEXAS  
1974




BROWN SHRIMP  
PENSACOLA TO THE MISSISSIPPI RIVER  
1974

CUMULATIVE  
PERCENT




BROWN SHRIMP  
TEXAS COAST  
1975

CUMULATIVE  
PERCENT






BROWN SHRIMP  
MISSISSIPPI RIVER TO TEXAS  
1975



BROWN SHRIMP  
PENSACOLA TO THE MISSISSIPPI RIVER  
1975


CUMULATIVE  
PERCENT

55




BROWN SHRIMP  
TEXAS COAST  
1976

CUMULATIVE  
PERCENT




BROWN SHRIMP  
MISSISSIPPI RIVER TO TEXAS  
1976

57



BROWN SHRIMP  
PENSACOLA TO THE MISSISSIPPI RIVER  
1976


58




BROWN SHRIMP  
TEXAS COAST  
1977

CUMULATIVE  
PERCENTAGE


59



# BROWN SHRIMP MISSISSIPPI RIVER TO TEXAS 1977




# BROWN SHRIMP PENSACOLA TO THE MISSISSIPPI RIVER 1977






BROWN SHRIMP  
TEXAS COAST  
1978

CUMULATIVE  
62




BROWN SHRIMP  
MISSISSIPPI RIVER TO TEXAS  
1978




BROWN SHRIMP  
PENSACOLA TO THE MISSISSIPPI RIVER  
1978

CUMULATIVE  
64




BROWN SHRIMP  
TEXAS COAST  
1979



# BROWN SHRIMP MISSISSIPPI RIVER TO TEXAS 1979


99



## 1979




BROWN SHRIMP  
TEXAS COAST  
1980




BROWN SHRIMP  
MISSISSIPPI RIVER TO TEXAS  
1980

69  
CUMULATIVE  
PERCENT






BROWN SHRIMP  
PENSACOLA TO THE MISSISSIPPI RIVER  
1980



BROWN SHRIMP  
TEXAS COAST  
1981

CUMULATIVE  
PERCENT



The graph displays the percentage of the population in different age groups across the months of the year. The Y-axis represents the percentage from 0 to 100. The X-axis lists the months from January (J) to December (D). The age groups are labeled as follows: <15, 15-20, 21-25, 26-30, 31-40, 41-50, 51-67, and  $\geq 68$ . All age groups reach 100% in May. The <15 group starts at approximately 90% in January and remains high. The 15-20 group starts at 22% in January and peaks at 42% in February. The 21-25 group starts at 10% in January and peaks at 27% in February. The 26-30 group starts at 5% in January and peaks at 12% in February. The 31-40 group starts at 2% in January and peaks at 3% in February. The 41-50 group starts at 1% in January and peaks at 2% in February. The 51-67 group starts at 0% in January and peaks at 1% in February. The  $\geq 68$  group starts at 0% in January and peaks at 1% in February.

CUMULATIVE  
PERCENT

72

BROWN SHRIMP  
PENSACOLA TO THE MISSISSIPPI RIVER  
1981

