




NOAA Technical Memorandum NMFS-F/NEC-31

Evidence of Nearshore Summer Upwelling Off Atlantic City, New Jersey



U.S. DEPARTMENT OF COMMERCE
National Oceanic and Atmospheric Administration
National Marine Fisheries Service
Northeast Fisheries Center
Woods Hole, Massachusetts
November 1984

NOAA TECHNICAL MEMORANDUM NMFS-F/NEC

Under the National Marine Fisheries Service's mission to "Achieve a continued optimum utilization of living resources for the benefit of the Nation," the Northeast Fisheries Center (NEFC) is responsible for planning, developing, and managing multidisciplinary programs of basic and applied research to: (1) better understand the living marine resources (including marine mammals) of the Northwest Atlantic, and the environmental quality essential for their existence and continued productivity; and (2) describe and provide to management, industry, and the public, options for the utilization and conservation of living marine resources and maintenance of environmental quality which are consistent with national and regional goals and needs, and with international commitments.

The timely need for such information by decision makers often precludes publication in formal journals. The *NOAA Technical Memorandum NMFS-F/NEC* series provides a relatively quick and highly visible outlet for documents prepared by NEFC authors, or similar material prepared by others for NEFC purposes, where formal review and editorial processing are not appropriate or feasible. However, documents within this series reflect sound professional work and can be referenced in formal journals. Any use of trade names within this series does not imply endorsement.

Copies of this and other *NOAA Technical Memorandums* are available from the National Technical Information Service, 5285 Port Royal Rd., Springfield, VA 22161. Recent issues of *NOAA Technical Memorandum NMFS-F/NEC* are noted below:

16. *Contaminants in New York Bight and Long Island Sediments and Demersal Species, and Contaminant Effects on Benthos, Summer 1980.* By Robert N. Reid, John E. O'Reilly, and Vincent S. Zdanowicz, eds. September 1982. x + 96 p., 36 figs., 21 tables. NTIS Access No. PB83-152116.

17. *Summary of the Physical Oceanographic Processes and Features Pertinent to Pollution Distribution in the Coastal and Offshore Waters of the Northeastern United States, Virginia to Maine.* By Merton C. Ingham (ed.), Reed S. Armstrong, J. Lockwood Chamberlin, Steven K. Cook, David G. Mountain, Ronald J. Schlitz, James P. Thomas, James J. Bisagni, John F. Paul, and Catherine E. Warsh. December 1982. vi + 166 p., 21 figs., 2 tables. NTIS Access. No. PB83-156364.

18. *Stock Discrimination of Summer Flounder (*Paralichthys dentatus*) in the Middle and South Atlantic Bights: Results of a Workshop.* By Michael J. Fogarty, Glenn DeLaney, John W. Gillikin, Jr., John C. Poole, Daniel E. Ralph, Paul G. Scarlett, Ronald W. Smith, and Stuart J. Wilk. January 1983. iii + 14 p., 2 figs., 3 tables. NTIS Access. No. PB83-168856.

(continued on inside back cover)



NOAA Technical Memorandum NMFS-F/NEC- 31

This TM series is used for documentation and timely communication of preliminary results, interim reports, or special purpose information, and has not received complete formal review, editorial control, or detailed editing.

Evidence of Nearshore Summer Upwelling Off Atlantic City, New Jersey

Merton C. Ingham¹ and James Eberwine²

¹Atlantic Environmental Group, National Marine Fisheries Service, Narragansett, RI 02882

²National Weather Service Office, Atlantic City, NJ 08405

U.S. DEPARTMENT OF COMMERCE

Malcolm Baldrige, Secretary

National Oceanic and Atmospheric Administration

John V. Byrne, Administrator

National Marine Fisheries Service

William G. Gordon, Assistant Administrator for Fisheries

Northeast Fisheries Center

Woods Hole, Massachusetts

November 1984

ABSTRACT

Analysis of nearshore ocean temperature data collected in the floatwell of a tide gauge on Steel Pier and wind data collected at the Atlantic City, New Jersey, weather station during summer months of 1981 shows a strong correlation between major cooling or warming events in the water and the strength of upwelling or downwelling wind components one day earlier.

EVIDENCE OF NEARSHORE SUMMER UPWELLING

OFF ATLANTIC CITY, NEW JERSEY

Merton C. Ingham and James Eberwine

One of the tenets of classical theory of coastal upwelling dynamics (Defant 1961, p. 652) holds that the most effective wind direction for transporting surface waters offshore, thereby bringing cooler water up into the surface layer, is one which is parallel to the coastline (with the land on the left in the northern hemisphere). Assuming that this is applicable to the waters over the continental shelf in the Middle Atlantic Bight, upwelling should occur along the Virginia - New Jersey coastline during the summer months when winds from the southwestern quadrant are common (Williams and Godshall 1977; Lettau et al. 1976).

Hicks and Miller (1980) reported indications of summer upwelling in response to southerly winds, bringing cold bottom water into the surf zone near Monterey Beach, New Jersey (about 80 km northeast of Atlantic City, New Jersey). They found upwelling in July 1976 to be vigorous enough, following two days of strong southerly winds, to bring waters of the "cold cell" into the surf zone, causing a drop of 9.3°C (from 18.3° to 9.0°C) in surf temperature. Their analysis of 1973 and 1974 data also showed large decreases of surf temperature during July, August, and September, each preceded one or two days earlier by strong winds from the south and southwest.

Recreational surf fishermen along the New Jersey shore also have frequently observed the upwelling effects of strong south winds, but haven't used the term "upwelling" or the concept in describing their experiences. Wilson (1983), for example, speaks of "cold currents" and

"cold waters" being brought in by southerly winds, and describes the effects of the cooler water on the behavior of several species of gamefish.

An unusual opportunity to gather evidence of wind-driven coastal upwelling exists in the area of Atlantic City, New Jersey, as the consequence of the availability of accurate wind data taken at the weather station (a first-order station of the National Weather Service at the Aviation Facilities Experimental Center) and water temperature data logged four times a day at the tide gauge station on the Steel Pier. Although the weather station is about 16 km inland from the pier, wind data collected there and reduced to daily averages realistically portray general wind field conditions along the coast and in the vicinity of the pier.

The trend of the local coastline is about WSW to ENE in the vicinity of the pier (Fig. 1), which means that a wind from about 250° would have the maximum upwelling effect, and one from 70° would produce the maximum convergence and "downwelling". During the summer months, the monthly mean resultant wind directions (based on 1941-70 data) observed at the weather station are: June - 140° , July - 260° , August - 200° , and September - 270° . (U.S. Department of Commerce 1975).

The wind data used in this study were daily mean vector resultant velocities taken from Local Climatological Data summaries for the Atlantic City weather station published monthly by the National Climatic Data Center.⁽¹⁾ Upwelling wind components are computed from these data

(1) National Climatic Data Center, National Environmental Satellite, Data, and Information Service, NOAA, Asheville, North Carolina 28801.

by multiplying them by the cosine of the angle between the published daily resultant and the 250° - 70° line and assigning the appropriate sign, positive for upwelling or negative for downwelling.

The seawater temperature data were obtained from manuscript records of 6-hourly observations recorded at the Atlantic City weather station by telephone link with a thermistor installed in the floatwell of the National Ocean Service tide station on Steel Pier. The average depth of the thermistor beneath the surface is 2.4 m, ranging from about 3.2 m at high tide to 1.9 m at low tide. The 6-hourly data were converted to daily averages, and daily changes in water temperature were calculated by subtracting each day's average from the next day's.

Visual comparison of time-series plots of daily water temperature and wind component data for summer 1981 (Fig. 2) was inconclusive, though there seemed to be some correspondence between major wind events and rapid changes in temperature. Linear regression analysis of daily water temperature changes and the upwelling wind component the day before yielded correlation coefficients which were too low to imply a functional relationship. It appeared that random small-amplitude noise in the water temperature data might be responsible for the poor correlation. Accordingly, we re-ran the linear regression analysis using just those water temperature changes which were at least one standard deviation above or below monthly means (high-pass filter). The results of this analysis were far more conclusive (Fig. 3), yielding correlation coefficients (R) of -0.777 , -0.855 , and -0.752 for June, July, and August, respectively. The coefficients are negative because upwelling wind components (+) produce decreases in water temperature and downwelling components (-) produce increases in water temperature.

Best-fit linear equations for the three months of high-pass data are: June, $Y = 4.18 - 0.89 X$; July, $Y = 2.27 - 1.16 X$; and August, $Y = 2.51 - 0.91 X$, where Y = the one-day change in water temperature ($^{\circ}\text{F}$) and X = upwelling wind component (mi/hr). These equations have quite similar slopes but different intercepts (Fig. 3), which suggests that changes in upwelling/downwelling wind strength produce comparable changes in water temperature in each of the three months, but that the wind strength required for a given change is greater in June than in July and August, probably because of smaller differences between the temperature of upwelled bottom water and surface water in June.

In an attempt to document and verify the upwelling events, a search of a file of infrared imagery from the GOES satellite for summer 1981 was conducted. It revealed that there was clear sky in the Atlantic City area both before and after just one of the upwelling events, the one which began about 13 July. Imagery from the Advanced Very High Resolution Radiometer on the NOAA 6 satellite revealed a narrow band of cooler water in the Atlantic City area on the 15th that wasn't present on the 11th (Figs. 4 and 5).

The results of this study of wind and water temperature data from just one summer suggest that wind-driven upwelling does occur along the central New Jersey coast. More sophisticated analyses of summertime conditions in other years should be conducted before more extensive quantification or modeling is undertaken, however. Specifically, it would be important to test the assumption that 250° winds are the most effective for upwelling and to determine more accurately the lag time between wind conditions and water temperature changes.

REFERENCES

Defant, A. 1961. Physical oceanography. Vol. 1.

MacMillan Co., New York. 729 pp.

Hicks, D.C., and J.R. Miller. 1980. Meteorological forcing and bottom water movement off the northern New Jersey coast. Estuarine Coastal Mar. Sci. 11: 563-571.

Lettau, B., W.A. Brower, Jr., and R.G. Quayle. 1976. Marine climatology. MESA N. Y. Bight Atlas Monogr. 7. New York Sea Grant Institute, Albany, N.Y. 239 pp.

U.S. Department of Commerce. 1975. Local climatological data -- annual summaries for 1974. Part 11. NEB-WY0. National Climatological Data Center, Asheville, N. C.

Williams, R.G., and F.A. Godshall. 1977. Summarization and interpretation of historical physical oceanographic and meteorological information for the Mid-Atlantic region. Report prepared for the Bureau of Land Management by the National Oceanic and Atmospheric Administration. NTIS # PB-277-104. 295 pp.

Wilson, Russ. 1983. Ill wind blows no good. The Fisherman -- The New Jersey, Delaware Bay Edition, 25 August: 20-21.




FIGURE 1. Location of the study area and points at which water temperature and wind velocity data were collected.




FIGURE 2. Trends in daily average water temperature (°F) measured at the Steel Pier and upwelling/downwelling wind component (mi/hr) from wind velocity measurements made at the Atlantic City weather station in 1981.




FIGURE 3. Best-fit linear regressions of water temperature change ($^{\circ}$ F) and upwelling (+)/downwelling (-) wind components (mi/hr) for high-pass filtered data (Δ water temperatures greater than 1 standard deviation from monthly mean).



FIGURE 4. Infrared imagery from the NOAA 6 satellite for 11 July 1981. Arrow shows location of Atlantic City, New Jersey.



FIGURE 5. Infrared imagery from the NOAA 6 satellite for 15 July 1981. Arrow shows location of Atlantic City, New Jersey.

(continued from inside front cover)

19. *Environmental Benchmark Studies in Casco Bay - Portland Harbor, Maine, April 1980.* By Peter F. Larsen, Anne C. Johnson, and Lee F. Doggett. January 1983. xi + 173 p., 39 figs., 12 tables, 2 app. NTIS Access. No. PB83-184069.

20. *Annual NEMP Report on the Health of the Northeast Coastal Waters of the United States, 1981.* Northeast Monitoring Program Report No. NEMP-IV-82-65. February 1983. xii + 86 p., 21 figs., 15 tables, 1 app. NTIS Access. No. PB83-193912.

21. *MARMAP Plankton Survey Manual.* By Jack W. Jossi and Robert R. Marak. March 1983. xvii + 260 p., 41 figs., 3 tables, 2 app. NTIS Access. No. PB83-210203.

22. *Status of the Fishery Resources Off the Northeastern United States for 1982.* By Resource Assessment Division, Northeast Fisheries Center. June 1983. iii + 128 p., 44 figs., 44 tables. NTIS Access. No. PB83-236554.

23. *Nantucket Shoals Flux Experiment Data Report I. Hydrography.* By W. Redwood Wright. June 1983. i + 105 p., 100 figs., 1 table. NTIS Access. No. PB83-236562.

24. *Residual Drift and Residence Time of Georges Bank Surface Waters with Reference to the Distribution, Transport, and Survival of Larval Fishes.* By John B. Colton, Jr., and Jacquelyn L. Anderson. June 1983. ix + 45 p., 22 figs., 2 tables, 1 app. NTIS Access. No. PB84-107820.

25. *Histological Techniques for Marine Bivalve Mollusks.* By Dorothy W. Howard and Cecelia S. Smith. June 1983.

26. *106-Mile Site Characterization Update.* By John B. Pearce, Don C. Miller, and Carl Berman, eds. August 1983. xxxi + 483 p., 180 figs., 32 tables, 1 app. NTIS Access. No. PB84-118363.

27. *Pelagic Distributions of Marine Birds Off the Northeastern United States.* By Kevin D. Powers. November 1983. xvi + 201 p., 116 figs., 5 tables, 9 app. NTIS Access. No. PB84-187871.

28. *Food of Seventeen Species of Northwest Atlantic Fish.* By Ray E. Bowman and William L. Michaels. January 1984. xx + 183 p., 2 figs., 61 tables, 19 app. NTIS Access. No. PB84-219195.

29. *Status of the Fishery Resources Off the Northeastern United States for 1983.* By Resource Assessment Division, Northeast Fisheries Center; Emory D. Anderson, ed. July 1984. iii + 132 p., 44 figs., 48 tables. NTIS Access. No. PB85-106847.

30. *Recent Estimates of Adult Spawning Stock Biomass Off the Northeastern United States from MARMAP Ichthyoplankton Surveys.* By Peter Berrien, Wallace Morse, and Michael Pennington. July 1984. ix + iii p., 25 figs., 25 tables. NTIS Access. No. PB85-108991.

Information & Publications Office
Northeast Fisheries Center
National Marine Fisheries Service, NOAA
Water St.
Woods Hole, MA 02543

Postage and Fees Paid
U.S. Department of Commerce
COM-210

THIRD CLASS