

NOAA Technical Report NMFS 18



A Temporal and Spatial Study of Invertebrate Communities Associated With Hard-Bottom Habitats in the South Atlantic Bight

E. L. Wenner, P. Hinde, D. M. Knott, and
R. F. Van Dolah

November 1984

U.S. DEPARTMENT OF COMMERCE
National Oceanic and Atmospheric Administration
National Marine Fisheries Service

NOAA TECHNICAL REPORTS NMFS

The major responsibilities of the National Marine Fisheries Service (NMFS) are to monitor and assess the abundance and geographic distribution of fishery resources, to understand and predict fluctuations in the quantity and distribution of these resources, and to establish levels for optimum use of the resources. NMFS is also charged with the development and implementation of policies for managing national fishing grounds, development and enforcement of domestic fisheries regulations, surveillance of foreign fishing off United States coastal waters, and the development and enforcement of international fishery agreements and policies. NMFS also assists the fishing industry through marketing service and economic analysis programs, and mortgage insurance and vessel construction subsidies. It collects, analyzes, and publishes statistics on various phases of the industry.

The NOAA Technical Report NMFS series was established in 1983 to replace two subcategories of the Technical Reports series: "Special Scientific Report—Fisheries" and "Circular." The series contains the following types of reports: Scientific investigations that document long-term continuing programs of NMFS, intensive scientific reports on studies of restricted scope, papers on applied fishery problems, technical reports of general interest intended to aid conservation and management, reports that review in considerable detail and at a high technical level certain broad areas of research, and technical papers originating in economics studies and from management investigations.

Copies of NOAA Technical Report NMFS are available free in limited numbers to governmental agencies, both Federal and State. They are also available in exchange for other scientific and technical publications in the marine sciences. Individual copies may be obtained from: U.S. Department of Commerce, National Technical Information Service, 5285 Port Royal Road, Springfield, VA 22161.

NOAA Technical Report NMFS 18



A Temporal and Spatial Study of Invertebrate Communities Associated With Hard-Bottom Habitats in the South Atlantic Bight

E. L. Wenner, P. Hinde, D. M. Knott, and
R. F. Van Dolah

November 1984

U.S. DEPARTMENT OF COMMERCE

Malcolm Baldrige, Secretary

National Oceanic and Atmospheric Administration

John V. Byrne, Administrator

National Marine Fisheries Service

William G. Gordon, Assistant Administrator for Fisheries

The National Marine Fisheries Service (NMFS) does not approve, recommend or endorse any propriety product or proprietary material mentioned in this publication. No reference shall be made to NMFS, or to this publication furnished by NMFS, in any advertising or sales promotion which would indicate or imply that NMFS approves, recommends or endorses any proprietary product or proprietary material mentioned herein, or which has as its purpose an intent to cause directly or indirectly the advertised product to be used or purchased because of this NMFS publication.

CONTENTS

Introduction	1
Materials and methods	1
Collection of samples	1
Data analysis	2
Results and discussion	3
Hydrographic measurements and description of study sites	3
Large macrofauna sampled by dredge and trawl	3
Biomass of larger macroinvertebrates	8
Smaller macroinvertebrates sampled by suction and grab	8
Conclusions	13
Acknowledgments	15
Literature cited	15
Appendices	17-104

Figures

1. Location of hard-bottom stations sampled during 1981	2
2. Normal cluster dendrogram showing station groups formed using the Jaccard similarity coefficient and flexible sorting	4
3. Inverse classification hierarchies and nodal diagram showing constancy and fidelity of station-species group coincidence based on seasonal dredge collections	6
4. Inverse classification hierarchies and nodal diagram showing constancy and fidelity of station-species group coincidence based on seasonal trawl collections	8
5. Normal cluster dendrogram of station groups formed using the Bray-Curtis similarity coefficient, log transformation, and flexible sorting	13
6. Inverse classification hierarchies and nodal diagram showing constancy and fidelity of station-species group coincidence based on seasonal suction and grab collections	15

Tables

1. Number of species and percent contribution for each taxonomic group collected by dredge and trawl at each station and season	3
2. Species groups resulting from numerical classification of data from samples collected by dredge during 1981	5
3. Species groups resulting from numerical classification of data from samples collected by trawl during 1981	7
4. Percent contribution to epifaunal biomass of taxonomic groups in dredge and trawl collections	9
5. Average density and percent of total number for 10 numerically dominant invertebrates collected by suction and grab	10
6. The 10 most abundant invertebrate species collected by suction and grab samplers at each station	11
7. Seasonal community structure values for pooled replicate samples of invertebrates at each station	12
8. Species groups resulting from numerical classification of data from samples collected by suction and grab samplers during all seasons, 1981	14

A Temporal and Spatial Study of Invertebrate Communities Associated With Hard-Bottom Habitats in the South Atlantic Bight¹

E. L. WENNER, P. HINDE, D. M. KNOTT, and R. F. VAN DOLAH²

ABSTRACT

Species composition, biomass, density, and diversity of benthic invertebrates from six hard-bottom areas were evaluated. Seasonal collections using a dredge, trawl, and suction and grab samplers yielded 432, 525, and 845 taxa, respectively. Based on collections with the different gear types, species composition of invertebrates was found to change bathymetrically. Inner- and middle-shelf sites were more similar to each other in terms of invertebrate species composition than they were to outer-shelf sites, regardless of season. Sites on the inner and outer shelf were grouped according to latitude; however, results suggest that depth is apparently a more important determinant of invertebrate species composition than either season or latitude. Sponges generally dominated dredge and trawl collections in terms of biomass. Generally, cnidarians, bryozoans, and sponges dominated at sites in terms of number of taxa collected.

The most abundant smaller macrofauna collected in suction and grab samples were polychaetes, amphipods, and mollusks. Densities of the numerically dominant species changed both seasonally and bathymetrically, with very few of these species restricted to a specific bathymetric zone.

The high diversity of invertebrates from hard-bottom sites is attributed to the large number of rare species. No consistent seasonal changes in diversity or number of species were noted for individual stations or depth zones. In addition, H' and its components showed no definite patterns related to depth or latitude. However, more species were collected at middle-shelf sites than at inner- or outer-shelf sites, which may be related to more stable bottom temperature or greater habitat complexity in that area.

INTRODUCTION

Although other studies have provided species lists of invertebrates from hard-bottom areas [see Continental Shelf Associates (1979) for literature review], Wenner et al. (1983) provided the first detailed information on structure of hard-bottom invertebrate communities in the South Atlantic Bight. Results indicated that the invertebrate fauna of these areas was more diverse than the surrounding sand biotope. Species composition was apparently related more to station depth than to season, although species associations did vary between winter and summer. Because sampling during their study was limited to only two seasons, it was not possible to determine seasonal trends in community composition, structure, and biomass which might be occurring over an annual cycle.

This paper presents results from a second-year study of invertebrates associated with hard-bottom habitats and focuses on seasonal as well as bathymetric changes in community parameters. Comparisons with results from the initial study of hard-bottom sites by Wenner et al. (1983) provide additional information on annual variations in the epibenthic invertebrate community.

MATERIALS AND METHODS

Collection of Samples

Two study sites were selected between lat. 31°24' and 32°49'N from each of three bathymetric zones: Inner shelf (IS01, 17-18 m; IS02, 17-22 m), middle shelf (MS02, 23-29 m; MS06, 32-36 m), and outer shelf (OS01, 59-67 m; OS06, 46-69 m) (Fig. 1). Four of the sites (IS01, IS02, MS02, OS01) were previously assessed during the initial study (Wenner et al. 1983). All sites were sampled quarterly during 1981 as follows: Winter (25 February to 17 March), spring (26 April to 28 May), summer (25 July to 13 August), and fall (6 October to 16 November).

Prior to sampling for invertebrates, hydrographic profiles of temperature, salinity, and dissolved oxygen were obtained at each site by Niskin bottles with reversing thermometers.

Qualitative samples of large and encrusting macroinvertebrates were taken quarterly with a heavy duty Cerame-Vivas dredge (Duke University Marine Laboratory 1968) having a mouth opening 90 cm × 37 cm and maximum mesh opening 40 mm × 30 mm. Two replicate dredge tows were attempted at each station with a total of 43 actually completed. Tow length was standardized to approximately 0.1 km using loran-C positioning.

Additional qualitative collections of large macroinvertebrates were obtained as incidental catch by a trawl (12 m headrope, 16 m footrope, 12.8 m vertical height) equipped with steel doors and rubber rollers. Six replicate tows (three day and three night) were attempted quarterly at all sites except station OS06, where high bottom relief prevented use of the trawl. Altogether, 119 trawl tows were completed. Trawl tows were standardized to a distance of about 1 km.

¹Contribution No. 178 from the Marine Resources Research Institute, Charleston, S.C.

²Marine Resources Research Institute, P.O. Box 12559, Charleston, SC 29412.




Figure 1.—Location of hard-bottom stations sampled during 1981.

Invertebrates collected by dredge and trawl were sorted into the following major taxonomic groups for wet-weight estimates: Porifera, Hydrozoa, Scleractinia/Octocorallia, Mollusca, Decapoda, Echinodermata, Ascidiacea, and other miscellaneous invertebrates. Encrusting organisms which could not easily be removed from rocks were not weighed. Representative specimens of each species were preserved in 10% buffered seawater Formalin and transferred to 70% isopropanol prior to identification.

Quantitative samples of smaller benthic invertebrates not adequately sampled by dredge and trawl were obtained at inner- and middle-shelf sites by scuba divers using an airlift suction sampling device similar to that described by Chess (1979). Divers obtained five replicate samples at each station from a suitable area chosen to avoid large patches of sand commonly found at the sites. A disc with five equally spaced radial marks was dropped to the bottom and a 3 m line, fastened to the center of the disc, was then used to place five open-ended quadrat boxes (0.1 m², 10 cm walls) equidistantly around the disc. Exact positioning of the quadrat boxes was accomplished by randomly selecting one of nine possible quadrat areas from a larger grid frame attached to the 3 m line. Fauna within the quadrat was sampled by scraping the area while simultaneously sucking with the airlift suction device to a depth of 10 cm or until rocky substrate was encountered. Suction was applied until all sediment within a quadrat was removed to the designated depth. All suction samples were collected in 1.0 mm mesh bags and brought to the surface for preservation. On deck,

each sample was narcotized with 10% magnesium chloride and preserved in a 10% seawater-Formalin solution.

At each outer-shelf station, where water depth precluded the use of a diver-operated suction device, five quantitative 0.1 m² samples were attempted with a Smith-McIntyre grab. Only 33 samples were collected due to ineffective sampling of the grab on hard substrates. After retrieval, each sample was placed on a 1.0 mm sieve and washed to remove the finer sediment prior to preservation in 10% buffered seawater Formalin. Macroinvertebrates collected by suction and grab samplers were identified and counted, except for encrusting fauna which were not easily quantified by counting (e.g., hydroids and colonial corals).

Data Analysis

Community diversity was estimated by the Shannon index (H') (Pielou 1975) in conjunction with species richness (SR) (Margalef 1958), and evenness (J') (Pielou 1975). These measures were computed on data from pooled replicates taken with suction or grab samplers at each station, since pooling the replicates provided a larger sample size and a more representative estimate of community diversity at a site. The degree of dominance at a station was quantified with the dominance index (DI) (McNaughton 1967). For qualitative collections by dredge and trawl, diversity was evaluated by comparing the number of species (s) among stations. The Kruskal-Wallis one-way analysis by ranks (Siegel 1956) was used to determine whether s differed significantly between stations or seasons.

Cluster analysis was used to determine patterns of similarity among stations (pooled replicated collections) and among species using qualitative binary data (i.e., species presence or absence) for dredge and trawl samples, and quantitative abundance data for suction and grab samples. Due to the large number of species in collections made by the various sampling devices, it was necessary to reduce data sets which contained > 200 species prior to cluster analysis in order to remain within the computational limits of available computer programs. Data sets were reduced by both elimination of species which were infrequently collected and elimination of taxa having undetermined or questionable identity. The elimination of these taxa from our data sets was justifiable because "rare" species usually do not have definable distribution patterns and can confuse interpretation of cluster analysis. Data sets subjected to cluster analysis were reduced to include only those species represented in 5 or more dredge collections, 9 or more trawl collections, and 12 or more suction and grab collections.

Species and collections were classified using flexible sorting (Lance and Williams 1967) with a cluster intensity coefficient (β) of -0.25 . The Jaccard similarity coefficient (Clifford and Stephenson 1975) was used with presence-absence data collected by dredge and trawl. Species abundance data collected by suction and grab were transformed using $\log_{10}(x + 1)$ and clustered using the Bray-Curtis measure of similarity (Bray and Curtis 1957). Since attribute values covered a wide range, a strong transformation was used to reduce the sensitivity of the Bray-Curtis measure to larger values (Clifford and Stephenson 1975).

Subsequent to cluster analysis, species and station groups were chosen using a variable stopping rule (Boesch 1977). Then nodal analyses (Williams and Lambert 1961; Lambert and Williams 1962) were employed to describe pooled collections at a station in terms of their characteristic species and to describe species groups resulting from inverse cluster analysis in terms of their patterns of occurrence at stations (Boesch 1977). Coincidence of station and

species groups was expressed by graded constancy and fidelity values, which were used to determine those species that were most consistently collected and most restricted to pooled collections at a station. The mathematical definitions for constancy and fidelity are found in Boesch (1977).

RESULTS AND DISCUSSION

Hydrographic Measurements and Description of Study Sites

Hydrographic parameters at the hard-bottom sites were similar to those reported by Wenner et al. (1983). Bottom temperatures on the inner shelf were more extreme (11.7°-27.3°C) than those on the outer shelf (17.0°-23.9°C). Temperatures at middle-shelf sites were similar to those encountered at outer-shelf sites (16.0°-23.4°C). Bottom salinity and dissolved oxygen were generally high at the sites during all seasons, ranging from 34.3 to 36.6‰ and 3.9-6.4 ml/l, respectively.

Television transect videotapes showed that the bottom at station IS01 was smooth, flat, and overlaid by a layer of sand. Sessile invertebrates at IS01 were numerous and uniformly covered the broad expanse of sand-covered hardpan. Station IS02 had a more irregular bottom than IS01 with numerous rock ledges and discontinuous outcroppings up to 1 m in height. These outcroppings were distributed throughout areas of sand and were covered with a dense assemblage of epifauna, particularly adjacent to the ledges.

Middle-shelf sites were characterized as having moderate relief ledges, although there were also areas of low rocky relief interspersed with sand. At MS01, several ledges up to 1 m in height were observed, whereas ledges with a vertical drop of nearly 2 m were observed at MS06. Epifaunal density was usually high on emergent rock of moderate to high relief at MS06.

Outer-shelf stations, which were located in the vicinity of the shelf break, were distinctly different from one another with

respect to their bottom topography. Station OS01 was a low relief hardground with only occasional exposed rock of little relief and relatively sparse epifaunal density. In contrast, OS06 was a typical shelf edge reef, with irregular bottom topography characterized by ledges and extensive rock outcropping of moderate to high relief which were often heavily encrusted with invertebrates.

Large Macrofauna Sampled by Dredge and Trawl

Seasonal collections of epifauna from hard-bottom sites sampled by dredge and trawl yielded 432 and 525 taxa, respectively. A list of the identified taxa, arranged phylogenetically for each station and sampling gear, is found in Appendices 1 and 2. The Bryozoa (91 taxa), Porifera (89 taxa), and Cnidaria (70 taxa), dominated dredge collections from all seasons in terms of numbers of taxa. Porifera (111 taxa) were also well represented in trawl collections, along with other taxa such as decapod crustaceans (86 taxa), Bryozoa (85 taxa), and Mollusca (85 taxa). Both the present and previous study (Wenner et al. 1983) of hard-bottom communities have shown that Porifera, Bryozoa, and Cnidaria are the most diverse taxonomic groups encountered.

The only invertebrate groups whose percent contribution of species in dredge and trawl collections changed noticeably across the continental shelf were the Porifera, Cnidaria, and Bryozoa (Table 1). In terms of number of taxa, sponges were most important at sites on the inner shelf. Their richness diminished at middle- and outer-shelf sites where cnidarian and bryozoan taxa dominated. No seasonal trends in percent composition among taxa were noted.

The number of invertebrate taxa (*s*) collected at each station by dredge and trawl did not change significantly between seasons (Kruskal-Wallis test, $P > 0.05$). These results were consistent with those reported by Wenner et al. (1983) who found no significant difference in *s* between winter and summer sampling periods. However, in the present study we did find a significant difference

Table 1.—Number of species and percent contribution for each taxonomic group collected by dredge (D) and trawl (T) at each station and season.

Station	Season	Number samples		Porifera		Cnidaria		Mollusca		Cirripedia		Decapoda		Echino- dermata		Bryozoa		Tunicata		Total number
		D	T	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	
IS01	Winter	2	6	45	28.1	26	16.2	16	10.0	4	2.5	15	9.4	8	5.0	29	18.1	17	10.6	160
	Spring	2	6	25	17.9	31	22.1	12	8.6	5	3.6	15	10.7	9	6.4	27	19.3	16	11.4	140
	Summer	2	6	33	20.8	36	22.6	17	10.7	4	2.5	12	7.5	10	6.2	29	18.2	18	11.3	159
	Fall	2	6	21	15.7	42	31.3	10	7.5	5	3.7	12	8.9	6	4.5	25	18.7	13	9.7	134
IS02	Winter	1	5	49	24.4	37	18.4	20	9.9	4	2.0	25	12.4	7	3.5	43	21.4	16	7.9	201
	Spring	2	6	49	25.5	25	13.0	23	11.9	5	2.6	22	11.5	13	6.8	37	19.3	18	9.4	192
	Summer	2	6	40	24.2	37	22.4	15	9.1	4	2.4	20	12.1	7	4.2	28	16.9	14	8.5	165
	Fall	2	6	51	27.4	37	19.8	20	10.8	4	2.1	15	8.1	8	4.3	35	18.8	16	8.6	186
MS02	Winter	1	6	29	17.7	38	23.2	16	9.8	4	2.4	15	9.1	15	9.1	32	19.5	15	9.1	164
	Spring	2	6	30	14.2	37	17.5	28	13.3	5	2.4	29	13.7	10	4.7	50	23.6	22	10.4	211
	Summer	2	6	15	8.8	43	25.1	14	8.2	4	2.3	22	12.9	7	4.1	50	29.2	16	9.4	171
	Fall	2	6	24	14.2	44	26.0	13	7.7	3	1.8	15	8.9	9	5.3	43	25.4	18	10.6	169
MS06	Winter	2	6	20	12.4	43	26.7	15	9.3	3	1.9	17	10.6	6	3.7	49	30.4	8	4.9	161
	Spring	2	6	38	16.2	49	20.8	34	14.5	3	1.3	17	7.2	16	6.8	63	26.8	15	6.4	235
	Summer	2	6	36	20.3	45	25.4	14	7.9	4	2.3	13	7.3	13	7.3	43	24.3	9	5.1	177
	Fall	2	6	38	19.7	46	23.8	16	8.3	3	1.6	18	9.3	8	4.1	54	27.9	10	5.2	193
OS01	Winter	2	6	26	13.7	54	28.4	17	8.9	4	2.1	25	13.2	9	4.7	44	23.2	11	5.8	190
	Spring	2	6	29	15.3	48	25.4	14	7.4	6	3.2	35	18.5	11	5.8	38	20.1	8	4.2	189
	Summer	2	6	28	14.0	53	26.5	16	8.0	5	2.5	39	19.5	13	6.5	38	19.0	8	4.0	200
	Fall	2	6	36	18.7	50	25.9	23	11.9	6	3.1	25	13.0	9	4.7	32	16.6	12	6.2	193
OS06	Winter	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	Spring	2	—	—	—	40	38.5	4	3.8	3	2.9	2	1.9	1	0.9	54	51.9	—	—	104
	Summer	1	—	3	3.6	22	26.5	4	4.8	—	—	6	7.2	8	9.6	34	40.9	6	7.2	83
	Fall	2	—	14	11.2	39	31.2	7	5.6	1	0.8	11	8.8	8	6.4	42	33.6	3	2.4	125

in s between stations sampled (Kruskal-Wallis test, $P < 0.01$) by trawl. Although the fewest species were collected at IS01, this is not necessarily indicative of generally lower richness at inner shelf stations, nor do these results imply that s is influenced solely by depth or latitude, since little difference was noted between numbers of species collected at IS02, middle-shelf sites, and OS01. Increases (or decreases) in s are probably also influenced by localized factors at each site such as topographic complexity and habitat heterogeneity. The extensive hard-bottom area of IS01 had little vertical relief compared with IS02, which was characterized by numerous rock ledges supporting dense epifaunal populations. These rock ledges provide complex microhabitats which allow many epifaunal species to coexist (Schoener 1974; Menge and Sutherland 1976). In addition, many sessile species settle in greater abundance on structurally complex substrata because there is more surface area to which they can securely attach (Seed 1976; Russ 1980; Dean 1981).

Normal cluster analysis indicated that pooled replicate dredge and trawl collections were classified into three station groups corresponding to inner-shelf sites (group 1), middle-shelf sites (group 2), and outer-shelf sites (group 3) (Fig. 2). Stations were not grouped in a manner which would indicate that species composition changed seasonally in a consistent pattern. Even within each site group, no grouping of stations by season was evident. This

suggests that the qualitative composition of the epifaunal community sampled by dredge and trawl was generally similar throughout the year. In addition, there was no indication that stations were grouped in accordance with their latitudinal location on the continental shelf. Comparison with results of normal cluster analysis from the initial study indicates that depth is apparently a more important determinant of faunal composition than season or latitude at hard-bottom sites off South Carolina and Georgia. These results suggest that the epifaunal community of hard-bottom areas are persistent throughout the year. Osman (1977) suggested that changes in the composition of epifaunal invertebrates occur slowly. Growth and reproduction rate of epifauna are reduced during the winter with the result that few new species are added, but several are lost.

Inverse cluster analysis of the 199 most frequently occurring species from seasonal dredge collections formed 10 groups (Table 2). Nodal analysis indicated that the three major branches of the inverse cluster dendrogram represent species assemblages found primarily at inner-shelf sites, those found across the shelf, and those found primarily at middle- and outer-shelf sites (Fig. 3).

Species which were most consistently collected at inner-shelf stations partially comprise groups A, C, and D. However, these assemblages were not restricted to sites on the inner shelf, as indicated by moderate fidelity values. Species in these groups which were collected only at inner-shelf sites included the sponges *Homaxinella waltonsmithi* and *Tedania ignis*; the echinoderms *Lytechinus variegatus*, *Encope michelini*, and *Luidia alternata*; and the mollusk *Pisania tineta*. Except for the sponges *H. waltonsmithi* and *T. ignis*, these species are found on a wide variety of bottom types and are not limited to hard substrates. Although the cluster hierarchy indicated that group B and C species were most similar in their occurrence at hard-bottom sites, the nodal diagram illustrates differences in the distribution of species in these two groups. Species in group B were most constant at station OS01; whereas species in group C were most constant at station IS02.

Species in groups E and F were consistently collected at most sites sampled. Group E species were highly constant at all sites and included the most ubiquitous species collected. Many of the species in this group were dominant epifaunal organisms in terms of their frequency of occurrence at hard-bottom sites across the shelf. However, because of their ubiquity they were not faithful to any site group. Species in group F were also fairly ubiquitous but were more commonly encountered at inner- and middle-shelf sites.

Species groups G-J were most consistently collected at middle- and outer-shelf sites. Group G species displayed high constancy at stations MS02 and MS06; however, the only species in this group which was collected exclusively at middle-shelf hard-bottom stations was the bryozoan *Schizoporella errata*. Group H species also were common at station MS06 but were most consistently encountered at station OS06. Species groups I and J displayed highest constancy and fidelity at outer shelf sites; however, only species in group J were restricted to these locations, as indicated by their very high fidelity values. Species in groups I and J which were collected exclusively at outer-shelf sites included the cnidarians *Balanophyllia floridana*, *Diodogorgia* sp., *Ellisella* sp., and *Srylatula* sp.; the decapod crustaceans *Galathea rostrata*, *Dardanus insignis*, and *Parthenope fraterculus*; the bryozoan *Membranipora arborescens*; and the echinoderm *Astroporpa annulata*.

Comparisons of species assemblages identified by cluster and nodal analyses between the present and the previous study (Wenner et al. 1983) have shown that several species are most frequently encountered in one depth zone. Those which typified




Figure 2.—Normal cluster dendrogram showing station groups formed using the Jaccard similarity coefficient and flexible sorting. Data from replicate dredge and trawl collections were pooled for each station prior to cluster analysis.

Table 2.—Species groups resulting from numerical classification of data from samples collected by dredge during 1981. (Ar = Arthropoda; Bry = Bryozoa; Ch = Chordata; Cn = Cnidaria; Ech = Echinodermata; Mo = Mollusca; Po = Porifera.)

Group A	<i>Floridina antiqua</i> (Bry) <i>Encope michelini</i> (Ech) <i>Pisania tincta</i> (Mo) <i>Hemectyon pearsei</i> (Po) <i>Dromidia antillensis</i> (Ar) <i>Oculina</i> sp. (Cn) <i>Halichondria bowerbanki</i> (Po) <i>Ciocalapata gibbsi</i> (Po) <i>Ircinia campana</i> (Po) <i>Ircinia felix</i> (Po) <i>Arbacia punctulata</i> (Ech) <i>Higginsia strigilata</i> (Po) <i>Ircinia ramosa</i> (Po) <i>Geodia gibberosa</i> (Po)	<i>Antropora tincta</i> (Bry) <i>Turbicellepora dichotoma</i> (Bry) <i>Clytia fragilis</i> (Cn) <i>Balanus venustus</i> (Ar) <i>Cribrilaria radiata</i> (Bry) <i>Microporella umbracula</i> (Bry) <i>Aetea anguina</i> (Bry) <i>Amathia distans</i> (Bry) <i>Celleporaria albirostris</i> (Bry) <i>Parasmitina spathulata</i> (Bry) <i>Telesto sanguinea</i> (Cn) <i>Cleidochasma porcellanum</i> (Bry) <i>Halecium tenellum</i> (Cn) <i>Sertularia plumulifera</i> (Cn) <i>Halecium dysymmetrum</i> (Cn) <i>Pseudomedeaeus agassizii</i> (Ar)	Group H <i>Parellisina latirostris</i> (Bry) <i>Nellia tenella</i> (Bry) <i>Nemertesia simplex</i> (Cn) <i>Caberia boryi</i> (Bry) <i>Microporella marsupiatia</i> (Bry) <i>Proboscina floridana</i> (Bry) <i>Gymnangium sinuosum</i> (Cn) <i>Cribrilaria floridana</i> (Bry) <i>Lichenopora radiata</i> (Bry) <i>Exochella longirostris</i> (Bry) <i>Clypeaster subdepressus</i> (Ech) <i>Floridina parvicella</i> (Bry) <i>Turritopsis nutricula</i> (Cn) <i>Hincksella cylindrica</i> (Cn)
<i>Lissodendoryx sigmata</i> (Po) <i>Clavelina gigantea</i> (Ch) <i>Scypha barbadensis</i> (Po) <i>Clathrina cpricaea</i> (Po) <i>Pilumnus dasypodus</i> (Ar) <i>Celleporina hassalli</i> (Bry) <i>Membranipora tenuis</i> (Bry) <i>Synalpheus minus</i> (Ar) <i>Distaplia bermudensis</i> (Ch) <i>Ocnus pygmaeus</i> (Ech) <i>Homaxinella waltonsmithi</i> (Po) <i>Lytechinus variegatus</i> (Ech) <i>Crepidula aculeata</i> (Mo) <i>Stryela partita</i> (Ch) <i>Celleporaria mordax</i> (Bry) <i>Bimeria humilis</i> (Cn) <i>Symplegma viride</i> (Ch) <i>Trididemnum savignii</i> (Ch) <i>Homaxinella</i> sp. A (Po) <i>Diplosoma macdonaldi</i> (Ch) <i>Eudistoma carolinense</i> (Ch) <i>Sertularia distans</i> (Cn)	Group D <i>Homaxinella rudis</i> (Po) <i>Cinachyra alloclada</i> (Po) <i>Spirastrella coccinea</i> (Po) <i>Cinachyra rhizophyta</i> (Po) <i>Luidia alternata</i> (Ech) <i>Tedania ignis</i> (Po) <i>Phakellia lobata</i> (Po) <i>Macrocoeloma camptocercum</i> (Ar) <i>Pseudochama radians</i> (Mo) <i>Haliclona</i> sp. C (Po) <i>Nolella stipata</i> (Bry) <i>Buskia</i> sp. (Bry) <i>Xytopseus griseus</i> (Po) <i>Sigmatocia caerulea</i> (Po)	Group F <i>Haliclona oculata</i> (Po) <i>Leptogorgia virgulata</i> (Cn) <i>Vermicularia knorrii</i> (Mo) <i>Hippaliosina rostrigera</i> (Bry) <i>Reptadeonella hastingsae</i> (Bry) <i>Conopea merrilli</i> (Ar) <i>Pilumnus sayi</i> (Ar) <i>Celleporaria magnifica</i> (Bry) <i>Styela plicata</i> (Ch) <i>Pteria colymbus</i> (Mo) <i>Molgula occidentalis</i> (Ch) <i>Modiolus americanus</i> (Mo) <i>Arca imbricata</i> (Mo) <i>Lophogorgia hebes</i> (Cn) <i>Mithrax pleuracanthus</i> (Ar) <i>Eudendrium carneum</i> (Cn)	Group I <i>Bougainvillia</i> sp. (Cn) <i>Sertularella areyi</i> (Cn) <i>Antenella secundaria</i> (Cn) <i>Eudendrium tenellum</i> (Cn) <i>Dynamena dalmasi</i> (Cn) <i>Balanus calidus</i> (Ar) <i>Aplousina gigantea</i> (Bry) <i>Filillum serratum</i> (Cn) <i>Stylopoma informata</i> (Bry) <i>Smittipora leviseni</i> (Bry) <i>Campanularia hincksii</i> (Cn) <i>Thesea</i> sp. (Cn) <i>Hippoporina contracta</i> (Bry) <i>Scandia mutabilis</i> (Cn) <i>Hebella venusta</i> (Cn) <i>Hebella scandens</i> (Cn) <i>Mithrax acuticornis</i> (Ar) <i>Galathea rostrata</i> (Ar) <i>Amathia convoluta</i> (Bry) <i>Alderina smitti</i> (Bry) <i>Ctenostomata</i> (Bry) <i>Scrupocellaria regularis</i> (Bry) <i>Aglaophenia elongata</i> (Cn)
Group B <i>Microcionia prolifera</i> (Po) <i>Teichaxinella corrugata</i> (Po) <i>Pseudaxinella lunaecharta</i> (Po) <i>Arca zebra</i> (Mo) <i>Axinella bookhouti</i> (Po) <i>Pandaros acanthifolium</i> (Po) <i>Diodora cayenensis</i> (Mo) <i>Murex florifer dilectus</i> (Mo) <i>Stenocionops furcata coelata</i> (Ar) <i>Leptogorgia</i> sp. (Cn) <i>Antipatharia</i> (Cn) <i>Membraniporella aragoi</i> (Bry) <i>Eucidaris tribuloides</i> (Ech) <i>Pachycheles rugimanus</i> (Ar)	Group E <i>Didemnum candidum</i> (Ch) <i>Chama congregata</i> (Mo) <i>Monostaechas quadridens</i> (Cn) <i>Microporella ciliata</i> (Bry) <i>Obelia dichotoma</i> (Cn) <i>Schizoporella cornuta</i> (Bry) <i>Trypostega venusta</i> (Bry) <i>Titanideum frauenfeldii</i> (Cn) <i>Balanus trigonus</i> (Ar) <i>Halecium</i> sp. (Cn) <i>Telesto fruticulosa</i> (Cn) <i>Ophiolithrix angulata</i> (Ech) <i>Dynamena cornicina</i> (Cn) <i>Crisia</i> sp. (Bry) <i>Sertularia marginata</i> (Cn) <i>Bugula rylandi</i> (Bry) <i>Aglaophenia latecarinata</i> (Cn) <i>Salacia desmoides</i> (Cn) <i>Clytia cylindrica</i> (Cn) <i>Aglaophenia trifida</i> (Cn) <i>Sertularella conica</i> (Cn) <i>Sertularella</i> sp. A (Cn) <i>Dynamena quadridentata</i> (Cn)	Group G <i>Synthecium tubithecra</i> (Cn) <i>Petraliella bisinuata</i> (Bry) <i>Bugula fulva</i> (Bry) <i>Schizoporella floridana</i> (Bry) <i>Aevertillia setigera</i> (Bry) <i>Thyroscyphus marginatus</i> (Cn) <i>Chaperia</i> sp. (Bry) <i>Diaperoecia floridana</i> (Bry) <i>Parellisina curvirostris</i> (Bry) <i>Cupuladria doma</i> (Bry) <i>Cupuladria canariensis</i> (Bry) <i>Crepidacantha poissoni teres</i> (Bry) <i>Megalobranchium soriatum</i> (Ar) <i>Smittina smittiella</i> (Bry) <i>Schizoporella errata</i> (Bry) <i>Leuconia aspera</i> (Po) <i>Lyonsia beana</i> (Mo) <i>Alcyonidium</i> sp. (Bry) <i>Podochela riisei</i> (Ar) <i>Gonodactylus bredini</i> (Ar)	Group J <i>Dardanus insignis</i> (Ar) <i>Membranipora arborescens</i> (Bry) <i>Halopteris clarkei</i> (Cn) <i>Cycloperella rubra</i> (Bry) <i>Astroporpa annulata</i> (Ech) <i>Stylatula</i> sp. (Cn) <i>Balanophyllia floridana</i> (Cn) <i>Ellisella</i> sp. (Cn) <i>Diodogorgia</i> sp. (Cn) <i>Parthenope fraterculus</i> (Ar)
Group C <i>Spheciospongia vesparium</i> (Po) <i>Lissodendoryx isodictyalis</i> (Po) <i>Leodia sexiesperforata</i> (Ech) <i>Cliona caribbaea</i> (Po) <i>Cinachyra kuekenthali</i> (Po) <i>Aplysina fistularis</i> (Po) <i>Aplousina tuberosa</i> (Bry) <i>Amphimedon compressa</i> (Po) <i>Thalysias juniperina</i> (Po) <i>Ircinia strobilina</i> (Po) <i>Synalpheus townsendi</i> (Ar) <i>Hiatella arctica</i> (Mo) <i>Amathia alternata</i> (Bry)			




Figure 3.—Inverse classification hierarchies and nodal diagram showing constancy and fidelity of station-species group coincidence based on seasonal dredge collections.

inner-shelf hard-bottom sites included the sponges *Homaxinella waltonsmithi*, *Spherospongia vesparium*, *Cliona caribbaea*, and *Halichondria bowerbanki*; the echinoderms *Encope michelini*, *Lytechinus variegatus*, *Ocnus pygmaeus*, and *Arbacia punctulata*; the bryozoan *Membranipora tenuis*; and the decapod crustacean *Synalpheus minus*. Middle-shelf hard-bottom sites were characterized by the bryozoans *Chaperia* sp., *Aevertillia setigera*, *Schizoporella floridana*, *Petraliella bisinuata*, and *Parellisina curvirostris*; and the hydroids *Thyroscyphus marginatus* and *Synthecium tubithecra*. Species which were consistently collected at outer-shelf sites during both years of sampling by dredge included the bryozoans *Smittipora levinseni*, *Stylopoma informata*, *Cribrilaria floridana*, and *Cycloperiella rubra*; the hydroids *Dynamensia dalmasi* and *Filellum serratum*; the echinoderm *Astroporpa annulata*; and the decapod crustacean *Mithrax acuticornis*. Few species assemblages identified by cluster analysis of dredge samples were restricted to any one depth zone or hard-bottom site. This suggests that few species assemblages, other than those displaying high fidelity values in nodal diagrams, could be considered characteristic of a particular depth zone in the South Atlantic Bight.

Inverse cluster analysis of 184 species from seasonal trawl collections produced seven groups. Species and site group coincidences were interpreted in terms of nodal constancy and fidelity (Table 3, Fig. 4). Group A contained species which were characteristic of inner-shelf stations and station MS06. Species in group B also displayed high constancy for inner-shelf stations; however, no species in either group were restricted to specific hard-bottom sites. Only the tunicate Ascidiacea A was collected exclusively at inner-shelf sites.

Groups C, D, and E were composed of species which were fairly ubiquitous at hard-bottom sites across the shelf. Species in group

C were highly constant at inner- and middle-shelf stations, whereas those in group D were consistently collected at every live-bottom site sampled. This group contained many of the most frequently occurring epifaunal species collected during the study. Group E contained species which displayed high constancy at stations IS02, MS02, and MS06. Species in group F were highly constant at station MS06, but were not restricted to this station. Those species in group G were most frequently collected at both middle-shelf sites and at the outer-shelf station. As indicated by the inverse dendrogram, species in group H, which were highly constant at station OS01, were least similar to other groups in terms of their distribution. Several species, such as the bryozoans *Aplousina gigantea* and *Cycloperiella rubra*, the octocoral *Ellisella* sp., and the decapod crustacean *Mesopenaeus tropicalis*, were collected only at station OS01.

No species assemblage identified from our study was restricted to specific hard-bottom sites. However, comparison of species assemblages from cluster analysis of winter and summer from the first-year study (Wenner et al. 1983) with seasonal data from the present study show that the following species formed a consistently recurring assemblage on the inner shelf: The bryozoans *Scrupocellaria regularis* and *Hippoporina contracta*; the echinoderm *Ocnus pygmaeus*; and the crustaceans *Synalpheus longicarpus*, *S. minus*, *Pilumnus pannosus*, *Megalobrachium soriatum*, *Trachypenaeus constrictus*, and *Conopea galeata*. Other species, such as the cnidarian *Sertularella areyi*; the echinoderm *Astroporpa annulata*; the sponges *Xytopses griseus* and *Sigmadocia caerulea*; and the decapod crustaceans *Mesopenaeus tropicalis*, *Sicyonia brevirostris*, and *Solenocera atlantidis*, were consistently collected during both years at outer-shelf sites. These findings suggest that there is no sharp bathymetric zonation of epifaunal species groups in hard-bottom habitats of the South

Table 3.—Species groups resulting from numerical classification of data from samples collected by trawl during 1981. (Ar = Arthropoda; Bry = Bryozoa; Ch = Chordata; Cn = Cnidaria; Ech = Echinodermata; Mo = Mollusca; Po = Porifera.)

Group A	Group C	<i>Titanideum frauenfeldii</i> (Cn) <i>Obelia dichotoma</i> (Cn) <i>Telesto fruticulosa</i> (Cn) <i>Didemnum candidum</i> (Ch) <i>Scandia mutabilis</i> (Cn) <i>Clytia cylindrica</i> (Cn) <i>Dromidia antillensis</i> (Ar)	<i>Parellisia curvirostris</i> (Bry) <i>Polyandrocarpa floridana</i> (Ch) <i>Thyroscyphus marginatus</i> (Cn) <i>Schizoporella floridana</i> (Bry) <i>Hebella scandens</i> (Cn) <i>Syntheconium tubithecra</i> (Cn) <i>Aglaophenia allmani</i> (Cn) <i>Metapenaeopsis goodei</i> (Ar) <i>Campanularia hincksi</i> (Cn) <i>Astrophyton munitatum</i> (Ech) <i>Hebella venusta</i> (Cn) <i>Petraliella bisinuata</i> (Bry) <i>Halecium tenellum</i> (Cn) <i>Celleporaria albirostris</i> (Bry) <i>Plumularia setacea</i> (Cn) <i>Hinckella cylindrica</i> (Cn) <i>Lissodendoryx sigmata</i> (Po) <i>Aglaophenia lateocarinata</i> (Cn) <i>Bugula rylandi</i> (Bry) <i>Chaperia</i> sp. (Bry) <i>Bugula fulva</i> (Bry) <i>Aeverillia setigera</i> (Bry) <i>Celleporina hassalli</i> (Bry) <i>Muricea pendula</i> (Cn) <i>Bugula microecia</i> (Bry)
<i>Anthosigmella varians</i> (Po) <i>Asterias forbesii</i> (Ech) <i>Cinachya alloclada</i> (Po) <i>Pandarus acanthifolium</i> (Po) <i>Leuconia aspera</i> (Po) <i>Crassimarginatella leucocypha</i> (Bry) <i>Trypotegea venusta</i> (Bry) <i>Microporella umbracula</i> (Bry) <i>Homaxinella waltonsmithi</i> (Po) <i>Higginsia strigilata</i> (Po) <i>Lytechinus variegatus</i> (Ech) <i>Ciocalapata gibbsi</i> (Po) <i>Cribrilaria radiata</i> (Bry) <i>Amphimedon compressa</i> (Po) <i>Reptadeonella hastingsae</i> (Bry) <i>Microcionia prolifera</i> (Po) <i>Myriastira fibrosa</i> (Po)	<i>Syrela plicata</i> (Ch) <i>Mithrax pleuracanthus</i> (Ar) <i>Molgula occidentalis</i> (Ch) <i>Pilumnus sayi</i> (Ar) <i>Leptogorgia virgulata</i> (Cn) <i>Spheciospongia vesparium</i> (Po) <i>Sertularella conica</i> (Cn) <i>Crisia</i> sp. (Bry) <i>Synalpheus townsendi</i> (Ar) <i>Distaplia bermudensis</i> (Ch) <i>Diodora cayenensis</i> (Mo) <i>Diplosoma macdonaldi</i> (Ch) <i>Lophogorgia hebes</i> (Cn) <i>Arbacia punctulata</i> (Ech) <i>Antropora tinca</i> (Bry) <i>Haliclona oculata</i> (Po) <i>Sertularia distans</i> (Cn) <i>Cinachya kuekenethali</i> (Po) <i>Arca imbricata</i> (Mo)	Group E	Group H
Group B	Group D	Group F	Group I
<i>Scrupocellaria regularis</i> (Bry) <i>Gymnangium sinuosum</i> (Cn) <i>Hemectyon pearsei</i> (Po) <i>Spirastrella coccinea</i> (Po) <i>Ascidacea A</i> (Ch) <i>Cliona caribbaea</i> (Po) <i>Clathrina coriacea</i> (Po) <i>Trachypenaeus constrictus</i> (Ar) <i>Nolella stipata</i> (Bry) <i>Synalpheus longicarpus</i> (Ar) <i>Synalpheus minus</i> (Ar) <i>Pilumnus dasypodus</i> (Ar) <i>Celleporaria mordax</i> (Bry) <i>Syrela partita</i> (Ch) <i>Pilumnus pannosus</i> (Ar) <i>Conopea galeata</i> (Ar) <i>Ocnus pygmaeus</i> (Ech) <i>Podochela riisei</i> (Ar) <i>Macrocoeloma trispinosum</i> (Ar) <i>Botryllus planus</i> (Ch) <i>Megalobrachium soriatum</i> (Ar) <i>Scypha barbadensis</i> (Po) <i>Sundanella sibogae</i> (Bry) <i>Hippoporina contracta</i> (Bry) <i>Symplegma viride</i> (Ch) <i>Botrylloides nigrum</i> (Ch) <i>Membranipora tenuis</i> (Bry)	<i>Salacia desmoides</i> (Cn) <i>Amathia distans</i> (Bry) <i>Eudendrium</i> sp. (Cn) <i>Parasmittina spathulata</i> (Bry) <i>Ophiolithrix angulata</i> (Ech) <i>Thalysia juniperina</i> (Po) <i>Halecium dysymmetrum</i> (Cn) <i>Hippaliosina rostrigera</i> (Bry) <i>Sertularia plumulifera</i> (Cn) <i>Telesto sanguinea</i> (Cn) <i>Balanus venustus</i> (Ar) <i>Dynamena cornicina</i> (Cn) <i>Dynamena quadridentata</i> (Cn) <i>Eudendrium carneum</i> (Cn) <i>Conopea merrilli</i> (Ar) <i>Aetea anguina</i> (Bry) <i>Turbicellepora dichotoma</i> (Bry) <i>Pteria colymbus</i> (Mo) <i>Balanus trigonus</i> (Ar) <i>Schizoporella cornuta</i> (Bry) <i>Microporella ciliata</i> (Bry) <i>Halecium</i> sp. (Cn) <i>Sertularia marginata</i> (Cn) <i>Monostaeas quadridens</i> (Cn) <i>Aglaophenia trifida</i> (Cn)	<i>Steganoporella magnilabris</i> (Bry) <i>Neofibularia noliangere</i> (Po) <i>Smittina smittiella</i> (Bry) <i>Strongylophora</i> sp. B (Po) <i>Caberia boryi</i> (Bry) <i>Callyspongia fallax</i> (Po) <i>Pilumnus floridanus</i> (Ar) <i>Geodia gibberosa</i> (Po) <i>Alpheus formosus</i> (Ar) <i>Schizoporella errata</i> (Bry) <i>Clavelina gigantea</i> (Ch) <i>Crepidula aculeata</i> (Mo)	<i>Ellisella</i> sp. (Cn) <i>Aplousina gigantea</i> (Bry) <i>Mesopenaeus tropicalis</i> (Ar) <i>Sertularella areyi</i> (Cn) <i>Sylatula</i> sp. (Cn) <i>Cladocarpus flexilis</i> (Cn) <i>Diodogorgia</i> sp. (Cn) <i>Halopteris clarkei</i> (Cn) <i>Cycloperella rubra</i> (Bry) <i>Sylopoma informata</i> (Bry) <i>Antipatharian</i> (Cn) <i>Sicyonia brevirostris</i> (Ar) <i>Alcyonidium</i> sp. (Bry) <i>Haliclona</i> sp. C (Po) <i>Leptogorgia</i> sp. (Cn) <i>Turritopsis nutricula</i> (Cn) <i>Thesea</i> sp. (Cn) <i>Astropora annulata</i> (Ech) <i>Solenocera atlantidis</i> (Ar) <i>Nellia tenella</i> (Bry) <i>Xylopsues griseus</i> (Po) <i>Teichaxinella corrugata</i> (Po) <i>Sigmatocia caerulea</i> (Po) <i>Axinella bookhouti</i> (Po) <i>Halichondria bowerbanki</i> (Po) <i>Balanus calidus</i> (Ar)
		Group G	
		<i>Eudendrium tenellum</i> (Cn) <i>Bimeria humilis</i> (Cn) <i>Bougainvillia</i> sp. (Cn) <i>Hiattella arctica</i> (Mo) <i>Sertularella</i> sp. A (Cn) <i>Clytia fragilis</i> (Cn) <i>Celleporaria magnifica</i> (Bry) <i>Ctenostomata</i> (Bry) <i>Amathia alternata</i> (Bry) <i>Stenocionops furcata coelata</i> (Ar) <i>Aglaophenia elongata</i> (Cn) <i>Filellum serratum</i> (Cn) <i>Diaperoecia floridana</i> (Bry) <i>Pseudomedusa agassizii</i> (Ar) <i>Dynamena dalmasi</i> (Cn) <i>Pachycheles rugimanus</i> (Ar) <i>Scyllarides nodifer</i> (Ar)	




Figure 4.—Inverse classification hierarchies and nodal diagram showing constancy and fidelity of station-species group coincidence based on seasonal trawl collections.

Atlantic Bight. However, the least similarity in species composition apparently occurs between inner- and outer-shelf sites while middle-shelf sites represent a transition zone between the other strata.

Biomass of Larger Macroinvertebrates

Comparisons of total biomass between taxa showed that Porifera were dominant at all stations during each season, except at station MS06 in spring, when biomass of Ascidiacea was highest (Table 4). These results are similar to those reported by Wenner et al. (1983) who found that Porifera accounted for a large percentage of the total invertebrate biomass from both the dredge and trawl collections. In the present study, the seasonal contribution of Porifera to the total biomass of invertebrates collected by dredge and trawl was as follows: Winter (78%), spring (59%), summer (67%), and fall (76%). The only taxa other than sponges and tunicates which formed a significant component of sample biomass in dredge or trawl collections were anthozoans and mollusks.

Analysis of variance of logarithmically transformed biomass values from replicate samples (OS06 excluded) indicated no significant difference in biomass between seasons for invertebrates collected by dredge or trawl ($P > 0.05$). Because many of the large macrofauna sampled by dredge and trawl, such as sponges, appear to persist throughout the year, major changes in biomass were not observed. Undoubtedly, our sampling methods were not refined enough to detect minor seasonal changes in biomass due to growth or mortality.

Significant differences in biomass were noted between stations ($P < 0.05$) for dredge-collected invertebrates, with the greatest average biomass recorded at stations IS02 ($\bar{x} = 22.3$ kg) and IS01

($\bar{x} = 13.6$ kg). The lowest biomass of invertebrates collected by dredge occurred at station MS06 ($\bar{x} = 4.9$ kg). Low biomass at this station is probably related to the patchy occurrence of invertebrates. Television transects showed that epifaunal organisms were sparse over parts of this station, with large expanses of sand interspersed with patches of live bottom. In addition, some sampling bias may have resulted from limiting dredge tows at MS02 and MS06 to areas of low relief. Since epifaunal growth was most dense on moderate to high relief ledges and rocks, an underestimation of biomass may have resulted from avoiding these areas.

Biomass values for trawl-collected invertebrates were greatest at an inner-shelf site, which is consistent with results from the initial study (Wenner et al. 1983). Among stations sampled by trawl (OS06 excluded), average biomass values were significantly greater ($P < 0.001$) at IS02 ($\bar{x} = 35.8$ kg), MS02 ($\bar{x} = 23.2$ kg), and IS01 ($\bar{x} = 10.6$ kg). The lowest average biomass was recorded at station OS01 ($\bar{x} = 3.2$ kg).

Smaller Macroinvertebrates Sampled by Suction and Grab

A total of 845 taxa was collected during seasonal sampling with suction sampler and grab. A listing of all fauna collected, including their ranks by abundance and density at each station, is presented in Appendix 3. The phylum Annelida, which was represented by more species than any other group during the initial study (Wenner et al. 1983), was also the most diverse group in the present study, where it contained 285 species. Annelids also dominated collections in terms of numerical abundance, accounting for 72.4% of the total 60,739 invertebrate specimens taken by suction sampler and grab. Mollusks ranked second among major

Table 4 —Percent contribution to epifaunal biomass of taxonomic groups in dredge (D) and trawl (T) collections. Percent of the total biomass for both gear types is also shown. The mean ($\bar{x}$), standard deviation (SD), and number of samples (n) for which biomass measurements were taken are shown.

Seaso.	IS01			IS02			MS02			MS06			OS01			OS06		
	D	T	Total	D	T	Total	D	T	Total	D	T	Total	D	T	Total	D	T	Total
Winter																		
Porifera	27	66	54	58	89	86	13	86	85	45	85	84	30	37	34			
Hydroidea	<1	<1	<1	<1	<1	<1	3	<1	<1	2	<1	<1	2	1	2			
Anthozoa	38	4	14	9	2	3	14	2	3	3	2	2	37	15	23			
Mollusca	6	1	3	18	<1	2	24	<1	1	36	1	2	15	5	8			
Decapoda	<1	<1	<1	<1	<1	<1	<1	<1	<1	1	<1	<1	2	2	2			
Echinodermata	5	1	3	7	1	2	21	1	2	6	<1	<1	14	13	14			
Ascidacea	24	26	25	7	7	7	25	9	9	7	11	11	<1	27	18			
Other																		
Invertebrata	0	0	0	0	<1	<1	0	0	0	0	0	0	0	0	0			
Total biomass (kg)	Σ	47.80	111.09	158.89	29.55	284.64	314.19	2.47	179.54	182.01	1.09	77.79	78.88	6.27	11.94	18.22		
	$\bar{x}$	23.90	18.50	19.86	—	56.93	52.36	—	29.90	26.00	0.55	12.96	9.86	3.14	1.99	2.28	Not sampled	
	SD	15.33	10.78	11.08	—	45.01	41.45	—	38.08	36.28	0.52	11.96	11.63	0.62	1.76	1.59		
	n	2	6	8	1	5	6	1	6	7	2	6	8	2	6	8		
Spring																		
Porifera	60	43	51	57	65	62	32	75	70	25	32	31	<1	65	57	0		
Hydroidea	<1	1	1	0	<1	<1	1	<1	<1	<1	<1	1	2	4	4	5		
Anthozoa	9	1	5	6	2	4	18	4	5	5	3	4	41	6	10	65		
Mollusca	7	1	4	11	1	4	8	2	3	35	10	13	1	<1	<1	17		
Decapoda	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	11	9	24	16	17	12		
Echinodermata	4	1	2	6	1	3	2	4	4	10	7	7	31	8	10	1		
Ascidacea	20	52	37	20	30	27	38	15	17	25	37	35	<1	1	1	0		
Total biomass (kg)	Σ	36.53	42.76	79.29	78.14	191.34	269.48	14.29	108.13	122.42	7.98	48.10	56.08	3.12	23.38	26.50	0.16	
	$\bar{x}$	18.26	7.13	9.91	39.07	31.90	33.68	7.14	18.02	15.30	3.99	8.02	7.01	0.27	3.99	3.55	0.08	
	SD	1.17	5.88	7.18	10.33	15.76	14.27	5.50	12.24	11.69	4.75	5.71	5.48	1.56	3.89	3.31	0.04	
	n	2	6	8	2	6	8	2	6	8	2	6	8	2	6	8	2	
Summer																		
Porifera	52	31	36	29	73	72	2	74	70	84	77	77	80	36	46	49		
Hydroidea	<1	2	1	1	<1	1	9	2	2	<1	<1	<1	0	2	2	12		
Anthozoa	5	4	5	12	2	2	1	10	10	6	2	2	14	13	13	21		
Mollusca	6	22	18	7	1	1	2	<1	<1	3	<1	1	<1	<1	<1	6		
Decapoda	<1	<1	<1	<1	<1	<1	<1	2	2	<1	2	2	<1	6	5	<1		
Echinodermata	4	4	4	6	1	1	3	2	2	<1	13	12	2	29	23	11		
Ascidacea	33	36	36	44	23	23	82	10	14	6	6	6	3	13	11	1		
Total biomass (kg)	Σ	11.45	37.74	49.19	6.22	288.89	295.09	5.12	88.74	93.87	2.90	42.71	45.61	4.62	14.90	19.52	1.54	
	$\bar{x}$	5.73	6.30	6.15	3.11	48.15	36.89	2.56	14.79	11.73	1.45	7.12	5.70	2.31	2.48	2.44	—	
	SD	4.86	3.10	3.22	3.35	33.45	35.15	3.30	19.73	17.66	1.17	10.41	9.20	2.91	1.73	1.83	—	
	n	2	6	8	2	6	8	2	6	8	2	6	8	2	6	8	1	
Fall																		
Porifera	20	76	66	66	45	54	68	88	86	95	93	93	40	54	50	36		
Hydroidea	1	<1	<1	<1	<1	<1	3	2	2	<1	<1	<1	<1	<1	1	3		
Anthozoa	38	2	9	4	5	5	1	3	3	<1	1	1	10	9	9	25		
Mollusca	8	1	2	6	1	4	6	<1	1	2	1	1	35	1	12	18		
Decapoda	<1	<1	<1	<1	<1	<1	<1	1	1	0	<1	<1	4	20	14	5		
Echinodermata	2	1	1	8	3	5	4	2	2	1	2	2	2	15	11	11		
Ascidacea	30	19	21	15	44	32	18	4	5	1	3	3	9	0	3	2		
Total biomass (kg)	Σ	13.08	61.63	74.72	42.34	56.31	98.64	16.53	157.30	173.83	27.18	90.30	117.48	12.42	25.47	37.89	0.99	
	$\bar{x}$	6.54	10.27	9.34	21.17	9.38	12.33	8.26	26.22	21.73	13.59	15.05	14.68	6.21	4.24	4.74	0.50	
	SD	5.10	8.54	7.67	7.95	6.64	8.38	4.04	34.04	29.98	14.17	16.09	14.64	1.59	4.23	3.74	0.37	
	n	2	6	8	2	6	8	2	6	8	2	6	8	2	6	8	2	

taxonomic groups with a total of 251 species; however, mollusks accounted for only 4.3% of the total number of invertebrate specimens and were outranked by amphipods (100 species) which represented 13.3% of total invertebrate abundance. Other groups represented in suction and grab collections included decapod crustaceans (111 species), arthropods other than amphipods and decapods (49 species), echinoderms (39 species), nemerteans (3 species), and sipunculans (7 species). Many of the taxa collected by suction and grab represent sand-bottom species which are also common at the hard-bottom sites, since there is generally a sand layer covering the rock bottom.

The 10 invertebrate species which numerically outranked others at the hard-bottom sites during all seasons were the polychaetes *Filograna implexa* ($n = 30,346$), *Pista palmata* ($n = 1,439$), *Syllis spongicola* ($n = 1,358$), *Spiophanes bombyx* ($n =$

650), *Chone americana* ($n = 543$), *Malacoceros glutaeus* ($n = 535$), and *Exogone dispar* ($n = 482$); and the amphipods *Luconacia incerta* ($n = 907$), *Photis* sp. ($n = 631$), and *Erichthonius brasiliensis* ($n = 523$). These species accounted for about 62% of all invertebrate specimens caught in suction and grab samples.

Densities of the 10 numerically dominant species changed seasonally and bathymetrically. In winter, the amphipods *Luconacia incerta*, *Photis* sp., and *Erichthonius brasiliensis* were important at inner-shelf sites (Table 5). Their densities at these sites decreased in spring; and during the rest of the year <5 individuals/0.1 m² were collected. The density of *F. implexa* was high in fall, when approximately 182 individuals/0.1 m² were collected at IS01. Another species whose density was highest in fall on the inner shelf was the polychaete *Syllis spongicola*. However,

Table 5.—Average density ($\bar{x}$) expressed as number per 0.1 m², and percent of total number (%) for 10 numerically dominant invertebrates collected by suction and grab. Number of collections (n) is indicated.

Species	IS01 ($n = 5$)		IS02 ($n = 5$)		MS02 ($n = 5$)		MS06 ($n = 5$)		OS01 ($n = 3$)		OS06 ($n = 5$)	
	$\bar{x}$	%	$\bar{x}$	%	$\bar{x}$	%	$\bar{x}$	%	$\bar{x}$	%	$\bar{x}$	%
----- Winter -----												
<i>Filograna implexa</i>	3.2	<1	0.4	<1	53.2	8.9	0.6	<1	—	—	—	—
<i>Pista palmata</i>	0.2	<1	0.8	<1	1.4	<1	2.2	<1	20.33	20.4	—	—
<i>Syllis spongicola</i>	6.6	1.6	1.4	<1	35.4	5.9	16.0	4.2	—	—	—	—
<i>Luconacia incerta</i>	22.6	5.6	68.0	15.8	17.8	2.9	45.0	11.9	—	—	—	—
<i>Spiophanes bombyx</i>	4.6	1.2	0.4	<1	3.0	<1	9.4	2.5	17.66	17.7	—	—
<i>Photis</i> sp.	24.8	6.2	20.4	4.7	31.8	5.3	2.0	<1	3.66	3.7	—	—
<i>Chone americana</i>	4.4	1.1	—	—	7.8	1.3	5.2	1.4	0.33	<1	—	—
<i>Malacoceros glutaeus</i>	0.4	<1	0.2	<1	19.0	3.2	6.2	1.6	—	—	—	—
<i>Erichthonius brasiliensis</i>	41.0	10.2	27.0	6.3	2.0	<1	0.2	<1	—	—	—	—
<i>Exogone dispar</i>	16.8	4.2	6.6	1.5	18.0	3.0	7.2	1.9	—	—	—	—
----- Spring -----												
<i>Filograna implexa</i>	—	—	0.2	<1	1,084.4	66.2	0.2	<1	—	—	34.6	14.2
<i>Pista palmata</i>	—	—	0.4	<1	1.8	<1	—	—	248.4	43.3	2.0	<1
<i>Syllis spongicola</i>	4.8	2.4	1.2	<1	17.2	1.0	0.2	<1	7.6	1.3	4.4	1.8
<i>Luconacia incerta</i>	4.6	2.2	2.0	<1	7.0	<1	3.4	1.8	—	—	1.0	<1
<i>Spiophanes bombyx</i>	0.4	<1	0.2	<1	2.4	<1	22.2	11.6	48.8	8.5	4.4	1.8
<i>Photis</i> sp.	0.8	<1	10.8	3.7	4.2	<1	2.3	1.3	15.8	2.7	0.6	<1
<i>Chone americana</i>	1.8	<1	0.4	<1	41.2	2.5	3.8	2.0	0.6	<1	2.8	1.1
<i>Malacoceros glutaeus</i>	0.2	<1	0.4	<1	11.6	<1	2.0	1.0	—	—	—	—
<i>Erichthonius brasiliensis</i>	3.4	1.7	16.8	5.7	1.6	<1	0.6	<1	1.4	<1	0.4	1
<i>Exogone dispar</i>	4.6	2.2	6.0	2.0	8.4	<1	0.6	<1	—	—	—	—
----- Summer -----												
<i>Filograna implexa</i>	0.2	<1	—	—	0.2	<1	1.6	<1	—	—	1.0	<1
<i>Pista palmata</i>	—	—	—	—	1.0	<1	0.8	<1	14.0	10.1	0.2	<1
<i>Syllis spongicola</i>	14.8	10.4	3.6	2.2	7.4	3.7	6.6	2.8	8.0	5.7	2.0	1.5
<i>Luconacia incerta</i>	0.8	<1	0.2	<1	0.8	<1	0.4	<1	—	—	—	—
<i>Spiophanes bombyx</i>	0.2	<1	0.2	<1	0.6	<1	1.6	<1	11.6	8.4	—	—
<i>Photis</i> sp.	0.8	<1	—	—	—	—	0.2	<1	—	—	3.0	2.2
<i>Chone americana</i>	1.4	1.0	—	—	1.0	<1	24.6	10.4	0.2	<1	4.6	3.4
<i>Malacoceros glutaeus</i>	2.8	1.9	1.8	1.1	2.4	1.2	8.8	3.7	0.2	<1	—	—
<i>Erichthonius brasiliensis</i>	4.0	2.8	—	—	0.2	<1	0.4	<1	0.4	<1	0.2	<1
<i>Exogone dispar</i>	1.6	1.1	1.2	<1	0.8	<1	2.0	<1	—	—	0.2	<1
----- Fall -----												
<i>Filograna implexa</i>	182.4	41.4	2.4	2.2	531.6	55.4	4,155.0	93.0	—	—	18.0	24.6
<i>Pista palmata</i>	0.4	<1	0.2	<1	0.4	—	1.2	<1	—	—	0.2	<1
<i>Syllis spongicola</i>	55.0	12.5	1.6	1.5	44.6	4.6	20.2	<1	10.0	13.4	3.0	4.1
<i>Luconacia incerta</i>	3.6	<1	0.4	<1	3.0	—	0.8	<1	—	—	—	—
<i>Spiophanes bombyx</i>	—	—	0.8	<1	1.8	—	1.6	<1	5.0	6.7	0.2	<1
<i>Photis</i> sp.	1.6	<1	0.4	<1	4.4	<1	—	—	—	—	—	—
<i>Chone americana</i>	1.8	<1	—	—	3.4	<1	2.4	<1	0.4	<1	0.6	<1
<i>Malacoceros glutaeus</i>	5.4	1.2	7.0	6.4	36.8	3.8	1.8	<1	—	—	—	—
<i>Erichthonius brasiliensis</i>	1.0	<1	0.4	<1	3.2	<1	0.4	<1	—	—	—	—
<i>Exogone dispar</i>	14.2	3.2	2.8	2.6	3.8	<1	1.6	<1	—	—	—	—

neither of these species was consistently abundant at both inner-shelf sites.

Filigrana implexa was the dominant species at middle-shelf sites (Table 6). The maximum density of *F. implexa* occurred in fall at MS06 when 4,155 individuals/0.1 m² were collected (Table 5). The high densities of *F. implexa* which were observed at mid-shelf sites did not occur on the outer shelf (Table 6). Species which were important at these deeper sites were the polychaetes *Pista palmata* and *Spiophanes bombyx*, with the highest densities for both occurring in spring.

Comparisons with results presented by Wenner et al. (1983) confirm that *F. implexa* is the numerically dominant invertebrate at middle-shelf sites. However, its importance at outer-shelf sites was considerably diminished during the second year of sampling when numbers of *Pista palmata* exceeded those of other species collected. *Filigrana implexa* was also important on the inner shelf, but densities did not approach those noted in previous collections from these sites.

Other investigators (Frankenberg 1971; Frankenberg and Leiper 1977; Tenore 1978) have noted that density variation over space and time is typical of numerically dominant species in soft-bottom communities on the continental shelf off Georgia. Wenner et al. (1983) speculated that variable abundances of benthic invertebrates could be influenced by 1) differentially successful and sequential recruitment by larvae of various species; 2) predation,

especially by fishes; and 3) differing microhabitats within and between sites.

Predation could influence the communities in this study since it has been found to be a major factor affecting benthic populations in soft- and hard-bottom communities of the intertidal zone (Paine 1966; Peterson 1977). However, studies on the effects of predation on the continental shelf are limited and largely conjectural. Frankenberg and Leiper (1977) hypothesized that fishes remove large numbers of small crustaceans, such as cumaceans and amphipods, from benthic assemblages. Predation on benthos can also be expected to alter species composition, resulting in greater abundance of those species having higher intrinsic rates of population increase (Slobodkin 1961); and, furthermore, a selective preference of predators for soft forms such as crustaceans or polychaetes may shift species composition toward forms with exoskeletons which are more resistant to predation (e.g., mollusks) (Richards 1963). Examination of stomach contents of 10 species of fishes known to inhabit live-bottom areas showed that amphipods, polychaetes, and decapods were most heavily preyed upon, including the numerically dominant species *Exogone dispar* and *Erichthonius brasiliensis* (G. Sedberry³).

³G. Sedberry, Assistant Marine Scientist, Marine Resources Research Institute, Box 12559, Charleston, SC 29412, pers. commun. January 1983.

Table 6.—The 10 most abundant invertebrate species collected by suction and grab samplers at each station. (An = Annelida; Ar = Arthropoda; Ech = Echinodermata; Si = Sipunculida.)

Species	Total number	Percent of total	Species	Total number	Percent of total
IS01			IS02		
<i>Filigrana implexa</i> (An)	929	15.6	<i>Luconacia incerta</i> (Ar)	353	7.1
<i>Syllis spongicola</i> (An)	406	6.8	<i>Elasmopus</i> sp. A (Ar)	225	4.5
<i>Erichthonius brasiliensis</i> (Ar)	247	4.2	<i>Erichthonius brasiliensis</i> (Ar)	221	4.4
<i>Axiobella mucosa</i> (An)	246	4.1	<i>Lembos smithi</i> (Ar)	179	3.6
<i>Exogone dispar</i> (An)	186	3.1	<i>Caprella equilibra</i> (Ar)	165	3.3
<i>Luconacia incerta</i> (Ar)	158	2.7	<i>Podocerus</i> sp. (Ar)	160	3.2
<i>Ampharete americana</i> (An)	143	2.4	<i>Photis</i> sp. (Ar)	158	3.2
<i>Photis</i> sp. (Ar)	140	2.4	<i>Leptochelia</i> sp. (Ar)	127	2.6
<i>Elasmopus</i> sp. A (Ar)	130	2.2	<i>Lumbrineris inflata</i> (An)	109	2.2
<i>Gammaropsis</i> sp. (Ar)	105	1.8	<i>Polycirrus carolinensis</i> (An)	84	1.7
MS02			MS06		
<i>Filigrana implexa</i> (An)	8,347	49.2	<i>Filigrana implexa</i> (An)	20,787	78.8
<i>Syllis spongicola</i> (An)	523	3.1	<i>Luconacia incerta</i> (Ar)	248	0.9
<i>Malacoceros glutaeus</i> (An)	349	2.1	<i>Syllis spongicola</i> (An)	215	0.8
<i>Chone americana</i> (An)	267	1.6	<i>Chone americana</i> (An)	180	0.7
<i>Ophiobothris angulata</i> (Ech)	203	1.2	<i>Spiophanes bombyx</i> (An)	174	0.7
<i>Photis</i> sp. (Ar)	202	1.2	<i>Syllis hyalina</i> (An)	173	0.7
<i>Acanthoastorius millsi</i> (Ar)	195	1.2	<i>Aspidosiphon gosnoldi</i> (Si)	120	0.5
<i>Megalobrachium soriatum</i> (Ar)	192	1.1	<i>Polydora colonia</i> (An)	107	0.4
<i>Lumbrineris inflata</i> (An)	165	1.0	<i>Autolytus</i> sp. (An)	96	0.4
<i>Exogone dispar</i> (An)	155	0.9	<i>Malacoceros glutaeus</i> (An)	94	0.4
OS01			OS06		
<i>Pista palmata</i> (An)	1,373	32.4	<i>Filigrana implexa</i> (An)	268	11.9
<i>Spiophanes bombyx</i> (An)	380	8.9	<i>Plakosyllis quadrioculata</i> (An)	112	5.0
<i>Erichthonius</i> sp. A (Ar)	170	4.0	<i>Rildardanus laminosa</i> (Ar)	104	4.6
<i>Syllis spongicola</i> (An)	128	3.0	<i>Protomedea</i> sp. A (Ar)	71	3.2
<i>Onuphis pallidula</i> (An)	114	2.7	<i>Ampharete americana</i> (An)	54	2.4
<i>Photis</i> sp. (Ar)	90	2.1	<i>Syllis spongicola</i> (An)	53	2.4
<i>Owenia fusiformis</i> (An)	84	2.0	<i>Sphaerosyllis</i> sp. (An)	47	2.1
<i>Photis longicaudata</i> (Ar)	67	1.6	Syllidae L. (An)	47	2.1
<i>Mesochaetopterus</i> sp. (An)	59	1.4	<i>Chone americana</i> (An)	40	1.8
<i>Ampharete acutifrons</i> (An)	57	1.4	<i>Syllis hyalina</i> (An)	40	1.8

Much of the variability in populations of infaunal organisms can be attributed to our sampling methods which included too few replicates to accurately determine whether changes in density were due only to seasonality. In addition, the complex physiography and diverse substrates of hard-bottom habitats create a habitat patchiness that could result in either selective settling by larvae or differential survival of different benthic invertebrate species. Other physical factors which have been reported to affect stability (i.e., persistence) of benthic populations are fluctuations in water temperature (Boesch et al. 1977) and disturbance of sediments by storms (McCall 1977). Boesch et al. (1977) attributed the temporal stability of macrofaunal species on the outer shelf of the Middle Atlantic Bight to decreasing thermal variation and reduced effects of storms in deeper water. Although upwelling of colder water does occur on the continental shelf in the South Atlantic Bight, most of the benthic infaunal species collected by suction sampler and grab have wide latitudinal ranges and are eurybathic, extending from the intertidal zone to the continental slope and beyond. Thus, most should be fairly tolerant of the environmental conditions found at hard-bottom sites on the shelf.

Values of Shannon diversity (H'), evenness (J'), richness (SR), dominance (DI), number of species (s), and number of individuals (n) showed no consistent patterns between depth zones and seasons (Table 7). During all seasons, H' was > 5 bits/individual at both inner-shelf stations, except IS01 in fall when H' was 4.58. This comparatively low H' value reflected an uneven distribution of individuals among species at IS01 and, consequently, an elevated dominance index (DI) value rather than low SR or s . Low evenness and high DI were attributed to large numbers of the colonial polychaete *Filograna implexa*.

More species were collected at mid-shelf stations than at any others sampled. Approximately 42 and 48% of all species collected at MS02 and MS06, respectively, were represented by one or two individuals and are consequently considered rare. Both s and SR were greatest at mid-shelf stations in winter; however, the

community-structure values were not seasonally consistent between stations. For example, values of H' were high (> 6) at MS06 during all seasons except fall when the lowest H' value for any station sampled occurred. At MS02, high H' values were noted only in winter and summer. As noted for inner-shelf stations, low H' values were affected by low evenness which reflected strong dominance by the polychaete *F. implexa*.

Community-structure values at outer-shelf stations were also not consistent between stations or seasons. At station OS06, H' and J' showed little seasonal variation and indicated that species were fairly diverse and evenly distributed. The dominance index also indicated that collections were not numerically dominated by one or two species. However, s and SR at OS06 did change with season, being much lower in fall. At station OS01, seasonal changes in H' reflected concomitant changes in evenness and dominance. Thus, although more species and greater SR occurred at OS01 in spring, H' was lower than in any other season at this station due to numerical dominance by the polychaete *Pista palmata*. Most community-structure values at outer-shelf sites were comparable with those obtained on the inner shelf; however, the fewest species obtained on the shelf were recorded at OS01 in winter and OS06 in fall. In addition, fewest individuals were collected at outer-shelf stations.

In general, results from the present study are consistent with those reported in the initial study of hard-bottom habitats (Wenner et al. 1983). Sampling during both years has shown that rare species are a conspicuous component of the hard-bottom community and contribute to elevated species richness at all of the sites sampled. In addition, no consistent seasonal changes in diversity or s (Kruskal-Wallis test, $P > 0.05$) were noted for stations or depth zones. The low H' values which occurred at hard-bottom sites sampled in both years of study apparently reflect localized changes in the population density of a few dominant species, such as *F. implexa*, and are probably not indicative of any seasonality.

Although H' and its components did not show any definite patterns with regard to depth or latitude, the number of species was found to be significantly different between sites (Kruskal-Wallis test, $P < 0.01$). Specifically, more species were collected at mid-shelf sites, which may be related to more stable bottom temperatures in this area of the shelf (Miller and Richards 1980). In terms of community structure, these sites were characterized by 1) a large number of species, many of which were rare, and 2) high numerical dominance by *Filograna implexa*. In contrast, inner-shelf sites, which were subject to more extreme temperature variations, generally had fewer species and less dominance than sites on the middle shelf. Outer-shelf sites are also thermally variable because of intrusions of colder upwelled water, which predominantly occurs in fall (Atkinson et al. 1978). Fewest species were recorded at outer-shelf sites, and many of those which occurred were rare. In addition, dominance was relatively low. These results suggest that fluctuating temperatures at inner- and outer-shelf sites may exert some control on community structure at live-bottom sites. However, it is also possible that the lower richness at outer-shelf sites was due to collection of samples with a grab which does not sample hard substrates as effectively as the suction device. Although within-reef habitat heterogeneity is one important factor in determining richness of benthic fauna, thermal stability is probably also important.

Normal cluster analysis of pooled replicate collections taken with suction and grab samplers grouped stations according to depth zones with inner-shelf stations forming group 1, middle-shelf stations forming group 2, and two outer-shelf sites forming

Table 7.—Seasonal community structure values [number of individuals, number of species, diversity (H'), evenness (J'), richness (SR), and dominance index (DI)] for pooled replicate samples of invertebrates at each station.

Station	Season	No. individuals	No. species	H'	J'	SR	DI
IS01	winter	2,003	245	6.09	0.77	32.10	17.2
	spring	1,020	178	5.90	0.79	25.55	24.2
	summer	713	172	6.17	0.83	26.03	20.5
	fall	2,204	222	4.58	0.59	28.71	53.8
IS02	winter	2,158	176	5.58	0.75	22.80	22.6
	spring	1,468	197	6.41	0.84	26.88	11.2
	summer	808	172	6.42	0.87	25.54	12.9
	fall	547	148	6.25	0.87	23.30	15.2
MS02	winter	2,987	313	6.33	0.79	39.00	14.8
	spring	8,190	306	3.22	0.39	33.85	68.7
	summer	990	261	7.19	0.90	37.70	8.5
	fall	4,799	307	3.99	0.48	36.10	60.0
MS06	winter	1,896	319	6.75	0.81	42.13	17.3
	spring	954	230	6.61	0.84	33.38	17.4
	summer	1,188	251	6.71	0.84	35.34	14.6
	fall	22,328	297	0.85	0.10	29.46	93.5
OS01	winter	299	77	4.88	0.78	13.33	38.1
	spring	2,869	255	4.48	0.56	31.90	51.8
	summer	694	135	5.78	0.82	20.48	18.4
	fall	373	125	6.10	0.88	20.94	20.1
OS06	winter	—	—	—	—	—	—
	spring	1,215	198	6.03	0.79	27.74	21.3
	summer	670	189	6.69	0.88	28.89	7.5
	fall	366	99	5.27	0.80	16.60	32.2

groups 3 and 4 (Fig. 5). The two outer-shelf station groups constituted a larger agglomeration in the dendrogram which was very dissimilar in species composition to the inner- and middle-shelf groups.

None of the stations were grouped together by season; however, inner- and outer-shelf sites were grouped according to latitude with IS01 and OS01 clustering separately from IS02 and OS06, respectively. Major agglomerations in the dendrogram indicated bathymetric separation of sites, suggesting that depth is a major factor influencing species distribution. During the previous study, quantitative collections of smaller epibenthic and infaunal organisms were separated according to bathymetric zones during winter but not during summer, suggesting that the quantitative composition of the hard-bottom community was much more homogeneous across the shelf during summer (Wenner et al. 1983). However, based on the present study, we have found no evidence to suggest consistent seasonal change in community composition. Similarly, seasonal differences in species assemblages were not observed by Boesch et al. (1977) in a study of benthic communities in the Middle Atlantic Bight.

Most species groups delineated by cluster analysis were not limited in their distribution to one depth zone (Table 8, Fig. 6). Species in group A were consistently collected at inner- and middle-shelf sites, whereas those in group B were highly constant at all sites. Group C species were also ubiquitous, but displayed high constancy for middle-shelf stations. The only species in this group which was collected solely at middle-shelf sites was the polychaete *Ceratonereis* sp. A. Species in group D, which were widely distributed across the shelf, displayed highest constancy at middle-shelf stations and OS01. Groups E, F, and G were most consistently collected at inner- and middle-shelf stations but were not restricted in their distribution to these sites. The tanaidacean *Apseudes* sp. A, which occurred in group G, was the only species which was restricted to inner-shelf sites. Species in group H were found at all stations but were most frequently encountered at middle-shelf stations and OS06.

Examination of inverse cluster groups and nodal diagrams from the initial survey of hard-bottom habitats also revealed that few

species were limited to specific bathymetric zones during both years of study (Wenner et al. 1983). No species was found to be highly constant at inner-shelf sites, whereas the polychaete *Eunice filamentosa*, the amphipod *Megaluropus* sp., and the mollusk *Chione grus* were most consistently collected at middle-shelf sites during both years. In addition, the amphipod *Erichthonius* sp. A, the polychaete *Syllis* sp. D, and the echinoderm *Genocidaris maculata* were characteristically found at sites sampled on the outer shelf in both studies. These results indicate that many of the smaller epifaunal and infaunal species comprising benthic assemblages at hard-bottom sites are relatively ubiquitous across the shelf, although differences in overall community composition were observed between depth zones.

CONCLUSIONS

1) A total of 1,307 invertebrate taxa was collected with all sampling devices documenting the diverse nature of offshore hard-bottom habitats. Most sites were inhabited by rare species which were represented by only one or two individuals. Numerical dominance was limited to the polychaetes *Filograna implexa* and *Pista palmata* which were prevalent at middle- and outer-shelf sites, respectively.

2) Diversity of hard-bottom sites was generally high during all seasons. No consistent seasonal changes in diversity were noted for stations or depth zones throughout the study area. Low H' values were attributable to low evenness caused by high numbers of species such as *Filograna implexa* and *Pista palmata*. Although H' and its components did not show any well defined changes with depth, season, or latitude, more species were collected at middle-shelf sites. This may be related to less variable bottom temperatures in this area of the shelf, coupled with relatively high habitat complexity due to the frequent occurrence of large sponges and octocorals, which offer refuge to other benthic invertebrates. The number of identifiable invertebrate taxa collected at each station by dredge and trawl did not change significantly between seasons. Changes in the number of species between stations were attributable to varying degrees of topographic complexity and habitat heterogeneity.

3) Porifera, Bryozoa, and Cnidaria were the most diverse taxonomic groups collected by dredge and trawl during both years in which sampling was conducted. Sponges were most diverse at sites on the inner shelf, but their diversity diminished in dredge and trawl collections at middle- and outer-shelf sites. Taxa accounting for the greatest number of species in the latter areas were cnidarians and bryozoans.

4) Comparisons of total biomass between taxa showed that Porifera were dominant at most stations. No significant difference in biomass estimates was observed between seasons; however, bathymetric differences did occur, with biomass being greatest on the inner shelf for dredge and trawl collections. This may reflect the more uniform distribution of epifaunal invertebrates at inner-shelf sites; however, differences between sites may also be caused by sampling bias since efforts were made not to sample over high relief ledges on the middle shelf in order to minimize loss of gear.

5) The hard-bottom sites sampled throughout the entire study area were distinguished from each other primarily by trends in faunal composition related to depth. No evidence of seasonal grouping was noted. Epifaunal species censused by dredge and trawl appeared to be more faithful to particular depth zones than did benthic organisms sampled by suction sampler and grab. However, most species assemblages were not restricted to a particular depth zone. Inner- and middle-shelf sites were more




Figure 5.—Normal cluster dendrogram of station groups formed using the Bray-Curtis similarity coefficient, log transformation, and flexible sorting. Data from replicated suction and grab collections were pooled for each station prior to cluster analysis.

Table 8.—Species groups resulting from numerical classification of data from samples collected by suction and grab samplers during all seasons, 1981. (Am = Amphipoda; Br = Branchiopoda; Cu = Cumacea; D = Decapoda; E = Echinodermata; I = Isopoda; M = Mollusca; My = Mysidacea; P = Polychaeta; Po = Porifera; Py = Pycnogonida; Si = Sipunculida; T = Tanaidacea.)

Group A	<i>Filograna implexa</i> (P) <i>Harmothoe</i> sp. A (P) <i>Caulieriella killariensis</i> (P) <i>Ampharete americana</i> (P) <i>Prionospio cristata</i> (P) <i>Axiognathus squamatus</i> (E) <i>Loimia medusa</i> (P) <i>Spio pettiboneae</i> (P) <i>Polydora caeca</i> (P) <i>Prionospio cirrifer</i> (P) <i>Microdeutopus myersi</i> (Am) <i>Armandia maculata</i> (P) <i>Erichthonius brasiliensis</i> (Am) <i>Photis</i> sp. (Am) <i>Gammaropsis</i> sp. (Am) <i>Caprella equilibra</i> (Am) <i>Leptochelia</i> sp. (T) <i>Anoplodactylus petiolatus</i> (Py) <i>Diplodonta punctata</i> (M) <i>Mitrella lunata</i> (M) <i>Suturoglypta iontha</i> (M) <i>Pherusa ehlersi</i> (P) <i>Caprella penantis</i> (Am)	<i>Varicorbula operculata</i> (M) <i>Syllis cornuta</i> (P) <i>Schistomeringos rudolphi</i> (P) <i>Brania clavata</i> (P)	<i>Botula fusca</i> (M) <i>Piitho lherminieri</i> (D) <i>Pagurus hendersoni</i> (D) <i>Ceratonereis mirabilis</i> (P) <i>Ophiostigma isacanthum</i> (E) Majidae A (D) <i>Arabella mutans</i> (P) <i>Anchialina typica</i> (My) <i>Pelia mutica</i> (D) <i>Lithophaga bisulcata</i> (M) <i>Gastrochaena hians</i> (M)
	Group C	Group D	Group G
<i>Ophiothrix angulata</i> (E) <i>Polycirrus carolinensis</i> (P) <i>Lumbrineris inflata</i> (P) <i>Exogone dispar</i> (P) <i>Ophiactis savignyi</i> (E) <i>Luconacia incerta</i> (Am) <i>Podocerus</i> sp. (Am) <i>Megalobrachium soriatum</i> (D) <i>Paracerceis caudata</i> (I) <i>Ampithoe</i> sp. A (Am) <i>Elasmopus</i> sp. A (Am) <i>Carpas bermudensis</i> (I) <i>Lembos smithi</i> (Am) <i>Websterinereis tridentata</i> (P) <i>Podarke obscura</i> (P) <i>Synalpheus townsendi</i> (D) Xanthidae A (D) <i>Melita appendiculata</i> (Am) Sipunculida A (Si) <i>Pherusa inflata</i> (P) <i>Maera</i> sp. A (Am) <i>Pista quadrilobata</i> (P) <i>Arca imbricata</i> (M) <i>Ampelisca schellenbergi</i> (Am) <i>Chama congregata</i> (M) <i>Gouldia cerina</i> (M) <i>Pseudomadaeus agassizii</i> (D) <i>Lysianopsis alba</i> (Am) <i>Crepidula aculeata</i> (M) <i>Syllis gracilis</i> (P)	<i>Marginella lavalleana</i> (M) <i>Eunice filamentosa</i> (P) Melpheippidae A (Am) <i>Diodora cayenensis</i> (M) <i>Musculus lateralis</i> (M) <i>Ceratonereis versipedata</i> (P) <i>Ceratonereis</i> sp. A (P) <i>Inachoides forceps</i> (D) <i>Latreutes parvulus</i> (D) <i>Trichobranchus glacialis</i> (P) <i>Marphysa</i> sp. A (P) <i>Phyllodoce fragilis</i> (P) <i>Alpheus normanni</i> (D) <i>Trypanosyllis zebra</i> (P) <i>Sicyonia laevigata</i> (D) <i>Branchiosyllis exilis</i> (P) <i>Websterinereis</i> sp. B (P) <i>Chevalia</i> sp. (Am) <i>Chione grus</i> (M) <i>Polycirrus eximius</i> (P) <i>Hiatella arctica</i> (M) <i>Tellina americana</i> (M) Diastylidae A (Cu) <i>Semele nuculoides</i> (M) <i>Acanthohaustorius millsi</i> (Am) <i>Megaluropus</i> sp. (Am) <i>Syllis regulata carolinae</i> (P)	<i>Leptochela papulata</i> (D) <i>Bowmaniella portoricensis</i> (My) <i>Ampelisca cristoides</i> (Am) <i>Paraprionospio pinnata</i> (P) <i>Mesochaetopterus</i> sp. (P) <i>Onuphis nebulosa</i> (P) <i>Batrachonotus fragosus</i> (D) <i>Ampharete acutifrons</i> (P) Spionidae B (P) <i>Unciola</i> sp. A (Am) <i>Onuphis pallidula</i> (P) <i>Ampelisca vadorum</i> (Am) <i>Aonides mayaguezensis</i> (P) <i>Automate evermanni</i> (D) Xanthidae B (D) <i>Glycera capitata</i> (P) <i>Nassarius albus</i> (M) <i>Pomatoceros americanus</i> (P) <i>Nassarina minor</i> (M) <i>Lysidice ninetta</i> (P) <i>Lumbrineris coccinea</i> (P) <i>Tellina sybaritica</i> (M) <i>Laonice cirrata</i> (P) <i>Nephtys squamosa</i> (P) Pelecypoda B (M)	<i>Phyllodoce castanea</i> (P) <i>Pseudovermilia occidentalis</i> (P) <i>Apseudes</i> sp. A (T) <i>Arbacia punctulata</i> (E) <i>Eunice antennata</i> (P) <i>Pilumnus sayi</i> (D) <i>Hyboscolex longiseta</i> (P) Amphipoda E (Am) <i>Paraphoxus spinosus</i> (Am) Amphipoda B (Am) <i>Cerithium atratum</i> (M) <i>Vermiliopsis annulata</i> (P) <i>Cumingia tellinoides</i> (M) <i>Pseudovermilia</i> sp. A (P) <i>Mithrax pleuracanthus</i> (D) Tanystylidae A (Py) <i>Paguristes tortugae</i> (D)
Group B		Group E	Group H
<i>Aspidosiphon gosnoldi</i> (Si) <i>Syllis hyalina</i> (P) <i>Chone americana</i> (P) <i>Eunice vittata</i> (P) <i>Leucothoe spinicarpa</i> (Am) <i>Syllis spongicola</i> (P) <i>Pagurus carolinensis</i> (D) <i>Laevicardium pictum</i> (M) <i>Malacoceros glutaeus</i> (P) <i>Megalomma bioculatum</i> (P) Chrysopetalidae A (P) <i>Amphiodia pulchella</i> (E) <i>Eulalia sanguinea</i> (P) <i>Crassinella lunulata</i> (M) <i>Axiothella mucosa</i> (P) <i>Spiophanes bombyx</i> (P) <i>Owenia fusiformis</i> (P) <i>Pista palmata</i> (P)		<i>Erichsonella filiformis</i> (I) <i>Marginella hartleyanum</i> (M) <i>Lucifer faxoni</i> (D) <i>Pseudeurythoe ambigua</i> (P) <i>Mangelia rugirima</i> (M) <i>Corbula contracta</i> (M) <i>Aspidosiphon albus</i> (Si) <i>Chione latilirata</i> (M) <i>Phyllodoce longipes</i> (P) <i>Glycera</i> sp. B (P) <i>Marginella roscida</i> (M) <i>Phyllochaetopterus socialis</i> (P) <i>Subadyte pellucida</i> (P)	Melitidae A (Am) Syllidae L (P) <i>Syllis</i> sp. D (P) <i>Genocidaris maculata</i> (E) <i>Glycera tessellata</i> (P) <i>Odontosyllis enopla</i> (P) <i>Syllis alternata</i> (P) <i>Syllis ferrugina</i> (P) <i>Rildardanus laminosa</i> (Am) <i>Erichthonius</i> sp. A (Am) <i>Glycera papillosa</i> (P) <i>Proiodorvillea kefersteini</i> (P) <i>Lembos unicornis</i> (Am) <i>Goniadides carolinae</i> (P) <i>Thyone pseudofusus</i> (E)
		Group F	
		<i>Hydroides crucigera</i> (P) <i>Hydroides</i> sp. A (P)	




Figure 6.—Inverse classification hierarchies and nodal diagram showing constancy and fidelity of station-species group coincidence based on seasonal suction and grab collections.

similar in species composition to each other than they were to outer-shelf sites.

6) The densities of some numerically dominant species collected by suction and grab changed seasonally and bathymetrically. Such density variations may result from: 1) Differentially successful and sequential recruitment among larvae of various species; 2) predation, especially by marine fishes; and 3) habitat complexity. Seasonal influences on densities of individual species were not large enough to cause detectable seasonal changes in overall community structure and composition.

ACKNOWLEDGMENTS

We are grateful to all persons associated with the joint South Carolina-Georgia project to study hard-bottom habitats which was funded by the Minerals Management Service (Contract AA551-CT1-18). Those persons who helped with collection of samples and identification of invertebrate taxa included: B. B. Boothe, D. R. Calder, A. L. Edwards, J. V. Miglarese, F. L. Nicholson, C. B. O'Rourke, E. C. Roland, and S. Shipman. Additional technical help at sea and in the laboratory was provided by W. T. Hodges, Jr., M. S. Maclin, G. H. Steele, A. C. Vitucci, Jr., H. L. Ansley, C. F. Cowman, C. D. Harris, R. Mahood, and S. P. Olsson. We also thank M. J. Clise and G. Gash for help with data processing and coordination; L. H. Hodges and N. M. Beaumont for secretarial support; K. Swanson for graphics; and C. K. Biernbaum and G. R. Sedberry for critical review of the manuscript.

LITERATURE CITED

- ATKINSON, L. P., R. T. BARBER, and M. WADE.
1978. Hydrography, nutrients, chlorophyll, and organic carbon. South Atlantic Benchmark Program. Vol. 3 Draft Report, p. 15-72. Bureau of Land Management, Outer Continental Shelf Office, New Orleans, La.
- BOESCH, D. F.
1977. Application of numerical classification in ecological investigations of water pollution. U.S. Environmental Protection Agency Office of Research and Development, Report No. EPA-600/3-77-033, 115 p.
- BOESCH, D. F., J. N. KRAEUTER, and D. K. SERAFY.
1977. Distribution and structure of communities of macrobenthos on the outer continental shelf of the middle Atlantic Bight: 1975-1976 investigations. Va. Inst. Mar. Sci., Spec. Rep. Appl. Mar. Sci. Ocean Eng. 175, 111 p.
- BRAY, J. R., and J. T. CURTIS.
1957. An ordination of the upland forest communities of southern Wisconsin. Ecol. Monogr. 27:325-349.
- CHESS, J. R.
1979. An airlift sampling device for *in situ* collecting of biota from rocky substrata. Mar. Technol. Soc. J. 12(3):20-23.
- CLIFFORD, H. T., and W. STEPHENSON.
1975. An introduction to numerical classification. Acad. Press, N.Y., 229 p.
- CONTINENTAL SHELF ASSOCIATES.
1979. South Atlantic hard bottom study. Prepared for Bureau of Land Management (Contract AA551-CT8-25), Washington, D.C., 356 p.
- DEAN, T. A.
1981. Structural aspects of sessile invertebrates as organizing forces in an estuarine fouling community. J. Exp. Mar. Biol. Ecol. 53:163-180.
- DUKE UNIVERSITY MARINE LABORATORY.
1968. Manual for use of R/V *Eastward*. Duke University Cooperative Research and Training Program in Biological Oceanography, Beaufort, N.C., 146 p.
- FRANKENBERG, D.
1971. The dynamics of benthic communities off Georgia, USA. Thalassia Jugosl. 7:49-55.
- FRANKENBERG, D., and A. S. LEIPER.
1977. Seasonal cycles in benthic communities of the Georgia continental shelf. In B. C. Coull (editor), Ecology of marine benthos, p. 383-397. Univ. South Carolina Press, Columbia.
- LAMBERT, J. M., and W. T. WILLIAMS.
1962. Multivariate methods in plant ecology. IV. Nodal analysis. J. Ecol. 50:775-802.
- LANCE, G. N., and W. T. WILLIAMS.
1967. A general theory of classificatory sorting strategies. I. Hierarchical systems. Comput. J. 9:373-380.
- MARGALEF, D. R.
1958. Information theory in ecology. Gen. Syst. 3:36-71.
- McCALL, P. L.
1977. Community patterns and adaptive strategies of the infaunal benthos of Long Island Sound. J. Mar. Res. 35:221-266.

- McNAUGHTON, S. J.
1967. Relationships among functional properties of California grasslands. *Nature* (Lond.) 216:168-169.
- MENGE, B. A., and J. P. SUTHERLAND.
1976. Species diversity gradients: synthesis of the roles of predation, competition, and temporal heterogeneity. *Am. Nat.* 110:351-369.
- MILLER, G. C., and W. J. RICHARDS.
1980. Reef fish habitat, faunal assemblages, and factors determining distributions in the South Atlantic Bight. *Proc. Gulf Caribb. Fish. Inst.* 32: 114-130.
- OSMAN, R. W.
1977. The establishment and development of a marine epifaunal community. *Ecol. Monogr.* 47:37-63.
- PAINE, R. T.
1966. Food web complexity and species diversity. *Am. Nat.* 100:65-75.
- PETERSON, C. H.
1977. Competitive organization of the soft-bottom macrobenthic communities of southern California lagoons. *Mar. Biol. (Berl.)* 43:343-359.
- PIELOU, E. C.
1975. *Ecological diversity*. John Wiley & Sons, N.Y., 165 p.
- RICHARDS, S. W.
1963. The demersal fish population of Long Island Sound. *Bull. Bingham Oceanogr. Collect. Yale Univ.* 18(2), 101 p.
- RUSS, G. R.
1980. Effects of predation by fishes, competition, and structural complexity of the substratum on the establishment of a marine epifaunal community. *J. Exp. Mar. Biol. Ecol.* 42:55-69.
- SEED, R.
1976. Ecology. In B. L. Bayne (editor), *Marine mussels: their ecology and physiology*, p. 13-66. Cambr. Univ. Press, Cambridge, Engl.
- SCHOENER, T. W.
1974. Resource partitioning in ecological communities. *Science* (Wash., D.C.) 185:27-39.
- SIEGEL, S.
1956. *Nonparametric statistics for the behavioral sciences*. McGraw-Hill, N.Y., 312 p.
- SLOBODKIN, L. B.
1961. *Growth and regulation of animal populations*. Holt, Rinehart and Winston, N.Y., 184 p.
- TENORE, K. R.
1978. Macroinfaunal benthos. South Atlantic OCS Benchmark Program, 1977 report. Vol. III, p. 257-288. Bureau of Land Management, New Orleans, La.
- WENNER, E. L., D. M. KNOTT, R. F. VAN DOLAH, and V. G. BURRELL, Jr.
1983. Invertebrate communities associated with hard bottom habitats in the South Atlantic Bight. *Estuarine Coastal Shelf Sci.* 17:143-158.
- WILLIAMS, W. T., and J. M. LAMBERT.
1961. Multivariate methods in plant ecology. III. Inverse association-analysis. *J. Ecol.* 49:717-729.

Appendix 1. Phylogenetic list of invertebrate taxa collected by dredge during 1981 cruises to the stations off South Carolina and Georgia.
w = winter, sp = spring, s = summer and f = fall. No dredge samples were collected during winter samples at station OS06.

	IS01	IS02	MS02	MS06	OS01	OS06
Porifera						
Demospongea - Ceractinomorpha						
Keratosa						
Aplysillidae						
<u>Chelonaplysilla</u> sp.	f	f	f	-	-	-
Darwinellidae						
<u>Darwinella</u> sp.	-	w,sp,f	-	f	f	-
Spongiidae						
<u>Aplysina fistularis</u>	-	w,sp,f	sp,s,f	f	f	-
<u>Euspongia officinalis</u>		f				
<u>Ircinia campana</u>	w	w,sp,s,f	f	-	-	f
<u>Ircinia felix</u>	w,sp	w,sp,s,f	sp,f	-	w,f	-
<u>Ircinia ramosa</u>	w,s	w,sp,s,f	sp,f	sp	f	-
<u>Ircinia</u> sp. B	-	-	-	-	-	s,f
<u>Ircinia strobilina</u>	-	sp,f	sp,f	sp	-	-
<u>Oligoceras</u> sp.	-	-	-	-	f	-
<u>Spongia tubulifera</u>	sp	sp	-	sp	-	-
Haplosclerida						
Adociidae						
<u>Sigmatocia caerulea</u>	w	w	w,sp	-	w,s,f	-
<u>Strongylophora</u> sp. B	-	f	-	f	f	-
Desmacidonidae						
<u>Xytopsues griseus</u>	w,s	w,s,f	sp	-	w,s,f	-
Halicionidae						
<u>Amphimedon compressa</u>	-	f	f	f	f	-
<u>Callyspongia fallax</u>	s	-	-	f	-	-
<u>Haliclona areolata</u>	-	w	-	-	-	-
<u>Haliclona oculata</u>	w,sp,s,f	w,sp,s,f	-	s,f	-	-
<u>Haliclona</u> sp. C	w,sp	w	sp	-	-	-
undetermined Halicionidae	w	w	-	-	-	-
Nepheliospongiidae						
<u>Xestospongia halichondrioides</u>	-	-	-	sp	-	-
Poecilosclerida						
Poecilosclerida H						
	-	-	w	-	w	-
Amphilectidae						
<u>Tenaciella obliqua</u>	f	-	-	-	-	-
Microcionidae						
<u>Microciona prolifera</u>	s	sp,f	-	f	w,s,f	-
<u>Microciona</u> sp. D	-	-	-	-	s	-
<u>Microciona spinosa</u>	-	f	-	-	w	-
<u>Pandaros acanthifolium</u>	w,sp,s,f	w,sp	-	f	w,f	-

Appendix 1. (Continued)

	IS01	IS02	MS02	MS06	OS01	OS06
<u>Thalyseurypon</u> sp.	-	-	-	-	-	f
<u>Thalyseurypon foliacea</u>	-	sp	-	-	-	-
<u>Thalysias juniperina</u>	-	w,sp,s,f	w,sp,f	sp	-	f
<u>Thalysias</u> sp.	-	f	-	-	f	-
Mycalidae						
<u>Neofibularia nolitangere</u>	-	-	sp	sp	-	-
Myxillidae						
undetermined Myxillidae	w	-	-	-	-	-
Plocamiidae						
<u>Holoplocamia</u> sp.	-	-	sp	-	-	-
<u>Holoplocamia</u> sp. A	-	-	-	-	w	-
Tedaniidae						
<u>Lissodendoryx isodictyalis</u>	f	sp	f	sp,f	-	s
<u>Lissodendoryx sigmata</u>	s	s,f	sp,f	w	-	-
<u>Tedania ignis</u>	w	w,sp,f	-	-	-	-
Halichondriida						
Halichondriidae						
<u>Ciocalapata gibbsi</u>	w,sp,f	w,sp,s,f	f	f	w	f
<u>Halichondria bowerbanki</u>	w,sp,f	w,sp,s,f	f	-	-	f
Demospongea - Tetractinomorpha						
Axinellida						
Axinellidae						
<u>Axinella bookhouti</u>	w,sp,f	w,sp,f	-	f	w,s,f	-
<u>Homaxinella rudis</u>	w	w,sp,f	-	f	-	-
<u>Homaxinella</u> sp. A	sp,f	s,f	-	sp	f	-
<u>Homaxinella waltonsmithi</u>	w,sp,s,f	w,sp,f	-	-	-	-
<u>Phakellia folium</u>	-	w	-	-	-	-
<u>Phakellia lobata</u>	w	w,sp,f	-	-	f	-
<u>Pseudaxinella lunaecharta</u>	w	-	-	f	w,f	-
<u>Pseudaxinella</u> sp. A	-	w	-	-	-	-
<u>Pseudaxinella</u> sp. B	sp	sp	-	-	-	-
<u>Pseudaxinella wilsoni</u>	-	sp	-	-	-	f
<u>Teichaxinella corrugata</u>	s	sp,f	-	-	w,f	-
<u>Teichaxinella morchella</u>	-	-	-	f	-	-
<u>Teichaxinella shoemakeri</u>	-	-	-	-	f	f
undetermined Axinellidae	sp	w,sp,s,f	-	-	-	-
Demoxiidae						
<u>Higginsia strigilata</u>	w	w,sp,s,f	f	sp	w	-
Euryponidae						
<u>Tricheurypon viride</u>	sp,s	-	-	-	-	-
Hemiasterellidae						
undetermined Hemiasterellidae	-	-	sp	-	-	-

Appendix 1. (Continued)

	IS01	IS02	MS02	MS06	OS01	OS06
Raspaliidae						
<u>Ectyoplasia ferox</u>	sp	-	-	-	-	-
<u>Hemectyon pearsei</u>	sp	w,sp,f	f	-	-	-
<u>Raspaciona</u> sp. A	-	sp	-	-	-	-
Hadromerida						
Clionidae						
<u>Cliona caribbaea</u>	w	w,sp,f	sp,f	f	w,s,f	-
Spirastrellidae						
<u>Anthosigmella varians</u>	w	w,sp	-	-	-	f
<u>Columnitis</u> sp.	w	w,sp,f	-	-	-	-
<u>Spheciospongia othella</u> (?)	sp	-	-	-	-	-
<u>Spheciospongia vesparium</u>	sp,f	w,sp	-	sp,f	-	s
<u>Spirastrella coccinea</u>	w	w,sp,s,f	-	f	-	-
<u>Timea mixta</u>	w	f	-	-	-	-
Tethyidae						
<u>Tethya crypta</u>	w	w	-	-	sp	-
Choristida - Astrophorina						
Ancorinidae						
<u>Myriastria fibrosa</u>	s	w,f	-	sp	-	-
<u>Myriastria incrustata</u>	-	-	-	-	f	-
<u>Myriastria kallitetilla</u>	w	sp	-	sp	f	-
<u>Myriastria</u> sp. A	w	-	-	-	-	-
Geodiidae						
<u>Erylus</u> sp.	-	-	-	-	-	f
<u>Geodia gibberosa</u>	s	w,sp,s,f	f	w,f	w	-
<u>Stelletta grubii</u>	-	-	f	-	f	-
<u>Stelletta</u> sp. A	-	-	-	-	w,s,f	-
Choristida - Spirophorina						
Craniellidae						
<u>Cinachyra alloclada</u>	w	w,sp,f	-	f	w	-
<u>Cinachyra kuekenthali</u>	w	w,sp,f	f	f	f	-
<u>Cinachyra rhizophyta</u>	w	w,sp,f	-	-	-	-
Choristida - Chondrosiina						
Chondrillidae						
<u>Chondrilla nucula</u>	-	sp	-	-	-	-
Chondrosiidae						
<u>Chondrosia collectrix</u>	w	sp,f	-	-	-	-
Calcarea						
Leucettidae						
<u>Leucetta</u> sp. A	-	-	-	sp	-	-

Appendix 1. (Continued)

	ISO1	ISO2	MSO2	MSO6	OSO1	OSO6
<u>Leucetta floridana</u>	-	-	-	-	-	f
Leuconiidae						
<u>Leuconia aspera</u>	w,s	sp	w,sp	w,f	s	f
Calcarea - Calcinea						
Clathrinida						
Clathrinidae						
<u>Clathria</u> sp.	-	w	-	-	w,f	-
<u>Clathrina coriacea</u>	s,f	-	sp	-	-	f
Calcarea - Calcaronea						
Scyettida						
Scyettidae						
<u>Scypha barbadensis</u>	s,f	f	w,sp	f	-	f
Cnidaria						
Hydrozoa						
Anthomedusa/Athecata						
Bougainvilliidae						
<u>Bimeria humilis</u>	f	s,f	s,f	f	s	f
<u>Bougainvillia</u> sp.	-	w	-	-	sp,s	sp
Clavidae						
<u>Turritopsis fascicularis</u>	-	s	-	sp,s	-	-
<u>Turritopsis nutricula</u>	s,f	-	s	sp,f	sp	sp
Eudendriidae						
<u>Eudendrium carneum</u>	w,s,f	sp,s,f	w,sp	w,sp,s,f	w,f	-
<u>Eudendrium</u> sp.	s,f	w,sp,s,f	s,f	-	sp,s,f	s,f
<u>Eudendrium tenellum</u>	-	s	-	s,f	w,sp,s	sp,f
Hydractiniidae						
<u>Clavactinia</u> sp.	sp	-	-	-	sp	-
<u>Hydractiniidae</u> B	-	-	sp	-	-	-
Leptomedusae/Thecata						
Campanulariidae						
<u>Campanularia hincksii</u>	-	w	w,sp,s,f	w,sp,f	w,sp,f	sp,s,f
<u>Clytia cylindrica</u>	s,f	w,s	w,sp,f	w,f	w,sp,s,f	sp,s,f
<u>Clytia fragilis</u>	-	w,sp,s	w,sp,s,f	w,f	w,sp,s	sp
<u>Obelia dichotoma</u>	w,sp,s,f	w,sp,s,f	w,sp,s,f	w,sp,s,f	w,sp,s	sp,s,f
Haleciidae						
<u>Halecium diminutivum</u>	s	-	-	-	-	-
<u>Halecium dysymmetrum</u>	w,sp,s	w,sp,s	-	w,sp	sp,s	sp,f
<u>Halecium</u> sp.	w,sp,s,f	w,sp,s,f	sp,s	sp,s,f	w,sp,s,f	sp,f
<u>Halecium tenellum</u>	w,sp	w,f	w,sp,s	w,sp,s	w,sp,s,f	sp,f

Appendix 1. (Continued)

	IS01	IS02	MS02	MS06	OS01	OS06
Lafoeidae						
<u>Filellum serratum</u>	-	w	s	sp	w,sp,s,f	sp,f
<u>Hebella scandens</u>	-	w	w,s,f	-	w,sp,f	sp,s,f
<u>Hebella venusta</u>	-	w	w,sp,s,f	sp,s,f	w,sp,f	sp
<u>Lafoea fruticulosa</u>	-	-	-	-	-	sp,f
<u>Scandia mutabilis</u>	sp	w,f	w,sp,s,f	f	w,f	sp,f
Laodiceidae						
<u>Modeeria rotunda</u>	-	-	-	f	w	-
Plumulariidae						
<u>Aglaophenia allmani</u>	-	-	sp,f	sp	-	-
<u>Aglaophenia elongata</u>	sp	-	f	-	w,sp	sp,s,f
<u>Aglaophenia latecarinata</u>	sp,s,f	f	sp,s,f	w,sp,f	s	sp,s
<u>Aglaophenia trifida</u>	w,sp,s,f	w	w,sp,s,f	w,sp,f	w,sp,s,f	sp,s,f
<u>Antenella secundaria</u>	-	w	-	sp	s	sp,f
<u>Gymnangium sinuosum</u>	s	f	-	s	-	sp,s,f
<u>Halopteris clarkei</u>	f	w	-	-	w,sp,s,f	sp,s,f
<u>Monostaechas quadridens</u>	w,sp,s,f	w,sp,s,f	w,sp,s,f	w,sp,s,f	w,s,f	sp,s,f
<u>Nemertesia simplex</u>	-	-	-	f	sp	sp,s,f
<u>Plumularia margaretta</u>	-	-	-	f	-	-
<u>Plumularia setacea</u>	s	-	-	-	w,sp	s
<u>Schizotricha tenella</u>	-	-	-	sp	-	-
Sertulariidae						
<u>Dynamena cornicina</u>	w,sp,s,f	w,sp,s,f	w,sp,s,f	w,sp,s,f	w,s	s,f
<u>Dynamena dalmasi</u>	-	-	-	s	sp,s	sp,f
<u>Dynamena quadridentata</u>	w,sp	w,sp,s,f	w,sp,s,f	sp,s,f	w,s	sp,f
<u>Salacia desmoides</u>	f	w,s	w,sp	w,sp,s,f	w,sp,s,f	sp,s,f
<u>Sertularella areyi</u>	-	w	-	-	w,sp,s	sp,f
<u>Sertularella conica</u>	w,sp,s,f	-	w,sp,s,f	sp,s,f	w,s	sp
<u>Sertularella pinnigera</u>	f	-	f	f	-	-
<u>Sertularella sp. A</u>	w,s,f	-	w,f	sp,s,f	sp,s,f	sp,f
<u>Sertularia distans</u>	w,sp,f	-	sp,s	w,sp,s,f	f	-
<u>Sertularia marginata</u>	w,sp,s,f	w,s,f	w,sp,s,f	w,sp,s,f	s	sp,s
<u>Sertularia plumulifera</u>	w,sp,f	w,sp	sp,s	w,sp	w,sp,s	sp,f
<u>Thyroscyphus marginatus</u>	s,f	w,f	w,sp,s,f	w,sp,s,f	w,f	-
Syntheciidae						
<u>Hincksella cylindrica</u>	s	-	s,f	sp,f	-	sp,f
<u>Synthecium tubitheca</u>	-	-	w,sp,s,f	w,sp,s,f	f	sp,f
Anthozoa - Octorallia						
Telestacea						
Telestidae						
<u>Telesto fruticulosa</u>	w,s,f	w,sp,s,f	w,sp,s,f	sp,s	w,sp,s,f	sp,f
<u>Telesto sanguinea</u>	w,s	w,sp,s,f	w,sp,s	w,f	w,sp,s,f	sp,s,f

Appendix 1. (Continued)

	IS01	IS02	MS02	MS06	OS01	OS06
Alcyonacea						
Nidaliidae						
<u>Nidalia occidentalis</u>	-	-	-	-	-	s
Gorgonacea						
Anthothelidae						
<u>Diodogorgia</u> sp.	-	-	-	-	w,f	s,f
<u>Titanideum frauenfeldii</u>	w,sp,s,f	w,sp,s,f	w,sp,f	sp,s,f	w,sp,s,f	sp,s,f
Ellisellidae						
<u>Ellisella</u> sp.	-	-	-	-	w,f	sp,s,f
Gorgoniidae						
<u>Leptogorgia virgulata</u>	w,sp,s,f	w,sp,s,f	f	w,sp,s,f	-	f
<u>Lophogorgia cardinalis</u>	-	-	f	-	-	sp,f
<u>Lophogorgia hebes</u>	w,sp,s,f	w,sp,s,f	w,sp	w,sp,s,f	sp	-
Paramuriceidae						
<u>Thesea</u> sp.	-	-	f	sp,f	w,sp,s,f	sp,s,f
Plexauridae						
<u>Muricea pendula</u>	-	f	-	-	w,sp	-
Pennatulacea						
Renillidae						
<u>Renilla reniformis</u>	-	-	-	-	w	-
Virgulariidae						
<u>Stylatula</u> sp.	-	-	-	-	w,s,f	sp,f
Anthozoa - Zoantharia						
Actiniaria						
undetermined Actiniaria	w,sp,f	w,sp,f	sp	w,sp	w,s	-
Scleractinia - Faviina						
Faviidae						
<u>Solenastria hyades</u>	w,f	-	-	-	-	-
Oculinidae						
<u>Oculina arbuscula</u>	-	w	-	-	-	-
<u>Oculina</u> sp.	w,sp	sp,f	f	-	sp,f	-
Rhizangiidae						
<u>Astrangia astreiformis</u>	-	-	-	s	-	-
<u>Phyllangia americana</u>	-	f	f	-	-	-
Scleractinia - Dendrophylliina						
Dendrophylliidae						
<u>Balanophyllia floridana</u>	-	-	-	-	w,s,f	f

Appendix 1. (Continued)

	IS01	IS02	MS02	MS06	OS01	OS06
Antipatharia						
undetermined Antipatharian	f	s	w,f	-	w,sp,f	-
Mollusca						
Gastropoda - Prosobranchia						
Archaeogastropoda						
Fissurellidae						
<u>Diodora cayenensis</u>	w	w,sp	-	w,sp	w,s,f	-
Trochidae						
<u>Calliostoma euglyptum</u>	-	sp	-	-	-	-
<u>Calliostoma roseolum</u>	-	-	w	-	-	-
Mesogastropoda						
Crepidulidae						
<u>Crepidula aculeata</u>	w,sp,s	w,sp,f	w,sp	-	-	-
<u>Crepidula plana</u>	-	-	-	-	-	sp
Cymatiidae						
<u>Cyphoma mcgintyi</u>	-	-	-	-	sp	-
<u>Distorsio constricta mcgintyi</u>	-	-	-	sp	-	sp,f
Cypraeidae						
<u>Cypraea spurca acicularis</u>	-	-	-	-	-	f
Eratoidae						
<u>Erato maugeriae</u>	-	-	-	-	s	-
Mytridae						
<u>Vexillum gemmatum</u>	-	-	-	w	-	-
Ovulidae						
<u>Simnia uniplicata</u>	w	-	-	-	-	-
Turritellidae						
<u>Vermicularia knorrii</u>	f	w,sp,s,f	-	sp,s,f	f	-
Velutinidae						
<u>Lamellaria perspicua</u>	-	-	-	-	s	-
Vermetidae						
<u>Spiroglyphus irregularis</u>	-	-	sp	-	-	-
Neogastropoda						
Buccinidae						
<u>Pisania tinctoria</u>	sp	w,sp,s,f	-	-	-	-
Columbellidae						
<u>Anachis avara</u>	-	w	-	-	w	-
<u>Anachis lafresnayi</u>	s	-	-	-	sp,f	sp
<u>Nassarina minor</u>	-	f	-	-	-	-
<u>Suturoglypta iontha</u>	-	f	-	s	-	-
Conidae						
<u>Conus floridanus</u>	-	-	-	-	-	sp

Appendix 1. (Continued)

	IS01	IS02	MS02	MS06	OS01	OS06
Fascioliariidae						
<u>Fasciolaria lilium hunteria</u>	s	sp	-	-	-	f
<u>Fasciolaria tulipa</u>	-	w,sp	-	-	-	-
<u>Pleuroploca gigantea</u>	-	-	-	s	-	-
Marginellidae						
<u>Marginella apicina</u>	-	-	-	-	w	-
Muricidae						
<u>Calotrophon ostrearum</u>	-	-	-	-	w	-
<u>Favartia cellulosa</u>	-	-	-	sp	-	-
<u>Murex florifer dilectus</u>	w	w,sp	-	-	w,f	-
<u>Murex pomum</u>	-	w	-	-	-	-
<u>Murex sp. B</u>	-	-	-	-	w	-
Nassariidae						
<u>Nassarius albus</u>	-	-	-	-	sp	-
Turridae						
<u>Kurtziella limonitella</u>	s	-	-	-	-	-
Pelecypoda						
Arcoida						
Arcidae						
<u>Anadara notabilis</u>	-	-	-	sp,f	-	f
<u>Arca imbricata</u>	w,sp,s,f	w,sp,s,f	w,sp,f	w,sp,s,f	f	s
<u>Arca zebra</u>	w,s	-	f	-	w,f	-
<u>Arcopsis adamsi</u>	-	-	-	-	w,s,f	s
<u>Barbatia candida</u>	-	-	f	-	s,f	f
Mytiloida						
Mytilidae						
<u>Lithophaga bisulcata</u>	sp	w,sp	-	-	f	-
<u>Modiolus americanus</u>	w,sp,s,f	w,sp,f	w,sp,f	w,sp,s,f	-	-
<u>Musculus lateralis</u>	f	-	-	-	-	-
Pterioida						
Anomiidae						
<u>Anomia squamula</u>	-	-	sp	-	-	-
<u>Pododesmus rudis</u>	-	f	sp	-	-	-
Pectinidae						
<u>Chlamys benedicti</u>	-	-	-	s	w,f	s
Plicatulidae						
<u>Plicatula gibbosa</u>	-	f	-	-	f	-
Pteriidae						
<u>Pteria colymbus</u>	w,sp,s,f	w,sp,s,f	w,sp,s	w,sp,f	s,f	-
Spondylidae						
<u>Spondylus americanus</u>	-	w	-	-	-	-

Appendix 1. (Continued)

	IS01	IS02	MS02	MS06	OS01	OS06
Ostreina						
Ostreidae						
<u>Lopha frons</u>	-	-	-	sp	-	-
<u>Ostrea permolis</u>	s	-	-	sp	-	-
Veneroida						
Cardiidae						
<u>Americardia media</u>	-	-	f	-	-	f
<u>Laevicardium pictum</u>	w	sp,f	-	-	-	-
Chamidae						
<u>Chama congregata</u>	w,s,f	w,sp,f	sp,f	w,sp,s	w,s,f	s
<u>Chama macerophylla</u>	-	w,f	-	-	-	-
<u>Chama sarda</u>	-	f	-	-	-	-
<u>Pseudochama radians</u>	w,sp	w,sp	sp	-	-	-
Tellinidae						
<u>Tellina americana</u>	-	w	-	-	-	-
Veneridae						
<u>Chione grus</u>	s,f	-	-	-	w	-
<u>Chione latilirata</u>	w	-	-	-	-	-
<u>Macrocallista maculata</u>	-	w	-	-	-	-
Myoida						
Corbulidae						
<u>Corbula chittyana</u>	-	-	-	-	-	f
Gastrochaenidae						
<u>Gastrochaena hians</u>	-	-	-	-	w,f	-
Hiatellidae						
<u>Hiatella arctica</u>	-	sp,s,f	sp	sp,s,f	w,s,f	-
Pholadomyoida						
Lyonsiidae						
<u>Lyonsia beana</u>	w,s,f	w,sp	w,sp	sp,s	-	-
Octopoda						
Octopodidae						
<u>Octopus joubini</u>	-	-	sp	-	f	-
<u>Octopus sp.</u>	sp	f	-	-	f	-
Arthropoda - Mandibulata						
Crustacea - Cirripedia						
Thoracia - Lepadomorpha						
Scalpellidae						
<u>Scalpellum diceratum</u>	-	-	-	-	w	-

Appendix 1. (Continued)

	IS01	IS02	MS02	MS06	OS01	OS06
Thoracia - Balanomorpha						
Archaeobalanidae						
<u>Conopea galeata</u>	w	s	-	-	s	-
<u>Conopea merrilli</u>	w,f	w,sp,s,f	w,sp,s,f	w,sp,s	w,sp,f	-
Balanidae						
<u>Balanus calidus</u>	sp,f	-	sp	s	sp,s	sp
<u>Balanus trigonus</u>	w,sp,s,f	w,sp,s,f	w,sp,s	w,sp,s,f	w,sp,s,f	sp,f
<u>Balanus venustus</u>	w,sp,f	w,s,f	w,sp	sp,f	w,sp	sp
Crustacea - Malacostraca						
Stomatopoda						
Gonodactylidae						
<u>Gonodactylus bredini</u>	-	w,s	w,sp	w	f	s
Decapoda - Pleocyemata - Caridea						
Alpheidae						
<u>Alpheus formosus</u>	-	-	sp	-	-	-
<u>Synalpheus fritzmuelleri</u>	sp	-	-	-	-	-
<u>Synalpheus longicarpus</u>	sp	s,f	-	sp	-	-
<u>Synalpheus minus</u>	sp,s,f	sp	sp,s	-	-	-
<u>Synalpheus townsendi</u>	-	w,sp,s,f	w,sp,s	sp,s	w,s,f	f
Decapoda - Pleocyemata - Palinura						
Scyllaridae						
<u>Scyllarus chacei</u>	-	-	-	-	f	-
<u>Scyllarus depressus</u>	-	-	-	-	-	f
Decapoda - Pleocyemata - Anomura						
Diogenidae						
<u>Dardanus fucosus</u>	-	-	sp	s	w	-
<u>Dardanus insignis</u>	-	-	-	-	w,sp,f	sp,f
<u>Paguristes sericeus</u>	-	sp	-	-	sp	-
<u>Paguristes tortugae</u>	sp	w,sp	-	-	-	-
<u>Petrochirus diogenes</u>	-	-	s	-	sp	-
Galatheididae						
<u>Galathea rostrata</u>	-	-	-	-	w,sp,s	s,f
Paguridae						
<u>Pagurus carolinensis</u>	-	sp	s	-	-	-
<u>Pylopagurus corallinus</u>	-	-	-	-	w	f
Porcellanidae						
<u>Megalobrachium soriatum</u>	sp	w	sp	w,sp	-	-
<u>Pachycheles rugimanus</u>	-	-	-	w,s	w,s,f	-
<u>Petrolisthes galathinus</u>	-	w	-	-	-	-
<u>Porcellana sigsbeiana</u>	-	-	-	-	sp	-

Appendix 1. (Continued)

	IS01	IS02	MS02	MS06	OS01	OS06
Decapoda - Pleocyemata - Brachyura						
Calappidae						
<u>Calappa angusta</u>	-	-	-	-	-	sp,f
<u>Osachila semilevis</u>	-	-	-	-	w	-
Cancridae						
<u>Cancer irroratus</u>	w	-	-	-	-	-
Dromiidae						
<u>Dromidia antillensis</u>	w,sp	sp,s	f	-	-	-
Grapsidae						
<u>Euchirograpsus americanus</u>	-	-	-	-	w,s,f	s
Majidae						
<u>Macrocoeloma camptocerum</u>	w	w,sp	sp	-	-	-
<u>Macrocoeloma trispinosum</u>	-	f	-	-	-	-
<u>Metaporaphis calcarata</u>	-	-	-	-	s	-
<u>Mithrax acuticornis</u>	-	-	f	w	w,s	s,f
<u>Mithrax forceps</u>	-	w,s	f	-	-	-
<u>Mithrax pleuracanthus</u>	w,sp,s,f	w,sp,f	w,sp,f	w,sp	-	-
<u>Pelia mutica</u>	-	-	sp	-	-	-
<u>Podochela gracilipes</u>	-	-	-	-	s	f
<u>Podochela riisei</u>	-	w,sp	w,sp	-	-	-
<u>Stenocionops furcata coelata</u>	f	-	f	-	w,s,f	s
<u>Stenorhynchus seticornis</u>	-	w	-	w	s	-
Parthenopidae						
<u>Parthenope fraterculus</u>	-	-	-	-	w,s,f	s,f
Pinnotheridae						
<u>Pinnotheres maculatus</u>	sp	w	-	-	-	-
<u>Pinnotheres ostreum</u>	-	w	-	w	-	-
Portunidae						
<u>Portunus ordwayi</u>	-	-	-	-	w,sp,f	-
Raninidae						
<u>Hypochonca sabulosa</u>	-	w,sp	-	-	f	-
<u>Symethis variolosa</u>	-	-	-	-	-	f
Xanthidae						
<u>Carpoporus papulosus</u>	-	-	-	-	-	f
<u>Glyptoxanthus erosus</u>	-	-	-	s	-	-
<u>Pilumnus dasypodus</u>	s	-	w,sp,s	s	-	-
<u>Pilumnus floridanus</u>	-	-	-	w,sp	s,f	-
<u>Pilumnus pannosus</u>	sp	w,sp,f	-	-	-	-
<u>Pilumnus sayi</u>	w,sp,s,f	w,sp,s,f	sp,s,f	w,sp,s	sp	-
<u>Pseudomedeus agassizii</u>	-	w,sp	sp,s,f	w,sp	sp,s	s,f
Xanthidae A	-	-	-	-	s	-

Appendix 1. (Continued)

	IS01	IS02	MS02	MS06	OS01	OS06
Bryozoa						
Stenolaemata						
Cyclostomata						
Crisiidae						
<u>Crisia</u> sp.	w,sp,f	w,sp,s,f	w,sp,s,f	w,sp,s,f	w,s	s,f
Diaperoeciidae						
<u>Diaperoecia</u> floridana	-	w	w,sp,f	w,sp,s,f	w,sp,s	sp,f
Lichenoporidae						
<u>Lichenopora</u> radiata	-	sp	s	w,sp,s,f	-	s,f
Oncousoeciidae						
<u>Proboscina</u> floridana	-	-	-	sp,s,t	-	sp,f
Plagioeciidae						
<u>Plagioecia</u> dispar	-	-	-	-	-	sp,s,f
Gymnolaemata						
Ctenostomata						
undetermined Ctenostomata	w,sp	w	sp,f	sp,s,f	w,sp	sp,s,f
Alcyonidiidae						
<u>Alcyonidium</u> sp.	s,f	w	w,sp	-	-	sp
Buskiidae						
<u>Buskia</u> sp.	w	w	w,sp,s	-	w	f
Nolleliidae						
<u>Nolella</u> stipata	w,sp	w	w,sp,s	-	w,s	-
Valkeriidae						
<u>Aeverrillia</u> setigera	-	sp	w,sp,s	w,sp,f	-	sp
Vesiculariidae						
<u>Amathia</u> alternata	-	w,sp,s,f	f	-	s	sp
<u>Amathia</u> convoluta	w	w	-	sp	w	sp,s
<u>Amathia</u> distans	-	w,s,f	w,sp,s,f	w,sp,s,f	w,s,f	sp,s,f
Victorellidae						
<u>Sundanella</u> sibogae	sp,s	f	s	-	-	-
Cheilostomata - Anasca						
Aeteidae						
<u>Aetea</u> anguina	w,sp	w,sp,f	w,sp,s	w,sp,s,f	sp	sp,s,f
Alderinidae						
<u>Alderina</u> smitti	w,sp	-	-	sp	w,s	sp,s,f
<u>Antropora</u> tincta	w,sp	w,sp,s,f	w,sp,s,f	sp,f	w,sp,s	sp,f
<u>Copidozoum</u> tenuirostre	-	-	-	sp,f	-	-
<u>Crassimarginatella</u> leucocypha	-	sp,f	-	-	-	-
<u>Parellisina</u> curvirostris	-	w	sp,f	w,sp,s,f	w,s	-
<u>Parellisina</u> latirostris	-	-	f	s,f	sp,f	sp,s,f

Appendix 1. (Continued)

	IS01	IS02	MS02	MS06	OS01	OS06
Bugulidae						
<u>Bugula fulva</u>	-	w	w,sp,s,f	w,sp,f	-	sp
<u>Bugula grayi</u>	-	-	f	f	-	-
<u>Bugula johnstonae</u>	-	-	-	-	-	sp,f
<u>Bugula microoecia</u>	-	w	sp	w,sp	-	-
<u>Bugula rylandi</u>	w,sp,s,f	w	w,sp,s,f	w,sp,f	-	s
<u>Bugula sp.</u>	-	-	-	f	-	-
<u>Bugula turrita</u>	-	-	sp	sp	-	-
<u>Caulibugula dendrograpta</u>	-	-	s,f	f	-	-
<u>Caulibugula pearsei</u>	sp	-	-	w,s	-	-
Cellariidae						
<u>Cellaria nodosa</u>	-	-	-	-	-	sp
Chaperiidae						
<u>Chaperia sp.</u>	s	-	w,sp,s,f	w,sp,s,f	w	-
Cupuladriidae						
<u>Cupuladria canariensis</u>	-	w	sp	w,sp,f	-	sp
<u>Cupuladria doma</u>	w	-	sp	w,sp,f	-	sp,f
<u>Discoporella umbellata</u>	-	-	-	w	f	sp
Farciminariidae						
<u>Nellia tenella</u>	-	-	f	s,f	-	sp,s,f
Hincksiniidae						
<u>Aplousina gigantea</u>	w	w,s	sp,s	sp	w,sp,s,f	sp,s,f
<u>Aplousina tuberosa</u>	-	-	f	w,f	f	s
Membraniporidae						
<u>Membranipora arborescens</u>	-	-	-	-	f	sp,f
<u>Membranipora tenuis</u>	w,sp,f	sp	s	-	-	-
Microporidae						
<u>Micropora coriacea</u>	-	-	-	-	-	sp,s,f
Onychocellidae						
<u>Floridina antiqua</u>	-	w,sp,s,f	-	sp	w,sp,s,f	sp
<u>Floridina parvicella</u>	-	s	s	sp	sp	sp,f
<u>Smittipora levinseni</u>	-	w,sp	sp,s	w,sp,f	w,sp,s,f	sp,s,f
Scrupocellariidae						
<u>Caberia boryi</u>	-	-	f	w,sp,s,f	-	sp,s
<u>Scrupocellaria bertholleti</u>	-	w	-	-	-	-
<u>Scrupocellaria regularis</u>	sp,s	w,f	sp,f	sp	w	sp,s,f
Steganoporellidae						
<u>Steganoporella magnilabris</u>	-	-	-	s,f	-	-
Cheilostomata - Cribrimorpha						
Cribrilinidae						
<u>Bellulopora bellula</u>	-	-	-	-	-	sp,s,f
<u>Cribrilaria floridana</u>	-	sp	s	w,sp,s,f	-	f
<u>Cribrilaria radiata</u>	w,sp	w,sp,s,f	sp,s	w,sp,s,f	w,sp	sp,s,f
<u>Membraniporella aragoi</u>	-	w	-	w	w,sp,f	-
undetermined Cribrilinidae	-	sp	-	-	-	-

Appendix 1. (Continued)

	IS01	IS02	MS02	MS06	OS01	OS06
Cheilostomata - Ascophora						
Adeonidae						
<u>Reptadeonella hastingsae</u>	w,sp,f	w,sp,s,f	w,sp,s,f	w,sp,s,f	w,sp,s,f	sp
Celleporariidae						
<u>Celleporaria albirostris</u>	-	w,f	w,sp,s,f	w,sp,s,f	w,sp,s,f	sp,s,f
<u>Celleporaria magnifica</u>	w,sp,s	w,sp,s,f	sp,s,f	w,sp,s,f	w,s	-
<u>Celleporaria mordax</u>	s,f	w,s,f	s	-	-	-
<u>Trematoecia turrita</u>	-	-	-	-	s	sp,f
Celleporinidae						
<u>Celleporina hassalli</u>	f	-	w,sp,s	-	-	-
<u>Turbicellepora dichotoma</u>	w,sp,s	w,sp,s	w,sp,s,f	w,sp,f	w,sp,s	sp
Cheiloporinidae						
<u>Hippaliosina rostrigera</u>	w,sp,s	w,sp,s,f	w,sp,s,f	w,sp,s,f	w,sp,f	-
Crepidacanthidae						
<u>Crepidacantha sp.</u>	-	-	-	-	-	sp,f
<u>Crepidacantha poissoni teres</u>	-	-	sp	w,sp	-	sp,s
Exochellidae						
<u>Exochella longirostris</u>	-	sp	s	sp	-	sp,s,f
<u>Triporula stellata</u>	-	-	-	-	-	sp,f
Hippoporinidae						
<u>Hippopleurifera mucronata</u>	-	-	s	-	-	-
<u>Hippoporidra calcarata</u>	-	-	-	-	w	sp
<u>Hippoporidra janthina</u>	-	-	-	-	-	sp
<u>Hippoporina contracta</u>	w,sp	w,sp,s,f	w,sp,s,f	-	w,f	sp,s,f
<u>Hippoporina verrilli</u>	-	f	s	-	sp	-
Hippothoidae						
<u>Tryposostega venusta</u>	w,sp,s,f	w,sp,s,f	sp,s,f	w,sp,f	w,sp,s,f	sp,s,f
Microporellidae						
<u>Microporella ciliata</u>	w,sp,f	w,sp,s,f	w,sp,s,f	w,sp,s,f	w,sp,s,f	sp,s,f
<u>Microporella marsupiata</u>	-	-	-	sp,f	-	sp,s
<u>Microporella umbracula</u>	w,sp,s	w,sp,s,f	sp,s,f	w,sp,s,f	w	sp,s,f
Petraliidae						
<u>Petraliella bisinuata</u>	-	-	w,sp,s,f	w,sp,s,f	f	sp,f
<u>Petraliella marginata</u>	-	-	-	-	f	f
Phylactellidae						
<u>Phylactella aviculifera</u>	-	sp	s	f	-	f
Schizoporellidae						
<u>Arthropoma cecili</u>	-	-	-	w,sp	-	-
<u>Cleidochasma porcellanum</u>	w	w,sp,s	s	w,sp,s,f	sp,s,f	sp,s,f
<u>Cycloperiella rubra</u>	-	w	-	-	w,sp,s,f	sp,s,f
<u>Escharina porosa</u>	-	-	-	sp	s	sp,s
<u>Schizoporella cornuta</u>	w,sp,s,f	w,sp,s,f	w,sp,s,f	w,sp,f	w,sp,s	sp,s,f
<u>Schizoporella errata</u>	-	-	w,sp,f	w,f	-	-
<u>Schizoporella floridana</u>	sp	-	w,sp,s,f	w,sp,s,f	-	sp
<u>Stylopoma informata</u>	-	w,f	s,f	w,sp,f	w,sp,s,f	sp,s,f

Appendix 1. (Continued)

	IS01	IS02	MS02	MS06	OS01	OS06
Sertellidae						
<u>Rhyncozoon rostratum</u>	-	-	sp	-	-	-
<u>Rhyncozoon tuberculatum</u>	-	sp	-	sp	-	-
Smittinidae						
<u>Parasmittina nitida</u>	-	-	s	-	w	s
<u>Parasmittina spathulata</u>	w	w,sp,s,f	w,sp,s,f	w,sp,s,f	w,sp,s,f	sp,s,f
<u>Smittina smittiella</u>	-	w	sp	w,f	s	-
Stomachetosellidae						
<u>Cigclisula serrulata</u>	-	-	-	-	-	sp,f
Echinodermata						
Crinoidea						
Comasteridae						
<u>Comactinia meridionalis</u>	-	-	sp,s,f	-	-	-
Stelleroidea - Asteroidea						
Platyasterida						
Luidiidae						
<u>Luidia alternata</u>	w	w,sp,f	-	-	-	-
<u>Luidia bernasconiae</u>	-	-	-	-	f	-
<u>Luidia clathrata</u>	-	-	w	sp	-	-
Paxillosida						
Astropectinidae						
<u>Astropecten articulatus</u>	-	-	s	-	w,sp	-
<u>Astropecten comptus</u>	-	-	-	-	w,s	-
<u>Astropecten duplicatus</u>	s	-	-	f	-	-
<u>Astropecten nitidus</u>	-	-	w	-	-	-
<u>Astropecten sp.</u>	-	sp,f	sp	f	sp,s	-
Valvatida						
Ophidiasteridae						
<u>Narcissia trigonaria</u>	-	-	-	-	-	s,f
Spinulosida						
Echinasteridae						
<u>Echinaster brasiliensis</u>	-	-	-	-	sp	-
<u>Echinaster modestus</u>	-	s	-	-	-	-
<u>Echinaster serpentarius</u>	-	w	-	-	-	-
<u>Echinaster sp.</u>	w,s,f	w,sp,f	sp,f	-	sp	-

Appendix 1. (Continued)

	IS01	IS02	MS02	MS06	OS01	OS06
Forcipulata						
Asteriidae						
<u>Asterias forbesii</u>	w,sp,s,f	sp	-	-	-	-
Phyronophiurida - Euryalina						
Gorgonocephalidae						
<u>Astrophyton muricatum</u>	-	-	f	w	-	-
<u>Astroporpa annulata</u>	-	-	-	-	w,sp,s,f	sp,s,f
Ophiurida						
Goniasteridae						
<u>Tosia parva</u>	-	-	-	-	-	f
Gorgonocephalidae						
<u>Astrocyclus caecilia</u>	-	-	-	-	sp	-
Ophiactidae						
<u>Ophiactis savignyi</u>	-	-	-	-	-	s,f
Ophiochitonidae						
<u>Ophionereis reticulata</u>	-	-	-	f	-	-
Ophiodermatidae						
<u>Ophioderma brevispinum</u>	-	-	f	-	-	f
Ophiothricidae						
<u>Ophiothrix angulata</u>	w,sp,s,f	sp,s,f	sp,s,f	s,f	w,sp,s,f	s,f
<u>Ophiothrix suensonii</u>	-	-	-	-	f	f
Echinoidea						
Cidaroida						
Cidaridae						
<u>Eucidaris tribuloides</u>	-	-	-	sp	w,sp,f	s
Arbacoida						
Arbaciidae						
<u>Arbacia punctulata</u>	w,sp,s	w,sp,s,f	f	-	w,f	s
Temnopleuroida						
Toxopneustidae						
<u>Lytechinus variegatus</u>	w,sp,s,f	w,sp,f	-	-	-	-
Clypeasteroida						
Clypeasteridae						
<u>Clypeaster prostratus</u>	-	sp	-	-	-	-
<u>Clypeaster subdepressus</u>	-	s	w,s	sp	-	f
Mellitidae						
<u>Encope michelini</u>	sp,s	w,sp,f	-	-	-	-
<u>Leodia sexiesperforata</u>	s	sp	f	sp,f	-	-
Cassiduloida						
Echinolampadidae						
<u>Echinolampas depressa</u>	-	-	-	-	-	s

Appendix 1. (Continued)

	IS01	IS02	MS02	MS06	OS01	OS06
Spatangoida						
undetermined Spatangonidae	-	-	-	-	-	f
Holothuroidea						
Dendrochirotida						
Cumariidae						
<u>Ocnus pygmaeus</u>	w,sp,s,f	w,sp,f	sp	sp	-	s
Aspidochirotida						
Synallactidae						
<u>Isostichopus badinotus</u>	-	-	f	-	-	-
Chordata						
Ascidiacea						
Ascidiacea A	-	w,f	-	-	-	-
undetermined Ascidiacea	-	-	w	-	-	-
Enterogona						
Asciidiidae						
<u>Ascidia curvata</u>	f	-	sp,s,f	-	-	-
Clavelinidae						
<u>Clavelina gigantea</u>	s	-	sp,f	-	-	-
<u>Clavelina picta</u>	sp	w	w	-	-	-
<u>Clavelina</u> sp.	w,sp,s,f	sp,s,f	sp,f	sp,s	f	-
<u>Distaplia bermudensis</u>	w,sp,s,f	w,sp,s,f	sp	-	f	s
<u>Eudistoma carolinense</u>	sp,s	-	sp,s	f	w,sp,s,f	-
<u>Eudistoma hepaticum</u>	-	sp,s	-	-	-	-
<u>Eudistoma</u> sp.	w,sp,s,f	-	sp,s,f	sp	s,f	f
Didemnidae						
<u>Didemnum candidum</u>	w,sp,s,f	w,sp,f	w,sp,f	sp,s	s,f	s,f
<u>Didemnum</u> sp.	-	-	f	-	-	-
<u>Didemnum speciosum</u>	s,f	sp	-	-	f	-
<u>Diplosoma macdonaldi</u>	w,sp,s,f	-	sp	f	s,f	-
<u>Tridemnum savignii</u>	sp	s	-	-	f	s
<u>Tridemnum</u> sp.	w,s	sp,f	f	-	s,f	s,f
Polyclinidae						
<u>Aplidium constellatum</u>	-	w	w	-	-	-
<u>Aplidium pellucidum</u>	w,sp	-	-	-	w	s
<u>Aplidium</u> sp. A	sp	-	-	-	-	-
<u>Aplidium</u> sp. B	sp	w	-	-	-	-
<u>Aplidium stellatum</u>	w	-	-	-	-	-

Appendix 1. (Continued)

	IS01	IS02	MS02	MS06	OS01	OS06
Pleurogona						
Molgulidae						
<u>Molgula occidentalis</u>	w,sp,s,f	w,sp,s,f	w,sp,f	w,sp,s,f	-	-
Styelidae						
<u>Botrylloides nigrum</u>	s	-	sp	-	f	-
<u>Botryllus planus</u>	sp	sp	-	-	f	-
<u>Polyandrocarpa floridana</u>	w	-	s	sp,s	-	-
<u>Polyandrocarpa gravei</u>	-	w,sp	f	-	-	s
<u>Polyandrocarpa tinctoria</u>	-	-	sp	-	-	-
<u>Styela partita</u>	w,sp,f	sp,s,f	sp	-	-	-
<u>Styela plicata</u>	w,sp,s,f	w,sp,s,f	w,sp,s	w,sp,f	-	-
<u>Symplegma viride</u>	f	f	-	f	w,sp	-

Appendix 2. Phylogenetic list of invertebrate taxa collected by trawl during 1981 cruises to the stations off South Carolina and Georgia.
w = winter, sp = spring, s = summer and f = fall. No trawling was done at station OS06 during any season.

	IS01	IS02	MS02	MS06	OS01
Porifera					
Demospongea - Ceractinomorpha					
Keratosa					
Keratosa D	-	-	-	-	f
Aplysillidae					
<u>Chelonaplysilla</u> sp.	-	sp,s	-	s	sp,f
Darwinellidae					
<u>Darwinella</u> sp.	s,f	w,f	w,sp,f	s	-
Dysideidae					
<u>Dysidea variabilis</u> (?)	w	-	-	-	-
<u>Polyfibrospongia echina</u>	w	-	-	-	-
Spongiidae					
<u>Aplysina fistularis</u>	w	w,sp,s,f	w,sp,s,f	w,sp,s	sp
<u>Aplysina lacunosa</u>	-	-	-	s	-
<u>Euspongia officinalis</u>	-	w	-	-	-
<u>Hyatella intestinalis</u>	-	f	-	-	-
<u>Ircinia campana</u>	f	w,sp,s,f	w,sp,s,f	w,sp,f	sp,s,f
<u>Ircinia felix</u>	w	w,sp,s,f	w,sp	sp,s	w,s
<u>Ircinia ramosa</u>	-	w,sp,s,f	w,sp,s,f	w,sp,f	w,s
<u>Ircinia strobilina</u>	-	w,s,f	w,sp,s,f	w,sp,f	sp,f
<u>Spongia tubulifera</u>	s	-	-	sp	-
Haplosclerida					
Adociidae					
<u>Sigmadocia caerulea</u>	w	-	-	s	w,sp,s,f
<u>Strongylophora</u> sp. B	-	f	f	w,sp,s,f	-
Desmacidonidae					
<u>Xytopsues griseus</u>	w,sp,s	sp	-	s	w,sp,s
<u>Xytopsues</u> sp.	-	-	-	-	s
Haliclonidae					
<u>Amphimedon compressa</u>	-	s	-	sp,s,f	s
<u>Callyspongia fallax</u>	s	w,sp	sp	w,sp,f	-
<u>Callyspongia vaginalis</u>	-	-	-	sp	-
<u>Haliclona areolata</u>	-	s	-	w,f	-
<u>Haliclona oculata</u>	w,sp,s,f	w,sp,s,f	w,sp	w,sp,s,f	-
<u>Haliclona palmata</u>	-	w	-	-	-
<u>Haliclona</u> sp. C	sp	-	w	w	w,sp,s
<u>Niphantes digitalis</u>	-	w	-	-	-
<u>Niphantes erecta</u>	-	-	-	s	-
Nepheliospongiidae					
<u>Xestospongia halichondrioides</u>	-	-	-	s,f	s,f
Poecilosclerida					
Poecilosclerida H	-	w	-	-	w,sp
Amphilectidae					
<u>Tenaciella obliqua</u>	w	-	-	f	-

Appendix 2. (Continued)

	IS01	IS02	MS02	MS06	OS01
Microcionidae					
<u>Microciona prolifera</u>	s	-	w,f	s,f	s,f
<u>Microciona</u> sp.	-	-	-	s	-
<u>Microciona spinosa</u>	s	-	w	s	w,sp
<u>Pandaros acanthifolium</u>	w,sp,s	sp	w	sp,s,f	s
<u>Thalyseurypon foliacea</u>	-	s,f	-	-	-
<u>Thalysias juniperina</u>	s	w,sp,s,f	w,sp,s	w,sp,s,f	sp,s
<u>Thalysias</u> sp.	-	-	-	-	f
Mycalidae					
<u>Mycale</u> sp.	-	-	-	-	s
<u>Mycale</u> sp. A	-	-	-	-	sp
<u>Neofibularia nolitangere</u>	-	-	sp	w,sp,s,f	sp
<u>Zygomycale parishii</u>	-	-	-	f	-
Plocamiidae					
<u>Holoplocamia</u> sp.	-	-	-	sp	sp
Tedaniidae					
<u>Acarnus innominatus</u>	-	-	-	s	-
<u>Iotrochota birotulata</u>	-	-	sp	sp	w,f
<u>Lissodendoryx isodictyalis</u>	w	sp,s,f	w,sp,s,f	w,sp,s,f	-
<u>Lissodendoryx sigmata</u>	s,f	sp,f	sp,s,f	w,sp,s,f	s
<u>Lissodendoryx</u> sp. B	-	w	-	-	-
<u>Tedania ignis</u>	-	w,sp,s	-	-	-
Halichondriida					
Halichondriidae					
<u>Ciocalapata gibbsi</u>	w,s,f	w,sp,s,f	-	s,f	sp,s,f
<u>Halichondria bowerbanki</u>	sp	w,sp,s	-	s	sp,s
<u>Halichondria</u> sp.	-	-	w	-	-
Demospongea - Tetractinomorpha					
Axinellida					
Axinellidae					
<u>Axinella bookhouti</u>	sp,f	w,sp,s	-	s,f	sp,s,f
<u>Homaxinella rudis</u>	-	w,s	w	sp,f	-
<u>Homaxinella</u> sp. A	w	w,sp,s,f	w,s,f	sp,s,f	f
<u>Homaxinella waltonsmithi</u>	w,sp,s,f	w,sp,s,f	s	sp,s,f	w
<u>Phakellia lobata</u>	w	w,s,f	w	-	-
<u>Pseudaxinella linaechara</u>	f	s,f	-	-	s
<u>Pseudaxinella</u> sp. A	w	w	-	-	-
<u>Pseudaxinella</u> sp. B	s	sp	-	-	-
<u>Pseudaxinella wilsoni</u>	s	-	-	-	f
<u>Teichaxinella corrugata</u>	s	sp,f	-	s	w,sp,s,f
<u>Teichaxinella morchella</u>	-	-	-	-	s
undetermined Axinellidae	-	s,f	-	-	-
Demoxiidae					
<u>Higginsia strigilata</u>	s,f	w,sp,s,f	-	sp,s,f	w,f

Appendix 2. (Continued)

	IS01	IS02	MS02	MS06	OS01
Raspaliidae					
<u>Ectyoplasia ferox</u>	-	s	-	-	-
<u>Ectyoplasia</u> sp. A	w	w	-	-	-
<u>Hemectyon pearsei</u>	w,s	w,sp,s,f	-	-	sp
<u>Raspaciona</u> sp. A	w	s	-	-	-
<u>Raspaciona</u> sp. B	-	f	-	-	-
Hadromerida					
Clionidae					
<u>Cliona caribbaea</u>	w,sp,s	w,sp,s,f	w,sp,s,f	-	w,sp,s
<u>Cliona celata</u>	-	-	-	-	s
<u>Cliona lampa</u>	-	-	-	sp	-
<u>Cliona</u> sp.	-	-	-	sp	-
Coppatidae					
<u>Spirasigma</u> sp. (?)	-	-	-	sp	-
Placospongiidae					
<u>Placospongia carinata</u>	-	-	-	-	sp
Podospongiidae					
<u>Podospongia</u> sp.	-	-	-	s	-
Spirastrellidae					
<u>Anthosigmella varians</u>	w,sp,f	sp,f	-	w,s,f	sp
<u>Spheciospongia othella</u> (?)	f	-	-	-	-
<u>Spheciospongia vesparium</u>	w,sp,s,f	w,sp,s,f	w,sp,s,f	s,f	-
<u>Spirastrella coccinea</u>	w,s	w,sp,s,f	-	sp	-
<u>Timea mixta</u>	w	w,sp	-	w,sp	-
Suberitidae					
<u>Polymastia</u> sp.	-	f	-	-	-
undetermined Suberitidae	-	w	-	-	-
Tethyidae					
<u>Aaptos aaptos</u>	-	f	-	-	f
<u>Tethya actinia</u>	-	-	s	-	-
<u>Tethya crypta</u>	-	-	sp	-	-
Choristida - Astrophorina					
Ancorinidae					
<u>Myriastria fibrosa</u>	s	s,f	w	w,s,f	w,f
<u>Myriastria incrustata</u>	-	f	sp	-	-
<u>Myriastria kallitetilla</u>	s	-	-	f	w,sp,s
<u>Myriastria</u> sp. A	-	-	-	-	sp
Geodiidae					
<u>Geodia gibberosa</u>	-	w,sp,s,f	sp	w,sp,s,f	w,sp
<u>Stelletta grubii</u>	-	-	s	-	s
<u>Stelletta</u> sp. A	-	-	-	-	w,s,f

Appendix 2. (Continued)

	IS01	IS02	MS02	MS06	OS01
Choristida - Spirophorina					
Cranelliidae					
<u>Cinachyra alloclada</u>	w,sp,s	w,sp,s,f	w	s,f	s
<u>Cinachyra kukenthali</u>	w,s,f	w,s,f	w,sp,s	w,sp,s,f	-
<u>Cinachyra rhizophyta</u>	w,s	s	-	-	-
Choristida - Chondrosiina					
Chondrillidae					
<u>Chondrilla nucula</u>	-	w	w,sp,f	-	-
Chondrosiidae					
<u>Chondrosia collectrix</u>	w	-	w,sp	sp	-
Lithistida					
Scleritodermidae					
<u>Scleritoderma paccardi</u> (?)	-	-	-	-	sp
Calcarea					
Leucettidae					
<u>Leucetta</u> sp. A	-	-	sp	sp	-
<u>Leucetta floridana</u>	-	-	-	sp	-
Leuconiidae					
<u>Leuconia aspera</u>	w,sp,s	-	w,sp	sp,f	sp
<u>Leuconia crustacea</u>	w	-	-	-	-
Calcarea - Calcinea					
Clathrinida					
Clathrinidae					
<u>Clathria</u> sp. A	-	s	-	-	-
<u>Clathrina coriacea</u>	w,sp,s,f	sp,s,f	w,sp,s	-	w,sp
<u>Leucosolenia canariensis</u>	-	-	-	-	w
<u>Leucosolenia cancellata</u>	s	-	-	-	-
Calcarea - Calcaronea					
Scyettida					
Scyettidae					
<u>Scypha barbadensis</u>	w,sp,s,f	sp,s,f	sp	w,sp	sp
<u>Scypha</u> sp. A	sp	-	-	-	-
Cnidaria					
Scyphozoa					
Semaestomeae					
Cyaneidae					
<u>Cyanea capillata</u>	-	w	w	-	-

Appendix 2. (Continued)

	IS01	IS02	MS02	MS06	OS01
Ulmaridae					
<u>Aurelia aurita</u>	f	-	-	-	-
Rhizostomeae					
Stomolophidae					
<u>Stomolophus meleagris</u>	-	w,s,f	-	-	-
Cubozoa					
Carybdeidae					
<u>Tamoya haplonema</u>	-	f	-	-	s,f
Hydrozoa					
Anthomedusa/Athecata					
Bougainvilliidae					
<u>Bimeria humilis</u>	s,f	s,f	s,f	w,sp,s,f	sp,s,f
<u>Bougainvillia</u> sp.	f	w,s,f	f	w,sp,s	w,sp,s,f
Clavidae					
<u>Turritopsis fascicularis</u>	-	s	-	w,sp	w
<u>Turritopsis nutricula</u>	f	sp	s,f	w,s,f	w,sp,f
Eudendriidae					
<u>Eudendrium carneum</u>	sp,s,f	w,sp,s,f	w,sp,s,f	w,sp,s,f	w,s,f
<u>Eudendrium</u> sp.	sp,s,f	w,sp,s,f	w,s,f	w,sp,s,f	sp,s,f
<u>Eudendrium tenellum</u>	f	s,f	s	w,sp,f	w,sp,s,f
Hydractiniidae					
<u>Clavactinia</u> sp.	sp	-	-	-	-
Hydractiniidae A	-	-	-	-	w
Tubulariidae					
<u>Ectopleura</u> sp.	-	-	-	-	w,sp
Leptomedusae/Thecata					
Campanulariidae					
<u>Campanularia hincksii</u>	sp	-	w,sp,s,f	w,sp,s,f	w,sp,s,f
<u>Clytia cylindrica</u>	w,sp,f	s,f	w,sp,s,f	w,sp,s,f	w,sp,s,f
<u>Clytia edwardsi</u>	-	s	-	sp	s
<u>Clytia fragilis</u>	w,s	w,s,f	w,sp,s,f	w,sp,f	w,sp,s,f
<u>Clytia paulensis</u>	-	-	-	-	w,sp
<u>Obelia dichotoma</u>	w,sp,s,f	w,sp,s,f	w,sp,s,f	w,sp,s,f	w,sp,s,f
Campulinidae					
<u>Cuspidella humilis</u>	-	-	w,sp	-	-
Haleciidae					
<u>Halecium diminutivum</u>	f	-	-	-	-
<u>Halecium dysymmetrum</u>	w,s	w,sp,s,f	sp,s,f	w,sp,s	sp,s
<u>Halecium</u> sp.	w,sp,s,f	w,sp,s,f	w,sp,s,f	w,sp,s,f	w,sp,s,f
<u>Halecium tenellum</u>	f	w	w,sp,s,f	w,sp,s,f	w,sp,s,f

Appendix 2. (Continued)

	IS01	IS02	MS02	MS06	OS01
Lafoeidae					
<u>Filellum serratum</u>	-	-	w,sp,s	w,sp,s	w,sp,s,f
<u>Hebella scandens</u>	sp,s,f	s	w,sp,s,f	w,s,f	w,sp,f
<u>Hebella venusta</u>	f	-	w,sp,s,f	w,sp,s,f	w,sp,s,f
<u>Scandia mutabilis</u>	w,sp,s,f	s,f	w,sp,s,f	w,sp,s,f	w,s,f
Laodiceidae					
<u>Modeeria rotunda</u>	-	-	-	-	s
Plumulariidae					
<u>Aglaophenia allmani</u>	s	s	w,sp,s	w,sp,s,f	w,sp,s,f
<u>Aglaophenia elongata</u>	sp,f	s	s	sp,s	w,sp,s,f
<u>Aglaophenia latecarinata</u>	s,f	sp,s,f	sp,s,f	w,sp,s,f	w,s,f
<u>Aglaophenia trifida</u>	w,sp,s,f	w,sp,s,f	w,sp,s,f	w,sp,s,f	w,sp,s,f
<u>Antenella secundaria</u>	-	-	-	sp	w,sp,f
<u>Cladocarpus flexilis</u>	-	-	-	f	w,sp,s
<u>Gymnangium sinuosum</u>	s,f	s,f	f	-	-
<u>Halopteris clarkei</u>	-	s	-	-	sp,s,f
<u>Halopteris diaphana</u>	-	-	-	sp	-
<u>Monostaechas quadridens</u>	w,sp,s,f	w,sp,s,f	w,sp,s,f	w,sp,s,f	w,sp,s,f
<u>Nemertesia simplex</u>	-	-	-	f	w,s,f
<u>Plumularia margaretta</u>	-	f	-	w	-
<u>Plumularia setacea</u>	s,f	-	s	w,sp,s,f	w,s,f
Sertulariidae					
<u>Diphasia tropica</u>	w	-	-	w	-
<u>Dynamena cornicina</u>	w,sp,s,f	w,sp,s,f	w,sp,s,f	w,sp,s,f	w,s,f
<u>Dynamena dalmasi</u>	-	w	w,s	w,sp,s,f	w,sp,s,f
<u>Dynamena quadridentata</u>	w,sp,s,f	w,sp,s,f	w,sp,s,f	w,sp,s,f	w,s,f
<u>Salacia desmoides</u>	sp,s	w,sp,s,f	w,sp,s,f	w,sp,s,f	w,sp,s,f
<u>Sertularella areyi</u>	-	-	sp	-	w,sp,s,f
<u>Sertularella conica</u>	w,sp,s,f	w,sp,s,f	w,sp,s,f	w,sp,s,f	w,sp
<u>Sertularella pinnigera</u>	-	-	s,f	f	-
<u>Sertularella sp. A</u>	sp,s	f	sp,s	w,sp,s,f	w,sp,s,f
<u>Sertularia distans</u>	w,sp,s,f	w,sp,s,f	w,sp	w,sp,s	s
<u>Sertularia marginata</u>	w,sp,s,f	w,sp,s,f	w,sp,s,f	w,sp,s,f	w,sp,s,f
<u>Sertularia plumulifera</u>	w,sp,s,f	w,sp,s	w,sp,s,f	w,sp,s	w,sp,s,f
<u>Thyroscyphus marginatus</u>	s,f	s	w,sp,s,f	w,sp,s,f	w,s,f
<u>Thyroscyphus ramosus</u>	-	-	-	f	-
Syntheciidae					
<u>Hincksella cylindrica</u>	s,f	-	s,f	w,sp,s,f	-
<u>Synthecium tubithec</u>	s	-	w,sp,s,f	w,sp,s,f	w,sp,s
Anthozoa - Octocorallia					
Telestacea					
Telestidae					
<u>Telesto fruticulosa</u>	w,sp,s,f	w,sp,s,f	w,sp,s,f	sp,s,f	w,sp,s,f
<u>Telesto riisei</u>	-	-	-	-	w,sp,s
<u>Telesto sanguinea</u>	w,sp,s,f	w,sp,s,f	sp,s,f	sp,s	w,sp,s,f

Appendix 2. (Continued)

	IS01	IS02	MS02	MS06	OS01
Gorgonacea					
Anthothelidae					
<u>Diodogorgia</u> sp.	-	-	-	f	w,s,f
<u>Titanicum rrauenfeldii</u>	w,sp,s,f	w,sp,s,f	w,sp,s,f	w,sp,s,f	w,sp,s,f
Ellisellidae					
<u>Ellisella</u> sp.	-	-	-	-	w,sp,s,f
Gorgoniidae					
<u>Leptogorgia setacea</u>	-	-	-	-	w,sp,s,f
<u>Leptogorgia</u> sp. A	-	-	-	sp	-
<u>Leptogorgia virgulata</u>	w,sp,s,f	w,sp,s,f	w,sp,s,f	w,sp,s,f	f
<u>Lophogorgia hebes</u>	w,sp,s,f	w,sp,s,f	w,sp,s,f	sp,s,f	s,f
Paramuriceidae					
<u>Thesea</u> sp.	-	-	s	sp,s,f	w,sp,s,f
Plexauridae					
<u>Muricea pendula</u>	sp,f	-	w,sp,s,f	w,sp,f	w
Pennatulacea					
Virgulariidae					
<u>Stylatula</u> sp.	-	-	-	f	w,sp,s,f
Anthozoa - Zoantharia					
Actiniaria					
undetermined Actiniaria	w,sp,s,f	w,sp,s,f	w,sp,s,f	w,sp,s,f	sp,s
Scleractinia - Faviina					
Faviidae					
<u>Solenastria hyades</u>	w	-	-	-	-
Oculinidae					
<u>Oculina arbuscula</u>	-	sp	-	-	-
<u>Oculina</u> sp.	sp,s	f	w	-	-
Rhizangiidae					
<u>Phyllangia americana</u>	-	-	w,sp	sp,s	-
Scleractinia - Dendrophylliina					
Dendrophylliidae					
<u>Balanophyllia floridana</u>	-	-	-	-	s,f
Antipatharia					
undetermined Antipatharian	-	f	f	-	sp,s,f
Arthropoda - Chelicerata					
Merostomata - Xiphosura					

Appendix 2. (Continued)

	IS01	IS02	MS02	MS06	OS01
Limulidae					
<u>Limulus polyphemus</u>	-	w	-	-	-
Arthropoda - Mandibulata					
Crustacea - Cirripedia					
Thoracia - Lepadomorpha					
Lepadidae					
<u>Lepas pectinata</u>	-	-	-	-	sp
Scalpellidae					
<u>Scalpellum diceratum</u>	-	-	-	-	sp,s,f
Thoracia - Balanomorpha					
Archaeobalanidae					
<u>Conopea galeata</u>	w,sp,s,f	w,sp,s,f	w,sp	-	f
<u>Conopea merrilli</u>	w,sp,s,f	w,sp,f	w,sp,s,f	w,sp,s,f	w,sp,s,f
Balanidae					
<u>Balanus calidus</u>	sp	sp	s	s	sp,s,f
<u>Balanus trigonus</u>	w,sp,s,f	w,sp,s,f	w,sp,s,f	w,sp,s,f	w,sp,s,f
<u>Balanus venustus</u>	w,sp,s,f	w,sp,f	w,sp,s,f	w,s	w,sp,f
Crustacea - Malacostraca					
Stomatopoda					
Gonodactylidae					
<u>Gonodactylus bredini</u>	-	-	w,f	w,s,f	-
Decapoda - Dendrobranchiata					
Penaeidae					
<u>Mesopenaeus tropicalis</u>	-	-	-	-	w,sp,s,f
<u>Metapenaeopsis goodei</u>	s	-	w,s,f	w,s,f	w,sp,s,f
<u>Penaeus duorarum duorarum</u>	-	f	s,f	-	s
<u>Trachypenaeus constrictus</u>	w,sp,f	w,sp,s,f	w,sp,s,f	w	w
Sicyonidae					
<u>Sicyonia brevirostris</u>	w	w,f	w,sp	-	w,sp,s,f
Solenoceridae					
<u>Solenocera atlantidis</u>	-	-	s	s,f	w,sp,s,f
Decapoda - Pleocyemata - Caridea					
Alpheidae					
Alpheidae A	-	-	sp	-	-
<u>Alpheus formosus</u>	-	w,s	w,sp	w,sp,f	sp
<u>Alpheus normanni</u>	-	-	-	s	s

Appendix 2. (Continued)

	IS01	IS02	MS02	MS06	OS01
<u>Alpheus</u> sp. A	-	-	-	-	s
<u>Automate evermanni</u>	-	-	-	s	-
<u>Synalpheus apioceros</u>	-	-	f	-	-
<u>Synalpheus fritzmuelleri</u>	f	sp,s,f	-	-	-
<u>Synalpheus longicarpus</u>	w,sp,f	w,sp,s,f	sp,s	sp,f	-
<u>Synalpheus minus</u>	sp,f	w,sp,s,f	sp	f	sp,s
<u>Synalpheus</u> sp. A	-	-	s	-	-
<u>Synalpheus townsendi</u>	w,sp,s,f	w,sp,s,f	w,sp,s	w,sp,f	w,sp,f
Hippolytidae					
<u>Lysmata wurdemanni</u>	-	-	-	f	-
<u>Tozeuma serratum</u>	-	-	sp	-	sp,s
Pandalidae					
<u>Parapandalus</u> sp.	-	-	-	-	f
<u>Plesionika tenuipes</u>	-	-	-	-	sp
Palaemonidae					
<u>Anchistioides antiguensis</u>	-	-	-	f	-
<u>Periclimenes perryae</u> (?)	-	-	-	sp	-
<u>Pontonia margarita</u>	-	-	s	-	-
Processidae					
<u>Processa tenuipes</u>	-	-	-	-	sp,s
Decapoda - Pleocyemata - Palinura					
Scyllaridae					
<u>Scyllarides nodifer</u>	w	-	w,sp,s,f	w,sp,s	sp,s,f
<u>Scyllarus chacei</u>	-	-	-	-	s
<u>Scyllarus depressus</u>	-	-	-	-	s,f
Decapoda - Pleocyemata - Anomura					
Albuneidae					
<u>Albunea gibbesii</u>	-	-	-	f	-
Axiidae					
<u>Axiopsis hirsutimana</u>	-	-	-	-	w,s
Diogenidae					
<u>Dardanus fucosus</u>	-	-	-	s,f	-
<u>Dardanus insignis</u>	-	-	-	-	sp,s
<u>Paguristes sericeus</u>	-	-	-	-	s
<u>Paguristes tortugae</u>	f	w,sp,s	-	-	-
Galatheididae					
<u>Galathea rostrata</u>	-	-	-	f	sp,s,f
Paguridae					
<u>Pagurus carolinensis</u>	-	-	w,sp	-	w
<u>Pagurus defensus</u>	-	-	-	-	w
<u>Pylopagurus corallinus</u>	-	-	-	-	sp,s
<u>Pylopagurus holthuisi</u>	-	-	-	f	-
Porcellanidae					
<u>Megalobrachium soriatum</u>	-	sp,s	sp,s	sp	sp
<u>Pachycheles rugimanus</u>	-	-	sp	w,sp,f	w,sp,s,f
<u>Petrolisthes galathinus</u>	-	w,s,f	w,sp	-	-

Appendix 2. (Continued)

	IS01	IS02	MS02	MS06	OS01
Decapoda - Pleocyemata - Brachyura					
Cancridae					
<u>Cancer irroratus</u>	w	sp	-	-	-
Dorippidae					
<u>Ethusa mascarone americana</u>	-	-	-	-	w
Dromiidae					
<u>Dromidia antillensis</u>	w,sp,s,f	sp,s,f	w,sp,f	w,sp,s,f	sp,s,f
Goneplacidae					
<u>Glyptoplax smithi</u>	-	-	-	-	f
Grapsidae					
<u>Euchirograpsus americanus</u>	-	-	-	-	sp,s
<u>Planes minutus</u>	-	-	-	-	f
Homolidae					
<u>Homola barbata</u>	-	-	-	-	sp,s
Homolidae A	-	-	-	-	sp
Leucosiidae					
<u>Iliacantha intermedia</u>	-	-	-	-	w
Majidae					
<u>Batrochonotus fragosus</u>	-	-	-	-	s
<u>Collodes trispinosus</u>	-	-	-	-	s
<u>Libinia dubia</u>	s	-	-	-	-
<u>Libinia emarginata</u>	w	-	-	-	-
<u>Macrocoeloma camptocerum</u>	w	w,sp	-	-	-
<u>Macrocoeloma trispinosum</u>	w,sp	sp,s,f	s,f	-	-
<u>Metaporaphis calcaratta</u>	sp	w,sp	sp	-	s
<u>Mithrax acuticornis</u>	-	-	sp	-	sp,s
<u>Mithrax forceps</u>	-	s	w,sp	w,sp	-
<u>Mithrax pleuracanthus</u>	w,sp,s,f	w,sp,s,f	w,sp,s,f	w,sp,s,f	-
<u>Pelia mutica</u>	-	sp	sp	-	sp
<u>Podochela gracilipes</u>	-	-	-	-	sp,s
<u>Podochela riisei</u>	w,sp	w,sp,s,f	w,sp	-	-
<u>Podochela sidneyi</u>	w	w	-	-	w,f
<u>Stenocionops furcata coelata</u>	sp	-	sp,s	sp	w,sp,s,f
<u>Stenorhynchus seticornis</u>	-	w	-	-	sp,s,f
Palicidae					
<u>Palicus faxoni</u>	-	-	-	-	s
Parthenopidae					
<u>Parthenope fraterculus</u>	-	-	-	-	sp,s
Portunidae					
<u>Ovalipes stephensoni</u>	w,s	s	s	sp	-
<u>Portunus gibbesii</u>	s	s	s	-	-
<u>Portunus ordwayi</u>	-	-	-	-	f
<u>Portunus spinicarpus</u>	-	-	-	-	sp,s
<u>Portunus spinimanus</u>	-	-	-	sp	s
Xanthidae					
<u>Carpoporos papulosus</u>	-	-	-	-	s
<u>Glyptoxanthus erosus</u>	-	-	f	sp	w
<u>Lobopilumnus agassizii</u>	-	-	-	-	s
<u>Pilumnus dasypodus</u>	sp,s,f	w,sp,s,f	w,sp,f	w	-

Appendix 2. (Continued)

	IS01	IS02	MS02	MS06	OS01
<u>Pilumnus floridanus</u>	s	w,sp,s	sp	w,sp,f	sp
<u>Pilumnus pannosus</u>	sp,s,f	w,sp,s,f	sp,s	-	-
<u>Pilumnus sayi</u>	w,sp,s,f	w,sp,s,f	w,sp,s,f	w,sp,s,f	sp,f
<u>Pseudomedeus agassizii</u>	-	w,sp	w,sp,s	w,sp,f	w,sp,s,f
Xanthidae A	-	-	sp	-	sp

Bryozoa

Stenolaemata

Cyclostomata

Crisiidae

Crisia sp.

w,sp,s,f w,sp,s,f w,sp,s,f w,sp,s,f w,sp

Diaperoeciidae

Diaperoecia floridana

- - w,sp,s w,sp,s,f w,sp,s,f

Lichenoporidae

Lichenopora radiata

- - - w,sp,f -

Oncousoeciidae

Proboscina floridana

- - - sp -

Plagioeciidae

Plagioecia dispar

- - - s,f -

Gymnolaemata

Ctenostomata

undetermined Ctenostomata

s w,f w,sp,s sp,f w,sp,s,f

Alcyonidiidae

Alcyonidium sp.

f - w,sp - w,s

Buskiidae

Buskia sp.

sp sp sp - -

Nolleliidae

Nolella stipata

w,sp w,s,f w,sp,s w w,sp

Valkeriidae

Aeverrillia setigera

f s,f w,sp,s,f sp,s,f w

Vesiculariidae

Amathia alternata

s s,f w,s,f f w,sp,s,f

Amathia convoluta

- - - sp -

Amathia distans

s,f w,sp,s,f w,sp,s,f w,sp,s,f w,sp,s,f

Victorellidae

Sundanella sibogae

sp,s,f sp,s,f sp,f sp sp,s,f

Cheilostomata - Anasca

Aeteidae

Aetea anguina

w,sp,f w,sp,f w,sp,s,f w,sp,s,f w,sp,s,f

Aetea truncata

w - sp w -

Appendix 2. (Continued)

	IS01	IS02	MS02	MS06	OS01
Alderinidae					
<u>Alderina smitti</u>	-	-	sp,f	w,s,f	-
<u>Antropora tincta</u>	sp,s,f	w,sp,s,f	w,sp,s	sp,s	sp,s
<u>Copidozum tenuirostre</u>	-	-	-	w,sp,f	-
<u>Crassimarginatella leucocypha</u>	w,sp,s	w,sp	w	w,sp	w
<u>Parellisina curvirostris</u>	-	-	w,sp,s,f	w,sp,s,f	sp,s,f
<u>Parellisina latirostris</u>	-	-	f	-	-
Bugulidae					
<u>Bugula fulva</u>	s	w,sp	w,sp,s,f	w,sp,s,f	w
<u>Bugula grayi</u>	-	-	sp,f	sp	-
<u>Bugula marcusii</u>	-	sp	-	sp	-
<u>Bugula microoecia</u>	-	-	w,sp,s	w,sp	w
<u>Bugula rylandi</u>	s,f	w,sp,f	w,sp,s,f	w,sp,s,f	w
<u>Bugula sp.</u>	-	-	sp,s,f	f	-
<u>Bugula turrita</u>	sp	w	-	-	-
<u>Caulibugula dendrograpta</u>	f	-	f	s	-
<u>Caulibugula pearsei</u>	sp,f	sp,f	-	sp,f	-
Cellariidae					
<u>Cellaria nodosa</u>	-	-	-	-	sp
Chaperiidae					
<u>Chaperia sp.</u>	s,f	sp,s,f	w,sp,s,f	w,sp,s,f	w
Cupuladriidae					
<u>Cupuladria doma</u>	-	-	-	w,sp	w
<u>Discoporella umbellata</u>	-	-	-	-	s,f
Electrinidae					
<u>Electra lacinosa</u>	-	-	-	w,sp,s	-
Epistomiidae					
<u>Synnotum aegyptiacum</u>	-	-	-	f	-
Farciminariidae					
<u>Nellia tenella</u>	-	-	s,f	s,f	sp,s
Hincksinidae					
<u>Aplousina gigantea</u>	-	-	-	-	w,sp,s,f
<u>Aplousina tuberosa</u>	-	-	-	w,sp	-
Membraniporidae					
<u>Membranipora savartii</u>	-	-	w,sp	w	sp
<u>Membranipora tenuis</u>	sp,s,f	-	sp,f	-	sp
Microporidae					
<u>Micropora coriacea</u>	-	-	-	-	sp
Onychocellidae					
<u>Floridina antiqua</u>	-	sp	-	s	sp,s
<u>Smittipora levinseni</u>	-	f	w,sp	w	s
Scrupocellariidae					
<u>Caberia boryi</u>	-	w	s	w,sp,s,f	-
<u>Scrupocellaria bertholleti</u>	f	f	-	-	-
<u>Scrupocellaria regularis</u>	w,s,f	w,s,f	f	-	-
Steganoporellidae					
<u>Steganoporella magnilabris</u>	-	w	sp	w,sp,s,f	-

Appendix 2. (Continued)

	IS01	IS02	MS02	MS06	OS01
Cheilostomata - Cribrimorpha					
Cribrilinidae					
<u>Cribrilaria floridana</u>	-	-	-	w,sp,s,f	-
<u>Cribrilaria radiata</u>	s	s,f	-	w,sp,f	s
<u>Membraniporella aragoi</u>	-	-	-	-	s,f
Cheilostomata - Ascophora					
Adeonidae					
<u>Reptadeonella hastingsae</u>	s	w,s	w,sp,f	w,sp,s,f	s,f
Celleporariidae					
<u>Celleporaria albirostris</u>	w,s,f	-	w,sp,s,f	w,sp,s,f	w,sp,s,f
<u>Celleporaria magnifica</u>	w,s	w,s,f	w,sp,s,f	w,sp,s,f	w,sp,s,f
<u>Celleporaria mordax</u>	sp,s,f	w,sp,s,f	s,f	w	-
Celleporinidae					
<u>Celleporina hassalli</u>	sp,s,f	s	w,sp,s,f	sp,s	-
<u>Turbicellepora dichotoma</u>	w,sp,s,f	w,sp,s,f	w,sp,s,f	w,sp,s,f	w,sp,s,f
Cheiloporinidae					
<u>Hippaliosina rostrigera</u>	w,sp,s	w,sp,s	w,sp,s,f	w,sp,s	sp,s
Crepidacanthidae					
<u>Crepidacantha poissoni teres</u>	-	-	-	s,f	s
Exochellidae					
<u>Exochella longirostris</u>	-	-	-	sp	s
Hippoporinidae					
<u>Hippopleurifera mucronata</u>	-	-	sp,f	-	-
<u>Hippoporella uvulifera</u>	-	s	-	-	-
<u>Hippoporidra janthina</u>	-	sp,s,f	s,f	s	-
<u>Hippoporina contracta</u>	w,sp,s,f	s,f	w,s,f	w,sp	sp,f
Hippoporinidae A	-	-	-	sp	-
Hippothoidae					
<u>Trypostega venusta</u>	w,sp,s	sp,s,f	-	w,sp,s,f	f
Microporellidae					
<u>Microporella ciliata</u>	w,sp,s,f	w,sp,s,f	w,sp,s,f	w,sp,s,f	w,sp,s,f
<u>Microporella marsupiata</u>	w	-	-	sp,f	-
<u>Microporella umbracula</u>	w,sp,s	w,s,f	sp,f	w,sp,s,f	f
Petralliidae					
<u>Petralliella bisinuata</u>	f	-	w,sp,s,f	w,sp,s,f	w,sp,s,f
<u>Petralliella marginata</u>	-	-	-	-	s,f
Phylactellidae					
<u>Phylactella aviculifera</u>	-	-	-	sp,f	-
Schizoporellidae					
<u>Arthropoma cecili</u>	-	-	-	w,s,f	-
<u>Cleidochasma porcellanum</u>	s	-	sp	sp	w,s,f
<u>Cycloperiella rubra</u>	-	-	-	-	sp,s,f
<u>Schizoporella cornuta</u>	w,sp,s,f	w,sp,s,f	w,sp,s,f	w,sp,s,f	w,sp,s,f
<u>Schizoporella errata</u>	-	-	w,sp	w,sp,f	sp
<u>Schizoporella floridana</u>	s,f	-	w,sp,s,f	w,sp,s,f	w,f
<u>Stylopoma informata</u>	-	-	-	w	sp,s,f

Appendix 2. (Continued)

	IS01	IS02	MS02	MS06	OS01
Sertellidae					
<u>Rhyncozoon rostratum</u>	-	f	f	sp	-
<u>Rhyncozoon tuberculatum</u>	-	-	-	sp, f	-
Smittinidae					
<u>Parasmittina spathulata</u>	s	w, sp, f	w, sp, s, f	w, sp, s, f	sp, s, f
<u>Smittina smittiella</u>	-	w, f	-	w, sp, s, f	-
Stomachetosellidae					
<u>Bracebridgia subsulcata</u>	-	-	-	f	-
Mollusca					
Gastropoda - Prosobranchia					
Archaeogastropoda					
Fissurellidae					
<u>Diodora cayenensis</u>	w, sp, s, f	w, sp, s, f	w, sp, s, f	w, sp, f	s
Trochidae					
<u>Calliostoma euglyptum</u>	-	s	-	f	-
<u>Calliostoma pulchrum</u>	-	-	sp	-	-
<u>Calliostoma roseolum</u>	-	-	-	-	sp
Mesogastropoda					
Crepidulidae					
<u>Crepidula aculeata</u>	s	w, s	w, sp	w, sp	-
<u>Crepidula fornicata</u>	-	-	-	sp	-
<u>Crepidula plana</u>	-	-	-	-	f
Cymatiidae					
<u>Cymatium labiosum</u>	-	-	-	s	-
<u>Distorsio constricta macgintyi</u>	-	-	-	-	f
Cypraeidae					
<u>Cypraea cervus</u>	-	w	-	-	w
<u>Cypraea cinerea</u>	-	-	-	w	-
<u>Cypraea spurca acicularis</u>	-	-	-	sp	f
Eratoidae					
<u>Erato maugeriae</u>	-	s	-	-	-
<u>Trivia maltbiana</u>	-	-	-	sp	s
<u>Trivia nix</u>	s	-	-	-	-
<u>Trivia pendiculus</u>	-	-	s	-	s
Ovulidae					
<u>Simnia acicularis</u>	-	-	s	-	-
<u>Simnia uniplicata</u>	-	sp	sp	-	-
Turritellidae					
<u>Vermicularia knorrii</u>	-	s, f	-	w, f	s, f
Triphoridae					
<u>Triphora nigrocincta</u>	-	-	-	sp	-
Vermetidae					
<u>Spiroglyphus irregularis</u>	-	-	f	-	-

Appendix 2. (Continued)

	IS01	IS02	MS02	MS06	OS01
Neogastropoda					
Buccinidae					
<u>Cantharus multangulus</u>	-	sp	-	-	-
<u>Pisania tinctoria</u>	f	w,sp,s,f	sp,s,f	sp,f	sp,f
Columbellidae					
<u>Anachis avara</u>	-	-	-	-	s
<u>Anachis lafresnayi</u>	sp	sp,s	-	f	s
<u>Mitrella lunata</u>	-	s	sp	-	-
<u>Nassarina minor</u>	-	-	-	-	s
<u>Suturoglypta iontha</u>	-	f	sp	-	-
Coralliophilidae					
<u>Coralliophila caribaea</u>	-	-	sp	-	-
Fasciariidae					
<u>Fasciolaria lilium hunteria</u>	s	-	-	-	-
<u>Fasciolaria tulipa</u>	-	-	s	-	-
<u>Pleuroploca gigantea</u>	-	-	-	sp	-
Marginellidae					
<u>Marginella roscida</u>	-	-	-	f	-
Muricidae					
<u>Calotrophon ostrearum</u>	-	-	sp	-	-
<u>Murex florifer dilectus</u>	s	-	-	-	-
<u>Murex pomum</u>	-	-	-	f	-
<u>Murex recurvirostris rubidus</u>	-	f	-	-	-
<u>Thais haemastoma florida</u>	-	-	sp	-	-
<u>Urosalpinx perrugata</u>	-	sp	-	-	-
Nassariidae					
<u>Nassarius albus</u>	-	s	f	-	-
<u>Nassarius vibex</u>	-	-	-	w	-
Gastropoda - Opisthobranchia					
Aplysiacea					
Aplysiidae					
<u>Aplysia willcoxi</u>	s	-	-	-	-
Tylodidae					
<u>Pleurobranchaea hedgpethi</u>	sp	-	-	-	-
Nudibranchia					
undetermined Nudibranchia	-	-	-	-	f
Bornellidae					
<u>Bornella calcarata</u>	-	-	w,sp	-	sp,s
Dendrodorididae					
<u>Dendrodoris krebsii</u>	-	-	s	-	-
<u>Dendrodoris warta</u>	-	-	-	w	w
Dorididae					
<u>Anisodoris worki</u>	-	-	w	-	-
<u>Hypselodoris edenticulata</u>	-	-	-	sp	-
<u>Taringa aivica</u>	sp	-	-	-	-

Appendix 2. (Continued)

	IS01	IS02	MS02	MS06	OS01
Dotodae					
<u>Doto doerga</u> (?)	-	-	s	-	-
Pelecypoda					
Arcoida					
Arcidae					
<u>Anadara notabilis</u>	-	-	-	sp	-
<u>Arca imbricata</u>	w,s	w,sp,s,f	w,sp,s	sp,s	-
<u>Arca zebra</u>	w	-	w	w,f	-
<u>Barbatia candida</u>	-	-	-	f	sp
<u>Barbatia domingensis</u>	-	-	-	sp	-
Mytiloida					
Mytilidae					
<u>Gregariella coralliophaga</u>	-	-	-	sp	-
<u>Lithophaga antillarum</u>	-	-	sp	sp	-
<u>Lithophaga aristata</u>	-	-	sp	sp	-
<u>Lithophaga bisulcata</u>	-	sp,f	s	sp	sp
<u>Modiolus americanus</u>	f	sp	-	w,s	-
<u>Musculus lateralis</u>	f	s,f	sp,s	-	-
Pinnidae					
<u>Pinna carnea</u>	-	-	w	-	-
Pterioida					
Anomiidae					
<u>Anomia simplex</u>	-	-	f	s	-
<u>Anomia squamula</u>	-	-	sp	-	-
<u>Pododesmus rudis</u>	-	sp	sp	-	-
Pectinidae					
<u>Chlamys benedicti</u>	-	-	-	sp	w,sp,s,f
<u>Lyropecten nodosus</u>	-	-	-	f	w,f
Plicatulidae					
<u>Plicatula gibbosa</u>	w	-	-	-	-
Pteriidae					
<u>Pteria colymbus</u>	w,sp,s,f	w,sp,s,f	w,sp,s,f	w,sp,s,f	w,sp,s,f
Ostreina					
Ostreidae					
<u>Crassostrea virginica</u>	-	-	-	w	-
<u>Lopha frons</u>	-	-	w,sp	s	-
<u>Ostrea equestris</u>	s	-	w,s	sp,s	-
<u>Ostrea frons</u>	-	-	w	-	-
<u>Ostrea permolis</u>	w	s	w	w	-
Veneroida					
Chamidae					
<u>Chama congregata</u>	w	w,sp,s,f	w,sp,s,f	w,sp	s
<u>Chama macerophylla</u>	-	-	sp	sp	-
<u>Pseudochama radians</u>	-	w	-	-	-

Appendix 2. (Continued)

	IS01	IS02	MS02	MS06	OS01
Veneridae					
<u>Chione</u> <u>grus</u>	-	-	-	sp	-
<u>Gouldia</u> <u>cerina</u>	-	-	-	f	-
Myoida					
Corbulidae					
<u>Corbula</u> <u>chittyana</u>	-	-	-	sp	-
Gastrochaenidae					
<u>Gastrochaena</u> <u>hians</u>	-	sp	-	sp	sp
<u>Spengleria</u> <u>rostrata</u>	-	-	-	sp	-
Hiatellidae					
<u>Hiatella</u> <u>arctica</u>	w,s	s,f	sp,s	w,sp,s,f	sp,s,f
Pholadomyoida					
Lyonsiidae					
<u>Lyonsia</u> <u>beana</u>	w,s,f	-	f	-	-
Cephalopoda					
Sepioidea					
Sepiolidae					
<u>Rossia</u> <u>sp.</u>	w	-	-	-	-
Teuthoidea					
Loliginidae					
<u>Loligo</u> <u>pealei</u>	w	w	-	s	s
<u>Loligo</u> <u>plei</u>	sp,s	w,sp,s	sp,s	-	s
undetermined Loliginidae	w,sp,s,f	w,sp,s,f	w,sp,s,f	w,sp,s,f	w,sp,s,f
Ommastrephidae					
<u>Illex</u> <u>illecebrosus</u>	-	-	f	-	-
Octopoda					
Octopodidae					
<u>Octopus</u> <u>sp.</u>	-	w	sp	sp	sp
Echinodermata					
Crinoidea					
Comasteridae					
<u>Comactinia</u> <u>meridionalis</u>	-	-	w,sp,f	w,sp	-
Stelleroidea - Asteroidea					
Platyasterida					
Luidiidae					
<u>Luidia</u> <u>alternata</u>	w,sp,s	sp,f	-	sp	-
<u>Luidia</u> <u>clathrata</u>	-	sp	-	sp	s

Appendix 2. (Continued)

	IS01	IS02	MS02	MS06	OS01
Paxillosida					
Astropectinidae					
<u>Astropecten articulatus</u>	-	-	w	-	-
<u>Astropecten comptus</u>	-	-	w,s	-	w,sp
<u>Astropecten</u> sp.	-	-	-	-	sp,s
Valvatida					
Ophidiasteridae					
<u>Narcissia trigonaria</u>	-	-	-	-	s
Spinulosida					
Echinasteridae					
<u>Echinaster brasiliensis</u>	-	-	-	-	s,f
<u>Echinaster modestus</u>	-	-	w	-	w
<u>Echinaster serpentarius</u>	w	w	-	-	-
<u>Echinaster</u> sp.	sp,s,f	sp,s,f	sp,s,f	sp,s	sp,s,f
Forcipulata					
Asteriidae					
<u>Asterias forbesii</u>	w,sp	sp	-	-	sp
Stelleroidea - Ophiuroidea					
Ophiuroidea A	-	-	w	-	-
Ophiuroidea B	-	-	w	-	-
Phyronophiurida - Euryalina					
Gorgonocephalidae					
<u>Astrophyton muricatum</u>	sp	s	w,sp,s,f	w,sp,s,f	w,sp,s,f
<u>Astroporpa annulata</u>	-	-	-	sp,s,f	w,sp,s,f
Ophiurida					
Amphiuridae					
<u>Amphiodia pulchella</u>	-	-	-	s	-
<u>Axiognathus squamatus</u>	-	-	-	sp,s	-
<u>Ophiostigma tsacanthum</u>	-	-	-	s	-
Ophiactidae					
<u>Ophiactis savignyi</u>	-	-	sp	sp,s	-
Ophiidermatidae					
<u>Ophioderma brevispinum</u>	-	-	w,sp	sp,s	-
Ophiothricidae					
<u>Ophiothrix angulata</u>	w,sp,s	w,sp,s,f	w,sp,s	w,sp,s,f	w,sp,s
<u>Ophiothrix suensonii</u>	-	-	-	-	w,s

Appendix 2. (Continued)

	IS01	IS02	MS02	MS06	OS01
Echinoidea					
Cidaroida					
Cidaridae					
<u>Eucidaris tribuloides</u>	-	-	-	w,sp,s	w,sp,f
Arbacoidea					
Arbaciidae					
<u>Arbacia punctulata</u>	w,sp,s,f	w,sp,s,f	w,sp,s	sp,s	s,f
Temnopleuroida					
Temnopleuridae					
<u>Genocidaris maculata</u>	-	-	-	s	s
Toxopneustidae					
<u>Lytechinus variegatus</u>	w,s,f	w,sp,s,f	-	w,sp,s,f	-
Holothuroidea					
Dendrochirotida					
Cumariidae					
<u>Ocnus pygmaeus</u>	w,sp,s,f	w,sp,s,f	w	w	-
Aspidochirotida					
Holothuridae					
<u>Holothuria dakarensis</u>	-	sp	-	-	-
<u>Holothuria princeps</u>	-	-	f	-	-
<u>Holothuria surinamensis</u>	-	-	sp	-	-
Synallactidae					
<u>Isostichopus badinotus</u>	-	-	w,f	-	s
Chordata					
Ascidiacea					
Ascidiacea A	s	w,sp,f	-	-	-
undetermined Ascidiacea	-	w,f	-	-	-
Enterogona					
Asciidiidae					
<u>Ascidia curvata</u>	f	f	s,f	s	sp
<u>Ascidia sp. A</u>	-	sp,s,f	-	-	-
Clavelinidae					
<u>Clavelina gigantea</u>	s	w	w,sp,s,f	w,sp,f	-
<u>Clavelina picta</u>	w	w	-	sp	-
<u>Clavelina sp.</u>	w,sp,s,f	w,sp,s,f	sp,s,f	w,sp,s	w
<u>Clavelina sp. D</u>	s	-	-	sp	-
<u>Distaplia bermudensis</u>	w,sp,s,f	w,sp,s,f	w,sp,s,f	sp,f	w
<u>Eudistoma carolinense</u>	-	-	sp	s	sp,s

Appendix 2. (Continued)

	IS01	IS02	MS02	MS06	OS01
<u>Euristoma olivaceum</u>	w	-	-	-	-
<u>Euristoma</u> sp.	w,sp,s,f	w,sp,s,f	w,sp,s,f	w,sp,s,f	w,s,f
Didemnidae					
<u>Didemnum candidum</u>	w,sp,s,f	w,s,f	w,sp,s,f	w,sp,s,f	w,sp,s,f
<u>Didemnum</u> sp.	-	-	sp	-	w
<u>Didemnum speciosum</u>	sp,s	-	f	-	s
<u>Diplosoma macdonaldi</u>	w,sp,f	w,sp,s,f	w,sp,s,f	sp,f	w,sp,s
<u>Trididemnum savignii</u>	sp	sp	-	s	sp,s
<u>Trididemnum</u> sp.	sp,s	sp,s,f	w,sp,s	sp	w,s,f
Polyclinidae					
<u>Aplidium bermudae</u>	w	-	-	-	-
<u>Aplidium constellatum</u>	w	w	w	-	-
<u>Aplidium exile</u>	-	-	-	w	-
<u>Aplidium pellucidum</u>	w,s	-	-	-	sp,s
<u>Aplidium</u> sp. A	-	-	sp	w	-
<u>Aplidium</u> sp. B	-	-	w,sp	-	w
<u>Aplidium</u> sp. D	-	-	-	sp	-
<u>Aplidium stellatum</u>	w	w	w	w	-
Pleurogona					
Molgulidae					
<u>Molgula occidentalis</u>	w,sp,s,f	w,sp,s,f	w,sp,s,f	w,sp,s,f	-
Pyuridae					
<u>Pyura vittata</u>	w	sp	sp,f	-	-
Styelidae					
<u>Botrylloides nigrum</u>	f	sp,f	sp,s	sp	f
<u>Botryllus planus</u>	s	sp,s,f	w,s,f	sp	-
<u>Polyandrocarpa floridana</u>	-	-	w,sp,s,f	w,sp,s,f	sp
<u>Polyandrocarpa gravei</u>	f	-	w,sp,s,f	-	-
<u>Polyandrocarpa tinctoria</u>	-	-	f	-	-
<u>Styela partita</u>	w,sp,s,f	w,sp,s,f	sp,f	-	-
<u>Styela plicata</u>	w,sp,s,f	w,sp,s,f	w,sp,s,f	w,sp,s,f	-
<u>Symplegma viride</u>	sp,s,f	s,f	sp,s	-	f

Appendix 3. Ranked abundance of invertebrate species collected by suction and grab samplers at each station off South Carolina and Georgia during 1981. Average ($\bar{x}$) density, expressed as number per 0.10 m², and ranked abundance for each season are indicated along with rank for all seasons combined.

OVERALL RANK	SPECIES	STATION		IS01		WINTER			SPRING			SUMMER			FALL		
						MEAN		ST ERR	MEAN		ST ERR	MEAN		ST ERR	MEAN		ST ERR
1.0	FILOGRANA IMPLEXA	3.2	2.9									0.2	0.2		182.4	124.8	
2.0	SYLLIS SPONGICOLA	6.6	5.6						4.8	3.2		14.8	14.3		55.0	53.3	
3.0	ERICHTHONIUS BRASILIENSIS	41.0	27.3						3.4	1.1		4.0	2.6		1.0	0.8	
4.0	AXIOHELLA MUCOSA	3.6	2.0						29.4	17.6		14.4	4.8		1.8	0.8	
5.0	EXOgone DISPAR	16.8	6.2						4.6	1.9		1.6	0.6		14.2	2.7	
6.0	LUCONACIA INCERTA	22.6	11.5						4.6	2.4		0.8	0.4		3.6	1.4	
7.0	AMPHARETE AMERICANA	28.0	9.5						0.2	0.2		0.4	0.2				
8.0	PHOTIS SP.	24.8	11.3						0.8	0.4		0.8	0.5		1.6	0.9	
9.0	ELASMOPUS SP. A	1.4	0.9						20.0	11.2		2.4	0.8		2.2	1.9	
10.0	GAMMAROPSIS SP.	9.0	3.6						11.2	8.0		0.4	0.2		0.4	0.2	
11.0	CAPRELLA EQUILIBRA	16.8	8.2						2.8	1.6							
12.0	PODOCERUS SP.	13.6	4.1						1.0	0.0		0.6	0.2		2.2	0.9	
13.0	LEUCOTHOE SPINICARPA	1.4	1.2						7.4	6.2		5.2	5.2		2.2	2.2	
14.0	RILDARDANUS LAMINOSA	6.2	1.3						5.8	3.0		1.2	0.2		2.2	0.6	
15.0	MICRODEUTOPUS MYERSI	14.2	6.9												0.2	0.2	
16.0	AMPELISCA SCHELLENBERGI	4.2	1.3						1.2	0.7		2.6	0.5		5.6	1.8	
17.0	OWENIA FUSIFORMIS	8.4	4.8						3.2	2.2		1.2	0.8		0.6	0.2	
18.0	CARPIAS BERMUDENSIS	0.6	0.4						7.6	2.8		1.6	1.1		3.4	3.1	
19.0	PRIONOSPPIO CRISTATA	12.0	8.0												1.0	0.8	
20.0	NEMATODA	4.8	1.9						2.6	1.2		1.6	0.7		3.8	1.1	
21.5	ASPIDOSIPHON GOSNOLDI	4.4	2.1						1.0	0.4		2.6	0.9		4.2	2.5	
21.5	MEGALOBRACHIUM SORIATUM	1.4	0.9						4.4	3.0		0.6	0.6		5.6	5.1	
23.0	SIPUNCULIDA	7.6	5.4						0.2	0.2		0.6	0.2		3.6	1.5	
24.0	GOULDIA CERINA	1.4	0.4						1.4	0.7		5.6	0.7		1.2	0.6	
25.0	CHONE AMERICANA	4.4	1.6						1.8	1.2		1.4	0.4		1.8	1.2	
26.5	MALACOCEROS GLUTAEUS	0.4	0.2						0.2	0.2		2.8	1.5		5.4	2.1	
26.5	PARACERCETIS CAUDATA	1.8	0.7						2.0	0.8		1.4	0.7		5.6	1.1	
28.0	OPHIOTHRIX ANGULATA	1.4	0.7						2.4	1.2		0.2	0.2		3.6	3.3	
29.0	SYLLIS HYALINA	1.8	1.0						1.8	1.0		0.8	0.4		3.2	1.5	
30.5	ERICHSONELLA FILIFORMIS	0.2	0.2						3.0	1.3		3.4	1.5		0.6	0.2	
30.5	PAGURUS CAROLINENSIS	0.4	0.2						1.2	1.0		0.4	0.4		5.2	2.8	
32.0	EUNICE VITTATA	1.4	0.5						0.6	0.4		2.6	2.6		2.4	0.8	
33.0	LUCIFER FAXONI								0.2	0.2		6.6	4.1				
34.0	AMPHIODIA PULCHELLA	5.0	4.1						0.6	0.6		0.2	0.2		0.6	0.4	
35.5	HYDROIDES CRUCIGERA	1.6	1.1						0.6	0.2		1.0	0.4		3.0	2.1	
35.5	MARGINELLA HARTLEYANUM	1.8	0.5						2.0	0.7		2.0	0.8		0.4	0.4	
38.0	CHAMA CONGREGATA	1.2	0.7						0.8	0.6		1.0	0.5		3.0	2.3	
38.0	LEPTOCHELIA SP.	3.0	2.0						1.8	1.0		0.8	0.4		0.4	0.4	
38.0	MELITA APPENDICULATA	1.0	0.3						5.0	4.3							
40.0	BOTULA FUSCA	1.6	1.0						0.8	0.5		0.2	0.2		3.2	1.0	
41.0	HYDROIDES SP. A	1.4	0.5						1.0	0.4		0.8	0.5		2.4	1.1	
42.0	TURBELLARIA	2.8	1.1									0.8	0.4		1.8	1.1	
43.5	NEMERTINEA	1.2	3.4						0.2	0.2		1.8	0.5		0.4	0.2	
43.5	SPIOPHANES BOMBYX	4.6	2.6						0.4	0.4		0.2	0.2				
45.0	LAEVICARDIUM PICTUM	0.2	0.2						1.4	0.7		1.0	0.5		2.4	0.8	
46.0	CHRYSOPETALIDAE A	2.0	1.5						0.8	0.6		1.2	0.6		0.8	0.4	
47.5	OPHIOSTIGMA ISACANTHUM	1.4	1.4						0.8	0.5		1.4	0.5		1.0	1.0	
47.5	PILUMNUS SP.	0.2	0.2						1.6	0.9		0.2	0.2		2.6	1.3	
50.0	LUMBRINEKIS INFLATA	1.8	1.0						0.6	0.4		1.2	0.6		0.8	0.4	
50.0	LYSIANOPSIS ALBA								0.6	0.2		2.6	1.4		1.2	0.6	
50.0	PRIONOSPPIO CIRRIFERA	4.0	2.4									0.2	0.2		0.2	0.2	
52.5	LITHOPHAGA BISULCATA	0.2	0.2						1.2	1.0		0.4	0.2		2.4	0.7	
52.5	POLYCIRRUS CAROLINENSIS	1.6	0.7						0.2	0.2		0.2	0.2		2.2	1.1	
54.5	CERATONEREIS MIRABILIS	0.6	0.2						0.2	0.2		0.4	0.4		2.8	1.1	
54.5	PAGURIDEA								0.4	0.2					3.6	2.6	

Appendix 3. (Continued)

STATION IS01

OVERALL RANK	SPECIES	WINTER		SPRING		SUMMER		FALL	
		MEAN	ST ERR	MEAN	ST ERR	MEAN	ST ERR	MEAN	ST ERR
57.0	AMPITHOE SP. A	0.6	0.4	0.6	0.2	1.4	0.9	1.2	0.6
57.0	PAGURUS HENDERSONI	1.0	0.4	0.4	0.4	1.0	0.8	1.4	0.7
57.0	PITHO LHERMINIERI	1.6	1.2	.	.	0.6	0.4	1.6	0.5
61.5	MEGALOMMA BIOCULATUM	0.8	0.6	1.6	0.5	0.6	0.4	0.6	0.4
61.5	PELIA MUTICA	0.2	0.2	1.4	0.7	0.6	0.6	1.4	1.4
61.5	POTAMILLA SP.	0.2	0.2	0.2	0.2	.	.	3.2	2.5
61.5	PSEUDOMEDAUS AGASSIZII	.	.	1.0	0.8	1.6	1.1	1.0	0.3
61.5	SPIO SP. A	3.2	1.9	0.4	0.4	.	.	.	.
61.5	SUTUROGLYPTA IONTHA	1.8	0.6	1.4	0.5	.	.	0.4	0.2
65.0	ARMANDIA MACULATA	3.4	3.1	.	.	.	.	.	.
66.0	EULALIA SANGUINEA	2.2	0.9	.	.	0.2	0.2	0.8	0.5
67.5	PAGURIDAE	2.4	0.9	0.6	0.4	.	.	.	.
67.5	SERPULIDAE	0.4	0.2	0.6	0.4	0.8	0.2	1.2	0.5
70.0	CREPIDULA ACULEATA	0.6	0.4	0.2	0.2	0.6	0.6	1.4	1.2
70.0	DIPLODONTA SEMIASPERA	0.4	0.2	.	.	0.4	0.2	2.0	0.6
70.0	PODARKE OBSCURA	0.6	0.4	1.6	0.7	0.2	0.2	0.4	0.4
73.0	CRASSINELLA LUNULATA	0.4	0.4	0.4	0.2	1.0	0.4	0.8	0.4
73.0	NOTOMASTUS SP.	0.8	0.6	.	.	0.6	0.4	1.2	0.7
73.0	PAGURUS SP.	1.2	0.5	0.6	0.4	0.2	0.2	0.6	0.6
76.0	AMPELISCA VADORUM	2.0	0.8	.	.	0.2	0.2	0.2	0.2
76.0	PAGURISTES SP.	0.6	0.4	0.6	0.4	0.4	0.2	0.8	0.5
76.0	PAGURISTES TORTUGAE	1.4	0.7	0.2	0.2	0.2	0.2	0.6	0.2
79.5	CERITHIUM ATRATUM	0.4	0.4	0.4	0.4	0.8	0.4	0.6	0.4
79.5	PHYLLOCHAETOPTERUS SOCIALIS	1.0	0.6	.	.	0.2	0.2	1.0	0.8
79.5	POTAMILLA RENIFORMIS	.	.	0.2	0.2	0.4	0.4	1.6	1.0
79.5	SIPUNCULIDA A	0.6	0.6	0.8	0.5	0.6	0.4	0.2	0.2
85.5	ACANTHOCHITONA PYGMAEA	1.4	1.0	.	.	0.2	0.2	0.4	0.4
85.5	HYDROIDES UNCINATA	0.2	0.2	0.8	0.4	0.2	0.2	0.8	0.6
85.5	LOIMIA MEDUSA	1.8	0.7	.	.	0.2	0.2	.	.
85.5	MAJIDAE A	.	.	0.8	0.6	.	.	1.2	0.6
85.5	MITHRAX PLEURACANTHUS	0.8	0.6	0.8	0.2	.	.	0.4	0.4
85.5	SICYONIA SP.	0.2	0.2	.	.	0.2	0.2	1.6	1.2
85.5	VERMILIOPSIS ANNULATA	0.4	0.4	.	.	0.2	0.2	1.4	0.5
85.5	WEBSTERINEREIS TRIDENTATA	0.4	0.4	0.6	0.6	0.2	0.2	0.8	0.4
93.0	ARABELLA MUTANS	0.6	0.2	0.6	0.4	0.4	0.2	0.2	0.2
93.0	ARCA IMBRICATA	.	.	1.4	0.9	0.4	0.2	.	.
93.0	BRACHYURA	.	.	1.4	0.2	.	.	0.4	0.2
93.0	GASTROCHAENA HIANIS	0.2	0.2	0.2	0.2	0.8	0.4	0.6	0.4
93.0	LEMBOS UNICORNIS	0.4	0.4	0.8	0.4	0.4	0.2	0.2	0.2
93.0	RUDILEMBOIDES NAGLEI	1.6	0.9	0.2	0.2	.	.	.	.
93.0	TANYSTYLUM CALICIROSTRE	0.8	0.6	.	.	.	.	1.0	0.8
100.0	DIPLODONTA PUNCTATA	0.2	0.2	1.0	0.8	0.2	0.2	0.2	0.2
100.0	OPHIACTIS SAVIGNYI	.	.	1.2	1.2	0.4	0.2	.	.
100.0	PARAPHOXUS SPINOSUS	1.2	0.2	0.4	0.2	.	.	.	.
100.0	PILUMNUS SAYI	1.2	0.6	0.4	0.2	.	.	.	.
100.0	POLYDORA WEBSTERI	0.8	0.6	0.2	0.2	0.2	0.2	0.4	0.2
100.0	SYLLIS GHACILIS	0.2	0.2	0.4	0.2	.	.	1.0	0.8
100.0	THELEPUS SETOSUS	0.6	0.6	.	.	.	.	1.0	0.6
110.0	AMPHIPODA B	0.2	0.2	.	.	0.8	0.5	0.4	0.4
110.0	ANCISTROSYLLIS RIGIDA	.	.	0.6	0.6	0.2	0.2	0.6	0.4
110.0	BOWMANIELLA PORTORICENSIS	.	.	.	.	.	.	1.4	0.6
110.0	CAPITELLIDAE D	0.4	0.2	.	.	.	.	1.0	0.5
110.0	CHIONE GHUS	0.2	0.2	0.6	0.6	.	.	0.6	0.6
110.0	CORBULA CONTRACTA	0.6	0.4	.	.	0.4	0.2	0.4	0.4
110.0	HYBOSCOLEX LONGISETA	1.0	0.5	.	.	.	.	0.4	0.4

Appendix 3. (Continued)

OVERALL RANK	SPECIES	STATION IS01									
		WINTER		SPRING		SUMMER		FALL			
		MEAN	ST ERR	MEAN	ST ERR	MEAN	ST ERR	MEAN	ST ERR		
110.0	LEMBOS SMITHI	1.2	0.6	0.2	0.2	.	.	.	.		
110.0	MAERA WILLIAMSI	0.2	0.2	.	.	1.2	1.2	0.8	0.6		
110.0	POLYDORA CAECA	0.6	0.2	.	.	.	.	0.2	0.2		
110.0	STROMBIFORMIS BILINEATUS	1.2	0.7	.	.	.	.	0.2	0.2		
110.0	SYLLIS CORNUTA	0.8	0.4	0.6	0.4	.	.	.	.		
110.0	SYNALPHEUS TOWNSENDI	0.2	0.2	0.6	0.4	0.2	0.2	0.4	0.4		
123.5	AMPHIPODA E	0.6	0.4	0.6	0.4	.	.	.	.		
123.5	ANCHIALINA TYPICA	.	.	0.2	0.2	1.0	0.4	.	.		
123.5	CANCER IRRORATUS	1.2	0.6	.	.	.	.	.	.		
123.5	CAPITELLIDAE	0.2	0.2	.	.	0.4	0.4	0.6	0.6		
123.5	CAULLERIELLA KILLARIENSIS	1.2	0.6	.	.	.	.	.	.		
123.5	CUMINGIA TELLINOIDES	0.2	0.2	0.2	0.2	0.6	0.4	0.2	0.2		
123.5	MAERA SP. A	0.2	0.2	.	.	1.0	1.0	.	.		
123.5	MANGELIA RUGIRIMA	0.4	0.2	0.2	0.2	0.2	0.2	0.4	0.4		
123.5	MICROPANOPE SP.	0.2	0.2	1.0	0.8	.	.	0.8	0.8		
123.5	PAGURUS HENDERSONI *	0.4	0.4	.	.	.	.	0.4	0.4		
123.5	PHERUSA INFLATA	0.2	0.2	.	.	0.6	0.2	0.4	0.4		
123.5	SPIO PETTIBONEAE	1.0	0.5	.	.	.	.	0.2	0.2		
123.5	SYNALPHEUS SP.	.	.	1.2	0.6	.	.	.	.		
123.5	TURBONILLA HEMPHILLI	0.2	0.2	.	.	.	.	1.0	1.0		
141.5	ASPIDOSIPHON ALBUS	0.4	0.4	.	.	0.4	0.2	0.2	0.2		
141.5	AUTOLYTUS SP.	.	.	.	.	0.8	0.6	0.2	0.2		
141.5	BODOTRIIDAE A	0.8	0.4	.	.	.	.	0.2	0.2		
141.5	CAPITELLA CAPITATA	0.2	0.2	.	.	0.8	0.5	.	.		
141.5	CARIDEA	0.6	0.6	0.2	0.2	0.2	0.2	0.2	0.2		
141.5	COLOMASTIX SP.	.	.	0.8	0.8	.	.	.	.		
141.5	INACHOIDES FORCEPS	0.8	0.4	0.2	0.2	.	.	.	.		
141.5	ISCHNOCHITONIDAE A	0.8	0.8	.	.	0.2	0.2	.	.		
141.5	LYSIANASSA SP.	.	.	1.0	1.0	.	.	0.2	0.2		
141.5	MARGINELLA AUREOCINCTA	.	.	0.6	0.4	0.2	0.2	0.2	0.2		
141.5	MARGINELLA ROSCIDA	0.6	0.4	.	.	0.4	0.4	.	.		
141.5	MEGALUROPOUS SP.	.	.	.	.	.	.	1.0	0.4		
141.5	MICROPHOLIS PACHYBACTERA	1.0	1.0	.	.	.	.	.	.		
141.5	MITRELLA LUNATA	0.8	0.4	.	.	.	.	0.2	0.2		
141.5	MUREX FLORIFER DILECTUS	0.2	0.2	.	.	.	.	0.8	0.5		
141.5	MUSCULUS LATERALIS	.	.	.	.	.	.	1.0	0.6		
141.5	OCNUS PYGMAEUS	0.2	0.2	0.4	0.2	0.4	0.4	0.4	0.2		
141.5	TELLINA AMERICANA	0.6	0.6	.	.	.	.	0.4	0.2		
141.5	THOR SP.	0.2	0.2	.	.	.	.	0.8	0.6		
141.5	TRICHOBRANCHUS GLACIALIS	0.8	0.8	0.2	0.2	.	.	.	.		
141.5	VERMICULARIA KNORRII	.	.	.	.	.	.	1.0	1.0		
141.5	XANTHIDAE	0.2	0.2	0.4	0.4	.	.	0.4	0.4		
166.5	AMPELISCA SP. C	.	.	0.4	0.2	.	.	0.8	0.6		
166.5	ARBACIA PUNCTULATA	.	.	0.2	0.2	0.2	0.2	0.2	0.2		
166.5	BATRACHONOTUS FRAGOSUS	0.4	0.4	.	.	.	.	0.2	0.2		
166.5	CRATENA KAORUAE	0.8	0.6	.	.	.	.	.	.		
166.5	DODECACERIA CORALLII	0.6	0.6	0.2	0.2	.	.	.	.		
166.5	ERATO MAUGERIAE	0.4	0.2	0.4	0.2	.	.	.	.		
166.5	EUNICE ANTENNATA	.	.	0.4	0.2	.	.	0.4	0.4		
166.5	GLYCYMERIS PECTINATA	0.2	0.2	0.4	0.4	0.2	0.2	.	.		
166.5	HARMOTHOE SP. A	0.8	0.5	.	.	.	.	.	.		
166.5	HIATELLA ARCTICA	0.2	0.2	0.4	0.4	.	.	0.2	0.2		
166.5	HYPOCONCHA SABULOSA	0.2	0.2	.	.	.	.	0.6	0.2		
166.5	LATREUTES PARVULUS	.	.	0.2	0.2	.	.	0.6	0.2		
166.5	LEPTOCHELA PAPULATA	0.2	0.2	.	.	0.2	0.2	0.4	0.2		

Appendix 3. (Continued)

STATION IS01

OVERALL RANK	SPECIES	WINTER		SPRING		SUMMER		FALL	
		MEAN	ST ERR	MEAN	ST ERR	MEAN	ST ERR	MEAN	ST ERR
166.5	MEDIOMASTUS CALIFORNIENSIS	0.8	0.4	.	.	.	.	.	.
166.5	MICROPANOPE SP. *	0.8	0.8	.	.	.	.	.	.
166.5	MICROPROTOPUS SP.	0.8	0.2	.	.	.	.	.	.
166.5	OLIGOCHAETA	0.2	0.2	0.4	0.4	.	.	0.2	0.2
166.5	OSTRACODA	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
166.5	PHYLLODOCE SP.	.	.	0.2	0.2	.	.	0.6	0.4
166.5	PINNIXA SP.	0.6	0.4	0.2	0.2	.	.	.	.
166.5	PLEUROMERIS TRIDENTATA	0.4	0.4	0.4	0.4	.	.	.	.
166.5	POLYCHERIA SP.	0.4	0.4	.	.	0.4	0.4	.	.
166.5	POLYPLACOPHORA A	.	.	.	.	.	.	0.8	0.4
166.5	PROCERAEA SP.	0.2	0.2	0.2	0.2	.	.	0.4	0.4
166.5	PROCESSA SP.	0.2	0.2	.	.	0.2	0.2	0.4	0.4
166.5	SABELLARIA VULGARIS BEAUFORTENSIS	0.8	0.4	.	.	.	.	.	.
166.5	TANYSTYLIDAE A	0.6	0.4	.	.	0.2	0.2	.	.
166.5	XANTHIDAE A	.	.	0.6	0.2	.	.	0.2	0.2
197.0	ALPHEUS NORMANNI	0.2	0.2	.	.	0.4	0.2	.	.
197.0	AMERICARDIA MEDIA	.	.	0.4	0.2	0.2	.	.	.
197.0	AMPHIPODA J	.	.	0.2	0.2	0.4	0.4	.	.
197.0	ANOPLDACTYLUS PETIOLATUS	0.2	0.2	0.2	0.2	.	.	0.2	0.2
197.0	CERATONEREIS SP.	.	.	0.6	0.4	.	.	.	.
197.0	CHRYSOPETALIDAE B	0.4	0.2	.	.	.	.	0.2	0.2
197.0	CORBULA BARRATTIANA	.	.	0.6	0.4	.	.	.	.
197.0	DIASTYLIDAE A	0.2	0.2	.	.	0.4	0.2	.	.
197.0	EULALIA MACROCEROS	.	.	.	.	.	.	0.6	0.4
197.0	GONIADIDES CAROLINAE	.	.	.	.	0.4	0.4	0.2	0.2
197.0	HYDROIDES SP. B	.	.	.	.	0.2	0.2	0.4	0.4
197.0	LEMBOS UNIFASCIATUS REDUCTUS	.	.	.	.	0.2	0.2	0.4	0.4
197.0	LITHOPHAGA ARISTATA	.	.	0.4	0.4	.	.	.	.
197.0	LUCAPINELLA LIMATULA	0.2	0.2	.	.	.	.	0.4	0.2
197.0	LYONSIA BEANA	.	.	.	.	0.2	0.2	0.4	0.2
197.0	MAJIDAE	.	.	.	.	.	.	0.6	0.4
197.0	MALDANIDAE	0.2	0.2	0.2	0.2	.	.	0.2	0.2
197.0	MARGINELLA CARNEA	.	.	0.2	0.2	0.2	0.2	0.2	0.2
197.0	MEGALOMMA SP. A	0.6	0.4	.	.	.	.	.	.
197.0	MONOCULODES SP.	0.4	0.2	0.2	0.2	.	.	.	.
197.0	NEOPONTONIDES BEAUFORTENSIS	.	.	0.2	0.2	.	.	0.4	0.2
197.0	PELECYPODA B	0.2	0.2	0.4	0.4	.	.	.	.
197.0	PETALOPROCTUS SOCIALIS	0.6	0.2	.	.	.	.	.	.
197.0	PISTA PALMATA	0.2	0.2	.	.	.	.	0.4	0.4
197.0	PSEUDEURYTHOE AMBIGUA	.	.	0.2	0.2	0.4	0.2	.	.
197.0	PSEUDOVERMILIA SP. A	0.2	0.2	0.2	0.2	.	.	0.2	0.2
197.0	SABELLIDAE	0.2	0.2	.	.	0.4	0.2	.	.
197.0	SERPULIDAE H	.	.	.	.	.	.	0.6	0.4
197.0	SERPULIDAE K	.	.	0.2	0.2	0.4	0.2	.	.
197.0	SICYONIA TYPICA	.	.	.	.	.	.	0.6	0.6
197.0	SPIONIDAE B	0.4	0.4	0.2	0.2	.	.	.	.
197.0	TIRON TROPAKIS	0.4	0.4	.	.	.	.	0.2	0.2
197.0	VARICORBULA OPERCULATA	0.6	0.2	.	.	.	.	.	.
244.5	ACANTHOHAUSTORIUS MILLSI	0.4	0.2	.	.	.	.	.	.
244.5	AMPHIODIA TRYCHNA	.	.	.	.	.	.	0.4	0.4
244.5	ANOPLDACTYLUS INSIGNIS	0.2	0.2	.	.	.	.	0.2	0.2
244.5	APANTHURA MAGNIFICA	.	.	.	.	0.4	0.2	.	.
244.5	AUSTRAEOLIS CATINA	0.4	0.2	.	.	.	.	.	.
244.5	CADULUS QUADRIDENTATUS	.	.	.	.	0.2	0.2	0.2	0.2
244.5	CHAETOPTERIDAE	0.4	0.2	.	.	.	.	.	.

Appendix 3. (Continued)

OVERALL RANK	SPECIES	STATION IS01											
		WINTER			SPRING			SUMMER			FALL		
		MEAN	ST	ERR	MEAN	ST	ERR	MEAN	ST	ERR	MEAN	ST	ERR
244.5	CHIONE LATILIRATA	.	.	.	.	.	.	.	.	.	0.4	.	0.2
244.5	CRASSISPIRA ALBOMACULATA	.	.	.	.	.	.	0.2	.	0.2	0.2	.	0.2
244.5	CREPIDULA PLANA	0.4	.	0.4	.	.	.	.	.	.	.	.	.
244.5	CREPIDULA SP. *	0.4	.	0.2	.	.	.	.	.	.	.	.	.
244.5	DIODORA CAYENENSIS	.	.	.	.	.	.	0.2	.	0.2	0.2	.	0.2
244.5	DROMIDIA ANTILLENIS	.	.	.	.	.	.	.	.	.	0.4	.	0.4
244.5	EBALIA STIMPSONI	.	.	.	.	.	.	.	.	.	0.4	.	0.4
244.5	EDEIS SPINOSA	.	.	.	.	.	.	0.2	.	0.2	0.2	.	0.2
244.5	ERICHTHONIUS SP. A	0.4	.	0.4	.	.	.	.	.	.	0.2	.	0.2
244.5	FAVARTIA CELLULOSA	0.2	.	0.2	.	.	.	.	.	.	.	.	.
244.5	HEXAPANOPUS SP. *	0.4	.	0.2	.	.	.	.	.	.	.	.	.
244.5	HOLOTHUROIDEA	.	.	.	.	.	.	0.2	.	0.2	0.2	.	0.2
244.5	HYALINA ALBOLINEATA	0.2	.	0.2	0.2	.	0.2	.	.	.	0.4	.	0.4
244.5	INACHOIDES FORCEPS *	.	.	.	.	.	.	.	.	.	0.4	.	0.4
244.5	LEMBOS SPINICARPUS INERMIS	.	.	.	.	.	.	.	.	.	0.4	.	0.4
244.5	LUMBRINERIS COCCINEA	0.2	.	0.2	.	.	.	0.4	.	0.2	0.2	.	0.2
244.5	LYONSIA SP.	.	.	.	.	.	.	.	.	.	0.4	.	0.4
244.5	LYSIANASSIDAE A	.	.	.	.	.	.	.	.	.	0.4	.	0.4
244.5	MARPHYSA SP. A	0.2	.	0.2	0.2	.	0.2	.	.	.	0.4	.	0.4
244.5	MICROPHRYS ANTILLENIS	.	.	.	.	.	.	.	.	.	0.4	.	0.4
244.5	MITHRAX SP.	.	.	.	.	.	.	.	.	.	0.4	.	0.4
244.5	NEREIDAE	.	.	.	.	.	.	.	.	.	0.4	.	0.2
244.5	NEREIS SP.	0.2	.	0.2	.	.	.	0.2	.	0.2	.	.	.
244.5	NICOMACHE TRISPINATA	0.2	.	0.2	0.2	.	0.2	.	.	.	.	.	.
244.5	NOTOMASTUS HEMIPODUS	0.4	.	0.4	.	.	.	.	.	.	.	.	.
244.5	OLIVELLA MINUTA	.	.	.	.	.	.	0.2	.	0.2	0.2	.	0.2
244.5	OPHIUROIDEA	0.4	.	0.4	.	.	.	.	.	.	0.4	.	0.4
244.5	PERICLIMENES LONGICAUDATUS	.	.	.	.	.	.	.	.	.	.	.	.
244.5	PERICLIMENES SP.	0.4	.	0.2	.	.	.	.	.	.	.	.	.
244.5	PERUSA EHLERSI	0.4	.	0.2	.	.	.	.	.	.	.	.	.
244.5	POLYCHAETA	.	.	.	.	.	.	0.4	.	0.2	.	.	.
244.5	POLYCIRRUS EXIMIUS	.	.	.	0.2	.	0.2	.	.	.	0.2	.	0.2
244.5	PONTOGENIA SERICOMA	0.2	.	0.2	.	.	.	.	.	.	0.2	.	0.2
244.5	PROCESSA BERMUDENSIS	0.4	.	0.4	.	.	.	.	.	.	.	.	.
244.5	SABELLARIA VULGARIS	0.4	.	0.2	.	.	.	.	.	.	.	.	.
244.5	SCOLOPLOS SP. B	.	.	.	.	.	.	0.4	.	0.4	.	.	.
244.5	SEMELE NUCULOIDES	.	.	.	0.2	.	0.2	0.2	.	0.2	.	.	.
244.5	SICYONIA BREVIROSTRIS	0.2	.	0.2	.	.	.	.	.	.	0.2	.	0.2
244.5	SPIROBIS SP.	.	.	.	.	.	.	.	.	.	0.4	.	0.4
244.5	STENOCHOE GEORGIANA	0.4	.	0.2	.	.	.	.	.	.	.	.	.
244.5	SYLLIDAE	.	.	.	.	.	.	.	.	.	0.4	.	0.2
244.5	SYLLIDAE L	.	.	.	.	.	.	0.2	.	0.2	0.2	.	0.2
244.5	SYLLIDAE N	.	.	.	.	.	.	0.2	.	0.2	0.2	.	0.2
244.5	SYLLIS ALTERNATA	.	.	.	.	.	.	0.2	.	0.2	0.2	.	0.2
244.5	SYLLIS FERRUGINA	.	.	.	.	.	.	0.2	.	0.2	0.2	.	0.2
244.5	SYLLIS REGULATA CAROLINAE	.	.	.	0.2	.	0.2	0.2	.	0.2	0.2	.	0.2
244.5	SYNCHELIDIUM AMERICANUM	0.4	.	0.4	.	.	.	.	.	.	.	.	.
244.5	TEREBELLIDAE	0.4	.	0.4	.	.	.	.	.	.	.	.	.
244.5	TRITONIA BAYERI MISA	.	.	.	0.2	.	0.2	.	.	.	0.2	.	0.2
244.5	TRYPANOSYLLIS SP.	0.4	.	0.4	.	.	.	.	.	.	.	.	.
244.5	TRYPANOSYLLIS ZEBRA	.	.	.	0.2	.	0.2	.	.	.	0.2	.	0.2
244.5	TURBO CASTANEA	0.2	.	0.2	.	.	.	.	.	.	0.2	.	0.2
244.5	TURRONILLA INCISA	0.4	.	0.4	.	.	.	.	.	.	.	.	.
244.5	WEBSTERINEREIS SP. A	.	.	.	.	.	.	0.2	.	0.2	0.2	.	0.2
244.5	XANTHIDAE B	.	.	.	.	.	.	0.2	.	0.2	0.2	.	0.2

Appendix 3. (Continued)

OVERALL RANK	SPECIES	STATION		IS01													
		WINTER		SPRING		SUMMER			FALL								
		MEAN	ST ERR	MEAN	ST ERR	MEAN	ST	ERR	MEAN	ST	ERR	MEAN	ST	ERR	MEAN	ST	ERR
351.0	ACROCIRRIDAE A	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.	.	.	.
351.0	ALPHEIDAE	.	.	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.	.
351.0	AMAEA RETIFERA	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.	.	.	.
351.0	AMPELISCA CRISTOIDES	.	.	.	.	.	.	.	.	.	.	0.2	.	0.2	.	.	.
351.0	AMPHARETE ACUTIFRONS	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.	.	.	.
351.0	AMPHILOCHUS SP.	.	.	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.	.
351.0	AMPHIPODA G	.	.	.	.	0.2	0.2	.	.	.	.	.	.	.	.	.	.
351.0	AMPHIPODA H	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.	.	.	.
351.0	ANACHIS AVARA *	.	.	.	.	0.2	0.2	.	.	.	.	.	.	.	.	.	.
351.0	ANACHIS FLORIDANA	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.	.	.	.
351.0	ARCA SP.	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.	.	.	.
351.0	ARCA ZEBRA	.	.	.	.	.	.	.	.	.	.	0.2	.	0.2	.	.	.
351.0	ARCOPSIS ADAMSI	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.	.	.	.
351.0	ASTERIAS FORBESII	.	.	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.	.
351.0	ASTHENOTHAERUS HEMPHILLI	.	.	.	.	0.2	0.2	.	.	.	.	0.2	.	0.2	.	.	.
351.0	AUTOMATE EVERMANI	.	.	.	.	.	.	.	.	.	.	0.2	.	0.2	.	.	.
351.0	AXIOGNATHUS SQUAMATUS	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.	.	.	.
351.0	BRANCHIOSYLLIS OCLATA	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.	.	.	.
351.0	BRANIA CLAVATA	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.	.	.	.
351.0	CALOTROPHON OSTREARUM	.	.	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.	.
351.0	CANTHARUS MULTANGULUS	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.	.	.	.
351.0	CERADOCUS SP. B	.	.	.	.	0.2	0.2	.	.	.	.	0.2	.	0.2	.	.	.
351.0	CERAPUS TUBULARIS	.	.	.	.	.	.	.	.	.	.	0.2	.	0.2	.	.	.
351.0	CERATONEREIS VERSIPEDATA	.	.	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.	.
351.0	CERODRILLIA CERVINA	.	.	.	.	0.2	0.2	.	.	.	.	.	.	.	.	.	.
351.0	CHRYSOPETALIDAE C	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.	.	.	.
351.0	CHRYSOPETALIDAE D	.	.	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.	.
351.0	CIROLANA PARVA	.	.	.	.	.	.	.	.	.	.	0.2	.	0.2	.	.	.
351.0	CIRRIFORMIA FILIGERA	.	.	.	.	.	.	.	.	.	.	0.2	.	0.2	.	.	.
351.0	CONUS JASPIDEUS STEARNSI	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.	.	.	.
351.0	COOPERELLA ATLANTICA	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.	.	.	.
351.0	COROPHIIDAE	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.	.	.	.
351.0	COROPHIUM SP.	.	.	.	.	.	.	.	.	.	.	0.2	.	0.2	.	.	.
351.0	CREPIDULA SP.	.	.	.	.	0.2	0.2	.	.	.	.	.	.	.	.	.	.
351.0	CRYOTURRIS ELATA	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.	.	.	.
351.0	DELECTOPECTEN SP. A	.	.	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.	.
351.0	DENDROCHIROTTIDA	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.	.	.	.
351.0	DENDRODORIS SP. A	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.	.	.	.
351.0	DIASTYLIDAE	.	.	.	.	.	.	.	.	.	.	.	.	.	0.2	.	0.2
351.0	DIODORA LISTERI	.	.	.	.	.	.	.	.	.	.	0.2	.	0.2	.	.	.
351.0	DONDICE OCCIDENTALIS	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.	.	.	.
351.0	DORIOPSISILLA ALBOPUNCTATA	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.	.	.	.
351.0	EBALIA CARIOSA	.	.	.	.	.	.	.	.	.	.	0.2	.	0.2	.	.	.
351.0	EUNICE WEBSTERI	.	.	.	.	.	.	.	.	.	.	0.2	.	0.2	.	.	.
351.0	EUNICIDAE	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.	.	.	.
351.0	EUPHROSINE TRILOBA	.	.	.	.	.	.	.	.	.	.	0.2	.	0.2	.	.	.
351.0	EURYDICE LITTORALIS	.	.	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.	.
351.0	EXOGONE SP.	.	.	.	.	0.2	0.2	.	.	.	.	.	.	.	.	.	.
351.0	GASTROPTERON RUBRUM	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.	.	.	.
351.0	GLYCERA TESSELATA	.	.	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.	.
351.0	GONEPLAX SIGSBEEI	.	.	.	.	0.2	0.2	.	.	.	.	.	.	.	.	.	.
351.0	HETEROMYSIS FORMOSA	.	.	.	.	.	.	.	.	.	.	0.2	.	0.2	.	.	.
351.0	HIPPOLYTIDAE	.	.	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.	.
351.0	JAEROPSIS CORALICOLA	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.	.	.	.
351.0	KURTZIELLA ATROSTYLA	.	.	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.	.

Appendix 3. (Continued)

OVERALL RANK	SPECIES	STATION IS01											
		WINTER			SPRING			SUMMER			FALL		
		MEAN	ST	ERR	MEAN	ST	ERR	MEAN	ST	ERR	MEAN	ST	ERR
351.0	KURTZIELLA LIMONITELLA	.	.	.	.	.	.	0.2	0.2	.	.	.	.
351.0	LAEVICARDIUM SP.	.	.	.	.	.	.	.	.	.	.	.	.
351.0	LAMELLARIA PERSPICUA	0.2	0.2	.	0.2	0.2	.	.	.	.	.	.	.
351.0	LATREUTES FUCORUM *	.	.	.	.	.	.	.	.	.	0.2	0.2	.
351.0	LEMBOS SP.	.	.	.	0.2	0.2	.	.	.	.	.	.	.
351.0	LEPTOCHELA SP.	.	.	.	.	.	.	0.2	0.2	.	.	.	.
351.0	LILJEBORGIA SP. B	.	.	.	0.2	0.2	.	.	.	.	.	.	.
351.0	LINGA AMIANTUS	.	.	.	.	.	.	0.2	0.2	.	.	.	.
351.0	LUMBRINERIS SP.	.	.	.	.	.	.	0.2	0.2	.	.	.	.
351.0	LYSIDICE NINETTA	.	.	.	.	.	.	.	.	.	0.2	0.2	.
351.0	MACROCOELOMA SP.	.	.	.	.	.	.	.	.	.	0.2	0.2	.
351.0	MACROCOELOMA CAMPTOCERUM	0.2	0.2	.	.	.	.	.	.	.	.	.	.
351.0	MARGINELLA LAVALLEEANA	.	.	.	.	.	.	.	.	.	0.2	0.2	.
351.0	MARGINELLA SP.	.	.	.	0.2	0.2	.	.	.	.	.	.	.
351.0	MARPHISA SP. B	.	.	.	.	.	.	0.2	0.2	.	.	.	.
351.0	MEGALOMMA SP.	.	.	.	0.2	0.2	.	.	.	.	.	.	.
351.0	MESOCHAETOPTERUS SP.	0.2	0.2	.	.	.	.	.	.	.	.	.	.
351.0	MICROPHRYS SP.	.	.	.	0.2	0.2	.	.	.	.	.	.	.
351.0	MICROPHRYS SP. *	.	.	.	0.2	0.2	.	.	.	.	.	.	.
351.0	MITHRAX ACUTICORNIS	.	.	.	.	.	.	.	.	.	0.2	0.2	.
351.0	MITHRAX SP. *	.	.	.	.	.	.	0.2	0.2	.	.	.	.
351.0	MUNIDA SP.	0.2	0.2	.	.	.	.	.	.	.	.	.	.
351.0	MUREX RECURVIOSTRIS RUBIDUS	.	.	.	.	.	.	0.2	0.2	.	.	.	.
351.0	MYSIDACEA	.	.	.	.	.	.	0.2	0.2	.	.	.	.
351.0	MYSIDOPSIS FURCA	.	.	.	0.2	0.2	.	.	.	.	.	.	.
351.0	NASSARIUS ALBUS	.	.	.	.	.	.	.	.	.	0.2	0.2	.
351.0	NASSARIUS VIBEX	.	.	.	0.2	0.2	.	.	.	.	.	.	.
351.0	NEMERTINEA C	.	.	.	.	.	.	0.2	0.2	.	.	.	.
351.0	NEMERTINEA E	0.2	0.2	.	.	.	.	.	.	.	.	.	.
351.0	NEODRILLIA CYDIA	.	.	.	0.2	0.2	.	.	.	.	.	.	.
351.0	NEPHTYIDAE	0.2	0.2	.	.	.	.	.	.	.	.	.	.
351.0	NEPHTYS SQUAMOSA	.	.	.	0.2	0.2	.	.	.	.	.	.	.
351.0	NEREIS SUCCINEA	.	.	.	.	.	.	.	.	.	0.2	0.2	.
351.0	NUCULA PROXIMA	.	.	.	.	.	.	.	.	.	0.2	0.2	.
351.0	NYMPHOPSIS DUODORSOSPINOSA	.	.	.	.	.	.	0.2	0.2	.	.	.	.
351.0	OCTOPUS SP.	.	.	.	.	.	.	0.2	0.2	.	.	.	.
351.0	ODONTOSYLLIS SP.	.	.	.	.	.	.	.	.	.	0.2	0.2	.
351.0	ODONTOSYLLIS ENOPLA	.	.	.	.	.	.	.	.	.	0.2	0.2	.
351.0	OLIVA SP.	.	.	.	0.2	0.2	.	.	.	.	.	.	.
351.0	OLIVELLA BULLULA	.	.	.	.	.	.	0.2	0.2	.	.	.	.
351.0	ONUPHIDAE	0.2	0.2	.	.	.	.	.	.	.	.	.	.
351.0	ONUPHIS SP.	.	.	.	.	.	.	0.2	0.2	.	.	.	.
351.0	OPHTHOCNIDA SCABRIUSCULA	0.2	0.2	.	.	.	.	.	.	.	.	.	.
351.0	OPHTHOLEPIS ELEGANS	.	.	.	.	.	.	.	.	.	0.2	0.2	.
351.0	OXYUROSTYLIS SMITHI	.	.	.	.	.	.	.	.	.	0.2	0.2	.
351.0	PENAEOIDEA	.	.	.	.	.	.	.	.	.	0.2	0.2	.
351.0	PERICLIMENAEUS SP. A	0.2	0.2	.	.	.	.	.	.	.	.	.	.
351.0	PHERUSA SP.	.	.	.	.	.	.	.	.	.	0.2	0.2	.
351.0	PHITISICA MARINA	.	.	.	.	.	.	.	.	.	0.2	0.2	.
351.0	PHYLLODOCE FRAGILIS	.	.	.	.	.	.	.	.	.	0.2	0.2	.
351.0	PHYLLODOCE LONGIPES	0.2	0.2	.	.	.	.	.	.	.	.	.	.
351.0	PHYLLODOCIDAE	.	.	.	0.2	0.2	.	.	.	.	.	.	.
351.0	PILUMNUS DASYPODUS	0.2	0.2	.	.	.	.	.	.	.	.	.	.
351.0	PILUMNUS FLORIDANUS	.	.	.	0.2	0.2	.	.	.	.	.	.	.
351.0	PILUMNUS PANNOSUS	.	.	.	0.2	0.2	.	.	.	.	.	.	.

Appendix 3. (Continued)

		STATION		IS01													
OVERALL RANK	SPECIES	WINTER		SPRING		SUMMER			FALL								
		MEAN	ST ERR	MEAN	ST ERR	MEAN	ST	ERR	MEAN	ST	ERR						
351.0	PINNIXA CYLINDRICA	.	.	0.2	0.2	.	.	.	.	.	.	.	.	.			
351.0	PINNOTHERES SP.	.	.	.	.	0.2	.	0.2	.	.	.	.	.	.			
351.0	PIONOSYLLIS SP. A	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.			
351.0	PISANIA TINCTA	.	.	0.2	0.2	.	.	.	.	.	.	.	.	.			
351.0	PLICATULA GIBBOSA	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.			
351.0	PODOCHELA SP.	.	.	.	.	.	.	.	.	.	.	0.2	0.2	.			
351.0	POLINICES SP.	.	.	0.2	0.2	.	.	.	.	.	.	.	.	.			
351.0	PORCELLANIDAE	0.2	0.2	.	.	.	.	.	.	.	.	0.2	0.2	.			
351.0	PORTUNIDAE	.	.	.	.	.	.	.	.	.	.	0.2	0.2	.			
351.0	PROCERAEA FASCIATA	0.2	0.2	.	.	.	.	.	0.2	0.2	.	0.2	0.2	.			
351.0	PROCESSA HEMPHILLI	.	.	.	.	.	.	.	.	.	.	.	.	.			
351.0	PROCESSA VICINA	.	.	.	.	.	.	.	.	.	.	0.2	0.2	.			
351.0	PROTULA TUBULARIA	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.			
351.0	PSEUDOVERMILIA OCCIDENTALIS	.	.	0.2	0.2	.	.	.	0.2	0.2	.	.	.	.			
351.0	PTERIA COLYMBUS	.	.	.	.	.	.	.	.	.	.	.	.	.			
351.0	SABELLARIA GRACILLIS	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.			
351.0	SABELLARIA VULGARIS VULGARIS	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.			
351.0	SABELLARIIDAE	.	.	.	.	0.2	.	0.2	.	.	.	.	.	.			
351.0	SCALIRREGMIDAE	.	.	.	.	0.2	0.2	.	.	.	.	.	.	.			
351.0	SCOLOPLOS SP. A	.	.	.	.	.	.	.	0.2	0.2	.	.	.	.			
351.0	SEILA ADAMSI	0.2	0.2	.	.	.	.	.	.	.	.	0.2	0.2	.			
351.0	SEMELE BELLASTRIATA	.	.	.	.	.	.	.	0.2	0.2	.	0.2	0.2	.			
351.0	SEMELE PURPURASCENS	.	.	.	.	.	.	.	0.2	0.2	.	0.2	0.2	.			
351.0	SERPULIDAE A	.	.	.	.	.	.	.	.	.	.	.	.	.			
351.0	SICYONIA LAEVIGATA	.	.	.	.	.	.	.	0.2	0.2	.	.	.	.			
351.0	SIPUNCULIDA C	.	.	0.2	0.2	.	.	.	.	.	.	.	.	.			
351.0	SIRAIUS BICOLOR	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.			
351.0	SPELOEOPHORUS ELEVATUS	.	.	.	.	.	.	.	.	.	.	0.2	0.2	.			
351.0	STENOTHOIDAE	.	.	0.2	0.2	.	.	.	.	.	.	.	.	.			
351.0	STENOTHOIDAE A	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.			
351.0	SYLLIDAE H	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.			
351.0	TELLINA AEQUISTRIATA	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.			
351.0	TELLINA SP.	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.			
351.0	THARYX MARIONI	.	.	.	.	.	.	.	.	.	.	0.2	0.2	.			
351.0	THYONE PSEUDOFUSUS	.	.	.	.	.	.	.	.	.	.	0.2	0.2	.			
351.0	TOZEUMA SP.	.	.	.	.	.	.	.	0.2	0.2	.	.	.	.			
351.0	TRITONIA BAYERI	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.			
351.0	TURBONILLA ABRUPTA	.	.	0.2	0.2	.	.	.	.	.	.	.	.	.			
351.0	TURBONILLA INTERRUPTA	.	.	0.2	0.2	.	.	.	.	.	.	.	.	.			
351.0	WILLIAMIA KREBSII	.	.	.	.	.	.	.	0.2	0.2	.	.	.	.			
351.0	XANTHIDAE *	.	.	0.2	0.2	.	.	.	.	.	.	.	.	.			

Appendix 3. (Continued)

OVERALL RANK	SPECIES	STATION		IS02										
		WINTER		SPRING		SUMMER			FALL					
		MEAN	ST ERR	MEAN	ST ERR	MEAN	ST	ERR	MEAN	ST	ERR	MEAN	ST	ERR
1.0	LUCONACIA INCERTA	68.0	19.5	2.0	0.4	0.2		0.2	0.4		0.4		0.4	
2.0	ELASMOPUS SP. A	14.0	4.5	13.8	7.2	12.2		4.7	5.0		2.2		2.2	
3.0	ERICHTHONIUS BRASILIENSIS	27.0	11.3	16.8	7.1				0.4		0.4		0.4	
4.0	LEMBOS SMITHI	15.2	7.8	6.8	2.9	8.6		5.6	5.2		2.6		2.6	
5.0	CAPRELLA EQUILIBRA	29.8	9.3	3.2	0.9									
6.0	PODOCERUS SP.	26.4	7.8	4.0	1.3	0.8		0.6	0.8		0.4		0.4	
7.0	PHOTIS SP.	20.4	5.0	10.8	4.9				0.4		0.2		0.2	
8.0	LEPTOCHELIA SP.	17.2	7.3	7.8	4.6	0.4		0.4						
9.0	LUMBRINERIS INFLATA	8.0	3.6	8.6	3.4	3.0		1.0	2.2		0.9		0.9	
10.0	POLYCIRRUS CAROLINENSIS	5.6	1.7	8.4	4.2	1.8		0.6	1.0		0.5		0.5	
11.0	EXOGONE DISPAR	6.6	2.6	6.0	3.9	1.2		0.6	2.8		1.4		1.4	
12.0	GAMMAROPSIS SP.	0.2	0.2	16.0	10.0									
13.0	SPIO SP. A	4.4	4.4	11.0	10.8									
14.5	AMPITHOE SP. A	3.2	1.1	6.2	3.4	2.2		1.3	2.4		1.1		1.1	
14.5	MEGALOBRACHIUM SORIATUM	7.2	2.1	2.0	0.9	4.4		2.5	0.4		0.2		0.2	
16.0	PSEUDOMEDAEUS AGASSIZII	4.2	1.7	1.6	0.9	7.0		4.5	0.8		0.6		0.6	
17.0	CARPIAS BERMUDENSIS	0.6	0.4	9.2	5.3	2.4		1.3	1.2		0.4		0.4	
18.0	ARCA IMBRICATA	2.6	1.0	4.2	1.3	5.8		2.1	0.6		0.6		0.6	
19.0	SYLLIS GRACILIS	6.4	1.6	1.8	0.7	3.2		1.3	0.8		0.5		0.5	
20.0	MICRODEUTOPUS MYERSI	10.0	1.9	0.4	0.4				1.4		0.5		0.5	
21.0	OXYUROSTYLIS SMITHI	0.2	0.2	1.6	1.0				9.6		0.7		0.7	
22.0	MAERA SP. A	4.0	1.9	4.4	1.3	2.0		1.8	0.6		0.2		0.2	
24.0	PILUMNUS SP.	4.2	1.3	3.0	1.3	2.0		1.1	1.6		0.7		0.7	
24.0	SIPUNCULIDA A	4.8	1.5	2.8	1.8	3.0		1.1	0.2		0.2		0.2	
24.0	SYLLIS HYALINA	3.0	0.9	2.2	1.0	2.4		0.6	3.2		1.1		1.1	
26.0	OPHIACTIS SAVIGNYI	6.2	1.5	2.4	0.7				1.6		1.0		1.0	
27.5	ASPIDOSIPHON GOSNOLDI	1.4	0.7	5.8	2.0	2.4		0.9						
27.5	OPHIOTHRIX ANGULATA	4.4	1.4	3.8	1.3	0.8		0.5	0.6		0.4		0.4	
29.0	MALACOCEROS GLUTAEUS	0.2	0.2	0.4	0.2	1.8		0.9	7.0		2.5		2.5	
30.5	NEMATODA	6.2	4.7	2.4	1.0	0.2		0.2	0.2		0.2		0.2	
30.5	PODARKE OBSCURA	0.4	0.2	4.8	1.7	3.4		0.9	0.4		0.4		0.4	
32.0	CHAMA CONGREGATA	3.2	1.2	1.6	0.7	3.2		1.4	0.4		0.4		0.4	
33.0	SYNALPHEUS TOWNSENDI	1.4	1.2	0.6	0.4	4.0		1.9	2.0		1.0		1.0	
34.5	ASTERIAS FORBESII			7.8	1.3									
34.5	SYLLIS SPONGICOLA	1.4	1.2	1.2	0.7	3.6		2.4	1.6		1.0		1.0	
36.5	APSEUDES SP. A	2.8	1.2	3.4	1.9	1.2		0.5	0.2		0.2		0.2	
36.5	CREPIDULA ACULEATA	3.4	1.9	1.6	1.1	2.6		1.0						
38.0	AMPELISCA SCHELLENBERGI	3.8	1.1	1.4	1.0	1.8		0.8	0.4		0.4		0.4	
39.0	EULALIA SANGUINEA	1.6	0.6	2.0	0.7	1.6		0.7	1.0		0.6		0.6	
40.0	PARACERCEIS CAUDATA	1.8	0.6	0.4	0.2	0.8		0.4	3.0		2.5		2.5	
42.0	MELITA APPENDICULATA			4.0	3.3	1.2		0.7	0.4		0.4		0.4	
42.0	PISTA QUADRILOBATA	1.0	0.6	2.8	1.4	1.4		0.7	0.4		0.2		0.2	
42.0	XANTHIDAE A	0.8	0.6	1.2	1.2	3.2		1.5	0.4		0.2		0.2	
44.5	ARMANDIA MACULATA	5.4	3.4											
44.5	TURBONILLA INTERRUPTA	3.2	2.9	1.2	0.6	1.0		1.0						
46.5	AMPHARETE AMERICANA	4.0	1.0	1.2	1.0									
46.5	JASSA FALCATA	5.2	2.3											
48.0	PRIONOSPION CIRRIFERA	4.0	1.6	0.4	0.4	0.4		0.4	0.2		0.2		0.2	
50.5	AXIOGNATHUS SQUAMATUS	1.2	0.7	3.0	2.5				0.4		0.4		0.4	
50.5	GOULDIA CERINA	0.8	0.8	0.8	0.5	3.0		2.0						
50.5	LEMBOS UNICORNIS	1.2	1.2	2.2	1.5	1.0		0.4	0.2		0.2		0.2	
50.5	TURRONILLA INCISA	4.6	4.3											
53.0	CHRYSOPETALIDAE A	0.4	0.2	2.6	1.7	0.8		0.6	0.4		0.4		0.4	
54.5	AMPHIODIA PULCHELLA			2.8	2.8				1.2		0.4		0.4	
54.5	LYSIANOPSIS ALBA	0.8	0.4	0.8	0.4	1.6		0.7	0.8		0.6		0.6	

Appendix 3. (Continued)

OVERALL RANK	SPECIES	STATION		IS02										
		WINTER		SPRING		SUMMER			FALL					
		MEAN	ST ERR	MEAN	ST ERR	MEAN	ST	ERR	MEAN	ST	ERR	MEAN	ST	ERR
57.0	ANOPILODACTYLUS PETIOLATUS	1.8	0.7	1.8	0.9	.	.	.	.	.	.	.	.	.
57.0	MITRELLA LUNATA	3.0	1.7	0.6	0.4	.	.	.	.	.	.	.	.	.
57.0	PHERUSA EHLERSI	1.4	0.5	1.0	0.8	0.8	0.5	0.4	0.4	0.4	0.4	0.4	0.4	0.4
60.5	BRACHYURA	2.8	1.8	0.4	0.4	0.2	0.2	.	.	.	.	.	.	.
60.5	ERICHTHONIUS SP. A	0.2	0.2	1.2	1.0	2.0	2.0	.	.	.	.	.	.	.
60.5	LEUCOTHOE SPINICARPA	0.2	0.2	.	.	2.4	1.5	0.8	0.6	0.6	0.6	0.6	0.6	0.6
60.5	SYNALPHEUS SP.	0.2	0.2	.	.	3.2	1.6	0.6	0.4	0.6	0.4	0.6	0.4	0.4
64.0	MEGALOMMA BIOCCULATUM	0.4	0.2	1.6	0.7	0.6	0.4	0.6	0.2	0.2	0.2	0.2	0.2	0.2
64.0	PILUMNUS FLORIDANUS	2.6	1.1	0.4	0.2	.	.	.	2.4	1.0	1.0	1.0	1.0	1.0
64.0	SYLLIS SP. B	.	.	0.8	0.4	.	.	.	.	.	.	.	.	.
67.0	AMPHIPODA E	0.4	0.4	2.0	1.3	0.6	0.6	0.6	2.4	2.4	2.4	2.4	2.4	2.4
67.0	FILOGRANA IMPLEXA	0.4	0.4	0.2	0.2	.	.	.	0.2	0.2	0.2	0.2	0.2	0.2
67.0	SERPULIDAE	0.2	0.2	2.2	1.3	0.4	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
69.5	APANTHURA MAGNIFICA	1.2	0.8	0.8	0.6	0.8	0.8	.	.	.	.	.	.	.
69.5	PHYLLODOCE CASTANEA	0.2	0.2	1.2	0.4	1.4	0.7	.	.	.	.	.	.	.
72.5	AMPELISCA SP. C	.	.	.	.	1.2	1.2	1.4	0.7	0.7	0.7	0.7	0.7	0.7
72.5	LYSIANASSA SP.	0.8	0.8	1.6	0.9	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
72.5	PHERUSA INFLATA	0.4	0.2	1.0	0.6	1.0	0.4	0.2	0.2	0.2	0.2	0.2	0.2	0.2
72.5	SYLLIS ALTERNATA	1.2	1.0	.	.	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6
78.5	CERATONEREIS MIRABILIS	.	.	.	.	1.2	0.6	1.2	0.6	1.2	0.6	1.2	0.6	1.2
78.5	JAEROPSIS CORALICOLA	.	.	1.6	1.0	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8
78.5	LATREUTES PARVULUS	0.2	0.2	1.6	0.9	0.2	0.2	0.2	0.4	0.4	0.4	0.4	0.4	0.4
78.5	MEGALUROPIUS SP.	.	.	0.2	0.2	.	.	.	2.2	1.0	1.0	1.0	1.0	1.0
78.5	NEMERTINEA	0.8	0.4	0.6	0.4	.	.	.	1.0	0.5	0.5	0.5	0.5	0.5
78.5	PAGURUS SP.	1.4	1.0	0.2	0.2	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4
78.5	POLYDORA TETRABRANCHIA	.	.	2.2	1.6	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
78.5	WEBSTERINEREIS TRIDENTATA	0.2	0.2	1.0	0.4	1.0	0.4	0.2	0.2	0.2	0.2	0.2	0.2	0.2
85.0	ALPHEUS NORMANNI	0.6	0.4	0.6	0.6	0.2	0.2	0.2	0.8	0.4	0.4	0.4	0.4	0.4
85.0	CAPRELLA PENANTIS	1.0	0.5	1.2	0.6	.	.	.	1.0	0.8	0.8	0.8	0.8	0.8
85.0	GONIADIDES CAROLINAE	.	.	0.8	0.8	0.4	0.4	0.4	1.0	0.8	0.8	0.8	0.8	0.8
85.0	PILUMNUS DASYPODUS	2.0	0.9	0.2	0.2	.	.	.	.	.	.	.	.	.
85.0	THARYX MARIONI	2.2	2.2	.	.	.	.	.	.	.	.	.	.	.
90.5	BRANCHIOSYLLIS EXILIS	0.6	0.4	0.2	0.2	0.4	0.2	0.2	0.8	0.4	0.4	0.4	0.4	0.4
90.5	DIPLODONTA PUNCTATA	1.4	0.7	0.6	0.4	.	.	.	.	.	.	.	.	.
90.5	GAMMARUS SP.	.	.	2.0	1.8	.	.	.	.	.	.	.	.	.
90.5	LOIMIA MEDUSA	0.4	0.4	0.8	0.6	.	.	.	0.8	0.4	0.4	0.4	0.4	0.4
90.5	PAGURUS CAROLINENSIS	.	.	0.6	0.4	1.4	1.2	0.2	0.6	0.4	0.4	0.4	0.4	0.4
90.5	PRIONOSPIO CRISTATA	0.6	0.4	0.6	0.4	0.2	0.2	0.2	0.6	0.4	0.4	0.4	0.4	0.4
98.5	AXIOTHELLA MUCOSA	.	.	0.2	0.2	0.8	0.8	0.8	0.8	0.5	0.5	0.5	0.5	0.5
98.5	CERADOCUS SP.	0.2	0.2	.	.	1.4	1.4	0.2	0.2	0.2	0.2	0.2	0.2	0.2
98.5	HETEROMYSIS FORMOSA	.	.	1.8	1.3	.	.	.	.	.	.	.	.	.
98.5	HEXAPANOPEUS SP. *	1.8	1.8	.	.	.	.	.	.	.	.	.	.	.
98.5	MAERA CAROLINIANA	.	.	0.8	0.8	0.4	0.4	0.4	0.6	0.4	0.4	0.4	0.4	0.4
98.5	PELIA MUTICA	0.4	0.2	1.4	0.9	.	.	.	.	.	.	.	.	.
98.5	PISANIA TINCTA	0.4	0.2	1.2	0.7	0.2	0.2	0.2	.	.	.	.	.	.
98.5	PSEUDOVERMILIA OCCIDENTALIS	.	.	1.0	0.5	0.8	0.6	0.6	0.6	0.4	0.4	0.4	0.4	0.4
98.5	THOR SP.	0.4	0.2	0.4	0.2	0.4	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
98.5	TURBELLARIA	0.6	0.4	0.6	0.4	0.4	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
106.0	AUTOLYTUS SP.	0.6	0.2	0.4	0.4	0.4	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
106.0	CRASSISPIRA ALBOMACULATA	0.2	0.2	0.2	0.2	1.2	1.0	.	.	.	.	.	.	.
106.0	DASYBRANCHUS SP.	1.2	0.5	0.2	0.2	.	.	.	0.2	0.2	0.2	0.2	0.2	0.2
106.0	SICYONIA LAEVIGATA	0.8	0.6	.	.	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4
106.0	SPIOPHANES BOMBYX	0.4	0.2	0.2	0.2	0.2	0.2	0.2	0.8	0.4	0.4	0.4	0.4	0.4
113.5	BRANCHIOSYLLIS OCULATA	0.2	0.2	.	.	.	.	.	1.2	0.5	0.5	0.5	0.5	0.5
113.5	INACHOIDES FORCEPS	0.4	0.2	0.4	0.2	0.4	0.4	0.4	0.2	0.2	0.2	0.2	0.2	0.2

Appendix 3. (Continued)

OVERALL RANK	SPECIES	STATION		IS02										
		WINTER		SPRING		SUMMER			FALL					
		MEAN	ST ERR	MEAN	ST ERR	MEAN	ST	ERR	MEAN	ST	ERR	MEAN	ST	ERR
113.5	MARGINELLA LAVALLEEANA	0.4	0.2	0.2	0.2	0.8	0.5		0.8			0.4		
113.5	PHYLLODOCE FRAGILIS	0.6	0.4						0.2			0.2		
113.5	PISTA PALMATA	0.8	0.8	0.4	0.4									
113.5	POLYCHERIA SP.	1.4	1.2											
113.5	POLYCIRRUS EXIMIUS	0.8	0.6			0.4	0.2		0.2			0.2		
113.5	POLYDORA CAECA	0.4	0.2	0.8	0.6	0.2	0.2							
113.5	SABELLARIA VULGARIS BEAUFORTENSIS	0.4	0.4	0.8	0.4				0.2			0.2		
113.5	SYLLIS REGULATA CAROLINAE								1.4			0.4		
127.5	AMPELISCA VADORUM	1.0	0.4						0.2			0.2		
127.5	ARABELLA MUTANS			0.6	0.4	0.4	0.4		0.2			0.2		
127.5	CHAMA MACEROPHYLLA	1.0	0.6			0.2	0.2							
127.5	DIODORA CAYENENSIS			0.4	0.4	0.6	0.2		0.2			0.2		
127.5	DODECACEA CORALLII	0.8	0.6	0.4	0.2									
127.5	EUNICE FILAMENTOSA	0.4	0.4	0.4	0.4	0.4	0.4							
127.5	HYDROIDES SP. B	0.2	0.2	0.6	0.6	0.2	0.2		0.2			0.2		
127.5	LITHOPHAGA BISULCATA	0.4	0.2	0.6	0.4	0.2	0.2							
127.5	MARPHYSIA SP. A	1.0	0.8			0.2	0.2							
127.5	OWENIA FUSIFORMIS	0.2	0.2	0.6	0.4	0.4	0.2							
127.5	PAGURISTES TORTUGAE	0.6	0.2	0.4	0.2				0.2			0.2		
127.5	PAGURUS HENDERSONI			0.2	0.2	0.2	0.2		0.8			0.8		
127.5	PILUMNUS PANNOSUS					0.6	0.4		0.6			0.4		
127.5	PSEUDOVERMILIA SP. A	0.6	0.2	0.2	0.2	0.4	0.4							
127.5	SPIO PETTIBONEAE			0.2	0.2				1.0			1.0		
127.5	SPIONIDAE	0.2	0.2	1.0	0.8									
127.5	TELLINA AMERICANA	0.2	0.2			0.2	0.2		0.8			0.8		
127.5	TRYPANOSYLLIS ZEBRA	0.6	0.4						0.6			0.4		
143.0	AMPHARETE ACUTIFRONS	0.2	0.2	0.8	0.6									
143.0	HOLOTHUROIDEA	0.2	0.2	0.8	0.8									
143.0	LUCIFER FAXONI			0.8	0.8	0.2	0.2							
143.0	LUMBRINERIS COCCINEA			0.4	0.4	0.4	0.4		0.2			0.2		
143.0	MEDIOMASTUS CALIFORNIENSIS	0.6	0.6	0.4	0.2									
143.0	MELPHIDIPPIDAE A			0.2	0.2	0.8	0.8							
143.0	NYMPHONIDAE A	0.8	0.8						0.2			0.2		
143.0	PAGURISTES SP.	0.2	0.2	0.8	0.6									
143.0	RILDARDANUS LAMINOSA	0.6	0.2	0.2	0.2	0.2	0.2							
143.0	SUTUROGLYPHA IONTHA	0.2	0.2	0.8	0.8									
143.0	THYONE PSEUDOFUSUS	0.2	0.2	0.4	0.4	0.4	0.2							
143.0	VERMICULARIA KNORRII	0.2	0.2			0.8	0.4							
143.0	XANTHIDAE			0.6	0.4	0.4	0.4							
158.0	ANOPLODACTYLUS INSIGNIS	0.8	0.5											
158.0	ARBACIA PUNCTULATA	0.6	0.2			0.2	0.2							
158.0	CHIONE GRUS	0.2	0.2			0.4	0.2		0.2			0.2		
158.0	CHRYSOPETALIDAE B	0.2	0.2	0.2	0.2				0.4			0.4		
158.0	CUMINGIA TELLINOIDES			0.2	0.2	0.6	0.2							
158.0	ENGINA TURBINELLA	0.2	0.2			0.6	0.6							
158.0	EUNICE VITTATA			0.2	0.2	0.4	0.4		0.2			0.2		
158.0	GLYCERA PAPILLOSA					0.6	0.4		0.2			0.2		
158.0	GNATHIA SP. A	0.4	0.2	0.2	0.2	0.2	0.2							
158.0	MALDANIDAE					0.6	0.4		0.2			0.2		
158.0	PILUMNUS SAYI			0.6	0.4				0.2			0.2		
158.0	PROCESSA SP.			0.6	0.6				0.2			0.2		
158.0	SABELLARIA FLORIDENSIS	0.4	0.2	0.4	0.4									
158.0	SABELLARIA VULGARIS	0.6	0.4						0.2			0.2		
158.0	SABELLARIA VULGARIS VULGARIS	0.6	0.4	0.2	0.2									
158.0	SEBA SP.			0.8	0.6									

Appendix 3. (Continued)

OVERALL RANK	SPECIES	STATION		IS02									
		WINTER		SPRING		SUMMER		FALL					
		MEAN	ST ERR	MEAN	ST ERR	MEAN	ST ERR	MEAN	ST ERR	MEAN	ST ERR	MEAN	ST ERR
158.0	XANTHIDAE D	.	.	.	.	0.8	0.8	.	.	.	.	.	.
183.5	ANCHIALINA TYPICA	.	.	.	.	0.6	0.4	.	.	.	.	.	.
183.5	ANOPLODACTYLUS LENTUS	0.2	0.2	.	.	0.4	0.4	.	.	.	.	.	.
183.5	BRANIA CLAVATA	.	.	0.4	0.4	0.2	0.2	.	.	.	.	.	.
183.5	CAPRELLIDAE A	0.4	0.4	.	.	0.2	0.2	.	.	.	.	.	.
183.5	CARINOBATEA CARINATA	0.6	0.4	.	.	.	.	.	.	.	.	.	.
183.5	CERAPUS TUBULARIS	.	.	0.6	0.4	.	.	.	.	.	.	.	.
183.5	CERITHIOPSIS EMERSONI	0.6	0.6	.	.	0.6	0.6	.	.	.	.	.	.
183.5	CERITHIOPSIS GREENI	.	.	.	.	.	.	.	.	.	.	.	.
183.5	CERITHIUM ATRATUM	.	.	0.6	0.6	.	.	.	.	.	.	.	.
183.5	COLOMASTIX SP.	.	.	0.2	0.2	0.4	0.4	.	.	.	.	.	.
183.5	COROPHIDAE	0.6	0.6	.	.	.	.	.	.	.	.	.	.
183.5	ERATO MAUGERIAE	0.2	0.2	.	.	0.4	0.4	.	.	.	.	.	.
183.5	EUNICE ANTENNATA	0.2	0.2	.	.	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
183.5	FAVARTIA CELLULOSA	.	.	0.2	0.2	0.4	0.4	.	.	.	.	.	.
183.5	GITANOPSIS SP.	.	.	0.6	0.2	.	.	.	.	.	.	.	.
183.5	GLYCERA TESSELATA	.	.	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
183.5	HIPPOLYTIDAE	0.2	0.2	.	.	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
183.5	HYBOSCOLEX LONGISETA	.	.	0.4	0.2	0.2	0.2	.	.	.	.	.	.
183.5	LAEVICARDIUM PICTUM	.	.	0.4	0.4	0.2	0.2	.	.	.	.	.	.
183.5	LEMBOS SP.	.	.	0.4	0.2	.	.	0.2	0.2	0.2	0.2	0.2	0.2
183.5	LUMBRINERIS LATREILLI	.	.	.	.	.	.	0.6	0.4	.	.	.	.
183.5	MICROPANOPE SP.	0.2	0.2	0.2	0.2	0.2	0.2	.	.	.	.	.	.
183.5	NEMERTINEA E	.	.	0.6	0.4	.	.	.	.	.	.	.	.
183.5	OPHIOSTIGMA SP. A	0.6	0.4	.	.	.	.	.	.	.	.	.	.
183.5	PAGURIDAE	0.6	0.6	.	.	.	.	.	.	.	.	.	.
183.5	PAGURIDEA	.	.	0.4	0.4	0.2	0.2	.	.	.	.	.	.
183.5	PHYLLOCHAETOPTERUS SOCIALIS	.	.	0.2	0.2	0.4	0.2	.	.	.	.	.	.
183.5	PHYLLODOCE LONGIPES	.	.	.	.	0.4	0.2	0.2	0.2	0.2	0.2	0.2	0.2
183.5	PILUMNUS SP. *	.	.	0.6	0.6	.	.	.	.	.	.	.	.
183.5	POLYDORA WEBSTERI	.	.	.	.	0.6	0.6	.	.	.	.	.	.
183.5	RUDILEMBOIDES NAGLEI	.	.	0.6	0.4	.	.	.	.	.	.	.	.
183.5	SIPUNCULIDA C	0.2	0.2	0.4	0.4	.	.	.	.	.	.	.	.
183.5	SYLLIDAE	.	.	.	.	0.4	0.4	0.2	0.2	0.2	0.2	0.2	0.2
183.5	TIMARETE SP.	0.6	0.4	.	.	.	.	.	.	.	.	.	.
220.5	ACCALATHURA CHENULATA	.	.	.	.	.	.	.	.	.	.	.	.
220.5	AMPHILOCHUS SP.	.	.	.	.	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4
220.5	AMPHINOMIDAE B	.	.	.	.	.	.	.	.	.	.	.	.
220.5	ANCISTROSYLLIS RIGIDA	.	.	.	.	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
220.5	BOWMANIELLA PORTORICENSIS	.	.	0.2	0.2	.	.	0.2	0.2	0.2	0.2	0.2	0.2
220.5	CAPITELLA CAPITATA	.	.	0.4	0.2	.	.	.	.	.	.	.	.
220.5	CARIDEA	.	.	0.4	0.2	.	.	.	.	.	.	.	.
220.5	CERATONEREIS SP.	.	.	0.2	0.2	0.2	0.2	.	.	.	.	.	.
220.5	CHAETOPLEURA APICULATA	.	.	.	.	.	.	0.4	0.4	0.4	0.4	0.4	0.4
220.5	CHONE AMERICANA	.	.	0.4	0.2	.	.	.	.	.	.	.	.
220.5	CRASSINELLA LUNULATA	.	.	0.2	0.2	0.2	0.2	.	.	.	.	.	.
220.5	DENDROCHIROTIDA	.	.	0.4	0.4	.	.	.	.	.	.	.	.
220.5	DIODURA CAYENENSIS *	0.4	0.4	.	.	.	.	.	.	.	.	.	.
220.5	EBALIA SP.	.	.	0.2	0.2	0.2	0.2	.	.	.	.	.	.
220.5	EUNICE SP.	.	.	0.4	0.4	.	.	.	.	.	.	.	.
220.5	EURYDICE LITTORALIS	0.2	0.2	.	.	.	.	0.2	0.2	0.2	0.2	0.2	0.2
220.5	HARMOTHOE SP. A	0.2	0.2	0.2	0.2	.	.	.	.	.	.	.	.
220.5	HIPPOLYTE NICHOLSONI *	.	.	.	.	.	.	0.4	0.4	0.4	0.4	0.4	0.4
220.5	HYPOCONCHA SABULOSA	.	.	0.2	0.2	0.2	0.2	.	.	.	.	.	.
220.5	MELITIDAE	0.2	0.2	0.2	0.2	.	.	.	.	.	.	.	.

Appendix 3. (Continued)

OVERALL RANK	SPECIES	STATION			IS02											
		WINTER			SPRING			SUMMER			FALL					
		MEAN	ST	ERR	MEAN	ST	ERR	MEAN	ST	ERR	MEAN	ST	ERR	MEAN	ST	ERR
220.5	METOPORHAPHIS CALCARATA	.	.	.	0.4	0.4	.	0.4	0.2	.	.	.	.	.	.	.
220.5	MICROPHRYS SP.	.	.	.	0.2	0.2	.	.	.	.	0.2	0.2	.	0.2	0.2	.
220.5	NASSARIUS ALBUS	.	.	.	.	.	.	0.4	0.4	.	.	.	.	.	.	.
220.5	NEMERTINEA C	.	.	.	0.4	0.4	.	.	.	.	.	.	.	.	.	.
220.5	NYMPHONIDAE	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
220.5	OLIGOCHAETA	0.2	0.2	.	.	.	.	.	.	.	0.2	0.2	.	0.2	0.2	.
220.5	PARAPHOXUS SPINOSUS	0.4	0.2	.	.	.	.	.	.	.	.	.	.	.	.	.
220.5	PERICLIMENES SP.	.	.	.	0.2	0.2	.	0.4	0.2	.	.	.	.	.	.	.
220.5	PHYLLODOCE SP.	.	.	.	0.2	0.2	.	0.2	0.2	.	.	.	.	.	.	.
220.5	PINNIXA SP.	.	.	.	0.4	0.2	.	.	.	.	.	.	.	.	.	.
220.5	PORTUNUS SP.	.	.	.	0.4	0.4	.	.	.	.	.	.	.	.	.	.
220.5	RUPELLARIA TYPICA	.	.	.	0.4	0.2	.	.	.	.	.	.	.	.	.	.
220.5	SABELLIDAE	.	.	.	.	.	.	.	.	.	0.4	0.2	.	0.4	0.2	.
220.5	SEILA ADAMSI	.	.	.	0.2	0.2	.	0.2	0.2	.	.	.	.	.	.	.
220.5	SICYONIA SP.	0.2	0.2	.	.	.	.	.	.	.	0.2	0.2	.	0.2	0.2	.
220.5	STENOCIONOPS FURCATA COELATA *	0.4	0.4	.	.	.	.	.	.	.	.	.	.	.	.	.
220.5	SYLLIS FERRUGINA	.	.	.	.	.	.	0.2	0.2	.	0.2	0.2	.	0.2	0.2	.
220.5	TIRON TROPAKIS	.	.	.	.	.	.	.	.	.	0.4	0.2	.	0.4	0.2	.
220.5	TRICHOBRANCHUS GLACIALIS	.	.	.	.	.	.	.	.	.	0.4	0.2	.	0.4	0.2	.
220.5	XANTHIDAE B	.	.	.	0.2	0.2	.	.	.	.	0.2	0.2	.	0.2	0.2	.
295.0	ACANTHOHAUSTORIUS BOUSFIELDI	.	.	.	.	.	.	.	.	.	0.2	0.2	.	0.2	0.2	.
295.0	AEOLIDIAE A	.	.	.	.	.	.	.	.	.	0.2	0.2	.	0.2	0.2	.
295.0	AMPHITHE SP.	.	.	.	0.2	0.2	.	.	.	.	.	.	.	.	.	.
295.0	ANACHIS LAFRESNAYI	.	.	.	.	.	.	0.2	0.2	.	.	.	.	.	.	.
295.0	ARABELLA TRICOLOR	.	.	.	0.2	0.2	.	.	.	.	.	.	.	.	.	.
295.0	AUSTRAEOLIS CATINA	.	.	.	.	.	.	0.2	0.2	.	.	.	.	.	.	.
295.0	BARBATIA CANDIDA	.	.	.	.	.	.	.	.	.	0.2	0.2	.	0.2	0.2	.
295.0	BATRACHONOTUS FRAGOSUS	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.	.	.
295.0	CANTHARUS MULTANGULUS	.	.	.	0.2	0.2	.	.	.	.	.	.	.	.	.	.
295.0	CAPITELLIDAE D	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.	.	.
295.0	CAULLERIELLA KILLARIENSIS	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.	.	.
295.0	CERATONEREIS VERSIPEDATA	.	.	.	.	.	.	.	.	.	0.2	0.2	.	0.2	0.2	.
295.0	CHEVALIA SP.	.	.	.	0.2	0.2	.	.	.	.	.	.	.	.	.	.
295.0	CHRYSOPETALIDAE	.	.	.	.	.	.	.	.	.	0.2	0.2	.	0.2	0.2	.
295.0	CLYPEASTER SP.	.	.	.	.	.	.	.	.	.	0.2	0.2	.	0.2	0.2	.
295.0	CRASSOSTREA VIRGINICA	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.	.	.
295.0	DONDICE OCCIDENTALIS	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.	.	.
295.0	DORVILLEA SOCIABILIS	.	.	.	.	.	.	0.2	0.2	.	.	.	.	.	.	.
295.0	DROMIDIA ANTILLENIS	.	.	.	.	.	.	0.2	0.2	.	.	.	.	.	.	.
295.0	ECHINASTER SP.	.	.	.	0.2	0.2	.	.	.	.	.	.	.	.	.	.
295.0	EPITONIUM HUMPHREYSI	.	.	.	.	.	.	0.2	0.2	.	.	.	.	.	.	.
295.0	ERICHSONELLA FILIFORMIS	.	.	.	0.2	0.2	.	.	.	.	.	.	.	.	.	.
295.0	EUNICE TENUIS	.	.	.	.	.	.	.	.	.	0.2	0.2	.	0.2	0.2	.
295.0	GLYCERIDAE	.	.	.	.	.	.	0.2	0.2	.	.	.	.	.	.	.
295.0	GONODACTYLUS BREDINI	.	.	.	.	.	.	0.2	0.2	.	.	.	.	.	.	.
295.0	HEMIPODUS ROSEUS	.	.	.	.	.	.	.	.	.	0.2	0.2	.	0.2	0.2	.
295.0	HESIONIDAE	.	.	.	.	.	.	0.2	0.2	.	.	.	.	.	.	.
295.0	HETEROCRYPTA GRANULATA	.	.	.	.	.	.	0.2	0.2	.	.	.	.	.	.	.
295.0	HIATELLA ARCTICA	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.	.	.
295.0	HOLOTHURIA DAKARENSIS	.	.	.	.	.	.	0.2	0.2	.	.	.	.	.	.	.
295.0	HYDROIDES SP. A	.	.	.	.	.	.	.	.	.	0.2	0.2	.	0.2	0.2	.
295.0	HYPsicOMUS PHAEOTAENIA	.	.	.	.	.	.	.	.	.	0.2	0.2	.	0.2	0.2	.
295.0	ISCHNOCHITONIDAE A	.	.	.	.	.	.	.	.	.	0.2	0.2	.	0.2	0.2	.
295.0	LAMELLARIA PERSPICUA	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.	.	.
295.0	LEPTOCHELA SP.	.	.	.	.	.	.	0.2	0.2	.	.	.	.	.	.	.

Appendix 3. (Continued)

OVERALL RANK	SPECIES	STATION IS02											
		WINTER			SPRING			SUMMER			FALL		
		MEAN	ST	ERR	MEAN	ST	ERR	MEAN	ST	ERR	MEAN	ST	ERR
295.0	LINGA AMIANUS	0.2		0.2	.		.	.		.	.		.
295.0	LUIDIA ALTERNATA	.		.	0.2		0.2	.		.	.		.
295.0	LUMBRINERIDAE	0.2		0.2	.		.	.		.	.		.
295.0	LUMBRINERIS IMPATIENS	0.2		0.2	.		.	.		.	.		.
295.0	LYONSIA SP.	.		.	.		.	0.2		0.2	.		.
295.0	LYSMATA INTERMEDIA	.		.	.		.	0.2		0.2	.		.
295.0	LYSMATA SP.	.		.	.		.	0.2		0.2	.		.
295.0	MACROCOELOMA TRISPINOSUM	.		.	0.2		0.2	.		.	0.2		0.2
295.0	MAJIDAE A	.		.	.		.	.		.	.		.
295.0	MESANTHURA DECORATA	0.2		0.2	0.2		0.2	.		.	.		.
295.0	MITHRAX FORCEPS	.		.	.		.	0.2		0.2	.		.
295.0	MITHRAX PLEURACANTHUS	.		.	.		.	.		.	.		.
295.0	MITRELLA RAVENELI	0.2		0.2	.		.	.		.	.		.
295.0	MODIOLUS MODIOLUS SQUAMOSUS	0.2		0.2	.		.	.		.	.		.
295.0	MONOCULODES SP.	.		.	0.2		0.2	.		.	.		.
295.0	MUREX FLORIFER DILECTUS	.		.	.		.	0.2		0.2	0.2		0.2
295.0	MUSCULUS LATERALIS	.		.	.		.	.		.	.		.
295.0	NAINERIS SP.	0.2		0.2	.		.	0.2		0.2	.		.
295.0	NEMATONEREIS SP.	.		.	0.2		0.2	.		.	.		.
295.0	NEREIS ACUMINATA	0.2		0.2	.		.	.		.	.		.
295.0	NEREIS LAMELLOSA	0.2		0.2	.		.	.		.	.		.
295.0	NEREIS SP.	.		.	.		.	0.2		0.2	.		.
295.0	NEREIS SUCCINEA	.		.	0.2		0.2	.		.	.		.
295.0	NOTOBRYON WARDI	.		.	0.2		0.2	.		.	.		.
295.0	NUCULA CRENULATA	0.2		0.2	.		.	.		.	.		.
295.0	ODONTOSYLLIS ENOPLA	.		.	.		.	.		.	0.2		0.2
295.0	OPHIOCNIDA SCABRIUSCULA	.		.	0.2		0.2	.		.	0.2		0.2
295.0	OPHIUROIDEA	.		.	.		.	0.2		0.2	.		.
295.0	ORBINIIDAE	.		.	.		.	0.2		0.2	.		.
295.0	PALAEMONIDAE	.		.	.		.	.		.	0.2		0.2
295.0	PALEANOTUS SP. B	.		.	.		.	0.2		0.2	.		.
295.0	PANANTHURA FORMOSA	.		.	0.2		0.2	.		.	.		.
295.0	PELECYPODA C	.		.	.		.	0.2		0.2	.		.
295.0	PERICLIMENES IRIDESCENS	.		.	.		.	.		.	0.2		0.2
295.0	PERICLIMENES SP. *	.		.	0.2		0.2	.		.	.		.
295.0	PETALOPROCTUS SOCIALIS	.		.	.		.	.		.	0.2		0.2
295.0	PETROLISTHES GALATHINUS	.		.	.		.	0.2		0.2	.		.
295.0	PHYLLODOCE SP. E	.		.	0.2		0.2	.		.	.		.
295.0	PINNOTHERES OSTREUM	0.2		0.2	.		.	.		.	.		.
295.0	PINNOTHERES SP.	.		.	0.2		0.2	.		.	.		.
295.0	PINNOTHERIDAE	.		.	0.2		0.2	.		.	.		.
295.0	PITHO LHERMINIERI	.		.	.		.	.		.	0.2		0.2
295.0	PLEUROPLOCA GIGANTEA	.		.	.		.	0.2		0.2	.		.
295.0	PODOCHELA RIISEI	0.2		0.2	.		.	.		.	.		.
295.0	PODOCHELA SP.	.		.	.		.	.		.	0.2		0.2
295.0	PODODESMUS SP.	.		.	.		.	0.2		0.2	.		.
295.0	POLYCHAETA G	.		.	.		.	.		.	0.2		0.2
295.0	POLYCIRRUS SP. C	.		.	.		.	.		.	0.2		0.2
295.0	POLYDORA COLONIA	.		.	.		.	0.2		0.2	.		.
295.0	POLYDORA SOCIALIS	.		.	0.2		0.2	.		.	.		.
295.0	POLYDORA SP.	0.2		0.2	.		.	.		.	.		.
295.0	POLYNOIDAE	.		.	0.2		0.2	.		.	.		.
295.0	POLYPLACOPHORA C	.		.	.		.	0.2		0.2	.		.
295.0	PORTUNIDAE	.		.	0.2		0.2	.		.	.		.

Appendix 3. (Continued)

OVERALL RANK	SPECIES	STATION		IS02									
		WINTER		SPRING		SUMMER		FALL					
		MEAN	ST ERR	MEAN	ST ERR	MEAN	ST ERR	MEAN	ST ERR	MEAN	ST ERR	MEAN	ST ERR
295.0	POTAMILLA RENIFORMIS	.	.	.	.	.	.	0.2	0.2	.	.	.	.
295.0	PRIONOSPIO SP.	0.2	0.2	.	.	.	.	.	.	.	.	.	.
295.0	PROCERAEA SP.	.	.	.	.	0.2	0.2	.	.	.	.	.	.
295.0	PYRGOSPIRA TAMPAENSIS	.	.	.	.	0.2	0.2	.	.	.	.	.	.
295.0	SCHISTOMERINGOS RUDOLPHI	.	.	0.2	0.2	.	.	.	.	.	.	.	.
295.0	SICYONIA BREVIROSTRIS	.	.	.	.	.	.	0.2	0.2	.	.	.	.
295.0	SICYONIA BREVIROSTRIS *	.	.	.	.	.	.	0.2	0.2	.	.	.	.
295.0	SICYONIIDAE	.	.	.	.	.	.	0.2	0.2	.	.	.	.
295.0	SPELOEOPHORUS PONTIFER	.	.	.	.	0.2	0.2	0.2	0.2	.	.	.	.
295.0	SPIRORBIS SP.	.	.	.	.	0.2	0.2	.	.	.	.	.	.
295.0	STENORHYNCHUS SETICORNIS	.	.	0.2	0.2	.	.	.	.	.	.	.	.
295.0	SYLLIDAE H	0.2	0.2	.	.	.	.	.	.	.	.	.	.
295.0	TELLINA SYBARITICA	.	.	.	.	0.2	0.2	.	.	.	.	.	.
295.0	TEREBELLIDAE	0.2	0.2	.	.	.	.	.	.	.	.	.	.
295.0	THOR MANNINGI *	.	.	.	.	.	.	0.2	0.2	.	.	.	.
295.0	TRIPHORA NIGROCINCTA	.	.	0.2	0.2	.	.	.	.	.	.	.	.
295.0	TURBONILLA SP. *	.	.	.	.	0.2	0.2	.	.	.	.	.	.
295.0	VERMILIOPSIS ANNULATA	.	.	.	.	0.2	0.2	.	.	.	.	.	.
295.0	VEXILLUM WANDUENSIS	.	.	0.2	0.2	.	.	.	.	.	.	.	.

Appendix 3. (Continued)

OVERALL RANK	SPECIES	STATION		MS02							
		WINTER		SPRING		SUMMER		FALL			
		MEAN	ST ERR	MEAN	ST ERR	MEAN	ST ERR	MEAN	ST ERR	MEAN	ST ERR
1.0	FILOGRANA IMPLEXA	53.2	33.2	1084.4	747.0	0.2	0.2	531.6	355.4		
2.0	SYLLIS SPONGICOLA	35.4	19.6	17.2	8.0	7.4	5.0	44.6	42.6		
3.0	MALACOCEROS GLUTAEUS	19.0	8.1	11.6	10.9	2.4	1.2	36.8	19.3		
4.0	CHONE AMERICANA	7.8	2.9	41.2	20.9	1.0	0.8	3.4	1.6		
5.0	OPHIOTHRIX ANGULATA	26.4	15.7	11.8	6.8	1.0	0.8	1.4	0.7		
6.0	PHOTIS SP.	31.8	13.6	4.2	3.4	.	.	4.4	3.7		
7.0	ACANTHOHAUSTORIUS MILLSI	4.2	2.7	20.4	13.5	2.0	1.2	12.4	10.7		
8.0	MEGALOBRACHIUM SORIATUM	5.6	4.2	5.8	5.1	7.0	4.8	20.0	10.8		
9.0	LUMBRINERIS INFLATA	11.2	6.3	10.0	4.1	3.4	1.7	8.4	3.3		
10.0	EXOGONE DISPAR	18.0	9.6	8.4	3.7	0.8	0.4	3.8	1.6		
11.0	OWENIA FUSIFORMIS	5.2	2.2	23.0	11.4	0.8	0.4	1.8	0.6		
12.0	CHEVALIA SP.	21.4	20.6	4.8	4.8	0.2	0.2	3.8	3.3		
13.0	LAEVICARDIUM PICTUM	1.8	1.1	21.6	2.9	2.0	0.8	4.2	2.0		
14.0	LUCONACIA INCERTA	17.8	7.7	7.0	5.3	0.8	0.6	3.0	1.5		
15.0	BRANCHIOSYLLIS EXILIS	9.4	4.9	8.4	4.1	3.2	2.0	2.8	1.2		
16.0	LEMNOS SMITHI	4.4	2.0	7.4	3.2	9.4	5.9	1.0	0.6		
17.0	MICRODEUTOPUS MYERSI	12.0	7.3	.	.	0.2	0.2	9.6	4.0		
18.0	AMPITHOE SP. A	6.0	2.4	12.4	6.6	0.8	0.6	1.6	1.1		
19.0	PAGURUS CAROLINENSIS	4.4	2.5	5.8	2.3	1.2	0.8	7.2	3.1		
20.0	GAMMAROPSIS SP.	11.0	5.0	6.4	3.4	.	.	0.8	0.8		
21.0	THOR SP.	4.2	3.2	7.4	3.7	2.8	1.9	3.6	2.3		
23.0	AMPHIODIA PULCHELLA	2.8	2.5	3.2	1.9	3.2	1.8	7.8	2.3		
23.0	OPHIACTIS SAVIGNYI	3.6	1.5	10.6	3.5	1.2	0.8	1.6	1.0		
23.0	PAGURUS SP.	9.6	5.7	1.6	0.9	0.4	0.4	5.4	2.5		
25.0	LEUCOTHOE SPINICARPA	3.2	1.8	3.8	2.4	7.2	4.5	2.4	1.2		
26.0	XANTHIDAE A	.	.	7.2	4.7	1.8	1.3	6.6	4.8		
27.0	MITRELLA LUNATA	10.8	4.6	3.2	1.6	.	.	1.4	0.5		
28.0	SUTUROGLYPHA IONTHA	7.2	3.0	6.4	2.6	0.2	0.2	1.0	0.6		
29.0	NEMERTINEA	7.2	3.1	3.8	1.3	0.2	0.2	2.6	1.3		
30.0	SYLLIS HYALINA	4.8	2.7	3.6	2.1	2.0	1.5	2.6	1.4		
31.0	PODARKE OBSCURA	2.6	1.2	4.2	2.0	2.2	1.7	3.8	1.4		
32.0	POLYCIRRUS CAROLINENSIS	2.4	1.3	7.4	4.0	1.0	0.8	1.8	1.1		
33.0	PILUMNUS SP.	2.6	1.4	1.0	0.4	2.4	1.9	6.4	2.6		
34.0	AUTOLYTUS SP.	2.6	1.5	3.8	1.6	1.4	0.7	4.0	1.9		
35.0	CAPRELLA PENANTIS	3.0	1.5	7.6	6.4	.	.	1.0	1.0		
36.0	CAPRELLA EQUILIBRA	2.0	0.7	9.4	8.4	.	.	.	.		
37.5	ASPIDOSIPHON GOSNOLDI	3.2	1.4	5.4	2.9	1.0	0.5	1.6	0.6		
37.5	WEBSTERINEREIS SP. B	3.2	1.5	3.4	1.3	2.0	0.9	2.6	1.0		
39.5	PSEUDOMEDAEUS AGASSIZII	0.4	0.4	6.8	4.9	2.6	1.7	1.0	0.6		
39.5	WEBSTERINEREIS TRIDENTATA	3.2	1.1	3.2	1.6	1.2	0.6	3.2	1.5		
41.5	HARMOTHOE SP. A	5.2	2.4	4.4	1.9	0.4	0.2	0.6	0.2		
41.5	PRIONOSPION CRISTATA	0.8	0.6	4.4	2.6	1.0	0.3	4.4	1.8		
43.5	SPHAERODORIDIUM CLAPAREDI	10.4	9.9	.	.	.	.	.	.		
43.5	SYNALPHEUS TOWNSENDI	1.8	1.1	3.6	3.3	2.8	1.7	2.2	1.3		
45.0	EUNICE VITTATA	2.2	1.0	2.2	0.7	1.6	1.0	4.2	1.6		
46.0	ELASMOPUS SP. A	2.6	1.7	3.8	2.8	0.8	0.4	2.8	2.3		
47.0	POLYDORA CAECA	0.4	0.2	0.6	0.4	.	.	8.6	3.6		
48.0	EULALIA SANGUINEA	1.6	0.5	5.8	3.4	1.0	0.8	1.0	0.8		
49.0	AMPHARETE AMERICANA	2.2	1.1	0.6	0.4	0.8	0.5	5.4	1.5		
50.0	HIATELLA ARCTICA	1.2	1.2	6.6	4.9	0.6	0.4	0.2	0.2		
51.5	MEGALOMMA BILOCULATUM	1.8	0.8	5.8	3.1	0.2	0.2	0.6	0.4		
51.5	PROCERAEA SP.	.	.	5.8	3.2	0.4	0.2	2.2	0.8		
53.0	CHIONE GRUS	1.4	0.6	3.0	1.8	1.2	0.7	2.4	0.8		
55.0	AXIOGNATHUS SQUAMATUS	2.4	1.2	1.8	1.1	0.6	0.6	3.0	1.3		
55.0	SEMELE NUCULOIDES	2.2	0.7	2.6	2.1	1.6	0.9	1.4	0.5		

Appendix 3. (Continued)

OVERALL RANK	SPECIES	STATION		MS02										
		WINTER		SPRING		SUMMER			FALL					
		MEAN	ST ERR	MEAN	ST ERR	MEAN	ST	ERR	MEAN	ST	ERR	MEAN	ST	ERR
55.0	SPIOPHANES BOMBYX	3.0	2.0	2.4	1.3	0.6		0.6	1.8		0.8			
57.0	TURBELLARIA	2.4	0.9	0.4	0.4	1.2		0.7	3.6		1.4			
58.0	PODOCERUS SP.	2.2	1.5	2.0	1.4	1.2		0.7	2.0		1.3			
59.0	ERICHTHONIUS BRASILIENSIS	2.0	1.4	1.6	0.5	0.2		0.2	3.2		1.6			
60.0	CERATONEREIS VERSIPEDATA	1.8	1.0	2.4	1.1	0.6		0.4	1.8		1.3			
61.0	POLYCIURUS EXIMIUS	1.8	0.9	2.0	1.4	0.8		0.6	1.8		0.9			
63.0	CERITHIUM ATRATUM	1.8	1.6	3.6	2.4	0.6		0.6	0.2		0.2			
63.0	MAJIDAE A	.	.	.	.	1.4		1.0	4.8		3.2			
63.0	TRAVISTIA PARVA	.	.	4.6	4.3	0.2		0.2	1.4		1.4			
67.0	DIASTYLIDAE A	2.2	0.8	0.2	0.2	.		.	3.6		0.6			
67.0	OPHIOSTIGMA SP. A	5.8	4.0	.	.	.		.	0.2		0.2			
67.0	PARACERCEIS CAUDATA	2.8	1.2	2.4	0.8	0.4		0.2	0.4		0.4			
67.0	PHERUSA EHLERSI	3.4	2.7	1.8	0.8	0.2		0.2	0.6		0.4			
67.0	TELLINA AMERICANA	1.6	1.0	1.2	0.7	1.4		0.5	1.8		1.3			
70.5	ELASMOPUS SP. B	0.4	0.4	5.4	5.4	.		.	.		.			
70.5	GOULDIA CERINA	0.4	0.4	0.2	0.2	4.4		2.1	0.8		0.4			
72.0	OPHIOSTIGMA ISACANTHUM	.	.	1.4	1.0	2.0		1.1	2.2		1.4			
73.0	NEMATODA	2.2	1.4	2.8	1.7	0.2		0.2	0.2		0.2			
75.0	BRACHYURA	0.6	0.4	2.0	1.4	1.4		0.5	1.2		0.7			
75.0	CHRYSOPETALIDAE A	1.8	1.6	1.2	0.4	0.4		0.2	1.8		0.7			
75.0	LUMBRINERIS COCCINEA	1.8	1.6	2.2	1.6	0.4		0.2	0.8		0.5			
77.5	LEPTOCHELA PAPULATA	.	.	0.4	0.2	2.2		1.6	2.4		1.2			
77.5	SYNALPHEUS SP.	.	.	1.6	1.0	2.4		1.7	1.0		0.4			
80.0	GONIADIDES CAROLINAE	0.6	0.6	0.8	0.4	2.6		2.3	0.8		0.4			
80.0	LEPTOCHELA SP.	.	.	4.8	4.1	.		.	.		.			
80.0	PAGURIDEA	.	.	0.6	0.4	1.6		1.0	2.6		1.0			
83.0	ODONTOSYLLIS ENOPLA	1.2	0.7	1.0	0.8	1.2		0.6	1.2		0.7			
83.0	PISTA PALMATA	1.4	1.0	1.8	0.9	1.0		0.4	0.4		0.2			
83.0	SIPUNCULIDA A	1.4	1.0	2.8	1.5	.		.	0.4		0.2			
85.5	MELITIDAE A	1.6	1.0	0.8	0.5	1.8		1.8	0.2		0.2			
85.5	XANTHIDAE	4.4	3.9	.	.	.		.	.		.			
87.0	CARPIAS BERMUDENSIS	1.2	0.7	1.4	0.9	0.8		0.5	0.8		0.6			
89.0	INACHOIDES FORCEPS	1.2	0.7	0.4	0.4	.		.	2.4		1.5			
89.0	LATREUTES PARVULUS	1.8	1.1	0.8	0.8	0.2		0.2	1.2		0.6			
89.0	PRIONOSPIO CIRRIFERA	1.6	1.0	0.8	0.6	.		.	1.6		0.7			
92.0	AXIOHELLA MUCOSA	0.2	0.2	2.0	1.4	0.6		0.4	1.0		0.4			
92.0	NOTOMASTUS SP.	1.4	0.7	1.6	0.9	.		.	0.8		0.4			
92.0	TRYPANOSYLLIS ZEBRA	2.0	1.5	0.4	0.2	.		.	1.4		0.7			
95.5	BODOTRIIDAE B	1.6	0.5	0.6	0.4	.		.	1.4		1.4			
95.5	CIRROPHORUS LYRIFORMIS	.	.	.	.	2.6		2.6	1.0		0.3			
95.5	MARPHYS A SP. A	1.8	0.9	0.6	0.6	0.6		0.2	0.6		0.6			
95.5	PHERUSA INFLATA	1.6	0.7	1.2	0.8	.		.	0.8		0.6			
101.5	CERATONEREIS SP.	0.2	0.2	0.6	0.4	1.2		1.0	1.4		1.0			
101.5	CERATONEREIS SP. A	1.2	0.6	0.8	0.4	.		.	1.4		0.7			
101.5	CERATONEREIS MIRABILIS	0.2	0.2	0.4	0.2	1.4		0.5	1.4		0.9			
101.5	COLOMASTIX SP.	.	.	.	.	0.6		0.6	2.8		2.5			
101.5	LOIMIA MEDUSA	2.2	1.5	1.0	0.3	.		.	0.2		0.2			
101.5	MELITA APPENDICULATA	1.4	0.9	0.4	0.2	0.6		0.6	1.0		1.0			
101.5	OPHIOSTIGMA SP.	.	.	3.4	2.1	.		.	.		.			
101.5	OPHIUROIDEA	2.2	1.0	.	.	1.2		1.2	.		.			
108.5	BOWMANIELLA PORTORICENSIS	.	.	1.4	0.7	1.6		0.6	0.2		0.2			
108.5	CIRRATULIDAE	3.0	3.0	0.2	0.2	.		.	.		.			
108.5	MAERA SP. A	1.4	0.6	1.4	1.4	0.4		0.4	.		.			
108.5	MUSCULUS LATERALIS	1.6	0.8	0.4	0.2	0.8		0.4	0.4		0.4			
108.5	NASSARIUS ALBUS	0.8	0.6	1.6	0.8	0.4		0.2	0.4		0.2			

Appendix 3. (Continued)

OVERALL RANK	SPECIES	STATION		MS02										
		WINTER		SPRING		SUMMER			FALL					
		MEAN	ST ERR	MEAN	ST ERR	MEAN	ST	ERR	MEAN	ST	ERR	MEAN	ST	ERR
108.5	XANTHIDAE B	0.6	0.6	0.8	0.6	.	.	.	1.8	.	.	1.8	1.0	.
115.5	ALPHEUS NORMANNI	0.8	0.8	0.6	0.2	0.6	0.4	.	1.0	0.5	.	.	.	.
115.5	ARCA IMBRICATA	2.0	1.3	1.0	0.4	.	.	.	.	.	.	.	.	.
115.5	CRASSINELLA LUNULATA	0.2	0.2	2.0	0.9	0.8	0.4	.	.	.	.	.	.	.
115.5	MEGALUROPOUS SP.	0.8	0.6	0.2	0.2	.	.	.	2.0	0.6	.	.	.	.
115.5	PISTA QUADRILOBATA	1.4	0.7	1.6	0.7	.	.	.	.	.	.	.	.	.
115.5	TANYSTYLIDAE A	1.2	0.4	1.6	0.7	.	.	.	0.2	0.2	.	.	.	.
115.5	TELLINA SYBARITICA	0.4	0.4	1.4	0.5	0.4	0.2	.	0.8	0.8	.	.	.	.
115.5	TRICHOBRANCHUS GLACIALIS	1.4	1.2	0.4	0.2	0.2	0.2	0.2	1.0	0.4	.	.	.	.
123.5	EUNICE FILAMENTOSA	1.0	0.8	0.4	0.4	0.4	0.2	0.2	1.0	1.0	.	.	.	.
123.5	GASTROCHAENA HIANIS	0.2	0.2	1.2	1.2	0.2	0.2	0.2	1.2	0.5	.	.	.	.
123.5	MARGINELLA LAVALLEEANA	1.0	0.6	1.6	0.7	0.2	0.2	0.2	.	.	.	.	.	.
123.5	NASSARINA MINOR	1.2	1.0	1.2	0.6	0.4	0.2	0.2	.	.	.	.	.	.
123.5	NICOMACHE TRISPINATA	1.8	1.6	.	.	0.4	0.4	0.4	0.6	0.4	.	.	.	.
123.5	PAGURUS HENDERSONI	0.2	0.2	0.4	0.2	0.6	0.2	0.2	1.6	0.9	.	.	.	.
123.5	PELIA MUTICA	1.0	0.8	0.8	0.6	0.4	0.4	0.4	0.6	0.6	.	.	.	.
123.5	SICYONIA SP.	0.6	0.4	0.8	0.5	.	.	.	1.4	0.9	.	.	.	.
129.0	AMPELISCA VADORUM	2.2	0.8	0.2	0.2	.	.	.	0.2	0.2	.	.	.	.
129.0	DIPLODONTA PUNCTATA	2.0	1.8	0.6	0.4	.	.	.	0.4	0.2	.	.	.	.
129.0	SYLLIS GRACILIS	1.8	1.1	0.2	0.2	0.2	0.2	0.2	0.4	0.2	.	.	.	.
134.0	ANCHIALINA TYPICA	.	.	0.8	0.5	1.2	0.4	0.2	0.4	0.2	.	.	.	.
134.0	LAONICE CIRRHATA	1.2	1.0	0.2	0.2	0.2	0.2	0.2	0.8	0.6	.	.	.	.
134.0	LEUCOTHOIDES POTTSI	.	.	1.8	1.3	0.4	0.2	0.2	0.2	0.2	.	.	.	.
134.0	PETALOPROCTUS SOCIALIS	.	.	1.8	1.8	0.6	0.4	0.4	.	.	.	.	.	.
134.0	PROCESSA SP.	.	.	.	.	.	.	.	2.4	1.0	.	.	.	.
134.0	SCOLOPLOS SP. A	.	.	.	.	0.4	0.4	0.4	2.0	1.5	.	.	.	.
134.0	SYNCHELIDIUM AMERICANUM	1.8	1.0	0.6	0.4	.	.	.	.	.	.	.	.	.
140.5	AMPHIPODA B	1.2	1.0	0.2	0.2	0.8	0.6	0.6	.	.	.	.	.	.
140.5	ANOPILODACTYLUS PETIOLATUS	1.2	0.7	0.6	0.4	0.2	0.2	0.2	0.2	0.2	.	.	.	.
140.5	HYDROIDES SP. A	0.8	0.6	.	.	0.8	0.5	0.5	0.6	0.6	.	.	.	.
140.5	LITHOPHAGA BISULCATA	0.2	0.2	0.2	0.2	0.4	0.4	0.4	1.4	0.5	.	.	.	.
140.5	MITHRAX PLEURACANTHUS	1.4	1.0	0.2	0.2	.	.	.	0.6	0.6	.	.	.	.
140.5	POMATOCEKOS AMERICANUS	0.6	0.2	1.2	0.7	0.2	0.2	0.2	0.2	0.2	.	.	.	.
151.0	AMPELISCA SCHELLENBERGI	.	.	1.2	1.0	.	.	.	0.8	0.5	.	.	.	.
151.0	ARABELLA MUTANS	0.2	0.2	1.0	0.8	0.4	0.2	0.2	0.4	0.2	.	.	.	.
151.0	CHONE SP.	.	.	.	.	2.0	2.0	2.0	.	.	.	.	.	.
151.0	COLOMASTIX SP. B	.	.	1.6	1.6	0.4	0.4	0.4	.	.	.	.	.	.
151.0	COMACTINIA MERIDIONALIS	0.6	0.4	0.2	0.2	0.6	0.6	0.6	0.6	0.6	.	.	.	.
151.0	LYSIANOPSIS ALBA	.	.	1.0	1.0	0.8	0.8	0.8	0.2	0.2	.	.	.	.
151.0	MALDANIDAE	0.6	0.2	0.4	0.4	0.8	0.6	0.6	0.2	0.2	.	.	.	.
151.0	MELPHIDIPPIIDAE A	0.4	0.2	0.4	0.2	0.6	0.4	0.4	0.6	0.4	.	.	.	.
151.0	NEPHTYS SQUAMOSA	0.4	0.2	0.2	0.2	0.6	0.6	0.6	0.8	0.5	.	.	.	.
151.0	PELECYPODA B	0.8	0.8	0.4	0.2	0.4	0.4	0.4	0.4	0.4	.	.	.	.
151.0	PHYLLODOCIDAE	0.4	0.2	1.2	1.2	0.2	0.2	0.2	0.2	0.2	.	.	.	.
151.0	SERPULIDAE	0.8	0.4	0.4	0.4	0.6	0.6	0.6	0.2	0.2	.	.	.	.
151.0	SIPUNCULIDA	0.2	0.2	0.6	0.4	0.2	0.2	0.2	0.8	0.6	.	.	.	.
151.0	SYLLIS ALTERNATA	0.2	0.2	1.6	0.9	.	.	.	.	.	.	.	.	.
151.0	SYLLIS REGULATA CAROLINAE	1.4	0.9	0.2	0.2	.	.	.	0.4	0.4	.	.	.	.
164.0	AMPHARETE ACUTIFRONS	0.2	0.2	0.6	0.4	1.0	0.6	0.6	.	.	.	.	.	.
164.0	CERITHIOPSIS GREENI	.	.	1.8	1.8	.	.	.	.	.	.	.	.	.
164.0	LEMBOS UNICORNIS	.	.	.	.	1.4	1.0	1.0	0.4	0.2	.	.	.	.
164.0	LYSIANASSIDAE A	0.4	0.2	0.8	0.4	0.2	0.2	0.2	0.4	0.2	.	.	.	.
164.0	LYSIDICE NINETTA	0.4	0.2	0.4	0.2	0.8	0.6	0.6	0.2	0.2	.	.	.	.
164.0	PAGURUS HENDERSONI *	1.8	1.8	.	.	.	.	.	.	.	.	.	.	.
164.0	PINNIXA SP.	0.4	0.4	0.8	0.2	.	.	.	0.6	0.4	.	.	.	.

Appendix 3. (Continued)

OVERALL RANK	SPECIES	STATION		MS02										
		WINTER		SPRING		SUMMER			FALL					
		MEAN	ST ERR	MEAN	ST ERR	MEAN	ST	ERR	MEAN	ST	ERR	MEAN	ST	ERR
164.0	SICYONIA LAEVIGATA	1.2	0.8	.	.	.	.	.	0.6	.	.	0.2	.	0.2
164.0	SPIO PETTIBONEAE	0.6	0.4	0.4	0.4	.	.	.	0.8	.	.	0.6	.	0.6
164.0	TIRON TROPAKIS	0.2	0.2	.	.	.	.	.	1.6	.	.	0.8	.	0.8
164.0	TURBONILLA INCISA	0.4	0.2	.	.	0.4	0.4	0.4	1.0	.	.	0.4	.	0.4
177.5	ASCORHYNCHUS CASTELLI	0.8	0.6	0.4	0.4	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
177.5	CAPRELLIDAE A	.	.	.	.	1.6	1.6	1.6	.	.	.	.	.	.
177.5	CAULLERIELLA KILLARIENSIS	1.0	0.5	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
177.5	CERITHIOPSIS EMERSONI	.	.	1.6	1.6	.	.	.	.	.	.	.	.	.
177.5	CREPIDULA ACULEATA	0.4	0.4	0.2	0.2	1.0	0.4	0.4	.	.	.	.	.	.
177.5	DIODORA CAYENENSIS	0.4	0.4	0.2	0.2	0.8	0.4	0.4	0.2	0.2	0.2	0.2	0.2	0.2
177.5	EULALIA MACROCEROS	0.4	0.4	0.6	0.4	0.4	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
177.5	GALATHEA ROSTRATA	0.4	0.2	1.2	1.2	.	.	.	.	.	.	.	.	.
177.5	GLYCERA CAPITATA	0.2	0.2	1.2	1.2	0.2	0.2	0.2	.	.	.	.	.	.
177.5	HYBOSCOLEX LONGISETA	0.4	0.2	0.8	0.4	.	.	.	0.4	.	.	0.2	.	0.2
177.5	LUMBRINERIS LATREILLI	0.6	0.6	0.8	0.6	.	.	.	0.2	.	.	0.2	.	0.2
177.5	MARGINELLA AUROECINCTA	.	.	.	.	1.0	1.0	1.0	0.6	0.6	0.6	0.4	0.4	0.4
177.5	ONUPHIS NEBULOSA	.	.	0.6	0.4	0.8	0.8	0.8	0.2	0.2	0.2	0.2	0.2	0.2
177.5	OSTRACODA	1.0	0.8	.	.	0.6	0.6	0.6	.	.	.	.	.	.
177.5	PERICLIMENES SP.	0.2	0.2	0.2	0.2	0.2	0.2	0.2	1.0	.	.	0.4	.	0.4
177.5	PHYLLODOCE FRAGILIS	0.8	0.4	0.2	0.2	0.4	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
190.0	CARIDEA	.	.	0.8	0.4	.	.	.	0.6	0.6	0.6	0.4	0.4	0.4
190.0	CHAMA CONGREGATA	0.4	0.2	.	.	0.2	0.2	0.2	0.8	0.8	0.8	0.6	0.6	0.6
190.0	LINGA AMIANTUS	.	.	0.6	0.6	0.2	0.2	0.2	0.6	0.6	0.6	0.4	0.4	0.4
190.0	NEREIDAE C	1.4	1.4	.	.	.	.	.	.	.	.	.	.	.
190.0	PAGURIDAE	0.4	0.4	1.0	0.6	.	.	.	.	.	.	.	.	.
190.0	PHYLLODOCE SP.	.	.	0.6	0.4	0.2	0.2	0.2	0.6	0.6	0.6	0.4	0.4	0.4
190.0	SEMELE BELLASTRIATA	.	.	.	.	0.2	0.2	0.2	1.2	1.2	1.2	0.6	0.6	0.6
190.0	SYLLIDAE I	1.0	1.0	0.2	0.2	.	.	.	0.2	0.2	0.2	0.2	0.2	0.2
190.0	SYLLIDAE L	.	.	0.2	0.2	0.4	0.2	0.2	0.8	0.8	0.8	0.4	0.4	0.4
203.0	BRANCHIOSYLLIS OCULATA	.	.	0.8	0.6	0.4	0.4	0.4	.	.	.	.	.	.
203.0	CHAMA MACEROPHYLLA	0.8	0.8	0.4	0.4	.	.	.	.	.	.	.	.	.
203.0	COLUBRARIA LANCEOLATA	.	.	.	.	1.2	1.2	1.2	.	.	.	.	.	.
203.0	GLYCERA TESSELATA	0.2	0.2	0.4	0.2	0.2	0.2	0.2	0.4	0.4	0.4	0.4	0.4	0.4
203.0	HOLOTHUROIDEA	0.4	0.2	.	.	0.4	0.4	0.4	0.4	0.4	0.4	0.2	0.2	0.2
203.0	HYDROIDES CRUCIGERA	0.6	0.4	.	.	0.2	0.2	0.2	0.4	0.4	0.4	0.2	0.2	0.2
203.0	LUCIFER FAXONI	.	.	.	.	1.2	0.4	0.4	.	.	.	.	.	.
203.0	MANGELIA RUGIRIMA	.	.	0.8	0.8	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
203.0	PHYLLOCARIDA	.	.	0.6	0.4	0.6	0.4	0.4	0.6	0.6	0.6	0.6	0.6	0.6
203.0	PHYLLODOCE CASTANEA	.	.	0.6	0.6	.	.	.	0.2	0.2	0.2	0.2	0.2	0.2
203.0	PSEUDOVERMILIA SP. A	0.8	0.5	0.2	0.2	.	.	.	1.0	1.0	1.0	0.5	0.5	0.5
203.0	PSEUDOVERMILIA OCCIDENTALIS	.	.	0.2	0.2	.	.	.	0.4	0.4	0.4	0.2	0.2	0.2
203.0	SABELLARIA VULGARIS VULGARIS	0.6	0.4	.	.	0.2	0.2	0.2	0.4	0.4	0.4	0.2	0.2	0.2
203.0	STENOTHOE SP. A	.	.	0.8	0.8	.	.	.	0.4	0.4	0.4	0.2	0.2	0.2
203.0	SYLLIS SP. D	0.4	0.4	0.4	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
203.0	SYLLIS SP. I	.	.	.	.	.	.	.	1.2	1.2	1.2	0.7	0.7	0.7
203.0	VARICORBULA OPERCULATA	0.4	0.2	0.2	0.2	0.2	0.2	0.2	0.4	0.4	0.4	0.4	0.4	0.4
229.0	APANTHURA MAGNIFICA	.	.	0.2	0.2	.	.	.	0.8	0.8	0.8	0.6	0.6	0.6
229.0	ARMANDIA MACULATA	0.4	0.2	.	.	.	.	.	0.6	0.6	0.6	0.2	0.2	0.2
229.0	CAPITELLIDAE A	0.4	0.4	0.2	0.2	.	.	.	0.4	0.4	0.4	0.4	0.4	0.4
229.0	CERAPUS TUBULARIS	0.2	0.2	0.8	0.6	.	.	.	.	.	.	.	.	.
229.0	CHRYSOPETALIDAE B	0.4	0.2	.	.	.	.	.	0.6	0.6	0.6	0.4	0.4	0.4
229.0	COLOMASTIX SP. A	1.0	0.8	.	.	.	.	.	.	.	.	.	.	.
229.0	CREPIDULA PLANA	.	.	1.0	1.0	.	.	.	.	.	.	.	.	.
229.0	CRUCIGERA WEBSTERI	0.2	0.2	0.4	0.2	0.4	0.2	0.2	.	.	.	.	.	.
229.0	CUMINGIA TELLINOIDES	0.8	0.6	.	.	.	.	.	0.2	0.2	0.2	0.2	0.2	0.2

Appendix 3. (Continued)

OVERALL RANK	SPECIES	STATION		MS02										
		WINTER		SPRING		SUMMER			FALL					
		MEAN	ST ERR	MEAN	ST ERR	MEAN	ST	ERR	MEAN	ST	ERR	MEAN	ST	ERR
229.0	EURYDICE LITTORALIS	0.2	0.2	0.2	0.2	0.6		0.6	1.0		0.8			
229.0	HAPLOSCOLOPLOS FRAGILIS	.	.	.	.	0.4		0.4	1.0		1.0			
229.0	HESIONIDAE	.	.	0.6	0.6	.		.	0.6		0.6			
229.0	HIPPOLYTE SP.	.	.	.	.	.		.	0.8		0.8			
229.0	HIPPOLYTIDAE	0.2	0.2	0.2	0.2	.		.	0.2		0.2			
229.0	HYPOCONCHA SABULOSA	0.2	0.2	0.4	0.4	0.4		0.4	.		.			
229.0	JAEROPSIS CURALICOLA	0.2	0.2	0.2	0.2	0.4		0.2	0.2		0.2			
229.0	LIMA PELLUCIDA	0.2	0.2	1.0	0.6	.		.	.		.			
229.0	LYONSIA HYALINA	.	.	0.8	0.6	0.2		0.2	0.6		0.4			
229.0	MAGELONA SP. A	0.4	0.4	.	.	0.6		0.6	0.4		0.4			
229.0	MAJIDAE	.	.	.	.	.		.	.		.			
229.0	MITHRAX SP.	.	.	1.0	0.5	0.4		0.4	0.2		0.2			
229.0	MODIOLUS AMERICANUS	0.4	0.4	.	.	.		.	1.0		0.6			
229.0	NEREIS SP.	.	.	.	.	.		.	0.8		0.6			
229.0	OPHELIA DENTICULATA	1.0	0.6	.	.	.		.	0.2		0.2			
229.0	PALICUS SP.	.	.	0.2	0.2	0.4		0.2	0.8		0.6			
229.0	PARAPRIONOSPION PINNATA	.	.	0.4	0.2	0.6		0.4	0.2		0.2			
229.0	PERSICULA CANTENATA	0.4	0.4	.	.	.		.	1.0		1.0			
229.0	POLYDORA SP.	.	.	0.2	0.2	0.2		0.2	0.4		0.2			
229.0	POLYDORA TETRABRANCHIA	0.2	0.2	0.2	0.2	0.2		0.2	.		.			
229.0	SCHISTOMERINGOS RUDOLPHI	0.6	0.4	0.2	0.2	0.2		0.2	1.0		1.0			
229.0	SPIONIDAE	.	.	.	.	.		.	0.4		0.2			
229.0	STREBLOSOMA SP.	0.6	0.6	.	.	1.0		1.0	.		.			
229.0	SYLLIDAE C	.	.	.	.	0.2		0.2	0.6		0.4			
229.0	SYNALPHEUS FRITZMUELLERI *	0.8	0.5	.	.	0.2		0.2	.		.			
229.0	VERMILIOPSIS ANNULATA	.	.	0.6	0.4	.		.	0.2		0.2			
263.5	AONIDES MAYAGUEZENSIS	0.2	0.2	.	.	.		.	0.8		0.6			
263.5	ASPIUDOSIPHON ALBUS	.	.	.	.	.		.	0.2		0.2			
263.5	AUTOMATE EVERMANNI	0.2	0.2	0.4	0.2	0.2		0.2	0.2		0.2			
263.5	BATRACHONOTUS FRAGOSUS	0.2	0.2	0.2	0.2	0.6		0.4	.		.			
263.5	CAPITELLIDAE	0.2	0.2	0.2	0.2	.		.	0.6		0.4			
263.5	DIPLODONTA SEMIASPERA	0.6	0.6	0.2	0.2	0.2		0.2	.		.			
263.5	EULALIA SP.	0.2	0.2	0.8	0.6	0.2		0.2	.		.			
263.5	EUNICIDAE	.	.	0.2	0.2	0.2		0.2	0.2		0.2			
263.5	GLYCERA SP. C	0.2	0.2	0.2	0.2	.		.	0.6		0.4			
263.5	GLYCERA PAPILLOSA	.	.	0.2	0.2	.		.	.		.			
263.5	HETEROCRYPTA GRANULATA	0.6	0.4	0.2	0.2	.		.	0.2		0.2			
263.5	HETEROMYSIS FORMOSA	0.8	0.6	.	.	.		.	.		.			
263.5	KURTZIELLA ATRUSTYLA	.	.	.	.	0.6		0.4	0.2		0.2			
263.5	LEMBOS UNIFASCIATUS REDUCTUS	.	.	0.4	0.4	0.4		0.4	.		.			
263.5	MELITIDAE	0.6	0.4	0.2	0.2	.		.	.		.			
263.5	MICROPANOPE SP.	.	.	0.4	0.2	.		.	.		.			
263.5	MICROPHOLIS PACHYRACTERA	0.4	0.4	0.4	0.2	.		.	.		.			
263.5	NEREIDAE	.	.	.	.	0.4		0.4	0.4		0.4			
263.5	NEREIS RIISEI	0.2	0.2	.	.	0.2		0.2	0.6		0.6			
263.5	OLIGOCHAETA	0.2	0.2	.	.	.		.	0.2		0.2			
263.5	PALAEONIDAE	.	.	0.6	0.6	.		.	0.8		0.8			
263.5	PARAONIDAE	.	.	.	.	0.2		0.2	0.2		0.2			
263.5	PERICLIMENAEUS SCHMITTI	.	.	0.4	0.4	.		.	.		.			
263.5	PHERUSA SP.	.	.	0.2	0.2	.		.	.		.			
263.5	PISANIA TINCTA	0.6	0.6	0.2	0.2	.		.	0.4		0.4			
263.5	PLEUROMERIS TRIDENTATA	0.2	0.2	0.2	0.2	.		.	.		.			
263.5	PRIONOSPION SP.	0.4	0.2	.	.	0.4		0.4	0.4		0.4			
263.5	PROTODORVILLEA KEFERSTEINI	.	.	.	.	.		.	.		.			
263.5	SCOLOPLOS RUBRA	.	.	.	.	.		.	.		.			

Appendix 3. (Continued)

OVERALL RANK	SPECIES	STATION		MS02										
		WINTER		SPRING		SUMMER			FALL					
		MEAN	ST ERR	MEAN	ST ERR	MEAN	ST	ERR	MEAN	ST	ERR	MEAN	ST	ERR
263.5	SIPUNCULIDA E	0.4	0.2	.	.	.	.	.	0.4	.	.	0.4	0.4	0.4
263.5	SPIONIDAE B	.	.	0.4	0.2	.	.	.	0.4	.	.	0.4	0.2	0.4
263.5	SYLLIS CORNUTA	.	.	0.2	0.2	0.2	0.2	0.2	0.4	.	.	0.4	0.4	0.4
263.5	TRYPTON SP.	0.2	0.2	0.4	0.4	0.2	0.2	0.2	.	.	.	.	.	.
263.5	WEBSTERINEREIS SP. A	.	.	0.6	0.4	0.2	0.2	0.2	.	.	.	.	.	.
310.5	ALPHEUS SP.	.	.	.	.	.	.	.	0.6	.	.	0.4	0.4	0.4
310.5	AMPELISCA CRISTOIDES	.	.	0.4	0.4	.	.	.	0.2	.	.	0.2	0.2	0.2
310.5	AMPHIPODA E	0.6	0.4	.	.	.	.	.	.	.	.	.	.	.
310.5	ANAMIXIS SP.	.	.	.	.	.	.	.	0.6	.	.	0.4	0.4	0.4
310.5	ANOMIA SQUAMULA	.	.	.	.	0.4	0.4	0.4	0.2	0.4	0.2	0.2	0.2	0.2
310.5	ARBACIA PUNCTULATA	.	.	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
310.5	ASTERIAS FORBESII	.	.	0.6	0.6	.	.	.	.	.	.	.	.	.
310.5	BRACHIOPODA A	0.2	0.2	0.2	0.2	0.2	0.2	0.2	.	.	.	.	.	.
310.5	BRANIA CLAVATA	0.2	0.2	0.2	0.2	.	.	.	0.2	.	.	0.2	0.2	0.2
310.5	CHIONE SP. *	0.6	0.6	.	.	.	.	.	.	.	.	.	.	.
310.5	CHIONE LATILIRATA	.	.	.	.	0.4	0.4	0.4	0.2	0.4	0.2	0.2	0.2	0.2
310.5	CIRRIFORMIA FILIGERA	.	.	0.6	0.4	.	.	.	.	.	.	.	.	.
310.5	CORBULA CONTRACTA	.	.	.	.	0.4	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
310.5	CUMACEA	0.6	0.6	.	.	.	.	.	.	.	.	.	.	.
310.5	CYCLASPIS VARIANS	0.6	0.4	.	.	.	.	.	.	.	.	.	.	.
310.5	DENDRODORIS KREBSII	0.2	0.2	0.4	0.2	.	.	.	.	.	.	.	.	.
310.5	DIASTYLIDAE	.	.	.	.	0.2	0.2	0.2	0.4	0.4	0.4	0.4	0.4	0.4
310.5	EBALIA STIMPSONI	0.2	0.2	.	.	.	.	.	0.4	0.4	0.4	0.4	0.4	0.4
310.5	ENGINA TURBINELLA	.	.	0.6	0.6	.	.	.	.	.	.	.	.	.
310.5	EUNICE SP.	0.2	0.2	.	.	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
310.5	EUNICE ANTENNATA	0.4	0.2	.	.	.	.	.	0.2	0.2	0.2	0.2	0.2	0.2
310.5	EUPHROSINE TRILOBA	0.2	0.2	.	.	.	.	.	0.4	0.4	0.4	0.4	0.4	0.4
310.5	EXOGONE SP.	0.6	0.6	.	.	.	.	.	.	.	.	.	.	.
310.5	FAVARTIA CELLULOSA	0.2	0.2	0.4	0.2	.	.	.	.	.	.	.	.	.
310.5	GLYCERA SP. B	.	.	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
310.5	HEMUS CRISTULIPES	.	.	0.2	0.2	.	.	.	0.4	0.4	0.4	0.4	0.4	0.4
310.5	HIPPOMEDON SP. A	0.2	0.2	0.4	0.4	.	.	.	.	.	.	.	.	.
310.5	HYDROIDES UNCINATA	.	.	0.2	0.2	0.4	0.4	0.4	.	.	.	.	.	.
310.5	LAEVICARDIUM MORTONI	0.6	0.6	.	.	.	.	.	.	.	.	.	.	.
310.5	LASAEIDAE	.	.	.	.	.	.	.	0.6	0.6	0.6	0.6	0.6	0.6
310.5	LEUCUTHOE SP. A	.	.	0.6	0.6	.	.	.	.	.	.	.	.	.
310.5	MAERA CAROLINIANA	0.6	0.4	.	.	.	.	.	.	.	.	.	.	.
310.5	METOPORHAPHIS CALCARATA	.	.	.	.	.	.	.	0.6	0.6	0.6	0.6	0.6	0.6
310.5	NEOPONTONIDES BEAUFORTENSIS *	0.6	0.6	.	.	.	.	.	.	.	.	.	.	.
310.5	NEPHTYS INCISA	0.2	0.2	.	.	.	.	.	0.4	0.4	0.4	0.4	0.4	0.4
310.5	ODONTOSYLLIS SP.	0.2	0.2	.	.	.	.	.	0.4	0.4	0.4	0.4	0.4	0.4
310.5	OGYRIDES SP.	.	.	.	.	0.2	0.2	0.2	0.4	0.4	0.4	0.4	0.4	0.4
310.5	ONUPHIS PALLIDULA	.	.	0.4	0.2	0.2	0.2	0.2	.	.	.	.	.	.
310.5	PAGURISTES TORTUGAE	0.2	0.2	0.2	0.2	.	.	.	0.2	0.2	0.2	0.2	0.2	0.2
310.5	PERICLIMENES AMERICANUS	.	.	.	.	.	.	.	0.6	0.6	0.6	0.6	0.6	0.6
310.5	PILUMNUS SAYI	0.4	0.4	0.2	0.2	.	.	.	0.4	0.4	0.4	0.4	0.4	0.4
310.5	PINNOTHERIDAE	.	.	0.2	0.2	.	.	.	0.4	0.4	0.4	0.4	0.4	0.4
310.5	PISTONE REMOTA	.	.	.	.	0.2	0.2	0.2	0.4	0.4	0.4	0.4	0.4	0.4
310.5	PITHO LHERMINIERI	.	.	.	.	0.2	0.2	0.2	0.4	0.4	0.4	0.4	0.4	0.4
310.5	SABELLIDAE	0.2	0.2	.	.	.	.	.	0.4	0.4	0.4	0.4	0.4	0.4
310.5	SEMELE PURPURASCENS	0.2	0.2	.	.	.	.	.	0.2	0.2	0.2	0.2	0.2	0.2
310.5	SEROLIS MGRAYI	.	.	.	.	0.4	0.4	0.4	0.2	0.2	0.2	0.2	0.2	0.2
310.5	SPEOCARCINUS CAROLINENSIS	.	.	.	.	.	.	.	0.6	0.6	0.6	0.6	0.6	0.6
310.5	STENORHYNCHUS SETICORNIS	0.2	0.2	0.2	0.2	.	.	.	0.2	0.2	0.2	0.2	0.2	0.2
310.5	STROMBIFORMIS BILINEATUS	0.2	0.2	0.2	0.2	.	.	.	0.2	0.2	0.2	0.2	0.2	0.2

Appendix 3. (Continued)

OVERALL RANK	SPECIES	STATION		MS02										
		WINTER		SPRING		SUMMER			FALL					
		MEAN	ST ERR	MEAN	ST ERR	MEAN	ST	ERR	MEAN	ST	ERR	MEAN	ST	ERR
310.5	SUBADYTE PELLUCIDA	.	.	0.6	0.6	.	.	.	.	.	.	.	.	.
310.5	SYLLIDAE	0.2	0.2	0.2	0.2	0.2	0.2	.	.	.	.	.	.	.
310.5	SYLLIS SP. B	0.2	0.2	.	.	.	.	.	0.4	0.4	.	.	.	.
310.5	SYLLIS SP. G	.	.	0.6	0.2	.	.	.	.	.	.	.	.	.
310.5	TANAIDACEA B	0.2	0.2	.	.	.	.	.	0.4	0.4	.	.	.	.
310.5	TEREBELLIDAE	0.2	0.2	0.2	0.2	.	.	.	0.2	0.2	.	.	.	.
310.5	THOR FLORIDANUS	.	.	0.6	0.6	.	.	.	.	.	.	.	.	.
310.5	THOR MANNINGI	.	.	.	.	0.2	0.2	.	0.4	0.2	.	.	.	.
310.5	TRIPHORA NIGROCINCTA	.	.	0.6	0.6	.	.	.	.	.	.	.	.	.
310.5	TURBONILLA INTERRUPTA	.	.	0.6	0.4	.	.	.	.	.	.	.	.	.
369.5	ACANTHOHAUSTORIUS SHOEMAKERI	0.4	0.2	.	.	.	.	.	.	.	.	.	.	.
369.5	AMPHIODIA TRYCHNA	0.4	0.2	.	.	.	.	.	.	.	.	.	.	.
369.5	AMPHIPODA F	0.2	0.2	.	.	.	.	.	0.2	0.2	.	.	.	.
369.5	ARCA ZERRA	0.2	0.2	.	.	0.2	0.2	.	.	.	.	.	.	.
369.5	ARICIDEA SP.	.	.	0.4	0.4	.	.	.	.	.	.	.	.	.
369.5	ASTROPECTEN SP.	.	.	0.4	0.4	.	.	.	.	.	.	.	.	.
369.5	BARNEA TRUNCATA	0.4	0.4	.	.	.	.	.	.	.	.	.	.	.
369.5	BODOTRIIDAE A	0.2	0.2	.	.	.	.	.	0.2	0.2	.	.	.	.
369.5	CHIONE INTAPURPUREA	0.4	0.4	.	.	.	.	.	.	.	.	.	.	.
369.5	CIRRATULUS SP.	.	.	0.4	0.4	.	.	.	.	.	.	.	.	.
369.5	DIODORA LISTERI	.	.	.	.	0.4	0.4	.	.	.	.	.	.	.
369.5	DORIOPSILLA ALBOPUNCTATA	0.4	0.4	.	.	.	.	.	.	.	.	.	.	.
369.5	DORIOPSILLA ROWENA	0.4	0.2	.	.	.	.	.	.	.	.	.	.	.
369.5	ETHUSA MASCARONE AMERICANA	0.2	0.2	.	.	.	.	.	0.2	0.2	.	.	.	.
369.5	EULALIA VIRIDIS	.	.	.	.	.	.	.	0.4	0.2	.	.	.	.
369.5	HARMOTHOE SP.	.	.	.	.	.	.	.	0.4	0.4	.	.	.	.
369.5	HEMIPODUS ROSEUS	0.4	0.4	.	.	.	.	.	.	.	.	.	.	.
369.5	HYDROIDES SP. C	.	.	0.4	0.4	.	.	.	.	.	.	.	.	.
369.5	LATREUTES SP.	.	.	.	.	.	.	.	0.4	0.4	.	.	.	.
369.5	LEPTOGNATHIIDAE A	0.2	0.2	0.2	0.2	.	.	.	.	.	.	.	.	.
369.5	LUMBRINERIOPSIS PARADOXA	.	.	.	.	0.2	0.2	.	0.2	0.2	.	.	.	.
369.5	LYONSIA SP.	.	.	.	.	0.4	0.4	.	.	.	.	.	.	.
369.5	LYONSIA SP. *	0.4	0.2	.	.	.	.	.	.	.	.	.	.	.
369.5	LYONSIA BEANA	.	.	.	.	.	.	.	.	.	.	.	.	.
369.5	LYSIDICE SP. A	.	.	.	.	0.4	0.4	.	.	.	.	.	.	.
369.5	MADRELLA SP.	0.4	0.4	.	.	.	.	.	.	.	.	.	.	.
369.5	MEDIOMASTUS CALIFORNIENSIS	0.2	0.2	.	.	.	.	.	0.2	0.2	.	.	.	.
369.5	MITHRAX FORCEPS	.	.	0.2	0.2	.	.	.	0.2	0.2	.	.	.	.
369.5	MUNIDA SP.	0.2	0.2	.	.	0.2	0.2	.	.	.	.	.	.	.
369.5	MUREX POMUM	0.4	0.2	.	.	.	.	.	.	.	.	.	.	.
369.5	NATICA PUSILLA	.	.	.	.	.	.	.	0.4	0.4	.	.	.	.
369.5	NOTOMASTUS HEMIPODUS	0.2	0.2	.	.	.	.	.	0.2	0.2	.	.	.	.
369.5	OCNUS PYGMAEUS	0.2	0.2	.	.	0.2	0.2	.	.	.	.	.	.	.
369.5	ONUPHIDAE	0.4	0.2	.	.	.	.	.	.	.	.	.	.	.
369.5	OPHIODERMA BREVISPINUM	0.2	0.2	0.2	0.2	.	.	.	.	.	.	.	.	.
369.5	PARAPHOXUS SPINOSUS	0.4	0.4	.	.	.	.	.	.	.	.	.	.	.
369.5	PELECYPODA G	.	.	.	.	.	.	.	0.4	0.2	.	.	.	.
369.5	PERICLIMENAEUS SP. A	0.4	0.4	.	.	.	.	.	.	.	.	.	.	.
369.5	PHIDIANA LYNCEUS	.	.	.	.	0.4	0.2	.	.	.	.	.	.	.
369.5	PILUMNUS DASYPODUS	.	.	0.4	0.4	.	.	.	.	.	.	.	.	.
369.5	PODOCHELA SP.	.	.	0.4	0.2	.	.	.	.	.	.	.	.	.
369.5	PODOCHELA GRACILIPES	.	.	.	.	.	.	.	0.4	0.4	.	.	.	.
369.5	POLYCERA AURISULA	.	.	0.2	0.2	.	.	.	0.2	0.2	.	.	.	.
369.5	PORTUNUS SP.	.	.	0.2	0.2	.	.	.	0.2	0.2	.	.	.	.
369.5	PRIONOSPIO SP. C	0.4	0.4	.	.	.	.	.	.	.	.	.	.	.

Appendix 3. (Continued)

OVERALL RANK	SPECIES	STATION		MS02									
		WINTER		SPRING		SUMMER		FALL					
		MEAN	ST ERR	MEAN	ST ERR	MEAN	ST ERR	MEAN	ST ERR	MEAN	ST ERR	MEAN	ST ERR
369.5	PSEUDEURYTHOE AMBIGUA	.	.	.	.	0.4	0.4	.	.	.	.	.	.
369.5	SCOLOPLOS ARMIGER	.	.	.	.	0.4	0.2	.	.	.	.	.	.
369.5	SIRAIUS BICOLOR	0.2	0.2	.	.	0.2	0.2	.	.	.	.	.	.
369.5	SPEOCARCINUS CAROLINENSIS *	.	.	.	.	.	.	0.4	0.4	.	.	.	.
369.5	STENOTHOE SP. B	.	.	0.4	0.4	.	.	.	.	.	.	.	.
369.5	STENOTHOE GEORGIANA	0.4	0.4	.	.	.	.	.	.	.	.	.	.
369.5	SYNALPHEUS MINUS	.	.	.	.	0.4	0.4	.	.	.	.	.	.
369.5	TELLINA ALTERNATA	0.4	0.4	.	.	.	.	.	.	.	.	.	.
369.5	THYONE PSEUDOFUSUS	0.2	0.2	0.2	0.2	.	.	.	.	.	.	.	.
369.5	TRICHOPOXUS FLORIDANUS	0.2	0.2	.	.	.	.	0.2	0.2	.	.	.	.
369.5	TURRONILLA SP. D	0.4	0.4	.	.	.	.	.	.	.	.	.	.
369.5	TURRIDAE F	0.4	0.2	.	.	.	.	.	.	.	.	.	.
369.5	ZEBINA BROWNIANA	.	.	.	.	0.4	0.2	.	.	.	.	.	.
488.5	ACTEON CANDENS	.	.	0.2	0.2	.	.	.	.	.	.	.	.
488.5	ALABA INCERTA	.	.	.	.	0.2	0.2	.	.	.	.	.	.
488.5	ALPHEIDAE	.	.	.	.	.	.	0.2	0.2	.	.	.	.
488.5	ALPHEUS SP. B	.	.	.	.	0.2	0.2	.	.	0.2	0.2	.	.
488.5	ALPHEUS FORMOSUS	0.2	0.2	.	.	.	.	.	.	.	.	.	.
488.5	AMBLYOSYLLIS FORMOSA	.	.	.	.	.	.	0.2	0.2	.	.	.	.
488.5	AMPELISCA VERRILLI	.	.	0.2	0.2	.	.	.	.	0.2	0.2	.	.
488.5	AMPHILOCHUS SP.	0.2	0.2	.	.	.	.	.	.	.	.	.	.
488.5	AMPHIPODA G	.	.	.	.	0.2	0.2	.	.	.	.	.	.
488.5	AMPHITHOIDEA A	.	.	.	.	0.2	0.2	.	.	.	.	.	.
488.5	AMYGDALUM SAGITTATUM	.	.	0.2	0.2	.	.	.	.	.	.	.	.
488.5	ANCISTROSYLLIS RIGIDA	.	.	.	.	.	.	0.2	0.2	.	.	.	.
488.5	ANOPLODACTYLUS INSIGNIS	.	.	0.2	0.2	.	.	.	.	.	.	.	.
488.5	ARABELLA IRICOLOR	.	.	0.2	0.2	.	.	.	.	.	.	.	.
488.5	ARCOPSIS ADAMSI	.	.	.	.	0.2	0.2	.	.	.	.	.	.
488.5	ARENE TRICARINATA	.	.	.	.	.	.	0.2	0.2	0.2	0.2	.	.
488.5	ARICIDEA FRAGILIS	.	.	.	.	.	.	0.2	0.2	0.2	0.2	.	.
488.5	ARICIDEA SUECICA	.	.	.	.	0.2	0.2	.	.	.	.	.	.
488.5	AUSTRAEOLIS CATINA	.	.	.	.	.	.	0.2	0.2	0.2	0.2	.	.
488.5	BARLEFIA SP. *	0.2	0.2	.	.	.	.	.	.	.	.	.	.
488.5	BODOTRIIDAE C	0.2	0.2	.	.	.	.	.	.	.	.	.	.
488.5	BODOTRIIDAE D	.	.	.	.	.	.	0.2	0.2	.	.	.	.
488.5	BOTULA FUSCA	.	.	.	.	0.2	0.2	.	.	.	.	.	.
488.5	BRANCHIOSYLLIS SP.	0.2	0.2	.	.	.	.	.	.	.	.	.	.
488.5	BRANIA SP.	.	.	0.2	0.2	.	.	.	.	.	.	.	.
488.5	CAECUM CUBITATUM	0.2	0.2	.	.	.	.	.	.	.	.	.	.
488.5	CALLIOSTOMA SP.	.	.	0.2	0.2	.	.	.	.	.	.	.	.
488.5	CALLIOSTOMA PULCHRUM *	.	.	0.2	0.2	.	.	.	.	.	.	.	.
488.5	CAPITELLIDAE B	.	.	0.2	0.2	.	.	.	.	.	.	.	.
488.5	CERADOCUS SP.	0.2	0.2	.	.	.	.	.	.	.	.	.	.
488.5	CERATONEREIS IRRITABILIS	.	.	0.2	0.2	.	.	.	.	.	.	.	.
488.5	CHAETOPTERIDAE	.	.	.	.	0.2	0.2	.	.	.	.	.	.
488.5	CHAETOZONE SETOSA	.	.	.	.	0.2	0.2	.	.	.	.	.	.
488.5	CHIONE CANCELLATA	0.2	0.2	.	.	.	.	.	.	.	.	.	.
488.5	CHRYSOPETALIDAE C	0.2	0.2	.	.	.	.	.	.	.	.	.	.
488.5	CHRYSOPETALIDAE D	.	.	.	.	.	.	0.2	0.2	.	.	.	.
488.5	CONUS DAUCUS	0.2	0.2	.	.	.	.	.	.	.	.	.	.
488.5	CRENELLA DIVARICATA	.	.	0.2	0.2	.	.	.	.	.	.	.	.
488.5	CUMINGIA COARCTATA	.	.	0.2	0.2	.	.	.	.	.	.	.	.
488.5	DASYBRANCHUS SP.	.	.	.	.	.	.	.	.	0.2	0.2	.	.
488.5	DELECTOPECTEN SP.	0.2	0.2	.	.	.	.	.	.	.	.	.	.
488.5	DENDRODURIS SP. A	.	.	0.2	0.2	.	.	.	.	.	.	.	.

Appendix 3. (Continued)

OVERALL RANK	SPECIES	STATION		MS02										
		WINTER		SPRING		SUMMER			FALL					
		MEAN	ST ERR	MEAN	ST ERR	MEAN	ST	ERR	MEAN	ST	ERR	MEAN	ST	ERR
488.5	DIODORA CAYENENSIS *	0.2	0.2	.	.	0.2	.	.	.	.	.	.	.	.
488.5	DIOPATRA CUPREA	.	.	.	.	.	.	.	0.2	.	.	0.2	.	.
488.5	DONDICE OCCIDENTALIS	.	.	.	.	.	.	.	.	.	.	.	.	.
488.5	DORTOPSILLA LEIA	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.
488.5	DORVILLEA SOCIABILIS	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.
488.5	DROMIDIA ANTILLENIS	.	.	.	.	0.2	.	.	.	.	.	.	.	.
488.5	DROMIDIA ANTILLENIS *	.	.	0.2	0.2	.	.	.	.	.	.	.	.	.
488.5	ECHINASTER SP.	.	.	.	.	0.2	.	.	0.2	.	.	0.2	.	.
488.5	EPITONIUM FOLIACEICOSTUM	.	.	.	.	0.2	.	.	.	.	.	.	.	.
488.5	ERATO MAUGERIAE	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.
488.5	ERICHSONELLA FILIFORMIS	.	.	.	.	0.2	.	.	0.2	.	.	0.2	.	.
488.5	ETEONE LACTEA	.	.	.	.	.	.	.	.	.	.	0.2	.	.
488.5	GALATHEIDAE	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.
488.5	GASTROPODA	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.
488.5	GENOCIDARIS MACULATA	.	.	.	.	.	.	.	0.2	.	.	0.2	.	.
488.5	GLOTTIDIA PYRAMIDATA	.	.	0.2	0.2	.	.	.	.	.	.	.	.	.
488.5	GLYCERA SP.	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.
488.5	GLYCYMERIS SP.	.	.	.	.	.	.	.	0.2	.	.	0.2	.	.
488.5	GONEPLACIDAE	.	.	.	.	.	.	.	0.2	.	.	0.2	.	.
488.5	GONEPLAX SIGSBEI	.	.	.	.	0.2	.	0.2	.	.	.	.	.	.
488.5	GONADIDAE	.	.	.	.	0.2	.	0.2	.	.	.	.	.	.
488.5	GONODACTYLUS BREDINI	.	.	.	.	.	.	.	0.2	.	.	0.2	.	.
488.5	GRAPSIDAE	.	.	.	.	.	.	.	0.2	.	.	0.2	.	.
488.5	GREGARIELLA CORALLIOPHAGA	.	.	0.2	0.2	.	.	.	.	.	.	.	.	.
488.5	GYTIS BREVI PALPA	.	.	0.2	0.2	.	.	.	.	.	.	.	.	.
488.5	HAPLOSCOLOPLOS FOLIUSUS	.	.	0.2	0.2	.	.	.	.	.	.	.	.	.
488.5	HERMODICE CARUNCULATA	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.
488.5	HYDROIDES SP. B	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.
488.5	HYDROIDES SP. D	.	.	0.2	0.2	.	.	.	.	.	.	.	.	.
488.5	HYPERIDEA	.	.	.	.	0.2	.	0.2	.	.	.	.	.	.
488.5	INACHOIDES FORCEPS *	.	.	0.2	0.2	.	.	.	.	.	.	.	.	.
488.5	IRIDOPAGURUS DISPAR *	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.
488.5	ISCHNOCHITONIDAE A	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.
488.5	LEIOCAPITELLA GLABRA	.	.	0.2	0.2	.	.	.	.	.	.	.	.	.
488.5	LEIOCHRIDES PALLIDIOR	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.
488.5	LEMROS SPINICARPUS INERMIS	.	.	.	.	.	.	.	0.2	.	.	0.2	.	.
488.5	LEPTOCHELA SP.	.	.	.	.	0.2	.	0.2	.	.	.	0.2	.	.
488.5	LUCAPINELLA LIMATULA	.	.	0.2	0.2	.	.	.	.	.	.	.	.	.
488.5	LUMBRINERIDES ACUTA	.	.	.	.	0.2	.	0.2	.	.	.	.	.	.
488.5	LYSIANASSA SP.	.	.	.	.	0.2	.	0.2	.	.	.	.	.	.
488.5	MACROCOELOMA SP. *	.	.	.	.	0.2	.	0.2	.	.	.	0.2	.	.
488.5	MACROCOELOMA EUTHECA *	.	.	.	.	.	.	.	0.2	.	.	0.2	.	.
488.5	MAGELONA SP.	.	.	.	.	0.2	.	0.2	.	.	.	.	.	.
488.5	MAGELONA SP. C	.	.	.	.	0.2	.	0.2	.	.	.	.	.	.
488.5	MALLUVIUM BENTHOPHILUM *	.	.	.	.	0.2	.	0.2	.	.	.	.	.	.
488.5	MARIONIA CUCULLATA	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.
488.5	MEGALOMMA SP.	0.2	0.2	.	.	.	.	.	0.2	.	.	0.2	.	.
488.5	MEIOSQUILLA QUADRIDENS	.	.	.	.	.	.	.	.	.	.	.	.	.
488.5	MELANELLA CONGIDEA	.	.	.	.	0.2	.	0.2	.	.	.	0.2	.	.
488.5	MUREX FLORIFER DILECTUS	.	.	.	.	.	.	.	0.2	.	.	0.2	.	.
488.5	MURICIDAE	.	.	.	.	.	.	.	0.2	.	.	0.2	.	.
488.5	NAINERIS SP.	.	.	0.2	0.2	.	.	.	.	.	.	.	.	.
488.5	NANOPLAX XANTHIFORMIS *	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.
488.5	NATANTIA	.	.	0.2	0.2	.	.	.	.	.	.	.	.	.
488.5	NEMATONEKEIS SP.	.	.	0.2	0.2	.	.	.	.	.	.	.	.	.

Appendix 3. (Continued)

OVERALL RANK	SPECIES	STATION MS02											
		WINTER			SPRING			SUMMER			FALL		
		MEAN	ST	ERR	MEAN	ST	ERR	MEAN	ST	ERR	MEAN	ST	ERR
488.5	NEMERTINEA E	0.2		0.2	.		.	.		.	.		.
488.5	NEOPONTONIDES BEAUFORTENSIS	.		.	.		.	.		.	0.2		0.2
488.5	NEREIDAE A	.		.	.		.	.		.	0.2		0.2
488.5	NEREIS SUCCINEA	.		.	0.2		0.2	.		.	.		.
488.5	NOTOMASTUS LATERICEUS	.		.	0.2		0.2	.		.	.		.
488.5	ONUPHIS EREMITA	.		.	.		.	.		.	0.2		0.2
488.5	OPHIOCNIDA SP.	.		.	.		.	.		.	0.2		0.2
488.5	OPHIOCNIDA SCABRIUSCULA	0.2		0.2	.		.	.		.	.		.
488.5	OPHIOPHRAGMUS PULCHER	0.2		0.2	.		.	.		.	.		.
488.5	ORBINIA RISERI	.		.	.		.	.		.	0.2		0.2
488.5	ORBINITIDAE	.		.	.		.	0.2		0.2	.		.
488.5	OSTREA PERMOLIS	0.2		0.2	.		.	.		.	.		.
488.5	OVALIPES SP.	.		.	.		.	.		.	0.2		0.2
488.5	PACHYCHELES KUGIMANUS	.		.	.		.	.		.	0.2		0.2
488.5	PAGURISTES SP.	.		.	0.2		0.2	.		.	.		.
488.5	PAGURISTES TRIANGULATUS	.		.	0.2		0.2	.		.	.		.
488.5	PALEANOTUS SP. B	.		.	.		.	0.2		0.2	.		.
488.5	PALOLO SICILIENSIS	.		.	.		.	.		.	0.2		0.2
488.5	PAPYRIDEA SOLENIFORMIS	0.2		0.2	.		.	.		.	.		.
488.5	PARAMETOPELLA CYPRIS	.		.	.		.	.		.	0.2		0.2
488.5	PARAPIONOSYLLIS SP. A	0.2		0.2	.		.	.		.	.		.
488.5	PECTINIDAE A	.		.	0.2		0.2	.		.	.		.
488.5	PELECYPODA	0.2		0.2	.		.	.		.	.		.
488.5	PELECYPODA C	.		.	.		.	0.2		0.2	.		.
488.5	PELECYPODA H	.		.	.		.	0.2		0.2	.		.
488.5	PENAEIDAE	.		.	.		.	.		.	0.2		0.2
488.5	PERICLIMENES IRIDESCENS	.		.	.		.	.		.	0.2		0.2
488.5	PETROLISTHES GALATHINUS	.		.	.		.	.		.	0.2		0.2
488.5	PHASCOLOPSIS GOULDI	.		.	.		.	.		.	0.2		0.2
488.5	PHORONIS ARCHITECTA	.		.	0.2		0.2	.		.	.		.
488.5	PHYTISICA MARINA	0.2		0.2	.		.	.		.	.		.
488.5	PHYLLODOCE SP. E	.		.	0.2		0.2	.		.	.		.
488.5	PHYLLODOCE ARENAE	.		.	.		.	0.2		0.2	.		.
488.5	PHYLLODOCE LONGIPES	.		.	.		.	0.2		0.2	.		.
488.5	PILUMNUS FLORIDANUS	.		.	0.2		0.2	.		.	.		.
488.5	PINNOTHERES SP.	.		.	0.2		0.2	.		.	.		.
488.5	PIONOSYLLIS SP.	.		.	.		.	0.2		0.2	.		.
488.5	PITAR FULMINATUS	.		.	.		.	0.2		0.2	.		.
488.5	POLYCHAETA	.		.	0.2		0.2	.		.	.		.
488.5	POLYCHAETA H	.		.	.		.	.		.	0.2		0.2
488.5	POLYCIRRUS SP. B	.		.	0.2		0.2	.		.	.		.
488.5	POLYCIRRUS EXIMIUS DUBIUS	0.2		0.2	.		.	.		.	.		.
488.5	POLYNOIDAE	.		.	.		.	0.2		0.2	.		.
488.5	PONTONINAE	.		.	0.2		0.2	.		.	.		.
488.5	POTAMILLA RENIFORMIS	0.2		0.2	.		.	.		.	.		.
488.5	PROSCOLOPLOS SP.	.		.	.		.	0.2		0.2	.		.
488.5	PSAMMOLYCE CTENIDOPHORA	.		.	.		.	0.2		0.2	.		.
488.5	PTERIA COLYMBUS	0.2		0.2	.		.	.		.	.		.
488.5	PTEROMERIS PERPLANA	.		.	.		.	.		.	0.2		0.2
488.5	PTILANTHURA TRICARINA	.		.	.		.	0.2		0.2	.		.
488.5	PYRUNCULUS CAELATUS	0.2		0.2	.		.	.		.	.		.
488.5	RILDARDANUS LAMINOSA	.		.	0.2		0.2	.		.	.		.
488.5	RISSOINA SP. A	.		.	0.2		0.2	.		.	.		.
488.5	RISSOINA CATESBYANA	0.2		0.2	.		.	.		.	.		.
488.5	SCOLOPLOS SP.	0.2		0.2	.		.	.		.	.		.

Appendix 3. (Continued)

OVERALL RANK	SPECIES	STATION MS02											
		WINTER			SPRING			SUMMER			FALL		
		MEAN	ST	ERR	MEAN	ST	ERR	MEAN	ST	ERR	MEAN	ST	ERR
488.5	SICYONIA BREVIROSTRIS	.	.	.	.	.	.	.	.	.	0.2	.	0.2
488.5	SICYONIA TYPICA	.	.	.	.	.	.	.	.	.	0.2	.	0.2
488.5	SICYONIA TYPICA *	.	.	.	.	.	.	.	.	.	0.2	.	0.2
488.5	SIGALIONIDAE	.	.	.	.	.	.	0.2	.	0.2	.	.	.
488.5	SIPUNCULUS NUDUS	.	.	.	.	.	.	0.2	.	0.2	.	.	.
488.5	SIRAIUS BICOLOR *	0.2	.	0.2	.	.	.	.	.	.	.	.	.
488.5	SPELOPHORUS PONTIFER	.	.	.	0.2	.	0.2	.	.	.	.	.	.
488.5	SPHAEROSYLLIS SP.	.	.	.	.	.	.	0.2	.	0.2	.	.	.
488.5	SPIONIDAE A	.	.	.	0.2	.	0.2	.	.	.	.	.	.
488.5	SPIOPHANES SP.	.	.	.	0.2	.	0.2	.	.	.	.	.	.
488.5	STENOCLIONOPS FURCATA CUELATA	.	.	.	.	.	.	0.2	.	0.2	.	.	.
488.5	STHENELAIS BOA	.	.	.	.	.	.	.	.	.	0.2	.	0.2
488.5	STROMBIFORMIS AURICINCTUS	.	.	.	0.2	.	0.2	.	.	.	.	.	.
488.5	SYLLIDAE D	.	.	.	.	.	.	.	.	.	0.2	.	0.2
488.5	SYLLIDAE J	0.2	.	0.2	.	.	.	.	.	.	0.2	.	0.2
488.5	SYLLIS FERRUGINA	.	.	.	.	.	.	.	.	.	0.2	.	0.2
488.5	TELLINA SP.	.	.	.	0.2	.	0.2	.	.	.	.	.	.
488.5	TEREBELLIDAE A	0.2	.	0.2	.	.	.	.	.	.	.	.	.
488.5	TEREBELLIDAE D	.	.	.	.	.	.	.	.	.	0.2	.	0.2
488.5	TEREBELLIDAE G	.	.	.	.	.	.	.	.	.	0.2	.	0.2
488.5	TEREBELLIDES STROEMI	0.2	.	0.2	.	.	.	.	.	.	.	.	.
488.5	THECACEA PENNIGERA	0.2	.	0.2	.	.	.	.	.	.	.	.	.
488.5	THELEPUS SETOSUS	.	.	.	.	.	.	0.2	.	0.2	.	.	.
488.5	THUNOR SP. *	0.2	.	0.2	.	.	.	.	.	.	.	.	.
488.5	TRIVIA CANDIDULA	.	.	.	.	.	.	.	.	.	0.2	.	0.2
488.5	TRIVIA PEDICULUS	.	.	.	.	.	.	0.2	.	0.2	.	.	.
488.5	TURRHONILLA SP. F	.	.	.	.	.	.	.	.	.	0.2	.	0.2
488.5	VOLVULELLA PERSIMILIS	.	.	.	0.2	.	0.2	.	.	.	.	.	.

Appendix 3. (Continued)

OVERALL RANK	SPECIES	STATION MS06									
		WINTER		SPRING			SUMMER			FALL	
		MEAN	ST ERR	MEAN	ST	ERR	MEAN	ST	ERR	MEAN	ST ERR
1.0	FILOGRANA IMPLEXA	0.6	0.4	0.2	0.2	0.2	1.6	1.0	1.0	4155.0	2605.6
2.0	LUCONACIA INCERTA	45.0	29.8	3.4	0.9	0.4	0.4	0.2	0.2	0.8	0.4
3.0	SYLLIS SPONGICOLA	16.0	8.8	0.2	0.2	0.2	6.6	3.8	3.8	20.2	13.3
4.0	CHONE AMERICANA	5.2	1.8	3.8	2.8	2.8	24.6	11.3	2.4	2.4	0.7
5.0	SPIOPHANES BOMBYX	9.4	3.0	22.2	13.4	1.6	1.6	0.8	1.6	1.6	1.1
6.0	SYLLIS HYALINA	12.8	4.2	5.8	3.0	3.0	4.6	0.7	11.4	4.5	4.5
7.0	ASPIDOSIPHON GOSNOLDI	7.6	1.6	1.0	0.5	0.5	9.8	5.3	5.6	5.6	2.8
8.0	POLYDORA COLONIA	21.4	21.4	.	.	.	.	.	.	.	.
9.0	AUTOLYTUS SP.	0.6	0.4	.	.	.	0.2	0.2	0.2	18.4	16.2
10.0	MALACOCEROS GLUTAEUS	6.2	5.3	2.0	1.2	1.2	8.8	4.4	1.8	1.8	0.6
11.0	LUMBRINEIS INFLATA	6.2	3.5	0.4	0.4	0.4	4.8	1.5	5.8	5.8	2.5
12.5	EUNICE VITTATA	3.0	2.0	4.0	1.9	1.9	4.2	1.4	3.8	0.9	0.9
12.5	WEBSTERINEREIS SP. B	5.8	3.9	1.2	0.6	0.6	2.8	0.9	5.2	3.0	3.0
14.0	CHRYSOPETALIDAE A	1.6	0.8	11.0	6.5	1.4	1.4	1.2	0.8	0.8	0.5
15.0	BRANIA CLAVATA	1.8	1.3	0.4	0.2	0.2	.	.	11.0	7.1	7.1
16.0	ACANTHOHAUSTORIUS MILLSI	4.8	2.1	7.6	4.1	4.1	0.6	0.4	1.2	1.2	0.6
17.0	OWENIA FUSIFORMIS	6.0	3.1	2.8	1.7	1.7	2.8	1.1	1.7	1.7	0.6
18.5	NEMERTINEA	3.0	1.1	2.4	1.5	1.5	3.6	0.9	2.6	0.7	0.7
18.5	PHYLLOCHAETOPTERUS SOCIALIS	.	.	0.2	0.2	0.2	10.0	5.5	1.4	1.4	0.7
20.0	EXOGONE DISPAR	7.2	2.6	0.6	0.4	0.4	2.0	0.8	1.6	1.6	1.0
21.0	NEMATODA	5.2	2.3	1.4	0.6	0.6	4.2	1.3	0.4	0.4	0.2
22.0	THOR SP.	0.8	0.6	.	.	.	.	.	10.2	3.8	3.8
23.0	OPHIACTIS SAVIGNYI	6.8	5.4	0.6	0.4	0.4	1.2	0.8	1.8	1.8	1.3
24.0	OPHIOTHRIX ANGULATA	2.0	1.0	0.6	0.2	0.2	0.6	0.4	6.0	2.5	2.5
25.0	PODOCERUS SP.	7.8	3.2	0.2	0.2	0.2	.	.	1.0	0.6	0.6
26.0	SIPUNCULIDA	3.2	1.0	0.2	0.2	0.2	2.0	0.7	3.2	2.1	2.1
27.0	LEUCOTHOE SPINICARPA	0.6	0.4	1.2	1.0	1.0	2.0	1.0	4.6	2.0	2.0
28.5	ELASMOPUS SP. B	0.6	0.6	.	.	.	4.0	3.5	3.4	1.6	1.6
28.5	EULALIA SANGUINEA	1.8	0.6	.	.	.	2.6	1.4	3.6	1.5	1.5
30.5	AMPHIODIA PULCHELLA	0.6	0.4	4.4	1.5	1.5	1.6	0.7	1.2	1.2	1.0
30.5	HYDROIDES CRUCIGERA	5.2	3.7	.	.	.	0.8	0.6	1.8	1.0	1.0
32.0	NOTOMASTUS SP.	3.0	2.3	0.4	0.2	0.2	1.6	0.5	2.4	0.9	0.9
33.0	CHIONE GKUS	1.0	0.4	0.4	0.2	0.2	5.2	1.9	0.6	0.6	0.4
34.5	SYLLIS SP. D	3.0	2.3	1.4	0.7	0.7	0.8	0.5	1.6	0.6	0.6
34.5	SYLLIS REGULATA CAROLINAE	5.4	3.1	.	.	.	.	.	1.4	0.9	0.9
36.0	SYLLIDAE	1.6	0.9	.	.	.	1.0	0.5	4.0	2.8	2.8
37.0	LAEVICARDIUM PICTUM	0.2	0.2	1.0	0.3	0.3	4.6	1.6	0.4	0.4	0.4
38.5	BRANCHIOSYLLIS EXILIS	3.2	1.9	0.6	0.4	0.4	0.4	0.2	1.8	0.8	0.8
38.5	PAGURUS CAROLINENSIS	0.4	0.4	0.8	0.8	0.8	2.4	1.2	2.4	1.7	1.7
40.0	PROTOMEDEIA SP. A	2.6	1.9	0.2	0.2	0.2	.	.	3.0	2.0	2.0
41.5	MEGALOMMA BIOCATULUM	1.4	0.9	0.2	0.2	0.2	3.0	1.7	1.0	0.8	0.8
41.5	UNCIOLA SP. A	0.2	0.2	5.4	3.4	3.4	.	.	.	.	.
43.0	OPHIOTISMA ISACANTHUM	.	.	.	.	.	3.8	2.8	1.6	1.2	1.2
44.0	POLYCIARRUS CAROLINENSIS	2.0	1.0	0.2	0.2	0.2	0.6	0.2	2.4	1.4	1.4
45.0	CRASSINELLA LUNULATA	1.4	1.0	1.4	0.7	0.7	1.8	0.7	0.4	0.4	0.4
47.5	ELASMOPUS SP. A	0.4	0.4	1.8	1.3	1.3	2.0	1.3	0.6	0.4	0.4
47.5	LEMBOS SMITHI	1.2	1.2	2.8	2.1	2.1	.	.	0.8	0.8	0.4
47.5	LEMBOS SPINICARPUS INERMIS	.	.	.	.	.	.	.	4.8	2.4	2.4
47.5	TELLINA AMERICANA	1.2	0.5	2.4	0.6	0.6	1.2	0.4	.	.	.
52.0	AMPELISCA SCHELLENBERGI	1.4	1.0	0.4	0.4	0.4	1.0	0.6	1.8	0.9	0.9
52.0	AXIOGNATHUS SQUAMATUS	1.8	1.1	0.6	0.2	0.2	1.8	1.3	0.4	0.4	0.4
52.0	MEGALUROPOUS SP.	4.6	2.7	.	.	.	.	.	.	.	.
52.0	PHOTIS SP.	2.0	1.5	2.4	0.8	0.8	0.2	0.2	.	.	.
52.0	SERPULIDAE	1.4	1.4	.	.	.	0.2	0.2	3.0	2.1	2.1
55.0	POLYCIARRUS EXIMIUS	1.6	0.5	1.4	0.4	0.4	0.4	0.2	1.0	0.4	0.4

Appendix 3. (Continued)

OVERALL RANK	SPECIES	STATION MS06											
		WINTER			SPRING			SUMMER			FALL		
		MEAN	ST	ERR	MEAN	ST	ERR	MEAN	ST	ERR	MEAN	ST	ERR
58.0	BOWMANIELLA PORTURICENSIS	0.8		0.5	1.8		1.1	1.2		0.4	0.4		0.2
58.0	GENOCIDARIS MACULATA	2.4		1.0	0.4		0.2	1.0		0.8	0.4		0.2
58.0	PISTA PALMATA	2.2		1.3	.		.	0.8		0.6	1.2		0.8
58.0	SALMACINA SP. A	.		.	.		.	.		.	4.2		4.2
58.0	XANTHIDAE A	0.4		0.4	0.6		0.2	.		.	3.2		1.5
62.5	ARCA IMBRICATA	.		.	0.2		0.2	3.2		2.0	0.6		0.6
62.5	HIATELLA ARCTICA	2.2		0.9	0.4		0.4	1.4		1.0	.		.
62.5	OPHIOSTIGMA SP. A	4.0		2.1	.		.	.		.	.		.
62.5	POLYDORA CAECA	2.8		1.7	0.4		0.4	0.4		0.4	0.4		0.4
66.0	ANCHIALINA TYPICA	.		.	0.8		0.4	2.4		1.5	0.6		0.6
66.0	SYLLIS ALTERNATA	0.4		0.4	1.0		1.0	1.0		0.8	1.4		0.9
66.0	TURBELLARIA	0.8		0.5	0.4		0.2	1.0		0.5	1.6		0.6
72.0	AMPELISCA VAOURUM	.		.	3.4		2.7	0.2		0.2	.		.
72.0	CHAMA CONGREGATA	1.6		1.2	0.2		0.2	1.6		1.6	0.2		0.2
72.0	LOIMIA MEDUSA	1.4		0.6	0.4		0.2	0.4		0.2	1.4		0.7
72.0	LYSIDICE NINETTA	1.0		1.0	0.4		0.2	1.4		0.4	0.8		0.6
72.0	MELITA APPENDICULATA	0.2		0.2	2.0		1.5	.		.	1.4		0.7
72.0	NASSARINA MINOR	1.2		0.8	.		.	1.2		0.5	1.2		1.2
72.0	PAGURUS SP.	1.4		0.7	0.2		0.2	1.6		0.9	0.4		0.4
72.0	SCHISTOMERINGOS RUDOLPHI	1.2		1.0	1.6		1.2	0.4		0.4	0.4		0.2
72.0	SYNALPHEUS TOWNSENDI	0.6		0.4	.		.	.		.	3.0		1.9
77.5	GONIADIDAE A	.		.	2.6		2.3	.		.	0.8		0.6
77.5	PARAPIONOSYLLIS SP.	0.4		0.4	.		.	.		.	3.0		3.0
82.5	AMPHARETE AMERICANA	1.0		0.8	0.2		0.2	1.0		0.5	1.0		0.3
82.5	CHEVALIA SP.	.		.	.		.	0.2		0.2	3.0		3.0
82.5	ERICHSONELLA FILIFORMIS	0.2		0.2	0.4		0.2	2.4		0.7	0.2		0.2
82.5	GAMMAROPSIS SP.	1.2		0.6	1.8		1.2	.		.	0.2		0.2
82.5	MUNIDA SP.	0.2		0.2	1.6		1.6	0.2		0.2	1.2		0.4
82.5	PERICLIMENES AMERICANUS	.		.	.		.	.		.	3.2		2.0
82.5	THOR MANNINGI	.		.	.		.	.		.	3.2		1.3
82.5	VARICORBULA OPERCULATA	1.6		1.4	0.4		0.4	1.2		0.6	.		.
89.0	HARMOTHOE SP. A	1.6		0.9	0.2		0.2	0.4		0.4	0.8		0.4
89.0	LEPTOCHELA PAPULATA	0.6		0.4	1.6		0.4	0.6		0.2	0.2		0.2
89.0	MITHRAX SP.	.		.	.		.	.		.	3.0		2.0
89.0	PARAPRIONOSPION PINNATA	1.2		0.6	0.8		0.4	1.0		1.0	.		.
89.0	PROTODORVILLEA KEFERSTEINI	2.8		1.6	0.2		0.2	.		.	.		.
94.5	AMPHIPODA G	0.6		0.6	0.8		0.6	0.2		0.2	1.2		1.2
94.5	ARABELLA MUTANS	.		.	1.4		1.0	0.8		0.2	0.6		0.4
94.5	GASTROCHAENA HANS	1.8		1.0	.		.	0.6		0.4	0.4		0.4
94.5	HYDROIDES SP. B	.		.	.		.	2.6		1.2	0.2		0.2
94.5	PAGURIDEA	.		.	.		.	0.8		0.5	2.0		0.5
94.5	SIPUNCULIDA A	2.2		1.0	.		.	0.4		0.4	0.2		0.2
100.0	BOTULA FUSCA	.		.	.		.	2.6		0.8	.		.
100.0	MELITIDAE A	.		.	0.6		0.4	.		.	2.0		1.0
100.0	ONUPHIS NEBULOSA	0.2		0.2	0.4		0.4	0.4		0.2	1.6		1.1
100.0	ONUPHIS PALLIDULA	.		.	2.6		0.9	.		.	.		.
100.0	SEMELE NUCULOIDES	0.6		0.6	1.0		0.3	0.8		0.8	0.2		0.2
106.0	CAPRELLA EQUILIBRA	2.4		1.7	.		.	.		.	.		.
106.0	CARPIAS BERMUDENSIS	0.6		0.4	1.8		1.1	.		.	.		.
106.0	CAULLERIELLA KILLARIENSIS	0.8		0.8	0.4		0.4	0.2		0.2	1.0		0.8
106.0	LITHOPHAGA BISULCATA	1.0		1.0	.		.	1.0		1.0	0.4		0.4
106.0	NASSARIUS ALBUS	1.0		0.3	0.4		0.2	1.0		0.8	.		.
106.0	ODONTOSYLLIS ENOPLA	0.8		0.4	.		.	0.6		0.2	1.0		0.8
106.0	PILUMNUS SP.	1.2		0.7	0.4		0.4	.		.	0.8		0.6
112.0	ASCORHYNCHUS CASTELLI	.		.	.		.	.		.	2.2		1.9

Appendix 3. (Continued)

STATION MS06

OVERALL RANK	SPECIES	WINTER		SPRING		SUMMER		FALL	
		MEAN	ST ERR	MEAN	ST ERR	MEAN	ST ERR	MEAN	ST ERR
112.0	DIASTYLIDAE A	0.8	0.4	0.8	0.6	0.2	0.2	0.4	0.4
112.0	HEMUS CRISTULIPES	1.0	0.6	0.2	0.2	.	.	1.0	0.8
112.0	PHERUSA INFLATA	1.8	0.9	.	.	0.2	0.2	0.2	0.2
112.0	POMATOCEROS AMERICANUS	0.8	0.5	.	.	1.4	0.9	.	.
122.5	AMPHIPODA F	0.4	0.4	1.6	1.6	.	.	.	.
122.5	ASPIDOSIPHON ALBUS	1.0	0.4	0.2	0.2	0.8	0.6	1.4	0.7
122.5	BRACHYURA	.	.	.	.	0.6	0.2	1.0	0.6
122.5	CERATONEREIS SP.	0.2	0.2	0.4	0.2	0.4	0.2	.	.
122.5	CHIONE LATILIRATA	0.4	0.4	0.2	0.2	1.4	0.4	0.4	0.2
122.5	GOULDIA CERINA	0.8	0.8	.	.	0.8	0.5	.	.
122.5	HOLOTHUROIDEA	2.0	1.5	.	.	.	.	.	.
122.5	KURTZIELLA LIMONITELLA	0.6	0.4	.	.	1.4	0.7	.	.
122.5	LUMBRINERIDES ACUTA	0.2	0.2	1.2	0.7	0.2	0.2	0.4	0.2
122.5	MALDANIDAE	0.2	0.2	0.8	0.6	0.8	0.4	0.2	0.2
122.5	MESOCOAETOPTERUS SP.	0.4	0.2	0.4	0.2	0.8	0.6	0.4	0.4
122.5	MUNIDIA PUSILLA	.	.	1.2	0.7	.	.	0.8	0.6
122.5	POLYDORA SP.	1.8	1.3	.	.	.	.	0.2	0.2
122.5	SPIO PETTIBONEAE	0.8	0.6	0.8	0.4	0.2	0.2	0.2	0.2
122.5	SYLLIS CORNUTA	1.2	0.4	0.4	0.4	.	.	0.4	0.2
122.5	XANTHIDAE B	.	.	0.2	0.2	0.8	0.5	1.0	0.4
135.0	AMPHITHOE SP. A	.	.	1.4	1.2	.	.	0.4	0.2
135.0	ARMANDIA MACULATA	1.0	0.6	0.4	0.4	.	.	0.4	0.2
135.0	EUPHKOSINE TRILOBA	0.6	0.6	.	.	.	.	1.2	0.7
135.0	MAJIDAE A	.	.	.	.	0.6	0.6	1.2	0.5
135.0	PAGURUS HENDERSONI	.	.	0.4	0.2	0.4	0.2	1.0	0.8
135.0	SCOLOPLOS SP. A	.	.	1.2	1.0	.	.	0.6	0.4
135.0	SYLLIDAE L	.	.	0.4	0.2	.	.	1.4	1.0
135.0	TEREBELLIDAE	0.4	0.4	.	.	.	.	1.4	1.0
135.0	TEREBELLIDAE C	0.2	0.2	.	.	.	.	1.6	1.4
146.0	AMPHARETE ACUTIFRONS	.	.	1.0	0.8	0.2	0.2	0.4	0.2
146.0	CARIDEA	0.2	0.2	0.4	0.4	.	.	1.0	0.4
146.0	CERATONEREIS MIRABILIS	0.2	0.2	0.2	0.2	0.2	0.2	1.0	0.0
146.0	ERICHTHONIUS BRASILIENSIS	0.2	0.2	0.6	0.2	0.4	0.2	0.4	0.2
146.0	GLYCERA CAPITATA	.	.	0.2	0.2	0.2	0.2	1.2	0.7
146.0	GLYCERA TESSELATA	0.6	0.4	.	.	0.6	0.4	0.4	0.4
146.0	MITRELLA LUNATA	1.4	1.0	0.2	0.2	.	.	.	.
146.0	NEPHTYS SQUAMOSA	0.8	0.4	0.4	0.2	.	.	0.4	0.2
146.0	PERICLIMENES SP.	.	.	0.6	0.6	.	.	1.0	0.5
146.0	PHYLLOCARIDA	0.4	0.4	.	.	0.2	0.2	1.0	1.0
146.0	PHYLLODOCIDAE	0.2	0.2	.	.	0.6	0.6	0.8	0.4
146.0	PRIONOSPIO CIRRIFERA	0.6	0.6	0.4	0.4	.	.	0.6	0.4
146.0	THYONE PSEUDOFUSUS	1.2	1.2	.	.	0.2	0.2	0.2	0.2
160.5	ALPHEUS NORMANNI	0.4	0.2	.	.	0.2	0.2	0.8	0.6
160.5	AMPELISCA CRISTOIDES	0.4	0.2	0.6	0.4	0.2	0.2	0.2	0.2
160.5	ANOPLODACTYLUS PETIOLATUS	0.4	0.2	0.2	0.2	0.8	0.6	.	.
160.5	ARCA ZEBRA	1.2	0.7	.	.	.	.	0.2	0.2
160.5	EUNICIDAE	.	.	1.0	0.8	0.2	0.2	0.2	0.2
160.5	HETEROCRYPTA GRANULATA	1.4	1.2	.	.	.	.	.	.
160.5	JAEROPSIS CORALICOLA	1.4	0.9	.	.	.	.	.	.
160.5	LEMOS UNICORNIS	0.6	0.4	0.4	0.4	0.4	0.4	.	.
160.5	LEUCOTHOE SP. A	0.2	0.2	.	.	0.2	0.2	1.0	0.6
160.5	LUMBRINERIOPSIS PARADUXA	.	.	1.2	0.8	.	.	0.2	0.2
160.5	LYONSIA SP. *	1.4	1.2	.	.	.	.	.	.
160.5	MARGINELLA HARTLEYANUM	0.2	0.2	1.0	0.8	0.2	0.2	.	.
160.5	NEMATONEREIS SP.	1.0	0.6	0.2	0.2	0.2	0.2	.	.

Appendix 3. (Continued)

OVERALL RANK	SPECIES	STATION		MS06										
		WINTER		SPRING		SUMMER			FALL					
		MEAN	ST ERR	MEAN	ST ERR	MEAN	ST	ERR	MEAN	ST	ERR	MEAN	ST	ERR
160.5	OPHIUROIDEA	1.2	0.4	.	.	.	.	.	0.2	.	.	0.2	.	0.2
160.5	SYLLIS FERRUGINA	.	.	.	.	.	.	.	1.4	.	.	1.0	.	1.0
160.5	TANYSTYLIDAE A	.	.	1.0	0.8	.	.	.	0.4	.	.	0.4	.	0.4
179.0	AONIDES MAYAGUEZENSIS	0.2	0.2	0.2	0.2	.	.	.	0.8	.	.	0.5	.	0.5
179.0	AXIOHELLA MUCOSA	0.4	0.4	0.4	0.2	.	.	.	0.4	.	.	0.2	.	0.2
179.0	CAPITELLIDAE	0.4	0.2	0.2	0.2	0.2	.	.	0.4	.	.	0.2	.	0.2
179.0	COLOMASTIX SP.	.	.	.	.	0.8	0.6	.	0.4	.	.	0.6	.	0.6
179.0	DIODORA CAYENENSIS	0.6	0.2	.	.	.	.	.	1.0	.	.	0.6	.	0.6
179.0	GALATHEA ROSTRATA	.	.	0.2	0.2	.	.	.	1.2	.	.	0.7	.	0.7
179.0	GONEPLACIDAE A	.	.	.	.	.	.	.	0.8	.	.	0.8	.	0.8
179.0	GREGARIELLA CORALLIOPHAGA	.	.	.	.	0.4	0.2	.	1.2	.	.	1.2	.	1.2
179.0	LYSIANASSIDAE A	.	.	.	.	.	.	.	.	.	.	.	.	.
179.0	MARGINELLA LAVALLEEANA	0.6	0.6	.	.	0.6	0.6	.	.	.	.	.	.	.
179.0	OSTRACODA	0.6	0.4	0.2	0.2	0.4	0.2	.	0.8	.	.	0.4	.	0.4
179.0	PELIA MUTICA	.	.	.	.	0.4	0.2	.	.	.	.	.	.	.
179.0	PLEUROMERIS TRIDENTATA	.	.	0.2	0.2	1.0	0.5	.	.	.	.	.	.	.
179.0	POLYDORA TETRABRANCHIA	1.2	0.8	.	.	.	.	.	.	.	.	.	.	.
179.0	PRIONOSPIO CRISTATA	0.2	0.2	0.6	0.4	.	.	.	0.4	.	.	0.2	.	0.2
179.0	PSEUDEURYTHOE AMBIGUA	.	.	0.6	0.4	0.4	0.2	.	0.2	.	.	0.2	.	0.2
179.0	PSEUDOMEDAEUS AGASSIZII	.	.	0.6	0.6	0.6	0.2	.	.	.	.	.	.	.
179.0	SICYONIA LAEVIGATA	0.4	0.4	.	.	0.8	0.4	.	0.6	.	.	0.6	.	0.6
179.0	SYLLIS GRACILIS	0.4	0.2	0.2	0.2	.	.	.	0.6	.	.	0.4	.	0.4
179.0	TRYPANOSYLLIS ZEBRA	0.4	0.2	.	.	0.2	0.2	.	0.6	.	.	0.2	.	0.2
179.0	WEBSTERINEREIS TRIDENTATA	0.6	0.4	0.4	0.2	.	.	.	0.2	.	.	0.4	.	0.4
200.5	ACANTHOHAUSTORIUS BOUSFIELDI	.	.	0.4	0.4	.	.	.	0.6	.	.	0.4	.	0.4
200.5	ALPHEUS SP.	.	.	.	.	0.2	0.2	.	0.8	.	.	0.4	.	0.4
200.5	ANCISTROSYLLIS RIGIDA	.	.	.	.	0.6	0.4	.	0.4	.	.	0.4	.	0.4
200.5	BATRACHONOTUS FRAGOSUS	.	.	0.2	0.2	0.6	0.4	.	0.2	.	.	0.2	.	0.2
200.5	BUDOTRIIDAE D	0.4	0.4	0.2	0.2	0.2	0.2	.	0.6	.	.	0.4	.	0.4
200.5	CHAETOZONE SETOSA	0.2	0.2	0.2	0.2	.	.	.	.	.	.	.	.	.
200.5	CUMACEA	0.6	0.4	0.4	0.4	.	.	.	1.0	.	.	0.4	.	0.4
200.5	DROMIDIA ANTILLENIS	.	.	.	.	.	.	.	.	.	.	.	.	.
200.5	LEPTOCHELIA SP.	0.8	0.4	.	.	0.2	0.2	.	.	.	.	.	.	.
200.5	LYONSIA SP.	.	.	.	.	1.0	0.3	.	0.6	.	.	0.4	.	0.4
200.5	MAGELONA SP. A	0.4	0.4	.	.	.	.	.	.	.	.	.	.	.
200.5	MODIOLUS AMERICANUS	0.2	0.2	.	.	0.8	0.4	.	0.6	.	.	0.6	.	0.6
200.5	MUSCULUS LATERALIS	0.2	0.2	.	.	0.2	0.2	.	0.4	.	.	0.2	.	0.2
200.5	NYMPHOPSIS DUODORSUSPINOSA	0.2	0.2	.	.	0.4	0.4	.	0.4	.	.	0.4	.	0.4
200.5	PHYLLODOCE LONGIPES	.	.	0.6	0.4	0.4	0.4	.	0.4	.	.	0.2	.	0.2
200.5	PODARKE OBSCURA	0.2	0.2	0.4	0.2	.	.	.	0.4	.	.	0.2	.	0.2
200.5	PROCERAEA SP.	0.2	0.2	0.2	0.2	0.4	0.2	.	0.2	.	.	0.2	.	0.2
200.5	SEROLIS MGRAYI	.	.	0.6	0.4	0.4	0.4	.	0.4	.	.	0.4	.	0.4
200.5	SICYONIA SP.	.	.	.	.	0.6	0.6	.	.	.	.	.	.	.
200.5	SPIONIDAE	1.0	0.5	.	.	.	.	.	.	.	.	.	.	.
200.5	SYNCHELIDIUM AMERICANUM	0.8	0.6	0.2	0.2	.	.	.	.	.	.	.	.	.
200.5	TANAIDACEA B	0.2	0.2	0.2	0.2	0.2	0.2	.	0.4	.	.	0.4	.	0.4
233.5	ACCALATHURA CRENULATA	.	.	0.4	0.2	0.4	0.2	.	.	.	.	.	.	.
233.5	ACTEON CANDENS	0.2	0.2	.	.	0.6	0.2	.	.	.	.	.	.	.
233.5	AMPHIPODA B	0.6	0.6	0.2	0.2	.	.	.	.	.	.	.	.	.
233.5	AMYGVALUM SAGITTATUM	.	.	0.8	0.8	.	.	.	.	.	.	.	.	.
233.5	ANAMIXIS SP.	.	.	.	.	.	.	.	0.8	.	.	0.8	.	0.8
233.5	ASTROPECTEN SP.	0.4	0.2	0.2	0.2	0.2	0.2	.	.	.	.	.	.	.
233.5	BARLEEIA SP. *	.	.	.	.	0.8	0.6	.	.	.	.	.	.	.
233.5	ENCOPE SP.	.	.	0.8	0.4	.	.	.	.	.	.	.	.	.
233.5	EUNICE FILAMENTOSA	.	.	.	.	0.4	0.2	.	0.4	.	.	0.2	.	0.2

Appendix 3. (Continued)

OVERALL RANK	SPECIES	STATION MS06									
		WINTER		SPRING		SUMMER		FALL			
		MEAN	ST ERR	MEAN	ST ERR	MEAN	ST ERR	MEAN	ST ERR		
233.5	EURYDICE LITTORALIS	.	.	0.4	0.2	0.2	0.2	0.2	0.2		
233.5	GLYCERA SP. B	0.2	0.2	0.6	0.2	.	.	.	.		
233.5	GONIADA MACULATA	.	.	0.8	0.8	.	.	.	.		
233.5	HYDROIDES SP. A	0.2	0.2	.	.	0.6	0.4	.	.		
233.5	LATREUTES PARVULUS	0.6	0.4	0.2	0.2	.	.	.	.		
233.5	LIMA PELLUCIDA	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2		
233.5	LITHOPHAGA ARISTATA	0.4	0.4	.	.	0.2	0.2	0.2	0.2		
233.5	LUCAPINELLA LIMATULA	0.8	0.6	.	.	.	.	.	.		
233.5	LYONIA BEANA	0.4	0.4	.	.	0.2	0.2	0.2	0.2		
233.5	MAERA CAROLINIANA	.	.	.	.	.	.	0.8	0.8		
233.5	MAJIDAE	.	.	0.2	0.2	0.2	0.2	0.4	0.4		
233.5	MELPHIDIPPIDAE A	0.2	0.2	.	.	.	.	0.6	0.6		
233.5	MICROPHYS ANTILLENIS	0.8	0.6	.	.	.	.	.	.		
233.5	MITHRAX PLEURACANTHUS	0.2	0.2	0.2	0.2	.	.	0.4	0.4		
233.5	MUSCULUS SP. *	0.8	0.8	.	.	.	.	.	.		
233.5	NEMATONEREIS SP. A	.	.	.	.	0.4	0.4	0.4	0.2		
233.5	NEMERTINEA E	0.2	0.2	.	.	.	.	0.6	0.4		
233.5	NEREIDAE	0.6	0.4	.	.	.	.	0.2	0.2		
233.5	NOTOMASTUS LATERICEUS	0.2	0.2	0.2	0.2	0.4	0.2	.	.		
233.5	OCNUS PYGMAEUS	0.6	0.6	.	.	0.2	0.2	.	.		
233.5	OLIGOCHAETA	0.4	0.4	0.2	0.2	.	.	0.2	0.2		
233.5	PALICUS SP.	0.6	0.4	0.2	0.2	.	.	.	.		
233.5	PELECYPODA A	0.4	0.2	0.4	0.4	.	.	.	.		
233.5	PELECYPODA B	0.6	0.4	0.2	0.2	.	.	.	.		
233.5	PENAEOIDEA	.	.	.	.	.	.	0.8	0.5		
233.5	PISONE REMOTA	.	.	0.6	0.4	.	.	0.2	0.2		
233.5	PODOCHELA SP.	.	.	0.2	0.2	0.6	0.6	.	.		
233.5	POLYDORA SP. D	0.8	0.8	.	.	.	.	.	.		
233.5	PSEUDOVERMILIA OCCIDENTALIS	.	.	.	.	0.6	0.6	0.2	0.2		
233.5	SCOLOPLOS RUBRA	0.6	0.4	.	.	.	.	0.2	0.2		
233.5	SIPUNCULIDA E	0.6	0.2	.	.	.	.	0.2	0.2		
233.5	SUBADYTE PELLUCIDA	0.2	0.2	0.2	0.2	0.4	0.2	.	.		
233.5	SYLLIS SP. G	0.8	0.6	.	.	.	.	.	.		
233.5	SYLLIS SP. I	.	.	.	.	.	.	0.8	0.6		
233.5	XANTHIDAE	.	.	.	.	0.2	0.2	0.6	0.4		
283.5	AMPHILOCHUS SP.	0.2	0.2	.	.	.	.	0.4	0.2		
283.5	AMPITHOIDAE A	0.2	0.2	0.4	0.4	.	.	.	.		
283.5	APSEUDES SP. B	0.4	0.4	0.2	0.2	.	.	.	.		
283.5	ASTACILLA SP. A	.	.	.	.	.	.	0.6	0.6		
283.5	AUTOMATE EVERMANNI	.	.	.	.	.	.	0.6	0.4		
283.5	BODOTRIIDAE B	0.4	0.2	0.2	0.2	.	.	.	.		
283.5	CAPITELLIDAE A	.	.	0.4	0.2	.	.	0.2	0.2		
283.5	CAPITELLIDAE D	.	.	.	.	.	.	0.6	0.6		
283.5	CERATONEREIS SP. A	0.2	0.2	.	.	.	.	0.4	0.2		
283.5	CHAETOPTERIDAE	.	.	.	.	0.6	0.4	.	.		
283.5	CHAMA MACEROPHYLLA	.	.	.	.	0.6	0.6	.	.		
283.5	CHLAMYS BENEDICTI	.	.	.	.	0.2	0.2	0.4	0.4		
283.5	CIRRATULIDAE	.	.	.	.	0.6	0.4	.	.		
283.5	CORBULA CONTRACTA	0.2	0.2	0.4	0.4	.	.	.	.		
283.5	COROPHIIDAE	0.4	0.4	.	.	.	.	0.2	0.2		
283.5	CRENELLA DECUSSATA	.	.	.	.	0.6	0.4	.	.		
283.5	DIASTYLIDAE	.	.	0.4	0.4	.	.	0.2	0.2		
283.5	DIPLODONTA NUCLEIFORMIS	0.6	0.6	.	.	.	.	.	.		
283.5	ERATO MAUGERIAE	0.2	0.2	.	.	0.4	0.4	.	.		
283.5	ETHUSA MASCARONE AMERICANA	.	.	.	.	.	.	0.6	0.6		

Appendix 3. (Continued)

		STATION MS06											
OVERALL RANK	SPECIES	WINTER			SPRING			SUMMER			FALL		
		MEAN	ST	ERR	MEAN	ST	ERR	MEAN	ST	ERR	MEAN	ST	ERR
283.5	GEUKENSIA DEMISSA	0.6		0.2	.		.	.		.	0.2		0.2
283.5	GONODACTYLUS BREDINI	0.4		0.4	.		.	.		.	0.6		0.6
283.5	HIPPOLYTIDAE	.		.	.		.	.		.	.		.
283.5	HIPPOMEDON SP. A	.		.	0.6		0.4	.		.	.		.
283.5	HYBOSCOLEX LONGISETA	0.4		0.2	.		.	0.2		0.2	.		.
283.5	LAONICE CIRRATA	0.4		0.2	.		.	.		.	0.2		0.2
283.5	LEMBOS SP.	.		.	.		.	.		.	0.6		0.6
283.5	LEPTOGNATHIIDAE A	0.2		0.2	.		.	0.2		0.2	0.2		0.2
283.5	LEUCOTHOIDES POTTSI	.		.	.		.	.		.	0.6		0.6
283.5	LYSIANOPSIS ALBA	.		.	0.2		0.2	.		.	0.4		0.2
283.5	MANGELIA RUGIRIMA	0.2		0.2	.		.	0.2		0.2	0.2		0.2
283.5	MARGINELLA ROSCIDA	.		.	0.4		0.2	0.2		0.2	.		.
283.5	MUREX FLORIFER DILECTUS	.		.	.		.	0.6		0.6	.		.
283.5	NEREIS SP.	.		.	.		.	0.2		0.2	0.4		0.2
283.5	NEREIS SUCCINEA	.		.	0.6		0.6	.		.	.		.
283.5	NOTOMASTUS AMERICANUS	.		.	0.6		0.6	.		.	.		.
283.5	OPHELIA DENTICULATA	.		.	.		.	.		.	0.6		0.4
283.5	PARAONIDAE	.		.	0.6		0.6	.		.	.		.
283.5	PELECYPODA C	0.2		0.2	.		.	0.4		0.4	.		.
283.5	PERICLIMENES IRIDESCENS	.		.	.		.	.		.	0.6		0.4
283.5	PHYLLODOCE SP.	.		.	.		.	0.4		0.2	0.2		0.2
283.5	PILUMNUS SAYI	0.2		0.2	.		.	0.2		0.2	0.2		0.2
283.5	PINNOTHERIDAE	.		.	0.2		0.2	0.2		0.2	0.2		0.2
283.5	POLYCIRRUS EXIMIUS DUBIUS	0.2		0.2	.		.	.		.	0.4		0.2
283.5	PONTUGENIA SERICOMA	0.4		0.2	.		.	0.2		0.2	.		.
283.5	PORTUNUS SP.	0.2		0.2	0.2		0.2	.		.	0.2		0.2
283.5	PROCESSA SP.	.		.	.		.	.		.	0.6		0.2
283.5	SEMELE PURPURASCENS	.		.	.		.	0.2		0.2	0.4		0.4
283.5	SIPUNCULIDA B	.		.	0.6		0.6	.		.	.		.
283.5	SPIO SP.	0.4		0.2	.		.	.		.	0.2		0.2
283.5	SYLLIDAE C	0.4		0.4	.		.	.		.	0.2		0.2
283.5	SYLLIDAE I	0.4		0.2	.		.	0.2		0.2	.		.
283.5	SYLLIS SP. F	0.4		0.2	0.2		0.2	.		.	.		.
283.5	TRACHYCARIS RESTRICTUS	.		.	.		.	.		.	0.6		0.4
283.5	VERTICORDIA ORNATA	0.4		0.4	0.2		0.2	.		.	.		.
283.5	WEBSTERINEREIS SP. A	.		.	.		.	.		.	0.6		0.6
353.5	ACANTHOCHITONA PYGMAEA	0.4		0.4	.		.	.		.	.		.
353.5	AEGIDAE A	.		.	0.2		0.2	0.2		0.2	.		.
353.5	ALPHEUS SP. B	.		.	.		.	.		.	0.4		0.2
353.5	AMPHIPODA E	0.2		0.2	.		.	0.2		0.2	.		.
353.5	ARBACIA PUNCTULATA	0.4		0.4	.		.	.		.	.		.
353.5	ARICIDEA CERRUTI	0.4		0.4	.		.	.		.	.		.
353.5	ASTHENOTHAERUS HEMPHILLI *	.		.	.		.	0.4		0.2	.		.
353.5	ATYS SANDERSONI	0.4		0.4	.		.	.		.	.		.
353.5	BODOTRIIDAE A	0.2		0.2	0.2		0.2	.		.	.		.
353.5	BRACHYCYTHARA BARBARAE LYONS	0.4		0.4	.		.	.		.	.		.
353.5	CADULUS QUADRIDENTATUS	.		.	.		.	0.4		0.2	.		.
353.5	CAPITELLA CAPITATA	0.2		0.2	0.2		0.2	.		.	.		.
353.5	CERADOCUS SP.	0.2		0.2	.		.	.		.	0.2		0.2
353.5	CERATONEREIS VERSIPEDATA	.		.	.		.	.		.	0.4		0.4
353.5	CHAETOPTERUS VARIOPEDATUS	0.4		0.4	.		.	.		.	.		.
353.5	CHIONE CANCELLATA	0.2		0.2	0.2		0.2	.		.	.		.
353.5	CYCLOPECTEN NANUS	.		.	0.4		0.2	.		.	.		.
353.5	DENTALIIDAE *	0.4		0.2	.		.	.		.	.		.
353.5	DIPLODONTA PUNCTATA	0.2		0.2	.		.	0.2		0.2	.		.

Appendix 3. (Continued)

OVERALL RANK	SPECIES	STATION MS06											
		WINTER			SPRING			SUMMER			FALL		
		MEAN	ST	ERR	MEAN	ST	ERR	MEAN	ST	ERR	MEAN	ST	ERR
353.5	DROMITIDAE	.	.	.	0.2	.	0.2	.	.	.	0.2	.	0.2
353.5	EBALIA CARIOSA	.	.	.	.	.	.	.	.	.	0.4	.	0.4
353.5	EBALIA STIMPSONI	.	.	.	.	.	.	.	.	.	0.4	.	0.4
353.5	ENGINA TURBINELLA	0.2	.	0.2	.	.	.	0.2	.	0.2	.	.	.
353.5	EUCIDARIS TRIBULOIDES	.	.	.	.	.	.	0.2	.	0.2	0.2	.	0.2
353.5	EUNICE ANTENNATA	0.4	.	0.4	.	.	.	.	.	.	.	.	.
353.5	GASTROPODA A	0.2	.	0.2	0.2	.	0.2	.	.	.	.	.	.
353.5	GLYCERA PAPILLOSA	0.2	.	0.2	.	.	.	.	.	.	0.2	.	0.2
353.5	HAPLOSCOLOPLOS ROBUSTUS	0.4	.	0.2	.	.	.	.	.	.	.	.	.
353.5	HESIONIDAE	0.2	.	0.2	.	.	.	.	.	.	0.2	.	0.2
353.5	HIPPOLYTIDAE *	0.4	.	0.4	.	.	.	.	.	.	.	.	.
353.5	HYALINA AVENA	.	.	.	0.4	.	0.4	.	.	.	.	.	.
353.5	HYDROIDES UNCINATA	0.4	.	0.4	.	.	.	.	.	.	.	.	.
353.5	INACHOIDES FORCEPS	0.2	.	0.2	0.2	.	0.2	.	.	.	.	.	.
353.5	INACHOIDES FORCEPS *	.	.	.	.	.	.	.	.	.	0.4	.	0.4
353.5	ISCHNOCHITONIDAE A	0.4	.	0.4	.	.	.	.	.	.	.	.	.
353.5	LUMBRINERIS SP.	0.4	.	0.4	.	.	.	.	.	.	.	.	.
353.5	LUMBRINERIS COCCINEA	0.4	.	0.4	.	.	.	.	.	.	.	.	.
353.5	LUMBRINERIS LATREILLI	.	.	.	0.2	.	0.2	.	.	.	0.2	.	0.2
353.5	LYONSTIA HYALINA	.	.	.	0.4	.	0.4	.	.	.	.	.	.
353.5	MAERA SP. A	0.2	.	0.2	.	.	.	0.2	.	0.2	.	.	.
353.5	MAERA WILLIAMSII	.	.	.	.	.	.	.	.	.	0.4	.	0.2
353.5	MICROPANOPE SCULPTIPES	.	.	.	.	.	.	.	.	.	0.4	.	0.2
353.5	MYSIDACEA	0.4	.	0.2	.	.	.	.	.	.	.	.	.
353.5	NANOPLAX XANTHIFORMIS	.	.	.	.	.	.	.	.	.	0.4	.	0.2
353.5	NEREIS ACUMINATA	0.4	.	0.2	.	.	.	.	.	.	.	.	.
353.5	NICOMACHE TRISPINATA	.	.	.	.	.	.	0.4	.	0.4	.	.	.
353.5	OPHIOCNIDA SCABRIUSCULA	0.4	.	0.4	.	.	.	.	.	.	.	.	.
353.5	OPHIOPHRAGMUS SP.	.	.	.	.	.	.	0.4	.	0.4	.	.	.
353.5	OPISTHOBRANCHIA *	0.4	.	0.2	.	.	.	.	.	.	.	.	.
353.5	PARAPINNIXA BOUVIERI	.	.	.	.	.	.	0.2	.	0.2	0.2	.	0.2
353.5	PARAPIONOSYLLIS LONGICIRRATA	0.4	.	0.4	.	.	.	.	.	.	.	.	.
353.5	PELECYPODA	.	.	.	0.4	.	0.2	.	.	.	.	.	.
353.5	PENAEIDAE	.	.	.	.	.	.	0.4	.	0.4	.	.	.
353.5	PERICLIMENAEUS SP.	.	.	.	.	.	.	.	.	.	0.4	.	0.2
353.5	PHERUSA EHLERSI	.	.	.	0.2	.	0.2	.	.	.	0.2	.	0.2
353.5	PISTA QUADRILOBATA	0.4	.	0.4	.	.	.	.	.	.	.	.	.
353.5	POECILOCHAETUS SP.	.	.	.	0.2	.	0.2	.	.	.	0.2	.	0.2
353.5	POTAMILLA RENIFORMIS	.	.	.	.	.	.	0.4	.	0.4	.	.	.
353.5	PROCESSA HEMPHILLI *	.	.	.	.	.	.	.	.	.	0.4	.	0.4
353.5	PSEUDOCHEMA RADIANUS	0.2	.	0.2	0.2	.	0.2	.	.	.	.	.	.
353.5	PSEUDOMEDEUS AGASSIZII *	.	.	.	.	.	.	0.4	.	0.4	.	.	.
353.5	PTEROMERIS PEKPLANA	.	.	.	.	.	.	0.4	.	0.2	.	.	.
353.5	PYRUNCULUS CAELATUS	.	.	.	0.4	.	0.4	.	.	.	.	.	.
353.5	RILDARDANUS LAMINOSA	.	.	.	.	.	.	0.2	.	0.2	0.2	.	0.2
353.5	SCOLELEPIS SP. B	.	.	.	.	.	.	.	.	.	0.4	.	0.2
353.5	SEMELE BELLASTRIATA	.	.	.	0.2	.	0.2	0.2	.	0.2	.	.	.
353.5	SPEOCARCINUS CAROLINENSIS	.	.	.	.	.	.	.	.	.	0.4	.	0.4
353.5	SPHAERODORIDIUM CLAPAREDI	0.4	.	0.4	.	.	.	.	.	.	.	.	.
353.5	SPIO SP. A	0.4	.	0.2	.	.	.	.	.	.	.	.	.
353.5	STHENELAIS BOA	.	.	.	.	.	.	0.2	.	0.2	0.2	.	0.2
353.5	STROMBIFORMIS BILINEATUS	.	.	.	.	.	.	0.4	.	0.2	.	.	.
353.5	SUTUROGLYPTA IONTHA	0.2	.	0.2	.	.	.	.	.	.	0.2	.	0.2
353.5	SYLLIDAE D	0.2	.	0.2	0.2	.	0.2	.	.	.	.	.	.
353.5	SYLLIDAE N	.	.	.	.	.	.	0.2	.	0.2	0.2	.	0.2

Appendix 3. (Continued)

OVERALL RANK	SPECIES	STATION MS06											
		WINTER			SPRING			SUMMER			FALL		
		MEAN	ST	ERR	MEAN	ST	ERR	MEAN	ST	ERR	MEAN	ST	ERR
353.5	SYNALPHEUS SP.	.	.	.	.	.	.	.	.	.	0.4	0.4	.
353.5	SYNELMIS ALBINI	0.2	.	0.2	.	.	.	.	.	.	0.2	0.2	.
353.5	TEREBELLIDAE A	.	.	.	0.2	.	0.2	0.2	.	0.2	.	.	.
353.5	THARYX MARIUNI	0.4	.	0.2	.	.	.	.	.	.	.	.	.
353.5	TIRON TROPAKIS	.	.	.	0.4	.	0.4	.	.	.	.	.	.
353.5	TRAVISIA PARVA	.	.	.	0.2	.	0.2	0.2	.	0.2	.	.	.
353.5	TRICHOPOXUS FLORIDANUS	0.2	.	0.2	.	.	.	0.2	.	0.2	.	.	.
353.5	TRIVIA MALTBIANA	.	.	.	.	.	.	.	.	.	0.4	0.2	.
353.5	TURBONILLA ABRUPTA	0.4	.	0.4	.	.	.	.	.	.	.	.	.
353.5	TURBONILLA INTERRUPTA	.	.	.	0.2	.	0.2	0.2	.	0.2	.	.	.
496.5	ALPHEIDAE	.	.	.	.	.	.	.	.	.	0.2	0.2	.
496.5	AMBLYOSYLLIS FORMOSA	.	.	.	.	.	.	0.2	.	0.2	.	.	.
496.5	AMERICARDIA MEDIA	.	.	.	.	.	.	0.2	.	0.2	.	.	.
496.5	AMPHILOCHIDAE	.	.	.	.	.	.	.	.	.	0.2	0.2	.
496.5	AMPHIURA FIBULATA	.	.	.	.	.	.	.	.	.	0.2	0.2	.
496.5	AMYGDALUM PAPHIUM	.	.	.	0.2	.	0.2	.	.	.	.	.	.
496.5	ANACHIS LAFRESNAYI	.	.	.	.	.	.	.	.	.	0.2	0.2	.
496.5	ANOMIA SIMPLEX	.	.	.	.	.	.	0.2	.	0.2	.	.	.
496.5	ANOPLODACTYLUS INSIGNIS	.	.	.	.	.	.	.	.	.	0.2	0.2	.
496.5	APANTHURA MAGNIFICA	.	.	.	.	.	.	0.2	.	0.2	.	.	.
496.5	ARCIDAE *	0.2	.	0.2	.	.	.	.	.	.	.	.	.
496.5	ARCOPSIS ADAMSI	.	.	.	.	.	.	.	.	.	0.2	0.2	.
496.5	ARENE TRICARINATA	0.2	.	0.2	.	.	.	.	.	.	.	.	.
496.5	ASTACILLA LAUFFI	.	.	.	.	.	.	0.2	.	0.2	.	.	.
496.5	ASTHENOTHAERUS HEMPHILLI	.	.	.	.	.	.	0.2	.	0.2	.	.	.
496.5	AUSTRAEOLIS CATINA	.	.	.	.	.	.	0.2	.	0.2	.	.	.
496.5	BARBATTIA DOMINGENSIS	0.2	.	0.2	.	.	.	.	.	.	.	.	.
496.5	BERGHIA SP.	.	.	.	.	.	.	.	.	.	0.2	0.2	.
496.5	BODOTRIIDAE	.	.	.	0.2	.	0.2	.	.	.	.	.	.
496.5	BODOTRIIDAE C	0.2	.	0.2	.	.	.	.	.	.	.	.	.
496.5	CADULUS TETRASCHISTUS	0.2	.	0.2	.	.	.	.	.	.	.	.	.
496.5	CAECUM COOPERI	0.2	.	0.2	.	.	.	.	.	.	.	.	.
496.5	CALOCARIS HIRSUTIMANA	.	.	.	.	.	.	.	.	.	0.2	0.2	.
496.5	CAPITELLIDAE B	.	.	.	.	.	.	.	.	.	0.2	0.2	.
496.5	CAPITELLIDAE E	.	.	.	0.2	.	0.2	.	.	.	.	.	.
496.5	CAPITELLIDAE G	.	.	.	.	.	.	.	.	.	0.2	0.2	.
496.5	CERAPUS TUBULARIS	.	.	.	0.2	.	0.2	.	.	.	.	.	.
496.5	CERITHIOPSIS METAXAE TAENIOLATA	0.2	.	0.2	.	.	.	.	.	.	.	.	.
496.5	CERITHIUM ATHATUM	.	.	.	.	.	.	0.2	.	0.2	.	.	.
496.5	CHAETOPLEURA APICULATA	.	.	.	.	.	.	.	.	.	0.2	0.2	.
496.5	CHAMA SP.	.	.	.	.	.	.	0.2	.	0.2	.	.	.
496.5	CHRYSOPETALIDAE	0.2	.	0.2	.	.	.	.	.	.	.	.	.
496.5	CIRROLANA PARVA	.	.	.	0.2	.	0.2	.	.	.	0.2	0.2	.
496.5	CIRRATULUS SP.	.	.	.	.	.	.	.	.	.	.	.	.
496.5	CIRRATULUS SP. A	0.2	.	0.2	.	.	.	.	.	.	.	.	.
496.5	CIRRIFORMIA FILIGERA	.	.	.	.	.	.	.	.	.	0.2	0.2	.
496.5	CLYPEASTER SP.	.	.	.	.	.	.	0.2	.	0.2	.	.	.
496.5	COLOMASTIX SP. B	.	.	.	0.2	.	0.2	.	.	.	.	.	.
496.5	CRENELLA DIVARICATA	.	.	.	0.2	.	0.2	.	.	.	.	.	.
496.5	CREPIDULA ACULEATA	.	.	.	.	.	.	.	.	.	0.2	0.2	.
496.5	CRONIUS SP.	0.2	.	0.2	.	.	.	.	.	.	.	.	.
496.5	CRONIUS RUBER	.	.	.	0.2	.	0.2	.	.	.	.	.	.
496.5	CUMINGIA TELLINOIDES	.	.	.	0.2	.	0.2	.	.	.	.	.	.
496.5	CYCLOPECTEN NANUS *	0.2	.	0.2	.	.	.	.	.	.	.	.	.
496.5	DELECTOPECTEN SP. *	0.2	.	0.2	.	.	.	.	.	.	.	.	.

Appendix 3. (Continued)

OVERALL RANK	SPECIES	STATION MS06							
		WINTER		SPRING		SUMMER		FALL	
		MEAN	ST ERR	MEAN	ST ERR	MEAN	ST ERR	MEAN	ST ERR
496.5	DENTALIUM OCCIDENTALE	.	.	.	.	.	.	0.2	0.2
496.5	DENTALIUM OCCIDENTALE *	0.2	0.2	.	.	.	.	.	.
496.5	DIODORA LISTERI	.	.	.	.	.	.	0.2	0.2
496.5	DIODORA LISTERI *	.	.	.	.	0.2	0.2	.	.
496.5	DIOPATRA TRIDENTATA	0.2	0.2	.	.	.	.	.	.
496.5	DONDICE OCCIDENTALIS	0.2	0.2	.	.	.	.	.	.
496.5	ELASMOPUS SP. D	.	.	0.2	0.2	.	.	.	.
496.5	EULALIA SP.	0.2	0.2	.	.	.	.	.	.
496.5	EULALIA MACROCEROS	.	.	0.2	0.2	.	.	.	.
496.5	EUNICE SP.	.	.	.	.	.	.	0.2	0.2
496.5	EUNICE SP. B	0.2	0.2	.	.	.	.	.	.
496.5	EURYDICE PIPERATA	.	.	.	.	0.2	0.2	.	.
496.5	FLABELLIGERA SP.	.	.	.	.	0.2	0.2	.	.
496.5	GASTROPODA E	0.2	0.2	.	.	.	.	.	.
496.5	GLOTTIDIA PYRAMIDATA	.	.	.	.	0.2	0.2	.	.
496.5	GLYCERA SP. C	0.2	0.2	.	.	.	.	.	.
496.5	GLYCYMERIS SP. *	0.2	0.2	.	.	.	.	.	.
496.5	GONIADIDES CAROLINAE	0.2	0.2	.	.	.	.	.	.
496.5	HARMOTHOE SP.	.	.	.	.	0.2	0.2	.	.
496.5	HESIONURA SP. A	.	.	.	.	.	.	0.2	0.2
496.5	HYALINA ALBOLINEATA	.	.	.	.	0.2	0.2	.	.
496.5	HYDROIDES SP. D	0.2	0.2	.	.	.	.	.	.
496.5	HYPsicOMUS PHAEOtaENIA	0.2	0.2	.	.	.	.	.	.
496.5	IRIDOPAGURUS DISPAR	.	.	.	.	0.2	0.2	.	.
496.5	KURTZIELLA ATROSTYLA	0.2	0.2	.	.	.	.	.	.
496.5	LASAEIDAE	.	.	.	.	.	.	0.2	0.2
496.5	LEIOCHRIDES PALLIDIOR	0.2	0.2	.	.	.	.	.	.
496.5	LEPTOCHELA SP.	.	.	.	.	.	.	0.2	0.2
496.5	LILJEBORGIA SP. A	.	.	0.2	0.2	.	.	.	.
496.5	LILJEBORGIA SP. B	.	.	.	.	.	.	0.2	0.2
496.5	LIMEA BRONNIANA	.	.	.	.	0.2	0.2	.	.
496.5	LINGA AMIANTUS	0.2	0.2	.	.	.	.	.	.
496.5	LISSA BICARINATA	.	.	.	.	0.2	0.2	.	.
496.5	LUCIFER FAXONI	.	.	.	.	0.2	0.2	.	.
496.5	LUCINA NASSULA	.	.	.	.	.	.	0.2	0.2
496.5	LYROPECTEN SP. *	.	.	0.2	0.2	.	.	.	.
496.5	LYSIANASSIDAE	0.2	0.2	.	.	.	.	.	.
496.5	LYSMATA SP. *	.	.	.	.	0.2	0.2	.	.
496.5	MACROCALLISTA MACULATA	.	.	0.2	0.2	.	.	0.2	0.2
496.5	MACROCOELOMA SP. *	.	.	.	.	.	.	.	.
496.5	MACROCOELOMA CAMPTOCERUM	.	.	.	.	0.2	0.2	.	.
496.5	MACROCOELOMA TRISPINOSUM	.	.	.	.	0.2	0.2	.	.
496.5	MARGINELLA SP.	.	.	.	.	0.2	0.2	.	.
496.5	MARGINELLA SP. 4	0.2	0.2	.	.	.	.	.	.
496.5	MARGINELLA AUREOCINCTA	.	.	.	.	0.2	0.2	.	.
496.5	MARGINELLA VIRGINIANA	0.2	0.2	.	.	.	.	.	.
496.5	MARGINELLIDAE *	0.2	0.2	.	.	.	.	.	.
496.5	MEGALOBRACHIUM SORIATUM	.	.	.	.	.	.	0.2	0.2
496.5	MESORHOEA SEXSPINOSA	.	.	0.2	0.2	.	.	.	.
496.5	MICRODEUTOPUS MYERSI	.	.	0.2	0.2	.	.	.	.
496.5	MICROPANOPE SP.	0.2	0.2	.	.	.	.	.	.
496.5	MICROPANOPE NUTTINGI	.	.	.	.	.	.	0.2	0.2
496.5	MICROPHRYS SP.	0.2	0.2	.	.	.	.	.	.
496.5	MICROPHRYS SP. *	.	.	.	.	0.2	0.2	.	.
496.5	MODIOLUS MODIOLUS SQUAMOSUS	0.2	0.2	.	.	.	.	.	.

Appendix 3. (Continued)

STATION MS06

OVERALL RANK	SPECIES	WINTER		SPRING		SUMMER		FALL	
		MEAN	ST ERR	MEAN	ST ERR	MEAN	ST ERR	MEAN	ST ERR
496.5	MUNIDA IRRASA *	0.2	0.2	.	.	.	.	.	.
496.5	MUSCULUS NIGER	.	.	.	.	0.2	0.2	.	.
496.5	NAINERIS SP.	0.2	0.2	.	.	.	.	.	.
496.5	NANNOSQUILLA SP. A	.	.	0.2	0.2	.	.	.	.
496.5	NATICA SP. A	.	.	0.2	0.2	.	.	.	.
496.5	NEMERTINEA C	.	.	.	.	0.2	0.2	.	.
496.5	NEOPONTONIDES BEAUFORTENSIS	.	.	.	.	0.2	0.2	.	.
496.5	NEPHTYS INCISA	0.2	0.2	.	.	.	.	.	.
496.5	NOTOBRYON WARDI	.	.	.	.	0.2	0.2	.	.
496.5	NOTOCIRRUS SPINIFERUS	0.2	0.2	.	.	.	.	.	.
496.5	NOTOMASTUS HEMIPODUS	.	.	.	.	.	.	0.2	0.2
496.5	NOTOMASTUS LOBATUS	.	.	.	.	.	.	0.2	0.2
496.5	NYMPHONIDAE A	.	.	.	.	.	.	0.2	0.2
496.5	NYMPHOPSIS SP. A	.	.	.	.	.	.	.	.
496.5	OLIVELLA BULLULA	.	.	.	.	0.2	0.2	.	.
496.5	ONUPHIS SP.	.	.	.	.	0.2	0.2	.	.
496.5	ONUPHIS EREMITA	.	.	0.2	0.2	.	.	.	.
496.5	OPALIA PUMILIO	0.2	0.2	.	.	.	.	.	.
496.5	OPHELIIDAE	.	.	0.2	0.2	.	.	.	.
496.5	OPHIOLEPIS ELEGANS	0.2	0.2	.	.	.	.	.	.
496.5	OPHIONEREIS RETICULATA	.	.	.	.	.	.	0.2	0.2
496.5	OPHIOSTIGMA SP.	.	.	.	.	.	.	0.2	0.2
496.5	OPHIUROIDEA A	0.2	0.2	.	.	.	.	.	.
496.5	ORBINIIDAE	0.2	0.2	.	.	.	.	.	.
496.5	ORBINIIDAE B	0.2	0.2	.	.	.	.	.	.
496.5	ORCHOMENE SP. A	.	.	.	.	0.2	0.2	.	.
496.5	OSACHILA TUBEROSA	.	.	.	.	.	.	0.2	0.2
496.5	OXYUROSTYLIS SMITHI	.	.	.	.	.	.	0.2	0.2
496.5	PAGURISTES TORTUGAE	.	.	.	.	.	.	0.2	0.2
496.5	PAGURUS HENDERSONI *	.	.	.	.	0.2	0.2	.	.
496.5	PANANTHURA FORMOSA	.	.	.	.	0.2	0.2	.	.
496.5	PAPYRIDEA SEMISULCATA	.	.	.	.	0.2	0.2	.	.
496.5	PAEULEPIS SP.	.	.	0.2	0.2	.	.	.	.
496.5	PARTHENOPE FRATERCULUS	.	.	.	.	0.2	0.2	.	.
496.5	PARVITURBOIDES INTERRUPTUS	0.2	0.2	.	.	.	.	.	.
496.5	PERICLIMENES SP. *	.	.	0.2	0.2	.	.	.	.
496.5	PHIDIANA LYNCEUS	0.2	0.2	.	.	.	.	.	.
496.5	PHORONIS ARCHITECTA	.	.	0.2	0.2	.	.	.	.
496.5	PHYTISICA MARINA	0.2	0.2	.	.	.	.	.	.
496.5	PHYLLODOCE ARENAE	.	.	.	.	.	.	0.2	0.2
496.5	PHYLLODOCE CASTANEA	.	.	.	.	0.2	0.2	.	.
496.5	PHYLLODOCIDAE C	.	.	0.2	0.2	.	.	.	.
496.5	PILUMNUS DASYPODUS	.	.	0.2	0.2	.	.	.	.
496.5	PINNIXA SP.	.	.	0.2	0.2	.	.	.	.
496.5	PINNOTHERES MACULATUS *	.	.	.	.	0.2	0.2	.	.
496.5	PISANIA TINCTA	.	.	0.2	0.2	.	.	.	.
496.5	PITHO LHERMINIERI	.	.	.	.	.	.	0.2	0.2
496.5	PPLICATULA GIBBOSA	.	.	.	.	.	.	0.2	0.2
496.5	PODOCHELA GRACILIPES	.	.	.	.	0.2	0.2	.	.
496.5	POLYCHAETA F	.	.	.	.	0.2	0.2	.	.
496.5	POLYDORA SP. C	0.2	0.2	.	.	.	.	.	.
496.5	POLYDORA SP. E	0.2	0.2	.	.	.	.	.	.
496.5	POLYNOIDAE A	.	.	.	.	.	.	0.2	0.2
496.5	POLYPLACOPHORA A	.	.	.	.	0.2	0.2	.	.
496.5	PORTUNIDAE	.	.	.	.	.	.	0.2	0.2

Appendix 3. (Continued)

		STATION MS06											
OVERALL RANK	SPECIES	WINTER			SPRING			SUMMER			FALL		
		MEAN	ST	ERR	MEAN	ST	ERR	MEAN	ST	ERR	MEAN	ST	ERR
496.5	PORTUNUS ORUWAYI	.	.	.	.	.	.	.	.	.	0.2	0.2	.
496.5	PORTUNUS ORUWAYI *	0.2	.	0.2	.	.	.	.	.	.	.	.	.
496.5	PSAMMOLYCE CTENIDOPHORA	.	.	.	0.2	0.2	.	.	.	.	.	.	.
496.5	PYLOPAGURUS SP. *	.	.	.	0.2	0.2	.	.	.	.	.	.	.
496.5	PYLOPAGURUS HOLTHUISI	.	.	.	.	.	.	.	.	.	0.2	0.2	.
496.5	RISSOINA SP. A	0.2	.	0.2	.	.	.	.	.	.	.	.	.
496.5	RUDILEMBOIDES NAGLEI	.	.	.	0.2	0.2	.	.	.	.	.	.	.
496.5	SABELLARIIDAE	.	.	.	.	.	.	.	.	.	0.2	0.2	.
496.5	SABELLIDAE	.	.	.	.	.	.	0.2	0.2	.	.	.	.
496.5	SALMONEUS ORTMANNI *	.	.	.	.	.	.	.	.	.	0.2	0.2	.
496.5	SCALIBREGMA INFLATUM	.	.	.	.	.	.	0.2	0.2	.	.	.	.
496.5	SIPUNCULIDA C	.	.	.	.	.	.	0.2	0.2	.	.	.	.
496.5	SPATANGOIDEA	0.2	.	0.2	.	.	.	.	.	.	.	.	.
496.5	SPELOEOPHORUS NODOSUS	.	.	.	.	.	.	.	.	.	0.2	0.2	.
496.5	SPIONIDAE B	.	.	.	0.2	0.2	.	.	.	.	.	.	.
496.5	SPIROGLYPTUS IRREGULARIS	0.2	.	0.2	.	.	.	.	.	.	.	.	.
496.5	SQUILLA HEPTACANTHA	.	.	.	.	.	.	.	.	.	0.2	0.2	.
496.5	STENOCLIONOPS FURCATA COELATA	0.2	.	0.2	.	.	.	.	.	.	.	.	.
496.5	STENOPLEUSTES SP. A	0.2	.	0.2	.	.	.	.	.	.	.	.	.
496.5	STENORHYNCHUS SETICORNIS	0.2	.	0.2	.	.	.	.	.	.	.	.	.
496.5	STENOTHOE SP. A	.	.	.	0.2	0.2	.	.	.	.	.	.	.
496.5	STHENELAIS SP.	.	.	.	0.2	0.2	.	.	.	.	.	.	.
496.5	SYLLIDAE K	0.2	.	0.2	.	.	.	.	.	.	.	.	.
496.5	SYLLIDES SP.	0.2	.	0.2	.	.	.	.	.	.	.	.	.
496.5	SYLLIS SP.	0.2	.	0.2	.	.	.	.	.	.	.	.	.
496.5	SYLLIS SP. B	.	.	.	0.2	0.2	.	.	.	.	.	.	.
496.5	SYLLIS SP. E	0.2	.	0.2	.	.	.	.	.	.	.	.	.
496.5	SYNALPHEUS MINUS	.	.	.	.	.	.	0.2	0.2	.	.	.	.
496.5	SYNAPTIDAE	.	.	.	0.2	0.2	.	.	.	.	.	.	.
496.5	TELLINA SP.	0.2	.	0.2	.	.	.	.	.	.	.	.	.
496.5	TELLINA SYBARITICA	.	.	.	.	.	.	0.2	0.2	.	.	.	.
496.5	THALENESSA SP. A	.	.	.	.	.	.	.	.	.	0.2	0.2	.
496.5	TRACHYPENAEUS CONSTRICTUS	0.2	.	0.2	.	.	.	.	.	.	.	.	.
496.5	TRICHOBRANCHUS GLACIALIS	0.2	.	0.2	.	.	.	.	.	.	.	.	.
496.5	TRICHOPOXUS EPISIOMUS	0.2	.	0.2	.	.	.	.	.	.	.	.	.
496.5	TRICOLIA THALLASSICOLA	.	.	.	.	.	.	0.2	0.2	.	.	.	.
496.5	TRIVIA CANDIOULA	.	.	.	.	.	.	.	.	.	0.2	0.2	.
496.5	TRIVIA NIX	.	.	.	.	.	.	0.2	0.2	.	.	.	.
496.5	TRYPTON CARNEUS	0.2	.	0.2	.	.	.	.	.	.	.	.	.
496.5	TURBONILLA HEMPHILLI	.	.	.	.	.	.	0.2	0.2	.	.	.	.
496.5	UHLIAS LIMBATUS	.	.	.	.	.	.	.	.	.	0.2	0.2	.
496.5	UNCIOLO SP.	0.2	.	0.2	.	.	.	.	.	.	.	.	.
496.5	UNCIOLO SPICATA	.	.	.	.	.	.	.	.	.	0.2	0.2	.
496.5	VERMILIOPSIS ANNULATA	.	.	.	.	.	.	0.2	0.2	.	.	.	.
496.5	VEXILLUM GEMMATUM	0.2	.	0.2	.	.	.	.	.	.	.	.	.
496.5	XANTHIDAE *	0.2	.	0.2	.	.	.	.	.	.	.	.	.
496.5	XANTHIDAE C	0.2	.	0.2	.	.	.	.	.	.	.	.	.

Appendix 3. (Continued)

OVERALL RANK	SPECIES	STATION 0501		WINTER		SPRING		SUMMER		FALL	
		MEAN	ST ERR	MEAN	ST ERR	MEAN	ST ERR	MEAN	ST ERR	MEAN	ST ERR
1.0	PISTA PALMATA	20.3	20.3	248.4	128.8	14.0	9.8				
2.0	SPIOPHANES BOMBYX	17.7	16.2	48.8	18.9	11.6	7.4				
3.0	ERICHTHONIUS SP. A	4.0	2.1	20.0	15.5	11.2	4.7				
4.0	SYLLIS SPONGICOLA			7.6	7.6	8.0	8.0				
5.0	ONUPHIS PALLICOLA	2.3	2.3	20.0	10.4	0.6	0.6				
6.0	PHOTIS SP.	3.7	3.7	15.8	12.0						
7.0	OWENIA FUSIFORMIS	2.0	2.0	7.4	5.5	6.0	5.8				
8.0	PHOTIS LONGICAUDATA			9.6	4.1	3.0	1.7				
9.0	MESOCHAETOPTERUS SP.	1.0	1.0	10.2	5.5	0.6	0.4				
10.0	AMPHARETE ACUTIFRONS	1.0	1.0	1.4	0.9	6.2	4.1				
11.0	CRASSINELLA LUNULATA			7.8	4.3	1.2	1.0				
12.0	UNCIOLOA SP. A	0.3	0.3	7.8	1.5	1.0	0.6				
13.0	ONUPHIS NERULOSA	2.0	2.0	4.8	1.8	1.2	1.0				
14.0	BOWMANIELLA PORTORICENSIS	0.3	0.3	4.8	2.1	2.4	1.3				
15.0	AMPELISCA VADORUM			4.8	2.0	1.4	0.6				
16.0	AMPHIODIA PULCHELLA			4.6	1.3	1.2	1.0				
17.0	MUNIDA PUSILLA			6.2	6.2						
18.0	AMPHARETE AMERICANA	0.3	0.3	1.8	1.8	3.2	1.8				
19.0	NEMATODA			5.0	4.8						
20.0	AGLAOPHAMUS VERRILLI	2.7	2.7	0.8	0.4	2.2	2.2				
21.5	AXIOHELLA MUCOSA			4.0	1.6	0.6	0.6				
21.5	NEMERTINEA	1.3	0.7	2.2	1.0	1.6	1.0				
24.0	APSEUDES SP. B			3.4	1.2	0.8	0.6				
24.0	PARAPRIONOSPIO PINNATA	0.3	0.3	3.2	1.0	1.4	0.9				
24.0	PROTOHAUSTORIUS WIGLEYI					5.0	3.1				
26.0	RILDARDANUS LAMINOSA	2.7	2.7	0.8	0.6	0.6	0.4				
27.5	PHITISICA MARINA			4.4	2.7	0.2	0.2				
27.5	SPIONIDAE B	1.7	1.7	0.8	0.6	1.8	1.3				
29.0	EURYDICE PIPERATA			0.4	0.4	3.4	1.9				
30.0	PRIONOSPIO CRISTATA	0.3	0.3	2.6	2.6	0.6	0.6				
31.5	HARMOTHOE SP. A			2.0	1.0	0.8	0.6				
31.5	MICROPANOPE SP. *			2.2	2.2						
33.5	AMYGDALUM SAGITTATUM			3.4	1.2						
33.5	EULALIA SANGUINEA			2.6	1.4	0.6	0.4				
35.0	EUNICE VITTATA			1.6	0.6	0.6	0.6				
36.0	TRICHOPOXUS FLORIDANUS			1.6	0.9	0.8	0.5				
38.5	ACANTHOHAUSTORIUS MILLSI			1.4	1.2	1.4	1.2				
38.5	BATRACHONOTUS FRAGOSUS			1.2	0.6	1.2	1.0				
38.5	LAONICE CIRKATA	1.0	1.0	1.4	0.5	0.4	0.4				
38.5	STHENELAIS BOA	0.3	0.3	0.6	0.4	1.0	1.0				
42.0	ARMANDIA MACULATA										
42.0	EUCEPAMUS PRAELONGUS	2.0	1.5	1.4	0.7						
42.0	XANTHIDAE B					2.0	1.3				
44.0	GLYCERA CAPITATA			1.2	1.0	0.2	0.2				
45.5	AUTOMATE EVERMANNI			0.6	0.4	0.4	0.2				
45.5	LOIMIA MEDUSA	2.0	1.1	0.2	0.2	0.8	0.8				
50.0	AMPELISCA CRISTOIDES			0.4	0.2	1.4	0.7				
50.0	AMPELISCA VENETIENSIS			1.2	1.2	0.2	0.2				
50.0	GAMMAROPSIS SP.			1.4	0.9	0.6	0.4				
50.0	GLOTTIDIA PYRAMIDATA			1.0	0.6						
50.0	PHYLLODOCE LONGIPES			1.4	0.6	0.6	0.4				
50.0	PSAMMOLYCE CTENIDOPHORA			0.4	0.2	1.2	1.0				
50.0	THYONE PSEUDOFUSUS			1.6	1.6	0.4	0.4				
57.5	ASYCHIS CAROLINAE	1.7	1.7	0.8	0.8						
57.5	ERICHTHONIUS BRASILIENSIS			1.4	0.9	0.4	0.4				

Appendix 3. (Continued)

		STATION OS01											
OVERALL RANK	SPECIES	WINTER			SPRING			SUMMER			FALL		
		MEAN	ST	ERR	MEAN	ST	ERR	MEAN	ST	ERR	MEAN	ST	ERR
57.5	HOLOTHURIDEA	.	.	.	1.8	1.6	.	1.2	1.0	.	0.2	0.2	.
57.5	LEPTOCHELA PAPULATA	.	.	.	0.4	0.2	.	.	.	.	.	.	.
57.5	NEPHTYS SQUAMOSA	.	.	.	0.8	0.6	.	1.0	0.4	.	.	.	.
57.5	NOTOMASTUS SP.	0.3	0.3	.	0.2	0.2	.	0.8	0.6	.	0.6	0.6	.
57.5	PAGURIDEA	.	.	.	0.6	0.2	.	0.8	0.5	.	0.4	0.2	.
57.5	XANTHIDAE A	.	.	.	.	.	.	1.8	1.8	.	.	.	.
66.5	AXIOGNATHUS SQUAMATUS	0.7	0.7	.	.	.	.	1.0	1.0	.	0.2	0.2	.
66.5	BRACHYURA	.	.	.	0.2	0.2	.	0.6	0.2	.	0.8	0.2	.
66.5	DIOPATRA CUPREA	.	.	.	1.4	0.5	.	0.2	0.2	.	.	.	.
66.5	GLYCERA SP. C	.	.	.	1.6	0.7	.	.	.	.	.	.	.
66.5	GONIADA MACULATA	1.0	1.0	.	0.6	0.4	.	.	.	.	0.4	0.4	.
66.5	GONIADIDAE A	.	.	.	1.0	1.0	.	.	.	.	0.6	0.6	.
66.5	GONIADIDES CAROLINAE	.	.	.	1.6	1.0	.	.	.	.	.	.	.
66.5	NOTOMASTUS AMERICANUS	.	.	.	0.2	0.2	.	.	.	.	1.4	0.9	.
66.5	OPHTHURIDEA	0.7	0.7	.	0.8	0.8	.	.	.	.	0.4	0.2	.
66.5	POMATOCEROS AMERICANUS	.	.	.	0.6	0.6	.	0.6	0.4	.	0.4	0.4	.
77.0	CHEVALIA SP.	.	.	.	.	.	.	1.4	1.2	.	.	.	.
77.0	CHONE AMERICANA	0.3	0.3	.	0.6	0.4	.	0.2	0.2	.	0.4	0.2	.
77.0	LUMBRINERIS COCCINEA	0.7	0.7	.	0.4	0.4	.	.	.	.	0.6	0.4	.
77.0	PAGURUS SP.	1.7	1.7	.	0.2	0.2	.	0.2	0.2	.	.	.	.
77.0	PARTHENOPE FRATERCULUS	.	.	.	1.4	1.2	.	.	.	.	.	.	.
77.0	PRIONOSPIO SP. B	0.3	0.3	.	1.0	0.5	.	.	.	.	0.2	0.2	.
77.0	PROCESSA SP.	.	.	.	1.4	0.4	.	.	.	.	.	.	.
77.0	PROTODORVILLEA KEFERSTEINI	.	.	.	1.4	0.9	.	.	.	.	.	.	.
77.0	STOMATOPODA A	.	.	.	1.4	0.4	.	.	.	.	.	.	.
77.0	TELLINA SYBARITICA	.	.	.	0.4	0.4	.	0.4	0.4	.	0.6	0.4	.
77.0	THARYX MARIONI	.	.	.	0.8	0.6	.	0.6	0.6	.	.	.	.
89.5	ASPIDOSIPHON GOSNOLDI	.	.	.	0.8	0.4	.	0.4	0.4	.	.	.	.
89.5	ASTROPECTEN SP.	.	.	.	1.2	0.6	.	.	.	.	.	.	.
89.5	CALOCARIS HIRSUTIMANA	.	.	.	1.2	0.7	.	.	.	.	.	.	.
89.5	CHIONE GRUS	0.7	0.3	.	0.4	0.2	.	0.2	0.2	.	0.2	0.2	.
89.5	CLYTHROCERUS PERPUSILLUS	0.7	0.7	.	0.2	0.2	.	0.2	0.2	.	0.4	0.4	.
89.5	GLYCERA SP. B	.	.	.	1.0	0.5	.	0.2	0.2	.	.	.	.
89.5	LEMROS UNICORNIS	.	.	.	1.0	0.5	.	0.2	0.2	.	.	.	.
89.5	MALDANIDAE	.	.	.	1.0	1.0	.	.	.	.	0.2	0.2	.
89.5	NASSARIUS ALBUS	.	.	.	0.6	0.4	.	0.4	0.2	.	0.2	0.2	.
89.5	NOTOPYGOS CRINITA	.	.	.	1.2	1.0	.	.	.	.	.	.	.
89.5	OLIGOCHAETA	.	.	.	0.6	0.6	.	.	.	.	0.6	0.4	.
89.5	PAGURUS HENDERSONI	1.7	1.7	.	0.2	0.2	.	.	.	.	.	.	.
89.5	PELECYPODA	.	.	.	1.0	0.5	.	0.2	0.2	.	.	.	.
89.5	TEREBELLIDAE F	.	.	.	.	.	.	0.4	0.4	.	0.8	0.8	.
103.5	CAPITELLIDAE	.	.	.	0.4	0.2	.	.	.	.	0.6	0.6	.
103.5	CAULLERIELLA KILLARIENSIS	.	.	.	0.4	0.2	.	0.4	0.4	.	0.2	0.2	.
103.5	CHAETOPTERUS VARIOPEDATUS	.	.	.	0.8	0.5	.	0.2	0.2	.	.	.	.
103.5	IDUNELLA SP.	.	.	.	.	.	.	0.6	0.4	.	0.4	0.4	.
103.5	LUCIFER FAXONI	.	.	.	0.2	0.2	.	0.2	0.2	.	0.6	0.2	.
103.5	MARGINELLA ROSCIDA	0.3	0.3	.	0.6	0.2	.	0.2	0.2	.	.	.	.
103.5	NEOPONTONIDES BEAUFORTENSIS	.	.	.	.	.	.	1.0	1.0	.	.	.	.
103.5	PHYLLOCHAETOPTERUS SOCIALIS	0.3	0.3	.	0.4	0.4	.	0.4	0.2	.	.	.	.
103.5	POLYDORA CAECA	.	.	.	0.4	0.2	.	0.6	0.2	.	.	.	.
103.5	PSEUDOMEDEAUS AGASSIZII	1.7	1.2	.	.	.	.	.	.	.	.	.	.
103.5	SPIOPHANES SP. A	.	.	.	1.0	1.0	.	.	.	.	.	.	.
103.5	SYLLIS HYALINA	.	.	.	0.6	0.6	.	0.4	0.2	.	.	.	.
103.5	THOR SP.	.	.	.	0.4	0.4	.	0.6	0.4	.	.	.	.
103.5	TURBELLARIA	.	.	.	1.0	0.3	.	.	.	.	.	.	.

Appendix 3. (Continued)

OVERALL RANK	SPECIES	STATION 0501															
		WINTER		SPRING		SUMMER		FALL									
		MEAN	ST ERR	MEAN	ST ERR	MEAN	ST ERR	MEAN	ST ERR								
118.5	ACTEON CANDENS	0.3	0.3	0.6	0.2	.	.	.	.	.	.	.	.	.	.	.	.
118.5	AONIDES MAYAGUEZENSIS	.	.	0.2	0.2	.	.	0.6	0.6	.	.	0.6	0.6	.	.	.	.
118.5	DELECTOPECTEN SP.	.	.	0.6	0.6	.	.	0.2	0.2	.	.	0.2	0.2	.	.	.	.
118.5	GLYCERA PAPILLOSA	0.7	0.7	0.4	0.4	.	.	.	.	.	.	.	.	.	.	.	.
118.5	GLYCERA SP. A	.	.	0.4	0.4	0.2	0.2	0.2	0.2	.	.	0.2	0.2	.	.	.	.
118.5	MACROCLYMENE ZONALIS	.	.	0.8	0.8	.	.	.	.	.	.	.	.	.	.	.	.
118.5	ODONTOSYLLIS ENOPLA	.	.	0.8	0.6	.	.	.	.	.	.	.	.	.	.	.	.
118.5	PALICUS SP.	0.3	0.3	0.6	0.2	.	.	.	.	.	.	.	.	.	.	.	.
118.5	PELECYPODA B	.	.	0.2	0.2	0.6	0.2	.	.	.	.	.	.	.	.	.	.
118.5	PODOCHELA GRACILIPES	0.7	0.7	.	.	0.4	0.2	.	.	.	.	.	.	.	.	.	.
118.5	POLYCIRRUS EXIMIUS	0.7	0.7	0.4	0.4	.	.	.	.	.	.	.	.	.	.	.	.
118.5	SABELLARIA VULGARIS VULGARIS	.	.	0.2	0.2	0.6	0.6	0.4	0.2	.	.	0.4	0.2	.	.	.	.
118.5	SEMELE BELLASTRIATA	.	.	.	.	0.4	0.2	.	.	.	.	0.2	0.2	.	.	.	.
118.5	SPIO PETTIBONEAE	.	.	0.6	0.4	.	.	.	.	.	.	0.6	0.2	.	.	.	.
118.5	SPIO SP. A	1.0	1.0	0.2	0.2	.	.	.	.	.	.	0.6	0.2	.	.	.	.
118.5	STOMATOPODA	.	.	0.2	0.2	0.6	0.6	.	.	.	.	.	.	.	.	.	.
142.5	AMPHINOMIDAE B	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
142.5	AMPHIPODA G	.	.	0.6	0.4	.	.	.	.	.	.	.	.	.	.	.	.
142.5	ANOPLODACTYLUS PETIOLATUS	.	.	0.6	0.4	.	.	.	.	.	.	.	.	.	.	.	.
142.5	CAPRELLA EQUILIBRA	.	.	0.6	0.6	.	.	.	.	.	.	.	.	.	.	.	.
142.5	GLYCERIDAE	.	.	.	.	0.4	0.4	0.2	0.2	.	.	0.2	0.2	.	.	.	.
142.5	GOULDIA CERINA	.	.	.	.	0.2	0.2	0.4	0.2	.	.	.	.	.	.	.	.
142.5	LILJEBORGIA SP. B	1.0	0.6	0.4	0.4	0.2	0.2	.	.	.	.	.	.	.	.	.	.
142.5	LIMA PELLUCIDA	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
142.5	LUMBRINERIS INFLATA	1.0	1.0	.	.	.	.	.	.	.	.	.	.	.	.	.	.
142.5	MEGALOBRACHIUM SORIATUM	.	.	0.6	0.6	.	.	.	.	.	.	.	.	.	.	.	.
142.5	MITRELLA LUNATA	.	.	0.6	0.2	.	.	.	.	.	.	.	.	.	.	.	.
142.5	NEPHTYS INCISA	.	.	0.6	0.2	.	.	.	.	.	.	.	.	.	.	.	.
142.5	NUCULA PROXIMA	.	.	0.6	0.4	.	.	.	.	.	.	.	.	.	.	.	.
142.5	ONUPHIS SP.	.	.	0.2	0.2	0.4	0.4	.	.	.	.	.	.	.	.	.	.
142.5	OSTRACODA	.	.	0.4	0.2	.	.	0.2	0.2	.	.	0.2	0.2	.	.	.	.
142.5	PAGURUS CAROLINENSIS	.	.	0.2	0.2	0.2	0.2	.	.	.	.	0.4	0.4	.	.	.	.
142.5	PALAEMONIDAE *	.	.	0.2	0.2	.	.	.	.	.	.	0.4	0.2	.	.	.	.
142.5	PARAONIDAE	.	.	0.2	0.2	.	.	.	.	.	.	.	.	.	.	.	.
142.5	PHILINE SAGHA	.	.	0.6	0.6	.	.	.	.	.	.	.	.	.	.	.	.
142.5	PHORONIS ARCHITECTA	.	.	0.6	0.4	.	.	.	.	.	.	.	.	.	.	.	.
142.5	PHYLLODOCE ARENAE	.	.	0.4	0.2	.	.	.	.	.	.	0.2	0.2	.	.	.	.
142.5	PHYLLODOCE SP.	.	.	0.4	0.2	.	.	.	.	.	.	0.2	0.2	.	.	.	.
142.5	PSEUDEURYTHOE AMBIGUA	.	.	.	.	.	.	.	.	.	.	0.6	0.4	.	.	.	.
142.5	PTILANTHURA TRICARINA	.	.	0.2	0.2	0.4	0.2	.	.	.	.	0.2	0.2	.	.	.	.
142.5	SCHISTOMERINGOS RUDOLPHI	.	.	0.4	0.4	.	.	.	.	.	.	.	.	.	.	.	.
142.5	SIPUNCULIDA	.	.	0.6	0.4	.	.	.	.	.	.	.	.	.	.	.	.
142.5	SIPUNCULIDA E	.	.	0.6	0.4	.	.	.	.	.	.	.	.	.	.	.	.
142.5	SPIONIDAE	0.3	0.3	.	.	.	.	.	.	.	.	0.4	0.4	.	.	.	.
142.5	SPIONIDAE A	.	.	0.6	0.4	.	.	.	.	.	.	.	.	.	.	.	.
142.5	SUBADYTE PELLUCIDA	0.3	0.3	0.4	0.4	0.4	0.2	.	.	.	.	.	.	.	.	.	.
142.5	TEREBELLIDAE A	0.3	0.3	.	.	.	.	.	.	.	.	.	.	.	.	.	.
142.5	TEREBELLIDAE H	.	.	.	.	.	.	.	.	.	.	0.6	0.6	.	.	.	.
189.5	ACCALATHURA CRENULATA	.	.	0.2	0.2	.	.	.	.	.	.	0.2	0.2	.	.	.	.
189.5	AMPHIODIA SP.	.	.	0.2	0.2	.	.	.	.	.	.	0.2	0.2	.	.	.	.
189.5	AMPHIPODA E	0.7	0.3	.	.	.	.	.	.	.	.	.	.	.	.	.	.
189.5	ARABELLA MUTANS	.	.	0.4	0.4	.	.	.	.	.	.	.	.	.	.	.	.
189.5	ARGOPECTEN GIBBUS	.	.	0.4	0.4	.	.	.	.	.	.	.	.	.	.	.	.
189.5	ARICIDEA FRAGILIS	.	.	0.2	0.2	.	.	0.2	0.2	.	.	0.2	0.2	.	.	.	.
189.5	ASPIDOSIPHON ALBUS	.	.	0.4	0.2	.	.	.	.	.	.	.	.	.	.	.	.

Appendix 3. (Continued)

		STATION 0501											
OVERALL RANK	SPECIES	WINTER			SPRING			SUMMER			FALL		
		MEAN	ST	ERR	MEAN	ST	ERR	MEAN	ST	ERR	MEAN	ST	ERR
189.5	CHRYSOPETALIDAE A	.	.	.	0.4	.	0.4	.	.	.	.	.	.
189.5	CHRYSOPETALIDAE B	.	.	.	0.4	.	0.4	.	.	.	.	.	.
189.5	CORBULA CONTRACTA	0.3	.	0.3	.	.	.	0.2	.	0.2	.	.	.
189.5	CRENELLA DIVARICATA	.	.	.	0.4	.	0.4	.	.	.	.	.	.
189.5	DAPHNELLA MORRA	.	.	.	0.4	.	0.4	.	.	.	.	.	.
189.5	EBALIA STIMPSONI	.	.	.	0.2	.	0.2	.	.	.	0.2	.	0.2
189.5	EPITONIUM NOVANGLIAE	.	.	.	0.4	.	0.4	.	.	.	.	.	.
189.5	ETHUSA MASCARONE AMERICANA	.	.	.	.	.	.	0.2	.	0.2	0.2	.	0.2
189.5	GASTROCHAENA HIANIS	0.7	.	0.7	.	.	.	0.2	.	0.2	.	.	.
189.5	GLYCERA TESSELATA	.	.	.	0.2	.	0.2	0.2	.	0.2	.	.	.
189.5	HIPPOMEDON SP. B	.	.	.	0.4	.	0.2	.	.	.	.	.	.
189.5	IRIDOPAGURUS DISPAR	0.3	.	0.3	.	.	.	.	.	.	0.2	.	0.2
189.5	LEMBOS SPINICARPUS INERMIS	.	.	.	.	.	.	0.4	.	0.4	.	.	.
189.5	LUMBRINERIDAE	.	.	.	0.4	.	0.4	.	.	.	.	.	.
189.5	LUMBRINERIDES ACUTA	.	.	.	0.4	.	0.2	.	.	.	.	.	.
189.5	LUMBRINERIDOPSIS PARADOXA	.	.	.	0.4	.	0.2	.	.	.	.	.	.
189.5	LYONSIA HYALINA	.	.	.	0.4	.	0.4	.	.	.	.	.	.
189.5	MAERA SP. A	0.7	.	0.7	.	.	.	.	.	.	.	.	.
189.5	MAGELONA SP. A	.	.	.	0.2	.	0.2	.	.	.	0.2	.	0.2
189.5	MANGELIA RUGIRIMA	0.3	.	0.3	.	.	.	0.2	.	0.2	.	.	.
189.5	MELITA APPENDICULATA	.	.	.	.	.	.	0.4	.	0.4	.	.	.
189.5	MICRODEUTOPUS MYERSI	0.7	.	0.7	.	.	.	.	.	.	.	.	.
189.5	MICROPANOPE PUSILLA *	.	.	.	0.4	.	0.4	.	.	.	.	.	.
189.5	MICROSPPIO PIGMENTATA	.	.	.	0.4	.	0.4	.	.	.	.	.	.
189.5	NAINERIS SP.	.	.	.	0.4	.	0.4	.	.	.	.	.	.
189.5	NEMERTINEA D	.	.	.	.	.	.	0.2	.	0.2	0.2	.	0.2
189.5	OCNUS PYGMAEUS	.	.	.	.	.	.	0.4	.	0.2	.	.	.
189.5	PAGURISTES TRIANGULATUS	.	.	.	.	.	.	0.4	.	0.4	.	.	.
189.5	PARAPHOXUS SPINOSUS	.	.	.	.	.	.	.	.	.	0.4	.	0.2
189.5	PECTINIDAE A	.	.	.	0.4	.	0.4	.	.	.	.	.	.
189.5	PHILINE QUADRATA	.	.	.	0.4	.	0.4	.	.	.	.	.	.
189.5	PHYLLODOCE SP. F	.	.	.	0.4	.	0.4	.	.	.	.	.	.
189.5	PHYLLODOCIDAE	0.3	.	0.3	0.2	.	0.2	.	.	.	.	.	.
189.5	PILUMNUS SP.	.	.	.	0.4	.	0.4	.	.	.	.	.	.
189.5	PISIONE REMOTA	.	.	.	0.2	.	0.2	.	.	.	0.2	.	0.2
189.5	PLEUROMERIS TRIDENTATA	.	.	.	0.4	.	0.4	.	.	.	.	.	.
189.5	PLICATULA GIBBOSA	.	.	.	0.4	.	0.4	.	.	.	.	.	.
189.5	PODOCHELA SP.	.	.	.	0.2	.	0.2	0.2	.	0.2	.	.	.
189.5	POECILOCHAETUS SP.	.	.	.	0.2	.	0.2	0.2	.	0.2	.	.	.
189.5	PRIONOSPPIO CIRRIFERA	.	.	.	0.2	.	0.2	0.2	.	0.2	.	.	.
189.5	PROCESSA GUYANAE	.	.	.	0.4	.	0.4	.	.	.	.	.	.
189.5	SCOLELEPIS TEXANA	.	.	.	0.4	.	0.4	.	.	.	.	.	.
189.5	SCOLOPLOS SP.	.	.	.	0.4	.	0.2	.	.	.	.	.	.
189.5	SICYONIA SP.	.	.	.	0.2	.	0.2	0.2	.	0.2	.	.	.
189.5	SIGALIONIDAE	.	.	.	.	.	.	.	.	.	0.4	.	0.2
189.5	SIPUNCULIDA B	.	.	.	.	.	.	0.4	.	0.4	.	.	.
189.5	SOLENOCERA SP.	.	.	.	0.4	.	0.4	.	.	.	.	.	.
189.5	STENOPLEUSTES SP. A	.	.	.	0.2	.	0.2	0.2	.	0.2	.	.	.
189.5	STROMBIFORMIS BILINEATUS	.	.	.	0.4	.	0.2	.	.	.	.	.	.
189.5	SYLLIDAE L	.	.	.	.	.	.	0.4	.	0.4	.	.	.
189.5	SYNCHELIDUM AMERICANUM	.	.	.	0.2	.	0.2	0.2	.	0.2	.	.	.
189.5	TEREBELLIDAE C	.	.	.	0.2	.	0.2	.	.	.	0.2	.	0.2
189.5	TEREBELLIDES STROEMI	.	.	.	.	.	.	0.4	.	0.4	.	.	.
189.5	THARYX SP.	.	.	.	.	.	.	.	.	.	0.4	.	0.2
189.5	TRICHOPHOXUS EPISTOMUS	.	.	.	0.2	.	0.2	0.2	.	0.2	.	.	.

Appendix 3. (Continued)

OVERALL RANK	SPECIES	STATION 0501											
		WINTER			SPRING			SUMMER			FALL		
		MEAN	ST	ERR	MEAN	ST	ERR	MEAN	ST	ERR	MEAN	ST	ERR
296.0	ALPHEUS SP. B	.	.	.	.	.	.	.	.	.	0.2	0.2	.
296.0	AMPELISCA SP. C	.	.	.	.	.	.	.	.	.	0.2	0.2	.
296.0	AMPELISCA VERRILLI	.	.	.	0.2	0.2	.	.	.	.	.	.	.
296.0	AMPHINOMIDAE A	.	.	.	0.2	0.2	.	.	.	.	.	.	.
296.0	AMPHIODIA TRYCHNA	.	.	.	0.2	0.2	.	.	.	.	.	.	.
296.0	ANOMURA	.	.	.	0.2	0.2	.	.	.	.	.	.	.
296.0	ARBACIA PUNCTULATA	0.3	0.3	.	.	.	.	.	.	.	.	.	.
296.0	ARGOPECTEN IRRADIANS	.	.	.	0.2	0.2	.	.	.	.	.	.	.
296.0	ARICIDEA SUECICA	.	.	.	0.2	0.2	.	.	.	.	.	.	.
296.0	ARMINA WATTLA	.	.	.	0.2	0.2	.	.	.	.	.	.	.
296.0	ASTERIAS FORBESII	.	.	.	0.2	0.2	.	.	.	.	.	.	.
296.0	ASTHENOTHAERUS HEMPHILLI	.	.	.	.	.	.	.	.	.	0.2	0.2	.
296.0	ASTROPECTEN COMPTUS	.	.	.	0.2	0.2	.	.	.	.	.	.	.
296.0	BODOTRIIDAE B	.	.	.	0.2	0.2	.	.	.	.	.	.	.
296.0	BRANCHIOSYLLIS EXILIS	0.3	0.3	.	.	.	.	.	.	.	0.2	0.2	.
296.0	CARIDEA	.	.	.	.	.	.	0.2	0.2	.	.	.	.
296.0	CAVOLINIA LONGIROSTRIS	.	.	.	.	.	.	0.2	0.2	.	.	.	.
296.0	CHAETOPTERIDAE	.	.	.	.	.	.	.	.	.	.	.	.
296.0	CHIONE LATILIRATA	.	.	.	0.2	0.2	.	.	.	.	.	.	.
296.0	CIRSOTREMA DALLI	.	.	.	0.2	0.2	.	.	.	.	.	.	.
296.0	COLLODES SP.	0.3	0.3	.	.	.	.	.	.	.	.	.	.
296.0	DENTALIUM OCCIDENTALE	.	.	.	.	.	.	.	.	.	0.2	0.2	.
296.0	DIACRIA QUADRIDENTATA	.	.	.	0.2	0.2	.	.	.	.	.	.	.
296.0	DIASTYLIDAE A	.	.	.	0.2	0.2	.	.	.	.	.	.	.
296.0	DIODORA CAYENENSIS *	0.3	0.3	.	.	.	.	.	.	.	.	.	.
296.0	DIPLOCIRRUS CAPENSIS	.	.	.	.	.	.	0.2	0.2	.	.	.	.
296.0	DONAX SP.	.	.	.	0.2	0.2	.	.	.	.	.	.	.
296.0	DROMIIDAE	.	.	.	.	.	.	.	.	.	0.2	0.2	.
296.0	EBALIA SP.	.	.	.	0.2	0.2	.	.	.	.	.	.	.
296.0	EBALIA SP. *	.	.	.	0.2	0.2	.	.	.	.	.	.	.
296.0	ECHINASTER SP.	.	.	.	0.2	0.2	.	.	.	.	.	.	.
296.0	ECHINOIDEA	.	.	.	0.2	0.2	.	.	.	.	.	.	.
296.0	ECHTURIDA	.	.	.	.	.	.	.	.	.	0.2	0.2	.
296.0	EPITONIUM KREBSII	.	.	.	.	.	.	.	.	.	0.2	0.2	.
296.0	EULALIA SP.	.	.	.	0.2	0.2	.	.	.	.	.	.	.
296.0	EUNICE ANTENNATA	0.3	0.3	.	.	.	.	.	.	.	.	.	.
296.0	EUNICE SP.	0.3	0.3	.	.	.	.	.	.	.	.	.	.
296.0	GALATHEA ROSTRATA	0.3	0.3	.	.	.	.	.	.	.	.	.	.
296.0	GONEPLAX SIGSBEI	.	.	.	0.2	0.2	.	.	.	.	.	.	.
296.0	GONIADA TERES	.	.	.	0.2	0.2	.	.	.	.	.	.	.
296.0	GONIADIDAE B	.	.	.	0.2	0.2	.	.	.	.	.	.	.
296.0	HIATELLA SP. A	.	.	.	0.2	0.2	.	.	.	.	.	.	.
296.0	HIPPOLYTE SP. *	0.3	0.3	.	.	.	.	.	.	.	.	.	.
296.0	HIPPOLYTIDAE	.	.	.	0.2	0.2	.	.	.	.	.	.	.
296.0	HYDROIDES PROTULICOLA	0.3	0.3	.	.	.	.	.	.	.	.	.	.
296.0	HYDROIDES UNCINATA	.	.	.	0.2	0.2	.	.	.	.	.	.	.
296.0	HYPOCONCHA ARCUATA	0.3	0.3	.	.	.	.	.	.	.	.	.	.
296.0	HYPOCONCHA SABULOSA	.	.	.	.	.	.	.	.	.	0.2	0.2	.
296.0	IRIDOPAGURUS DISPAR *	.	.	.	0.2	0.2	.	.	.	.	.	.	.
296.0	LAEVICARDIUM PICTUM	.	.	.	0.2	0.2	.	.	.	.	.	.	.
296.0	LEPTOCHELA BERMUDENSIS	.	.	.	0.2	0.2	.	.	.	.	.	.	.
296.0	LEPTOCHELA SP.	.	.	.	.	.	.	0.2	0.2	.	.	.	.
296.0	LEUCOTHOE SPINICARPA	.	.	.	.	.	.	.	.	.	0.2	0.2	.
296.0	LYONSIA SP.	.	.	.	.	.	.	0.2	0.2	.	.	.	.
296.0	LYSIDICE NINETTA	.	.	.	.	.	.	0.2	0.2	.	.	.	.

Appendix 3. (Continued)

		STATION 0501									
OVERALL RANK	SPECIES	WINTER		SPRING		SUMMER		FALL			
		MEAN	ST ERR	MEAN	ST ERR	MEAN	ST ERR	MEAN	ST ERR		
296.0	MAGFLONA SP. B	.	.	.	.	.	.	0.2	0.2		
296.0	MALACOCEROS GLUTAEUS	.	.	.	.	0.2	0.2	.	.		
296.0	MARGINELLA HARTLEYANUM	.	.	0.2	0.2	.	.	.	.		
296.0	MARGINELLA SP.	.	.	0.2	0.2	.	.	.	.		
296.0	MEDIOMASTUS CALIFORNIENSIS	.	.	.	.	0.2	0.2	.	.		
296.0	MEGALOMMA BIOCULATUM	.	.	0.2	0.2	.	.	.	.		
296.0	MICROPANOPE SP.	.	.	.	.	.	.	0.2	0.2		
296.0	MICROPANOPE URINATOR *	.	.	0.2	0.2	.	.	.	.		
296.0	MUNIDA IRRASA	.	.	.	.	0.2	0.2	.	.		
296.0	MUNIDA SP.	.	.	0.2	0.2	.	.	.	.		
296.0	MUREX FLORIFER DILECTUS	.	.	0.2	0.2	.	.	.	.		
296.0	MUSCULUS LATEHALIS	.	.	0.2	0.2	.	.	.	.		
296.0	NANNODIELLA MELANITICA	0.3	0.3	.	.	.	.	.	.		
296.0	NANNOSQUILLA SP. A	.	.	0.2	0.2	.	.	.	.		
296.0	NASSARINA MINOR	.	.	0.2	0.2	.	.	.	.		
296.0	NASSARIUS VIBEX	0.3	0.3	.	.	.	.	.	.		
296.0	NEPHTYIDAE	.	.	0.2	0.2	.	.	.	.		
296.0	NEREIDAE	.	.	.	.	.	.	0.2	0.2		
296.0	NEREIS ACUMINATA	.	.	0.2	0.2	.	.	.	.		
296.0	NEREIS SP.	.	.	.	.	.	.	0.2	0.2		
296.0	NOTOMASTUS HEMIPODUS	.	.	0.2	0.2	.	.	.	.		
296.0	ODOSTOMIA LAEVIGATA	.	.	0.2	0.2	.	.	.	.		
296.0	ONUPHIDAE	.	.	0.2	0.2	.	.	.	.		
296.0	ONUPHIS SP. A	.	.	.	.	.	.	0.2	0.2		
296.0	OPHIOCNIDA SP.	.	.	0.2	0.2	.	.	.	.		
296.0	OPHIOTHRIX ANGULATA	.	.	0.2	0.2	.	.	.	.		
296.0	PAGURIDAE	.	.	0.2	0.2	.	.	.	.		
296.0	PAGURUS HENDERSONI *	.	.	.	.	.	.	0.2	0.2		
296.0	PAGURUS PIERCEI	0.3	0.3	.	.	.	.	.	.		
296.0	PAGURUS SP. *	.	.	.	.	0.2	0.2	.	.		
296.0	PALICUS ALTERNATUS	.	.	.	.	0.2	0.2	.	.		
296.0	PALLIOLUM RETICULUM	.	.	0.2	0.2	.	.	.	.		
296.0	PARANAITIS POLYNOIDES	.	.	.	.	0.2	0.2	.	.		
296.0	PARANAITIS SPECIOSA	.	.	0.2	0.2	.	.	.	.		
296.0	PARAPINNIXA BOUVIERI	.	.	.	.	0.2	0.2	.	.		
296.0	PARDALISCIDAE A	.	.	0.2	0.2	.	.	.	.		
296.0	PASIPHAIDAE	.	.	0.2	0.2	.	.	.	.		
296.0	PECTINIDAE	.	.	.	.	0.2	0.2	.	.		
296.0	PELECYPODA F	.	.	0.2	0.2	.	.	.	.		
296.0	PELIA MULICA	.	.	0.2	0.2	.	.	.	.		
296.0	PERICLIMENES AMERICANUS	.	.	.	.	.	.	0.2	0.2		
296.0	PERICLIMENES SP.	.	.	.	.	0.2	0.2	.	.		
296.0	PETROLISTHES GALATHINUS	.	.	0.2	0.2	.	.	.	.		
296.0	PHERUSA EHLESI	.	.	0.2	0.2	.	.	.	.		
296.0	PHYLLODOCE MADEIRENSIS	.	.	.	.	0.2	0.2	.	.		
296.0	PHYLLODOCE PANAMENSIS	.	.	0.2	0.2	.	.	.	.		
296.0	PINNIXA SP.	.	.	0.2	0.2	.	.	.	.		
296.0	PINNOTHERIDAE	.	.	0.2	0.2	.	.	.	.		
296.0	PISTA QUADRILOBATA	0.3	0.3	.	.	.	.	.	.		
296.0	PLAKOSYLLIS QUADRIOCULATA	.	.	.	.	0.2	0.2	.	.		
296.0	PODARKE OBSCURA	.	.	.	.	.	.	0.2	0.2		
296.0	PODOCHELA RIISEI	.	.	0.2	0.2	.	.	.	.		
296.0	POLINICES UBERINUS	.	.	0.2	0.2	.	.	.	.		
296.0	POLYCHAETA E	.	.	.	.	0.2	0.2	.	.		
296.0	PORTUNUS SP.	.	.	0.2	0.2	.	.	.	.		

Appendix 3. (Continued)

		STATION 0501											
OVERALL RANK	SPECIES	WINTER			SPRING			SUMMER			FALL		
		MEAN	ST	ERR	MEAN	ST	ERR	MEAN	ST	ERR	MEAN	ST	ERR
296.0	PORTUNUS SPINICARPUS	.	.	.	.	.	.	0.2	.	0.2	.	.	.
296.0	PRIMNO LATREILLEI	.	.	.	0.2	.	0.2	.	.	.	.	.	.
296.0	PRIONOSPIO SP.	.	.	.	0.2	.	0.2	.	.	.	.	.	.
296.0	PYLOPAGURUS HOLTHUISI	.	.	.	.	.	.	.	.	.	0.2	.	0.2
296.0	PYLOPAGURUS SP.	.	.	.	0.2	.	0.2	.	.	.	.	.	.
296.0	PYRUNCULUS CAELATUS	.	.	.	0.2	.	0.2	.	.	.	.	.	.
296.0	RANINIDAE	.	.	.	0.2	.	0.2	.	.	.	.	.	.
296.0	SABELLIDAE	.	.	.	.	.	.	.	.	.	0.2	.	0.2
296.0	SCALIBREGMA INFLATUM	.	.	.	.	.	.	.	.	.	0.2	.	0.2
296.0	SCOLELEPIS SP. B	.	.	.	.	.	.	.	.	.	0.2	.	0.2
296.0	SCOLOPLOS SP. A	.	.	.	.	.	.	.	.	.	0.2	.	0.2
296.0	SEMELF NUCULOIDES	.	.	.	0.2	.	0.2	.	.	.	.	.	.
296.0	SERPULIDAE	0.3	.	0.3	.	.	.	.	.	.	.	.	.
296.0	SICYONIA BREVIROSTRIS *	0.3	.	0.3	.	.	.	.	.	.	.	.	.
296.0	SIGATICA CAROLINENSIS	0.3	.	0.3	.	.	.	.	.	.	.	.	.
296.0	SIPUNCULIDA A	.	.	.	.	.	.	0.2	.	0.2	.	.	.
296.0	SIPUNCULIDA C	.	.	.	0.2	.	0.2	.	.	.	.	.	.
296.0	SOLEMYA VELUM *	0.3	.	0.3	.	.	.	.	.	.	.	.	.
296.0	SOLENOCERA ATLANTIDIS	.	.	.	.	.	.	0.2	.	0.2	.	.	.
296.0	STENOCIONOPS FURCATA CUELATA	0.3	.	0.3	.	.	.	.	.	.	.	.	.
296.0	STENORHYNCHUS SETICORNIS	.	.	.	.	.	.	0.2	.	0.2	.	.	.
296.0	STHENELANELLA SP. A	.	.	.	.	.	.	.	.	.	0.2	.	0.2
296.0	SYLLIDAE	.	.	.	0.2	.	0.2	.	.	.	0.2	.	0.2
296.0	SYLLIS CORNUTA	.	.	.	.	.	.	.	.	.	.	.	.
296.0	SYNALPHEUS SP.	0.3	.	0.3	.	.	.	.	.	.	.	.	.
296.0	TANAIDACEA B	.	.	.	0.2	.	0.2	.	.	.	.	.	.
296.0	TELLINA SP.	.	.	.	.	.	.	0.2	.	0.2	.	.	.
296.0	TELLINA SP. *	.	.	.	0.2	.	0.2	.	.	.	.	.	.
296.0	TEREBELLIDAE	.	.	.	.	.	.	.	.	.	0.2	.	0.2
296.0	THOR FLORIDANUS	.	.	.	0.2	.	0.2	.	.	.	.	.	.
296.0	THOR MANNINGI	.	.	.	.	.	.	.	.	.	0.2	.	0.2
296.0	TRACHYCARDIUM EGMONTIANUM *	0.3	.	0.3	.	.	.	.	.	.	.	.	.
296.0	TRACHYCARIS RESTRICTUS	.	.	.	.	.	.	0.2	.	0.2	.	.	.
296.0	TRAVISIA PARVA	.	.	.	.	.	.	0.2	.	0.2	.	.	.
296.0	TRICHOBRANCHUS GLACIALIS	.	.	.	.	.	.	.	.	.	0.2	.	0.2
296.0	TURBONILLA ABRUPTA	.	.	.	0.2	.	0.2	.	.	.	.	.	.
296.0	VARICORBULA OPERCULATA	.	.	.	.	.	.	.	.	.	0.2	.	0.2
296.0	VOLVULELLA PAUPERCULA	.	.	.	0.2	.	0.2	.	.	.	.	.	.
296.0	VOLVULELLA PERSIMILIS	.	.	.	0.2	.	0.2	.	.	.	.	.	.
296.0	WEBSTERINEREIS SP.	.	.	.	.	.	.	.	.	.	0.2	.	0.2
296.0	XANTHIDAE	.	.	.	0.2	.	0.2	.	.	.	.	.	.

Appendix 3. (Continued)

OVERALL RANK	SPECIES	STATION 0506							
		WINTER		SPRING		SUMMER		FALL	
		MEAN	ST ERR	MEAN	ST ERR	MEAN	ST ERR	MEAN	ST ERR
1.0	FILOGRANA IMPLEXA	.	.	34.6	30.7	1.0	1.0	18.0	6.6
2.0	PLAKOSYLLIS QUADRIOCULATA	.	.	17.2	16.9	5.2	2.3	.	.
3.0	RILDARDANUS LAMINOSA	.	.	15.8	8.8	4.0	1.2	1.0	0.4
4.0	HESIONIDAE	.	.	13.4	13.4	0.8	0.6	.	.
5.0	PROTOMEDEIA SP. A	.	.	5.0	3.1	0.2	0.2	5.6	4.7
6.0	AMPHAPETE AMERICANA	.	.	5.0	4.0	4.4	3.5	1.2	0.6
7.5	SPHAEROSYLLIS SP.	.	.	9.0	7.8	.	.	0.4	0.4
7.5	SYLLIS SPONGICOLA	.	.	4.4	2.9	2.0	0.5	3.0	1.5
9.5	CHONE AMERICANA	.	.	2.8	1.5	4.6	2.6	0.6	0.2
9.5	SYLLIDAE L	.	.	3.6	3.3	4.0	2.3	0.4	0.2
11.0	SYLLIS HYALINA	.	.	3.8	2.6	1.2	0.7	2.0	0.8
12.5	COROPHIDIIDAE	.	.	6.6	6.6	.	.	.	.
12.5	GENOCIDARIS MACULATA	.	.	1.8	1.6	4.2	2.6	0.6	0.2
14.0	SERPULIDAE N	.	.	2.6	2.3	3.8	2.3	.	.
15.0	PISONE REMOTA	.	.	2.2	2.2	3.2	1.8	.	.
16.5	CHRYSOPETALIDAE A	.	.	1.8	1.3	3.4	1.8	.	.
16.5	NEMATODA	.	.	0.4	0.4	4.8	3.0	.	.
19.5	GLYCERA TESSELATA	.	.	2.2	1.7	1.4	0.7	1.0	0.6
19.5	MUNIDA SP.	.	.	2.4	2.1	1.6	1.1	0.6	0.4
19.5	OLIGOCHAETA	.	.	1.6	1.6	2.8	1.7	0.2	0.2
19.5	SPIOPHANES BUMBYX	.	.	4.4	2.9	.	.	0.2	0.2
23.0	GLYCERA PAPILLOSA	.	.	1.8	1.0	2.4	2.1	.	.
23.0	MELITIDAE A	.	.	2.0	2.0	0.8	0.5	1.4	0.9
23.0	SYLLIS ALTERNATA	.	.	2.0	1.1	1.4	0.9	0.8	0.6
25.5	MUNIDA PUSILLA	.	.	2.2	1.3	0.2	0.2	1.4	0.9
25.5	PRIONOSPIO CRISTATA	.	.	2.6	2.3	0.8	0.6	0.2	0.2
27.5	PHOTIS SP.	.	.	0.6	0.4	3.0	1.9	.	.
27.5	TANAIDACEA B	.	.	0.8	0.6	2.8	2.5	.	.
30.5	CAULLERIELLA KILLARIENSIS	.	.	2.0	1.8	1.0	0.3	0.4	0.4
30.5	EUNICE VITTATA	.	.	2.2	1.5	0.6	0.4	0.6	0.6
30.5	NOTOMASTUS LATERICEUS	.	.	.	.	3.0	1.8	0.4	0.4
30.5	SYLLIDAE	.	.	2.8	2.5	0.6	0.6	.	.
34.0	GLYCERIDAE	.	.	2.8	2.8	0.2	0.2	.	.
34.0	ODONTOSYLLIS ENOPLA	.	.	1.6	1.4	1.0	0.3	0.4	0.2
34.0	SYLLIS SP. D	.	.	1.8	0.8	0.6	0.4	0.6	0.6
36.0	GLYCERA SP. B	.	.	.	.	.	.	2.8	1.3
37.5	PROTODORVILLEA KEFERSTEINI	.	.	1.2	1.0	1.4	0.7	.	.
37.5	TURBELLARIA	.	.	2.4	1.9	.	.	0.2	0.2
39.5	EURYDICE PIPERATA	.	.	0.2	0.2	2.2	1.1	.	.
39.5	PISTA PALMATA	.	.	2.0	1.3	0.2	0.2	0.2	0.2
44.0	ERICHTHONIUS SP. A	.	.	2.2	1.2	.	.	.	.
44.0	GONIADIDES CAROLINAE	.	.	0.2	0.2	2.0	1.3	.	.
44.0	HESIONIDAE A	.	.	0.6	0.6	.	.	1.6	1.4
44.0	LANICE SP.	.	.	2.0	1.8	.	.	0.2	0.2
44.0	LEUCOTHOE SPINICARPA	.	.	0.6	0.4	0.6	0.6	1.0	1.0
44.0	SYLLIS FERRUGINA	.	.	1.4	0.9	0.2	0.2	0.6	0.4
44.0	THOR SP.	.	.	0.2	0.2	0.8	0.6	1.2	0.4
50.0	HARMOTHOE SP. A	.	.	1.0	1.0	1.0	0.4	.	.
50.0	PAGURIDEA	.	.	0.4	0.4	1.2	0.6	0.4	0.2
50.0	PERICLIMENES AMERICANUS	.	.	.	.	0.4	0.4	1.6	0.8
50.0	SPHAEROSYLLIS PIRIFERA	.	.	.	.	0.6	0.4	1.4	1.2
50.0	XANTHIDAE A	.	.	0.2	0.2	0.4	0.2	1.4	1.0
54.0	CERATONEREIS MIRABILIS	.	.	.	.	1.6	0.2	0.2	0.2
54.0	PAGURISTES SP.	.	.	1.6	0.7	0.2	0.2	.	.
54.0	SABELLIDAE	.	.	0.6	0.4	1.2	0.8	.	.

Appendix 3. (Continued)

STATION 0506

OVERALL RANK	SPECIES	WINTER		SPRING		SUMMER		FALL	
		MEAN	ST ERR	MEAN	ST ERR	MEAN	ST ERR	MEAN	ST ERR
60.0	AMPHIPODA G	.	.	.	.	1.6	1.4	.	.
60.0	INGOLFIELLA SP. A	.	.	1.4	1.4	0.2	0.2	.	.
60.0	LEPTOCHELIA SP.	.	.	1.0	0.4	0.4	0.2	0.2	0.2
60.0	NEMERTINEA	.	.	0.6	0.6	0.4	0.4	0.6	0.6
60.0	NOTOMASTUS SP.	.	.	0.8	0.6	0.6	0.4	0.2	0.2
60.0	PAGURIDAE	.	.	1.6	1.1	.	.	.	.
60.0	SEMELE PURPURASCENS	.	.	1.2	1.0	0.4	0.4	.	.
60.0	SPATANGOIDEA	.	.	.	.	1.6	0.9	.	.
60.0	XANTHIDAE	.	.	1.2	1.2	0.2	0.2	0.2	0.2
67.5	ARCOPSIS ADAMI	.	.	1.0	0.4	0.2	0.2	0.2	0.2
67.5	CRASSINELLA LUNULATA	.	.	1.2	1.2	.	.	0.2	0.2
67.5	OPHIOLEPIS ELEGANS	.	.	1.4	1.4	.	.	.	.
67.5	PAGURUS SP.	.	.	1.0	0.8	.	.	0.4	0.4
67.5	SERPULIDAE	.	.	0.4	0.2	0.4	0.4	0.6	0.4
67.5	SIPUNCULIDA	.	.	0.2	0.2	1.0	0.4	0.2	0.2
75.5	ASPIDOSIPHON GOSNOLDI	.	.	0.6	0.6	0.2	0.2	0.4	0.4
75.5	AUTOLYTUS SP.	.	.	0.2	0.2	0.2	0.2	0.8	0.6
75.5	GLYCERA CAPITATA	.	.	0.2	0.2	1.0	0.8	.	.
75.5	LEUCOTHOE SP. A	.	.	.	.	.	.	1.2	1.0
75.5	OPHIUROIDEA	.	.	0.8	0.8	0.2	0.2	0.2	0.2
75.5	PHYLLOCHAETOPTERUS SOCIALIS	.	.	.	.	1.0	0.6	0.2	0.2
75.5	PRIONOSPIO SP.	.	.	1.2	1.2	.	.	.	.
75.5	PYLOPAGURUS HOLTHUISI	.	.	0.4	0.4	0.6	0.6	0.2	0.2
75.5	SIPUNCULIDA B	.	.	1.2	0.8	.	.	.	.
75.5	SUBADYTE PELLUCIDA	.	.	0.4	0.2	0.6	0.4	0.2	0.2
86.0	ALPHEUS NORMANNI	.	.	.	.	0.2	0.2	0.8	0.5
86.0	AONIDES MAYAGUEZENSIS	.	.	.	.	0.6	0.6	0.4	0.2
86.0	ARBACIA PUNCTULATA	.	.	0.6	0.4	.	.	0.4	0.4
86.0	ELASMOPUS SP. A	.	.	.	.	1.0	0.5	.	.
86.0	LAONICE CIRRATA	.	.	0.2	0.2	0.6	0.2	0.2	0.2
86.0	LUCONACIA INCERTA	.	.	1.0	1.0	.	.	.	.
86.0	MICROPANOPE SCULPTIPES	.	.	0.6	0.6	0.2	0.2	0.2	0.2
86.0	PAGURISTES TORTUGAE	.	.	1.0	0.6	.	.	.	.
86.0	PHYLLOCARIDA	.	.	0.4	0.4	.	.	0.6	0.4
86.0	TEREBELLIDES STROEMI	.	.	0.8	0.4	0.2	0.2	.	.
86.0	XANTHIDAE B	.	.	.	.	1.0	0.6	.	.
104.5	ALPHEIDAE	.	.	0.8	0.8	.	.	.	.
104.5	AMPELISCA VADORUM	.	.	.	.	0.8	0.6	.	.
104.5	AMPHIPODA K	.	.	0.4	0.4	0.4	0.2	.	.
104.5	BRANCHIOSYLLIS EXILIS	.	.	0.8	0.5	.	.	.	.
104.5	CAPITELLIDAE A	.	.	0.6	0.4	0.2	0.2	.	.
104.5	ECHINOIDEA	.	.	0.8	0.5	.	.	.	.
104.5	GLANS DOMINGUENSIS	.	.	.	.	0.8	0.8	.	.
104.5	ISCHNOCHITONIDAE A	.	.	0.8	0.8	.	.	.	.
104.5	LEMBOS UNICORNIS	.	.	0.4	0.4	0.4	0.4	.	.
104.5	LEPTOCHELIA PAPULATA	.	.	.	.	0.8	0.6	.	.
104.5	MAJIDAE A	.	.	.	.	0.6	0.6	0.2	0.2
104.5	MARGINELLA HARTLEYANUM	.	.	0.6	0.6	0.2	0.2	.	.
104.5	NANOPLAX XANTHIFORMIS	.	.	0.2	0.2	.	.	0.6	0.6
104.5	NEPHTYS SQUAMOSA	.	.	.	.	0.6	0.2	0.2	0.2
104.5	ODONTOSYLLIS SP.	.	.	0.8	0.8	.	.	.	.
104.5	OWENIA FUSIFORMIS	.	.	0.6	0.4	0.2	0.2	.	.
104.5	PAGURUS HENDERSONI	.	.	0.4	0.4	0.2	0.2	0.2	0.2
104.5	PHYLLODOCIDAE	.	.	0.8	0.8	.	.	.	.
104.5	POLYDORA CAECA	.	.	0.4	0.2	0.4	0.4	.	.

Appendix 3. (Continued)

		STATION 0506							
OVERALL RANK	SPECIES	WINTER		SPRING		SUMMER		FALL	
		MEAN	ST ERR	MEAN	ST ERR	MEAN	ST ERR	MEAN	ST ERR
104.5	POLYDORA SP.	.	.	0.4	0.4	0.2	0.2	0.2	0.2
104.5	POMATOCEROS AMERICANUS	.	.	0.6	0.4	0.2	0.2	.	.
104.5	PRIONOSPIO CIRRIFERA	.	.	0.6	0.6	.	.	0.2	0.2
104.5	PRIONOSPIO SP. B	.	.	0.6	0.6	0.2	0.2	.	.
104.5	SCHISTOMERINGOS RUDOLPHI	.	.	0.8	0.8	.	.	.	.
104.5	SPIO PETTIBONEAE	.	.	0.4	0.4	0.4	0.4	.	.
104.5	WEBSTERINEREIS SP. B	.	.	0.8	0.5	.	.	.	.
132.5	AMPELISCA VENETIENSIS	.	.	0.2	0.2	0.2	0.2	0.2	0.2
132.5	AMPHIURA STIMPSONI	.	.	0.6	0.4	.	.	.	.
132.5	ARMANDIA MACULATA	.	.	0.2	0.2	0.4	0.4	.	.
132.5	AUTOMATE EVERMANNI	.	.	.	.	0.6	0.6	.	.
132.5	BOWMANIELLA PORTORICENSIS	.	.	.	.	0.6	0.6	.	.
132.5	BRACHYURA	.	.	0.2	0.2	0.4	0.2	.	.
132.5	CHEVALIA SP.	.	.	0.6	0.6	.	.	.	.
132.5	ERICHTHONIUS BRASILIENSIS	.	.	0.4	0.4	0.2	0.2	.	.
132.5	EULALIA SANGUINEA	.	.	0.4	0.2	0.2	0.2	.	.
132.5	GONIADIDAE A	.	.	0.2	0.2	0.4	0.4	.	.
132.5	IRIDOPAGURUS DISPAR	.	.	0.6	0.4	.	.	.	.
132.5	LEMBOS SPINICARPUS INERMIS	.	.	0.2	0.2	0.2	0.2	0.2	0.2
132.5	LOIMIA MEDUSA	.	.	0.4	0.2	0.2	0.2	.	.
132.5	LUCIFER FAXONI	.	.	0.2	0.2	0.2	0.2	0.4	0.2
132.5	LYSIDICE NINETTA	.	.	.	.	0.2	0.2	.	.
132.5	MARGINELLA ROSCIDA	.	.	.	.	0.2	0.2	0.2	0.2
132.5	OPHIOMUSIUM SP.	.	.	.	.	0.6	0.4	.	.
132.5	OPHIOSTIGMA ISACANTHUM	.	.	.	.	0.6	0.6	.	.
132.5	PALICUS SP.	.	.	0.4	0.2	0.2	0.2	.	.
132.5	PERICLIMENES SP. *	.	.	0.4	0.4	0.2	0.2	.	.
132.5	POECILOCHAETUS SP.	.	.	.	.	0.6	0.4	.	.
132.5	PROCESSA SP.	.	.	.	.	0.6	0.4	.	.
132.5	PHOTOMEDEIA SP.	.	.	.	.	0.6	0.6	.	.
132.5	PYLOPAGURUS SP. *	.	.	.	.	.	.	0.6	0.6
132.5	SYLLIS REGULATA CAROLINAE	.	.	.	.	0.2	0.2	0.4	0.2
132.5	SYNALPHEUS SP.	.	.	0.4	0.2	.	.	0.2	0.2
132.5	TEREBELLIDAE C	.	.	0.4	0.4	.	.	0.2	0.2
132.5	THYONE PSEUDOFUSUS	.	.	0.4	0.4	0.2	0.2	.	.
132.5	TRYPANOSYLLIS ZEBRA	.	.	0.2	0.2	.	.	0.4	0.4
132.5	UNCIOLA SP. A	.	.	.	.	0.6	0.2	.	.
173.5	ALPHEOPSIS SP. *	.	.	0.2	0.2	0.2	0.2	.	.
173.5	AMPHIPODA J	.	.	.	.	0.4	0.4	.	.
173.5	AMPHITHOE SP. A	.	.	.	.	0.4	0.4	.	.
173.5	AMPHITHOIDAE A	.	.	0.2	0.2	0.2	0.2	.	.
173.5	ANACHIS LAFRESNAYI	.	.	.	.	0.4	0.2	.	.
173.5	ANOPLODACTYLUS PETIOLATUS	.	.	0.4	0.4	.	.	.	.
173.5	ASTROPORPA ANNULATA	.	.	0.4	0.4	.	.	.	.
173.5	AXIOTHELLA MUCOSA	.	.	0.4	0.4	.	.	.	.
173.5	BRANCHIOSYLLIS OCULATA	.	.	.	.	.	.	0.4	0.4
173.5	CHLAMYS BENEDICTI	.	.	.	.	0.4	0.4	.	.
173.5	CORBULA CHITTYANA	.	.	.	.	0.4	0.4	.	.
173.5	EUNICE ANTENNATA	.	.	.	.	.	.	0.4	0.2
173.5	EUPHROSINE SP.	.	.	0.4	0.4	.	.	.	.
173.5	GALATHEA ROSTRATA	.	.	.	.	0.2	0.2	0.2	0.2
173.5	GASTROPODA K	.	.	0.4	0.4	.	.	.	.
173.5	LILJEBORGIA SP. A	.	.	.	.	0.4	0.2	.	.
173.5	LILJEBORGIA SP. B	.	.	0.4	0.4	.	.	.	.
173.5	LUMBRINERIS LATREILLI	.	.	.	.	0.4	0.2	.	.

Appendix 3. (Continued)

OVERALL RANK	SPECIES	STATION OS06							
		WINTER		SPRING		SUMMER		FALL	
		MEAN	ST ERR	MEAN	ST ERR	MEAN	ST ERR	MEAN	ST ERR
173.5	MAJIDAE	.	.	0.2	0.2	0.2	0.2	.	.
173.5	MELITA APPENDICULATA	.	.	0.4	0.4	.	.	.	.
173.5	MITHRAX ACUTICORNIS	.	.	.	.	0.2	0.2	0.2	0.2
173.5	NASSARINA MINOR	.	.	0.2	0.2	0.2	0.2	.	.
173.5	NASSARIUS ALBUS	.	.	.	.	0.2	0.2	0.2	0.2
173.5	NEREIS SUCCINEA	.	.	.	.	0.4	0.4	.	.
173.5	ONUPHIS PALLICULA	.	.	0.2	0.2	0.2	0.2	.	.
173.5	ONUPHIS SP.	.	.	0.2	0.2	.	.	0.2	0.2
173.5	OPHIACTIS SAVIGNYI	.	.	0.2	0.2	.	.	0.2	0.2
173.5	OPHIOCNIDA SP.	.	.	0.4	0.4	.	.	.	.
173.5	OPHIOTHRIX ANGULATA	.	.	0.2	0.2	0.2	0.2	.	.
173.5	OSACHILA TUBEROSA	.	.	.	.	0.4	0.4	.	.
173.5	PAGURUS CAROLINENSIS	.	.	0.4	0.2	.	.	.	.
173.5	PALICUS FAXONI	.	.	0.2	0.2	0.2	0.2	.	.
173.5	PARACERCEIS CAUDATA	.	.	0.2	0.2	.	.	0.2	0.2
173.5	PARTHENOPE FRATERCULUS	.	.	.	.	0.4	0.2	.	.
173.5	PELECYPODA G	.	.	.	.	0.4	0.4	.	.
173.5	PERICLIMENES IRIDESCENS	.	.	.	.	.	.	0.4	0.4
173.5	PILUMNUS SP.	.	.	0.4	0.4	.	.	.	.
173.5	PITAR SIMPSONI	.	.	.	.	0.4	0.2	.	.
173.5	POLYCIRRUS EXIMIUS	.	.	0.4	0.2	.	.	.	.
173.5	POLYPLACOPHORA A	.	.	.	.	0.4	0.4	.	.
173.5	RIMULA FRENULATA	.	.	0.4	0.4	.	.	.	.
173.5	SPIONIDAE	.	.	.	.	0.4	0.4	.	.
173.5	STHENELAIS BOA	.	.	.	.	0.4	0.4	.	.
173.5	STOMATOPODA	.	.	.	.	0.4	0.4	.	.
173.5	SYLLIS CORNUTA	.	.	.	.	.	.	0.4	0.4
173.5	TEREBELLIDAE	.	.	0.2	0.2	0.2	0.2	.	.
173.5	TETRATHYRUS SP.	.	.	.	.	0.4	0.4	.	.
173.5	THARYX MARIGNI	.	.	.	.	0.4	0.4	.	.
173.5	TRICHOBRANCHUS GLACIALIS	.	.	.	.	.	.	0.4	0.2
173.5	VERMILIOPSIS ANNULATA	.	.	0.2	0.2	0.2	0.2	.	.
173.5	WEBSTERINEREIS SP.	.	.	.	.	.	.	0.4	0.4
173.5	WEBSTERINEREIS TRIDENTATA	.	.	0.2	0.2	.	.	0.2	0.2
257.5	AGLAOPHAMUS VERRILLI	.	.	.	.	0.2	0.2	.	.
257.5	ALBUNEIDAE	.	.	.	.	0.2	0.2	.	.
257.5	ALPHEUS SP. B	.	.	.	.	.	.	0.2	0.2
257.5	AMPELISCA SCHELLENBERGI	.	.	0.2	0.2	.	.	.	.
257.5	AMPHINOMIDAE B	.	.	.	.	0.2	0.2	.	.
257.5	AMPHIPODA	.	.	.	.	0.2	0.2	.	.
257.5	AMPHIPODA B	.	.	.	.	0.2	0.2	.	.
257.5	AMYGDALUM SAGITTATUM	.	.	0.2	0.2	.	.	.	.
257.5	ANOPLODACTYLUS INSIGNIS	.	.	.	.	.	.	0.2	0.2
257.5	ARCA IMBRICATA	.	.	0.2	0.2	.	.	.	.
257.5	ASTERIAS FORBESII	.	.	0.2	0.2	.	.	.	.
257.5	AUTOMATE SP.	.	.	0.2	0.2	.	.	.	.
257.5	AXIOGNATHUS SQUAMATUS	.	.	.	.	0.2	0.2	.	.
257.5	CALLIOSTOMA PULCHRUM	.	.	0.2	0.2	.	.	.	.
257.5	CALOCARIS HIRSUTIMANA	.	.	.	.	.	.	0.2	0.2
257.5	CAPITELLA CAPITATA	.	.	.	.	.	.	0.2	0.2
257.5	CERADOCUS SP. B	.	.	.	.	0.2	0.2	.	.
257.5	CERATONEREIS SP.	.	.	0.2	0.2	.	.	.	.
257.5	CERATONEREIS SP. B	.	.	0.2	0.2	.	.	.	.
257.5	CERODRILLIA PERRYAE	.	.	.	.	.	.	0.2	0.2
257.5	CHAETOPTERUS VARIOPELATUS	.	.	0.2	0.2	.	.	.	.

Appendix 3. (Continued)

OVERALL RANK	SPECIES	STATION OS06							
		WINTER		SPRING		SUMMER		FALL	
		MEAN	ST ERR	MEAN	ST ERR	MEAN	ST ERR	MEAN	ST ERR
257.5	CHAETOZONE SETOSA	.	.	.	.	.	.	0.2	0.2
257.5	CHAMA CONGREGATA	.	.	0.2	0.2	.	.	.	.
257.5	CHIONE SP.	.	.	.	.	0.2	0.2	.	.
257.5	CLYTHROCERUS PERPUSILLUS	.	.	0.2	0.2	.	.	.	.
257.5	COLOMASTIX SP.	.	.	0.2	0.2	.	.	.	.
257.5	CRASSINELLA MARTINICENSIS	.	.	0.2	0.2	.	.	.	.
257.5	CRENELLA DECUSSATA	.	.	0.2	0.2	.	.	.	.
257.5	CREPIDULA ACULEATA	.	.	0.2	0.2	.	.	.	.
257.5	DARDANUS INSIGNIS	.	.	0.2	0.2	.	.	.	.
257.5	DENDROCHIROTIDA	.	.	.	.	0.2	0.2	.	.
257.5	DIPLOCIRKUS CAPENSIS	.	.	.	.	0.2	0.2	.	.
257.5	ERICHSONELLA FILIFORMIS	.	.	.	.	0.2	0.2	.	.
257.5	EULALIA SP.	.	.	.	.	0.2	0.2	.	.
257.5	EULALIA SP. A	.	.	.	.	0.2	0.2	.	.
257.5	EXOgone DISPAR	.	.	.	.	0.2	0.2	.	.
257.5	FAVARTIA CELLULOSA	.	.	.	.	0.2	0.2	.	.
257.5	GAMMAROPSIS SP.	.	.	.	.	.	.	0.2	0.2
257.5	GASTROCHAENA HIANIS	.	.	0.2	0.2	.	.	.	.
257.5	GONEPLACIDAE *	.	.	0.2	0.2	.	.	.	.
257.5	GONIADIDAE	.	.	0.2	0.2	.	.	.	.
257.5	GOULDIA CERINA	.	.	.	.	0.2	0.2	.	.
257.5	HETEROMYSIS FORMOSA	.	.	0.2	0.2	.	.	.	.
257.5	HIPPOMEDON SP. A	.	.	.	.	0.2	0.2	.	.
257.5	HOLOTHUROIDEA	.	.	.	.	0.2	0.2	.	.
257.5	HYDROIDES SP. D	.	.	.	.	0.2	0.2	.	.
257.5	HYPERIIDAE	.	.	.	.	0.2	0.2	.	.
257.5	HYPsicOMUS PHAEOTAENIA	.	.	.	.	0.2	0.2	.	.
257.5	LAEVICARDIUM PICTUM	.	.	0.2	0.2	.	.	.	.
257.5	LEPTOCHELA SP.	.	.	.	.	0.2	0.2	.	.
257.5	LIMEA SCABRA	.	.	0.2	0.2	.	.	.	.
257.5	LIMEA BRUNNIANA	.	.	.	.	0.2	0.2	.	.
257.5	LUMBRINERIDAE	.	.	.	.	0.2	0.2	.	.
257.5	LYTECHINUS VARIEGATUS	.	.	0.2	0.2	.	.	.	.
257.5	MACRURA	.	.	.	.	0.2	0.2	.	.
257.5	MAERA SP.	.	.	0.2	0.2	.	.	.	.
257.5	MAGELONA SP. B	.	.	.	.	0.2	0.2	.	.
257.5	MEIOSQUILLA QUADRIDENS	.	.	.	.	0.2	0.2	.	.
257.5	MELITIDAE	.	.	.	.	0.2	0.2	.	.
257.5	MESOCOAETOPTERUS SP.	.	.	0.2	0.2	.	.	.	.
257.5	MICROJASSA SP. A	.	.	0.2	0.2	.	.	.	.
257.5	MICROPANOPE SP.	.	.	.	.	.	.	0.2	0.2
257.5	MICROSPID PIGMENTATA	.	.	0.2	0.2	.	.	.	.
257.5	MITHRAX PLEURACANTHUS	.	.	0.2	0.2	.	.	.	.
257.5	NATICA CANRENA	.	.	.	.	0.2	0.2	.	.
257.5	NEODRILLIA CYDIA	.	.	.	.	0.2	0.2	.	.
257.5	OPHIOPHRAGMUS SP.	.	.	.	.	0.2	0.2	.	.
257.5	PAGURUS PIERCEI	.	.	0.2	0.2	.	.	.	.
257.5	PALEANOTUS SP. A	.	.	.	.	0.2	0.2	.	.
257.5	PARTHENOPE SP.	.	.	.	.	0.2	0.2	.	.
257.5	PELECYPODA A	.	.	.	.	0.2	0.2	.	.
257.5	PERICLIMENAEUS ASCIDIARUM	.	.	0.2	0.2	.	.	.	.
257.5	PERICLIMENAEUS ATLANTICUS	.	.	0.2	0.2	.	.	.	.
257.5	PERICLIMENES SP.	.	.	.	.	.	.	0.2	0.2
257.5	PHYSISICA MARINA	.	.	0.2	0.2	.	.	.	.
257.5	PHYLLODOCE CASTANEA	.	.	.	.	0.2	0.2	.	.

Appendix 3. (Continued)

		STATION 0506											
OVERALL RANK	SPECIES	WINTER			SPRING			SUMMER			FALL		
		MEAN	ST	ERR	MEAN	ST	ERR	MEAN	ST	ERR	MEAN	ST	ERR
257.5	PHYLLODOCE MUCOSA	.	.	.	0.2	.	0.2	.	.	.	.	.	.
257.5	PILUMNUS SAYI *	.	.	.	0.2	.	0.2	.	.	.	.	.	.
257.5	PINNA CARNEA	.	.	.	.	.	.	0.2	.	0.2	.	.	.
257.5	PINNIXA SP.	.	.	.	0.2	.	0.2	.	.	.	.	.	.
257.5	PINNOTHERIDAE	.	.	.	0.2	.	0.2	.	.	.	.	.	.
257.5	PODARKE OBSCURA	.	.	.	0.2	.	0.2	.	.	.	.	.	.
257.5	PODOCERUS SP.	.	.	.	0.2	.	0.2	.	.	.	.	.	.
257.5	PODOCHELA GRACILIPES	.	.	.	.	.	.	0.2	.	0.2	.	.	.
257.5	POLYCHAEIA G	.	.	.	.	.	.	0.2	.	0.2	.	.	.
257.5	POLYCIRRUS CAROLINENSIS	.	.	.	0.2	.	0.2	.	.	.	.	.	.
257.5	PRIONOSPIO CIRROBRANCHIATA	.	.	.	0.2	.	0.2	.	.	.	.	.	.
257.5	PROCERAEA SP.	.	.	.	.	.	.	.	.	.	0.2	.	0.2
257.5	PROCESSA PROFUNDA	.	.	.	.	.	.	.	.	.	0.2	.	0.2
257.5	PROTULA TUBULARIA	.	.	.	0.2	.	0.2	.	.	.	.	.	.
257.5	PSAMMOLYCE CTENIDOPHORA	.	.	.	0.2	.	0.2	.	.	.	.	.	.
257.5	PSEUDEURYTHOE AMBIGUA	.	.	.	0.2	.	0.2	.	.	.	.	.	.
257.5	PSEUDOMEDAUS AGASSIZII	.	.	.	.	.	.	0.2	.	0.2	.	.	.
257.5	PSEUDOVERMILIA SP.	.	.	.	0.2	.	0.2	.	.	.	.	.	.
257.5	PSEUDOVERMILIA OCCIDENTALIS	.	.	.	0.2	.	0.2	.	.	.	.	.	.
257.5	SABELLA MICROPHALMA	.	.	.	.	.	.	0.2	.	0.2	.	.	.
257.5	SCALIBREGMA INFLATUM	.	.	.	.	.	.	0.2	.	0.2	.	.	.
257.5	SEMELE SP.	.	.	.	.	.	.	0.2	.	0.2	.	.	.
257.5	SERPULA VERMICULARIS GRANULOSA	.	.	.	0.2	.	0.2	.	.	.	.	.	.
257.5	SICYONIA SP.	.	.	.	.	.	.	0.2	.	0.2	.	.	.
257.5	SOLASTER CARIBBAEUS	.	.	.	0.2	.	0.2	.	.	.	0.2	.	0.2
257.5	SPATIANGONIDAE	.	.	.	.	.	.	.	.	.	0.2	.	0.2
257.5	SPIONIDAE B	.	.	.	0.2	.	0.2	.	.	.	.	.	.
257.5	STHENELAIS SP.	.	.	.	.	.	.	0.2	.	0.2	.	.	.
257.5	SYLLIDAE D	.	.	.	0.2	.	0.2	.	.	.	.	.	.
257.5	SYLLIDAE E	.	.	.	0.2	.	0.2	.	.	.	.	.	.
257.5	SYLLIDAE I	.	.	.	0.2	.	0.2	.	.	.	.	.	.
257.5	SYLLIS GRACILIS	.	.	.	.	.	.	0.2	.	0.2	.	.	.
257.5	SYNALPHEUS LONGICARPUS	.	.	.	0.2	.	0.2	.	.	.	.	.	.
257.5	SYNALPHEUS MINUS	.	.	.	.	.	.	0.2	.	0.2	.	.	.
257.5	SYNALPHEUS SP. *	.	.	.	.	.	.	0.2	.	0.2	.	.	.
257.5	TANYSTYLIDAE A	.	.	.	0.2	.	0.2	.	.	.	.	.	.
257.5	TRACHYCARIS RESTRICTUS	.	.	.	.	.	.	0.2	.	0.2	.	.	.
257.5	TRITONIA BAYERI MISA	.	.	.	.	.	.	.	.	.	0.2	.	0.2
257.5	TRIVIA CANDIDULA	.	.	.	0.2	.	0.2	.	.	.	.	.	.
257.5	VITRINELLIDAE C	.	.	.	0.2	.	0.2	.	.	.	.	.	.