



NOAA Technical Memorandum NMFS-NE-310

**Northeast Regional Action Plan
to implement the NOAA Fisheries
Climate Science Strategy through 2024**

**US DEPARTMENT OF COMMERCE
National Oceanic and Atmospheric Administration
National Marine Fisheries Service
Northeast Fisheries Science Center
Woods Hole, Massachusetts
November 2023**



NOAA Technical Memorandum NMFS-NE-310

This series represents a secondary level of scientific publishing. All issues employ thorough internal scientific review; some issues employ external scientific review. Reviews are transparent collegial reviews, not anonymous peer reviews. All issues may be cited in formal scientific communications.

Northeast Regional Action Plan to implement the NOAA Fisheries Climate Science Strategy through 2024

Vincent Saba¹, Diane Borggaard², Joseph Caracappa³, Patricia Clay⁴, Lisa Colburn⁵, Mathias Collins⁶, Jennifer Cudney⁸, Jonathan Deroba³, Geret DePiper⁷, Paula Fratantoni³, Marianne Ferguson², Sean Hayes³, Kimberly Hyde⁵, Kirsten Jabanoski⁹, Michael Johnson², John Kocik¹⁰, Ellen Keane², Dan Kircheis², Scott Large³, Andrew Lipsky³, Sean Lucey³, Anna Mercer⁵, Shannon Meseck⁹, Timothy Miller³, Christopher Orphanides⁵, Julie Reichert-Nguyen¹¹, Ronald Vogel¹², Bruce Vogt¹¹, Gary Wikfors⁹

- ¹NOAA Fisheries Northeast Fisheries Science Center, Geophysical Fluid Dynamics Laboratory, Princeton University, 201 Forrestal Road, Princeton, NJ, 08540
- ²NOAA Fisheries, Greater Atlantic Regional Fisheries Office, 55 Great Republic Drive, Gloucester, MA, 01930
- ³NOAA Fisheries, Northeast Fisheries Science Center, 166 Water Street, Woods Hole, MA, 02543
- ⁴NOAA Fisheries, Northeast Fisheries Science Center, % F/ST5, 1315 East West Highway, Building SSMC3, Silver Spring, MD 20910
- ⁵NOAA Fisheries, Northeast Fisheries Science Center, 28 Tarzwell Drive, Narragansett, RI 02882
- ⁶NOAA Fisheries, Office of Habitat Conservation, Greater Atlantic Regional Fisheries Office, 55 Great Republic Drive, Gloucester, MA 01930
- ⁷NOAA Fisheries, Northeast Fisheries Science Center, 15 Carlson Lane, Falmouth, MA 02540
- ⁸NOAA Fisheries, Office of Sustainable Fisheries, Atlantic Highly Migratory Species Division, 1315 East-West Highway 13th Floor, Silver Spring, MD 20910
- ⁹NOAA Fisheries, Northeast Fisheries Science Center, 212 Rogers Avenue, Milford, CT 06460
- ¹⁰NOAA Fisheries, Northeast Fisheries Science Center, 17 Godfrey Drive, Orono, ME 04473
- ¹¹NOAA Fisheries, Office of Habitat Conservation, Chesapeake Bay Office, 200 Harry S Truman Parkway, Annapolis, MD 21401
- ¹²NOAA Fisheries, National Environmental Satellite Data and Information Service, 5830 University Research Ct., Building NCWCP, College Park, MD 20740

**US DEPARTMENT OF COMMERCE
National Oceanic and Atmospheric Administration
National Marine Fisheries Service
Northeast Fisheries Science Center
Woods Hole, Massachusetts
November 2023**

Editorial Notes

Information Quality Act Compliance: In accordance with section 515 of Public Law 106-554, the Northeast Fisheries Science Center (NEFSC) completed both technical and policy reviews for this report. These pre-dissemination reviews are on file at the NEFSC Editorial Office.

Species Names: The NEFSC Editorial Office's policy on the use of species names in all technical communications is generally to follow the American Fisheries Society's lists of scientific and common names for fishes, mollusks, and decapod crustaceans and to follow the Society for Marine Mammalogy's guidance on scientific and common names for marine mammals. Exceptions to this policy occur when there are subsequent compelling revisions in the classifications of species, resulting in changes in the names of species.

Statistical Terms: The NEFSC Editorial Office's policy on the use of statistical terms in all technical communications is generally to follow the International Standards Organization's handbook of statistical methods.

Citation: Saba V, Borggaard D, Caracappa J, Clay P, Colburn L, Collins M, Cudney J, Deroba J, DePiper G, Fratantoni P, et al. 2023. Northeast Regional Action Plan to implement the NOAA Fisheries Climate Science Strategy through 2024. US Dept Commer Northeast Fish Sci Cent Tech Memo 310. 39 p.

TABLE OF CONTENTS

EXECUTIVE SUMMARY.....	2
Build and Maintain Infrastructure and Track Change (NCSS Objectives 6-7).....	2
Understand Mechanisms and Project Future Conditions (NCSS Objectives 4-5).....	3
Inform Management (NCSS Objectives 1-3).....	3
1. Introduction.....	3
2. NERAP Accomplishments (FY16-FY21).....	4
2.1 Maintaining infrastructure and tracking change (NCSS Objectives 6-7)	4
2.2 Understanding mechanisms and projecting future conditions (NCSS Objectives 4-5)	6
2.3 Informing management (NCSS Objectives 1-3).....	7
3. Key Needs	9
3.1 Maintain infrastructure and track change (NCSS Objectives 6-7)	9
3.2 Understand mechanisms and project future conditions (NCSS Objectives 4-5)	11
3.3 Inform management (NCSS Objectives 1-3).....	11
4. Action Plan	12
4.1 Maintain infrastructure and track change (NCSS Objectives 6-7)	13
4.2 Understand mechanisms and project future conditions (NCSS Objectives 4-5)	13
4.3 Inform management (NCSS Objectives 1-3).....	13
4.4 Descriptions of NERAP Priority Actions.....	14
4.4.1 Maintain infrastructure and track change (NCSS Objectives 6-7).....	14
4.4.2 Understand mechanisms and project future conditions (NCSS Objectives 4-5).....	21
4.4.3 Inform management (NCSS Objectives 1-3).....	22
5. Metrics for measuring success	25
FIGURES AND TABLE	26
REFERENCES.....	33

EXECUTIVE SUMMARY

Changing climate and oceans are affecting the nation's valuable living marine resources and the many people, businesses, and communities that depend on them. Warming oceans, melting sea ice, rising seas, extreme events, and acidification are impacting the distribution and abundance of species, the structure of marine and coastal ecosystems, and, by extension, coastal communities in many regions. These impacts are expected to continue and there is much at risk.

To prepare for and respond to climate impacts on marine and coastal resources and resource users, the 2015 NOAA Fisheries Climate Science Strategy (NCSS) identified 7 key objectives to increase the production, delivery, and use of climate-related information needed to fulfill the agency's mandates (e.g., fisheries management, protected resources conservation) in a changing climate. Beginning in 2016, NOAA Fisheries developed Regional Action Plans (RAPs) to implement the NCSS in each region based on regional needs and capabilities. This Northeast Regional Action Plan (NERAP) builds on previous efforts and describes proposed actions to continue to implement the NCSS in this region through 2024. These activities support NOAA's effort to build a climate-ready nation.

Climate change is directly impacting the ocean and watersheds throughout the Northeast U.S. Over the last 2 decades, ocean temperature in the region has warmed faster than any other marine region in the nation. This warming impacts species distributions, abundance, growth rates, and many other aspects of fish, invertebrate, and marine mammal populations.

Since the publication of the NERAP in 2016, the NOAA Northeast Fisheries Science Center (NEFSC), the Greater Atlantic Regional Fisheries Office (GARFO), and the Chesapeake Bay Office (CBO) have worked with many partners to make progress in addressing the priority actions identified in the 2016 NERAP. A summary of accomplishments can be found in the NCSS 5-year Progress Report. However, there are still many actions needed to better inform fisheries management, protected species conservation, and community adaptation with relevant climate-based information.

This update to the NERAP describes actions needed to continue to address the goals and objectives of the NCSS in this region through 2024.

The following is a list of 10 priority actions and the NCSS objectives they address.

Build and Maintain Infrastructure and Track Change (NCSS Objectives 6-7)

- Maintain and expand ecosystem survey and data collection efforts in the Northeast U.S. Continental Shelf ecosystem including the Bottom Trawl Survey, Ecosystem Monitoring Program, Sea Scallop Survey, Northern Shrimp Survey, Clam Survey, Gulf of Maine Bottom Longline Survey, and Protected Species surveys. This also includes the Gulf of Maine longline survey, data poor species surveys, North Atlantic right whale (*Eubalaena glacialis*) prey sampling, ocean acidification monitoring, and the cooperative shark tagging program.
- Coordinate with other NOAA Programs, Line Offices, and partners to link living marine resource data, science, and management to climate science and research activities.
- Continue to build industry-based fisheries and ocean observing capabilities.

- Continue production of the NEFSC State of the Ecosystem reports and other related products that include climate-relevant information that is useful to management, such as historical, forecasted, and projected biophysical conditions; marine heatwaves; species distribution and abundance shifts; biogeochemical indices; coastal community vulnerability to sea level rise and storm surge; and fishing community vulnerability due to dependence on climate-vulnerable species.

Understand Mechanisms and Project Future Conditions (NCSS Objectives 4-5)

- Conduct laboratory and field research on the mechanistic effects of multiple climate factors (e.g., temperature, ocean acidification, dissolved oxygen) on living marine resources with the goal of informing process-based models for single species, multispecies, and the ecosystem.
- Work with NOAA Oceanic and Atmospheric Research, National Weather Service, National Ocean Service, and academic partners to improve regional hindcasts, forecasts, and projections of ocean and estuarine/river physics and biogeochemistry to develop and improve climate-ready management of living marine resources.

Inform Management (NCSS Objectives 1-3)

- Improve spatial management of living marine resources through an increased utility of spatial and temporal distributions, abundance, productivity, migration, and phenology in management decisions.
- Develop and use Vulnerability Analyses, Scenario Planning, and Management Strategy Evaluations to examine the effects of different management strategies under various climate change scenarios.
- Increase social, economic, and ecosystem scientist involvement in climate change research through multidisciplinary work—including the Northeast Integrated Ecosystem Assessment—that examines relationships between various ecosystem components with the goal of enhancing ecosystem-based management with climate information.
- Develop stock assessment models (e.g., the Woods Hole Assessment Model [WHAM]) that include environmental terms (e.g., temperature, ocean acidification) with a priority for stocks that have upcoming research track assessments.

1. INTRODUCTION

The NOAA Fisheries National Climate Science Strategy (NCSS) seeks to increase the production, delivery, and use of the climate-related information required to fulfill NOAA Fisheries mandates regarding fisheries, protected species, aquaculture, habitats, and ecosystems (Link et al. 2015). Seven interdependent objectives were defined with the goal of informing and fulfilling these mandates in a changing climate (Figure 1). Each marine region across the U.S. is unique in terms of oceanography, marine resource availability, fisheries, protected species, management, and socioeconomic characteristics. Therefore, the national strategy required Regional Action Plans

(RAPs) from the various science centers and regional offices across NOAA Fisheries in order to address each region's unique set of goals and priorities.

Since the publication of the Northeast Regional Action Plan (NERAP) in December of 2016 (Hare et al. 2016a), the Northeast Fisheries Science Center (NEFSC), the Greater Atlantic Regional Fisheries Office (GARFO), the NOAA Chesapeake Bay Office (CBO), and other NOAA Fisheries offices have made substantial progress in addressing the NERAP's priority actions (Peterson et al. 2021). Progress on the NERAP can be tracked in a variety of ways, but a primary metric for tracking research success is to track publications relative to the 7 objectives outlined in the NCSS (Figure 2). From 2016 to 2020, over 60 peer-reviewed reports and journal articles have been published that focus on the Northeast U.S. Continental Shelf ecosystem and directly address the 7 NCSS objectives. These publications were authored or co-authored by NEFSC, GARFO, and CBO staff, and many were the result of collaborations with academic institutions and non-governmental organizations (NGOs).

Reducing the impacts of changing climate and ocean conditions on living marine resources and resource-dependent communities is contingent upon both mitigation and adaptation. Mitigation measures to reduce current and future greenhouse gas emissions or remove carbon dioxide from the atmosphere are key factors in determining which future climate scenario we are headed toward and the degree and types of adaptation needed to increase the resilience of fisheries and communities. NOAA is working to reduce greenhouse gas emissions from its facilities and ships (DOC 2021), as well as taking action to increase resiliency and adapt to current and anticipated impacts of changing climate and ocean conditions on living marine resources. This plan discusses actions to increase the resilience of the nation's valuable living marine resources and the people, businesses, and communities that depend on them. These activities support NOAA's effort to build a climate-ready nation.

2. NERAP ACCOMPLISHMENTS (FY16-FY21)

2.1 Maintaining infrastructure and tracking change (NCSS Objectives 6-7)

Much of the progress made on the NERAP would not be possible without the NEFSC's survey infrastructure. While the NEFSC has maintained its critical surveys, there has been a decline in the number of days at sea available (Figure 3), which has led to a decrease in the amount of data being collected. This is a major concern given that the ecosystem has been changing rapidly; for example, both the trend and variability of ocean temperature within the region has been increasing. These changes in the ocean have been associated with abundance, productivity, and distribution shifts in the living marine resources of the region. Therefore, it is critical that we not only maintain our surveys but increase our survey and data collection activities to effectively track changes in the ocean both seasonally and annually. Many NEFSC programs collect biological and environmental data simultaneously on these surveys, and this interdisciplinary data collection has supported science to understand the effect of climate change on living marine resources. Due to the COVID-19 pandemic, NEFSC surveys were impacted for the spring and summer of 2020, resulting in substantial data gaps (Figure 3). Ocean acidification (OA) monitoring is still limited spatially and temporally, but proposals are being considered to enhance sampling. The use of satellite data is essential for increasing the spatial and temporal coverage of ocean data. Progress

has been made toward regional optimization of phytoplankton size class/functional type algorithms and long-term trend analyses of the 20+ year time series of ocean color data (Turner et al. 2021).

NEFSC scientists have collaborated with academic and NGO groups on various NERAP-related projects. The majority of these projects are funded by various NOAA programs but also stem from NASA, the Lenfest Ocean Program, and the National Science Foundation. Some of our key academic and NGO partners include the Woods Hole Oceanographic Institution (WHOI), Gulf of Maine Research Institute, University of Massachusetts, Rutgers University, University of Connecticut – Stony Brook, Princeton University, University of Rhode Island, and Monmouth University. Increased collaboration with Canada’s Department of Fisheries and Oceans (DFO) has also occurred and resulted in peer-reviewed publications (Greenan et al. 2019; Richardson et al. 2020). The Northwest Atlantic component of the DFO-NOAA Climate and Fisheries Vulnerability Science Collaboration Framework is currently being developed with the goal of enhancing collaborations between NOAA and DFO on climate research.

The recently formed NOAA Fisheries Northeast Climate Science Team can fulfill several roles in advancing the ongoing climate science work of the NEFSC and GARFO. These include promoting integration of various climate science activities across the NEFSC and GARFO; coordinating strategic engagement with partners; promoting awareness of NEFSC, GARFO, and partner-based climate science activities; tracking progress toward climate science goals; being a representative body of the northeast region’s climate science community; interfacing with NOAA leadership on climate science work, support, and prioritization; and targeting relevant funding opportunities toward NERAP priorities. The NEFSC is tracking and communicating climate research through the ecosystems and climate change webpage and the New England groundfish in a changing climate webpage.

With support from the NOAA’s Ocean Acidification Program (OAP), the NEFSC has conducted experimental studies to understand the resilience to OA in the following shellfish species with both fisheries and aquaculture relevance: the Eastern Oyster (*Crassostrea virginica*), the northern surf clam (*Spisula solidissima*), and the Atlantic sea scallop (*Placopecten magellanicus*). Data from laboratory experimental studies are supporting dynamic energy budget (DEB) models that project commercially-relevant performance variables, including growth rate, time to harvest size, and reproductive potential under OA scenarios projected by Earth system models. Another OA Program-supported project includes assessments of impacts to fishing communities due to impacts on Atlantic sea scallops.

The NERAP identified the need for a Northeast Watershed Program: a coordinated, multidisciplinary effort comparable to the Northwest Fisheries Science Center Watershed Program for developing basic and applied science—including climate science—in support of the management of diadromous species in freshwater and estuarine environments. While a dedicated, funded program has not yet been achieved, a group of interested individuals representing multiple offices (NEFSC, GARFO, and National Marine Fisheries Service [NMFS] Office of Habitat Conservation) meet regularly to identify, fund, and implement priority studies (e.g., Collins 2019; Lombard et al. 2021) and advance program development.

Changes in the ecosystem are tracked and reported through the annual state of the ecosystem reports for New England and the Mid-Atlantic. Ecological and biological indicators are routinely updated along with climate indicators for fishing communities (Clay et al. 2020). Tracking change in the physical and chemical state of the ocean through variables such as temperature (Chen et al. 2020; Friedland et al. 2020a; Gawarkiewicz et al. 2019), ocean circulation (Saba et al. 2016; Caesar et al. 2018), and acidification (Meseck et al. 2018; Poach et al. 2019) are

key to developing indicators of climate change in marine ecosystems. The close collaboration between the NEFSC and NOAA's Geophysical Fluid Dynamics Laboratory (GFDL) has streamlined the use of high-resolution global climate models that are used for projecting long-term change to the Northeast U.S. Continental Shelf. The shelf is an oceanographically complex region that is challenging to model due its fine-scale bathymetry and circulation. GFDL's CM2.6 global climate model has been an integral part of our NERAP progress on projecting future conditions.

Studies that track contemporary change and project future change in marine taxa have almost exclusively focused on changes in species distributions. Impacts of warming ocean temperature on marine species distributions is a very common research theme within the Northeast U.S. This is due to the high spatial and temporal availability of ocean temperature data combined with the long-term time series of the NEFSC's fall and spring bottom trawl survey, which is a fisheries-independent survey that samples shelf waters from Cape Hatteras, NC, to the Gulf of Maine and collects temperature and species data concomitantly. These studies have analyzed a broad suite of marine taxa in the contemporary period from the 1970's onward (Kleisner et al. 2016; Friedland et al. 2019) to document observed distribution shifts associated with warming ocean temperature. Other studies have relied on the NEFSC EcoMon survey data to document shifts in zooplankton (Morse et al. 2017) and ichthyoplankton (Walsh et al. 2015; McManus et al. 2018).

2.2 Understanding mechanisms and projecting future conditions (NCSS Objectives 4-5)

The majority of NERAP research has addressed objectives 4-5, which focus primarily on understanding mechanisms of change and developing long-term projections of future conditions (Figure 2).

Decadal-scale projections of species distribution shifts under climate change have relied on GFDL's CM2.6 (Kleisner et al. 2017; McHenry et al. 2019), which projects an enhanced warming of the U.S. Northeast Continental Shelf due to a change in regional circulation (Saba et al. 2016). The mechanisms resolved in CM2.6 corroborate contemporary observations over the last decade (Neto et al. 2021). This high-resolution global climate model has been utilized extensively for projections of protected species distributions such as loggerhead sea turtles (*Caretta caretta*; Patel et al. 2021) and the prey of critically endangered North Atlantic right whales (*Eubalaena glacialis*; Grieve et al. 2017). Additionally, a project is underway to link available data to describe and understand how socioeconomic conditions affect U.S. loggerhead nesting beach success rates. Major land threats to the loggerhead nesting sites in the U.S. include construction (armoring), ecosystem alterations (beach erosion), pollution (light), and species interactions (predation). As the range of these turtles changes due to climate change, additional socioeconomic data will be needed for any new nesting beach sites. Moreover, assessing local light pollution ordinances in connection with nesting data may yield insight into the relative effectiveness of different rules.

The skill of seasonal to annual (S2A) forecasts of ocean conditions in the U.S. Northeast Continental Shelf, such as sea surface temperature, is relatively low compared to other large marine ecosystems (Stock et al. 2015). The reason for the poor skill in this region is because SST forecasts are derived from global models that have coarse resolution in their ocean and atmosphere components. Tactical fishery and protected species management may greatly benefit from more skillful S2A forecasts of ocean conditions. Given the majority of commercial species within the

region are demersal, bottom temperature forecasts may be more relevant to stock assessments and management advice. A new statistical model that forecasts bottom temperature in the U.S. Northeast Continental Shelf has significant skill for lead times up to 5 months in the Mid-Atlantic Bight and up to 10 months in the Gulf of Maine, although the prediction skill varies notably by season (Chen et al. 2021).

Estuaries and rivers are important habitats for many marine species in the U.S. Northeast, and thus there is a need to understand and predict changes to these watersheds associated with climate. Many of these habitats are listed as critical or essential under the Endangered Species Act. Historical, downscaled model hindcasts of Long Island Sound (Georges et al. 2016; Schulte et al. 2017; Schulte et al. 2018) and the Chesapeake Bay (Muhling et al. 2017; Muhling et al. 2018; Ross et al. 2021) have been developed and analyzed relative to large scale climate forcing.

A characterization of river flood seasonality and trends in 90 watersheds across the northeast U.S. was also completed (Collins 2019). Northeast U.S. rivers were evaluated in a national study focusing on large-flood seasonality and historical occurrence trends (Collins et al. 2022).

Understanding mechanisms of climate change impacts on marine ecosystems is a critical component of studies that utilize observed relationships between the environment and marine taxa to model historical and projected change. Laboratory-based process studies at the NEFSC are at the forefront of this research at both the Milford, CT, and Sandy Hook, NJ, facilities. The Milford lab primarily focuses on economically and ecologically important bivalve shellfish while Sandy Hook focuses on finfish and some invertebrates. In the laboratory, physiological processes (including feeding rates) have been documented for bivalve species including oysters, scallops, and surf clams (*Spisula solidissima*) under OA conditions (Meseck et al. 2020; Pousse et al. 2020). In the field, Milford scientists documented that total benthic bivalve abundance is correlated with sediment carbonate chemistry parameters (Meseck et al. 2018).

These studies demonstrate that from the laboratory to the field, bivalves are sensitive to OA. Research projects in collaboration with our academic partners have analyzed the impacts of changing water temperature on black sea bass (*Centropristis striata*) aerobic scope and hypoxia tolerance (Slesinger et al. 2019); estimates of CO₂ and co-stressor effects on early life-stages of finfish such as winter flounder (*Pseudopleuronectes americanus*), summer flounder (*Paralichthys dentatus*), mid-Atlantic forage fishes, and New England groundfish; and individual-based process models of CO₂ effects on winter flounder. The Sandy Hook Lab has developed a novel apparatus for testing plasticity of responses to thermal, CO₂, and dissolved oxygen regimes. These studies capture how rate processes are affected by environmental drivers and are precisely the kinds of quantitative descriptions needed in order to model climate change impacts. All of these laboratory and field studies are needed to inform single, multispecies, and ecosystem models. The improved mechanistic understanding of climate impacts on species, habitats, and communities combine with forecasts of future conditions to provide climate vulnerability assessments (CVAs) and assess which ecosystem components are at greatest risk. Vulnerability assessments are available for fish/invertebrates (Hare et al. 2016b), habitat (Farr et al. 2021), and fishing communities (Colburn et al. 2016), and methodologies have been developed for conducting assessments for marine mammals and sea turtles (Lettrich et al. 2019, 2020).

2.3 Informing management (NCSS Objectives 1-3)

A primary way management can use climate information is through climate-enhanced fishery stock assessments (Holsman et al. 2019). Progress on climate-enhanced stock assessment

variables (e.g., demographics, recruitment, population growth) has been moving forward on key commercial and recreational species including southern New England yellowtail flounder (*Limanda ferruginea*; Miller et al. 2016a; Xu et al. 2018), summer flounder (O’Leary et al. 2019; O’Leary et al. 2020), winter flounder (Bell et al. 2018), northern shrimp (*Pandalus borealis*; Cao et al. 2017), Atlantic cod (*Gadus morhua*; Miller et al. 2018), surf clam (Hennen et al. 2018), and black sea bass (Miller et al. 2016b). The Woods Hole Assessment Model (WHAM) was developed by scientists at the NEFSC, and it can be used to support climate-enhanced stock assessments via the incorporation of time-varying processes with links to environmental covariates (Stock and Miller 2021). A framework has been developed for incorporating climate and habitat information into fisheries management using risk assessment and management strategy evaluation (MSE; Gaichas et al. 2016). Support was provided to the Mid-Atlantic Fisheries Management Council (MAFMC) risk assessment (Gaichas et al. 2018), which included the results from the CVA and habitat shifts into a conceptual model for high-risk summer flounder fisheries in 2019 (DePiper et al. 2021).

Further progress has been made on social science research that connects changes in levels of dependence on fishing to levels of social well-being in fishing communities and vulnerability to sea level rise and storm surge. Additional fishery-level performance metrics allow the calculation of changes in revenue by management plan over time that can be discussed in relation to changes in changing availability of species related to climate change. All revenue figures are in 2021 dollars based on the gross domestic product (GDP) implicit price deflator, allowing comparison of revenue between time periods using a consistent dollar value. Simulation models have been developed that address various climate impacts to single species and have evaluated climate-informed reference points. This work is coupled to new research that links climate- and stock-related projections for groundfish to economic outcomes for fishermen and fishing communities. Statistical models are being developed that explain how fishermen select target stocks and landing locations. These models can then be used to understand how these behaviors will change under various climate and policy scenarios. Other social science research projects, the first of which has been funded by the New England groundfish/climate program, include: 1) climate vulnerabilities and adaptation pathways for northeast U.S. fishing communities, including developing indices of vulnerability to climate change for groundfish at the fishing community level; 2) stakeholder engagement in MSE of New England groundfish in a changing ocean; 3) developing northeast fishing community indices of vulnerability to climate change based on sea surface temperature, stock size/status, and OA using the 82 species in Hare et al. (2016b); 4) developing indicators of climate vulnerability, specifically to OA, for northeast fishing communities dependent on landings of Atlantic sea scallops (*Placopecten magellanicus*); and 5) tracking community-level socioeconomic conditions impacting loggerhead nesting as nesting beaches move further north.

Important advancements have been made specific to climate and protected species. Climate scenario planning workshops and reports were completed for endangered North Atlantic right whales (Borggaard et al. 2020) and Atlantic salmon (*Salmo salar*; Borggaard et al. 2019). A range-wide salmon habitat synthesis is ongoing and will describe habitat conditions suitable, including preferences and tolerances, for Atlantic salmon by life stage in freshwater and marine systems. Additional work is aimed at identifying climate-resilient habitats throughout Maine watersheds that are listed as critical habitat for Atlantic salmon. Part of this effort includes understanding where rivers are naturally high in baseflow, which is streamflow that has a relatively high proportion of inputs from groundwater and/or lake/wetland outflow. This is important because

rivers with high baseflow tend to be cooler and have greater flow during summer. These areas thus provide refuge during thermal stress periods for Atlantic salmon. A model was developed to predict and map relative baseflow quantities at a high spatial resolution (~2.5 km²), providing valuable information to managers making decisions about where to conserve or restore habitat (Lombard et al. 2021). Through the Collaborative Management Strategy for the Gulf of Maine Atlantic Salmon Recovery Program, work has been done with tribal partners, the state of Maine, U.S. Fish and Wildlife Service (USFWS), and NGOs to identify and protect climate-resilient habitats important to Atlantic salmon (CMS 2019).

For North Atlantic right whales, short-duration zooplankton sampling trips were conducted in the southern New England region during the winter and early spring (January-April) of 2019, 2020, and 2021 when right whales are in the area. Our goal is to describe vertical distribution patterns of right whale prey in relation to physical features to better understand the mesoscale processes that result in super aggregations of right whale prey. Correlations between ocean warming and right whale prey availability suggest an inverse relationship between *Calanus* spp. and ocean temperature (Sorochnan et al. 2019).

Additional progress has been made to advance regional watershed science through continued coordination across the region. To better understand the impacts of historical and projected climate change in freshwater and estuarine environments, synthesis work on the diadromous fish community and ecosystem interactions of 12 NOAA trust species started in 2020. This synthesis will integrate the structure and function of diadromous fish communities in freshwater ecosystems while synthesizing their ecosystem roles and interactions in the northeast U.S. This synthesis will encompass changing environmental conditions in rivers, estuaries, and the coastal ocean. Differing rates of change and species distributions caused by climate but mitigated with habitat improvements (enhanced fish passage and restoration of stream functions) require comprehensive analyses. Through this synthesis, the research gaps and needs in 2 rapidly changing systems will be highlighted. Additionally, the NEFSC continues to monitor phenology changes in Atlantic salmon to better understand climate impacts and drivers on this protected species.

3. KEY NEEDS

3.1 Maintain infrastructure and track change (NCSS Objectives 6-7)

None of the NERAP research can be successful without a solid infrastructure for ocean observations that includes both physical and biological surveys. There is concern about the decline in the number of observations per year, which is a direct result of the declining number of days at sea (Figure 3). This observation decline is occurring at a time of both increasing trend and variability of many ocean and biological variables. Skillful models, whether for single species or the entire ocean ecosystem including human dimensions, can only be developed and validated if observations exist over sufficient temporal and spatial scales that capture seasonal, annual, and decadal variability. Therefore, there is a critical need to increase the number of days at sea for the NEFSC's EcoMon survey and to broaden the suite of biogeochemical variables that are measured and include depth profiles of nutrients, plankton productivity, dissolved oxygen, and pH.

Further complicating the continuation of these ecosystem surveys is planned offshore wind development that will result in major adverse impacts on NOAA Fisheries scientific surveys along much of the shelf from Massachusetts south through North Carolina. Wind energy development

impacts scientific surveys in 4 ways: 1) preclusion of sampling platforms from wind energy areas due to operational and safety limitations; 2) impacts on the random-stratified statistical design that is the basis for scientific assessments, advice, and analysis; 3) alteration of benthic, pelagic, and airspace habitats in and around wind energy areas; and 4) impacts on sampling efforts outside wind energy areas by increasing vessel transit time and challenges to aircraft operations. To address the major adverse impacts from offshore wind development on NOAA Fisheries survey enterprise, this objective will also include implementation of a Federal Survey Mitigation Program for the northeast region as called for in the draft Federal Survey Mitigation Implementation Strategy (Hare et al. 2022).

Progress has been made toward regional optimization of phytoplankton size class/functional type algorithms (Turner et al. 2021), but more work is needed to use surface satellite measurements of ocean variables as indicators of ecosystem change and variability. New biological surveys, such as the Gulf of Maine longline survey and right whale prey surveys, need to continue to track changes in key ecosystem indicators from lower to higher trophic levels. Additional dedicated funding for social and economic surveys and other research will also be needed to build up the time series data needed for effective modeling.

Increased collaboration with the fishing industry, through cooperative research, is also needed to enhance observed datasets of targeted and non-targeted catch, as well as physical measurements such as ocean temperature (e.g., eMOLT Program, Study Fleet Program; Manning et al. 2009; Jones et al. 2022). Continued collaboration with academic institutions and NGOs needs to continue, as well as research and data coordination with Canada's DFO and NOAA's Southeast Fisheries Science Center (SEFSC) regarding commercial species, protected species, and ecosystem indicators. Continued collaborations with tribal partners, the state of Maine, the USFWS, and NGOs are also important in identifying and protecting climate-resilient Atlantic salmon habitats. Further science is needed on full life cycle habitat resilience for salmon that connects changes in rivers to associated estuary and marine habitats. As such, a salmon-specific CVA and MSE would be a capstone project of the NOAA climate strategy and benefit both NOAA and partner agency managers. Better integration with the Northeast Integrated Ecosystem Assessment (IEA) Program can help translate climate information into ecosystem assessments, state of the ecosystem reports, ecosystem and socioeconomic profiles, and ecosystem status reports.

A major gap in the present infrastructure is the lack of a centralized, regional online catalog that contains ocean and climate data and model output for the northeast. Presently, these data and model outputs are maintained at various web portals and require additional resources to render them for regional analyses. Climate research funding also needs to continue so that targeted research projects can continue to address the priorities of the NERAP. Most funding that has addressed NERAP priorities derives from NOAA's Oceanic and Atmospheric Research (OAR) Climate Program Office (CPO) and OAP, as well as NEFSC funding from NERAP/Ecosystem-Based Fisheries Management (EBFM)/Stock Assessment Improvement Plan (SAIP)/IEA, and the New England groundfish/climate program. In collaboration with external partners, some projects are funded by NASA, Lenfest, and the National Science Foundation (NSF). While this piecemeal funding process has enabled critical research that addresses NERAP priorities, a more solidified, permanent funding infrastructure designed to produce and maintain climate research is needed. The recent NOAA Climate, Ecosystems, and Fisheries Initiative (CEFI) addresses the ocean modeling and communities of practice component of this research but has not been fully funded.

3.2 Understand mechanisms and project future conditions (NCSS Objectives 4-5)

Projections of future change have largely focused on marine species habitat and distribution using NOAA's high-resolution global climate model CM2.6. While these long-term projections (20-80 years) could be useful for fishery management plans, MSEs, scenario planning, and vulnerability assessments over decadal periods, they are not useful for tactical management decisions that are made on a year-to-year to decadal time scale. S2A forecasts of ocean conditions that are tied to stock assessments would be more useful to tactical fisheries management. However, the skill of ocean forecasting models for even the most standard ocean variable—sea surface temperature—is relatively low in the U.S. Northeast marine ecosystem compared to other coastal large marine ecosystems. Improved S2A skill of bottom temperature forecasts using statistical methods has been completed, but the improved skill is specific to certain regions and time periods (Chen et al. 2020). Through NEFSC and OAR CPO funding, pilot studies have started the process of a dynamical approach, which is to develop regional ocean models for the Northwest Atlantic that can run in hindcast, forecast, and projection mode. This dynamical approach directly addresses the recent CEFI, and the goal is to have multiple regional ocean model simulations for the region that are based on NOAA's state-of-the-art ocean model, MOM6. NOAA OAR CPO has also funded the development of annual to decadal ocean forecasts using the Scripps Coupled Ocean-Atmosphere Regional (SCOAR) modeling system.

While more skillful S2A ocean forecasts are a critical need, high-resolution biophysical ocean hindcasts are an essential need to fill temporal and spatial gaps in the observed time-series of key ocean variables, such as bottom temperature, circulation, pH, primary and secondary productivity, and dissolved oxygen. Tracking change to the northeast U.S. continental shelf doesn't need to be limited to in situ observations, which have been declining in the northeast U.S. (Figure 3). Ocean model hindcasts, after validation, can also be used to track historical changes at a much higher spatial and temporal resolution than in situ measurements. These ocean hindcasts are also critical to understand mechanisms between ocean change and living marine resources through the association of changes in survey data to key biophysical ocean variables in validated ocean models. While there have been previous efforts to develop regional ocean model hindcasts and projections that resolve biogeochemistry (Zhang et al. 2019), more models, including those that assimilate data, are needed to assess model uncertainty and to create model ensembles of historical, near-term, and long-term ocean conditions.

Laboratory and field process studies that determine mechanistic links between the ocean environment and marine species are needed to inform process-based single species, multispecies, and ecosystem models. While the NEFSC and its partners have made progress on identifying mechanistic underpinnings between temperature, OA, and marine species, more studies are needed on key species that were identified to be highly vulnerable to climate change (Figure 4). These studies need to focus on multiple life history stages from larval to adult and investigate synergistic impacts of warming ocean temperature and OA.

3.3 Inform management (NCSS Objectives 1-3)

There is a critical need for more focused research that can inform and enhance living marine resource tactical management decisions for commercial, recreational, and protected species. This is a very challenging task not just for the northeast region but for all U.S. regions. The northeast U.S. fisheries management system consists of 2 Federal councils, the Atlantic States

Marine Fisheries Commission (ASMFC), the NOAA Office of Sustainable Fisheries – Atlantic Highly Migratory Species Division, state agencies, and GARFO. Working with management partners is essential for informing management. Moreover, there are very few operational fishery stock assessments in the U.S. and worldwide that use environmental or ecosystem data quantitatively or qualitatively to inform year-to-year management decisions or even longer-term fishery management plans. Finally, there are no living marine resource management decisions in the U.S. that are based on forecasted (S2A) or projected (decadal) ocean conditions. Therefore, it is essential to produce new research results that support the use of climate and environmental information for upcoming research track stock assessments, which may be the primary mechanism to inform management with climate-enhanced stock assessments. The recent Northeast Climate Integrated Modeling (NCLIM) project is an example of this approach with the goals of: 1) identifying and anticipating major ecosystem changes that influence multiple stocks or management decisions; 2) informing decision-making around impacts of shifting species; and 3) informing decision-making around changes in stock productivity. Climate information has also been used in scenario planning for Atlantic salmon (Borggaard et al. 2019) and North Atlantic right whales (Borggaard et al. 2020), but more of these exercises are needed to help inform MSEs for harvested species and conservation planning for protected species.

Progress on informing management is also based on studies that analyze human dimension variables, such as socioeconomic- and climate-informed reference points and climate vulnerability indices for fishing communities in the region. Social vulnerability indicators for fishing communities, meanwhile, provide an important context for understanding the impact of climate change; for example, highly vulnerable communities may be more likely to have difficulty responding effectively to climate changes. End-to-end ecosystem models, such as the northeast U.S. version of ATLANTIS (NEUS Atlantis), attempts to simulate the entire ecosystem from fundamental physical and chemical processes to food webs to fisheries management and social and economic factors. The NEUS Atlantis can be used to model human dimensions variables (management strategies, fleet behavior, and market responses) under various climate change scenarios over decadal periods. However, these sub-models have not been implemented in the current version of NEUS Atlantis. Therefore, dedicated research is needed to parameterize and validate fishery behavior, management scenarios, and market responses. Moreover, social vulnerability indicators and cultural factors impacting fishery decision making are not easily included in NEUS Atlantis, and region-wide comparable data, especially for cultural factors, are currently lacking. Ultimately, with the future integration of global climate and regional ocean models, NEUS Atlantis will be able to inform management through full ecosystem-level projections under multiple climate scenarios and under an array of human behaviors.

4. ACTION PLAN

NERAP priority actions for FY22-FY24 are listed and described by NCSS objectives. Many NERAP Actions are relevant to multiple NCSS objectives but are aligned with the most relevant objective. In total, 10 NERAP priority actions are identified, and each action has equal priority to others. Many of these priorities address multiple NOAA Fisheries mission elements (sustainable fisheries, protected resources, aquaculture, habitat, and ecosystem), and this plan works across all mission elements. The public comments on section 216(c) of Executive Order 14008 (Making Fisheries and Protected Resources More Resilient to Climate Change) were considered in the development of these priority actions.

4.1 Maintain infrastructure and track change (NCSS Objectives 6-7)

NERAP Priority Action 1: Maintain and expand ecosystem survey and data collection efforts in the Northeast U.S. Continental Shelf ecosystem including the Bottom Trawl Survey, Ecosystem Monitoring Program, Sea Scallop Survey, Northern Shrimp Survey, Clam Survey, Gulf of Maine Bottom Longline Survey, and Protected Species surveys. This also includes the Gulf of Maine longline survey, data poor species surveys, right whale prey sampling, ocean acidification monitoring, and the cooperative shark tagging program.

NERAP Priority Action 2: Coordinate with other NOAA Programs, Line Offices, and partners to link living marine resource data, science, and management to climate science and research activities.

NERAP Priority Action 3: Continue to build industry-based fisheries and ocean observing capabilities.

NERAP Priority Action 4: Continue production of the NEFSC State of the Ecosystem reports and other related products that include climate relevant information that is useful to management, such as historical, forecasted, and projected biophysical conditions; marine heatwaves; species distribution and abundance shifts; biogeochemical indices; coastal community vulnerability to sea level rise and storm surge; and fishing community vulnerability due to dependence on climate-vulnerable species.

4.2 Understand mechanisms and project future conditions (NCSS Objectives 4-5)

NERAP Priority Action 5: Conduct laboratory and field research on the mechanistic effects of multiple climate factors (e.g., temperature, ocean acidification, dissolved oxygen) on living marine resources with the goal of informing process-based models for single species, multispecies, and the ecosystem.

NERAP Priority Action 6: Work with NOAA OAR, National Weather Service (NWS), National Ocean Service (NOS), and academic partners to improve regional hindcasts, forecasts, and projections of ocean and estuarine/river physics and biogeochemistry to develop and improve climate-ready management of living marine resources.

4.3 Inform management (NCSS Objectives 1-3)

NERAP Priority Action 7: Improve spatial management of living marine resources through an increased utility of spatial and temporal distributions, abundance, productivity, migration, and phenology in management decisions.

NERAP Priority Action 8: Develop and use Vulnerability Analyses, Scenario Planning, and Management Strategy Evaluations to examine the effects of different management strategies under various climate change scenarios.

NERAP Priority Action 9: Increase social, economic, and ecosystem scientist involvement in climate change research through multidisciplinary work—including the Northeast Integrated Ecosystem Assessment—that examines relationships

between various ecosystem components with the goal of enhancing ecosystem-based management with climate information.

NERAP Priority Action 10: Develop stock assessment models (e.g., WHAM) that include environmental terms (e.g., temperature, ocean acidification) with a priority for stocks that have upcoming research track assessments.

4.4 Descriptions of NERAP Priority Actions

4.4.1 Maintain infrastructure and track change (NCSS Objectives 6-7)

NERAP Priority Action 1: Maintain and expand ecosystem survey and data collection efforts in the Northeast U.S. Continental Shelf ecosystem including the Bottom Trawl Survey, Ecosystem Monitoring Program, Sea Scallop Survey, Northern Shrimp Survey, Clam Survey, Gulf of Maine Bottom Longline Survey, and Protected Species surveys. This also includes the Gulf of Maine longline survey, data poor species surveys, right whale prey sampling, ocean acidification monitoring, and the cooperative shark tagging program.

The NEFSC has a long history of conducting surveys of the Northeast U.S. Shelf ecosystem ranging from chemistry through marine mammals and seabirds. This effort should be maintained and is fundamental to the success of the NERAP. The decreasing trend in the number of days at sea (Figure 3) during a period of increasing trends and variability in ocean conditions is very concerning. Further complicating the continuation of these ecosystem surveys is planned offshore wind development that will disrupt NOAA scientific surveys along much of the shelf from Massachusetts south through North Carolina, with future impacts to surveys in the southeast region. The NEFSC will explore opportunities to measure ocean change utilizing the proposed wind platforms in the region as resources become available to implement a Northeast Regional Survey Mitigation Program (Hare et al. 2022). Enhanced survey data coordination and expansion with the SEFSC is also needed. While increasing the number of ocean observations is beyond the scope of the NERAP, it is still a critical priority to maintain the timeliness, precision, and accuracy of these surveys and maximize the number of observations each year. These surveys include:

- Spring and Fall Bottom Trawl Survey – 2 times per year (including Ecosystem Monitoring Program operations)
- Ecosystem Monitoring Survey – 4 times per year
- Scallop Survey – 1 time per year
- Northern Shrimp Survey – 1 time per year
- Gulf of Maine Cooperative Bottom Long Line Survey
- North Atlantic Right Whale Aerial Survey
- Marine Mammal and Sea Turtle Shipboard and Aerial Surveys
- Large Coastal Shark Bottom Longline Survey
- Cooperative Atlantic States Shark Pupping and Nursery Longline/Gillnet Survey
- Clam and Ocean Quahog Survey – 1 time per year

Protected species surveys and telemetry tagging are also critical to maintain (e.g., North Atlantic right whale, sea turtles, seals, Atlantic salmon, Atlantic Marine Assessment Program for Protected Species). To the extent possible, climate, ecosystem, and habitat information should be collected on all surveys, thereby allowing simultaneous environmental and biological data to be collected and used in a number of analyses related to other actions described here in the NERAP.

Continued collection of fishery-dependent data is also critical to living marine resource management, and these data can be used to improve the scientific understanding of the effect of climate change on fisheries in the northeast U.S.

NERAP Priority Action 2: Coordinate with other NOAA Programs, Line Offices, and partners to link living marine resource data, science, and management to climate science and research activities. These include:

- **NOAA Integrated Ecosystem Assessment Program:** The northeast IEA team provides several products to the region's fishery management councils. Recognizing the importance of climate in the region, the second priority goal of the northeast IEA work plan is to integrate climate and earth system modeling to further understand past ecosystem variability and assess predictions across management-relevant time scales. This priority goal will be achieved through 4 related objectives. The first will be to analyze high-resolution regional physical and physical-biological simulations to understand past ecosystem variability and indicator robustness. This will include the development of diagnostics for the representation of IEA indicators and associated ecosystem status report measures in high-resolution regional earth system simulations. For those indicators successfully captured by the simulations, the IEA team will try to understand the drivers of indicator variability and the physical and biogeochemical changes associated with each indicator shift. The second objective will be to assess the seasonal to multi-decadal predictability and robustness of IEA indicators in the U.S. northeast shelf. The predictability of IEA-associated indicators will be assessed within the seasonal to multi-annual predictions associated with the first objective. In addition, CMIP6-driven climate change projections will be analyzed to understand the robustness of climate change indicators. The third objective will be to integrate regional earth system predictions and projections into the region's ecosystem status reports and other associated living marine resource applications. This objective builds off the first 2 and will bring the climate and earth system modeling into a product that goes before the region's fisheries management councils. The final objective is to continue development of estuarine forecasting capabilities that will complement the shelf-scale hindcast, prediction, and projection capabilities in the other objectives.
- **NOAA Chesapeake Bay Office:** The CBO is supporting research to understand the impacts of changing habitat and climate conditions on key species, such as striped bass (*Morone saxatilis*), summer flounder, and forage fish in the Chesapeake Bay. Specific projects include development of an estuarine habitat index for summer flounder, a nursery habitat assessment for striped bass, and seasonal summaries for the Mid-Atlantic State of the Ecosystem Report. The seasonal summaries currently include CBO's Chesapeake Bay Interpretive Buoy System, data as well as satellite data from CBO's cross Line Office collaboration with the NOAA Environmental Satellite Data and Information Service (NESDIS) CoastWatch program. The CBO is enhancing observational capacity to track changing habitat conditions and impacts on living resources by deploying 2 vertical high-frequency hypoxia profilers in the Chesapeake Bay mainstem in fall 2021 and establishing a mainstem telemetry receiver array to track fish movement. The hypoxia profilers will provide

data to develop fish habitat condition products over the next few years. A fishery biologist has been hired to analyze existing telemetry receiver data and develop a science plan for the new array. The CBO is also part of the NOAA-funded project to develop a shellfish Regional Vulnerability Assessment (RVA) in the Chesapeake Bay, which aims to: 1) assess the vulnerability of the oyster aquaculture industry and oyster restoration to OA and other co-stressors; and 2) produce the information required by regional communities to aid in adaptation to these stressors. In achieving these goals, we will better understand which shellfish stakeholders will be able to successfully adapt, which will seek alternative livelihoods, and what specifically causes the difference between these 2 disparate outcomes. Social, cultural, and economic factors will all play a role. The CBO is also supporting efforts to synthesize information on rising water temperatures, their effects on Chesapeake Bay fisheries and habitat, and the development of a Bay Water Temperature Change Indicator related to these effects as part of a larger partnership with the Chesapeake Bay Program. Information from the NOAA Northeast Fish and Shellfish CVA and Habitat CVA are being applied to inform the synthesis. Buoy data from the NOAA Chesapeake Bay Interpretive Buoy System (CBIBS) and satellite data from the NESDIS are being evaluated for the Bay Water Temperature Change Indicator, as well as other long-term data sources. Workshops with the Chesapeake Bay Program's Scientific and Technical Advisory Committee (STAC) were scheduled for 2022 to inform discussions to develop recommendations on management responses to ecological impacts from rising water temperatures.

- NOAA Fisheries Office of Aquaculture: The following key research and action needs relevant to the interests of the Office of Aquaculture have been identified by NEFSC scientists: 1) breeding for climate resilience in oysters and other cultivated shellfish; 2) host-parasite or pathogen interactions under climate-change conditions, including biogeographic range expansion of parasites; 3) range expansion of shellfish predators (e.g., blue crabs [*Callinectes sapidus*] and cownose rays [*Rhinoptera bonasus*]) that will affect regional shellfish cultivation practices; 4) diversification in the shellfish farming sector (e.g., sea scallop farming development in Maine) and hatchery production of surfclams for resource enhancement of offshore fisheries; 5) continued improvements in dependability of shellfish hatchery production, including probiotics and microbiome optimization, application of monitoring and control technologies, and improved nutrition; 6) cultivation strategies for offshore blue mussels (*Mytilus edulis*) consistent with protected species restoration goals; 7) use of cultivated shellfish in comprehensive, living shoreline restoration projects; 8) improved understanding of the factors which affect fishing community attitudes toward the development of aquaculture in their communities and of fisher attitudes toward their possible involvement in aquaculture as an alternative or a supplement to fishing (Johnson et al. 2019).
- NOAA Ocean Acidification Program: A key research and action need of the OA Office has been identified by NEFSC scientists as improved understanding of the impacts of OA on affected species and the communities that harvest them. The following research and action needs relevant NOAA OAP for the Northeast Continental Shelf have been identified: 1) improved forecasting of OA across daily to decadal timescales that better quantifies the primary drivers of carbonate

dynamics in context with other environmental changes (i.e., temperature and riverine influence); 2) determine how OA in concert with other stressors impacts ecologically and/or economically important marine species, with a focus on understanding the effects on aquaculture stocks; 3) evaluate costs and benefits of mitigation and adaptation strategies for communities, ecosystems, and economics; and 4) promote integration of OA understanding into regional planning and management. Components to this research include: 1) biological samples (pteropods) with EcoMon monitoring cruises for in situ indicators of climate change; 2) laboratory multi-generational experiments to explore climate change adaptation potential of marine bivalves; 3) dynamic energy budget models of surfclams, oysters, and sea scallops to determine individual responses to climate change; 4) combination laboratory and field studies on Atlantic surf clams to evaluate aquaculture practices; and 5) assessment of Atlantic sea scallop vulnerability to OA and the relative social vulnerability of scallop-dependent communities to future OA conditions.

- NOAA Ocean and Atmospheric Research, National Ocean Service (NOS), and National Weather Service (NWS): Collaboration with these 3 NOAA Line Offices is needed to achieve the ocean modeling objectives outlined in the NOAA Climate and Fisheries Initiative. Regional ocean and watershed hindcasts, forecasts, and projections of physical and biogeochemical conditions require dedicated regional ocean model development by OAR's GFDL, coastal ocean and watershed model development by NOS, and S2A forecasting models developed by NWS. Coordination with NOAA OAR's CPO is needed to develop targeted research requests for annual funding opportunities within the Modeling, Analysis, Predictions and Projections (MAPP) Program, Climate and Fisheries Adaptation (CAFA) Program, and Climate Variability & Predictability (CVP) Program.
- NOAA Fisheries Atlantic Highly Migratory Species Management Division: Work with the Atlantic Highly Migratory Species Division and the Southeast Fisheries Science Center to conduct a CVA for Atlantic highly migratory species (HMS). Several shark stocks were included in previous assessments, but the need remains for a more complete Atlantic HMS CVA. The inclusion of a comprehensive Atlantic HMS CVA as a near-future project would address various priorities of the NERAP and the southeast regional action plan (SERAP). When HMS CVA discussions occurred in 2014-2015, it was decided to wait to conduct an HMS CVA until the regions had finished CVAs for council-managed fisheries. Council-managed CVAs are mostly done, which has led to some excellent ongoing work applying results to management as a key tool in addressing climate change. Without an HMS CVA, HMS fisheries and their stakeholders cannot be included as effectively in these discussions of management approaches.
- State agencies: Continued collaborations with state agencies are important to leverage resources, coordinate on research and monitoring, and share goals of climate initiatives (e.g., Maine Department of Marine Resources and Maine Climate Council).
- Academic institutions and non-governmental organizations: Research collaborations are needed with key academic and non-governmental partners such as Rutgers University, WHOI, University of Massachusetts, University of

Connecticut, University of Rhode Island, University of Maryland, University of Maine, Princeton University, Gulf of Maine Research Institute, and the Lenfest Ocean Program. The Cooperative Institute for the North Atlantic Region (CINAR) is one vehicle for these partnerships.

- Watershed Program for the East Coast: Diadromous species are important in the region for a variety of reasons (e.g., protected species, commercial and recreational harvest, ecosystem interactions). These species include Atlantic salmon, Atlantic sturgeon (*Acipenser oxyrinchus oxyrinchus*), shortnose sturgeon (*Acipenser brevirostrum*), rainbow smelt (*Osmerus mordax*), alewife (*Alosa pseudoharengus*), blueback herring (*Alosa aestivalis*), American eel (*Anguilla rostrata*), hickory shad (*Alosa mediocris*), American shad (*Alosa sapidissima*), striped bass, sea-run brook trout (*Salvelinus fontinalis*), sea lamprey (*Petromyzon marinus*), white perch (*Morone americana*), and tomcod (*Microgadus tomcod*). These species are included in the larger group of species considered in many of the actions prioritized here, but there are also a number of specific needs that exceed the scope of the NCSS and this NERAP. On the West Coast, the Northwest Fisheries Science Center hosts a Watershed Program, which investigates the ecology of freshwater and estuarine ecosystems to assist with the management and recovery of Pacific salmon (*Oncorhynchus* spp.) and other NOAA trust resources. The program provides technical support to NOAA Fisheries policy makers and regulatory staff, and collaborates with other federal and state agencies (e.g., USGS, USFWS), tribes, and educational institutions on research and outreach related to the management of Pacific salmon and other diadromous fishes. NOAA Fisheries should consider developing and funding such a program, hosted by the Northeast Fisheries Science Center, to work with partners on the East Coast.
- Canada's Department of Fisheries and Oceans: In 2021, the DFO-NOAA Climate and Fisheries Sciences Collaboration Framework was developed to identify priority topics and issues of common interest and possible activities to address them over the next 3 years. The work plan is organized around 3 general thematic areas: 1) detecting climate-related changes/impacts (observations, monitoring); 2) understanding mechanisms of climate-related changes/impacts (research); and 3) forecasting and projecting and responding to climate-related changes (modeling and adaptation). In the Atlantic region, there is already a rich history of bilateral collaboration. The Canada-U.S. Ecosystem Science (CAUSES) Working Group's mission is to conduct ecosystem research to inform management advice for shared stocks in the eastern U.S. and Canadian marine ecosystems through communication, discussion, and shared expertise and tools between DFO and the NEFSC and other regional research groups. In 2021, CAUSES incorporated the North Atlantic component of the bilateral DFO-NOAA Climate and Fisheries Sciences Collaboration Framework. There is a corresponding DFO-NOAA OA collaboration group (Collaborative Framework for Joint DFO/NOAA Ocean Acidification Research and Monitoring) that operates independently of CAUSES.
- Tribal governments: In April 2022, NOAA climate scientists and fisheries managers participated in the New England Federal Partners-Tribal Climate Summit. This summit identified a number of needs, including several related to NOAA science and management. These include, among others, needs related to impact

assessments, monitoring systems, and improved coordination between natural resource managers and tribal representatives. We will continue to build these partnerships and explore how traditional ecological knowledge can inform and enrich this work.

NERAP Priority Action 3: Continue to build industry-based fisheries and ocean observing capabilities.

Industry-based fisheries and ocean observing can be accomplished both through fishery-independent research (e.g., surveys) and fishery-dependent research (e.g., commercial catch and environmental data collection). The common thread in these approaches is the engagement of the fishing community in the development and design of research projects, execution of field work, and interpretation of results. The NEFSC Cooperative Research Branch leads a suite of collaborative surveys and research, but many other institutions and organizations across the northeast region support industry-based fisheries and ocean observing capabilities.

Industry-based surveys in the northeast region include, but are not limited to, the Maine/New Hampshire Trawl Survey, Northeast Area Monitoring and Assessment Program inshore trawl survey, scallop research-set-aside surveys, and the NEFSC Gulf of Maine Bottom Longline Survey. These surveys are conducted by teams of scientists and fishermen working onboard commercial fishing vessels. Each of these surveys collects fishery-independent data on the distribution and abundance of fish and invertebrate species and the environment in which they are living. Data from these industry-based surveys are used in stock assessments and, in many cases, provides information about data poor species (e.g., cusk [*Brosme brosme*] and thorny skate [*Amblyraja radiata*]) and their habitats.

Fishery-dependent ocean observing initiatives in the region include, but are not limited to, the NEFSC Study Fleet; the Commercial Fisheries Research Foundation's (CFRF) Lobster, Jonah Crab, and Black Sea Bass Research Fleets; the NEFSC environmental Monitors on Lobster Traps and Large Trawlers Program (eMOLT); the WHOI and CFRF's Shelf Research Fleet; and the NEFSC Industry-Based Biological Sampling Program (InBios). The Study Fleet engages fishermen in collecting fine-scale catch, effort, and environmental data during routine fishing practices to precisely characterize fishing effort, spatiotemporal trends in resources species catch, and associated environmental conditions (Jones et al. 2022). Study Fleet data have been used to develop habitat models for mackerel, butterfish (*Peprilus triacanthus*), and shortfin squid (*Illex illecebrosus*), and have been integrated in catch-per-unit effort indices in the summer flounder, scup (*Stenotomus chrysops*), haddock (*Melanogrammus aeglefinus*), and shortfin squid stock assessments. The CFRF's Lobster, Jonah Crab, and Black Sea Bass Research Fleets apply a similar approach but instead focus fishermen's efforts on collecting biological (e.g., size, sex) data from commercial catch, as well as paired bottom water temperatures (Mercer et al. 2018). These data are used to characterize commercial catch for stock assessments and to understand environmental drivers of life history characteristics and population dynamics. The WHOI/CFRF Shelf Research Fleet and eMOLT focus on leveraging fishermen's time on the water to collect oceanographic data (i.e. temperature, depth, salinity) from across the northeast region throughout the year (Manning et al. 2009; Gawarkiewicz and Mercer 2019). These data provide a more complete picture of the seasonal and fine-scale dynamics of the subsurface ocean environment than traditional semi-annual surveys. Data from these programs feed into regional oceanographic models (ROMs, Finite Volume Community Ocean Model [FVCOM]) and can be paired with fishery catch and survey data to understand environmental drivers of resources species, climate impacts, and other factors.

Finally, InBios engages the fishing industry in collecting fish and invertebrate samples from areas and times of year otherwise not accessible to scientists but important for understanding life history. In this way, InBios engages the fishing industry in data gaps related to age, growth, and maturity of species, which are impacted by a changing climate.

The potential for industry vessels to collect ecological and oceanographic data could increase observing capacity in the region by at least an order of magnitude and provide critical observations of the water column, near surface atmosphere, and resource species. These observations can contribute to ocean modeling and prediction, but can also help fishermen make decisions with regard to limiting their incidental catch and their ability to adapt to changing conditions. Facilitating these interactions in short-term (days to years) applications would help develop the relationships necessary to make adaptive decisions in the medium term (years to decades).

NERAP Priority Action 4: Continue production of the NEFSC State of the Ecosystem reports and other related products that include climate-relevant information that is useful to management, such as historical, forecasted, and projected biophysical conditions; marine heatwaves; species distribution and abundance shifts; biogeochemical indices; coastal community vulnerability to sea level rise and storm surge; and fishing community vulnerability due to dependence on climate-vulnerable species.

The NEFSC Ecosystem Status Report, Ecosystem Advisories, and State of the Ecosystem reports meet one of the immediate-term actions defined in the NCSS. These products provide information on the current and past states of the Northeast U.S. Continental Shelf ecosystem. The information in these products is also provided to the NEFMC and MAFMC in State of the Ecosystem reports designed specifically for the Councils. The report draws on information collected across the NEFSC and academic partners from oceanographic to social indicators.

Due to the long, data-rich time series of the NEFSC bottom trawl survey, the majority of climate-fisheries research in the region has focused on the effects of warming ocean temperature on species distribution shifts. While these studies are important and need to continue, there is also a need for research focused on process-based (e.g., food availability, growth, mortality, species interactions) distribution shifts. Understanding the synergistic impacts of warming temperature and OA on species distributions and abundance is also a critical research need. Tracking, forecasting, and projecting species abundance is not a simple task, but progress is needed to help inform management.

Research should be conducted using other datasets including other NEFSC surveys, state surveys, Northeast Area Monitoring and Assessment Program (NEAMAP) surveys, Canadian DFO surveys, and SEFSC surveys. In addition, cooperative work with industry is underway and will be continued (e.g., NEFSC Observer Program, Study Fleet, Cooperative Catchability studies). Tagging data should also be incorporated into this effort where appropriate. Changes in the distribution of commercial and recreational catches and discards should also be examined as spatial changes in fishing may have important implications for assessments and management. Further, most work has focused on adult stages, and thus work should be conducted on understanding distribution changes of early life stages: eggs to juveniles. In particular, the connections between life stages through the availability of appropriate habitat should be examined. Finally, most work has been completed on commercially exploited fish and invertebrates, so emphasis should also be given to other species including recreationally important fish, protected species, and forage species.

4.4.2 Understand mechanisms and project future conditions (NCSS Objectives 4-5)

NERAP Priority Action 5: Conduct laboratory and field research on the mechanistic effects of multiple climate factors (e.g., temperature, ocean acidification, dissolved oxygen) on living marine resources with the goal of informing process-based models for single species, multispecies, and the ecosystem.

A mechanistic understanding of the effect of climate change on behavioral, physiological, ecological, and biophysical processes is critical to inform process-based models and improve scientific advice to managers. The NEFSC currently has seawater laboratory facilities in Sandy Hook, NJ, and Milford, CT. Both facilities have the ability to manipulate temperature, carbonate chemistry, and other factors and the ability to examine interactive effects of multiple stressors. Scientists at these facilities have experience working with phytoplankton, mollusks, crustaceans, and fish. Joint investments by the NOAA OAR OA Program and the NEFSC are supporting climate-related work at these facilities focused on the effect of OA on the all life stages of fish and mollusks, including biochemical, physiological, behavioral, and ecological responses.

Most work has focused on the larval stage, but more focus on all life stages from egg to adult should be investigated. These experiments should focus on collecting physiological data that can be used in DEB models, individual-based models, and ecosystem-based models. In particular, the potential for adaptation/acclimation to climate change needs to be investigated. Laboratory experiments allow for determining the extreme endpoint of the organism, but these experiments should be conducted concurrently with field sampling.

Laboratory and field experiments will provide much needed information to: 1) understand the effects of OA and other environmental changes on marine bivalves (e.g., oysters, surf clams, sea scallops, and bay scallops [*Argopecten irradians*]) and finfish in the New England and Mid-Atlantic regions; 2) understand biological response and potential for adaptation (genetic/acclimation/plasticity) of marine bivalves; and 3) provide data on biological and ecological processes (e.g., growth, consumption, metabolism) needed by modelers to improve predictions of long-term effects that will assist managers.

NERAP Priority Action 6: Work with NOAA OAR, NWS, NOS, and academic partners to develop and improve regional hindcasts, forecasts, and projections of ocean and estuarine/river physics and biogeochemistry to develop and improve climate-ready management of living marine resources.

The recent NOAA CEFI addresses the ocean modeling and communities of practice component of this priority action, but permanent funding for the CEFI has not yet been approved. However, funding has been provided for pilot studies to begin the development and validation of regional ocean model simulations for the entire U.S. east coast, shelf, and slope seas using NOAA GFDL's state-of-art ocean model MOM6 (Adcroft et al. 2019) coupled to GFDL's biogeochemical model COBALT (Stock et al. 2020). NOAA OAR CPO has also funded the development of annual to decadal ocean forecasts using the SCOAR modeling system.

The CEFI is primarily focused on regional ocean modeling that doesn't resolve the small-scale dynamics of estuaries and rivers, which are critical habitat for many recreational, commercial, and protected species in the region. Therefore, there is a continued need for new and improved watershed model development (Georges et al. 2016; Bever et al. 2021), as well as statistical downscaling efforts that utilize atmospheric variables as proxies for historical, forecasted, and projected watershed conditions (Muhling et al. 2018; Ross et al. 2020; Ross et al. 2021).

There is a critical need to continue these pilot studies and start new research that improves historical, forecasted, and projected biophysical ocean, estuary, and river conditions in the region. Work is needed to test and validate these new regional models and assess model uncertainty. An end goal is to have an ensemble of regional model simulations, much like the model assessment of the Intergovernmental Panel on Climate Change (IPCC), that can be used to assess model uncertainty in historical, forecasted, and projected ocean and watershed conditions. Building a capacity to distribute model output to various communities of practice is also a priority and a major component of the CEFI. There is a need for a centralized database that maintains and distributes in situ ocean data, watershed data, ocean reanalysis, ocean color data, and climate and ocean model output, which is also linked to NERAP Priority Action 2 and will require collaboration with other NOAA Line Offices. This modeling effort needs to be linked to NOAA Fisheries scientific-advice processes.

4.4.3 Inform management (NCSS Objectives 1-3)

NERAP Priority Action 7: Improve spatial management of living marine resources through an increased utility of spatial and temporal distributions, abundance, productivity, migration, and phenology in management decisions.

Research is needed that translates observed and forecasted shifts in species distributions, abundance, productivity, migration, and phenology into management advice. Contemporary and projected species distribution shifts have been one of the more predominant climate-fisheries research themes in the northeast U.S. However, stock structure, which is largely defined spatially, needs to be reevaluated in light of documented distribution changes. Process-based models of species distributions at multiple life history stages are also needed that explore impacts of climate change that are not exclusive to ocean temperature effects on adults. Other variables that need to be considered are OA, species interactions, fishery mortality, dissolved oxygen (coastal species), and food availability. Species distribution models that estimate abundance or biomass are not common because they are difficult to develop and validate. However, management decisions are based on abundance estimates, and thus understanding the spatial structure of a stock biomass historically and into the future is an essential component of climate-ready fisheries management.

NERAP Priority Action 8: Develop and use Vulnerability Analyses, Scenario Planning, and Management Strategy Evaluations to examine the effects of different management strategies under various climate change scenarios.

MSE is a simulation technique that allows the evaluation of a range of management options and identifies tradeoffs in performance across the range of options. An operating model is developed to represent the “true” dynamics of the system based on current understanding. An estimation model is used to assess the state of the system based on various observing or sampling processes. Finally, the effect of different management strategies can be examined in the context of the operating and estimation models. The NEFSC should continue to develop MSEs and seek external funding to apply the approach to climate-related issues. The NEFSC should also continue and expand work with academic scientists involved in MSE work in the region. Finally, the NEFSC and GARFO should continue to work with the NEFMC, MAFMC, ASMFC, and other offices within NOAA Fisheries to incorporate climate factors into management frameworks. The NEFSC is currently engaged in a Summer Flounder MSE for the MAFMC which focuses on translating discards into landings in order to increase the value generated by the fishery. One of the primary drivers of the MSE is climate change, and in particular, the shifting summer flounder distribution and the implications of state-level allocations based on historical fishing patterns. These types of projects should continue to be identified and supported.

CVAs can help to understand the sensitivity of marine species, freshwater and marine habitat, and fishing communities to climate change. These assessments help management prepare for future changes in marine ecosystems from lower trophic levels to keystone predators and protected species up through local seafood markets and fishing communities. NOAA Fisheries is currently collaborating with the New England Fisheries Management Council (NEFMC) and MAFMC on a synthesis product for the U.S. NES marine species and habitat CVAs to better understand the potential vulnerability of fish stocks to climate change. Further applications of these vulnerability assessments are needed to help inform management of high risk species and habitat under continued climate change. A CVA is also needed for Atlantic HMS, which requires collaborations with the NOAA Fisheries Atlantic Highly Migratory Species Division and the SEFSC.

Scenario planning offers a structured process that, amongst other outcomes, can help evaluate and prioritize actions associated with adapting to climate change. The NEFSC and GARFO should continue to develop scenario planning capabilities and continue to apply and/or support this framework as needs arise given its applicability to planning for a changing climate. Additionally, climate-related needs identified through completed scenario planning efforts should be tracked and furthered. These needs include assessing OA impacts on prey, conducting modeling studies (present conditions and projected into future) focused on spatial and temporal movement of right whales and copepods (e.g., current and future whale habitat use and distribution) and climate, and collecting long-term monitoring data on plankton.

The Northeast Regional Coordinating Council (NRCC) conducted a scenario planning project focusing on 2 components of climate change and its impact on fisheries management. The first focused on how climate change might affect stock distribution, availability, and other aspects of East Coast marine fisheries, and the second identified how those impacts will affect future governance and fisheries management. This effort is coastwide with the core team comprised of representatives from the various management bodies in the region (New England, Mid-Atlantic, and South Atlantic fisheries management councils, the ASMFC, and state agencies) and NOAA Fisheries (Greater Atlantic and Southern Atlantic regional offices, NEFSC, and NOAA headquarters). The project worked iteratively with stakeholders to develop a series of different scenarios that will then be used to develop a better understanding of the future challenges and opportunities facing fishery management along with a set of near-term and long-term management priorities under a range of different future conditions. In addition, the project developed policy recommendations for broader governance changes that should improve the ability to adapt to future scenarios. The project also generated a list of data gaps, research priorities, and monitoring needs for changing conditions along the east coast of the U.S.

NERAP Priority Action 9: Increase social, economic, and ecosystem scientist involvement in climate change research through multidisciplinary work—including the Northeast Integrated Ecosystem Assessment—that examines relationships between various ecosystem components with the goal of enhancing ecosystem-based management with climate information.

Ecosystems include humans and climate change acts on human communities both directly (e.g., sea level rise) and indirectly (e.g., species range shifts). There is an ongoing effort in the NEFSC to conduct multidisciplinary work in the northeast U.S. region that better integrates social and natural sciences. Major changes have been made to our NEUS Atlantis ecosystem model to better capture the dynamics of individual fisheries, and as a result, significant updates to the human dimension sub-models need to be made. These sub-models are specific to the fishing communities and socioeconomic characteristics of the northeast U.S., and dedicated social sciences research is

needed for model parameterization and validation. This multidisciplinary model development, and the background research to support it, will improve these end-to-end ecosystem model simulations and provide insight into the relationship between fishermen and fishing fleet behavior (and the underlying social, economic, and cultural motivations for behavior at both individual and community levels) and socioeconomic responses to changing ecosystem conditions due to climate change.

The development and potential use of Ecosystem and Socioeconomic Profiles (ESPs) is a critical component of this action. For example, the development of ESPs for black sea bass and bluefish are now part of the research track process. The goals of ESPs are to provide relevant ecosystem and socioeconomic information for fisheries management, work with management bodies to identify on-ramps where ESP information can fill knowledge gaps, and work toward an operational ecosystem approach to fisheries management (EAFM).

Shifting species distribution and other impacts of climate change also highlight the need for behavioral models of fishing activity in order to predict likely future responses to both changing drivers and management strategies. For example, state-level stock allocations in the Mid-Atlantic that are based on historical fishing patterns have become increasingly contentious given the shifts in stock distributions observed over the past decade. Effective management necessitates an understanding of how recreational and commercial fisheries are likely to respond to these dynamics in the future.

Additional human dimensions projects are needed in a number of areas, including OA projects that connect impacts to marine species to human community vulnerability; habitat studies that connect fishers' local ecological knowledge to climate studies of oceanographic and biological changes of habitat structure and function; and continued, new, and expanded work on MSEs and risk assessments with the MAFMC and the NEFMC. Some such projects are funded and at various stages of completion. Others, such as the habitat studies, are not yet funded.

It is important to understand not just the impacts of climate change in general on fishing communities but also the impacts of specific aspects of climate change. Sea level rise risk and storm surge risk indicators for fishing communities in the Northeast and other NMFS regions are available online. Depending on the species commonly caught there, human communities may be more or less vulnerable to changes in ocean temperature, OA, or both. Currently, a project is underway looking specifically at the impacts of OA on fishing communities that land species strongly impacted by OA, using an approach based on Hare et al. 2016b and Colburn et al. 2016. In another study that is reaching its conclusion, the NEFSC developed and tested a methodology to classify northeast U.S. fishing communities according to their vulnerability to specific climate change or climate change-related factors, including temperature, OA, and stock size and status.

Moreover, the vulnerability and resilience of fishing communities to the effects of ocean warming and OA on northeast species is dependent on the fishing community's adaptive capacity in relation to both social and environmental exposure and sensitivity factors. Measures of social well-being, sustainability, vulnerability, and resilience for fishing communities are already available. Viable measures of social well-being, sustainability, vulnerability, and resilience for the fishing industry would also be beneficial to coastal communities and have yet to be applied to OA or warming specifically.

NERAP Priority Action 10: Develop stock assessment models (e.g., the Woods Hole Assessment Model [WHAM]) that include environmental terms (e.g., temperature, ocean acidification) with a priority for stocks that have upcoming research track assessments.

Moving forward, multiple alternative stock assessment models and approaches need to be developed and evaluated. To be incorporated into operational assessments, these models and approaches need to undergo a formal scientific peer-review process. Assessments are prepared during a Northeast Regional Stock Assessment Workshop (SAW) and then reviewed by an independent panel of stock assessment experts called the Stock Assessment Review Committee (SARC). Further, both the ability to forecast environmental factors and better estimate historical environmental factors are necessary to include environmental terms in stock assessment models. An example of this approach is currently underway within the NCLIM project, which is a collaboration of the Gulf of Maine Research Institute, NEFSC, and Rutgers University. The approach focuses on stocks that have upcoming research track assessments and combines climate models, ecosystem/population models, and human dimension models to help inform the management process with a climate-informed assessment (Figure 5). A Northeast Ecosystem and Socioeconomic Profile Workshop is planned, with the aim to develop a flexible, standardized framework that helps integrate ecosystem and socioeconomic factors into fisheries decision-making. The framework is intended to provide supplemental data that will directly inform both the stock assessment process and the scientific advice for fishery management.

The WHAM is a tool being developed for the region that can help address this priority action. The WHAM currently has the capability to link recruitment trends or natural mortality to environmental covariates and carry these relationships and other correlated processes forward into projections. The research track working group for Atlantic butterfish is also currently considering climate effects on the distribution of the stock, and the recently formed research track working group on state-space assessment models will be exploring WHAM and its applications in climate-enhanced stock assessments for peer review in the fall of 2023.

5. METRICS FOR MEASURING SUCCESS

In response to a report from the Government Accountability Office (GAO-16-827) on the initial Regional Action Plans for climate science, the metrics used to measure success in implementing this NERAP are selected to be concrete, quantifiable, and time-bound. They are detailed in Table 1. Broadly, metrics fall into the following categories: reports and peer-reviewed publications, computational models, biophysical sampling/data collection, planning and coordination meetings, laboratory studies, and management advice. These metrics will be used in annual progress updates to the NOAA Fisheries Science Board.

FIGURES AND TABLE




Figure 1. The NOAA Fisheries Climate Science Strategy is organized around the 7 priority science objectives.




Figure 2. Number of Northeast Fisheries Science Center, Greater Atlantic Regional Fisheries Office, and Chesapeake Bay Office peer-reviewed publications from 2016-2020 that address the 7 NOAA Fisheries Climate Science Strategy objectives and 15 Northeast Regional Action Plan (NERAP) priorities.




Figure 3. Number of ocean stations sampled each year by NOAA Northeastern Fisheries Science Center surveys (e.g., Ecosystem Monitoring; bottom trawl; Marine Resources Monitoring, Assessment, and Prediction; scallop; protected species).




Figure 4. Directional effect of climate change from Hare et al. 2016b. Colors represent expected negative (red), neutral (tan), and positive (green) effects. Certainty in score is denoted by text font and text color: very high certainty (>95%, black, bold font), high certainty (90–95%, black, italic font), moderate certainty (66–90%, white or gray, bold font), low certainty (<66%, white or gray, italic font).




Figure 5. Schematic of the Northeast Climate Integrated Modeling (NCLIM) approach to informing management with climate-enhanced stock assessment information for the upcoming research track assessments for Atlantic cod (*Gadus morhua*), southern New England yellowtail flounder (*Limanda ferruginea*), and black sea bass (*Centropristis striata*).

Table 1. Tracking NERAP progress.

Objective	NERAP Priority Action	Metric	
<p align="center">Maintaining Infrastructure (NCSS Objectives 6-7)</p>	<p>NERAP Priority Action 1: Maintain and expand ecosystem survey and data collection efforts in the Northeast U.S. Continental Shelf ecosystem including the Bottom Trawl Survey, Ecosystem Monitoring Program, Sea Scallop Survey, Northern Shrimp Survey, Clam Survey, Gulf of Maine Bottom Longline Survey, and Protected Species Surveys. This also includes the Gulf of Maine longline survey, data poor species surveys, right whale prey sampling, ocean acidification monitoring, and the cooperative shark tagging program.</p>	<p>Maintain existing physical and biological surveys. Increase the number of ocean observations for both biological and physical variables. Improved monitoring of ocean acidification through collaborations with academic partners that are using slocum gliders to measure pH and aragonite saturation throughout the water column.</p>	
	<p>NERAP Priority Action 2: Coordinate with other NOAA Programs, Line Offices, and partners to link living marine resource data, science, and management to climate science and research activities.</p>	<p>NOAA Integrated Ecosystem Assessment Program</p>	<p>NOAA Chesapeake Bay Office (Data sharing, SOE reporting, research collaborations)</p>
		<p>NOAA Ocean Acidification Program (proposals, research priorities)</p>	<p>NOAA Ocean and Atmospheric Research, National Ocean Service (NOS), and National Weather Service (NWS) (modeling collaborations)</p>
		<p>NOAA Fisheries Atlantic Highly Migratory Species Division</p>	<p>State agencies (survey data sharing and coordination)</p>
		<p>Academic and non-governmental organizations (research proposals, workshops)</p>	<p>Watershed Program for the East Coast</p>
		<p>Canada Department of Fisheries and Oceans (DFO) (Enhanced survey data coordination, research collaboration, joint use of new ocean models, workshops)</p>	<p>Tribal governments</p>
		<p>NERAP Priority Action 3: Continue to build industry-based fisheries and ocean observing capabilities.</p>	<p>Maintain existing relationships and build new relationships with the fishing industry to continue the collection of fisheries-dependent biological and physical data.</p>
		<p>NERAP Priority Action 4: Continue production of the NEFSC State of the Ecosystem reports and other related products that include climate relevant information that is useful to management such as historical, forecasted, and projected biophysical conditions, marine heatwaves, species distribution and abundance shifts, biogeochemical indices, coastal community vulnerability to sea level rise and storm surge, and fishing community vulnerability due to dependence on climate-vulnerable species.</p>	<p>Production of annual State of the Ecosystem reports that include existing and new climate indicators that are relevant to fisheries and protected species management. Work with management councils to include new climate-relevant indicators that help inform annual management advice.</p>

Objective	NERAP Priority Action	Metric
<p>Tracking and projecting change, understanding mechanisms (NCSS Objectives 4-6)</p>	<p>NERAP Priority Action 5: Conduct laboratory and field research on the mechanistic effects of multiple climate factors (e.g. temperature, ocean acidification, dissolved oxygen) on living marine resources with the goal of informing process-based models for single species, multi-species, and the ecosystem.</p>	<p>Process-based research on the impacts of temperature and OA on finfish and invertebrates (multiple life stages). Utilization of laboratory results to inform species process-based models</p>
	<p>NERAP Priority Action 6: Work with NOAA Oceanic and Atmospheric Research, National Weather Service, National Ocean Service, and academic partners to develop and improve regional hindcasts, forecasts, and projections of ocean and estuarine/river physics and biogeochemistry to develop and improve climate-ready management of living marine resources.</p>	<p>Improved ocean hindcasts, forecasts, and projections for the U.S. northeast shelf that include both physical and biogeochemical variables. More skillful hindcasts and forecasts of ocean variables such as SST and bottom temperature. High-resolution decadal scale projections that include biogeochemistry.</p>
	<p>NERAP Priority Action 7: Improve spatial management of living marine resources through an increased utility of spatial and temporal distributions, abundance, productivity, migration, and phenology in management decisions.</p>	<p>Test, refine, develop, and where appropriate, transition into operations, multi-spectral and hyperspectral remote sensing models and products for NOAA Fisheries and CoastWatch. This will include the compilation of a comprehensive dataset by data mining existing observations and collection of additional observations on planned NEFSC Ecosystem Monitoring cruises. Where feasible, research will be in synergy with NASA PACE activities.</p>
	<p>NERAP Priority Action 8: Develop and use Vulnerability Analyses, Scenario Planning, and Management Strategy Evaluations to examine the effect of different management strategies under various climate change scenarios.</p>	<p>Spatial metrics incorporated into management advice for fisheries, protected species, and habitat either qualitatively or quantitatively.</p>
<p>Informing Management (NCSS Objectives 1-3)</p>	<p>NERAP Priority Action 9: Increase social, economic, and ecosystem scientist involvement in climate change research through multidisciplinary work, including the Northeast Integrated Ecosystem Assessment, that examines relationships between various ecosystem components with the goal of enhancing ecosystem-based management with climate information.</p>	<p>Application of species, habitat, and social vulnerability analyses into management advice or use.</p> <p>Development of new scenario planning activities.</p> <p>Development of management strategy evaluations that consider climate change scenarios.</p>
	<p>NERAP Priority Action 10: Development of stock assessment models (e.g. WHAM) that include environmental terms (e.g., temperature, ocean acidification) with a priority for stocks that have upcoming research track assessments.</p>	<p>Human dimension and ecosystem variables incorporated into climate science research and management advice or use.</p>
	<p>NERAP Priority Action 10: Development of stock assessment models (e.g. WHAM) that include environmental terms (e.g., temperature, ocean acidification) with a priority for stocks that have upcoming research track assessments.</p>	<p>Development of stock assessment models or variables that include environmental terms with a focus on stocks that up for research track assessments.</p>
	<p>NERAP Priority Action 10: Development of stock assessment models (e.g. WHAM) that include environmental terms (e.g., temperature, ocean acidification) with a priority for stocks that have upcoming research track assessments.</p>	<p>Incorporation of qualitative or quantitative climate information into management advice for a fishery or protected species. Inform research track assessments with new information on climate variables.</p>

REFERENCES

- Adcroft A, Anderson W, Balaji V, Blanton C, Bushuk M, Dufour CO, Dunne JP, Griffies SM, Hallberg R, Harrison MJ, et al. 2019. The GFDL Global Ocean and Sea Ice Model OM4.0: model description and simulation features. *J Adv Model Earth Syst.* 11:3167-3211.
- Bell RJ, Wood A, Hare J, Richardson D, Manderson J, Miller T. 2018. Rebuilding in the face of climate change. *Can J Fish Aquat Sci.* 75:1405-1414.
- Bever AJ, Friedrichs MAM, St-Laurent P. 2021. Real-time environmental forecasts of the Chesapeake Bay: model setup, improvements, and online visualization. *Environ Model Softw.* 140:105036.
- Borggaard DL, Dick DM, Star J, Alexander M, Bernier M, Collins M, Damon-Randall K, Dudley R, Griffis R, Hayes S, et al. 2019. Atlantic Salmon (*Salmo salar*) Scenario Planning Pilot Report. US Dept Commer Greater Atlantic Regional Fisheries Office Policy Series 19-05. 89 p.
- Borggaard DL, Dick DM, Star J, Zoodsma B, Alexander MA, Asaro MJ, Barre L, Bettridge S, Burns P, Crocker J, et al. 2020. North Atlantic Right Whale (*Eubalaena glacialis*) Scenario Planning Summary Report. US Dept Commer NOAA Tech Memo NMFS-OPR-68. 88 p.
- Caesar L, Rahmstorf S, Robinson A, Feulner G, Saba V. 2018. Observed fingerprint of a weakening Atlantic Ocean overturning circulation. *Nature.* 556:191-196.
- Cao J, Thorson JT, Richards RA, Chen Y. 2017. Spatiotemporal index standardization improves the stock assessment of northern shrimp in the Gulf of Maine. *Can J Fish Aquat Sci.* 74:1781-1793.
- Chen Z, Kwon YO, Chen K, Fratantoni P, Gawarkiewicz G, Joyce TM. 2020. Long-term SST variability on the Northwest Atlantic Continental Shelf and Slope. *Geophys Res Lett.* 47.
- Chen Z, Kwon YO, Chen K, Fratantoni P, Gawarkiewicz G, Joyce TM, Miller TJ, Nye JA, Saba VS, Stock BC. 2021. Seasonal prediction of bottom temperature on the Northeast U.S. Continental Shelf. *J Geophys Res Oceans.* 126(5):e2021JC017187.
- Clay PM, Howard J, Busch DS, Colburn LL, Himes-Cornell A, Rumrill SS, Zador SG, Griffis RB. 2020. Ocean and coastal indicators: understanding and coping with climate change at the land-sea interface. *Clim Change.* 163:1773-1793.
- Colburn LL, Jepson M, Weng C, Seara T, Weiss J, Hare JA. 2016. Indicators of climate change and social vulnerability in fishing dependent communities along the Eastern and Gulf Coasts of the United States. *Mar Pol.* 74:323-333.
- Collins MJ. 2019. River flood seasonality in the Northeast United States: characterization and trends. *Hydrol Process.* 33:687-698.

- Collins MJ, Hodgkins GA, Archfield SA, Hirsch RM. 2022. The occurrence of large floods in the United States in the modern hydroclimate regime: seasonality, trends, and large-scale climate associations. *Water Resour Res.* 58(2):e2021WR030480.
- [DOC] Department of Commerce. 2021. Department of Commerce 2021 Climate Action Plan for Adaptation and Resilience. Washington (DC): DOC.
- DePiper G, Gaichas S, Muffley B, Ardini G, Brust J, Coakley J, Dancy K, Elliott GW, Leaning DC, Lipton D, et al. 2021. Learning by doing: collaborative conceptual modelling as a path forward in ecosystem-based management. *ICES J Mar Sci.* 78:1217-1228.
- Friedland KD, McManus MC, Morse RE, Link JS. 2019. Event scale and persistent drivers of fish and macroinvertebrate distributions on the Northeast US Shelf. *ICES J Mar Sci.* 76:1316-1334.
- Friedland KD, Morse RE, Manning JP, Melrose DC, Miles T, Goode AG, Brady DC, Kohut JT, Powell EN. 2020. Trends and change points in surface and bottom thermal environments of the US Northeast Continental Shelf Ecosystem. *Fish Oceanogr.* 29:396-414.
- Gaichas SK, DePiper GS, Seagraves RJ, Muffley BW, Sabo MG, Colburn LL, Loftus AJ. 2018. Implementing ecosystem approaches to fishery management: risk assessment in the US Mid-Atlantic. *Front Mar Sci.* 5:442.
- Gaichas SK, Seagraves RJ, Coakley JM, DePiper GS, Guida VG, Hare JA, Rago PJ, Wilberg MJ. 2016. A framework for incorporating species, fleet, habitat, and climate interactions into fishery management. *Front Mar Sci.* 3:105.
- Gawarkiewicz G and Mercer AM. 2019. Partnering with fishing fleets to monitor ocean conditions. *Ann Rev Mar Sci.* 11:391-411.
- Gawarkiewicz G, Chen K, Forsyth J, Bahr F, Mercer AM, Ellertson A, Fratantoni P, Seim H, Haines S, Han L. 2019. Characteristics of an advective marine heatwave in the Middle Atlantic Bight in early 2017. *Front Mar Sci.* 6:712.
- Georgas N, Yin L, Jiang Y, Wang Y, Howell P, Saba V, Schulte J, Orton P, Wen B. 2016. An open-access, multi-decadal, three-dimensional, hydrodynamic hindcast dataset for the Long Island sound and New York/New Jersey Harbor estuaries. *J Mar Sci Eng.* 4.
- Gonçalves Neto A, Langan JA, Palter JB. 2021. Changes in the Gulf Stream preceded rapid warming of the Northwest Atlantic Shelf. *Commun Earth Environ.* 2.
- Greenan BJW, Shackell NL, Ferguson K, Greyson P, Cogswell A, Brickman D, Wang Z, Cook A, Brennan CE, Saba VS. 2019. Climate change vulnerability of American lobster fishing communities in Atlantic Canada. *Front Mar Sci.* 6:579.

- Grieve BD, Hare JA, Saba VS. 2017. Projecting the effects of climate change on *Calanus finmarchicus* distribution within the U.S. Northeast Continental Shelf. *Sci Rep.* 7:6264.
- Hare JA, Blythe B, Ford K, Hooker B, Jensen B, Lipsky A, Nachman C, Pfeiffer L, Rasser M, Renshaw K. 2022. NOAA Fisheries and BOEM Federal Survey Mitigation Implementation Strategy – Northeast U.S. region. US Dept Commer Northeast Fish Sci Cent Tech Memo 292. 29 p.
- Hare JA, Borggaard DL, Friedland KD, Anderson J, Burns P, Chu K, Clay PM, Collins MJ, Cooper P, Fratantoni PS, et al. 2016a. Northeast Regional Action Plan – NOAA Fisheries Climate Science Strategy US Depart Commer Northeast Fish Sci Cent Tech Memo 239. 96 p.
- Hare JA, Morrison WE, Nelson MW, Stachura MM, Teeters EJ, Griffis RB, Alexander MA, Scott JD, Alade L, Bell RJ, et al. 2016b. A vulnerability assessment of fish and invertebrates to climate change on the Northeast U.S. Continental Shelf. *PLOS ONE.* 11:e0146756.
- Hennen DR, Mann R, Munroe DM, Powell EN. 2018. Biological reference points for Atlantic surf clam (*Spisula solidissima*) in warming seas. *Fish Res.* 207:126-139.
- Holsman KK, Ianelli JN, Aydin K. 2019. 2019 Climate-enhanced multi-species Stock Assessment for walleye pollock, Pacific cod, and arrowtooth flounder in the Eastern Bering Sea. North Pacific Fisheries Management Council Bering Sea and Aleutian Islands Stock Assessment and Fishery Evaluation (SAFE) Report. Seattle (WA): Alaska Fisheries Science Center, National Marine Fisheries Service.
- Johnson TR, Beard K, Brady DC, Byron CJ, Cleaver C, Duffy K, Keeney N, Kimble M, Miller M, Moeykens S, et al. 2019. A social-ecological system framework for marine aquaculture research. *Sustainability.* 11(9):2522.
- Jones AW, Burchard KA, Mercer AM, Hoey JJ, Morin MD, Gianesin GL, Wilson JA, Alexander CR, Lowman BA, Duarte DG, et al. 2022. Learning from the study fleet: maintenance of a large-scale reference fleet for Northeast U.S. fisheries. *Front Mar Sci.* 9.
- Kleisner KM, Fogarty MJ, McGee S, Barnett A, Fratantoni P, Greene J, Hare JA, Lucey SM, McGuire C, Odell J, et al. 2016. The effects of sub-regional climate velocity on the distribution and spatial extent of marine species assemblages. *PLOS ONE.* 11:e0149220.
- Kleisner KM, Fogarty MJ, McGee S, Hare JA, Moret S, Perretti CT, Saba VS. 2017. Marine species distribution shifts on the U.S. Northeast Continental Shelf under continued ocean warming. *Prog Oceanogr.* 153:24-36.
- Lettrich MD, Asaro MJ, Borggaard DL, Dick DM, Griffis RB, Litz JA, Orphanides CD, Palka DL, Pendleton DE, Soldevilla MS. 2019. A method for assessing the vulnerability of marine mammals to a changing climate. US Dept Commer NOAA Tech Memo NMFS-F/SPO-196. 75 p.

- Lettrich MD, Dick DM, Fahy CC, Griffis RB, Haas HL, Jones TT, Kelly IK, Klemm D, Lauritsen AM, Sasso CR, et al. 2020. A method for assessing the vulnerability of sea turtles to a changing climate. US Dept Commer NOAA Tech Memo NMFS-F/SPO-211. 83 p.
- Link JS, Griffis R, Busch S, editors. 2015. NOAA Fisheries climate science strategy. US Dept Commer NOAA Tech Memo NMFS-F/SPO-155. 70p.
- Lombard P J, Dudley RW, Collins MJ, Saunders R, Atkinson E. 2021. Model estimated baseflow for streams with endangered Atlantic salmon in Maine, USA. *River Res Appl.* 37(9):1254-1264.
- Manning J and Pelletier E. 2009. Environmental Monitors on Lobster Traps (eMOLT): long-term observations of New England's bottom-water temperatures. *J Oper Oceanogr.* 2(1):25-33.
- McHenry J, Welch H, Lester SE, Saba VS. 2019. Projecting marine species range shifts from only temperature can mask climate vulnerability. *Glob Chang Biol.* 25:4208-4221.
- McManus MC, Hare JA, Richardson DE, Collie JS. 2018. Tracking shifts in Atlantic mackerel (*Scomber scombrus*) larval habitat suitability on the Northeast U.S. Continental Shelf. *Fish Oceanogr.* 27:49-62.
- Mercer AM, Ellertson A, Spencer D, Heimann T. 2018. Fishers fill data gaps for American lobster (*Homarus americanus*) and Jonah crab (*Cancer borealis*) in the Northeast USA. *Bull Mar Sci.* 94(3):1121-1135.
- Meseck SL, Mercaldo-Allen R, Kuropat C, Clark P, Goldberg R. 2018. Variability in sediment-water carbonate chemistry and bivalve abundance after bivalve settlement in Long Island Sound, Milford, Connecticut. *Mar Pollut Bull.* 135:165-175.
- Meseck SL, Sennefelder G, Krisak M, Wikfors GH. 2020. Physiological feeding rates and cilia suppression in blue mussels (*Mytilus edulis*) with increased levels of dissolved carbon dioxide. *Ecol Indic.* 117:106675.
- Miller AS, Shepherd GR, Fratantoni PS. 2016a. Offshore habitat preference of overwintering juvenile and adult black sea bass, *Centropristis striata*, and the relationship to year-class success. *PLOS ONE.* 11:e0147627.
- Miller TJ, Hare JA, Alade LA. 2016b. A state-space approach to incorporating environmental effects on recruitment in an age-structured assessment model with an application to southern New England yellowtail flounder. *Can J Fish Aquat Sci.* 73:1261-1270.
- Miller TJ, O'Brien L, Fratantoni PS. 2018. Temporal and environmental variation in growth and maturity and effects on management reference points of Georges Bank Atlantic cod. *Can J Fish Aquat Sci.* 75:2159-2171.

- Moore SS, Seavy NE, Gerhart M. 2013. Scenario planning for climate change adaptation: a guidance for resource managers. Petaluma (CA): Point Blue Conservation Science and California Coastal Conservancy; Oakland (CA): California Coastal Conservancy.
- Morse RE, Friedland KD, Tommasi D, Stock C, Nye J. 2017. Distinct zooplankton regime shift patterns across ecoregions of the U.S. Northeast continental shelf Large Marine Ecosystem. *J Mar Syst.* 165:77-91.
- Muhling BA, Gaitán CF, Stock CA, Saba VS, Tommasi D, Dixon KW. 2018. Potential salinity and temperature futures for the Chesapeake Bay using a statistical downscaling spatial disaggregation framework. *Estuaries Coast.* 41:349-372.
- Muhling BA, Jacobs J, Stock CA, Gaitan CF, Saba VS. 2017. Projections of the future occurrence, distribution, and seasonality of three *Vibrio* species in the Chesapeake Bay under a high-emission climate change scenario. *GeoHealth.* 1:278-296.
- [NMFS] National Marine Fisheries Service. 2021. Fisheries of the United States, 2019. US Dept Commer NOAA NMFS Current Fishery Statistics No. 2019. 164 p.
- O’Leary CA, Miller TJ, Thorson JT, Nye JA. 2018. Understanding historical summer flounder (*Paralichthys dentatus*) abundance patterns through the incorporation of oceanography-dependent vital rates in Bayesian hierarchical models. *Can J Fish Aquat Sci.* 76:1275-1294.
- O’Leary CA, Thorson JT, Miller TJ, Nye JA. 2020. Comparison of multiple approaches to calculate time-varying biological reference points in climate-linked population-dynamics models. *ICES J Mar Sci.* 77:930-941.
- Patel SH, Winton MV, Hatch JM, Haas HL, Saba VS, Fay G, Smolowitz RJ. 2021. Projected shifts in loggerhead sea turtle thermal habitat in the Northwest Atlantic Ocean due to climate change. *Sci Rep.* 11.
- Poach M, Munroe D, Vasslides J, Abrahamsen I, Coffey N. 2019. Monitoring coastal acidification along the U.S. East coast: concerns for shellfish production. *Bull Jap Fish Res Edu Agen.* 49:53-64.
- Pousse E, Poach ME, Redman DH, Sennefelder G, White LE, Lindsay JM, Munroe D, Hart D, Hennen D, Dixon MS, et al. 2020. Energetic response of Atlantic surf clam *Spisula solidissima* to ocean acidification. *Mar Pollut Bull.* 161:111740.
- Richardson DE, Carter L, Curti KL, Marancik KE, Castonguay M. 2020. Changes in the spawning distribution and biomass of Atlantic mackerel (*Scomber scombrus*) in the western Atlantic ocean over 4 decades. *Fish Bull.* 118:120-134.
- Ross AC, Stock CA, Adams-Smith D, Dixon KW, Findell KL, Saba VS, Vogt B. 2021. Anthropogenic influences on extreme annual streamflow into Chesapeake Bay from the Susquehanna River. *Bull Am Meteorol Soc.* 102(1):S25-S32.

- Ross AC, Stock CA, Dixon KW, Friedrichs MAM, Hood RR, Li M, Pegion K, Saba V, Vecchi GA. 2020. Estuarine forecasts at daily weather to subseasonal time scales. *Earth Space Sci.* 7.
- Saba VS, Griffies SM, Anderson WG, Winton M, Alexander MA, Delworth TL, Hare JA, Harrison MJ, Rosati A, Vecchi GA, et al. 2016. Enhanced warming of the Northwest Atlantic Ocean under climate change. *J Geophys Res Oceans.* 121:118-132.
- Schulte JA, Georgas N, Saba V, Howell P. 2017. Meteorological aspects of the eastern North American pattern with impacts on Long Island Sound salinity. *J Mar Sci Eng.* 5.
- Schulte JA, Georgas N, Saba V, Howell P. 2018. North Pacific influences on Long Island Sound temperature variability. *J Clim.* 31:2745-2769.
- Slesinger E, Andres A, Young R, Seibel B, Saba V, Phelan B, Rosendale J, Wieczorek D, Saba G. 2019. The effect of ocean warming on black sea bass (*Centropristis striata*) aerobic scope and hypoxia tolerance. *PLOS ONE.* 14:e0218390.
- Sorochan KA, Plourde S, Morse R, Pepin P, Runge J, Thompson C, Johnson CL. 2019. North Atlantic right whale (*Eubalaena glacialis*) and its food: (II) interannual variations in biomass of *Calanus* spp. on western North Atlantic shelves. *J Plankton Res.* 41:687-708.
- Stock BC and Miller TJ. 2021. The Woods Hole Assessment Model (WHAM): a general state-space assessment framework that incorporates time- and age-varying processes via random effects and links to environmental covariates. *Fish Res.* 240:10597.
- Stock CA, Dunne JP, Fan S, Ginoux P, John J, Krasting JP, Laufkötter C, Paulot F, Zadeh N. 2020. Ocean biogeochemistry in GFDL's Earth System Model 4.1 and its response to increasing atmospheric CO₂. *J Adv Model Earth Syst.* 12: e2019MS002043
- Stock CA, Pegion K, Vecchi GA, Alexander MA, Tommasi D, Bond NA, Fratantoni PS, Gudgel RG, Kristiansen T, O'Brien TD, et al. 2015. Seasonal sea surface temperature anomaly prediction for coastal ecosystems. *Prog Oceanogr.* 137:219-236.
- Turner KJ, Mouw CB, Hyde KJW, Morse R, Ciochetto AB. 2021. Optimization and assessment of phytoplankton size class algorithms for ocean color data on the Northeast U.S. continental shelf. *Remote Sens Environ.* 267:112729.
- Walsh HJ, Richardson DE, Marancik KE, Hare JA. 2015. Long-term changes in the distributions of larval and adult fish in the northeast U.S. shelf ecosystem. *PLOS ONE.* 10:e0137382.
- Xu H, Miller TJ, Hameed S, Alade LA, Nye JA. 2018. Evaluating the utility of the Gulf Stream Index for predicting recruitment of Southern New England-Mid Atlantic yellowtail flounder. *Fish Oceanogr.* 27:85-95.

Zhang S, Stock CA, Curchitser EN, Dussin R. 2019. A numerical model analysis of the mean and seasonal nitrogen budget on the Northeast U.S. Shelf. *J Geophys Res Oceans*. 124:2969-2991.

Procedures for Issuing Manuscripts in the Northeast Fisheries Science Center Reference Document (CRD) and the Technical Memorandum (TM) Series

The mission of NOAA's National Marine Fisheries Service (NMFS) is "stewardship of the nation's ocean resources and their habitat." As the research arm of the NMFS's Greater Atlantic Region, the Northeast Fisheries Science Center (NEFSC) supports the NMFS's mission by "conducting ecosystem-based research and assessments of living marine resources, with a focus on the Northeast Shelf, to promote the recovery and long-term sustainability of these resources and to generate social and economic opportunities and benefits from their use." Results of NEFSC research are largely reported in primary scientific media (e.g., anonymously peer-reviewed scientific journals). However, to assist itself in providing data, information, and advice to its constituents, the NEFSC occasionally releases its results in its own series.

NOAA Technical Memorandum NMFS-NE – This series is issued irregularly. The series typically includes: data reports of long-term field or lab studies of important species or habitats; synthesis reports for important species or habitats; annual reports of overall assessment or monitoring programs; manuals describing program-wide surveying or experimental techniques; literature surveys of important species or habitat topics; proceedings and collected papers of scientific meetings; and indexed and/or annotated bibliographies. All issues receive internal scientific review, and most issues receive technical and copy editing.

Northeast Fisheries Science Center Reference Document – This series is issued irregularly. The series typically includes: data reports on field and lab studies; progress reports on experiments, monitoring, and assessments; background papers for, collected abstracts of, and/or summary reports of scientific meetings; and simple bibliographies. Issues receive internal scientific review, and most issues receive copy editing.

CLEARANCE

All manuscripts submitted for issuance as CRDs must have cleared the NEFSC's manuscript/abstract/webpage review process. If your manuscript includes material from another work which has been copyrighted, you will need to work with the NEFSC's Editorial Office to arrange for permission to use that material by securing release signatures on the "NEFSC Use-of-Copyrighted-Work Permission Form."

For more information, NEFSC authors should see the NEFSC's online publication policy manual, "Manuscript/Abstract/Webpage Preparation, Review, & Dissemination: NEFSC Author's Guide to Policy, Process, and Procedure."

STYLE

The CRD series is obligated to conform with the style contained in the current edition of the United States Government Printing Office Style Manual; however, that style manual is silent on many

aspects of scientific manuscripts. The CRD series relies more on the CSE Style Manual. Manuscripts should be prepared to conform with both of these style manuals.

The CRD series uses the Integrated Taxonomic Information System, the American Fisheries Society's guides, and the Society for Marine Mammalogy's guide for verifying scientific species names.

For in-text citations, use the name-date system. A special effort should be made to ensure all necessary bibliographic information is included in the list of references cited. Personal communications must include the date, full name, and full mailing address of the contact.

PREPARATION

Once your document has cleared the review process, the Editorial Office will contact you with publication needs—for example, revised text (if necessary) and separate digital figures and tables if they are embedded in the document. Materials may be submitted to the Editorial Office as email attachments or intranet downloads. Text files should be in Microsoft Word, tables may be in Word or Excel, and graphics files may be in a variety of formats (JPG, GIF, Excel, PowerPoint, etc.).

PRODUCTION AND DISTRIBUTION

The Editorial Office will perform a copy edit of the document and may request further revisions. The Editorial Office will develop the inside and outside front covers, the inside and outside back covers, and the title and bibliographic control pages of the document.

Once the CRD is ready, the Editorial Office will contact you to review it and submit corrections or changes before the document is posted online. A number of organizations and individuals in the Northeast Region will be notified by e-mail of the availability of the document online.