




NOAA Technical Memorandum NMFS-SEFC-147

FISHERIES LIBRARY
75 VIRGINIA BEACH DR.
MIAMI, FL 33149

Bibliographies of the National Marine Fisheries Service's Assessments of Impacts of the Buccaneer Gas and Oil Field and of Brine Disposal from Salt Domes of the Strategic Petroleum Reserve.

**SOUTHEAST FISHERIES CENTER
GALVESTON LABORATORY**



OCTOBER 1984

U.S. DEPARTMENT OF COMMERCE
National Oceanic and Atmospheric Administration
National Marine Fisheries Service
Southeast Fisheries Center
Galveston Laboratory
Galveston, Texas 77550





NOAA Technical Memorandum NMFS-SEFC-147

Bibliographies of the National Marine Fisheries Service's Assessments of Impacts of the Buccaneer Gas and Oil Field and of Brine Disposal from Salt Domes of the Strategic Petroleum Reserve.

BY

Charles Wax Caillouet, Jr.

**U. S. DEPARTMENT OF COMMERCE
Malcolm Baldrige, Secretary**

**National Oceanic and Atmospheric Administration
John V. Byrne, Administrator**

**National Marine Fisheries Service
William G. Gordon, Assistant Administrator for Fisheries**

OCTOBER 1984

Technical Memorandums are used for documentation and timely communication of preliminary results, interim reports, or special-purpose information, and have not received complete formal review, editorial control, or detailed editing.

CONTENTS

Notice..... ii

Bibliography of the National Marine Fisheries Service's
Assessment of Impacts of Buccaneer Gas and Oil Field..... 1

Bibliography of the National Marine Fisheries Service's
Assessment of Impacts of Brine Disposal from Salt Domes
of the Strategic Petroleum Reserve..... 21

NOTICE

This Technical Memorandum should be cited as follows:

Caillouet, C. W., Jr. 1984. Bibliographies of the National Marine Fisheries Service's assessments of impacts of the Buccaneer gas and oil field and of brine disposal from salt domes of the Strategic Petroleum Reserve. NOAA Technical Memorandum NMFS-SEFC-147, ii + 35 p.

It can be obtained by writing to the National Technical Information Service, 5258 Port Royal Road, Springfield, Virginia 22161.

Bibliography of the National Marine Fisheries Service's

Assessment of Impacts of Buccaneer Gas and Oil Field

The environmental assessment of the Buccaneer gas and oil field in the northwestern Gulf of Mexico was conducted during 1975-1980. The project, funded by the Environmental Protection Agency through Interagency Agreement with NOAA/NMFS and managed by the NMFS Galveston Laboratory, was an inter-disciplinary investigation of environmental impacts of the Buccaneer gas and oil field located about 50 km south southeast of Galveston, Texas. The investigation covered sediments, suspended particulates, hydrocarbons, biocides, elemental sulfur, avian populations, benthic communities, microbial communities, fishes (pelagic, demersal and reef), macro-crustaceans, bio-fouling communities, currents and hydrography, trace metals, ichthyoplankton, and mathematical modeling. The journal articles, reports, dissertations and theses from this study are listed below (many of the reports are available from the National Technical Information Service, U.S.

Department of Commerce, 5285 Port Royal Road, Springfield, VA 22161:

Anderson, J. B., R. R. Schwarzer, and H. C. Clark. 1977. Sedimentology, geochemistry and trace metal analysis, p. 89-174. In: Jackson, W. B. (ed.), Environmental assessment of an active oil field in the northwestern Gulf of Mexico, 1976-1977. NOAA Annual Report to EPA, NTIS Accession No. PB 283890.

Anderson, J. B., R. R. Schwarzer, H. C. Clark, R. B. Wheeler, C. L.

- Hokanson, J. E. Anderson, and J. Okafor. 1979. Describe the fine sediments and nepheloid layer of the oil field, focusing upon their relationship to heavy metal adsorption/determine levels, pathways, and bioaccumulation of heavy metals in the marine ecosystem in the oil field, p. 2.3.2/2.4.2-1 to 2.3.2/2.4.2-118. In: Jackson, W. B. (ed.), Environmental assessment of an active oil field in the northwestern Gulf of Mexico, 1977-1978. Vol. III, Chemical and physical investigations. NOAA NMFS Annual report to EPA, NTIS Accession No. PB80 107899.
- Anderson, J. B., R. B. Wheeler, and R. R. Schwarzer. 1981. Sedimentologic and geochemical results of the Buccaneer oil/gas field study, p. 721-724. In: 1981 Oil Spill Conference (EPA, API, USCG), Atlanta, Georgia.
- Anderson, J. B., R. B. Wheeler, and R. R. Schwarzer. 1981. Sedimentology and geochemistry of recent sediments, p. 59-67. In: Middleditch, B. S. (ed.). Environmental effects of offshore oil production: the Buccaneer gas and oil field study. Marine Science Vol. 14, Plenum Press, New York, xvii + 446 p.
- Armstrong, R. S. 1980. Current patterns and hydrography of the Buccaneer field and adjacent waters. Vol. IV In: Jackson, W. B. and E. P. Wilkens (eds.), Environmental assessment of Buccaneer gas and oil field in the northwestern Gulf of Mexico, 1975-1980. NOAA Tech. Memo. NMFS-SEFC-50, xxvi + vi + 31 p., NTIS Accession No. PB81 188351.

- Armstrong, R. S. and R. C. Hamilton. 1979. Describe seasonal circulation patterns in the oil field, p. 2.3.9-1 to 2.3.9-250. In: Jackson, W. B. (ed.), Environmental assessment of an active oil field in the northwestern Gulf of Mexico, 1977-1978. Vol. III (Parts A & B), Chemical and physical investigations. NOAA NMFS Annual Report to EPA, NTIS Accession No. PB80 107899.
- Aumann, G. D. 1977. The effect of structures on migratory and local marine birds, p. 551-574. In: Jackson, W. B. (ed.), Environmental assessment of an active oil field in the northwestern Gulf of Mexico, 1976-1977. NOAA Annual Report to EPA, NTIS Accession No. PB 283890.
- Behrens, E. W. 1977. Total organic carbon and carbon isotopes of sediments, p. 737-759. In: Jackson, W. B. (ed.), Environmental assessment of an active oil field in the northwestern Gulf of Mexico, 1976-1977. NOAA Annual Report to EPA, NTIS Accession No. PB 283890.
- Boland, G. S. 1980. Morphological parameters of the barnacle, Balanus tintinnabulum antillensis, as indicators of physiological and environmental conditions. M. S. Thesis, Texas A&M University, College Station, Texas.
- Brooks, J. M., E. L. Estes, and W. H. Huang. 1980. Investigations of surficial sediments and suspended particulates at Buccaneer field. Vol. II In: Jackson, W. B. and E. P. Wilkens (eds.), Environ-

mental assessment of Buccaneer gas and oil field in the northwestern Gulf of Mexico, 1978-1979. NOAA Tech. Memo. NMFS-SEFC-36, xxiv + 261 p., NTIS Accession No. PB81 192403.

Brooks, J. M., E. L. Estes, D. A. Wiesenburg, C. R. Schwab, and H. A. Abdel-Reheim. 1980. Investigations of surficial sediments, suspended particulates and volatile hydrocarbons at Buccaneer gas and oil field. Vol. I In: Jackson, W. B. and E. P. Wilkens (eds.), Environmental assessment of Buccaneer gas and oil field in the northwestern Gulf of Mexico, 1975-1980. NOAA Tech. Memo. NMFS-SEFC-47, xxvi + xii + 89 p., NTIS Accession No. PB81 157778.

Caillouet, C. W., W. B. Jackson, G. R. Gitschlag, E. P. Wilkens, and G. M. Faw. 1981. Review of the environmental assessment of the Buccaneer gas and oil field in the northwestern Gulf of Mexico. Proc. Thirty-Third Annual Gulf and Caribb. Fish. Inst., p. 101-124.

Chang, E. S. 1978. Sterols in the marine food web. M.S. Thesis, University of Houston, Houston, Texas.

Chi-Hsin-Hsu. 1981. The mode of action of the bacteriocin produced by Vibrio harveyi strain SY. M.S. Thesis, University of Houston, Houston, Texas.

Collins, L. A., J. H. Finucane, and L. E. Barger. 1980. Description of larval and juvenile red snapper, Lutjanus campechanus. Fishery Bulletin, U.S. 77(4):965-974.

Danek, L. J. and M. S. Tomlinson. 1980. Currents and hydrography of the Buccaneer field and adjacent waters. Vol. VI In: Jackson, W. B. and E. P. Wilkens (eds.), Environmental assessment of Buccaneer gas and oil field in the northwestern Gulf of Mexico, 1978-1979. NOAA Tech. Memo. NMFS-SEFC-40, xxiv + v + 33 p., NTIS Accession No. PB81 192445.

Davis, J. 1981. The bacterial flora associated with the blue crab, Callinectes sapidus. M. S. Thesis, University of Houston, Houston, Texas.

Davis, J. W. and R. K. Sizemore. 1981. Non-selectivity of Rimler-Schotts medium for Aeromonas hydrophila in estuarine environments. Appl. Environ. Microbiol. 42:544-545.

Davis, J. W. and R. K. Sizemore. 1982. The incidence of Vibrio species associated with blue crabs (Callinectes sapidus) collected from Galveston Bay, Texas. Appl. Environ. Microbiol. 43:1092-1097.

Devereux, R. 1981. The role of plasmids in hydrocarbon utilization by marine Pseudomonas species. M. S. Thesis, University of Houston, Houston, Texas.

Devereux, R. and R. K. Sizemore. 1981. Incidence of degradative plasmids in hydrocarbon-utilizing bacteria isolated from the Gulf of Mexico, p. 409-414. In: Developments in Industrial Microbiology, Vol. 22, Society for Industrial Microbiology.

Devereux, R. and R. K. Sizemore. 1982. Plasmid incidence in marine bacteria isolated from petroleum polluted sites on different petroleum hydrocarbons. Mar. Pollut. Bull. 13(6):198-202.

Emiliani, D., K. N. Baxter, W. B. Jackson, J. Morgan, G. Zamora, and L. Lansford. 1977. Demersal finfish and macro-crustaceans collected in the vicinity of Buccaneer oil field during 1976-1977, p. 275-310. In: Jackson, W. B. (ed.), Environmental assessment of an active oil field in the northwestern Gulf of Mexico, 1976-1977. NOAA Annual Report to EPA, NTIS Accession No. PB 283890.

Faw, G. M. 1981. Crude oil biodegradation studies in Buccaneer gas and oil field. M.S. Thesis, University of Houston, Houston, Texas.

Finucane, J. H. 1977. Ichthyoplankton of the Buccaneer oil field off Galveston, Texas. Paper presented at the Fourth Biennial International Estuarine Research Conference, Mt. Pocono, Pa. (abstract published in proceedings).

Finucane, J. H. and L. A. Collins. 1977. Ichthyoplankton, p. 339-486. In: Jackson, W. B. (ed.), Environmental assessment of an active oil field in the northwestern Gulf of Mexico, 1976-1977. NOAA Annual Report to EPA, NTIS Accession No. PB 282890.

Finucane, J. H., L. A. Collins, and L. E. Barger. 1979. Determine the effects of discharges on seasonal abundance, distribution and composition of ichthyoplankton in the oil field, p. 2.3.6-1 to

- 2.3.6-157. In: Jackson, W. B. (ed.), Environmental assessment of an active oil field in the northwestern Gulf of Mexico, 1976-1977. Vol. II, Data management and biological investigations. NOAA Annual Report to EPA, NTIS Accession No. PB80 165970.
- Fotheringham, N. 1977. Effects of offshore oil field structures on their biotic environment: Benthos and plankton, p. 487-550. In: Jackson, W. B. (ed.), Environmental assessment of an active oil field in the northwestern Gulf of Mexico, 1976-1977. NOAA Annual Report to EPA, NTIS Accession No. PB 283890.
- Fucik, K. and I. T. Show. 1980. Sources, fate and effects modeling. Vol. IX In: Jackson, W. B. and E. P. Wilkens (eds.), Environmental assessment of Buccaneer gas and oil field in the northwestern Gulf of Mexico, 1978-1979. NOAA Tech. Memo. NMFS-SEFC-43, xxiv + v + 84 p. (plus 32 p. of Appendices), NTIS Accession No. PB82 161845.
- Gallaway, B. J. 1980. Pelagic, reef and demersal fishes and macrocrustaceans/biofouling communities. Vol II In: Jackson, W. B. and E. P. Wilkens (eds.), Environmental assessment of Buccaneer gas and oil field in the northwestern Gulf of Mexico, 1975-1980. NOAA Tech. Memo. NMFS-SEFC-48, xxvi + vi + 82 p. NTIS Accession No. PB81 188344.
- Gallaway, B. J., R. Howard, K. Green, and L. Martin. 1976. A study plan for the Buccaneer oil field assessment program. A report to

the NMFS Galveston Laboratory from LGL Limited-U.S., Inc., Bryan, Texas, 69 pp.

Galloway, B. J., M. F. Johnson, R. L. Howard, L. R. Martin, and G. S. Boland. 1979. A study of the effects of Buccaneer oil field structures and associated effluents on biofouling communities and the Atlantic spadefish (Chaetodipterus faber), p. 2.3.8-1 to 2.3.8-126. In: Jackson, W. B. (ed.), Environmental assessment of an active oil field in the northwestern Gulf of Mexico, 1977-1978. Vol. II, Data management and biological investigations. NOAA NMFS Annual Report to EPA, NTIS Accession No. PB80 165970.

Galloway, B. J. and F. J. Margraf. 1979. Simulation modeling of biological communities associated with a production platform in the Buccaneer oil and gas field, p. 2.5.1-1 to 2.5.1-67. In: Jackson, W. B. (ed.), Environmental assessment of an active oil field in the northwestern Gulf of Mexico, 1977-1978. Vol. II, Data management and biological investigations. NOAA NMFS Annual Report to EPA, NTIS Accession No. PB80 165970.

Galloway, B. J. and L. R. Martin. 1980. Effects of gas and oil field structures and effluents on pelagic and reef fishes, demersal fishes and macrocrustaceans. Vol. III In: Jackson, W. B. and E. P. Wilkens (eds.), Environmental assessment of Buccaneer gas and oil field in the northwestern Gulf of Mexico, 1978-1979. NOAA Tech. Memo. NMFS-SEFC-37, xxiv + 49 p., NTIS Accession No. PB81

192411.

Hada, H. 1981. A survey of the incidence and function of plasmids in marine Vibrio spp. Ph.D. Dissertation, University of Houston, Houston, Texas.

Hada, H. S., M. I. Kirchevsky, and R. K. Sizemore. 1983. The use of cluster analysis to help determine the function of plasmids in marine Vibrio species. *J. Microbiol. Methods* 1:229-237.

Hada, H. S. and R. K. Sizemore. 1981. Incidence of plasmids in marine Vibrio spp. isolated from an oil field in the northwestern Gulf of Mexico. *Appl. Environ. Microbiol.* 41:199-202.

Hada, H. S. and R. K. Sizemore. (in press). Plasmids in marine Vibrio spp.: Incidence and determination of potential functions using numerical taxonomic methods. In: Vibrios in the Environment, John Wiley & Sons, publishers.

Harper, D. E., Jr. 1977. Distribution and abundance of macrobenthic and meiobenthic organisms, p. 175-274. In: Jackson, W. B. (ed.), Environmental assessment of an active oil field in the northwestern Gulf of Mexico, 1976-1977. NOAA Annual Report to EPA, NTIS Accession No. PB 283890.

Harper, D. E., Jr., D. L. Potts, R. R. Salzer, R. J. Case, R. L. Jaschok, and C. M. Walker. 1981. Distribution and abundance of macrobenthic and meiobenthic organisms, p. 133-177. In:

Middleditch, B. S. (ed.), Environmental Effects of Offshore Oil Production: The Buccaneer gas and oil field study, Marine Science Vol. 14, Plenum Press, N.Y.

Harper, D. E., Jr., R. J. Scudato, and C. S. Giam. 1976. A preliminary environmental assessment of the Buccaneer oil/gas field (pilot study of the Buccaneer oil field; benthos and sediments). A report from Texas A&M University at Galveston to the National Marine Fisheries Service, Galveston, Texas (Contract No. 03-6-042-35110), 63 p. (plus 83 p. of appendices).

Hollaway, S. L., G. M. Faw, and R. K. Sizemore. 1980. The bacterial community composition of an active oil field in the northwestern Gulf of Mexico. Mar. Pollut. Bull. 11:153-156.

Holley, H. and K. Savastano. 1977. Data management system development and monitoring, p. 11-88. In: Jackson, W. B. (ed.), Environmental assessment of an active oil field in the northwestern Gulf of Mexico, 1976-1977. NOAA Annual report to EPA, NTIS Accession No. PB 283890.

Holley, H. and K. Savastano. 1979. Buccaneer oil field study data management and system development final report, p. 2.2.3-1 to 2.2.3-159. In: Jackson, W. B. (ed.), Environmental assessment of an active oil field in the northwestern Gulf of Mexico, 1977-1978. Vol. II, Data management and biological investigations. NOAA NMFS Annual Report to EPA, NTIS Accession No. PB80 165970.

- Howard, R. L., G. S. Boland, B. J. Gallaway, and G. D. Dennis. 1980. Effects of gas and oil field structures and effluents on fouling community production and function. Vol. V In: Jackson, W. B. and E. P. Wilkens (eds.), Environmental assessment of Buccaneer gas and oil field in the northwestern Gulf of Mexico, 1978-1979. NOAA Tech. Memo. NMFS-SEFC-39, xxiv + v + 60 p., NTIS Accession No. PB81 192437.
- Hoyt, P. 1981. Characterization of harveyicin and demonstration of competitive dominance in the bacteriocin producing strain of Beneckea harveyi; SY. M.S. Thesis, University of Houston, Houston, Texas.
- Hoyt, P. R. and R. K. Sizemore. 1982. Demonstration of competitive dominance by a bacteriocin producing strain of Vibrio harveyi. Appl. Environ. Microbiol. 44:(3):653-658.
- Jackson, W. B. (ed.). 1979. Synopsis. Vol. I In: Environmental assessment of an active oil field in the northwestern Gulf of Mexico, 1977-1978. NOAA NMFS Annual Report to EPA, xvii + 62 p., NTIS Accession No. PB 301339/AS.
- Jackson, W. B., K. N. Baxter, and C. W. Caillouet. 1978. Environmental assessment of the Buccaneer oil and gas field off Galveston, Texas: An overview. Proc. Offshore Technol. Conf., May, 1978, Houston, Texas, OTC 3081:277-284.
- Jackson, W. B., K. N. Baxter, and C. W. Caillouet. 1979. Environ-

mental assessment of an offshore oil field. *Ecolibrium* 8(1):
7-9.

Jackson, W. B. and E. P. Wilkens (eds.). 1980. Synopsis, Part A.
Vol. I In: Environmental assessment of Buccaneer gas and oil
field in the northwestern Gulf of Mexico, 1978-1979. NOAA Tech.
Memo. NMFS-SEFC-35, xxiii + 72 p., NTIS Accession No. PB82
161167.

Klima, E. F., C. Caillouet, N. Baxter, and W. Jackson. 1977.
Environmental assessment of an active oil field in the north-
western Gulf of Mexico, 1977-1978; a report of NMFS/LGL Workshop
I, 1977-1978. NMFS SEFC Galveston Laboratory, Galveston, Texas.

Martin, J. A. 1977. Hydrography of the Buccaneer oil field and
adjacent waters, p. 575-632. In: Jackson, W. B. (ed.),
Environmental assessment of an active oil field in the north-
western Gulf of Mexico, 1976-1977. NOAA Annual Report to EPA,
NTIS Accession No. PB 283890.

McCall, J. O. 1979. Isolation and characterization of a bac-
teriocinogenic strain of the bioluminescent marine bacterium
Beneckea harveyi. M.S. Thesis, University of Houston, Houston,
Texas.

McCall, J. O. and R. K. Sizemore. 1979. Description of a bac-
teriocinogenic plasmid in Beneckea harveyi. *Appl. Environ.*
Microbiol. 38(5):974-979.

- McKinney, L. 1979. Liljeborgiid amphipods from the Gulf of Mexico and Caribbean Sea. Bull. Mar. Sci. 29(2):140-154.
- McKinney, L. 1980. The genus Photis from the Texas coast with a description of a new species, Photis melanicus. Contrib. Mar. Sci. Univ. Texas (in press).
- Middleditch, B. S. 1980. Hydrocarbons, biocides, and sulfur. Vol. V
In: Jackson, W. B. and E. P. Wilkens, (eds.), Environmental assessment of Buccaneer gas and oil field in the northwestern Gulf of Mexico, 1975-1980. NOAA Tech. Memo. NMFS-SEFC-51, xxvi + ix + 70 p., NTIS Accession No. PB81 178451.
- Middleditch, B. S. (ed.). 1981. Environmental effects of offshore oil production: The Buccaneer gas and oil field study. Mar. Sci. Vol. 14, Plenum Press, New York, xvii + 446 p.
- Middleditch, B. S. 1982. Alkanes in shrimp from Buccaneer oil field. Bull. Environ. Toxicol. 29:18-23.
- Middleditch, B. S. 1982. Volatile constituents of the produced water effluent from the Buccaneer gas and oil field. J. Chromatog. 239:159-171.
- Middleditch, B. S. and B. Basile. 1976. Deuteriated analogs as internal standards for the quantitation of environmental alkanes by gas chromatography. Analyt. Letters 9(11):1031-1034.
- Middleditch, B. S. and B. Basile. 1979. Sulfur-34 as an internal

standard for quantitation of environmental sulfur by combined gas chromatography-mass spectrometry. *Analyt. Letters* 12(A7):777-781.

Middleditch, B. S. and B. Basile. 1980. Alkanes in benthic organisms from the Buccaneer oil field. *Bull. Environ. Contam. Toxicol.* 24:945-952.

Middleditch, B. S. and B. Basile. 1980. Discharge of elemental sulfur and its distribution in surficial sediments in the region of the Buccaneer oil field. *J. Chromatog.* 199:161-169.

Middleditch, B. S. and B. Basile. 1984. The mass spectrometer as a gas chromatographic detector: environmental applications. In: Nicholson, A. J. C. and I. G. McWilliams (eds.). *Proceedings of International Conference on Detectors and Chromatography*, May 30-June 3, 1983, Australian Scientific Industry Association, Melbourne, Australia.

Middleditch, B. S., B. Basile, and E. S. Chang. 1977a. Hydrocarbon levels and pathways in the major components of a marine ecosystem, p. 633-736. In: Jackson, W. B. (ed.), *Environmental assessment of an active oil field in the northwestern Gulf of Mexico*, 1976-1977. NOAA Annual Report to EPA, NTIS Accession No. PB 283890.

Middleditch, B. S., B. Basile, and E. S. Chang. 1977b. Environmental effects of offshore oil production: alkanes in the region of the Buccaneer oil field. *J. Chromatog.* 142:777-785.

- Middleditch, B. S., B. Basile, and E. S. Chang. 1978. Alkanes in surficial sediments from the region of the Buccaneer oil field. *J. Chromatog.* 158:449-463.
- Middleditch, B. S., B. Basile, and E. S. Chang. 1978. Discharge of alkanes during offshore oil production in the Buccaneer oil field. *Bull. Environ. Contam. Toxicol.* 20:59-65.
- Middleditch, B. S., B. Basile, and E. S. Chang. 1979. Alkanes in seawater in the vicinity of the Buccaneer oil field. *Bull. Environ. Contam. Toxicol.* 21:413-420.
- Middleditch, B. S., B. Basile, and E. S. Chang. 1982. Alkanes in shrimp from the Buccaneer oil field. *Bull. Environm. Contam. Toxicol.* 29:18-23.
- Middleditch, B. S., B. Basile, and S. R. Missler. 1979. Determine levels, pathways, and bioaccumulation of selected discharge constituents (non-metals) in the marine ecosystem in the oil field/hydrocarbon modeling, p. 2.4.1-1 to 2.4.1-302. In: Jackson, W. B. (ed.), Vol.III (Parts A & B), Chemical and physical investigations. Environmental assessment of an active oil field in the northwestern Gulf of Mexico, 1977-1978. NOAA NMFS Annual Report to EPA, NTIS Accession No. PB80 107889.
- Middleditch, B. S., E. S. Chang, and B. Basile. 1979a. Alkanes in plankton from the Buccaneer oil field. *Bull. Environ. Contam. Toxicol.* 21:421-427.

- Middleditch, B. S., E. S. Chang, and B. Basile. 1979b. Alkanes in barnacles (Balanus tintinnabulum) from the Buccaneer oil field. Bull. Environ. Contam. Toxicol. 23:6-12.
- Middleditch, B. S., E. S. Chang, B. Basile, and S. R. Missler. 1979. Alkanes in fish from the Buccaneer oil field. Bull. Environ. Contam. Toxicol. 22:249-257.
- Middleditch, B. S. and D. West. 1979. Bicyclohexyl interference in the quantitation of environmental alkanes by gas chromatography. Analyt. Letters 12(A12):1279-1284.
- Middleditch, B. S. and D. L. West. 1980. Hydrocarbons, biocides, and sulfur. Vol. VII In: Jackson, W. B. and E. P. Wilkens (eds.). Environmental assessment of Buccaneer gas and oil field in the northwestern Gulf of Mexico, 1978-1979. NOAA Tech. Memo. NMFS-SEFC-41, xxiii + 112 p., NTIS Accession No. PB81 225443.
- Mitchell, F. 1977. Accessible data base management, p. 7-9. In: Jackson, W. B. (ed.), Environmental assessment of an active oil field in the northwestern Gulf of Mexico, 1976-1977. NOAA Annual Report to EPA, NTIS Accession No. PB 283890.
- Olsen, K. D. and R. K. Sizemore. 1981. Effects of an established offshore oil platform on the autochthonous bacteria community, p. 685-694. In: Developments in Industrial Microbiology, Vol. 22, Society for Industrial Microbiology.

- Rose, C. D. and T. J. Ward. 1981. Principles of aquatic hazard evaluation as applied to ocean-disposed wastes. American Society for Testing and Materials, Special Technical Publication 737, Aquatic Toxicology: Fourth Conference, p. 138-158.
- Savastano, K. and H. Holley. 1980. Implement, monitor and modify data management system, Part B. Vol. I In: Jackson, W. B. and E. P. Wilkens (eds.), Environmental assessment of Buccaneer gas and oil field in the northwestern Gulf of Mexico, 1978-1979, NOAA Tech. Memo. NMFS-SEFC-35, xxiii + 72 p., NTIS Accession No. PB82 161167.
- Show, I. T., Jr. 1979. An application of compartmental models to mesoscale marine ecosystems, pages 73-97. In: Matis, J. H., B. C. Patten, and G. C. White (eds.), Compartmental Analysis of EcoSystem Models. Vol. 10. Statistical Ecology Series. Internat. Coop. Publ. House, Fairland, Maryland.
- Sizemore, R. K., S. L. Hollaway, and G. M. Faw. 1979. Determine and compare groups of bacteria from an active oil field and unaltered area, p. 2.3.7-1 to 2.3.7-41. In: Jackson, W. B. (ed.), Environmental assessment of an active oil field in the northwestern Gulf of Mexico, 1977-1978. Vol. II. Data management and biological investigations. NOAA NMFS Annual Report to EPA, NTIS Accession No. PB80 165970.
- Sizemore, R. K., G. H. Hsu, and K. D. Olsen. 1981. Bacterial commu-

nity composition and activity, p. 223-235. In: Middleditch, B. S. (ed.), Environmental Effects of Offshore Oil Production: The Buccaneer gas and oil field study. Mar. Sci. Vol. 14, Plenum Press, N.Y.

Sizemore, R. K. and K. Olsen. 1980. Bacterial communities. Vol. IV In: Jackson, W. B. and E. P. Wilkens (eds.), Environmental assessment of Buccaneer gas and oil field in the northwestern Gulf of Mexico, 1978-1979. NOAA Tech. Memo. NMFS-SEFC-38, xxiv + iv + 32 p., NTIS Accession No. PB81 192429.

Sizemore, R. K. and K. Olsen. 1980. Bacteriology of a Gulf of Mexico gas and oil field. Vol. III In: Jackson, W. B. and E. P. Wilkens (eds.), Environmental assessment of Buccaneer gas and oil field in the northwestern Gulf of Mexico, 1975-1980. NOAA Tech. Memo. NMFS-SEFC-49, xxvi + v + 21 p., NTIS Accession No. PB81 192189.

Smedes, G., J. Calman, and J. Beebe. 1980. Hydrodynamic modeling. Vol. X In: Jackson, W. B. and E. P. Wilkens (eds.), Environmental assessment of Buccaneer gas and oil field in the northwestern Gulf of Mexico, 1978-1979. NOAA Tech. Memo. NMFS-SEFC-44, xxiv + vi + 18 p. (plus 61 p. of Appendices), NTIS Accession No. PB81 193229.

Tillery, J. B. 1980. Trace metals. Vol. VIII. In: Jackson, W. B. and E. P. Wilkens (eds.). Environmental assessment of Buccaneer gas and oil field in the northwestern Gulf of Mexico, 1978-1979.

- NOAA Tech. Memo. NMFS-SEFC-42, xxiv + viii + 93 p., NTIS Accession No. PB81 192452.
- Tillery, J. B. 1980. Trace metals. Vol. VI. In: Jackson, W. B. and E. P. Wilkens (eds.), Environmental assessment of Buccaneer gas and oil field in the northwestern Gulf of Mexico, 1975-1980. NOAA Tech. Memo. NMFS-SEFC-52, xxvi + v + 39 p., NTIS Accession No. PB81 192197.
- Trent, L., I. K. Workman, S. Dime, and C. Jones. 1977. Recreational fisheries and the distribution of predatory pelagic fishes, p. 311-338. In: Jackson, W. B. (ed.), Environmental assessment of an active oil field in the northwestern Gulf of Mexico, 1976-1977. NOAA NMFS Annual Report to EPA, NTIS Accession No. PB 283890.
- West, D. L. 1979. Hydrocarbons, sulfur and acrolein in the marine ecosystem associated with offshore oil production. M.S. Thesis, University of Houston, Houston, Texas.
- Wheeler, R. B. 1979. Environmental trace metal geochemistry of the Buccaneer oil and gas field. M.S. Thesis, Rice University, Houston, Texas. 176 pp.
- Wheeler, R. B., J. B. Anderson, R. R. Schwarzer, and C. L. Hokanson. 1980. Sedimentary processes and trace metal contaminants in Buccaneer oil/gas field, northwestern Gulf of Mexico. Environ. Geol. 3:163-175.

Wheeler, R. B., R. R. Schwarzer, and J. B. Anderson. 1978.

Assessment of environmental impact of offshore production in the
Buccaneer oil field: sedimentologic and geochemical results.
Proceedings of the Offshore Technology Conference, May 1978,
Houston, Texas OTC 3082:285-290.

Workman, I. K. and C. E. Jones. 1979. Determine effects of oil field
discharges on species composition and abundance of pelagic fishes
and demersal fishes and macro-crustaceans in the oil field, p.
2.3.5-1 to 2.3.5-149. In: Jackson, W. B. (ed.), Environmental
assessment of an active oil field in the northwestern Gulf of
Mexico, 1977-1978. Vol. II. Data management and biological
investigations. NOAA NMFS Annual Report to EPA, NTIS Accession
No. PB80 165970.

Zein-Eldin, Z. P. and P. M. Keney. 1979. Bioassay of Buccaneer oil
field effluents with penaeid shrimp, p. 2.3.4-1 to 2.3.4-25. In:
Jackson, W. B. (ed.), Environmental assessment of an active oil
field in the northwestern Gulf of Mexico, 1977-1978. Vol. II.
Data management and biological investigations. NOAA NMFS Annual
Report to EPA, NTIS Accession No. PB80 165970.

Bibliography of the National Marine Fisheries Service's Assessment
of Impacts of Brine Disposal from Salt Domes of the
Strategic Petroleum Reserve

The Energy Policy and Conservation Act of 1975 (PL 94-163) required the development of a SPR (Strategic Petroleum Reserve) to avert future energy crises such as that experienced by the United States in the early 1970's. As a consequence, the DOE (U.S. Department of Energy) developed a SPR program that involved solution-mining of land-based salt domes in Texas and Louisiana to create caverns for the storage of petroleum. The program called for discharge of concentrated brine, produced as a by-product of solution-mining of salt domes, into the waters of the northwestern Gulf of Mexico. The EPA (Environmental Protection Agency), NMFS (National Marine Fisheries Service), the commercial shrimp industry, and environmental groups expressed concern regarding potential effects of brine disposal on shrimp (Penaeus spp.) and other biota of the northwestern Gulf of Mexico. As required by EIS's (Environmental Impact Statements), brine disposal monitoring plans were developed by DOE, EPA and NOAA (National Oceanic and Atmospheric Administration). These plans encompassed environmental impact research managed by NMFS and others.

Among the salt dome brine disposal sites originally proposed by DOE, only two are currently operational: the Bryan Mound brine disposal site off Freeport, Texas, and the West Hackberry brine disposal

site off Southwest Louisiana (Fig. 1). At the Bryan Mound brine disposal site, brine discharge was initiated on March 10, 1980. At the West Hackberry brine disposal site, brine discharge began May 10, 1981. It has been proposed by DOE that an additional brine disposal site called Big Hill be located off the upper coast of Texas, and it is projected to become operational in 1985.

The NMFS brine disposal impact assessment consisted of four major projects (Table 1) carried out during the years 1978-1982 by NMFS and its contractors (Table 2). The research, managed by the NMFS SEFC (Southeast Fisheries Center) Galveston Laboratory in Galveston, Texas, was funded by DOE through interagency agreements with NOAA. Program management, which covered the NMFS projects and other projects carried out by the NOS (National Ocean Survey) and DOE contractors, was provided by NOAA's NESDIS (National Environmental Satellite, Data and Information Service) in Washington, D.C. Data management for the NMFS projects was provided by the NMFS SEFC NFEL (National Fisheries Engineering Laboratory) in Bay St. Louis, Mississippi, the NMFS SEFC FIMD (Fisheries Information Management Division) in Miami, Florida, and the NESDIS. Project data files are archived by FIMD and NESDIS.

For the most part, the projects employed state-of-the-art fisheries assessment techniques. The rationale for such an approach was (1) that the results would be expressed in terms familiar to the shrimp industry, and (2) any failure to detect impacts would suggest that the impacts were not of major consequence to the shrimp fisheries.




Figure 1. Locations of study sites in the assessment of impacts of brine disposal in the northwestern Gulf of Mexico.

Table 1. Sequences of four projects and their work units along with responsible organization or agency (in parentheses) contributing to the assessment of impacts of brine disposal in the northwestern Gulf of Mexico (See Table 2 for explanation of acronyms; see Fig. 1 for locations of study sites).

Projects and Work Units (responsible organization or agency)	1978	1979	1980	1981	1982
<u>Biological/Chemical Survey of Texoma (West Hackberry) and Capline (Weeks Island) Sector Salt Dome Brine Disposal Sites off Louisiana</u>					
Macro- and Meiobenthos (CEM).....	X	X			
Zooplankton (LGL).....	X	X			
Bacteria (TAMU).....	X	X			
Demersal Fishes & Macro-crustaceans (TAMU).....	X	X			
Sediments & Suspended Particulates (ERCO).....	X	X			
Hydrocarbons (ERCO).....	X	X			
Trace Metals (SRI).....	X	X			
Inorganic Nutrients (TAMU).....	X	X			
Shrimp Catch and Effort Data Analysis (LGL).....	X	X			
<u>Shrimp and Redfish Studies; Bryan Mound Brine Disposal Site off Freeport, Texas</u>					
Shrimp Catch and Effort Data Analysis (SAI).....	X				X
Analysis of Shrimping Success, Recruitment & Environmental Variables (SAI).....	X				X
Shrimp Mark-Release Investigations (LGL).....	X				X
Shrimp Spawning Site Survey (LGL).....	X				X
Shrimp Catch-Effort Sampling Survey (LGL).....	X				X
Brine Toxicity Bioassays on Shrimp (UH).....	X				X
Brine Avoidance/Attraction Bioassays on Shrimp (UH).....	X				X
Brine Toxicity Bioassays on Redfish (TAMU/LGL).....	X				X
Brine Avoidance/Attraction Bioassays on Redfish (TAMU/LGL).....	X				X
<u>Shrimp Population Studies; West Hackberry and Big Hill Brine Disposal Sites off Southwest Louisiana and upper Texas Coast</u>					
Shrimp Catch and Effort Data Analysis (TAMU).....	X				X
Analysis of Shrimping Success, Recruitment & Environmental Variables (URI).....	X				X
Shrimp Mark-Release and Port Interview Sampling Survey (LGL)	X				X
Shrimp Spawning Site Survey (LGL).....	X				X
<u>Shrimp Population Studies; Bryan Mound Brine Disposal Site off Freeport, Texas</u>					
Shrimp Mark-Release and Port Interview Sampling Survey (LGL).....	X				X
Brine Disposal Impact Analysis & Assessment (TAMU).....	X				X

Table 2. List of participating principal investigators contributing to the assessment of impacts of brine disposal in the northwestern Gulf of Mexico.

Boehm, P. D., ERCO (Energy Resources Company, Inc.), Cambridge, MA

Brooks, J. M., Department of Oceanography, TAMU (Texas A & M University), College Station, TX

Comiskey, C. E., SAI (Science Applications, Inc.), Oak Ridge, TN

Foreman, J., NESDIS (National Environmental Satellite, Data and Information Service), NOAA (National Oceanic and Atmospheric Administration), Washington, D.C.

Gallaway, B. J., LGL (LGL Ecological Research Associates, Inc.), Bryan, TX

Gazey, W. J., LGL, Bryan, TX

Grant, W. E., Department of Wildlife & Fisheries Sciences, TAMU, College Station, TX

Hausknecht, K. A., ERCO, Cambridge, MA

Holley, H., NFEL (National Fisheries Engineering Laboratory), SEFC (Southeast Fisheries Center), NMFS (National Marine Fisheries Service), NOAA, Bay St. Louis, MS

Howe, N. R., Marine Science Program, UH (University of Houston), Galveston, TX

Table 2.(continued)

Johnson, M. F., LGL, Bryan, TX

Landry, A. M., Department of Marine Biology, Mitchell Campus, TAMU,
Galveston, TX

Margraf, F. J., LGL, Bryan, TX

Matis, J. H., School of Applied Statistics, TAMU, College Station, TX

Neff, J. M., Department of Biology, TAMU, College Station, TX

Owens, D. W., Department of Biology, TAMU, College Station, TX

Parker, R. H., CEM (Coastal Ecosystems Management, Inc.), Fort Worth,
TX

Reitsema, L. A., LGL, Bryan, TX

Saila, S. B., Graduate School of Oceanography, URI (University of
Rhode Island), Kingston, RI

Savastano, K., NFEL, SEFC, NMFS, NOAA, Bay St. Louis, MS

Schwarz, J. R., Department of Marine Biology, Mitchell Campus, TAMU,
Galveston, TX

Tidwell, D., FIMD (Fisheries Information Management Division), SEFC,
NMFS, NOAA, Miami, FL

Tillery, J. B., SRI (Southwest Research Institute), San Antonio, TX

Major products of NMFS's assessment were (1) final reports to DOE (see bibliography), copies of which are available from the DOC (Department of Commerce), NTIS (National Technical Information Service), 5285 Port Royal Road, Springfield, Virginia 22161, (2) data files archived by the NESDIS and the FIMD, and (3) research papers by participating investigators (see bibliography).

BIBLIOGRAPHY

- Boehm, P. D. and D. L. Fiest. 1980. Determine hydrocarbon composition and concentration in major components of the ecosystem, Vol. VI In: Jackson, W. B. and G. M Faw (eds.). Biological/chemical survey of Texoma and Capline sector salt dome brine disposal sites off Louisiana, 1978-1979. NOAA Tech. Memo. NMFS-SEFC-30, xxi + v + 138 p. DOC NTIS Accession No. PB81 174971.
- Brooks, J. M. 1980. Determine seasonal variations in inorganic nutrients composition and concentration in the water column, Vol. VIII In: Jackson, W. B. and G. M. Faw (eds.). Biological/ chemical survey of Texoma and Capline sector salt dome brine disposal sites off Louisiana, 1978-1979. NOAA Tech. Memo. NMFS-SEFC-32, xx + iv + 31 p. DOC NTIS Accession No. PB81 182685.
- CEAS (Center for Environmental Assessment Services). 1981. Brine disposal in the Gulf of Mexico: projected impacts for West Hackberry based on Bryan Mound experience. CEAS, Marine Environmental Assessment Division, Environmental Data and Information Service, NOAA, U.S. Dept. of Commerce, Wash., D.C., 51 p. plus appendices.
- CEDDA (Center for Experiment Design and Data Analysis). 1978. Analysis of brine disposal in the Gulf of Mexico: Bioassay results. Report to the Department of Energy, Strategic Petroleum

Reserve Program, Salt Dome Storage. CEDDA, Marine Assessment Division, Environmental Data Service, NOAA, U.S. Dept. of Commerce, Wash., D.C., 116 p.

Comiskey, C. E. , R. McCord, D. Bozarth, S. Grady, C. Hall, C. Brandt and T. Farmer. 1982a. Analysis of data on shrimping success, shrimp recruitment and associated environmental variables. Vol. I(A) In: Klima, E. F. (Contracting Officer's Technical Representative). Shrimp and redfish studies; Bryan Mound brine disposal site off Freeport, Texas, 1979-1981. NOAA Tech. Memo. NMFS-SEFC-65, xix + li + 449 p. DOC NTIS Accession No. PB83 139279.

Comiskey, C. E., R. McCord, D. Bozarth, S. Grady, C. Hall, C. Brandt, and T. Farmer. 1982b. Texas coast shrimp catch and effort data analysis. Vol. 1(B) In: Klima, E. F. (Contracting Officer's Technical Representative). Shrimp and redfish studies; Bryan Mound brine disposal site off Freeport, Texas, 1979-1981. NOAA Tech. Memo. NMFS-SEFC-65, xxii + xxiv + 217 p. DOC NTIS Accession No. PB83 122705.

Foreman, J., K. Pechmann, F. Mitchell, L. May, J. Butler, and L. Bowser. 1983. Brine disposal in the Gulf of Mexico: Inventory of the Bryan Mound site pre-disposal data set, 9/15/77-3/31/80. National Environmental Satellite, Data, and Information Service, NOAA, U.S. Dept. of Commerce, Wash., D.C., 38 p.

Galloway, B. J. and L. A. Reitsema. 1981. Shrimp spawning site survey. Vol. III In: Jackson, W. B. and E. P. Wilkens (eds.). Shrimp and redfish studies; Bryan Mound brine disposal site off Freeport, Texas, 1979-1981. NOAA Tech. Memo. NMFS-SEFC-67, xviii + viii + 84 p. DOC NTIS Accession No. PB81 249591.

Gazey, W. J., B. J. Galloway, R. C. Fechhelm, L. R. Martin and L. A. Reitsema. 1982a. Shrimp mark-release and port interview sampling survey of catch and effort with recovery of recaptured tagged shrimp. Vol. III In: Jackson, W. B. (ed.). Shrimp population studies; West Hackberry and Big Hill brine disposal sites off southwest Louisiana and upper Texas coasts, 1980-1982. NOAA NMFS Final Report to DOE, xxiv + 288 p. DOC NTIS Accession No. PB83 172270.

Gazey, W. J., B. J. Galloway, R. C. Fechhelm, L. R. Martin and L. A. Reitsema. 1982b. Shrimp mark-release and port interview sampling survey of shrimp catch and effort with recovery of recaptured tagged shrimp. Vol. I In: Jackson, W. B. (ed.). Shrimp population studies; Bryan Mound brine disposal site off Freeport, Texas, 1981-1982. NOAA NMFS Final Report to DOE, xxv + 291 p. DOC NTIS Accession No. PB83 175869.

Hausknecht, K. A. 1980. Describe surficial sediments and suspended particulate matter. Vol. V In: Jackson, W. B. and G. M. Faw (eds.). Biological/chemical survey of Texoma and Capline Sector salt dome brine disposal sites off Louisiana, 1978-1979. NOAA

Tech. Memo. NMFS-SEFC-29, xx + vii + 56 p. DOC NTIS Accession No. PB81 174963.

Howe, N. R. 1981. Brine toxicity and avoidance/attraction bioassays on shrimp. Vol. VI In: Jackson, W. B. and E. P. Wilkens (eds.). Shrimp and redfish studies; Bryan Mound brine disposal site off Freeport, Texas, 1979-1981. NOAA Tech. Memo. NMFS-SEFC-70, xviii + iv + 60 p. DOC NTIS Accession No. PB81 249609.

Howe, N. R., W. D. Quast and L. M. Cooper. 1982. Lethal and sub-lethal effects of a simulated salt brine effluent on adults and subadults of the shrimps Penaeus setiferus and P. aztecus. Mar. Biol. 68:37-47.

Johnson, M. F. 1981. Shrimp mark-release investigations. Vol. II In: Jackson, W. B. and E. P. Wilkens (eds.). Shrimp and redfish studies; Bryan Mound brine disposal site off Freeport, Texas, 1979-1981. NOAA Tech. Memo. NMFS-SEFC-66, xx + x + 110 p. DOC NTIS Accession No. PB81 249583.

Johnson, M. F. 1981. Interview sampling survey of shrimp catch and effort. Vol. IV In: Jackson, W. B. and J. R. Bennett (eds.). Shrimp and redfish studies; Bryan Mound brine disposal site off Freeport, Texas, 1979-1981. NOAA Tech. Memo. NMFS-SEFC-68, xxiii + 38 p. DOC NTIS Accession No. PB82 116062.

Landry, A. M. and H. W. Armstrong. 1980. Determine seasonal abun-

dance, distribution and community composition of demersal finfishes and macrocrustaceans. Vol. IV In: Jackson, W. B. and G. M. Faw (eds.). Biological/chemical survey of Texoma and Capline sector salt dome brine disposal sites off Louisiana, 1978-1979. NOAA Tech. Memo. NMFS-SEFC-28, xx + xvi + 180 p. DOC NTIS Accession No. PB81 174955.

Margraf, F. J. 1980. Analysis of variance of Gulf coast shrimp data. Vol. IX In: Jackson, W. B. and G. M. Faw (eds.). Biological/chemical survey of Texoma and Capline sector salt dome brine disposal sites off Louisiana, 1978-1979. NOAA Tech. Memo. NMFS-SEFC-33, xx + xxii + 293 p. DOC NTIS Accession No. PB81 133803.

Matis, J. H., P. E. Carothers and W. E. Grant. 1982. Brine disposal impact analysis and assessment. Vol. II In: Jackson, W. B. (ed.). Shrimp population studies; Bryan Mound brine disposal site off Freeport, Texas, 1981-1982. NOAA NMFS Final Report to DOE, viii + 91 p. DOC NTIS Accession No. PB83 216465.

Matis, J. H. and W. E. Grant. 1982. Shrimp catch and effort data analysis. Vol. II In: Jackson, W. B. (ed.). Shrimp population studies; West Hackberry and Big Hill brine disposal sites off southwest Louisiana and upper Texas coasts, 1980-1982. NOAA NMFS Final Report to DOE, xi + 195 p. DOC NTIS Accession No. PB83 172262.

- Neff, J. M., M. P. Coglianesi, W. McCulloch, L. A. Reitsema and S. Anderson. 1982. Brine toxicity bioassays on redbfish. Vol. V (Part A) In: Jackson, W. B. (ed.). Shrimp and redbfish studies; Bryan Mound brine disposal site off Freeport, Texas, 1979-1981. NOAA Tech. Memo. NMFS-SEFC-69, xvii + x + 82 p. DOC NTIS Accession No. PB83 159459.
- Owens, D. W. and K. A. Jones. 1982. Acclimation temperature and orientation to salinity gradients by the red drum. Amer. Zool. 22:864 (Abstract #82).
- Owens, D. W., K. A. Jones and B. J. Gallaway. 1982. Brine avoidance/attraction bioassays on redbfish. Vol. V (Part B) In: Jackson, W. B. (ed.). Shrimp and redbfish studies; Bryan Mound brine disposal site off Freeport, Texas, 1979-1981. NOAA Tech. Memo. NMFS-SEFC-69, xvii + ix + 58 p. DOC NTIS Accession No. PB82 215914.
- Parker, R. H., A. L. Crowe and L. S. Bohme. 1980. Describe living and dead benthic (macro- and meio-) communities. Vol. I In: Jackson, W. B. and G. M. Faw (eds.). Biological/chemical survey of Texoma and Capline sector salt dome brine disposal sites off Louisiana, 1978-1979. NOAA Tech. Memo. NMFS-SEFC-25, xx + vi + 103 p. DOC NTIS Accession No. PB81 133795.
- Reitsema, L. A. 1980. Determine seasonal abundance, distribution and community composition of zooplankton. Vol. II In: Jackson, W. B.

and G. M. Faw (eds.). Biological/chemical survey of Texoma and Capline sector salt dome brine disposal sites off Louisiana, 1978-1979. NOAA Tech. Memo. NMFS-SEFC-26, xx + ix + 133 p. DOC NTIS Accession No. PB81 175838.

Reitsema, L. A., B. J. Gallaway and G. S. Lewbel. 1982. Shrimp spawning site survey. Vol. IV In: Jackson, W. B. (ed.). Shrimp population studies; West Hackberry and Big Hill brine disposal sites off southwest Louisiana and upper Texas coasts, 1980-1982. NOAA NMFS Final Report to DOE, x + 82 p. DOC NTIS Accession No. PB83 172288.

Saila, S. B., H. A. Walker, E. Lorda, J. Kelly and M. Prager. 1982. Analysis of data on shrimping success, shrimp recruitment and associated environmental variables. Vol. I In: Jackson, W. B. (ed.). Shrimp population studies; West Hackberry and Big Hill brine disposal sites off southwest Louisiana and upper Texas coasts, 1980-1982. NOAA NMFS Final Report to DOE, xviii + 238 p. DOC NTIS Accession No. PB83 172254.

Schwarz, J. R., S. K. Alexander, S. J. Schropp and V. L. Carpenter. 1980. Describe bacterial communities. Vol. III In: Jackson, W. B. and G. M. Faw (eds.). Biological/chemical survey of Texoma and Capline sector salt dome brine disposal sites off Louisiana, 1978-1979. NOAA Tech. Memo. NMFS-SEFC-27, xx + vi + 48 p. DOC NTIS Accession No. PB81 174948.

Tillery, J. B. 1980. Determine trace metal composition and concentration in major components of the ecosystem. Vol. VII In: Jackson, W. B. and G. M. Faw (eds.). Biological/chemical survey of Texoma and Capline sector salt dome brine disposal sites off Louisiana, 1978-1979. NOAA Tech. Memo. NMFS-SEFC-31, xx + viii + 72 p. DOC NTIS Accession No. PB81 174989.

Turgeon, K. 1981. Synthesis of Texoma/Capline chemical and biological survey results: June 1978-May 1979. Marine Environmental Assessment Division, Center for Environmental Assessment Services, Environmental Data and Information Services, NOAA, DOC, Wash., D.C., xii + 87 p.