

1
2 Received Date : 12-Feb-2016
3 Revised Date : 12-Sep-2016
4 Accepted Date : 19-Sep-2016
5 Article type : Original Article
6
7

8 **Title: How old are you? Genet age estimates in a clonal animal.**

9 Authors: M.K. Devlin-Durante¹, M.W. Miller², Caribbean *Acropora* Research Group, W.F.
10 Precht³, I.B. Baums^{1*}
11

12 ¹ Dept of Biology, The Pennsylvania State University, 208 Mueller Lab, University Park, PA
13 16802 USA

14 ² National Marine Fisheries Service, Southeast Fisheries Science Center 75 Virginia Beach Dr.,
15 Miami, FL 33149 USA

16 ³ Marine & Coastal Programs, Dial Cordy & Associates, 90 Osceola Ave, Jacksonville Beach,
17 FL 32250 USA
18

19 Caribbean *Acropora* Research Group:

20 Lisa Carne, Fragments of Hope, Placencia Village Stann Creek District, Belize, isas@btl.net

21 Tyler B. Smith, Center for Marine and Environmental Studies, University of the Virgin Islands, St.

22 Thomas, VI 00802, USA, tsmith@uvi.edu

23 Anastazia T. Banaszak, Unidad Académica de Sistemas Arrecifales, Puerto Morelos, Instituto de Ciencias
24 del Mar y Limnología, Universidad Nacional Autónoma de México, banaszak@cmarl.unam.mx

25 Lisa Greer, Washington and Lee University, Lexington, VA 24450, greerl@wlu.edu

26 Adele Irwin, Washington and Lee University, Lexington, VA 24450, irwina15@mail.wlu.edu

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/mec.13865](https://doi.org/10.1111/mec.13865)

This article is protected by copyright. All rights reserved

27 Nicole D. Fogarty, Department of Marine and Environmental Sciences Halmos College of Natural
28 Science and Oceanography Nova Southeastern University, 8000 N. Ocean Drive, Dania Beach, FL
29 33004-3078, nicole.fogarty@nova.edu

30 Dana E. Williams, Cooperative Institute for Marine and Atmospheric Studies, University of Miami, 4600
31 Rickenbacker Causeway. Miami FL, 33149 USA; and ², dana.williams@noaa.gov

32
33 * Corresponding author: Iliana Baums, Assoc Prof, Dep of Biology, Pennsylvania State
34 University, 208 Mueller Lab, University Park, PA 16802 USA, Fax 814.865.9131,
35 baums@psu.edu

36 **Keywords**

37 Somatic mutations, microsatellite, longevity, clonal, population dynamics

38
39 **Running Head:** Coral genet age estimates

40 **Abstract**

41 Foundation species such as redwoods, seagrasses and corals are often long-lived and
42 clonal. Genets may consist of hundreds of members (ramets) and originated hundreds to
43 thousands of years ago. As climate change and other stressors exert selection pressure on species,
44 the demography of populations change. Yet, because size does not indicate age in clonal
45 organisms, demographic models are missing data necessary to predict the resilience of many
46 foundation species. Here, we correlate somatic mutations with genet age of corals and provide
47 the first, preliminary estimates of genet age in a colonial animal. We observed somatic mutations
48 at 5 microsatellite loci in range wide samples of the endangered coral, *Acropora palmata*
49 (n=3352). Colonies harbored 342 unique mutations in 147 genets. Genet age ranged from 30-838
50 years old (y/o) assuming a mutation rate of $1.195^{-04} \text{ locus}^{-1} \text{ year}^{-1}$ based on colony growth rates
51 and 236-6500 y/o assuming a mutation rate of $1.542^{-05} \text{ locus}^{-1} \text{ year}^{-1}$ based on sea level changes
52 to habitat availability. Long-lived *A. palmata* genets imply a large capacity to tolerate past
53 environmental change and yet recent mass mortality events in *A. palmata* suggest that capacity is
54 now being frequently exceeded.

This article is protected by copyright. All rights reserved

55 **Introduction**

56 The population dynamics of a species depend in part on the longevity of each individual.
57 However, in colonial organisms such as corals neither “individual” nor “age” are easy to define,
58 making longevity the least accessible demographic trait to study for these organisms. Coral
59 colonies consist of genetically identical polyps that each fulfill the function of an individual
60 (reproduction, growth, defense), yet it is the collection of polyps in a colony that represent the
61 ecologically significant unit (Santelices 1999). Hence, studies of coral population dynamics
62 often track the fate of colonies rather than that of individual polyps. The very nature of the
63 clonality of corals allows colonies to survive partial mortality (Hughes & Jackson 1980),
64 propagate asexually through fragmentation (Highsmith 1982), and partake in clonal fission and
65 fusion (Hughes & Jackson 1980). The result is independent colonies (ramets) not connected by
66 live tissue that share the same genotype (clonemates of the same genet). Coral species where
67 clonemates constitute a significant proportion of local populations are found in at least nine coral
68 genera (Supplementary Table 1). Ramets are produced throughout the lifetime of the genet and
69 hence they can be of different chronological age and size although their genetic age (i.e. the time
70 since meiosis and zygote formation) remains the same. Taken together these processes have the
71 net effect of decoupling size of a ramet from its age (Hughes & Jackson 1980).

72 In non-colonial multicellular organisms, size is often a good proxy of genet age until
73 adult size is attained. After adult size is reached, age determination becomes more challenging
74 but the incorporation of environmental signals into tissues (Prouty *et al.* 2011), the shortening of
75 telomeres with increasing numbers of cell divisions (Barrett *et al.* 2013), decreasing reproductive
76 output, and phenotypic changes (Caspari & Lee 2004) can be quantified as indicators of age in a
77 wide range of multicellular organisms. Many of these approaches are not useful in plants and
78 colonial invertebrates: Radiocarbon or U-series dating (Radtke *et al.* 2003) is an alternative to
79 using size or phenotypic changes as a proxy for genetic age, however this requires the
80 identification and continued existence of the oldest portion of a genet because, as such,
81 environmental signals reflect ramet age, not genet age (Eggins *et al.* 2005). This may be
82 possible in some clonal plant species in which ramet attachment persists and the center, typically
83 the oldest portion of a genet, can be identified (Vasek 1980), and perhaps for coral species not

84 prone to fragmentation (Table 1, Supplementary Table 1). Furthermore, reproduction is tied to
85 colony size so recently fragmented ramets belonging to previously fecund colonies might not
86 produce gametes themselves (Okubo *et al.* 2007) and phenotypic changes are not obvious
87 because a genetically old but small coral colony is not visually distinguishable from a genetically
88 young and small colony.

89 A possible method for determining genet age is to use mutation accumulation in somatic
90 tissues to estimate longevity. Despite their asexual origin, clonemates are not always exactly
91 genetically identical. The concept is based on “the somatic mutation theory of clonality”
92 (Klekowski 1997) which reasons that continuous division of mitotic cells in a clonal organism
93 will lead to the accumulation of somatic mutations over time. Somatic mutations convert a
94 genetically homogenous individual into a mosaic with divergent cell lineages (mosaicism). Due
95 to the stochastic nature of somatic mutations, the incidence of genetic mosaicism would be
96 expected to increase with increasing longevity of the organism and also with a higher prevalence
97 of asexual reproduction; gain in ramet number or size increases the total number of dividing cells
98 available for mutation (Orive 2001). Thus, it should be possible to relate the accumulation of
99 somatic mutations to genet age.

100 Utilizing genetic divergence generated by somatic mutations is a novel approach for
101 calculating lifespans in clonal organisms (Heinze & Fussi 2008). The use of neutral
102 microsatellites is ideal for divergence estimates due to their high mutation rates that range from
103 10^{-2} to 10^{-6} per sexual generation (Shimoda *et al.* 1999; Ellegren 2000; Peery *et al.* 2012).
104 Genetic divergence in microsatellite loci has been used to model clonal age in the aspen tree
105 *Populus tremuloides* (Ally *et al.* 2008) and the water flea *Daphnia magna* (Robinson *et al.*
106 2012).

107 Limitations of life span estimates based on genetic divergence include the necessity of
108 clonality, the low frequency or absence of mutations in some species (Lanner & Connor 2001;
109 Cloutier *et al.* 2002) and difficulties in measuring mutational rates that are often variable among
110 loci (Chakraborty *et al.* 1997; Schug *et al.* 1998). It can also be challenging to distinguish
111 somatic mutations from allelic variation (Heinze & Fussi 2008) if the species under
112 consideration is inbred.

113 Furthermore, the rate of somatic mutational divergence not only differs between species
114 (Klekowski & Godfrey 1989), but also among individuals (Haag-Liautard *et al.* 2007; Conrad *et*
115 *al.* 2011) with intraspecies variation partly due to varying exposure to environmental stress (de
116 Witte & Stöcklin 2010). Genetic homogeneity can be restored from a mosaic state through
117 sexual reproduction, but also through parallel back mutations or lineage selection (Klekowski &
118 Kazarinova-Fukshansky 1984) which would lead to underestimates of mutational load and thus
119 clonal age. Despite the limitations, genetic divergence estimates are the most promising
120 technique to estimate genet age in colonial marine invertebrates.

121 To demonstrate the potential of using somatic divergence estimates to estimate genet
122 longevity, we used genetic divergence in 5 microsatellite loci to calculate the age of 90 genets of
123 the elkhorn coral, *Acropora palmata*. *A. palmata* is an ideal species for determining genet age
124 based on somatic mutations because this species relies heavily on fragmentation for local
125 population maintenance (Highsmith 1982; Baums *et al.* 2006a; Williams & Miller 2012) and
126 some genets have > 30 members (Baums *et al.* 2014). The process of fragmentation and re-
127 growth of colonies from fragments has been documented photographically via quarterly surveys
128 over the past decade or so (Supplemental Figure 1, Williams & Miller 2012) and fragments
129 match donor colony genotypes. Furthermore, in a previous range-wide study of population
130 genetic structure in *A. palmata* we noticed the occasional occurrence of three alleles per locus in
131 this otherwise diploid species (Baums *et al.* 2005a). *A. palmata* is a self-incompatible
132 hermaphrodite (Szmant 1986a; Baums *et al.* 2005a) and population genetic data show that the
133 species is genetically diverse and outbred (Baums *et al.* 2005b). Here, we investigate whether
134 third alleles in *A. palmata* arose from somatic mutations and then use somatic mutations to
135 estimate genet age in this species.

136 **Methods**

137 *Study System*

138 *Acropora palmata* is a fast-growing, branching coral that once dominated coral reefs in
139 the Caribbean and North-West Atlantic. Adult colonies release egg-sperm bundles once a year
140 after the August full moon during a synchronized mass-spawning events (Szmant 1986b). Egg-

141 sperm bundles float to the surface where they break apart. Successful fertilization requires the
142 union of egg and sperm from different genets, i.e. *A. palmata* is a self-incompatible
143 hermaphrodite (Baums *et al.* 2005a). Gametes develop into non-feeding planula larvae during a 3
144 day – 2 week planktonic period. Mature larvae search for suitable habitat and metamorphose into
145 primary polyps during a process generally referred to as settlement (Fig 1). Once the primary
146 polyp is established, it will bud repeatedly, a type of asexual reproduction, and eventually form a
147 colony of genetically identical polyps. In some cases, two genetically distinct primary polyps
148 (recently settled larvae) can fuse, resulting in colonies with mixtures of polyps of different
149 genotypes (chimerism, Barki *et al.* 2002; Puill-Stephan *et al.* 2009; Work *et al.* 2011). Signals
150 and resources are shared across the colony. There is also division of labor to a degree with some
151 polyps primarily engaged with defense, reproduction or growth (Soong & Lang 1992). Because
152 of this integration, the colony is usually considered as the ecologically significant unit. We refer
153 to an assemblage of genetically identical colonies that are descendants of a single zygote as a
154 “genet” (Harper 1977; Hughes 1989; Carvalho 1994). Physiologically distinct colonies, formed
155 from fragmentation, that can function and survive on their own but belong to the same genet are
156 termed “ramets” (Kays & Harper 1974).

157 Samples of *A. palmata* were collected from Florida and the Caribbean (2001 to 2012,
158 n=3352, Fig 2, Table 2). The time range of sample collection lends an error rate of +/- 12 years
159 to the age calculations. Previous population genetic evidence (Baums *et al.* 2005b) divided *A.*
160 *palmata* samples into two largely isolated populations, the eastern Caribbean (including Bonaire,
161 Curacao, St Vincent and the Grenadines, the US Virgin Islands) and the western Caribbean
162 (including the Bahamas, Belize, Cuba, Dominican Republic, Florida, Mexico, Mona, Navassa
163 and Panama). Samples from Puerto Rico were assigned to the eastern Caribbean but show some
164 degree of admixture between the east and the west. A subset of the total dataset (n = 430 from 14
165 reefs in the Bahamas, Bonaire, Curacao, Florida, Panama, the US Virgin Islands and Navassa)
166 were sampled using a stratified, random sampling approach, as described in Baums *et al.*
167 (2006a). Most colonies within our collection were only sampled once, however 11 colonies from
168 Florida were resampled in 2011 and 2014 at 2-8 locations within the colony (Supplemental Table
169 1).

170 Microsatellite scoring.

This article is protected by copyright. All rights reserved

171 All samples were genotyped at five (166, 181, 182, 192, and 207) previously published,
172 polymorphic microsatellite loci with Mendelian inheritance as shown by experimental crosses
173 (Baums *et al.* 2005b). All 5 microsatellite loci are AAT trinucleotide repeats. Two 10 μ l
174 multiplex PCR reactions (M-I and M-II) were performed per sample. M-I consisted of 0.2 μ l
175 each of primer pairs 166-PET (5 μ M), 192-6FAM (5 μ M) and 181-NED (5 μ M), 1 μ l 10 \times PCR
176 Reaction Buffer (Promega), 0.8 μ l of MgCl₂ (25 mM), 0.2 μ l of dNTPs (10 mM), 0.3 μ l of Taq-
177 Polymerase (5 U μ l⁻¹, Storage Buffer B, Promega) and 6.1 μ l H₂O. M-II consisted of 0.2 μ l each
178 of primer pairs 207-PET (5 μ M) and 182-6FAM (5 μ M), 1 μ l Promega 10 \times PCR Reaction
179 Buffer, 1.2 μ l of MgCl₂ (25 mM), 0.2 μ l of dNTPs (10 mM), 0.2 μ l of Taq-Polymerase (5 U μ l⁻
180 1) and 6 μ l H₂O. DNA (100 to 200 ng, 1 μ l) was added to each reaction. Thermal cycling was
181 carried out with Eppendorf Mastercylers with an initial denaturation step at 95°C for 5 min
182 followed by 35 cycles of 95°C for 20 s, 50°C for 20 s, 72°C for 30 s. A final extension of 30 min
183 at 72°C ensured that the majority of amplicons were +A (Brownstein *et al.* 1996). PCR products
184 were visualized using an ABI 3730. An internal size standard (Gene Scan 500-Liz, Applied
185 Biosystems) was used for accurate sizing. Electropherograms were analyzed with GeneMapper
186 Software 5.0 (Applied Biosystems).

187 Unique clonal IDs for a genet were assigned to corals that have exact matching
188 multilocus genotypes or have exact matching multilocus genotypes (share all the same diploid
189 state ancestral alleles) and have an additional allele(s). The exceptions to this rule were 4% of
190 mutations that were either a full mutation (e.g. ancestral state 166/175 to 166/178), or a loss of
191 heterozygosity (e.g. to 166/166, Table 3), but at the other 4 loci all alleles were shared with other
192 members of the genet (see Supplementary Table 2 for an example genet).

193 Loci had an average of 19.6 alleles (StDev \pm 2.3). This level of polymorphism translated
194 into a high power of distinguishing closely related (i.e. inbred) multilocus genotypes (MLGs)
195 from those that were the product of asexual reproduction (i.e. clonemates) where the probability
196 of identity = 10^{-5} (Baums *et al.* 2005b) (See Supplementary Figure 2). When considering only
197 genotypes with 2 alleles per locus (n=2643, i.e. those without somatic mutations) the average
198 probability of encountering a genotype more than once by chance (psex) was 2.23^{-07} (MLGsim
199 2.0, <http://www.rug.nl/research/gelifes/tres/software>), indicating that identical genotypes were
200 the result of asexual reproduction. Once asexually produced, identical MLGs are removed from

201 the dataset, no heterozygote deficits are detected (i.e. all loci adhere to Hardy-Weinberg
202 expectations (Baums *et al.* 2005a)) and thus *A. palmata* shows no sign of inbreeding (Halkett *et*
203 *al.* 2005).

204 *Mutation Step Analysis*

205 For all genets with at least two ramets each novel mutation was reported (referred to as a
206 unique mutation, UM). A total of 342 unique mutations were found in 147 genets with 1387
207 ramets (Table 2, Fig 3). In order to discriminate between a mutated allele and a PCR error, a
208 singleplex PCR was performed for all unique mutations. Following a step-wise mutation model
209 (Kimura & Ohta 1978) the smallest possible mutational step that could have resulted in the new
210 allele was used to determine which of the two ancestral alleles mutated and the size of the
211 mutation step (in repeat units). Mutations were excluded if there were no other samples within
212 the genet that were bi-allelic at that locus making it impossible to determine the mutation step.
213 However, sometimes a genet had only two ramets and both ramets had different mutations at the
214 same locus. In that case the ancestral allele state was determined to consist of the two alleles
215 found in both ramets (Table 2). The mutational-step analysis contained a reduced sample size of
216 $n=1387$ (Table 3).

217 *Clustering analysis*

218 To determine whether the samples with three alleles could be attributed to somatic
219 mutations or chimerism, we applied a Bayesian clustering analysis using the program
220 STRUCTURE 2.3.4 (Pritchard *et al.* 2000) to all genets with at least 5 ramets ($n_{\text{genets}} = 90$, Table 2).
221 We forced a diploid state by replacing the ancestral allele with the 3rd allele mutation. There was
222 no missing genotype data. We assumed that ramets should only diverge from the ancestral
223 genotype in one or two loci or alleles if somatic mutations were the cause, following previous
224 studies (Puill-Stephan *et al.* 2009; Maier *et al.* 2011). Alternatively, colonies were defined as
225 chimeras if genotypes differed by more than 60% in their major cluster assignment probability
226 from other members of their genet as defined by Schweinsberg *et al.* (2015). STRUCTURE 2.3.4
227 (Pritchard *et al.* 2000) was run with a burn-in period of 100,000 and 1,000,000 MCMC repeats
228 with 3 iterations per K, without a prior (Fig 4). Because of their large number, Florida genets
229 were run in two separate groups each containing 22 genets, with $K=22$. The eastern Caribbean

230 samples (23 genets, K=23) and all other western Caribbean samples (23 genets, K=23) were run
231 in two additional groups. Results of the three runs per group were merged with CLUMPAK
232 (Kopelman *et al.* 2015).

233 *Clonal Richness vs. Mosaicism*

234 We evaluated whether somatic mutations were found more often on reefs where little
235 sexual recruitment was evident (and thus were presumably inhabited by older individuals) by
236 tallying all mutations in all samples and comparing the number of mutations detected with the
237 number of genets present. This was expressed as clonal richness. We did this analysis on two
238 datasets. We compared the proportion of non-mosaic samples to clonal richness on reefs with
239 ≥ 10 samples, with no limitations placed on the genet size (Table 2). Therefore, clonal and non-
240 clonal samples were included in this analysis (i.e. all genotype samples n=3352, Table 2). Then,
241 we only compared reefs that were sampled with similar sampling effort. (see Table 1 in Baums *et al.*
242 (2006a). The clonal richness R is calculated as the number of genets G relative to the number
243 of analyzed ramets N with the modification by Dorken and Eckert (2001):

$$R = \frac{G - 1}{N - 1}$$

244

245 A monoclonal stand has a clonal richness of R=0 whereas the maximum clonal richness
246 of R=1 is reached when all samples from a reef are of a different MLG. We chose clonal richness
247 as an indicator for clonal diversity because other measures assume a constant ploidy level (most
248 often diploidy e.g. G_o/G_e) and were not designed for samples with somatic mutations.

249 *Estimates of Genet Age using Genetic Divergence*

250 The methods for calculation clonal age utilizing genetic divergence are described in (Ally
251 *et al.* 2008). In brief, there are two statistics, π_k and S_k , that describes genetic divergence within
252 a clone (Slatkin 1996). We calculated the average number of pairwise differences per locus for

253 the kth clone: $\pi_k = \frac{1}{n} \sum_{i=1}^{n-1} \sum_{j=i+1}^n s_{ij}$

254 where n is the number of sampled ramets, s_{ij} is the number of genetic differences between ramet
255 i and j averaged across loci, and $\frac{n-1}{2}$ is the total number of pairwise differences (Ally *et al.*

256 2008). We chose π_k to measure the level of genetic divergence because it has been shown to be

257 more robust to deviations from a star-like phylogeny than S_k (the observed proportion of
258 polymorphic loci) (Ally *et al.* 2008). Two demographic models were contrasted: one of constant
259 ramet population size (as in the classic Wright–Fisher model), while the second demographic
260 model is one of population growth. The ramet population growth model was determined by
261 plotting both π_k vs. $S_k / \sum_{i=1}^{n-1} (\frac{1}{i})$, which should exhibit a 1:1 slope if a population has been
262 constant in size, and π_k vs. $2S_k/n$ in which a 1:1 slope would be predicted for a clonal growth
263 model. The fit of the models was determined by regression analysis obtained in Sigmaplot 10.0.

264 Further restrictions, to the sample set, were applied for clonal age estimates, with ramet
265 size of $n \geq 5$ resulting in $n=90$ genets used in this analysis (Table 2). While most colonies were
266 sampled once, we captured the allelic variation within a genet by restricting age calculations to
267 those genets with at least 5 ramets. We still may have missed some somatic mutations at these
268 loci leading to an underestimation of the minimum genet age. Note that ramets lacking mutations
269 but belonging to a genet that had other ramets with mutations (ramet number 5 or greater) were
270 included (Table 2). If the genet had at least 5 ramets but no ramets had mutations then
271 microsatellite divergence and therefore age could not be calculated.

272 There are currently no direct estimates for microsatellite mutation rates in *A. palmata*.
273 We assumed the same mutation rate for all samples, but we were uncertain about that rate.
274 Hence, we used a range by setting a maximum and a minimum. The upper bound for the
275 mutation rate (relatively fast mutation rate) implies that a shorter amount of time has passed to
276 accumulate the observed variation relative to the lower bound of the estimate (relatively slow
277 mutation rate). Genet P1028 from Elbow reef in Florida had the smallest microsatellite
278 divergence rate. This genet had 55 ramets, among which the largest single colony was 270 x 170
279 x 70 cm (L x W x H). The branch extension rate was measured directly on three ramets of this
280 genet (P1028) during Jan-July 2006. A small beaded cable tie was deployed on each of three
281 branches of each ramet as a benchmark. The length of the branch tip from this benchmark was
282 measured in situ over this six-month period, averaged over branches and ramets, and converted
283 to an annualized rate of linear branch extension equal to 4.441 (± 2.64 cm Stdev) cm/year. The
284 annual increment in colony diameter was assumed to be twice the branch extension rate, 8.882
285 cm/year. The maximum measured diameter of a ramet of this genet was 270 cm then the colony

286 must have been growing for at least 30.4 years. This results in a maximum mutation rate of
287 1.195^{-04} per locus per year.

288 We turned to the geological record to establish a minimum mutation rate. C14 dates from
289 cores taken at Looe Key in Florida put the start-up of *A. palmata* reef growth at the base of
290 present-day shallow spur & reef zone at around 6,500 ybp (Lidz *et al.* 1985). Our clone with the
291 highest π_k value is from Looe Key in Florida (Supplemental Table 2), thus assumed to be the
292 oldest, and the minimum mutation rate can be calculated by setting this clone at a maximum age
293 of 6,500 years. This results in a minimum mutation rate of 1.542^{-05} per locus per year. This is
294 likely a maximal estimate because reef growth may not have been continuous at Looe Key.

295

296 Results

297 *Identification of mutation type (somatic vs. chimera)*

298 There were three samples in two genets (2 samples in genet P2445 from Looe Key,
299 Florida and 1 sample in genet P2151 from Molasses Reef, Florida) out of 90 genets with at least
300 5 ramets (comprising 1294 samples), that differed by more than 60% in their major cluster
301 assignment from other ramets of the genet (Fig 4). Therefore, the majority of samples (98%)
302 showing three alleles were determined to be the result of somatic mutations rather than
303 chimerism (Fig 4).

304 *Somatic Mutations*

305 Genets with at least two ramets were included in the mutational step analysis. Of the
306 3352 samples genotyped, 1387 ramets of 147 genets satisfied this requirement across the
307 Caribbean and Florida. We found 342 unique mutational changes across the 5 microsatellite loci
308 (Table 3). Of the 342 somatic mutations, 305 involved a one-step increase (150) or decrease
309 (155), with an additional 14 one-step mutations in which direction could not be determined due
310 to the mutated allele size being equidistant from each parental allele (for example 163/169
311 parental genotype with mutated allele 166). This results in 93% of the mutations being either a
312 one-step increase or decrease further supporting the explanation of somatic mutation for the 3rd
313 alleles. The remaining 22 mutations were the result of either multi-step changes or, in one case,
314 involved the loss of heterozygosity.

This article is protected by copyright. All rights reserved

315 An important factor contributing to a microsatellite mutation rate is the repeat length; the
316 more repeat units, the greater the opportunity for replication slippage. The five loci used here had
317 repeat lengths from 10 to 28 trinucleotide repeats (Fig 3A). As expected, with increasing repeat
318 length the number of unique mutations observed at a locus increased linearly (slope =6.465 +/-
319 0.473 SE, $F_{1,3}=186.633$, $p<0.001$, adjusted (adj.) $R^2=0.979$, Fig 3A). [This result has also been
320 confirmed in experiments with trinucleotides in humans where the mutation rate for 28-31 repeat
321 lengths was more than 4 times that seen for 20-22 repeat lengths (Zhang *et al.* 1994).] When
322 considering all loci together, and designating allele 1 as the smaller allele in an individual and
323 allele 2 as the larger, there are more mutations found in allele 2 (213) than allele 1 (97) (Fig 3B,
324 excluding the 14 mutations in which the mutated allele could not be determined, 17 mutations in
325 homozygotes, and the 1 mutation determined to be a loss of heterozygosity).

326 Most colonies within our collection were only sampled once, however 11 colonies from
327 Florida were resampled in 2011 and 2014 at 2-8 locations within the colony (these samples were
328 not included in any other analysis, Supplementary Table 3). There were five colonies from Sand
329 Island and Molasses reefs in Florida that had no mutations when initially sampled from 2005-
330 2009 and re-analysis in 2011 and 2014 also showed no mutations (average $n=4.6$ samples per
331 colony). One colony from Sand Island had multiple alleles at locus 166 of 149/173/176 bp in
332 2007. The same three alleles were found in the additional sampling throughout the colony ($n=4$)
333 in 2011. In two colonies, multiple alleles were not recovered when resampled ($n=8$). In three
334 colonies intracolony variation was observed: in one case a mutation was found in only half the
335 samples from one colony. In the other two colonies, a new mutation was recovered in some
336 samples, with the original mutation(s) varying throughout replicate samples (Supplementary
337 Table 2, Supplementary Fig 4). Thus, sampling a colony once may cause an underestimation of
338 mutational load due to intracolony variation in some colonies (Supplementary Table 2).

339 *Clonal Richness vs. Mosaicism*

340 Clonal richness ranged from 0 to 1 and is directly proportional to the number of sexual
341 recruits. The proportion of non-mosaic genotypes (i.e. those with only bi-allelic loci) increased
342 with increasing genotypic diversity of the *A. palmata* stand (Fig. 5A) considering a total sample
343 size of 3352 from 13 regions. However, we were concerned that this result may be due to a

344 greater power of detection in genets with more ramets. Therefore, we limited our analysis to
345 colonies that were sampled on three spatial scales (5, 10, and 15 m radii) using a random
346 sampling procedure (Baums *et al.* 2006a) to detect both common and rare genets, resulting in
347 486 total samples from 7 regions. Again the proportion of non-mosaic genotypes increased with
348 increasing genotypic diversity when only considering reefs sampled with similar sampling effort
349 (Fig. 5B). Therefore, mosaicism appears to be more common on reefs dominated by asexual
350 reproduction than those dominated by sexual recruitment.

351 A previous study showed that genotypic richness was greater and more homogeneous
352 (mean $N_g/N=0.64 \pm 0.17$) in the eastern (US Virgin Islands, St. Vincent and the Grenadines,
353 Bonaire, and Curaçao) than the western province (Florida, Bahamas, Panama, and Mexico) with
354 the exclusion of Navassa (Baums *et al.* 2006b). When comparing the proportion of non-mosaic
355 genotypes per reef between western (also including Belize, the Dominican Republic, Mona, and
356 Navassa) and eastern populations, the east had significantly more non-mosaic genets than the
357 west (Mann-Whitney U-Test, east $n=38$, west $n=48$, $p<0.001$).

358 *Growth Models*

359 The regression of π_k vs. $Sk/\sum_{i=1}^{n-1} \left(\frac{1}{i}\right)$ (Fig 6A) for the western population had a slope of
360 1.027 ± 0.1037 SE ($F_{1,66}=98.088$, $p<0.0001$, adj. $R^2=0.594$) and was not significantly different
361 from the value expected (1:1 relationship of π_k vs. $Sk/\sum_{i=1}^{n-1} \left(\frac{1}{i}\right)$) if genet size were
362 approximately constant over time with continuous ramet turnover (ANCOVA, $p=0.468$).
363 Whereas, the regression of π_k vs $2S_k/n$ (Fig 6B) for the western population had a slope of 1.194
364 ± 0.222 SE ($F_{1,66}=29.059$, $p<0.0001$, adj. $R^2=0.295$) and was significantly different from the
365 value expected (1:1 relationship of π_k vs $2S_k/n$) if the genet had been spatially expanding
366 continuously since larval settlement (ANCOVA, $p<0.0001$)

367 The regression of π_k vs $2S_k/n$ (Fig 6C) for the eastern population had a slope of $1.069 \pm$
368 0.109 SE ($F_{1,14}=95.471$, $p<0.0001$, adj. $R^2=0.863$) and was significantly different from the value
369 expected (1:1 relationship of π_k vs. $Sk/\sum_{i=1}^{n-1} \left(\frac{1}{i}\right)$) (Fig 6D) if genet size were approximately
370 constant over time with continuous ramet turnover (ANCOVA, $p<0.01$). The regression of π_k vs.
371 $Sk/\sum_{i=1}^{n-1} \left(\frac{1}{i}\right)$ for the eastern population had a slope of 0.818 ± 0.111 SE ($F_{1,14}=54.372$,

372 $p < 0.0001$, adj. $R^2 = 0.781$) and was not significantly different from the value expected (1:1
373 relationship of π_k vs $2S_k/n$) if the genet had been spatially expanding continuously since larval
374 settlement (ANCOVA, $p = 0.17$).

375 *Microsatellite Divergence Estimate of Age*

376 Estimated age calculations in the western Caribbean reefs ranged from 30-838 years old
377 (y/o) from the maximum mutation rate and 236-6500 y/o from the minimum mutation rate. Both
378 the youngest genet and the oldest genet were from reefs in Florida (Elbow and Looe Key, Table
379 4). Genets in the eastern Caribbean were from 76-627 y/o to 590-4865 y/o. An age comparison
380 between the eastern and western populations, including only genets with somatic mutations (west
381 $n = 61$, east $n = 15$) yielded no significant differences (Kruskal-Wallis Test, $p > 0.05$).

382 **Discussion**

383 Determination of genet age distribution in coral populations is important for
384 understanding demographic changes in response to environmental perturbation and ultimately for
385 understanding the evolutionary potential of these foundation species. *A. palmata*, the now
386 endangered but previously dominant shallow reef-builder in the Caribbean, lends itself to
387 somatic mutation analyses because of the importance of asexual reproduction via fragmentation
388 resulting in genets with many members. Here, we show that some *A. palmata* genets are
389 apparently of substantial age (Table 4). This was surprising, as previously only cold-water corals
390 were found to be > 1000 y/o (Table 1).

391 The Quaternary fossil record of *A. palmata* assemblages suggests that their habitat
392 tolerances and preferences have remained relatively constant through time and space (Goreau
393 1959; Shinn 1963; Gischler 2015). Consequently, the distribution of *A. palmata* on shallow-
394 water reefs has persisted through repeated glacial–interglacial cycles. Thus, at scales from
395 decades to millennia, the persistence of *A. palmata* and the assemblages they comprise has been
396 met through the capacity of those corals incrementally to track favorable environments that have
397 shifted spatially over time (Precht and Aronson personal correspondence). These geological data
398 point to the possibility of potentially millennial-age (or older) genets within modern-day
399 populations of *A. palmata*.

400 We stress that absolute genet ages derived from somatic mutations as presented here have
401 to be interpreted cautiously. Because direct measurements of microsatellite mutation rates in
402 corals are not available and probably will not be for some time, we used other evidence to
403 bracket minimum and maximum mutation rates. We assigned the highest mutation rate to the
404 genet with the smallest microsatellite divergence rate among clone members and measured the
405 growth rate of the largest colony. Growth rates of *A. palmata* can vary with season, latitude and
406 reef location, and the measured linear extension rate of 4.44 cm/year of this colony was
407 somewhat slower than published growth rate measurements of 6 – 9 cm/year from Florida and
408 across the Caribbean (Gladfelter *et al.* 1978; Lirman 2000; Bak *et al.* 2009). We set the
409 minimum mutation rate to the genet with the largest microsatellite divergence rate among clone
410 members and asked how long this genet could have existed in this location (Looe Key, Florida).
411 By turning to the published fossil record, we ascertained that *A. palmata* colonies at this location
412 could not have been more than 6,500 years old (Lidz *et al.* 1985). While it is perhaps unlikely
413 that this genet is 6,500 years old because *A. palmata* presence at this location may not have been
414 continuous over this time frame, it is a maximal estimate. The resulting mutation rates (1.195^{-04} -
415 1.542^{-05} per locus per year) fall within reported microsatellite mutation rates from 10^{-2} to 10^{-6} per
416 sexual generation (Kruglyak *et al.* 1998; Shimoda *et al.* 1999; Ellegren 2000; Hoekert *et al.*
417 2002; O'Connell & Ritland 2004; Peery *et al.* 2012) when adjusted to generational times of
418 acroporids (4-8 years, Wallace 1985). An analysis of environmental markers in extant *A.*
419 *palmata* skeletons could substantiate genet age estimates (however the oldest portion of the genet
420 may no longer exist).

421 Despite the uncertainties surrounding absolute genet age determination, relative genet age
422 comparisons across the range of *A. palmata* should still be valid and are presented here for the
423 first time.

424 *Range edge populations and dominance of asexual reproduction*

425 Sessile organisms capable of asexual reproduction are often largely clonal at the edge of
426 the species' range, both in terrestrial and marine ecosystems (Eckert 2002; Baums 2008).
427 Populations at the range margins of the marine angiosperm *Zostera marina* had clonal richness
428 values of less than 0.2 and sexual reproduction was rare or absent (Reusch & Boström 2011).

429 The marginal *A. palmata* population of Florida averaged 3.7 unique mutations per multilocus
430 genotype whereas eastern, lower latitude populations such as Bonaire, Curacao and USVI ranged
431 from 1.2-1.3 UMs per MLG, $n=1387$ (Table 3). This would mean that the Florida genets are
432 older. Nevertheless, when considering only the large clonal stands the ages were not significantly
433 different between the eastern and western populations (Table 4) suggesting a more or less similar
434 historical presence of *A. palmata* in both populations but a higher frequency of sexual renewal in
435 the East.

436 *Mosaicism due to somatic Copy Number Variations*

437 At first glance, the appearance of three alleles per locus in *A. palmata* multilocus
438 genotypes is puzzling. One explanation is gene or genome duplication (Wang *et al.* 2009;
439 Richards & Oppen 2012). However, several lines of evidence argue against this interpretation.
440 Preliminary assembly of 2 lanes of genomic sequencing data (Illumina) showed no evidence of
441 genome duplication (I.Baums pers. observ). Additionally, a chromosomal spread analysis of *A.*
442 *palmata* larvae revealed a count of $n=24$ (supplemental Fig. 5), a diploid state. The basic
443 scleractinian chromosome number is $x=14$ and $x=12$ (Kenyon 1997). Inherited, duplicated
444 genomic regions are also unlikely. In the latter case, all 5 microsatellite loci would have to be
445 located in duplicated regions as all five loci show tri-allelic genotypes, albeit usually only one
446 locus was mutated in any given sample: for genets with $n \geq 5$ ramets, 15.56% had 0 mutated loci,
447 58.89% had 1 mutated locus, 20% had 2 mutated loci, and 5.56% had 3 mutated loci. Four of the
448 five loci amplify a similar range of allele sizes in the Caribbean sister species, *A. cervicornis*.
449 Fossil records date back 6.6 (Budd & Johnson 1999) and 2.6-3.6 (McNeill *et al.* 1997) million
450 years, respectively for *A. cervicornis* and *A. palmata*. Thus, the duplication events would have to
451 have occurred before the speciation event because tri-allelic genotypes were found in both
452 species across the entire Caribbean range. Such duplicated genomic regions would have been
453 mutating separately for several million years making it unlikely that the majority of mutations
454 are just one mutation step away as observed here.

455 Genomic instability is a mechanism of aging with somatic copy number variations (CNV)
456 prevalent in many human cancers (Shlien & Malkin 2009) and somatic CNVs increase with age
457 in human blood cell genomes (Forsberg *et al.* 2012). We posit that *A. palmata* genomes

458 accumulate somatic duplications with age, resulting in multiple copies of the microsatellite loci
459 available for replication slippage (Fig 7). This led to some ramets having up to 4 alleles at a
460 single locus.

461 Recovery of tri-allelic genotypes was robust to repeated DNA extractions, and repeated
462 PCR reactions, and has been observed in other coral species (Wang *et al.* 2009) and the marine
463 angiosperm *Zostera marina* (Reusch & Boström 2011). Baums *et al.* (2005a) found triploid
464 larvae in some experimental crosses, ranging from 7 to 36% of the larvae genotyped. Larvae did
465 survive to 90 hours post fertilization but it is unknown if they would settle and grow into
466 reproductive adults. The most likely explanation for the triploid status was having a second
467 maternal allele, either due to retention of a polar body, self-fertilization or mitotic
468 parthenogenesis. Multiple alleles (3-5) were detected in 15% of Pacific Acroporids at a single
469 locus due to inherited gene duplication; in this study, all alleles in the example chromatogram
470 were greater than a one mutation step difference (130, 140, 150, 162 bp, Richards & Oppen
471 2012). Interestingly, predominately sexually reproducing coral species on the Great Barrier Reef
472 show somatic mutation in the form of two-alleles per locus (presumably generated by a single
473 slippage event without duplication) rather than three alleles (Schweinsberg *et al.* 2015). This
474 leads us to hypothesize that highly fragmenting coral species such as *A. palmata* accumulate
475 somatic CNVs over the long lifetime of the genet. Independent evidence for or against somatic
476 CNV would have to come from Fluorescent In Situ Hybridization (FISH, Langersafer *et al.*
477 1982) or through controlled crosses of gametes from a tri-allelic genet and a genet without
478 mutations within the 5 microsatellite loci, if there is not a sequestered germline. A triploid (or
479 tetraploid) state at a microsatellite locus could also stem from the mutation of cells that are able
480 to proliferate, such as stem-like cells (Reyes-Bermudez & Miller 2009), resulting in two (or
481 more) diploid cell lineages found throughout the colony.

482 *Mosaicism versus Chimerism*

483 Genetic diversity within a colony could stem from the fusion of two or more larvae or
484 juvenile corals, producing a chimera (Fig 7). Such fusion in early life stages has been observed in
485 scleractinian corals and is generally attributed to an immature immune system that is not yet able
486 to distinguish between self and non-self (Frank *et al.* 1997; Permata & Hidaka 2005; Puill-

487 Stephan *et al.* 2009). However, the prevalence of chimerism in adult colonies in the genus
488 *Acropora* is generally low (2-5%, Schweinsberg *et al.* 2015). Retrieval of genotypes that vary at
489 several loci among branches from one colony may indicate chimerism (Fig 7). A colony was
490 classified as a chimera if it differed by more than 60% in its major cluster assignment probability
491 from other members of its genet as defined by Schweinsberg *et al.* (2015). Only 0.2% of samples
492 from the 90 genets (n=1296) were classified as possible chimeras, thus making mosaicism the
493 more likely explanation for most of the observed intracolony genetic variation.

494 *Evolutionary and ecological consequences of genet longevity*

495 The presence of large, potentially centennial-aged genets within a population begs
496 questions with regard to their history as well as their adaptive potential over the coming decades
497 of rapid environmental change. It is likely that the environmental conditions in most shallow
498 coastal habitats over the lifespan of these very old genets were quite different from today, which
499 implies that these old genets 1) possess a great degree of plasticity enabling them to persist
500 throughout these environmental variations (Barshis *et al.* 2013) and/or 2) that they have in fact
501 ‘migrated’ among nearby coastal habitats over the centuries. For example, it is possible that our
502 current observation of a very old clone is in a distinct location from where it originally recruited
503 with fragments ‘migrating’ upslope in tracking slow holocene sea level change (Gischler 2015).

504 Alternatively, the General-Purpose Genotype model (Baker 1965) explains the ubiquity
505 of clonal organisms by their ability to retain the most competent genotypes over time; favoring
506 the absence of sexual reproduction once an optimal genotype is found. For example, (Van
507 Doninck *et al.* 2002) showed much higher ecological tolerances of a ubiquitous asexual ostracod
508 in comparison with additional species that were asexual and narrowly distributed or that had
509 mixed reproductive modes. If *A. palmata* genets have persisted over hundreds to thousands of
510 years, it implies persistence through substantial environmental changes, and possibly gives hope
511 that they can survive additional anticipated climate change. The overall recent declines of *A.*
512 *palmata* including declines of certain moderate-sized clones in particular (Banks *et al.* 2010)
513 suggest there is a limit to this tolerance, which may be exceeded soon.

514 However, *A. palmata* is not entirely asexual and there is also the possibility that a
515 preponderance of large, old genets is not necessarily adaptive. Potts (1984) suggested that

516 because of corals' extreme longevity, many species (or populations) have not had the
517 opportunity, since current coastal habitats became habitable, to complete adequate sexual
518 generations to reach evolutionary equilibrium. Because fecundity of corals increases with genet
519 size (senescence notwithstanding), there may be a tendency for large old clones to dominate the
520 gene pool and diminish the chances for newer genets, possibly even those better-adapted to
521 current environmental conditions, to expand. If this is true, it implies that the presence of large
522 old clones (possibly of General Purpose Genotypes) may impair the rapid adaptation needed for
523 persistence under climate change.

524 The occurrence of somatic mutations raises the question of whether they can be the target
525 of selection and rapid adaptation. Mosaicism is thought to be favored in plants because it offers
526 an advantage in the Red Queen race against pests and parasites by increasing the standing genetic
527 diversity that prevents the evolution of specific metabolic pathways that could be used to
528 overcome the defenses of the plant (Valen 1974; Gill *et al.* 1995). Mutations in the soma are
529 available for immediate selection pressure from the environment as they compete with other
530 wild-type and mutated lineages within the organism. The selection of somatic cell lineages,
531 termed intra-organismal selection (also called somatic, diplontic, or cell-lineage selection; see
532 (Buss 1983; Hughes 1989; Otto & Hastings 1998; Clarke 2011)) may have the potential for rapid
533 evolutionary change in a modular organism by allowing within-organism gene frequency
534 changes within a single generation (Klekowski & Kazarinova-Fukshansky 1984). Through the
535 displacement of the wild-type lineage, the mutation of regenerating cells can be considered
536 evolution since they are potentially heritable in clonal Cnidaria through both sexual and asexual
537 routes. Alternatively, the coexistence of multiples lineages within an organism may result in
538 intra-organismal competition or cell parasitism leading to the decrease of overall fitness (Michod
539 & Roze 1999; Pineda-Krch & Lehtilä 2004). A theoretical population model suggested that
540 strong negative selection against intra-individual mutations keeps changes of allele frequencies
541 due to somatic mutations very low (Orive 2001).

542 Currently, empirical confirmation of somatic selection has been limited. However, there
543 are many organisms that have been evolving in the absence of sex including rotifers (Welch &
544 Meselson 2000), *Artemia* (Perez *et al.* 1994) and salamanders in the genus *Ambystoma* (Hedges
545 *et al.* 1992). [See Van Oppen *et al.* (2011) for a review on somatic mutations as fuel for

546 adaptation in invertebrates]. Somatic selection has also been demonstrated experimentally in
547 plants (Breese *et al.* 1965; Whitham & Slobodchikoff 1981; Monro & Poore 2009). Somatic
548 mutations may be widespread in corals (Levitan *et al.* 2011; Schweinsberg *et al.* 2015) and
549 within mosaic *Acropora hyacinthus* colonies it was shown that transfer of intercolonial genetic
550 variation to the next generation via gametes is possible (Schweinsberg *et al.* 2013) albeit this was
551 not the case in *Orbicella* (Barfield *et al.* 2016).

552 The ability of the coral host to respond to a changing environment occurs not only
553 through genetic adaptation but also through acclimatization by varying phenotypic responses. It
554 has recently become apparent that some environmentally induced nongenetic or epigenetic
555 changes are also heritable through a process known as transgenerational acclimatization (van
556 Oppen *et al.* 2015). Epigenetic changes include histone modifications, DNA methylation,
557 chromatin remodeling, and gene regulatory mechanisms involving small noncoding RNAs
558 (Danchin *et al.* 2011). A recent study in the clonal tree poplar showed the persistent influence of
559 geographic origin on the ability to respond to stress within a common garden experiment,
560 showing that the older the clone (longer clones of the same genet lived in different
561 environmental conditions) the more divergent the transcriptomic response was to drought and the
562 greater the variation in genome methylation patterns (Raj *et al.* 2011). Although not directly
563 linked to epigenetic changes, the pacific coral *Acropora hyacinthus* (cryptic species E) was able
564 to acclimatize to new microenvironments by increasing bleaching resistance, as measured
565 through transcriptomic responses and chlorophyll A changes, without altering their abundances
566 of symbiont type (Palumbi *et al.* 2014). This imprinted “memory” of past stress responses could
567 have profound implications for asexually reproducing corals in that ramets distributed across a
568 reef could have divergent epigenetic “memories” due to varying environmental conditions such
569 as water flow, light and pathogen exposure. In addition, epigenetic changes along with somatic
570 mutations have the ability to be passed on to the next generation in organisms without segregated
571 germ lines.

572 The current paucity of clonal age estimates impairs our understanding of the ecology and
573 evolution of marine foundation fauna. These estimates are difficult to come by because size and
574 age are not related in colonial, asexually reproducing organisms. Significant asexual colony
575 reproduction occurs in at least nine coral genera and thus the decoupling of size and genet age is

576 a widespread phenomenon in corals (Supplementary Table 1). Alternative methods to estimating
577 genet age include the use of somatic mutations but without direct mutation rate measurements,
578 the uncertainty of the age estimates is considerable. Regardless, when applied to a fragmenting
579 Caribbean coral, the results point towards genet ages that rival those of the most ancient
580 organisms on earth alive today. This raises questions about their adaptive potential to a rapidly
581 changing climate. Does their past ability to survive environmental change predict future success?
582 The answer will come from experimental studies combined with demographic and theoretical
583 models.

584 **Acknowledgements**

585 Field collection in Mexico was funded by the Consejo Nacional de Ciencias y Tecnología
586 grant number 153260 (to ATB). Funding was provided by the National Science Foundation grant
587 OCE 0928764, OCE-1516763, and NOAA - National Marine Fisheries Service to IB. Samples
588 were obtained under permit numbers AN001,US107A, 0385, MH-HR-010-MEX, 3235,
589 12US784243/g.

591 **References**

- 592 Ally D, Ritland K, Otto SP (2008) Can clone size serve as a proxy for clone age? An exploration
593 using microsatellite divergence in *Populus tremuloides*. *Molecular Ecology* **17**, 4897-
594 4911.
- 595 Bak RPM, Nieuwland G, Meesters EH (2009) Coral growth rates revisited after 31 years: what is
596 causing lower extension rates in *Acropora palmata*? *Bulletin of Marine Science* **84**, 287-
597 294.
- 598 Baker HG (1965) Characteristics and modes of origin of weeds, pp. 147-168 pp. Academic
599 Press, New York & London.
- 600 Banks SC, Ling SD, Johnson CR, *et al.* (2010) Genetic structure of a recent climate change-
601 driven range extension. *Molecular Ecology* **19**, 2011-2024.

- 602 Barfield S, Aglyamova GV, Matz MV (2016) Evolutionary origins of germline segregation in
603 Metazoa: evidence for a germ stem cell lineage in the coral *Orbicella faveolata* (Cnidaria,
604 Anthozoa). *Proceedings of the Royal Society of London B: Biological Sciences* **283**.
- 605 Barki Y, Gateno D, Graur D, Rinkevich B (2002) Soft-coral natural chimerism: a window in
606 ontogeny allows the creation of entities comprised of incongruous parts. *Marine Ecology-
607 Progress Series* **231**, 91-99.
- 608 Barrett ELB, Burke TA, Hammers M, Komdeur J, Richardson DS (2013) Telomere length and
609 dynamics predict mortality in a wild longitudinal study. *Molecular Ecology* **22**, 249-259.
- 610 Barshis DJ, Ladner JT, Oliver TA, *et al.* (2013) Genomic basis for coral resilience to climate
611 change. *Proc Natl Acad Sci U S A* **110**, 1387-1392.
- 612 Baums IB (2008) A restoration genetics guide for coral reef conservation. *Molecular Ecology* **17**,
613 2796-2811.
- 614 Baums IB, Devlin-Durante MK, LaJeunesse TC (2014) New insights into the dynamics between
615 reef corals and their associated dinoflagellate endosymbionts from population genetic
616 studies. *Molecular Ecology* **23**, 4203-4215.
- 617 Baums IB, Hughes CR, Hellberg MH (2005a) Mendelian microsatellite loci for the Caribbean
618 coral *Acropora palmata*. *Marine Ecology - Progress Series* **288**, 115-127.
- 619 Baums IB, Miller MW, Hellberg ME (2005b) Regionally isolated populations of an imperiled
620 Caribbean coral, *Acropora palmata*. *Molecular Ecology* **14**, 1377-1390.
- 621 Baums IB, Miller MW, Hellberg ME (2006a) Geographic variation in clonal structure in a reef
622 building Caribbean coral, *Acropora palmata*. *Ecological Monographs* **76**, 503-519.
- 623 Baums IB, Paris CB, Cherubin LM (2006b) A bio-oceanographic filter to larval dispersal in a
624 reef-building coral. *Limnology and Oceanography* **51**, 1969-1981.
- 625 Breese E, Hayward M, Thomas A (1965) Somatic selection in perennial ryegrass. *Heredity* **20**,
626 367-379.
- 627 Brownstein MJ, Carpten JD, Smith JR (1996) Modulation of non-templated nucleotide addition
628 by tag DNA polymerase: Primer modifications that facilitate genotyping. *Biotechniques*
629 **20**, 1004-1010.
- 630 Budd AF, Johnson KG (1999) Origination preceding extinction during late Cenozoic turnover of
631 Caribbean reefs. *Paleobiology* **25**, 188-200.

- 632 Buss LW (1983) Evolution, development, and the units of selection. *Proceedings of the National*
633 *Academy of Sciences* **80**, 1387-1391.
- 634 Carvalho GR (1994) Genetics of aquatic clonal organisms. In: *Genetics and Evolution of Aquatic*
635 *Organisms*. (ed. Beaumont AR), pp. 291-323. Chapman and Hall, London.
- 636 Caspari R, Lee SH (2004) Older age becomes common late in human evolution. *Proceedings of*
637 *the National Academy of Sciences of the United States of America* **101**, 10895-10900.
- 638 Chakraborty R, Kimmel M, Stivers DN, Davison LJ, Deka R (1997) Relative mutation rates at
639 di-, tri-, and tetranucleotide microsatellite loci. *Proceedings of the National Academy of*
640 *Sciences* **94**, 1041-1046.
- 641 Clarke E (2011) Plant individuality and multilevel selection theory. *The major transitions in*
642 *evolution revisited*. MIT Press, Cambridge, 227-250.
- 643 Cloutier D, Rioux D, Beaulieu J, Schoen DJ (2002) Low rate of somatic mutation at
644 microsatellite loci in Eastern White Pine, *Pinus strobus*.
- 645 Conrad DF, Keebler JEM, DePristo MA, *et al.* (2011) Variation in genome-wide mutation rates
646 within and between human families. *Nature* **201**, 1.
- 647 Danchin É, Charmantier A, Champagne FA, *et al.* (2011) Beyond DNA: integrating inclusive
648 inheritance into an extended theory of evolution. *Nat Rev Genet* **12**, 475-486.
- 649 de Witte LC, Stöcklin J (2010) Longevity of clonal plants: why it matters and how to measure it.
650 *Annals of Botany* **106**, 859-870.
- 651 Dorken ME, Eckert CG (2001) Severely reduced sexual reproduction in northern populations of
652 a clonal plant, *Decodon verticillatus* (Lythraceae). *Journal of Ecology* **89**, 339-350.
- 653 Eckert CG (2002) The loss of sex in clonal plants. *Evolutionary Ecology* **15**, 501-520.
- 654 Eggins SM, Grün R, McCulloch MT, *et al.* (2005) In situ U-series dating by laser-ablation multi-
655 collector ICPMS: new prospects for Quaternary geochronology. *Quaternary Science*
656 *Reviews* **24**, 2523-2538.
- 657 Ellegren H (2000) Microsatellite mutations in the germline: implications for evolutionary
658 inference. *Trends in Genetics* **16**, 551-558.
- 659 Forsberg LA, Rasi C, Razzaghian HR, *et al.* (2012) Age-related somatic structural changes in the
660 nuclear genome of human blood cells. *American Journal of Human Genetics* **90**, 217-
661 228.

- 662 Frank U, Oren U, Loya Y, Rinkevich B (1997) Alloimmune maturation in the coral *Stylophora*
663 *pistillata* is achieved through three distinctive stages, 4 months post-metamorphosis.
664 *Proceedings of the Royal Society B-Biological Sciences* **264**, 99-104.
- 665 Gill DE, Chao L, Perkins SL, Wolf JB (1995) Genetic mosaicism in plants and clonal animals.
666 *Annual Review of Ecology and Systematics* **26**, 423-444.
- 667 Gischler E (2015) Quaternary reef response to sea-level and environmental change in the western
668 Atlantic. *Sedimentology* **62**, 429-465.
- 669 Gladfelter EH, Monahan RK, Gladfelter WB (1978) Growth rates of five reef-building corals in
670 the northeastern Caribbean. *Bulletin of Marine Science* **28**, 728-734.
- 671 Goreau TF (1959) The ecology of Jamaican coral reefs: Species composition and zonation.
672 *Ecology* **40**, 67-90.
- 673 Haag-Liautard C, Dorris M, Maside X, *et al.* (2007) Direct estimation of per nucleotide and
674 genomic deleterious mutation rates in *Drosophila*. *Nature* **445**, 82-85.
- 675 Halkett F, Simon JC, Balloux F (2005) Tackling the population genetics of clonal and partially
676 clonal organisms. *Trends in Ecology & Evolution* **20**, 194-201.
- 677 Hall-Spencer J, Allain V, Fossa JH (2002) Trawling damage to Northeast Atlantic ancient coral
678 reefs. *Proceedings of the Royal Society B-Biological Sciences* **269**, 507-511.
- 679 Harper JL (1977) *Population Biology of Plants* Academic Press, London.
- 680 Hedges SB, Bogart JP, Maxson LR (1992) Ancestry of unisexual salamanders.
- 681 Heinze B, Fussi B (2008) Somatic mutations as a useful tool for studying clonal dynamics in
682 trees. *Molecular Ecology* **17**, 4779-4781.
- 683 Highsmith RC (1982) Reproduction by fragmentation in corals. *Marine Ecology-Progress Series*
684 **7**, 207-226.
- 685 Hoekert WE, Neufeglise H, Schouten AD, Menken SB (2002) Multiple paternity and female-
686 biased mutation at a microsatellite locus in the olive ridley sea turtle (*Lepidochelys*
687 *olivacea*). *Heredity (Edinb)* **89**, 107-113.
- 688 Hughes RN (1989) *A functional biology of clonal animals*. Chapman and Hall, London and New
689 York.
- 690 Hughes TP, Jackson JBC (1980) Do corals lie about their age? Some demographic consequences
691 of partial mortality, fission, and fusion. *Science* **209**, 713-715.

- 692 Kays S, Harper JL (1974) The regulation of plant and tiller density in a grass sward. *Journal of*
693 *Ecology* **63**, 97-105.
- 694 Kenyon JC (1997) Models of reticulate evolution in the coral genus *Acropora* based on
695 chromosome numbers: parallels with plants. *Evolution* **51**, 756-767.
- 696 Kimura M, Ohta T (1978) Stepwise mutation model and distribution of allelic frequencies in a
697 finite population. *Proceedings of the National Academy of Science of the United States of*
698 *America* **75**, 2868-2872.
- 699 Klekowski EJ (1997) Somatic mutation theory of clonality. In: *The ecology and evolution of*
700 *clonal growth in plants* (eds. de Kroon H, van Groenendael J), pp. 227–241. Backhuys
701 Publishers, Leiden, The Netherlands.
- 702 Klekowski EJ, Godfrey PJ (1989) Ageing and mutation in plants. *Nature* **340**, 389-391.
- 703 Klekowski EJ, Jr., Kazarinova-Fukshansky N (1984) Shoot Apical Meristems and Mutation:
704 Selective Loss of Disadvantageous Cell Genotypes. *American Journal of Botany* **71**, 28-
705 34.
- 706 Kopelman NM, Mayzel J, Jakobsson M, Rosenberg NA, Mayrose I (2015) Clumpak: a program
707 for identifying clustering modes and packaging population structure inferences across K.
708 *Molecular Ecology Resources* **15**, 1179-1191.
- 709 Kruglyak S, Durrett RT, Schug MD, Aquadro CF (1998) Equilibrium distributions of
710 microsatellite repeat length resulting from a balance between slippage events and point
711 mutations. *Proceedings of the National Academy of Sciences of the United States of*
712 *America* **95**, 10774-10778.
- 713 Langersafer PR, Levine M, Ward DC (1982) Immunological method for mapping genes on
714 *Drosophila* polytene chromosomes. *Proceedings of the National Academy of Sciences of*
715 *the United States of America-Biological Sciences* **79**, 4381-4385.
- 716 Lanner RM, Connor KF (2001) Does bristlecone pine senesce? *Experimental Gerontology* **36**,
717 675-685.
- 718 Levitan DR, Fogarty ND, Jara J, Lotterhos KE, Knowlton N (2011) Genetic, spatial and
719 temporal components of precise spawning synchrony in reef building corals of the
720 *Montastraea annularis* species complex. *Evolution* **65**, 1254-1270.

- 721 Lidz BH, Robbin DM, Shinn EA (1985) Holocene carbonate sedimentary petrology and facies
722 accumulation, Looe-Key-National-Marine-Sanctuary, Florida. *Bulletin of Marine Science*
723 **36**, 672-700.
- 724 Lirman D (2000) Fragmentation in the branching coral *Acropora palmata* (Lamarck): growth,
725 survivorship, and reproduction of colonies and fragments. *Journal of Experimental*
726 *Marine Biology and Ecology* **251**, 41-57.
- 727 Maier E, Buckenmaier A, Tollrian R, Nürnberger B (2011) Intracolony genetic variation in the
728 scleractinian coral *Seriatopora hystrix*. *Coral Reefs* **31**, 505-517.
- 729 McNeill DF, Budd AF, Borne PF (1997) Earlier (Late Pliocene) first appearance of the
730 Caribbean reef-building coral *Acropora palmata*: Stratigraphic and evolutionary
731 implications. *Geology* **25**, 891-894.
- 732 Michod RE, Roze D (1999) Cooperation and conflict in the evolution of individuality. III.
733 Transitions in the unit of fitness. *Lectures on Mathematics in the Life Sciences*, 47-92.
- 734 Monroe K, Poore AG (2009) The Potential for Evolutionary Responses to Cell-Lineage Selection
735 on Growth Form and Its Plasticity in a Red Seaweed. *The American Naturalist* **173**, 151-
736 163.
- 737 O'Connell LM, Ritland K (2004) Somatic mutations at microsatellite loci in western redcedar
738 (*Thuja plicata* : Cupressaceae). *Journal of Heredity* **95**, 172-176.
- 739 Okubo N, Motokawa T, Omori M (2007) When fragmented coral spawn? Effect of size and
740 timing on survivorship and fecundity of fragmentation in *Acropora formosa*. *Marine*
741 *Biology* **151**, 353-363.
- 742 Orive ME (2001) Somatic mutations in organisms with complex life histories. *Theoretical*
743 *Population Biology* **59**, 235-249.
- 744 Otto SP, Hastings IM (1998) Mutation and selection within the individual. *Genetica* **102**, 507-
745 524.
- 746 Palumbi SR, Barshis DJ, Traylor-Knowles N, Bay RA (2014) Mechanisms of reef coral
747 resistance to future climate change. *Science* **344**, 895-898.
- 748 Peery MZ, Kirby R, Reid BN, *et al.* (2012) Reliability of genetic bottleneck tests for detecting
749 recent population declines. *Molecular Ecology* **21**, 3403-3418.

- 750 Perez ML, Valverde JR, Batuecas B, *et al.* (1994) Speciation in the *Artemia* genus:
751 mitochondrial DNA analysis of bisexual and parthenogenetic brine shrimps. *Journal of*
752 *Molecular Evolution* **38**, 156-168.
- 753 Permata WD, Hidaka M (2005) Ontogenetic changes in the capacity of the coral *Pocillopora*
754 *damicornis* to originate branches. *Zoological Science* **22**, 1197-1203.
- 755 Pineda-Krch M, Lehtilä K (2004) Costs and benefits of genetic heterogeneity within organisms.
756 *Journal of Evolutionary Biology* **17**, 1167-1177.
- 757 Potts DC (1984) Generation Times and the Quaternary Evolution of Reef-Building Corals.
758 *Paleobiology* **10**, 48-58.
- 759 Pritchard JK, Stephens M, Donnelly P (2000) Inference of population structure using multilocus
760 genotype data. *Genetics* **155**, 945-959.
- 761 Prouty NG, Roark EB, Buster NA, Ross SW (2011) Growth rate and age distribution of deep-sea
762 black corals in the Gulf of Mexico. *Marine Ecology Progress Series* **423**, 101-U121.
- 763 Puill-Stephan E, Willis BL, van Herwerden L, van Oppen MJH (2009) Chimerism in Wild Adult
764 Populations of the Broadcast Spawning Coral *Acropora millepora* on the Great Barrier
765 Reef. *PLoS ONE* **4**, e7751.
- 766 Radtke U, Schellmann G, Scheffers A, *et al.* (2003) Electron spin resonance and radiocarbon
767 dating of coral deposited by Holocene tsunami events on Curaçao, Bonaire and Aruba
768 (Netherlands Antilles). *Quaternary Science Reviews* **22**, 1309-1315.
- 769 Raj S, Bräutigam K, Hamanishi ET, *et al.* (2011) Clone history shapes *Populus* drought
770 responses. *Proceedings of the National Academy of Sciences* **108**, 12521-12526.
- 771 Reusch TH, Boström C (2011) Widespread genetic mosaicism in the marine angiosperm *Zostera*
772 *marina* is correlated with clonal reproduction. *Evolutionary Ecology* **25**, 899-913.
- 773 Reyes-Bermudez A, Miller DJ (2009) In vitro culture of cells derived from larvae of the staghorn
774 coral *Acropora millepora*. *Coral Reefs* **28**, 859-864.
- 775 Richards ZT, Oppen M (2012) Rarity and genetic diversity in Indo-Pacific *Acropora* corals.
776 *Ecology and Evolution* **2**, 1867-1888.
- 777 Richards ZT, Shen C-C, Hobbs J-PA, *et al.* (2015) New precise dates for the ancient and sacred
778 coral pyramidal tombs of Leluh (Kosrae, Micronesia). *Science Advances* **1**, e1400060.

779 Roark EB, Guilderson TP, Dunbar RB, Fallon SJ, Mucciarone DA (2009) Extreme longevity in
780 proteinaceous deep-sea corals. *Proceedings of the National Academy of Sciences* **106**,
781 5204-5208.

782 Roark EB, Guilderson TP, Flood-Page S, *et al.* (2005) Radiocarbon-based ages and growth rates
783 of bamboo corals from the Gulf of Alaska. *Geophysical Research Letters* **32**.

784 Robinson JD, Haag CR, Hall DW, Pajunen I, Wares JP (2012) Genetic Estimates of Population
785 Age in the Water Flea, *Daphnia magna*. *Journal of Heredity* **103**, 887-897.

786 Santelices B (1999) How many kinds of individual are there? *Trends in Ecology & Evolution* **14**,
787 152-155.

788 Schug MD, Hutter CM, Wetterstrand KA, *et al.* (1998) The mutation rates of di-, tri- and
789 tetranucleotide repeats in *Drosophila melanogaster*. *Molecular Biology and Evolution* **15**,
790 1751-1760.

791 Schweinsberg M, González Pech RA, Tollrian R, Lampert KP (2013) Transfer of intracolony
792 genetic variability through gametes in *Acropora hyacinthus* corals. *Coral Reefs*, 1-11.

793 Schweinsberg M, Weiss LC, Striewski S, Tollrian R, Lampert KP (2015) More than one
794 genotype: how common is intracolony genetic variability in scleractinian corals?
795 *Molecular Ecology* **24**, 2673-2685.

796 Shimoda N, Knapik EW, Ziniti J, *et al.* (1999) Zebrafish Genetic Map with 2000 Microsatellite
797 Markers. *Genomics* **58**, 219-232.

798 Shinn EA (1963) Spur and groove formation on the Florida Reef Tract. *Journal of Sedimentary*
799 *Petrology* **33**.

800 Shlien A, Malkin D (2009) Copy number variations and cancer. *Genome Medicine* **1**.

801 Slatkin M (1996) Gene genealogies within mutant allelic classes. *Genetics* **143**, 579-587.

802 Soong K, Lang JC (1992) Reproductive Integration in Reef Corals. *The Biological Bulletin* **183**,
803 418-431.

804 Szmant AM (1986a) Reproductive ecology of Caribbean reef corals. *Coral Reefs* **5**.

805 Szmant AM (1986b) Reproductive ecology of Caribbean reef corals. *Coral Reefs* **5**, 43-53.

806 Valen L (1974) Molecular evolution as predicted by natural selection. *Journal of Molecular*
807 *Evolution* **3**, 89-101.

- 808 Van Doninck K, Schön I, De Bruyn L, Martens K (2002) A general purpose genotype in an
809 ancient asexual. *Oecologia* **132**, 205-212.
- 810 Van Oppen MJ, Souter P, Howells EJ, Heyward A, Berkelmans R (2011) Novel genetic diversity
811 through somatic mutations: fuel for adaptation of reef corals? *Diversity* **3**, 405-423.
- 812 van Oppen MJH, Oliver JK, Putnam HM, Gates RD (2015) Building coral reef resilience
813 through assisted evolution. *Proceedings of the National Academy of Sciences* **112**, 2307-
814 2313.
- 815 Vasek FC (1980) Creosote Bush: Long-Lived Clones in the Mojave Desert. *American Journal of*
816 *Botany* **67**, 246-255.
- 817 Wallace CC (1985) Reproduction, recruitment and fragmentation in 9 sympatric species of the
818 coral genus *Acropora*. *Marine Biology* **88**, 217-233.
- 819 Wang S, Zhang LL, Meyer E, Matz MV (2009) Construction of a high-resolution genetic linkage
820 map and comparative genome analysis for the reef-building coral *Acropora millepora*.
821 *Genome Biology* **10**.
- 822 Welch DBM, Meselson M (2000) Evidence for the evolution of bdelloid rotifers without sexual
823 reproduction or genetic exchange. *Science* **288**, 1211-1215.
- 824 Whitham TG, Slobodchikoff C (1981) Evolution by individuals, plant-herbivore interactions,
825 and mosaics of genetic variability: the adaptive significance of somatic mutations in
826 plants. *Oecologia* **49**, 287-292.
- 827 Williams DE, Miller MW (2012) Attributing mortality among drivers of population decline in
828 *Acropora palmata* in the Florida Keys (USA). *Coral Reefs* **31**, 369-382.
- 829 Work TM, Forsman ZH, Szabó Z, *et al.* (2011) Inter-Specific Coral Chimerism: Genetically
830 Distinct Multicellular Structures Associated with Tissue Loss in *Montipora capitata*.
831 *PLoS ONE* **6**, e22869.
- 832 Zhang L, Leeflang EP, Yu J, Arnheim N (1994) Studying human mutations by sperm typing:
833 instability of CAG trinucleotide repeats in the human androgen receptor gene. *Nature*
834 *Genetics* **7**, 531-535.
- 835

836 **Data Accessibility**

837 Multilocus genotypes are available at DRYAD: <http://dx.doi.org/10.5061/dryad.f6600>

838 **Author contributions**

839 MD and IB designed the study and wrote the manuscript with key input from MM and WP. MD
840 analyzed and interpreted the data. Funding was provided and samples were collected by IB and
841 the Caribbean *Acropora* Research Group.

842

843

844 **Tables**

845 *Table 1* Published age estimates of coral colonies.

Species	Age Estimate (years)	Method	Region	Depth	Year Collected	Reference
<i>Leiopathes</i>	70 - 2040	C14 and growth ring measurements	Gulf of Mexico	304-317m	Not stated	(Prouty <i>et al.</i> 2011)
<i>Gerardia sp.</i> <i>Leiopathes</i>	300 - 2700 350 - 4200	$\delta^{13}C$	Hawaii	400-500m	2004	(Roark <i>et al.</i> 2009)
<i>Keratoisis</i> , <i>Isidella</i> , or <i>Acanella spp.</i>	75–126	C14	Gulf of Alaska	634-720m	2002	(Roark <i>et al.</i> 2005)
<i>Lophelia pertusa</i>	451 \pm 36	C14	West Ireland	840–1300m	1995-1997	(Hall-Spencer <i>et al.</i> 2002)
<i>Pocillopora</i>	3.69 \pm 0.48	U/Th	Kosrae and Lelu	unknown	2012	(Richards <i>et al.</i>

<i>verruoca</i>	3.82 ± 0.55		Island			2015)
	3.89 ± 0.42					

846

847

848

849

850 **Table 2** Summary table of *Acropora palmata* samples used in the various analyses. MLG = multilocus
851 genotype. UM =Unique Mutations. *Puerto Rico contains admixed *A. palmata* genets between the eastern
852 and western Caribbean.

		Clonal Richness vs Non-mosaic samples: MLGs with n≥1 ramets	Mutational Analysis: MLGs with n≥2 ramets				Genet Age Analysis: MLGs with n≥5 ramets	
	Region	Samples	Samples	MLGs	UM	UM/MLG	Samples	MLGs
East	Bonaire	43	8	3	4	1.3	0	0
	Curacao	286	73	17	20	1.2	55	7
	Puerto Rico*	308	41	12	16	1.3	46	7
	SVG	210	33	12	18	1.5	10	2
	USVI	464	65	9	14	1.6	64	7
West	Bahamas	259	134	23	46	2.0	131	17
	Belize	152	16	4	8	2.0	5	1
	Cuba	2	0	0	0	NA	0	0
	Dom. Rep.	49	4	1	2	2.0	0	0
	Florida	1036	892	47	175	3.7	931	44
	Mexico	180	33	3	7	2.3	0	0
	Mona	70	18	3	11	3.7	0	0
	Navassa	176	21	8	12	1.5	0	0
	Panama	117	49	5	9	1.8	52	5
	TOTAL	3352	1387	147	342		1294	90

853

854

855 **Table 3** Ancestral alleles could be determined for some *A. palmata* genets with only two ramets. A =
 856 allele size, bp = basepairs.

Clonal ID	Database ID	Locus	A1 (bp)	A2 (bp)	Mutated allele (bp)	2nd mutated allele (bp)
P2635	4597	192	166	175	169	
P2635	4602	192	166	175	172	178
P2634	1643	192	166	181	163	
P2634	1644	192	166	181	178	
P1084	1601	192	160	181	178	
P1084	1602	192	160	181	157	

857

858

859 **Table 4** Calculated age of *Acropora palmata* genets from throughout the Caribbean and north-west
 860 Atlantic. N is the number of ramets, π_k is microsatellite divergence. CI = confidence interval. SVG = St.
 861 Vincent and the Grenadines. USVI = US Virgin Islands.

Region	Reef	Clonal ID	N	π_k	Oldest Age (years)	Youngest Age (years)	Within a 5% CI around growth model
Bahamas	BlackBouy	P1100	5	0.000	< 254	< 30	Yes
	BockCay	P1106	10	0.080	1397	167	Yes
		P1110	5	0.160	2794	335	Yes
	CharliesBeach	P1089	15	0.053	931	112	Yes
	GreatIguana	P1042	11	0.145	2540	304	Yes
		P1043	7	0.267	4657	558	No
	HallsPond	P1130	6	0.173	3027	363	Yes

Region	Reef	Clonal ID	N	π_k	Oldest Age (years)	Youngest Age (years)	Within a 5% CI around growth model
	LittleDarby	P1112	12	0.067	1164	139	Yes
	MiddleBeach	P1079	9	0.100	1746	209	No
		P1080	5	0.120	2095	251	Yes
	NairnCay	P2365	6	0.000	< 254	< 30	Yes
	PerryShallow	P1073	7	0.057	998	120	Yes
		P1075	6	0.067	1164	139	Yes
		P2475	5	0.080	1397	167	Yes
		P1148	6	0.293	5122	613	No
		P1123	7	0.057	998	120	Yes
		P1122	9	0.000	< 254	< 30	Yes
Belize	GSTF12	P2276	5	0.120	2095	251	Yes
Curacao	BlueBay	P2161	5	0.080	1397	167	Yes
		P1200	11	0.036	635	76	Yes
	EastPoint	P1258	5	0.080	1397	167	Yes
		P1244	11	0.073	1270	152	Yes
	SeaAquarium	P1199	7	0.000	< 254	< 30	Yes
		P1232	5	0.000	< 254	< 30	Yes
		P2194	11	0.109	1905	228	Yes
Florida	Boomerang	P1040	10	0.040	698	84	Yes
	Carrysfort	P2115	17	0.092	1609	193	No
		P2118	41	0.137	2385	286	No
		P2121	24	0.049	848	102	Yes
		P2591	11	0.102	1778	213	No
	Elbow	P1028	55	0.015	254	30	Yes
		P1029	6	0.067	1164	139	Yes

Region	Reef	Clonal ID	N	π_k	Oldest Age (years)	Youngest Age (years)	Within a 5% CI around growth model
		P1030	7	0.000	< 254	< 30	Yes
		P1033	7	0.152	2661	319	Yes
		P1032	30	0.256	4469	535	No
		P2122	8	0.136	2370	284	No
		P2123	16	0.265	4628	554	No
		P2126	27	0.135	2357	282	No
	French	P2539	6	0.067	1164	139	Yes
		P2538	20	0.261	4559	546	No
		P2128	54	0.126	2206	264	No
		P2564	24	0.178	3113	373	No
	GrecianRocks	P2582	19	0.042	735	88	Yes
		P1034	14	0.057	998	120	Yes
	Horseshoe	P1000	25	0.113	1967	236	No
		P2559	7	0.114	1996	239	Yes
	KeyLargoDR	P2132	14	0.202	3531	423	No
		P2134	13	0.254	4433	531	No
		P2138	14	0.110	1919	230	Yes
		P2139	6	0.133	2328	279	No
		P2597	5	0.200	3492	418	No
	LittleGrecian	P1026	5	0.080	1397	167	Yes
		P1001	24	0.032	557	67	Yes
	LooeKey	P2427	28	0.052	915	110	Yes
		P2429	31	0.401	7000	838	No
		P2445	29	0.140	2452	294	No
	Marker3	P1039	52	0.046	801	96	Yes

Region	Reef	Clonal ID	N	π_k	Oldest Age (years)	Youngest Age (years)	Within a 5% CI around growth model
Panama	Molasses	P2151	25	0.207	3621	434	No
		P2146	32	0.150	2619	314	No
	RockKey	P1018	5	0.080	1397	167	Yes
		P1017	16	0.115	2008	241	No
	SandIsland	P1007	9	0.044	776	93	Yes
		P1002	96	0.094	1641	196	No
		P1003	29	0.216	3776	452	No
		P1021	5	0.000	< 254	< 30	Yes
	Triangle	P2416	38	0.087	1525	183	No
	WesternSambo	P1012	8	0.044	776	93	Yes
		P1011	11	0.108	1881	225	Yes
		P1008	8	0.000	< 254	< 30	Yes
	BastimentosI	P1150	16	0.065	1135	136	Yes
		Bocas Del Drago	P1168	15	0.076	1330	159
	P1167		5	0.220	3842	460	No
Tobobe West I	P1183	6	0.107	1863	223	No	
Wild Cayne	P1177	10	0.040	698	84	Yes	
Puerto Rico	Cayo Ron	P2286	6	0.173	3027	363	Yes
		P2294	5	0.000	< 254	< 30	Yes
	La Cordillera	P2301	8	0.000	< 254	< 30	Yes
		P2334	5	0.000	< 254	< 30	Yes
		P2339	5	0.000	< 254	< 30	Yes
	San Cristobal	P1857	10	0.204	3570	428	Yes
P1878		7	0.000	< 254	< 30	Yes	
SVG	Mustique	P1667	5	0.080	1397	167	Yes

Region	Reef	Clonal ID	N	π_k	Oldest Age (years)	Youngest Age (years)	Within a 5% CI around growth model
		P1668	5	0.300	5239	627	Yes
USVI	Grounding VI	P1430	5	0.000	< 254	< 30	Yes
	Hawksnest Bay	P1399	30	0.076	1325	159	Yes
		P1403	5	0.080	1397	167	Yes
		P1402	6	0.120	2095	251	Yes
		P1406	6	0.133	2328	279	Yes
	Salt Pond	P1555	5	0.120	2095	251	Yes
	Tague Bay	P2504	7	0.095	1663	199	Yes

Author Manuscript

862 **Figures**

863 **Figure 1** Diagram depicting (A) the formation of a chimera from the settlement and fusion of
864 gametes of different genets. (B) An illustration of asexual reproduction by fragmentation and the
865 accumulation of mutations with age. See Supplemental Figure 1 for a photo time series of
866 fragmentation. Example alleles at one locus are given in basepairs (three digit numbers separated
867 by forward slashes). Diagram not to scale.

868
869 **Figure 2** Samples of *Acropora palmata* were collected throughout Florida and the Caribbean.
870 DR = Dominican Republic, USVI = U.S. Virgin Islands, SVG = St. Vincent and the Grenadines.
871 See (Baums *et al.* 2005b, 2006a) for sampling location details.

872
873 **Figure 3** Mutation Step Analysis. In panel (A), as the repeat length of a microsatellite locus
874 increases, the total number of unique mutations found within each locus increases linearly (slope
875 = 6.47 ± 0.47 SD, $F_{2,3}=186.6328$, $p=0.0008$, adj. $R^2=0.98$). (B) Most mutations were one step
876 away from the ancestral allele size (i.e. +/- 3 bp) with allele 1 (the smaller of the two alleles)
877 showing more repeat unit losses than gains and the larger allele (allele 2) showing more gains
878 than losses of repeat units. 29 mutations were excluded from (B). 28 mutations were excluded
879 because the mutation step was equidistant for allele 1 and 2 so that the mutated allele could not
880 be determined; 1 mutation was a dropped allele.

881
882 **Figure 4** Assignment of ramets to genets using Bayesian clustering analysis in *A. palmata*.
883 Included were all genets with $n \geq 5$ ramets (Table 2). Black lines above graphs indicate samples
884 that have mutations. An asterisk indicates colonies that have a <40% assignment probability to
885 the most closely related genet. These colonies are possible chimeras. Probability of membership
886 to a given cluster (Y-axis) is plotted for each sample (X-axis). Colors indicate cluster
887 membership for each panel (A – D). Genets from the eastern Caribbean are shown panel A,
888 genets from the western Caribbean in panels B – D. Florida was split into two groups (C, D)
889 because of the large number of genets from this region.

890

891 **Figure 5** The proportion of non-mosaic genets per reef as a function of clonal richness at each
892 reef. A) Total sample size of 3352 colonies from 86 reefs within 13 different regions across the
893 Caribbean with $n \geq 10$ colonies reef⁻¹. Exponential Rise to Maximum, Single, 2 Parameter
894 equation: $f = 0.8763 * (1 - \exp(-3.9422 * x))$ (adjusted $R^2 = 0.6495$). B) Including only colonies that
895 were sampled on three spatial scales (5, 10, and 15 m radii plots) using a random sampling
896 procedure (described in Baums *et al.* 2005a) for a total of 486 total samples from 7 regions.
897 Exponential Rise to Maximum, Single, 2 Parameter equation: $f = 1.0192 * (1 - \exp(-2.4822 * x))$
898 (adjusted $R^2 = 0.7575$).

899

900 **Figure 6** A comparison of two growth models for the western (Panel A,B) and eastern (Panel
901 C,D) Caribbean. The western Caribbean population included Florida, Bahamas, Panama and
902 Belize. The eastern Caribbean population included Curacao, US Virgin Islands, and St. Vincent
903 and the Grenadines. Panel (A,C): In a constant population model with continuous ramet turn-
904 over, the slope of π_k vs. $S_k / \sum_{i=1}^{n-1} (\frac{1}{i})$ would exhibit a 1:1 relationship (dotted line). Panel
905 (B,D): In a population that is growing in size, the slope of π_k vs $2S_k/n$ should exhibit a 1:1
906 relationship (dotted line). See text for statistical analysis.

907


908 **Figure 7** Diagram depicting how duplication of a microsatellite (msat) locus (yellow) leads to
909 copy number variation (CNV) on chromosomes (blue) in a diploid species. Once a locus is
910 duplicated, the microsatellite repeats (orange/white) may mutate through slippage of the DNA
911 polymerase during mitotic replication leading to the detection of three alleles in
912 electropherograms. With time, alleles on both chromosomes may duplicate and mutate leading to
913 detection of four alleles per samples (not shown). Allele sizes are given in basepairs. Diagram
914 not to scale.

915


916

917

A)




B)




918
919
920
921

Figure 1




922
923
924
925

Figure 2




926

927 **Figure 3**




928
929
930

Figure 4



931
932

Figure 5



933

934

Figure 6

935

936

937


938

939

940

941

942




943



944


Figure 7

Author Manuscript




mec_13865_f2.eps







mec_13865_f4.eps

A.



B.





mec_13865_f6.eps

