

1 Variability in fin whale (*Balaenoptera physalus*) occurrence in the Bering Strait
2 and southern Chukchi Sea in relation to environmental factors

3
4 Erica Escajeda^{a,*}, Kathleen M. Stafford^b, Rebecca A. Woodgate^b, Kristin L. Laidre^{a,b}

5
6
7 ^a*School of Aquatic and Fishery Sciences, University of Washington, Seattle, WA, USA*

8 ^b*Applied Physics Laboratory, University of Washington, Seattle WA, USA*

9
10 *Corresponding author. Tel: +1 303-483-3717
11 E-mail address: escajeda@uw.edu (E. Escajeda)

12
13

14 **Abstract**

15

16 Fin whales (*Balaenoptera physalus*) are common summer visitors to the Pacific Arctic,
17 migrating through the Bering Strait and into the southern Chukchi Sea to feed on seasonally-
18 abundant prey. The abundance and distribution of fin whales in the Chukchi Sea varies from
19 year-to-year, possibly reflecting fluctuating environmental conditions. We hypothesized that fin
20 whale calls were most likely to be detected in years and at sites where productive water masses
21 were present, indicated by low temperatures and high salinities, and where strong northward
22 water and wind velocities, resulting in increased prey advection, were prevalent. Using acoustic
23 recordings from three moored hydrophones in the Bering Strait region from 2009–2015, we
24 identified fin whale calls during the open-water season (July–November) and investigated
25 potential environmental drivers of interannual variability in fin whale presence. We examined
26 near-surface and near-bottom temperatures (T) and salinities (S), wind and water velocities
27 through the strait, water mass presence as estimated using published T/S boundaries, and
28 satellite-derived sea surface temperatures and sea-ice concentrations. Our results show

29 significant interannual variability in the acoustic presence of fin whales with the greatest
30 detections of calls in years with contrasting environmental conditions (2012 and 2015). Colder
31 temperatures, lower salinities, slower water velocities, and weak southward winds prevailed in
32 2012 while warmer temperatures, higher salinities, faster water velocities, and moderate
33 southward winds prevailed in 2015. Most detections (96%) were recorded at the mooring site
34 nearest the confluence of the nutrient-rich Anadyr and Bering Shelf water masses, ~35 km north
35 of Bering Strait, indicating that productive water masses may influence the occurrence of fin
36 whales. The disparity in environmental conditions between 2012 and 2015 suggests there may be
37 multiple combinations of environmental factors or other unexamined variables that draw fin
38 whales into the Pacific Arctic.

39

40 *Keywords:* Arctic, Bioacoustics, Fin whales, Chukchi Sea, Bering Strait, Interannual variability,
41 Environmental factors, Marine ecology, Marine mammals

42

43

44 **1. Introduction**

45

46 The Arctic has undergone unprecedented environmental shifts as a result of climate
47 warming (Post et al., 2019). Prominent among these shifts is the loss of sea-ice cover during the
48 summer (Comiso et al., 2008; Cavalieri and Parkinson, 2012; Vaughan et al., 2013; Wood et al.,
49 2015a,b; Walsh et al., 2017) along with earlier melting in the spring and delayed onset of
50 freezing in the fall (Markus et al., 2009; Stroeve et al., 2014; Frey et al., 2015; Stabenon et al.,
51 2019; Baker et al., this issue). Environmental shifts as a result of climate change are especially

52 evident in the Chukchi Sea where annual sea-ice cover has declined by ~13 days each decade
53 from 1979 to 2013 (Laidre et al., 2015), extending the open-water season (Grebmeier et al.,
54 2010; Stroeve et al., 2014; Wood et al., 2015b; Woodgate, 2018). Declining sea ice is expected
55 to result in range expansions of temperate and subarctic species into the Arctic (Root et al., 2003;
56 Wassmann et al., 2011; Laidre and Heide-Jørgensen, 2012; Woodgate et al., 2015). Subarctic
57 cetaceans, such as fin whales (*Balaenoptera physalus*), are thought to be expanding their range
58 and residence time in the Chukchi Sea (Woodgate et al., 2015), which could lead to increased
59 competition with Arctic cetaceans (Clarke et al., 2013).

60 Fin whales are a cosmopolitan mysticete whose range extends through most of the
61 world's oceans (Mizroch et al., 1984). Though their exact migration patterns are unclear, fin
62 whales are thought to breed in lower latitudes during winter and migrate to high-latitude areas,
63 such as the Bering and Chukchi seas, in summer to feed on seasonally abundant prey (Mizroch et
64 al., 1984; Mizroch et al., 2009). Fin whale diets vary seasonally and spatially across the North
65 Pacific, but typically include euphausiids and forage fish species (Pike, 1950; Nemoto, 1959;
66 Nemoto and Kasuya, 1965; Mizroch et al., 1984; Flinn et al., 2002; Witteveen and Wynne,
67 2016). Fin whales are generally thought to avoid sea ice, though they have been observed
68 swimming along the ice edge in the Arctic (Sleptsov, 1961; Mizroch et al., 1984).

69 Fin whales produce low frequency signals (< 100 Hz), with high intensities (source levels
70 up to 189 dB re 1 μ Pa at 1m) and short durations (\leq 1 s; Watkins, 1981; Watkins et al., 1987;
71 Širović et al., 2007). The most commonly documented call is a short (~1 s) down-sweep
72 generally starting around 25 Hz and ending at 15 Hz with peak energy centered near 20 Hz
73 (Watkins, 1981; Watkins et al., 1987). The fin whale “20-Hz pulse” can occur in regular
74 sequences, forming a stereotyped song that lasts from < 1 hr to ~33 hrs (Watkins et al., 1987).

75 Such sequences are believed to be produced by males as a mating display starting in the fall and
76 lasting through spring (Watkins et al., 2000; Croll et al., 2002; Stafford et al., 2007). Fin whales
77 also produce 20-Hz and higher frequency pulses in short, irregular sequences that may serve as
78 contact calls (Watkins, 1981; McDonald et al., 1995; Edds-Walton, 1997), especially during the
79 summer months (Širović et al., 2013).

80 Historical records dating back to the early 20th century suggest fin whales commonly
81 occurred in the southwest Chukchi Sea during the summer (Mizroch et al., 2009). Soviet and
82 Japanese whaling expeditions in the 1930–1940s captured fin whales as far west as Cape
83 Schmidt (68°55'18.3"N 179°27'42.7"W), and as far north as the central Chukchi Sea (69°04'N,
84 171°06'W) and Wrangel Island (Tomilin, 1957; Nemoto, 1959; Sleptsov, 1961; Mizroch et al.,
85 2009; Fig. 1). Fin whales were observed in the Chukchi Sea as early as June (Nikulin, 1946) and
86 stayed in the area until October (Nikulin, 1946; Nasu, 1960; Votrogov and Ivashin, 1980).
87 Sleptsov (1961) describes fin whales as ‘one of the numerous baleen whales that inhabit the
88 Chukchi Sea’ and reported seeing hundreds of fin whales in the span of six days between the
89 Bering Strait and Cape Serdtse-Kamen in September 1939. By the mid-20th century, intense
90 whaling in the North Pacific had taken a toll on fin whale populations and fin whales were rarely
91 seen in the Chukchi Sea. Only a few sightings of fin whales were recorded between 1958 and
92 1981 (Nasu, 1960; Votrogov and Ivashin, 1980; Ljungblad et al., 1982). More recent visual and
93 acoustic observations of fin whales chart their presence in portions of the northeastern Chukchi
94 Sea (Delarue et al., 2013), southcentral Chukchi Sea (Clarke et al., 2015; Brower et al., 2018),
95 and the southern Chukchi Sea north of the Bering Strait (Tsujii et al., 2016).

96 We hypothesize that observed spatial variability in fin whale presence may be connected
97 to environmental variability in the study region. In addition to the seasonal cycle of sea ice, the

98 Chukchi Sea is characterized by the presence of distinct water masses defined by differences in
99 temperature and salinity which vary from year to year (Coachman et al., 1975). The water
100 masses in the Chukchi Sea have varying levels of nutrients and chlorophyll-*a* (chl-*a*), leading to
101 distinct phytoplankton and zooplankton communities (Hopcroft et al., 2010; Eisner et al., 2013;
102 Pisareva et al., 2015; Danielson et al., 2017; Sigler et al., 2017). Large, chain-forming diatoms
103 are found in areas with high chl-*a* concentrations, such as the productive Anadyr Water (AW) in
104 the western Chukchi Sea, whereas smaller phytoflagellates occur in low-nutrient areas, such as
105 the less productive Alaskan Coastal Water (ACW) in the eastern Chukchi Sea (Springer and
106 McRoy, 1993; Eisner et al., 2013; Danielson et al., 2017). Consequently, large copepods and
107 other zooplankton groups are found in the AW while smaller copepods are ubiquitous in the
108 ACW zooplankton community (Eisner et al., 2013; Sigler et al., 2017). It might be therefore
109 expected that fin whales would occupy areas where the AW, or similarly productive water
110 masses, dominate.

111 The Chukchi Sea is a highly advective ecosystem that is heavily influenced by the inflow
112 of Pacific Water which enters through the Bering Strait (Woodgate et al., 2005a; Fig. 1).
113 Advection from the northern Bering Sea provides the main source of zooplankton for the
114 Chukchi Sea and is an important factor in determining zooplankton biomass and secondary
115 production (Weingartner, 1997; Kitamura et al., 2017). High northward water velocities through
116 the strait likely translate to increased advection of Pacific-origin prey into the Chukchi Sea.
117 Therefore, we hypothesize that years with high detections of fin whale calls will have high
118 northward (along-channel) water velocities.

119 The Bering Strait is divided into two channels by the Diomed Islands roughly mid-strait
120 (Fig. 1). The western channel of Bering Strait is comparatively cold and salty due to the

121 prevalence of the AW, while the eastern channel tends to be warmer and fresher due to the
122 presence of the ACW (Coachman et al., 1975; Woodgate et al., 2005b; Woodgate et al., 2015).
123 The cold and salty Bering Shelf Water (BSW) passes through the central strait (Coachman et al.,
124 1975; Woodgate et al., 2005b). Variability in wind strength and direction can influence the
125 position of these water masses and overall transport in the strait. Strong along-channel
126 (northward) winds through Bering Strait may push the less productive surface ACW against the
127 Alaskan coast via Ekman transport, allowing the more productive AW to shift east and replace it
128 in the surface waters (Woodgate, 2018). Similarly, southward winds spread the ACW westwards
129 across the surface of the strait and draw AW to the east at depth (Woodgate et al., 2015). Thus,
130 wind changes could affect feeding opportunities for fin whales at different depths across the
131 strait. Northward winds are linked to northward flow through the strait (Woodgate et al., 2005a),
132 which leads to higher advection of prey into the Chukchi Sea, in general. Therefore, we
133 hypothesize that fin whale occurrence may be related to northward wind velocity through the
134 strait.

135 Given that the Bering Strait is the only gateway from the Pacific Ocean into the Chukchi
136 Sea (Fig. 1), the region is an ideal study area for recording the occurrence of migrating fin
137 whales. In this paper, we investigate whether fin whales exhibited any interannual variation in
138 their acoustic presence during the open-water season (July–November) from 2009–2015 and
139 explore correlations between the acoustic presence of fin whales and environmental variation in
140 the Bering Strait region. We hypothesize that high levels of fin whale calls occur in the years
141 when and at the mooring sites where the highly productive Bering Shelf/Anadyr waters are
142 prevalent, when/where there are higher northward water velocities (and thus primarily northward

143 winds) through the strait, and in years when sea ice forms later in the fall, allowing fin whales to
144 remain in the Chukchi Sea longer into the season.

145

146

147 **2. Methods**

148

149 *2.1. Acoustic Data*

150

151 Acoustic data were collected from three AURAL-M2 hydrophones (Autonomous
152 Underwater Recorder for Acoustic Listening-Model 2, Multi-Électronique, Inc.; sensitivity of
153 -154 dB re 1 V/ μ Pa and 16-bit resolution) attached to oceanographic moorings positioned within
154 the eastern channel of the Bering Strait (A2 in the center of the eastern channel, and A4 in the
155 Alaskan Coastal Current on the east side of the channel), and a central strait location ~ 35 km
156 north of the strait in the southern Chukchi Sea (A3; Fig. 1). Hydrophones were first installed on
157 the moorings in September 2009 and recorded through 2015. Each hydrophone was positioned
158 4–8 m above the seafloor and sampled at 8192 Hz or 16384 Hz with various hourly duty cycles
159 and recording start dates (Table 1). We assume that calls recorded during the hydrophones' duty
160 cycle are representative of fin whale acoustic activity for the entire hour in which the calls were
161 recorded.

162 We quantified fin whale calling activity as the number of hours per day with fin whale
163 calls present, hereafter referred to as 'fin whale hours' (FWH). Note that since we were only able
164 to detect calling whales, we could not assume the absence of fin whales during any hour nor
165 could we estimate the abundance of fin whales using call abundance alone. The term 'recording

166 years' refers to years that each hydrophone actively recorded data. Analysis of the recordings
167 was restricted to the recording start date (typically July) until the end of November, called here
168 as the 'recording period.' Given the shallow depth of the study area, it is likely that all calls from
169 individuals within 10–20 km of the hydrophones were recorded (Woodgate et al., 2015). If we
170 use the conservative call detection range of 10 km, the hydrophones cover a total of 892 km², or
171 ~3% of the study area (Fig.1). Hydrophones at A2 and A4 cover ~64% of the eastern channel
172 area (~900 km²), while the width of the A3 10-km call detection buffer covers ~10% of the
173 across-strait distance at its latitude north of the strait.

174 We identified hours with fin whale 20-Hz pulses using the spectrogram correlation tool
175 implemented in Ishmael (2014 version; Mellinger and Clark 2000; Mellinger, 2002). Detector
176 parameters included a threshold of 10 to reduce the number of false detections and a smoothing
177 time constant of 0.3 s. Each hour identified by the detector was then manually verified to contain
178 fin whale calls by inspecting the spectrogram in Ishmael (FFT 4096, Hanning window,
179 spectrogram equalization enabled with a time constant of 30 s) and eliminating any false
180 positives from the dataset. The hour before and after a true positive FWH were examined to
181 capture any hours that were not picked up by the detector, adding a total of 269 FWH to our
182 detections (~11% of the total number of FWH for all three sites).

183 To investigate spatial and temporal patterns in the presence of fin whales, we compared
184 FWH between years and sites using a nonparametric two-sample Wilcoxon rank-sum test under
185 the null hypothesis of equal distributions. Since all hydrophones recorded in October, we
186 restricted our interannual comparisons of FWH within each mooring site and between the three
187 sites to October only to avoid issues with unequal recording period lengths. We also compared
188 the date of departure of calling fin whales from the study region by calculating the 95% quantile

189 of the cumulative distribution of days with fin whale calls starting on 1 October of each year,
190 following the procedure of Hauser et al. (2017). We used a significance threshold of 0.05 for all
191 statistical tests and assumed independence between daily values.

192

193 2.2. *Environmental Data Collection*

194

195 Six environmental variables were recorded *in-situ* by other sensors on the same
196 moorings, including: near-bottom temperature and salinity (40–55 m depth) measured by Sea-
197 Bird (SBE) SBE16 and SBE37 sensors; near-surface temperature and salinity (14–19 m depth)
198 measured by the ISCAT system developed at the University of Washington (e.g. Woodgate et al.,
199 2015), which includes a SBE37 temperature-salinity-pressure sensor in an ice-resistant housing;
200 and water velocity (cm s^{-1}) and direction ($^{\circ}$) measured by Teledyne’s Workhorse Acoustic
201 Doppler Current Profilers (ADCPs). The ADCPs measured water velocity in 2-m bins from ~15
202 m to ~45 m depth (see Supplemental Tables S1–S3 for instrument depths). For simplicity, we
203 used only data from the ADCP bin closest to ~30 m depth. Note that henceforth the term ‘near-
204 surface’ refers to measurements taken by the ISCATs and ‘near-bottom’ refers to those taken by
205 the SBEs. Some ISCAT recorders were lost/stopped recording before the 30 November cut-off
206 date (see Woodgate et al., 2015 and Supplemental Tables S1–S3 for data gaps along with other
207 mooring sensor information). Note that the ISCAT for A3 stopped recording in August 2014, 45
208 days after deployment, thus near-surface temperature and salinity data are not available for fall
209 2014.

210 In addition to the *in-situ* data, we examined northward wind velocity, and satellite-
211 derived sea surface temperatures (SST) and sea-ice concentrations. Wind velocity data were

212 obtained from the National Center for Environmental Prediction (NCEP) R1 dataset, with a
213 spatial resolution at the Bering Strait of 2.5°. We used the National Oceanic and Atmospheric
214 Administration's (NOAA) Optimum Interpolation satellite sea surface temperature (OISST)
215 gridded product with a 0.25° resolution (<https://www.esrl.noaa.gov/psd/>; Reynolds et al., 2007).
216 Daily mean SSTs were extracted from the cell containing each mooring's position.

217 For sea-ice concentrations, we sought datasets with the highest resolution available. We
218 required data from different passive microwave sea-ice satellites to cover the entire duration of
219 the study. For years 2009 and 2010, we obtained Advanced Microwave Scanning Radiometer –
220 Earth Observing System (AMSR-E) sea-ice concentration data with a resolution of 6.25 km from
221 the Integrated Climate Data Center (ICDC, icdc.cen.uni-hamburg.de; Kaleschke et al., 2001;
222 Spreen et al., 2008). The AMSR-E satellite failed in early October 2011, consequently for 2011
223 and 2012 we used data from the Special Scanning Microwave/Imager (SSM/I) with a spatial
224 resolution of 25 km (Cavalieri 1996). High resolution Advanced Microwave Scanning
225 Radiometer 2 (AMSR-2) data with a grid resolution of 6.25 km were used for 2013–2015
226 (Beitsch et al. 2014).

227 We derived daily mean sea-ice concentration for the area of the Chukchi Sea as defined
228 by the International Hydrographic Organization (IHO;
229 <http://www.marinegions.org/gazetteer.php?p=details&id=4257>), and for a custom study area
230 polygon (Fig. 1). The study area polygon was defined by the bounds set by Cape Serdtse-Kamen,
231 Russian Federation, in the northwest; Nunyamo, Russian Federation, to the southwest; York,
232 Alaska, USA, on the Seward Peninsula to the southeast; and Cape Espenberg, Alaska, USA, to
233 the northwest (Fig. 1). We determined the study area polygon by estimating where sea ice, if
234 present, could potentially create a migration barrier for fin whales. All satellite-derived data were

235 visualized in ArcMap (v. 10.1) using the WGS 1984 datum and projected in a custom polar
236 stereographic projection with a central meridian of -171°W.

237 *Environmental Data Analysis*

238 To ensure consistency when comparing the environmental data over time, we calculated
239 summary statistics for October data since there were no data gaps in the *in-situ* temperature and
240 salinity data in this month (except for a gap in the near-surface data for 2014 at A3). For the
241 ADCP data, we elected to compare the monthly mean northward water velocities for June to
242 November to capture the summertime peak in transport through the Bering Strait (Woodgate et
243 al., 2005b). We investigated correlations between days with fin whale calls present (i.e. FWH >
244 0) and select individual environmental variables using non-parametric Kendall's rank correlation
245 tests. The Kendall's rank coefficient, tau (τ), indicates the direction of association ($-1 < \tau < 1$)
246 and the resulting *p*-value indicates presence of a statistically significant correlation under the null
247 hypothesis of non-correlation between the samples.

248 We tested for interactions between fin whale presence and along-channel (northward)
249 wind patterns within the Bering Strait by comparing the daily mean northward wind velocity on
250 days when the number of FWH reached above a certain threshold (≥ 1 hr, ≥ 6 hrs, ≥ 12 hrs, and
251 ≥ 18 hrs) and days without any FWH. We calculated summary statistics for northward wind
252 velocities in October only, including an overall mean along-channel wind velocity as well as
253 mean wind velocity for days with no FWHs and days with FWHs above a threshold (see
254 categories above). We then compared the overall October mean along-channel wind velocity to
255 the mean wind velocities for days with and without FWHs using a Wilcoxon rank sum test.

256 For the sea ice analysis, we calculated the melt-out and freeze-up dates as the day of the
257 year when the sea ice concentration within the study area decreased/increased below/above 80%,

258 respectively, following Markus et al. (2009) and Stroeve et al. (2014). We defined an area as
259 'ice-free' if the mean sea ice concentration was $\leq 15\%$, a threshold commonly used to indicate
260 the presence of sea ice (Serreze et al., 2009; Stroeve et al., 2012; Serreze et al., 2016). We
261 calculated the melt period length using the number of days between the initiation of melting (\leq
262 80% concentration) and when the study area was ice-free ($\leq 15\%$ concentration). For the freeze-
263 up period length, we calculated the number of days between the first day sea ice concentration
264 reached $\geq 15\%$ and the first day the sea ice reached $\geq 80\%$ concentration in the fall. We compared
265 the calculated fin whale departure date and sea ice freeze-up date for each year using a two-sided
266 Pearson correlation test after testing for normality.

267

268 2.3. *Water Masses*

269

270 Water mass presence for each day was estimated for the near-surface and near-bottom
271 using temperature and salinity (T/S) bounds suggested by Danielson et al. (2017). These authors
272 distinguish five water mass categories: the Alaskan Coastal Water (ACW), Bering Chukchi
273 Summer Water (BCSW), Bering Chukchi Winter Water (BCWW), Melt Water (MW), and water
274 from the Atlantic layer in the Arctic (AtlW). Danielson et al. (2017) combine the ACW, BSSW,
275 and CSSW into one water mass, the BCSW, since the T/S properties of these three water masses
276 are often indistinguishable from each other. Similarly, Danielson et al., (2017) do not distinguish
277 between Bering Sea Winter Water and Chukchi Sea Winter Water, and instead combine the two
278 into one water mass, the BCWW. Note that since the T/S bounds of these waters vary
279 interannually (Coachman et al., 1975), there are limitations to the representativeness of the above
280 water mass identifications.

281 Chi-squared tests of independence were performed for each mooring site using pooled
282 presence/absence of fin whale calls for each day across all recording years along with the daily
283 water mass designations to determine whether there was a significant association between the
284 presence of fin whale calls and water mass. If a chi-squared test was inappropriate (e.g. in the
285 case of small sample sizes), a Fisher's exact test was applied instead. Fisher's exact test
286 evaluates the significance of association, or contingency between two categorical variables, and
287 is insensitive to sample sizes. All analyses were performed using in the statistical software R (v.
288 3.5.3; R Core Team 2019).

289

290

291 **3. Results**

292

293 *3.1. Fin Whale Detections*

294

295 We processed a total of 52,272 audio files collected from ~July to November 2009–2015
296 (Table 1). Fin whales were detected at all three sites, with the highest frequency and abundance
297 of fin whale hours (FWH) at site A3 by a large margin (Fig. 2; Supplemental Figs. S1–S3).
298 About one third (34.4%) of the total recording days at A3 had at least one hour with fin whale
299 calls, compared to only 4.6% at A2 and 1.5% at A4. Calling fin whales were detected in all
300 recording years at A2 and A3, but were only detected in 2014 and 2015 at A4. October had the
301 highest occurrence of FWH across all sites (68.4%), and given the hydrophones all had data from
302 October, we restricted our statistical tests to this month. Wilcoxon rank-sum tests revealed
303 statistically significant differences in the distribution of FWH in October at the three mooring

304 sites (A2 and A3: $W = 5259.5$, $p < 0.001$, $n = 186$ days; A2 and A4: $W = 8423$, $p = 0.006$, $n =$
305 124 days; A3 and A4: $W = 1709$, $p < 0.001$, $n = 124$ days). The earliest detection of fin whale
306 calls across all sites and years occurred on 23 July 2013 at A3, and the latest fin whale detection
307 occurred on 20 November 2015 at A3 (Table 2). Annual fin whale departure dates using the 95%
308 quantile were only calculated for A3 given the lack of data at A2 and A4. (See Supplemental Fig.
309 S4 for the cumulative distribution of days with fin whale calls at A3.) Fin whale departure dates
310 at A3 did not show any statistically significant trend ($R^2 = 0.20$, $p = 0.311$; Fig. 3).

311 At A3, fin whale calling activity was highest in 2012 and 2015 (52 and 71 days with at
312 least one FWH, respectively), while calling activity was the lowest in 2010 (22 days) followed
313 by 2011 and 2013 (28 days). The Wilcoxon tests comparing FWH in October between years at
314 A3 show significant differences in the distributions fin whale detections across years, with
315 significant values ($p < 0.01$) between all consecutive years except 2009 and 2010 ($p = 0.736$) and
316 2010 and 2011 ($p = 0.463$; Table 3). Wilcoxon tests comparing FWH in 2012 and 2015 to the
317 other years detected significantly different distributions ($p < 0.01$), except for the test between
318 2012 and 2014 ($p = 0.614$; Table 3).

319 Fin whale calls were less common at A2, though 2015 had relatively higher call activity
320 with 40 hrs with fin whale calls compared to 2–10 hrs in each of the other six years. At A4, fin
321 whale calls were only detected in 2014 (1 h) and 2015 (19 h). Insufficient sample sizes precluded
322 any statistical comparisons of fin whale vocal activity between years for A2 and A4.

323

324 3.2. *Sea-ice conditions and analyses*

325 Sea-ice conditions within the study area were highly variable from year to year. Melt-out
326 dates ranged from as early as 27 April (2011) to as late as 20 May (2010; Table 4). The number

327 of days between the initiation of melting (< 80% concentration) and ice-free conditions in the
328 study area (< 15% concentration) ranged from 21 days (2015) to 41 days (2013; Table 4). The
329 study area was typically ice-free starting in late May to early June, with the earliest ice-free date
330 occurring on 24 May 2015 and the latest on 17 June 2010. On average, freeze-up dates ($\geq 80\%$
331 concentration) occurred in early to mid-December, with the earliest freeze-up on 28 November
332 2009 and the latest on 25 December 2010. The freeze-up periods for each year were typically
333 much shorter than the melt periods, with the number of days between the ice-free date and
334 freeze-up initiation ranging from five days (2014) to 23 days (2010 and 2012; Table 4).

335 Fin whale departure dates for each year at A3 were compared to the sea ice freeze-up date
336 for the study area and the Chukchi Sea, as well as the day of the year when the daily mean near-
337 surface and near-bottom temperatures first reached $\leq 0^{\circ}\text{C}$ (Fig. 4). Two-sided Pearson correlation
338 tests indicated no significant correlation between fin whale departure date and sea ice freeze-up
339 date for the study area ($t = -1.046$, $p = 0.344$) or the Chukchi Sea ($t = -0.308$, $p = 0.771$). The
340 latest fin whale departure date occurred on 17 November 2011 and 2015 when the mean sea ice
341 concentrations were $\sim 0.8\%$ and 4.9% in the study area, and 21.0% and 18.2% in the Chukchi
342 Sea, respectively (Table 4).

343

344 *3.3. Environmental conditions at the moorings*

345

346 Environmental data at the three mooring sites exhibited strong interannual and spatial
347 variation. The highest temperatures and lowest salinities on average were seen at A4 (e.g. 2013
348 October near-surface mean temperature = 3.5°C , $\text{SD} = 0.7^{\circ}\text{C}$; near-surface mean salinity = 30.3
349 psu , $\text{SD} = 1.3 \text{ psu}$). Conversely, A2 and A3 had lower temperatures and higher salinities than A4

350 (A2: 2013 October near-surface mean temperature = 3.3°C, SD = 0.7°C, near-surface mean
351 salinity = 31.1 psu, SD = 1 psu; A3: 2013 October near-surface mean temperature = 2.9°C, SD =
352 0.8°C, near-surface mean salinity = 31.7 psu, SD = 0.8 psu; Fig. 5). This spatial structure, with
353 warm fresh waters near the Alaskan Coast, is typical for the presence of the Alaskan Coastal
354 Current (see discussion in Woodgate et al., 2015). There were also significant interannual
355 differences across all three sites. The lowest near-surface and near-bottom temperatures occurred
356 in 2012 while the highest temperatures occurred in 2015 (Fig. 5; Woodgate, 2018).

357 Northward water velocities were on average the highest at sites A2 and A4 during the
358 open-water season (Fig. 6), consistent with known seasonality in the flow due to weaker
359 opposing southward winds in summer (Woodgate et al., 2005b). The year 2012 had the weakest
360 northward water velocity throughout the open-water season while 2014 had sustained high
361 northward velocities throughout the season (Fig. 6; Woodgate, 2018). Overall, northward water
362 velocities weakened over the period between July and November with the slowest northward
363 water velocities occurring in November, except in 2012 and 2014 when the seasonal minimum
364 velocities were seen in September and October (Fig. 6). Direction of flow at all three sites was
365 primarily northward during the open-water season (see Supplemental Figs. S5–S11 for plots of
366 the water and wind velocity vectors along with fin whale acoustic presence at A3 during the
367 open-water season). For a more detailed overview of variation in Bering Strait transport through
368 2015, see Woodgate (2018).

369 Due to low fin whale detections at A2 and A4, we focused our wind analysis on site A3
370 and used wind data from the grid point closest to the mooring (67.5°N, 190°W, ~140 km to the
371 northwest of A3). On average, along-channel winds were mainly southward during the month of
372 October, with the strongest mean winds occurring in 2013 (October \bar{x} = -6.2 m/s, SD = 5.4 m/s)

373 and the weakest mean winds in 2012 (October $\bar{x} = -0.4$ m/s, SD = 8.1 m/s; Table 5). Note that the
374 negative sign indicates a southward direction.

375

376 3.4. *Environmental Analyses*

377

378 We focused our environmental analyses on the A3 mooring site due to the relative lack of
379 fin whale detections at A2 and A4. The Kendall's rank correlation tests between FWH on days
380 with fin whale calls (i.e. FWH > 0) and the environmental variables produced statistically
381 significant ($p < 0.05$) though small correlations for daily mean water speed, and along-channel
382 wind and water velocities pooled for all seven years (2009–2015; Table 6). We ran a second test
383 using October data only and found similar results, as well as the addition of significant
384 correlations between FWH and near-surface temperature and SST at site A3 (Table 6).

385 Days with fin whale calls mostly had southward mean wind velocities while days without
386 calls (i.e. FWH = 0) mostly had northward overall mean winds (Table 5; Fig. 7). The Wilcoxon
387 test comparing the overall mean along-channel wind velocity for October of each year against
388 the means for days with and without FWHs revealed that days without FWH and days with FWH
389 ≥ 6 and 12 hrs had statistically significant differences in along-channel wind velocities in 2011
390 and 2014 only (Table 5). Insufficient data precluded any tests for days with FWH ≥ 18 hrs.

391

392 3.5. *Water mass composition at the moorings*

393 Water mass composition at A2 and A3 during the open water season was dominated by
394 the presence of the Bering Chukchi Summer Water (BCSW) at both the near-surface (>70% of
395 days at both sites) and near-bottom levels (>90% of days at both sites) for all recording years

396 (see Supplemental Figs. S12-S14 for plots with the water mass composition at the three sites
397 during the open water season). The water mass composition at A4 was similarly dominated by
398 BCSW at the near-bottom (73% of days in July–November) and to a lesser extent in the near-
399 surface (51% of days in July–November). The cold and salty Bering Chukchi Winter Water
400 (BCWW) appeared in both levels in the water column in November at all three sites, when it is
401 assumed that fin whales are beginning their migration south. A fresher, colder signal, that falls
402 within the Melt Water (MW) category as defined by Danielson et al., (2017), appeared in the
403 near-surface at all three sites in September and October 2012 and 2013, with the strongest signal
404 in 2012. However, since the sea-ice edge is far away from the mooring sites in September and
405 October, the freshening observed in 2012 and 2013 was likely due to fresh waters from either the
406 Alaskan Coastal Current or the Siberian Coastal Current (SCC). The SCC is a cold, fresh current
407 present seasonally in the Chukchi Sea only in some years (Weingartner et al., 1999). Also
408 noteworthy was a warm ACW signal in the near-surface at A2 in 2013, 2014, and 2015 and at
409 A3 in 2010 and 2015.

410 We conducted a side-by-side comparison of the daily water mass designations for A2 and
411 A3 and noted the number of days when at least one of the water mass designations at A2 did not
412 match those from A3. Out of 726 days when both moorings were recording and had data for both
413 instruments, 14 days (~2% of total days) had different water mass composition in the near-
414 bottom water and 69 days (~10% of total days) for the near-surface water. In contrast, A2 and A4
415 had different water mass compositions on 203 days (~39%) for the near-surface and 127 days
416 (24%) for the near-bottom. The comparison between A3 and A4 yielded 311 days (60%) with
417 different water mass composition at the near-surface and 136 days (26%) at the near-bottom.

418 These results indicate that despite close spatial proximity, A2 and A4 had very different water
419 mass composition while A2 and A3 had similar water mass composition.

420

421 3.6. *Water Mass Analyses*

422

423 The chi-squared tests of independence between the pooled FWH and the near-
424 surface/near-bottom water mass designations at site A3 suggest that the occurrence of fin whale
425 calls during the study period was statistically dependent on the occurrence of water masses (both
426 tests using near-surface and near-bottom water mass designations: $p < 0.001$). We repeated the
427 tests of independence for each recording year at A3, using the Fisher's Exact Test to compare the
428 daily near-surface and near-bottom water mass designations to the total FWH for each day. The
429 results show a significant relationship for 2009, 2011, 2012, and 2015 (all $p < 0.02$), signifying
430 that fin whale presence was statistically dependent on water mass presence for these years. We
431 were unable to execute the Fisher's Exact test for 2013 (near-bottom water mass) and 2014 (both
432 near-surface and near-bottom) due to the fact that only one water mass (BCSW) was present at
433 both levels in the water column, resulting in zeros in both the expected and observed columns of
434 the test's contingency tables.

435 We were unable to perform a chi-squared test for independence for A2 and A4 due to the
436 presence of small expected values ($E_{i,j} < 5$) in the contingency tables generated by the test. At
437 A4, fin whale calls were only heard on days when the BCSW was present at both levels of the
438 water column. Calling fin whales were only heard at A2 on days when the BCSW was present in
439 the near-bottom waters. We applied a Fisher's Exact Test to the A2 near-surface water mass

440 designations and found that fin whale calls and water mass occurrence in the near-surface waters
441 were statistically independent of each other ($p = 0.48$).

442

443

444 **4. Discussion**

445

446 The results of this study show a pattern of interannual and spatial variation in the
447 presence of acoustically-active fin whales in the Bering Strait region. Across all three sites, the
448 year 2015 had the most fin whale detections followed by 2012, though these years had
449 contrasting temperatures and salinities, sea-ice conditions, water velocity and wind patterns. Site
450 A3, where the Anadyr and Bering Shelf waters were most prevalent, had the most hours with fin
451 whale calls, supporting our hypothesis that water masses may affect the occurrence of fin whales.
452 We found small but significant correlations between FWH and northward wind and water
453 velocities, near-surface temperatures and SST at site A3. However, our p -values for the
454 correlation tests were potentially too low and likely overestimated the real significance of the
455 tests given that the days with fin whale calls were likely not independent of each other. In
456 addition, the statistically significant correlations between FWH and environmental variables
457 were small (< 0.25). Thus, we conclude that it is not possible to prove a strong relationship
458 between individual environmental parameters and FWH with our data. More data and greater
459 spatial coverage are necessary to prove any significant association between days with fin whale
460 calls and environmental factors in the Bering Strait region.

461 Most fin whale calls were heard in October, potentially due to fact that fin whale 20-Hz
462 pulses serve primarily a reproductive purpose (Watkins et al., 2000; Croll et al., 2002; Stafford et

463 al., 2007) and tend to be heard closer to the winter mating season (Stafford et al., 2007).
464 Consequently, fin whale vocalizations may not be a reliable indication of when fin whales first
465 pass northwards through the Bering Strait. Additionally, the dates of departure from the Bering
466 Strait region presented here only apply to vocal fin whales since we could not detect non-vocal
467 whales, which could have remained in the area beyond these dates. Due to this inherent bias, the
468 departure dates presented in this study only provide an approximation for when fin whales leave
469 the region. The departure dates from the A3 mooring site did not exhibit a significant trend (Fig.
470 3), therefore it is not possible to determine whether fin whales are extending their residence time
471 in the Chukchi Sea from our data. Perhaps this is not surprising given that we only have seven
472 years of data, and interannual variability is substantial. In general, the fin whale departure dates
473 at A3 occurred in early November, ranging from 31 October (2010) to 17 November (2011 and
474 2015). What signaled the fin whales to leave the Chukchi Sea is not clear. Sea-ice concentrations
475 in the study area around the last detection dates were well below 'ice-free' levels ($< 15\%$; Table
476 4), indicating that the Bering Strait was still navigable and free of sea ice. It is possible, though,
477 that fin whales respond to cooling water temperatures since all departure dates occurred before
478 near-surface and near-bottom water temperatures at A3 reached below 0°C (Fig. 4).

479 The overwhelming majority of fin whale calls were detected at site A3, where calling fin
480 whales were heard every year. There are multiple possible explanations for the spatial variability
481 observed in fin whale detections. First, site A3 is situated at the confluence of two productive
482 water masses, the Anadyr Water (AW) and Bering Shelf Water (BSW), which likely provide
483 better feeding opportunities for fin whales. The dominant water mass detected at A3 was the
484 Bering Chukchi Summer Water (BCSW), which is composed of the AW and BSW, and thus has
485 high nutrient levels and larger zooplankton (Eisner et al., 2013; Ershova et al., 2015; Danielson

486 et al., 2017). Though fin whale calls were also detected on days when other fresher water masses
487 were present at A3, including days in 2015 when Alaskan Coastal Water (ACW) was present in
488 the near-surface (Fig. S13). Fin whale calls were also detected on days in September 2012 when
489 a fresh, cold signal appeared in the near-surface waters at A3, possibly indicating the presence of
490 the Siberian Coastal Current (SCC).

491 The SCC occasionally flows into the Bering Strait during periods with strong or
492 persistent southward winds (Weingartner et al., 1999). Ershova et al., (2015) detected the
493 presence of the SCC in the central Chukchi Sea in September 2012, therefore it is possible that
494 the reach of the SCC extended to the A3 site that month. Fig. S8 shows that winds measured in
495 September 2012 were predominantly southward, which has been shown to cause the ACW to
496 deviate away from the Alaskan coast and towards the western Chukchi Sea (Woodgate et al.,
497 2015; Pisareva, 2018; Morris, 2019). Often the presence of the cold and fresh SCC creates a
498 front (Weingartner et al., 1999), which could isolate and cluster prey. In 1992-1993, Moore et al.,
499 (1995) observed bowhead whales (*Balaena mysticetus*) feeding in close association with salinity
500 and thermal fronts along the Chukotka coast. Moreover, *Thysanoessa inermis*, a common fin
501 whale prey (Nemoto 1959; Witteveen and Wynne, 2016), was found to be the dominant
502 zooplankton species collected from a dense prey patch near a front, lending support to the
503 potential importance of the SCC in creating favorable feeding conditions for fin whales at A3.

504 In addition to its proximity to productive water masses, A3 may be situated close to
505 oceanographic features created by currents, such as island wake eddies, that are known to create
506 favorable foraging opportunities for baleen whales (Johnston et al., 2005a; Chenoweth et al.,
507 2011). Eddies create upwelling zones which promote phytoplankton blooms (Hasegawa et al.,
508 2009) and have been shown to be important feeding habitat for auklets and other planktivores in

509 the Bering and Chukchi seas (Piatt and Springer, 2003). In the Bay of Fundy, Canada, island
510 wake eddy systems were found to be important feeding grounds for fin whales as well as minke
511 whales (*B. acutorostrata*) and harbor porpoises (*Phocoena phocoena*; Johnston et al., 2005a,b).
512 Currents moving past the Diomede Islands generate island wake eddies (Coachman et al., 1975;
513 Woodgate et al., 2015) that are then carried northwards towards A3, according to satellite SST
514 data (Woodgate, pers. comm.). The island wake eddies may create opportune feeding conditions
515 for fin whales at A3.

516 In contrast, site A2 had fin whale detections in all recording years but in lower
517 abundance, while fin whale calls were largely absent from site A4. Given its position in the less-
518 productive Alaskan Coastal Current, A4 may present lower quality feeding areas for fin whales
519 than the other two sites. Though A2 had similar water mass composition as A3, water velocities
520 were higher at A2, potentially transporting prey out of the area. Therefore, fin whales may be
521 less inclined to stay at in the region around A2 due to fewer feeding opportunities. Also, the
522 position of site A3 north and towards the middle of the Bering Strait gives it an advantage over
523 A2 in capturing the calls of fin whales migrating through the western strait. Whereas A2 and A4
524 can only record the calls of fin whales passing through the east channel of the strait, A3 can
525 potentially record calling whales migrating through both channels.

526 While the spatial variability in fin whale detections may be explained, the exact
527 environmental mechanisms for the observed temporal variability are less clear. Both 2012 and
528 2015 stand out as years with the highest number of fin whale detections at A3, yet the two years
529 had very different environmental conditions. The year 2012 had the coldest October mean
530 temperatures (near-bottom October mean at A3 = 1.0°C), late sea ice breakup (16 May),
531 anomalously low flow (Woodgate, 2018), and weak mean northward wind velocities in the fall.

532 On the other hand, 2015 had a very warm annual mean temperature (near-bottom October mean
533 at A3 = 3.6°C), earlier sea ice breakup (4 May), variable northward wind velocities, and high
534 flow (Woodgate, 2018). Our results suggest that at A3, the occurrence of fin whale calls is more
535 strongly related to southward winds than northward winds, but this relation does not hold for all
536 years (Table 5). Thus, we cannot attribute interannual variation in the acoustic presence of fin
537 whales to any one environmental predictor. Instead, we believe that a combination of conditions
538 not only in the Chukchi Sea, but also in the Bering Sea, contributes to the abundance of fin
539 whales in the study area. We hypothesize a series of ‘push’ and ‘pull’ factors below that may
540 have influenced the observed interannual variation in the presence of acoustically-active fin
541 whales.

542 Pull factors imply that conditions in the Chukchi Sea were favorable for zooplankton and
543 other fin whale prey in 2012 and 2015, thus drawing more fin whales into the area to feed. The
544 abundance of hours with fin whale calls at A3 in 2012 may point to the fact that the year was
545 particularly cold, and thus, productive. Colder temperatures are more favorable for the secondary
546 production of *Calanus* copepods (Kimmel et al., 2017), a prominent constituent of the Chukchi
547 Sea zooplankton. Cold years in the Bering and Chukchi seas have been also found to have higher
548 zooplankton biomass and abundance (Ohashi et al., 2013; Ershova et al., 2015; Pinchuk and
549 Eisner, 2017), and thus stronger recruitment for walleye pollock (*Gadus chalcogrammus*) and
550 Pacific cod (*G. macrocephalus*; Stabeno et al., 2012), as well as zooplankton predators like fin
551 whales. Friday et al., (2013) observed twice as many fin whales along the eastern Bering Sea
552 shelf in 2008 and 2010 when temperatures were cold than they did in 2002, a warm year. In their
553 August–September 2012 sampling of the Chukchi Sea, Danielson et al., (2017) observed an
554 abnormally high biomass of large copepods as well as a predominance of the BCSW in the

555 bottom water at multiple sampling stations. During the same sampling period, Pinchuk and
556 Eisner (2017) report a high abundance of *Calanus glacialis* and widespread distribution of
557 Pacific-origin zooplankton in 2012, adding evidence to our theory that 2012 was a favorable year
558 for fin whale prey.

559 Conversely, 2015 was a warm year with high salinities. High salinities are usually
560 indicative of high Anadyr Water content and thus are typically associated with high nutrient
561 levels (Danielson et al., 2017). Consequently, 2015 may have had higher zooplankton abundance
562 due to a nutrient-rich environment. Pinchuk and Eisner (2017) found a strong correlation
563 between the biomass of Pacific-origin zooplankton and high salinities associated with the
564 BCSW, which was the dominant water mass at A3 in 2015 (Supplementary Figs. S12-S14). It is
565 also possible that the earlier sea-ice retreat and warmer water temperatures observed in 2015
566 created better conditions for Pacific-origin copepods and euphausiids. Matsuno et al. (2011)
567 found that Pacific copepod species (e.g. *Eucalanus bungii*) expanded into the Chukchi Sea in
568 2007, a year with relatively early sea-ice retreat and abnormally high sea surface temperatures,
569 similar to 2015. A notable pull factor for 2015 could also have been the strong water velocities
570 measured in the Bering Strait. Strong velocities likely led to higher transport of both nutrients
571 and zooplankton from the Bering Sea into the Chukchi Sea, creating better feeding opportunities
572 for summer migrant fin whales.

573 In contrast to pull factors, potential push factors consist of poorer conditions in other
574 reaches of the fin whale range, thereby sending fin whales into the Chukchi Sea in search of
575 better conditions. Such areas include the Bering Sea and Gulf of Alaska, where fin whales are
576 known to occur in the summer months (Moore et al., 1998, 2000; Stafford et al., 2007). Both
577 2014 and 2015 were significantly warmer in comparison to historical records for the Bering Sea

578 (Duffy-Anderson et al., 2017). Warm years in the Bering Sea result in poor recruitment in
579 walleye pollock due to the prevalence of small, lipid-poor copepods (Kimmel et al., 2017). In
580 2015, an anomalously warm water mass, nicknamed the “Blob,” pervaded the North Pacific,
581 leading to declines in krill and to northward distribution shifts of multiple marine species
582 (Cavole et al., 2016). Concurrent with the appearance of the Blob were reports of a mass
583 mortality event of common murrelets (*Uria aalga*) in the Gulf of Alaska (Piatt et al., 2018).
584 Additionally, 12 fin whales stranded on Kodiak Island between May and June 2015 (Savage et
585 al., 2017). Though the causes of death for the whales were not determined, ecological conditions
586 rather than anthropogenic factors (e.g. ship strikes) are thought to be the culprit (Savage et al.,
587 2017). Warmer temperatures observed in 2015 may have affected prey availability in other fin
588 whale summer feeding grounds, pushing fin whales into the Chukchi Sea in search of better
589 feeding opportunities.

590 Another possible explanation for the increased observation of fin whale calls in 2015 is
591 that the North Pacific population of fin whales is increasing (Zerbini et al., 2006), and thus may
592 be reclaiming portions of its previous range (Clarke et al., 2013; Brower et al., 2018). An
593 increased number of fin whales observed during annual surveys conducted by the Aerial Surveys
594 of Arctic Marine Mammals Project (ASAMM) from 2008–2016 in comparison to 1982–1991
595 support this theory (Brower et al., 2018). Brower et al. (2018) report seeing the most fin whales
596 in the south-central Chukchi Sea in 2014 (44% of observations) and in 2015 (27%). However, it
597 is difficult to evaluate habitat reclamation of fin whales using their calls alone given that only
598 males are thought to produce the 20-Hz pulse and we could only detect vocal fin whales.

599 Limitations of the present study include limited spatial coverage of the study area with
600 hydrophones located in only the east channel and north of the Bering Strait. Since there are no

601 recent surveys on the western side of the Bering Strait or Chukchi Sea, our knowledge of fin
602 whale habitat use in this region is limited. Given that the productive AW is typically found
603 mainly in the west channel of the Bering Strait, it is possible that most fin whales may traverse
604 through the strait on the western side. However, without adequate observation platforms
605 covering both sides of the strait, the exact migration path of fin whales in the region remains
606 unknown.

607 The results of this study corroborate patterns of interannual variation in fin whale
608 presence observed by previous studies. Like the present study, Delarue et al., (2013) noted low
609 fin whale detections in the northeast Chukchi Sea in 2009 and 2010, attributing diminished vocal
610 activity to poorer feeding conditions. In contrast, more fin whales were heard in 2007, a
611 particularly warm year in the Chukchi Sea with early ice retreat and low sea-ice extent, as well as
612 high transport through the Bering Strait (Delarue et al., 2013; Woodgate et al., 2010). The
613 conditions in 2007 described by Delarue et al., (2013) are very similar to those we observed in
614 2015, when fin whale calls were the most abundant.

615 Our results present a preliminary examination of how environmental variations in the
616 southern Chukchi Sea and Bering Strait may lead to interannual variability in the acoustic
617 presence of fin whales. Though we were unable to identify a single environmental driver that
618 explained the variation, differences in temperature, salinity, wind and water velocities likely
619 played a role. There are potentially numerous combinations of environmental variables that
620 create preferential feeding opportunities for fin whales. Delarue et al., (2013) hypothesize that
621 perhaps the combination of environmental variables observed in 2007 (warm SSTs, low sea-ice
622 concentrations, and high transport) created favorable conditions for the fin whales' zooplankton
623 prey. However, the abundance of fin whales heard in 2012, a period with colder water

624 temperatures, low transport, and high spring sea-ice concentrations, suggests that alternative
625 environmental drivers are also favorable for fin whale feeding.

626 Conditions in the Bering Sea may also be an important factor in determining fin whale
627 occurrence in the Chukchi Sea. Comparing fin whale detections in the southern Chukchi Sea
628 with those in the Bering Sea could help indicate whether fin whale presence in one region results
629 in higher fin whale presence in the other. Examining the environmental conditions in the Bering
630 Sea for 2009–2015 could shed light on the patterns of fin whale occupation found in the present
631 study. Continued monitoring of fin whale presence in the southern Chukchi and Bering seas in
632 relation to oceanographic features is necessary for composing a more complete picture of how
633 fin whale presence in the Pacific Arctic is changing in response to environmental shifts over
634 time.

635 **Acknowledgements**

636

637 This work was supported by the National Science Foundation (NSF) 495 Graduate
638 Research Fellowship (grant #DGE-1256082), and by the NSF Polar Programs Arctic Observing
639 Network (R. Woodgate grant numbers PLR-1304052 and 1758565; K. Stafford grant number
640 1107106). A portion of the mooring work was supported by the Russian-American Long-term
641 Census of the Arctic (RUSALCA), funded by the U.S. National Oceanic and Atmospheric
642 Administration (NOAA) (grant numbers NA10OAR4320148, AM105, and AM133), the Applied
643 Physics Laboratory at the University of Washington, and the National Science Foundation (grant

644 number ARC-1107106). The Bering Strait mooring data are available at the permanent archives
645 of the U.S. National Centers for Environmental Information/National Oceanographic Data
646 Center, www.nodc.noaa.gov, and at: psc.apl.washington.edu/BeringStrait.html. This manuscript
647 may be referenced through the North Pacific Research Board Arctic Integrated Ecosystem
648 Research Program archives as ArcticIERP-13. We thank the four reviewers for their suggestions
649 which improved the manuscript.

650 **References**

651

652 Baker, M., Kivva, K., Pisareva, M., Watson, J., Selivanova, J., this issue. Shifts in the physical
653 environment in the Pacific Arctic and implications for ecological timing and conditions.
654 *Deep Sea Res. Part II Top. Stud. Oceanogr.*

655 Beitsch, A., Kaleschke, L., Kern, S., 2014. Investigating high-resolution AMSR2 sea ice
656 concentrations during the February 2013 fracture event in the Beaufort Sea. *Remote Sens.*
657 *6*, 3841–3856. <https://doi.org/10.3390/rs6053841>

658 Berline, L., Spitz, Y.H., Ashjian, C.J., Campbell, R.G., Maslowski, W., Moore, S.E., 2008.
659 Euphausiid transport in the Western Arctic Ocean. *Mar. Ecol. Prog. Ser.* 360, 163–178.

660 Brower, A.A., Clarke, J.T., Ferguson, M.C., 2018. Increased sightings of subArctic cetaceans in
661 the eastern Chukchi Sea, 2008–2016: population recovery, response to climate change, or
662 increased survey effort? *Polar Biol.* 41, 1033–1039. [https://doi.org/10.1007/s00300-018-](https://doi.org/10.1007/s00300-018-2257-x)
663 [2257-x](https://doi.org/10.1007/s00300-018-2257-x)

664 Cavalieri, D., 1996. Sea ice concentrations from Nimbus-7 SMMR and DMSP SSM/I-SSMIS
665 passive microwave data, Version 1. <https://doi.org/10.5067/8GQ8LZQVL0VL>. [Date
666 Accessed: 25 April 2017].

667 Cavalieri, D.J., Parkinson, C.L., 2012. Arctic sea ice variability and trends, 1979-2010.
668 Cryosphere 6, 881–889. <https://doi.org/10.5194/tc-6-881-2012>

669 Cavole, L., Diner, R., Giddings, A., Koester, I., Pagniello, C., Paulsen, M.-L., Ramirez-Valdez,
670 A., Schwenck, S., Yen, N., Zill, M., Franks, P., 2016. Biological impacts of the 2013–2015
671 warm-water anomaly in the Northeast Pacific: Winners, Losers, and the Future.
672 Oceanography 29. <https://doi.org/10.5670/oceanog.2016.32>

673 Chenoweth, E., Gabriele, C., Hill, D., 2011. Tidal influences on humpback whale habitat
674 selection near headlands. Mar. Ecol. Prog. Ser. 423, 279–289.
675 <https://doi.org/10.3354/meps08891>

676 Clarke, J., Stafford, K., Moore, S., Rone, B., Aerts, L., Crance, J., 2013. Subarctic cetaceans in
677 the southern Chukchi Sea: Evidence of recovery or response to a changing ecosystem.
678 Oceanography 26, 136–149. <https://doi.org/10.5670/oceanog.2013.81>

679 Clarke, J.T., Ferguson, M.C., Curtice, C., Harrison, J., 2015. 8. Biologically important areas for
680 cetaceans within U.S. waters – Arctic Region. Aquat. Mamm. 41, 94–103.
681 <https://doi.org/10.1578/AM.41.1.2015.94>

682 Coachman, L.K., Aagaard, K., Tripp, R.B., 1975. Bering Strait: the regional physical
683 oceanography. University of Washington Press, Seattle.

684 Comiso, J.C., Parkinson, C.L., Gersten, R., Stock, L., 2008. Accelerated decline in the Arctic sea
685 ice cover. Geophys. Res. Lett. 35. <https://doi.org/10.1029/2007GL031972>

686 Croll, D.A., Clark, C.W., Acevedo, A., Tershy, B., Flores, S., Gedamke, J., Urban, J., 2002. Only
687 male fin whales sing loud songs. *Nature* 417, 809.

688 Danielson, S.L., Eisner, L., Ladd, C., Mordy, C., Sousa, L., Weingartner, T.J., 2017. A
689 comparison between late summer 2012 and 2013 water masses, macronutrients, and
690 phytoplankton standing crops in the northern Bering and Chukchi Seas. *Deep Sea Res. Part*
691 *II Top. Stud. Oceanogr.* 135, 7–26. <https://doi.org/10.1016/j.dsr2.2016.05.024>

692 Delarue, J., Martin, B., Hannay, D., Berchok, C.L., 2013. Acoustic occurrence and affiliation of
693 fin whales detected in the northeastern Chukchi Sea, July to October 2007–10. *Arctic* 66,
694 159–172.

695 Duffy-Anderson, J.T., Stabeno, P.J., Siddon, E.C., Andrews, A.G., Cooper, D.W., Eisner, L.B.,
696 Farley, E.V., Harpold, C.E., Heintz, R.A., Kimmel, D.G., Sewall, F.F., Spear, A.H.,
697 Yasumishii, E.C., 2017. Return of warm conditions in the southeastern Bering Sea:
698 Phytoplankton - Fish. *Plos One* x12, e0178955.
699 <https://doi.org/10.1371/journal.pone.0178955>

700 Edds-Walton, P.L., 1997. Acoustic communication signals of mysticete whales. *Bioacoustics* 8,
701 47–60. <https://doi.org/10.1080/09524622.1997.9753353>

702 Eisner, L., Hillgruber, N., Martinson, E., Maselko, J., 2013. Pelagic fish and zooplankton species
703 assemblages in relation to water mass characteristics in the northern Bering and southeast
704 Chukchi seas. *Polar Biol.* 36, 87–113. <https://doi.org/10.1007/s00300-012-1241-0>

705 Ershova, E., Hopcroft, R., Kosobokova, K., Matsuno, K., Nelson, R.J., Yamaguchi, A., Eisner,
706 L., 2015. Long-Term Changes in Summer Zooplankton Communities of the Western
707 Chukchi Sea, 1945–2012. *Oceanography* 28, 100–115.
708 <https://doi.org/10.5670/oceanog.2015.60>

709 Flinn, R.D., Trites, A.W., Gregr, E.J., Perry, R.I., 2002. Diets of fin, sei, and sperm whales in
710 British Columbia: an analysis of commercial whaling records, 1963-1967. *Mar. Mamm.*
711 *Sci.* 18, 663–679. <https://doi.org/10.1111/j.1748-7692.2002.tb01065.x>

712 Frey, K.E., Moore, G.W.K., Cooper, L.W., Grebmeier, J.M., 2015. Divergent patterns of recent
713 sea ice cover across the Bering, Chukchi, and Beaufort seas of the Pacific Arctic Region.
714 *Prog. Oceanogr.* 136, 32–49. <https://doi.org/10.1016/j.pocean.2015.05.009>

715 Friday, N.A., Zerbini, A.N., Waite, J.M., Moore, S.E., Clapham, P.J., 2013. Cetacean
716 distribution and abundance in relation to oceanographic domains on the eastern Bering Sea
717 shelf, June and July of 2002 2008, and 2010. *Deep Sea Res. Part II Top. Stud. Oceanogr.*
718 94, 244–256. <https://doi.org/10.1016/j.dsr2.2013.03.011>

719 Grebmeier, J.M., Moore, S.E., Overland, J.E., Frey, K.E., Gradinger, R., 2010. Biological
720 response to recent Pacific Arctic sea ice retreats, *Eos, Trans.*, pp. 161–162.
721 <https://doi.org/10.1029/2010EO180001>

722 Hasegawa, D., Lewis, M.R., Gangopadhyay, A., 2009. How islands cause phytoplankton to
723 bloom in their wakes. *Geophys. Res. Lett.* 36. <https://doi.org/10.1029/2009GL039743>

724 Hauser, D.D.W., Laidre, K.L., Stafford, K.M., Stern, H.L., Suydam, R.S., Richard, P.R., 2017.
725 Decadal shifts in autumn migration timing by Pacific Arctic beluga whales are related to
726 delayed annual sea ice formation. *Glob. Change Biol.* 23, 2206–2217.
727 <https://doi.org/10.1111/gcb.13564>

728 Hopcroft, R.R., Kosobokova, K.N., Pinchuk, A.I., 2010. Zooplankton community patterns in the
729 Chukchi Sea during summer 2004. *Deep Sea Res. Part II Top. Stud. Oceanogr.* 57, 27–39.
730 <https://doi.org/10.1016/j.dsr2.2009.08.003>

731 Jakobsson, M., L. A. Mayer, B. Coakley, J. A. Dowdeswell, S. Forbes, B. Fridman, H.
732 Hodnesdal, R. Noormets, R. Pedersen, M. Rebesco, H.-W. Schenke, Y. Zarayskaya A, D.
733 Accettella, A. Armstrong, R. M. Anderson, P. Bienhoff, A. Camerlenghi, I. Church, M.
734 Edwards, J. V. Gardner, J. K. Hall, B. Hell, O. B. Hestvik, Y. Kristoffersen, C. Marcussen,
735 R. Mohammad, D. Mosher, S. V. Nghiem, M. T. Pedrosa, P. G. Travaglini, Weatherall, P.,
736 The International Bathymetric Chart of the Arctic Ocean (IBCAO) Version 3.0, *Geophys.*
737 *Res. Lett.*, doi: 10.1029/2012GL052219.

738 Johnston, D., Thorne, L., Read, A., 2005a. Fin whales *Balaenoptera physalus* and minke whales
739 *Balaenoptera acutorostrata* exploit a tidally driven island wake ecosystem in the Bay of
740 Fundy. *Mar. Ecol. Prog. Ser.* 305, 287–295. <https://doi.org/10.3354/meps305287>

741 Johnston, D., Westgate, A., Read, A., 2005b. Effects of fine-scale oceanographic features on the
742 distribution and movements of harbour porpoises *Phocoena phocoena* in the Bay of Fundy.
743 *Mar. Ecol. Prog. Ser.* 295, 279–293. <https://doi.org/10.3354/meps295279>

744 Kaleschke, L., Lüpkes, C., Vihma, T., Haarpaintner, J., Bochert, A., Hartmann, J., Heygster, G.,
745 2001. SSM/I Sea Ice Remote Sensing for Mesoscale Ocean-Atmosphere Interaction
746 Analysis. *Canadian Journal of Remote Sensing* 27, 526–537.
747 <https://doi.org/10.1080/07038992.2001.10854892>

748 Kaleschke, L., Tian-Kunze, X., 2016. "AMSR2 ASI 3.125 km Sea Ice Concentration Data,
749 V0.1". Institute of Oceanography, University of Hamburg, Germany, digital media (ftp-
750 projects.zmaw.de/seaice/) [1/1/2013 - 31/12/2015].

751 Kimmel, D.G., Eisner, L.B., Wilson, M.T., Duffy-Anderson, J.T., 2018. Copepod dynamics
752 across warm and cold periods in the eastern Bering Sea: Implications for walleye pollock

753 (*Gadus chalcogrammus*) and the Oscillating Control Hypothesis. *Fish. Oceanogr.* 27, 143–
754 158. <https://doi.org/10.1111/fog.12241>

755 Kitamura, M., Amakasu, K., Kikuchi, T., Nishino, S., 2017. Seasonal dynamics of zooplankton
756 in the southern Chukchi Sea revealed from acoustic backscattering strength. *Cont. Shelf*
757 *Res.* 133, 47–58. <https://doi.org/10.1016/j.csr.2016.12.009>

758 Laidre, K.L., Heide-Jørgensen, M.P., 2012. Spring partitioning of Disko Bay, West Greenland,
759 by Arctic and Subarctic baleen whales. *ICES J. Mar. Sci.* 69, 1226–1233.
760 <https://doi.org/10.1093/icesjms/fss095>

761 Laidre, K.L., Stern, H., Kovacs, K.M., Lowry, L., Moore, S.E., Regehr, E.V., Ferguson, S.H.,
762 Wiig, Ø., Boveng, P., Angliss, R.P., Born, E.W., Litovka, D., Quakenbush, L., Lydersen,
763 C., Vongraven, D., Ugarte, F., 2015. Arctic marine mammal population status, sea ice
764 habitat loss, and conservation recommendations for the 21st century: Arctic Marine
765 Mammal Conservation. *Conserv. Biol.* 29, 724–737. <https://doi.org/10.1111/cobi.12474>

766 Matsuno, K., Yamaguchi, A., Hirawake, T., Imai, I., 2011. Year-to-year changes of the
767 mesozooplankton community in the Chukchi Sea during summers of 1991, 1992 and 2007,
768 2008. *Polar Biol.* 34, 1349–1360. <https://doi.org/10.1007/s00300-011-0988-z>

769 Markus, T., Stroeve, J.C., Miller, J., 2009. Recent changes in Arctic sea ice melt onset, freezeup,
770 and melt season length. *J. Geophys. Res.* 114. <https://doi.org/10.1029/2009JC005436>

771 McDonald, M.A., Hildebrand, J.A., Webb, S.C., 1995. Blue and fin whales observed on a
772 seafloor array in the Northeast Pacific. *J. Acoust. Soc. Am.* 98, 712–721.
773 <https://doi.org/10.1121/1.413565>

774 Mellinger, D.K., Clark, C.W., 2000. Recognizing transient low-frequency whale sounds by
775 spectrogram correlation. *J. Acoust. Soc. Am.* 107, 3518–29.

776 Mellinger, D., 2002. Ishmael 1.0 User's Guide. NOAA Technical Memorandum OAR PMEL-
777 120. <http://www.pmel.noaa.gov/pubs/PDF/mell2434/mell2434.pdf>.

778 Mizroch, S.A., Rice, D.W., Breiwick, J.M., 1984. The fin whale, *Balaenoptera physalus*. Mar.
779 Fish. Rev. 46, 20–24.

780 Mizroch, S.A., Rice, D.W., Zwiefelhofer, D., Waite, J., Perryman, W.L., 2009. Distribution and
781 movements of fin whales in the North Pacific Ocean. Mammal Rev. 39, 193–227.
782 <https://doi.org/10.1111/j.1365-2907.2009.00147.x>

783 Moore, S.E., George, J.C., Coyle, K.O., Weingartner, T.J., 1995. Bowhead whales along the
784 Chukotka Coast in autumn. Arctic 48, 155–160.

785 Moore, S.E., Stafford, K.M., Dahlheim, M.E., Fox, C.G., Braham, H.W., Polovina, J.J., Bain,
786 D.E., 1998. Seasonal variation in reception of fin whale calls at five geographic areas in the
787 North Pacific. Mar. Mamm. Sci. 14, 617–627.

788 Moore, S.E., Waite, J., Mazzuca, L., Hobbs, R., 2000. Mysticete whale abundance and
789 observations of prey associations on the central Bering Sea shelf. J. Cetac. Res. Manage. 2,
790 227–234.

791 Morris, B.A., 2019. Seasonality and forcing factors of the Alaskan Coastal Current in the Bering
792 Strait from July 2011 to July 2012 (Thesis). University of Washington, Seattle, WA.

793 Nasu, K., 1960. Oceanographic investigation in the Chukchi Sea during the summer of 1958.
794 Scientific Reports of the Whales Research Institute 15, 143–157.

795 Nemoto, T., 1959. Food of baleen whales with reference to whale movements. Scientific Reports
796 of the Whales Research Institute 14, 149–290.

797 Nemoto, T., Kasuya, T., 1965. Foods of baleen whales in the Gulf of Alaska of the North Pacific.
798 Scientific Reports of the Whales Research Institute 19, 45–51.

799 Nikulin, P.G., 1946. Distribution of cetaceans in the seas surrounding the Chukchi Peninsula.
800 Izv. TINRO 22.

801 Ohashi, R., Yamaguchi, A., Matsuno, K., Saito, R., Yamada, N., Iijima, A., Shiga, N., Imai, I.,
802 2013. Interannual changes in the zooplankton community structure on the southeastern
803 Bering Sea shelf during summers of 1994–2009. *Deep Sea Res. Part II Top. Stud.*
804 *Oceanogr.* 94, 44–56. <https://doi.org/10.1016/j.dsr2.2013.03.018>

805 Piatt, J., Springer, A., 2003. Advection, pelagic food webs and the biogeography of seabirds in
806 Beringia. *Mar. Ornithol.* 31, 141–154.

807 Piatt, J., Jones, T., Kuletz, K., et al., 2018. Unprecedented scale of seabird mortality in the NE
808 Pacific during the 2015-2016 marine heatwave. Presented at the Alaska Marine Science
809 Symposium, Anchorage, Alaska.

810 Pike, G., 1950. Stomach contents of whales caught off the coast of British Columbia. *Fisheries*
811 *Research Board Canadian Pacific Progress Report* 83, 27–28.

812 Pinchuk, A.I., Eisner, L.B., 2017. Spatial heterogeneity in zooplankton summer distribution in
813 the eastern Chukchi Sea in 2012–2013 as a result of large-scale interactions of water
814 masses. *Deep Sea Res. Part II Top. Stud. Oceanogr.* 135, 27–39.
815 <https://doi.org/10.1016/j.dsr2.2016.11.003>

816 Pisareva, M.N., Pickart, R.S., Spall, M.A., Nobre, C., Torres, D.J., Moore, G.W.K., Whitley,
817 T.E., 2015. Flow of pacific water in the western Chukchi Sea: results from the 2009
818 RUSALCA expedition. *Deep Sea Res. Part I Oceanogr. Res. Pap.* 105, 53–73.
819 <https://doi.org/10.1016/j.dsr.2015.08.011>

820 Pisareva, M.N., 2018. An overview of the recent research on the Chukchi Sea water masses and
821 their circulation. *Russ. J. Earth Sci.* 18, 1–13.

822 Post, E., Alley, R.B., Christensen, T.R., Macias-Fauria, M., Forbes, B.C., Gooseff, M.N., Iler,
823 A., Kerby, J.T., Laidre, K.L., Mann, M.E., Olofsson, J., Stroeve, J.C., Ulmer, F., Virginia,
824 R.A., Wang, M., 2019. The polar regions in a 2°C warmer world. *Sci. Adv.* 5, eaaw9883.
825 <https://doi.org/10.1126/sciadv.aaw9883>

826 Reynolds, R.W., Smith, T.M., Liu, C., Chelton, D.B., Casey, K.S., Schlax, M.G., 2007. Daily
827 High-Resolution-Blended Analyses for Sea Surface Temperature. *J. Clim.* 20, 5473–5496.
828 <https://doi.org/10.1175/2007JCLI1824.1>

829 Root, T.L., Price, J.T., Hall, K.R., Schneider, S.H., Rosenzweig, C., Pounds, J.A., 2003.
830 Fingerprints of global warming on wild animals and plants. *Nature* 421, 57–60.
831 <https://doi.org/10.1038/nature01333>

832 Savage, K., 2017. Alaska and British Columbia large whale unusual mortality event summary
833 report. Protected Resources Division, National Oceanic and Atmospheric Administration,
834 Juneau, AK.

835 Serreze, M.C., Barrett, A.P., Stroeve, J.C., Kindig, D.N., Holland, M.M., 2009. The emergence
836 of surface-based Arctic amplification. *Cryosphere* 3, 11–19. [https://doi.org/10.5194/tc-3-](https://doi.org/10.5194/tc-3-11-2009)
837 [11-2009](https://doi.org/10.5194/tc-3-11-2009)

838 Serreze, M.C., Crawford, A.D., Stroeve, J.C., Barrett, A.P., Woodgate, R.A., 2016. Variability,
839 trends, and predictability of seasonal sea ice retreat and advance in the Chukchi Sea. *J.*
840 *Geophys. Res.: Oceans* 121, 7308–7325. <https://doi.org/10.1002/2016JC011977>

841 Sigler, M.F., Mueter, F.J., Bluhm, B.A., Busby, M.S., Cokelet, E.D., Danielson, S.L., Robertis,
842 A.D., Eisner, L.B., Farley, E.V., Iken, K., Kuletz, K.J., Lauth, R.R., Logerwell, E.A.,
843 Pinchuk, A.I., 2017. Late summer zoogeography of the northern Bering and Chukchi seas.

844 Deep Sea Res. Part II Top. Stud. Oceanogr. 135, 168–189.
845 <https://doi.org/10.1016/j.dsr2.2016.03.005>

846 Širović, A., Hildebrand, J.A., Wiggins, S.M., 2007. Blue and fin whale call source levels and
847 propagation range in the Southern Ocean. J. Acoust. Soc. Am. 122, 1208–1215.
848 <https://doi.org/10.1121/1.2749452>

849 Širović, A., Williams, L.N., Kerosky, S.M., Wiggins, S.M., Hildebrand, J.A., 2013. Temporal
850 separation of two fin whale call types across the eastern North Pacific. Mar. Biol. 160, 47–
851 57. <https://doi.org/10.1007/s00227-012-2061-z>

852 Sleptsov, M.M., 1961. O kolebaniakh chislennosti kitov v Chukotskom more v raznye gody. (On
853 fluctuations in the number of whales in the Chukchi Sea in different years), in: Proceedings
854 of the A.N. Severtsov Institute of Animal Morphology. Nauka, Moscow, pp. 54–64.

855 Spreen, G., Kaleschke, L., Heygster, G., 2008. Sea ice remote sensing using AMSR-E 89-GHz
856 channels. J. Geophys. Res. 113, C02S03. <https://doi.org/10.1029/2005JC003384>

857 Springer, A.M., McRoy, C.P., 1993. The paradox of pelagic food webs in the northern Bering
858 Sea—III. Patterns of primary production. Cont. Shelf Res. 13, 575–599.
859 [https://doi.org/10.1016/0278-4343\(93\)90095-F](https://doi.org/10.1016/0278-4343(93)90095-F)

860 Stabeno, P.J., Kachel, N.B., Moore, S.E., Napp, J.M., Sigler, M., Yamaguchi, A., Zerbini, A.N.,
861 2012. Comparison of warm and cold years on the southeastern Bering Sea shelf and some
862 implications for the ecosystem. Deep Sea Res. Part II Top. Stud. Oceanogr. 65–70, 31–45.
863 <https://doi.org/10.1016/j.dsr2.2012.02.020>

864 Stabeno, P.J., Bell, S.W., Bond, N.A., Kimmel, D.G., Mordy, C.W., Sullivan, M.E., 2019.
865 Distributed Biological Observatory Region 1: Physics, chemistry and plankton in the

866 northern Bering Sea. *Deep Sea Res. Part II Top. Stud. Oceanogr.* 162, 8–21.
867 <https://doi.org/10.1016/j.dsr2.2018.11.006>

868 Stafford, K.M., Mellinger, D.K., Moore, S.E., Fox, C.G., 2007. Seasonal variability and
869 detection range modeling of baleen whale calls in the Gulf of Alaska, 1999–2002. *J.*
870 *Acoust. Soc. Am.* 122, 3378–3390. <https://doi.org/10.1121/1.2799905>

871 Stroeve, J.C., Serreze, M.C., Holland, M.M., Kay, J.E., Malanik, J., Barrett, A.P., 2012. The
872 Arctic’s rapidly shrinking sea ice cover: a research synthesis. *Climatic Change* 110, 1005–
873 1027. <https://doi.org/10.1007/s10584-011-0101-1>

874 Stroeve, J.C., Markus, T., Boisvert, L., Miller, J., Barrett, A., 2014. Changes in Arctic melt
875 season and implications for sea ice loss. *Geophys. Res. Lett.* 41, 1216–1225.
876 <https://doi.org/10.1002/2013GL058951>

877 Tomilin, A.G. 1957. *Zveri SSSR i Prilezhashchikh Stran (The Mammals of the USSR and*
878 *Adjacent Countries)*. Vol. IX. *Kitoobraznye (Cetacea)*. Proceedings of the USSR Academy
879 of Sciences. Nauk USSR, Moscow. 756 pp. [In Russian]

880 Tsujii, K., Otsuki, M., Akamatsu, T., Matsuo, I., Amakasu, K., Kitamura, M., Kikuchi, T.,
881 Miyashita, K., Mitani, Y., 2016. The migration of fin whales into the southern Chukchi Sea
882 as monitored with passive acoustics. *ICES J. Mar. Sci.: Journal du Conseil* 73, 2085–2092.
883 <https://doi.org/10.1093/icesjms/fsv271>

884 Vaughan, D. G., Comiso, J., Allison, I., Carrasco, J., Kaser, G., Kwok, R., et al., 2013.
885 Observations: Cryosphere. In T. F. Stocker, D. Qin, G. K. Plattner, M. Tignor, S. K. Allen,
886 J. Boschung, A. Nauels, et al., (Eds.), *Climate change 2013: The physical science basis.*
887 Contribution of working group I to the fifth assessment report of the intergovernmental

888 panel on climate change Cambridge, United Kingdom and New York, NY, USA:
889 Cambridge University Press.

890 Votrogov, L.M., and M.V. Ivashin. 1980. Sightings of fin and humpback whales in the Bering
891 and Chukchi Seas. Report of the International Whaling Commission 30:247–248.

892 Walsh, J.E., Fetterer, F., Scott Stewart, J., Chapman, W.L., 2017. A database for depicting Arctic
893 sea ice variations back to 1850. Geogr. Rev. 107, 89–107. [https://doi.org/10.1111/j.1931-](https://doi.org/10.1111/j.1931-0846.2016.12195.x)
894 [0846.2016.12195.x](https://doi.org/10.1111/j.1931-0846.2016.12195.x)

895 Wassmann, P., Duarte, C.M., Agustí, S., Sejr, M.K., 2011. Footprints of climate change in the
896 Arctic marine ecosystem. Glob. Change Biol. 17, 1235–1249.
897 <https://doi.org/10.1111/j.1365-2486.2010.02311.x>

898 Watkins, W.A., 1981. Activities and underwater sounds of fin whales. Scientific Reports of the
899 Whales Research Institute 33, 83–117.

900 Watkins, W.A., Tyack, P., Moore, K.E., Bird, J.E., 1987. The 20-Hz signals of finback whales
901 (*Balaenoptera physalus*). J. Acoust. Soc. Am. 82, 1901–1912.

902 Watkins, W.A., Daher, M.A., Repucci, G.M., George, J.E., Martin, D.L., Dimarzio, N.A.,
903 Gannon, D.P., 2000. Seasonality and distribution of whale calls in the North Pacific.
904 Oceanography 13, 62–67.

905 Weingartner, T., 1997. A review of the physical oceanography of the northeastern Chukchi Sea.
906 In: Reynolds, J. (Ed.), Fish Ecology in Arctic North America, American Fisheries Society
907 Symposium. Bethesda, MD, pp. 40–59.

908 Weingartner, T.J., Danielson, S., Sasaki, Y., Pavlov, V., Kulakov, M., 1999. The Siberian
909 Coastal Current: A wind- and buoyancy-forced Arctic coastal current. J. Geophys. Res.
910 Oceans 104, 29697–29713. <https://doi.org/10.1029/1999JC900161>

911 Witteveen, B.H., Wynne, K.M., 2016. Trophic niche partitioning and diet composition of
912 sympatric fin (*Balaenoptera physalus*) and humpback whales (*Megaptera novaeangliae*) in
913 the Gulf of Alaska revealed through stable isotope analysis. *Mar. Mamm. Sci.* 32, 1319–
914 1339. <https://doi.org/10.1111/mms.12333>

915 Wood, K., Wang, J., Salo, S., Stabeno, P., 2015a. The climate of the Pacific Arctic during the
916 first RUSALCA decade 2004–2013. *Oceanography* 28, 24–35.
917 <https://doi.org/10.5670/oceanog.2015.55>

918 Wood, K.R., Bond, N.A., Danielson, S.L., Overland, J.E., Salo, S.A., Stabeno, P.J., Whitefield,
919 J., 2015b. A decade of environmental change in the Pacific Arctic region. *Prog. Oceanogr.*
920 136, 12–31. <https://doi.org/10.1016/j.pocean.2015.05.005>

921 Woodgate, R.A., Aagaard, K., Weingartner, T.J., 2005a. A year in the physical oceanography of
922 the Chukchi Sea: Moored measurements from autumn 1990–1991. *Deep Sea Res. Part II*
923 *Top. Stud. Oceanogr.* 52, 3116–3149. <https://doi.org/10.1016/j.dsr2.2005.10.016>

924 Woodgate, R.A., Aagaard, K., Weingartner, T.J., 2005b. Monthly temperature, salinity, and
925 transport variability of the Bering Strait through flow. *Geophys. Res. Lett.* 32, n/a-n/a.
926 <https://doi.org/10.1029/2004GL021880>

927 Woodgate, R., Stafford, K., Prahl, F., 2015. A synthesis of year-round interdisciplinary mooring
928 measurements in the Bering Strait (1990–2014) and the RUSALCA years (2004–2011).
929 *Oceanography* 28, 46–67. <https://doi.org/10.5670/oceanog.2015.57>

930 Woodgate, R.A., 2018. Increases in the Pacific inflow to the Arctic from 1990 to 2015, and
931 insights into seasonal trends and driving mechanisms from year-round Bering Strait
932 mooring data. *Prog. Oceanogr.* 160, 124–154. <https://doi.org/10.1016/j.pocean.2017.12.007>

933 Zerbini, A.N., Waite, J.M., Laake, J.L., Wade, P.R., 2006. Abundance, trends and distribution of
934 baleen whales off Western Alaska and the central Aleutian Islands. Deep Sea Res. Part I
935 Oceanogr. Res. Pap. 53, 1772–1790. <https://doi.org/10.1016/j.dsr.2006.08.009>

936 **Figure Captions**

937

938 **Fig 1.** Map of the study region with typical annual mean flow patterns of the three dominant
939 water masses in the Bering Strait region and 20-m bathymetric contours (International
940 Bathymetry Chart of the Arctic Ocean, v. 3). Positions of the three moorings along with the
941 boundaries of the study area polygon used in the sea ice concentration analysis are also
942 displayed. Inset map shows estimated call detection range buffers around each mooring (10 and
943 20 km). Note that the Alaskan Coastal Water is only present seasonally.

944 **Fig. 2.** Histograms of monthly sum of hours with fin whale calls ('FWH') recorded at the three
945 mooring sites (A2, A3, and A4) within the Bering Strait region from 2009–2015. The gray-
946 shaded boxes indicate periods when the hydrophones were not recording.

947 **Fig. 3.** Fin whale departure day of the year (DOY) for each year at the A3 mooring site, north of
948 the Bering Strait, along with the line of best fit ($R^2 = 0.203$, $p = 0.311$).

949 **Fig. 4.** Calculated fin whale departure days for each year at site A3 (light blue, solid line) with
950 other non-solid lines indicating the day of the year (DOY) when the daily mean near-surface
951 (ISCAT; red, medium-dashed line) and near-bottom (SBE; blue, dotted line) temperatures first
952 reached $\leq 0^\circ\text{C}$ at the A3 mooring site. The light gray, long-dashed line represents the DOYs
953 when sea ice concentration in the study area first reached $\geq 15\%$ in each year, and the dark gray,
954 dot-dashed line represents when sea ice concentration in the Chukchi Sea reached $\geq 15\%$. See
955 Fig. 1 for boundaries of study area.

956 **Fig. 5.** Plots of the mean temperatures ($^\circ\text{C}$) and salinities (psu) for October of each year for both
957 the near-surface and near-bottom levels of the water column at each mooring site in the study

958 area (A2, A3, and A4; see key for colors, symbols, and line styles). The vertical lines represent
959 the standard deviation of the monthly means.

960 **Fig. 6.** Monthly mean northward water velocity (cm/s) for the June through November at each
961 mooring site in the Bering Strait region (A2, A3, and A4; see key for colors, symbols, and line
962 styles). See Fig. 1 for mooring locations.

963 **Fig. 7.** Daily mean northward wind velocity for days with fin whale calls at site A3 ('FW Days',
964 black squares) and days without fin whale calls ('No FW Days', white triangles) in October.
965 Note that negative values signify southward wind velocities. The number of FW Days and No
966 FW Days is included for reference.

967 **Tables**

968

969 **Table 1.** Recording settings and positions of the three hydrophones. Dates are in the format
 970 'mm/dd/yyyy.'

Mooring	Year	Latitude N	Latitude W	Record Start Date	Record End Date	Sampling Rate (Hz)	Hourly Duty Cycle
A2	2009	65.80°	168.80°	9/1/2009	1/16/2010	16384	12 min
	2010	65.80°	168.80°	8/11/2010	12/8/2010	16384	15 min
	2012	65.80°	168.80°	9/1/2012	5/15/2013	16384	10 min
	2013	65.78°	168.57°	7/15/2013	7/1/2014	8192	20 min
	2014	65.78°	168.57°	7/10/2014	7/4/2015	8192	20 min
	2015	65.78°	168.57°	7/5/2015	7/8/2016	8192	20 min
A3	2009	66.33°	168.97°	9/1/2009	3/3/2010	16384	12 min
	2010	66.33°	168.97°	8/11/2010	2/19/2011	16384	15 min
	2011	66.33°	168.97°	10/1/2011	5/25/2012	8192	10 min
	2012	66.33°	168.97°	9/1/2012	5/17/2013	16384	10 min
	2013	66.33°	168.97°	7/15/2013	7/2/2014	8192	20 min
	2014	66.33°	168.97°	7/10/2014	7/2/2015	8192	20 min
	2015	66.33°	168.97°	7/5/2015	7/8/2016	8192	20 min
A4	2012	65.75°	168.37°	9/1/2012	6/24/2013	16384	10 min
	2013	65.75°	168.26°	7/15/2013	7/2/2014	8192	20 min
	2014	65.75°	168.25°	7/10/2014	7/2/2015	8192	20 min
	2015	65.75°	168.25°	7/5/2015	7/8/2016	8192	20 min

971 **Table 2.** Fin whale detection data for the three moorings, including the dates of the first and last
 972 detection, and total number of days with fin whale calls present ('FW Days').

Year	A2			A3			A4		
	First Detection Date	Last Detection Date	FW Days	First Detection Date	Last Detection Date	FW Days	First Detection Date	Last Detection Date	FW Days
2009	1 Oct	5 Nov	4	23 Sep	8 Nov	33	.	.	.
2010	14 Oct	17 Oct	2	29 Sep	5 Nov	22	.	.	.
2011	.	.	.	1 Oct	18 Nov	28	.	.	.
2012	28 Oct	2 Nov	3	1 Sep	7 Nov	52	None	None	0
2013	22 Sep	15 Nov	7	23 Jul	9 Nov	28	None	None	0
2014	17 Oct	19 Oct	3	9 Aug	13 Nov	37	2 Nov	2 Nov	1
2015	30 Sep	19 Nov	14	8 Aug	20 Nov	71	11 Oct	8 Nov	7

973 **Table 3.** Wilcoxon rank-sum test results comparing fin whale hours (FWH) recorded at A3 in
 974 October of each year. The p -values are listed in the upper section above the diagonal, and the
 975 gray shaded area below the diagonal are the W statistics from the Wilcoxon rank-sum tests (**bold**
 976 W values indicate significant results). Significant p -values ($p < 0.05$) are in **bold*** and indicate
 977 that the distribution of FWHs significantly differed between the two years.

Year	2009	2010	2011	2012	2013	2014	2015
2009	.	0.736	0.147	0.002*	0.42	0.026*	0.000*
2010	174.5	.	0.463	0.002*	0.554	0.062	0.000*
2011	186	220	.	0.002*	0.566	0.003*	0.000*
2012	69	77	76	.	0.003*	0.614	0.007*
2013	223	185	98.5	399.5	.	0.004*	0.000*
2014	96	120.5	57.5	257.5	47.5	.	0.006*
2015	30	14	14.5	110	21	106	.

978 **Table 4.** Sea ice statistics calculated for 2009–2015 for the study area and Chukchi Sea.
 979 Statistics for the study area include: melt initiation date (melt-out date), melt period (number of
 980 days between 80% and 15% sea ice conc.), date when the study area was ice-free (< 15% conc.;
 981 ice-free date), freeze-up period (number of days between 15% and 80% sea ice conc.), and mean
 982 sea ice concentration (%) in the study area on the last date fin whale calls were recorded ('Last
 983 FW'). Statistics for the Chukchi Sea include mean sea ice concentration (%) on the last date fin
 984 whale calls were recorded ('Last FWH').

Year	Study Area							Chukchi Sea
	Melt-Out Date	Ice-Free Date	Melt Period (# of days)	Freeze-up Date	Freeze-up Period (# of days)	Mean Nov. sea ice conc.	Last FW \bar{x} sea ice conc.	Last FW \bar{x} sea ice conc.
2009	14 May	5 June	23	28 Nov	12	30.2%	0.9%	1.5%
2010	20 May	17 June	29	25 Dec	23	3.4%	1.3%	4.5%
2011	27 April	30 May	34	4 Dec	12	13.3%	0.8%	21.0%
2012	16 May	10 June	25	11 Dec	23	21.3%	1.9%	18.8%
2013	5 May	14 June	41	18 Dec	19	6.3%	3.1%	12.8%
2014	1 May	31 May	31	17 Dec	5	5.4%	4.1%	7.07%
2015	4 May	24 May	21	10 Dec	17	20.1%	4.9%	18.2%

985 **Table 5.** Summary of the overall monthly mean along-channel wind velocities (m/s) for October
 986 along with overall means for days with and without fin whale hours (FWH) in October. Wind
 987 velocities were measured at the data point at 67.5°N and 190°W. Values in parentheses are the
 988 Wilcoxon rank-sum *p*-values for the comparison between the overall October mean for each year
 989 (**bold***: significant *p* < 0.05).

Year	All October (m/s)	Days without FWH (m/s)	Days with ≥ 1 FWH (m/s)	Days with ≥6 FWH (m/s)	Days with ≥12 FWH (m/s)	Days with ≥ 18 FWH (m/s)
2009	-2.7	2.5 (0.03*)	-5.2 (0.19)	-6.2 (0.07)	-6.6 (0.21)	NA
2010	-5.1	-3.2 (0.46)	-6.2 (0.58)	-7.3 (0.37)	-7.9 (0.25)	-9.5 (NA)
2011	-2.1	0.6 (0.12)	-4 (0.21)	-6.1 (0.04*)	-8.7 (0.02*)	NA
2012	-0.4	1.3 (0.97)	-0.5 (0.99)	-2.5 (0.45)	-3 (0.33)	-5.8 (NA)
2013	-6.2	-5.8 (0.88)	-6.7 (0.88)	-5.8 (0.84)	-7 (0.82)	NA
2014	-4.8	0.2 (0.04*)	-6.2 (0.35)	-6.8 (0.25)	-7.9 (0.14)	-9.2 (NA)
2015	-4.0	5.5 (0.18)	-4.3 (0.86)	-4.7 (0.68)	-6 (0.255.3)	-7.1 (NA)

990 **Table 6.** Summary table of the Kendall's rank correlation test results for site A3. Correlation
 991 tests were conducted between the number of fin whale hours (FWH) recorded on days with fin
 992 whale calls (FWH > 0) and the daily means of: near-surface and near-bottom temperatures,
 993 along-channel wind and water velocities, water speeds, and SST. Two sets of tests were carried
 994 out: pooled data for all months for all years (2009–2015), and on October data only for all years
 995 at A3 (2009–2015).

Environmental Variable (Daily Means)	Pooled data - all months (n = 271)		Oct only - all years pooled (n = 156)	
	<i>p</i>	τ	<i>p</i>	τ
Near-surface Temperature	0.674	0.019	0.012*	0.151
Near-surface Salinity	0.053	-0.087	0.851	-0.011
Near-bottom Temperature	0.82	0.01	0.129	0.084
Near-bottom Salinity	0.29	-0.044	0.507	0.037
Water Speed	< 0.001*	-0.167	< 0.001*	-0.28
SST	0.202	-0.054	0.044*	0.111
Along-channel water velocity	< 0.001*	-0.15	< 0.001*	-0.207
Along-channel wind velocity	< 0.001*	-0.194	< 0.001*	-0.231

Figures




Fig 1. Map of the study region with typical annual mean flow patterns of the three dominant water masses in the Bering Strait region and 20-m bathymetric contours (International Bathymetry Chart of the Arctic Ocean, v. 3). Positions of the three moorings along with the boundaries of the study area polygon used in the sea ice concentration analysis are also displayed. Inset map shows estimated call detection range buffers around each mooring (10 and 20 km). Note that the Alaskan Coastal Water is only present seasonally.




Fig. 2. Histograms of monthly sum of hours with fin whale calls (‘FWH’) recorded at the three mooring sites (A2, A3, and A4) within the Bering Strait region from 2009–2015. The gray-shaded boxes indicate periods when the hydrophones were not recording.




Fig. 3. Fin whale departure day of the year (DOY) for each year at the A3 mooring site, north of the Bering Strait, along with the line of best fit ($R^2 = 0.203$, $p = 0.311$).




Fig. 4. Calculated fin whale departure days for each year at site A3 (light blue, solid line) with other non-solid lines indicating the day of the year (DOY) when the daily mean near-surface (ISCAT; red, medium-dashed line) and near-bottom (SBE; blue, dotted line) temperatures first reached $\leq 0^{\circ}\text{C}$ at the A3 mooring site. The light gray, long-dashed line represents the DOYs when sea ice concentration in the study area first reached $\geq 15\%$ in each year, and the dark gray, dot-dashed line represents when sea ice concentration in the Chukchi Sea reached $\geq 15\%$. See Fig. 1 for boundaries of study area.




Fig. 5. Plots of the mean temperatures (°C) and salinities (psu) for October of each year for both the near-surface and near-bottom levels of the water column at each mooring site in the study area (A2, A3, and A4; see key for colors, symbols, and line styles). The vertical lines represent the standard deviation of the monthly means.




Fig. 6. Monthly mean northward water velocity (cm/s) for the June through November at each mooring site in the Bering Strait region (A2, A3, and A4; see key for colors, symbols, and line styles). See Fig. 1 for mooring locations.




Fig. 7. Daily mean northward wind velocity for days with fin whale calls at site A3 ('FW Days', black squares) and days without fin whale calls ('No FW Days', white triangles) in October. Note that negative values signify southward wind velocities. The number of FW Days and No FW Days is included for reference.