

Pelagic Plastic Pollution within the Surface Waters of Lake Michigan, USA

Sherri A. Mason,^{a,*} Laura Kammin,^b Marcus Eriksen,^c Ghadah Aleid,^a Stiv Wilson,^{c,1} Carolyn Box,^c Nick Williamson,^a Anjanette Riley^b

^aState University of New York at Fredonia, Department of Chemistry and Biochemistry, 280 Central Ave., Science Complex 340, Fredonia, NY 14063, USA.

^bIllinois-Indiana Sea Grant, 1101 W. Peabody Drive, Urbana, IL 61801, USA.

^c5 Gyres Institute, 3131 Olympic Blvd #302, Santa Monica, CA, 90404, USA.

*Corresponding author: E-mail: mason@fredonia.edu; voice: 716.673.3292

¹Present Address: The Story of Stuff Project, 1442 A Walnut Street #272, Berkeley, CA 94709, USA.

Abstract

During the summer of 2013, a total of 59 surface water samples were collected across Lake Michigan making it the best surveyed for pelagic plastics of all the Laurentian Great Lakes. Consistent with other studies within the Great Lakes, Mantra-trawl samples were dominated by particles less than 1 mm in size. Enumeration of collected plastics under a microscope found fragments to be the most common anthropogenic particle type, followed by fibers, with more minor contributions from pellets, films and foams. The majority of these pelagic plastic particles were found to be polyethylene, with polypropylene being the second most common polymeric type, which is consistent with manufacturing trends and beach survey results. The pelagic plastic was found to be fairly evenly distributed across the entire Lake Michigan surface, despite the formation of a seasonal gyre at the southern end of the lake. We found an average plastic abundance of $\sim 17,000$ particles/km², which when multiplied by the total surface area, gives on the order of 1 billion plastic particles floating on the surface of Lake Michigan. As the majority of these particles are extremely small, less than 1 mm in size, which allows for easy ingestion, these results highlight the need for additional studies with regard to possible impacts upon aquatic organisms.

Keywords

Plastic Pollution, Marine Debris, Great Lakes, Lake Michigan, Microplastics, Microfibers

Introduction

Global plastic production exceeded 300 million tonnes per year in 2014 (Plastics Europe, 2015). Estimates vary as to how much plastic ultimately reaches the oceans (Thompson, 2006; Jambeck et al., 2015), but without proper waste management and products designed for recovery or environmental degradation, the amount is expected to continue to grow over the next few decades as production continues to increase. While the presence of plastic debris in marine ecosystems has been well documented (Colton et al., 1974; Law et al., 2010; Moore et al., 2001; Thompson et al., 2004; Cózar et al., 2014), freshwater ecosystems, such as rivers and lakes, have only recently begun to be investigated (Eriksen et al., 2013; Hoellein et al., 2014; Free et al., 2014; Corcoran et al., 2015; Dris et al., 2015; Eerkes-Medrano et al., 2015). The Laurentian Great Lakes are the largest surface freshwater ecosystem in the world, containing nearly 20% of the world supply of surface freshwater. The water from the lakes and the channels that connect them eventually flows to the Atlantic Ocean; and thus the presence, transport and fate of microplastics in the Laurentian Great Lakes is of concern not only for freshwater systems, but also for the marine environment. Recent shoreline, stream and open-water surveys of the Great Lakes have found densities of plastics as high as those reported in oceanic gyres (Eriksen et al., 2013; Zybyszewski and Corcoran, 2011; McCormick et al., 2014; Zybeszewski et al., 2014).

Microplastics are defined as plastic particles less than 5mm in size (Thompson et al., 2009). Given their small size, mobility and widespread distribution, microplastics have a high potential to be ingested by aquatic organisms (Browne et al., 2008; Graham and Thompson, 2009; Murray and Cowie, 2011; Lusher et al., 2013). Direct effects of ingestion, such as abrasions or blockages and subsequent starvation, are likely to be less pronounced with micro- as compared to macro-

plastics. Of more concern for microplastics are potential secondary effects, such as the ability of the plastic to transfer inherent or adsorbed persistent organic pollutants (POPs) into the organism. Studies have shown that POPs will preferentially partition onto plastic reaching concentrations up to a million times greater than in the surrounding aqueous environment (Mato et al., 2001; Hirai et al., 2011). Once ingested, these chemicals can desorb, leading to a variety of negative impacts (Browne et al., 2013; Rochman et al., 2013; Wright et al., 2013; Chua et al. 2014; Rochman et al., 2014; de Sa et al., 2015; Tanaka et al., 2015). The longer the particles remain within the organism the greater the potential for this desorption and transfer of toxins (Andrady 2011). While this study was not focused on the impacts of plastic pollution, understanding the implications of such marine debris on aquatic organisms does provide context for its results.

Relative to oceans there is limited research on the distribution and abundance of plastic debris in the Great Lakes (Driedger et al., 2015) and only one other open-water study of pelagic plastics (Eriksen et al., 2013). In this study, we conducted the first open-water survey for plastic pollution in Lake Michigan in order to examine the abundance, composition, distribution and potential sources of plastic debris. This study is the most comprehensive single-lake, open-water study conducted in the Great Lakes to date.

Materials and Methods

Sample collection

During the summer of 2013, a total of 59 open-water samples were obtained across the surface of Lake Michigan (Figure 1). All samples were collected with the use of a manta trawl, which

consists of an aluminum frame with a rectangular opening 16 cm high by 61 cm wide attached to a 3m long, 333 micron-mesh net with a 30 x 10 cm² collecting bag. The trawl was towed along the lake surface and was positioned with the towline sitting outside of the ship's wake. The sea state for all sample sites was fairly calm with values between 1 and 3 on the Beaufort Wind Scale. While the sample sites were not equidistant from one another nor were the transects of equal length, all samples were collected for 30 minutes (timed using a stopwatch) and the tow speed was kept under 2.0 knots. The tow length was determined by three independent measures, (utilizing GPS coordinates, a flowmeter suspended within the mouth of the trawl net and via an onboard knotmeter), and averaged. The surface area sampled could then be calculated by multiplying the tow length by the width of the trawl mouth opening. All samples were rinsed from the collection bag into a sample container and immediately preserved in 70% isopropyl alcohol for later laboratory processing and analysis.

Sample Processing

Open-water samples were processed in a laboratory for isolation of plastic debris using a modified National Oceanic and Atmospheric Administration marine debris protocol (Masura et al., 2015), briefly described here. Each sample is filtered through a series of 8-inch diameter Tyler sieves of 4.75 mm, 1.00 mm and 0.355 mm stainless steel mesh, which separates the solid material into 3 size classifications (0.355-0.999 mm, 1.00-4.749 mm and ≥ 4.75 mm). Solids within the largest size fraction (≥ 4.75 mm) were manually sorted to remove visible plastic debris from organic material. In order to collect any microplastics initially stuck to the labile organic material, all were rinsed with DI water with rinse water running through the stacked sieves. The solids in the smaller two size classifications (0.355-0.999 mm and 1.00-4.749 mm) were

subjected to a wet peroxide oxidation (WPO), which digests labile organic material using 30% hydrogen peroxide in the presence of an iron (II) catalyst. Plastic debris is resistant to this wet peroxide oxidation (WPO) processing (Masura et al., 2015). After processing, samples were once again filtered through a stacked sieve set (1.00 mm and 0.355 mm). Within each size classification (0.355-0.999 mm and 1.00-4.749 mm) all microplastic particles were removed, enumerated and categorized as fragment, pellet, line/fiber, film or foam with a dissection microscope (Free et al., 2014) and later archived. Six blank samples in which DI water was stored within sample containers for periods of 1- 14 days were processed concurrently with the open-water samples and none were found to have any microplastic particulate indicating that the risk of sample contamination from the containers, lab or processing was negligible.

Spectroscopic Analysis

A scanning electron microscope with an elemental detection system (SEM/EDS) was used to analyze the pelagic plastic particles within the smallest size fraction (i.e., 0.355-0.999 mm) in order to distinguish between organic and inorganic materials owing to the more reflective nature of minerals as compared to carbon-based materials (Eriksen et al. 2013). Based upon availabilities at the time of analysis, particles within this size range were not amenable to other spectroscopic techniques (namely FTIR discussed below). Additionally, the SEM can provide detailed images of the particle surface, which are not available via other analytical techniques. Of the 59 samples, 52 contained particles within this smallest size classification. Of these 52 sites, a 20% random selection of sites (11 sites total) were chosen for analysis. For these 11 sites, all particles within the smallest size classification were analyzed using the SEM/EDS system. Samples were prepared for SEM/EDS analysis by securing the particles onto double-sided

carbon tape prior to imaging using a Phenom ProX desktop SEM/EDS system operating at 15 keV in backscatter mode.

Fourier Transform Infrared (FTIR) analysis was used to determine the polymeric composition of retrieved pelagic particles whose diameter (or longest particle edge) were greater than (or equal to) 4.75 mm. Of the 59 sampled sites, 30 sites contained a total of 122 particles within this largest size classification. A total of 72 particles (59%) were analyzed. For this analysis, particles were dissolved in a small amount of dichlorobenzene (~1 mL), which usually necessitated the use of some heat (100°C). A small amount of this solution was then transferred and allowed to dry on a Real Crystal (NaCl) IR Card (International Crystal Laboratories). Samples were analyzed using a Matteson Polaris FTIR operating at 32 scans and 4 cm⁻¹ resolution. Background scans using a Real Crystal IR Card with only dichlorobenzene dried to its surface were obtained every fifth spectra or each new day, whichever came first. Background spectra are automatically subtracted from each raw spectra to yield sample spectra. A library of spectra from known polymers was created by using in-house polymer samples or from common consumer products, such as plastic bottles, containers and cigarette filters. Sample spectra were compared to this library for polymeric composition identification to support visual interpretation.

As the instrumental equipment available at the time of analysis are both 'destructive' techniques (e.g., dissolution to form films for FTIR and bound to carbon tape for SEM/EDS), retrieved pelagic plastic particles within the middle size classification (1.00-4.749 mm) were not analyzed by either technique. Rather these particles were archived for future research endeavors.

Results and Discussion

During the summer of 2013, a total of 59 surface water samples were collected across Lake Michigan (Figure 1). As the only previous pelagic plastics survey in the Great Lakes collected at most 8 samples in each of Lakes Superior, Huron, and Erie (Eriksen et al., 2013), our collection effort on Lake Michigan constitutes the most intensive such survey in the Laurentian Great Lakes to-date. Plastic particles extracted from the samples were categorized and counted (Table 1). Of the 59 samples, all but one contained plastic (station 86, indicated in Figure 1). Using these counts and the surface area sampled (tow length * trawl net width), the plastic abundance (no. of particles per square kilometer, particles/km²) for each sample was determined (Table 1; Figure 1) in order to standardize the data due to the varying tow lengths. Aside from the one sample which did not contain any plastic, abundances varied from ~1400 to 100,000 particles/km².

Plastic particle distribution

Numerical modeling of circulation currents in Lake Michigan notes the creation of an anticyclonic gyre within the southern basin, which develops over the course of the summer and is especially prominent in August (Beletsky et al., 2006). As a result of this seasonal gyre, we expected that we might find greater particle abundances within the southern end of Lake Michigan as compared to the northern end during our August 2013 expedition. Our results, however, do not indicate any particular aggregation of plastic particles as a result of this temporary gyre formation (Figure 1). Rather, our results show a fairly even distribution of plastic particles across the lake surface. We reason the fairly even distribution may be a consequence of: (1) the long residence time of waters within Lake Michigan, (2) the high variability of circulation

currents, especially within the southern end of the lake, and/or (3) wind acting to move any debris with an above-surface profile differently than surface currents carrying subsurface debris. Such 'windage' effects were noted in the movement of debris after the Japanese tsunami (Desforges et al., 2013). Beletsky et al. (2006, 2007) both note a strong interannual variability within the circulation currents within Lake Michigan with warmer years exhibiting anticyclonic currents, which would aggregate particles, but colder years demonstrating cyclonic currents, which would act to transport plastic particles downward. NOAA NowCast data (<http://www.glerl.noaa.gov/res/glcf/>) indicates that August 2013 was a cyclonic year, rather than anticyclonic, which would help explain the lack of aggregation. Additionally, of the Great Lakes, Lake Michigan is second only to Lake Superior in terms of residence time (99 years) owing to its cul-de-sac formation and equal surface elevation to its sister Great Lake, Lake Huron. Water entering the lake circulates slowly and tends to remain for nearly a hundred years (Beletsky et al., 2006) before discharging to Lake Huron, which can also allow for a more even distribution of plastic pollution across the lake surface.

Plastic particle abundances averaged 17,276 particles/km² across the Lake Michigan surface, with the 95% confidence interval of the mean ranging from 12,898 to 21,655 particles/km². Extrapolating across the surface area of Lake Michigan, our data suggest there are on the order of one billion particles across the surface of just this one Great Lake, with 95% confidence values ranging from nearly 750 million up to 1.26 billion particles.

Consistent with the 2012 Great Lakes survey (Eriksen et al., 2013), the vast majority of plastic particles are found within the smallest size classification (0.355-0.999 mm; 59%; Table 2).

However, the distribution of plastic particles differed substantially between these two studies (Table 3). While pellets were the dominant particle type in the 2012 study, fragments dominate the pelagic plastic obtained from Lake Michigan (79%), with pellets making a more minor contribution (4%; Table 3). Interestingly, fibers were the second most abundant type (14%; Table 2), which stands in contrast to the other Great Lakes study, in which they were only a minor component (Table 3; Eriksen et al., 2013).

This difference in the distribution of pelagic plastic particle types could be attributed simply to spatio-temporal variations or may be a factor of differences in methodology. The sources of microplastics are diverse and include both primary and secondary sources. Primary sources include preproduction pellets and powders (Mato et al., 2001), as well as polyethylene and polypropylene microbeads used in many personal care products, such as facial scrubs and toothpastes (Gregory, 1996, Fendall and Sewell, 2009). Secondary sources are produced by mechanical and photo-oxidative degradation (Singh and Sharma, 2008) of plastic bags, bottles, fishing line and nets, and other litter types into smaller fragments (Browne et al. 2007; Cole et al., 2011). The prominence of fragments within the Lake Michigan samples as compared to pellets seems to indicate that secondary sources outweigh primary in this waterbody. This could be affiliated with the longer residence time of waters within Lake Michigan, which allow for longer exposure of primary plastics to UV light and freeze-thaw cycles that lead to brittleness, as well as the mechanical abrasion of wind and waves. The prominence of pellets within the Eriksen et al. 2012 study, largely due to Lake Erie samples, as compared to these Lake Michigan samples, could also be owing to a greater concentration of primary sources (pellet producers and

consumers) in the Sarnia area of Lake Huron and around Lake Erie (Zbyszewski and Corcoran, 2011; Zbyszewski et al., 2014) as compared to the Lake Michigan watershed.

In addition to these possible spatio-temporal variations, the sample processing methodology in the 2012 study differed from that employed here. Eriksen et al. (2013) separated plastic from the associated seston via density-flotation separation using salt-water (a common technique in oceanic studies), but here we utilized the wet peroxide oxidation technique. Fibers are typically composed of more negatively buoyant polymeric materials, such as polyamides, polyethylene terephthalate and polyester. Plastic fragments and pellets, however, are typically composed of polyethylene and polypropylene, which are positively buoyant polymers. Because of this difference in density, plastic fragments and pellets may be more easily separated via the density-flotation method as compared to the fibers causing the fibers to not be as prominent in the 2012 samples as compared to the Lake Michigan samples.

Plastic composition analysis

Several studies have confirmed the need to analyze suspected particulates obtained in environmental sampling in order to confirm their identification as plastic as compared to natural materials (Fillela, 2015; Eerkes-Medrano et al., 2015; Eriksen et al., 2013). Similar to the results from Eriksen et al. (2013), an average of 16% of the particles within the 0.355-0.999 mm size class were found to be misidentified as plastic when they were really of mineral origin based upon SEM/EDS analysis (Table 4). All counts within this size class were adjusted accordingly (Table 1). FTIR results of the 72 particles (59%) analyzed within the largest size classification (Table 5) found thirty-three (46%) to be polyethylene, with high-density polyethylene being

more prevalent (84%) as compared to low-density. Polypropylene was found to be the second most abundant polymeric type (33%) within these samples. These findings are consistent with global trends in the mass production of plastics, with polyethylene, followed by polypropylene, being the most widely manufactured polymers (Plastics-The Facts 2015) and with the findings of beach surveys conducted along the shores of the Laurentian Great Lakes (Zbyszewski and Corcoran, 2011). The remaining 20% of plastic particles analyzed are believed to be either copolymers (13%) or could not be identified (8%) given our instrumentation and polymer libraries (Table 5).

Conclusions

Open-water sampling of Lake Michigan revealed a fairly even distribution of pelagic plastic particles across the lake surface, despite the usual formation of a summer seasonal gyre at the southern end of the lake. This seasonal gyre formation is, however, highly variable, depending largely on water temperatures, with warmer years leading to the anticyclonic currents that would be associated with particle aggregation. As August 2013 was a cooler year, which resulted in cyclonic current formation that is associated with downward mobility of water and dispersal of surface particles, this may explain the lack of a plastic particulate ‘hot spot’ during this sampling expedition. Long residence time (99 years) of waters within Lake Michigan and ‘windage’ effects may also be factors in the surface distribution. Consistent with previous studies within the Laurentian Great Lakes, plastic particles less than 1 mm were the dominant size classification. This size class could have significant consequences for aquatic organisms as these smaller particles have greater surface area for possible adsorption of persistent organic pollutants, which are of concern in the Great Lakes, and are more easily ingested whether intentionally or not.

Fragments were the most abundant type of plastic particle, indicating secondary sources being more significant than primary. Ultimately, understanding the sources, spatio-temporal trends, eventual sinks and possible food web impacts of this pelagic plastic pollution will require additional study.

Acknowledgements

The authors wish to thank the financial support of Illinois-Indiana Sea Grant and the Burning River Foundation. Partial support was also provided by 5 Gyres Institute and the SUNY Fredonia Department of Chemistry. G. Aleid was supported through a Saudi National Government fellowship. The authors also wish to express their extreme gratitude to the captains and crew of the *Niagara*, *Inland Seas* and *Free at Last*, without whose support this research would not have been possible. Additional thanks to Katie Donnelly, an undergraduate research student at SUNY Fredonia, who provided additional assistance with regard to the GIS maps. Lastly, the authors wish to thank the two peer-reviewers, whose insightful comments helped to strengthen this manuscript.

References

- Andrady, A. L. 2011. Microplastics in the marine environment. *Mar. Pollut. Bull.* 62, 1596-1605.
- Beletsky, D.; Schwab, D.; McCormick, M. 2006. Modeling the 1998–2003 summer circulation and thermal structure in Lake Michigan, *J. Geophys. Res.* 111, 2156-2202.
- Beletsky, D.; Mason, D. M.; Schwab, D. J.; Rutherford, E. S.; Janssen, J.; Clapp, D. F.; Dettmers, J. M. 2007. Biophysical model of larval yellow perch advection and settlement in Lake Michigan, *J. Gt. Lakes Res.* 33, 842-866.
- Browne, M.A.; Galloway, T.; Thompson, R. 2007. Microplastics—An emerging contaminant of potential concern? *Integr. Environ. Assess. Manag.* 3, 559-566.
- Browne, M. A.; Dissanayake, A.; Galloway, T. S.; Lowe, D. M.; Thompson, R. C. 2008. Ingested microscopic plastic translocates to the circulatory system of the mussel, *Mytilus edulis* (L.). *Environ. Sci. Technol.* 42, 5026-5031.
- Browne, M. A.; Niven, S. J.; Galloway, T. S.; Rowland S. J.; Thompson, R.C. 2013. Microplastic moves pollutants and additives to worms, reducing functions linked to health and biodiversity. *Curr. Biol.* 23, 2388-2392.
- Chua, E. M.; Shimeta, J.; Nugegoda, D.; Morrison, P. D.; Clarke, B. O. 2014. Assimilation of polybrominated diphenyl ethers from microplastics by the marine amphipod, *allorchestes compressa*. *Environ. Sci. Technol.* 48, 8127 – 8134.
- Cole, M.; Lindeque, P.; Halsband, C.; Galloway, T. S. 2011. Microplastics as contaminants in the marine environment: A review. *Mar. Pollut. Bull.* 62, 2588-2597.
- Colton, J. B.; Knapp, F. D.; Burns, B. R. 1974. Plastic particles in surface waters of the northwestern Atlantic. *Sci.* 185, 491-497.
- Corcoran, P. L.; Norris, T.; Ceccanese, T.; Walzak, M. J.; Helm, P. A.; Marvin, C. H. 2015. Hidden plastics of Lake Ontario, Canada and their potential preservation in the sediment record. *Environ. Pollut.* 204, 17-25.
- Cózar, A.; Echevarría, F.; González-Gordillo, J. I.; Irigoien, X.; Ubeda, B.; Hernández-León, S.; Palma, A. T.; Navarro, S.; García-de-Lomas, J.; Ruiz, A.; Fernández-dePuelles, M.L.; Duarte, C. M. 2014. Plastic debris in the open ocean. *Proc. Natl. Acad. Sci.* 111, 10239-10244.

de Sa, L.C.; Luis, L.G.; Guilhermino, L. 2015. Effects of microplastics on juveniles of the common goby (*Pomatoschistus microps*): Confusion with prey, reduction of the predatory performance and efficiency, and possible influence of developmental conditions. *Environ. Pollut.* 196, 359-362.

Desforges, J-P. W.; Galbraith, M.; Dangerfield, N.; Ross, P. S. 2013. Widespread distribution of microplastics in subsurface seawater in the NE Pacific Ocean. *Mar. Pollut. Bull.* 79, 94-99.

Drieger, A. G. J.; Durr, H. H.; Mitchell, K.; Van Cappellen, P. 2015. Plastic debris in the Laurentian Great Lakes: A review. *J. Gt. Lakes Res.* 41, 9-19.

Dris, R.; Imhof, H.; Sanchez, W.; Gasperi, J.; Galgani, F.; Tassin, B.; Laforsch, C. 2015. Beyond the ocean: contamination of freshwater ecosystems with (micro-)plastic particles. *Environ. Chem.* 12, 539-550.

Eerkes-Medrano, D.; Thompson, R. C.; Aldridge, D. C. 2015. Microplastics in freshwater systems: A review of the emerging threats, identification of knowledge gaps and prioritization of research needs. *Water Res.* 75, 63-82.

Eriksen, M.; Lebreton, L. C. M.; Carson, H. S.; Thiel, M.; Moore, C. J.; Borerro, J. C.; Galgani, F.; Ryan, P.G.; Reisser, J. 2014. Plastic pollution in the world's oceans: More than 5 trillion plastic pieces weighing over 250,000 tons afloat at sea. *PLOS One* 9(12), <http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0111913>

Eriksen, M.; Mason, S.; Wilson, S.; Box, C.; Zellers, A.; Edwards, W.; Farley, H.; Amato, S. 2013. Microplastic pollution in the surface waters of the Laurentian Great Lakes. *Mar. Pollut. Bull.* 77, 177-182.

Fendall, L. S.; Sewell, M. A. 2009. Contributing to marine pollution by washing your face: Microplastics in facial cleaners. *Mar. Pollut. Bull.* 58, 1225-1228.

Fillela, M. 2015. Questions of size and numbers in environmental research on microplastics: methodological and conceptual aspects. *Environ. Chem.* 12, 527-538.

Free, C. M., Jensen, O. P.; Mason, S. A.; Eriksen, M.; Williamson, N. J.; Boldgiv, B. 2014. High-levels of microplastic pollution in a large, remote, mountain lake. *Mar. Pollut. Bull.*, 85, 156.

Graham, E. R.; Thompson, J. T. 2009. Deposit- and suspension-feeding sea cucumbers (*Echinodermata*) ingest plastic fragments. *J. Exper. Mar. Biol. Ecol.* 368, 22-29.

Gregory, M. 1996. Plastic 'scrubbers' in hand cleansers: A further (and minor) source for marine pollution identified. *Mar. Pollut. Bull.* 32, 867-871.

Hirai, H.; Takada, H.; Ogata, Y.; Yamashita, R.; Mizukawa, K.; Saha, M.; Kwan, C.; Moore, C.; Gray, H.; Laursen, D.; Zettler, E. R.; Farrington, J. W.; Reddy, C. M.; Peacock, E. E.; Ward, M. W. 2011. Organic micropollutants in marine plastic debris from the open ocean and remote and urban beaches. *Mar. Pollut. Bull.* 62, 1683-1692.

Hollein, T.; Rojas, M.; Pink, A.; Gasiior, J.; Kelly, J. 2014. Anthropogenic litter in urban freshwater ecosystems: Distribution and microbial interactions. *PLOS One* 9, e98485.

Jambeck, J. R.; Geyer, J.; Wilcox, C.; Siegler, T. R.; Perryman, M.; Andrady, A.; Narayan, R.; Law, K. L. 2015. Plastic waste inputs from land into the ocean. *Sci.* 347, 768-771.

Law, K. L.; Morét-Ferguson, S.; Maximenko, N.; Proskurowski, G.; Peacock, E. E.; Hafner, J.; Reddy, C. M. 2010. Plastic accumulation in the North Atlantic subtropical gyre. *Sci.* 329 1185-1188.

Lusher, A. L.; McHugh, M.; Thompson, R. C. 2013. Occurrence of microplastics in the gastrointestinal track of pelagic and demersal fish from the English Channel. *Mar. Pollut. Bull.* 76, 94-99.

Masura, J.; Baker, J.; Foster, G.; Arthur, C. 2015. Laboratory methods for the analysis of microplastics in the marine environment: recommendations for quantifying synthetic particles in waters and sediments. NOAA Tech. Memo. NOS-OR&R-48.

Mato, Y.; Isobe, T.; Takada, H.; Kanehiro, H.; Ohtake, C.; Kaminuma, T. 2001. Plastic resin pellets as a transport medium for toxic chemicals in the marine environment. *Environ. Sci. Technol.* 35, 318-324.

McCormick, A.; Hoellein, T. J.; Mason, S. A.; Schluep, J.; Kelly, J. J., 2014. Microplastic is an abundant and distinct microbial habitat in an urban river. *Environ. Sci. Technol.* 48, 11863-11871.

Moore, C. J.; Moore, S. L.; Leecaster, M. K.; Weisberg, S. B. 2001. A comparison of plastic and plankton in the North Pacific central gyre. *Mar. Pollut. Bull.* 42, 1297-1300.

Murray, F.; Cowie, P.R. 2011. Plastic contamination in the decapod crustacean *Nephrops norvegicus* (Linnaeus, 1758). *Mar. Pollut. Bull.* 62, 1207-1217.

Plastics Europe. 2015. *Plastics – the Facts 2014/2015: An analysis of European plastics production, demand and waste data*. Brussels, Belgium.

Rochman, C.M.; Hoh, E.; Kurobe, T.; Teh, S.J., 2013. Ingested plastic transfers hazardous chemicals to fish and induces hepatic stress. *Sci. Rep.* 3, 3263.

Rochman, C.M.; Kurobe, T.; Flores, I.; Teh, S.J., 2014. Early warning signs of endocrine disruption in adult fish from the ingestion of polyethylene with and without sorbed chemical pollutants from the marine environment. *Sci. Total Environ.* 493C, 656-661.

Singh, B.; Sharma, N. 2008. Mechanistic implications of plastic degradation. *Poly. Degrad. Stabil.* 440, 561-584.

Tanaka, K.; Takada, H.; Yamashita, R.; Mizukawa, K.; Fukuwaka, M.; Watanuki, Y. 2015. Facilitated Leaching of Additive-Derived PBDEs from Plastic by Seabirds' Stomach Oil and Accumulation in Tissues. *Environ. Sci. Technol.* 49, 11799 – 11807.

Thompson, R. C. 2006. Plastic debris in the marine environment: consequences and solutions, in: Krause, J.C., von Nordheim, H., Bräger, S. (Eds.), *Marine Nature Conservation in Europe*, Stralsund, Germany, pp. 107-116.

Thompson, R. C.; Olsen, Y.; Mitchell, R. P.; Davis, A.; Rowland, S. J.; John, A. W. G.; McGonigle, D.; Russell, A. E. 2004. Lost at sea: Where is all the plastic? *Sci.* 304, 838.

Thompson, R. C.; Moore, C. J.; vom Saal, F. S.; Swan, S. H. 2009. Plastics, the environment and human health: current consensus and future trends. *Philos. Trans. R. Soc. B* 364, 2153-2166.

Wright, S.L.; Rowe, D.; Thompson, R. C.; Galoway, T.S., 2013. Microplastic ingestion decreases energy reserves in marine worms. *Curr. Biol.* 23: R1031-R1033.

Zbyszewski, M., Corcoran, P., 2011. Distribution and degradation of fresh water plastic particles along the beaches of Lake Huron, Canada. *Water Air Soil Pollut.* 220, 365-372.

Zbyszewski, M., Corcoran, P.L.; Hockin, A. 2014. Comparison of the distribution and degradation of plastic debris along shorelines of the Great Lakes, North America. *J. Gt. Lakes Res.* 40, 288-299.

Tables

Table 1. Locations, counts and abundances of pelagic plastic for all 59 Lake Michigan samples.

Sample ID	Date	Latitude	Longitude	Particle Counts						Tow Length (km)	Abundance (count/km ²)
				Fragments	Pellets	Fibers/Lines	Films	Foams	Total		
GL13 #23	17-Jun-2013	44.99	-85.63	10	--	6	--	--	16	1.54	17,082
GL13 #24	18-Jun-2013	44.96	-85.58	6	--	--	1	--	7	2.05	5,593
GL13 #25	26-Jun-2013	44.79	-85.61	10	--	--	1	--	11	2.36	7,657
GL13 #26	8-Jul-2013	45.29	-85.73	5	1	7	1	--	14	2.01	11,414
GL13 #27	8-Jul-2013	45.42	-86.25	108	--	23	6	--	137	2.25	100,016
GL13 #28	24-Jul-2013	45.34	-85.77	21	--	4	5	--	30	2.29	21,474
GL13 #29	25-Jul-2013	45.20	-85.48	32	2	17	--	--	51	2.33	35,811
GL13 #40	2-Aug-2013	45.83	-85.00	31	--	23	--	--	53	2.26	38,773
GL13 #41	2-Aug-2013	45.79	-85.12	18	--	4	--	--	22	2.69	13,430
GL13 #42	2-Aug-2013	45.58	-85.39	6	1	2	--	--	9	2.08	7,084
GL13 #43	3-Aug-2013	45.57	-86.15	20	--	--	--	--	20	2.09	15,670
GL13 #44	3-Aug-2013	45.26	-86.23	33	3	5	1	--	42	2.51	27,457
GL13 #45	3-Aug-2013	45.14	-86.25	2	5	--	--	--	7	2.28	5,023
GL13 #46	3-Aug-2013	45.08	-86.53	5	1	--	1	--	7	2.59	4,433
GL13 #47	3-Aug-2013	45.00	-86.83	14	--	7	--	--	21	2.90	11,853
GL13 #48	4-Aug-2013	44.75	-86.64	4	--	--	--	--	4	4.66	1,407
GL13 #49	4-Aug-2013	44.34	-86.60	15	--	1	--	--	16	1.57	16,662
GL13 #50	4-Aug-2013	44.34	-86.99	11	--	2	--	--	14	1.92	11,629
GL13 #51	4-Aug-2013	44.22	-87.31	14	--	--	1	2	17	1.92	14,523
GL13 #52	4-Aug-2013	43.92	-87.04	31	--	1	--	--	32	2.40	21,890
GL13 #53	4-Aug-2013	43.72	-86.77	18	--	2	2	--	22	2.19	16,449
GL13 #54	4-Aug-2013	43.56	-87.07	84	4	7	--	--	95	2.04	76,284
GL13 #55	5-Aug-2013	43.66	-87.43	25	2	7	--	--	34	2.15	25,869
GL13 #56	5-Aug-2013	43.36	-87.36	48	14	3	1	--	66	2.17	49,838
GL13 #57	5-Aug-2013	43.21	-87.61	5	1	5	--	--	11	2.11	8,553
GL13 #58	5-Aug-2013	43.03	-87.87	36	--	3	--	--	39	2.41	26,516
GL13 #59	6-Aug-2013	43.03	-87.89	33	--	4	--	--	37	2.07	29,241
GL13 #60	6-Aug-2013	42.99	-87.82	29	3	4	1	--	37	2.09	29,072
GL13 #61	6-Aug-2013	42.88	-87.49	4	2	--	--	--	6	2.28	4,315
GL13 #62	6-Aug-2013	42.76	-87.21	3	--	--	--	--	3	2.30	2,138
GL13 #63	6-Aug-2013	42.54	-87.43	10	--	--	1	2	13	1.92	11,109
GL13 #64	7-Aug-2013	42.33	-87.62	4	--	--	1	--	5	2.51	3,261
GL13 #65	7-Aug-2013	42.30	-87.35	12	1	--	--	--	13	1.76	12,113
GL13 #66	7-Aug-2013	41.97	-87.49	9	1	--	--	--	9	2.11	7,311
GL13 #67	7-Aug-2013	41.89	-87.55	21	--	3	--	--	24	2.04	18,968
GL13 #68	7-Aug-2013	41.89	-87.59	20	2	4	--	--	26	2.21	19,321
GL13 #69	8-Aug-2013	42.15	-87.60	12	--	3	--	--	15	1.98	12,441
GL13 #70	8-Aug-2013	42.29	-87.59	4	--	3	--	--	7	1.57	7,290
GL13 #71	9-Aug-2013	42.63	-87.65	20	--	3	--	--	23	1.74	21,613
GL13 #72	9-Aug-2013	42.79	-87.47	8	1	--	--	--	9	1.79	8,264
GL13 #73	9-Aug-2013	42.77	-87.21	12	--	6	--	--	18	1.62	18,215
GL13 #74	9-Aug-2013	42.75	-86.97	11	1	1	--	1	13	1.29	16,877
GL13 #75	9-Aug-2013	42.72	-86.78	25	--	--	--	--	25	1.91	21,465
GL13 #76	9-Aug-2013	42.57	-86.55	11	4	4	--	5	24	1.89	20,821
GL13 #77	10-Aug-2013	42.38	-86.45	7	--	--	1	--	8	2.14	6,133
GL13 #78	10-Aug-2013	42.34	-86.71	17	--	2	1	--	20	1.62	20,228
GL13 #79	10-Aug-2013	42.30	-86.92	14	--	--	1	--	15	1.81	13,618
GL13 #80	10-Aug-2013	42.23	-87.18	10	--	--	--	--	10	1.77	9,656
GL13 #81	10-Aug-2013	42.19	-87.29	13	2	1	--	--	16	1.81	14,496
GL13 #82	10-Aug-2013	42.15	-87.39	24	4	8	--	--	36	2.45	24,052
GL13 #83	10-Aug-2013	42.12	-87.50	15	--	--	--	--	15	2.13	11,528
GL13 #84	10-Aug-2013	42.10	-87.60	27	--	6	--	--	33	2.05	26,403
GL13 #85	19-Aug-2013	44.76	-87.77	7	--	1	1	--	9	3.28	4,504
GL13 #86	19-Aug-2013	45.10	-87.45	--	--	--	--	--	0	2.79	0
GL13 #87	20-Aug-2013	45.45	-86.90	14	--	2	--	--	16	2.41	10,894
GL13 #88	20-Aug-2013	45.58	-86.42	7	--	--	--	--	7	3.66	3,136
GL13 #89	20-Aug-2013	45.74	-86.17	14	--	1	--	--	15	2.33	10,561
GL13 #90	20-Aug-2013	45.83	-85.86	7	--	1	--	--	8	2.48	5,282
GL13 #91	20-Aug-2013	45.89	-85.47	4	--	--	--	--	4	2.53	2,592

Table 2. Plastic abundances (count/km²) averaged over all 59 samples, distinguished by size and particle type.

	0.355—	1.000—	% of Total	
	0.999mm	4.749mm > 4.75mm		
Fragment	8,381.3	4,418.4	866.3	79%
Pellet	540.8	141.9	23.8	4%
Fiber/Line	1,006.3	858.9	549.6	14%
Film	105.3	97.0	139.0	2%
Foam	118.7	29.0	0.0	1%
count/km ²	10,152.5	5,545.2	1,578.8	
% of total	59%	32%	9%	

Table 3. Comparison of results of this study to a previous Laurentian Great Lakes study in percentages by size and particle type.

	Size Classification			Particle Type				
	0.355—	1.000—	% of total	Fragment	Pellet	Fiber/ Line	Film	Foam
	0.999mm	4.749mm > 4.75mm						
Eriksen et al. 2013	81%	17%	2%	42%	48%	0%	1%	8%
This Study	59%	32%	9%	79%	4%	14%	2%	1%

Table 4. Summary of SEM/EDS analysis on retrieved particles in smallest size classification (0.355 – 0.999 mm).

Sample ID	Number of Particles: 0.355 - 0.999 mm	Percent of Total	
		Plastic	Mineral
GL13 #24	7	100%	0%
GL13 #40	37	93%	7%
GL13 #45	7	100%	0%
GL13 #49	3	67%	33%
GL13 #50	16	73%	27%
GL13 #56	23	67%	33%
GL13 #60	30	100%	0%
GL13 #66	8	60%	40%
GL13 #67	20	93%	7%
GL13 #74	15	75%	25%
GL13 #80	8	80%	20%
Total/Average	174	84%	16%

Table 5. Summary of FTIR results on retrieved pelagic plastic particles in largest size classification (≥ 4.75 mm).
PE = polyethylene; PP = Polypropylene

Sample ID	Number of Particles: > 4.75 mm	Type of Polymer			
		PE	PP	Copolymer	unknown
GL13 #23	1	0	1	0	0
GL13 #26	1	0	1	0	0
GL13 #27	6	3	1	1	1
GL13 #28	3	1	2	0	0
GL13 #29	2	1	1	0	0
GL13 #40	1	1	0	0	0
GL13 #44	2	0	1	1	0
GL13 #49	3	0	1	0	2
GL13 #51	1	0	0	0	1
GL13 #54	5	1	1	3	0
GL13 #55	3	3	0	0	0
GL13 #56	15	12	3	0	0
GL13 #59	4	2	1	0	1
GL13 #60	2	0	1	0	1
GL13 #62	3	2	1	0	0
GL13 #65	2	1	1	0	0
GL13 #67	2	0	2	0	0
GL13 #68	3	0	0	3	0
GL13 #75	2	2	0	0	0
GL13 #76	3	0	3	0	0
GL13 #83	3	2	1	0	0
GL13 #85	2	2	0	0	0
GL13 #88	3	0	2	1	0
Total	72	33	24	9	6
Percentage	100%	46%	33%	13%	8%

Figure Captions

Figure 1. Distribution of plastic abundance (particles/km²) for the 59 open-water surface samples obtained across Lake Michigan during the summer 2013. All but one sample- station 86, indicated in the figure- contained plastic.

