




NOAA Technical Memorandum NMFS - SEFC - 250

Utilization of Marsh and Associated Habitats Along a Salinity Gradient in Galveston Bay.



GALVESTON LABORATORY

SOUTHEAST FISHERIES CENTER

NATIONAL MARINE FISHERIES SERVICE

FEBRUARY 1990

NATIONAL OCEANIC AND ATMOSPHERIC
ADMINISTRATION

DEPARTMENT OF COMMERCE



**NOAA TECHNICAL MEMORANDUM
NMFS-SEFC- 250**

**Utilization of Marsh and Associated Habitats
Along a Salinity Gradient in Galveston Bay.**

By

R.J. Zimmerman, T.J. Minello,
M.C. Castiglione, and D.L. Smith

U.S. DEPARTMENT OF COMMERCE
Robert Mosbacher, Secretary

NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION
John A. Knauss, Administrator

NATIONAL MARINE FISHERIES SERVICE
William W. Fox, Jr., Assistant Administrator for Fisheries

FEBRUARY 1990

This Technical Memorandum series is used for documentation and timely communication of preliminary results, interim reports, or similar special-purpose information. Although the memoranda are not subject to complete formal review, editorial control, or detailed editing, they are expected to reflect sound professional work.

ACKNOWLEDGEMENTS

This project was the result of cooperative research between NOAA's National Marine Fisheries Service/Southeast Fisheries Center Galveston Laboratory and the Texas Parks and Wildlife Department and the Texas Water Development Board. The state agencies were mandated to study the effects and needs of freshwater inflow to the States's estuaries by House Bill 2 (1985) and Senate Bill 683 (1987) enacted by the Texas Legislature. As part of the program, this research was funded through the Texas Water Development Board's Water Research and Planning Fund, authorized under Texas Water Code Sections 15.402 and 16.058 (e), and administered by the Texas Parks and Wildlife Department under interagency cooperative contracts Nos. IAC(86-87)1590, IAC(88-89)0821 and IAC(88-89)1457. P. Barrick, T. Czapla, T. Delaney, C. Jackson, J. Kostera, E. Martinez, T. McTigue, and J. Thomas are due special thanks for their assistance in field work.

The National Marine Fisheries Service (NMFS) does not approve, recommend or endorse any proprietary or material mentioned in this publication. No reference shall be made to NMFS, or to this publication furnished by NMFS, in any advertising or sales promotion which would indicate or imply that NMFS approves, recommends, or endorses any proprietary product or proprietary material mentioned herein or which has as its purpose any intent to cause directly or indirectly the advertised product to be used or purchased because of this NMFS publication.

This report should be cited as follows:

**Zimmerman, R. J. , T. J. Minello, M.C. Castiglione, and D. L. Smith. 1990.
Utilization of marsh and associated habitats along a salinity gradient in
Galveston Bay. NOAA Technical Memorandum NMFS-SEFC-250, 68 pp.**

Copies may be obtained by writing:

National Marine Fisheries Service
Galveston Laboratory
4700 Ave. U
Galveston, TX 77551

or

National Technical Information Service
5258 Port Royal Road
Springfield, VA 22161

ABSTRACT

Nursery utilization of estuarine marshes by fishery species was studied in relation to salinity in Galveston Bay. The investigation revealed that effects of salinity on foods may explain distributions of fishery juveniles among marshes. Juvenile shrimp, crab and fish predators follow their prey, and environmental factors, such as the long-term effects of freshwater flows into an estuary, appeared to affect distributions and abundances of prey. Predator and prey abundances varied significantly among marshes in Galveston Bay, thus varying the nursery value of marshes to fishery juveniles.

The highest numbers of penaeid shrimps, blue crab and commercial fishes were in marshes of the middle and lower bay. These abundances were associated with high abundances of benthic peracarid crustaceans (amphipods and tanaidaceans) which have been shown to be used as foods through feeding experiments and gut analyses. Other foods such as annelid worms and bivalve mollusks were less utilized and less related to distributions of fishery juveniles. This cause-and-effect relationship may partly explains differences in utilization of marshes by fishery species.

Habitats in upper Galveston Bay were dominated by long-term effects of low salinity from the Trinity and San Jacinto rivers. Marshes and submerged vegetation at the Trinity River delta were characterized by brackish water plants having highly seasonal growth patterns with complete winter defoliation. This environment was not favorable for development of populations of small estuarine invertebrates, nor to the growth of epiphytic algae. Infauna consisted of a few species of osmoconforming oligochaete worms and bivalve mollusks. Transient species such as juveniles of shrimps, crabs, and fishes had ready access to these marshes but did not use them extensively, despite their own abilities to osmoregulate. This lack of attractiveness was apparently due to the absence of preferred foods, especially epiphytic algae and peracarid crustaceans. Hence the value of upper bay oligohaline marshes was not through direct utilization but was attributed to large quantities of organic detritus exported and utilized downstream.

Organic detritus from the upper bay was an apparent energy source for food chains in the

middle bay. Here, vascular plant detritus and reformed particles from dissolved organics, colonized and conditioned by bacteria and fungi, provided a nitrogen rich food resource for epibenthic detritivores. Stimulated by favorable food and salinity conditions, large numbers of epibenthic fauna developed (as indicated by abundances of peracarids and annelids), providing a rich feeding ground in both marsh and subtidal habitats. Importantly, this mid-bay region was the frontal zone where nutrients from the upper bay mixed with immigrating recruits from the lower bay.

Detritus in the lower bay was important, but perhaps to a lesser extent than in the middle bay. Reduced turbidities and marine salinities of the lower bay fostered the development of epiphytic algae and grazers as another carbon source. *Spartina* marshes in the lower bay persisted year-around and were regularly inundated by marine waters, thus offering perennial substrata for epiphyte colonization. Young shrimp, crabs and fishes were more abundant in marsh habitats, compared to bare subtidal habitat, in the lower bay than in the middle and upper bay.

These collective findings revealed how marshes of the Galveston Bay system are utilized by consumers. Low salinity (oligohaline) marshes in the upper bay (especially at the Trinity Delta) exported large amounts of organic material to the middle bay. The plants of the river delta defoliate each winter and the entire standing crop is exported downstream. Enriched plant detritus in the middle system increases the productivity of epibenthic detritus feeders (such as peracarid crustaceans) and these were foraged by juveniles of commercially valuable fishes, shrimps and crabs. Because both the marsh and subtidal bottom in the middle bay had high abundances of forage organisms, the entire area was valuable nursery habitat. The moderate influence of mesohaline to polyhaline salinities in the middle bay also encouraged utilization by consumers. In the lower bay, algal carbon was another base for secondary productivity in marsh and seagrass habitats heavily epiphytized by algae. Finally, the interconnections between the different systems of the bay appeared to be critical to maintaining overall fishery productivity.

[THIS PAGE INTENTIONALLY LEFT BLANK]

INTRODUCTION

Purpose

The purpose of this study was to characterize marsh use by fishery species relative to salinity regime. Several hypotheses were proposed. The central hypothesis was that marshes in the mid-range salinity regimes are more utilized by the estuarine aquatic fauna than marshes in low or high salinity regimes. Subhypotheses proposed, a) that habitats with mid-range salinities have higher densities of fishery organisms, and b) that habitats with mid-range salinities have greater abundances of foods foraged by juveniles of fishery species.

The Galveston Bay System

The Physical Environment. The physical environment of the Trinity-Galveston Bay system has been reviewed by Wermund et al. (1989). Descriptions from surveys are in Reid (1955), Chambers and Sparks (1959), Pullen et al. (1969, 1971), Diener (1975), the Texas Department of Water Resources (TDWR 1981a and b), Fisher (1983), and White et al. (1985). In his 1963-66 study, Pullen (1971) reported temperature ranges between 0.4 and 36.0° C, salinity ranges between 0.1 and 36.6 ppt, and dissolved O₂ between 0.2 and 13.6 ml/l. From salinity averages, the 10 ppt isohaline line was placed through the middle of Trinity Bay (north to south), the 15 ppt line crossed through the middle of Galveston Bay (east to west) extending the length of East Bay, and the 20 and 25 ppt lines were confined to lower Galveston Bay near the pass into the Gulf of Mexico at Bolivar Roads. The lower bays West Bay and Christmas Bay were not included in early surveys, but salinities are generally known to be higher than the upper and middle bays (including East Bay) due to proximity to major

passes into the Gulf (White et al. 1985).

The Galveston Bay system has about 1,554 km² of open water, intertidal marshes and flats representing 23% of the total estuarine area in Texas (Armstrong 1987). Pullen estimated that the largest bays in the system, Trinity Bay, Galveston Bay, East Bay and West Bay, covered approximately 1,360 km². Despite the relatively large size the system is very shallow with mean depths generally under 2m. Diener (1975) reported on acreage of open water and maximum and mean depth at mean low water for Trinity Bay (337 km²; 5.2m max. and 1.6m mean), upper Galveston Bay (283 km²; 12.8m max. and 1.7m mean), lower Galveston Bay (362 km²; 13.4m max. and 2m mean), East Bay (135 km²; 3.7m max. and 1m mean), West Bay (180 km²; 7.6m max. and 1.2m) and Christmas-Bastrop Bay (39.2 km²; 6.1m max. and 1m mean).

Surface sediments in the Galveston Bay system were described by White et al. (1985) as composed of mud, muddy sand and sandy mud. In general, the upper areas in the system are muddy and the lower areas are sandy. Fine grained mud predominates in Trinity Bay, upper Galveston Bay and East Bay. The Trinity river delta and the passes at either end of Galveston Island are sandy. Bay margins along Bolivar peninsula and Galveston Island are muddy sand. Marsh sediments in the system reflect open bay characteristics; thus Trinity delta marshes are sandy to muddy, upper and middle Galveston Bay and East Bay marshes are muddy, and lower Galveston Bay, West Bay and Christmas Bay marshes are sandy to muddy sand.

The major river inputs in the system are the Trinity and San Jacinto Rivers, contributing 5 million and 1.4 million ac/ft of freshwater per year respectively. About 2.5 million ac/ft/yr is added from local rainfall of 50 inches (127 cm) rainfall/yr (Wermund et al. 1988).

The system includes 603 km² of wetlands (TDWR 1981a). The Trinity River delta is the largest river delta in Texas, comprised of 54 km² of marshes, 68 km² of cypress swamps, and 35 km² of shallow fresh to brackish lakes. Salt marshes cover 140 km² and brackish marshes occupy 230 km² of intertidal wetlands throughout the remainder of the system (Fisher et. al 1972). The balance is freshwater and terrestrial marsh.

Normal tides in the system have a relatively low diurnal amplitude (about 30 cm) as compared to a seasonal range of about 1 m (Hicks et. al 1983). However, because the bay is shallow, meteorological forces of wind and barometric pressure often override tidal forces (Smith 1982). Strong weather fronts from the west and northwest, during the winter months, drive water away from the coast thus lowering water-level in the bay. The opposite effect occurs during the warm season when southeast winds and tropical depressions move water toward the coast and elevate water levels. These forces cause tidal variations that routinely exceed the predicted values, often beyond the annual range. Freshwater inflow from high rainfall also has an effect on elevating water-levels. Trinity delta marshes and other marshes in the upper bay and in East Bay (lower bay) are inundated for extended periods due to flood events (Borey 1979; Texas Dept. Water Resources 1981a; Borey et. al 1983).

Biological Components. The biological components of Galveston Bay have been reviewed by Sheridan et. al (1988). The major fisheries have been described generally and their relationships to freshwater inflow modeled by TDWR (1981a and b). Relationships between benthic invertebrates and sediments in the bay have been characterized by White et. al (1985). Other descriptions of the biota include marsh vegetation (Fisher et. al 1972), benthic algae (Lowe et. al 1978), phytoplank-

ton (TDWR 1981a), zooplankton (Holt and Strawn 1983), molluscan distributions (Harry 1976), oyster reefs (Hofstetter 1977 and 1983), penaeid shrimp populations (Chin 1960; Baxter and Renfro 1967; Parker 1970; Zimmerman and Minello 1984), the blue crab population (More and Moffett 1964; More 1969; Hammerschmitt 1985), and the fish community (Parker 1965; Sheridan 1983).

The biota is dominated by subtropical to temperate estuarine species, including populations of considerable economic value. Penaeid shrimp (*Penaeus aztecus* and *P. setiferus*) lead the economically important species followed by oysters (*Crassostrea virginica*), blue crabs (*Callinectes sapidus*) and finfishes (Sheridan et. al 1988). Commercial and recreational fishes in order of kg landed are spotted seatrout (*Cynoscion nebulosus*), southern flounder (*Paralichthys lethostigma*), sand seatrout (*Cynoscion* spp.), Atlantic croaker (*Micropogonias undulatus*), and red drum (*Sciaenops ocellatus*). All of the commercially important species require the estuary at least as a nursery and many species, commercial and otherwise, are closely associated with marsh habitats. Examples such as grass shrimp (*Palaemonetes pugio*), mud fish (*Fundulus grandis*), and the naked goby (*Gobiosoma boscii*) are marsh residents, and juveniles of brown shrimp, blue crabs and spotted seatrout have been shown to select tidally flooded marsh in preference to non-vegetated mud bottom (Zimmerman and Minello 1984).

Most faunal species occur throughout the system, although abundances may be unevenly distributed depending on location and season. Prior studies indicate a coarse relationship between distributions and salinity. For instance, the clams *Rangia cuneata* and *R. flexuosa* are more abundant in the upper and middle subsystem (fresher), oyster reefs are prevalent in the middle subsystem and hard clams (*Mercenaria mercenaria*) and

bay scallops (*Argopecten*) only occur in small populations in the lower subsystem (more saline).

Emergent marshes are the dominant vegetation throughout the system (Fisher et. al 1972). Submerged aquatic vegetation (SAV) is currently limited to small stands mostly in Trinity Bay and Christmas Bay (West 1972). Sheridan et al. (1988) reports that SAV has declined in the system from about 21 km² in 1960 to <1 km² in 1979. We report on the present composition and seasonal dynamics of the Trinity delta and Christmas Bay grass beds.

Influences of Freshwater Inflow

Recruitment to the Nursery.

Gulf of Mexico species that require estuarine nurseries usually have postlarvae that follow salinity gradients from saline to brackish conditions. Most of these species use freshwater as a cue for directional movement. Planktonic larvae of barnacles and oysters detect salinity differences in water masses and respond behaviorally to effect their transport within estuaries. Swimming behavior by oyster larvae is stimulated by increased salinities and suppressed by decreased salinities (Haskin 1964). This helps larvae position themselves for favorable tidal transport on salinity wedges. Likewise, megalopae of blue crabs and postlarvae of penaeid shrimps move vertically in the water column responding to salinity changes that signal transport into an estuary.

Recruits also depend upon freshwater inflow to sustain the quality of nursery habitat. Since primary production in estuaries is driven by nutrient availability (Nixon 1981), high production depends on nutrients resupplied by freshwater inflow (Flint et. al 1983). A close relationship between estuarine chlorophyll *a* level and river flow (Bennett et. al 1986)

exemplifies this relationship. In northwestern Gulf of Mexico estuaries, nutrients and suspended organic solids are largely imported via riverine flow through freshwater marshes (Stern et. al 1986). In *Spartina* salt marshes nitrogen levels and soil hydrology interact to determine production levels (Mendelssohn 1979; Conner et. al 1987). Salt marshes are also benefited by freshwater through moderation of detrimental high soil salinities (Webb 1983). At the consumer level, high numbers of estuarine infauna (useful as food for fishery juveniles) have been attributed to rainfall and floods (Flint 1985). Increased recruitment and survival of red drum in the Laguna Madre has been related to moderation of hypersaline conditions through floods after hurricanes (Matlock 1987). River transported sediments also supply turbidity and soft substrates that are good refuges from predation for juvenile recruits (Minello et. al 1987).

Nurseries are usually located along the shallow edges of an estuary. The most effective nurseries are vegetated, such as emergent marshes, mangroves, seagrasses, and algae beds. In Galveston Bay, nursery habitat consists of extensive areas of brackish and salt marshes and limited areas of submerged vegetation. Parker (1970) reported that postlarval brown shrimp in Galveston Bay move directly to the marshes after they immigrate through the passes (Baxter and Renfro 1967). Zimmerman et. al (1984) showed that juvenile brown shrimp, from 15 mm to 60 mm total length (TL), were strongly attracted to salt marsh habitat in West Bay. The selective value of the attraction was increased abundances of foods (Gleason and Zimmerman 1984; Gleason 1986; Zimmerman et. al) and greater protective cover from *Spartina alterniflora* (Minello and Zimmerman 1983). Juvenile blue crabs exhibited a similar strong attraction for marsh and seagrass habitats, in West Bay and Christmas Bay, apparently for the same reasons (Thomas 1989; Thomas et. al 1990).

Other important transient species using the West Bay salt marsh as a nursery (in order of abundances) were white shrimp, pinfish, spot (*Leiostomus xanthurus*), bay anchovy (*Anchoa mitchilli*), Atlantic croaker, Gulf menhaden (*Brevoortia patronus*), spotted seatrout, southern flounder, striped mullet (*Mugil cephalus*), and red drum (Zimmerman and Minello 1984). The only estuarine fishes of commercial interest not found as juveniles in the West Bay salt marsh nursery were sheepshead (*Archosargus probatocephalus*) and black drum (*Pogonias cromis*).

Oysters are recruited throughout Galveston Bay forming reefs in areas with salinities ranging between about 10 and 30 ppt (Hopkins 1931; Hofsetter 1977 and 1983; Sheridan et. al 1988). Salinities above 7 ppt are required for spawning (Loosanoff 1953) and spat grow best in salinities above 12 ppt (Davis and Calabrese 1964). Salinities above 20 ppt in Galveston Bay favor populations of oyster drills (*Thais haemastoma*), a predator, and a disease (*Perkinsus marinus*) that reduces oyster numbers (Sheridan et. al, 1988). As a consequence of predation and disease at higher salinities and of physiological limitations at lower salinities, the most productive oyster reefs are in the middle of Galveston Bay where salinities are 10 to 20 ppt (Sheridan et. al 1988).

Fishery Yields. Relationships between freshwater flow into estuaries and fishery production are poorly established and not well understood. An overall review of the influence of freshwater inflows on estuarine productivity is provided by Turek et al. (1987) with citations of case studies and previous reviews by Copeland (1966), Baxter (1977), Armitage (1978), Pandian (1980), Benson (1981) and Peters (1982).

Our present concept of relationships between freshwater input and fisheries yields

arises from inferences based upon correlations. Estuaries are by definition mixtures of fresh and marine waters (Pritchard, 1967) and 69 percent of all finfish and shellfish landings in the U.S. are from estuarine-dependent species (McHugh, 1966 and 1976). A simplified view of estuarine-dependent productivity is dependence upon the freshwater flow which creates estuaries. In this view, large estuarine areas, supported by freshwater inflow, would produce greater fishery yields. This inference is based upon a few studies that show a positive correlation between fishery yield and estuarine area. The most often cited studies are Turner (1977) and Nixon (1982).

The estuarine dependency of fisheries in the Gulf of Mexico is about 98 percent (Gunter, 1967). The Texas Department of Water Resources (1981b) has produced 115 significant multiple regressions from models of Texas estuaries relating fishery yields to the amount of freshwater inflow. Most of these are linked to spring and late fall inflows indicating important seasonal relationships. In addition, a major drought in Texas during the 1950s caused low fishery yields and adverse effects on estuarine populations (Powell, 1985). Estuarine-dependent populations apparently recovered quickly after spring and fall rains in 1957 at the end of the drought (Hoese, 1960).

Habitat Modification. The intertidal marsh surface and shallow water without vegetation (bare bottom) in northwestern Gulf of Mexico estuaries comprise the principal nursery habitats for immigrating postlarvae of fishery species. In the NW Gulf, these habitats occur together in a reticulated pattern with a high degree of interfacing. This habitat mosaic is caused by marsh deterioration resulting from subsidence, loss of sediment input and saltwater intrusion (Craig et al. 1980; Reidenbaugh et al. 1983; Hatton et al. 1983). The condition increases both shore-

line complexity and the opportunity for habitat selection by recruiting animals. This benefits species like brown shrimp whose juveniles select flooded marsh in preference to non-vegetated subtidal bottom (Zimmerman and Minello 1984). In support of this observation, the offshore catch of brown shrimp has been positively correlated with the amount of intertidal marsh area (Turner 1977), shoreline complexity (Faller 1979) and the ratio of marsh to open-water (Browder 1985). However, similar observations have not been made for white shrimp. Young white shrimp demonstrate no consistent preference between marsh surface and bare bottom habitats. The findings suggest differences in the usage and value of marsh for the principal two fishery species in the Gulf of Mexico.

Juvenile brown shrimp are frequently associated with vegetated habitats such as marshes and seagrasses and young white shrimp are commonly identified with open-water, nonvegetated, muddy bottom habitats (Loesch 1965; Christmas et al. 1976; Stokes 1974; and Zimmerman et al. 1984). White shrimp have also been associated with detritus rich sediments (Williams 1955). Recent evidence may explain these different habitat associations through feeding (Zimmerman et al.). Brown shrimp are highly effective in feeding on benthic infauna and epifauna, while white shrimp are much less so. The high numbers of small benthic macrofauna sought by carnivorous brown shrimp are most abundant in vegetated habitats. In the NW Gulf, these habitats are predominantly intertidal marshes. White shrimp have been shown to exploit epiphytes and possibly planktonic resources that brown shrimp do not (McTigue and Zimmerman, in prep). Growth of white shrimp was not different when held in separate cages in marsh and nonvegetated habitats. By contrast, brown shrimp grew more slowly on nonvegetated bottom than in marsh (Zimmerman et al.). Apparently, habitat requirements differ for each species, and habi-

tat changes, such as marsh loss or nutrient enrichment, do not equally affect both species.

Marshes in the NW Gulf are currently not accreting enough sediment to offset subsidence and are sustaining increased salt water intrusion due to diversion of freshwater riverflow (Craig et al., 1980; and Hatton et al., 1983). Although these processes ultimately destroy marsh habitat, the short-term effect may make more habitat available for exploitation. Microtidal diurnal amplitudes in the Gulf are dominated by high seasonal tides (Provost 1976). This effect increases the duration of marsh inundation during spring and fall seasons. A mild climate allows development of high abundances of epifauna and infauna during the winter season (Flint and Younk, 1983). These phenomena provide an abundance of foods, available for spring exploitation, to incoming brown shrimp recruits. Increased accessibility to intertidal habitats appears to be the key to production in brown shrimp and other estuarine-dependent animals that use the marsh surface as a nursery.

METHODS

Study Sites

Marshes in three parts of the Galveston Bay system were chosen for study based on salinity characteristics (Fig.1). The upper, middle and lower parts of the system corresponded to oligohaline (0.5 to 5 ppt), mesohaline (5 to 18 ppt) and polyhaline (18 to 30 ppt) salinity regimes based on classification by Cowardin et al. (1979). Two marsh sites were chosen in each regime based on observed similarity to other marshes in the area and on accessibility for sampling. The salinity regimes were characterized using Texas Parks and Wildlife Department (TPWD) records taken over the past 10 years within 1 km of each site, as well as from salinities measured in 1987 during the study. Marshes were




FIGURE 1. Map of sites.

compared to open water habitats in the adjacent bay throughout the study. The open water habitats were either nonvegetated (barren) mud or sand bottom, or submerged aquatic vegetation (SAV), such as seagrasses, or both.

In the upper bay, two marsh sites (Sites 1 and 2) were studied on the Trinity River delta located at 94° 42' W, 29° 44' 36" N and at 94° 43' 18" W, 29° 45' 30" N (Fig.1). The marshes had mixed emergent vegetation but the dominant plant near the marsh edge was *Scirpus* spp.. Submerged aquatic vegetation (SAV) was present at both sites during the summer and was mostly comprised of *Ruppia maritima*, *Najas* sp. and *Vallisneria americana*. Both marshes were situated along coves that opened into Trinity Bay. The site closest to the bay near the navigation channel was designated the outer site (Site 2; OTD, outer Trinity Delta) and the inland delta site, near southwest pass, was designated the inner site (Site 1; ITD, inner Trinity Delta). Ten year monthly mean salinities from TPWD ranged from 3.0 to 18.9 ppt with an overall mean of 9.2 ppt at the outer site. Mean monthly salinities at the inner marsh site ranged from 1.7 to 14.4 ppt at the inner site with an overall mean of 6.0 ppt. Because of the low salinity occurrences, the inner site was designated as oligohaline. The dominance at the inner site of *Najas* and *Vallisneria*, plants which do not tolerate long-term salinities above 6 ppt, confirm the validity of the classification. Because of its slightly higher salinities, the outer site was classified as a transition from oligohaline to mesohaline.

In the middle of the bay, mesohaline marshes were selected at Smith Point (Site 3; SP) and at Moses Lake (Site 4; ML) at 94° 45' 24" W, 29° 33' 18" N and 94° 55' 30" W, 29° 26' 24" N, respectively. At Smith Point, the marsh was mostly composed of *Spartina alterniflora* with *Juncus roemerianus* and *Spartina*

cynosuroides mixed in. At Moses Lake, the marsh was *Spartina alterniflora*, *Juncus roemerianus* and *Distichlis spicata*. There was no SAV in the area; open water bottoms adjacent to the marsh varied from hard clay and soft mud to muddy sand with broken *Rangia* shell. The ten year mean of salinities was 11.7 ppt for Smith Point and 15.7 ppt for Moses Lake.

In the lower bay, polyhaline marsh sites were selected in West Bay, at the Galveston Island State Park, (Site 5;WB) and in Christmas Bay (Site 6;CB). They were located at 94° 59' W, 29° 12' N and 95° 10' W, 29° 2' 48" N, respectively. These marshes were composed of monotypic stands of *Spartina alterniflora* with some *Salicornia virginica* and *Batis maritima* at higher elevations. The subtidal bottom next to the marsh in West Bay was sandy mud without SAV habitat present; but at Christmas Bay the bottom was sandy and SAV habitat was present. The stand of SAV was mostly *Halodule wrightii* with traces of *Ruppia maritima*, *Halophila engelmannii* and *Thalassia testudinum*. Ten year mean salinities from TPWD were 23.8 ppt in West Bay and 26.4 ppt in Christmas Bay.

Field Procedures:

The principal method of sampling animal abundances on the marsh surface and in nearby shallow-water subtidal habitats was drop trap sampling (Fig.2). Drop trap sampling was developed to compare animal densities among a variety of shallow-water habitats. The method employs a large cylinder (1.8 m dia.) dropped from a boom on a boat to entrap organisms within a prescribed area (2.6 m²). Most of the mobile fauna are captured by using dip nets while the water is pumped out of the sampler. When the sampler is completely drained, animals remaining on the bottom are picked up by hand. The technique is designed to sample fishes, crabs and shrimps in marshes, seagrass beds and




FIGURE 2. Drawing of the drop trap sampler.

oyster reefs where methods such as trawls and seines are ineffective. The technique improves on conventional methods because catch efficiency is very high (85 to 100 %) and measurements approach actual densities (numbers/unit area) of target organisms; hence, with drop trap sampling, quantitative comparisons of organism abundances within and between marshes and among a variety of other habitats are possible. The technique has been used in water depths up to 1.1 meter in marshes, SAV beds, mangroves, oyster reefs, and bare sand and mud bottoms. The methodology was described by Zimmerman et al. (1984).

In Galveston Bay, drop trap sampling was employed to assess utilization of intertidal marshes and subtidal bottoms by fishery species along a salinity gradient. Four replicate samples (2.6 m² each) of each habitat type at each site were taken during the spring, summer and fall seasons of 1987. Sampling always included marsh and bare mud bottom habitats (subtidal open-water adjacent to the marsh edge) in sample pairs (4 replicate pairs per site) and SAV habitat (4 additional replicates) when present. Thus, without SAV, 8 samples of marsh and 8 samples of adjacent mud bottom were taken in each the upper, middle and lower system (48 total) during April, July and November (144 overall total). This balanced set of replicates among habitats and season constituted the basis for our main comparisons. Since SAV was only at the Trinity delta and Christmas Bay sites and was seasonally present, this habitat was compared between sites and with other habitats separately. The main observations from drop trap samples were fish and decapod crustacean densities. The organisms were collected in the field and preserved with 10% Formalin, then taken to the laboratory for identification, measurement and enumeration.

Other observations included densities

of infauna and epifauna, vegetation type and biomass, and measurements of water depth, temperature, salinity, dissolved oxygen and turbidity. Infauna and epifauna were sieved from a single 10 cm dia. x 5 cm sediment core taken within each drop trap. These small macrofauna were retained on a 500 micron square mesh screen, then placed in zip-lock plastic bags with 10% Formalin with Rose Bengal stain, and stored for sorting at the laboratory. All emergent plants in marsh samples were cut and placed in plastic garbage bags, without preservation, for laboratory processing. Maximum and minimum water depth was measured in each drop trap with a meter rule. Water temperature and dissolved oxygen was measured using a YSI Model 51B meter, and salinity was measured using an American Optical refractometer. Water samples (500 cm²) were taken to measure turbidity (HR Instruments Model DRT 15) and to check conductivity/salinity with a Hydrolab Data Sonde at the laboratory.

Laboratory Procedures:

In the laboratory, fishes and crustaceans were sorted to species (using identifications based on guides, keys and taxonomic papers listed in Appendix I). Fish were measured to nearest mm total length and counted in groups of 10 mm size intervals (1 to 10 mm, 11 to 20 mm, etc.). Decapod crustaceans were measured to nearest mm total length for shrimps and carapace width for crabs and counted in groups of 5 mm size intervals (1 to 5 mm, 6 to 10 mm, etc.). The data were recorded on printed forms and entered in DBASE III Plus files using a microcomputer. Infauna and epifauna were processed similarly except they were not measured, and individuals were identified to species only in 1 of each 4 replicates; in the other 3 replicates, they were counted as peracarid crustaceans, annelid worms, mollusks or other fauna. Marsh plants were first weighed wet, then air dried for two months and weighed dry. After drying,

the number of culms in each sample were counted to calculate density, then discarded. All faunal samples were stored in 5% Formalin (with seawater) or 70% ETOH for reference. These will be kept for at least 3 years from the date of collection. All field sheets and data entry forms are on file and will be kept for at least 5 years.

Analytical Procedures:

We used analysis of variance (ANOVA) to test for significance of observations among habitats, areas of the bay, and seasons. In the main design, marsh and nonvegetated habitats were considered subsamples since they were always sampled together. Sites were combined within upper, middle, and lower areas of the bay to test for effect of location. Seasons were the spring, summer, and fall. Data were transformed using $\log x + 1$ since variances were usually proportional to the means (see means and standard errors in Appendices II through V). Differences between observation means were tested at the 0.05 significance level. The main observations were densities of selected faunal groups and taxa, including all fishes, all decapods, game fishes, bait fishes, penaeid shrimp, economically important and most abundant species. The game fish were comprised of southern flounder, spotted seatrout and red drum. Bait fish were bay anchovy, pinfish, and striped mullet. Economically important decapods, analyzed as individual species, were brown shrimp, white shrimp, pink shrimp, and blue crab. Other observations included physical parameters, densities of forage organisms (annelid worms and peracarid crustaceans) and vegetational parameters. SAV, marsh, and nonvegetated habitats were compared only between the two sites where SAV was always present (Christmas Bay and the Inner Trinity Delta). Because most species were transient and highly seasonal, occurrences or high abundances within species were often confined to one or two seasons.

This weakened our justification for testing across all seasons (including seasons as a level in the ANOVA design) in all taxa. It also increased interaction of season with habitat and bay area. Therefore, many tests at the family or species level were limited to within seasons. In all ANOVAs, where probabilities were equal to or greater than 0.05, and interactions were not significant, we used LSD multiple range tests to identify differences. In some cases where season, area of the bay and habitat interacted significantly, we used paired t-tests to independently analyze for difference between habitats. We also analyzed for differences in selection of marsh versus nonvegetated habitat between sites, using percent abundance in the marsh (calculated from animal densities in pairs of samples of marsh and nonvegetated bottom) as the observation. The observations in this case were arcsine transformed. All analyses were executed on a micro-computer using SAS/STAT programs. The untransformed means and standard errors of species densities were calculated by season/site/habitat and are tabulated in Appendices II through V.

Total abundances within species were tabulated for each site. Since sites were located within characteristic salinity regimes, abundances within species at each site roughly corresponded to relationship with salinity. Total from marsh and nonvegetated habitats were combined, but SAV was not included since it did not occur at all sites. The distribution center was used to characterize the most closely associated salinity regime for each species. Salinity regimes at each site were calculated as 1987 mean (our data, taken during drop trap sampling) and as the ten year historical mean (data from random sampling by the Texas Parks and Wildlife Department within 1 km of each site). The sites and their corresponding salinities (1987 and historical, respectively) were: Site 1 - Inner Trinity Delta (3.6 and 6.0 ppt), Site 2 - Outer Trinity Delta (3.4 and 9.2 ppt), Site 3 - Smith Point (9.8 and 11.7 ppt), Site 4 - Moses Lake (15.5 and 15.7

ppt), Site 5 - West Bay (27.2 and 23.8 ppt) and Site 6 - Christmas Bay (27.9 and 26.4 ppt).

represents mean salinities in different parts of the bay.

RESULTS

Physical Parameters

Salinity: Salinities in Galveston Bay during the 1987 survey are graphically compared to 10 year TPWD averages in Figure 3, with means and standard errors are given in Appendix II. The unequal sample sizes among sites for the TPWD 10 year historical database should be noted. The 10 year (historical) mean at the Trinity Delta inner site (Site 1) is based on 26 measurements and 25 measurements at the outer site (Site 2). Since the measurements are few and they were taken randomly over time, not all the monthly means are available (June is missing for the inner site and February is missing for the outer site). The remaining sites, in the middle and lower bay, were based on more observations; eg., Smith Point (Site 3), 125; Moses Lakes (Site 4), 241; West Bay (Site 5), 156; Christmas Bay (Site 6), 87. Withstanding some imprecision for the upper bay, the TPWD record

The salinity gradient was evident both in 1987 and historically (Fig. 3). The salinity values classify the bay into oligohaline, mesohaline and polyhaline environments that correspond to the upper, middle and lower divisions of the bay (Fig. 4). Seasonal differences are evident with steeper gradients occurring in the spring and summer due to lower salinities in the upper bay and higher salinities in the lower system (Fig. 3). During the fall, salinities reach their annual peak in the upper system, thus reducing the slope of the gradient. These seasonal variations in salinity impose short-term influences on the environment. During 1987, in particular, short term influence of lowering of salinity was observed in the middle bay. In Table 1, the short-term differences in salinity are depicted between sites seasonally and at the same time the overall integrity of the gradient is demonstrated. There was no difference in salinity between marsh and open water habitats (paired-t tests within sites and seasons, $n = 4$, $P > 0.05$).




FIGURE 3. Salinities in marsh and adjacent nonvegetated open water at sites in Galveston Bay during a drop trap sampling survey in 1987, and TPWD sampling within 1 km of each site between 1977-87.




Figure 4. Salinity regimes in Galveston Bay.

TABLE 1. Mean salinities (ppt salinity) in upper, middle, and lower Galveston Bay during 1987. Underline denotes no significant difference between values (ANOVA, df = 5, P > 0.05; LSD, df = 42).

SEASON	UPPER BAY		MIDDLE BAY		LOWER BAY	
Spring (April-May)	Site 2 <u>0.01</u>	Site 1 <u>0.02</u>	Site 3 <u>8.5</u>	Site 4 <u>15.5</u>	Site 6 <u>22.1</u>	Site 5 <u>33.3</u>
Summer (July)	Site 1 <u>0.0</u>	Site 2 <u>0.5</u>	Site 3 <u>0.8</u>	Site 4 <u>9.0</u>	Site 5 <u>27.8</u>	Site 6 <u>29.4</u>
Fall (November)	Site 2 <u>9.6</u>	Site 1 <u>10.8</u>	Site 3 <u>20.0</u>	Site 5 <u>20.5</u>	Site 4 <u>22.1</u>	Site 6 <u>32.1</u>

Sites: Site 1 = Inner Trinity Delta; Site 2 = Outer Trinity Delta; Site 3 = Smith Point; Site 4 = Moses Lake; Site 5 = West Bay; Site 6 = Christmas Bay (see Fig. 1). For dates and time of day refer to Appendix II.

Water Depths: Mean water depth at all sites was always less than 1 m and was always deeper in open water (near the edge of the marsh) than on the marsh surface (Appendix II). However, due to variability in water depths (they were changing with the tides) during sampling, differences in depth were not always significant between habitats (paired-t tests within sites, n = 4, P > 0.05). For the same reason, differences between marsh and open water depths among sites were usually not significantly different (ANOVAs, df = 5, P > 0.05; Table 2).

Other Parameters: Water temperature, dissolved oxygen and water turbidity values rarely differed between habitats within sites (paired t-tests within sites, n = 4, P > 0.05), but often differed between sites (Table 3; ANOVA, df = 5, P < 0.05; LSD multiple range test, df = 42). However, gradient-related patterns in temperature and dissolved oxygen were not apparent. A weak pattern of higher turbidities at upper bay sites and lower turbidities at lower bay sites was evident. Mean temperatures were lowest during the fall sampling (18.8 to 25.2° C) and highest during the summer sampling (27.6 to 32.0° C). Dissolved oxygen was lowest during fall sampling (4.0 to 9.4 ppm) and highest during spring sampling (7.0 to 12.4 ppm). Turbidities were generally lower during the spring sam-

pling (13.4 to 44.3 ppm) and highest during fall sampling (22.0 to 89.5 ppm).

Demersal Organisms

All Fishes: During 1987, 49 species of fishes among 2030 individuals were captured in 144 drop trap samples (2.6 m² each) from marsh and adjacent nonvegetated open water habitats in Galveston Bay (Appendix III). The number of fishes from marshes was 1,410 (7.5/m²) compared to 620 (3.3/m²) from nonvegetated open water. Abundances were significantly higher in marshes across all areas of the bay in all seasons (ANOVA, df = 108, P < 0.05). Densities were significantly different between seasons with lowest densities in the spring and highest in the fall (Fig. 5). Over all seasons, sites in the middle bay had significantly higher fish densities than the upper or lower bay. Within seasons, spring densities were not significantly different between sites in either habitat, summer densities were significantly different between sites in both marsh and open water, and fall densities were significantly different between sites only in the marsh (ANOVA, df = 5, P < 0.05). The main pattern, mostly due to summer and fall occurrences, was one of higher abundances at the middle bay sites (Smith Point and Moses Lake)(Fig. 5).

TABLE 2. Difference in water depth (cm difference between habitats) between marsh and adjacent subtidal nonvegetated habitats at sites in upper, middle and lower Galveston Bay during 1987. Values are means marsh depths minus adjacent open water depth from 4 pairs of samples at each site during flood tide. Underline denotes no significant difference among values (ANOVA, df = 5, P > 0.05; LSD multiple range test, df = 42).

SEASON	SITES					
Spring (April-May)	Site 1 <u>7.1</u>	Site 4 10.2	Site 3 <u>18.0</u>	Site 5 <u>18.5</u>	Site 1 30.6	Site 6 <u>51.0</u>
Summer (July)	Site 2 <u>9.4</u>	Site 1 10.9	Site 5 <u>19.9</u>	Site 4 22.0	Site 5 <u>24.4</u>	Site 3 <u>33.5</u>
Fall (November)	Site 5 <u>4.5</u>	Site 4 12.0	Site 1 <u>16.5</u>	Site 3 19.1	Site 5 <u>23.9</u>	Site 2 <u>30.1</u>

Sites are identified in Table 1. For exact dates and time of day refer to Appendix II.

TABLE 3. Means of temperature, dissolved oxygen and turbidity at sites along an environmental gradient in Galveston Bay during drop sampling in 1987. Mean value at each site is from combined measurements in marsh and open water (n = 8).

PARAMETER SEASON	SITES					
TEMPERATURE (°C)						
Spring (April-May)	Site 6 <u>23.7</u>	Site 1 <u>28.0</u>	Site 2 <u>28.6</u>	Site 5 <u>28.8</u>	Site 4 <u>29.5</u>	Site 3 <u>30.5</u>
Summer (July)	Site 4 <u>27.6</u>	Site 2 <u>30.6</u>	Site 6 30.7	Site 1 31.2	Site 3 <u>31.4</u>	Site 5 <u>32.0</u>
Fall (November)	Site 1 <u>18.8</u>	Site 5 <u>20.9</u>	Site 4 <u>22.4</u>	Site 2 <u>22.7</u>	Site 3 <u>22.9</u>	Site 6 <u>25.2</u>
DISSOLVED OXYGEN (ppm)						
Spring (April-May)	Site 6 <u>7.0</u>	Site 5 <u>7.6</u>	Site 1 <u>8.3</u>	Site 2 9.5	Site 3 <u>11.7</u>	Site 4 <u>12.4</u>
Summer (July)	Site 6 <u>6.0</u>	Site 5 <u>7.0</u>	Site 4 7.1	Site 1 <u>7.4</u>	Site 3 <u>8.2</u>	Site 2 <u>9.4</u>
Fall (November)	Site 1 <u>4.0</u>	Site 2 <u>7.9</u>	Site 5 7.9	Site 4 8.0	Site 3 <u>8.4</u>	Site 6 <u>9.4</u>
TURBIDITY (FTUs)						
Spring (April-May)	Site 5 <u>13.4</u>	Site 6 14.6	Site 3 <u>17.0</u>	Site 4 <u>29.3</u>	Site 2 <u>33.1</u>	Site 1 <u>44.3</u>
Summer (July)	Site 5 <u>10.3</u>	Site 4 <u>26.8</u>	Site 3 <u>30.5</u>	Site 2 <u>30.9</u>	Site 5 <u>32.0</u>	Site 1 <u>46.4</u>
Fall (November)	Site 6 <u>22.0</u>	Site 5 24.4	Site 1 <u>50.8</u>	Site 2 <u>51.5</u>	Site 3 <u>70.9</u>	Site 4 <u>89.5</u>

Sites: (Identified in Table 1). Underline denotes no significant difference among values (ANOVA, df = 5, P > 0.05; LSD multiple range test, df = 42).




FIGURE 5. Densities of fishes in marsh and adjacent nonvegetated habitats at sites along a salinity gradient in Galveston Bay.

Game Fishes: Spotted seatrout, southern flounder, and red drum followed occurred in a pattern similar to that of all fishes, but abundances were not significantly different between habitats, areas of the bay and seasons (ANOVA, $df = 108$, $P > 0.05$). Nevertheless, peak abundances occurred in the summer and fall, and at middle bay sites (Fig.6). Within habitats, peak densities were in marsh habitat at the outer Trinity Delta and Smith Point, and in open water at Smith Point, Christmas Bay and West Bay.

All Decapod Crustaceans: During 1987, 18,051 decapod crustaceans among 28 species were caught in 144 drop trap samples from marsh and nonvegetated open water (subtidal) habitats in Galveston Bay (Appendix III). Of these, 16,914 individuals ($90/m^2$) were on marsh surface and 1,137 ($6.1/m^2$) were on nonvegetated bottom. Like fishes, decapod abundances were significantly higher in marshes than open water across all areas of the bay in all seasons (ANOVA, $df = 108$, $P > 0.05$). The pattern was one of highest abundances at the middle bay sites (Smith Point and Moses Lake) and lowest abundances at the two upper bay sites (Trinity Delta) (Fig. 7). Lowest densities occurred in the spring and highest densities occurred in the summer and fall (Fig. 7). Densities were

significantly different among sites within both habitats within all seasons (ANOVAs, $df = 5$, $P < 0.05$).

All Penaeid Shrimps: Shrimp densities were significantly higher in marshes than open water across all areas of the bay in all seasons (ANOVA, $df = 108$, $P < 0.05$). The middle and lower bay did not differ in abundances, but the upper bay was significantly lower. Spring and fall densities of penaeid shrimps were highest at lower bay sites (West Bay and Christmas Bay) declining toward the upper bay (Fig. 8). Summer densities were highest in the middle bay (Smith Point), declined sharply in the upper bay, and were intermediate in the lower bay. The overall pattern indicates highest abundances in the lower bay and lowest abundances in the upper bay. Moreover, the lower bay sites (West Bay and Christmas Bay) were the only sites where densities were always significantly higher in the marsh as compared to nonvegetated open water (paired t-tests, $n = 4$, $P < 0.05$).

Brown Shrimp: Spring and summer densities of brown shrimp were highest, and fall densities were lowest (Fig. 9). Densities were usually greater in the marsh than in nonvegetated open water. Densities were significantly different among areas of the bay





FIGURE 6. Densities of game fishes in marsh and adjacent nonvegetated habitats at sites along a salinity gradient in Galveston Bay.





FIGURE 7. Densities of decapod crustaceans in marsh and adjacent nonvegetated habitats at sites along a salinity gradient in Galveston Bay.




FIGURE 8. Densities of all penaeid shrimps in marsh and adjacent nonvegetated habitats at all sites along a salinity gradient in Galveston Bay.




FIGURE 9. Densities of brown shrimp (*Penaeus aztecus*) in marsh and adjacent nonvegetated habitats at sites along a salinity gradient in Galveston Bay.

and habitats but not between seasons (ANOVA, $df = 108$, $P < 0.05$). Densities were significantly higher in the lower bay than the middle or upper bay. Accordingly, brown shrimp were mostly in the lower bay (West Bay and Christmas Bay) during the spring, and in the middle bay and lower bay during the summer and the fall (Smith Point and West Bay) (Fig. 9). Brown shrimp were absent from upper bay Trinity Delta sites during the spring (the period of peak seasonal abundance) and only a few were present at these sites during the summer and fall. Within marsh habitat highest abundances were also in the lower bay (Fig. 9).

White Shrimp: White shrimp were not present during the spring. Peak annual densities occurred in the summer the middle bay (Smith Point), and highest fall densities occurred in the lower bay (Christmas Bay) (Fig. 10). Densities were significantly different between seasons and areas of the bay, although the lower and middle bay did not differ (ANOVA, $df = 108$, $P < 0.05$). Like brown shrimp, abundances of white shrimp were sharply (significantly) reduced in the upper bay. Mean densities in the marsh were often much higher than in nonvegetated open water (Fig. 10), but differences were not significant. This occurred because of aggregation behavior (clumping) in white shrimp.

Pink Shrimp: Pink shrimp were only present during the summer and fall, and peak annual densities occurred in the fall (Fig. 11). In the summer pink shrimp were only in the lower bay, but in the fall they occurred throughout the system. The highest fall densities were in the middle and lower bay (Moses Lake, West Bay, and Christmas Bay). Densities were always greater in the marsh than in nonvegetated open water, but significant interaction occurred between habitat and season (ANOVA, $df = 108$, $P < 0.05$) primarily because of low densities in the summer. For the same reason, significant interaction occurred

between area and season. Analysis of the fall season alone revealed significant differences between habitats and areas of the bay (ANOVA, $df = 18$, $P < 0.05$).

Blue Crab: Blue crabs were distributed throughout the Galveston Bay system in all seasons. Densities were lowest in the spring and highest in the fall (Fig. 12) and significantly different among seasons (ANOVA, $df = 108$, $P < 0.05$). The overall pattern in marsh habitat indicated highest abundances in the middle bay, intermediate abundances in the lower bay, and lowest abundances in the upper bay (Fig. 12). Densities of open water were approximately equivalent throughout the bay, except during the fall when densities were higher in the middle bay (Fig. 12). However, significant interaction occurred between area and habitat. This was primarily due to habitat selection differences between different parts of the bay. Blue crabs were always more abundant in marsh in the lower and middle bay, but in the upper bay densities were often higher in open water. For instance, during the spring, crabs were significantly higher in open water at the inner Trinity Delta site, significantly higher in marsh at the Smith Point site, and not different between habitats at any of the other sites (paired t-tests within sites, $n = 4$, $P > 0.05$).

Grass Shrimp: Grass shrimp occurred in all seasons as the most abundant decapod crustacean in marsh habitat. Densities peaked during the summer and fall, in the middle bay (Smith Point and Moses Lake) (Fig. 13). Densities were consistently higher in marsh compared to nonvegetated open water, but significant interactions occurred between habitat and season, and between habitat and area of the bay (ANOVA, $df = 108$, $P < 0.05$). The interaction effect was due to the extremely low numbers, approaching zero, of nearly all the nonvegetated habitat samples (Fig. 13; Appendix II).





FIGURE 10. Densities of white shrimp (*Penaeus setiferus*) in marsh and adjacent nonvegetated habitats at sites along a salinity gradient in Galveston Bay.





FIGURE 11. Densities of pink shrimp (*Penaeus duorarum*) in marsh and adjacent nonvegetated habitats at sites along a salinity gradient in Galveston Bay.





FIGURE 12. Densities of blue crab (*Callinectes sapidus*) in marsh and adjacent nonvegetated habitats at sites along a salinity gradient in Galveston Bay.





FIGURE 13. Densities of grass shrimp (*Palaemonetes pugio*) in marsh and adjacent nonvegetated habitats at sites along a salinity gradient in Galveston Bay.

Forage Animals

All Epifauna and Infauna: All macrofauna taken from sediment cores (10 cm dia.; 78.5 cm² each) were considered to be potential forage organisms (prey) for demersal fishes and decapod crustaceans. In order of abundance, the main taxa included annelid worms, peracarid crustaceans (mostly amphipods and tanaidaceans), and small mollusks. Densities of forage taxa were highest in the middle and lower bay (particularly, Moses Lake and West Bay) during the spring, and highest in the upper bay during the summer (Fig. 14). Marsh always had higher forage densities, but means were not significantly different from open water (ANOVA, $df = 108$, $P > 0.05$). Densities were highly dependent on variations in abundances of annelid worms and peracarid crustaceans.

Annelid Worms: Infaunal annelid worms (polychaeta and oligochaeta) were the most abundant group among the forage taxa (Appendix IV). Densities of annelids were highest during the spring in the middle bay, and during the summer and fall in the upper

bay (Fig.15). Middle bay abundances declined from spring to summer, but the upper bay abundances increased. Densities were not significantly different among seasons, habitats, or areas of the bay (ANOVAs, $df = 108$, $P > 0.05$). Overall, however, highest abundances occurred in the upper bay (Fig. 15).

Peracarid crustaceans: Peracarideans (amphipods and tanaids) were second in abundance to annelid worms as forage animals (Appendix IV). Like annelids, seasonal densities were highest during the spring declining to lowest levels the fall (Fig. 16). In contrast to annelids, peracarids were virtually absent from the upper system in all seasons. Also, in the middle system peracarid abundances were comparatively high (Fig. 16). Overall, densities were significantly different among seasons and areas of the bay (ANOVA, $df = 108$, $P > 0.05$), but not between habitats.

Overall Distributions

Among 47 species of fishes, 8 species were mostly in the upper bay (Sites 1 and 2),




FIGURE 14. Densities of forage taxa for small fishes and decapod crustaceans in marsh and adjacent nonvegetated habitats at sites along a salinity gradient in Galveston Bay.





FIGURE 15. Densities of annelid worms in marsh and adjacent nonvegetated habitats at sites along a salinity gradient in Galveston Bay.





FIGURE 16. Densities of peracarid crustaceans in marsh and adjacent nonvegetated habitats at sites along a salinity gradient in Galveston Bay.

24 species were mostly in the middle bay (Sites 3 and 4) and 15 species were mostly in the lower system (Sites 5 and 6) (Table 4). Overall abundances of fishes were highest in the middle bay. Of 2030 individuals, 394 (19.4 %) were in the upper bay, 1168 (57.5 %) were in the middle bay and 468 (23.0 %) were in the lower system (Table 4).

Among 28 species of decapod crustaceans, 1 species was mostly in the upper bay (Sites 1 and 2), 14 species were mostly in the middle bay (Sites 3 and 4), and 13 were mostly in the lower bay (Sites 4 and 5) (Table 5). Of 18,051 individuals, 756 (4.2 %) were in the upper bay, 12433 (68.9 %) were in the middle bay, and 4862 (26.9 %) were in the lower bay (Table 5).

The abundance centers for each of the eleven fishery species, as related to 1987 and historical salinities, respectively, were:

- 1) common croaker - 6.6 and 10.4 ppt
- 2) red drum - 10.6 and 12.2 ppt
- 3) spotted seatrout - 15.1 and 15.4 ppt
- 4) blue crab - 15.5 and 15.7 ppt
- 5) white shrimp - 16.1 and 16.1 ppt
- 6) southern flounder - 18.1 and 17.5 ppt
- 7) menhaden - 20.2 and 19.0 ppt
- 8) pink shrimp - 20.6 and 19.3 ppt
- 9) brown shrimp - 23.2 and 21.1 ppt
- 10) sheepshead - 27.2 and 23.8 ppt
- 11) stone crab - 27.2 and 23.8 ppt

The most common salinity regimes for fishery species during 1987 ranged from mesohaline (6.6 ppt) to polyhaline (27.2 ppt); moreover, of the 11 fishery species, 6 were mesohaline and 5 were polyhaline.

Among 42 forage species, 6 species were mostly in the upper bay (Sites 1 and 2), 19 species were mostly in the middle bay (Sites 3 and 4) and 17 species were mostly in the lower bay (Sites 5 and 6) (Table 6). Of 33,897 individuals, 8,356 (24.7 %) were in

upper bay, 18,260 (53.9 %) were in the middle bay and 7,281 (21.5 %) were in the lower bay (Table 6).

Effect of SAV Habitat

SAV habitat occurred only at Trinity Delta and in Christmas Bay. In both areas, SAV was in the low intertidal zone, exposed only during extremely low winter tides, adjacent to marsh. Bayward was subtidal non-vegetated sand. Animal densities within Trinity Delta and Christmas Bay sites were usually not different between marsh and SAV habitats. But, marsh and SAV habitats at Christmas Bay nearly always had higher animal densities than those at the Trinity Delta (Figs. 17 through 23). The outer Trinity Delta site had some, albeit sparse, SAV year-around, while the inner site had SAV only during the summer.

Highest fish densities occurred in the SAV habitats, at Christmas Bay during the spring and fall, and at the Trinity Delta outer site during the summer (Figure 17). In the spring, fish densities were significantly higher in SAV at Christmas Bay than in any other habitat, including those of the outer Trinity Delta (ANOVA, $df = 18$, $P < 0.05$). During the summer and fall, marsh and SAV fish densities did not differ. Game fishes were consistently more abundant in Christmas Bay, but as a group did not differ among sites in densities between marsh, SAV or nonvegetated habitats (Fig. 18).

Decapod crustacean densities were significantly higher in Christmas Bay marsh and/or SAV habitats (ANOVA, $df = 18$, $P < 0.05$), than habitats at the Trinity Delta (Figure 19). Moreover, decapod densities did not differ significantly in Christmas Bay between marsh and SAV in any season (ANOVA, $df = 18$, $P > 0.05$). Penaeid shrimps, as a group, did not differ in density between marsh and SAV habitats, but densities between sites

TABLE 4. Total fishes by site and in relation to salinity in Galveston Bay.

GALVESTON BAY STUDY								
TOTAL ABUNDANCES OF FISHES								
ALL SEASONS AND HABITATS COMBINED								
2.6 m sq. Drop samples n = 24 per site SPE CIES	OLIGOHALINE		MESOHALINE		POLYHALINE		1987 SALINITY	HISTORICAL SALINITY
	SITE 1	SITE 2	SITE 3	SITE 4	SITE 5	SITE 6		
1 <i>Fundulus jenkinsi</i>	4	0	0	0	0	0	3.6	6.0
2 <i>Pomoxis annularis</i>	3	0	0	0	0	0	3.6	6.0
3 <i>Lucania parva</i>	14	0	1	1	0	0	3.5	7.0
4 <i>Fundulus pulvereus</i>	8	2	0	0	1	0	3.5	7.8
5 <i>Elops saurus</i>	2	0	2	0	0	0	3.4	9.2
6 <i>Ictalurus punctatus</i>	0	1	0	0	0	0	3.4	9.2
7 <i>Cyprinodon variegatus</i>	150	0	0	4	0	39	3.8	9.4
8 <i>Micropogonias undulatus</i>	14	3	5	2	4	2	6.6	10.4
9 <i>Fundulus grandis</i>	32	35	8	20	10	7	7.6	10.8
10 <i>Gambusia affinis</i>	1	0	0	0	1	0	9.8	11.7
11 <i>Gobiosox strumosus</i>	0	0	3	0	0	0	9.8	11.7
12 <i>Oligoplites saurus</i>	0	0	3	0	0	0	9.8	11.7
13 <i>Membras martinica</i>	0	0	2	0	0	0	9.8	11.7
14 <i>Syngnathus louisianae</i>	1	0	2	0	1	0	9.8	11.7
15 <i>Arius felis</i>	0	0	1	0	0	0	9.8	11.7
16 <i>Hyporhamphus unifasciatus</i>	0	0	1	0	0	0	9.8	11.7
17 <i>Stellifer lanceolatus</i>	0	0	1	0	0	0	9.8	11.7
18 <i>Mugil cephalus</i>	16	16	5	17	2	9	9.8	11.7
19 <i>Sciaenops ocellatus</i>	0	1	5	0	1	0	10.6	12.2
20 <i>Anchoa mitchilli</i>	7	37	129	69	24	1	11.3	12.7
21 <i>Myrophis punctatus</i>	13	2	8	41	3	3	12.1	13.3
22 <i>Citharichthys spilopterus</i>	2	0	1	0	0	2	12.1	13.3
23 <i>Symphurus plagiusa</i>	4	0	55	0	16	15	14.2	14.8
24 <i>Leiostomus xanthurus</i>	1	5	8	0	4	6	14.3	14.8
25 <i>Gobiosoma boscii</i>	0	1	165	483	29	1	14.4	14.9
26 <i>Cynoscion nebulosus</i>	0	2	11	4	9	2	15.1	15.4
27 <i>Gobiosoma robustum</i>	0	0	0	2	0	0	15.5	15.7
28 <i>Sphoeroides parvus</i>	0	0	0	2	0	0	15.5	15.7
29 <i>Menidia beryllina</i>	4	5	1	40	39	1	17.8	17.3
30 <i>Paralichthys lethostigma</i>	0	1	3	1	1	3	18.1	17.5
31 <i>Brevoortia patronus</i>	3	0	0	9	23	0	20.2	19.0
32 <i>Microgobius thalassinus</i>	0	0	1	0	4	0	22.5	20.6
33 <i>Opsanus beta</i>	0	0	0	1	2	0	23.3	21.1
34 <i>Syngnathus scovelli</i>	0	2	0	6	4	7	24.2	21.7
35 <i>Lagodon rhomboides</i>	1	0	20	9	17	43	25.9	22.9
36 <i>Archosargus probatocephala</i>	0	0	0	0	1	0	27.2	23.8
37 <i>Chaetodipterus faber</i>	0	0	0	0	1	0	27.2	23.8
38 <i>Fundulus similis</i>	0	0	0	0	2	0	27.2	23.8
39 <i>Gobionellus boleosoma</i>	0	1	15	0	5	95	27.6	25.2
40 <i>Adinia xenica</i>	0	0	0	1	0	4	27.6	25.4
41 <i>Achirus lineatus</i>	0	0	0	0	0	1	27.9	26.4
42 <i>Dasyatis sabina</i>	0	0	0	0	0	1	27.9	26.4
43 <i>Eucinostomus argenteus</i>	0	0	0	0	0	1	27.9	26.4
44 <i>Orthopristis chrysoptera</i>	0	0	0	0	0	1	27.9	26.4
45 <i>Synodus foetens</i>	0	0	0	0	0	1	27.9	26.4
46 <i>Trinectes maculatus</i>	0	0	0	0	0	1	27.9	26.4
47 <i>Eucinostomus spp.</i>	0	0	0	0	0	18	27.9	26.4
FISH TOTALS:	280	114	456	712	204	264		

TABLE 5. Total decapod crustaceans by site and in relation to salinity in Galveston Bay.

GALVESTON BAY STUDY
TOTAL ABUNDANCES OF DECAPOD CRUSTACEANS
ALL SEASONS AND HABITATS COMBINED

2.6 m sq. Drop samples n = 24 SPECIES	OLIGOHALINE		MESOHALINE		POLYHALINE		1987 SALINITY	HISTORICAL SALINITY
	SITE 1	SITE 2	SITE 3	SITE 4	SITE 5	SITE 6		
1 <i>Sesarma reticulatum</i>	2	0	0	0	1	0	5.5	10.0
2 <i>Uca pugnax</i>	2	0	7	0	0	0	7	10.6
3 <i>Xanthidae, unknown species</i>	0	0	4	0	0	0	9.8	11.7
4 <i>Eurypanopeus depressus</i>	0	0	3	0	0	0	9.8	11.7
5 <i>Neopanope texana</i>	1	4	31	2	1	3	10.8	12.4
6 <i>Rhithropanopeus harrisi</i>	0	1	32	4	0	2	11.1	12.6
7 <i>Palaemonetes pugio</i>	187	339	6276	2792	956	1708	14	14.6
8 <i>Palaemonetes vulgaris</i>	0	2	358	132	94	74	14.5	15.0
9 <i>Callinectes sapidus</i>	65	104	390	1243	333	200	15.3	15.6
10 <i>Uca rapax</i>	0	0	0	1	0	0	15.5	15.7
11 <i>Palaemonetes intermedius</i>	0	0	128	92	24	58	16	16.0
12 <i>Penaeus setiferus</i>	0	8	378	14	103	163	16.1	16.1
13 <i>Penaeus duorarum</i>	0	29	46	152	134	82	20.6	19.3
14 <i>Eurypanopeus abbreviatus</i>	0	0	0	2	2	0	21.4	19.8
15 <i>Panopeus herbstii</i>	0	0	2	0	0	2	21.4	19.8
16 <i>Penaeus aztecus</i>	2	10	248	94	478	259	23.2	21.1
17 <i>Libinia dubia</i>	0	0	0	0	1	0	27.2	23.8
18 <i>Pinnixa chaetoptera</i>	0	0	0	0	1	0	27.2	23.8
19 <i>Uca spp.</i>	0	0	0	0	1	0	27.2	23.8
20 <i>Menippe mercenaria</i>	0	0	0	1	0	1	27.2	23.8
21 <i>Sesarma cinereum</i>	0	0	0	0	3	0	27.2	23.8
22 <i>Petrolisthes armatus</i>	0	0	0	0	5	0	27.2	23.8
23 <i>Alpheus heterochaelis</i>	0	0	0	1	27	31	27.6	25.2
24 <i>Clibanarius vittatus</i>	0	0	0	0	40	58	27.6	25.4
25 <i>Pagurus spp.</i>	0	0	0	0	0	2	27.9	26.4
26 <i>Panopeus turgidus</i>	0	0	0	0	0	3	27.9	26.4
27 <i>Hippolyte zostericola</i>	0	0	0	0	0	5	27.9	26.4
28 <i>Uca minax</i>	0	0	0	0	0	7	27.9	26.4
CRUSTACEAN TOTALS:	259	497	7903	4530	2204	2658		

TABLE 6. Total epifauna and infauna by site and in relation to salinity in Galveston Bay.

GALVESTON BAY STUDY								
TOTAL ABUNDANCES OF EPI-IFAUNA								
ALL SEASONS AND HABITATS COMBINED	OLIGOHALINE		MESOHALINE		POLYHALINE		1987	HISTORICAL
78.5 cm sq. cores n = 6 per site	SAMPLING SITES						SALINITY	SALINITY
SPECIES	1	2	3	4	5	6		
ANNELIDS								
1 <i>Laeonereis culveri</i>	285	134	5	1	0	20	3.5	7.8
2 <i>Oligochaete spp.</i>	580	396	39	154	13	39	3.4	9.1
3 <i>Nereidae sp.</i>	0	1	0	0	0	0	3.4	9.2
4 <i>Parandalia fauveli</i>	0	0	1	0	0	0	9.8	11.7
5 <i>Hobsonia gunneri</i>	6	21	4	28	7	1	10.8	12.4
6 <i>Polydora ligni</i>	0	9	19	33	3	0	11.6	12.9
7 <i>Marphysa sanguinea</i>	0	0	0	1	0	0	15.5	15.7
8 <i>Steninerereis martini</i>	0	0	0	3	0	0	15.5	15.7
9 <i>Mediomastus spp.</i>	0	0	0	6	0	0	15.5	15.7
10 <i>Mediomastus ambiseta</i>	0	0	0	9	0	0	15.5	15.7
11 <i>Eteone lactea</i>	0	0	0	17	2	0	16.8	16.6
12 <i>Streblospio benedicti</i>	3	29	15	769	316	47	18.8	18
13 <i>Nereis (Neanthes) succinea</i>	0	0	4	1	2	4	21.9	20.2
14 <i>Capitella capitata</i>	0	0	30	49	81	72	25.3	22.5
15 <i>Asychis elongatus</i>	0	0	0	0	1	0	27.2	23.8
16 <i>Scolecopsis sp.</i>	0	0	0	0	1	0	27.2	23.8
17 <i>Glycera dibranchiata</i>	0	0	0	0	3	0	27.2	23.8
18 <i>Mediomastus californiensis</i>	0	0	0	0	3	1	27.4	24.5
19 <i>Tharyx setigera</i>	0	0	0	0	14	6	27.4	24.6
20 <i>Scoloplos fragilis</i>	0	0	0	0	1	3	27.7	25.8
21 <i>Heteromastis filiformis</i>	0	0	0	1	6	44	27.8	26.1
22 <i>Aricidea (Acmira) philibinae</i>	0	0	0	0	1	6	27.8	26.1
23 <i>Axiiothella mucosa</i>	0	0	0	0	0	1	27.9	26.4
24 <i>Capitellidae sp.</i>	0	0	0	0	0	1	27.9	26.4
25 <i>Melinna maculata</i>	0	0	0	0	0	1	27.9	26.4
ANNELID TOTALS: Identified (n = 6):	874	590	117	1072	454	246		
Not Identified (n = 24):	5074	2663	971	4800	2567	1923		
CRUSTACEANS								
1 <i>Corophium sp. B</i>	0	1	0	0	0	0	3.4	9.2
2 <i>Callinectes sapidus</i>	0	0	1	0	0	0	9.8	11.7
3 <i>Xanthidae sp.</i>	0	0	1	0	0	0	9.8	11.7
4 <i>Gammarus mucronatus</i>	0	0	24	2	3	0	11.4	15.7
5 <i>Hargeria rapax</i>	0	0	281	603	99	12	14.6	15.1
6 <i>Corophium sp.</i>	0	6	7	1065	0	0	15.4	15.6
7 <i>Grandiclerella bonneroides</i>	0	0	14	37	13	0	15.4	15.6
8 <i>Ampelisca abdita</i>	1	0	16	598	43	0	16	16
9 <i>Mysidopsis bahia</i>	0	0	0	0	3	0	27.2	23.8
10 <i>Edotea montosa</i>	0	0	0	0	4	0	27.2	23.8
CRUSTACEAN TOTALS: Identified (n = 6):	1	7	344	2305	165	12		
Not Identified (n = 24):	6	16	1174	10835	2315	211		
MOLLUSKS:								
1 <i>Amygdalum papyrium</i>	0	0	1	0	0	0	9.8	11.7
2 <i>Odostomia sp.</i>	0	0	0	1	0	0	15.5	15.7
3 <i>Tellina sp.</i>	0	0	1	0	0	1	21.4	19.8
4 <i>Mulinia lateralis</i>	0	0	0	0	2	0	27.2	27.2
5 <i>Acteocina canaliculata</i>	0	0	0	0	5	0	27.2	27.2
6 <i>Pandora (Clidophora) trilineata</i>	0	0	0	0	0	1	27.9	26.4
7 <i>Gastropod sp. A</i>	0	0	0	0	0	3	27.9	26.4
MOLLUSCAN TOTALS: Identified (n = 6):			2	1	7	5		
Not Identified (n = 24):	157	35	32	8	14	46		
OTHERS:								
1 <i>Odonata sp. A</i>	2	0	0	0	0	0	3.6	6
2 <i>Odonata sp. B</i>	1	0	0	0	0	0	3.6	6
3 <i>Chironomid sp.</i>	1	24	2	12	0	0	5.7	10.8
4 <i>Nemertean sp.</i>	0	0	0	2	0	0	15.5	15.7
OTHER TOTALS: Categorized (n = 6):	4	24	2	14				
Not Categorized (n = 24):	231	174	49	391	157	48		




FIGURE 17. Comparative densities of fishes in marsh, submerged aquatic vegetation (SAV), and nonvegetated open water between the upper (Trinity River delta) and lower (Christmas Bay) parts of the Galveston Bay system, during 1987.




FIGURE 18. Comparative densities of decapod crustaceans in marsh, submerged aquatic vegetation (SAV), and nonvegetated open water between the upper (Trinity River delta) and lower (Christmas Bay) parts of the Galveston Bay system, during 1987.




FIGURE 19. Comparative densities of game fishes in marsh, submerged aquatic vegetation (SAV), and nonvegetated open water between the upper (Trinity River delta) and lower (Christmas Bay) parts of the Galveston Bay system, during 1987.






FIGURE 20. Comparative densities of brown shrimp in marsh, submerged aquatic vegetation (SAV), and nonvegetated open water between the upper (Trinity River delta) and lower (Christmas Bay) parts of the Galveston Bay system, during 1987.

**WHITE SHRIMP
Spring**



Summer



Fall




FIGURE 21. Comparative densities of white shrimp in marsh, submerged aquatic vegetation (SAV), and nonvegetated open water between the upper (Trinity River delta) and lower (Christmas Bay) parts of the Galveston Bay system, during 1987.




FIGURE 22. Comparative densities of pink shrimp in marsh, submerged aquatic vegetation (SAV), and non-vegetated open water between the upper (Trinity River delta) and lower (Christmas Bay) parts of the Galveston Bay system, during 1987.




FIGURE 23. Comparative densities of blue crab in marsh, submerged aquatic vegetation (SAV), and non-vegetated open water between the upper (Trinity River delta) and lower (Christmas Bay) parts of the Galveston Bay system, during 1987.

were always significantly higher in Christmas Bay (ANOVA, $df = 18$, $P < 0.05$) (Fig. 20). This pattern was similarly repeated in brown shrimp (Fig. 21) and pink shrimp (Fig. 22). Blue crab did not differ between habitats except in the fall (ANOVA, $df = 18$, $P < 0.05$) (Fig. 23).

Characterization of Marshes

The Upper Bay: Marshes in the upper system (Trinity Bay) were dominated by the Trinity River and other streams flowing into the estuary. Overall salinities in 1987 were lower than historical averages in the upper system revealing a wet year (Fig. 4). Both the inner and outer marsh at the Trinity River Delta were strictly oligohaline during the spring and summer of 1987. By fall, salinities had increased to low mesohaline range. Responses of the marsh community reflected both the 1987 conditions and the general characteristics of the delta environment.

Plant cover was very sparse at the beginning of spring as a result of the previous winter die-back. Marsh bulrush (*Scirpus* spp.), the dominant plant, emerged in April along with the subdominants, arrowheads (*Sagittaria lancifolia*, *S. latifolia*), alligator weed (*Alternanthera philoxeroides*), pickerel weed (*Pontederia cordata*), water hyssop (*Bacopa monnieri*), and switchgrass (*Panicum* sp.). All were under heavy grazing pressure by nutria (personal observation). Grazing and tidal and floodwater export previous production left the intertidal zone virtually bare. Subtidal areas adjacent to the marsh were also barren. By July, the plants had recovered to near maximum annual biomass on the marsh surface as well as in subtidal areas. Bulrush cover in the marsh was dense and lush, and subtidal areas were covered with submerged aquatic vegetation (SAV) to a water depth of about 80 cm deep. The dominant SAV species at the inner site (Site 1) were quillwort (*Isoetes* sp.) and widgeongrass (*Ruppia maritima*) in shallow water, with naiads (*Najas* spp.) and tape-

grass in deeper water. Quillwort was only in very shallow water (less than 20 cm deep and often exposed) next to the marsh edge. It formed a dense short turf year-around, and at the outer site (Site 2) coverage was more extensive. Large beds of tapegrass were present in water 30 to 80 cm deep at the inner site but not at the outer site. Further examination revealed that tapegrass beds covered many hectares extending westward for at least 2 kilometers. Tapegrass beds appeared to be a seasonally persistent vegetational feature that has not been previously reported for the delta. Most of the vegetation experienced a die-back during the fall (September and October) that was associated with increased salinities; but, it was not known whether salinity caused the die-back. Almost all of the fall standing crop of plants was exported as detritus into Trinity Bay during the ensuing winter months.

Of 8 species of fishes in delta marshes, 4 were cyprinodontidae (killifishes), two were freshwater species (crappie and channel catfish) and one was an estuarine species of commercial and recreational value (Atlantic croaker) (Table 4). During the spring, these fishes were mostly found in open water (not much vegetation present), but in the summer and fall they shifted into marsh habitat. Hence, the movements of fishes between habitats corresponded to seasonal changes in plant cover. McIvor and Odum (1988) point out that such differences in selection for the marsh surface may be controlled by the differences in the quality of nearby subtidal habitat that fishes must use when the marsh is drained. Fishes that seek high quality subtidal bottom for food and protection at low tide simply move onto the nearest marsh surface at flood tide. The single estuarine fish of commercial value (Atlantic croaker) associated with Trinity delta marshes also has been reported in abundance under low salinity conditions (0 to 11 ppt) in upper Barataria Bay, Louisiana (Rogers and Herke 1987). This species was

apparently one of the few commercial species able to use oligohaline, nonvegetated bottom as a nursery habitat.

Only one decapod crustacean (a crab) was more abundant at the upper bay sites than other areas, although 10 of the 28 species in the bay used the upper bay at some time during the year (Table 5). All 3 penaeid shrimps and the blue crab used the delta marshes, but not in large numbers. Baldauf (1970) also noted, from monthly trawl surveys taken in 1967, 1968 and 1969, that brown shrimp, white shrimp and blue crab use the delta as a nursery. He concluded that brown shrimp abundances were less during years when Trinity River flow was high and that white shrimp abundances were not influenced by differences in annual river flow. His comparison of catches in open water deep channels with shallow water yielded fewer shrimp next to the the marsh. These data suggest that the delta marsh surface may not be as important as shrimp habitat as the deeper water in the upper bay. We might add that the nursery roles of marsh surface and open water appeared to reverse in importance from the upper to the lower bay. Therefore, direct utilization of the marsh surface became increasingly evident toward the lower system.

Small macroinvertebrates, useful as forage organisms, were comprised almost entirely of annelids worms at the delta during 1987. A nereid polychaete (*Laeonereis culveri*) and several unidentified oligochaete species were the dominant infauna (Table 6). Nereids and oligochaetes are reported detritivores (Tenore et al. 1977; Tenore 1977). Epifaunal peracarid crustaceans were essentially absent. Since peracarids are highly utilized and often are preferred (or more available) as prey by small fishes and decapod crustaceans, their absence may have affected the distributions of these predators. A least the absence of peracarids would have lessened the feeding value of delta marshes

for exploiting predators. We propose that the lack of peracarids was directly attributable to low salinities, since estuarine peracarids have poor ability to osmoregulate and cannot accommodate freshwater conditions for very long.

The Middle Bay: The marshes in the middle part of Galveston Bay were greatly influenced by mixing of freshwater from the upper system and seawater from the lower system. This was clearly demonstrated during 1987. Salinities in the middle bay (Smith Point and Moses Lake sites) varied more than any other part of the system, with values from near 0 ppt to above 20 ppt (Fig. 4). Seasonal values were similar to either those of the upper system or lower system depending upon circumstances; eg. spring salinities were mid-range (8 to 15 ppt); summer salinities were similar to the upper system (0.8 to 9 ppt) following several months of high freshwater inflow; fall salinities were like those of the lower system (20 to 22 ppt) following reduced freshwater inflow and high equinoxtides. Over the long term, the middle system was unquestionably mesohaline, despite short-term salinities that varied between oligohaline and polyhaline.

Marshes in the middle bay were mixed stands of smooth cordgrass (*Spartina alterniflora*), black rush (*Juncus roemerianus*), saltgrass (*Distichlis spicata*) and marsh hay (*Spartina patens*). Smooth cordgrass dominated the outer fringe (low zone). Subtidal SAV was not present, possibly due to the extreme variations in salinity. But the presence of expansive subtidal oyster reefs provided ample shell for periphytic green and bluegreen algal colonization. These small algae were dense enough at Smith Point to be seen during aerial surveillance and initially mistaken for SAV beds.

In the middle bay, fishes were more numerous (57.5 % of all individuals) and had

more species with higher abundances (24 of 47 species) than any other part of the bay (Table 4). Moreover, they were nearly as diverse as those in the lower bay (32 versus 34 of 47 species). The most abundant species included the most valuable of the commercial and recreational fishes in the bay, menhaden, spotted seatrout, southern flounder, and red drum, as well as, many fishes important in food chains (bait fishes), bay anchovy, spot, silversides and mullet. The salinity regimes of these species were 9.8 to 20.2 ppt in 1987 and from 11.7 to 19.0 ppt historically. This suggested that the bay area with mid-mesohaline to low polyhaline salinities was an optimal environment for fishes.

Decapod crustaceans were less diverse in the middle bay (17 versus 24 of 28 species) than in the lower system, but they were more numerous (68.9 % of all individuals) and had more of the most abundant species (14 of 28) (Table 5). Like fishes, the list of most abundant decapods in the middle bay included important commercial species, white shrimp, pink shrimp and blue crab, and food chain species, grass shrimps and xanthid crabs. The 1987 salinity regime of these species ranged from 9.8 to 21.4 ppt, and the historical salinity regime ranged from 11.7 to 19.8 ppt. Thus, optimal conditions for these decapod crustaceans of fishery value were mid-mesohaline to low polyhaline regimes.

Most of the forage species (25 of 42) occurred in the middle bay, and of these, 21 were more abundant in the middle bay than elsewhere (Table 6). Moreover, 53.9 % of all individuals occurred at the middle bay sites. Abundances of peracarid crustaceans were strikingly higher in the middle bay, and this association with high abundances fish and decapod predators strongly suggested a food chain connection. It has been well established that peracarids are a key component in the diets of many small estuarine fishes (Stoner 1982; Huh and Kitting 1985; Whitfield 1988).

Gut analyses of fishes from Galveston Bay (Sheridan 1983) and other Texas bays (Minello et al. 1987) support this observation. Furthermore, small juveniles of brown shrimp, pink shrimp, and blue crab have been shown to prefer amphipods and tanaids over other benthos (Leber 1979; Thomas 1989, Zimmerman et al).

The Lower Bay: Historical and 1987 salinity regimes in the lower bay marshes (West Bay and Christmas Bay) were polyhaline, with short term incursions of mesohaline to hypersaline conditions. Gulf water normally dominates through tides. But evaporation often produces a hypersaline environment during dry summers, and this condition can be alleviated or abruptly reversed by high rainfall caused by tropical depressions. In general, however, the lower bay was more saline and less variable than the middle and upper bay due to moderation from the Gulf.

Lower bay marshes were almost entirely smooth cordgrass in the lower zone which gradually changed to mixed stands of smooth cordgrass, glasswort (*Salicornia* spp.), and saltwort (*Batis maritima*) in the upper zone. A salt pan, without rooted vegetation but a bluegreen algal mat (Sage and Sullivan 1978; Pulich and Rabalais 1986), occurred between the marsh and terrestrial environment. Epiphytic algae on smooth cordgrass (Sullivan 1978, 1981) and macroalgae (Conover 1964; Williams-Cowper 1978) were more abundant in the lower bay than elsewhere. SAV occurred in Christmas Bay including, shoal grass (*Halodule wrightii*), widgeon grass (*Ruppia maritima*), turtle grass (*Thalassia testudinum*) and *Halophila engelmannii*. In West Bay, SAV beds were present as late as 1975, but have since disappeared.

A similar number of fish species occurred in the lower bay as compared to the middle bay (34 of 47 overall), but abundances

were lower (23.0 % of all individuals). A relatively low proportion of fish species occurring in the lower bay were most abundant there (12 of 34). Of commercial and recreational fishes, only the sheepshead (*Archosargus probatocephalus*) was more abundant in the lower bay (Table 4). Under most circumstances, the proportion of fully marine species could be expected to dominate as the salinities become increasingly euhaline, somewhere between about 20 and 35 ppt (Remane 1934). This was not evident, thus indicating the polyhaline nature of the the lower system.

Decapod crustaceans were most diverse in lower bay marshes (24 of 28 species), but with only 26.9 % of all individuals. Of the 24 species, 13 were more abundant than in the upper and middle bay. Among commercially important species that were most abundant in the lower bay were brown shrimp and stone crab (Table 5).

Of 42 forage species, 28 occurred in the lower bay among 21.5 % of all individuals, again indicating relatively higher diversity than the middle and upper bay. Of 28 species in the lower bay, only 12, a low proportion, were more abundant there than elsewhere. Peracarids were numerous but not as abundant as in the middle bay. Annelid worm abundances were intermediate to those of the other areas. The presence of algae and seagrasses provided additional food and structure, and less variable estuarine salinities afforded more stability to forage species in the lower system. In addition, smooth cordgrass remained in place throughout the year even though a die back occurred in the winter (dead stems remain erected for several years before they deteriorated). The grass culms provided a year-around surface for an epiphytic algal community. Both epiphytic algae and dead cordgrass are available as food and shelter for annelids, amphipods, tanaids and other organisms. This epiphytic community was well developed at

West Bay and, like barrier island salt marshes elsewhere, had significantly higher numbers of epifauna among grass culms than on the surrounding bottom (Rader 1984; Zimmerman et al.). This greatly increased the nursery value of lower bay marshes for foraging estuarine fishes, shrimps and crabs.

DISCUSSION

The Salinity Gradient in Galveston Bay

The salinity gradient is clearly apparent in the Galveston Bay system and reflects the dominating influence of freshwater inflow on characteristics of marsh communities in the system. During 1987, the salinity gradient was steeper than usual, as salinities were lower in the upper bay and higher in the lower bay than historical means (see Fig. 4). The gradient was steepest in the summer (July) when salinities in the upper bay and part of the middle bay approached zero. These low salinities are short-term phenomena that are within the range of annual variability; likewise, the higher salinities in marshes of the lower system during 1987 were short term events (within a season) that occur normally. Data from 1982 through 1988 from a salt marsh in West Bay (the Jamaica Beach site) reveal that short term conditions are often hypersaline in the late summer. Our record shows that during August at the Jamaica Beach marsh salinities were 38 ppt in 1982 (Zimmerman and Minello, 1984) and 41 ppt in 1985 (unpublished) over a period of several weeks. Because of this variability, the gradient within the Galveston Bay system can be expected to range, at least on the short term, from fresh (0 ppt) to hypersaline (40+ ppt). The historical means at the sites along the gradient perhaps best describe salinity regimes in the system. In Figure 3, we have compared different parts of the system using 1987 and historical salinity regimes. These are long term attributes of the environmental gradient. Both long term

(annual) and short term (seasonal) variations in salinity influence the responses of organisms.

Effect of Salinity on Organisms

Deviation in 1987 salinities from the historical means together with distributional responses of organisms provided insight into the short term versus long term effects of salinity. Under short term low salinity stress, the larger mobile fauna have the option to leave an area or to stay and accommodate. Less mobile organisms under the same circumstances, such as small epifauna, infauna and plants, cannot leave and thus must accommodate, at least temporarily, or suffer mortalities.

Many, if not most, estuarine species can temporarily accommodate oligohaline salinities below 5 ppt. Decapod crustaceans, such as brown shrimp, white shrimp and blue crab, are notable for their ability to accommodate low salinities (Zein-Elden 1989; Gifford 1962; Tagatz 1971). For example, we have observed responses of these and other estuarine species to abrupt lowering of salinities from mesohaline (7 to 15 ppt) to oligohaline (less than 1 ppt) during flooding of the Lavaca River delta in June of 1987. Freshwater flooding did not reduce densities of brown shrimp, white shrimp, grass shrimp, blue crabs in the delta marshes. Of fishes, bay anchovies and menhaden actually significantly increased their densities during the flooding. Similar results were observed in the middle of Galveston Bay in 1987 where faunal abundances were not depressed during short term lowering of salinities (a few days to several weeks, but less than a month) during the summer.

By contrast, species are known to suffer mortalities due to abrupt lowering of salinity (reviewed by Brongersma-Sanders 1957). In lower Texas bays mortalities occur when

populations acclimated to euhaline conditions (30 to 36 ppt) are exposed to rapid lowering of salinities due to rainfall from tropical depressions. Molluscan bivalves suffered mass mortalities in Redfish Bay after Hurricane Beulah in 1967 (Zimmerman and Chaney 1969). Salinities, in this instance, were reduced from 30 ppt to less than 1 ppt within about a week. Hedgpeth (1953) reported mortalities after a similar event in Nueces Bay. Low salinity limitations are known for many estuarine species. The restriction of oyster populations to salinities above 5 ppt (reviewed by Van Sickle et al. 1976) and their predator, the oyster drill, to salinities above 15 ppt (Gunter 1979) are well known examples. Even among euryhaline species, such as red drum, white shrimp and brown shrimp, low salinities and temperature extremes that do not restrict juveniles and adults can be limiting to postlarvae (Holt et al. 1981; Zein-Eldin 1989).

There are good physiological reasons for such limitations. In some crustacea, the size of antennal gland is larger in animals that must maintain an internal fluid concentration that is hypotonic relative to the environment. The larger size is due to longer nephridial canals providing more surface area for salt resorption and dilute urine production. This occurs in crayfish and some shrimp (Barnes 1980) and in freshwater amphipods (Green 1968). In marine, estuarine and terrestrial amphipods, the antennal glands are smaller than in comparable freshwater species (Schlieper 1930; Bousfield 1973). This restricts many, if not most, estuarine amphipods from oligohaline environments and may account for their paucity at the Trinity delta during 1987. Most decapod crustaceans, like fishes, osmoregulate through their gills (not antennal glands) in brackish waters (Barnes 1980). Adaptation to resident living under oligohaline conditions is difficult in any case. Few aquatic fauna are well adapted to survive and reproduce in this transition zone between rivers

and estuaries over the long term (Remane and Schlieper 1958). Those that do, such as some bivalves (*Rangia*) and annelids (nereids), usually exhibit specialized adaptations (Hopkins et al. 1973; Oglesby 1965a, 1965b). The capitellid and oligochaete infaunal worms that were abundantly found at the Trinity River delta are so adapted.

Marsh Utilization By Fishery Species

Our hypothesis was that marshes under mid-range salinity regimes are more utilized by fishery species. The test of the null hypothesis was to disprove that utilization at sites in the middle bay, in the middle of the salinity gradient, was not different from sites of the upper and lower subsystems. Using abundances, our results showed that fishery species were more abundant overall in the middle bay than in the other parts of the bay, thus disproving the null hypothesis. Indeed, most commercial and recreational species, including white shrimp, pink shrimp, blue crab, spotted seatrout, southern flounder, and red drum, had highest overall abundances in the middle bay. As previously discussed and expected, salinities of the middle bay overlapped extensively with those of the upper and lower bay (especially for short periods of time). This underscores the evidence that it is not salinity alone, but a complex of associated factors that create this attractive mid-bay environment. It is safe to say that the favorable conditions in marshes of the middle bay are influenced by or derived from the inputs of the upper and the lower bay.

Fishery species were not greatly attracted to the oligohaline marshes of the lower Trinity River delta during 1987. Although these delta marshes were not directly utilized, they nonetheless may be of substantial indirect importance to fishery species. Nearly the entire annual production of plants from the delta marshes at our sites is exported into the bay each year. This dead plant material

becomes particulate detritus that fuels detritus based food chains in at least the middle subsystem and perhaps the lower subsystem.

Distributions of Foods

Annelid worms and peracarid crustaceans (amphipods and tanaids) constituted the most abundant macrofaunal benthos in sediments in Galveston Bay. Evidence from our feeding experiments (Thomas 1989; Zimmerman et al.) and gut analyses (Minello et al. 1989) indicate these small animals are the principal foods of small fishes, shrimps and crabs in the estuary. Moreover, the literature cites numerous examples of the importance of these forage organisms in estuarine food chains (Kikuchi 1974; Young et al. 1976; Bell and Coull 1978; Nelson 1981; Stoner 1982; Huh and Kitting 1985; Whitfield 1988).

However, benthic foods (both plant and animal) appeared to be differentially abundant throughout the bay and highly dependent upon location. Among plants, vascular plant detritus appeared more abundant in the upper and middle subsystems, while epiphytic and macro-algae was most abundant in the lower subsystem. Annelid worms were numerous throughout, but most abundant in the upper subsystem. Peracarid crustaceans were most abundant in the middle subsystem and nearly absent in the upper subsystem.

Since larger predators (fishes, crabs and shrimps) were exceptionally numerous in the middle subsystem, a food chain relationship with forage organisms can be inferred. We propose that the relationship is based upon the input of detritus and abundances of peracarids. As detritus from delta marshes is exported down the salinity gradient, it breaks up into smaller particles, is colonized and enriched with nitrogen by microflora, thus becoming ideal food for detritivorous annelids

(Tenore 1977; Findlay and Tenore 1982), peracarids (Hargrave 1970; Monk 1977; Zimmerman et al. 1979) and molluscs (Newell 1964). Since very large populations of annelids and peracarids occurred in the middle subsystem, detritus availability and conditioning appears to be most favorable in this area. These small prey are available to support large numbers of small fishes and decapod crustaceans, and many of these, in turn, serve as ready food for larger fishes and crustaceans. Thus, a classical detritus-based benthic food web (Odum and Heald 1975; Odum 1980) is created in the middle bay. Among the forage animals, peracarids appear to be more preferred and are more available than annelids (Huh and Kitting 1985; Leber 1985; Luczkovich 1988; Thomas 1989; Zimmerman et al.). The relative absence of peracarids from the delta marshes was striking and we predict it may have been a reason that so few predators were attracted there.

Effect of Salinity on Fishery Habitat

The direct effect of salinity (that is, salinity per se) appears to have little influence on distributions of demersal fishes, crabs and shrimps except under extreme circumstances. Even then, most estuarine species tolerate very low salinities (less than 1 ppt) for short periods of time (days to weeks). Large natant decapods and fishes in Texas estuaries commonly move across salinity gradients into low salinities (Baldauf 1970; Renfro 1960). Their presence or absence in low salinity situations appears to be a behavior of choice. Species such as brown shrimp, white shrimp, blue crab, grass shrimp, menhaden, bay anchovies, striped mullet, red drum, southern flounder and Atlantic croaker are often noted in very low salinity waters. During the summer of 1987, we obtained all of these species in the mid-bay marsh at Smith Point with salinity of 0.8 ppt. The salinity was similar (0.5 ppt) at the delta marsh sites, yet these estuarine species were virtually absent. We submit that

the reason for these differences in abundances was not due to the short term effect of salinity itself, but to habitat differences that developed from long term exposure to low salinity.

One difference we noted was the effect of salinity on distribution of forage organisms. The absence of amphipods and tanaidaceans in the delta marshes compared to their exceptional abundances in mid-bay marshes suggests that this is at least one long term salinity effect. It has been known that oligohaline salinity regimes (< 5 ppt) diminish the number of residents of small less mobile estuarine species (Remane and Schlieper 1958). Estuarine amphipods and tanaids are among fauna whose species are limited to only a few adapted to tolerate oligohaline conditions for long periods of time. Since they are highly useful forage organisms, their absence diminishes the value of a low salinity marsh for predators. However, we know little about these kinds of effects and how they may control the relationships between salinity and fishery productivity. This is a fertile and necessary area of further research.

CONCLUSIONS

Salinity Characteristics of Galveston Bay Marshes

The environment in the Galveston Bay system is characterized by a strong salinity gradient. Salinities along the gradient range from fresh (0 ppt) to hypersaline (> 40 ppt) depending upon seasonal and annual rainfall. Normally, the upper system (Trinity Bay) is oligohaline to mesohaline, the middle system (Galveston Bay proper) is mesohaline to polyhaline, and the lower system (West Bay and Christmas Bay) is polyhaline. High rainfall during the spring and summer of 1987 reduced the salinities, causing in oligohaline conditions (< 1 ppt) throughout the upper

conditions (< 1 ppt) throughout the upper system and highly variable conditions (< 1 to 15 ppt) in the middle system. Salinities of the lower system (22 to 33 ppt) were relatively unaffected. The resulting summer salinity gradient was the steepest of the year. As freshwater input diminished in the fall and equinox tides caused salinities in the upper system to increase to near 10 ppt, the slope of the gradient lessened across the system. These long term and short term salinity characteristics reflect freshwater inflow effects that determine the nature of marsh communities in the system.

Biological Characteristics of Galveston Bay Marshes

Marsh communities are clearly different between the upper, middle and lower subsystems in Galveston Bay. Biological attributes uniquely characterize each subsystem, inferring relationships to salinity. At the same time, the subsystems are interconnected and depend on one another through materials flow. These interrelationships appear to have a large effect on determining how the different marshes function for fishery species.

The upper subsystem, represented by the lower Trinity River delta, is oligohaline and strongly reflects freshwater influences. Emergent marsh plants (*Scirpus* and *Sagittaria*) are those commonly associated with active deltaic environments. This is one of the few areas in Galveston Bay supporting large stands of submerged aquatic vegetation (SAV). Part of the deltaic SAV is an extensive area of previously unreported *Vallisneria* habitat. During the winter months most of the emergent marsh and subtidal SAV dies back and is exported. SAV growth is essentially limited to the summer months. Among forage organisms present in the marshes and SAV habitat, peracarid crustaceans are few, but annelid worms are abundant. This pattern corresponds to relatively low useage of del-

taic marsh and SAV by fishes and decapod crustaceans (usually not significantly different from useage of nonvegetated open water). As a result, since it is continuously available, nonvegetated subtidal bottom appears to be more directly useful as nursery habitat in the upper subsystem compared to the marsh surface and SAV. Even so, overall abundances of animals are significantly lower in the upper subsystem compared to the middle and lower subsystems.

By contrast, peracarids are exceptionally abundant in the middle subsystem and abundances of fishes and decapods are also high. The relationship exists because the large numbers of peracarids, in both marsh and open water, are useful as food to juveniles of many demersal species. Consequently, marsh and nonvegetated bottom in the middle subsystem serve equally as nursery habitats that contribute to high production in fishery species. However, this productivity appears to be directly related to organic materials flow from the upper subsystem. We propose that the middle region receives most of its dead plant material, that is highly useful to peracarid detritivores such as amphipods and tanaids, from the deltaic marshes of upper region.

In the lower subsystem, marshes appear to be proportionately more important as nurseries compared to nonvegetated bottom. Forage organisms are significantly more abundant on the marsh surface and the structure of *Spartina* culms offers stable year-around shelter. In addition, epiphytic algae populations are well developed in lower subsystem marshes. These factors improve the direct value of these marshes to exploiting juveniles of fishes and decapods crustaceans. The salinity regime, however, is not necessarily less stressful than in other parts of the bay, since hypersaline conditions are not uncommon in the lower system.

The Relationship Between Salinity and Marsh Utilization

Over time, each part of the Galveston Bay system incurs salinities that may cause physiological stress to organisms. However most of the higher estuarine animals (such as fishery juveniles) are adapted to accommodate these stresses, and therefore, most distributions are probably due to other factors.

Fishery species were more abundant as species and individuals in marshes with mesohaline to polyhaline salinity regimes. This occurred primarily in the middle area of Galveston Bay where freshwater and saltwater mixing characteristics were strong. Material imports and physical mixing processes here stimulated food chain responses. Thus, cause-and-effect relationships leading to high utilization were related to salinity, but not necessarily controlled by salinity. Nevertheless, salinity parameters may be viewed as an indicator of physical mixing and marsh utilization characteristics.

LITERATURE CITED

- Armitage, P. D. 1978.** Downstream changes in the composition, numbers, and biomass of fauna in the Tees below Cow Green Reservoir and in an unregulated tributary, Maize Beck, in the first five years after impoundment. *Hydrobiologia* 58:145-156.
- Armstrong, N. E. 1987.** The ecology of open-bay bottoms of Texas: a community profile. U.S. Fish. Wildl. Serv. Biol. Rep. 85(7.12) 104 pages.
- Baldauf, R. J. 1970.** A study of selected chemical and biological conditions of the lower Trinity River and the upper Trinity Bay. Tech. Rep. No. 26, Water Resources Institute, Texas A&M Univ., College Station, Texas. 168 pp.
- Barnes, R. D. 1980** (4th edition). Invertebrate zoology. Saunders College, Philadelphia, PA. 1089 pp.
- Baxter, K. N. and W. C. Renfro 1967.** Seasonal occurrence and size distribution of postlarval brown and white shrimp near Galveston, Texas, with notes on species identification. *Fish. Bull.* 66:149-158.
- Baxter, R. M. 1977.** Environmental effects of dams and impoundments. *Ann. Rev. Ecol. Syst.* 8:255-283.
- Bell, S.S. and B.C. Coull 1978.** Field evidence that shrimp predation regulates meiofauna. *Oecologia* 35:141-148.
- Bennett, J. P., J. W. Woodward, and D. J. Shultz 1986.** Effect of discharge on the chlorophyll a distribution in the tidally influenced Potomac River. *Estuaries* 9:250-260.
- Benson, N. G. 1981.** The freshwater-inflow-to estuaries issue. *Fisheries* 6 (5):8-10.
- Borey, R. B. 1979.** Hydrology and organic carbon export of an irregularly flooded brackish marsh on the upper Texas coast. Dissertation. Rice Univ., Houston, Texas. 142 pp.
- Borey, R. B., P. A. Harcombe and F. M. Fisher 1983.** Water and organic fluxes from an irregularly flooded brackish marsh of the upper Texas coast, U.S.A. *Estuar. Coast Shelf Sci.* 16:379-402.
- Bousfield, E. L. 1973.** Shallow-water gammaridean amphipoda of New England. Cornell University Press, Ithaca, N.Y. 312 pp.
- Brongersma-Sanders, M. 1957.** Mass mortality in the sea. pp.941-1010. In: J.W. Hedgpeth (ed.) *Treatise on Marine Ecology and Paleoecology* Vol. 1 Ecology. Geol. Soc. Amer. Mem. 67.
- Browder, J. A., H. A. Bartley and K. E. Davis 1985.** A probabilistic model of the relationship between marshland-water interface and marsh disintegration. *Ecol. Model.* 29:245-260.
- Chambers, G. V., and A. K. Sparks 1959.** An ecological survey of the Houston Ship Channel and adjacent waters. *Univ. Tex. Inst. Mar. Sci. Publ.* 6:215-250.
- Chin, E. 1960.** The bait fishery of Galveston Bay, Texas. *Trans. Am. Fish. Soc.* 89:135-141.
- Christmas, J.Y., W. Langley, and T. Van Devender 1976.** Investigations of commercially important penaeid shrimp in Mississippi. *Gulf Coast Res. Lab., Ocean Springs, Miss.* 66 pp.
- Conner, W. H., J. W. Day, Jr., J. G. Gosselink, C. S. Hopkinson, Jr., and W. C. Stowe 1987.** Vegetation: composition and production. Chapter 4. In: Conner, W. H., and J. W. Day, Jr., (eds.) *The ecology of Barataria Basin, Louisiana: an estuarine profile.* U.S. Fish Wildl. Serv. Biol. Rep. 85(7.13). 165 pp.
- Conover, J. T. 1964.** The ecology, seasonal periodicity, and distribution of benthic plants in some Texas lagoons. *Bot. Mar.* 7:4-41.
- Copeland, B. J. 1966.** Effects of decreased river flow on estuarine ecology. *J. Water Pollut. Control Fed.* 38:1831-1839.
- Cowardin, L.M., V. Carter, F.C. Golet and E.T. LaRoe 1979.** Classification of wetlands and deepwater habitats of the United States. U.S. Fish Wildl. Serv. Prog. FWS/OBS-79/31. 103 pp.
- Craig, N. J., R. E. Turner and J. W. Day, Jr. 1980.** Wetland losses and their consequence in coastal Louisiana. *Z. Geomorph. N. F.* 34:255-241.
- Davis, H. C., and A. Calabrese 1964.** Combined effects of temperature and salinity on development of eggs and growth of larvae of *M. mercenaria* and *C. virginica*. U.S. Fish Wildl. Serv. *Fish. Bull.* 63:643-655.
- Diener, R. A. 1975.** Cooperative Gulf of Mexico estuarine inventory and study - Texas: area description. NOAA Tech. Rep., NMFS CIRC-393. 129 pp.

- Faller, K. H. 1979.** Shoreline as a controlling factor in commercial shrimp production. NASA Tech. Memo 72-732, 27 pp.
- Findlay, S. and K. Tenore 1982.** Nitrogen source for a detritivore: detritus substrate versus associated microbes. *Science* 218:371-373.
- Fisher, J. C. 1983.** Chemical and physical characteristics of water in estuaries of Texas, October 1976 - September 1978. Tex. Dept. Water Res. Rpt. 282. 215 pp. Tex. Dept. Water Res., P.O. Box 13087, Austin, TX 78711.
- Fisher, W. L., J. H. McGowen, L. F. Brown, Jr., and C. G. Groat 1972.** Environmental geological atlas of the Texas coastal zone-Galveston-Houston area. Bureau of Economic Geology, Univ. Texas, Austin, TX. 91 pp.
- Flint, R. W. 1985.** Long-term estuarine variability and associated biological response. *Estuaries* 8:158-169.
- Flint, R. W., and J. A. Younk 1983.** Estuarine benthos: long-term community structure variations, Corpus Christi bay, Texas. *Estuaries* 6:126-141.
- Flint, R. W., G. L. Powell, and R. D. Kalke 1983.** Ecological effects from the balance between new and new recycled nitrogen in Texas coastal waters. *Estuaries* 9:284-294.
- Gifford, C. A. 1962.** Some aspects of osmotic and ionic regulation in the blue crab, *Callinectes sapidus*, and the ghost crab, *Ocypode albicans*. *Publ. Inst. Mar. Sci.* 8:97-125.
- Gleason, D.F. 1986.** Utilization of salt marsh plants by postlarval brown shrimp: carbon assimilation rates and food preferences. *Mar. Ecol. Prog. Ser.* 31:151-158.
- Gleason, D. F. and R. J. Zimmerman 1984.** Herbivory potential of postlarval brown shrimp associated with salt marshes. *J. Exp. Mar. Biol. Ecol.* 84:235-246.
- Green, J. 1968.** The biology of estuarine animals. University Washington Press, London. 401 pp.
- Gunter, G. 1967.** Some relationships of estuaries to fisheries of the Gulf of Mexico. pp. 621-637. In: G.H. Lauff (ed). *Estuaries*. Amer. Assoc. Adv. Sci. Publ. No. 83.
- Gunter, G. 1979.** Studies of the southern oyster borer, *Thais haemastoma*. *Gulf Res. Rep.* 6:249-260.
- Hammerschmidt, P. C. 1985.** Relative blue crab abundance in Texas coastal waters. *J. Shellfish Res.* 5:9-11.
- Hargrave, B. T. 1970.** The effect of a deposit-feeding amphipod on the metabolism of benthic microflora. *Limnol. Oceanogr.* 15:21-30.
- Harry, H.W. 1976.** Correlation of benthic mollusca with substrate composition in lower Galveston Bay, Texas. *Veliger* 19:135-152.
- Haskin, H. H. 1964.** The distribution of oyster larvae. Proc. of a symposium on experimental marine ecology. Occas. Pap. 2, Grad. School Oceanogr., Univ. R.I.:76-80.
- Hatton, R. S., R. D. Delune and W. H. Patrick, Jr. 1983.** Sedimentation, accretion, and subsidence in marshes of Barataria Basin, Louisiana. *Limnol. Oceanogr.* 28:494-502.
- Hedgpeth, J. W. 1953.** An Introduction to the zoogeography of the northwestern Gulf of Mexico with reference to invertebrate fauna. *Publ. Inst. Mar. Sci. Texas* 3:107-224.
- Hicks, S. D., H. A. Debaugh Jr. and L. E. Hickman 1983.** Sea level variations for the United States 1855-1980. NOAA/NOS Rep., National Ocean Survey, Tides and Water Levels Branch, Rockville, MD. 170 pp.
- Hoese, H. D. 1960.** Biotic changes in a bay associated with the end of a drought. *Limnol. Oceanogr.* 5:326-336.
- Hofstetter, R. P. 1977.** Trends in population levels of the American oyster *Crassostrea virginica* Gmelin on public reefs in Galveston Bay, Texas. Texas Parks and Wildl. Dept., Tech. Ser. No. 24. 90 pp.
- Hofstetter, R. P. 1983.** Oyster population trends in Galveston Bay 1973-1978. Texas Parks and Wildl. Dept. Manag. Data Ser. No. 51. 33 pp.
- Holt, J., R. Godbout and C. R. Arnold 1981.** Effects of temperature and salinity on egg hatching and larval survival of red drum, *Sciaenops ocellata*. *Fish. Bull.* 79:569-573.
- Holt, J. and K. Strawn 1983.** Community structure of macrozooplankton in Trinity and upper Galveston Bays. *Estuaries* 6:66-75.

- Hopkins, A. E. 1931.** Factors influencing the spawning and setting of oysters in Galveston Bay, Texas. Bull. U.S. Bur. Fish. 47:57-83.
- Hopkins, S. H., J. W. Anderson and K. Horvath 1973.** The brackish water clam *Rangia cuneata* as indicator of the ecological effects of salinity changes in coastal waters. Contract Rep. H-73-1, U.S. Army Engineer Waterways Experiment Station, Vicksburg, Mississippi. 250 pp.
- Huh, S. and C. Kitting 1985.** Trophic relationships among concentrated populations of small fishes in seagrass meadows. J. Exp. Mar. Biol. Ecol. 92:29-43.
- Kikuchi, T. 1974.** Japanese contributions on consumer ecology in eelgrass (*Zostera marina* L.) beds, with special reference to trophic relationships and resources inshore fisheries. Aquaculture 4:145-160.
- Leber, K. M. 1985.** The influence of predatory decapods, refuge, and microhabitat selection on seagrass communities. Ecology 66:1951-1964.
- Loesch, H. 1965.** Distribution and growth of penaeid shrimp in Mobile Bay, Alabama. Publ. Inst. Mar. Sci., Univ. Tex. 10:41-58.
- Loosanoff, V. L. 1953.** Gonadal development and spawning of oysters (*C. virginica*) in low salinities. Anat. Rec. 101:705 (Abstract).
- Lowe, G. C., Jr. and E. R. Cox 1978.** Species composition and seasonal periodicity of the marine benthic algae of Galveston Island, Texas. Contrib. Mar. Sci. 21:9-24.
- Luczkovich, J. J. 1988.** The role of prey detection in the selection of prey by pinfish *Lagodon rhomboides* (Linnaeus). J. Exp. Mar. Biol. Ecol. 123:15-30.
- Matlock, G. C. 1987.** The role of hurricanes in determining year-class strength of red drum. Contrib. Mar. Sci. 30:39-47.
- McHugh, J. L. 1966.** Management of estuarine fisheries. pp. 133-154. In: R.F. Smith, A.H. Schwartz and W.H. Massmann (eds.). A Symposium on Estuarine Fisheries. Amer. Fish. Soc. Spec. Publ. 3.
- McHugh, J. L. 1976.** Estuarine fisheries: are they doomed? Vol. I. pp. 15-27. In: M. Wiley (ed.) Estuarine Processes. Academic Press, NY.
- McIvor, C. C. and W. E. Odum 1988.** Food, predation risk, and microhabitat selection in a marsh fish assemblage. Ecology 69:1341-1351.
- Mendelssohn, I. A. 1979.** Nitrogen metabolism in the height forms of *Spartina alterniflora* in North Carolina. Ecology 60:574-584.
- Minello, T. J., and R. J. Zimmerman 1983.** Fish predation on juvenile brown shrimp, *Penaeus aztecus* Ives: the effect of simulated *Spartina* structure on predation rates. J. Exp. Mar. Biol. Ecol. 72:211-231.
- Minello, T.J., R.J. Zimmerman and T.E. Czaplak 1989.** Habitat-related differences in diets of small fishes from Lavaca Bay, Texas, 1985-1986. NOAA Tech. Memo. NMFS-SEFC-236. 16p.
- Minello, T. J., R. J. Zimmerman, and E. X. Martinez 1987.** Fish predation on juvenile brown shrimp, *Penaeus aztecus* Ives: effects of turbidity and substratum on predation rates. Fish. Bull. 85:59-70.
- Monk, D. C. 1977.** The digestion of cellulose and other dietary components, and pH of the gut in the amphipod *Gammarus pulex* (L.). Freshw. Biol. 7:431-440.
- More, W. R. 1969.** A contribution to the biology of the blue crab (*Callinectes sapidus*, Rathbun) in Texas, with a description of the fishery. Tex. Parks Wildl. Dept. Tech. Ser. 1:1-13.
- More, W. R. and A. W. Moffett 1964.** Population studies of the blue crabs of Galveston Bay system. Coastal Fish. Proj. Reps., Tex. Parks Wildl. Dept., pp. 551-575.
- Nelson, W.G. 1981.** Experimental studies of decapod and fish predation on seagrass macrobenthos. Mar. Ecol. Prog. Ser. 5:141-149.
- Newell, R. 1964.** The role of detritus in the nutrition of two marine deposit feeders, the prosobranch, *Hydrobia ulvae* and the bivalve, *Macoma balthica*. Proc. Zool. Soc. Lond. 144:25-45.
- Nixon, S. W. 1981.** Remineralization and nutrient cycling in coastal marine ecosystems. pp. 111-138. In: B.J. Neilson and L.E. Cronin (eds.), Estuaries and nutrients. Humana Press, Clifton, N. J.
- Nixon, S. W. 1982.** Nutrient dynamics, primary production and fishery yields of lagoons. Oceanol. Acta. Proc. Internat. Symp. Coastal Lagoons.

- Odum, E. P. 1980.** The status of three ecosystem-level hypotheses regarding salt marsh estuaries: tidal subsidy, outwelling, and detritus-based food chains. pp. 485-495. In: V. S. Kennedy (ed.), *Estuarine Perspectives*. Academic Press, New York, N.Y.
- Odum, W. E. and E. J. Heald 1975.** Trophic analyses of an estuarine mangrove community. *Bull. Mar. Sci.* 22:671-738.
- Oglesby, L. C. 1965 a.** Steady-state parameters of water and chloride regulation in estuarine nereid polychaetes. *Comp. Biochem. Physiol.* 14:621-640.
- Oglesby, L.C. 1965b.** Water and chloride fluxes in estuarine nereid polychaetes. *Comp. Biochem. Physiol.* 16:437-455.
- Pandian, T. J. 1980.** Impact of dam-building on marine life. *Helgol. Meeresunters.* 33:415-421.
- Parker, J. C. 1965.** An annotated checklist of the fishes of the Galveston Bay system, Texas. *Publ. Inst. Mar. Sci. Univ.* 10:201-220.
- Parker, J. C. 1970.** Distribution of juvenile brown shrimp (*Penaeus aztecus* Ives) in Galveston Bay, Texas, as related to certain hydrographic features and salinity. *Contrib. Mar. Sci.* 15:1-12.
- Peters, J. C. 1982.** Effects of river and streamflow alteration on fishery resources. *Fisheries* 7(2):20-22.
- Powell, G. L. 1985.** Water quality management problems in Texas bays and estuaries. Papers from the Estuarine Management Practices Symposium. Louisiana Sea Grant Program. LSU Center for Wetland Resources, Baton Rouge, LA 70803. pp. 56-71.
- Pritchard, D. 1967.** What is an estuary: physical viewpoint. pp. 3-5. In: G.H. Lauff (ed.). *Estuaries*. Amer. Assoc. Adv. Sci. Publ. No. 83.
- Provost, M.W. 1976.** Tidal datum planes circumscribing salt marshes. *Bull. Mar. Sci.* 26:558-563.
- Pulich, W., Jr. and S. Rabalais 1986.** Primary production potential of blue-green algal mats on southern Texas tidal flats. *Southw. Nat.* 31:39-47.
- Pullen, E. J. and L. Trent 1969.** Hydrographic observations from the Galveston Bay system, Texas, 1959-67. *Contrib. No. 280*, Bureau of Com. Fish. Biol. Lab., Galveston, TX. U.S. Fish Wildl. Serv. Data Rep. 31. 151 pp. (XFDR-B 31 1-151).
- Pullen, E. J., W. L. Trent and G. B. Adams 1971.** A hydrographic survey of the Galveston Bay system, Texas, 1963-66. NOAA Tech. Rep. NMFS SSRF-639. 13 pp.
- Rader D. N. 1984.** Salt-marsh benthic invertebrates: small-scale patterns of distribution and abundance. *Estuaries* 7:413-420.
- Reid, G. K., Jr. 1955.** A summer study of the biology and ecology of East Bay, Texas. I. Introduction, description of the area, some aspects of the fish community, the invertebrate community. *Tex. J. Sci.* 7:316-343.
- Reidenbaugh, T. G., W. C. Banta, M. Varricchio, R. P. Strieter and S. Mendoza. 1983.** Short-term accretional and erosional patterns in a Virginia salt marsh. *Gulf Res. Rep.* 7:211-215.
- Remane, A. 1934.** Die brackwasserfauna. *Verh. dt. zool. Ges.* 36:34-74.
- Remane, A. and C. Schlieper 1958.** The biology of brackish water. Translated 1971, Wiley-Interscience, New York, N.Y. 372 pp.
- Renfro, W. C. 1960.** Salinity relations of some fishes in the Aransas River, Texas. *Tulane Studies in Zoology* 8:83-91.
- Rogers, B. D., and W. H. Herke 1987.** Diel otter trawl catch of Atlantic croaker, *Micropogonias undulatus*, in a Louisiana estuary. *Northeast Gulf Science* 9:147-152.
- Sage, W. W. and M. J. Sullivan 1978.** Distribution of blue-green algae in a Mississippi Gulf Coast salt marsh. *J. Phycol.* 14:333-337.
- Schlieper, C. 1930.** Die osmoregulation wasserlebender tiere. *Biol. Rev.* 5:309-356.
- Sheridan, P. F. 1983.** Abundance and distribution of fishes in the Galveston Bay system, 1963-1964. *Contrib. Mar. Sci.* 26:143-163.
- Sheridan, P. F., R. D. Slack, S. M. Ray, L. W. McKinney, E. F. Klima and T. R. Calnan 1988.** Biological components of Galveston Bay. pp. 23-51. In: *An Estuary-of-the Month Seminar, Galveston Bay, Issues, Resources, Status and Management*. Presented by NOAA/ Estuarine Programs Office and the U.S. Environmental Protection Agency, March 14, 1988. Published by the U.S. Printing Office, Washington, D.C.

- Smith, N. 1982.** Tidal flushing of intracoastal bays. *Contrib. Mar. Sci.* 25:1-12.
- Stern, M. K., J. W. Day, Jr., and K. G. Teague 1986.** Seasonality of materials transport through a coastal freshwater marsh: riverine versus tidal forcing. *Estuaries* 9:301-308.
- Stokes, G.M. 1974.** The distribution and abundance of penaeid shrimp in the lower Laguna Madre of Texas, with a description of the live bait shrimp fishery. *Coastal Fish. Proj. Rep., Texas Parks Wildl. Dept. Tech. Ser. No. 15.* 32 pp.
- Stoner, A.W. 1982.** The influence of benthic macrophytes on the foraging behaviour of pinfish, *Lagodon rhomboides* (Linnaeus). *J. Exp. Mar. Biol. Ecol.* 58:271-284.
- Sullivan, M. J. 1978.** Diatom community structure: taxonomic and statistical analyses of a Mississippi salt marsh. *J. Phycol.* 14:468-475.
- Sullivan, M. J. 1981.** A preliminary checklist of marine benthic diatoms of Mississippi. *Gulf Res. Rep.* 7:13-18.
- Tagatz, M. E. 1971.** Osmoregulatory ability of blue crabs in different temperature-salinity combinations. *Chesapeake Science* 12:14-17.
- Tenore, K. R. 1977.** Utilization of aged detritus derived from different sources by the polychaete *Capitella capitata*. *Mar. Biol.* 44:51-55.
- Tenore, K. R., J. H. Tietjen and J. J. Lee. 1977.** Effect of meiofauna on the incorporation of aged eelgrass detritus by the polychaete, *Nephtys incisa*. *J. Fish. Res. Bd. Can.* 34:563-567.
- Texas Department of Water Resources 1981a.** Trinity-San Jacinto estuary: a study of the influence of freshwater inflows. *Tex. Dept. Water Res. LP-113.* Tex. Dept. Water Res., P.O. Box 13087, Austin, TX 78711.
- Texas Department of Water Resources 1981b.** The influence of freshwater inflows upon the major bays and estuaries of the Texas coast. Executive Summary. LP-115. *Tex. Dept. Water Res., P.O. Box 1307, Austin, Tx 78711.*
- Thomas, J.L. 1989.** A comparative evaluation of *Halodule wrightii* Aschers, *Spartina alterniflora* Loisel and bare sand as nursery habitats for juvenile *Callinectes sapidus*. M.S. Thesis, Texas A&M University. 119 pages.
- Thomas, J. T., R. J. Zimmerman and T. J. Minello 1990.** Abundance patterns of juvenile blue crabs (*Callinectes sapidus*) in nursery habitats of two Texas bays. *Bull. Mar. Sci.* 46(1):(in press).
- Turek, J. G., T. E. Goodger, T. E. Bigford and J. S. Nichols 1987.** Influence of freshwater inflows on estuarine productivity. NOAA Tech. Memo. NMFS-F/NEC-46. 26 pp.
- Turner, R. E. 1977.** Intertidal vegetation and commercial yields of penaeid shrimp. *Trans. Am. Fish. Soc.* 106:411-416.
- Van Sickle, V. R., B. B. Barrett, T. B. Ford and L. J. Gulick 1976.** Barataria Basin: salinity changes and oyster distribution. Sea Grant Publication No. LSU-T-76-02. Louisiana State University, Baton Rouge, LA 70803.
- Webb, J. W. 1983.** Soil salinity variations and their effects on *Spartina alterniflora*. *Contrib. Mar. Sci.* 26:1-13.
- Wermund, E.G., R.A. Morton and G. Powell 1989.** Geology, climate and water circulation of the Galveston Bay System. pp. 3-22. In: An Estuary-of-the Month Seminar, Galveston Bay, Issues, Resources, Status and Management. Presented by NOAA/Estuarine Programs Office and the U.S. Environmental Protection Agency, March 14, 1988. Published by the U.S. Printing Office, Washington, D.C.
- West, R. L. 1972.** Inventory of aquatic vegetation. Texas Parks Wildl. Dept., Coastal Waterfowl project No. W-29-R-25, Job No. 20. 6 pp.
- White, W. A., T. R. Calnan, R. A. Morton, R. S. Kimble, T. G. Littleton, J. H. McGowen, H. S. Nance and K. E. Schmedes 1985.** Submerged lands of Texas, Galveston-Houston area: sediments, geochemistry, benthic macroinvertebrates, and associated wetlands. Report, 145 pp. Bureau of Economic Geology, Univ. Tex., Austin, TX 78713.
- Whitfield, A.K. 1988.** The fish community of the Swartvlei Estuary and the influence of food availability on resource utilization. *Estuaries* 11:160-170.
- Williams, A.B. 1955.** A contribution to the life histories of commercial shrimps (Penaeidae) in North Carolina. *Bull. Mar. Sci.* 5:116-146.
- Williams-Cowper, S. 1978.** The drift algae community of seagrass beds in Redfish Bay, Texas. *Contrib. Mar. Sci.* 21:126-132.

Young, D.K., M.A. Buzas and M.W. Young 1976. Species densities of macrobenthos associated with seagrass: a field experimental study of predation. J. Mar. Res. 34:577-592.

Zein-Eldin, Z. P. 1989. Growth and survival of postlarval *Penaeus setiferus* and *P. aztecus* under controlled temperature and salinity combinations. (unpublished manuscript) 25 pp.

Zimmerman, R. J. and A. H. Chaney 1969. Salinity decrease as an affector of molluscan density levels in a turtle grass (*Thalassia testudinum* Konig) bed in Redfish Bay, Texas. Texas A&I University Studies 2: 5-10.

Zimmerman, R. J., R. Gibson and J. Harrington 1979. Herbivory and detritivory among gammaridean amphipods from a Florida seagrass community. Mar. Biol. 54:41-47.

Zimmerman, R. J. and T. J. Minello 1984. Densities of *Penaeus aztecus*, *Penaeus setiferus*, and other natant macrofauna in a Texas salt marsh. Estuaries 7:421-433.

Zimmerman, R. J., T. J. Minello, and G. Zamora 1984. Selection of vegetated habitat by *Penaeus aztecus* in a Galveston Bay salt marsh. Fish. Bull. 82:325-336.

Zimmerman, R. J., T. J. Minello, and S. Dent. Habitat-related growth and resource partitioning by penaeid shrimp in a salt marsh. Mar. Ecol. Prog. Ser. (conditionally accepted).

APPENDIX I: Principal Keys and References Used to Identify Galveston Bay Aquatic Fauna.

Fishes:

Hoese, H.D. and R.H. Moore 1977. Fishes of the Gulf of Mexico, Texas, Louisiana, and adjacent waters. Texas A&M Press, College Station, Texas. 327 pp.

Murdy, E.O. 1983. Saltwater fishes of Texas: a dichotomous key. Texas A&M Sea Grant College Program TAMU-SG-83-607, College Station.

U.S. Fish and Wildlife Service 1978. Development of fishes of the Mid-Atlantic Bight: an atlas of egg, larval and juvenile stages. Volumes I-VII. U.S. Fish Wildl. Serv., Biol. Serv. Program, FWS/OBS-78/12.

Crustaceans:

Bousfield, E.L. 1973. Shallow-water gammaridean Amphipoda of New England. Cornell University Press, Ithaca, New York. 312 pp.

Chaney, A.H. 1983. Key to the common inshore crabs of Texas. pp. 1-30 In: A.H. Chaney, Keys to selected marine invertebrates of Texas. Caesar Kleberg Wildlife Research Institute Tech. Bull. No. 4, Kingsville, Texas. 86 pp.

Felder, D.L. 1973. An annotated key to crabs and lobsters (Decapoda, Reptantia) from coastal waters of the northwestern Gulf of Mexico. Center for Wetland Resources, Louisiana State University. LSU-SG-73-02. Baton Rouge, Louisiana. 103 pp.

Heard, R.W. 1982. Guide to common tidal marsh invertebrates of the northeastern Gulf of Mexico. Mississippi-Alabama Sea Grant Consortium. MASGP-79-004. Ocean Springs, Mississippi. 82 pp.

Schultz, G.A. 1969. The marine isopod crustaceans. William C. Brown Co. Publ., Dubuque, Iowa. 359 pp.

Williams, A.B. 1984. Shrimps, lobsters and crabs of the Atlantic coast of the eastern United States, Maine to Florida. Smithsonian Institution Press. Washington, D.C. 550 pp.

Molluscs:

Andrews, J. 1981. Texas shells. University of Texas Press. Austin, Texas. 175 pp.

Annelids:

Fauchald, K. 1977. The polychaete worms. Definitions and keys to the orders, families and genera. Natural History Museum of Los Angeles County in conjunction with the Allan Hancock Foundation. Science Series 28, University of Southern California, Los Angeles, California. 188 pp.

Uebelacker, J.M. and P.G. Johnson (eds.) 1984. Taxonomic guide to the polychaetes of the northern Gulf of Mexico. Vol. I - VI. Minerals Management Service, U.S. Dept. Interior, Gulf of Mexico Regional Office, Metairie, Louisiana.

Plants:

Charbreck, R.H. and R.E. Condrey 1979. Common vascular plants of the Louisiana marsh. Sea Grant Pub.No. LSU-T-79-003. Louisiana State Center for Wetland Resources, Baton Rouge, Louisiana. 116 pp.

Edwards, P. 1976. Illustrated guide to the seaweeds and seagrasses in the vicinity of Port Aransas, Texas. Univ. Texas Press, Austin, Texas. 126 pp.

Eleuterius, L.N. 1980. Tidal marsh plants of Mississippi and adjacent states. Mississippi-Alabama Sea Grant Consortium Pub. No. MASGP-77-039. Gulf Coast Research Laboratory, Ocean Springs, Mississippi. 130 pp.

Tarver, D.P., J.A. Rodgers, M.J. Mahler and R. L. Lazor 1986.

Aquatic and wetland plants of Florida. Published by the Bureau of Aquatic Plant Research and Control, Florida Department of Natural Resources, Tallahassee, Florida. 127p.

[THIS PAGE INTENTIONALLY LEFT BLANK]

APPENDIX II: PHYSICAL MEASUREMENTS, SPRING.

ENVIRONMENTAL PARAMETERS SPRING SAMPLING SET	Site 1 TRINITY RIVER INNER DELTA				Site 2 TRINITY RIVER OUTER DELTA			
	Vegetated		Non-vegetated		Vegetated		Non-vegetated	
	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.
Temperature (Deg. C)	27.4	0.43	28.5	0.22	28.7	0.93	28.6	1.27
Salinity (ppt)	0	0.02	0	0	0	0	0	0.01
Dissolved Oxygen (ppm)	8.1	0.42	8.5	0.29	9.2	0.51	9.7	0.74
Turbidity (FTU)	45	5.58	43.5	10.44	28	2.04	38.3	3.12
Median Depth (cm)	7.4	0.92	38	13.5	13.6	1.09	20.8	3.68
Maximum Depth (cm)	9.5	0.87	44	15.44	17	1.41	21.3	3.68
Minimum Depth (cm)	5.3	1.03	32	11.61	10.3	3.42	20.3	3.68
Time Interval: (date time)	(April 21: 1835 - 1929 hrs)				(April 20: 1610 - 1854 hrs)			

	Site 3 SMITH POINT				Site 4 MOSES LAKE			
	Vegetated		Non-vegetated		Vegetated		Non-vegetated	
	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.
Temperature (Deg. C)	31.1	0.24	29.8	0.53	29.5	0.67	29.6	0.78
Salinity (ppt)	8.8	0.25	8.3	0.25	15.5	0.29	15.5	0.29
Dissolved Oxygen (ppm)	11.1	0.36	12.3	0.36	12.7	2.1	12.1	2.16
Turbidity (FTU)	13	5.43	21	4.81	31.5	10.99	27	7.55
Median Depth (cm)	22.5	2.61	40.5	3.85	8.5	1.67	18.8	1.61
Maximum Depth (cm)	25	2.52	41.3	3.97	16	3.19	19.5	1.66
Minimum Depth (cm)	20	2.86	39.8	3.75	1	0.71	18	1.58
Time Interval: (date time)	(April 21: 1457 - 1613 hrs)				(April 30: 1440 - 1551 hrs.)			

	Site 5 JAMAICA BEACH				Site 6 CHRISTMAS BAY			
	Vegetated		Non-vegetated		Vegetated		Non-vegetated	
	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.
Temperature (Deg. C)	28.8	0.47	28.8	0.23	23.7	0.25	23.6	0.2
Salinity (ppt)	33.3	0.14	33.3	0.32	23	1.35	21.3	0.25
Dissolved Oxygen (ppm)	7.5	0.13	7.7	0.38	7.7	0.23	6.4	0.19
Turbidity (FTU)	12.6	1.75	14.1	1.65	18.3	1.49	11	4.06
Median Depth (cm)	13.4	1.42	31.9	2.23	18.5	3.58	69.5	1.14
Maximum Depth (cm)	18.4	1.14	33.6	2.38	23.5	2.18	70.3	1.11
Minimum Depth (cm)	8.4	1.85	30.3	2.09	13.5	5.12	68.8	1.18
Time Interval: (date time)	(May 1: 1310 - 1725 hrs)				(May 6: 1147 - 1515 hrs)			

Drop samples; 2.6 m sq. each; N = 4;

APPENDIX II (continued): PHYSICAL MEASUREMENTS, SUMMER.

GALVESTON BAY STUDY ENVIRONMENTAL PARAMETERS SUMMER SAMPLING SET	Site 1 TRINITY RIVER INNER DELTA				Site 2 TRINITY RIVER OUTER DELTA			
	Vegetated		Non-vegetated		Vegetated		Non-vegetated	
	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.
Temperature (Deg. C)	31.4	0.24	31	0.41	30.4	0.69	30.8	0.48
Salinity (ppt)	0	0	0	0	0.4	0.08	0.5	0.03
Dissolved Oxygen (ppm)	7.3	0.41	7.6	0.42	9.2	0.47	9.5	0.27
Turbidity (FTU)	46.8	0.75	46	9.69	30.8	6.52	31	6.67
Median Depth (cm)	28.8	2.79	52.5	14.27	37.8	2.05	47.1	5.3
Maximum Depth (cm)	35.5	4.65	58.5	17.21	40	2.42	48	5.43
Minimum Depth (cm)	22	4.14	46.5	11.65	35.5	2.06	46.3	5.17
Time Interval: (date time)	(July 21: 1425 - 1630 hrs)				(July 21: 1115 - 1333 hrs)			
	Site 3 SMITH POINT				Site 4 MOSES LAKE			
	Vegetated		Non-vegetated		Vegetated		Non-vegetated	
	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.
Temperature (Deg. C)	31.5	0.29	31.3	0.25	26.4	2.79	28.8	0.25
Salinity (ppt)	0.8	0.03	0.7	0.02	9	0	9	0
Dissolved Oxygen (ppm)	8.6	0.46	7.9	0.25	6.8	0.76	7.5	0.42
Turbidity (FTU)	34.8	7.97	26.3	1.11	28	5.58	25.5	2.1
Median Depth (cm)	34.3	4.99	67.8	4.51	40.8	3.5	62.8	3.82
Maximum Depth (cm)	41.5	5.56	69	4.45	49	3.83	64	4.06
Minimum Depth (cm)	27	5.08	66.5	4.57	32.5	4.5	61.5	3.57
Time Interval: (date time)	(July 22: 1320 - 1450 hrs)				(July 20: 0954 - 1113 hrs)			
	Site 5 JAMAICA BEACH				Site 6 CHRISTMAS BAY			
	Vegetated		Non-vegetated		Vegetated		Non-vegetated	
	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.
Temperature (Deg. C)	31.9	0.22	32.2	0.15	31.4	1.55	30	0
Salinity (ppt)	27.9	0.13	27.8	0.14	29.5	0.5	29.3	0.48
Dissolved Oxygen (ppm)	7.3	0.21	6.7	0.32	5.7	0.48	6.3	0.55
Turbidity (FTU)	32.3	3.22	31.8	4.33	13.8	2.93	6.8	1.25
Median Depth (cm)	16.7	1.64	36.6	1.33	32.6	1.6	57	2.46
Maximum Depth (cm)	22.3	1.75	38.8	1.33	34.8	2.5	60	2.8
Minimum Depth (cm)	11.1	1.78	34.5	1.34	30.5	0.87	54	2.45
Time Interval: (date time)	(July 17: 1035 - 1347 hrs)				(July 24: 0946 - 1156 hrs)			

Drop samples; 2.6 m sq. each; N = 4;

APPENDIX II (continued): PHYSICAL MEASUREMENTS, FALL.

ENVIRONMENTAL PARAMETERS FALL SAMPLING SET	Site 1 TRINITY RIVER INNER DELTA				Site 2 TRINITY RIVER OUTER DELTA			
	Vegetated		Non-vegetated		Vegetated		Non-vegetated	
	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.
Temperature (Deg. C)	18.5	0.46	19	0.31	23	0.44	22.3	0.35
Salinity (ppt)	11	0	10.5	0.29	9.8	0.25	9.5	0.29
Dissolved Oxygen (ppm)	3.7	0.41	4.3	0.38	7.9	0.56	7.8	0.31
Turbidity (FTU)	68.8	10.08	32.8	8.37	64.8	19.6	38.3	12.56
Median Depth (cm)	8.9	1.88	25.4	8.5	5.6	0.43	35.8	4.62
Maximum Depth (cm)	12.3	2.21	27.8	10.16	9	0.71	39.8	5.07
Minimum Depth (cm)	5.5	2.1	23	6.86	2.3	0.85	31.8	4.23
Time Interval: (date time)	(November 3: 0725 - 0921 hrs)				(November 2: 1017 - 1245 hrs)			

	Site 3 SMITH POINT				Site 4 MOSES LAKE			
	Vegetated		Non-vegetated		Vegetated		Non-vegetated	
	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.
Temperature (Deg. C)	23.1	0.46	22.8	0.29	22.3	0.51	22.4	0.28
Salinity (ppt)	20	0	20	0	22	0.41	22.3	0.48
Dissolved Oxygen (ppm)	8.6	0.13	8.1	0.11	7.6	1.83	8.5	1.86
Turbidity (FTU)	90.5	38.2	51.3	13.44	111.3	18.19	67.8	14.79
Median Depth (cm)	25	3.17	44.1	1.42	18.8	2.92	30.8	2.25
Maximum Depth (cm)	29.5	2.72	45.3	1.6	36.3	5.02	32	2.48
Minimum Depth (cm)	20.5	3.66	43	1.29	1.3	1.25	29.5	2.02
Time Interval: (date time)	(November 3: 1158 - 1315 hrs)				(November 4: 0921 - 1115 hrs)			

	Site 5 JAMAICA BEACH				Site 6 CHRISTMAS BAY			
	Vegetated		Non-vegetated		Vegetated		Non-vegetated	
	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.
Temperature (Deg. C)	20.9	0.06	20.9	0.06	25.3	1.4	25.1	0.66
Salinity (ppt)	20.5	0.29	20.5	0.29	32.5	0.87	31.8	0.25
Dissolved Oxygen (ppm)	8	0.15	7.8	0.18	9.4	0.27	9.4	1.5
Turbidity (FTU)	25.9	4.71	22.9	1.59	18	2.86	26	11.8
Median Depth (cm)	22.1	1.01	46.1	2.9	17.5	2.07	22	5.94
Maximum Depth (cm)	27	0.94	48.8	3.01	18.5	1.85	24.3	5.36
Minimum Depth (cm)	17.3	1.11	43.4	3.4	16.5	2.33	19.8	6.66
Time Interval: (date time)	(October 23: 0823 - 1209 hrs)				(November 5: 1015 - 1240 hrs)			

Drop samples; 2.6 m sq. each; N = 4;

APPENDIX III: FISH AND DECAPOD CRUSTACEAN DENSITIES, UPPER BAY, SPRING.

SPECIES	Site 1				Site 2			
	TRINITY RIVER INNER DELTA				TRINITY RIVER OUTER DELTA			
	Vegetated		Non-vegetated		Vegetated		Non-vegetated	
	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.
FISHES:								
<i>Microgogonias undulatus</i>	0	0	3.5	2.02	0	0	0.8	0.75
<i>Fundulus grandis</i>	0	0	0.3	0.25	1.5	0.87	0	0
<i>Myrophis punctatus</i>	0	0	2.8	1.6	0	0	0.5	0.29
<i>Anchoa mitchilli</i>	0	0	0	0	0	0	2	1.68
<i>Leiostomus xanthurus</i>	0	0	0	0	0	0	0.5	0.5
<i>Fundulus pulvereus</i>	1.8	1.75	0	0	0	0	0	0
<i>Mugil cephalus</i>	0	0	0.8	0.75	0	0	0	0
<i>Elops saurus</i>	0	0	0.5	0.5	0	0	0	0
<i>Brevoortia patronus</i>	0	0	0.3	0.25	0	0	0	0
<i>Citharichthys spilopterus</i>	0	0	0.3	0.25	0	0	0	0
<i>Gambusia affinis</i>	0.3	0.25	0	0	0	0	0	0
<i>Paralichthys lethostigma</i>	0	0	0	0	0.3	0.25	0	0
<i>Symphurus plagiusa</i>	0	0	0.3	0.25	0	0	0	0
<i>Syngnathus floridae</i>	0	0	0.3	0.25	0	0	0	0
Cyprinodontidae	1.8	1.75	0.3	0.25	1.5	0.87	0	0
Sciaenidae	0	0	3.5	2.02	0	0	1.3	0.75
Commercial/Sports Fishes	0	0	0	0	0.3	0.25	0	0
FISH TOTALS:	2	1.68	8.8	3.04	1.8	1.03	3.8	2.46
CRUSTACEANS:								
<i>Callinectes sapidus</i>	1.3	0.48	8	1.58	1.3	0.75	1.8	1.18
<i>Palaemonetes pugio</i>	0	0	0.5	0.29	0	0	0	0
CRUSTACEAN TOTALS:	1.3	0.48	8.5	1.32	1.3	0.75	1.8	1.18

APPENDIX III (continued): FISH AND DECAPOD CRUSTACEAN DENSITIES, MIDDLE BAY, SPRING.

SPECIES	Site 3 SMITH POINT				Site 4 MOSES LAKE			
	Vegetated		Non-vegetated		Vegetated		Non-vegetated	
	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.
FISHES:								
<i>Lagodon rhomboides</i>	3.8	1.7	0	0	1	1	0	0
<i>Gobionellus boleosoma</i>	2.3	0.85	0.8	0.48	0	0	0	0
<i>Myrophis punctatus</i>	0.3	0.25	0	0	1	1	1	0.71
<i>Leiostomus xanthurus</i>	0	0	2	1.08	0	0	0	0
<i>Gobiosoma boscii</i>	1.3	1.25	0.3	0.25	0	0	0	0
<i>Mugil cephalus</i>	0	0	0.3	0.25	1	0.71	0.3	0.25
<i>Fundulus grandis</i>	0.5	0.29	0	0	0.3	0.25	0	0
<i>Symphurus plagiusa</i>	0	0	0.8	0.48	0	0	0	0
<i>Elops saurus</i>	0	0	0.5	0.5	0	0	0	0
<i>Adina xenica</i>	0	0	0	0	0.3	0.25	0	0
<i>Lucania parva</i>	0.3	0.25	0	0	0	0	0	0
<i>Menidia beryllina</i>	0	0	0	0	0	0	0.3	0.25
<i>Micropogonias undulatus</i>	0	0	0.3	0.25	0	0	0	0
<i>Paralichthys lethostigma</i>	0	0	0.3	0.25	0	0	0	0
Cyprinodontidae	0.8	0.48	0	0	0.5	0.29	0	0
Gobiidae	3.5	1.71	1	0.41	0	0	0	0
Sciaenidae	0	0	2.3	0.95	0	0	0	0
Commercial/Sports Fishes	0	0	0.3	0.25	0	0	0	0
FISH TOTALS:	8.3	3.45	5	1.08	3.5	2.02	1.5	1.19
CRUSTACEANS:								
<i>Palaemonetes pugio</i>	290.5	48.05	94.3	93.92	37.5	37.17	0	0
<i>Penaeus aztecus</i>	2.5	1.32	10	2.48	9	9	0.5	0.29
<i>Callinectes sapidus</i>	13.3	2.1	2.8	1.25	2.8	2.43	0.5	0.5
<i>Rhithropanopeus harrisi</i>	1	0.58	0.3	0.25	0	0	0	0
Penaeidae	2.5	1.32	10	2.48	9	9	0.5	0.29
CRUSTACEAN TOTALS:	307.3	51.36	107.3	93.26	49.3	48.58	1	0.41

APPENDIX III (continued): FISH AND DECAPOD CRUSTACEAN DENSITIES, LOWER BAY, SPRING.

GALVESTON BAY STUDY LOWER BAY SYSTEM Macrofauna/2.6 m sq. (n=4) May 1st & 6th, 1987 SPECIES	Site 5 JAMAICA BEACH				Site 6 CHRISTMAS BAY			
	Vegetated		Non-vegetated		Vegetated		Non-vegetated	
	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.
FISHES:								
<i>Lagodon rhomboides</i>	1.3	0.75	0.5	0.29	3	1.68	5	4.02
<i>Menidia beryllina</i>	7.3	6.92	0.5	0.29	0	0	0	0
<i>Brevoortia patronus</i>	0	0	5.8	5.11	0	0	0	0
<i>Gobionellus boleosoma</i>	0	0	0	0	2.3	0.85	0.3	0.25
<i>Leiostomus xanthurus</i>	0	0	0.5	0.29	0	0	1.5	0.65
<i>Micropogonias undulatus</i>	0	0	1	0	0	0	0.5	0.5
<i>Myrophis punctatus</i>	0	0	0.5	0.29	0.5	0.5	0.3	0.25
<i>Fundulus grandis</i>	0	0	0	0	1	0.71	0	0
<i>Mugil cephalus</i>	0	0	0	0	0.3	0.25	0.3	0.25
<i>Paralichthys lethostigma</i>	0	0	0	0	0	0	0.5	0.5
<i>Symphurus plagiosa</i>	0	0	0	0	0	0	0.5	0.5
<i>Citharichthys spilopterus</i>	0	0	0	0	0	0	0.3	0.25
<i>Dasyatis sabina</i>	0	0	0	0	0	0	0.3	0.25
<i>Orthopristis chrysoptera</i>	0	0	0	0	0	0	0.3	0.25
<i>Synodus foetens</i>	0	0	0	0	0	0	0.3	0.25
Cyprinodontidae	0	0	0	0	1	0.71	0	0
Gobiidae	0	0	0	0	2.3	0.85	0.3	0.25
Sciaenidae	0	0	1	0	0	0	2	1.08
Commercial/Sports Fishes	0	0	0	0	0	0	0.5	0.5
FISH TOTALS:	8.5	7.53	7.8	4.82	7	2.65	9.8	6.3
CRUSTACEANS:								
<i>Palaemonetes pugio</i>	15	8.36	0.5	0.29	143.3	63.73	0	0
<i>Penaeus aztecus</i>	41.5	8.37	10	1.15	39.3	8.36	7	1.87
<i>Callinectes sapidus</i>	6	2.38	1.5	0.29	9	2.58	0.3	0.25
<i>Clibanarius vittatus</i>	2.5	1.26	0.5	0.29	4.8	1.7	0	0
<i>Hippolyte zostericola</i>	0	0	0	0	1.3	1.25	0	0
<i>Pagurus spp.</i>	0	0	0	0	0	0	0.5	0.5
<i>Neopanope texana</i>	0	0	0	0	0.3	0.25	0	0
Unknown crustacean species	0	0	0	0	0	0	0.3	0.25
Penaeidae	41.5	8.37	10	1.15	39.3	8.36	7	1.87
CRUSTACEAN TOTALS:	64.8	10.62	11.8	1.32	200.3	70.63	8	1.68

APPENDIX III (continued): FISH AND DECAPOD CRUSTACEAN DENSITIES, UPPER BAY, SUMMER.

SPECIES	Site 1				Site 2			
	TRINITY RIVER				TRINITY RIVER			
	Vegetated		Non-vegetated		Vegetated		Non-vegetated	
	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.
FISHES:								
<i>Fundulus grandis</i>	6.8	2.63	0.3	0.25	2.8	0.25	0	0
<i>Mugil cephalus</i>	3	3	0.3	0.25	3.3	0.85	0.3	0.25
<i>Cyprinodon variegatus</i>	5	4.36	0	0	0	0	0	0
<i>Lucania parva</i>	2.8	2.43	0	0	0	0	0	0
<i>Anchoa mitchilli</i>	0	0	1.8	1.03	0	0	0.5	0.29
<i>Menidia beryllina</i>	1	0.41	0	0	1.3	0.75	0	0
<i>Leiostomus xanthurus</i>	0	0	0.3	0.25	0.3	0.25	0.5	0.29
<i>Conodon nobilis</i>	0.8	0.75	0	0	0	0	0	0
<i>Symphurus plagiusa</i>	0.8	0.75	0	0	0	0	0	0
<i>Myrophis punctatus</i>	0.3	0.25	0.3	0.25	0	0	0	0
<i>Brevoortia patronus</i>	0.5	0.29	0	0	0	0	0	0
<i>Fundulus jenkinsi</i>	0.5	0.29	0	0	0	0	0	0
<i>Fundulus pulvereus</i>	0.3	0.25	0	0	0.3	0.25	0	0
<i>Syngnathus scovelli</i>	0	0	0	0	0.5	0.29	0	0
<i>Citharichthys spilopterus</i>	0	0	0.3	0.25	0	0	0	0
<i>Gobionellus boleosoma</i>	0	0	0	0	0	0	0.3	0.25
<i>Gobiosoma bosci</i>	0	0	0	0	0	0	0.3	0.25
<i>Ictalurus punctatus</i>	0	0	0	0	0	0	0.3	0.25
<i>Lagodon rhomboides</i>	0.3	0.25	0	0	0	0	0	0
<i>Sciaenops ocellatus</i>	0	0	0	0	0.3	0.25	0	0
Unknown fish species	0.3	0.25	0	0	0	0	0	0
Cyprinodontidae	15.3	6.39	0.3	0.25	3	0.41	0	0
Gobiidae	0	0	0	0	0	0	0.5	0.29
Sciaenidae	0	0	0.3	0.25	0.5	0.29	0.5	0.29
Commercial/Sports Fishes	0	0	0	0	0.3	0.25	0	0
FISH TOTALS:	22	5.34	3	1.22	8.5	1.04	2	0.91
CRUSTACEANS:								
<i>Palaemonetes pugio</i>	45.8	24.35	0	0	16.3	8	0	0
<i>Callinectes sapidus</i>	2.3	1.31	1.3	0.75	4	1.47	0	0
<i>Penaeus setiferus</i>	0	0	0	0	1.3	0.75	0	0
<i>Palaemonetes vulgaris</i>	0	0	0	0	0.5	0.5	0	0
<i>Penaeus aztecus</i>	0	0	0	0	0.5	0.29	0	0
<i>Sesarma reticulatum</i>	0.5	0.29	0	0	0	0	0	0
<i>Uca pugnax</i>	0.5	0.5	0	0	0	0	0	0
<i>Neopanope texana</i>	0.3	0.25	0	0	0	0	0	0
Penaeidae	0	0	0	0	1.8	0.75	0	0
CRUSTACEAN TOTALS:	49.3	27.78	1.3	0.75	22.5	7.51	0	0

APPENDIX III (continued): FISH AND DECAPOD CRUSTACEAN DENSITIES, MIDDLE BAY, SUMMER.

GALVESTON BAY STUDY MID-BAY SYSTEM Macrofauna/2.6 m sq. (n=4) July 20 & 22, 1987 SPECIES	Site 3 SMITH POINT				Site 4 MOSES LAKE			
	Vegetated		Non-vegetated		Vegetated		Non-vegetated	
	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.
FISHES:								
<i>Gobiosoma boscii</i>	32.8	9.31	0.8	0.48	9.5	5.84	3.3	2.93
<i>Anchoa mitchilli</i>	1.8	1.75	30.5	13.99	0	0	7	6.04
<i>Myrophis punctatus</i>	0.3	0.25	1.3	0.63	2.8	2.14	0.5	0.29
<i>Mugil cephalus</i>	1	0.58	0	0	2.8	1.31	0	0
<i>Lagodon rhomboides</i>	1.3	0.95	0	0	1.3	0.25	0	0
<i>Brevoortia patronus</i>	0	0	0	0	0	0	2.3	2.25
<i>Fundulus grandis</i>	0.8	0.75	0	0	1.5	0.87	0	0
<i>Cynoscion nebulosus</i>	1.5	0.29	0	0	0.5	0.5	0	0
<i>Menidia beryllina</i>	0.3	0.25	0	0	1.5	1.19	0	0
<i>Syngnathus scovelli</i>	0	0	0	0	1.3	1.25	0	0
<i>Cyprinodon variegatus</i>	0	0	0	0	1	0.71	0	0
<i>Oligoplites saurus</i>	0.8	0.48	0	0	0	0	0	0
<i>Paralichthys lethostigma</i>	0.3	0.25	0.3	0.25	0	0	0	0
<i>Membras martinica</i>	0	0	0.5	0.5	0	0	0	0
Unknown fish species	0.5	0.5	0	0	0	0	0	0
<i>Arius felis</i>	0	0	0.3	0.25	0	0	0	0
<i>Citharichthys spilopterus</i>	0	0	0.3	0.25	0	0	0	0
<i>Gobiesox strumosus</i>	0.3	0.25	0	0	0	0	0	0
<i>Hyporhamphus unifasciatus</i>	0.3	0.25	0	0	0	0	0	0
<i>Lucaania parva</i>	0	0	0	0	0.3	0.25	0	0
<i>Sciaenops ocellatus</i>	0.3	0.25	0	0	0	0	0	0
<i>Sphoeroides parvus</i>	0	0	0	0	0	0	0.3	0.25
<i>Stellifer lanceolatus</i>	0.3	0.25	0	0	0	0	0	0
Cyprinodontidae	0.8	0.75	0	0	2.8	1.11	0	0
Gobiidae	32.8	9.31	0.8	0.48	9.5	5.84	3.3	2.93
Sciaenidae	2	0.41	0	0	0.5	0.5	0	0
Commercial/Sports Fishes	2	0	0.3	0.25	0.5	0.5	0	0
FISH TOTALS:	42	10.97	33.7	13.92	22.3	7.11	13.3	10.97
CRUSTACEANS:								
<i>Palaemonetes pugio</i>	590	167.07	0.3	0.25	242	16.52	0.3	0.25
<i>Penaeus setiferus</i>	79	41.29	4.3	1.75	0.5	0.29	1.3	0.63
<i>Penaeus aztecus</i>	33.8	13.85	7.5	0.5	2.3	0.63	0.8	0.48
<i>Callinectes sapidus</i>	10.8	4.27	2.3	1.31	6.3	1.03	0.8	0.48
<i>Rhithropanopeus harrisi</i>	3.3	1.97	0.5	0.5	0.3	0.25	0	0
<i>Palaemonetes vulgaris</i>	1.3	0.95	0	0	0.5	0.5	0	0
<i>Uca pugnax</i>	1.8	1.75	0	0	0	0	0	0
<i>Neopanope texana</i>	1.3	1.25	0	0	0	0	0	0
<i>Palaemonetes intermedius</i>	0	0	0	0	1	0.41	0	0
<i>Eurypanopeus depressus</i>	0.3	0.25	0	0	0	0	0	0
Penaeidae	112.5	53.16	11.8	2.17	2.8	0.48	2	0.71
CRUSTACEAN TOTALS:	721	183.37	14.8	3.77	252.8	17.76	3	0.58

APPENDIX III (continued): FISH AND DECAPOD CRUSTACEAN DENSITIES, LOWER BAY, SUMMER.

LOWER BAY SYSTEM Macrofauna/2.6 m sq. (n=4). July 17 & 24, 1987 SPECIES	Site 5 JAMAICA BEACH				Site 6 CHRISTMAS BAY			
	Vegetated		Non-vegetated		Vegetated		Non-vegetated	
	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.
FISHES:								
<i>Cyprinodon variegatus</i>	0	0	0	0	8.5	8.5	0	0
<i>Lagodon rhomboides</i>	2	0.91	0.5	0.5	2	1	0.5	0.29
<i>Eucinostomus spp.</i>	0	0	0	0	0	0	4.3	4.25
<i>Gobiosoma boscii</i>	3.5	1.44	0.5	0.29	0.3	0.25	0	0
<i>Anchoa mitchilli</i>	0	0	1.8	0.75	0	0	0	0
<i>Menidia beryllina</i>	0.3	0.25	1.3	0.48	0.3	0.25	0	0
<i>Mugil cephalus</i>	0	0	0.5	0.29	1.3	0.95	0	0
<i>Cynoscion nebulosus</i>	0.5	0.29	0.5	0.29	0.5	0.29	0	0
<i>Symphurus plagiusa</i>	0	0	0.5	0.29	0.3	0.25	0.5	0.5
<i>Adinia xenica</i>	0	0	0	0	1	0.71	0	0
<i>Fundulus grandis</i>	1	0.41	0	0	0	0	0	0
<i>Syngnathus scovelli</i>	1	0.58	0	0	0	0	0	0
<i>Fundulus similis</i>	0.5	0.29	0	0	0	0	0	0
<i>Microgobius thalassinus</i>	0	0	0.5	0.29	0	0	0	0
<i>Opsanus beta</i>	0.5	0.29	0	0	0	0	0	0
<i>Archosargus probatocephalus</i>	0	0	0.3	0.25	0	0	0	0
<i>Chaetodipterus faber</i>	0	0	0.3	0.25	0	0	0	0
<i>Citharichthys spilopterus</i>	0	0	0	0	0	0	0.3	0.25
<i>Eucinostomus argenteus</i>	0	0	0	0	0	0	0.3	0.25
<i>Gambusia affinis</i>	0.3	0.25	0	0	0	0	0	0
<i>Leiostomus xanthurus</i>	0	0	0.3	0.25	0	0	0	0
<i>Myrophis punctatus</i>	0	0	0.3	0.25	0	0	0	0
<i>Paralichthys lethostigma</i>	0	0	0	0	0	0	0.3	0.25
<i>Syngnathus louisianae</i>	0.3	0.25	0	0	0	0	0	0
Unknown fish species	0	0	0	0	0	0	0.3	0.25
Cyprinodontidae	1.3	0.63	0	0	9.5	8.19	0	0
Gobiidae	3.5	1.44	1	0.41	0.3	0.25	0	0
Sciaenidae	0.5	0.29	0.5	0.29	0.5	0.29	0	0
Commercial/Sports Fishes	0.5	0.29	0.5	0.29	0.5	0.29	0.3	0.25
FISH TOTALS:	8.5	2.53	5.5	0.65	14	7.01	6.3	3.92
CRUSTACEANS:								
<i>Palaemonetes pugio</i>	180.3	39.73	0.5	0.29	70.3	16.24	0.5	0.5
<i>Penaeus aztecus</i>	41.5	6.24	8.3	2.02	5.3	2.02	0.8	0.48
<i>Callinectes sapidus</i>	27.8	2.29	1.8	0.63	2.8	1.8	3	0.71
<i>Penaeus setiferus</i>	12	3.49	5.8	2.25	2.8	0.48	0	0
<i>Penaeus duorarum</i>	12.3	3.09	1.5	0.87	1.3	0.63	0	0
<i>Alpheus heterochaelis</i>	6.5	4.63	0	0	0	0	0	0
<i>Clibanarius vittatus</i>	4	1.22	0	0	1	0.58	0.8	0.48
<i>Palaemonetes intermedius</i>	3.3	2.59	0	0	1.8	1.44	0	0
<i>Uca minax</i>	0	0	0	0	1.8	0.85	0	0
<i>Neopanope texana</i>	0.3	0.25	0	0	0.5	0.5	0	0
<i>Petrolisthes armatus</i>	0.8	0.75	0	0	0	0	0	0
<i>Palaemonetes vulgaris</i>	0.3	0.25	0.3	0.25	0	0	0	0
<i>Eurypanopeus abbreviatus</i>	0.5	0.5	0	0	0	0	0	0
<i>Libinia dubia</i>	0.3	0.25	0	0	0	0	0	0
<i>Menippe mercenaria</i>	0	0	0	0	0.3	0.25	0	0
<i>Panopeus herbstii</i>	0	0	0	0	0.3	0.25	0	0
Unknown crustacean species	0.3	0.25	0	0	0	0	0	0
Penaeidae	65.5	6.69	15	3.03	9.3	2.29	0.8	0.48
CRUSTACEAN TOTALS:	287.5	39.89	17	3.08	87.8	20.5	5	1.41

APPENDIX III (continued): FISH AND DECAPOD CRUSTACEAN DENSITIES, UPPER BAY, FALL

SPECIES	Site 1				Site 2			
	TRINITY RIVER				TRINITY RIVER			
	Vegetated		Non-vegetated		Vegetated		Non-vegetated	
GALVESTON BAY STUDY								
UPPER BAY SYSTEM								
Macrofauna/2.6 m sq. (n = 4)								
November 2-3, 1987								
	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.
FISHES:								
<i>Cyprinodon variegatus</i>	31.5	31.5	1	0.71	0	0	0	0
<i>Anchoa mitchilli</i>	0	0	0	0	0	0	6.8	6.75
<i>Fundulus grandis</i>	0.8	0.48	0	0	3.5	3.18	1	1
<i>Lucania parva</i>	0.5	0.29	0.3	0.25	0	0	0	0
<i>Cynoscion nebulosus</i>	0	0	0	0	0	0	0.5	0.5
<i>Fundulus jenkinsi</i>	0.5	0.5	0	0	0	0	0	0
<i>Mugil cephalus</i>	0	0	0	0	0	0	0.5	0.5
<i>Fundulus pulvereus</i>	0	0	0	0	0.3	0.25	0	0
Cyprinodontidae	33.3	31.59	1.3	0.63	3.8	3.12	1	1
Sciaenidae	0	0	0	0	0	0	0.5	0.5
Commercial/Sports Fishes	0	0	0	0	0	0	0.5	0.5
FISH TOTALS:	33.3	31.59	1.3	0.63	3.8	3.12	8.8	6.37
CRUSTACEANS:								
<i>Palaemonetes pugio</i>	0.5	0.29	0	0	68.3	18.67	0.3	0.25
<i>Callinectes sapidus</i>	0.3	0.25	3.3	2.29	7.8	2.25	11.3	1.7
<i>Penaeus duorarum</i>	0	0	0	0	7	5.7	0.3	0.25
<i>Penaeus aztecus</i>	0	0	0.5	0.29	1.3	0.95	0.8	0.25
<i>Neopanope texana</i>	0	0	0	0	0.5	0.29	0.5	0.29
<i>Penaeus setiferus</i>	0	0	0	0	0	0	0.8	0.48
<i>Rhithropanopeus harrisi</i>	0	0	0	0	0.3	0.25	0	0
Penaeidae	0	0	0.5	0.29	8.3	6.64	1.8	0.63
CRUSTACEANS TOTALS:	0.8	0.48	3.8	2.5	85	26.71	13.8	1.18

APPENDIX III (continued): FISH AND DECAPOD CRUSTACEAN DENSITIES, MIDDLE BAY, FALL.

GALVESTON BAY STUDY MIDDLE BAY SYSTEM Macrofauna/2.6 m sq. (n=4) November 3-4, 1987 SPECIES	Site 3 SMITH POINT				Site 4 MOSES LAKE			
	Vegetated		Non-vegetated		Vegetated		Non-vegetated	
	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.
FISHES:								
<i>Gobiosoma boscii</i>	6.3	1.44	0	0	106.8	18.75	1.3	0.75
<i>Symphurus plagiatus</i>	7	2.68	6	2.8	0	0	0	0
<i>Anchoa mitchilli</i>	0	0	0	0	0.8	0.75	9.5	8.51
<i>Menidia beryllina</i>	0	0	0	0	1	1	7.3	6.6
<i>Myrophis punctatus</i>	0	0	0.3	0.25	4.5	0.87	0.5	0.29
<i>Fundulus grandis</i>	0.8	0.48	0	0	3.3	1.38	0	0
<i>Cynoscion nebulosus</i>	1.3	0.63	0	0	0.5	0.5	0	0
<i>Micropogonias undulatus</i>	0.3	0.25	0.8	0.75	0	0	0.5	0.5
<i>Sciaenops ocellatus</i>	0	0	1	0.41	0	0	0	0
<i>Gobionellus boleosoma</i>	0	0	0.8	0.75	0	0	0	0
<i>Gobiosox strumosus</i>	0.5	0.5	0	0	0	0	0	0
<i>Syngnathus louisianae</i>	0.5	0.5	0	0	0	0	0	0
<i>Gobiosoma robustum</i>	0	0	0	0	0.3	0.25	0.3	0.25
<i>Microgobius thalassinus</i>	0	0	0.3	0.25	0	0	0	0
<i>Mugil cephalus</i>	0	0	0	0	0.3	0.25	0	0
<i>Opsanus beta</i>	0	0	0	0	0.3	0.25	0.3	0.25
<i>Paralichthys lethostigma</i>	0	0	0	0	0	0	0	0
<i>Sphoeroides parvus</i>	0	0	0	0	0	0	0.3	0.25
<i>Syngnathus scovelli</i>	0	0	0	0	0.3	0.25	0	0
Unknown fish species	0	0	0	0	0.3	0.25	0	0
Cyprinodontidae	0.8	0.48	0	0	3.3	1.38	0	0
Gobiidae	6.3	1.44	1	0.71	107	18.57	1.3	0.75
Sciaenidae	1.5	0.5	1.8	0.48	0.5	0.5	0.5	0.5
Commercial/Sports Fishes	1.3	0.63	1	0.41	0.5	0.5	0.3	0.25
FISH TOTALS:	16.5	3.86	9	2.94	118	19.01	19.8	9.46
CRUSTACEANS:								
<i>Palaemonetes pugio</i>	593.8	35.73	0.3	0.25	417	89.09	1.3	0.75
<i>Callinectes sapidus</i>	57.5	9.4	11	3.39	269.3	55.61	31.3	17.63
<i>Palaemonetes vulgaris</i>	88.3	59.49	0	0	32.5	17.96	0	0
<i>Palaemonetes intermedius</i>	32	32	0	0	22	7.82	0	0
<i>Penaeus duorarum</i>	10	4.26	1.5	0.87	35.3	14.64	2.8	0.95
<i>Penaeus aztecus</i>	8.3	4.61	0.3	0.25	5.8	3.84	5.3	1.44
<i>Penaeus setiferus</i>	4	1.68	7.3	1.49	0.3	0.25	1.5	0.87
<i>Neopanope texana</i>	6.5	6.5	0	0	0.5	0.5	0	0
<i>Rhithropanopeus harrisi</i>	3	2.38	0	0	0.8	0.75	0	0
Xanthidae, unknown species	1	1	0	0	0	0	0	0
<i>Eurypanopeus abbreviatus</i>	0	0	0	0	0.5	0.5	0	0
<i>Eurypanopeus depressus</i>	0.5	0.5	0	0	0	0	0	0
<i>Panopeus herbstii</i>	0.5	0.5	0	0	0	0	0	0
<i>Alpheus heterochaelis</i>	0	0	0	0	0.3	0.25	0	0
<i>Menippe mercenaria</i>	0	0	0	0	0.3	0.25	0	0
<i>Uca rapax</i>	0	0	0	0	0.3	0.25	0	0
Penaeidae	22.3	10.26	9	1.22	41.3	12.51	9.5	2.4
CRUSTACEAN TOTALS:	805.3	108.13	20.3	3.68	784.5	70.91	42	19.73

APPENDIX III (continued): FISH AND DECAPOD CRUSTACEAN DENSITIES, LOWER BAY, FALL.

SPECIES	Site 5				Site 6			
	JAMAICA BEACH		Non-vegetated		CHRISTMAS BAY		Non-vegetated	
	Vegetated	Non-vegetated	Vegetated	Non-vegetated	Vegetated	Non-vegetated	Vegetated	Non-vegetated
	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.
FISHES:								
<i>Gobionellus boleosoma</i>	1	0.41	0.3	0.25	19.8	11.88	1.5	0.96
<i>Symphurus plagiosa</i>	2.3	0.75	1.3	0.25	1	0.71	1.5	0.29
<i>Anchoa mitchilli</i>	0	0	4.3	2.21	0.3	0.25	0	0
<i>Gobiosoma boscii</i>	3	1.58	0.3	0.25	0	0	0	0
<i>Fundulus grandis</i>	1.5	0.87	0	0	0.8	0.48	0	0
<i>Syngnathus scovelli</i>	0	0	0	0	1.8	0.75	0	0
<i>Cynoscion nebulosus</i>	1.3	0.25	0	0	0	0	0	0
<i>Cyprinodon variegatus</i>	0	0	0	0	1.3	0.75	0	0
<i>Menidia beryllina</i>	0.3	0.25	0.3	0.25	0	0	0	0
<i>Microgobius thalassinus</i>	0	0	0.5	0.5	0	0	0	0
<i>Mugil cephalus</i>	0	0	0	0	0	0	0.5	0.5
<i>Achirus lineatus</i>	0	0	0	0	0	0	0.3	0.25
<i>Eucinostomus spp.</i>	0	0	0	0	0	0	0.3	0.25
<i>Fundulus pulvereus</i>	0.3	0.25	0	0	0	0	0	0
<i>Lagodon rhomboides</i>	0	0	0	0	0	0	0.3	0.25
<i>Leiostomus xanthurus</i>	0	0	0.3	0.25	0	0	0	0
<i>Paralichthys lethostigma</i>	0	0	0.3	0.25	0	0	0	0
<i>Sciaenops ocellatus</i>	0	0	0.3	0.25	0	0	0	0
<i>Trinectes maculatus</i>	0	0	0	0	0	0	0.3	0.25
Cyprinodontidae	1.5	0.87	0	0	2	1.15	0	0
Gobiidae	3.8	1.55	1	0.71	19.8	11.88	1.5	0.96
Sciaenidae	1.3	0.25	0.3	0.25	0	0	0	0
Commercial/Sports Fishes	1.3	0.25	0.5	0.29	0	0	0	0
FISH TOTALS:	8	1.87	6.5	2.02	24.8	12.75	4.5	1.85
CRUSTACEANS:								
<i>Palaemonetes pugio</i>	42.8	7.36	0	0	212.5	76.32	0.5	0.29
<i>Callinectes sapidus</i>	41.5	6.51	4.8	0.25	32	14.46	3	1.08
<i>Penaeus setiferus</i>	7.3	1.11	0.8	0.48	36	22.87	2	2
<i>Palaemonetes vulgaris</i>	23	12.77	0	0	18.5	18.5	0	0
<i>Penaeus duorarum</i>	17.5	4.84	2.3	0.63	17.5	5.66	1.8	0.25
<i>Penaeus aztecus</i>	16.3	5.07	2	0.71	11.8	9.17	0.8	0.25
<i>Palaemonetes intermedius</i>	2.5	0.87	0	0	11	7.08	0	0
<i>Clibanarius vittatus</i>	2.8	0.48	0.3	0.25	7.5	3.01	0.5	0.5
<i>Alpheus heterochaelis</i>	0.3	0.25	0	0	7.8	7.42	0	0
<i>Sesarma cinereum</i>	0.8	0.75	0	0	0	0	0	0
<i>Petrolisthes armatus</i>	0.5	0.5	0	0	0	0	0	0
<i>Rhithropanopeus harrissi</i>	0	0	0	0	0.5	0.5	0	0
<i>Panopeus herbstii</i>	0	0	0	0	0.3	0.25	0	0
<i>Pinnixa chaetoptera</i>	0.3	0.25	0	0	0	0	0	0
<i>Sesarma reticulatum</i>	0.3	0.25	0	0	0	0	0	0
<i>Uca spp.</i>	0.3	0.25	0	0	0	0	0	0
Penaeidae	40.8	6.84	4.8	1.03	65.3	22.84	4.5	2.18
CRUSTACEANS TOTALS:	153.3	28.25	9.5	1.04	355.3	84.05	8.5	1.19

APPENDIX IV: EPIFAUNA AND INFAUNA DENSITIES, SPRING.

GALVESTON BAY MARSH STUDY		SITE/HABITAT				SITE/HABITAT			
Epi-Infaua/78.5 cm sq. (n=4)									
April 20 - May 6, 1987									
UPPER BAY:									
Taxonomic Group	TRINITY RIVER OUTER DELTA				TRINITY RIVER INNER DELTA				
	Vegetated		Non-vegetated		Vegetated		Non-vegetated		
	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.	
Annelids	148	29.819	63.5	37.529	133.5	57.77	86.25	22.103	
Crustaceans	0	0	0	0	0.5	0.5	0.25	0.25	
Molluscs	0	0	3.5	2.843	0.5	0.289	1.25	1.25	
Others	8.75	2.594	25.5	23.514	4.25	2.016	4.25	3.924	
Totals	156.75	31.006	92.5	61.907	138.75	60.401	92	25.72	
MIDDLE BAY:									
Taxonomic Group	SMITH POINT				MOSES LAKE				
	Vegetated		Non-vegetated		Vegetated		Non-vegetated		
	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.	
Annelids	46	14.071	78.25	32.294	282	39.684	182	36.681	
Crustaceans	218.5	85.927	2.25	0.629	1193.5	261.25	1302.25	169.075	
Molluscs	3.25	3.25	1.25	0.25	0.75	0.75	0	0	
Others	4.5	2.843	3	1.225	48.25	32.281	22.75	7.565	
Totals	272.25	96.28	84.75	32.255	1524.5	285.066	1507	190.195	
LOWER BAY:									
Taxonomic Group	JAMAICA BEACH				CHRISTMAS BAY				
	Vegetated		Non-vegetated		Vegetated		Non-vegetated		
	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.	
Annelids	144.5	42.822	65.25	17.983	150.5	41.242	17.5	3.329	
Crustaceans	483.25	211.775	72.5	18.554	16.5	6.958	3	1.08	
Molluscs	0.5	0.5	1	0.707	0.25	0.25	7.25	3.683	
Others	23.25	18.346	3.75	2.496	2	1.354	0.25	0.25	
Totals	651.5	249.324	142.5	16.983	169.25	49.123	29	4.262	

APPENDIX IV (continued): EPIFAUNA AND INFAUNA DENSITIES, SUMMER.

GALVESTON BAY MARSH STUDY
Epi-Infaua/78.5 cm sq. (n=4)
July 17 - 24, 1987

Taxonomic Group	TRINITY RIVER OUTER DELTA				TRINITY RIVER INNER DELTA			
	Vegetated		Non-vegetated		Vegetated		Non-vegetated	
	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.
Annelids	166.75	43.991	91	19.101	381.75	78.742	140.75	47.776
Crustaceans	1.75	1.75	0	0	0.25	0.25	0	0
Molluscs	0.5	0.5	0.75	0.479	6	3.894	23	22.668
Others	3.5	1.19	2	1.08	36	18.353	5.75	3.449
Totals	172.5	43.963	93.75	18.277	424	99.25	169.5	72.882

Taxonomic Group	SMITH POINT				MOSES LAKE			
	Vegetated		Non-vegetated		Vegetated		Non-vegetated	
	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.
Annelids	28	13.681	34.75	12.652	183.75	133.982	185.75	98.68
Crustaceans	60.5	36.999	4.25	3.591	98	87.358	29.75	17.853
Molluscs	0.5	0.289	1.75	0.75	0.5	0.289	0	0
Others	1.5	0.5	1.25	0.946	15	13.385	3	2.041
Totals	90.5	48.086	42.25	15.971	297.5	234.225	218.5	116.963

Taxonomic Group	JAMAICA BEACH				CHRISTMAS BAY			
	Vegetated		Non-vegetated		Vegetated		Non-vegetated	
	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.
Annelids	131	56.254	120.75	55.253	116.5	51.745	43.5	14.086
Crustaceans	4.25	1.25	7.75	2.136	28.25	26.597	0.5	0.5
Molluscs	0	0	1.75	0.25	0	0	1.25	0.946
Others	7	7	3.25	2.926	3.75	3.75	1	0.707
Totals	142.25	53.4	133.5	55.468	148.5	77.881	46.25	15.451

APPENDIX IV (continued): EPIFAUNA AND INFAUNA DENSITIES, FALL

GALVESTON BAY MARSH STUDY

Epi-Infauuna/78.5 cm sq. (n=4)

October 23 - November 5, 1987

UPPER BAY:

Taxonomic Group	TRINITY RIVER OUTER DELTA				TRINITY RIVER INNER DELTA			
	Vegetated		Non-vegetated		Vegetated		Non-vegetated	
	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.
Annelids	192.25	20.621	63	14.566	305	36.03	221.25	8.938
Crustaceans	0.75	0.479	1.5	0.866	0.5	0.289	0	0
Molluscs	4	3.674	0	0	0.25	0.25	8.25	4.973
Others	3	2.345	1.75	1.031	2	0.816	5.5	3.227
Totals	200	24.742	66.25	13.937	307.75	36.954	235	10.48

MIDDLE BAY:

Taxonomic Group	SMITH POINT				MOSES LAKE			
	Vegetated		Non-vegetated		Vegetated		Non-vegetated	
	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.
Annelids	6.25	2.056	49.5	7.577	241.5	87.463	125	59.611
Crustaceans	2	1.414	6	3.83	32.75	7.307	52.5	20.234
Molluscs	0	0	1.25	0.479	0.25	0.25	0.5	0.5
Others	0.75	0.479	1.25	1.25	6.5	3.428	2.25	1.652
Totals	9	1.414	58	6.671	281	92.416	180.25	78.715

LOWER BAY:

Taxonomic Group	JAMAICA BEACH				CHRISTMAS BAY			
	Vegetated		Non-vegetated		Vegetated		Non-vegetated	
	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.
Annelids	78	32.357	102.25	39.205	109.25	24.178	43.5	11.701
Crustaceans	4	1.581	7	3.674	3.75	1.652	0.75	0.479
Molluscs	0	0	0.25	0.25	0.75	0.479	2	1.225
Others	0.5	0.289	1.5	0.5	0.75	0.479	4.25	3.924
Totals	82.5	33.908	111	40.663	114.5	23.869	50.5	14.192

APPENDIX V: FISH AND DECAPOD CRUSTACEAN DENSITIES AT SITES WITH SAV HABITAT.
GALVESTON BAY STUDY

SPECIES	SPRING				SUMMER				FALL					
	SITE 2 OUTER DELTA		SITE 6 CHRISTMAS BAY		SITE 1 INNER DELTA		SITE 2 OUTER DELTA		SITE 6 CHRISTMAS BAY		SITE 2 OUTER DELTA		SITE 6 CHRISTMAS BAY	
	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.	MEAN	S.E.
FISH:														
<i>Gobionellus boleosoma</i>	0	0	10.5	1.443	0	0	0	0	1	0.707	0	0	42	12.871
<i>Lagodon rhomboides</i>	0	0	27	5.845	0	0	0	0	6.5	1.848	0	0	0.25	0.25
<i>Cyprinodon variegatus</i>	0.25	0.25	0	0	26	3.24	0	0	0	0	0	0	0	0
<i>Lucania parva</i>	0	0	0	0	18	3.028	0	0	0	0	0	0	0	0
<i>Gobiosoma robustum</i>	0	0	1	1	0	0	0	0	1.5	1.5	0	0	7	4.123
<i>Gobiosoma bosci</i>	0	0	2	2	0	0	0.25	0.25	5	2.345	0	0	2	1.08
<i>Symphurus plagiusa</i>	0	0	0.5	0.289	0	0	0	0	1.5	0.957	0	0	2	0.913
<i>Syngnathus scovelli</i>	0	0	0	0	0	0	0	0	1	0.408	0	0	2.5	0.645
<i>Fundulus grandis</i>	0.75	0.75	0	0	0	0	0	0	0	0	2.25	1.315	0	0
<i>Myrophis punctatus</i>	0	0	0.75	0.479	0.5	0.5	1.25	0.479	0	0	0	0	0.25	0.25
<i>Anchoa mitchilli</i>	0.25	0.25	0	0	0	0	0	0	0.75	0.75	1.25	1.25	0	0
<i>Cynoscion nebulosus</i>	0	0	0.75	0.75	0	0	0	0	0.5	0.289	0	0	1	0.707
<i>Micropogonias undulatus</i>	2	0.707	0	0	0	0	0	0	0	0	0	0	0.25	0.25
<i>Bairdiella chrysoura</i>	0	0	0	0	0	0	0	0	1.5	0.645	0	0	0	0
<i>Leiostomus xanthurus</i>	0.75	0.75	0	0	0	0	0	0	0.25	0.25	0	0	0.25	0.25
<i>Adina xenica</i>	0	0	0	0	0	0	0	0	0	0	0.25	0.25	0	0
<i>Arius felis</i>	0	0	0	0	0	0	0	0	0.25	0.25	0	0	0	0
<i>Menidia beryllina</i>	0	0	0	0	0	0	0	0	0	0	0	0	0.25	0.25
<i>Opsanus beta</i>	0	0	0	0	0	0	0	0	0.25	0.25	0	0	0	0
<i>Synodus foetens</i>	0	0	0.25	0.25	0	0	0	0	0	0	0	0	0	0
Cyprinodontidae	1	1	0	0	44	3.7193	0	0	0	0	2.5	1.1902	0	0
Gobiidae	0	0	13.5	2.0207	0	0	0.25	0.25	7.5	2.3274	0	0	51	12.0623
Sciaenidae	2.75	1.0308	0.75	0.75	0	0	0	0	2.25	0.4787	0	0	1.5	0.5
Bait Fishes	0.25	0.25	27	5.8452	0	0	0	0	7.25	2.1747	1.25	1.25	0.25	0.25
Commercial/Sports Fishes	0	0	0.75	0.75	0	0	0	0	0.5	0.2887	0	0	1	0.7071
FISH TOTALS:	4		42.75		44.5		1.5		20		3.75		57.75	
CRUSTACEANS:														
<i>Palaemonetes pugio</i>	0.25	0.25	38.25	6.237	0.5	0.289	0.25	0.25	55	16.558	9.5	8.17	139.75	57.469
<i>Callinectes sapidus</i>	0.75	0.25	6.25	1.031	1.75	0.629	0.75	0.25	7.75	2.496	15.5	3.403	45.5	5.545
<i>Penaeus aztecus</i>	0	0	40	5.672	0	0	0	0	28.5	3.884	1.25	0.75	8.25	3.092
<i>Penaeus duorarum</i>	0	0	0	0	0	0	0	0	11	3.808	1	0.408	51	12.537
<i>Palaemonetes vulgaris</i>	0	0	0	0	0	0	0	0	0	0	0	0	33.5	13.865
<i>Hippolyte zostericola</i>	0	0	6.25	2.529	0	0	0	0	5.25	2.72	0	0	8.5	5.545
<i>Alpheus heterochaelis</i>	0	0	0	0	0	0	0	0	4.5	2.533	0	0	9.75	2.75
<i>Palaemonetes intermedius</i>	0	0	0.5	0.5	0	0	0	0	2.75	2.428	0	0	11	3.189
<i>Penaeus setiferus</i>	0	0	0	0	0	0	0	0	1.75	0.479	1.25	0.75	5.25	3.924
<i>Tozeuma carolinense</i>	0	0	0	0	0	0	0	0	0	0	0	0	2.5	2.179
<i>Rhithropanopeus harrissi</i>	0	0	0.75	0.479	0	0	0.25	0.25	0	0	0.75	0.75	0	0
<i>Clibanarius vittatus</i>	0	0	0	0	0	0	0	0	0	0	0	0	1.25	0.75
<i>Neopanope texana</i>	0	0	0	0	0	0	0	0	0	0	1.25	0.946	0	0
<i>Panopeus turgidus</i>	0	0	0	0	0	0	0	0	0.75	0.75	0	0	0.25	0.25
<i>Panopeus herbstii</i>	0	0	0	0	0	0	0	0	0	0	0.75	0.75	0	0
Grass Shrimp	0.25	0.25	38.75	5.9214	0.5	0.2887	0.25	0.25	57.75	18.9225	9.5	8.1701	184.25	66.1279
Penaeid Shrimp	0	0	40	5.6716	0	0	0	0	41.25	5.391	3.5	1.0408	64.5	8.5878
CRUSTACEAN TOTALS:	1	0.4082	92	12.4833	2.25	0.8539	1.25	0.25	117.25	22.9578	31.25	9.4989	316.5	61.0389