


Cell Chemical Biology

Shipworm symbiosis ecology-guided discovery of an antibiotic that kills colistin-resistant *Acinetobacter*

Graphical abstract



Authors

Bailey W. Miller, Albebson L. Lim, Zhenjian Lin, ..., Colin Manoil, Eric W. Schmidt, Margo G. Haygood

Correspondence

ews1@utah.edu (E.W.S.),
margo.haygood@utah.edu (M.G.H.)

In brief

The symbiotic bacteria associated with marine shipworms are prolific producers of bioactive metabolites. Miller et al. use an ecology-guided approach to isolate the turnercyclamycins, proposed to be important for this association.

Turnercyclamycins are described as a group of lipopeptide antibiotics with selective activity against Gram-negative pathogens, including drug-resistant *Acinetobacter*.

Highlights

- Microbial competition in a symbiosis leading to antibiotics
- Group of lipopeptide antibiotics that targets drug-resistant *Acinetobacter*



Article

Shipworm symbiosis ecology-guided discovery of an antibiotic that kills colistin-resistant *Acinetobacter*

Bailey W. Miller,¹ Albebson L. Lim,¹ Zhenjian Lin,¹ Jeannie Bailey,² Kari L. Aoyagi,³ Mark A. Fisher,³ Louis R. Barrows,⁴ Colin Manoil,² Eric W. Schmidt,^{1,5,*} and Margo G. Haygood^{1,*}

¹Department of Medicinal Chemistry, University of Utah, Salt Lake City, UT 81112, USA

²Department of Genome Sciences, University of Washington, Seattle, WA 98195, USA

³Department of Pathology and ARUP Laboratories, University of Utah, Salt Lake City, UT 84112, USA

⁴Department of Pharmacology and Toxicology, University of Utah, Salt Lake City, UT 84112, USA

⁵Lead contact

*Correspondence: ews1@utah.edu (E.W.S.), margo.haygood@utah.edu (M.G.H.)

<https://doi.org/10.1016/j.chembiol.2021.05.003>

SUMMARY

Teredinibacter turnerae is an intracellular bacterial symbiont in the gills of wood-eating shipworms, where it is proposed to use antibiotics to defend itself and its animal host. Several biosynthetic gene clusters are conserved in *T. turnerae* and their host shipworms around the world, implying that they encode defensive compounds. Here, we describe turnercyclamycins, lipopeptide antibiotics encoded in the genomes of all sequenced *T. turnerae* strains. Turnercyclamycins are bactericidal against challenging Gram-negative pathogens, including colistin-resistant *Acinetobacter baumannii*. Phenotypic screening identified the outer membrane as the likely target. Turnercyclamycins and colistin operate by similar cellular, although not necessarily molecular, mechanisms, but turnercyclamycins kill colistin-resistant *A. baumannii*, potentially filling an urgent clinical need. Thus, by exploring environments that select for the properties we require, we harvested the fruits of evolution to discover compounds with potential to target unmet health needs. Investigating the symbionts of shipworms is a powerful example of this principle.

INTRODUCTION

Wood-eating shipworms (Teredinidae), a family of marine bivalve mollusks, have a close symbiotic relationship with the intracellular bacteria living in their gills. These bacteria, largely represented by the gamma-proteobacterial genus *Teredinibacter*, support their hosts both through the fixation of atmospheric nitrogen and by contributing cellulolytic enzymes necessary for wood degradation (Distel et al., 2002; Ekborg et al., 2007; Lechene et al., 2007). While these microbes are located in bacteriocytes within the gill filaments of the shipworm, their enzymes are transported to and can be detected in the cecum, where wood is digested (O'Connor et al., 2014). Interestingly, this organ has been shown to be nearly devoid of bacterial inhabitants, despite a presumed abundance of nutritionally accessible carbon released by digested lignocellulose (Betcher et al., 2012). Thus, in contrast to other wood-eating symbioses wherein the microbes consume energy-rich glucose and secrete lower-energy metabolites, this strategy allows the host shipworm to directly consume glucose by hosting the bacteria in the gills and not in the digestive organ. To make this remarkable lifestyle possible, it has been hypothesized that shipworm symbionts secrete potent antibiotics into the host, where they would have

to be compatible with animal metabolism (Elshahawi et al., 2013; Yang et al., 2009). If so, the symbiosis might provide a ready source of new antibiotics to target the rise of multidrug resistance, a crucial problem in human health. Typically, intracellular symbiotic bacteria are difficult to cultivate, making it hard to discover the important antibiotics that result. However, a number of *Teredinibacter turnerae* strains have been successfully cultured, facilitating genome sequencing and large-scale fermentation for natural product isolation. Significantly, some of the potent compounds isolated from *Teredinibacter* in culture are also found in host shipworm animals, where they presumably help to keep out competing microbes (Han et al., 2013; Elshahawi et al., 2013). The compounds known so far include tartrolon, an exceptionally potent antiparasitic compound (Elshahawi et al., 2013; O'Connor et al., 2020), and a siderophore, turnerbactin (Han et al., 2013). However, the key antibacterial compounds, which would potentially be responsible for shaping the shipworm gill communities and the axenic cecum, have yet to be described.

To more rapidly hone in on ecologically important antibacterial compounds, we analyzed the genomes of cultivated symbiont strains and the metagenomes of shipworm gills, focusing on antibiotic-like biosynthetic gene clusters (BGCs) (Altamia



found this cluster to be ubiquitous (Figure 1A); however, the region including *turD* and *turE* appeared to be disrupted in six of these strains. In correctly assembled sequences, this region contained two repeats of 6k bp in length that were 98% identical, while in the misassembled sequences this region was scrambled. Therefore, it is likely that this is a result of bioinformatic assembly error, and that the gene cluster is intact and functional in all strains. All of the intact NRPSs were further compared by BLASTp, and it was found that each megasynthetase was greater than 85% similar to all of the other corresponding proteins from the homologous clusters, demonstrating a very high degree of conservation (Figure 1B).

In the *T. turnerae* T7901 genome, the five genes within this cluster include *turA* (three modules, 11.2 kbp), *turB* (three modules, 12.8 kbp), *turC* (two modules, 6.6 kbp), *turD* (two modules, 8.1 kbp), and *turE* (three modules, 12.1 kbp). Each module contains a condensation (C), adenylation (A), and thiolation (T) domain, and five of the modules also contain epimerization domains. The first domain begins with a so-called C-starter domain, which is known to acylate the initial amino acid in a process known as lipoinitiation (Rausch et al., 2007). The final module terminates in a thioesterase domain, which is responsible for product release, often by means of macrocyclization (Keating et al., 2001). This domain architecture indicates that the product of the *tur* cluster is a cyclic lipopeptide, a known class of antibacterial scaffolds.

Often, NRPS genes are naturally combinatorialized to create families of compounds that consist of structural analogs distributed throughout related producing organisms (Zan et al., 2019). To look for the potential of variants with divergent peptide backbones, the adenylation domains for the cluster in each of the eight sequenced strains were compared by BLASTp. In some of the misassembled sequences, the A domains in *turD* and *turE* were not intact and were thus not included in the analysis. The results show that *tur* is an exceptionally well-conserved cluster, with each A domain retaining greater than 90% identity to all of the analogous domains across all eight strains (Figure 1C). The substrate prediction, as determined by antiSmash 5.0, was also consistent for each domain (Blin et al., 2019). Importantly, the strains included in this analysis were isolated from a variety of shipworm genera and species collected in disparate locations around the world, indicating that this gene cluster, and thus its biosynthetic product, are well preserved, and thus likely important to the association between host and symbiont (Table S4).

Based upon the ubiquitous nature of this potential antibiotic, which is proposed to function in the symbiosis to defend the host and symbiont from other bacteria, we prioritized discovering the products of the *tur* pathway, seeking a lipopeptide containing 13 amino acids in the cultures of *T. turnerae* T7901.

Isolation and planar structure elucidation

The liquid culture of *T. turnerae* T7901 (6.6L) was extracted and subjected to a series of partitions. Initially, the expected peptides were not observed in crude extracts, and indeed the crude extract was not toxic to *Acinetobacter*. However, after several trials we observed a particularly robust boundary layer between the water and ethyl acetate fractions, which was enriched in a series of lipopeptides. Semi-preparative high-performance liquid

chromatography (HPLC) yielded two major compounds **1** and **2**, along with minor analogs **3** and **4**.

Turnercyclamycin A (**1**) had a molecular formula of $C_{71}H_{125}N_{15}O_{24}$ based upon high-resolution mass spectrometry ($[M+2H]^{2+}$ $m/z = 786.9618$), inherent in which are 17 degrees of unsaturation. The 1H and ^{13}C spectra were consistent with a lipopeptide based on numerous amide NH resonances between δ_H 7.6–8.5, alpha proton signals from δ_H 4.03–5.05, and the presence of a large methylene envelope centered at δ_H 1.23 (Figure S1). The gradient heteronuclear single quantum coherence (gHSQC) experiment revealed the presence of 11 methyl groups, 17 distinct methylene proton pairs outside of the lipid envelope, and 19 methine protons (Table S1).

The amide NH and alpha methine protons for eight amino acids could clearly be correlated through well-separated gradient correlation spectroscopy (gCOSY) crosspeaks, and verified by gradient heteronuclear multiple bond correlation (gHMBC) experiments. A total correlation spectroscopy (zTOCSY) experiment further confirmed these eight correlations and identified an additional five residues. A combination of gCOSY, zTOCSY, and gHMBC experiments were used to elucidate the side chain from each alpha methine proton (Figure 2A), revealing the presence of ornithine, serine, valine, threonine, glycine, two isoleucines, glutamine, alanine, homoserine, β -hydroxyaspartic acid, β -hydroxyisoleucine, and an intriguing 2,4-diamino-3-hydroxybutanoic acid (DAHB) moiety. The presence of all amino acids was confirmed through subsequent Marfey's analysis, which also revealed masses consistent with the FDLA-derivatized β -hydroxyisoleucine and DAHB moieties, despite the lack of standards for retention time comparison.

gHMBC correlations from amide NH protons to carbonyls, alpha methine protons to carbonyls, and rotating frame Overhauser spectroscopy (ROESY) correlations from amide NH protons to alpha methine protons were used to determine the sequence of residues in the peptide portion of the molecule. These data converged on the linear sequence of Orn–Ser–Val– β -OH–Asp–DAHB–Thr–Gly– β -OH–Ile–Ile–hSer–Ala–Ile–Glu, which was consistent with the order of substrate predictions for the adenylation domains of each NRPS module in the proposed biosynthetic pathway.

The formula of the peptide backbone left a remainder of $C_{14}H_{21}O$, indicating a tetradecanoic acid, and two degrees of unsaturation. TOCSY and gHMBC correlations verified the fatty acid moiety, and gHMBC and ROESY correlations confirmed that it was linked through the terminal ornithine residue. The final remaining degree of unsaturation indicated macrocyclization, a common feature of lipopeptides. A key HMBC correlation from the β -H of the threonine residue (δ_H 5.06) to the carbonyl of the glutamic acid moiety (δ_C 170.7) indicated that macrocyclization is via ester linkage through the threonine oxygen. This was further supported by the downfield shifted β -oxymethine carbon at δ_C 70.2.

Turnercyclamycin B (**2**) was obtained as the second major compound. Based on the observed $[M+2H]^{2+}$ ion at $m/z = 799.9693$, its molecular formula was $C_{73}H_{127}N_{15}O_{24}$, differing from that of **1** by C_2H_2 and an additional degree of unsaturation. Comparison of the proton and HSQC spectra showed that all chemical shifts and correlations associated with the peptide portion of the molecule were identical to those of **1** (Figure S2;




Figure 2. Structure elucidation of turnercyclamycins A–D

(A) COSY/TOCSY (bold bonds), key ROESY (purple, dashed arrows), and key HMBC (red, solid arrows) correlations used to solve the planar structure. (B) Structures of turnercyclamycins with assigned absolute configurations.

based structural characterization. However, the observed $[M+2H]^{2+}$ ions at $m/z = 772.9473$ for **3** and $m/z = 793.9621$ for **4** indicated formulas of $C_{69}H_{121}N_{15}O_{24}$ and $C_{72}H_{127}N_{15}O_{24}$, respectively. The 1H NMR and COSY spectra for both compounds are superimposable with those of **2** (Figures S3). MS-MS fragmentation localized the differences in mass to the lipid tail (Figures S4A and S4B), indicating that **3** and **4** are identical to **1** and **2**, except that they contain unbranched C12:0 and C15:0 fatty acid tails, respectively.

A number of other minor analogs were identifiable by LC-MS analysis but were inseparable by HPLC. MS-MS fragmentation revealed the inclusion of a number of other lipid tails on the same peptide core, as well as two structural modifications to the peptide portion of the molecule. One appears to be an ornithine-to-lysine substitution, which is well supported by multiple fragment ions (Figure S4D). The other is a loss of 14 amu that is localized to the cyclic portion of the peptide, but the fragmentation pattern did not clearly identify which amino acid was altered. The explanation most consistent with the data is that an isoleucine residue is substituted by valine.

Table S1). However, an HSQC correlation between δ_H 5.32 (m, 2H) and δ_C 129.6, as well as a distinct TOCSY spin system including a methylene envelop and the olefinic protons, indicated the presence of an unsaturation within the fatty acid side chain. The 1H and ^{13}C chemical shifts of both olefinic protons were coincident. Both allylic carbons were observed at δ_C 26.7, a value that is consistent with a *cis* configuration. Typically, *trans* olefins in fatty acids result in a chemical shift approximately 5 ppm downfield of this value (Gunstone et al., 1977).

To determine the location of the double bond in the lipid side chain of **2**, ozonolysis was performed on intact **2** followed by a reductive workup and liquid chromatography-mass spectrometry (LC-MS). A major detected $[M+2H]^{2+}$ ion of $m/z = 758.91$, corresponding to a monoisotopic mass of 1,515.80, matches the mass of the intact peptide with a 9-formyl-nonanoic acid moiety (Figures S5H and S5I). Therefore, **2** was determined to contain a *cis*-9-hexadecanoic fatty acid tail.

The minor analogs turnercyclamycin C (**3**) and D (**4**) were obtained as pure compounds, but in insufficient yield for complete NMR-

Finally, we detected by LC-MS an assortment of analogs that represent ring-open linear peptides, methyl esters, and linear methyl esters. We propose that these represent hydrolysis artifacts and transesterification byproducts of the methanolic extraction and isolation work up (Figure S4K).

Proposed biogenesis

The adenylation domain substrate predictions of each NRPS module map very well to the NMR-elucidated structure. The only adenylation domains for which no substrate was predicted correspond to the incorporation of DAHB and homoserine, both unusual amino acids. The prediction for the modules incorporating β -OH-Asp and β -OH-Ile were aspartic acid and isoleucine, respectively. In addition, the presence or absence of epimerase domains perfectly matched the results of the Marfey's analysis, with E domains present in all *D*-configuration amino acid extensions for which standards were available (Figure 3).

Upstream of the five NRPS megasynthase genes are four open reading frames that are predicted to be involved in the tailoring




Figure 3. Proposed biogenesis

Red amino acid labels indicate D-configuration, green indicates L-configuration, and blue hydroxyl groups are proposed to be installed by tailoring enzymes.

and export of the compound (Table S3). Two, *turF* and *turH*, encode α -ketoglutarate-dependent dioxygenases (Fe/ α KG), which are well characterized for performing hydroxylation reactions (Wu et al., 2016). Another, *turG*, encodes a cupin domain-containing protein with similarity to the protein JmjC. This class of enzyme also uses non-heme Fe and α KG to demethylate histones via hydroxylation, and related enzymes are responsible for the β -hydroxylation of lysine (Markolovic et al., 2018; Tsukada et al., 2006). It is likely that these three enzymes are performing the three β -hydroxylation reactions for the final product (Figure 3).

Stereochemical assignment by advanced Marfey's analysis

Compound **1** was hydrolyzed, and the constituent amino acids were derivatized with L-1-fluoro-2,4-dinitrophenyl-5-leucine-amide (FDLA) then analyzed by RP-ultra performance liquid chromatography-(UPLC)-HRESIMS. The result was compared against derivatized standards, when available, and D-FDLA-derivatized L-configuration standards were used to confirm retention times for D-configuration amino acids. This analysis confirmed the serine, valine, threonine, homoserine, alanine, and glutamic acid residues are all L-configuration. The ornithine residue was determined to be in the D-configuration, and both isoleucine residues were confirmed to be D-*allo*-configured by comparison with D- and L-FDLA-derivatized standards of L-Ile and L-*allo*-Ile, which all separated by UPLC. These results were in accordance with the presence or absence of an epimerase domain within their associated NRPS modules in the proposed BGC.

The β -hydroxyaspartic acid was resolved based on comparison of the elution order with previously reported values (Fuji et al., 1997) and the elution pattern from the potashchalins (Li et al., 2020). The hydrolysate of **1** was derivatized with both L-FDLA and D-FDLA, and after considerable LC method development a small but distinct change in retention time was observed.

The L-FDLA-derivatized peak elutes earlier, which corresponds to the D-configured alpha carbon. This is further supported by the presence of an epimerase domain within the module incorporating this residue into the growing peptide chain. The L- and D-FDLA-derivatized hydrolysate peaks were then compared with the L-FDLA-derivatized D/L-*threo*- β -hydroxyaspartic acid standard. The elution of the standard matches well with that of the hydrolysate, and thus the absolute configuration is deemed to be D-*threo*- β -hydroxyaspartic acid. EICs and retention times for the derivitized hydrolysate and amino acid standards are supplied in Figures S5A–S5G and Table S2.

As no standards are available for DAHB, the amino acid was synthesized following the methods employed in the total synthesis of odorhabdin, in which the base hydrolysis and subsequent deacetylation of hydroxyectoine was reported to result in a 70:30 mixture of the 2*S*,3*S* and 2*R*,3*S* diastereomers, respectively (Sarciaux et al., 2018). The crude reaction mixture was derivitized with L- and D-FDLA, and the four diastereomers were clearly distinguishable based on retention time and peak height. The minor peak in the L-FDLA-derivatized reaction mixture, corresponding to 2*R*,3*S* DAHB, matched the L-FDLA-derivitized peak in the hydrolysate of **1**.

Stereochemical prediction by biosynthetic logic

The remaining residue, β -hydroxyisoleucine, presented a challenge due to the lack of any commercially available standards. β -Hydroxyisoleucine is rarely found in natural products; it was reported without stereochemical assignments in stalobactin (Matsui et al., 2020), and the C2 and C3 configurations of this residue were only resolved in phomopsin A through X-ray crystallography (Culvenor et al., 1989). Attempts to grow diffraction-quality crystals were unsuccessful, so investigation of the NRPS domain architecture for the proposed biosynthesis of the turnercyclamycins was used to predict the C2 stereochemistry. The L- or D-configuration of all 11 residues that were assigned via Marfey's analysis perfectly correlated to the absence or

Table 1. MICs of 1 and 2 bacterial and human (HEK293) cells

Pathogen or cell line	MIC ($\mu\text{g/mL}$)	
	1	2
<i>Acinetobacter baumannii</i>	8	8
<i>Escherichia coli</i>	1	1
<i>Enterococcus faecalis</i>	>64	>64
<i>Klebsiella pneumoniae</i>	2	2
<i>Staphylococcus aureus</i>	>64	>64
HEK-293	>64	>64

MIC, minimum inhibitory concentration.

presence of an epimerase domain. Thus, it is predicted that the β -hydroxyisoleucine is in the L-configuration due to the lack of an epimerase domain. The C3 position, however, remains ambiguous due to a lack of knowledge of the stereoselectivity of the hydroxylating enzyme for this position.

Antibiotic activity of 1 and 2

Compounds **1** and **2** were initially tested against a panel of Gram-positive and Gram-negative pathogens in liquid broth assays. Both **1** and **2** were significantly active against *Escherichia coli* (minimum inhibitory concentration [MIC] 1 $\mu\text{g/mL}$), *Acinetobacter baumannii* (MIC 8 $\mu\text{g/mL}$), and *Klebsiella pneumoniae* (MIC 2 $\mu\text{g/mL}$), and neither compound showed activity against *Enterococcus faecalis* up to 64 $\mu\text{g/mL}$ (Table 1). Neither **1** nor **2** were bactericidal to *Staphylococcus aureus* at concentrations up to 64 $\mu\text{g/mL}$, but at 4 $\mu\text{g/mL}$ and above they slowed growth, decreasing it by 50% during the period of the assay (Figure S7A). Since many antibiotics are relatively inactive against *A. baumannii*, we further investigated this activity, substituting the nonpathogenic relative *A. baylyi*. **1** and **2** showed the same MIC against *A. baylyi*, with similar phenotypic effects on the organism, validating the use of the strain. Plating of inhibited wells from *A. baylyi* broth cultures determined that the growth inhibition was a result of bactericidal, not bacteriostatic, activity (Figure S6).

Lipopeptide antibiotics often meet hurdles in development due to toxicity against mammalian cells and the lysing of erythrocytes owing to their amphipathic nature (Agner et al., 2000). Thus, **1** and **2** were tested for cytotoxicity against the human kidney-derived HEK293 cell line and for hemolytic activity in freshly harvested murine erythrocytes. Both compounds showed no activity in either assay up to 64 $\mu\text{g/mL}$, in comparison with control compounds that behaved as expected.

The combination of bactericidal activity against *Acinetobacter* and a lack of observed toxicity led us to further investigate the mechanism of action. A phenotypic assay was employed in which *A. baylyi* was cultivated in the presence or absence of drug, and the morphology and growth characteristics were compared with knockout mutants that are deficient in various essential proteins (Bailey et al., 2019; Gallagher et al., 2020). A total of 43 genes corresponding to 14 essential processes have been deleted, and the terminal morphologies of the mutant cells are distinct and depend on the process inactivated (Figure 4). In the presence of either **1** or **2** at the MIC, cells would divide a few times, then arrest growth as clumps of rounded

cells. This death phenotype was strikingly similar to that observed in mutants deficient in outer membrane protein and LPS synthesis (*bamA*, *lptAB*, *tamAB*, and *lolCD*) (Figure 4). Treatment with colistin, a clinically used lipopeptide targeting the outer membrane, also produced clumped, rounded cells. At concentrations 4 times the MIC, both **1** and **2** killed cells immediately and without the same morphological changes, potentially indicating a second mechanism at higher concentrations. Overall, the findings imply that the turnercyclamycins compromise outer membrane integrity.

If colistin and turnercyclamycins have the same molecular mechanism, it might be expected that colistin-resistant strains would also evade turnercyclamycins. To test this, we obtained clinical *A. baumannii* complex isolates from the Associated Regional and University Pathologists (ARUP) laboratory and the *A. baumannii* panel from the Centers for Disease Control and Prevention (CDC). We tested six of the CDC strains, including two each of strains susceptible, moderately resistant, and highly resistant to colistin. The three ARUP strains also showed a range of colistin resistance levels. Strikingly, turnercyclamycins retained their potency against these strains, including against highly colistin-resistant *Acinetobacter* strains (Table 2). This implies that the molecular mechanisms of these two lipopeptide classes may differ.

Since colistin/polymyxin resistance is increasingly important in diverse Gram-negative bacteria, we also investigated the effectiveness of **1** and **2** against *Yersinia pestis*. This belongs to a subset of bacteria in which the linkage of 4-aminoarabinose to the LPS confers polymyxin resistance (Aoyagi et al., 2015). *Y. pestis* susceptibility to **1** and **2** could be neatly correlated to a series of genetic mutants containing and lacking 4-aminoarabinose (Table 3). Thus, while **1** and **2** are effective against bacteria such as *Acinetobacter*, which resist polymyxins by adding phosphoethanolamine to their LPS (Beceiro et al., 2011), among other mechanisms, they may not be as effective against bacteria that resist polymyxins primarily by adding 4-aminoarabinose.

To investigate the potential resistance of *A. baumannii* to turnercyclamycins, we incubated dilutions of wild-type cells in the presence of **2** at $2\times$ MIC (16 $\mu\text{g/mL}$) for 4 days. Cells were added in 10 concentrations from 86 to 1.6×10^8 colony-forming units (CFU) per mL, and growth was monitored daily. Wells starting at densities of 10,752 CFU/mL and above were visibly turbid after 24 h. At 96 h, a well containing 2,150 CFU/mL was slightly turbid in one replicate, but not in a second replicate. By contrast, colistin cleared cultures containing up to 1.3×10^6 CFU/mL. Broth was obtained from a well that turned turbid on the fourth day of incubation and plated on medium containing **2** at $2\times$ MIC. Strikingly, only five colonies grew, when a lawn might be expected if cells in the well were resistant to the drug. A surviving colony was used for MIC determination, and interestingly the maximal activity of **2** was reached at the same concentration as the original wild-type strain (8 $\mu\text{g/mL}$), but only 70% inhibition was achieved, even up to 64 $\mu\text{g/mL}$ (Figure S7P). This same strain showed no changes in its MIC toward colistin (Figure S7Q). This indicates an inoculum effect in Mueller Hinton broth for **2** starting at $\sim 10^4$ CFU/mL (Brook, 1989). To investigate this effect on solid medium, dense (10^8 CFU/mL) cultures of wild-type *A. baumannii* were inoculated and spread on Mueller Hinton




Figure 4. Terminal morphologies of mutant and antibiotic-treated *A. baylyi*

Each panel is a compilation of representative cells from the same condition. Cells are moved closer to each other than they appeared under the microscope so that they fit within each panel, but the overlaid images do not obscure other cells. The images show bacteria whose growth has been inhibited by essential gene deletion or treatment with antibiotics at approximately the MICs (turnercyclamycin A and B, 8 $\mu\text{g}/\text{mL}$; colistin, 0.5 $\mu\text{g}/\text{mL}$). The deletion mutations inactivate the following processes: *dnaA*, DNA replication; *rpoC*, transcription; *ftsI*, cell division; *bamA*, outer membrane protein localization; *lptAB*, LPS localization. Scale bar, 2 μm . wt, wild type; TurnA, turnercyclamycin A (1); Turn B, turnercyclamycin B (2).

lipopeptides that are still effective against colistin-resistant *Acinetobacter*. Since we used clinical isolates resistant to colistin, as well as all other widely used anti-*Acinetobacter* drugs, this provides a chemical entity that might guide development of treatments for multidrug-resistant *Acinetobacter*.

Symbiotic relationships represent unique environments that are promising for antimicrobial natural products discovery. These systems are typically described

and understood based on the primary metabolic and nutritional interactions involved. However, exchanges of secondary metabolites also play important roles in these relationships. Microbes living within animal host tissues are of particular interest for antibiotic discovery because compounds made within these contexts must be non-toxic to the host cells and possess properties that facilitate distribution throughout the host tissues, which can translate to favorable pharmacokinetics. The shipworm-*Teredinibacter* symbiotic relationship was investigated as a particularly promising source of antibiotic compounds due to: (1) previous studies showing a nearly axenic cecum, despite this being the site of cellulose degradation, likely resulting in an abundance of free glucose; (2) the presence of bacterial cellulases in this same organ, suggesting a route of transport from symbiont-containing gills to the cecum; and (3) a number of highly conserved BGCs shared throughout the *Teredinibacter* lineage that could be producing complex PKS and NRPS compounds.

DISCUSSION

Antimicrobial compounds, particularly those with unique structural scaffolds and that function through mechanisms of action that have not yet been employed in the clinic, are in high demand. Each year, at least 35,000 people die from antibiotic-resistant bacterial infections in the United States (Howard et al., 2012; Lee et al., 2017; Piperaki et al., 2019; Wong et al., 2017). This is a number that keeps increasing, as bacteria are increasingly resistant to available drugs. *Acinetobacter* is an especially daunting pathogen, with more than 60% of patient-isolated strains showing resistance to standard treatments derived from β -lactams or fluoroquinolones. An increasing number of strains is extremely drug resistant, being untreatable even by the last-line lipopeptide agent, colistin (polymyxin E). Because there are very few *Acinetobacter* treatments in development, the US Centers for Disease Control considers carbapenem-resistant *Acinetobacter* to be an urgent threat (Centers for Disease Control and Prevention (U.S.), 2019). Here, we describe a group of

Following our ecology-guided rationale, a family of cyclic lipopeptides, the turnercyclamycins, was isolated and found to have activity against multiple Gram-negative pathogens, including colistin-resistant *Acinetobacter*. Last-line antibiotic resistance is a widespread problem beyond *Acinetobacter*, including in *E. coli* and other Gram-negative pathogens, so with further study this family of lipopeptides may find an impact on therapy. This family consists of a conserved, NRPS-derived 13 amino acid core sequence and a variable fatty acyl tail. The peptide portion is cyclized through an ester linkage via the threonine hydroxyl group, resulting in an eight amino acid

Table 2. MICs against ARUP and CDC panels of *Acinetobacter* strains with varying degrees of resistance against colistin

Pathogen	MIC ($\mu\text{g/mL}$)		Colistin
	1	2	
<i>Acinetobacter baumannii</i>	8	8	2
ARUP ABC 1	4	4	4
ARUP ABC 2	8	8	4
ARUP ABC 3	16	8	>32
CDC AB282	4	4	1
CDC AB286	4	4	1
CDC AB299	2	4	2
CDC AB302	8	4	2
CDC AB303	8	8	>32
CDC AB307	8	8	8

macrocycle and five amino acid linear portion. A number of features are unique to this structural family, most notably the inclusion of three β -hydroxy amino acid residues, two of which, β -hydroxyisoleucine and 2,4-diamino-3-hydroxy-butanoic acid, are exceptionally rare. There is a precedent for β -hydroxyaspartic acid in the literature, as it is a common feature of microbial siderophores. Notably, in BGC_1-encoding turnercyclamycins, all of the needed proteins and protein domains are present, including those needed for β -hydroxylation (Reitz et al., 2019; Markolovic et al., 2018; Hibi et al., 2011), strongly supporting the biogenetic hypothesis.

The chemical properties of the turnercyclamycins are also intriguing when considering their mechanism of action, especially in relation to colistin. Both compounds are cyclic lipopeptides that affect outer membrane integrity, but the *Acinetobacter* strains with resistance to colistin are still susceptible to turnercyclamycins. It is believed that colistin works largely due to the highly cationic peptide portion, which gains its positive charges through the side chains of five DAB moieties. This charged peptide displaces magnesium and calcium counter ions in the LPS, while the lipid tail works to solubilize the membrane through a detergent-like mechanism (Dixon and Chopra, 1986). The turnercyclamycins, on the other hand, are not rich in DAB residues or other cationic peptide side chains, and indeed maintain a neutral charge balance. This chemical property may be important in explaining how the turnercyclamycins retain activity against colistin-resistant strains, and likely points to a different molecular mechanism of action that accomplishes the same goal of outer membrane disruption. By contrast, some Gram-negative bacteria, including *Y. pestis*, incorporate 4-aminoarabinose into their LPS to evade polymyxins and other cationic antimicrobial peptides. This mode of resistance is clearly effective against turnercyclamycins, revealing that the compounds may exhibit therapeutic selectivity. In addition, the lack of toxicity or hemolytic activity of turnercyclamycins suggest that they may make useful therapeutics. Although we did not observe emergence of classic resistance mutations, the bacteria could evade inhibition by other means under certain conditions. An inoculum effect was observed, which warrants further study. This effect is similar to that reported for daptomycin, and further knowledge of the molecular mechanism of action and ion dependence of turnercyclamycins

Table 3. MICs against *Y. pestis* strains with varying levels of 4-amino-4-deoxy-L-arabinose modification of the LPS

Pathogen	Ara-4N modification level ^a	MIC ($\mu\text{g/mL}$)		
		1	2	Polymyxin B
<i>Y. pestis</i> KIM6+	High	≥ 256	≥ 256	≥ 256
<i>Y. pestis</i> ΔarnB	Mid	8	8	16
<i>Y. pestis</i> ΔarnOP	Low	4	8	0.0625
<i>Y. pestis</i> $\Delta\text{arnOP-Comp}$	High	≥ 256	≥ 256	≥ 256

^aFrom Aoyagi et al. (2015).

mycin will be necessary to allow for more accurate resistance studies (Quinn et al., 2007; Silverman et al., 2001). Further *in vivo* evaluation is required to determine the potential clinical utility of the compounds.

Secondary metabolism plays a large role in the association between *T. turnerae* and their shipworm hosts. Thus far, of the ubiquitous biosynthetic pathways found in all *T. turnerae* strains and in all shipworm gill metagenomes in which *T. turnerae* is a symbiont, three have been characterized. Because of their ubiquity, all three likely play crucial roles in symbiosis biology. Tartrolon D/E was originally isolated based on antibacterial activity, but its exquisite potency against apicomplexan parasites is hypothesized to protect the mollusk from gregarines, which are known to be pathogens of mollusks (Elshahawi et al., 2013; O'Connor et al., 2020; Rueckert et al., 2019). Turnerbactin, the triscatecholate siderophore, may be important both for the acquisition of iron for the symbiont and host, as well as for the sequestration of iron to limit the growth of pathogenic or opportunistic bacteria (Han et al., 2013). This most recent discovery of the turnercyclamycins represents the addition of a potent bactericidal agent to the molecular arsenal of *T. turnerae*, capable of directly killing bacteria that pose threats to its shipworm host, their shared food supply, or the ecological niche of the symbiont itself. This finding underscores the importance of closely examining the chemical biology of symbiotic systems with a particular focus on ecological rationale and metagenomics in the pursuit of new pharmacological agents for improved human health.

SIGNIFICANCE

Drug-resistant *Acinetobacter* are notoriously difficult to treat. Here, we describe a natural antibiotic that maintains potency against *Acinetobacter* strains that have resistance against a last-line therapeutic. To find the compound, we investigated the widespread association between shipworms and their symbiotic bacteria, and isolated the product of the most prevalent biosynthetic gene cluster in shipworms. This reinforces the role of better understanding ecology in the discovery of new drug leads.

STAR★METHODS

Detailed methods are provided in the online version of this paper and include the following:

- KEY RESOURCE TABLE

- RESOURCE AVAILABILITY
 - Lead contact
 - Materials availability
 - Data and code availability
- EXPERIMENTAL MODEL AND SUBJECT DETAILS
 - Bacterial strains and culture media
 - Mammalian cell lines and culture media
- METHOD DETAILS
 - Chemical extraction of *T. turnerae* and purification of turnercyclamycins
 - Synthesis of 2,4-diamino-3-hydroxy-butanoic acid
 - Determination of amino acid stereochemistry
 - Antimicrobial microdilution assay
 - Mammalian antiproliferative assay
 - Hemolysis assay
 - Dilution broth inhibition experiments
 - Microscopy of growth-inhibited bacteria
 - *tur* gene clusters analysis in *T. turnerae* strains
 - Spectroscopy
- QUANTIFICATION AND STATISTICAL ANALYSIS

SUPPLEMENTAL INFORMATION

Supplemental information can be found online at <https://doi.org/10.1016/j.chembiol.2021.05.003>.

ACKNOWLEDGMENTS

We thank Xuchen Zhan and Jon Rainier (University of Utah) for assistance with ozonolysis and the Center for High Performance Computing, University of Utah, for computational resources. This work was funded by grants U01TW008163 (NIH Fogarty) to M.G.H., NIH R35GM122521 to E.W.S., NOAA NA19OAR0110303 to E.W.S. and M.G.H., and R01AI148208 to C.M. We are grateful for key infrastructure support provided by the ALSAM Foundation.

AUTHOR CONTRIBUTIONS

Conceptualization, B.W.M., M.G.H., and E.W.S.; methodology, B.W.M., A.L.L., Z.L., and C.M.; investigation, B.W.M., A.L.L., Z.L., J.B., and K.L.A.; resources, M.A.F., L.R.B., C.M., E.W.S., and M.G.H.; writing – original draft, B.W.M., J.B., K.L.A., E.W.S., and M.G.H.; writing – review & editing, B.W.M., A.L.L., E.W.S., and M.G.H.; visualization, B.W.M., A.L.L., Z.L., J.B., and K.L.A.; supervision, M.A.F., L.R.B., C.M., and M.G.H.; funding acquisition, M.G.H., E.W.S., and C.M.

DECLARATION OF INTERESTS

The authors have filed a provisional patent application for the turnercyclamycins.

Received: December 28, 2020

Revised: April 19, 2021

Accepted: May 4, 2021

Published: June 18, 2021

REFERENCES

- Agner, G., Kaulin, Y.A., Gurnev, P.A., Szabo, Z., Schagina, L.V., Takemoto, J.Y., and Blasko, K. (2000). Membrane-permeabilizing activities of cyclic lipopeptides, syringopeptin 22A and syringomycin E from *Pseudomonas syringae* pv. *syringae* in human red blood cells and in bilayer lipid membranes. *Bioelectrochemistry* 52, 161–167.
- Altamia, M.A., Lin, Z., Trindade-Silva, A.E., Uy, I.D., Shipway, J.R., Wilke, D.V., Concepcion, G.P., Distel, D.L., Schmidt, E.W., and Haygood, M.G. (2020). Secondary metabolism in the gill microbiota of shipworms (Teredinidae) as revealed by comparison of metagenomes and nearly complete symbiont genomes. *mSystems* 5, e00261–20.
- Aoyagi, K.L., Brooks, B.D., Bearden, S.W., Monteneri, J.A., Gage, K.L., and Fisher, M.A. (2015). LPS modification promotes maintenance of *Yersinia pestis* in fleas. *Microbiology* 161, 628–638.
- Bailey, J., Cass, J., Gasper, J., Ngo, N.-D., Wiggins, P., and Manoil, C. (2019). Essential gene deletions producing gigantic bacteria. *PLoS Genet.* 15, e1008195.
- Beceiro, A., Llobet, E., Aranda, J., Bengoechea, J.A., Doumith, M., Hornsey, M., Dhanji, H., Chart, H., Bou, G., Livermore, D.M., et al. (2011). Phosphoethanolamine modification of lipid A in colistin-resistant variants of *Acinetobacter baumannii* mediated by the *pmrAB* two-component regulatory system. *Antimicrob. Agents Chemother.* 55, 3370–3379.
- Betcher, M.A., Fung, J.M., Han, A.W., O'Connor, R., Seronay, R., Concepcion, G.P., Distel, D.L., and Haygood, M.G. (2012). Microbial distribution and abundance in the digestive system of five shipworm species (Bivalvia: teredinidae). *PLoS One* 7, e45309.
- Blin, K., Shaw, S., Steinke, K., Villebro, R., Ziemert, N., Lee, S.Y., Medema, M.H., and Weber, T. (2019). antiSMASH 5.0: updates to the secondary metabolite genome mining pipeline. *Nucleic Acids Res.* 47, W81–W87.
- Boucher, H.W., Talbot, G.H., Bradley, J.S., Edwards, J.E., Gilbert, D., Rice, L.B., Scheld, M., Spellberg, B., and Bartlett, J. (2009). Bad bugs, no drugs: no ESKAPE! An update from the Infectious Diseases Society of America. *Clin. Infect. Dis.* 48, 1–12.
- Brook, I. (1989). Inoculum effect. *Rev. Infect. Dis.* 11, 361–368.
- Centers for Disease Control and Prevention (U.S.) (2019). Antibiotic Resistance Threats in the United States, 2019 (Centers for Disease Control and Prevention (U.S.)).
- Culvenor, C.C.J., Edgar, J.A., Mackay, M.F., Gorst-Allman, P., Marasas, C.F.O., Steyn, W.S., Vlegaar, P.R., and Wessels, L.P. (1989). Structure elucidation and absolute configuration of phomopsisin a, a hexapeptide mycotoxin produced by *Phomopsis leptostromiformis*. *Tetrahedron* 45, 2351–2372.
- Distel, D.L., Beaudoin, D.J., and Morrill, W. (2002). Coexistence of multiple proteobacterial endosymbionts in the gills of the wood-boring bivalve *Lyrodus pedicellatus* (Bivalvia: teredinidae). *Appl. Environ. Microbiol.* 68, 6292–6299.
- Dixon, R.A., and Chopra, I. (1986). Leakage of periplasmic proteins from *Escherichia coli* mediated by polymyxin B nonapeptide. *Antimicrob. Agents Chemother.* 29, 781–788.
- Ekborg, N.A., Morrill, W., Burgoyne, A.M., Li, L., and Distel, D.L. (2007). CelAB, a multifunctional cellulase encoded by *Teredinibacter turnerae* T7902T, a culturable symbiont isolated from the wood-boring marine bivalve *Lyrodus pedicellatus*. *Appl. Environ. Microbiol.* 73, 7785–7788.
- Elshahawi, S.I., Trindade-Silva, A.E., Hanora, A., Han, A.W., Flores, M.S., Vizzoni, V., Schrago, C.G., Soares, C.A., Concepcion, G.P., Distel, D.L., et al. (2013). Boronated tartrolon antibiotic produced by symbiotic cellulose-degrading bacteria in shipworm gills. *Proc. Natl. Acad. Sci. U S A* 110, E295–E304.
- Fujii, K., Ikai, Y., Mayumi, T., Oka, H., Suzuki, M., and Harada, K. (1997). A nonempirical method using LC/MS for determination of the absolute configuration of constituent amino acids in a peptide: elucidation of limitations of Marfey's method and of its separation mechanism. *Anal. Chem.* 69, 3346–3352.
- Gallagher, L.A., Bailey, J., and Manoil, C. (2020). Ranking essential bacterial processes by speed of mutant death. *Proc. Natl. Acad. Sci. U S A* 117, 18010–18017.
- Gunstone, F.D., Pollard, M.R., Scrimgeour, C.M., and Vedanayagam, H.S. (1977). Fatty acids. Part 50. ¹³C nuclear magnetic resonance studies of olefinic fatty acids and esters. *Chem. Phys. Lipids* 18, 115–129.
- Han, A.W., Sandy, M., Fishman, B., Trindade-Silva, A.E., Soares, C.A.G., Distel, D.L., Butler, A., and Haygood, M.G. (2013). Turnerbactin, a novel trisaccholate siderophore from the shipworm endosymbiont *Teredinibacter turnerae* T7901. *PLoS One* 8, e76151.

- Hibi, M., Kawashima, T., Kodera, T., Smirnov, S.V., Sokolov, P.M., Sugiyama, M., Shimizu, S., Yokozeki, K., and Ogawa, J. (2011). Characterization of *Bacillus thuringiensis* L-isoleucine dioxygenase for production of useful amino acids. *Appl. Environ. Microbiol.* **77**, 6926–6930.
- Howard, A., O'Donoghue, M., Feeney, A., and Sleator, R.D. (2012). *Acinetobacter baumannii*: an emerging opportunistic pathogen. *Virulence* **3**, 243–250.
- Keating, T.A., Ehmann, D.E., Kohli, R.M., Marshall, C.G., Trauger, J.W., and Walsh, C.T. (2001). Chain termination steps in nonribosomal peptide synthetase assembly lines: directed acyl-S-enzyme breakdown in antibiotic and siderophore biosynthesis. *ChemBioChem* **2**, 99–107.
- Lechene, C.P., Luyten, Y., McMahon, G., and Distel, D.L. (2007). Quantitative imaging of nitrogen fixation by individual bacteria within animal cells. *Science* **317**, 1563–1566.
- Lee, C.-R., Lee, J.H., Park, M., Park, K.S., Bae, I.K., Kim, Y.B., Cha, C.-J., Jeong, B.C., and Lee, S.H. (2017). Biology of *Acinetobacter baumannii*: pathogenesis, antibiotic resistance mechanisms, and prospective treatment options. *Front. Cell. Infect. Microbiol.* **7**, 55.
- Li, Y., Liu, L., Zhang, G., He, N., Guo, W., Hong, B., and Xie, Y. (2020). Potashchelins, a suite of lipid siderophores bearing both L-threo and L-erythro beta-hydroxyaspartic acids, acquired from the potash-salt-ore-derived extremophile *Halomonas* sp. MG34. *Front. Chem.* **8**, 197.
- Markolovic, S., Zhuang, Q., Wilkins, S.E., Eaton, C.D., Abboud, M.I., Katz, M.J., McNeil, H.E., Leśniak, R.K., Hall, C., Struwe, W.B., et al. (2018). The Jumonji-C oxygenase JMJD7 catalyzes (3S)-lysyl hydroxylation of TRAFAC GTPases. *Nat. Chem. Biol.* **14**, 688–695.
- Matsui, K., Kan, Y., Kikuchi, J., Matsushima, K., Takemura, M., Maki, H., Kozono, I., Ueda, T., and Minagawa, K. (2020). Stalobacin: discovery of novel lipopeptide antibiotics with potent antibacterial activity against multidrug-resistant bacteria. *J. Med. Chem.* **63**, 6090–6095.
- O'Connor, R.M., Fung, J.M., Sharp, K.H., Benner, J.S., McClung, C., Cushing, S., Lamkin, E.R., Fomenkov, A.I., Henrissat, B., Londer, Y.Y., et al. (2014). Gill bacteria enable a novel digestive strategy in a wood-feeding mollusk. *Proc. Natl. Acad. Sci. U S A* **111**, E5096–E5104.
- O'Connor, R.M., V, F.J.N., Abenoja, J., Bowden, G., Reis, P., Beaushaw, J., Relat, R.M.B., Driskell, I., Gimenez, F., Riggs, M.W., et al. (2020). A symbiotic bacterium of shipworms produces a compound with broad spectrum anti-apicomplexan activity. *PLoS Pathog.* **16**, e1008600.
- Piperaki, E.-T., Tzouveleakis, L.S., Miriagou, V., and Daikos, G.L. (2019). Carbapenem-resistant *Acinetobacter baumannii*: in pursuit of an effective treatment. *Clin. Microbiol. Infect.* **25**, 951–957.
- Quinn, B., Hussain, S., Malik, M., Drlica, K., and Zhao, X. (2007). Daptomycin inoculum effects and mutant prevention concentration with *Staphylococcus aureus*. *J. Antimicrob. Chemother.* **60**, 1380–1383.
- Rausch, C., Hoof, I., Weber, T., Wohlleben, W., and Huson, D.H. (2007). Phylogenetic analysis of condensation domains in NRPS sheds light on their functional evolution. *BMC Evol. Biol.* **7**, 78.
- Reitz, Z.L., Hardy, C.D., Suk, J., Bouvet, J., and Butler, A. (2019). Genomic analysis of siderophore β -hydroxylases reveals divergent stereocontrol and expands the condensation domain family. *Proc. Natl. Acad. Sci. U S A* **116**, 19805–19814.
- Rueckert, S., Betts, E.L., and Tsaousis, A.D. (2019). The symbiotic spectrum: where do the gregarines fit? *Trends Parasitol.* **35**, 687–694.
- Sarciaux, M., Pantel, L., Midrier, C., Serri, M., Gerber, C., Marcia de Figueiredo, R., Campagne, J.-M., Villain-Guillot, P., Gualtieri, M., and Racine, E. (2018). Total synthesis and structure-activity relationships study of odorhabdins, a new class of peptides showing potent antibacterial activity. *J. Med. Chem.* **61**, 7814–7826.
- Silverman, J.A., Oliver, N., Andrew, T., and Li, T. (2001). Resistance studies with daptomycin. *Antimicrob. Agents Chemother.* **45**, 1799–1802.
- Tsakada, Y., Fang, J., Erdjument-Bromage, H., Warren, M.E., Borchers, C.H., Tempst, P., and Zhang, Y. (2006). Histone demethylation by a family of JmjC domain-containing proteins. *Nature* **439**, 811–816.
- Waterbury, J.B., Calloway, C.B., and Turner, R.D. (1983). A cellulolytic nitrogen-fixing bacterium cultured from the gland of *Deshayes* in shipworms (*Bivalvia*: *teredinidae*). *Science* **221**, 1401–1403.
- Wong, D., Nielsen, T.B., Bonomo, R.A., Pantapalangkoor, P., Luna, B., and Spellberg, B. (2017). Clinical and pathophysiological overview of *Acinetobacter* infections: a century of challenges. *Clin. Microbiol. Rev.* **30**, 409–447.
- Wu, L.-F., Meng, S., and Tang, G.-L. (2016). Ferrous iron and α -ketoglutarate-dependent dioxygenases in the biosynthesis of microbial natural products. *Biochim. Biophys. Acta Proteins Proteom.* **1864**, 453–470.
- Yang, J.C., Madupu, R., Durkin, A.S., Ekborg, N.A., Peadarallu, C.S., Hostetler, J.B., Radune, D., Toms, B.S., Henrissat, B., Coutinho, P.M., et al. (2009). The complete genome of *Teredinibacter turnerae* T7901: an intracellular endosymbiont of marine wood-boring bivalves (Shipworms). *PLoS One* **4**, e6085.
- Zan, J., Li, Z., Tianero, M.D., Davis, J., Hill, R.T., and Donia, M.S. (2019). A microbial factory for defensive kahalalides in a tripartite marine symbiosis. *Science* **364**, eaaw6732.

STAR★METHODS

KEY RESOURCE TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Bacterial and virus strains		
<i>T. turnerae</i> T7901 ATCC 39867	Prof. Margo G. Haygood, University of Utah, USA	N/A
<i>Staphylococcus aureus</i> ATCC 1600	Prof. Louis R. Barrows, University of Utah, USA	N/A
<i>Enterococcus faecalis</i> ATCC 29212	Prof. Louis R. Barrows, University of Utah, USA	N/A
<i>Klebsiella pneumoniae</i> ATCC BAA-1705	Prof. Louis R. Barrows, University of Utah, USA	N/A
<i>Acinetobacter baumannii</i> ATCC 19606	Prof. Louis R. Barrows, University of Utah, USA	N/A
<i>Acinetobacter baylyi</i> strain ADP1	Prof. Colin Manoil, University of Washington, USA	N/A
AB282	CDC	N/A
AB286	CDC	N/A
AB299	CDC	N/A
AB302	CDC	N/A
AB303	CDC	N/A
AB307	CDC	N/A
ABC1	Prof. Mark A. Fisher, ARUP Laboratories, USA	N/A
ABC2	Prof. Mark A. Fisher, ARUP Laboratories, USA	N/A
ABC3	Prof. Mark A. Fisher, ARUP Laboratories, USA	N/A
<i>Escherichia coli</i> ATCC 23724	Prof. Louis R. Barrows, University of Utah, USA	N/A
Chemicals, peptides, and recombinant proteins		
5-hydroxyectoine	Sigma Aldrich	Cat#70709-1G-F
L-FDLA	TCI Chemicals	Cat#A5523
D-FDLA	TCI Chemicals	Cat#A5524
Mueller Hinton Broth 2	Sigma Aldrich	Cat#90922
Luria Bertani Broth (Lennox)	Sigma Aldrich	Cat#L3022
Thiazolyl Blue Tetrazolium Bromide (MTT)	Sigma Aldrich	Cat#M2128
Experimental models: cell lines		
HEK-293	ATCC	Cat#CRL-1573
Software and algorithms		
AntiSMASH 5.0	(Blin et al., 2019)	https://antismash.secondarymetabolites.org/#/start
BLASTp	U.S. National Center for Biotechnology Information	https://blast.ncbi.nlm.nih.gov/Blast.cgi
R	https://www.r-project.org/	N/A

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, Eric W. Schmidt (ews1@utah.edu).

Materials availability

Reagents generated in this study are available from the Lead Contact with a completed Materials Transfer Agreement.

Data and code availability

The code generated during this study is available in the [supplementary information](#) provided with this manuscript.

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Bacterial strains and culture media

T. turnerae T7901 was grown in shipworm basal medium (SBM) (Waterbury et al., 1983). *Staphylococcus aureus* (*S. aureus* subsp. *aureus* ATCC 1600), *Enterococcus faecalis* (ATCC 29212), *Klebsiella pneumoniae* (ATCC BAA-1705), *Acinetobacter baumannii* (ATCC 19606), and the CDC strains AB282, AB286, AB299, and AB302 were grown in Mueller Hinton Broth II (MHBII). *E. coli* was grown in Luria-Bertani broth. Each strain was grown from a single colony. *T. turnerae* was grown at 30°C with 180 rpm shaking, while all other strains were grown at 37°C with 220 rpm shaking.

Mammalian cell lines and culture media

HEK-293 (ATCC CRL-1573) cells were cultured in RPMI 1640 medium supplemented with 10% fetal bovine serum, 100 units of penicillin, and 100 µg/mL of streptomycin under a humidified environment with 5% CO₂ at 37°C.

METHOD DETAILS

Chemical extraction of *T. turnerae* and purification of turnercyclamycins

A glycerol stock of *T. turnerae* T7901 was revived by streaking 5 µL on a plate of SBM. 5 mL liquid SBM cultures were inoculated with single colonies and incubated at 30°C with shaking at 180 RPM for 4 days. These seed cultures were used to inoculate six liquid SBM (with phosphate f/c = 15 µM) 1.1 L cultures in 2.8L baffled Fernbach flasks, which were incubated at 30°C with shaking at 180 RPM for 7 days. Cells were removed from culture media by centrifugation at 7,068 x g, 4°C, for 30 min. The supernatant was decanted, and the cell pellet was frozen and lyophilized to dryness.

The dry cell pellet was extracted three times with 400 mL each of 2:1 DCM:MeOH. The extracts were combined and concentrated *in vacuo* to remove DCM, and water was added to a composition of 5% v/v in MeOH, then partitioned 3 times with an equal volume of hexanes. The methanolic fraction was dried *in vacuo*, then partitioned 3 times between equal volumes of water and DCM. The aqueous layer was then subsequently partitioned 3 times with equal volumes of ethyl acetate. This step formed a particularly well-defined insoluble boundary layer, which was collected separately.

The boundary layer was dried *in vacuo* then resuspended in MeOH. This was centrifuged to pellet insoluble material, then the supernatant was filtered (Nylon, 0.45 µm) and subjected to semipreparative RPHPLC (Phenomenex Luna C₁₈ column, 250 x 10 mm, 5 µm, flow rate 4.0 mL/min). Isocratic conditions (20% ACN/ 80% H₂O/ 0.01% TFA) were held for 5 min followed by a linear gradient to 75% ACN/ 25% H₂O/ 0.01% TFA for 15 min to yield turnercyclamycins A-D (**1**, *t_R* = 15.3 min, 5.6 mg; **2**, *t_R* = 15.8 min, 6.8 mg; **3**, *t_R* = 14.2 min, 0.9 mg; **4**, *t_R* = 16.8 min, 0.6 mg).

Turnercyclamycin A (1):

Off-white amorphous solid; UV/vis: λ_{max} 250, 270 nm; IR (neat) ν_{max} 3296, 3067, 2930, 1645, 1538, 1204, 1140 cm⁻¹; ¹H and ¹³C NMR data in DMSO-*d*₆, Table 1; HRESIMS [M+2H]²⁺ ion at *m/z*=786.9619 (calculated for C₇₁H₁₂₅N₁₅O₂₄: 786.9584).

Turnercyclamycin B (2):

Off-white amorphous solid; UV/vis: λ_{max} 250, 270 nm; IR (neat) ν_{max} 3296, 3065, 2921, 1645, 1540, 1204, 1139 cm⁻¹; ¹H and ¹³C NMR data in DMSO-*d*₆, Table 1; HRESIMS [M+2H]²⁺ ion at *m/z*=799.9693 (calculated C₇₃H₁₂₇N₁₅O₂₄= 799.9662).

Turnercyclamycin C (3):

Off-white amorphous solid; ¹H NMR in DMSO-*d*₆, Figure S3; HRESIMS [M+2H]²⁺ ion at *m/z*=772.9473 (calculated for C₆₉H₁₂₁N₁₅O₂₄: 772.9427).

Turnercyclamycin D (4):

Off-white amorphous solid; ¹H NMR in DMSO-*d*₆, Figure S3; HRESIMS [M+2H]²⁺ ion at *m/z*=793.9621 (calculated for C₇₂H₁₂₇N₁₅O₂₄: 793.9662).

Synthesis of 2,4-diamino-3-hydroxy-butanoic acid

5-Hydroxyectoine (12.1 mg, 0.0765 mmol) was dissolved in water (2 mL). NaOH (2 equiv., 0.153 mmol, 6.12 mg) was added and the mixture was stirred at 50°C for 6 h. The solvent was removed under a stream of N₂ gas, and the resulting residue was dissolved in aqueous HCl (6N, 2mL), and the mixture was heated at 110°C for 3 h. The solution was diluted with water (4 mL), frozen at -80°C, then lyophilized. The resulting residue was dissolved in water (1.53 mL) and analyzed by advanced Marfey's method, described below.

Determination of amino acid stereochemistry

1 mL 6 N HCl was added separately to 1 mg of **1** and 1 mg of **2** and the reaction vessels were heated to 100°C with stirring for 24 h. Solvent was evaporated under a stream of nitrogen gas and the resulting residues were dissolved in 250 µL of H₂O. Aliquots (50 µL) of

each hydrolysate solution were transferred to a clean glass vial, to which 20 μL aqueous 1 M NaHCO_3 and 100 μL Marfey's reagent (L-FDLA, 1% solution in acetone, TCI Chemicals) were added. The reactions were incubated for 1 h at 40°C, then quenched with 20 μL 1 N HCl. This solution was filtered through a polypropylene filter (0.45 μm), then diluted into MeOH for UPLC-MS analysis. Analysis was performed on an analytical Waters Acquity UPLC equipped with a Acquity HSS T3 column (100 x 2.1 mm, 1.8 μm , Waters, flow rate = 0.3 mL min^{-1} , method: 0-2 min: 20% (v/v) ACN in water containing 0.1% formic acid; 2-30 min: linear gradient 20%-65% ACN in water containing 0.1% formic acid; 30-31 min: linear gradient 65%-100% ACN in water containing 0.1% formic acid). Amino acid standards were derivatized in the same manner for comparison, and D-FDLA was used to derivatize L-configured amino acid standards to obtain the retention times for D-amino acids. Retention times for amino acids derived from **1**, **2**, and amino acid standards are summarized in Table S2.

Antimicrobial microdilution assay

Glycerol stocks of *Staphylococcus aureus* (*S. aureus* subsp. *aureus* ATCC 1600), *Enterococcus faecalis* (ATCC 29212), *Klebsiella pneumoniae* (ATCC BAA-1705), *Acinetobacter baumannii* (ATCC 19606), and the CDC strains AB282, AB286, AB299, and AB302 were streaked on Mueller Hinton Agar (MHA). ARUP *Acinetobacter* complex strains ABC1, ABC2, and ABC3, and CDC strains AB303 and AB307 were streaked on MHA including 2 $\mu\text{g}/\text{mL}$ colistin to remove sensitive bacteria. *Escherichia coli* (ATCC 23724) was streaked on Luria-Bertani agar (LBA) plates. Plates were incubated at 37°C for 8-12 h. Single colonies from the plates were then transferred into Mueller Hinton Broth II (MHB II) (Luria-Bertani broth for *E. coli*) and incubated for 6-8 h at 30°C, 150 rpm. The turbidity of the broth culture was then adjusted to match 0.5 McFarland standard (1×10^8 cells/mL). The adjusted broth culture was diluted 200-fold and used as inoculum for the assay. Each test organism (200 μL) was added to each well of a 96-well flat plate. Compounds were then added using a two-fold dilution scheme starting at 64 $\mu\text{g}/\text{mL}$, with 8 dilutions each. Following 18-20 h incubation, MTT (10 μL ; 5 mg/mL) was added to wells and incubated for 2 h. DMSO (100 μL) was then added to wells and incubated for 1 h. The A_{570} was then measured using a Biotek-Synergy 2 Microplate Reader (Biotek). For *Y. pestis*, testing was performed at 28°C as previously described (ref: Aoyagi et al., 2015) except cation-adjusted Mueller Hinton broth was used instead of HI broth.

Mammalian antiproliferative assay

HEK-293 cells were grown under conditions described above. Cells were seeded (10,000 cells/well) in 96-well plates and treated after 24 h with varying concentrations of the compound. After 72 h, the media was removed and MTT (15 μL ; 5 mg/mL) was added to each well. The plates were then incubated for 3 h at 37°C, 5% CO_2 . After incubation, DMSO (100 μL) was added and absorbance was read at 570 nm using a Biotek-Synergy 2 Microplate Reader (Biotek).

Hemolysis assay

Hemolytic activity of the compounds was measured using freshly purified 0.25% Cgrp grp mouse red blood cell suspension in 1x phosphate buffered saline (PBS). Various concentrations of the compounds were then added to the red blood cell suspension in 2-fold serial dilutions, and incubated for 1 h at 37°C. The suspension was then centrifuged at 1000 x g for 5 min, and the supernatant was collected. Supernatant (100 μL) was then transferred to wells of a 96-well plate. Absorbance was read at 540 nm using a Biotek-Synergy 2 Microplate Reader (Biotek). Results are reported as concentration of compound resulting in 10% hemolysis. Triton X-100 (0.01-10% v/v) served as positive control.

Dilution broth inhibition experiments

A. baumannii (ATCC 19606) was grown overnight, as described above. The correlation between OD_{600} and CFU/mL was determined by plating serial dilutions and counting colonies. Five plates at each concentration were used to count colonies. At the same time, this initial seed culture was used to initiate dilution experiments and was serially diluted 5-fold 10 times, for a final dilution series (1.6×10^8 , 3.36×10^7 , 6.72×10^6 , 1.34×10^6 , 268,800, 53,760, 10,752, 2,150, 430, 86 CFU / mL). Each dilution (200 μL) was placed in 5 replicate wells, for a total of 50 wells, to which was added **2** at 2xMIC (16 $\mu\text{g}/\text{mL}$). In addition, each dilution (200 μL) was added to the plate in triplicate without antibiotic. Two biological replicates were performed, with 96-well plates at 37°C for 96 h. Plates were monitored every 24 h by observing turbidity. At 96 h, plates were removed from incubation, and an MTT test was performed. One plate was used with colistin at 2xMIC in an identical manner. From one of the broth dilutions that was turbid at 2,150 CFU/mL, broth (100 μL) was plated on MHBII in the presence of 2xMIC, yielding 5 colonies. One of these was picked, and its MIC was determined in the same manner described above.

Microscopy of growth-inhibited bacteria

A. baylyi strain ADP1 overnight cultures grown in M9 minimal-succinate medium were diluted 1:4 and grown with shaking for one additional hour at 30°C., followed by spotting on thin minimal-succinate agar pads containing different drugs [turnercyclamycin (8 $\mu\text{g}/\text{mL}$) or colistin (0.5 $\mu\text{g}/\text{mL}$)] (Bailey et al., 2019). Bacteria were imaged after 8 h incubation at 30°C. Terminal morphologies of essential gene deletion mutants were evaluated after transformation with kanamycin-resistance marked PCR fragments as described earlier (Bailey et al., 2019; Gallagher et al., 2020). Phase contrast imaging was performed using a Nikon Eclipse 90 microscope using a 100x oil objective.

***tur* gene clusters analysis in *T. turnerae* strains**

A full or partial *tur* gene cluster was extracted from antiSMASH 5.0 output of each *T. turnerae* genome assembly using multigeneblast with *tur* in T7901 as reference. Protein sequences of a full NRPS genes (according to “CDS” feature) or A-domains (according to “aSDomain” feature) were extracted from the antiSMASH output genbank file using a bash script (supplied in [supplementary information](#)). The identity between NRPS genes or A-domains were calculated by BLASTp search (-outfmt "6 qseqid sseqid pident length evalue qcovs qlen slen"). An identity matrix table was made and plotted in R 3.5 using the pheatmap package.

Spectroscopy

UV-vis spectra were obtained using a Molecular Devices SpectraMax M2 spectrophotometer. IR spectra were obtained using a Nicolet iS50 FT-IR (Thermo Scientific). High resolution mass spectra were acquired using a Waters Xevo G2-XS QToF mass spectrometer equipped with a Zspray ESI source and fed by an Acquity H class UPLC system with a Waters Acquity CSH C18 column (2.1x50mm, 1.8 μ m). NMR data were collected using a Varian 500 MHz NMR spectrometer with 5 mm Varian HCN Oneprobe for proton detected experiments and a 3 mm Varian inverse probe for carbon detected experiments (¹H 500 MHz, ¹³C 125 MHz). Residual signals from solvents were used for referencing. ECD spectra were obtained on an AVIV Biomedical, Inc. CD Spec Model 410 (Lake-wood, NJ, USA). Analytical and semi-preparative HPLC was performed on a Thermo UltiMate 3000 system with a DAD detector.

QUANTIFICATION AND STATISTICAL ANALYSIS

All MIC data was graphed in GraphPad Prism 6.0 software. Error bars on MIC graphs were indicated as mean \pm SD of 3 replicate wells. The dose response curves for **1** and **2** against the HEK-293 cell line were plotted in GraphPad Prism 6.0 software, and non-linear regression failed to calculate an IC₅₀ value due to a lack in inhibitory activity. Error bars were indicated as mean \pm SD of 3 replicate wells. Hemolysis data were graphed in GraphPad Prism 6.0 software. Error bars were indicated as mean \pm SD of 3 replicate wells.