



NOAA- Technical Memorandum NMFS-AFSC-35

**Relative Abundance and
Size Composition of Sablefish
(*Anoplopoma fimbria*) in the
Coastal Waters of California and
Southern Oregon, 1984-1991**

by
N. B. Parks and F. R. Shaw

U.S. DEPARTMENT OF COMMERCE
National Oceanic and Atmospheric Administration
National Marine Fisheries Service
Alaska Fisheries Science Center

March 1994

NOAA Technical Memorandum NMFS

The National Marine Fisheries Service's Alaska Fisheries Science Center uses the NOAA Technical Memorandum series to issue informal scientific and technical publications when complete formal review and editorial processing are not appropriate or feasible. Documents within this series reflect sound professional work and may be referenced in the formal scientific and technical literature.

The NMFS-AFSC Technical Memorandum series of the Alaska Fisheries Science Center continues the NMFS-F/NWC series established in 1970 by the Northwest Fisheries Center. The new NMFS-NWFSC series will be used by the Northwest Fisheries Science Center.

This document should be cited as follows:

Parks, N. B., and F. R. Shaw. 1994. Relative abundance and size composition of sablefish (Anoplopoma fimbria) in the coastal waters of California and Southern Oregon, 1984-1991. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-35, 38 p.

Reference in this document to trade names does not imply endorsement by the National Marine Fisheries Service, NOAA.

GENERAL DISCLAIMER

This document may be affected by one or more of the following statements

- **This document has been reproduced from the best copy furnished by the sponsoring agency. It is being released in the interest of making available as much information as possible.**
- **This document may contain data which exceeds the sheet parameters.. It was furnished in this condition by the sponsoring agency and is the best copy available.**
- **This document may contain tone-on-tone or color graphs, charts and/or pictures which have been reproduced in black and white.**
- **This document is paginated as submitted by the original source.**
- **Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.**

BIBLIOGRAPHIC INFORMATION

PB94-184298

Report Nos: NOAA-TM-NMFS-AFSC-35

Title: Relative Abundance and Size Composition of Sablefish ('Anoplopoma fimbria') in the Coastal Waters of California and Southern Oregon, 1984-1991.

Date: Mar 94

Authors: N. B. Parks and F. R. Shaw

Performing Organization: Alaska Fisheries Science Center, Seattle, WA.

Type of Report and Period Covered: Technical memo.

Supplemental Notes: See also PB90-103961.

NTIS Field/Group Codes: 98F (Fisheries & Aquaculture), 47D (Biological Oceanography)

Price: PC A03/MF A01

Availability : Available from the National Technical Information Service, Springfield, VA. 22161

Number of Pages: 44p

Keywords: *Abundance, *Size determination, *Fisheries, Coastal waters, California, Graphs(Charts), Surveys, *Anoplopoma fimbria, *Sablefish, Southern region (Oregon), Catch statistics.

Abstract: Trap survey results indicate that the relative abundance of sablefish (*Anoplopoma fimbria*) off the coasts of California and southernmost Oregon decreased by 66% in numbers and 64% in kilograms per trap between 1988 and 1991. Catch rates decreased at all nine sites sampled. The largest decreases in catch per unit of effort were in the central part of the survey area; declines of between 70% and 84% were observed at the Point Arena, Half Moon Bay, Carmel Bay, and Morro Bay sites. Smaller declines in catch rate (24% to 57%) were observed at the northern (Cape Sebastian, Point Saint George, and Cape Mendocino) and Southern (Cartes Bank) sites. Small and undersized sablefish made up 83% of the catch, down from 90% in 1988, whereas medium and large sablefish comprised 14% and 3% of the catch, respectively. Highest catch rates (number/trap) occurred at the depths of 225 and 300 fathoms. Mean length increased with depth and generally decreased from north to south within the sampling area.



NOAA Technical Memorandum NMFS-AFSC-35

**Relative Abundance and
Size Composition of Sablefish
(Anhplopoma fimbria) in the
Coastal Waters of California and
Southern Oregon, 1984-1991.**

**b y
N. B. Parks and F. R. Shaw**

**Alaska Fisheries Science Center
7600 Sand Point Way N.E., BIN C-I 5700
Seattle, WA 981156070**

**U.S. DEPARTMENT OF COMMERCE
Ronald H. Brown, Secretary
National Oceanic and Atmospheric Administration
D. James Baker, Under Secretary and Administrator
National Marine Fisheries Service
Rolland A. Schmitten, Assistant Administrator for Fisheries**

March 1994

ABSTRACT

Trap survey results indicate that the relative abundance of sablefish (Anoplonoma fimbria) off the coasts of California and southernmost Oregon decreased by 66% in numbers and 64% in kilograms per trap between 1988 and 1991. Catch rates decreased at all nine sites sampled. The largest decreases in catch per unit of effort were in the central part of the survey area; declines of between 70% and 84% were observed at the Pt. Arena, Half Moon Bay, Cannel Bay, and Morro Bay sites. Smaller declines in catch rate (24% to 57%) were observed at the northern (Cape Sebastian, Pt. St. George, and Cape Mendocino) and southern (Cortes Bank) sites. Small and undersized sablefish made up 83% of the catch, down from 90% in 1988, whereas medium and large sablefish comprised 14% and 3% of the catch, respectively. Highest catch rates (number/trap) occurred at the depths of 225 and 300 fathoms'. Mean length increased with depth and generally decreased from north to south within the sampling area.†

THIS PAGE INTENTIONALLY LEFT BLANK

CONTENTS

	Page
INTRODUCTION	1
SURVEY METHODS AND- GEAR	2
RESULTS AND DISCUSSION	7
Catch Rates and Size Composition ;	8
Mean Length by Depth and by Latitude	16
Percentage of Females by 'Depth	16
* 8 Length-Weight Relationship	21
SUMMARY AND CONCLUSIONS	24.
CITATIONS :	26.
APPENDIX	31

INTRODUCTION

The sablefish (Anoplonoma fimbria) has been the most valuable commercial groundfish species landed on the West Coast in most years from 1985 to 1990 with an estimated ex-vessel value averaging \$11.5 million.' High demand and relatively high prices have resulted in heavy fishing pressure on the stocks. The 1989 optimum yield was set at 10,400 metric tons (t) and was reduced to 8,900 t in 1990 and 1991 (Methot and Hightower 1990). Methot (1991) estimated the long-term potential yield to be 6,900 t.

Alaska Fisheries Science Center (AFSC) trap surveys to measure changes in sablefish relative abundance were begun in Oregon and Washington waters in 1979 and extended to waters off California in 1980 (Parks and Hughes 1981; Parks 1982, 1984; and Parks and Shaw 1983, 1985, 1987, 1988a, 1988b, 1989, 1990). These surveys measure catch per unit effort (CPUE) at standard sites to use as indices of sablefish abundance for assessing the condition of the stock. Collecting biological data such as the state of maturity and the weight, length, and age composition of the stock are other important objectives of these surveys. A 1 1 fish released alive were tagged as part of an age validation study in which 75% were injected with oxytetracycline -(OTC) to mark the bony structures used to age the fish (otoliths), while

¹ Pacific Coast Fisheries Information Network (PacFin) as of 30 December 1991. Pacific Marine Fisheries Commission, Metro Center - Suite 170, 2000 S.W. First Ave., Portland, Oregon 97201-5346.

25% were not injected to be used as controls. This report presents results of the 1991 survey off California and southernmost Oregon and compares them with results of previous trap surveys and other assessments of the condition of the West Coast sablefish stock(s).

SURVEY METHODS AND GEAR

The 1991 survey methods were identical to the 1984 survey methods (Parks and Shaw 1985) except that conical traps have been used exclusively since 1986. Methods and trap gear used in earlier surveys are described in Parks and Shaw (1983).

The conical traps employed in this study were constructed with a 137 cm-diameter bottom ring and an 65 cm diameter top ring (Fig. 1). The traps were 71 cm high with a tunnel entrance on the side: The framework was covered with 5.7 cm 42-thread nylon webbing,. Tunnels were constructed of 5.1 cm nylon knotless web and rigged with a 'noose arrangement which closed by a magnesium alloy time-release device 24 (+ 1) hours after setting. Ten traps were attached with C-hooks to gangions at 50 fathom (fm) (91 m) intervals on a 550 fm groundline of 1.59 cm synthetic line (Fig. 2). A perforated plastic bait jar containing approximately 2 lb of chopped herring was hung in each trap.

Sampling was conducted during 7 September to 7,October 1991; from north to south at nine index sites off California and southern Oregon (Fig. 3). These sites were also sampled in 1984, 1986, and 1988. Ten-trap groundlines were fished along the 225,




Figure 1. --Conical trap used in recent sablefish abundance indexing surveys.




Figure 2. --A pictorial view of a string of conical trap gear used in sablefish indexing surveys.




Figure 3. --Sites sampled off California and southern Oregon (International North Pacific Fisheries Commission areas Eureka, Monterey, and Conception) during the 1991 sablefish abundance indexing survey.

300, 375; 450, and 525 fm isobaths (standard depths) at all sites. Additional strings were set opportunistically at 150 fm' and between 600 and 900 fm to obtain information on the portions of the population in depths shallower and deeper than those routinely sampled. Two sets were made at each depth at most sites. Loran C, Global Positioning System (GPS), and depth sounders were used to position replicate sets as near as possible to the locations of the original sets.

At each site, we collected data on the number and weight of each species captured in each trap, lengths of all sablefish, and sablefish biological specimens. and data including otoliths, individual weight, sex, and sexual maturity from a random sample of the first 20 fish taken at each depth at each site.

An age-validation study was initiated on the 1991 survey in which sablefish were injected with OTC to mark the age structure at the point in time that the fish was tagged. Approximately 75% of all sablefish tagged were injected with an intraperitoneal dose of 30 mg of OTC per kilogram of weight. The remaining 25% constituted the control group to determine to what extent the OTC injection affects survival. Sablefish not required for biological samples were placed into live tanks supplied with fresh running seawater immediately upon capture. Usually within 15 minutes of the completion of each haul, all viable sablefish were measured to the nearest centimeter in a padded tagging

cradle, injected, and tagged with a single blue Floy² anchor tag implanted just below the first dorsal fin. Anesthetic was not used.

RESULTS AND DISCUSSION

All nine index sites were sampled at the 150, 225, 300, 375, 450, and 525 fm depths. The 600-900 fm depth interval was sampled at all sites except those off Cape Sebastian and Pt. Arena. Adverse weather prevented fishing at these depths and also prevented the second replicate of the 375, 450, and 525 fm sets off Cape Sebastian.

Same data standardization was necessary at sites where traps were lost or conditions precluded replicating a set. If some traps on a string were lost, the observed catch for the string was adjusted by assigning the average catch in the remaining traps to lost traps. Occasionally, due to weather, gear conflict, or time constraints, the replicate set at a site could not be completed. In those cases, the catch in the replicate set (needed because the index is based on the average of both sets) was estimated by calculating the rate of decline in CPUE between the first and second sets at all other sites and depths and applying that rate to the catch from the single set; in 1991, the catch on the second set averaged 0.701 of the catch on the first

² Reference to trade name does not imply endorsement by the National Marine Fisheries Service, NOAA.

set,, Catch and size data from individual sampling sites are given in the Appendix.

Catch Rates and Size Composition

Catch rates (average number of fish per trap), based on data from standard survey depths (225-525 fm) at the nine abundance index sites, decreased 66% from 1988 to 1991. The mean catch rate of 3.6 sablefish. per trap also represents a decrease of 22% from 1986 and 63% from 1984 (Fig. 4). Standard errors and coefficients of variation associated with 1984, 1986, 1988, and 1991. mean catches per trap are shown in Table 1. Pair-wise comparisons between the catch rates (average number of sablefish, per trap). among survey years were tested at a level of significance of $c = 0.05$. The, Bonferroni joint' testing procedure (Neter et al. 1985) was used to adjust for the multiple between-. ' year comparisons. Only two of the six possible between-year comparisons were not significantly different (1984-88 and 1986-91, Table 2).

The mean catch rate of small sablefish (<1.93 kg) in 1991 decreased by 69% since 1988; The CPUE of large fish (>3.18 kg) held relatively steady at just under 0.1 per trap, while the CPUE of medium-sized sablefish (1.93 to 3.18 kg) declined 48%. The 1991 catch rates for all size groups are lower than in any of the previous surveys., Catch rates in 1991 were lower at all sites with the largest decreases occurring at Pt. Arena, Half Moon Bay,




Figure 4. --Mean catch rates of sablefish (number/trap) by size category at standard depths (225-525 fathoms) at two index sites off California in 1980-82 and nine index sites off California and southern Oregon in later years.

Table 1. --Mean catch rates, standard errors, and coefficients of variation for sablefish at index sites off California and southern Oregon from 1984 to 1991 (standard depths 225-525 fathoms).

Year	Mean catch rate	Standard error	Coefficients of variation
1984	8.8	0.663	7.5
1986	4.6	0.345	7.5
1988	10.6	1.164	11.0
1991	3.6	0.481	13.4

Table 2. --Pair-wise -comparisons between the catch rates (average numbers of sablefish/trap) from various survey years. Significance judged at 0.05% probability level.

Year	Difference between average catch rates,	Standard error of difference	95% confidence interval	Statistically significant
1984-86.	4.2	0.658	(1.46, 6.94)	Yes
1984 - 88	-1.8	1.340	(-0.94, 4.54)	no
1984-91	5.2	0.819	(2.46, 7.94)	y e s
1986-88	-6.0	1.214	(3.26, 8.74)	yes
1986-91	1.0	0.592	(-1.74, .3.74)	no'
1980-91	7.0	1.260	(4.26, 9.74)	yes

Carmel Bay, Morro Bay, and Cape Mendocino (-84%, -83%, -76%, -75%, and -70%, respectively). Catch rates declined less at Cape Sebastian, Cortes Bank, and Pt. St. George (-24%; -46%, and -57%, respectively). Average sablefish catch rates in weight (kilograms per trap) from the standard depths decreased 64% between 1988 and 1991 (Fig. 5). The 1991 value (5.4 kg/trap) was down 33% from the previous low in 1986 (8.0 kg/trap).

In 1991, the highest mean catch rates (number/trap) within the standard sampling depths (225-525 fm) occurred at 225 and 300 fm (Table 3). Catch rates were lowest at 450, 525, and >600 fm at 1.3, 1.2, and 1.5 fish/trap, respectively.

The sablefish size composition from the 1991 survey was unimodal, peaking at 48 cm (Fig. 6). Mean length declined from 53.1 and 53.7 cm in 1984, and 1986 to 50.8 and 51.7 cm in 1988 and 1991, respectively. The proportion of small and undersized sablefish (<1.93 kg) was 75% of the catch in 1986, 90% in 1988, and 83% in 1991, whereas the proportions of medium (1.93 to 3.18 kg) and large (>3.18 kg) sablefish in 1991 were up slightly from 1988 to 14% and 3%, respectively (Table 3, Fig. 6). These proportions are still well below those seen in 1984 and 1986.

The sex-specific size composition for sablefish taken at the standard depths during the 1991 survey is shown in 'Figure 7. The mean sizes of males and females were 50.8 and 56.0 cm,

respectively. Based upon calculations of length at 50% maturity from 1986 survey data (51.9 cm for males and 57.9 cm for females,




Figure. 5.--Mean catch rates of sablefish (kg/trap) at standard depths (225-525 fathoms) at the two original index sites off California 1980-82, and at nine sites off the coasts of California and southernmost Oregon in later years..

Table 3.-- The numbers of sablefish, average numbers and weight by trap, and percentage of sablefish by size^a captured at southernmost Oregon and California by depth in 1984, 1986, 1988, and 1991.

DEPTH (FM)	YEAR	TOTAL NO. SABLEFISH	AVG. NO. SABLEFISH PER TRAP	AVG. WT. (KG) PER TRAP	% LARGE SABLEFISH	% MEDIUM SABLEFISH	% SMALL SABLEFISH	% UNDERSIZED SABLEFISH
150 ^d	1984	420	5.2	7.3	1	11	26	62
	1986	286 ^c	2.4	3.1	3	9	21	67
	1988	1,104 ^c	6.3	7.7	<1	5	17	77
	1991	435	2.4	2.9	2	9	14	75
225	1984	2,736	15.2	23.1	2	16	32	50
	1986	1,262 ^c	7.0	10.6	2	18	25	55
	1988	4,136 ^c	23.0	28.5	<1	5	16	79
	1991	1,338	7.4	9.2	<1	7	13	79
300	1984	2,571	14.3	22.2	2	14	32	52
	1986	1,105 ^c	6.1	10.5	4	20	31	45
	1988	2,382 ^c	13.2	17.8	<1	7	18	74
	1991	913	5.1	7.8	1	15	22	62
375	1984	1,499	8.3	12.9	3	16	36	45
	1986	689	3.8	7.4	7	20	37	36
	1988	1,053 ^c	5.8	8.4	<1	9	24	66
	1991	492 ^c	2.7	4.6	4	19	27	50
450	1984	1,263 ^c	7.0	11.6	2	17	41	40
	1986	595 ^c	3.3	5.9	3	19	38	40
	1988	932 ^c	5.2	8.2	2	14	29	55
	1991	236 ^c	1.3	2.4	7	22	32	39
525	1984	757	4.2	8.5	5	34	44	17
	1986	479	2.7	5.4	4	33	45	18
	1988	1,075 ^c	6.0	11.5	2	28	42	27
	1991	223 ^c	1.2	2.7	10	41	37	12
> 600 ^e	1986	189	2.4	8.4	47	49	4	0
	1988	502 ^c	3.1	9.8	33	54	11	2
	1991	214 ^c	1.5	5.7	48	48	4	0
All depths combined	1984	9,246 ^c	9.4	15.0	2	17	35	46
	1986	4,605 ^c	4.2	7.5	6	21	31	42
	1988	11,094 ^c	9.1	13.3	2	10	21	67
	1991	3,851 ^c	3.2	5.0	5	16	19	60
Standard depths combined	1984	8,826 ^c	9.8 ^f	15.7	2	17	35	46
	1986	4,130 ^c	4.6 ^f	8.0	4	21	33	42
	1988	9,579 ^c	10.6 ^f	14.9	<1	9	22	68
	1991	3,202 ^c	3.6	5.4	3	14	21	62

^aLarge = over 3.18 kg round weight (rd. wt.); medium = 1.93-3.18 kg rd. wt.; small = 1.36-1.93 kg rd. wt.; and undersized = less than 1.36 kg rd. wt.

^bThe 1984 numbers are the result of upward adjustment on rectangular trap catches by 1.265 (a scaling factor to compensate for the lesser fishing efficiency of the rectangular traps) plus conical trap catches.

^cAdjusted upward for missing string(s) of gear, second repetitions not made, or lost traps, based on the average decline of catch from the first set to the second set or the average catch in remaining traps.

^d150 fathom depth fished at the four southern sites only in 1984, at the three northern and three southern sites only in 1986; and at eight sites in 1988.

^e>600 fathom depth not fished in 1984, fished at the four northern sites only in 1986, and fished at the eight southern sites in 1988, and Cape Sebastian and Pt. Arena sites not fished in 1991.

^fStandard error = 0.750 for 1984, 0.345 for 1986, and 1.16 for 1988.




Figure 6. --Sablefish length compositions for all southern Oregon and California index sites combined at standard depths (225-525 fathoms) for 1984, 1986, 1988, and 1991.




Figure 7.--Sablefish length compositions by sex of random samples taken from standard depths (225-525 fathoms) during the 1991 abundance index survey.

Parks and Shaw 1988a) we calculated that 65% of the males and 68% of the females 'within the standard depths (225-525 fm) were immature.

Mean Length by Depth and by Latitude

Sablefish length compositions were summarized by depth for the 1991 survey. (Fig. 8). Mean lengths typically increase with depth (Parks and Shaw 1989) and this was again-true in 1991. The largest increases occurred between 450 and 525 fm, where mean lengths increased by just over 3 cm, and between 525 and >600 fm, where mean lengths increased approximately 9 cm'. The mean lengths of sablefish captured at depths greater than 600 fm. averaged about 19 cm greater-than sablefish captured at 150 fm. In the 1984 and 1986 surveys, mean length, generally decreased from north to south (Parks and Shaw 1989). In the 1988 and 1991 surveys, mean length was generally higher at the four northern sites with the largest fish at the Pt. Arena site -and lower at the four southern sites (Fig. 9).

Percentage of Females by Depth .

The percentage of females of our random samples by depth in all four survey years showed that females were generally most abundant at the. shallowest and deepest depths and least abundant; at the intermediate depths (Fig. 10). These trends in the percentage of females were relatively consistent for most survey




Figure 8.--Sablefish length compositions and mean lengths by depth for all sites combined during the 1991 abundance index survey (>600 fathoms fished at all sites except Cape Sebastian and Pt. Arena).




Figure 9.--Sablefish mean fork length by site from southern Oregon and California abundance index surveys in 1984, 1986, 1988, and 1991 (standard depths 225-525 fathoms).




Figure 10. --Percent of females in random samples of sablefish from National Marine Fisheries Service abundance index surveys by depth and by survey year.

years, especially at depths greater than 600 fm where the percent of females ranged from 89 to 97%. In the 1991 survey, the percentage of females at 375 and 450 fm was much higher than in the three- earlier survey years. In standard depths (225-525 fm), females made up 52 to 62% of our random samples in all four survey years (Fig. 11). For all depths 'combined (150->600 fm), females made up 57 to 67% of our samples from the 1986, 1988, and 1991 surveys. This predominance of females, which also occurs in landings of the fixed-gear fishery, may be a result of the disproportional harvest of males by the trawl fishery. In trawl-caught samples of sablefish in the small size category (less than 58 centimeters fork length) from the Eureka and Monterey International North Pacific Fisheries Commission areas, a 5,000 fish sample (1,000 fish per year from 1986 through 1991) showed 56.4% to be males.

Length-Weight Relationship

The length-weight, relationship for male and female sablefish from the 1991 survey is shown in Figure 12. We found no significant difference in the average weight at length between males and females, as was the case for surveys conducted off Oregon and Washington (August and September 1987) and off southern Oregon and California (September to mid-November 1986) (Parks and Shaw 1988a, b). In the 1988 survey, the male and female length-weight relationships were significantly different




Figure 11.--Percentage of females in random samples of sablefish from the National Marine Fisheries Service abundance index surveys at the standard depths. (225-525, fathoms) and at all depths sampled (150->600 fathoms) by survey year.. Only standard depths were sampled in 1984.




Figure 12. --Length-weight relationship derived for sablefish (sexes combined) collected off southern Oregon and California in September and October 1991.

at the 99.9% level (Parks and Shaw 1989). The 1991 length-weight relationship for sexes combined is $W = .0017812 \times L^{3.4300R}$, where W = weight, in grams and L = length in centimeters (Fig. 12).

SUMMARY AND CONCLUSIONS

Sablefish catch rates (mean number per trap) in surveys off California and southern Oregon decreased by 66% between 1988 and **199:L**. Catch rates of small and undersized sablefish (<1.93 kg) declined 69% from 1988 to 1991, whereas catch rates of medium-size sablefish (1.93-3.18 kg) declined 48% and large-size sablefish (>3.18 kg) declined 3%. Catch rates decreased at all sites in 1991, with the largest decreases generally occurring from Pt. Delgada southward to Morro Bay. The three southern sites had the highest catch rates. At the Cortes Banksite, where fishing pressure is probably the lowest, the catch rate was over twice as great as the rate at the next highest site, Carmel Bay (Appendix-Fig.1). The lowest catch rates were found at the four sites between Cape Mendocino and Half Moon Bay; Tagging studies indicate that approximately 29% of the sablefish tagged in the Conception INPFC area were recovered north of the Conception INPFC area where fishing effort is greater (Shaw 1984). Because fishing pressure is lower at the two southern sites, a higher percentage of sablefish tagged in the Conception INPFC area was recovered outside that area than in other INPFC areas.

In 1991, over 62% of our catch consisted of undersized (<1.36 kg) sablefish, down from 68% in 1988. The proportion of small (1.36 to 1.93 kg) sablefish changed only slightly from 22% in 1988 to 21% in 1991, while the proportion of medium and large fish rose 9% to 14% and from 1% to 3%, respectively (Appendix Table 1). The mean length of males was 50.8 cm and the mean length of females was 56.0 cm (Fig. 7). Mean lengths increased with depth as, in previous surveys (Parks and Shaw 1989). Sablefish mean lengths were highest at the northern sites, peaked at the Pt. Arena site, and were lowest the four southern sites (Appendix Fig. 2).

While the 1988 survey indicated good recruitment of 2+ and 3+ year old (40-50 cm) sablefish, the low catches of small and medium fish in 1991 suggest that this substantial group of fish has apparently been removed from the population since the 1988 survey. As in the 1988 survey, the mean lengths for both sexes are shorter than our calculated lengths at 50% maturity (51.9 cm for males and 57.9 cm for females) (Parks and Shaw 1988a). We estimate that only about one-third of the population surveyed in both 1988 and 1991 is sexually mature. The size composition of the commercial catches indicate that large numbers of sablefish are being harvested before they have an opportunity to spawn.

CITATIONS

- Methot, R. D. 1991. Sablefish. In L. L. Low (editor),, Status of living marine resources off the Pacific Coast of the United States as assessed in 1991% U.S. Dep. Commer., NOAA Tech. Memo. NMFS F/NWC-210, 69 p.
- Methot, R. 1992. Assessment of the west coast sablefish stock in 1992. In Status of the Pacific coast groundfish fishery through 1992 and recommended acceptable biological catches for 1993. Pacific Fishery Management Council, Metro Center, Suite 420, 2000 S.W. First Avenue, Portland, OR 97201.
- Methot, R., and J. Hightower. 1990. Assessment of the west coast sablefish stock in 1990 ,and recommendations to management in 1991, 67 p. In Status of the Pacific-coast groundfish fishery through 1990. and recommended acceptable biological catches for 1991. Stock assessment and fishery evaluation, Appendix Vol. I. Pacific Fishery Management Council, Metro Center, Suite 420, 2000 S.W. First Avenue, Portland, OR 97201.
- Neter, J., W. Wasserman, and M. H. Kutner. 1985. Applied linear statistical models. Richard D. Irwin, Inc.; Homewood, IL, 1 , 1 2 7 p .

Parks, N. B. 1982.. "Changes in relative abundance and size composition of sablefish in coastal waters of Washington and Oregon, 1979-81, and California, 1980-81. U.S. Dep. Commer., NOM Tech. Memo. NMFS F/NWC-26, 28 p.

Parks, N. B. 1984. Changes in relative abundance and size composition of sablefish in coastal waters of Washington and Oregon, 1979-83.. U.S. Dep. Commer'. , NOAA Tech. Memo. NMFS F/NWC-61, 23 p.

Parks, N. B., and S. E. Hughes. 1981. Changes in relative abundance and size composition of sablefish in coastal waters of Washington and Oregon, 1979-80. U.S. Dep. Commer. NOM Tech. Memo., NMFS F/NWC-8, 25 p.

Parks, N. B., and F. R. Shaw. 1983. Changes in relative abundance and size composition of sablefish (Anoploma fimbria) in coastal waters of California, 1980-82. U.S. Dep. Commer., NOM Tech.. Memo. NMFS F/NWC-51, 16p.


Parks, N. B., and F. R. Shaw. 1985. Abundance and size composition of sablefish (Anoploma fimbria) in the coastal waters of California and Oregon in 1984. NWAFC Processed Rep. 85-16, 21 p. Alaska Fish. Sci. Cent. Natl. Mar. Fish. Serv., NOAA, 7600 Sand Point Way NE., Seattle, WA 98115.

- Parks, N. B., and F. R. Shaw. 1987: Changes in relative abundance and size composition of sablefish in coastal waters of Washington and Oregon 1979-85. U.S. Dep. Commer., NOM Tech. Memo NMFS F/NWC-124, 41 p.
- Parks, N. B., and F. R. Shaw. 1988a. Abundance and size composition of sablefish (Anoplonoma fimbria) in the coastal waters of California and southern Oregon, 1984-86. U.S.-Dep. Commer., NOM Tech., Memo. F/NWC-125, 29 p.
- Parks, N. B., and F. R. Shaw. 1988b: Changes in relative abundance and size composition of sablefish in coastal waters of Washington and Oregon, 1979-87. U.S. Dep. Commer., NOAA Tech. Memo. F/NWC-141, 33 p.
- Parks, N. B.; and F. R. Shaw. 1989. Relative abundance and size composition of sablefish- (Anoplonoma fimbria) in the coastal waters of California and southern Oregon, 1984-1988. 'U.S. Dep. Commer., NOAA Tech. Memo. F/NWC-167, 32 p.
- Parks, N. B., and F. R. Shaw. 1990. Changes in relative abundance and size composition of sablefish in coastal waters of Washington and Oregon, 1979-89. U.S. Dep. Commer., NOM Tech. Memo. F/NWC-188, 38 p.

Shaw, F. R. 1984. Data report: Results of sablefish tagging in waters off the coast of Washington, Oregon, and California, 1979-83. U.S. Dep. Commer., NOM Tech. Memo. F/NWC-69, 79 p.

APPENDIX







Individual Index Site Data



1984	7.5	8.4	8.2	7.4	9.3	15.2	10.9	6.6	14.7
1986	6.1	4.0	4.5	4.3	3.2	2.3	5.8	6.2	4.9
1988	4.7	8.1	4.6	7.3	10.8	8.9	18.8	15.5	17.2
1991	3.6	3.4	1.8	2.2	1.8	1.6	4.5	3.8	9.3

s i t e s

Appendix Figure 1.-- Mean catch rates of sablefish at nine index sites fished off California and southern Oregon in 1984, 1986, 1988, and 1991 (225525 fathoms),



34

Appendix Figure 2.--Sablefish length compositions by abundance indexing site at the standard depths (225 - 525 fathoms) during the 1991 abundance indexing survey.

Appendix Table 1.-- Sablefish catch rates and percentage by size category from southern Oregon and California index sites by depth in 1984, 1986, 1988, and 1991 (standard depths 225-525 fathoms).

	YEAR	TOTAL NO. SABLEFISH	AVG. NO. SABLEFISH PER TRAP	AVG. WT. (KG) PER TRAP	% LARGE ^b SABLEFISH	% MEDIUM ^c SABLEFISH	% SMALL ^d SABLEFISH	% UNDERSIZED ^e SABLEFISH
Cape Sebastian, OR	1984	748 ^f	7.5	14.3	7	19	25	49
	1986	613 ^f	6.1	11.7	9	21	32	38
	1988	472 ^f	4.7	7.8	2	16	32	50
	1991	359 ^f	3.6	5.7	3	14	18	65
Pt. St. George, CA	1984	843 ^f	8.4	14.4	3	22	34	41
	1986	405 ^f	4.0	7.6	7	28	35	30
	1988	807 ^f	8.1	9.7	2	6	17	75
	1991	343	3.4	6.1	4	20	26	50
Cape Mendocino, CA	1984	819 ^f	8.2	14.9	5	23	30	42
	1986	448 ^f	4.5	8.2	5	29	31	35
	1988	458 ^f	4.6	6.9	3	14	27	56
	1991	182 ^f	1.8	3.0	4	18	28	50
Pt. Delgada, CA	1984	735 ^f	7.4	13.2	4	23	39	34
	1986	427 ^f	4.3	8.3	5	26	32	37
	1988	726 ^f	7.3	10.5	2	11	25	62
	1991	220	2.2	3.7	6	21	22	51
Pt. Arena, CA	1984	928	9.3	15.4	1	20	41	38
	1986	325	3.2	6.0	4	25	33	39
	1988	1,079 ^f	10.8	17.6	2	17	33	48
	1991	178 ^f	1.8	3.5	10	35	27	28
Half Moon Bay, CA	1984	1,525	15.2	23.4	<1	15	40	44
	1986	226 ^f	2.3	3.9	1	20	38	41
	1988	892 ^f	8.9	12.0	<1	7	20	73
	1991	155	1.6	2.8	<1	11	24	64
Carmel Bay, CA	1984	1,094	10.9	16.6	<1	14	31	51
	1986	580	5.8	9.4	1	17	36	46
	1988	1,875	18.8	24.8	<1	7	18	75
	1991	453	4.5	6.4	1	14	20	65
Morro Bay, CA	1984	664	6.6	11.1	2	17	37	44
	1986	619 ^f	6.2	9.1	<1	13	27	59
	1988	1,546 ^f	15.5	21.0	<1	7	16	77
	1991	383	3.8	5.2	<1	9	14	76
Cortes Bank, CA	1984	1,470	14.7	20.0	<1	10	30	59
	1986	487 ^f	4.9	7.7	1	19	36	44
	1988	1,724 ^f	17.2	23.5	<1	9	22	69
	1991	929	9.3	12.2	1	9	20	70
All sites combined	1984	8,826 ^f	9.8	15.9	2	17	35	46
	1986	4,130 ^f	4.6	7.9	4	21	33	42
	1988	9,579 ^f	10.6	14.9	<1	9	22	68
	1991	3,202 ^f	3.6	5.3	3	14	21	62

^a These numbers are the result of upward adjustment on rectangular trap catches by 1.265 plus conical trap catches.

^b Over 3.18 kg round weight.

^c 1.93-3.18 kg round weight.

^d 1.36-1.93 kg round weight.

^e Less than 1.36 kg round weight.

^f Adjusted upward for missing string(s) of gear, second repetition which was not made, or individual traps which were lost, based on the average decline of catch from the first set to the second set.

Appendix Table 2.-- Sablefish catch rates and percentage above the minimum size limit' by index site during the 1991 abundance index survey.

DEPTH (FM)	Cape Sebastian, OR				Pt. St. George, CA				Cape Mendocino, CA			
	NO. OF FISH	NO. PER TRAP	WT. PER TRAP (KG)	% ABOVE SIZE LIMIT	NO. OF FISH	NO. PER TRAP	WT. PER TRAP (KG)	% ABOVE SIZE LIMIT	NO. OF FISH	NO. PER TRAP	WT. PER TRAP (KG)	% ABOVE SIZE LIMIT
150	3	0.2	0.1	50	80	4.0	6.2	42	108	5.4	6.9	32
225	105	5.2	7.6	30	131	6.6	9.9	37	87	4.4	6.1	31
300	100	5.0	7.7	29	123	6.2	10.2	44	28	1.4	1.8	32
375	124 ^b	6.2	10.4	44	53	2.6	6.0	70	27	1.4	2.9	70
450	20 ^b	1.0	1.6	50	23	1.2	2.9	87	17	0.8	1.7	88
525	10 ^b	0.5	1.4	83	13	0.6	1.4	85	23 ^b	1.2	2.7	100
650-850	--	--	--	--	20	1.0	3.4	100	20 ^b	1.0	4.8	100
All Depths	362	3.0	4.8	35	443	3.2	5.7	51	310	2.2	3.8	47
	Pt. Delgada, CA				Pt. Arena, CA				Half Moon Bay, CA			
150	14	0.7	1.1	57	19	0.9	1.2	32	42	2.1	2.4	31
225	103	5.2	7.9	37	42 ^b	2.1	3.2	28	33	1.6	1.9	15
300	51 ^b	2.6	4.0	44	38	1.9	3.8	76	33	1.6	1.9	21
375	32	1.6	2.5	50	36 ^b	1.8	3.8	71	49	2.4	3.2	29
450	21	1.0	2.5	90	17	0.8	1.9	82	8	0.4	0.6	38
525	13	0.6	1.5	92	45	2.2	4.9	93	32	1.6	3.4	84
650-850	36	1.8	6.3	100	--	--	--	--	58 ^b	2.9	11.9	100
All Depths	270	1.9	3.7	56	197	1.6	3.1	68	255	1.8	3.6	49

Appendix Table 2.--(Continued).

DEPTH (FM)	Carmel Bay, CA				Morro Bay, CA				Cortes Bank, CA			
	NO. OF FISH	NO. PER TRAP	WT. PER TRAP (KG)	% ABOVE SIZE LIMIT	NO. OF FISH	NO. PER TRAP	WT. PER TRAP (KG)	% ABOVE SIZE LIMIT	NO. OF FISH	NO. PER TRAP	WT. PER TRAP (KG)	% ABOVE SIZE LIMIT
150	50 ^b	2.5	2.1	6	30	1.5	1.5	3	89	4.4	4.4	8
225	201	10.0	12.1	20	150	7.5	7.7	5	486	24.3	25.9	14
300	139	7.0	10.2	34	133	6.6	9.1	22	268	13.4	21.6	47
375	49	2.4	3.5	43	30	1.5	2.5	43	92	4.6	7.0	47
450	23	1.2	2.1	65	41	2.0	3.5	54	66	3.3	5.0	38
525	41	2.0	4.4	85	29	1.4	2.9	76	17	0.8	1.6	88
630-850	23	1.2	4.2	100	46	2.3	7.7	100	11	0.6	1.5	100
All Depths	526	3.8	5.5	36	459	3.3	5.0	31	1,029	7.4	9.6	28

All sites combined				
150	435 ^b	2.4	2.9	25
225	1,338 ^b	7.4	9.2	21
300	913 ^b	5.1	7.8	39
375	492 ^b	2.7	4.6	50
450	236 ^b	1.3	2.4	61
525	223 ^b	1.2	2.7	88
630-850 ^c	214	1.5	5.6	100
All Depths	3,851	3.2	5.0	40

^a Regulation set by the Pacific Fishery Management Council and includes sablefish smaller than 22 inches total length (52.4 cm fork length) or 15.5 inches dorsal length (origin of dorsal fin to the tip of the tail if beheaded).

^b Adjusted upward for second set not made or for lost traps.

^c 630-850 fathom depth fished at seven sites.

RECENT TECHNICAL MEMORANDUMS

Copies of this and other NOAA Technical Memorandums are available from the National Technical Information Service, 5285 Port Royal Road, Springfield, VA 22167. Paper copies vary in price. Microfiche copies cost \$3.50.

AFSC-

- 34 SAMPLE, T. M., and D. G. NICHOL. 1994. Results of the 1990 U.S.-U.S.S.R. cooperative bottom trawl survey of the eastern and northwestern Bering Sea continental shelf, 183 p. NTIS number pending.
- 33 WEINBERG, K. L., M. E. WILKINS, R. R. LAUTH, and P. A. RAYMORE, JR. 1994. The 1989 Pacific west coast bottom trawl survey of groundfish resources: Estimates of distribution, abundance, and length and age composition, 168 p. plus Appendices. NTIS number pending.
- 32 PEREZ, M. A. 1994. Calorimetry measurements of energy value of some Alaskan fishes and squids, 32 p. NTIS No. PB94-152907.
- 31 RONHOLT, L. L., K. TESHIMA, and W. D. KESSLER. 1994. The groundfish resources of the Aleutian Islands region and southern Bering Sea 1980, 1983, and 1986, 351 p. NTIS No. PB94-152915
- 30 LORENZ, J. M. 1994. Distribution and habitat of adult salmon in the Situk River, Alaska: Potential impacts of flooding from Russell Fiord, 27 p. NTIS number pending.
- 29 IGNELL, S. E., L. J. SIMON, and M. L. DAHLBERG. 1994. Estimation of salmonid bycatch in the 1989 Japanese squid driftnet fishery, 45 p. NTIS No. PB94-152105.
- 28 MORTENSEN, D. G., and H. SAVIKKO. 1993. Effects of water temperature on growth of juvenile pink salmon (Oncorhynchus gorbuscha), 12 p. NTIS No. PB94-145349.
- 27 LOW, L-L. (coordinator). 1994. Status of living marine resources off Alaska, 1993, 110 p. NTIS No. PB94-149846.
- 26 LOW, L-L. (coordinator). 1993. Status of living marine resources off the Pacific coast of the United States for 1993, 90 p. NTIS No. PB94-133733.
- 25 KINOSHITA, R. K., A. GREIG, J. D. HASTIE, and J. M. TERRY. 1993. Economic status of the groundfish fisheries off Alaska, 1992, 102 p. PB94-142759.
- 24 SINCLAIR, E. H. (editor) 1993. Fur seal investigations, 1991, 142 p. NTIS No. PB94-118171.
- 23 PARKS, N. B., F. R. SHAW, and R. L HENRY. 1993. Results of a 1988 trawl survey of groundfish resources of the upper continental slope off Oregon, 164 p. NTIS No. PB94-118163.
- 22 YANG, M-S. 1993. Food habits of the commercially important groundfishes in the Gulf of Alaska in 1990, 150 p. NTIS No. PB94-112463.
- 21 KINOSHITA, R. K., and J. M. TERRY. 1993. Oregon, Washington, and Alaska exports of edible fishery products, 1992, 52 p. NTIS No. PB93-226652.
- 20 REEVES, J. E. 1993. Use of lower minimum size limits to reduce discards in the Bristol Bay red king crab (Paralithodes camtschaticus) fishery, 16 p. NTIS No. PB93-228187.