

## Phylogenetic Relationships Among Eastern Pacific Common Dolphins (*Delphinus* spp.)

M. Fernanda Urrutia-Osorio,<sup>1</sup> Thomas A. Jefferson,<sup>2,3</sup> Phillip A. Morin,<sup>2</sup>  
Brittany L. Hancock-Hanser,<sup>2</sup> Michael McGowen,<sup>4</sup>  
Madeleine A. Becker,<sup>5</sup> and Frederick I. Archer<sup>1,2</sup>

<sup>1</sup>Scripps Institution of Oceanography, University of California San Diego, 8622 Kennel Way, La Jolla, CA 92037, USA  
E-mail: [mfurruiti@ucsd.edu](mailto:mfurruiti@ucsd.edu)

<sup>2</sup>Southwest Fisheries Science Center, National Marine Fisheries Service, NOAA,  
8901 La Jolla Shores Drive, La Jolla, CA 92037, USA

<sup>3</sup>Clymene Enterprises, 13037 Yerba Valley Way, Lakeside, CA 92040, USA

<sup>4</sup>Division of Mammals, Department of Vertebrate Zoology, National Museum of Natural History,  
Smithsonian Institution, 1000 Madison Drive NW, Washington, DC 20560, USA

<sup>5</sup>Smithsonian-Mason School of Conservation, 1500 Remount Road, Front Royal, VA 22630, USA

### Abstract


Common dolphins in the eastern Pacific are currently recognized as separate subspecies, a long-beaked form (*Delphinus delphis bairdii*) and a short-beaked form (*Delphinus delphis delphis*), which are sympatric across portions of their ranges in waters off California. While the taxonomic status of these forms is still unclear, several studies have shown that they represent different evolutionary trajectories. Most of this work has been conducted on common dolphins in the Eastern North Pacific. In this study, the phylogenetic relationship of these two forms was examined, in addition to long-beaked common dolphins occurring off Peru in the Eastern South Pacific, using complete mitochondrial genomes. Peruvian and Californian long-beaked common dolphins formed a monophyletic clade with respect to short-beaked common dolphins from California and common dolphins from Senegal and the Black Sea. This clade of eastern Pacific long-beaked common dolphins was estimated to have zero connectivity with the Californian short-beaked form. Within the long-beaked clade, Peruvian long-beaked common dolphins were monophyletic, although Californian long-beaked dolphins were paraphyletic. These results are consistent with long-beaked common dolphins in the eastern Pacific being a species distinct from short-beaked common dolphins in the same region. Furthermore, evidence showed that Peruvian long-beaked common dolphins may represent a separate subspecies.

**Key Words:** *Delphinus*, phylogenetics, taxonomy, mitogenomes

### Introduction

The common dolphin (*Delphinus delphis*) is one of the most abundant, widespread, and genetically diverse dolphins in the world (Evans, 1994; Heyning & Perrin, 1994; Rosel et al., 1994; Amaral et al., 2007; Perrin, 2018). For years, phylogenetic studies have attempted to elucidate questions regarding the evolution and classification of this species (Rosel et al., 1994; LeDuc et al., 1999; Natoli et al., 2006; Cunha et al., 2015; Perrin, 2018). Most of this work has been focused on understanding relationships among forms that differ largely by length of the rostrum, often referred to as long- and short-beaked common dolphins. Currently, the Society for Marine Mammalogy (SMM) recognizes a single global species of common dolphins, *Delphinus delphis* (Committee on Taxonomy, 2024), composed of four subspecies: (1) *D. d. delphis* Linnaeus, 1758 (the nominate common dolphin, a globally distributed form that includes both long- and short-beaked populations), (2) *D. d. bairdii* Dall, 1873 (the Eastern North Pacific long-beaked common dolphin), (3) *D. d. ponticus* Barabash, 1935 (the Black Sea common dolphin, a short-beaked form), and (4) *D. d. tropicalis* van Bree, 1971 (the Indo-Pacific common dolphin, an exceedingly long-beaked form).

Both long- and short-beaked common dolphins are present in warm-temperate and tropical waters of the Pacific Ocean (Perrin, 2018), where they are often found sympatrically in the eastern Pacific along California, western Baja California, and Peru (Figure 1). In the Eastern North Pacific (ENP), both forms can be found in California and northern Mexico, where long-beaked common



**Figure 1.** Distribution of long- and short-beaked common dolphins (*Delphinus* spp.) in the eastern Pacific

dolphins (*D. d. bairdii*) are commonly distributed in coastal waters and short-beaked common dolphins (*D. d. delphis*) tend to occur more offshore (Banks & Brownell, 1969; van Bree & Purves, 1972; Heyning & Perrin, 1994; Rosel et al., 1994; Cunha et al., 2015; Perrin, 2018). A similar trend seems to be evident in the Eastern South Pacific (ESP), where both long- and short-beaked common dolphins are observed along Peru (Llapapasca et al., 2018; Santillán et al., 2023).

Long-beaked common dolphins in the ENP were first described as a distinct species of common dolphin by Dall (1873), referred to as *D. bairdii* (though Dall's reason for considering them a different species had more to do with geography than morphology). The differentiation of this form was based on its rostrum length and slenderness (length ratio of rostrum/greatest skull width), which was different from specimens from

the Atlantic Ocean (Dall, 1873; Miller, 1936; Banks & Brownell, 1969). Additionally, Banks & Brownell (1969) suggested that a “*bairdii*-like” population inhabited the ESP, where long-beaked common dolphins are mainly found in Peruvian waters, between 4° and 18° S (Llapapasca et al., 2018; Pacheco et al., 2019; Santillán et al., 2023).

Based largely on an examination of a series of skulls and color pattern differences, Heyning & Perrin (1994) described ENP long-beaked common dolphins as *Delphinus capensis* Gray, 1828, as skull measurements and tooth counts of the type specimen for that species, collected from the Cape of Good Hope (Eastern South Atlantic), were similar to long-beaked common dolphins from California and the name had priority over other named long-beaked common dolphin species. Species-level differentiation of the ENP long- and short-beaked common dolphins was also supported by an

analysis of mitochondrial DNA control region and cytochrome *b* sequences (Rosel et al., 1994).

However, later genetic analyses with a greater number of samples and geographical coverage indicate that ENP long-beaked common dolphins are not closely related to those occurring in the Eastern South Atlantic and South Africa from where the type specimen of *D. capensis* was originally described (Natoli et al., 2006; Amaral et al., 2007; Cunha et al., 2015). Thus, although the accumulated evidence suggests that the two forms of common dolphins occurring in the ENP are highly differentiated and might be considered a different species (Amaral et al., 2007; Cunha et al., 2015), the ENP long-beaked common dolphin form should not be referred to as *D. capensis* (Perrin, 2018).

Based on an examination of 351 skulls, 755 control regions, and 370 cytochrome *b* sequences, along with a review of color pattern differences and distribution of the forms, Jefferson et al. (2024) proposed that long-beaked common dolphins in the ENP be recognized as *D. bairdii*, a species distinct from the nominate short-beaked common dolphin *D. delphis*. However, even in light of the evidence presented, this proposal did not receive sufficient support by the SMM Committee on Taxonomy to recognize long-beaked common dolphins as a distinct species. The primary concerns were the lack of supporting nuclear data and a need for inclusion of samples from a wider geographic range. Thus, the SMM Committee on Taxonomy (2024) still recognizes the ENP long-beak common dolphin as the subspecies *D. d. bairdii*.


Furthermore, the evolutionary lineage of the ENP counterpart form, the ESP long-beaked common

dolphins, is also unclear. Due to the limited availability of biopsy samples and the absence of prior taxonomic studies, the relationship between long-beaked common dolphins in the ENP and the ESP remains unknown. Given the lack of connectivity between their distributions and the substantial distance separating them, it is unclear whether these two forms, despite both being long-beaked common dolphins, should be classified within the same taxon.

The aim of this study was to evaluate whether the long-beaked common dolphins from the ENP and the ESP share a common evolutionary lineage. In addition, the study determined the phylogenetic relationships and degree of genetic differentiation between ENP long- and short-beaked common dolphins using full mitogenome sequences.

## Methods

Skin samples from 28 long- and short-beaked common dolphins were collected from stranded and biopsied individuals. Based on the identification of color pattern by experienced observers, 12 long-beaked common dolphins from California (LB-Cali), currently known as *D. d. bairdii*, and six short-beaked common dolphins from California (SB-Cali) were identified from stranded individuals by the Southwest Fisheries Science Center (SWFSC), the Santa Barbara Natural Museum of Natural History, and Sea World San Diego. Samples from ten dolphins from the coast of Peru (LB-Peru) were identified as long-beaked common dolphins from individuals biopsied by SWFSC (Figure 2). No samples from Peruvian short-beaked common dolphins were available for the study.



**Figure 2.** Location and number of sample collections: Black Sea (pink), LB-Cali (yellow), LB-Peru (green), SB-Cali (blue), and Senegal (orange).

Tissues from biopsies and stranded animals were preserved frozen at  $-20^{\circ}$  or  $-80^{\circ}\text{C}$  in a 20% salt-saturated DMSO solution or in 100% ETOH in the SWFSC's Marine Mammal and Sea Turtle Research Collection (MMaSTR). Genomic DNA was extracted using sodium chloride precipitation (Miller et al., 1988) or the NucleoMag Tissue Extraction Kit (Macherey-Nagel, Inc., Bethlehem, PA, USA).

A capture array was designed as in Hancock-Hanser et al. (2013), with mitogenomes of 11 species, for use in capture enrichment of mitogenome DNA from multiple odontocete species (Table S1; this supplemental table is available in the "Supplemental Material" section of the *Aquatic Mammals* website), each at 20 copies on the array. The 60 bp probes were spaced every 15 bp so that every position in the sequences was represented four times in the probes. All the mitogenome sequences were modified to include a 40 bp overlap at both ends of the linearized sequence to allow for probe design across the artificial break point in the circular mitogenome.

DNA library preparation and array capture were performed according to Hancock-Hanser et al. (2013) with a few exceptions. Libraries were dual indexed using all unique i7 indexes and shared i5 indexes for groups of eight to 14 libraries to reduce impact of index-hopping during post-capture amplification (Kircher et al., 2011). Individual libraries were quantified using real time qPCR utilizing Bio-Rad iTaq universal SYBR supermix (Bio-Rad Laboratories, Inc., Hercules, CA, USA) and KAPA standards (KAPA Biosystems, Wilmington, MA, USA) made for Illumina primers. After hybridization and final amplification of the post hybridized product, the library was quantified as above and diluted to 4nM for loading on a MiSeq 1  $\times$  150 v3 kit flow cell per manufacturer's instructions (Illumina, San Diego, CA, USA). The library was sequenced using paired end reads with 75 cycles each.

Assembly of the reads for each sample was done through the custom pipeline in R, Version 4.0.2 (R Core Team, 2020), described in Archer et al. (2013) using default settings. All reads were assembled to a *Delphinus* mitogenome from GenBank (Accession No. NC\_012061). The first approximately 400 bp of the control region was replaced with Sanger sequenced versions (for methods, see Archer et al., 2013) to eliminate ambiguities from alignment and base calling in the pipeline.

The 'strataG' package for R (Archer et al., 2017) was used to calculate molecular diversity indices, such as haplotype and nucleotide diversity, proportion of unique haplotypes, and number of variable sites. In addition, pairwise levels of population differentiation were estimated with an

analysis of molecular variance (AMOVA),  $\Phi_{ST}$ ,  $F_{ST}$ , and  $\chi^2$  using 1,000 permutations.

To provide further clarification of relationships among long- and short-beaked individuals worldwide, four Black Sea short-beaked common dolphin (*Delphinus delphis ponticus*) mitogenomes from Biard et al. (2017) were downloaded from Genbank (Accession Nos. MF669495, MF669496, MF669497, and MF669498), and three long-beaked common dolphin mitogenomes from Senegal were obtained from Becker et al. (2024). Phylogenetic relationships and divergence times in million years (Ma) among the three common dolphin populations (LB-Peru, LB-Cali, and SB-Cali) were reconstructed from a total of 35 mitogenome sequences under a Bayesian framework using *BEAST*, Version 1.10.4 (Suchard et al., 2018). Mitogenomes from the Black Sea ( $n = 4$ ) and Senegal ( $n = 3$ ) were only used in the phylogenetic analysis, with the purpose of showing global patterns of common dolphin differentiation and, therefore, were not used in the population structure analysis, which focused on the eastern Pacific.

The program *jModelTest*, Version 2.1.10 (Guindon & Gascuel, 2003; Darriba et al., 2012) was used to select HKY + I as the best-fit model for nucleotide substitution for the data. Trees were generated using a Yule speciation process under a strict clock scenario. All trees were rooted to a reference sequence of striped dolphins (*Stenella coeruleoalba*; Genbank Accession No. NC\_012053.1). A prior distribution on the time since most recent common ancestor (TMRCA) was specified between the striped dolphin and the genus *Delphinus* estimated by McGowen et al. (2009). The program was run with ten million MCMC iterations, with samples taken every 1,000 iterations and a "burn in" process of 10%. The software *Tracer*, Version 1.7.1 (Rambaut et al., 2018), was used to evaluate effective mixing and convergence to the posterior distribution, and the program *TreeAnnotator*, Version 1.10.4 (Suchard et al., 2018), was used to summarize the information from the produced trees into a single tree.

Bidirectional migration rates between three populations in the eastern Pacific were estimated under a HKY substitution model using an Isolation with Migration (IM) analysis with the software *IMa2* (Hey & Nielsen, 2004; Hey, 2010). Several short runs were conducted to improve mixing and to optimize parameters. The final run included one million burn-in steps, followed by ten million MCMC iterations saved every 100 steps, resulting in 100,000 saved genealogies under a geometric heating scheme. Effective sample size (ESS) values and plot trends were used to evidence good mixing and convergence.

Following the guidelines for cetacean sub-species and species delimitation (Taylor et al., 2017; Morin et al., 2023), net nucleotide divergence (Nei's  $d_n$ ) was computed using 'strataG.' Percent diagnosability (PD) was calculated using a Random Forest model following Archer et al. (2017), with 10,000 trees, and sampling half of the smallest sample size per strata without replacement for each tree to create a balanced model.

## Results

### Genetic Diversity

Full mitochondrial DNA (16,376 bp) sequences were obtained from 28 common dolphin samples from California and Peru. The alignment of the 28 mitogenomes showed 347 variable sites, and a total of 25 different haplotypes were detected among the sequenced individuals in which almost all the samples had unique haplotypes (Table 1). The three putative populations showed high haplotype diversity (0.954 to 1.000), where the highest value was found within the SB-Cali population (Table 1). Only two haplotypes were shared: one between two individuals from Peru (LB-Peru-11880 and LB-Peru-11885), and the other among three long-beaked common dolphins from California (LB-Cali-65370, LB-Cali-79974, and LB-Cali-191186). Nucleotide diversity was greatest within the SB-Cali population (0.004), followed by the LB-Cali (0.002). The LB-Peru nucleotide diversity (0.0005) was an order of magnitude lower than the other two populations.

### Phylogenetic Reconstruction

The Bayesian phylogenetic tree showed strong support for most nodes, with a posterior distribution greater than 0.90 in 77% of the nodes (Figure 3). The phylogenetic tree identified a well-supported clade (posterior probability = 1.0) containing all the sequences from the LB-Peru population. The LB-Cali population was paraphyletic, in which most sequences were more closely related to LB-Peru individuals than to other LB-Cali dolphins.

The SB-Cali population grouped together with high confidence, except for SB-Cali-78511 (Figure 3). The phylogenetic tree also revealed that short-beaked common dolphins from the eastern Pacific share a closer genetic relationship with long-beaked common dolphins from Senegal and short-beaked common dolphins from the Black Sea than with long-beaked common dolphins from their own ocean basin. In addition, the most surprising aspect of the phylogenetic analysis is that sequence SB-Cali-78511 was more distantly related to the other five SB-Cali sequences than individuals from the Black Sea and Senegal.

Thus, sample SB-Cali-78511 was further analyzed to find any assembly errors, but none were found. The six SB-Cali sequences showed a total of 201 variable sites. When the SB-Cali-78511 sequence was removed from the group, the number of variable sites decreased to 126, meaning that the SB-Cali-78511 accounted for 37% of all the variable sites in the SB-Cali clade. Specifically, SB-Cali-78511 showed 78 differences to the other five SB-Cali sequences, 76 of which were transitions and two transversions. For the 78 sites where differences were found, the average coverage was 70 reads, and the bases for those sites were called with an average frequency of 99%.


The time since most common recent ancestor (TMRCA) between both common dolphin forms and striped dolphins was estimated at around 3.44 Ma (95% HDP = 2.65 to 4.23 Ma), while the long- and short-beaked forms were estimated to have diverged around 1.54 Ma (95% HDP = 1.13 to 1.95 Ma).

### Population Differentiation and Taxonomy

The three populations of common dolphins from the eastern Pacific showed significant genetic differentiation in the AMOVA analysis ( $\Phi_{st} = 0.3669$ ;  $p < 0.0009$ ), but not for the  $F_{st}$  and  $\chi^2$  analyses (Table 2). The strongest genetic differentiation resulted between the LB-Peru and SB-Cali samples, followed by LB-Cali

**Table 1.** Mitogenome (16,376 bp) summaries for the three common dolphin (*Delphinus* spp.) populations. N = number of sequences, N<sub>H</sub> = number of unique haplotypes, P<sub>H</sub> = proportion of unique haplotypes,  $h$  = haplotype diversity,  $\pi$  = nucleotide diversity, and VS = number of variable sites.

	N	N <sub>H</sub>	P <sub>H</sub>	$h$	$\pi$	VS
LB-Cali	12	10	0.750	0.954	0.002	91
LB-Peru	10	9	0.800	0.978	0.0005	28
SB-Cali	6	6	1.000	1.000	0.004	201
Total	28	25	0.821	0.989	0.004	347



**Figure 3.** Bayesian phylogenetic tree and TMRCA (time since most recent common ancestor) derived from *BEAST*. Numbers at each node correspond to the posterior probability.

**Table 2.** Population differentiation between three common dolphin populations in the eastern Pacific:  $\chi^2$ ,  $F_{ST}$ ,  $\Phi_{ST}$ , Nei's  $d_A$ , and percent diagnosability (PD). For  $\chi^2$ , only  $p$  values are shown. The last column shows the 95% confidence interval for the PD.

	$\chi^2$ $p$ value	$F_{ST}$ $p$ value	$F_{ST}$ $p$ value	$\Phi_{ST}$ $p$ value	$\Phi_{ST}$ $p$ value	$d_A$	PD (%)	PD (%) 95% CI
LB-Cali vs LB-Peru	0.096	0.034	0.068	0.344	0.001	0.0007	100	69-100
LB-Cali vs SB-Cali	0.269	0.025	0.269	0.622	0.001	0.0042	100	69-100
LB-Peru vs SB-Cali	0.487	0.012	0.487	0.713	0.001	0.0046	100	66-100

and SB-Cali, and then between LB-Peru and LB-Cali. Bidirectional migration between the three common dolphin populations was close to zero. Since *IMa2* estimates mutation rate-scaled parameters, migration rates were later converted to demographic estimates as the number of migrants per generation, assuming a generation time for common dolphins of 15 y (Taylor et al., 2007) and the mutation rate of  $1.18 \times 10^{-3}$  substitutions per site/generation estimated by the *BEAST* analysis conducted previously (Table 3).

For cetacean taxonomic delimitation using mitogenomes, Morin et al. (2023) proposed Nei's  $d_A$  values greater than 0.0006 for subspecies, and 0.008 for species classification and percent diagnosability (PD) greater than 95% for both delimitations. All pairwise comparisons for nucleotide divergence and diagnosability were found to fall within the suggested subspecies threshold values. However, for species delineation, all pairwise comparisons failed to meet the proposed threshold value for Nei's  $d_A$  but met the threshold for PD (Table 2).

**Table 3.** Bidirectional scaled migration rates ( $m$ ) and number of migrants per generation between the three eastern Pacific populations of common dolphins. Parentheses show the 95% highest posterior density interval.

	$m$	Migrants per generation
LB-Peru $\leftrightarrow$ LB-Cali	0.009 (0.001-0.104)	0.000
LB-Peru $\leftrightarrow$ SB-Cali	0.027 (0.001-0.120)	0.000
SB-Cali $\leftrightarrow$ LB-Cali	0.000 (0.000-0.135)	0.000

## Discussion

Common dolphins in the ENP are currently recognized as two different subspecies, the long-beaked (*D. d. bairdii*) and the short-beaked (*D. d. delphis*) (Committee on Taxonomy, 2024). In this analysis, long-beaked common dolphins from Peru and California form a monophyletic clade with respect to short-beaked common dolphins from California and the Black Sea, and long-beaked common dolphins from Senegal. Additionally, although Nei's  $d_A$  is lower than the subspecies-species threshold suggested for mitogenomes by Morin et al. (2023), Pacific long- and short-beaked common dolphins are 100% diagnosable based on mtDNA. The estimated divergence time between long- and short-beaked common dolphins in the North Pacific based on mitogenome sequences (1.54 Ma; 95% HPDI = 1.13 to 1.95 Ma) is similar to that found in other recently diverged odontocete sister species (McGowen et al., 2009; Amaral et al., 2012; Braulik et al., 2019; Morin et al., 2023), and the estimated migration rate between the forms is zero. Thus, this study supports recognizing long- and short-beaked common dolphins in the Pacific as separate species, with both the Peruvian and Californian long-beaked form belonging to the same taxon, likely representing *Delphinus bairdii* Dall, 1873 (Banks & Brownell, 1969; Jefferson et al., 2024).

The present article is limited by its exclusive use of mitochondrial DNA, requiring the inclusion of nuclear DNA for result clarity. The small sample size compromises statistical robustness, emphasizing the need for a larger sample. While species-level insights are offered, the study suggests that additional samples and nuclear data are essential to refine the conclusions, ensuring a more comprehensive and reliable understanding of the different forms of common dolphins in the eastern Pacific.

Previous studies have shown that ENP long-beaked common dolphins are genetically differentiated from common dolphins in other parts of the world (Rosel et al., 1994; Kingston & Rosel,

2004; Natoli et al., 2006; Amaral et al., 2007; Jefferson et al., 2024). Using 1,140 bp of cytochrome *b*, Cunha et al. (2015) found ENP long-beaked common dolphins to be paraphyletic with respect to the ENP short-beaked form. The authors attributed this finding to ancestral shared polymorphisms or incomplete lineage sorting between recently diverged species. However, the use of the full mitogenome in our study provides greater phylogenetic resolution at this scale (Morin et al., 2010, 2023; Archer et al., 2013).

Within the long-beaked clade, Peruvian long-beaked common dolphins were monophyletic, while Californian long-beaked common dolphins were paraphyletic with nine of 12 Californian long-beaked samples more closely related to Peruvian long-beaked common dolphins than to the other three Californian long-beaked dolphins. This pattern would be consistent with California as the source population for Peruvian long-beaked common dolphins was estimated to have diverged approximately 400 Kya (Figure 3). Additionally, our study indicates that there is no current gene flow between long-beaked common dolphins in these two regions, which is understandable given the great geographic distance (> 6,000 km) between them (see Hamilton et al., 2009). Thus, the monophyletic clade formed by Peruvian long-beaked common dolphins (Figure 3), in addition to meeting the subspecies threshold values for subspecies delimitation between the long-beaked form from Peru and California (Table 2), suggests that Peruvian long-beaked common dolphins may represent at a distinct evolutionary lineage at least at the subspecies level within the Pacific long-beaked taxon.

Other than the examination of two specimens from the coast of Peru (Banks & Brownell, 1969), taxonomic studies on Peruvian common dolphins have been scarce. The present study provides the first insights into the population structure and genetic differentiation of Peruvian long-beaked common dolphins. The nutrient-rich waters of the Humboldt Current make Peru's fisheries among the most important and productive fisheries in the world. The intense extraction of marine resources results in high mortality rates for Peruvian

dolphins due to bycatch (Van Waerebeek et al., 2002; Mangel et al., 2010; Ayala et al., 2019). Ayala et al. (2019) reported that 25 vessels off the coast of Peru incidentally caught more than 1,000 dolphins in 2009. The results of this study suggest that Peruvian long-beaked common dolphins represent a distinct evolutionary lineage. Follow-up studies on this population are needed to assess its taxonomic and conservation status and to establish mitigation measures in this highly impacted area.

While differentiation between long- and short-beaked common dolphins in the eastern Pacific Ocean is clear, it is less so for relationships between the eastern Pacific short-beaked form and common dolphins in the Atlantic Ocean. The presence of the SB-Cali-78511 sequence at the base of a clade containing all other Californian short-beaked dolphins as well as samples of the endemic *D. d. ponticus* subspecies from the Black Sea and long-beaked common dolphins from Senegal suggests that there is considerable ancestral polymorphism within the eastern Pacific short-beaked form. Within California, Oregon, and Washington waters, the short-beaked common dolphin population is estimated at 1,056,308 (CV = 0.207) individuals (Carretta et al., 2022); thus, a large amount of genetic diversity is expected, which is unlikely to be fully represented by the six samples present in this study.

Within the Atlantic, Cunha et al. (2015) showed that long- and short-beaked cytochrome *b* sequences from the western South Atlantic had no genetic differences, with all corresponding to *D. delphis*. Natoli et al. (2006) found genetic differences in the mtDNA control region between populations in different sides of the Atlantic Ocean but no significant genetic differences between populations located on the same side of that ocean. A recent study of full mitogenome sequences showed significant population structure within Senegal as well as differentiation between common dolphins in Senegal and the rest of the North Atlantic (Becker et al., 2024). As the authors note, improved resolution of phylogenetic patterns in this region and globally will likely require nuclear genomic level data.

In summary, this study has demonstrated that Peruvian long-beaked common dolphins should be considered within the same taxon as long-beaked common dolphins from the ENP, which are all currently regarded as the subspecies *Delphinus delphis bairdii* (Committee on Taxonomy, 2024). Additionally, within that taxon, they represent a distinct evolutionary lineage. It is clear that full elucidation of the taxonomy of these forms awaits further studies.

**Note:** The supplemental table for this article is available in the “Supplemental Material” section of the *Aquatic Mammals* website: <https://www.aquatic-mammalsjournal.org/supplemental-material>.

### Acknowledgements

This work was supported by the Southwest Fisheries Science Center Marine Mammal Genetics Program. The authors would like to thank the Santa Barbara Museum of Natural History, the Peruvian National Institute of Natural Resources (INRENA), Sea World San Diego, and the National Marine Fisheries Service West Coast Region Observer Program for providing samples for the study. Peruvian samples were imported under CITES Permit 99US774223/9. Samples from California were collected under NMFS Permits 14097, 18786, 0774-1437, and 932-1905.

### Literature Cited

- Amaral, A. R., Sequeira, M., Martínez-Cedeira, J., & Coelho, M. M. (2007). New insights on population genetic structure of *Delphinus delphis* from the Northeast Atlantic and phylogenetic relationships within the genus inferred from two mitochondrial markers. *Marine Biology*, 151(5), 1967-1976. <https://doi.org/10.1007/s00227-007-0635-y>
- Amaral, A. R., Beheregaray, L. B., Bilgmann, K., Freitas, L., Robertson, K., Sequeira, M., Stockin, K. A., Coelho, M. M., & Möller, L. M. (2012). Influences of past climatic changes on historical population structure and demography of cosmopolitan marine predator, the common dolphin (genus *Delphinus*). *Molecular Ecology*, 21(19), 4854-4871. <https://doi.org/10.1111/j.1365-294X.2012.05728.x>
- Archer, F. I., Adams, P. E., & Schneiders, B. B. (2017). stratag: An R package for manipulating, summarizing and analyzing population genetic data. *Molecular Ecology Resources*, 17(1), 5-11. <https://doi.org/10.1111/1755-0998.12559>
- Archer, F. I., Morin, P. A., Hancock-Hanser, B. L., Robertson, K. M., Leslie, M. S., Bérubé, M., Panigada, S., & Taylor, B. L. (2013). Mitogenomic phylogenetics of fin whales (*Balaenoptera physalus* spp.): Genetic evidence for revision of subspecies. *PLOS ONE*, 8(5), e63396. <https://doi.org/10.1371/journal.pone.0063396>
- Ayala, L., Ortiz, M., & Gelcich, S. (2019). Exploring the role of fishers knowledge in assessing marine megafauna bycatch: Insights from the Peruvian longline artisanal fishery. *Animal Conservation*, 22(3), 251-261. <https://doi.org/10.1111/acv.12460>
- Banks, R. C., & Brownell, R. L. (1969). Taxonomy of the common dolphins of the eastern Pacific Ocean. *Journal of Mammalogy*, 50(2), 262-271. <https://doi.org/10.2307/1378342>
- Becker, M. A., Murphy, K. R., Archer, F. I., Jefferson, T. A., Ndong, I., Potter, C. W., Urrutia-Orsorio, M. F., Keith-Diagne, L. W., & McGowen, M. R. (2024). Common dolphin (*Delphinus delphis*) mitochondrial genomes



- from Senegal reveal geographic structure across the North Atlantic but provide no support for global long-beaked clade. *Marine Mammal Science*, 40(4), e13144. <https://doi.org/10.1111/mms.13144>
- Biard, V., Gol'din, P., Gladilina, E., Vishnyakova, K., McGrath, K., Vieira, F. G., Wales, N., Fontaine, M. C., Speller, C., & Olsen, M. T. (2017). Genomic and proteomic identification of Late Holocene remains: Setting baselines for Black Sea odontocetes. *Journal of Archaeological Science: Reports*, 15, 262-271. <https://doi.org/10.1016/j.jasrep.2017.07.008>
- Braulik, G., Natoli, A., Kiszka, J., Parra, G., Plön, S., & Smith, B. D. (2019). *Tursiops aduncus*. In International Union for Conservation of Nature (Ed.), *The IUCN red list of threatened species 2019* (e.T41714A50381127). IUCN. <https://doi.org/10.2305/IUCN.UK.2019-3.RLTS.T41714A50381127.en>
- Carretta, J. V., Oleson, E. M., Forney, K. A., Muto, M. M., Weller, D. W., Lang, A. R., Baker, J., Hanson, B., Orr, A. J., Barlow, J., Moore, J. E., & Brownell, R. L., Jr. (2022). *U.S. Pacific marine mammal stock assessments: 2021* (NOAA Technical Memorandum NMFS-SWFSC-663). U.S. Department of Commerce, National Oceanic and Atmospheric Administration. <https://doi.org/10.25923/246k-7589>
- Committee on Taxonomy. (2024). *List of marine mammal species and subspecies*. Society for Marine Mammalogy. [www.marinemammalscience.org](http://www.marinemammalscience.org)
- Cunha, H. A., Loizaga De Castro, R., Secchi, E. R., Crespo, E. A., Lailson-Brito, J., Azevedo, A. F., Lazoski, C., & Sole-Cava, A. M. (2015). Molecular and morphological differentiation of common dolphins (*Delphinus* sp.) in the Southwestern Atlantic: Testing the two species hypothesis in sympatry. *PLOS ONE*, 10(11), 1-15. <https://doi.org/10.1371/journal.pone.0140251>
- Dall, W. H. (1873). Preliminary descriptions of three new species of Cetacea from the coast of California. *Annals and Magazine of Natural History*, 11(64), 316-317. <https://doi.org/10.1080/00222937308696820>
- Darriba, D., Taboada, G. L., Doallo, R., & Posada, D. (2012). jModelTest 2: More models, new heuristics and parallel computing. *Nature Methods*, 9, 772. <https://doi.org/10.1038/nmeth.2109>
- Evans, W. E. (1994). Common dolphin, white-bellied porpoise *Delphinus delphis* Linnaeus, 1758. In S. H. Ridgway & R. Harrison (Eds.), *Handbook of marine mammals: Vol. 5. The first book of dolphins* (pp. 191-224). Academic Press.
- Guindon, S., & Gascuel, O. (2003). A simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood. *Systematic Biology*, 52(5), 696-704. <https://doi.org/10.1080/10635150390235520>
- Hamilton, T. A., Redfern, J. V., Barlow, J., Ballance, L. T., Gerrodette, T., Holt, R. S., Forney, K. A., & Taylor, B. L. (2009). *Atlas of cetacean sightings for Southwest Fisheries Science Center Cetacean and Ecosystem surveys, 1986-2005* (NOAA Technical Memorandum NMFS-SWFSC 440). U.S. Department of Commerce, National Oceanic and Atmospheric Administration. 70 pp.
- Hancock-Hanser, B. L., Frey, A., Leslie, M. S., Dutton, P. H., Archer, F. I., & Morin, P. A. (2013). Targeted multiplex next-generation sequencing: Advances in techniques of mitochondrial and nuclear DNA sequencing for population genomics. *Molecular Ecology Resources*, 13(2), 254-268. <https://doi.org/10.1111/1755-0998.12059>
- Hey, J. (2010). Isolation with migration models for more than two populations. *Molecular Biology and Evolution*, 27(4), 905-920. <https://doi.org/10.1093/molbev/msp296>
- Hey, J., & Nielsen, R. (2004). Multilocus methods for estimating population sizes, migration rates and divergence time, with applications to the divergence of *Drosophila pseudoobscura* and *D. persimilis*. *Genetics*, 167(2), 747-760. <https://doi.org/10.1534/genetics.103.024182>
- Heyning, J. E., & Perrin, W. F. (1994). Evidence for two species of common dolphins (genus *Delphinus*) from the eastern North Pacific. *Contributions in Science, Natural History Museum of Los Angeles County*, 442, 1-35. <https://doi.org/10.5962/p.226804>
- Jefferson, T. A., Archer, F. I., & Robertson, K. M. (2024). The long-beaked common dolphin of the eastern Pacific Ocean: Taxonomic status and redescription of *Delphinus bairdii*. *Marine Mammal Science*, 40(4), e13133. <https://doi.org/10.1111/mms.13133>
- Kingston, S. E., & Rosel, P. E. (2004). Genetic differentiation among recently diverged delphinid taxa determined using AFLP markers. *Journal of Heredity*, 95(1), 1-10. <https://doi.org/10.1093/jhered/esh010>
- Kircher, M., Sawyer, S., & Meyer, M. (2011). Double indexing overcomes inaccuracies in multiplex sequencing on the Illumina platform. *Nucleic Acids Research*, 40(1), e3. <https://doi.org/10.1093/nar/gkr771>
- LeDuc, R. G., Perrin, W. F., & Dizon, A. E. (1999). Phylogenetic relationships among the delphinid cetaceans based on full cytochrome *b* sequences. *Marine Mammal Science*, 15(3), 619-648. <https://doi.org/10.1111/j.1748-7692.1999.tb00833.x>
- Llapasasca, M. A., Pacheco, A. S., Fiedler, P., Goya, E., Ledesma, J., Peña, C., & Vásquez, L. (2018). Modeling the potential habitats of dusky, commons and bottlenose dolphins in the Humboldt Current System off Peru: The influence of non-El Niño vs. El Niño 1997-98 conditions and potential prey availability. *Progress in Oceanography*, 168, 169-181. <https://doi.org/10.1016/j.pocan.2018.09.003>
- Mangel, J. C., Alfaro-Shigueto, J., Van Waerebeek, K., Cáceres, C., Bearhop, S., Witt, M. J., & Godley, B. J. (2010). Small cetacean captures in Peruvian artisanal fisheries: High despite protective legislation. *Biological Conservation*, 143(1), 136-143. <https://doi.org/10.1016/j.biocon.2009.09.017>
- McGowen, M. R., Spaulding, M., & Gatesy, J. (2009). Divergence date estimation and a comprehensive molecular tree of extant cetaceans. *Molecular Phylogenetics and Evolution*, 53(3), 891-906. <https://doi.org/10.1016/j.ympev.2009.08.018>

- Miller, G. S. (1936). The status of *Delphinus bairdii* Dall. *Proceedings of the Biological Society of Washington*, 49, 145-146.
- Miller, S. A., Dykes, D. D., & Polesky, H. (1988). A simple salting out procedure for extracting DNA from human nucleated cells. *Nucleic Acids Research*, 16(3), 1215. <https://doi.org/10.1093/nar/16.3.1215>
- Morin, P. A., Martien, K., Lang, A. R., Hancock-Hanser, B., Pease, V. L., Roberston, K. M., Sattler, M., Slikas, E., Rosel, P. E., Baker, C. S., Taylor, B. L., & Archer, F. I. (2023). Guidelines and quantitative standards for improved cetacean taxonomy using full mitochondrial genomes. *Journal of Heredity*, 114(6), 612-624. <https://doi.org/10.1093/jhered/esad049>
- Morin, P. A., Archer, F. I., Foote, A. D., Vilstrup, J., Allen, E. E., Wade, P., Durban, J., Parsons, K., Pitman, R., Li, L., Bouffard, P., Abel Nielsen, S. C., Rasmussen, M., Willerslev, M., Gilbert, M. T. P., & Harkins, T. (2010). Complete mitochondrial genome phylogeographic analysis of killer whales (*Orcinus orca*) indicates multiple species. *Genome Research*, 20(7), 908-916. <https://doi.org/10.1101/gr.102954.109>
- Natoli, A., Cañadas, A., Peddemors, V. M., Aguilar, A., Vaquero, C., Fernández-Piqueras, P., & Hoelzel, A. R. (2006). Phylogeography and alpha taxonomy of the common dolphin (*Delphinus* sp.). *Journal of Evolutionary Biology*, 19(3), 943-954. <https://doi.org/10.1111/j.1420-9101.2005.01033.x>
- Pacheco, A. S., Silva, S., Alcorta, B., Gubbins, S., Guidino, C., Sanchez-Salazar, F., Petit, A., Llapapasca, M. A., Balducci, N., Larrañaga, E., Zapata, M. A., Grados, E., Valdivia, C., Pinasco, G., Garcia Cegarra, A. M., Caceres, D., Biffi, D., Silva, L., Auger, A., Borda, D., . . . Jaramillo-Calle, V. (2019). Cetaceans diversity revealed from whale-watching. *Aquatic Mammals*, 45(1), 116-122. <https://doi.org/10.1578/AM.45.1.2019.116>
- Perrin, W. F. (2018). Common dolphin: *Delphinus delphis*. In B. Würsig, J. G. M. Thewissen, & K. M. Kovacs (Eds.), *Encyclopedia of marine mammals* (3rd ed., pp. 205-209). Academic Press. <https://doi.org/10.1016/B978-0-12-804327-1.00095-9>
- R Core Team. (2020). R: A language and environment for statistical computing. R Foundation for Statistical Computing. <https://www.R-project.org>
- Rambaut, A., Drummond, A. J., Xie, D., Baele, G., & Suchard, M. A. (2018). Posterior summarization in Bayesian phylogenetics using *Tracer 1.7*. *Systematic Biology*, 67(5), 901-904. <https://doi.org/10.1093/sysbio/syy032>
- Rosel, P. E., Dizon, A. E., & Heyning, J. E. (1994). Genetic analysis of sympatric morphotypes of common dolphins (genus *Delphinus*). *Marine Biology*, 119(2), 159-167. <https://doi.org/10.1007/BF00349552>
- Santillán, L., Pacheco, A. S., Martínez, J. N., Fupuy, J., Barboza, R., Cárdenas-Alayza, S., Torres, D. A., López, E., Medina, C., Pacheco, V., Canto-Hernández, J., Alfaro-Shigueto, J., & Ramírez, P. (2023). Distribution of short and long-beaked common dolphin morphs (*Delphinus* spp.) based on skull's rostrum index analysis along the Peruvian and Chilean coast. *Mastozoología Neotropical*, 30(1), e0908. <https://doi.org/10.31687/saremMN.23.30.1.07.e0908>
- Suchard, M. A., Lemey, P., Baele, G., Ayres, D. L., Drummond, A. J., & Rambaut, A. (2018). Bayesian phylogenetic and phylodynamic data integration using *BEAST 1.10*. *Virus Evolution*, 4(1), 1-5. <https://doi.org/10.1093/ve/vey016>
- Taylor, B. L., Chivers, S. J., Larese, J., & Perrin, W. F. (2007). *Generation length and percent mature estimates for IUCN assessments of cetaceans* (Southwest Fisheries Science Center Administrative Report LJ-07-01). National Oceanic and Atmospheric Administration, National Marine Fisheries Service.
- Taylor, B. L., Archer, F. I., Martien, K. K., Rosel, P. E., Hancock-Hanser, B. L., Lang, A. R., Leslie, M. S., Mesnick, S. L., Morin, P. A., Pease, V. L., Perrin, W. F., Robertson, K. M., Parsons, K. M., Viricel, A., Vollmer, N. L., Cipriano, F., Reeves, R. R., Krützen, M., & Baker, C. S. (2017). Guidelines and quantitative standards to improve consistency in cetacean subspecies and species delimitation relying on molecular genetic data. *Marine Mammal Science*, 33(S1), 132-155. <https://doi.org/10.1111/mms.12411>
- van Bree, P. J. H., & Purves, P. E. (1972). Remarks on the validity of *Delphinus bairdii* (Cetacea: Delphinidae). *Journal of Mammalogy*, 53(2), 372-374. <https://doi.org/10.2307/1379178>
- Van Waerebeek, K., Alfaro-Shigueto, J., Montes, D., Onton-Silva, K., Santillán, L., Van Bresseem, M-F., & Vega, D. (2002). *Fisheries related mortality of small cetaceans in neritic waters of Peru in 1999-2001* (SC/54/Sm10). Unpublished paper prepared for the IWC Scientific Committee.

**Phylogenetic Relationships Among Eastern Pacific  
Common Dolphins (*Delphinus* spp.)**

**Supplemental Table**

M. Fernanda Urrutia-Osorio,<sup>1</sup> Thomas A. Jefferson,<sup>2,3</sup> Phillip A. Morin,<sup>2</sup>  
Brittany L. Hancock-Hanser,<sup>2</sup> Michael McGowen,<sup>4</sup>  
Madeleine A. Becker,<sup>5</sup> and Frederick I. Archer<sup>1,2</sup>

<sup>1</sup>*Scripps Institution of Oceanography, University of California San Diego, 8622 Kennel Way, La Jolla, CA 92037, USA*

*E-mail: mfuurruti@ucsd.edu*

<sup>2</sup>*Southwest Fisheries Science Center, National Marine Fisheries Service, NOAA,  
8901 La Jolla Shores Drive, La Jolla, CA 92037, USA*

<sup>3</sup>*Clymene Enterprises, 13037 Yerba Valley Way, Lakeside, CA 92040, USA*

<sup>4</sup>*Division of Mammals, Department of Vertebrate Zoology, National Museum of Natural History,  
Smithsonian Institution, 1000 Madison Drive NW, Washington, DC 20560, USA*

<sup>5</sup>*Smithsonian-Mason School of Conservation, 1500 Remount Road, Front Royal, VA 22630, USA*

## Supplemental Table

**Table S1.** Mitogenomes used as templates in array creation with Genbank Accession No. and Southwest Fisheries Science Center Marine Mammal and Sea Turtle Research Collection (MMaSTR) identification numbers

Species	Genbank Accession No.	MMaSTR ID
<i>Delphinus delphis delphis</i>	MF669498	
<i>Delphinus delphis bairdii</i>	NC_012061	
<i>Pseudorca crassidens</i>	JF289173.1	
<i>Globicephala melas</i>	HM060334.1	
<i>Globicephala macrorhynchus</i>	HM060333.2	
<i>Peponocephala electra</i>	JF289175.1	
<i>Berardius minimus</i>		z0007696
<i>Berardius bairdii</i>		z0017152
<i>Phocoena dioptrica</i>		z0000981
<i>Phocoena phocoena</i>		z0000161
<i>Phocoena spinnipinnis</i>		z0052776