

QC  
475  
1812  
51  
1992  
C.2


NOAA Technical Memorandum NWS NHC-31

Updated February 1992

THE DEADLIEST, COSTLIEST, AND MOST INTENSE  
UNITED STATES HURRICANES OF THIS CENTURY  
(AND OTHER FREQUENTLY REQUESTED HURRICANE FACTS)

Prepared by:

Paul J. Hebert, WSFO, Miami, Florida  
Jerry D. Jarrell, NHC, Miami, Florida  
Max Mayfield, NHC, Miami, Florida



National Hurricane Center  
Coral Gables, Florida  
February 1992



## PREFACE

This version of the Deadliest, Costliest, and Most Intense United States Hurricanes of This Century is an update through the 1991 hurricane season. Hurricane Bob, with about \$1.5 billion damage to the United States East Coast has caused some of the damage tables to be revised. Also, 1990 Census figures have influenced several sections of the text. This is, nevertheless, a fairly minor update of work done by Hebert, Taylor and Case (1984).

THE DEADLIEST, COSTLIEST, AND MOST INTENSE  
UNITED STATES HURRICANES OF THIS CENTURY  
(AND OTHER FREQUENTLY REQUESTED HURRICANE FACTS)

by

Paul J. Hebert, Weather Service Forecast Office  
Jerry D. Jarrell & Max Mayfield, National Hurricane Center  
National Weather Service  
Miami, Florida.

ABSTRACT

Lists of United States hurricanes which have caused 25 or more deaths and more than one hundred million dollars in damages (unadjusted) during this century have been compiled from all data sources available at the National Hurricane Center (NHC). In addition, all major<sup>1</sup> hurricanes which have made landfall in the United States during this century are listed. Some additional statistics on United States hurricanes of this century and tropical cyclones in general are also presented.

1. INTRODUCTION

Numerous requests are received at the National Hurricane Center for statistical information on deaths, damages, and severity of hurricanes which have affected the United States. Various reference materials gave different estimates of these statistics so that decisions had to be constantly made as to which information should be given out by NHC as "official" from the National Hurricane Information Center (another function of NHC). Requests to other Weather Service Offices posed the same dilemma. These lists are being published in the hope of presenting a single source of the best currently available estimates of deaths, damages, and intensity of major U.S. hurricanes which have made landfall in this century. In some instances, data in these lists present revised estimates for individual hurricanes based on more complete information received after earlier published values, including the previous versions of this technical memorandum. There are other frequently asked questions about hurricanes. What is the average number of hurricanes per year? What year(s) had the most and least hurricanes? What hurricane had the longest life? When did the earliest and latest hurricane occur? What was the most intense Atlantic hurricane? What was the largest number of hurricanes in existence on the same day? When was the last time a major hurricane or any hurricane hit a given community directly<sup>2</sup>? Answers to these and several other questions are provided in Section 3.

<sup>1</sup> A major hurricane is a category 3, 4, or 5 on the Saffir/Simpson Hurricane Scale (see Table 1), and is comparable to a Great Hurricane in other publications.

<sup>2</sup> A direct hit means experiencing the core of strong winds and high tides of a hurricane.

Table 1. Saffir/Simpson Hurricane Scale Ranges.

Scale Number (Category)	Central Pressure (Millibars)	OR (Inches)	Winds (Mph)	OR (Feet)	Surge	Damage
1	≥ 980	≥ 28.94	74-95	4-5	Minimal	
2	965-979	28.50-28.91	96-110	6-8	Moderate	
3	945-964	27.91-28.47	111-130	9-12	Extensive	
4	920-944	27.17-27.88	131-155	13-18	Extreme	
5	< 920	< 27.17	> 155	> 18	Catastrophic	

## 2. CRITERIA

The statistics in most of the tables and figures in this publication depend directly on the criteria used in preparing another study, Hurricane Experience Levels of Coastal County Populations-Texas to Maine (Hebert, Taylor, and Case, 1984). The primary purpose of that study was to demonstrate, county by county, the low hurricane experience level of a large majority of the population. Statistics show that the largest loss of life and, for the most part, property occur in locations experiencing the core of a category 3 or higher hurricane. Unless a given population has experienced this core, or direct hit, with its very strong winds and high tides, it would defeat the primary purpose of the study on hurricane experience levels to so categorize it.

The central pressure ranges of hurricanes on the Saffir/Simpson scale will usually agree quite well with the wind ranges in that category. In the absence of other information, this is normally the best estimate of a hurricane's winds. However, some hurricanes which have developed from winter-type or subtropical low pressure systems occasionally have a minimum central pressure lower than the corresponding observed winds would suggest. In this instance, the wind criteria are used.

On the other hand, the surge is strongly dependent on the slope of the continental shelf (shoaling factor). This can change the height of the surge by a factor of two for a given central pressure and/or maximum wind.

Heavy rainfall associated with a hurricane is not one of the criteria for categorizing it.

The subjective determination of which category number to assign to a hurricane, as well as its direct or indirect effect, is made on a county by county basis with the intent of the study on hurricane experience levels foremost in mind.

As with the assignment of scale numbers, a certain amount of subjectivity was inescapable at times in determining which counties received direct or indirect hits during the various hurricane situations. However, certain arbitrary guidelines for these classifications as used in Hurricane Experience Levels, etc., are indicated below:

Direct Hit - When the innermost core region or "eye" moved over a county, it was counted as a direct hit. Using "R" as the radius of maximum winds in a hurricane (the distance in miles from the storm's center to the circle of maximum winds around the center), all or parts of counties falling within approximately 2R to the right and R to the left of a storm's landfall point were considered to have received a direct hit. (This assumes an observer at sea looking toward the

shore). On the average, this direct hit zone extended about 50 miles along the coastline ( $R \approx 15$  miles). Of course, some hurricanes were smaller than this and some, particularly at higher latitudes, were much larger. Cases were judged individually, and many borderline situations had to be resolved.

Indirect Hit - These were based primarily on a hurricane's strength and size, and on the configuration of the individual county coastline. Here again, much subjectivity was necessary in many cases which were complicated by storm paths and geography. Generally, those areas on either side of the direct hit zone which received at least wind gusts of hurricane force and/or tides of 4 to 5 feet or more above normal were considered to have had an indirect hit.

It is realized that the effect of an indirect hit by a large category 4 hurricane might be greater than that of a small category 1 affecting the same county. However, trying to account for these differences would hopelessly complicate the use of this system.

The study by Simpson and Lawrence (1971) gives climatological probabilities of the total number of storms and hurricanes to affect the U.S. coastline by fifty-mile wide coastal segments, as well as only hurricanes, and major (or great) hurricanes. While this 50 miles approximates that of the "core" used for direct hits, there are some differences. In the Simpson and Lawrence study, a storm/hurricane/great hurricane was counted in the segment where it crossed the coast plus the next segment to the right. As indicated earlier, the "core" used in Hebert, Taylor, and Case (1984) can be smaller or larger than 50 miles, and could also affect one of the segments in Simpson and Lawrence to the left of a coastline crossing which that study would not count.

The foregoing two studies and their associated criteria are climatological with their primary purpose being for use in assessing risk based on past experience. On the other hand, the National Weather Service's Hurricane Probability Program has as its purpose the assessment of risk based on a present hurricane threat to the United States coastline. It does this by arbitrarily defining a "strike" as the center of a hurricane moving through a zone within approximately 50 nautical miles to the right or 75 nautical miles to the left of the site of interest (Sheets, 1984). The asymmetry is to allow for the strongest winds in a hurricane frequently being further to the right of the center than the left - a consideration reflected also in the earlier studies discussed. This 125 nautical mile diameter circle approximates the region of hurricane force winds for a "typical" hurricane. It will usually be larger than the "core", and is fixed, like the segments in Simpson and Lawrence. **HURRICANE STRIKE PROBABILITIES HAVE NO RELATION TO HURRICANE INTENSITY.** Users of these probabilities must take this into account when assessing risk, based on the forecast time before landfall and strength. The reader is urged to refer to The

National Weather Service Hurricane Probability Program (Sheets, 1984) for a more thorough explanation of forecast probabilities.

Statistics on total storm/hurricane activity in the North Atlantic Ocean (which includes the Gulf of Mexico and the Caribbean Sea) can be found in Neumann, et al (1987). A detailed breakdown of hurricanes by category which have affected coastal counties of the Gulf of Mexico and North Atlantic Ocean both directly and indirectly can be found in Hebert, Taylor, and Case (1984), which has been updated where necessary for this technical memorandum. The best source of how a hurricane affected individual localities or states can be found in the annual articles on the hurricane season in the Monthly Weather Review (1990 for example), the Diagnostic Report of the National Hurricane Center (1991 for example) and Storm Data (1990 for example).

THIS PAGE LEFT BLANK INTENTIONALLY

### 3. DISCUSSION

#### Part I

- (1) What have been the deadliest hurricanes of this century in the United States?

Table 2 lists the 31 deadliest hurricanes to strike the U.S. in this century. Although technically incorrect, we have included 1900 in "this century". Three hurricanes prior to 1900 as well as a tropical storm which affected southern California in 1939 are listed as an addendum because of their large death tolls.

- (2) What have been the costliest hurricanes of this century in the United States?

Table 3 lists the 32 costliest hurricanes to strike the U.S. in this century. Figures are unadjusted for inflation. Table 3a re-orders some of these plus several other hurricanes after adjusting to 1990 dollars<sup>3</sup>.

- (3) What have been the most intense hurricanes to strike the United States during this century?

Table 4 lists the 60 major hurricanes which have struck the U.S. during this century. Hurricanes are ordered by the lowest estimated central pressure and/or highest category to affect the United States at time of landfall.

A look at the lists of deadliest and costliest hurricanes of this century reveals several striking facts: (1) The twelve deadliest hurricanes were all the equivalent of a category 4 or higher, if the excessive forward speed is considered as raising the category of a hurricane by one. (2) Large death totals were primarily a result of the 15 to 20 feet or more rise of the ocean (storm surge) associated with many of these major hurricanes. All but four of the thirty-one deadliest hurricanes were major hurricanes. Two of the four were the inland flood-producing hurricanes Agnes and Diane. (3) A large portion of the damage in three of the twelve costliest tropical cyclones (Table 3) resulted from inland flooding caused by torrential rainfall in mountainous areas. (4) Nearly three-fifths of the deadliest hurricanes were the equivalent of a category four or higher, but only two-fifths of the costliest hurricanes (Table 3) met this criterion. (5) Only one of the deadliest hurricanes has occurred during the past twenty years in contrast to about half of the costliest hurricanes.

---

<sup>3</sup> Adjusted to 1990 dollars on basis of U.S. Department of Commerce composite construction cost indexes.

Table 2. The deadliest United States hurricanes of this century.

DEADLIEST HURRICANES, UNITED STATES 1900-1991  
(25 or more deaths)

HURRICANE	YEAR	CATEGORY	DEATHS
1. TX (Galveston)	1900	4	6000+
2. FL (Lake Okeechobee)	1928	4	1836
3. FL (Keys)/S TX	1919	4	600#
4. New England	1938	3*	600
5. FL (Keys)	1935	5	408
6. AUDREY (SW LA/N TX)	1957	4	390
7. NE U.S.	1944	3*	390@
8. LA (Grand Isle)	1909	4	350
9. LA (New Orleans)	1915	4	275
10. TX (Galveston)	1915	4	275
11. CAMILLE (MS/LA)	1969	5	256
12. FL (Miami)	1926	4	243
13. DIANE (NE U.S.)	1955	1	184
14. SE FL	1906	2	164
15. MS/AL/Pensacola	1906	3	134
16. AGNES (NE U.S.)	1972	1	122
17. HAZEL (SC/NC)	1954	4*	95
18. BETSY (SE FL/SE LA)	1965	3	75
19. CAROL (NE U.S.)	1954	3*	60
20. SE FL/LA/MS	1947	4	51
21. DONNA (FL/Eastern U.S.)	1960	4	50
22. GA/SC/NC	1940	2	50
23. CARLA (TX)	1961	4	46
24. TX (Velasco)	1909	3	41
25. TX (Freeport)	1932	4	40
26. S TX	1933	3	40
27. HILDA (LA)	1964	3	38
28. SW LA	1918	3	34
29. SW FL	1910	3	30
30. CONNIE (NC)	1955	3	25
31. LA	1926	3	25

\* Moving more than 30 miles per hour.

# Over 500 of these lost on ships at sea; 600-900 estimated deaths.

@ Some 344 of these lost on ships at sea.

ADDENDUM

LA	1893	-	2000
SC/GA	1893	-	1000-2000
GA/SC	1881	-	700
SOUTHERN CALIFORNIA	1939	-	45

Table 3. The costliest United States hurricanes of this century.  
(Unadjusted)

COSTLIEST HURRICANES, UNITED STATES 1900-1991  
(More than \$100,000,000 damage)

	HURRICANE	YEAR	CATEGORY	DAMAGE (U.S.)
1.	HUGO (SC)	1989	4	\$7,000,000,000
2.	FREDERIC (AL/MS)	1979	3	2,300,000,000
3.	AGNES (NE U.S.)	1972	1	2,100,000,000
4.	ALICIA (N TX)	1983	3	2,000,000,000
5.	BOB (NC, NE U.S.)	1991	2	1,500,000,000
6.	JUAN (LA)	1985	1	1,500,000,000
7.	CAMILLE (MS/AL)	1969	5	1,420,700,000
8.	BETSY (SE FL/SE LA)	1965	3	1,420,500,000
9.	ELENA (MS/AL/NW FL)	1985	3	1,250,000,000
10.	GLORIA (Eastern U.S.)	1985	3*	900,000,000
11.	DIANE (NE U.S.)	1955	1	831,700,000
12.	ALLISON (N TX)	1989	T.S.®	500,000,000
13.	ELOISE (NW FL)	1975	3	490,000,000
14.	CAROL (NE U.S.)	1954	3*	461,000,000
15.	CELIA (S TX)	1970	3	453,000,000
16.	CARLA (TX)	1961	4	408,000,000
17.	CLAUDETTE (N TX)	1979	T.S.®	400,000,000
18.	DONNA (FL/Eastern U.S.)	1960	4	387,000,000
19.	DAVID (FL/Eastern U.S.)	1979	2	320,000,000
20.	New England	1938	3*	306,000,000
21.	KATE (FL Keys/NW FL)	1985	2	300,000,000
22.	ALLEN (S TX)	1980	3	300,000,000
23.	HAZEL (SC/NC)	1954	4*	281,000,000
24.	DORA (NE FL)	1964	2	250,000,000
25.	BEULAH (S TX)	1967	3	200,000,000
26.	AUDREY (LA/N TX)	1957	4	150,000,000
27.	CARMEN (LA)	1974	3	150,000,000
28.	CLEO (SE FL)	1964	2	128,500,000
29.	HILDA (Central LA)	1964	3	125,000,000
30.	FL (Miami)	1926	4	112,000,000
31.	SE FL/LA/MS	1947	4	110,000,000
32.	NE U.S.	1944	3*	100,000,000+

\* Moving more than 30 miles per hour.

® Only of Tropical Storm intensity, but included because of high damage amount.

Table 3a. The costliest United States hurricanes of this century.  
(Adjusted to 1990 dollars) \*\*

COSTLIEST HURRICANES, UNITED STATES 1900-1991  
(More than \$400,000,000 damage)

HURRICANE	YEAR	CATEGORY	DAMAGE (U.S.)
1. HUGO (SC)	1989	4	\$7,155,120,000
2. BETSY (FL/LA)	1965	3	6,461,303,000
3. AGNES (NE U.S.)	1972	1	6,418,143,000
4. CAMILLE (MS/AL)	1969	5	5,242,380,000
5. DIANE (NE U.S.)	1955	1	4,199,645,000
6. New England	1938	3*	3,593,853,000
7. FREDERIC (AL/MS)	1979	3	3,502,942,000
8. ALICIA (N TX)	1983	3	2,391,854,000
9. CAROL (NE U.S.)	1954	3*	2,370,215,000
10. CARLA (TX)	1961	4	1,926,731,000
11. DONNA (FL/Eastern U.S.)	1960	4	1,823,605,000
12. JUAN (LA)	1985	1	1,671,232,000
13. CELIA (S TX)	1970	3	1,560,440,000
14. BOB (NC and NE U.S.)	1991	2	1,500,000,000
15. HAZEL (SC/NC)	1954	4*	1,444,752,000
16. ELENA (MS/AL/NW FL)	1985	3	1,392,693,000
17. FL (Miami)	1926	4	1,315,397,000
18. N TX (Galveston)	1915	4	1,177,937,000 <sup>1</sup>
19. DORA (NE FL)	1964	2	1,157,596,000
20. ELOISE (NW FL)	1975	3	1,081,854,000
21. GLORIA (Eastern U.S.)	1985	3*	1,002,739,000
22. NE U.S.	1944	3*	925,055,000
23. BEULAH (S TX)	1967	3	844,304,000
24. N TX (Galveston)	1900	4	706,762,000 <sup>2</sup>
25. SE FL/LA/MS	1947	4	703,859,000
26. AUDREY (LA/N TX)	1957	4	696,091,000
27. CLAUDETTE (N TX)	1979	T.S. @	609,207,000
28. CLEO (SE FL)	1964	2	595,004,000
29. SW FL/NE FL	1944	3	582,785,000
30. HILDA (LA)	1964	3	578,798,000
31. SE FL	1945	3	539,087,000
32. ALLISON (N TX)	1989	T.S. @	511,080,000
33. DAVID (FL/Eastern U.S.)	1979	2	487,366,000
34. IONE (NC)	1955	3	444,353,000
35. ALLEN (S TX)	1980	3	410,908,000

<sup>1</sup> Considered too high in 1915 reference.

<sup>2</sup> Using 1915 cost adjustment base - none available prior to 1915.

\* Moving more than 30 miles per hour.

@ Only of Tropical Storm intensity, but included because of high damage.

\*\* Adjusted to 1990 dollars on basis of U.S. Department of Commerce composite construction cost indexes.

THIS PAGE LEFT BLANK INTENTIONALLY

Table 4. The most intense United States hurricanes of this century.  
(at time of landfall)

MOST INTENSE HURRICANES, UNITED STATES 1900-1991  
(At time of landfall)

	HURRICANE	YEAR	CATEGORY	MILLIBARS	INCHES
1.	FL (Keys)	1935	5	892	26.35
2.	CAMILLE (LA/MS)	1969	5	909	26.84
3.	FL (Keys)/S TX	1919	4	927	27.37
4.	FL (Lake Okeechobee)	1928	4	929	27.43
5.	DONNA (FL/Eastern U.S.)	1960	4	930	27.46
6.	TX (Galveston)	1900	4	931	27.49
7.	LA (Grand Isle)	1909	4	931	27.49
8.	LA (New Orleans)	1915	4	931	27.49
9.	CARLA (TX)	1961	4	931	27.49
10.	HUGO (SC)	1989	4	934	27.58
11.	FL (Miami)	1926	4	935	27.61
12.	HAZEL (SC/NC)	1954	4*	938	27.70
13.	SE FL/LA/MS	1947	4	940	27.76
14.	N TX	1932	4	941	27.79
15.	GLORIA (Eastern U.S.)	1985	3*&	942	27.82
16.	AUDREY (LA/N TX)	1957	4#	945	27.91
17.	TX (Galveston)	1915	4#	945	27.91
18.	CELIA (S TX)	1970	3	945	27.91
19.	ALLEN (S TX)	1980	3@	945	27.91
20.	New England	1938	3*	946	27.94
21.	FREDERIC (AL/MS)	1979	3	946	27.94
22.	NE U.S.	1944	3*	947	27.97
23.	SC/NC	1906	3	947	27.97
24.	BETSY (SE FL/SE LA)	1965	3	948	27.99
25.	SE FL/NW FL	1929	3	948	27.99
26.	SE FL	1933	3	948	27.99
27.	S TX	1916	3	948	27.99
28.	MS/AL	1916	3	948	27.99
29.	DIANA (NC)	1984	3+	949	28.02
30.	S TX	1933	3	949	28.02
31.	BEULAH (S TX)	1967	3	950	28.05
32.	HILDA (Central LA)	1964	3	950	28.05
33.	GRACIE (SC)	1959	3	950	28.05
34.	TX (Central)	1942	3	950	28.05
35.	SE FL	1945	3	951	28.08

Continued on next page

- \* Moving more than 30 miles per hour.
- & Winds and tides did not justify 4.
- # Classified 4 because of extreme tides.
- @ Reached Cat. 5 intensity three times along its path through the Caribbean and Gulf of Mexico. The lowest pressure reported was 899 mb (26.55 in.) at 1742 UTC 8/7/80 off the northeastern tip of the Yucatan Peninsula.
- + Cape Fear, NC area only; was a category 2 at final landfall.

Table 4 continued.

	HURRICANE	YEAR	CATEGORY	MILLIBARS	INCHES
36.	FL (Tampa Bay)	1921	3	952	28.11
37.	CARMEN (Central LA)	1974	3	952	28.11
38.	EDNA (New England)	1954	3*	954	28.17
39.	SE FL	1949	3	954	28.17
40.	ELOISE (NW FL)	1975	3	955	28.20
41.	KING (SE FL)	1950	3	955	28.20
42.	Central LA	1926	3	955	28.20
43.	SW LA	1918	3	955	28.20
44.	SW FL	1910	3	955	28.20
45.	NC	1933	3	957	28.26
46.	FL (Keys)	1909	3	957	28.26
47.	EASY (NW FL)	1950	3	958	28.29
48.	N TX	1941	3	958	28.29
49.	NW FL	1917	3	958	28.29
50.	N TX	1909	3	958	28.29
51.	MS/AL	1906	3	958	28.29
52.	ELENA (MS/AL/NW FL)	1985	3	959	28.32
53.	CAROL (NE U.S.)	1954	3*	960	28.35
54.	IONE (NC)	1955	3	960	28.35
55.	ALICIA (N TX)	1983	3	962	28.41
56.	CONNIE (NC/VA)	1955	3	962	28.41
57.	SW FL/NE FL	1944	3	962	28.41
58.	Central LA	1934	3	962	28.41
59.	SW FL/NE FL	1948	3	963	28.44
60.	NW FL	1936	3	964	28.47

\* Moving more than 30 miles per hour.

DIRECT HITS BY HURRICANES  
U.S. GULF & ATLANTIC COASTS  
1900-1991

Category 5:	2
4:	14
3:	44
2:	35
1:	57
<u>TOTAL</u>	<u>152</u>

Major hurricanes (categories 3,4,5): 60

This means that during the period 1900-1991, an average of 2 major hurricanes every 3 years made landfall somewhere along the U.S. Gulf or Atlantic coast. (All categories combined average about 5 hurricanes every 3 years for the same period.)

One of the greatest concerns of the National Weather Service's (NWS) hurricane preparedness officials is that the statistics in tables 1-4 will mislead people into thinking that no more large loss of life will occur in a hurricane because of our advanced technology. Dr. Robert Sheets, spokesman for the NWS hurricane warning service and Director of NHC, as well as former Director, Dr. Neil Frank, have repeatedly emphasized the great danger of a catastrophic loss of life in a future hurricane if proper preparedness plans for vulnerable areas are not formulated.

The study by Hebert, Taylor, and Case (1984), now being updated with 1990 census data, shows that 85% of U.S. coastal residents from Texas to Maine have never experienced a direct hit by a major hurricane. Many of those 45 million residents had moved to coastal sections during the past twenty-five years. Even the landfall of Hugo has not lessened an ever increasing concern brought by the continued increase in coastal populations.

A look at Table 5 which lists hurricanes by decades in this century shows that during the twenty year period 1960-1979 both the number and intensity of landfalling U.S. hurricanes decreased sharply! Based on 1900-1959 statistics from the same study, the expected number of hurricanes and major hurricanes during the period 1960-1979 was 36 and 15, respectively. In fact, only 27 or 75% of the expected number of hurricanes struck the U.S. with only 10 major hurricanes or 67% of that expected number. The decade of the eighties showed little change to this trend.

Hurricane Hugo became the first category 4 or 5 hurricane to strike the U.S. since Camille in 1969, a period of 20 years. On the average a category 4 or greater hurricane strikes the U.S. once every 6 years.

Fewer hurricanes do not necessarily mean a lesser threat of disaster, however. The 1919 hurricane which was both the third deadliest and third most intense of this century to strike the U.S. occurred in a year which had a total of only three storms/hurricanes. The most intense U.S. hurricane of record in 1935 and the eighth costliest in 1965 (Betsy) occurred in years which had a total of only six storms/hurricanes.

The conclusions are obvious. A large death toll in a U.S. hurricane is still possible. The decreased death totals in recent years may be as much a result of lack of major hurricanes striking the most vulnerable areas as they are of any fail-proof forecasting, warning, and observing systems. Continued coastal growth and inflation will almost certainly result in every future major landfalling hurricane (and even weaker hurricanes and tropical storms) replacing one of the current costliest hurricanes. If warnings are heeded and preparedness plans developed, the death toll can be reduced, but, in the absence of a change of attitude or laws restricting building near the ocean, large property losses are inevitable.

Table 5. Number of hurricanes of various categories to strike the United States each decade. Updated from Hebert and Case (1990).

DECADE	CATEGORY					ALL	MAJOR
	1	2	3	4	5	1,2,3,4,5	3,4,5
1900-1909	5	5	4	2		16	6
1910-1919	8	3	5	3		19	8
1920-1929	6	4	3	2		15	5
1930-1939	4	5	6	1	1	17	8
1940-1949	7	8	7	1		23	8
1950-1959	8	1	7	2		18	9
1960-1969	4	5	3	2	1	15	6
1970-1979	6	2	4			12	4
1980-1989	9	1	5	1		16	6
1990-1991		1				1	0
1900-1991	57	35	44	14	2	152	60

Note: Only the highest category to affect the U.S. has been used.

## Part II

In addition, to information about U.S. hurricanes, this section will also include statistics on total tropical storm and hurricane activity.

- (1) What is the average number of hurricanes per year?

Table 6 gives the average number of tropical cyclones which reached storm strength and hurricane strength for various time periods. A total of ten tropical cyclones reaching storm strength with six of these becoming hurricanes appears to be the best averages to use based on the past 40 to 50 year time period. The averages of eight tropical cyclones and five hurricanes for the longer 106-year period is a reflection of less detection and fewer actual storms prior to 1930.

- (2) What year(s) have had the most and least hurricanes?

Table 7 shows the years of maximum and minimum tropical cyclone and hurricane activity for the entire Atlantic Ocean. The only years when a hurricane failed to strike the U.S. coast were 1902, 1905, 1907, 1914, 1922, 1927, 1930, 1931, 1937, 1951, 1958, 1962, 1973, 1978, 1981, 1982 and 1990. Note that only twice has the U.S. gone as long as two years without a hurricane. The most hurricanes to strike the U.S. in one year were six in 1916 and 1985. There were five in 1933, and four in 1906, 1909, and 1964. Three hurricanes have struck the U.S. in one year a total of fifteen times. Ten of these fifteen times occurred during the sixteen years from 1944 to 1959! A chronological list of all hurricanes to strike the U.S. during this century through 1982 including month, category by states affected, and minimum sea level pressure at landfall can be found in Hebert, Taylor, and Case (1984). That publication is targeted for updating during 1992.

Table 6. Average number of tropical cyclones which reached storm strength and hurricane strength for various time periods. Updated from Neumann, et al (1987).

PERIOD	NUMBER OF YEARS	AVERAGES (PER YEAR)	
		TROPICAL CYCLONES	HURRICANES
1886-1991	106	8.4	4.9
1942-1991	50	9.8	5.7
1952-1991	40	9.7	5.6
1962-1991	30	9.6	5.4
1972-1991	20	9.4	5.1
1977-1991	15	9.5	5.3
1982-1991	10	9.2	4.8

Includes subtropical storms after 1967.

Table 7. Years of maximum and minimum tropical cyclone and hurricane activity in the North Atlantic Ocean, Caribbean Sea, and Gulf of Mexico during the period 1871-1991 (from Neumann, et al, 1987).

MAXIMUM ACTIVITY

TROPICAL CYCLONES <sup>1</sup>		HURRICANES <sup>2</sup>	
NUMBER	YEAR(S)	NUMBER	YEAR(S)
21	1933	12	1969
18	1969	11	1916, 1950
17	1887	10	1887, 1893, 1933
16	1936	9	1955, 1980

MINIMUM ACTIVITY

TROPICAL CYCLONES <sup>1</sup>		HURRICANES <sup>2</sup>	
NUMBER	YEAR(S)	NUMBER	YEAR(S)
1	1890, 1914	0	1907, 1914
2	1925, 1930	1	1890, 1905, 1919, 1925
		2	1895, 1897, 1904, 1917, 1922, 1930, 1931, 1982

<sup>1</sup> Includes subtropical storms after 1967. See Neumann, et al (1987).

<sup>2</sup> After 1885.

- (3) When did the earliest and latest hurricane occur?

The hurricane season is defined as June 1 through November 30. An early hurricane can be defined as occurring in the three months prior to the start of the season, and a late hurricane can be defined as occurring in the three months after the season. With these criteria the earliest observed hurricane in the Atlantic was on March 7, 1908, while the latest observed hurricane was on December 31, 1954. The earliest hurricane to strike the U.S. in this century was Alma which struck northwest Florida on June 9, 1966. The latest hurricane to strike the U.S. was late on November 30, 1925 near Tampa, Florida.

- (4) What were the longest-lived and shortest-lived hurricanes?

Ginger in 1971 holds the record for both the most number of days as a hurricane (20) and tropical cyclone (31). There have been many tropical cyclones which attained hurricane intensity for periods of 12 hours or less.

- (5) What were the strongest and weakest Atlantic hurricanes? To strike the United States?

In terms of central pressure (and probably winds) the strongest observed hurricane in the Atlantic basin was Gilbert in 1988 with a pressure of 888 millibars while located in the northwest Caribbean. The 1935 Labor Day hurricane in the Florida Keys with a pressure of 892 millibars was the strongest hurricane to strike the U.S. Numerous hurricanes have reached only the minimum wind speed near 74 miles per hour and struck the U.S.

- (6) How many hurricanes have there been in each month?

Table 8 adapted from Neumann, et al (1987) shows the total and average number of tropical cyclones and those which became hurricanes by months for the period 1886-1991. In addition, the monthly total and average number of hurricanes to strike the U.S. in this century (updated from Hebert and Case, 1990) are given.

Table 8. Total and average number of tropical cyclones and hurricanes in the North Atlantic Ocean, Caribbean Sea, and Gulf of Mexico by month of origin for the period 1886-1991 (from Neumann, et al, 1987), and for hurricanes striking the U.S. coast in this century (updated from Hebert and Case, 1990).

MONTH	TROPICAL CYCLONES <sup>1</sup>		HURRICANES		U.S. HURRICANES <sup>2</sup>	
	TOTAL	AVG.	TOTAL	AVG.	TOTAL	AVG.
JAN-APRIL	3	*	1	*	0	0.0
MAY	14	0.1	3	*	0	0.0
JUNE	56	0.5	23	0.2	11	0.1
JULY	68	0.7	35	0.3	16	0.2
AUGUST	216	2.0	150	1.4	37	0.3
SEPTEMBER	304	2.9	191	1.8	61	0.6
OCTOBER	187	1.8	95	0.9	23	0.2
NOVEMBER	43	0.4	22	0.2	4	*
DECEMBER	6	0.1	3	*	0	0.0
YEAR	897	8.5	523	4.9	152	1.4

<sup>1</sup> Includes subtropical storms after 1967. See Neumann, et al (1987) for details.

<sup>2</sup> 1900-1991.

\* Less than 0.05.

- (7) What was the largest number of hurricanes in existence in the Atlantic Ocean at the same time?

According to information on the current version of the master data file of Neumann, et al (1987), there have never been four hurricanes in existence in the North Atlantic at the same time in this century. On August 22, 1893 four hurricanes co-existed, one of them being the hurricane which killed an estimated 2,000 people in Georgia-South Carolina several days later. On September 11, 1961 three hurricanes and possibly a fourth existed. The only other years in this century with three hurricanes on the map at the same time were 1950 and 1967.

- (8) How many direct hits by hurricanes of various categories have affected each state?

Table 9, updated from Hebert and Case (1990), shows the number of hurricanes (direct hits) affecting the U.S. and individual states. The table shows that on the average close to two hurricanes per year strike the U.S., while two major hurricanes cross the U.S. coast somewhere every three years. Other noteworthy facts, updated from Hebert and Case (1990), are: 1.) Thirty-six percent of all U.S. hurricanes hit Florida; 2.) Sixty-nine percent of category 4 or higher hurricanes have hit either Florida or Texas; 3.) Along the middle Gulf coast, southern Florida, and New York and southern New England, approximately one out of every two hurricanes is major one.

Table 9. Number of hurricanes (direct hits) affecting the U.S. and individual states 1900-1991 according to Saffir/Simpson scale (updated from Hebert and Case, 1990).

AREA	CATEGORY NUMBER					ALL	MAJOR HURRICANES (≥3)
	1	2	3	4	5		
U.S. (Texas to Maine)	57	35	44	14	2	152	60
Texas	12	9	9	6	0	36	15
(North)	7	3	3	4	0	17	7
(Central)	2	2	1	1	0	6	2
(South)	3	4	5	1	0	13	6
Louisiana	8	5	7	3	1	24	11
Mississippi	1	1	5	0	1	8	6
Alabama	4	1	5	0	0	10	5
Florida	17	15	16	5	1	54	22
(Northwest)	9	7	6	0	0	22	6
(Northeast)	1	7	0	0	0	8	0
(Southwest)	6	3	5	2	1	17	8
(Southeast)	4	10	7	3	0	24	10
Georgia	1	4	0	0	0	5	0
South Carolina	6	4	2	2	0	14	4
North Carolina	10	3	8	1*	0	22	9
Virginia	2	1	1*	0	0	4	1*
Maryland	0	1*	0	0	0	1*	0
Delaware	0	0	0	0	0	0	0
New Jersey	1*	0	0	0	0	1*	0
New York	3	1*	5*	0	0	9	5*
Connecticut	2	3*	3*	0	0	8	3*
Rhode Island	0	2*	3*	0	0	5*	3*
Massachusetts	2	2*	2*	0	0	6	2*
New Hampshire	1*	1*	0	0	0	2*	0
Maine	5	0	0	0	0	5	0

\*. Indicates all hurricanes in this category were moving greater than 30 mph.

Note: State totals will not equal U.S. totals and Texas and Florida sectional totals will not necessarily equal state totals.

- (9) When are the major hurricanes likely to strike given areas?

Table 10 shows the incidence of major hurricanes by months for the U.S. and individual states. For the United States as a whole, September has had more major hurricanes than all other months combined. However, three of the most devastating hurricanes did not occur in September--AUDREY (June 1957), CAMILLE (August 1969), and HAZEL (October 1954). Only in Texas and Louisiana are major hurricanes in August and September almost an equal threat. Most major October hurricanes occur in southern Florida.

- (10) How long has it been since a major hurricane directly hit a given community? Any hurricane? Indirectly?

Table 11 summarizes the occurrence of the last major hurricane or of any hurricane to directly hit the more populated coastal communities from Brownsville, Texas to Eastport, Maine. In addition, if a hurricane indirectly affected a community after the last direct hit, it is listed in the last column of the table.

Perhaps the most illustrative example of the uncertainty of when a hurricane might strike a given locality is Pensacola, Florida. Although Dunn (1967) listed Pensacola as the city with the second highest frequency of hurricane force winds in the United States (1 in 10), it has been more than 60 years since any hurricane directly struck Pensacola!

In order to obtain the same type of information listed in Table 11 for the remaining coastal communities, the reader is referred to Hebert, Taylor, and Case (1984).

- (11) What is the total United States damage (unadjusted and adjusted) and death toll for each year of this century?

Table 12 summarizes this information. Tables 12a-c rank the years. In most years the death and damage totals are usually the result of a single, major hurricane. Gentry (1966) gives damages adjusted to 1957-59 costs as a base for the period 1915-1965. For the most part, death and damage totals for the period 1915-1965 were taken from Gentry's paper, and for the remaining years from the Monthly Weather Review. Adjusted damages were calculated to 1990 dollars by the same factors as used in Table 3a.

Table 10. Incidence of major hurricanes (direct hits) by months to affect the United States and individual states (1900-1991) according to the Saffir/Simpson Scale (updated from Hebert and Case, 1990).

AREA	MONTH					ALL
	JUNE	JULY	AUG.	SEPT.	OCT.	
U.S. (Texas to Maine)	2	3	13	35	7	60
Texas	1	1	7	6		15
(North)	1	1	3	2		7
(Central)			1	1		2
(South)			3	3		6
Louisiana	2		3	5	1	11
Mississippi		1	1	4		6
Alabama		1		4		5
Florida		1	1	15	5	22
(Northwest)		1		5		6
(Northeast)						0
(Southwest)				5	3	8
(Southeast)			1	7	2	10
Georgia				7		0
South Carolina				3	1	4
North Carolina			1	7	1	9
Virginia				1		1
Maryland						0
Delaware						0
New Jersey						0
New York			1	4		5
Connecticut			1	2		3
Rhode Island			1	2		3
Massachusetts				2		2
New Hampshire						0
Maine						0

Note: State totals will not equal U.S. totals and Texas and Florida sectional totals will not necessarily equal state totals.

Table 11. Last occurrence of a direct or indirect hit by any hurricane and/or by a major hurricane at the more populated coastal communities from Texas to Maine (updated from Hebert and Case, 1990). Category is in parentheses.

STATE	CITY	DIRECT			INDIRECT		
		LAST MAJOR		LAST ANY		LAST ANY	
Texas	Brownsville	1980(3)	Allen	1980(3)	Allen		
	Corpus Christi	1970(3)	Celia	1971(1)	Fern	1980(3)	Allen
	Port Aransas	1970(3)	Celia	1971(1)	Fern	1980(3)	Allen
	Matagorda	1961(4)	Carla	1971(1)	Fern	1983(3)	Alicia
	Freeport	1983(3)	Alicia	1983(3)	Alicia		
	Galveston	1983(3)	Alicia	1989(1)	Jerry		
	Houston	1941(3)		1989(1)	Jerry		
	Beaumont	<1900		1986(1)	Bonnie		
Louisiana	Cameron	1957(4)	Audrey	1985(1)	Danny	1985(1)	Juan
	Morgan City	1974(3)	Carmen	1985(1)	Juan		
	Houma	1974(3)	Carmen	1985(1)	Juan		
	New Orleans	1965(3)	Betsy	1965(3)	Betsy	1969(5)	Camille
Mississippi	Bay St. Louis	1985(3)	Elena	1985(3)	Elena		
	Biloxi	1985(3)	Elena	1985(3)	Elena		
	Pascagoula	1985(3)	Elena	1985(3)	Elena		
Alabama	Mobile	1985(3)	Elena	1985(3)	Elena		
Florida	Pensacola	1926(3)		1926(3)		1979(3)	Frederic
	Panama City	1975(3)	Eloise	1985(2)	Kate		
	Apalachicola	1985(3)	Elena	1985(2)	Kate		
	Homosassa	1950(3)	Easy	1968(2)	Gladys		
	St. Petersburg	1921(3)		1946(1)		1968(2)	Gladys
	Tampa	1921(3)		1946(1)		1968(2)	Gladys
	Sarasota	1944(3)		1946(1)		1966(2)	Alma
	Fort Myers	1960(3)	Donna	1960(3)	Donna	1966(2)	Alma
	Naples	1960(4)	Donna	1964(2)	Isbell	1965(3)	Betsy
	Key West	1948(3)		1987(1)	Floyd		
	Miami	1950(3)	King	1964(2)	Cleo	1965(3)	Betsy
	Fort Lauderdale	1950(3)	King	1964(2)	Cleo	1965(3)	Betsy
	West Palm Beach	1949(3)		1979(2)	David		
	Stuart	1949(3)		1979(2)	David		
	Fort Pierce	1933(3)		1979(2)	David		
	Vero Beach	<1900		1979(2)	David		
	Cocoa	<1900		1979(2)	David		

Table 11 continued.

STATE	CITY	DIRECT			INDIRECT	
		LAST MAJOR		LAST ANY	LAST ANY	
Florida (cont.)						
	Daytona Beach	<1900		1960(2)	Donna	1979(2) David
	St. Augustine	<1900		1964(2)	Dora	
	Jacksonville	<1900		1964(2)	Dora	
	Fernandina Beach	<1900		1928(2)		1964(2) Dora
Georgia	Brunswick	<1900		1928(1)		
	Savannah	<1900		1979(2)	David	
South Carolina	Hilton Head	1959(3)	Gracie	1979(2)	David	1985(1) Bob
	Charleston	1989(4)	Hugo	1989(4)	Hugo	
	Myrtle Beach	1954(4*)	Hazel	1954(4*)	Hazel	1989(4) Hugo
North Carolina	Wilmington	1960(3*)	Donna	1984(2)	+Diana	
	Morehead City	1960(3*)	Donna	1960(3*)	Donna	1985(3*) Gloria
	Cape Hatteras	1985(3*)	Gloria	1986(1)	Charley	
Virginia	Virginia Beach	1944(3*)		1986(1)	Charley	
	Norfolk	<1900		1955(1)	Connie	1985(3*) Gloria
Maryland	Ocean City	<1900		<1900		1985(3*) Gloria
	Baltimore	<1900		<1900		1954(2*) Hazel
Delaware	Rehoboth Beach	<1900		<1900		1985(3*) Gloria
	Wilmington	<1900		<1900		1954(2*) Hazel
New Jersey	Cape May	<1900		1903(1)		1985(3*) Gloria
	Atlantic City	<1900		1903(1)		1985(3*) Gloria
New York	New York City	<1900		1903(1)		1976(1) Belle
	Westhampton	1985(3*)	Gloria	1985(3*)	Gloria	
Connecticut	New London	1938(3*)		1991(2*)	Bob	
	New Haven	1938(3*)		1985(2*)	Gloria	
	Bridgeport	1954(3*)	Carol	1985(2*)	Gloria	
Rhode Island	Providence	1954(3*)	Carol	1991(2*)	Bob	
Mass.	Cape Cod	1954(3*)	Edna	1991(2*)	Bob	
	Boston	<1900		1960(1*)	Donna	1991(1*) Bob
New Hamp.	Portsmouth	<1900		1985(2*)	Gloria	
Maine	Portland	<1900		1985(1*)	Gloria	
	Eastport	<1900		1969(1)	Gerda	1985(1*) Gloria

+ Cape Fear only - Direct 3.

\* Moving more than 30 miles per hour.

&lt;1900 means before 1900

Table 12. Estimated annual deaths and damages (unadjusted and adjusted<sup>1</sup>) in the United States for each year of this century.

YEAR	DEATHS	DAMAGE (\$MILLIONS)		YEAR	DEATHS	DAMAGE (\$MILLIONS)	
		UNADJUSTED	ADJUSTED			UNADJUSTED	ADJUSTED
1900	6000+	30	(735) <sup>2</sup>	1946	0	5	38
1901	10	1	*	1947	53	136	870
1902	0	Minor	Minor	1948	3	18	105
1903	15	1	*	1949	4	59	344
1904	5	2	*	1950	19	36	206
1905	0	Minor	Minor	1951	0	2	10
1906	298	3+	*	1952	3	3	15
1907	0	0	0	1953	2	6	31
1908	0	0	0	1954	193	756	3887
1909	406	8	*	1955	218	985	4974
1910	30	1	*	1956	19	27	129
1911	17	1+	*	1957	400	152	705
1912	1	Minor	Minor	1958	2	11	51
1913	5	3	*	1959	24	23	108
1914	0	0	0	1960	65	396	1866
1915	550	63	1544 <sup>3</sup>	1961	46	414	1955
1916	107	33	672	1962	3	2	9
1917	5	Minor	Minor	1963	10	12	55
1918	34	5	66	1964	49	515	2385
1919	287	22	259	1965	75	1445	6572
1920	2	3	28	1966	54	15	65
1921	6	3	35	1967	18	200	844
1922	0	0	0	1968	9	10	40
1923	0	Minor	Minor	1969	256	1421	5244
1924	2	Minor	Minor	1970	11	454	1564
1925	6	Minor	Minor	1971	8	213	686
1926	269	112	1316	1972	121	2100	6418
1927	0	0	0	1973	5	3	8
1928	1836	25	293	1974	1	150	361
1929	3	1	11	1975	21	490	1082
1930	0	Minor	Minor	1976	9	100	213
1931	0	0	0	1977	0	10	19
1932	0	0	0	1978	36	20	35
1933	63	47	652	1979	22	3045	4638
1934	17	5	63	1980	2	300	411
1935	414	12	152	1981	0	25	32
1936	9	2	26	1982	0	Minor	Minor
1937	0	Minor	Minor	1983	22	2000	2392
1938	600	306	3594	1984	4	66	77
1939	3	Minor	Minor	1985	30	4000	4457
1940	51	5	61	1986	9	17	18
1941	10	8	91	1987	0	8	8
1942	8	27	266	1988	6	9	9
1943	16	17	157	1989	56	7670	7840
1944	64	165	1526	1990	13	57	57
1945	7	80	719	1991	16	1500	1500

<sup>1</sup> Adjusted to 1990 dollars on the basis of U.S. Department of Commerce composite construction cost index.

<sup>2</sup> Using 1915 cost adjustment - none available prior to 1915.

<sup>3</sup> Considered too high in 1915 reference.

\* Not available.

Table 12a. Estimated deaths in the United States for each year of this century ranked according to annual total.

RANK	YEAR	TOTAL	RANK	YEAR	TOTAL
1	1900	6000+	37	1934	17
2	1928	1836	39	1991	16
3	1938	600	39	1943	16
4	1915	550	41	1903	15
5	1935	414	42	1990	13
6	1909	406	43	1970	11
7	1957	400	44	1901	10
8	1906	298	44	1941	10
9	1919	287	44	1963	10
10	1926	269	47	1936	9
11	1969	256	47	1968	9
12	1955	218	47	1976	9
13	1954	193	47	1986	9
14	1972	121	53	1942	8
15	1916	107	51	1971	8
16	1965	75	51	1945	7
17	1960	65	54	1921	6
18	1944	64	54	1925	6
19	1933	63	54	1988	6
20	1989	56	57	1904	5
21	1966	54	57	1913	5
22	1947	53	57	1917	5
23	1940	51	57	1973	5
24	1964	49	61	1949	4
25	1961	46	61	1984	4
26	1978	36	63	1929	3
27	1918	34	63	1939	3
28	1910	30	63	1948	3
28	1985	30	63	1952	3
30	1959	24	63	1962	3
31	1979	22	68	1920	2
31	1983	22	68	1924	2
33	1975	21	68	1953	2
34	1950	19	68	1958	2
34	1956	19	68	1980	2
36	1967	18	73	1912	1
37	1911	17	73	1974	1

Note 1: There were 18 years in which no deaths occurred...1902, 1905, 1907, 1908, 1914, 1922, 1923, 1927, 1930, 1931, 1932, 1937, 1946, 1951, 1977, 1981, 1982, 1987.

Note 2: Death totals in this table do not agree with those in table 2 because the latter included deaths on ships at sea.

Table 12b. Estimated damages (unadjusted) in the United States for each year of this century ranked according to annual total.

RANK	YEAR	DAMAGE (\$ millions)	RANK	YEAR	DAMAGE (\$ millions)
1	1989	7670	37	1981	25
2	1985	4000	39	1959	23
3	1979	3045	40	1919	22
4	1972	2100	41	1978	20
5	1983	2000	42	1948	18
6	1991	1500	43	1943	17
7	1965	1445	43	1986	17
8	1969	1421	45	1966	15
9	1955	985	46	1935	12
10	1954	756	46	1963	12
11	1964	515	48	1958	11
12	1975	490	49	1968	10
13	1970	454	49	1977	10
14	1961	414	51	1988	9
15	1960	396	52	1909	8
16	1938	306	52	1941	8
17	1980	300	52	1987	8
18	1971	213	55	1953	6
19	1967	200	56	1918	5
20	1944	165	56	1934	5
21	1957	152	56	1940	5
22	1974	150	56	1946	5
23	1947	136	60	1906	3+
24	1926	112	61	1913	3
25	1976	100	61	1920	3
26	1945	80	61	1921	3
27	1984	66	61	1952	3
28	1915	63	61	1973	3
29	1949	59	66	1904	2
30	1990	57	66	1936	2
31	1933	47	66	1951	2
32	1950	36	66	1962	2
33	1916	33	70	1911	1+
34	1900	30	71	1901	1
35	1942	27	71	1903	1
35	1956	27	71	1910	1
37	1928	25	71	1929	1

Note 1: Eleven years had minor damage...1902, 1905, 1912, 1917, 1923, 1924, 1925, 1930, 1937, 1939, and 1982.

Note 2: Seven years had no damage...1907, 1908, 1914, 1922, 1927, 1931, and 1932.

Table 12c. Estimated damages (adjusted to 1990 dollars)<sup>1</sup> in the U.S. for each year of this century ranked according to annual total.

RANK	YEAR	DAMAGE (\$ millions)	RANK	YEAR	DAMAGE (\$ millions)
1	1989	7840	34	1976	213
2	1965	6572	35	1950	206
3	1972	6418	36	1943	157
4	1969	5244	37	1935	152
5	1955	4974	38	1956	129
6	1979	4638	39	1959	108
7	1985	4457	40	1948	105
8	1954	3887	41	1941	91
9	1938	3594	42	1984	77
10	1983	2392	43	1918	66
11	1964	2385	44	1966	65
12	1961	1955	45	1934	63
13	1960	1866	46	1940	61
14	1970	1564	47	1990	57
15	1915	1544 <sup>2</sup>	48	1963	55
16	1944	1526	49	1958	51
17	1991	1500	50	1968	40
18	1926	1316	51	1946	38
19	1975	1082	52	1921	35
20	1947	870	52	1978	35
21	1967	844	54	1981	32
22	1900	735 <sup>3</sup>	55	1953	31
23	1945	719	56	1920	28
24	1957	705	57	1936	26
25	1971	686	58	1977	19
26	1916	672	59	1986	18
27	1933	652	60	1952	15
28	1980	411	61	1929	11
29	1974	361	62	1951	10
30	1949	344	63	1962	9
31	1928	293	64	1988	9
32	1942	266	65	1973	8
33	1919	259	65	1987	8

Note 1: Eleven years with minor damage...1902, 1905, 1912, 1917, 1923, 1924, 1925, 1930, 1937, 1939, and 1982.

Note 2: Seven years with no damage...1907, 1908, 1914, 1922, 1927, 1931, 1932.

Note 3: Eight years prior to 1915 not adjusted...1901, 1903, 1904, 1906, 1909, 1910, 1911, 1913.

<sup>1</sup> Adjusted to 1990 dollars on basis of U.S. Department of Commerce composite construction cost index.

<sup>2</sup> Considered too high in 1915 reference.

<sup>3</sup> Using 1915 cost adjustment base - none available prior to 1915.

(12) Are there hurricane cycles?

Figures 1 through 10 show the landfalling portions of the tracks of major hurricanes which have struck any portion of the United States during this century.

The reader might note the tendency of the major hurricanes to cluster in certain areas during certain decades. Another interesting point is the general tendency for this clustering to occur in the latter half of individual decades in one area and in the first half of individual decades in another area. During the very active period of the thirties this clustering is not apparent.

A comparison of twenty-year periods beginning in 1900 indicates that the major hurricanes tended to be in the western Gulf Coast states at the beginning of the century, shifting to the eastern Gulf Coast states and Florida during the next twenty years, then to Florida and the Atlantic Coast states during the forties and fifties, and back to the western Gulf Coast states in the sixties and seventies. Do figures 9 and 10 indicate a shift to the eastern Gulf Coast state Florida, and the Atlantic Coast states in the eighties and nineties?

THIS PAGE LEFT BLANK INTENTIONALLY




Figure 1. Major landfalling United States hurricanes (greater than or equal to a category 3) during the period 1901-1910.




Figure 2. Major landfalling United States hurricanes (greater than or equal to a category 3) during the period 1911-1920.




Figure 3. Major landfalling United States hurricanes (greater than or equal to a category 3) during the period 1921-1930.




Figure 4. Major landfalling United States hurricanes (greater than or equal to a category 3) during the period 1931-1940.




Figure 5. Major landfalling United States hurricanes (greater than or equal to a category 3) during the period 1941-1950.




Figure 6. Major landfalling United States hurricanes (greater than or equal to a category 3) during the period 1951-1960.




Figure 7. Major landfalling United States hurricanes (greater than or equal to a category 3) during the period 1961-1970.




Figure 8. Major landfalling United States hurricanes (greater than or equal to a category 3) during the period 1971-1980.




Figure 9. Major landfalling United States hurricanes (greater than or equal to a category 3) during the period 1981-1990.




Figure 10. Major landfalling United States hurricanes (greater than or equal to a category 3) during the period 1991-2000.

(13) Are there hurricane cycles evident in certain years regardless of category or geographical area?

Table 13 gives a tabulation of hurricanes of all categories to affect the U.S. by individual years within each decade.

Table 13. Major and all category landfalling hurricanes in the United States by individual years.

	<u>MAJOR HURRICANES</u>										
	00	01	02	03	04	05	06	07	08	09	Total
1900-09	1						2			3	6
1910-19	1					2	2	1	1	1	8
1920-29		1					2		1	1	5
1930-39			1	3	1	1	1		1		8
1940-49		1	1		2	1		1	1	1	8
1950-59	2				3	2		1		1	9
1960-69	1	1			1	1		1		1	6
1970-79	1				1	1				1	4
1980-89	1			1	1	2				1	6
TOTAL	7	3	2	4	9	10	7	4	4	10	60
1990-99		1									

	<u>ALL HURRICANES</u>										
	00	01	02	03	04	05	06	07	08	09	Total
1900-09	1	2		2	1		4		1	4	15
1910-19	2	2	2	2		3	6	1	1	1	20
1920-29	2	2		1	2	1	3		2	2	15
1930-39			2	5	2	2	3		2	1	17
1940-49	2	2	2	1	3	3	1	3	3	3	23
1950-59	3		1	3	3	3	1	1		3	18
1960-69	2	1		1	4	1	2	1	1	2	15
1970-79	1	3	1		1	1	1	1		3	12
1980-89	1			1	1	6	2	1	1	3	16
TOTAL	14	12	8	16	17	20	23	8	11	22	151
1990-99		1									

Figures 1 through 10 certainly support the existence of a cyclical nature of major hurricanes affecting given regions. Table 13 is also suggestive of preferred periods. However, it is left to the reader to decide what weight should be given to these statistics.

## SUMMARY

In virtually every coastal city of any size from Texas to Maine, both present National Hurricane Center Director, Dr. Robert Sheets, and former Director, Dr. Neil Frank, have stated that the United States is building toward a hurricane disaster. The population growth versus low hurricane experience levels indicated in Hebert, Taylor, and Case (1984), together with updated statistics presented in the discussion section of this paper, form the basis for their statements. Stated simply, the areas of the United States where 9 out of 10 persons have lost their lives by drowning from the storm surge during hurricanes (along the immediate Gulf of Mexico and Atlantic shorelines) are the very areas where the most dramatic increases in population have occurred in recent years. This situation, in combination with continued building on low coastal elevations, will lead to serious problems for many areas in future hurricanes. Since it is likely that people will always live along the immediate shoreline, a pleasant way of life, the solution to the problem lies in education and preparedness.

The message to coastal residents is this: Become familiar with what hurricanes can do, and when a hurricane threatens your area, increase your chances of survival by moving away from the water until the hurricane has passed! Unless this message is clearly understood by coastal residents through a thorough and continuing preparedness effort, a future disaster is inevitable.

Acknowledgements: The work of J.G. Taylor and R.A. Case, co-authors of previous versions of this paper, contributed much. Joan David drafted the figures. Linda Kremkau provided the 1990 census data.

## REFERENCES

- Dunn, G.E., and Staff, 1967: Florida Hurricanes, ESSA Technical Memorandum WBTM-SR-38, 28 pp.
- Gentry, R.C., 1966: Nature and Scope of Hurricane Damage, American Society for Oceanography, Hurricane Symposium, Publication Number One, 344 pp.
- Hebert, P.J., J.G. Taylor, and R.A. Case, 1984: Hurricane Experience Levels of Coastal County Populations - Texas to Maine, NOAA, Technical Memorandum NWS-NHC-24, 127 pp.
- Hebert, P.J., and R.A. Case, 1990: The Deadliest, Costliest and Most Intense United States Hurricanes of This Century (and Other Frequently Requested Hurricane Facts), NOAA, Technical Memorandum NWS-NHC-31, 27 pp.
- Neumann, C.J., B.R. Jarvinen, A.C. Pike, and J.D. Elms, 1987: Tropical Cyclones of the North Atlantic Ocean, 1871-1986, NOAA, Historical Climatology Series 6-2, 186 pp.
- Saffir, H.S., 1977: Design and Construction Requirements for Hurricane Resistant Construction, American Society of Civil Engineers, New York, Preprint Number 2830, 20 pp.
- Sheets, R.C., 1984: The National Weather Service Hurricane Probability Program, NOAA, Technical Report NWS-37, 70 pp.
- Simpson, R.H., and M.B. Lawrence, 1971: Atlantic Hurricane Frequencies Along the U.S. Coastline, NOAA, Technical Memorandum NWS-SR-58, 14 pp.
- Simpson, R.H., 1971: A Proposed Scale for Ranking Hurricanes by Intensity, Minutes of the Eighth NOAA, NWS, Hurricane Conference, Miami, Fla.
- U.S. Weather Bureau: Monthly Weather Review, 1872-1970 (National Weather Service 1971-1973, and American Meteorological Society 1974-1991).
- U.S. Weather Bureau: Climatological Data and Storm Data, various volumes, various periods, National and State Summaries (National Weather Service 1971-1991).

THIS PAGE LEFT BLANK INTENTIONALLY

- NWS NHC 22 A Tropical Cyclone Data Tape for the North Atlantic Basin, 1886-1983: Contents, Limitations, and Uses. Brian R. Jarvinen, Charles J. Neumann, and Mary A. S. Davis - March 1984
- NWS NHC 23 Frequency and Motion of Western North Pacific Tropical Cyclones. Congyuan Xue and Charles J. Neumann - May 1984 (PB85106466)
- NWS NHC 24 Hurricane Experience Levels of Coastal County Populations - Texas to Maine - June 1984 (PB85111383)
- NWS NHC 25 A Tropical Cyclone Data Tape for the Eastern and Central North Pacific Basins, 1949-1983: Contents, Limitations, and Uses - September 1984 (PB85110054)
- NWS NHC 26 Annual Data and Verification Tabulation Atlantic Tropical Cyclones 1984. Gilbert B. Clark and Robert A. Case, NHC - February 1985
- NWS NHC 27 A Storm Surge Atlas for Corpus Christi, Texas. Brian R. Jarvinen, A. Barry Damiano, and Gloria J. D. Lockett - August 1985
- NWS NHC 28 A Statistical Model for the Prediction of Western North Pacific Tropical Cyclone Motion (WPCLPR). Yiming Xu and Charles J. Neumann, November 1985
- NWS NHC 29 Annual Data and Verification Tabulation - Atlantic Tropical Cyclones 1985. Gilbert B. Clark and Robert A. Case, NHC - March 1986
- NWS NHC 30 A Storm Surge Atlas for the Sabine Lake (Texas/Louisiana) Area. Victor Wiggert and Brian Jarvinen - April 1986
- NWS NHC 31 The Deadliest, Costliest, and Most Intense United States Hurricanes of This Century (and Other Frequently Requested Hurricane Facts)(Rev. of NOAA TM 18). Paul J. Hebert, WSFO Miami; Glenn Taylor and Robert A. Case, NHC - Revised February 1992
- NWS NHC 32 Comparison of Observed Versus SLOSH Model Computed Storm Surge Hydrographs Along the Delaware and New Jersey Shorelines for Hurricane Gloria, September 1985. Brian Jarvinen, NHC; Jeff Gebert, Army COE, Philadelphia, PA - September 1986
- NWS NHC 33 Hurricane Tracking Using an Envelope Approach - Impacts Upon Forecasts. Robert C. Sheets - October 1986
- NWS NHC 34 An Experiment in Statistical Prediction of Tropical Cyclone Intensity Change - Robert T. Merrill - February 1987
- NWS NHC 35 Annual Data and Verification Tabulation - Atlantic Tropical Cyclones 1986 - Gilbert B. Clark and Robert A. Case - March 1987
- NWS NHC 36 Observed Versus SLOSH Model Storm Surge for Connecticut, New York and Upper New Jersey in Hurricane Gloria, September 1985 - Brian Jarvinen and Jeff Gebert - August 1987
- NWS NHC 37 Observed Versus SLOSH Model Storm Surge for North Carolina in Hurricane Gloria, September 1985 - Brian Jarvinen and Allan McDuffie - August 1987
- NWS NHC 38 The National Hurricane Center Risk Analysis Program (HURISK) - Charles J. Neumann - November 1987
- NWS NHC 39 Satellite Interpretation Messages - A Users' Guide - James S. Lynch - May 1987
- NWS NHC 40 Annual Data and Verification Tabulation - Atlantic Tropical Cyclones 1987. Gilbert B. Clark and Robert A. Case - March 1988
- NWS NHC 41 The National Hurricane Center NHC83 Model. - Charles J. Neumann - May 1988
- NWS NHC 42 Annual Data and Verification Tabulation - Atlantic Tropical Cyclones 1988
- NWS NHC 43 Annual Data and Verification Tabulation - Eastern Pacific Tropical Cyclones 1988
- NWS NHC 44 A Revised National Hurricane Center NHC83 Model (NHC90) - Charles J. Neumann (Science Applications International Corporation) and Colin J. McAdie (NHC) - November 1991

## NOAA SCIENTIFIC AND TECHNICAL PUBLICATIONS

NOAA, the *National Oceanic and Atmospheric Administration*, was established as part of the Department of Commerce on October 3, 1970. The mission responsibilities of NOAA are to monitor and predict the state of the solid Earth, the oceans and their living resources, the atmosphere, and the space environment of the Earth, and to assess the socioeconomic impact of natural and technological changes in the environment.

The six Major Line Components of NOAA regularly produce various types of scientific and technical information in the following kinds of publications:

**PROFESSIONAL PAPERS** — Important definitive research results, major techniques, and special investigations.


**TECHNICAL REPORTS** — Journal quality with extensive details, mathematical developments, or data listings.

**TECHNICAL MEMORANDUMS** — Reports of preliminary, partial, or negative research or technology results, interim instructions, and the like.

**CONTRACT AND GRANT REPORTS** — Reports prepared by contractors or grantees under NOAA sponsorship.

**TECHNICAL SERVICE PUBLICATIONS** — These are publications containing data, observations, instructions, etc. A partial listing: Data serials; Prediction and outlook periodicals; Technical manuals, training papers, planning reports, and information serials; and Miscellaneous technical publications.

**ATLAS** — Analysed data generally presented in the form of maps showing distribution of rainfall, chemical and physical conditions of oceans and atmosphere, distribution of fishes and marine mammals, ionospheric conditions, etc.



Information on availability of NOAA publications can be obtained from:

**ENVIRONMENTAL SCIENCE INFORMATION CENTER  
ENVIRONMENTAL DATA SERVICE  
NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION  
U.S. DEPARTMENT OF COMMERCE**

**3300 Whitehaven Street, N.W.  
Washington, D.C. 20235**