



PRELIMINARY GUIDE TO THE IDENTIFICATION OF THE EARLY LIFE HISTORY STAGES OF
CORYPHAENID FISHES OF THE WESTERN CENTRAL ATLANTIC

BY

JAMES G. DITTY



U.S. DEPARTMENT OF COMMERCE
National Oceanic and Atmospheric Administration
National Marine Fisheries Service
Southeast Fisheries Science Center
Galveston Laboratory
4700 Avenue U
Galveston, TX 77551-5997

July 2001



**PRELIMINARY GUIDE TO THE IDENTIFICATION OF THE EARLY LIFE HISTORY STAGES OF
CORYPHAENID FISHES OF THE WESTERN CENTRAL ATLANTIC**

BY

JAMES G. DITTY

**U.S. DEPARTMENT OF COMMERCE
Donald L. Evans, Secretary**

**National Oceanic and Atmospheric Administration
Scott B. Gudes, Acting Under Secretary for Oceans and Atmosphere**

**National Marine Fisheries Service
William T. Hogarth, Acting Assistant Administrator for Fisheries**

July 2001

This Technical Memorandum series is used for documentation and timely communication of preliminary results, interim reports, or similar special-purpose information. Although the memoranda are not subject to complete formal review, editorial control, or detailed editing, they are expected to reflect sound professional work.

NOTICE

The National Marine Fisheries Service (NMFS) does not approve, recommend or endorse any proprietary product or material mentioned in this publication. No reference shall be made to NMFS or to this publication furnished by NMFS, in any advertising or sales promotion which would imply that NMFS approves, recommends, or endorses any proprietary product or proprietary material mentioned herein or which has as its purpose any intent to cause directly or indirectly the advertised product to be used or purchased because of this NMFS publication.

This report should be cited as follows:

Ditty, James G. 2001. Preliminary guide to the identification of the early life history stages of coryphaenid fishes of the western central Atlantic. NOAA Technical Memorandum NMFS-SEFSC-459, 6 p.

W. J. Richards, Editor. NOAA Fisheries, 75 Virginia Beach Drive, Miami, FL

This report will be posted on the Bethune Cookman College NOAA Cooperative web site later in 2001 at URL: <http://208.152.233.21/NOAA/> and will also appear on the SEFSC web site at URL: <http://www.sefsc.noaa.gov/> It will be a chapter entitled Coryphaenidae in the "Guide to the early life history stages of fishes of the western central Atlantic".

Copies may be obtained by writing:

The author at
NOAA Fisheries
4700 Avenue U
Galveston, TX 77551-4551

National Technical Information Center
5825 Port Royal Road
Springfield, VA 22161
(800) 553-6847 or (703) 605-6000
<<http://www.ntis.gov/numbers.htm>>

The dolphinfish family Coryphaenidae contains two cosmopolitan and epipelagic species, common dolphin (*Coryphaena hippurus*) and pompano dolphin (*C. equiselis*). Coryphaenids exhibit sexual dimorphism with mature males characterized by having a prominent bony crest on the front of the head (Palko et al. 1982). The dolphinfishes are elongate and compressed with very long dorsal and anal fin bases and deeply forked tails. Fins lack spines.

Highly prized and widely distributed game fish of both recreational and commercial importance, dolphinfish are found primarily in tropical and subtropical waters of the world (Briggs 1960) though *C. hippurus* also occurs in the Mediterranean Sea (Alemany and Massuti 1998). Two populations of common dolphinfish have been tentatively identified in the western Atlantic, a southern Caribbean and a northern Atlantic Ocean population, each with different life history parameters (Bentivoglio 1989). Dolphinfish grow rapidly but are short-lived (Beardsley 1967). Dolphinfish often associate with floating or drifting objects and are an important member of the *Sargassum* community. Both species spawn from spring through late fall in the northern Gulf of Mexico and possibly year-round in tropical Atlantic (Gibbs and Collette 1959) and southern Gulf waters where water temperatures remain above 24°C.

Lack of pigment along the caudal peduncle and over the caudal finfold separates common dolphin from pompano dolphin <4.5-mm SL. Morphometrics are generally greater in pompano dolphin than in common dolphin with pompano dolphin possessing both a deeper-body and larger eye by 9-mm. Pompano dolphin also has a larger mouth and longer preanal length by 13-mm. Common dolphin possess four spines along the outer margin of the preopercular shelf while pompano dolphin has five, although these spines are often difficult to count on larvae not cleared and stained. Pigmented pelvic fins and bands of pigment laterally on both the body and median fins of common dolphin are diagnostic for separating species >8-mm (Ditty et al. 1994).

Information on larval morphology of pompano dolphin corroborates Johnson's hypothesis of a relationship between Coryphaenidae and Rachycentridae (Johnson 1984) rather than that previously hypothesized between Rachycentridae and Echeineididae. Larvae of dolphinfishes and cobia share similar patterns of head and preopercular spination. Dolphinfishes lack the supracleithral spine found in cobia but possess a small spine on the maxillary that cobia lack. Dolphinfishes also differ from cobia by lacking dorsal and anal spines and having more vertebrae and soft dorsal rays. Echeineids lack head spines and have large hooked teeth anteriorly on the lower jaw (Ditty et al. 1994).

CORYPHAENIDAE

MERISTICS

Vertebrae

Precaudal	13-14
Caudal	19-20
Total	33-34

Number of fin spines and rays

Dorsal (no spines)	55 (52-59)
Anal	26 (23-29)
Pectoral	18-21
Pelvic	I, 5

Caudal

Dorsal Secondary	10-13
Principal	9+8
Ventral Secondary	10-14
Total	41 (37-44)

Gillrakers on first arch

Upper	1-2
Lower	9-10 (8-11)
Total	9-13

(reduce in number with growth)

Branchiostegals	7
-----------------	---

LIFE HISTORY

Range: throughout area

Habitat: epipelagic, shelf and oceanic

ELH pattern: oviparous, pelagic eggs and larvae

Spawning

Season: probably year-round at
water temps >21°C

Area: oceanic waters

Mode: serial spawner

Migration: north-south

Age at first maturity: 3-4 months

Literature: Gibbs and Collette 1959; Collette
et al. 1969; Potthoff 1971; 1980; Hagood
& Rothwell 1979; Johnson 1984;
Ditty et al. 1994

Coryphaena equiselis (Linnaeus, 1758)

EARLY LIFE HISTORY DESCRIPTION

EGGS:

Diameter: mean: 1.35 mm

No. of Oil Globules:

Oil Globule Diameter:

Yolk:

Shell:

Hatch Size:

Incubation: 38 hrs at 26°C

Pigment: oil globule and embryo

LARVAE:

Length at flexion: 7.5-9.0 mm SL


Length at transformation: 25-30 mm SL

Sequence of fin development: C, A, D, P₂, P₁

Pigment: heavy

Diagnostic: from *C. hippurus* <4.5 mm SL by
pigment in caudal finfold and number of spines
along outer shelf of preopercle; >8 mm SL
by lack of pigment on pelvic fins and
absence of lateral bands; from *Rachycentron*
by number of soft dorsal rays

ILLUSTRATIONS: After Ditty et al. 1994. A: 4.7
mm NL, B: 5.5 mm NL, C: 7.5 mm SL, D: 9.7
mm SL, E: 11.5 mm SL, F: 15.0 mm SL.



CORYPHAENIDAE

MERISTICS

Vertebrae

Precaudal	13-14
Caudal	17-18
Total	30-31

Number of fin spines and rays

Dorsal (no spines)	61 (58-66)
Anal	28 (25-31)
Pectoral	18-21
Pelvic	I, 5
Caudal	
Dorsal Secondary	10-14
Principal	9+8
Ventral Secondary	11-14
Total	41 (37-45)

Gillrakers on first arch

Upper	0-2
Lower	7-10
Total	8-12

(reduce in number with growth)

Branchiostegals	7
-----------------	---

LIFE HISTORY

Range: throughout area

Habitat: epipelagic, shelf and oceanic

ELH pattern: oviparous, pelagic eggs and larvae

Spawning

Season: probably year-round at water temps >21°C

Area: oceanic waters

Mode: serial spawner

Migration: north-south

Age at first maturity: 3-4 months in Gulf of Mexico, 6-7 months in N. E. Atlantic

Longevity: 4 years but usually <2 years

Literature: Gibbs and Collette 1959; Collette et al. 1969; Hassler & Rainville 1975; Potthoff 1980; Hagood & Rothwell 1979; Bentivoglio 1989; Ditty et al. 1994

Coryphaena hippurus (Linnaeus, 1758)

EARLY LIFE HISTORY DESCRIPTION

EGGS:

Diameter: mean: 1.3 mm

No. of Oil Globules: one

Oil Globule Diameter: 0.3-0.4 mm

Yolk: segmented

Shell: Smooth, clear

Hatch Size: 3.0 mm TL

Incubation: 38 hrs at 25°C; 26 hrs at 30°C

Pigment: oil globule and embryo

LARVAE:

Length at flexion: 7.5-9.0 mm SL

Length at transformation: 25-30 mm SL

Sequence of fin development: C, A, D, P₂, P₁,

Pigment: heavy

Diagnostic: from *C. equiselis* <4.5 mm SL by

lack of pigment along peduncle and in

caudal finfold and number of spines along

outer shelf of preopercle; >8 mm SL by

pigment on pelvic fins and presence of lateral

bands; from *Rachycentron* by number of soft

dorsal rays

Illustrations: Ditty et al. 1994


A: 3.5 mm NL, B: 5.0 mm NL, C: 7.1 mm SL,

D: 9.5 mm SL, E: 11.0 mm SL, F: 14.0 mm

SL.

Coryphaena hippurus

CORYPHAENIDAE



Literature Cited

- Aleman, F. & E. Massuti. 1998. First record of larval stages of *Coryphaena hippurus* (Pisces: Coryphaenidae) in the Mediterranean Sea. *Scientia. Mar.* 62(1-2): 181-184.
- Beardsley, G. L. 1967. Age, growth, and reproduction of the dolphin, *Coryphaena hippurus*, in the straits of Florida. *Copeia* 1967: 441-451.
- Bentivoglio, A. A. 1989. Investigations into the growth, maturity, mortality rates and occurrence of the dolphin (*Coryphaena hippurus*, Linnaeus) in the Gulf of Mexico. Ph. D. dissertation, Univ. College of North Wales, Bangor, U. K., 37 p.
- Briggs, J. C. 1960. Fishes of worldwide (circumtropical) distribution. *Copeia* 1960: 171-180.
- Collette, B. B., R. H. Gibbs, & G. E. Clipper. 1969. Vertebral numbers and identification of the two species of dolphin (*Coryphaena*). *Copeia* 1969: 630-631.
- Ditty, J. G., R. F. Shaw, C. B. Grimes, & J. S. Cope. 1994. Larval development, distribution, and abundance of common dolphin, *Coryphaena hippurus*, and pompano dolphin, *C. equiselis* (family: Coryphaenidae), in the northern Gulf of Mexico. *Fish. Bull. U. S.* 92: 275-291.
- Gibbs, R. H., Jr., & B. B. Collette. 1959. On the identification, distribution, and biology of the dolphins, *Coryphaena hippurus* and *C. equiselis*. *Bull. Mar. Sci.* 9: 117-152.
- Hagood, R. W., & G. N. Rothwell. 1979. Sea Grant interim project report - 1979. Aquaculture in tropical ocean - *Coryphaena* sp. Oceanic Inst., Makapuu Point, Waimanalo, HI 96795.
- Hassler, W. W., & R. P. Rainville. 1975. Techniques for hatching and rearing dolphin, *Coryphaena hippurus*, through larval and juvenile stages. Univ. N. Carolina Sea Grant Prog., UNC-SG-75-31, 17 p.
- Johnson, G. D. 1984. Percoidei: Development and relationships. Pages 464-498 in H. G. Moser, W. J. Richards, D. M. Cohen, M. P. Fahay, A. W. Kendall, Jr., and S. L. Richardson (eds.), Ontogeny and systematics of fishes, Amer. Soc. Ichthy. Herp., Spec. Publ. (1): 760 p.
- Palko, B. J., G. L. Beardsley, & W. J. Richards. 1982. Synopsis of the biological data on dolphin-fishes, *Coryphaena hippurus* Linnaeus and *Coryphaena equiselis* Linnaeus. NOAA Tech. Rep. NMFS Circ. 443, 28 p.
- Potthoff, T. 1971. Observations on two species of dolphin (*Coryphaena*) from the tropical mid-Atlantic. *Fish. Bull. U. S.* 69: 877-879.
- 1980. Development and structure of fins and fin supports in dolphin fishes *Coryphaena hippurus* and *Coryphaena equiselis* (Coryphaenidae). *Fish. Bull. U. S.* 78(2): 277-312.