

The Use of
Pb-210, Th-234 and Cs-137
as Tracers of Sedimentary Processes
in San Francisco Bay, California

CIRCULATED COPY
Sea Grant Collection

by

Christopher Channing Fuller

USCSG-TD-01-82

SEA GRANT
Dissertation Series

UNIVERSITY OF SOUTHERN CALIFORNIA
Institute for Marine and Coastal Studies
University Park, Los Angeles, CA 90007



The Use of
Pb-210, Th-234 and Cs-137
as Tracers of Sedimentary Processes
in San Francisco Bay, California

by
Christopher Channing Fuller

USCSG-TD-01-82

This work is the result of research sponsored by the National Oceanic and Atmospheric Administration (NOAA) of the Department of Commerce through a grant to the University of Southern California Sea Grant Program (NOAA Grant # NA 79-AA-D-0133).

All rights reserved. No part of this publication may be reproduced or transmitted in any form or by any means without permission in writing from the publisher. The U.S. Government, however, is authorized to produce and distribute reprints for governmental purposes.

Published and distributed by:

USC Sea Grant Institutional Program
Institute for Marine and Coastal Studies
University of Southern California
University Park
Los Angeles, CA 90089-0341

10/82

THE USE OF Pb-210, Th-234 AND Cs-137
AS TRACERS OF SEDIMENTARY PROCESSES
IN SAN FRANCISCO BAY, CALIFORNIA

by

Christopher Channing Fuller

A Thesis Presented to the
FACULTY OF THE GRADUATE SCHOOL
UNIVERSITY OF SOUTHERN CALIFORNIA

In Partial Fulfillment of the
Requirements for the Degree

MASTER OF SCIENCE
(Geological Sciences)

December, 1982

UNIVERSITY OF SOUTHERN CALIFORNIA
THE GRADUATE SCHOOL
UNIVERSITY PARK
LOS ANGELES, CALIFORNIA 90007

This thesis, written by

..... Christopher Channing Fuller

*under the direction of his Thesis Committee,
and approved by all its members, has been pre-
sented to and accepted by the Dean of The
Graduate School, in partial fulfillment of the
requirements for the degree of*

..... Master of Science

B. G. O'Neil

.....
Dean

Date..... SEPTEMBER 8, 1982

THESIS COMMITTEE

Douglas E. Hammond

.....
Chairman

W. J. Stinson

Al. Lee

ACKNOWLEDGMENTS

This thesis was made possible by the efforts of many people. Funding for the study was provided by the Sea Grant Program. (NOAA Grant # NA 79AA-D-0133), which is gratefully acknowledged. Many individuals have contributed support and guidance leading up to and during this work. It is impossible to acknowledge everyone, so many thanks to those whom are not mentioned.

First, to my family, who provided guidance and support throughout my academic career, despite my being on the other side of the continent for this part.

I am very grateful to the many people in the USC Geological Sciences department who have aided my efforts in this project. Will Benson, Joe Donoghue, Ethan Grossman, Blayne Hartman, Chi-An Huh, Masashi Kusakabe, Melody Lin, Larry Miller and Steve Warren provided valuable assistance in the field and laboratory, useful discussions, and help in maintaining the proper perspective for this study. Additional thanks to Blayne and Will for their patient tutoring in computer programming.

Special thanks to Dr. Richard Ku for his generosity in providing alpha and beta counter time and the thorium spike, as well as for valuable discussions throughout this project. To Dr. Donn Gorsline, through his lectures, who taught me the value of considering systems as a whole.

This study would not have been possible without ship time and facilities provided by Drs. John Conomos and Dave Peterson of the U. S. Geological Survey. Thanks to Dr. Brian Atwater of the U.S.G.S.

for valuable discussions and assistance in selecting a suitable site in the Bay's salt marshes for use as a natural fallout collector. Dr. Fred Nichols and Sally Wenke assisted in coring and sample preparation. The crew of the RV Polaris is credited for their skillful seamanship and memorable shipboard meals. Dana Harmon and Becky Rea braved great heights in providing invaluable assistance in the fallout collection program.

Drs. Jim Simpson and Curt Olsen are acknowledged for their assistance in setting up and calibrating the Cs-137 counting system and for their participation in the coring expeditions.

Thanks to Gloria Lee, Sue Turnbow and Sharon Wallace for the many hours of typing. Joceline Boucher drafted many of the figures.

Finally, to one swell guy, Doug Hammond, to whom I am very grateful for the years of employment, tutoring, patience, guidance and most of all, friendship. Without his association, I would not have learned the true value of data "good to within 5%."

DEDICATION

This thesis is dedicated to the memory of my father, John Channing Fuller, whose wisdom and love of nature inspired me to pursue studies of the environment.

TABLE OF CONTENTS

	Page
ACKNOWLEDGEMENTS.	ii
DEDICATION.	iv
LIST OF ILLUSTRATIONS	vii
LIST OF TABLES.	x
ABSTRACT.	1
CHAPTER 1. INTRODUCTION	6
General Statement.	6
Isotopes Used.	7
Environmental Setting.	10
Outline of Field and Laboratory Work	15
Outline of Thesis.	18
CHAPTER 2. URANIUM AND THORIUM.	19
Uranium in the Water Column.	19
Th-234 in the Water Column	23
Suspended Particle Residence Times	33
Th-234 in Sediments.	42
CHAPTER 3. Pb-210 AND Po-210.	57
Introduction	57
Chemistry of Lead in Natural Waters.	58
Water Column Results and Discussion.	59
Sediment Results and Discussion.	78
Atmospheric Fallout of Pb-210.	101
CHAPTER 4. Cs-137	123
Introduction	123

	Page
Geochemistry of Cesium and Cs-137.	126
Cs-137 Sediment Results and Discussion	131
CHAPTER 5. Pb-210 SEDIMENT PROFILE NUMERICAL MODELLING. . . .	144
Introduction	144
Model Description.	146
Model Results and Implications	150
CHAPTER 6. Pb-210, Cs-137 AND SEDIMENT BUDGETS AND IMPLICATIONS	
TOWARDS BAY-WIDE SEDIMENTATION	163
Introduction	163
Bay Wide Integrated Excess Pb-210 Activities	164
Sources of Pb-210 to the San Francisco Bay System. . . .	165
Sediment input to the Bay.	170
Bay Wide Cs-137 Inventories.	176
CHAPTER 7. SUMMARY AND CONCLUSIONS.	182
REFERENCES.	191
APPENDIX I. SAMPLE COLLECTION AND ANALYSIS METHODS.	198
APPENDIX II. SEDIMENT DATA.	237
APPENDIX III. NUMERICAL MODEL PROGRAM	248

LIST OF ILLUSTRATIONS

Figure	Page
1. Location Map.	8
2. South and Central San Francisco Bays station locations	16
3. San Pablo and Central Bays station locations.	17
4. U-238 (dpm/l) versus salinity ($^{\circ}/\text{‰}$).	20
5. Dissolved Th-234 to U-238 activity ratio versus Th-234 excess activity on suspended particles	29
6. Particle residence time (τ_p) versus suspended particle concentration.	36
7. Th-234 sediment activity profiles	45
8. Th-234 sediment activity profiles	46
9. Th-234 sediment activity profiles	47
10. Th-234 sediment activity profiles	48
11. Dissolved Pb-210 and Po-210 versus suspended sediment concentration (E).	62
12. Dissolved Po-210 activity (dpm/l) versus suspended particle excess Po-210 activity (dpm/gm).	65
13. Dissolved Pb-210 activity (dpm/gm) versus suspended particle excess Pb-210 activity (dpm/gm).	66
14. Total Pb-210 activity (dpm/l) versus suspended sediment concentration (mg/l)	70
15. Total Pb-210 activity (dpm/l) versus suspended concentration (mg/l).	71
16. Pb-210 and Cs-137 activity profiles	80
17. Pb-210 and Cs-137 activity profiles	81
18. Pb-210 and Cs-137 activity profiles	82
19. Pb-210 and Cs-137 activity profiles	83
20. Pb-210 and Cs-137 activity profiles	84

Figure	Page
21. Pb-210 and Cs-137 activity profiles	85
22. Pb-210 and Cs-137 activity profiles	86
23. Station 27 Pb-210 activity profiles	89
24. Pb-210 activity in rain (dpm/l) versus centimeter of precipitation for Los Angeles fallout samples. . . .	111
25. Pb-210 atmospheric deposition rate versus precipitation	112
26. Newark Slough Marsh Cs-137 sediment activity (pCi/gm) versus depth in sediment (cm).	118
27. Newark Slough Marsh Pb-210 sediment activity (dpm/gm) versus depth in sediment (cm).	119
28. Yearly Cs-137 fallout delivery to the San Francisco Bay area.	124
29. Cs-137 activity profiles.	132
30. Cs-137 activity profiles.	133
31. Cs-137 activity profiles.	134
32. Cs-137 activity profiles.	135
33. Cs-137 activity profiles.	136
34. Cs-137 activity profiles.	137
35. Numerical model generated Pb-210 profiles for Station 28C	152
36. Numerical model generated Pb-210 profiles for Station 28C	153
37. Numerical model generated Pb-210 profiles for Station 28C	154
38. Numerical model generated Pb-210 profiles for Station 27.	156
39. Numerical model generated Pb-210 profiles for Station 27.	158
40. Numerical model generated Pb-210 profiles for Station 27.	159

Figure	Page
41. Numerical model generated Pb-210 profiles for Station 27.	160
42. Numerical model generated Pb-210 profiles for Station 27.	161
43. Numerical model generated Pb-210 profiles for Station 13.13S.	162
44. Pressure filtration system schematic.	207
45. Correction method for overlap of alpha peaks.	218
46. Beta count rate versus absorber density	221
47. Beta count rate for Th-228 spike.	228

LIST OF TABLES

Table	Page
1. Uranium water column data	21
2. Th-234 water column data summary.	24
3. Dissolved Th-234 data	28
4. Comparison of particulate activities for Th-234, Pb-210, and Po-210 (dpm/gm total) from filter and total samples	31
5. Th-234 model B results.	39
6. Th-234 sediment data.	44
7. Comparison of integrated Th-234 activity in sediments ($\bar{A}_{Th}$) to production rate in overlying water column ($\bar{A}_{wc}$)	52
8. Summary of sediment reworking rates from Th-234 sediment profiles	55
9. Dissolved Pb-210 and Po-210 water column activity summary	61
10. Pb-210 and Po-210 water column data	73
11. Pb-210 and Po-210 suspended particle activities	76
12. Pb-210 and Cs-137 sediment data summary	91
13. Calculated sediment accumulation rates (w) from Pb-210 profiles (gm/cm ² yr).	95
14. San Francisco Bay area Pb-210 fallout data summary. . .	106
15. Los Angeles area Pb-210 fallout data summary.	108
16. Newark Slough Marsh sediment data summary	117
17. Cs-137 fallout data	127
18. Numerical model equations	147
19. Bay-wide integrated Pb-210 excess activities for South and San Pablo Bays with comparison to direct atmospheric fallout	166
20. Pb-210 budget for South Bay sediments	167

Table	Page
21. Sediment accumulation in San Francisco Bay from Pb-210 balance	174
22. Bay-wide Cs-137 inventories and comparison to direct fallout supported activity	178
23. Porosity and sediment density.	201
24. Radiochemical procedures for Th-234, U-238, Pb-210 and Po-210 for sediment samples and water sample precipitates	202
25. Pb-210 activity as a function of grain size	210
26. Comparison of Pb-210 sediment activities for HCl and HF leaching methods.	212
27. Th-228/Th-232 activity ratios in unspiked sediment and water column samples	214
28. Equation for Th-234 data reduction	223
29. Uranium precipitation yield (Y).	226
30. Th-234 blank analyses summary.	227
31. Alpha and Beta detector intercalibration: detector efficiency ratios.	230
32. Pb-210 and Po-210 blank analyses summary	232
33. Statistical functions from Barford (1967).	236
34. Summary of core descriptions - visual and X-radiograph	238
35. Pb-210 sediment data summary	241
36. Cs-137 sediment data summary	244
37. Numerical model program.	249

ABSTRACT

The partitioning of Pb-210, Po-210, and Th-234 between the dissolved ($<0.4\mu$) and solid phases in the water column of San Francisco Bay is presented. Profiles of Pb-210, Th-234, and Cs-137 in sediments are also presented. These data were used to determine residence times and resuspension rates of suspended sediment particles, sediment accumulation and mixing rates, and to further the knowledge of the geochemical behavior of these isotopes in estuarine environments.

An uptake rate constant for the scavenging of Th-234 from solution onto particles of $1.5 \pm 0.6 \text{ day}^{-1}$ is calculated from the average dissolved Th-234 activity, $.04 \pm .01 \text{ dpm/l}$, and its production rate by the decay of U-238. It is unclear whether this removal is due to reversible or irreversible adsorption. Residence times of suspended particles in surface waters ranging from 1 to 21 days are calculated from the mass balance of Th-234 in the water column and average 6.8 ± 4.5 days for the South Bay and 11 ± 10 days for San Pablo Bay. If the input of excess Th-234 by resuspension of bottom sediments is included, particle residence times of 3.5 ± 2.2 days for the South Bay and 5 ± 3 days for San Pablo Bay are calculated. Water column samples from near the bottom yield longer residence times, which probably reflect near bottom transport of particles from shallow to deep regions. Resuspension rates of $1.1 \pm 1.4 \text{ gm/cm}^2\text{yr}$ for deep areas of South Bay, 0.7 ± 1.0 for shallow areas of South Bay, and 0.7 ± 1.0 for the deep areas of San Pablo Bay are calculated from the distribution of Th-234 in the water column and sediment. Comparison

of these rates to accumulation rates suggest sediments are resuspended at least 2 to 5 times before final burial in the deep areas of South Bay.

Surface sediment mixing rate constants calculated from Th-234 sediment activities range from 1 to 10×10^{-6} cm²/sec at deep stations of the South Bay. The shallow reaches yield rate constants of less than 1×10^{-7} cm²/sec. Live polychaetes, observed to 60 cm in cores from the deep area, may be responsible for much of the sediment reworking. In San Pablo Bay sediments, where live benthic macrofauna were not observed, Th-234 indicates little or no sediment mixing. Integrated excess Th-234 activities in sediments at the deep stations of the South Bay have a 2 to 18-fold greater activity than can be supported by the decay of U-238 in the overlying water column. This suggests that these areas accumulate Th-234 and sediments from the adjacent shoal areas.

An inverse relationship between dissolved and suspended particle Po-210 activities suggests an irreversible scavenging of Po-210 from solution. Pb-210, however, shows an opposite trend, which is modelled as an equilibrium partitioning. Overall distribution coefficients of $1.3 \pm 1 \times 10^5$ cm³/gm for high salinities ($>24\text{‰}$) and $2.8 \pm 3 \times 10^5$ for low salinities ($<18\text{‰}$) are calculated and are used for determining suspended particle excess activities from total (unfiltered) water samples. A seasonal variation in excess Pb-210 activity on suspended particles ranging from 1.9 dpm/gm during low fresh water inflows and 3.8 dpm/gm during high flows is observed. The elevated activities during high flows

are similar to those measured in river water above the null zone of the estuary. Po-210 and Pb-210 activity ratios in total water samples range from 0.6 ± 1 to 1.2 ± 1 in both bays. The lower values measured during high fresh water inflows probably reflect the atmospheric source of excess Pb-210. Suspended particle activity ratios are generally higher than total samples indicating a preferential removal of Po-210 over Pb-210 from solution.

Excess Pb-210 in the sediments of the deep areas of South and San Pablo Bays extends much farther down core (up to 130 cm) than in the shallow reaches (10 to 20 cm). Upper limits of recent sediment accumulation rates are estimated from the integrated excess Pb-210 in the sediment column and the excess Pb-210 activity of suspended particles. These range from 0.03 to 0.10 gm/cm²yr in the shoals and 0.13 to 0.53 gm/cm²yr in the deep areas of the South Bay., Rates in San Pablo Bay range from 0.06 to 0.11 gm/cm²yr in the shoals and 0.9 to 1.0 gm/cm²yr in the deeper areas. However, numerical modelling of Pb-210 profiles using mixing and resuspension rates derived from Th-234 data shows that the combination of deposition, resuspension, and mixing can produce profiles observed in the shallow areas. In the deep areas modelling shows that some net deposition is required. These results indicate that an influx of new sediment particles to the sediment column can be maintained without net deposition and that the accumulation rates based on a Pb-210 mass balance are upper limits.

An atmospheric Pb-210 flux of 0.15 dpm/cm²yr for the San Francisco Bay area was measured with artificial collectors. This

rate agrees well with the rate of 0.12 determined by integrating the Pb-210 excess in a salt marsh core. A flux of 0.23 dpm/cm²yr was measured in the Los Angeles area with collectors. The higher flux in Los Angeles is probably due to a two fold greater rainfall during the collection period than in the Bay area. Both fluxes are significantly lower than Turekian et al's (1979) global model predicts. Bay-wide sediment and Pb-210 budgets suggest that the measured flux is more realistic than the model predicted flux.

The depth of Cs-137 activity in sediments and their integrated activities show geographic variability similar to that observed for Pb-210. An enhanced penetration of Cs-137 to equal or greater depths than excess Pb-210 is observed in cores with abundant active polychaete burrows. This may be the result of incigation and/or radial diffusion out of the burrows. Cs-137 activity profiles show no distinct maxima that can be correlated with temporal variations in fallout delivery. Their absence is probably due to sediment mixing and the temporal input of Cs-137 bearing particles from the drainage basin. The inventory of Cs-137 in the South Bay is 60% of the amount supplied by direct fallout. In San Pablo Bay, the inventory is twice the amount delivered by fallout. In the Central Bay, the inventory is equal to the activity delivered by direct fallout. Since a significant fraction of the fallout Cs-137 is flushed out of the bay in the dissolved phase, a large input of Cs-137 derived from the drainage basin is required to support the observed activities.

Bay-wide Pb-210 inventories indicate that direct fallout supplies 55% of the excess Pb-210 in South Bay sediment, 30% in San

Pablo Bay and 40% in the Central Bay. Assuming that the input of suspended sediment from local rivers and streams (with an excess Pb-210 activity equal to that measured in river water) is the only other source of Pb-210, $1.0 \pm .4 \times 10^{11}$ gm/yr of sediment must be deposited in the South Bay to maintain the observed Pb-210 activity. This is 33% of the sediment supplied by local streams. For San Pablo Bay, $2.2 \pm .3 \times 10^{11}$ gm/yr of sediment from local streams and rivers and from delta sources is required. This input accounts for 85% of the sediment delivered by the local sources. The Central Bay requires $0.8 \pm .3 \times 10^{11}$ gm/yr of sediment from local stream and delta sources. These required inputs result in overall accumulation rates of 0.03 gm/cm²yr for the South bay, 0.10 gm/cm²yr for muddy areas of the Central Bay and 0.11 gm/cm²yr in San Pablo Bay. Earlier sediment budgets, which suggested net erosion in the South Bay based on changes in bathymetry, may be in error because they fail to account for recent sea level rise and local subsidence.

